# Ergodic properties of one-dimensional incommensurate bilayer materials

Nathan J. Essner, Alexander B. Watson

March 12, 2024

#### Abstract

We review ergodic properties of one-dimensional incommensurate bilayer materials, especially the convergence of the density of states in the large-volume limit, from the perspective of the theory of ergodic Schrödinger operators. More precisely, we first provide a short introduction to ergodic Schrödinger operators as a unifying concept in spectral theory at a level accessible for nonspecialists. We then present two natural tight-binding models of incommensurate bilayer materials in one dimension and prove convergence of their density of states measures in the large volume limit using ideas from the theory of ergodic Schrödinger operators.

# Contents

1	Introduction	1
	1.1 Motivation	1
	1.2 Structure of work and description of results	2
	1.3 Related work and perspectives	3
<b>2</b>	Ergodic operators: basic concepts	3
	2.1 Definition of an ergodic operator	3
	2.2 Examples of ergodic Schrödinger operators	4
	2.2.1 The almost-Mathieu operator	4
	2.2.2 The Anderson model	5
	2.3 Existence of density of states measure for ergodic Schrödinger operators	5
3	Incommensurate bilayer materials: 1D tight-binding models	9
	3.1 Model of incommensurate coupled chains	9
	3.2 Reduced model of incommensurate coupled chains	9
4	Incommensurate bilayer materials: ergodic properties	10
	4.1 Reduced incommensurate coupled chain model as an ergodic operator	10
	4.2 Convergence of the density of states measure for the incommensurate coupled chain model	11
A	Proofs for Section 2	17

# 1 Introduction

## 1.1 Motivation

The field of two-dimensional materials began in 2005 with the isolation and characterization of graphene, a single layer of carbon atoms [24]. Since then, many other two-dimensional materials have been isolated, such as hexagonal Boron Nitride (hBN), and the transition metal dichalcogenides (TMDCs). The family of

two-dimensional materials displays a rich variety of material properties. For example, while graphene is a semimetal (gapless at the Fermi level, but with a small Fermi surface), hBN is an insulator (large band gap), and the TMDCs are generally semiconductors (small band gap) [15].

In recent years, attention has shifted to stackings of such materials with a relative twist. For general twist angles, such materials may be aperiodic at the atomic scale because of incommensurability of the layer Bravais lattices. However, for relatively small twist angles, many properties of such materials can be captured by effective periodic models over the lattice of interlayer disregistry oscillation known as the moiré pattern. For this reason, such materials are known as moiré materials.

Moiré materials have especially attracted attention since the observation of correlated insulating and superconducting electronic phases in twisted bilayer graphene twisted to the "magic angle"  $\approx 1^{\circ}$  [8,9] in 2018. This discovery was anticipated in 2011 with the observation that the Floquet-Bloch "moiré bands" of twisted bilayer graphene become nearly flat at this twist angle [6]. Recent years have seen many other quantum many-body electronic phases realized in moiré materials [18].

The present work is concerned with fundamental atomic-scale models of moiré materials. More precisely, we are concerned with the case where the individual layer Bravais lattices are incommensurate, so that the model has no exact periodic cell. For example, we could consider twisted bilayer graphene at an irrational relative twist angle [17, 22].

Models involving incommensurate periodicities have long been known to display rich spectral properties. Perhaps the canonical model of this type is the almost-Mathieu operator (also known as Harper's operator); see, e.g., [27]. This operator arises in the study of the quantum Hall effect, where a two-dimensional material is subjected to a perpendicular magnetic field [4]. The model depends on two independent lengthscales: the magnetic length (proportional to the strength of the magnetic field), and the material's lattice constant. The study of this operator when the ratio of these lengthscales is irrational has inspired many deep mathematical works; see, e.g., [3].

The conceptual similarities between atomic-scale models of incommensurate moiré materials and the almost-Mathieu operator suggest that the theory developed for understanding the almost-Mathieu operator may have applications in the study of incommensurate moiré materials. In fact, such a link is already known: both models have an ergodic structure which allows, for example, for proof of convergence of the density of states measure in the large-volume limit [7,22].

The goal of this work is to further clarify this connection by, first, reviewing the general concept of an ergodic Schrödinger operator [1, 14], and, then, establishing the sense in which models of incommensurate bilayer materials share this structure. This structure allows, for example, for a very straightforward proof of convergence of the density of states measure in the thermodynamic limit [1, 14]. We hope that this contribution will stimulate further interactions between the mathematical theories of ergodic Schrödinger operators and of incommensurate bilayer materials.

For the sake of clarity, we restrict attention in this work to one-dimensional models, where the same effect as an interlayer twist can be obtained by shifting the lattice constants of each layer with respect to each other. Our results do not depend on this restriction and would readily generalize to two-dimensional models, such as the tight-binding models of twisted bilayer graphene considered in [7,22].

## **1.2** Structure of work and description of results

The structure of this work, and a brief description of our results, are as follows. In Section 2, we review the basic concepts of ergodic Schrödinger operators for the benefit of nonspecialists. We provide the rigorous definition of such operators, provide examples of such operators, and then show how the properties of an ergodic Schrödinger operator allow for a straightforward proof of convergence of the density of states measure.

In Section 3, we describe a typical tight-binding model of an incommensurate moiré material specialized to one dimension. In such a case, incommensurability of the layers is guaranteed by choosing the ratio of layer lattice constants to be an irrational number. We refer to this model as the incommensurate coupled chain model. We then introduce a simplification of this model where electrons in a single layer can "hop" along the layer via interaction with a fictitious second layer with an irrationally-related lattice constant. We term this model the reduced incommensurate coupled chain model.

In Section 4, we investigate the ergodic properties of these models. First, we prove that the reduced incommensurate coupled chain model is an ergodic Schrödinger operator in sense of the definition provided

in Section 2. Then, we clarify the ergodic structure of the full incommensurate coupled chain model (which does not satisfy the definition of an ergodic Schrödinger operator) to the point that we can prove existence of the density of states measure by a similar argument as in the case of an ergodic Schrödinger operator.

#### **1.3** Related work and perspectives

We emphasize that we are not the first to recognize the ergodic structure of incommensurate bilayer materials, or even to prove convergence of the density of states of such materials in the large-volume limit; see [7, 22]. The work [7], in particular, considered the ergodic properties of the same coupled chain model we consider here, although we believe that the *reduced* coupled chain model we also consider is original to the present work. Other works which exploit ergodicity to obtain efficient numerical methods for computing properties of moiré materials include [10-12, 19-21].

Nevertheless, we hope that the particular formulation of these ergodic properties in the present work will be useful in connecting the mathematical theories of incommensurate bilayer materials and of ergodic Schrödinger operators. Such a connection should facilitate, for example, the study of localization properties of the models considered here. Such results have already been obtained for a model of bilayer graphene where one layer is strained with respect to the other, creating an effectively one-dimensional moiré pattern; see [5, 28].

## Acknowledgements

This work began as an independent study project by NJE during Summer and Fall 2023 supervised by ABW. ABW's research was supported in part by DMREF Award No. 1922165 through the National Science Foundation. ABW would like to thank Daniel Massatt, Eric Cancès and Simon Becker for stimulating conversations.

## 2 Ergodic operators: basic concepts

#### 2.1 Definition of an ergodic operator

In this section, we recall the theory and terminology necessary to provide the rigorous definition of an ergodic operator following [1].

First, recall that matrix elements of real-valued and measurable functions f of self-adjoint operators H acting in a Hilbert space  $\mathcal{H}$  can be conveniently defined through the spectral measure

$$\langle \varphi, f(H)\psi \rangle := \int f(\lambda) \,\mathrm{d}\mu_{\varphi,\psi},$$
(2.1)

where  $\mu_{\varphi,\psi}$  is the spectral measure of H associated to the vectors  $\varphi, \psi \in \mathcal{H}$ , and  $\langle, \rangle$  denotes the  $\mathcal{H}$ -inner product [1]. We then say that a function  $\omega \mapsto H(\omega)$  mapping a probability space  $(\Omega, \mathcal{A}, \mathbb{P})$  to the set of self-adjoint operators on  $\mathcal{H}$  is *weakly measurable* if, for all  $f \in L^{\infty}(\mathbb{R})$  and all  $\varphi, \psi \in \mathcal{H}$ , the functions  $\omega \mapsto \langle \varphi, f(H(\omega))\psi \rangle$  are  $\mathbb{P}$ -measurable [1]. We can now define a random operator.

**Definition 2.1.** A self-adjoint random operator is an operator-valued weakly measurable function defined on a probability space  $(\Omega, \mathcal{A}, \mathbb{P})$  which assigns to every  $\omega \in \Omega$  a self-adjoint operator  $H(\omega)$  acting in some common (separable) Hilbert space  $\mathcal{H}$ .

Recall that a transformation  $T: X \to X$  on a measure space  $(X, \mathcal{A}, \mu)$  is called *measure-preserving* if for all  $A \in \mathcal{A}$ ,  $\mu(T^{-1}A) = \mu(A)$ . We can now define an ergodic group action.

**Definition 2.2.** The action of a group of measure-preserving transformations  $(T_x)_{x \in I}$  on a probability space  $(\Omega, \mathcal{A}, \mathbb{P})$  is *ergodic* if all events  $A \in \mathcal{A}$  which are invariant under the group,  $T_x^{-1}A = A$  for all  $x \in I$ , are of probability zero or one.

Recall that a graph is a pair  $\mathbb{G} = (V, E)$ , where V is the set of vertices, and E is a set of pairs of vertices, known as the set of edges. A graph automorphism is a permutation  $\sigma$  of the graph's vertices such that, for all vertices  $v_1, v_2 \in V$ , if  $(v_1, v_2) \in E$  then  $(\sigma(v_1), \sigma(v_2)) \in E$ . A graph  $\mathbb{G}$  is called *vertex-transitive* if, for any vertices  $v_1, v_2 \in V$ , there exists a graph automorphism  $f : \mathbb{G} \to \mathbb{G}$  such that  $f(v_1) = v_2$ . Below, we will abuse notation to write  $\mathbb{G}$  both for the graph and for the set of vertices of  $\mathbb{G}$ .

We can now define an ergodic operator.

**Definition 2.3.** From [1], an *ergodic operator* is a random operator  $H(\omega)$  such that

- 1. The operators act in  $\mathcal{H} = \ell^2(\mathbb{G})$  where  $\mathbb{G}$  is a vertex set of a graph endowed with a vertex-transitive group of graph automorphisms  $\mathcal{S} = (S_x)_{x \in I}$ .
- 2. The group S can be represented by a group of measure-preserving transformations  $(T_x)_{x\in I}$  whose action on a probability space  $(\Omega, \mathcal{A}, \mathbb{P})$  is ergodic. Moreover, for every  $x \in I$  and  $\omega \in \Omega$ ,

$$H(T_x\omega) = U_{x,\omega}H(\omega)U_{x,\omega}^{\dagger}$$
(2.2)

with a unitary operator  $U_{x,\omega}$  for which all  $\psi \in \ell^2(\mathbb{G})$ 

$$(U_{x,\omega}\psi)(\xi) = \psi(S_x\xi)e^{i\phi_{x,\omega}(\xi)}$$
(2.3)

having some real-valued phase  $\phi_{x,\omega}(\xi)$ .

In what follows, we will discuss various examples of ergodic operators and of operators with similar properties. We will standardize notation so that, in each case, the relevant graph automorphisms will be denoted by S, the transformations of the operator's parameter space by T, and the unitary operators transforming the random operator by U.

**Remark 2.4.** The above definition of an ergodic operator is technically a "standard ergodic operator" as in [1]. Meanwhile, an "ergodic operator" is a random operator without the additional structure of each lattice shift corresponding to each transformation of the probability space. The focus of study will not be on this weaker definition of "ergodic operators." Although, despite the weaker condition, the ergodic random operators in [1] still have some of the desirable properties that relate to the discussion below.

### 2.2 Examples of ergodic Schrödinger operators

In this section, we give two examples of commonly studied ergodic operators, the almost-Mathieu operator and the Anderson model. For simplicity, we consider the Anderson model only in dimension one.

Both of these models act in  $\mathcal{H} = \ell^2(\mathbb{Z})$ , where we consider the lattice  $\mathbb{G} = \mathbb{Z}$  as endowed with the automorphism group of shifts  $x \in I = \mathbb{Z}$  so that

$$S_x n := n - x \quad \forall n \in \mathbb{Z}, \tag{2.4}$$

which is clearly vertex-transitive. We will write wave-functions  $\psi \in \mathcal{H}$  as  $\psi = (\psi_n)_{n \in \mathbb{Z}}$ , and the discrete Laplacian  $\Delta$  acting on wave-functions by

$$(\Delta \psi)_n = \psi_{n+1} + \psi_{n-1}.$$
 (2.5)

Note that we define the discrete Laplacian without the diagonal term  $-2\psi_n$  which does not affect the operator's spectral properties.

#### 2.2.1 The almost-Mathieu operator

The almost-Mathieu operator is famous for its remarkable spectral properties, such as the Hofstadter butterfly [16], and Aubry–André duality [1].

**Definition 2.5.** Let  $\theta \in [0, 2\pi)$ . For a fixed irrational  $\alpha \in (0, 1)$  and  $\lambda > 0$ , the *almost-Mathieu operator* is the deterministic operator

$$(H(\theta)\psi)_n = -(\Delta\psi)_n + \lambda\cos(2\pi\alpha n + \theta)\psi_n, \quad n \in \mathbb{Z}.$$
(2.6)

The almost-Mathieu operator is an ergodic operator in the following sense.

**Theorem 2.6.** For each  $x \in \mathbb{Z}$ , let

$$T_x\theta := \theta - 2\pi\alpha x \mod 2\pi, \quad (U_x\psi)_n = \psi_{n-x}, \quad n \in \mathbb{Z}.$$
(2.7)

Then,  $\{T_x\}_{x\in\mathbb{Z}}$  is a group of measure-preserving transformations whose action on the probability space  $([0, 2\pi), \mathcal{B}, \mathbb{P})$ , where  $\mathcal{B}$  denotes the Borel sets, and  $\mathbb{P}$  denotes the normalized Lebesgue measure, is ergodic. In addition, we have that  $\{U_x\}_{x\in\mathbb{Z}}$  is a group of unitary operators such that

$$H(T_x\theta) = U_x H(\theta) U_x^{\dagger}.$$
(2.8)

Thus, the almost-Mathieu operator is ergodic in the sense of Definition 2.3.

The most difficult step in the proof of Theorem 2.6 is the proof that "irrational rotations" of the circle, i.e., the transformations  $\{T_x\}_{x\in\mathbb{Z}}$  in (2.7), are ergodic. We prove this in Appendix A. The irrational rotations are notable here since we will use a similar form of ergodic transformations to discuss the ergodic properties of almost-periodic operators coming from incommensurate lattice shifts.

#### 2.2.2 The Anderson model

The Anderson model is another famous ergodic operator, introduced by Anderson [2] to model the effect of disorder on electrical conductivity.

**Definition 2.7.** Consider the probability space  $(\Omega, \mathcal{A}, \mathbb{P})$ , where  $\Omega = \mathbb{R}^{\mathbb{Z}}$ ,  $\mathcal{A}$  is the Borel sets generated by the product topology, and  $\mathbb{P}$  is a probability measure on this space. The *Anderson model* is the random Schrödinger operator

$$(H(\omega)\psi)_n = -(\Delta\psi)_n + V_n(\omega)\psi_n, \qquad (2.9)$$

where  $V_n(\omega) = \omega_n$ , where  $\omega = (\omega_n)_{n \in \mathbb{Z}} \in \Omega$ .

The following establishes sufficient conditions such that the Anderson model is ergodic.

**Theorem 2.8.** Fix a probability measure  $\mathbb{P}_0$  on the probability space  $(\mathbb{R}, \mathcal{B}, \mathbb{P}_0)$ , where  $\mathcal{B}$  is the Borel sets in  $\mathbb{R}$ . Then, suppose that the probability measure  $\mathbb{P}$  in Definition 2.7 is the product measure corresponding to  $\mathbb{P}_0$ . For each  $x \in \mathbb{Z}$ , let

$$(T_x\omega)_n := \omega_{n-x}, \quad (U_x\psi)_n := \psi_{n-x}, \quad n \in \mathbb{Z}.$$
(2.10)

Then,  $\{T_x\}_{x\in\mathbb{Z}}$  is a group of measure-preserving transformations whose action on the probability space  $(\Omega, \mathcal{A}, \mathbb{P})$  is ergodic, and  $\{U_x\}_{x\in\mathbb{Z}}$  is a group of unitary operators such that

$$H(T_x\omega) = U_x H(\omega) U_x^{\dagger}.$$
(2.11)

Thus, the Anderson model is ergodic in the sense of Definition 2.3.

As concrete examples,  $\mathbb{P}_0$  could be taken as the uniform measure on any interval, or corresponding to the Gaussian distribution. The proof of ergodicity of the transformations  $\{T_x\}_{x\in\mathbb{Z}}$  (2.10) follows from the observation that the transformations are *mixing* on the probability space; see [23, 26], or Exercise 3.2 of [1] for the specific application to the Anderson model.

#### 2.3 Existence of density of states measure for ergodic Schrödinger operators

In order to demonstrate the power of the concepts introduced in the previous section, we now show that they suffice to prove convergence of the density of states measure in the thermodynamic limit. The density of states measure is a key concept when studying electronic properties of materials.

We start by recalling the *Riesz representation theorem*. Recall that a linear functional I on  $C_c(\mathbb{R})$ (continuous functions on  $\mathbb{R}$  with compact support) is called *positive* if  $I(f) \ge 0$  whenever  $f \ge 0$ . A Borel measure is called *outer regular* on a set E if

$$\mu(E) = \inf\{\mu(U) : U \supset E, U \text{ open}\},\tag{2.12}$$

and *inner regular* on E if

$$\mu(E) = \sup\{\mu(K) : K \subset E, K \text{ compact}\}.$$
(2.13)

A Borel measure is called a *Radon measure* if it is finite on compact sets, outer regular on Borel sets, and inner regular on all open sets. The Riesz representation theorem states that if I is a positive linear functional on  $C_c(\mathbb{R})$ , there is a unique Radon measure  $\mu$  on  $\mathbb{R}$  such that

$$I(f) = \int_{\mathbb{R}} f \, \mathrm{d}\mu, \quad \forall f \in C_c(\mathbb{R}).$$
(2.14)

In particular, Radon measures can be defined via their action on test functions, and vice versa.

Throughout this section, let  $H(\omega)$  be an ergodic Schrödinger operator in the sense of Definition 2.3, acting on the lattice  $\mathbb{Z}$  endowed with the automorphism group (2.4), with probability space  $(\Omega, \mathcal{A}, \mathbb{P})$ . Our goal is to define the *density of states measure* for this operator. It is straightforward to define the density of states measure for finite truncations of this operator as follows.

**Definition 2.9.** For positive integers L, let  $\Lambda_L := [-L, L] \cap \mathbb{Z}$ , so that  $|\Lambda_L| = 2L + 1$ . Then, for each  $\omega \in \Omega$ , define the truncated operator  $H_L(\omega) = 1_{\Lambda_L} H(\omega) 1_{\Lambda_L}$ , where  $1_{\Lambda_L}$  is the characteristic function for the set  $\Lambda_L$ . We define the density of states measure for  $H_L(\omega)$  as the unique Radon measure  $dk_{\omega,L}$  corresponding to the linear functional

$$g \mapsto \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \langle \delta_x, g(H_L(\omega)) \delta_x \rangle = \int_{\mathbb{R}} g \, \mathrm{d}k_{\omega,L}, \qquad (2.15)$$

where, for each  $x \in \mathbb{Z}$ ,  $\delta_x$  is the vector whose xth entry is 1 and all others are 0.

We can derive a more intuitive formula for the density of states measure (2.15) as follows. Using the spectral theorem for Hermitian matrices [25] we have that

$$g(H_L(\omega)) = \sum_{j=1}^{2L+1} g(E_{j,\omega,L}) P_{j,\omega,L}, \quad \forall g \in C_c(\mathbb{R}),$$
(2.16)

where  $E_{j,\omega,L}$  and  $P_{j,\omega,L}$  denote the eigenvalues and associated eigenprojections of  $H_L(\omega)$ . But now we have

$$\frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \langle \delta_x, g(H_L(\omega)) \delta_x \rangle = \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \sum_{j=1}^{2L+1} g(E_{j,\omega,L}) \langle \delta_x, P_{j,\omega,L} \delta_x \rangle = \frac{1}{|\Lambda_L|} \sum_{j=1}^{2L+1} g(E_{j,\omega,L}),$$
(2.17)

where we used the fact that the trace of a projection matrix equals its rank, and hence we have

$$dk_{\omega,L} = \frac{1}{|\Lambda_L|} \sum_{j=1}^{2L+1} \delta_{E_{j,\omega,L}},$$
(2.18)

where  $\delta_{E_{j,\omega,L}}$  denotes the Dirac measure for the point  $E_{j,\omega,L}$ . In short, the density of states measure for the finite-dimensional matrix  $H_L(\omega)$  is a normalized sum of delta measures at each of its eigenvalues.

Our goal is to prove convergence of this measure for truncations of ergodic operators in the limit  $L \to \infty$ . The limiting measure is as follows.

**Definition 2.10.** The density of states measure (DOSM) is the measure dk defined by the linear functional

$$\int_{\mathbb{R}} g \, \mathrm{d}k := \mathbb{E}(\langle \delta_0, g(H(\omega))\delta_0 \rangle) = \int_{\Omega} \langle \delta_0, g(H(\omega))\delta_0 \rangle \, \mathrm{d}\mathbb{P}, \quad \forall g \in C_c(\mathbb{R}).$$
(2.19)

Note that the measure (2.15) depends on  $\omega$  but the limiting measure (2.19) does not. To show this is the correct limit, we first need the following result called Birkhoff's ergodic theorem.

**Theorem 2.11** (Birkhoff's Ergodic Theorem). Let  $(\Omega, \mathcal{A}, \mathbb{P})$  be a probability space equipped with a group of measuring preserving transformation,  $\{T_i\}_{i \in \mathbb{N}}$ , which acts ergodically. Then for each  $f \in L^1(\Omega, \mathbb{P})$  the following limit exists for  $\mathbb{P}$ -almost all  $\omega \in \Omega$  and satisfies

$$\lim_{N \to \infty} \frac{1}{N} \sum_{n=0}^{N-1} f(T_n \omega) = \mathbb{E}[f], \qquad (2.20)$$

where  $\mathbb{E}$  denotes the expectation of f with respect to  $\mathbb{P}$ , i.e.,

$$\mathbb{E}[f] := \int_{\Omega} f \, \mathrm{d}\mathbb{P}.$$
(2.21)

We refer to the full proof of Birkhoff's ergodic theorem to that of Theorem 3.3.1 in [14]. Following [22], we provide a proof in Appendix A for the simple case of Birkhoff's ergodic theorem below where f is smooth on the circle  $\mathbb{T} = [0, 1)$  and  $T_n$  are irrational rotations of the circle.

**Proposition 2.12.** Let  $\Omega = \mathbb{T}$  be the circle with  $\mathcal{A}$  the Borel sets and  $\mathbb{P}$  the uniform probability measure on  $\Omega$ . Further, let  $\{T_i\}_{i \in \mathbb{Z}}$  be the irrational rotations of the circle. Then if  $f \in C^{\infty}(\mathbb{T})$  (smooth functions), the following limit exists for all  $\omega \subset \Omega$  and satisfies

$$\lim_{N \to \infty} \frac{1}{2N+1} \sum_{n=-N}^{N} f(T_n \omega) = \mathbb{E}[f].$$
(2.22)

With such choice of functions f, note that the above proposition removes the stipulation that only  $\omega$  up to a full measure set in  $\Omega$  may satisfy the convergence. Furthermore, the proof mechanism uses the Fourier series of f to show the desired equality, which by uniform density of  $C^{\infty}(\mathbb{T})$  in  $C(\mathbb{T})$ , it further suffices that f need only be continuous and periodic.

Another result needed in the proof technique is the use of the Stone-Weierstrass theorem for dense subsets of  $C_0(\mathbb{R})$ , continuous functions decaying at infinity. We present the following form referenced in [1] relevant to our functions of an operator. Recall an *involutive subalgebra*  $\mathcal{A}$  of  $C_0(\mathbb{R})$  is a linear subspace and if  $f, g \in \mathcal{A}$ , then the (pointwise) product  $f \cdot g$  and the  $\overline{f}$  are in  $\mathcal{A}$ . Further, a subalgebra  $\mathcal{A} \subset C_0(\mathbb{R})$  separates points if for any  $x, y \in \mathbb{R}$ , there is a function  $f \in \mathcal{A}$  such that  $f(x) \neq f(y)$  with f(x) and f(y) non-zero.

**Theorem 2.13.** If  $\mathcal{A}$  is an involutive subalgebra of  $C_0(\mathbb{R})$  which separates points, then  $\mathcal{A}$  is dense in  $C_0(\mathbb{R})$  with respect to the topology of uniform convergence.

Equipped with Birkhoff's ergodic theorem and the above form of Stone-Weierstrass, we now state the theorem for the convergence of the truncated density of states for ergodic operators on  $\ell^2(\mathbb{Z})$ .

**Theorem 2.14.** Assume that  $H(\omega)$  is such that  $H_L(\omega) - 1_{\Lambda_L} H(\omega)$  is trace-class, and that

$$\operatorname{tr}|H_L(\omega) - 1_{\Lambda_L} H(\omega)| \le \varepsilon(L)|\Lambda_L|, \qquad (2.23)$$

where  $\varepsilon(L) \to 0$  as  $L \to \infty$ . Then there exists a full measure set  $\Omega_0 \subset \Omega$  such that for all  $\omega \in \Omega_0$  and all functions  $g \in C_0(\mathbb{R})$  the following limit exists and satisfies

$$\lim_{L \to \infty} \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \langle \delta_x, g(H_L(\omega)) \delta_x \rangle = \mathbb{E} \left( \langle \delta_0, g(H(\omega)) \delta_0 \rangle \right).$$
(2.24)

The condition that  $H_L(\omega) - 1_{\Lambda_L} H(\omega)$  must be trace-class is satisfied as long as the matrix elements of  $H(\omega)$  decay sufficiently rapidly away from the diagonal, e.g., exponentially fast. Before proving Theorem 2.14, note that it clearly implies the following corollary.

**Corollary 2.15.** For almost-all  $\omega \in \Omega$ , the measures  $dk_{\omega,L}$  converge weakly to dk, i.e.,

$$\lim_{L \to \infty} \int f \, \mathrm{d}k_{\omega,L} = \int f \, \mathrm{d}k \tag{2.25}$$

for all  $f \in C_b(\mathbb{R})$  (continuous and uniformly bounded functions), as  $L \to \infty$ .

*Proof of Theorem 2.14.* The following proof is similar to those of Theorem 3.15 of [1] and Theorem 4.3.8 in [14].

We start by establishing (2.24) for the specific function g(x) = 1/(x-z) for fixed  $z \in \mathbb{C} \setminus \mathbb{R}$ . Later we will apply Stone-Weierstrass to obtain convergence for general  $g \in C_0(\mathbb{R})$ . First, note that one can use the resolvent identity to get that

$$\begin{aligned} \left| \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \langle \delta_x, g(H_L(\omega)) \delta_x \rangle - \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \langle \delta_x, g(H(\omega)) \delta_x \rangle \right| \\ &= \frac{1}{|\Lambda_L|} \left| \sum_{x \in \Lambda_L} \langle \delta_x, \left( (H_L(\omega) - z)^{-1} - (H(\omega) - z)^{-1} \right) \delta_x \rangle \right| \\ &= \frac{1}{|\Lambda_L|} \left| \sum_{x \in \Lambda_L} \langle \delta_x, (H_L(\omega) - z)^{-1} (H(\omega) - H_L(\omega)) (H(\omega) - z)^{-1} \delta_x \rangle \right|. \end{aligned}$$

Using the triangle inequality, we can estimate

$$\frac{1}{|\Lambda_L|} \left| \sum_{x \in \Lambda_L} \langle \delta_x, (H_L(\omega) - z)^{-1} (H(\omega) - H_L(\omega)) (H(\omega) - z)^{-1} \delta_x \rangle \right|$$
  
$$\leq \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \left| \langle \delta_x, (H_L(\omega) - z)^{-1} (H(\omega) - H_L(\omega)) (H(\omega) - z)^{-1} \delta_x \rangle \right|.$$

We now estimate the right-hand side by the trace norm (2.23) multiplied by the operator norms of the resolvents [13], using the fact that  $|(A - z)^{-1}| \leq |\operatorname{Im}(z)|^{-1}$  for a self-adjoint operator A with  $z \in \mathbb{C} \setminus \mathbb{R}$  [1],

$$\frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \left| \langle \delta_x, (H_L(\omega) - z)^{-1} (H(\omega) - H_L(\omega)) (H(\omega) - z)^{-1} \delta_x \rangle \right|$$
  
$$\leq \frac{1}{|\Lambda_L| \cdot |\operatorname{Im}(z)|^2} \operatorname{tr} |H_L(\omega) - 1_{\Lambda_L} H(\omega)| \leq \frac{\varepsilon(L)}{|\operatorname{Im}(z)|^2}.$$

We thus conclude that

$$\lim_{L \to \infty} \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \langle \delta_x, g(H_L(\omega)) \delta_x \rangle = \lim_{L \to \infty} \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \langle \delta_x, g(H(\omega)) \delta_x \rangle.$$
(2.26)

To calculate the limit on the right-hand side, we note that, using (2.2)-(2.3),

$$\langle \delta_x, g(H(\omega))\delta_x \rangle = \langle U_x \delta_0, g(H(\omega))U_x \delta_0 \rangle = \langle \delta_0, U_x^{-1}g(H(\omega))U_x \delta_0 \rangle = \langle \delta_0, g(H(T_x\omega))\delta_0 \rangle.$$
(2.27)

But now Theorem 2.11 implies that

$$\lim_{L \to \infty} \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \langle \delta_x, g(H(\omega)) \delta_x \rangle = \lim_{L \to \infty} \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \langle \delta_0, g(H(T_x\omega)) \delta_0 \rangle = \mathbb{E}\left( \langle \delta_0, g(H(\omega)) \delta_0 \rangle \right)$$
(2.28)

for all  $\omega$  in a full measure subset  $\Omega_z \subset \Omega$  depending on z.

Let  $\mathcal{B} := \mathbb{Q} + i\mathbb{Q}_{\neq 0} \subset \mathbb{C} \setminus \mathbb{R}$ , where  $\mathbb{Q}$  denotes the set of rational numbers and  $\mathbb{Q}_{\neq 0}$  denotes the set of rational numbers omitting 0. Since  $\mathcal{B}$  is countable, the set  $\Omega_0 := \bigcap_{z \in \mathcal{B}} \Omega_z$  is still of full measure in  $\Omega$  with respect to  $\mathbb{P}$ . Then, let  $\mathcal{A}$  denote the involutive subalgebra generated by the functions (i.e., the smallest involutive subalgebra containing the functions)

$$\{x \mapsto 1/(x-z) : z \in \mathcal{B}\}.$$

Note that (2.24) holds for all  $g \in \mathcal{A}$  and  $\omega \in \Omega_0$  since g only involves finite linear combinations and/or finite products of resolvent functions, and the argument above for a single resolvent generalizes easily to these cases. Moreover, the set  $\mathcal{A}$  is dense with respect to the uniform topology within the involutive subalgebra generated by the functions  $\{x \mapsto 1/(x-z) : z \in \mathbb{C} \setminus \mathbb{R}\}$ . Since the latter subalgebra separates points, it follows from Stone-Weierstrass that  $\mathcal{A}$  is dense with respect to the uniform topology within  $C_0(\mathbb{R})$ . But now it follows that (2.24) holds for all  $g \in C_0(\mathbb{R})$  for all  $\omega \in \Omega_0$ .

# 3 Incommensurate bilayer materials: 1D tight-binding models

#### 3.1 Model of incommensurate coupled chains

In this section, we introduce the incommensurate coupled chain model, a 1D model which shares many of the features of tight-binding models of twisted bilayer graphene and other moiré materials. The model describes an electron hopping along two 1D chains, one with lattice constant 1, and another with lattice constant  $1-\theta$ , where  $0 < \theta < 1$  is irrational. Irrationality of  $\theta$  implies that the system has no exact periodic cell, so we call the chains *incommensurate*.

The coupled chain Hilbert space consists of infinite vectors defined on the lattices

$$\phi = \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}, \qquad \phi_i = (\phi_{n_i})_{n_i \in \mathbb{Z}}, \qquad \phi_{n_i} \in \mathbb{C},$$
(3.1)

with the natural inner product

$$\langle \phi \mid \psi \rangle = \sum_{n_1 \in \mathbb{Z}} \overline{\phi_{n_1}} \psi_{n_1} + \sum_{n_2 \in \mathbb{Z}} \overline{\phi_{n_2}} \psi_{n_2}.$$
(3.2)

We define the coupled chain Hamiltonian as follows.

**Definition 3.1.** For each  $b \in \mathbb{R}$ , known as the *interlayer shift*, let H(b) be the self-adjoint operator

$$H(b)\psi = \begin{pmatrix} (H(b)\psi)_1\\ (H(b)\psi)_2 \end{pmatrix}, \qquad (H(b)\psi)_i = ((H(b)\psi)_{n_i})_{n_i \in \mathbb{Z}},$$
(3.3)

where

$$(H(b)\psi)_{n_1} = \psi_{n_1+1} + \psi_{n_1-1} + \sum_{n_2 \in \mathbb{Z}} h(n_1 - (1-\theta)n_2 - b)\psi_{n_2}$$
  

$$(H(b)\psi)_{n_2} = \psi_{n_2+1} + \psi_{n_2-1} + \sum_{n_1 \in \mathbb{Z}} h(n_1 - (1-\theta)n_2 - b)\psi_{n_1},$$
(3.4)

where h is the interlayer hopping function: an even  $(h(-\eta) = h(\eta))$ , real, smooth, exponentially-decaying function.

A physically realistic choice for the interlayer hopping function is

$$h(\eta) := A e^{-B\sqrt{\eta^2 + L^2}},$$
(3.5)

where A, B, L > 0 are constants. Here, L > 0 represents the interchain distance. With this choice, the hopping amplitude between site  $n_1$  on chain 1, and site  $n_2$  on chain 2, depends only on the intersite distance  $\sqrt{(n_1 - (1 - \theta)n_2 - b)^2 + L^2}$ . However, we do not expect the specific form of h to modify the essential features of the model. We restrict to nearest-neighbor hopping for the intralayer hopping for simplicity.

For fixed b, the model (3.4)-(3.5) describes an electron hopping along a chain of atoms with positions  $\left\{\left(n, \frac{L}{2}\right)^{\top}\right\}_{n \in \mathbb{Z}}$ , which can hop to a second chain of atoms with positions  $\left\{\left((1-\theta)n+b, -\frac{L}{2}\right)^{\top}\right\}_{n \in \mathbb{Z}}$ . Note that we fix the origin in the x direction to coincide with the position of an atom in the top layer without loss of generality.

## 3.2 Reduced model of incommensurate coupled chains

We will also consider a simplified, or reduced, incommensurate coupled chain model defined as follows. In this model, we consider a single chain with lattice constant 1, such that electrons can hop along the chain directly or via a fictitious second chain with lattice constant  $1-\theta$ . More precisely, the *reduced incommensurate coupled chain* model is as follows.

**Definition 3.2.** For each  $b \in \mathbb{R}$ , let  $H_r(b)$  be the self-adjoint operator

$$H_{r}(b)\psi = (H_{r}(b)\psi_{n})_{n\in\mathbb{Z}}$$

$$\equiv \left(\psi_{n+1} + \psi_{n-1} + \sum_{n'\in\mathbb{Z}} \left(h(n-(1-\theta)n'-b)\sum_{n''\in\mathbb{Z}} h(n''-(1-\theta)n'-b)\psi_{n''}\right)\right)_{n\in\mathbb{Z}}$$

$$= \left(\psi_{n+1} + \psi_{n-1} + \sum_{n'\in\mathbb{Z}} \sum_{n''\in\mathbb{Z}} h(n-(1-\theta)n'-b)h(n''-(1-\theta)n'-b)\psi_{n''}\right)_{n\in\mathbb{Z}},$$
(3.6)

where h is an even, smooth, exponentially-decaying function.

We consider Definition 3.2 as a simplified version of Definition 3.1. Thus, the discussion below Definition 3.1 applies to the interlayer hopping function here as well.

## 4 Incommensurate bilayer materials: ergodic properties

#### 4.1 Reduced incommensurate coupled chain model as an ergodic operator

The following theorem asserts that the reduced incommensurate coupled chain model is an ergodic operator in the sense of Definition 2.3.

**Theorem 4.1.** For each  $x \in \mathbb{Z}$ , let

$$T_x b = b - x \mod (1 - \theta), \quad (U_x \psi)_n = \psi_{n-x}, \quad n \in \mathbb{Z}.$$
(4.1)

Then,  $\{T_x\}_{x\in\mathbb{Z}}$  is a group of measure-preserving transformations whose action on the probability space  $([0, 1-\theta), \mathcal{A}, \mathbb{P})$ , where  $\mathcal{A}$  denotes the Borel sets and  $\mathbb{P}$  denotes the normalized Lebesgue measure, is ergodic, and  $\{U_x\}_{x\in\mathbb{Z}}$  is a group of unitary operators such that

$$H_r(T_x b) = U_x H_r(b) U_x^{\dagger}. \tag{4.2}$$

Thus, the reduced incommensurate coupled chain model is an ergodic operator in the sense of Definition 2.3.

We provide the proof of Theorem 4.1 in Appendix B. The argument is similar to the proof that the almost-Mathieu family of operators is ergodic. In fact, the argument uses that n' in (3.6) can be reordered under its summation to get b shifted by an integer multiple of  $(1 - \theta)$ , akin to cosine being equivalent up to a shift of  $2\pi$  used in Definition 2.5.

With ergodicity established, we can apply Theorem 2.14 to obtain weak convergence of the truncated density of states measure.

**Corollary 4.2.** For positive integers L, let  $\Lambda_L := [-L, L] \cap \mathbb{Z}$ . Then, for each  $b \in [0, 1 - \theta)$ , define the truncated reduced coupled chain operator  $H_{r,L}(b) = 1_{\Lambda_L} H_r(b) 1_{\Lambda_L}$ , where  $1_{\Lambda_L}$  is the characteristic function for the set  $\Lambda_L$ . Then

$$\lim_{L \to \infty} \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \langle \delta_x, g(H_{r,L}(b)) \delta_x \rangle = \mathbb{E}(\langle \delta_0, g(H_r(b)) \delta_0 \rangle), \tag{4.3}$$

where  $\mathbb{E}$  denotes expectation with respect to the normalized Lebesgue measure, i.e.,

$$\mathbb{E}(\langle \delta_0, g(H_r(b))\delta_0 \rangle) := \frac{1}{|1-\theta|} \int_0^{1-\theta} \langle \delta_0, g(H_r(b))\delta_0 \rangle \,\mathrm{d}b.$$
(4.4)

Result (4.4) has a simple interpretation: the density of states for the reduced incommensurate coupled chain model is obtained by averaging the inter-layer shift b over the unit cell of the fictitious second layer. We will see the same idea in the following section.

## 4.2 Convergence of the density of states measure for the incommensurate coupled chain model

In this section, we will prove convergence of the truncated density of states measure for the full incommensurate coupled chain model. The fact that the model has two chains makes the proof slightly more involved than the case of the reduced coupled chain model. In particular, we will not prove this model is ergodic in the sense of Definition 2.3 as an intermediate step. However, we will see that the proof follows from the same essential ideas.

We start by defining the density of states for the incommensurate coupled chain truncated to the interval [-L, L] for L > 0 simultaneously for each chain, subject to Dirichlet boundary conditions.

**Definition 4.3.** For each fixed  $b \in \mathbb{R}$ , and each positive L > 0, let  $\Lambda_L^1 := \mathbb{Z} \cap [-L, L]$  and  $\Lambda_L^2 := \mathbb{Z} \cap [-\frac{L+b}{1-\theta}, \frac{L-b}{1-\theta}]$ . For L > |b|, we have

$$|\Lambda_L| := |\Lambda_L^1| + |\Lambda_L^2|, \text{ where } |\Lambda_L^1| = 1 + 2\lfloor L \rfloor, \quad |\Lambda_L^2| = 1 + \left\lfloor \frac{L+b}{1-\theta} \right\rfloor + \left\lfloor \frac{L-b}{1-\theta} \right\rfloor, \tag{4.5}$$

where  $\lfloor \cdot \rfloor$  denotes the floor function, i.e., the mapping of a positive number  $\eta$  to the largest integer less than or equal to  $\eta$ . Let

$$1_{\Lambda_L}\psi := \begin{pmatrix} (1_{\Lambda_L}\psi)_1\\ (1_{\Lambda_L}\psi)_2 \end{pmatrix}, \quad (1_{\Lambda_L}\psi)_i = ((1_{\Lambda_L}\psi)_{n_i})_{n_i \in \mathbb{Z}}, \quad i \in \{1, 2\},$$
(4.6)

where

$$(1_{\Lambda_L}\psi)_{n_1} = \begin{cases} \psi_{n_1} & n_1 \in \Lambda_L^1 \\ 0 & \text{else,} \end{cases} \quad (1_{\Lambda_L}\psi)_{n_2} = \begin{cases} \psi_{n_2} & n_2 \in \Lambda_L^2 \\ 0 & \text{else.} \end{cases}$$
(4.7)

Then, let  $H_L(b) := 1_{\Lambda_L} H(b) 1_{\Lambda_L}$ . We define the density of states measure for  $H_L(b)$  as the unique Radon measure  $dk_{b,L}$  corresponding to the linear functional

$$g \mapsto \frac{1}{|\Lambda_L|} \left( \sum_{x_1 \in \Lambda_L^1} \langle \delta_{x_1}, g(H_L(b)) \delta_{x_1} \rangle + \sum_{x_2 \in \Lambda_L^2} \langle \delta_{x_2}, g(H_L(b)) \delta_{x_2} \rangle \right) = \int_{\mathbb{R}} g \, \mathrm{d}k_{b,L}, \tag{4.8}$$

where, for each  $x_1 \in \mathbb{Z}$ ,  $\delta_{x_1}$  is the vector defined by

$$\delta_{x_1} := \begin{pmatrix} (\delta_{x_1})_1 \\ (\delta_{x_1})_2 \end{pmatrix}, \quad (\delta_{x_1})_i = ((\delta_{x_1})_{n_i})_{n_i \in \mathbb{Z}}, \quad i \in \{1, 2\},$$
(4.9)

where

$$(\delta_{x_1})_{n_1} = \delta_{n_1, x_1}, \quad n_1 \in \mathbb{Z}, \quad (\delta_{x_1})_{n_2} = 0, \quad n_2 \in \mathbb{Z},$$

$$(4.10)$$

where  $\delta_{n_1,x_1}$  is the Kronecker delta, and  $\delta_{x_2}$  is defined similarly, but for the other layer.

An identical calculation to the calculation below Definition 2.9 shows that the measure (4.8) is exactly a normalized sum of delta functions at the eigenvalues of the truncated operator  $H_L(b)$ .

We now introduce the ergodic structure which will allow us to prove convergence of the density of states measure (4.8) in the limit  $L \to \infty$ . We start with the following lemma, which follows immediately from the definitions.

**Lemma 4.4.** For each  $x \in \mathbb{Z}$ , define the operator  $U_x^1$ , which shifts in layer 1 by x while leaving layer 2 unchanged, as

$$U_x^1 \psi = \begin{pmatrix} (U_x^1 \psi)_1 \\ (U_x^1 \psi)_2 \end{pmatrix}, \quad (U_x^1 \psi)_i = \left( (U_x^1 \psi)_{n_i} \right)_{n_i \in \mathbb{Z}}, \quad i \in \{1, 2\},$$
(4.11)

where

$$(U_x^1\psi)_{n_1} = \psi_{n_1-x}, \quad (U_x^1\psi)_{n_2} = \psi_{n_2}, \quad n_1, n_2 \in \mathbb{Z}.$$
 (4.12)

Define operators  $U_x^2$  which shift by x in layer 2 and leave layer 1 unchanged analogously. Then, for each  $x \in \mathbb{Z}$  and  $b \in \mathbb{R}$ , let

$$T_x^1 b := b + x, \quad T_x^2 b := b - (1 - \theta)x.$$
 (4.13)

Then, we have

$$H(T_x^1 b) = U_x^1 H(b)(U_x^1)^{\dagger}, \quad H(T_x^2 b) = U_x^2 H(b)(U_x^2)^{\dagger}.$$
(4.14)

Identities (4.14) parallel (2.2) in Definition 2.3. We next require the following lemma, which establishes that the transformations (4.13) act ergodically when restricted to the unit cells of the other lattice with the uniform probability measure.

**Lemma 4.5.** Let  $\tilde{T}_x^1$  and  $\tilde{T}_x^2$  denote the transformations (4.14) restricted to the unit cells of layers 2 and 1 respectively, i.e.,

$$\begin{aligned}
\dot{T}_x^1 b &= b + x \qquad \mod (1 - \theta), \\
\tilde{T}_x^2 b &= b - (1 - \theta)x \qquad \mod 1
\end{aligned}$$
(4.15)

for each  $b \in \mathbb{R}$  and  $x \in \mathbb{Z}$ . Then, if  $0 < \theta < 1$  is irrational, the operators  $(\tilde{T}^1_x)_{x \in \mathbb{Z}}$  form a group of measurepreserving transformations which acts ergodically on the probability space  $([0, 1-\theta), \mathcal{A}, \mathbb{P}_2)$ , where  $\mathcal{A}$  denotes the Borel sets and  $\mathbb{P}_2$  is the normalized Lebesgue measure on  $[0, 1-\theta)$ . Similarly, the operators  $(\tilde{T}^2_x)_{x \in \mathbb{Z}}$  act ergodically on the probability space  $([0, 1), \mathcal{A}, \mathbb{P}_1)$ , where  $\mathbb{P}_1$  is the normalized Lebesgue measure on [0, 1).

*Proof.* The proof follows from ergodicity of irrational rotations of the circle; see Appendix A.  $\Box$ 

Note that the notation  $\mathbb{P}_1$  for the measure on [0, 1) and  $\mathbb{P}_2$  for the measure on  $[0, 1 - \theta)$  is natural given that these are the unit cells of layers 1 and 2, respectively.

We can now define the limiting density of states measure concisely as follows.

**Definition 4.6.** The density of states measure for H(b) is the measure dk defined by the linear functional

$$g \mapsto \frac{1-\theta}{1+(1-\theta)} \mathbb{E}_2[\langle \delta_{0_1}, g(H(\cdot))\delta_{0_1}\rangle] + \frac{1}{1+(1-\theta)} \mathbb{E}_1[\langle \delta_{0_2}, g(H(\cdot))\delta_{0_2}\rangle] = \int_{\mathbb{R}} g \,\mathrm{d}k, \tag{4.16}$$

where

$$\mathbb{E}_{2}[\langle \delta_{0_{1}}, g(H(\cdot))\delta_{0_{1}}\rangle] = \int_{[0,1-\theta)} \langle \delta_{0_{1}}, g(H(b_{2}))\delta_{0_{1}}\rangle \,\mathrm{d}\mathbb{P}_{2} = \frac{1}{1-\theta} \int_{[0,1-\theta)} \langle \delta_{0_{1}}, g(H(b_{2}))\delta_{0_{1}}\rangle \,\mathrm{d}b_{2}, \tag{4.17}$$

and

$$\mathbb{E}_{1}[\langle \delta_{0_{2}}, g(H(\cdot))\delta_{0_{2}}\rangle] = \int_{[0,1)} \langle \delta_{0_{2}}, g(H(b_{1}))\delta_{0_{2}}\rangle \,\mathrm{d}\mathbb{P}_{1} = \int_{[0,1)} \langle \delta_{0_{2}}, g(H(b_{1}))\delta_{0_{2}}\rangle \,\mathrm{d}b_{1}.$$
(4.18)

In short, the limiting density of states measure is obtained by averaging the interlayer shift b over the unit cells of each layer, up to normalization factors reflecting the excess of atoms in layer 2 compared with layer 1 within any interval [-L, L].

We now state the theorem for the convergence of traces to the desired integral using Birkhoff's theorem for the incommensurate chain operators.

**Theorem 4.7.** Let H(b) be the incommensurate chain operators defined in (3.3) equipped with the chainshifting operators  $U_x^1$  and  $U_x^2$  in Lemma 4.4. If the difference  $H_{\Lambda_L} - 1_{\Lambda_L} H(b)$  is trace-class and

$$\operatorname{tr}|H_L(b) - 1_{\Lambda_L} H(b)| \le \varepsilon(L)|\Lambda_L| \tag{4.19}$$

with  $\varepsilon(L) \to 0$  as  $L \to \infty$ , then

$$\frac{1}{|\Lambda_L|} \left( \sum_{x_1 \in \Lambda_L^1} \langle \delta_{x_1}, g(H_L(b)) \delta_{x_1} \rangle + \sum_{x_2 \in \Lambda_L^2} \langle \delta_{x_2}, g(H_L(b)) \delta_{x_2} \rangle \right) \rightarrow \frac{1-\theta}{1+(1-\theta)} \int_{[0,1-\theta]} \langle \delta_{0_1}, g(H(b_2)) \delta_{0_1} \rangle \, \mathrm{d}\mathbb{P}_2 + \frac{1}{1+(1-\theta)} \int_{[0,1)} \langle \delta_{0_2}, g(H(b_1)) \delta_{0_2} \rangle \, \mathrm{d}\mathbb{P}_1 \quad (4.20)$$

for almost-all  $b \in \mathbb{R}$  with respect to the Lebesgue measure and all  $g \in C_0(\mathbb{R})$  as  $L \to \infty$ .

As an immediate consequence, this may be restated in the following form concerning the convergence of measures for the truncated operators.

**Corollary 4.8.** For almost-all  $b \in \mathbb{R}$ , the measures  $dk_{b,L}$  defined in (4.8) converge weakly to dk as  $L \to \infty$ , i.e.

$$\lim_{L \to \infty} \int g \, \mathrm{d}k_{b,L} = \int g \, \mathrm{d}k$$

for all  $g \in C_b(\mathbb{R})$ .

Proof of Theorem 4.7. Similar to the proof of Theorem 2.14, the goal will be to first establish (4.20) for the choices of g(x) = 1/(x-z) with  $z \in \mathbb{C} \setminus \mathbb{R}$ , then extending the choice of functions to  $g \in C_0(\mathbb{R})$  from a countable set of such resolvent functions by Stone-Weierstrass.

However, there are extra considerations needed for the incommensurate coupled chain operators above those for ergodic operators. First, a determination of the relevant probability spaces and the ergodic transformations is needed. The choice of these is made in Lemma 4.5. Further, the calculation of the limiting constants in Definition 4.6 is required to convert from the finite density of states in (4.8) to a form on which Birkhoff's ergodic theorem will apply. Finally, there is an technical issue to resolve for simultaneously applying Birkhoff's ergodic theorem on each chain, restricting the choice of the b interlayer shift to probability spaces of different periodicites.

To start, we again consider the choice of function  $g : \mathbb{R} \to \mathbb{C}$  to exploit the resolvent formula with g(x) = 1/(x-z) with fixed  $z \in \mathbb{C} \setminus \mathbb{R}$ . Thus, one can similarly produce the following estimate

$$\frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \langle \delta_x, g(H_L(b)) \delta_x \rangle - \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \langle \delta_x, g(H(b)) \delta_x \rangle \right| \le \frac{\varepsilon(L)}{|\operatorname{Im}(z)|^2}, \tag{4.21}$$

where we use the shorthand notation

.

$$\sum_{x \in \Lambda_L} \langle \delta_x, (\cdot) \delta_x \rangle = \sum_{x_1 \in \Lambda_L^1} \langle \delta_{x_1}, (\cdot) \delta_{x_1} \rangle + \sum_{x_2 \in \Lambda_L^2} \langle \delta_{x_2}, (\cdot) \delta_{x_2} \rangle.$$

Indeed, this once again follows from a successive application of the resolvent identity, the triangle inequality, and passing to simple estimates on the operator and trace norms, using (4.19) for this last step. From this, we may conclude

$$\lim_{L \to \infty} \left| \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \langle \delta_x, g(H_L(b)) \delta_x \rangle - \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \langle \delta_x, g(H(b)) \delta_x \rangle \right| = 0$$
(4.22)

for any choice of  $z \in \mathbb{C} \setminus \mathbb{R}$ .

Next, the second limit of (4.22) can be factored into a usable form for the application of Birkhoff's ergodic theorem: note that we have that

$$\frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \langle \delta_x, g(H(b)) \delta_x \rangle = \frac{|\Lambda_L^1|}{|\Lambda_L^1| + |\Lambda_L^2|} \left( \frac{1}{|\Lambda_L^1|} \sum_{x_1 \in \Lambda_L^1} \langle \delta_{x_1}, g(H(b)) \delta_{x_1} \rangle \right) \\
+ \frac{|\Lambda_L^2|}{|\Lambda_L^1| + |\Lambda_L^2|} \left( \frac{1}{|\Lambda_L^2|} \sum_{x_2 \in \Lambda_L^2} \langle \delta_{x_2}, g(H(b)) \delta_{x_2} \rangle \right).$$
(4.23)

The terms in parentheses are the expressions we will use Birkhoff's ergodic theorem on.

First calculating the prefactors of the parentheses terms in (4.23) above, we may simply calculate express the size of each truncation to each chain in (4.5) as

$$|\Lambda_L^1| = 1 + 2L + \eta(L)$$
 and  $|\Lambda_L^2| = 1 + \frac{2L}{1-\theta} + \nu(L)$ 

where  $\eta : \mathbb{R} \to \mathbb{R}$  and  $\nu : \mathbb{R} \to \mathbb{R}$  are bounded functions of L. The limit of the size ratio can be calculated using these bounded functions to get

$$\lim_{L \to \infty} \frac{|\Lambda_L^1|}{|\Lambda_L^2|} = \lim_{L \to \infty} \frac{2 + \frac{1 + \eta(L)}{L}}{\frac{2}{1 - \theta} + \frac{1 + \nu(L)}{L}} = 1 - \theta,$$

from which it can be derived that

$$\lim_{L \to \infty} \frac{|\Lambda_L^1|}{|\Lambda_L^1| + |\Lambda_L^2|} = \frac{1 - \theta}{1 + (1 - \theta)} \quad \text{and} \quad \lim_{L \to \infty} \frac{|\Lambda_L^2|}{|\Lambda_L^1| + |\Lambda_L^2|} = \frac{1}{1 + (1 - \theta)}.$$
(4.24)

This gives the prefactors of the summands on the right-hand side of (4.23) in the limit as  $L \to \infty$ .

Meanwhile, the limits of each summand in (4.23) can be modified by shifting the  $\delta_x$ -vectors to a chosen fixed point, say to  $\delta_{0_1}$  and  $\delta_{0_2}$ :

$$\sum_{x_1 \in \Lambda_L^1} \langle \delta_{x_1}, g(H(b)) \delta_{x_1} \rangle = \sum_{x_1 \in \Lambda_L^1} \langle U_x^1 \delta_{0_1}, g(H(b)) U_x^1 \delta_{0_1} \rangle$$
$$= \sum_{x_1 \in \Lambda_L^1} \langle \delta_{0_1}, (U_x^1)^{\dagger} g(H(b)) U_x^1 \delta_{0_1} \rangle,$$

where  $U_x^i$  is defined in Lemma 4.4 and satisfies  $(U_x^i)^{\dagger} = U_{-x}^i$  for each  $x \in \mathbb{Z}$ . Furthermore, since the  $\delta_{x_1}$  (any  $x \in \mathbb{Z}$ ) vectors are fixed under  $U_y^2$  for all  $y \in \mathbb{Z}$ , we have

$$\sum_{x_1 \in \Lambda_L^1} \langle \delta_{x_1}, g(H(b)) \delta_{x_1} \rangle = \sum_{x_1 \in \Lambda_L^1} \langle \delta_{0_1}, (U_y^2 U_x^1)^{\dagger} g(H(b)) U_x^1 U_y^2 \delta_{0_1} \rangle$$
(4.25)

for any  $y \in \mathbb{Z}$ . In particular, note that for any  $b \in \mathbb{R}$  and any  $x \in \mathbb{Z}$ , there is a  $y(x) \in \mathbb{Z}$  such that  $b - x + (1 - \theta)y(x) \in [0, 1 - \theta)$ . We now set y in (4.25) to equal this y(x). Since g is a resolvent function, we get

$$\sum_{x_1 \in \Lambda_L^1} \langle \delta_{x_1}, g(H(b)) \delta_{x_1} \rangle = \sum_{x_1 \in \Lambda_L^1} \langle \delta_{0_1}, g\left( (U_x^1 U_{y(x)}^2)^{\dagger} H(b) U_x^1 U_{y(x)}^2 \right) \delta_{0_1} \rangle$$
$$= \sum_{x_1 \in \Lambda_L^1} \langle \delta_{0_1}, g(H(\tilde{T}_x^1 b)) \delta_{0_1} \rangle.$$
(4.26)

Note that because we chose y to equal y(x), in the argument of H we can replace  $T_x^1 b$ , which appeared in (4.13), by  $\tilde{T}_x^1 b$ , the ergodic transformation of the unit cell of layer 2 appearing in Lemma 4.5. Likewise,

$$\sum_{x_2 \in \Lambda_L^2} \langle \delta_{x_2}, g(H(b)) \delta_{x_2} \rangle = \sum_{x_2 \in \Lambda_L^2} \langle \delta_{0_2}, g(H(\tilde{T}_x^2 b)) \delta_{0_2} \rangle$$
(4.27)

for the summation over the second chain.

We may now combine (4.24), (4.26), and (4.27) into (4.23), getting

$$\lim_{L \to \infty} \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \langle \delta_x, g(H(b)) \delta_x \rangle = \frac{1 - \theta}{1 + (1 - \theta)} \left( \lim_{L \to \infty} \frac{1}{|\Lambda_L^1|} \sum_{x_1 \in \Lambda_L^1} \langle \delta_{0_1}, g(H(\tilde{T}_x^1 b)) \delta_{0_1} \rangle \right) + \frac{1}{1 + (1 - \theta)} \left( \lim_{L \to \infty} \frac{1}{|\Lambda_L^2|} \sum_{x_2 \in \Lambda_L^2} \langle \delta_{0_2}, g(H(\tilde{T}_x^2 b)) \delta_{0_2} \rangle \right)$$

Ignoring for a moment the question of the set of b for which this holds, Birkhoff's ergodic theorem may be applied to each limit in the above expression to yield, together with (4.22),

$$\lim_{L \to \infty} \frac{1}{|\Lambda_L|} \sum_{x \in \Lambda_L} \langle \delta_x, g(H_L(b)) \delta_x \rangle = \frac{1-\theta}{1+(1-\theta)} \int_{[0,1-\theta)} \langle \delta_{0_1}, g(H(b_2)) \delta_{0_1} \rangle \, \mathrm{d}\mathbb{P}_2 + \frac{1}{1+(1-\theta)} \int_{[0,1)} \langle \delta_{0_2}, g(H(b_1)) \delta_{0_2} \rangle \, \mathrm{d}\mathbb{P}_1, \tag{4.28}$$

which is the desired expression in (4.20) but with the choice of g(x) = 1/(x-z) for any  $z \in \mathbb{C} \setminus \mathbb{R}$ .

To expand the expression in (4.28) to all  $g \in C_0(\mathbb{R})$ , let  $b_2 \in \Omega_z^2 \subset [0, 1 - \theta)$  and  $b_1 \in \Omega_z^1 \subset [0, 1)$  be the sets afforded by Birkhoff's ergodic theorem such that (4.28) holds for all  $b \in \mathbb{R}$  such that  $b \mod 1 \in \Omega_z^1$ and  $b \mod (1 - \theta) \in \Omega_z^2$  with g(x) = 1/(x - z) for some  $z \in \mathbb{C} \setminus \mathbb{R}$ . Through choosing a dense countable subset  $\mathcal{B} \subset \mathbb{C} \setminus \mathbb{R}$  (e.g.  $\mathcal{B} = \{a + bi : a, b \in \mathbb{Q}, b \neq 0\}$  as in Theorem 2.14), let  $\mathcal{A}$  be an involutive subalgebra generated by the set of resolvents according to  $\mathcal{B}$ :

$$\{x \mapsto 1/(x-z) : z \in \mathcal{B}\}$$

Furthermore, let

$$\Omega^2 = \bigcap_{z \in \mathcal{B}} \Omega_z^2 \quad \text{and} \quad \Omega^1 = \bigcap_{z \in \mathcal{B}} \Omega_z^1,$$

which are the full-measure subsets  $[0, 1 - \theta)$  and [0, 1) with respect to  $\mathbb{P}_2$  and  $\mathbb{P}_1$  such that the limits on chain 1 and 2 converge via Birkhoff's ergodic theorem for any  $g \in \mathcal{A}$ , respectively.

To find the allowed initial choices of interlayer shift  $b \in \mathbb{R}$ , consider first the following subset of  $\mathbb{R}$ ,

$$T_x^2 \Omega^2 = \{ a \in [0,1) : a = T_x^2 b_2 \text{ for some } b_2 \in \Omega^2 \}$$

for each  $x \in \mathbb{Z}$ . From this set, we define

$$\tilde{\Omega}^2 = \bigcup_{x \in \mathbb{Z}} T_x^2 \Omega^2 \subset \mathbb{R}.$$

Likewise,

$$\tilde{\Omega}^1 = \bigcup_{x \in \mathbb{Z}} T^1_x \Omega^1 \subset \mathbb{R}.$$

In effect, we are trying to "undo" our choice of integrating over the interlayer shifts according to the lattice unit cells, extending the sets by periodicity from the lattice-unit interval back to the entire real numbers. Hence, as long as  $b \in \Omega := \tilde{\Omega}^1 \cap \tilde{\Omega}^2$ , the convergence in (4.28) may hold for any  $g \in \mathcal{A}$ . Lastly, note the set  $\Omega$  clearly has full Lebesgue measure as a subset of  $\mathbb{R}$  ( $\mathbb{R} \setminus \Omega$  has zero Lebesgue measure on all intervals).

From the involutive subalgebra  $\mathcal{A}$ , we may expand the choice of  $g \in \mathcal{A}$  to  $g \in C_0(\mathbb{R})$  via Stone-Weierstrass in Theorem 2.13 as in Theorem 2.14. This gives the desired result of (4.20) for any function  $g \in C_0(\mathbb{R})$  and almost-all initial choices of interlayer shift  $b \in \Omega \subset \mathbb{R}$ , completing the proof.

Of particular note in the above proof, the condition in Lemma 4.4 and specifically (4.14) is insufficient in creating a summation over transformations of an ergodic transformation as in (4.25) leading to (4.26). In particular, the invariance of vectors on one chain with respect to lattice shifts on the opposite chain provides the periodicity needed to create an ergodic transformation.

**Remark 4.9.** Note that one can show the respective limits in (4.20) exist and attains their correct limit directly following the technique of Proposition 2.12 whenever the observable  $\langle \delta_{0_i}, g(H(\cdot))\delta_{0_i} \rangle$  is e.g. smooth and periodic (or therefore merely continuous and periodic and using uniform density). The strategy follows from a similar result for continuous functions over incommensurate *atomic sheets*, the two dimensional analogue of the above results for the incommensurate chains (r.f. [22]). Furthermore, this strategy avoids the loss of choice for  $b \in \mathbb{R}$  from applying the full version of Birkhoff's ergodic theorem, but needs an extra requirement/argument to show that the observable above is continuous. The proof in 2.12 in Appendix A can be adapted to show Theorem 4.7 under such conditions.

# References

- [1] Michael Aizenman and Simone Warzel. Random operators, volume 168 of Graduate Studies in Mathematics.
- [2] P. W. Anderson. Absence of diffusion in certain random lattices. *Phys. Rev.*, 109:1492–1505, 3 1958.

- [3] Artur Avila and Svetlana Jitomirskaya. Solving the Ten Martini Problem, pages 5–16. Springer Berlin Heidelberg, Berlin, Heidelberg, 2006.
- [4] J. Avron. Topological quantum states and the 2016 Nobel prize. IAMP News Bulletin, 2017.
- [5] Simon Becker, Lingrui Ge, and Jens Wittsten. Hofstadter butterflies and metal/insulator transitions for moiré heterostructures. arxiv.org/abs/2206.11891.
- [6] Rafi Bistritzer and Allan H. MacDonald. Moiré bands in twisted double-layer graphene. Proceedings of the National Academy of Sciences of the United States of America, 108:12233–12237, 2011.
- [7] Eric Cancès, Paul Cazeaux, and Mitchell Luskin. Generalized Kubo formulas for the transport properties of incommensurate 2D atomic heterostructures. *Journal of Mathematical Physics*, 58, 2017.
- [8] Yuan Cao, Valla Fatemi, Ahmet Demir, Shiang Fang, Spencer L. Tomarken, Jason Y. Luo, Javier D. Sanchez-Yamagishi, Kenji Watanabe, Takashi Taniguchi, Efthimios Kaxiras, Ray C. Ashoori, and Pablo Jarillo-Herrero. Correlated insulator behaviour at half-filling in magic-angle graphene superlattices. Nature, 556:80–84, 2018.
- [9] Yuan Cao, Valla Fatemi, Shiang Fang, Kenji Watanabe, Takashi Taniguchi, Efthimios Kaxiras, and Pablo Jarillo-Herrero. Unconventional superconductivity in magic-angle graphene superlattices. *Nature*, 556:43–50, 2018.
- [10] S Carr, D Massatt, S B Torrisi, P Cazeaux, M Luskin, and E Kaxiras. Relaxation and domain formation in incommensurate two-dimensional heterostructures. *Phys. Rev. B*, 98:224102, 2018.
- [11] Stephen Carr, Daniel Massatt, Shiang Fang, Paul Cazeaux, Mitchell Luskin, and Effhimios Kaxiras. Twistronics: Manipulating the electronic properties of two-dimensional layered structures through their twist angle. *Physical Review B*, 95:1–6, 2017.
- [12] Paul Cazeaux, Mitchell Luskin, and Daniel Massatt. Energy minimization of two dimensional incommensurate heterostructures. Archive for Rational Mechanics and Analysis, 235:1289 – 1325, February 2020.
- [13] J.B. Conway. A Course in Functional Analysis. Graduate Texts in Mathematics. Springer New York, 1994.
- [14] David Damanik. One-dimensional ergodic Schrödinger operators. Graduate studies in mathematics, volume 221. American Mathematical Society, Providence, Rhode Island, 2022.
- [15] A. K. Geim and I. V. Grigorieva. Van der Waals heterostructures. Nature, 499:419–425, 2013.
- [16] Douglas R Hofstadter. Energy levels and wave functions of Bloch electrons in rational and irrational magnetic fields. *Phys. Rev. B*, 14:2239–2249, 9 1976.
- [17] Michael Hott, Alexander B. Watson, and Mitchell Luskin. From incommensurate bilayer heterostructures to Allen-Cahn: An exact thermodynamic limit. arxiv.org/abs/2305.18186.
- [18] Dante M. Kennes, Martin Claassen, Lede Xian, Antoine Georges, Andrew J. Millis, James Hone, Cory R. Dean, D. N. Basov, Abhay N. Pasupathy, and Angel Rubio. Moiré heterostructures as a condensed-matter quantum simulator. *Nature Physics*, 17:155–163, 2021.
- [19] Daniel Massatt, Stephen Carr, and Mitchell Luskin. Electronic observables for relaxed bilayer 2D heterostructures in momentum space. arxiv.org/abs/2109.15296.
- [20] Daniel Massatt, Stephen Carr, and Mitchell Luskin. Efficient computation of Kubo conductivity for incommensurate 2D heterostructures. *European Physical Journal B*, 93, 2020.
- [21] Daniel Massatt, Stephen Carr, Mitchell Luskin, and Christoph Ortner. Incommensurate heterostructures in momentum space. *Multiscale Modeling & Simulation*, 16:429–451, 2018.

- [22] Daniel Massatt, Mitchell Luskin, and Christoph Ortner. Electronic density of states for incommensurate layers. *Multiscale Model. Simul.*, 15(1):476–499, 2017.
- [23] M. G. Nadkarni. Basic ergodic theory. Birkhauser Verlag, Basel, 2nd edition, 1998.
- [24] K S Novoselov, A K Geim, S V Morozov, D Jiang, M I Katsnelson, I V Grigorieva, S V Dubonos, and A A Firsov. Two-dimensional gas of massless Dirac fermions in graphene. *Nature*, 438:197–200, 2005.
- [25] M Reed and B Simon. Methods of Modern Mathematical Physics, I: Functional Analysis. Academic press, 1972.
- [26] César Ernesto Silva. Invitation to ergodic theory. Student mathematical library, v. 42. American Mathematical Society, Providence, R.I, 2008.
- [27] Barry Simon. Almost periodic Schrödinger operators: A review. Advances in Applied Mathematics, 3(4):463–490, 1982.
- [28] A. Timmel and E. J. Mele. Dirac-Harper theory for one-dimensional moiré superlattices. *Physical Review Letters*, 125, 10 2020.

## A Proofs for Section 2

This section contains proofs that show the well-studied examples of random operators are ergodic. The primary inclusion here is the almost-Mathieu operator ergodicity, contained below. To show this, we first need the following result from [26] showing that the irrational rotations are well-defined candidates of ergodic transformations. Let  $T_x$  for any  $x \in \mathbb{Z}$  be the *irrational rotation* transformations defined as, similar to (2.7),

$$T_x \omega = (\omega - \alpha x) \mod 1,$$
 (A.1)

for  $\omega \in \Omega = \mathbb{T}$  on the probability space  $(\Omega, \mathcal{A}, \mu)$  with  $\alpha \in (0, 1)$  irrational,  $\mathcal{A}$  the Borel sets, and  $\mu$  the Lebesgue measure on [0, 1). Here,  $\mathbb{T}$  is identified properly as  $\mathbb{R} \mod 1$  and pointwise as [0, 1).

**Lemma A.1.** The irration rotation  $T_1$  generates a group of measure-preserving transformations.

*Proof.* Note that it suffices to check that  $T_1$  is an invertible, measure-preserving transformation on  $\Omega$ , Theorem 3.2.1 in [26]. Clearly the inverse transformation of  $T_1$ ,  $(T_1)^{-1}$ , is given by

$$(T_1)^{-1}\omega = (\omega + \alpha) \mod 1,$$

which is another irrational rotation in (A.1),  $(T_1)^{-1} = T_{-1}$ . Furthermore, note that the transformation  $(T_1)^x$  is also an irrational rotation for any  $x \in \mathbb{Z}$  and given by  $(T_1)^x = T_x$ . So, one can create the group of transformations from  $\{T_x\}_{x\in\mathbb{Z}}$  using associativity.

Next, consider any subset  $A \in \mathcal{A}$  and let  $A_0 = A \cap [0, \alpha)$  and  $A_1 = A \cap [\alpha, 1)$  partition  $\mathbb{T}$ . The sets  $T_1(A_0)$  and  $T_1(A_1)$  are clearly measurable by constituting shifts of  $A_0, A_1 \subset \mathbb{R}$ . Indeed, the measures are therefore the same by the shift-invariance of  $\mu$ , the Lebesgue measure on  $\mathbb{T} = [0, 1)$ . Therefore, the constructed group  $\{T_x\}_{x\in\mathbb{Z}}$  is measure preserving from composition, completing the proof.

Now that the irrational rotation  $T_1$  is well-defined by Lemma A.1, we next show that the orbit of any point is dense under the action of the group  $\{T_x\}_{x\in\mathbb{Z}}$ .

**Lemma A.2.** From the group of irrational rotations generated by  $T_1$  with rotation  $\alpha$ ,  $\{T_x\}_{x\in\mathbb{Z}}$ , the orbit  $\{T_x\omega: x\in\mathbb{Z}\}$  of any  $\omega\in\mathbb{T}$  is dense in  $\mathbb{T}$ .

*Proof.* This proof follows from Theorem 3.2.3 in [26], which itself is due to Kronecker. First one shows that each point in the orbit is unique. Suppose that  $T_n\omega = T_m\omega$  for any element  $\omega \in \mathbb{T}$  and some integers m, n. Then by construction of  $T_1, x + \alpha n \equiv (x + \alpha m) \mod 1$ , or  $\alpha(n-m) \equiv 0 \mod 1$ . Therefore,  $\alpha(n-m)$  being an integer implies that n = m as  $\alpha$  is irrational, making each point in the orbit of  $\omega$  unique.

Next, consider the sequence  $\{T_n\omega\}_{n\geq 0}$  for any element  $\omega \in \mathbb{T}$ . By the Bolzano-Weierstrass theorem, there is a convergent subsequence of  $\{T_n\omega\}_{n\geq 0}$  in  $\mathbb{T}$ . Thus, given any  $1/2 > \varepsilon > 0$  there exists nonnegative

integers p > q such that  $0 < d_{\mathbb{T}}(T_p\omega, T_q\omega) < \varepsilon$ . (Here,  $d_{\mathbb{T}}$  denotes the induced metric on  $d_{\mathbb{T}}$  as the infimum of the distance between any two representatives in  $\mathbb{R} \mod 1$ , which pointwise has the same definition  $d_{\mathbb{T}}(\omega_1, \omega_2) = \min\{|\omega_1 - \omega_2|, 1 - |\omega_1 - \omega_2|\}$ .) Furthermore, fixed rotations of the circle preserve distances, or  $d_{\mathbb{T}}(\omega_1, \omega_2) = d_{\mathbb{T}}(T_x\omega_1, T_x\omega_2)$  for any rotation  $T_x$  and  $\omega_1, \omega_2 \in \mathbb{T}$ . Therefore,

$$0 < d_{\mathbb{T}}(T_r\omega, \omega) < \varepsilon, \tag{A.2}$$

where r = p - q.

Finally, consider the sequence  $\{T_{lr}\omega\}_{l\geq 0}$ . Then consecutive terms are the same distance apart:

$$d_{\mathbb{T}}(T_{(l+1)r}\omega, T_{lr}\omega) = d_{\mathbb{T}}(T_r\omega, \omega) \tag{A.3}$$

using the same result as above. Since each point is distinct by the first paragraph, this implies that the points of  $\{T_{lr}\omega\}_{l\geq 0}$  divide  $[0,1) = \mathbb{T}$  at least into subintervals of length  $< \varepsilon$ . It follows that this subsequence and, hence, the sequence  $\{T_n\omega\}_{n\geq 0}$  are dense in  $\mathbb{T}$ .

Lastly, we can show that the irrational rotations are ergodic. There are several equivalent ways to show this, but, for brevity without proving equivalent conditions on the measurable sets in Definition 2.2, one can show the ergodicity using the outer regularity of the Lebesgue measure induced on  $\mathbb{T}$ .

**Lemma A.3.** The group of irrational rotations  $\{T_x\}_{x\in\mathbb{Z}}$  is an ergodic group of transformations on the space  $(\mathbb{T}, \mathcal{A}, \mu)$ .

*Proof.* The following proof is from [23]. Suppose E is a Borel set invariant under the group  $\{T_x\}_{x\in\mathbb{Z}}$  of positive Lebesgue measure,  $\mu(E) > 0$ . The desire is to show that  $\mu(E) = 1$ . Given  $\varepsilon > 0$ , cover E-a.e. by a sequence of disjoint open intervals  $C_1, C_2, C_3, \ldots$  such that  $\frac{\mu(E)}{1-\varepsilon} > \sum_{n=1}^{\infty} \mu(C_n)$  by the outer regularity of the Lebesgue measure. Furthermore, one may take each having Lebesgue measure less than  $\varepsilon$ . Since

$$(1-\varepsilon)\sum_{n=1}^{\infty}\mu(C_n) \le \mu(E) \le \sum_{n=1}^{\infty}\mu(C_n),\tag{A.4}$$

there must exist some n such that  $\mu(E \cap C_n) \ge (1 - \varepsilon)\mu(C_n)$ . Let C be this  $C_n$ .

Next, since E is invariant under the group and the rotations preserve the Lebesgue measure,  $\mu(E \cap T_m C) \ge (1-\varepsilon)\mu(C)$  for all integers m. By the density of the orbit in Lemma A.2, there exists integers  $m_1, m_2, \ldots, m_k$  such that the sets  $T_{m_1}C, T_{m_2}C, \ldots, T_{m_k}C$  are pairwise disjoint and their union has Lebesgue measure greater than  $1-2\varepsilon$ . Therefore, the disjoint property of the  $T_{m_i}C$  and the condition on C by the above paragraph,

$$\mu(E) \ge \sum_{i=1}^{k} \mu(E \cap T_{m_i}C) \ge (1-\varepsilon) \left(\sum_{i=1}^{k} T_{m_i}C\right) > (1-\varepsilon)(1-2\varepsilon),$$

so  $\mu(E) = 1$  as  $\varepsilon$  was arbitrary.

Finally, one can prove the ergodicity of the almost-Mathieu operators. The following proof notes the connection to the above lemmas.

*Proof of Theorem 2.6.* To begin, the first part is to show that the transformations

$$T_x \theta = (\theta - 2\pi\alpha x) \mod 2\pi$$

are ergodic as a group  $\{T_x\}_{x\in\mathbb{Z}}$ . This follows from applying Lemmas A.1-A.3 when  $\alpha$  is irrational, but the probability measure  $\mu$  is scaled down by  $2\pi$  for a uniform probability measure while  $T_x$  is scaled up by  $2\pi$  so as to rotate  $\mathbb{R}/2\pi$ . This yields that  $\{T_x\}_{x\in\mathbb{Z}}$  is an ergodic transformation over the probability space ( $\mathbb{R} \mod 2\pi, \mathcal{A}, \mu$ ), where  $\mu = d\lambda/2\pi$  with  $d\lambda$  the Lebesgue measure and  $\mathcal{A}$  the Borel sets on  $[0, 2\pi)$ .

Lastly, to show the almost-Mathieu operator  $H_{\alpha,\lambda}$  is ergodic, the indeed unitary operators

$$(U_x\psi)(\xi) = \psi(\xi - x),\tag{A.5}$$

corresponding to lattice shifts on  $\mathbb{Z}$  given by  $(U_x\psi)(\xi) = \psi(S_x\xi)$ , have that

$$(U_x H_{\alpha,\lambda}(\theta) U_x^{-1})(\psi)(\xi) = (U_x H_{\alpha,\lambda}(\omega))(\psi)(\xi + x)$$
  
=  $(U_x) ((-\Delta)(\psi)(\xi + x) + \lambda \cos(2\pi\alpha\xi + \theta)\psi(\xi + x))$   
=  $(-\Delta)(\psi)(\xi) + \lambda \cos(2\pi\alpha(\xi - x) + \theta)\psi(\xi)$   
=  $(-\Delta + \lambda \cos(2\pi\alpha\xi + \theta - 2\pi\alpha x))(\psi)(\xi)$   
=  $H_{\alpha,\lambda}(T_x \theta)(\psi)(\xi),$ 

constructing the desired unitary transformation  $T_x$  via direct correspondence with the graph automorphism  $S_x$  for all  $x \in \mathbb{Z}$ . Thus,  $H_{\alpha,\lambda}(\theta)$  is an ergodic operator.

Proof of Proposition 2.12. The following proof of the simple case of Birkhoff's ergodic theorem which follows the mechanism in [22] (see Remark 4.9). Recall the goal is to show that for  $f \in C^{\infty}(\mathbb{T})$  and any  $\omega \in \Omega = \mathbb{T}$ , we have that

$$\lim_{N \to \infty} \frac{1}{2N+1} \sum_{n=-N}^{N} f(T_n \omega) = \int_0^1 f(x) \, \mathrm{d}x,\tag{A.6}$$

with  $\{T_n\}_{n\in\mathbb{Z}}$  the irrational rotations of the circle (see (A.1)). The idea is to decompose f into its Fourier series,

$$f(x) = \sum_{m \in \mathbb{Z}} c_m e^{2\pi i m x}$$

with  $c_m \in \mathbb{C}$  and bound all  $m \neq 0$  terms to zero on both sides of (A.6).

First, for  $m \neq 0$ , the right-hand side in (A.6) is zero by integrating over the interval of the Fourier mode. Note that since  $f \in C^{\infty}(\mathbb{T})$ , the afforded uniform convergence of the Fourier series of the left-hand side in (A.6) allows one to independently describe each limit for each Fourier mode. To bound the left-hand side for a fixed  $m \neq 0$ , the summation can be written explicitly with each Fourier mode as

$$\frac{c_m}{2N+1} \sum_{n=-N}^{N} e^{2\pi i m(\omega - \alpha n \mod 1)} = \frac{c_m}{2N+1} \sum_{n=-N}^{N} e^{2\pi i m(\omega - \alpha n + k)}$$
$$= \frac{c_m e^{2\pi i m(\omega + k)}}{2N+1} \sum_{n=-N}^{N} e^{-2\pi i m \alpha n}$$
$$= \frac{c_m e^{2\pi i m(\omega + k)}}{2N+1} \cdot \frac{e^{-2\pi i m \alpha (N+1)} - e^{2\pi i m \alpha N}}{1 - e^{-2\pi i m \alpha}},$$

exploiting the finite geometric series. Since  $\alpha$  is irrational, the second denominator is nonzero  $1 - e^{-2\pi i m \alpha} \neq 0$  for every  $m \neq 0$ . Thus, there is a simple bound on each  $m \neq 0$  Fourier mode of the left-hand side of (A.6) as follows:

$$\left| \frac{c_m}{2N+1} \sum_{n=-N}^{N} e^{2\pi i m(T_n x)} \right| \le \left| \frac{c_m e^{2\pi i m(\omega+k)}}{2N+1} \cdot \frac{e^{-2\pi i m\alpha(N+1)} - e^{2\pi i m\alpha N}}{1 - e^{-2\pi i m\alpha}} \right| \\ \le \frac{4|c_m|}{(2N+1)|1 - e^{2\pi i m\alpha}|}.$$

Thus, as  $N \to \infty$ , the left-hand side of (A.6) vanishes for all  $m \neq 0$  as desired. Lastly, if m = 0, then both left-hand and right-hand sides in (A.6) are  $c_0 = \mathbb{E}[f]$  by direct computation. As all other Fourier modes vanish under the limit, this completes the proof.

# **B** Proofs for Section 4

This section contains extra proofs for ergodicity and related results for the family of operators defined with (effective) interlayer interactions.

Proof of Theorem 4.1. First, the calculation to see how  $H_r(b)$ , the reduced chain operators in (3.6), will transform under the indeed unitary operator  $U_1$  as follows:

$$\begin{aligned} (U_1)^{\intercal} H_r(b) U_1 \psi &= U_{-1} H_r(b) \left( \psi_{n-1} \right)_{n \in \mathbb{Z}} \\ &= U_{-1} \left( \psi_n + \psi_{n-2} + \sum_{n' \in \mathbb{Z}} \sum_{n'' \in \mathbb{Z}} h(n - (1 - \theta)n' - b)h(n'' - (1 - \theta)n' - b)\psi_{n''-1} \right)_{n \in \mathbb{Z}} \\ &= \left( \psi_{n+1} + \psi_{n-1} + \sum_{n' \in \mathbb{Z}} \sum_{n'' \in \mathbb{Z}} h(n + 1 - (1 - \theta)n' - b)h(n'' - (1 - \theta)n' - b)\psi_{n''-1} \right)_{n \in \mathbb{Z}} \\ &= \left( \psi_{n+1} + \psi_{n-1} + \sum_{n' \in \mathbb{Z}} \sum_{n'' \in \mathbb{Z}} h(n + 1 - (1 - \theta)n' - b)h(n'' + 1 - (1 - \theta)n' - b)\psi_{n''} \right)_{n \in \mathbb{Z}} \\ &= H_r(b - 1)\psi, \end{aligned}$$

or  $(U_1)^{\dagger}H_r(b)U_1 = H_r(b-1)$ , where  $(U_1)^{\dagger} = U_{-1}$ . The above definition of the Hamiltonian in (3.6) affords periodicity through the shifting of the n' summation index:

$$\begin{aligned} H_{r}(b+(1-\theta))\psi &= \left(\psi_{n+1}+\psi_{n-1}+\sum_{n'\in\mathbb{Z}}\sum_{n''\in\mathbb{Z}}\left(h(n-(1-\theta)n'-(b+(1-\theta)))\right)\\ &\quad \cdot h(n''-(1-\theta)n'-(b+(1-\theta)))\psi_{n''}\right)\Big)_{n\in\mathbb{Z}}\\ &= \left(\psi_{n+1}+\psi_{n-1}+\sum_{n'\in\mathbb{Z}}\sum_{n''\in\mathbb{Z}}\left(h(n-(1-\theta)(n'+1)-b)\right)\\ &\quad \cdot h(n''-(1-\theta)(n'+1)-b)\psi_{n''}\right)\Big)_{n\in\mathbb{Z}}\\ &= \left(\psi_{n+1}+\psi_{n-1}+\sum_{n'\in\mathbb{Z}}\sum_{n''\in\mathbb{Z}}h(n-(1-\theta)n'-b)h(n''-(1-\theta)n'-b)\psi_{n''}\right)_{n\in\mathbb{Z}}\\ &= H_{r}(b)\psi.\end{aligned}$$

From the previous two calculations,  $(U_1)^{\dagger}H_r(b)U_1 = H_r(T_1b)$ . By the idenification  $(T_1)^x = T_x$ , one gets that  $(U_x)^{\dagger}H_r(b)U_x = H_r(T_xb)$ . The choice of  $\theta$  as an irrational number and  $[0, (1-\theta))$  being equipped with the Lebesgue measure gives that the set  $\{T_x\}_{x\in\mathbb{Z}}$  constructs a vertex-transitive group of measure-preserving ergodic transformations similar to almost-Mathieu in Theorem 2.6 (applying Lemmas A.1-A.3). Therefore,  $H_r(b)$  is an ergodic random operator.