

Adiabatic versus instantaneous transitions from a harmonic oscillator to an inverted oscillator

Viktor V. Dodonov and Alexandre V. Dodonov

Institute of Physics and International Center of Physics, University of Brasilia, Brasilia, DF, Brazil

Correspondence: vicdod@gmail.com, adodonov@unb.br

1 Introduction

We consider a quantum system described by means of one of the simplest examples of Hamiltonians with time-dependent parameters:

$$H = p^2/2 + \gamma(t)x^2/2. \quad (1)$$

For the sake of simplicity, we assume the unit value of particle's mass. One of many brilliant results of classical and quantum mechanics is the existence of *adiabatic invariants* in the case when parameters of a system vary slowly with time. If $\gamma(t) = \omega^2(t)$ with a *real* frequency $\omega(t)$, the well known adiabatic invariant is the ratio of the energy of a harmonic oscillator $\mathcal{E}(t)$ to its time-dependent frequency $\omega(t)$ [1]:

$$\mathcal{E}(t)/\omega(t) = \text{const}. \quad (2)$$

The condition of validity of Eq. (2) can be written as

$$\dot{\gamma}^2/|\gamma(t)|^3 \ll 1. \quad (3)$$

This condition is obviously broken, if function $\gamma(t)$ passes through zero value. Recently [2], we studied the case when this function always remains non-negative, even after attaining zero values. In particular, it was shown that formula (2) remains valid after the frequency crosses zero value, but the value of the constant in the right-hand side is different from the value before the crossing. If function $\gamma(t)$ behaves as some power function ($\sim t^n$) in the vicinity of time instant when $\gamma(t) = 0$, this new value depends on the power n in the law of crossing zero.

Now, our main goal was to study, what can happen if function $\gamma(t)$ slowly passes through zero value and becomes *negative*. This means that the initial *harmonic oscillator* becomes an *inverted oscillator*. Note that Schwabl and Thirring [3] and especially Glauber [4, 5] used the Hamiltonian $H = -p^2/2 - \omega^2 x^2/2$ (or $H = -\hbar\omega a^\dagger a$) as a simple model of a quantum amplifier. In order to avoid a confusion, we call this model as a “totally inverted oscillator”. We consider here the “partially inverted oscillator”, with the positive term $p^2/2$ in the Hamiltonian, using a short name “inverted oscillator”.

The history of studies of the inverted oscillator in the quantum regime (called also sometimes as an “upside-down oscillator”, or “repulsive oscillator”, or “reversed oscillator”) is surprisingly long [6–78] (the “totally inverted oscillator” was considered in [79–81]). However, it seems that the energy evolution in the adiabatic regime was not considered in all cited papers. Another interesting (and exactly solvable) case corresponds to a sudden jump of function $\gamma(t)$ at $t = 0$ from the value ω_0^2 to some constant negative value $-\kappa^2$. An instantaneous frequency jump between two positive values of coefficient γ was studied in numerous papers [82–92]. However, the jump to the negative value of γ was not considered before. Therefore, we hope that the results of our study can be interesting for many readers.

2 Evolution of the mean oscillator energy

We assume that function $\gamma(t)$ equals zero at $t = 0$. Then, the initial conditions can be chosen at some negative value $t = -\tau$ of the time variable. If the evolution is governed by Hamiltonian (1), mean values of the coordinate and momentum can be expressed in terms of two fundamental solutions of the classical equation

$$\ddot{\varepsilon} + \gamma(t)\varepsilon = 0. \quad (4)$$

We suppose that $\gamma(t) = \omega_0^2 = \text{const}$ for $t \leq -\tau$, with $\omega_0 > 0$. If $\gamma(t)$ is a real function, it is convenient, following the seminal papers [93–96] (see also, e.g., a recent review [97]), to choose the fundamental

solutions in the form of mutually conjugated complex functions $\varepsilon(t)$ and $\varepsilon^*(t)$, satisfying the initial conditions

$$\varepsilon(-\tau) = \omega_0^{-1/2}, \quad \dot{\varepsilon}(-\tau) = i\omega_0^{1/2}, \quad (5)$$

so that the Wronskian identity for the solutions $\varepsilon(t)$ and $\varepsilon^*(t)$ has the form

$$\dot{\varepsilon}(t)\varepsilon^*(t) - \dot{\varepsilon}^*(t)\varepsilon(t) \equiv 2i. \quad (6)$$

Then, we can write at $t \geq -\tau$ the following expressions for the first-order mean values (which have the same form in the classical and quantum cases):

$$x(t) = x_0\sqrt{\omega_0} \operatorname{Re}[\varepsilon(t)] + \frac{p_0}{\sqrt{\omega_0}} \operatorname{Im}[\varepsilon(t)], \quad p(t) = x_0\sqrt{\omega_0} \operatorname{Re}[\dot{\varepsilon}(t)] + \frac{p_0}{\sqrt{\omega_0}} \operatorname{Im}[\dot{\varepsilon}(t)], \quad (7)$$

where $x_0 = x(-\tau)$ and $p_0 = p(-\tau)$.

The second-order moments of the canonical operators evolve at $t \geq -\tau$ as follows:

$$\langle x^2 \rangle_t = \langle x^2 \rangle_0 \omega_0 (\operatorname{Re}[\varepsilon_t])^2 + \frac{\langle p^2 \rangle_0}{\omega_0} (\operatorname{Im}[\varepsilon_t])^2 + \langle xp + px \rangle_0 (\operatorname{Re}[\varepsilon_t]) (\operatorname{Im}[\varepsilon_t]), \quad (8)$$

$$\langle p^2 \rangle_t = \langle x^2 \rangle_0 \omega_0 (\operatorname{Re}[\dot{\varepsilon}_t])^2 + \frac{\langle p^2 \rangle_0}{\omega_0} (\operatorname{Im}[\dot{\varepsilon}_t])^2 + \langle xp + px \rangle_0 (\operatorname{Re}[\dot{\varepsilon}_t]) (\operatorname{Im}[\dot{\varepsilon}_t]). \quad (9)$$

$$\langle xp + px \rangle_t = 2\langle x^2 \rangle_0 \omega_0 \operatorname{Re}[\varepsilon_t] \operatorname{Re}[\dot{\varepsilon}_t] + \frac{2\langle p^2 \rangle_0}{\omega_0} \operatorname{Im}[\varepsilon_t] \operatorname{Im}[\dot{\varepsilon}_t] + \langle xp + px \rangle_0 \operatorname{Im}[\dot{\varepsilon}_t \varepsilon_t]. \quad (10)$$

These expressions can be easily obtained, if one considers $x(t)$ and $p(t)$ as the Heisenberg operators and calculates the average values of their powers and products over the initial state. Such an approach works well for any quadratic Hamiltonian.

The time-dependent mean energy is given by the formula

$$\mathcal{E}(t) = \frac{1}{2} [\langle p^2 \rangle_t + \gamma(t) \langle x^2 \rangle_t]. \quad (11)$$

It is worth remembering that for systems with *quadratic* Hamiltonians with respect to x and p , the dynamics of the first-order mean values $\langle x \rangle$ and $\langle p \rangle$ are *totally independent* from the dynamics of the variances $\sigma_x = \langle x^2 \rangle - \langle x \rangle^2$, $\sigma_p = \langle p^2 \rangle - \langle p \rangle^2$ and $\sigma_{xp} = \langle xp + px \rangle / 2 - \langle x \rangle \langle p \rangle$. This means that the equations of the same form as (8) and (9) exist for the sets $(\langle x^2 \rangle, \langle p^2 \rangle, \langle x \rangle \langle p \rangle)$ and $(\sigma_x, \sigma_p, \sigma_{xp})$.

2.1 Adiabatic evolution if the oscillator remains harmonic

The adiabatic (quasiclassical) approximate complex solution to Eq. (4), satisfying the initial conditions (5), has the following form at $t < 0$ [when $\gamma(t) > 0$ and $\omega(t) > 0$]:

$$\varepsilon(t) \approx [\omega(t)]^{-1/2} e^{i\phi_\tau(t)}, \quad \dot{\varepsilon}(t) \approx i[\omega(t)]^{1/2} e^{i\phi_\tau(t)}, \quad \phi_\tau(t) = \int_{-\tau}^t \omega(z) dz. \quad (12)$$

Putting the solution (12) in the equations (8)-(11). we arrive immediately at the adiabatic invariant

$$\mathcal{E}(t)/\omega(t) = \mathcal{E}(-\tau)/\omega_0, \quad (13)$$

for arbitrary initial values at $t = -\tau$.

The solution (12) obviously loses its sense if $\omega(t) = 0$ at some time instant t_* (taken as $t = 0$ in our paper). However, if the function $\gamma(t)$ slowly passes through zero value and becomes positive (and not too small) again, the condition of validity of the quasiclassical approximation (3) is reestablished again. If $\gamma(t) = \omega^2(t) > 0$ for $t > 0$, the solution for $t > 0$ (outside some interval near $t = 0$) can be written as [2]

$$\varepsilon(t) \approx [\omega(t)]^{-1/2} [u_+ e^{i\phi(t)} + u_- e^{-i\phi(t)}], \quad \dot{\varepsilon}(t) \approx i[\omega(t)]^{1/2} [u_+ e^{i\phi(t)} - u_- e^{-i\phi(t)}], \quad (14)$$

where

$$\phi(t) = \int_0^t \omega(z) dz, \quad d\phi(t)/dt = \omega(t). \quad (15)$$

Constant complex coefficients u_{\pm} must obey the condition

$$|u_+|^2 - |u_-|^2 = 1, \quad (16)$$

which is the consequence of Eq. (6). Then, Eq. (11) assumes the form [2]

$$\frac{\langle \mathcal{E} \rangle_t}{\langle \mathcal{E} \rangle_{-\tau}} = \frac{\omega(t)}{\omega_0} (\beta + \Delta\beta), \quad (17)$$

where

$$\beta = |u_+|^2 + |u_-|^2 = 1 + 2|u_-|^2, \quad (18)$$

$$\Delta\beta = \{ [\omega_0^2 \langle x^2 \rangle_{-\tau} - \langle p^2 \rangle_{-\tau}] \operatorname{Re}(u_+ u_-) + \omega_0 \langle xp + px \rangle_{-\tau} \operatorname{Im}(u_+ u_-) \} / \langle \mathcal{E} \rangle_{-\tau}. \quad (19)$$

Eq. (17) can be interpreted as a generalized adiabatic formula for the energy after the frequency passes *slowly* through zero value and remains real. It shows that the quantum mechanical mean energy is proportional to the instant frequency $\omega(t)$ in the adiabatic regime. However, the proportionality coefficient strongly depends on the initial conditions in the most general case. For this reason, we pay special attention to the case when

$$\langle p^2 \rangle_{-\tau} = \omega_0^2 \langle x^2 \rangle_{-\tau}, \quad \langle xp + px \rangle_{-\tau} = 0. \quad (20)$$

It includes the vacuum, thermal and Fock initial quantum states. Then, $\Delta\beta = 0$. In addition, many formulas can be simplified:

$$\begin{aligned} \langle x^2 \rangle_t &= \omega_0 \langle x^2 \rangle_{-\tau} |\varepsilon(t)|^2, & \langle p^2 \rangle_t &= \langle p^2 \rangle_{-\tau} |\dot{\varepsilon}(t)|^2 / \omega_0, & \langle xp + px \rangle_t &= 2 \langle x^2 \rangle_{-\tau} \omega_0 \operatorname{Re}[\dot{\varepsilon}(t) \varepsilon^*(t)], \\ \langle \mathcal{E} \rangle_t &= \frac{\langle \mathcal{E} \rangle_{-\tau}}{2\omega_0} [\gamma(t) |\varepsilon(t)|^2 + |\dot{\varepsilon}(t)|^2]. \end{aligned} \quad (21)$$

Note that the combination

$$D \equiv \langle x^2 \rangle \langle p^2 \rangle - \langle xp + px \rangle^2 / 4 \quad (22)$$

does not depend on time due to identity (6). This is the simplest example of *universal quantum invariants*, which hold for all systems with quadratic Hamiltonians [98].

In the adiabatic regime, the second-order moments (8)-(10) can be written in terms of coefficients u_{\pm} as follows [in the case of the special initial conditions (20)],

$$\begin{aligned} \langle x^2 \rangle_t &= \langle x^2 \rangle_{-\tau} \frac{\omega_0}{\omega(t)} \left[|u_+|^2 + |u_-|^2 + 2 \operatorname{Re}(u_+ u_-^* e^{2i\phi(t)}) \right], \\ \langle p^2 \rangle_t &= \langle p^2 \rangle_{-\tau} \frac{\omega(t)}{\omega_0} \left[|u_+|^2 + |u_-|^2 - 2 \operatorname{Re}(u_+ u_-^* e^{2i\phi(t)}) \right], \\ \langle xp + px \rangle_t &= -4 \langle x^2 \rangle_{-\tau} \omega_0 \operatorname{Im}(u_+ u_-^* e^{2i\phi(t)}). \end{aligned}$$

Defining the “degree of squeezing” s as the ratio of the coordinate second-order moment to the variance $\hbar/(2\omega)$ in the ground state of the oscillator with the instant frequency ω (remember that we assume the unit mass), we obtain the relations

$$s_t / s_{-\tau} = |u_+|^2 + |u_-|^2 + 2 \operatorname{Re}(u_+ u_-^* e^{2i\phi(t)}), \quad (|u_+| + |u_-|)^{-2} \leq s_t / s_{-\tau} \leq (|u_+| + |u_-|)^2. \quad (23)$$

Hence, the initial ground state ($s_{-\tau} = 1$) can be squeezed when the frequency passes through zero (remaining real).

3 Exact solutions for the power profile of the frequency

The simplest function $\gamma(t)$ which changes its sign at the point $t = 0$ is the linear one: $\gamma(t) = -\omega_0^2 t/\tau$. We consider a more general power dependence:

$$\gamma(t) = \omega_0^2 \times \begin{cases} [(-t)/\tau]^n, & t \leq 0 \\ -[t/\tau]^n, & t \geq 0 \end{cases}, \quad n > 0. \quad (24)$$

For $t < 0$, making the transformation

$$\varepsilon(t) = \sqrt{|t|} Z[y(t)], \quad y(t) = g \left| \frac{t}{\tau} \right|^b, \quad (25)$$

one can reduce Eq. (4) to the Bessel equation

$$\frac{d^2 Z}{dy^2} + \frac{1}{y} \frac{dZ}{dy} + \left(1 - \frac{\nu^2}{y^2}\right) Z = 0, \quad (26)$$

with the following parameters:

$$\nu = \frac{1}{n+2} < \frac{1}{2}, \quad b = \frac{1}{2\nu}, \quad g = 2G\nu, \quad G = \omega_0\tau. \quad (27)$$

For $t > 0$, the same transformation (25) reduces Eq. (4) to the modified Bessel equation

$$\frac{d^2 Z}{dy^2} + \frac{1}{y} \frac{dZ}{dy} - \left(1 + \frac{\nu^2}{y^2}\right) Z = 0, \quad (28)$$

with the same parameters (27).

Hence, the function $\varepsilon(t)$ can be written as a superposition of the usual Bessel functions $J_\nu(y)$ and $J_{-\nu}(y)$ for $t < 0$, and a superposition of the modified Bessel functions $I_\nu(y)$ and $I_{-\nu}(y)$ for $t > 0$:

$$\varepsilon(t) = \sqrt{|t|} \times \begin{cases} \{A_- J_\nu[y(t)] + B_- J_{-\nu}[y(t)]\}, & t < 0 \\ \{A_+ I_\nu[y(t)] + B_+ I_{-\nu}[y(t)]\}, & t > 0 \end{cases}. \quad (29)$$

Constant complex coefficients A_- and B_- can be found from the initial conditions (5). Remembering that $d|t|/dt = -1$ for $t < 0$, one obtains the following equations:

$$A_- J_\nu(g) + B_- J_{-\nu}(g) = 1/\sqrt{G}, \quad A_- J'_\nu(g) + B_- J'_{-\nu}(g) = -\frac{1}{\sqrt{G}} \left(i + \frac{1}{2G}\right),$$

where $J'_{\pm\nu}(z)$ means the derivative of the Bessel function $J_{\pm\nu}(z)$ with respect to its argument z . Using the known Wronskian [99, 100]

$$J_\nu(z)J'_{-\nu}(z) - J_{-\nu}(z)J'_\nu(z) = -2 \sin(\nu\pi)/(z\pi),$$

one can obtain the following expressions:

$$A_- = -\frac{\nu\pi\sqrt{G}}{\sin(\nu\pi)} \left[J'_{-\nu}(g) + \left(i + \frac{1}{2G}\right) J_{-\nu}(g) \right], \quad B_- = \frac{\nu\pi\sqrt{G}}{\sin(\nu\pi)} \left[\left(i + \frac{1}{2G}\right) J_\nu(g) + J'_\nu(g) \right].$$

Using the known identities (see, e.g., formulas 7.2(54) and 7.2(55) in [100])

$$J_\nu(z) \pm \frac{z}{\nu} J'_\nu(z) = \frac{z}{\nu} J_{\nu\mp 1}(z), \quad (30)$$

one can simplify formulas for the coefficients A_- and B_- :

$$A_- = \frac{\nu\pi\sqrt{G}}{\sin(\nu\pi)} [J_{1-\nu}(g) - iJ_{-\nu}(g)], \quad B_- = \frac{\nu\pi\sqrt{G}}{\sin(\nu\pi)} [iJ_\nu(g) + J_{\nu-1}(g)]. \quad (31)$$

Coefficients A_+ and B_+ can be obtained from the conditions of the continuity of function $\varepsilon(t)$ and its time derivative at $t = 0$. The leading terms of the Bessel function $J_p(z)$ and the modified Bessel function $I_p(z)$ coincide at $z \rightarrow 0$:

$$J_p(z) \approx I_p(z) \approx z^p/[2^p \Gamma(p+1)], \quad z \rightarrow 0. \quad (32)$$

This means that $\sqrt{|t|}J_\nu(y) \rightarrow 0$ and $\sqrt{|t|}I_\nu(y) \rightarrow 0$ when $t \rightarrow 0$. On the other hand, the products $\sqrt{|t|}J_{-\nu}(y)$ and $\sqrt{|t|}I_{-\nu}(y)$ tend to identical finite values in this limit. Consequently, the continuity of function $\varepsilon(t)$ at $t = 0$ implies the condition $B_+ = B_-$.

The time derivative of function (29) at $t < 0$ (when $d|t|/dt = -1$) can be written with the aid of identities (30) as follows:

$$d\varepsilon/dt = \frac{y}{2\nu\sqrt{|t|}} [B_- J_{1-\nu}(y) - A_- J_{\nu-1}(y)], \quad t \leq 0. \quad (33)$$

On the other hand, using the special cases of Eqs. 7.11(19) and 7.11(20) from [100],

$$I_\nu(z) \pm \frac{z}{\nu} I'_\nu(z) = \pm \frac{z}{\nu} I_{\nu \mp 1}(z), \quad (34)$$

we can write

$$d\varepsilon/dt = \frac{y}{2\nu\sqrt{t}} [A_+ I_{\nu-1}(y) + B_+ I_{1-\nu}(y)], \quad t \geq 0. \quad (35)$$

Now, we notice that $yJ_{1-\nu}(y)/\sqrt{|t|} \rightarrow 0$ and $yI_{1-\nu}(y)/\sqrt{|t|} \rightarrow 0$ at $t \rightarrow 0$, while the products $yJ_{\nu-1}(y)/\sqrt{|t|}$ and $yI_{\nu-1}(y)/\sqrt{|t|}$ tend to identical finite values in this limit. Hence, the continuity of derivative $d\varepsilon/dt$ at $t = 0$ results in the condition $A_+ = -A_-$. Then, one can verify that the Wronskian identity (6) is satisfied both for $t \leq 0$ and $t \geq 0$, in view of the relations [100]

$$J_\nu(z)J_{1-\nu}(z) + J_{-\nu}(z)J_{\nu-1}(z) = 2\sin(\nu\pi)/(z\pi), \quad (36)$$

$$I_\nu(z)I_{1-\nu}(z) - I_{-\nu}(z)I_{\nu-1}(z) = -2\sin(\nu\pi)/(z\pi). \quad (37)$$

Using Eqs. (29) and (31), one can write down formula (21) for the mean energy ratio $\mathcal{R}(t) = \mathcal{E}(t)/\mathcal{E}(-\tau)$ as follows:

$$\mathcal{R}(t < 0) = \frac{1}{8} \left[\frac{g\pi}{\sin(\nu\pi)} \right]^2 \left| \frac{t}{\tau} \right|^{n+1} [K_-(g)K_+(y) + K_+(g)K_-(y) - 2K_0(g)K_0(y)], \quad (38)$$

$$\mathcal{R}(t > 0) = \frac{1}{8} \left[\frac{g\pi}{\sin(\nu\pi)} \right]^2 \left| \frac{t}{\tau} \right|^{n+1} [K_-(g)\tilde{K}_+(y) + K_+(g)\tilde{K}_-(y) - 2K_0(g)\tilde{K}_0(y)], \quad (39)$$

where

$$K_+(z) = J_{\nu-1}^2(z) + J_\nu^2(z), \quad K_-(z) = J_{1-\nu}^2(z) + J_{-\nu}^2(z),$$

$$\tilde{K}_+(z) = I_{\nu-1}^2(z) - I_\nu^2(z), \quad \tilde{K}_-(z) = I_{1-\nu}^2(z) - I_{-\nu}^2(z),$$

$$K_0(z) = J_{\nu-1}(z)J_{1-\nu}(z) - J_\nu(z)J_{-\nu}(z).$$

$$\tilde{K}_0(z) = I_{\nu-1}(z)I_{1-\nu}(z) - I_\nu(z)I_{-\nu}(z).$$

3.1 Adiabatic evolution of energy at $t < 0$

If $t < 0$, the known leading term of the asymptotic formula for the Bessel functions of large arguments [99, 100],

$$J_\nu(z) \sim \sqrt{\frac{2}{\pi z}} \cos\left(z - \frac{\nu\pi}{2} - \frac{\pi}{4}\right), \quad (40)$$

results in the following simple expressions at $z \gg 1$ (with corrections of the order of z^{-2}):

$$K_\pm(z) = \frac{2}{\pi z}, \quad K_0(z) = \frac{2\cos(\nu\pi)}{\pi z}. \quad (41)$$

Hence, in the adiabatic limit ($g \gg 1$ and $y \gg 1$), we obtain

$$\mathcal{R}(t < 0) = \frac{\omega(t) [1 - \cos^2(\nu\pi)]}{\omega_0 \sin^2(\nu\pi)} = \frac{\omega(t)}{\omega_0}. \quad (42)$$

This adiabatic formula holds for any value of the power n if $t < 0$.

In view of formula (32), the only nonzero contribution to the right-hand side of Eq. (38) at $t \rightarrow 0$ is given by the function $K_+(y) \approx J_{\nu-1}^2(y) \approx [(y/2)^{\nu-1}/\Gamma(\nu)]^2 \sim |t|^{-(n+1)}$, whereas the contributions of $K_-(y) \sim |t|^{-1}$ and $K_0(y) \sim |t|^0$ are eliminated by the term $|t|^{(n+1)}$. Hence,

$$\mathcal{R}(t = 0) = \frac{\pi g^{2\nu-1}}{[2^\nu \Gamma(\nu) \sin(\pi\nu)]^2}, \quad g \gg 1. \quad (43)$$

This means, in particular, that the adiabatic formula (42) holds under the condition $\omega(t)/\omega_0 \gg g^{2\nu-1}$ (provided $g = 2\nu\omega_0\tau \gg 1$). Remember that $2\nu - 1 < 0$.

3.2 The harmonic oscillator revival

The case when function $\gamma(t)$ returns to positive values $\omega^2(t)$ for $t > 0$ (when it is given by the expression in the second line of Eq. (24) with the positive sign) was studied in Ref. [2]. In this case, the energy ratio in the adiabatic regime is given by the formula

$$\mathcal{R}(t > 0) = \frac{\omega(t) [1 + \cos^2(\nu\pi)]}{\omega_0 \sin^2(\nu\pi)}. \quad (44)$$

Constant coefficients u_\pm depend on parameter ν as follows

$$u_+ = [\sin(\nu\pi)]^{-1}, \quad u_- = i \cot(\nu\pi).$$

Consequently, the squeezing coefficient satisfies the inequalities

$$\tan^2(\nu\pi/2) \leq s_t/s_{-\tau} \leq \cot^2(\nu\pi/2).$$

4 Evolution of the inverted oscillator energy

4.1 Asymptotic evolution with the power function $\gamma(t)$

If the oscillator becomes inverted with a slowly varying function $\gamma(t)$, given by Eq. (24) under the condition $\omega_0\tau \gg 1$, we need asymptotic formulas for the modified Bessel functions of large arguments. Remembering that function $I_\nu(z)$ is real for real values of its argument, we take the arithmetic average value (or the real part) between two equivalent expressions in formula 8.451.5 from the reference book [99]:

$$I_\nu(z) \sim \frac{e^z}{(2\pi z)^{1/2}} \sum_{k=0}^{\infty} \frac{\Gamma(\nu + k + 1/2)}{(2z)^k k! \Gamma(\nu - k + 1/2)} [(-1)^k - \sin(\nu\pi) e^{-2z}]. \quad (45)$$

Taking into account only terms with $k = 0$, one would obtain slowly varying (and incorrect) expressions for the functions $\tilde{K}_\pm(z)$ and $\tilde{K}_0(z)$, of the order of z^{-1} . To obtain correct asymptotic expressions for these functions, one has to take into account the higher order terms with $k = 1$ in Eq. (45). Then, the exponentially small corrections containing e^{-2z} can be neglected if $z \gg 1$. The correct expressions are as follows,

$$\tilde{K}_\pm(z) \approx \tilde{K}_0(z) \approx \frac{2\nu - 1}{2\pi z^2} e^{2z}. \quad (46)$$

Hence, the absolute value of the mean energy grows exponentially in the asymptotic adiabatic regime:

$$\mathcal{R}(t > 0) \approx \frac{(2\nu - 1) \exp[2y(t)]}{8g(t/\tau) \cos^2(\nu\pi/2)}, \quad g \gg 1, \quad y(t) = g \left(\frac{t}{\tau}\right)^{1+n/2} \gg 1. \quad (47)$$

Since $2\nu - 1 < 0$, this asymptotic time dependent ratio is always negative (for any value of parameter n).

4.2 If function $\gamma(t)$ becomes negative and constant

If $\gamma(t) = -\kappa^2$ with $\kappa = \text{const} > 0$ for $t > t_* > 0$, Eq. (4) has an exact solution:

$$\varepsilon(t) = [2\kappa]^{-1/2} [v_+ e^{\kappa t} + v_- e^{-\kappa t}], \quad \dot{\varepsilon}(t) = [\kappa/2]^{1/2} [v_+ e^{\kappa t} - v_- e^{-\kappa t}]. \quad (48)$$

In view of Eq. (6), constant complex coefficients v_{\pm} must satisfy the condition

$$\text{Im}(v_+ v_-^*) = 1. \quad (49)$$

In this case, the time-independent mean energy equals

$$\langle \mathcal{E} \rangle(t > t_*) = -\kappa \left[\langle x^2 \rangle_{-\tau} \omega_0 \text{Re}(v_+) \text{Re}(v_-) + \frac{\langle p^2 \rangle_{-\tau}}{\omega_0} \text{Im}(v_+) \text{Im}(v_-) + \frac{1}{2} \langle xp + px \rangle_{-\tau} \text{Im}(v_+ v_-) \right]. \quad (50)$$

All the second-order moments grow unlimitedly with time. Under the conditions (20), we can write

$$\begin{aligned} \langle x^2 \rangle_t &= \langle x^2 \rangle_{-\tau} \frac{\omega_0}{2\kappa} [|v_+|^2 e^{2\kappa t} + |v_-|^2 e^{-2\kappa t} + 2\text{Re}(v_+ v_-^*)], \\ \langle p^2 \rangle_t &= \langle p^2 \rangle_{-\tau} \frac{\kappa}{2\omega_0} [|v_+|^2 e^{2\kappa t} + |v_-|^2 e^{-2\kappa t} - 2\text{Re}(v_+ v_-^*)], \\ \langle xp + px \rangle_t &= \langle x^2 \rangle_{-\tau} \omega_0 [|v_+|^2 e^{2\kappa t} - |v_-|^2 e^{-2\kappa t}]. \end{aligned}$$

Nonetheless, the exponentially increasing terms are canceled in the formula for the mean energy, which does not depend on time if $\kappa = \text{const}$. For the initial N th Fock state, we obtain

$$\langle \mathcal{E} \rangle = -\kappa \text{Re}(v_+ v_-^*) (N + 1/2).$$

4.3 Instantaneous jump to the inverted oscillator

The coefficients v_{\pm} can be easily calculated in the case of a sudden jump of function $\gamma(t)$ at $t = 0$ from the value ω_0^2 to the constant value $-\kappa^2$, using the conditions of continuity of functions $\varepsilon(t)$ and $\dot{\varepsilon}(t)$:

$$v_{\pm} = \frac{1}{\sqrt{2}} \left(\sqrt{\frac{\kappa}{\omega_0}} \pm i \sqrt{\frac{\omega_0}{\kappa}} \right), \quad v_+ v_-^* = i + \frac{1}{2} \left(\frac{\kappa}{\omega_0} - \frac{\omega_0}{\kappa} \right). \quad (51)$$

Then, formula (50) yields the following mean energy after the jump for any initial state:

$$\langle \mathcal{E} \rangle_{t>0} = \frac{1}{2} [\langle p^2 \rangle_{-\tau} - \kappa^2 \langle x^2 \rangle_{-\tau}]. \quad (52)$$

This result can seem obvious, because the wave function and its mean values do not change during the instantaneous transformation of the Hamiltonian. However, it is interesting, because the mean energy after the jump can assume any value, depending on the ratio κ/ω_0 and initial conditions. In particular, under the initial conditions (20), the mean energy turns into zero if $\kappa = \omega_0$. For the initial N -th Fock state we have

$$\langle \mathcal{E} \rangle_{t>0}^{Fock} = \frac{\hbar\omega_0}{4} (2N + 1) \left(1 - \frac{\kappa^2}{\omega_0^2} \right). \quad (53)$$

5 Energy fluctuations

It can be interesting to know the strength of the *energy fluctuations* after the frequency passes through zero. These fluctuations can be characterized by the variance $\sigma_E = \langle E^2 \rangle - \langle E \rangle^2$. Using the solutions (7) of the Heisenberg equations of motion, one can write σ_E in terms of the initial fourth- and second-order moments of the canonical variables x and p and various products of functions $\varepsilon(t)$, $\dot{\varepsilon}(t)$ and their complex conjugated partners. The complete formula is rather cumbersome in the most general case. For this reason, we consider here (following [2]) the case of the initial Fock quantum state $|N\rangle$. In this special case (as well as for arbitrary diagonal mixtures of the Fock states), the nonzero statistical moments are those containing *even powers* of each variable, x or p . After some algebra, one can obtain the following formula (using dimensionless variables and assuming $\hbar = m = \omega_0 = 1$, so that $\langle x^4 \rangle = \langle p^4 \rangle$) [2]:

$$16\langle E^2 \rangle_t = 2\langle x^4 \rangle_{-\tau} (A^2 + B^2) + \langle x^2 p^2 + p^2 x^2 \rangle_{-\tau} (A^2 - B^2) + \langle (xp + px)^2 \rangle_{-\tau} C^2,$$

where

$$A(t) = \gamma(t) |\varepsilon(t)|^2 + |\dot{\varepsilon}(t)|^2, \quad B(t) = \text{Re} [\gamma(t) \varepsilon^2(t) + \dot{\varepsilon}^2(t)], \quad C(t) = \text{Im} [\gamma(t) \varepsilon^2(t) + \dot{\varepsilon}^2(t)].$$

5.1 The case of harmonic oscillator at $t > 0$

In the adiabatic regime (14) we have

$$A = 2\omega(t) (|u_+|^2 + |u_-|^2), \quad B = 4\omega(t)\text{Re}(u_+u_-), \quad C = 4\omega(t)\text{Im}(u_+u_-). \quad (54)$$

For the initial Fock state $|N\rangle$ we have

$$\begin{aligned} \langle x^4 \rangle_{-\tau} &= \frac{3}{4} (2N^2 + 2N + 1), & \langle x^2 p^2 + p^2 x^2 \rangle_{-\tau} &= \frac{1}{2} (2N^2 + 2N - 1), \\ \langle (xp + px)^2 \rangle_{-\tau} &= 2(N^2 + N + 1). \end{aligned}$$

Hence,

$$\langle E^2 \rangle_t / \omega^2(t) = (|u_+|^2 + |u_-|^2)^2 (N + 1/2)^2 + 2|u_+u_-|^2 (N^2 + N + 1).$$

Remembering that the mean energy equals $\langle E \rangle_t = \omega(t) (|u_+|^2 + |u_-|^2) (N + 1/2)$, we arrive at the following formula for the energy variance:

$$\sigma_E(t) = 2\omega^2(t)|u_+u_-|^2 (N^2 + N + 1), \quad \frac{\sigma_E(t)}{\langle E \rangle_t^2} = 2|u_+u_-|^2 \frac{N^2 + N + 1}{N^2 + N + 1/4}. \quad (55)$$

In the absence of zero frequency values we have $u_- = 0$. In this case, $\sigma_E(t) \equiv 0$, in accordance with the Born–Fock adiabatic theorem. However, this theorem is broken when the frequency passes through zero value. For the initial vacuum state ($N = 0$) and the power index $n = 2$ of the single frequency transition through zero value, we obtain $\sigma_E(t)/\langle E \rangle_t^2 = 16$. This ratio can be four times smaller if $N \gg 1$.

5.2 The case of inverted oscillator

In the case of inverted oscillator with a constant parameter κ , one should replace $\omega(t)$ with $i\kappa$ and use the solution (48) instead of (14). Then, we have the following expressions instead of (54):

$$A = -2\kappa\text{Re}(v_+v_-^*), \quad B = -2\kappa\text{Re}(v_+v_-), \quad C = -2\kappa\text{Im}(v_+v_-). \quad (56)$$

In the case of a sudden jump (51), we obtain

$$A = \omega_0 \left(1 - \frac{\kappa^2}{\omega_0^2}\right), \quad B = -\omega_0 \left(1 + \frac{\kappa^2}{\omega_0^2}\right), \quad C = 0.$$

Hence,

$$\langle E^2 \rangle_{t>0}^{Fock} = \frac{1}{16} (\hbar\omega_0)^2 \left[3(2N^2 + 2N + 1) \left(1 + \frac{\kappa^4}{\omega_0^4}\right) - 2\frac{\kappa^2}{\omega_0^2} (2N^2 + 2N - 1) \right], \quad (57)$$

$$\sigma_E^{Fock}(t > 0) = \frac{1}{8} (\hbar\omega_0)^2 (N^2 + N + 1) \left(1 + \frac{\kappa^2}{\omega_0^2}\right)^2. \quad (58)$$

6 Probability density distribution over energy eigenstates of the inverted oscillator

It is interesting to calculate the probability density of measuring the energy eigenvalues of the inverted oscillator with a fixed parameter κ . The evolution of the initial Fock state $|n\rangle$ is determined completely by the function $\varepsilon(t)$ introduced in Sec. 2. The time dependent wave function has the following form [93,94,96] (we use dimensionless units with $\hbar = m = 1$ in this section):

$$\psi_n(x, t) = (n! \varepsilon \sqrt{\pi})^{-1/2} \left(\frac{\varepsilon^*}{2\varepsilon}\right)^{n/2} \exp\left(\frac{i\dot{\varepsilon}}{2\varepsilon} x^2\right) H_n\left(\frac{x}{|\varepsilon|}\right), \quad (59)$$

where $H_n(z)$ is the Hermite polynomial. The wave functions $\psi(x; E)$ of energy eigenstates are solutions of the stationary Schrödinger equation

$$d^2\psi/dx^2 + (\kappa^2 x^2 + 2E)\psi = 0. \quad (60)$$

It is known that the energy spectrum of the inverted oscillator is continuous and doubly degenerate, with $-\infty < E < \infty$. Two independent normalized solutions to Eq. (60) can be found in papers [32, 74] (we use a slightly different notation):

$$\psi_{\pm}(x; E) = \mathcal{N} D_{\mu} \left(\mp x \sqrt{-2i\kappa} \right), \quad (61)$$

where $D_{\mu}(z)$ is the parabolic cylinder function defined as in books [99, 100]. The index μ and the normalization constant \mathcal{N} are related to the normalized energy $\tilde{E} = E/\kappa$ as follows (we assume $\kappa > 0$):

$$\mu = -\frac{1}{2} + i\tilde{E}, \quad \mathcal{N} = \frac{\Gamma(-\mu) \exp(\pi\tilde{E}/4)}{\pi(8\kappa)^{1/4}}, \quad (62)$$

where $\Gamma(z)$ is the Euler gamma function. With this normalization, functions (61) obey the relations

$$\int_{-\infty}^{\infty} \psi_{\pm}^*(x; E) \psi_{\pm}(x; E') dx = \delta(E - E'). \quad (63)$$

In view of the double degeneracy of energy levels in the inverted regime, the instantaneous energy probability distribution with respect to the normalized energy can be written as follows,

$$P_n(\tilde{E}) = \kappa \left| \int_{-\infty}^{\infty} \psi_n^*(x, t) |\psi_+(x; E) dx \right|^2 + \kappa \left| \int_{-\infty}^{\infty} \psi_n^*(x, t) |\psi_-(x; E) dx \right|^2. \quad (64)$$

Hence, one has to calculate the integrals of the following structure:

$$I_{\mu, n}(a, b, c) = \int_{-\infty}^{\infty} e^{-ax^2} D_{\mu}(cx) H_n(bx) dx. \quad (65)$$

Although explicit expressions for the integral (65) are absent in the main reference books, such as [99–101], it can be calculated analytically with the aid of the integral representation of the parabolic cylinder function (see formula 9.241.2 from [99]),

$$D_{\mu}(z) = \frac{e^{-z^2/4}}{\Gamma(-\mu)} \int_0^{\infty} \exp(-zy - y^2/2) y^{-\mu-1} dy. \quad (66)$$

Putting the integral (66) in Eq. (65) and changing the order of integrations, we arrive at the integral

$$\int_{-\infty}^{\infty} \exp[-(a + c^2/4)x^2 - ycx] H_n(bx) dx,$$

which can be calculated with the aid of formula 7.374.8 from [99],

$$\int_{-\infty}^{\infty} e^{-(\xi-\eta)^2} H_n(\alpha\xi) d\xi = \sqrt{\pi} (1 - \alpha^2)^{n/2} H_n \left(\frac{\alpha\eta}{\sqrt{1 - \alpha^2}} \right), \quad (67)$$

with

$$\alpha = \frac{b}{\sqrt{a + c^2/4}}, \quad \eta = -\frac{yc}{2\sqrt{a + c^2/4}}.$$

Hence, we can write

$$I_{\mu, n} = \frac{(-1)^n \sqrt{\pi} (a + c^2/4 - b^2)^{n/2}}{\Gamma(-\mu) (a + c^2/4)^{(n+1)/2}} \int_0^{\infty} \exp(-By^2) H_n(Ay) y^{-\mu-1} dy, \quad (68)$$

where

$$A = \frac{bc/2}{\sqrt{(a + c^2/4)(a + c^2/4 - b^2)}}, \quad B = \frac{a - c^2/4}{2(a + c^2/4)}.$$

The value of the integral in (68) depends on the parity of the Hermite polynomials, according to formulas 7.376.2 and 7.376.3 from [99]:

$$\int_0^{\infty} e^{-2\gamma x^2} x^{\nu} H_{2k}(x) dx = (-1)^k 2^{2k - \frac{3}{2} - \frac{\nu}{2}} \frac{\Gamma(\frac{\nu+1}{2}) \Gamma(k + \frac{1}{2})}{\sqrt{\pi\gamma} \frac{\nu+1}{2}} F \left(-k, \frac{\nu+1}{2}; \frac{1}{2}; \frac{1}{2\gamma} \right),$$

$$\int_0^\infty e^{-2\gamma x^2} x^\nu H_{2k+1}(x) dx = (-1)^k 2^{2k-\frac{\nu}{2}} \frac{\Gamma(\frac{\nu}{2}+1) \Gamma(k+\frac{3}{2})}{\sqrt{\pi}\gamma^{\frac{\nu}{2}+1}} F\left(-k, \frac{\nu}{2}+1; \frac{3}{2}; \frac{1}{2\gamma}\right),$$

where $F(a, b; c; z)$ is the Gauss hypergeometric function. In our case, we have

$$\nu = -\mu - 1 = -\frac{1}{2} - i\tilde{E}, \quad 2\gamma = B/A^2 = 2(a - c^2/4)(a + c^2/4 - b^2)/(bc)^2.$$

If $\varepsilon(t)$ is given by Eq. (48), we have

$$a = \frac{i\tilde{\varepsilon}^*}{2\varepsilon^*} = \frac{i\kappa v_+^* e^{\kappa t} - v_-^* e^{-\kappa t}}{2 v_+^* e^{\kappa t} + v_-^* e^{-\kappa t}}, \quad b = 1/|\varepsilon| = \frac{\sqrt{2\kappa}}{|v_+ e^{\kappa t} + v_- e^{-\kappa t}|}, \quad c^2/4 = -i\kappa/2,$$

$$a - c^2/4 = \frac{i\kappa v_+^* e^{\kappa t}}{v_+^* e^{\kappa t} + v_-^* e^{-\kappa t}}, \quad a + c^2/4 = -\frac{i\kappa v_-^* e^{-\kappa t}}{v_+^* e^{\kappa t} + v_-^* e^{-\kappa t}}.$$

Hereafter, we consider the special case of a sudden jump, with coefficients (51). Since the energy distribution cannot depend on time for the stationary Hamiltonian at $t > 0$, the coefficients a and b can be taken at the initial instant $t = 0+$. Hence, we can use more simple expressions

$$a = \frac{i\kappa v_+^* - v_-^*}{2 v_+^* + v_-^*} = \omega_0/2, \quad b = \frac{\sqrt{2\kappa}}{|v_+ + v_-|} = \sqrt{\omega_0},$$

$$a \pm c^2/4 = \frac{1}{2}(\omega_0 \mp i\kappa), \quad a + c^2/4 - b^2 = -\frac{1}{2}(\omega_0 + i\kappa),$$

so that

$$2\gamma = \frac{(\omega_0 + i\kappa)^2}{4i\kappa\omega_0} = \frac{(1 + i\rho)^2}{4i\rho}, \quad \rho \equiv \kappa/\omega_0.$$

Final formulas for the energy probability densities after the jump are as follows,

$$P_{2k}(\tilde{E}; \rho) = \frac{\Gamma(k + \frac{1}{2})\rho^{1/2}}{2\pi^2 k! (1 + \rho^2)^{1/2}} e^{2\tilde{E}\Phi(\rho)} \left| \Gamma\left(\frac{1}{4} - i\frac{\tilde{E}}{2}\right) F\left(-k, \frac{1}{4} - i\frac{\tilde{E}}{2}; \frac{1}{2}; z(\rho)\right) \right|^2, \quad (69)$$

$$P_{2k+1}(\tilde{E}; \rho) = \frac{8\Gamma(k + \frac{3}{2})\rho^{3/2}}{\pi^2 k! (1 + \rho^2)^{3/2}} e^{2\tilde{E}\Phi(\rho)} \left| \Gamma\left(\frac{3}{4} - i\frac{\tilde{E}}{2}\right) F\left(-k, \frac{3}{4} - i\frac{\tilde{E}}{2}; \frac{3}{2}; z(\rho)\right) \right|^2, \quad (70)$$

$$\Phi(\rho) = \frac{\pi}{4} - \tan^{-1}(\rho) = \tan^{-1}\left(\frac{1-\rho}{1+\rho}\right), \quad z(\rho) = \frac{4i\rho}{(1+i\rho)^2}.$$

If $\rho = 1$, the formulas can be simplified as follows,

$$P_{2k}(\tilde{E}) = \frac{\Gamma(k + 1/2)}{2^{3/2}\pi^2 k!} \left| \Gamma\left(\frac{1}{4} - i\frac{\tilde{E}}{2}\right) F\left(-k, \frac{1}{4} - i\frac{\tilde{E}}{2}; \frac{1}{2}; 2\right) \right|^2, \quad (71)$$

$$P_{2k+1}(\tilde{E}) = \frac{2^{3/2}\Gamma(k + 3/2)}{\pi^2 k!} \left| \Gamma\left(\frac{3}{4} - i\frac{\tilde{E}}{2}\right) F\left(-k, \frac{3}{4} - i\frac{\tilde{E}}{2}; \frac{3}{2}; 2\right) \right|^2. \quad (72)$$

We used the known formula $\Gamma(2z) = 2^{2z-1}\pi^{-1/2}\Gamma(z)\Gamma(z+1/2)$. All probability distributions (71) and (72) are even functions of energy, because the mean energy after the jump with $\rho = 1$ equals zero. In particular, for $k = 0$ we have

$$P_0(\tilde{E}) = (2\pi)^{-3/2} \left| \Gamma\left(\frac{1}{4} - i\frac{\tilde{E}}{2}\right) \right|^2, \quad P_1(\tilde{E}) = \left(\frac{2}{\pi^3}\right)^{1/2} \left| \Gamma\left(\frac{3}{4} - i\frac{\tilde{E}}{2}\right) \right|^2. \quad (73)$$

In these special cases, the correct normalization $\int_{-\infty}^{\infty} P(\tilde{E}) d\tilde{E} = 1$ can be checked with the aid of formula 2.2.4.1 from [101],

$$\int_0^\infty |\Gamma(a + ix)|^2 dx = 2^{-2a} \pi \Gamma(2a). \quad (74)$$

Using the relations

$$(a + ix)\Gamma(a + ix) = \Gamma(a + 1 + ix), \quad (a^2 + x^2) |\Gamma(a + ix)|^2 = |\Gamma(a + 1 + ix)|^2,$$

we obtain the integral

$$\int_0^\infty x^2 |\Gamma(a + ix)|^2 dx = 2^{-2a-1} \pi a \Gamma(2a). \quad (75)$$

Hence, writing $P(\tilde{E}) = N_a |\Gamma(a + i\tilde{E}/2)|^2$, we obtain the following formula for the mean value $\langle \tilde{E}^2 \rangle$ with respect to the probability densities $P_0(\tilde{E})$ and $P_1(\tilde{E})$: $\langle \tilde{E}^2 \rangle = 16\pi a N_a 2^{-2a-1} \Gamma(2a)$. For $a = 1/4$ we have $\langle \tilde{E}^2 \rangle = 1/2$, while for $a = 3/4$ we have $\langle \tilde{E}^2 \rangle = 3/2$. These values coincide exactly with those given by Eq. (57) for $N = 0$ and $N = 1$ (provided $\kappa = \omega_0$).

For $k = 1$ and $k = 2$, we obtain the following expressions:

$$P_2(\tilde{E}) = (2\pi^3)^{-1/2} \tilde{E}^2 \left| \Gamma\left(\frac{1}{4} - i\frac{\tilde{E}}{2}\right) \right|^2, \quad P_3(\tilde{E}) = \left(\frac{2}{\pi}\right)^{3/2} \frac{\tilde{E}^2}{3} \left| \Gamma\left(\frac{3}{4} - i\frac{\tilde{E}}{2}\right) \right|^2.$$

$$P_4(\tilde{E}) = \frac{(1 - 2\tilde{E}^2)^2}{6(2\pi)^{3/2}} \left| \Gamma\left(\frac{1}{4} - i\frac{\tilde{E}}{2}\right) \right|^2, \quad P_5(\tilde{E}) = \frac{6}{5(2\pi)^{3/2}} \left(1 - \frac{2}{3}\tilde{E}^2\right)^2 \left| \Gamma\left(\frac{3}{4} - i\frac{\tilde{E}}{2}\right) \right|^2.$$




Figure 1: Functions $P_0(\tilde{E})$, $P_4(\tilde{E})$ and $P_8(\tilde{E})$ after the jump with $\rho = 1$.




Figure 2: Functions $P_1(\tilde{E})$, $P_5(\tilde{E})$ and $P_9(\tilde{E})$ after the jump with $\rho = 1$.

Looking at plots of functions $P_n(\tilde{E})$ in Figures 1–4 (where we consider the arguments $\tilde{E} \geq 0$ only, due to the symmetry of distributions), one can notice that distributions $P_{2k}(\tilde{E})$ and $P_{2k+1}(\tilde{E})$ behave




Figure 3: Functions $P_6(\tilde{E})$, $P_{10}(\tilde{E})$ and $P_{14}(\tilde{E})$ after the jump with $\rho = 1$.




Figure 4: Functions $P_7(\tilde{E})$, $P_{11}(\tilde{E})$ and $P_{15}(\tilde{E})$ after the jump with $\rho = 1$.

in a similar way, in spite of slightly different analytical expressions (71) and (72) for the initial even and odd Fock states. On the other hand, the parity of number k turns out to be important. If this number is even ($k = 0, 2, 4, \dots$), the probability densities at $\tilde{E} = 0$ are different from zero. In contrast, $P_{2k}(0) = P_{2k+1}(0)$ for odd values $k = 1, 3, 5, \dots$: see Figures 3 and 4. All probability distributions have $[k/2]$ zeros at positive values of argument \tilde{E} (and symmetric zeros at negative values). The heights of the first $[k/2]$ maximas slowly increase with \tilde{E} , whereas the last maximum is significantly higher (by approximately 50%) and wider. This last maximum of $P_n(\tilde{E})$ is attained when \tilde{E} is a little smaller than n . The function $P_n(\tilde{E})$ rapidly goes to zero at $\tilde{E} > n$. Its asymptotic behavior for $|\tilde{E}| > n$ can be described by the formula $P_n(\tilde{E}) \sim g_n(\tilde{E}) \exp(-\pi|\tilde{E}|/2)$, where the pre-factor $g_n(\tilde{E})$ contains some powers of $|\tilde{E}|$. This is the consequence of formula 1.18(6) from [100]: $\lim_{|y| \rightarrow \infty} |\Gamma(x + iy)|^2 e^{\pi|y|} |y|^{1-2x} = 2\pi$. Numerical calculations confirm the asymptotic formula with high accuracy.

If $\rho \neq 1$, the distributions (69) and (70) lose the symmetry with respect to the point $\tilde{E} = 0$, in accordance with formula (53). However, due to the relations

$$\Phi(1/\rho) = -\Phi(\rho), \quad z(1/\rho) = z^*(\rho)$$

we have the following reciprocity formula:

$$P_n(\tilde{E}; \rho) = P_n(-\tilde{E}; 1/\rho). \quad (76)$$

Some examples of distributions with different values of ρ are shown in Figures 5 and 6.




Figure 5: Functions $P_8(\tilde{E})$ after the jumps with $\rho = 1/2, 1, 2$.




Figure 6: Functions $P_{10}(\tilde{E})$ after the jumps with $\rho = 1/2, 1, 2$.

7 Discussion

Let us discuss briefly the main results of this work. We have obtained explicit analytical formulas for the mean energy and its variance (characterizing the energy fluctuations) in the adiabatic regimes after the frequency passes through zero. The behavior of energy turns out to be quite different in two cases: when the frequency remains real and when it becomes imaginary. In the first case, the mean energy always increases when the frequency returns to its initial value, and the increment coefficient is determined by the exponent in the power law of the frequency crossing zero. On the other hand, if the frequency becomes imaginary, the absolute value of mean energy increases exponentially, even in the adiabatic regime, unless the Hamiltonian becomes time independent. It is worth emphasizing that even small corrections to the leading terms of simple adiabatic approximate formulas are crucial in this case, due to the unstable nature of the motion.

The mean energy does not depend on time after a sudden transformation of a harmonic oscillator to an inverted oscillator with constant parameters. In this case, the mean energy can be both positive and negative (or zero), depending on the ratio of the initial and final frequencies.

Comparing the results of this study with those obtained in paper [2], we can expect that the adiabatic behavior of energy and other observables can be even more intricate when the function $\gamma(t)$ in the Hamiltonian (1) passes through zero many times, changing its signs. Probably, some kinds of quasi-chaotic behaviors could be observed. However, this complicated problem deserves a separate study.

Acknowledgments

This work received partial financial support from Brazilian funding agencies National Council for Scientific and Technological Development (CNPq) and Fundação de Apoio à Pesquisa do Distrito Federal (FAPDF), grant number 00193-00001817/2023-43.

References

- [1] L. D. Landau and E. M. Lifschitz, *Mechanics*, Mir: Moscow, 1966.
- [2] V. V. Dodonov and A. V. Dodonov, Adiabatic amplification of the harmonic oscillator energy when the frequency passes through zero, *Entropy* **25**, 2 (2023).
- [3] F. Schwabl and W. Thirring, Quantum theory of laser radiation, *Ergeb. Exakten Naturwiss.* **36**, 219-242 (1964).
- [4] R. Glauber, Quantum amplifiers, attenuators and their irreversible behaviour, in *Proceedings of the Second International Seminar on Group Theoretical Methods in Physics (Zvenigorod, USSR, 24-26 November 1982)*, edited by M. A. Markov, V. I. Man'ko and A. E. Shabad, Harwood Academic, New York, 1985, Vol. I, pp. 137-148.
- [5] R. J. Glauber, Amplifiers, attenuators, and Schrödinger's cat, *Ann. New York Acad. Sci.* **480**, 336-372 (1986).
- [6] M. Moshinsky and T. H. Seligman, Canonical transformations to action and angle variables and their representations in quantum mechanics, *Ann. Phys.* **114**, 243-272 (1978).
- [7] V. V. Dodonov and V. I. Man'ko, Coherent states and the resonance of a quantum damped oscillator, *Phys. Rev. A* **20**, 550-560 (1979).
- [8] P. G. L. Leach, $Sl(3, \mathbb{R})$ and the repulsive oscillator, *J. Phys. A: Math. Gen.* **13**, 1991-2000 (1980).
- [9] A. H. Guth and S.-Y. Pi, Quantum mechanics of the scalar field in the new inflationary universe, *Phys. Rev. D* **32**, 1899-1920 (1985).
- [10] G. Barton, Quantum mechanics of the inverted oscillator potential, *Ann. Phys.* **166**, 322-363 (1986).
- [11] W. Elberfeldt, M. Klebert, W. Becker, and M. O. Scully, Laser-assisted tunnelling, *J. Phys. B: At. Mol. Phys.* **19**, 2589-2597 (1986).
- [12] N. L. Balazs and A. Voros, Wigner's function and tunneling, *Ann. Phys.* **199**, 123-140 (1990).
- [13] G. J. Papadopoulos, Time-dependent quantum tunneling via crossover processes, *J. Phys. A: Math. Gen.* **23**, 935-947 (1990).
- [14] V. V. Dodonov and D. E. Nikonov, Application of Feynman propagators to the motion of one-dimensional wave packets, *J. Sov. Laser Res.* **12**, 461-464 (1991).
- [15] S. Baskoutas, A. Jannussis, and R. Mignani, Dissipative tunnelling of the inverted Caldirola-Kanai oscillator, *J. Phys. A: Math. Gen.* **27**, 2189-2196 (1994).
- [16] M. Kleber, Exact solutions for time-dependent phenomena in quantum mechanics, *Phys. Rep.* **236**, 331-393 (1994).
- [17] Y. Nogami and F. M. Toyama, Nonlinear Schrödinger soliton in a time-dependent quadratic potential, *Phys. Rev. E* **49**, 4497-4501 (1994).
- [18] W. H. Zurek and J. P. Paz, Decoherence, chaos, and the second law, *Phys. Rev. Lett.* **72**, 2508-2511 (1994).
- [19] A. Albrecht, P. Ferreira, M. Joyce, and T. Prokopec, Inflation and squeezed quantum states, *Phys. Rev. D* **50**, 4807-4820 (1994).

- [20] R. K. Bhaduri, A. Khare, S. M. Reimann, and E. L. Tomusiak, The Riemann Zeta function and the inverted harmonic oscillator, *Ann. Phys.* **254**, 25-40 (1997).
- [21] M. Castagnino, R. Diener, L. Lara, and G. Puccini, Rigged Hilbert spaces and time asymmetry: the case of the upside-down simple harmonic oscillator, *Int. J. Theor. Phys.* **36**, 2349-2369 (1997).
- [22] M. M. Nieto and D. R. Truax, Displacement-operator squeezed states. II. Examples of time-dependent systems having isomorphic symmetry algebras, *J. Math. Phys.* **38**, 98-114 (1997).
- [23] G. J. Papadopoulos, Microwave-assisted tunnelling in the presence of dissipation, *J. Phys. A: Math. Gen.* **30**, 5497-5510 (1997).
- [24] S. Nonnenmacher and A. Voros, Eigenstate structures around a hyperbolic point, *J. Phys. A: Math. Gen.* **30**, 295-315 (1997).
- [25] G. G. Adamian, N. V. Antonenko, and W. Scheid, Tunneling with dissipation in open quantum systems, *Phys. Lett. A* **244**, 482-488 (1998).
- [26] P. A. Miller and S. Sarkar, Fingerprints of classical instability in open quantum dynamics, *Phys. Rev. E* **58**, 4217-4225 (1998).
- [27] A. Isar, A. Sandulescu, and W. Scheid, Dissipative tunneling through a parabolic potential in the Lindblad theory of open quantum systems, *Eur. Phys. J. D* **12**, 3-10 (2000).
- [28] T. Shimbori and T. Kobayashi, Stationary flows of the parabolic potential barrier in two dimensions, *J. Phys. A: Math. Gen.* **33**, 7637-7652 (2000).
- [29] S. Matsumoto and M. Yoshimura, Dynamics of barrier penetration in a thermal medium: Exact result for the inverted harmonic oscillator, *Phys. Rev. A* **63**, 012104 (2000).
- [30] R. Blume-Kohout and W. H. Zurek, Decoherence from a chaotic environment: An upside-down “oscillator” as a model, *Phys. Rev. A* **68**, 032104 (2003).
- [31] J.-R. Choi, Exact quantum state for the time-dependent repulsive oscillator with a driving force and with a singular perturbation, *Phys. Scr.* **70**, 271-275 (2004).
- [32] D. Chruściński, Quantum mechanics of damped systems. II. Damping and parabolic potential barrier, *J. Math. Phys.* **45**, 841-854 (2004).
- [33] I. A. Pedrosa and I. Guedes, Quantum states of a generalized time-dependent inverted harmonic oscillator, *Int. J. Mod. Phys. B* **18**, 1379-1385 (2004).
- [34] F. C. Lombardo and P. I. Villar, Decoherence induced by zero-point fluctuations in quantum Brownian motion, *Phys. Lett. A* **336**, 16-24 (2005).
- [35] C. Yuce, A. Kilic, and A. Coruh, Inverted oscillator, *Phys. Scr.* **74**, 114-116 (2006).
- [36] M. Combescure and A. Combescure, Quantum and classical fidelity for singular perturbations of the inverted and harmonic oscillator, *J. Math. Anal. Appl.* **326**, 908-928 (2007).
- [37] J. R. Choi and K. H. Yeon, Coherent states of the inverted Caldirola-Kanai oscillator with time-dependent singularities, *Ann. Phys.* **323**, 812-826 (2008).
- [38] V. I. Man’ko, G. Marmo, and C. Stornaiolo, Tomographic entropy and cosmology, *Gen. Relativ. Gravit.* **40**, 1449-1465 (2008).
- [39] K. B. Wolf, El oscilador repulsivo, *Rev. Mex. Fis.* **56**, 83-91 (2010).
- [40] K. K. Kim and S. P. Kim, Tunneling Gaussian wave packets for an inverted harmonic potential, *J. Kor. Phys. Soc.* **56**, 1055-1062 (2010).
- [41] G.-J. Guo, Z.-Z. Ren, G.-X. Ju, and X.-Y. Guo, Quantum tunneling effect of a time-dependent inverted harmonic oscillator, *J. Phys. A: Math. Theor.* **44**, 185301 (2011).

- [42] G.-J. Guo, Z.-Z. Ren, G.-X. Ju, and X.-Y. Guo, Quantum tunneling effect of a driven inverted harmonic oscillator, *J. Phys. A: Math. Theor.* **44**, 305301 (2011).
- [43] Z. Lewis and T. Takeuchi, Position and momentum uncertainties of the normal and inverted harmonic oscillators under the minimal length uncertainty relation, *Phys. Rev. D* **84**, 105029 (2011).
- [44] J. B. Geloun, J. Hnybida, and J. R. Klauder, Coherent states for continuous spectrum operators with non-normalizable fiducial states, *J. Phys. A: Math. Theor.* **45**, 085301 (2012).
- [45] H. C. Rosu, O. Cornejo-Pérez, and P. Chen, Nonsingular parametric oscillators Darboux-related to the classical harmonic oscillator, *EPL* **100**, (2012) 60006.
- [46] D. M. Heim, W. P. Schleich, P. M. Alsing, J. P. Dahl, and S. Varro, Tunneling of an energy eigenstate through a parabolic barrier viewed from Wigner phase space, *Phys. Lett. A* **377**, 1822-1825 (2013).
- [47] V. G. Bagrov, D. M. Gitman, E. S. Macedo, and A. S. Pereira, Coherent states of inverse oscillators and related problems, *J. Phys. A: Math. Theor.* **46**, 325305 (2013).
- [48] D. Bermudez and D. J. Fernández C., Factorization method and new potentials from the inverted oscillator, *Ann. Phys.* **333**, 290-306 (2013).
- [49] O. Steuernagel, Equivalence between free quantum particles and those in harmonic potentials and its application to instantaneous changes, *Eur. Phys. J. Plus* **129**, 114 (2014).
- [50] S. Gentilini, M. C. Braidotti, G. Marcucci, E. DelRe, and C. Conti, Physical realization of the Glauber quantum oscillator, *Sci. Rep.* **5**, 15816 (2015).
- [51] I. A. Pedrosa, A. L. de Lima, and A. M. M. Carvalho, Gaussian wave packet states of a generalized inverted harmonic oscillator with time-dependent mass and frequency, *Can. J. Phys.* **93**, 841-845 (2015).
- [52] A. Schulze-Halberg, Supersymmetric partners and confinement of a spiked inverted oscillator model, *Eur. Phys. J. Plus* **130**, 228 (2015).
- [53] M. Maamache, Y. Bouguerra, and J. R. Choi, Time behavior of a Gaussian wave packet accompanying the generalized coherent state for the inverted oscillator, *Prog. Theor. Exp. Phys.* **2016**, 063A01 (2016).
- [54] G. Marcucci and C. Conti, Irreversible evolution of a wave packet in the rigged-Hilbert-space quantum mechanics, *Phys. Rev. A* **94**, 052136 (2016).
- [55] A. Rauh, Coherent states of harmonic and reversed harmonic oscillator, *Symmetry* **8**, 46 (2016).
- [56] C. Yuce, Self-accelerating parabolic cylinder waves in 1-D, *Phys. Lett. A* **380**, 3791-3795 (2016).
- [57] F. Finster and J. M. Isidro, L^p -spectrum of the Schrödinger operator with inverted harmonic oscillator potential, *J. Math. Phys.* **58**, 092104 (2017).
- [58] M. Maamache and J. R. Choi, Quantum-classical correspondence for the inverted oscillator, *Chin. Phys. C* **41**, 113106 (2017).
- [59] K. Rajeev, S. Chakraborty, and T. Padmanabhan, Inverting a normal harmonic oscillator: physical interpretation and applications, *Gen. Relat. Grav.* **50**, 116 (2018).
- [60] V. V. Chistyakov, Quantum analogue of unstable limit cycles of a periodically perturbed inverted oscillator, *Theor. Math. Phys.* **198**, 17-28 (2019).
- [61] T. Morita, Thermal emission from semiclassical dynamical systems, *Phys. Rev. Lett.* **122**, 101603 (2019).
- [62] R. D. Mota, D. Ojeda-Guillén, M. Salazar-Ramírez, and V. D. Granados, Non-Hermitian inverted harmonic oscillator-type Hamiltonians generated from supersymmetry with reflections, *Mod. Phys. Lett. A* **34**, 1950028 (2019).

- [63] T. Ali, A. Bhattacharyya, S. S. Haque, E. H. Kim, N. Moynihan, and J. Murugan, Chaos and complexity in quantum mechanics, *Phys. Rev. D* **101**, 026021 (2020).
- [64] K. Aouda, N. Kanda, S. Naka, and Haruki Toyoda, Ladder operators in repulsive harmonic oscillator with application to the Schwinger effect *Phys. Rev. D* **102**, 025002 (2020).
- [65] P. A. Golovinski, Dynamics of driven Brownian inverted oscillator, *Phys. Lett. A* **384**, 126203 (2020).
- [66] B. M. Villegas-Martínez, H. M. Moya-Cessa, and F. Soto-Eguibar, Exact and approximated solutions for the harmonic and anharmonic repulsive oscillators: Matrix method, *Eur. Phys. J. D* **74**, 137 (2020).
- [67] A. G. Nikitin, Symmetries of Schrodinger equation with scalar and vector potentials, *J. Phys. A: Math. Theor.* **53**, 455202 (2020).
- [68] V. Subramanyan, S. S. Hegde, S. Vishveshwara, B. Bradlyn, Physics of the inverted harmonic oscillator: From the lowest Landau level to event horizons, *Ann. Phys.* **435**, 168470 (2021).
- [69] A. Bhattacharyya, W. Chemissany, S. S. Haque, J. Murugan, and B. Yan, The multi-faceted inverted harmonic oscillator: Chaos and complexity, *SciPost Phys. Core* **4**, 002 (2021).
- [70] K. Gietka and T. Busch, Inverted harmonic oscillator dynamics of the nonequilibrium phase transition in the Dicke model, *Phys. Rev. E* **104**, 034132 (2021).
- [71] S. A. Hojman, H. M. Moya-Cessa, F. Soto-Eguibar, F. A. Asenjo, Time-dependent harmonic oscillators and SUSY in time domain, *Phys. Scr.* **96**, 125218 (2021).
- [72] C. Yuce, Quantum inverted harmonic potential, *Phys. Scr.* **96**, 105006 (2021).
- [73] F. Bagarello, A Swanson-like Hamiltonian and the inverted harmonic oscillator, *J. Phys. A: Math. Theor.* **55**, 225204 (2022).
- [74] F. Ullinger, M. Zimmermann, and W. P. Schleich, The logarithmic phase singularity in the inverted harmonic oscillator, *AVS Quantum Sci.* **4**, 024402 (2022).
- [75] O. V. Man'ko and V. I. Man'ko, Inverted oscillator quantum states in the probability representation, *Entropy* **25**, 217 (2023).
- [76] R. Zerimeche, R. Moufok, N. Amaouche and M. Maamache, Inverted oscillator: pseudo hermiticity and coherent states, *Rev. Mex. Fis.* **69**, 010402 (2023).
- [77] A. Sacchetti, Quantum forced oscillator via Wigner transform, *Bollettino dell'Unione Matematica Italiana* **16**, 103-134 (2023).
- [78] V. N. Chernega and O. V. Man'ko, Dynamics of system states in the probability representation of quantum mechanics, *Entropy* **25**, 785 (2023).
- [79] W. Wöger, H. King, R.J. Glauber, and J. W. Haus, Spontaneous generation of coherent optical beats, *Phys. Rev. A* **34**, 4859-4868 (1986).
- [80] S. Tarzi, The inverted harmonic oscillator: some statistical properties, *J. Phys. A: Math. Gen.* **21**, 3105-3111 (1988).
- [81] R. Grimaudo, V. I. Man'ko, M. A. Man'ko, and A. Messina, Dynamics of a harmonic oscillator coupled with a Glauber amplifier, *Phys. Scr.* **95**, 024004 (2020).
- [82] L. Parker, Quantized fields and particle creation in expanding universes, II, *Phys. Rev. D* **3**, 346-356 (1971).
- [83] J. Janszky and Y. Y. Yushin, Squeezing via frequency jump, *Opt. Commun.* **59**, 151-154 (1986).
- [84] R. Graham, Squeezing and frequency changes in harmonic oscillations, *J. Mod. Opt.* **34**, 873-879 (1987).

- [85] X. Ma and W. Rhodes, Squeezing in harmonic oscillators with time-dependent frequencies, *Phys. Rev. A* **39**, 1941–1947 (1989).
- [86] C. F. Lo, Squeezing by tuning the oscillator frequency, *J. Phys. A: Math. Gen.* **23**, 1155–1165 (1990).
- [87] J. Janszky and P. Adam, Strong squeezing by repeated frequency jumps, *Phys. Rev. A* **46**, 6091–6092 (1992).
- [88] V. V. Dodonov, A. B. Klimov, and D. E. Nikonov, Quantum phenomena in nonstationary media, *Phys. Rev. A* **47**, 4422–4429 (1993).
- [89] T. Kiss, J. Janszky, and P. Adam, Time evolution of harmonic oscillators with time-dependent parameters: a step-function approximation, *Phys. Rev. A* **49**, 4935–4942 (1994).
- [90] E. Sassaroli, Y. N. Srivastava, and A. Widom, Photon production by the Dynamical casimir effect, *Phys. Rev. A* **50**, 1027–1034 (1994).
- [91] J. T. Mendonça, A. Guerreiro, and A. M. Martins, Quantum theory of time refraction, *Phys. Rev. A* **62**, 033805 (2000).
- [92] D. Martínez-Tibaduiza, L. Pires, and C. Farina, Time-dependent quantum harmonic oscillator: a continuous route from adiabatic to sudden changes, *J. Phys. B: At. Mol. Opt. Phys.* **54**, 205401 (2021).
- [93] K. Husimi, Miscellanea in elementary quantum mechanics. II. *Prog. Theor. Phys.* **9**, 381–402 (1953).
- [94] V. S. Popov and A. M. Perelomov, Parametric excitation of a quantum oscillator, *Zhurn. Eksp. Teor. Fiz.* **56**, 1375–1390 (1969) [*Sov. Phys. – JETP* **29**, 738–745] (1969).
- [95] H. R. Lewis Jr. and W. B. Riesenfeld, An exact quantum theory of the time-dependent harmonic oscillator and of a charged particle in a time-dependent electromagnetic field, *J. Math. Phys.* **10**, 1458–1473 (1969).
- [96] I. A. Malkin, V. I. Man’ko, and D. A. Trifonov, Coherent states and transition probabilities in a time-dependent electromagnetic field, *Phys. Rev. D* **2**, 1371–1385 (1970).
- [97] R. Cordero-Soto, E. Suazo, and S. K. Suslov, Quantum integrals of motion for variable quadratic Hamiltonians, *Ann. Phys.* **325**, 1884–1912 (2010).
- [98] V. V. Dodonov, Universal integrals of motion and universal invariants of quantum systems, *J. Phys. A: Math. Gen.* **33**, 7721–7738 (2000).
- [99] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*, Academic: Amsterdam, 2007.
- [100] A. Erdélyi, (Ed.) *Bateman Manuscript Project: Higher Transcendental Functions*, McGraw-Hill: New York, 1953.
- [101] A. P. Prudnikov, Y. A. Brychkov, and O. I. Marichev, *Integrals and Series. Volume 2. Special Functions*, Taylor & Francis: London, UK, 2002.