Data-Driven Tuning Parameter Selection for High-Dimensional Vector Autoregressions^{*}

Anders Bredahl Kock[†] Rasmus Søndergaard Pedersen[‡] Jesper Riis-Vestergaard Sørensen[§]

March 12, 2024

Abstract

Lasso-type estimators are routinely used to estimate high-dimensional time series models. The theoretical guarantees established for Lasso typically require the penalty level to be chosen in a suitable fashion often depending on *unknown* population quantities. Furthermore, the resulting estimates and the number of variables retained in the model depend crucially on the chosen penalty level. However, there is currently no theoretically founded guidance for this choice in the context of high-dimensional time series. Instead one resorts to selecting the penalty level in an ad hoc manner using, e.g., information criteria or cross-validation. We resolve this problem by considering estimation of the perhaps most commonly employed multivariate time series model, the linear vector autoregressive (VAR) model, and propose a weighted Lasso estimator with penalization chosen in a *fully data-driven* way. The theoretical guarantees that we establish for the resulting estimation and prediction error match those currently available for methods based on infeasible choices of penalization. We thus provide a first solution for choosing the penalization in high-dimensional time series models.

Keywords: High-dimensional time series, vector autoregression model, ℓ_1 -penalized estimation, tuning parameter selection, data-driven, sparsity, weak functional dependence.

^{*}First version: October 2022. We thank Abhishek Chakrabortty, Bent Nielsen, Lan Gao, Anders Rahbek, Mikkel Sølvsten, and participants at the Aarhus Workhop in Econometrics 2022 for insightful comments and discussion. Moreover, we thank Zifan Li for providing details about the simulation experiment in Wong et al. (2020). Pedersen gratefully acknowledges financial support from Independent Research Fund Denmark ("Robust Econometric Methods in Empirical and Quantitative Finance", FSE 0133-00162B).

[†]University of Oxford; e-mail: anders.kock@economics.ox.ac.uk.

[‡]University of Copenhagen; e-mail: rsp@econ.ku.dk.

[§]University of Copenhagen; e-mail: jrvs@econ.ku.dk.

1 Introduction

Multivariate time series play a fundamental role in many areas of research. The quintessential approach to modelling these series is based on the linear VAR model; see, e.g., Lütkepohl (2005). In order to fully capture the dynamics of the time series at hand and to decrease the risk of omitting variables, one frequently includes many explanatory variables. These considerations result in a high-dimensional VAR model wherein the number of parameters can exceed the number of observations. Consequently, the parameters can no longer be estimated by least squares, and as an alternative there has been a surge of research on Lassotype estimators [Tibshirani (1996)]. For instance, assuming independent and identically distributed (i.i.d.) Gaussian innovations in the VAR, consistency and oracle inequalities for the Lasso have been studied by Han and Liu (2013), Basu and Michailidis (2015), Kock and Callot (2015) and Davis et al. (2016) among others. The assumption of Gaussian innovations has been relaxed in several papers, including Song and Bickel (2011), Wong et al. (2020) and Miao et al. (2023). In particular, Wong et al. (2020) derive consistency results for the Lasso in VAR models under suitable mixing and moment conditions, whereas Miao et al. (2023) derive rates of convergence and oracle properties for factor augmented VAR models, relying on results for weakly dependent processes in the sense of Wu (2005). Masini et al. (2022) establish oracle inequalities in the case of martingale difference innovations that are not necessarily mixing. Chernozhukov et al. (2021) consider Lasso-driven inference for time series and spatial models, while Adamek et al. (2023) consider inference based on the desparsified Lasso when the data-generating process allows near-epoch dependence. Further related papers are Han et al. (2015), Guo et al. (2016) and Wu and Wu (2016).

To implement the Lasso one must choose the penalization level λ , and the theoretical results listed above are derived under a suitable choice of this level. However, the "right" choice of this tuning parameter typically depends on *unknown* population quantities such as mixing coefficients or other coefficients quantifying the dependence structure of the data generating process [Wong et al. (2020), Babii et al. (2022), Masini et al. (2022)], the population covariance matrix of the observed variables and innovations [Kock and Callot (2015), Medeiros and Mendes (2016)], the population coefficient matrix [Basu and Michailidis (2015)], or other quantities depending on the data generating process [Chernozhukov et al. (2021), Adamek et al. (2023), Miao et al. (2023)]. Consequently, despite the large amount of research on penalized estimation of VAR models, there is currently no firm guidance on how to choose λ in practice. Instead, one often resorts to information criteria [e.g. Kock and Callot (2015), Medeiros and Mendes (2016) and Masini et al. (2022)], cross-validation [e.g. Wong et al. (2020), Babii et al. (2022) and Miao et al. (2023)] or other ad hoc methods without providing theoretical guarantees for these. Crucially, such choices of penalization do not necessarily satisfy the conditions imposed in the theoretical results. Therefore, strictly speaking, the theoretical guarantees provided are only valid for fortuitous choices of λ . In this paper we resolve this problem by proposing a data-driven way of choosing λ along with prediction and estimation error guarantees for the resulting weighted Lasso estimator of the parameters in large VAR models. As upper bounds on the estimation error play a crucial role in establishing the validity of inference based on debiasing in high-dimensional models [see Javanmard and Montanari (2014), van de Geer et al. (2014) and Zhang and Zhang (2014)], our results open the door to doing so in VAR models with a data-driven choice of λ .

The algorithm we study originates from Belloni et al. (2012) who consider regressions in high dimensions with independent data. There are several challenges in adapting this algorithm to time series data: First, the analysis of Belloni et al. (2012) relies on the independence as well as certain high-level conditions on the explanatory variables and their relation to the model errors. In VAR models one cannot impose such conditions as their validity is completely determined by the model (which also generates the explanatory variables). Instead, imposing only primitive conditions, we carefully take into account the inherent dependence in the VAR. Second, as heavy tails are omnipresent in many time series, we allow for the possibility of certain types of heavy-tailed innovation distributions, that is, so-called sub-Weibull innovations. To accommodate dependence and heavy tails, we establish a novel maximal inequality for centered sums of dependent sub-Weibull random variables, that may be of independent interest. The validity of the penalty loadings (i.e. Lasso weights) proposed in the algorithm is then established by proving that these are close to certain infeasible ideal loadings that are constructed by means of blocking-based self-normalization. In order to verify that the ideal loadings are well-behaved, we rely on recent moderate deviation theory for self-normalized block-sums of weakly dependent processes (as quantified by a functional dependence measure) derived by Chen et al. (2016) and Gao et al. (2022).

In a simulation experiment, we show that the Lasso and post-Lasso estimators based on the proposed algorithm for tuning parameter choice perform quite satisfactory compared to Lasso estimators obtained from penalization using the Bayesian Information Criterion (BIC) [as used in, e.g., Kock and Callot (2015)]. As an empirical illustration, we consider estimation of VAR models based on monthly observations of 127 US macroeconomic variables. We find that the Lasso and post-Lasso estimators deliver superior out-of-sample forecasts compared to BIC-based Lasso estimators.

Outline

In Section 2 we present the weighted Lasso estimator and describe the data-driven tuning parameter selection. Section 3 presents the assumptions and performance guarantees for the weighted Lasso. Section 4 outlines the key steps in the proof of Theorem 1 and the challenges involved in establishing each step. Sections 5 and 6 contain simulations and an empirical illustration, respectively. Proofs and technical lemmas are contained in the supplementary appendices.

Notation

For $k \in \mathbb{N}$, we write $[k] := \{1, \ldots, k\}$. For $\boldsymbol{\delta} \in \mathbb{R}^k$, we denote its ℓ_r -norm, by $\|\boldsymbol{\delta}\|_r := (\sum_{j=1}^k |\delta_j|^r)^{1/r}, r \in [1, \infty)$, and $\|\boldsymbol{\delta}\|_{\ell_{\infty}} := \max_{j \in [k]} |\delta_j|$. For a matrix $\mathbf{A} \in \mathbb{R}^{k \times l}$, we use $\mathbf{A}_{1:m,1:n}$ to denote the first *n* entries of its first *m* rows (i.e. its $m \times n$ "upper left"). For a square matrix $\mathbf{A} \in \mathbb{R}^{k \times k}$, we denote all eigenvalues of \mathbf{A} by $\Lambda(\mathbf{A})$ and write $\rho(\mathbf{A}) := \max\{|\lambda| : \lambda \in \Lambda(\mathbf{A})\}$ for its spectral radius (with $|\cdot|$ the complex modulus). If $\mathbf{A} \in \mathbb{R}^{k \times k}$ is symmetric, we write $\Lambda_{\min}(\mathbf{A})$ and $\Lambda_{\max}(\mathbf{A})$ for the minimum and maximum eigenvalues, respectively.

For a random scalar X, we denote its L_r -norm by $||X||_r := (\mathbb{E}[|X|^r])^{1/r}$, $r \in [1, \infty)$, with $\mathbb{E}[\cdot]$ denoting the expectation operator. For $\alpha \in (0, \infty)$, we define the sub-Weibull(α) norm of X as $||X||_{\psi_{\alpha}} := \sup_{r \in [1,\infty)} r^{-1/\alpha} ||X||_r$. A random scalar X is said to be sub-Weibull(α) if $||X||_{\psi_{\alpha}} < \infty$.¹ For a k-dimensional random vector \mathbf{X} , we define its joint sub-Weibull(α) norm $||\mathbf{X}||_{\psi_{\alpha}} := \sup\{||\mathbf{u}^{\top}\mathbf{X}||_{\psi_{\alpha}} : ||\mathbf{u}||_{\ell_2} = 1\}$ and call \mathbf{X} jointly sub-Weibull(α) if $||\mathbf{X}||_{\psi_{\alpha}} < \infty$.

For non-random numbers a_n and positive numbers $b_n, n \in \mathbb{N}$, we write $a_n = o(1)$ if $a_n \to 0$ as $n \to \infty$, and $a_n \leq b_n$, if the sequence a_n/b_n is bounded. For random variables V_n and positive numbers b_n , we write $V_n \leq_P b_n$, if the sequence V_n/b_n is bounded in probability. We reserve the word "constant" for non-random quantities that do not depend on n.

2 Model and Penalization Algorithm

We study the q^{th} -order $(q \in \mathbb{N})$ VAR model given by

$$\boldsymbol{Y}_{t} = \sum_{j=1}^{q} \boldsymbol{\Theta}_{0j} \boldsymbol{Y}_{t-j} + \boldsymbol{\varepsilon}_{t}, \quad t \in \mathbb{Z},$$
(2.1)

yielding a stochastic process $\{\mathbf{Y}_t\}_{t\in\mathbb{Z}}$ which is (strictly) stationary under the assumptions in Section 3. Here $\mathbf{Y}_t := (Y_{t,1}, \ldots, Y_{t,p})^{\top}$ is a random vector of length p, $\{\mathbf{\Theta}_{0j}\}_{j=1}^q$ are $p \times p$

¹One may show that the space of sub-Weibull(α) random variables is complete with respect to $\|\cdot\|_{\psi_{\alpha}}$. Consequently, it holds that the norm $\|\cdot\|_{\psi_{\alpha}}$ is countably sub-additive, which we will use repeatedly.

(unknown) coefficient matrices, and $\boldsymbol{\varepsilon}_t := (\varepsilon_{t,1}, \ldots, \varepsilon_{t,p})^{\top}$ is a sequence of i.i.d. innovations. Given observations $\{\boldsymbol{Y}_t\}_{t=-(q-1)}^n$ from (2.1), the objective is to estimate $\{\boldsymbol{\Theta}_{0j}\}_{j=1}^q$, while allowing (but not requiring) that the number of elements p^2q in $\{\boldsymbol{\Theta}_{0j}\}_{j=1}^q$ is larger than the (effective) sample size *n*. Throughout we take $p \ge 2$ and $n \ge 3$.²

The process in (2.1) may be written in companion form,

$$\boldsymbol{Z}_{t} = \widetilde{\boldsymbol{\Theta}}_{0} \boldsymbol{Z}_{t-1} + \widetilde{\boldsymbol{\varepsilon}}_{t}, \quad t \in \mathbb{Z},$$
(2.2)

for

$$\underbrace{\boldsymbol{Z}_{t}}_{pq \times 1} := \begin{pmatrix} \boldsymbol{Y}_{t} \\ \boldsymbol{Y}_{t-1} \\ \vdots \\ \boldsymbol{Y}_{t-(q-1)} \end{pmatrix}, \quad \underbrace{\widetilde{\boldsymbol{\Theta}}_{0}}_{pq \times pq} := \begin{pmatrix} \boldsymbol{\Theta}_{01} & \boldsymbol{\Theta}_{02} & \cdots & \boldsymbol{\Theta}_{0q} \\ \boldsymbol{I}_{p} & \boldsymbol{0}_{p \times p} & \cdots & \boldsymbol{0}_{p \times p} \\ \boldsymbol{0}_{p \times p} & \boldsymbol{I}_{p} & \boldsymbol{0}_{p \times p} & \vdots \\ \vdots & \ddots & \ddots & \ddots & \\ \boldsymbol{0}_{p \times p} & \cdots & \boldsymbol{0}_{p \times p} & \boldsymbol{I}_{p} & \boldsymbol{0}_{p \times p} \end{pmatrix}, \quad \underbrace{\widetilde{\boldsymbol{\varepsilon}}_{t}}_{pq \times 1} := \begin{pmatrix} \boldsymbol{\varepsilon}_{t} \\ \boldsymbol{0}_{p \times 1} \\ \vdots \\ \boldsymbol{0}_{p \times 1} \end{pmatrix},$$

and $\mathbf{I}_p \in \mathbb{R}^{p \times p}$ being the identity matrix. In the special case of q = 1, we interpret $\widetilde{\Theta}_0$ as Θ_{01} and $\widetilde{\varepsilon}_t$ as ε_t . For $i \in [p]$, the i^{th} row in (2.1) is given by

$$Y_{t,i} = \mathbf{Z}_{t-1}^{\top} \boldsymbol{\beta}_{0i} + \varepsilon_{t,i}, \qquad (2.3)$$

where the vector $\boldsymbol{\beta}_{0i}^{\top}$ of length pq is the *i*th row of the companion matrix $\tilde{\boldsymbol{\Theta}}_{0}$. In a lowdimensional setting where $\boldsymbol{\beta}_{0i}^{\top}$ is of fixed length, estimation is often done by equationwise least squares with the same $n \times pq$ regressor matrix $\mathbf{X} := [\mathbf{Z}_0 : \cdots : \mathbf{Z}_{n-1}]^{\top}$ for all $i \in [p]$. However, when pq > n, the $pq \times pq$ Gram matrix $\mathbf{X}^{\top}\mathbf{X}$ has reduced rank, and Lasso-type estimators have been studied as an alternative under various sparsity assumptions. Nevertheless, as discussed in the introduction, the theoretical guarantees hitherto established for these are valid only for specific tuning parameter choices depending on unknown population quantities. As a result—despite the sensitivity of shrinkage estimators to the tuning parameter choice—one must currently resort to choosing this parameter by methods without theoretical guarantees when implementing shrinkage estimators in time series models. We provide a solution to this problem in the context of the weighted Lasso by proposing an algorithm for tuning parameter selection and explicitly incorporate the data-driven tuning

²We consider the VAR model in (2.1), allowing the number of output variables p, the (common) distribution of ε_t , and, hence, that of \mathbf{Y}_t to depend on the sample size n. That is, we consider an array $\{\{\mathbf{Y}_t^{(n)}\}_{t\in\mathbb{Z}}\}_{n\in\mathbb{N}}$ of stochastic processes, each process $\{\mathbf{Y}_t^{(n)}\}_{t\in\mathbb{Z}}$ presumed strictly stationary, in which each $\mathbf{Y}_t^{(n)} = (\mathbf{Y}_{t,1}^{(n)}, \dots, \mathbf{Y}_{t,p_n}^{(n)})^{\top}, t \in \mathbb{Z}$, is a random element of \mathbb{R}^{p_n} . To ease notation, we henceforth suppress the n superscript.

parameter choice into our theoretical guarantees.

A weighted Lasso estimator satisfies

$$\widehat{\boldsymbol{\beta}}_{i} := \widehat{\boldsymbol{\beta}}_{i}(\lambda, \widehat{\boldsymbol{\Upsilon}}_{i}) \in \operatorname*{argmin}_{\boldsymbol{\beta} \in \mathbb{R}^{pq}} \left\{ \widehat{Q}_{i}(\boldsymbol{\beta}) + \frac{\lambda}{n} \| \widehat{\boldsymbol{\Upsilon}}_{i} \boldsymbol{\beta} \|_{\ell_{1}} \right\},$$
(2.4)

where \hat{Q}_i is the sample average square error loss function

$$\widehat{Q}_{i}(\boldsymbol{\beta}) := \frac{1}{n} \sum_{t=1}^{n} \left(Y_{t,i} - \boldsymbol{Z}_{t-1}^{\top} \boldsymbol{\beta} \right)^{2}, \quad \boldsymbol{\beta} \in \mathbb{R}^{pq},$$
(2.5)

for $i \in [p]$, $\lambda \in (0, \infty)$ is a penalty level and each $\widehat{\Upsilon}_i := \operatorname{diag}(\widehat{v}_{i,1}, \ldots, \widehat{v}_{i,pq})$ is a diagonal matrix of data-dependent penalty loadings $\widehat{v}_{i,j} \in (0, \infty), j \in [pq]$. Both the penalty level and loadings will be specified in Algorithm 1 below. Although the minimization problem in (2.4) may have multiple solutions, we sometimes refer to such a $\widehat{\beta}_i$ as the Lasso estimator. The results established below apply to any such minimizer.

Note that much of the literature on the Lasso and its variants employs the same penalty loadings for all regressors (e.g. $\hat{v}_{i,j} \equiv 1$). Equal weighting implicitly treats the regressors as either being on the same scale or having been brought onto the same scale by some preliminary transformation of the data, which is then typically abstracted from in the theoretical analysis. Taking serious the effect of such a preliminary transformation is not trivial as it can alter the already intricate dependence structure of the time series at hand. By incorporating data-dependent loadings, our algorithm is not subject to this caveat.

Adapting Belloni et al. (2012, Algorithm A.1) for independent data to our setting, we consider the following data-driven penalization. Let

$$c := 1.1 \text{ and } \gamma_n = .1/\ln(\max\{n, pq\})$$
 (2.6)

and let $K \in \mathbb{N}_0$ denote a fixed number of iterations (loading updates). Following Belloni et al. (2012), we fix K at 15 in our simulations and empirical illustration. To state Algorithm 1 below, let Φ be the standard Gaussian cumulative distribution function (CDF).

Algorithm 1 (Data-Driven Penalization).

Initialize: Specify the penalty level λ in (2.4) as

$$\lambda_n^* := 2c\sqrt{n}\Phi^{-1} (1 - \gamma_n / (2p^2 q)), \qquad (2.7)$$

and specify the initial penalty loadings as

$$\widehat{v}_{i,j}^{(0)} := \sqrt{\frac{1}{n} \sum_{t=1}^{n} Y_{t,i}^2 Z_{t-1,j}^2}, \quad i \in [p], \quad j \in [pq].$$
(2.8)

Use λ_n^* and $\widehat{\Upsilon}_i^{(0)} := \operatorname{diag}(\widehat{v}_{i,1}^{(0)}, \dots, \widehat{v}_{i,pq}^{(0)})$ to compute a Lasso estimate $\widehat{\beta}_i^{(0)} := \widehat{\beta}_i(\lambda_n^*, \widehat{\Upsilon}_i^{(0)})$ via (2.4) for each $i \in [p]$. Store the residuals $\widehat{\varepsilon}_{t,i}^{(0)} := Y_{t,i} - \mathbb{Z}_{t-1}^{\top} \widehat{\beta}_i^{(0)}, t \in [n], i \in [p]$, and set k = 1.

Update: While $k \leq K$, specify the penalty loadings as

$$\widehat{\upsilon}_{i,j}^{(k)} := \sqrt{\frac{1}{n} \sum_{t=1}^{n} (\widehat{\varepsilon}_{t,i}^{(k-1)})^2 Z_{t-1,j}^2}, \quad i \in [p], \quad j \in [pq].$$

$$(2.9)$$

Use λ_n^* and $\widehat{\mathbf{\Upsilon}}_i^{(k)} := \operatorname{diag}(\widehat{v}_{i,1}^{(k)}, \dots, \widehat{v}_{i,pq}^{(k)})$ to compute a Lasso estimate $\widehat{\boldsymbol{\beta}}_i^{(k)} := \widehat{\boldsymbol{\beta}}_i(\lambda_n^*, \widehat{\mathbf{\Upsilon}}_i^{(k)})$ via (2.4) for each $i \in [p]$. Store the residuals $\widehat{\varepsilon}_{t,i}^{(k)} := Y_{t,i} - \mathbf{Z}_{t-1}^{\top} \widehat{\boldsymbol{\beta}}_i^{(k)}, t \in [n], i \in [p]$, and increment $k \leftarrow k+1$.

Note that Algorithm 1 does not require any knowledge of the degree of dependence in the process $\{Y_t\}_{t\in\mathbb{Z}}$ (as quantified by mixing coefficients or other unknown population quantities). This feature of the penalization is in contrast to current recommendations in the literature.

To discuss error rates for the implied Lasso estimator, let $\|\cdot\|_{2,n}$ be the prediction norm

$$\|\boldsymbol{\delta}\|_{2,n} := \sqrt{rac{1}{n}\sum_{t=1}^{n} \left(\boldsymbol{Z}_{t-1}^{ op} \boldsymbol{\delta}
ight)^2}$$

attached to the vector $\boldsymbol{\delta} \in \mathbb{R}^{pq}$, and let

$$s := \max_{i \in [p]} \sum_{j=1}^{pq} \mathbf{1} \left(\beta_{0i,j} \neq 0 \right), \qquad (2.10)$$

denote the sparsity number, that is, the largest cardinality of the support of β_{0i} across

 $i \in [p]$. Without loss of generality, we take $s \ge 1$. Then, under primitive conditions given in Section 3, any Lasso estimators $\widehat{\beta}_i := \widehat{\beta}_i(\lambda_n^*, \widehat{\Upsilon}_i^{(K)}), i \in [p]$, in (2.4) based on the penalty level λ_n^* in (2.7) and the (final) penalty loadings $\widehat{\Upsilon}_i^{(K)}$ arising from Algorithm 1 satisfy

$$\max_{i \in [p]} \|\widehat{\boldsymbol{\beta}}_i - \boldsymbol{\beta}_{0i}\|_{2,n} \lesssim_{\mathrm{P}} \sqrt{\frac{s \ln (pn)}{n}},$$
(2.11)

$$\max_{i \in [p]} \|\widehat{\boldsymbol{\beta}}_i - \boldsymbol{\beta}_{0i}\|_{\ell_1} \lesssim_{\mathrm{P}} \sqrt{\frac{s^2 \ln (pn)}{n}} \quad \text{and}$$
(2.12)

$$\max_{i \in [p]} \|\widehat{\boldsymbol{\beta}}_{i} - \boldsymbol{\beta}_{0i}\|_{\ell_{2}} \lesssim_{\mathrm{P}} \sqrt{\frac{s \ln (pn)}{n}}.$$
(2.13)

The formal statement, including a set of sufficient conditions, is given in Theorem 1 below. Importantly, the performance guarantees in (2.11)-(2.13) match those currently available based on *infeasible* penalty level choices.

Remark 1 (Including Intercepts). In applications one typically includes model intercepts in an unpenalized manner. In Section A.1, we modify Algorithm 1 for this purpose.

3 Assumptions and Performance Guarantees for Algorithm **1**

For the remainder of the paper we invoke (a subset of) the following assumptions.

Assumption 1 (Innovations). (1) The innovations $\{\boldsymbol{\varepsilon}_t\}_{t\in\mathbb{Z}}$ are *i.i.d.* centered random elements of \mathbb{R}^p with covariance matrix $\boldsymbol{\Sigma}_{\boldsymbol{\varepsilon}} \in \mathbb{R}^{p \times p}$. (2) There are constants $a_1 \in (0, \infty)$ and $\alpha \in (0, 2]$ such that $\|\boldsymbol{\varepsilon}_0\|_{\psi_{\alpha}} \leq a_1$. (3) There is a constant $a_2 \in (0, \infty)$ such that $\Lambda_{\min}(\boldsymbol{\Sigma}_{\boldsymbol{\varepsilon}}) \geq a_2$.

Assumption 1.1 is imposed in, e.g., Kock and Callot (2015), Wong et al. (2020, Example 1) and Miao et al. (2023). Assumption 1.2 states that ε_0 belongs to the sub-Weibull family of distributions with tail parameter (at least) α , and is also used in Wong et al. (2020) and Masini et al. (2022). Special cases are the sub-exponential ($\alpha = 1$) and sub-Gaussian ($\alpha = 2$) families. If ε_0 is Gaussian—as imposed in, e.g., Han and Liu (2013), Basu and Michailidis (2015), Kock and Callot (2015) and Davis et al. (2016)—then Assumption 1.2 holds if $\Lambda_{\max}(\Sigma_{\varepsilon})$ is bounded from above. Requiring that $\alpha \leq 2$ is innocuous and can be relaxed at the expense of some additional notation. Assumption 1.3 is part of Masini et al. (2022, Assumption A2) and parallels Miao et al. (2023, Assumption A.1(i)).

Assumption 2 (Companion Matrix). (1) The spectral radius $\rho(\Theta_0) < 1$. (2) There are constants $b_1, b_2 \in (0, \infty)$, such that $\max_{i \in [p]} \|(\widetilde{\Theta}_0^h)_{i,1:p}\|_{\ell_2} \leq b_1 e^{-b_2 h}$ for each $h \in \mathbb{N}$. (3) The lag-order q is constant.

Assumption 2.1 is standard and, together with Assumption 1.1, ensures that the process $\{Y_t\}_{t\in\mathbb{Z}}$ in (2.1) [as well as $\{Z_t\}_{t\in\mathbb{Z}}$ in (2.2)] is strictly stationary. In particular, $\{Y_t\}_{t\in\mathbb{Z}}$ has the moving average representation

$$oldsymbol{Y}_t = \sum_{\ell=0}^\infty (\widetilde{oldsymbol{\Theta}}_0^\ell)_{1:p,1:p} oldsymbol{arepsilon}_{t-\ell},$$

where the series converges absolutely almost surely (and, under Assumption 2.2, in sub-Weibull norm).

Assumption 2.2 is milder than Masini et al. (2022, Assumption A1) which imposes an exponential decay on the ℓ_1 -norms of the rows of $(\widetilde{\Theta}_0^{\ell})_{1:p,1:p}$ as ℓ grows (as opposed to our decay placed on the ℓ_2 -norms). Likewise, the assumption is also milder than Miao et al. (2023, Assumption A.1(vi)), which imposes an exponential decay on the spectral norm of $(\widetilde{\Theta}_0^{\ell})_{1:p,1:p}$ (exceeding the row-wise ℓ_2 -norms).³

Assumption 2.3 states that the lag-order q is constant. This appears to be reasonable for most practical purposes as q is often thought of as small relative to n.⁴ Indeed, in our application in Section 6 we have n = 758 and as a result of monthly sampling $q = 12 \ll n$ often suffices. We use that q does not depend on n in order to exploit recent moderate deviation results for self-normalized sums based on geometric moment contracting processes [Gao et al. (2022)], and the assumption therefore seems hard to relax with currently available methods.

We make the following assumption on the matrix $\Sigma_{\mathbf{Z}} := \mathrm{E}[\mathbf{Z}_0 \mathbf{Z}_0^{\top}]$.

Assumption 3 (Covariance). There is a constant $d \in (0, \infty)$ such that $\Lambda_{\min}(\Sigma_{\mathbf{Z}}) \ge d$.

Assumption 3 ensures that $\Sigma_{\mathbf{Z}}$ has full rank and is implied by, e.g., Miao et al. (2023, Assumption A.1). As can be inferred from Masini et al. (2022, p. 536), Assumption 3 follows from additional (boundedness) conditions on $\widetilde{\Theta}_0$, although these are not necessary. In the special case of a stationary VAR(1), Assumption 3 is actually implied by Assumption 1.⁵

Assumption 4 (Growth and Sparsity). For a constant $\alpha \in (0, 2]$ satisfying Assumption 1.2, we have $(\ln p)^{40/(3\alpha)} = o(n)$ and $s^2(\ln(pn))^{1+4/\alpha} = o(n)$, where s is given in (2.10).

Assumption 4 imposes sparsity on each row β_{0i}^{\top} of the parameter matrix $[\Theta_{01} : \cdots : \Theta_{0q}]$ as in, e.g., Kock and Callot (2015), Chernozhukov et al. (2021), and Masini et al. (2022).

³See Masini et al. (2022, Lemma 4) for further sufficient conditions for Assumption 2.2.

⁴This point was also made by Kock and Callot (2015), although their analysis allows for growing q.

⁵For ease of presentation we choose to work with Assumption 3, and note that this assumption is only used in the proof of Lemma 5. Inspection of this proof shows that one may replace Assumption 3 by the milder requirement that only certain *restricted* eigenvalues of $\Sigma_{\mathbf{Z}}$ are bounded away from zero. The latter condition is used in, e.g., Kock and Callot (2015) and Medeiros and Mendes (2016).

We can now state the properties of the weighted Lasso estimator with data-driven tuning parameter selection based on Algorithm 1.

Theorem 1 (Convergence Rates for Lasso with Data-Driven Penalization). Let Assumptions 1, 2, 3 and 4 hold. Then Lasso estimators $\hat{\beta}_i(\lambda_n^*, \hat{\Upsilon}_i^{(K)}), i \in [p]$, arising from the data-driven penalization in Algorithm 1 satisfy (2.11)–(2.13).

The estimation and prediction error rates guaranteed by Theorem 1 match those currently available in the literature building on infeasible penalty parameter choices. Algorithm 1 is thus the first data-driven tuning parameter method for which theoretical guarantees have been provided in the context of dependent data. Upper bounds on the estimation error are also a crucial ingredient in establishing uniformly valid inference based on the debiasing/desparsification methodology originating in Javanmard and Montanari (2014), van de Geer et al. (2014) and Zhang and Zhang (2014). This methodology has been used in the context of time series in Adamek et al. (2023). Thus, Theorem 1 opens the door to conducting inference in high-dimensional VAR models with data-driven penalty parameter choice.

In establishing Theorem 1, we cannot call upon the large existing body of maximal and concentration inequalities for independent random variables. Instead, we establish a new maximal inequality for sums of dependent random variables (Lemma B.2 in the appendix). This result is then used to show that the penalty loadings of Algorithm 1 are close to certain ideal, yet infeasible, blocking-based loadings (Lemma 6). Note that although these ideal loadings are based on blocking, Algorithm 1 is not, so no choice of block size is needed in practice. The ideal loadings can, in turn, be used as self-normalizing factors in an application of the recent moderate deviation theorems of Gao et al. (2022) for dependent random variables. Using these results, we show that λ_n^* is a high-probability upper bound on the maximum of such self-normalized sums. Section 4 explains further steps and challenges involved in establishing Theorem 1, while details are deferred to the appendix.

4 Main Steps in Proving Theorem 1

To sketch the proof of Theorem 1, we start with a "master lemma" that provides nonasymptotic error guarantees for the Lasso under standard high-level conditions, well-known in the literature. To state this lemma, let $C \in (0, \infty)$ and $\emptyset \neq T \subseteq [pq]$ and define

$$\mathcal{R}_{C,T} := \left\{ \boldsymbol{\delta} \in \mathbb{R}^{pq} : \left\| \boldsymbol{\delta}_{T^c} \right\|_{\ell_1} \leqslant C \left\| \boldsymbol{\delta}_T \right\|_{\ell_1} \right\},$$
(4.1)

where $\boldsymbol{\delta}_T \in \mathbb{R}^{pq}$ denotes the vector with coordinates $\delta_{T,j} := \delta_j$ if $j \in T$ and zero otherwise. Define the *restricted eigenvalue* of the (scaled) Gram matrix $\mathbf{X}^{\top}\mathbf{X}/n$ as

$$\widehat{\kappa}_{C}^{2} := \widehat{\kappa}_{C}^{2} \left(s, \mathbf{X}^{\top} \mathbf{X}/n \right) := \min_{\substack{T \subseteq [pq]: \\ |T| \in [s]}} \inf_{\substack{\delta \in \mathcal{R}_{C,T}: \\ \delta \neq \mathbf{0}}} \frac{\delta^{\top} (\mathbf{X}^{\top} \mathbf{X}/n) \delta}{\|\delta\|_{\ell_{2}}^{2}}.$$
(4.2)

Similar to Belloni et al. (2012), we introduce certain infeasible ideal penalty loadings. However, to account for the dependence in the data, the infeasible loadings are based on dividing the data into blocks of carefully chosen size $m_n := n^{1/5}$, the number of which is $l_n := n/m_n = n^{4/5.6}$ Such blocking is not necessary for independent data. We stress that the introduction of a block size m_n is solely for the purpose of the theoretical analysis, as our Algorithm 1 does not require any user-chosen block size. The indices in the k^{th} block are

$$H_{n,k} := \{ (k-1)m_n + 1, (k-1)m_n + 2, \dots, km_n \}, \quad k \in [l_n].$$
(4.3)

We define our blocking-based *ideal penalty loadings*

$$\widehat{\Upsilon}_{i}^{0} := \operatorname{diag}\left(\widehat{\upsilon}_{i,1}^{0}, \dots, \widehat{\upsilon}_{i,pq}^{0}\right), \quad i \in [p],$$

$$(4.4)$$

with

$$\widehat{v}_{i,j}^{0} := \sqrt{\frac{1}{n} \sum_{k=1}^{l_n} \left(\sum_{t \in H_{n,k}} \varepsilon_{t,i} Z_{t-1,j} \right)^2}, \quad i \in [p], \quad j \in [pq].$$

$$(4.5)$$

We define the score vector $\mathbf{S}_{n,i}$ as the (negative) gradient of the loss \widehat{Q}_i in (2.5) at $\boldsymbol{\beta}_{0i}$ upon normalizing by the ideal penalty loadings $\widehat{\boldsymbol{\Upsilon}}_i^0$,

$$\boldsymbol{S}_{n,i} := (\widehat{\boldsymbol{\Upsilon}}_{i}^{0})^{-1} (-\nabla \widehat{Q}_{i}(\boldsymbol{\beta}_{0i})) = \frac{2}{n} \sum_{t=1}^{n} (\widehat{\boldsymbol{\Upsilon}}_{i}^{0})^{-1} \boldsymbol{Z}_{t-1} \varepsilon_{t,i}, \quad i \in [p].$$
(4.6)

The score $S_{n,i}$ represents the estimation noise in problem (2.4), with the ideal penalty loadings yielding a form of self-normalization of the score. The self-normalization of the $S_{n,i}$ allows us to invoke recent moderate deviation theory for sums of dependent random variables developed in Gao et al. (2022). Specifically, with c and γ_n given in (2.6), and the penalty level λ_n^* given in (2.7), we show in Lemma 2 that $\lambda_n^*/n \ge c \max_{i \in [p]} ||S_{n,i}||_{\ell_{\infty}}$ with high probability.

The ideal penalty loadings in (4.4)–(4.5) are infeasible, since the innovations $\{\varepsilon_t\}_{t\in\mathbb{Z}}$ are unobservable. To overcome this problem, we prove that the loadings arising from Algorithm

⁶For simplicity of notation, we treat m_n as a divisor of n, such that both $m_n \in \mathbb{N}$ and $l_n \in \mathbb{N}$.

1 are "close" to the ideal loadings. More precisely, akin to Belloni et al. (2012, p. 2387), we refer to an arbitrary collection of penalty loadings $\widehat{\Upsilon}_i = \text{diag}(\widehat{v}_{i,1}, \ldots, \widehat{v}_{i,pq}), i \in [p]$, as asymptotically valid, if there are non-negative random scalars $\widehat{\ell} := \widehat{\ell}_n$ and $\widehat{u} := \widehat{u}_n, n \in \mathbb{N}$, and a constant $u \in [1, \infty)$, such that

$$\widehat{\ell}\widehat{v}_{i,j}^{0} \leqslant \widehat{v}_{i,j} \leqslant \widehat{u}\widehat{v}_{i,j}^{0} \text{ for all } (i,j) \in [p] \times [pq],$$

$$(4.7)$$

with probability approaching one as well as

$$\hat{\ell} \xrightarrow{\mathrm{P}} 1 \quad \text{and} \quad \hat{u} \xrightarrow{\mathrm{P}} u.$$
 (4.8)

We now present a master lemma that provides prediction and estimation error guarantees. The proof is adapted from Belloni et al. (2012, Lemma 6) to a form suitable for our purposes. Similar statements can be found in Bickel et al. (2009), Loh and Wainwright (2012) and Negahban et al. (2012) among others. To state the lemma, denote the smallest and largest ideal penalty loading $\hat{v}_{\min}^0 := \min_{(i,j) \in [p] \times [pq]} \hat{v}_{i,j}^0$ and $\hat{v}_{\max}^0 := \max_{(i,j) \in [p] \times [pq]} \hat{v}_{i,j}^0$, respectively, and their ratio $\hat{\mu}_0 := \hat{v}_{\max}^0 / \hat{v}_{\min}^0$.

Lemma 1 (Master Lemma). With the constant c given in (2.6) and the score vectors $\{S_{n,i}\}_{i\in[p]}$ given by (4.6), suppose that the penalty level satisfies $\lambda/n \ge c \max_{i\in[p]} \|S_{n,i}\|_{\ell_{\infty}}$, and that for some random scalars $\hat{\ell}$ and \hat{u} , the penalty loadings $\{\hat{\Upsilon}_i\}_{i=1}^p$ satisfy (4.7) with $1/c < \hat{\ell} \le \hat{u}$. Then the Lasso estimates $\hat{\beta}_i := \hat{\beta}_i(\lambda, \hat{\Upsilon}_i), i \in [p]$, arising from λ and $\{\hat{\Upsilon}_i\}_{i=1}^p$ via (2.4) satisfy the error bounds

$$\begin{aligned} \max_{i\in[p]} \|\widehat{\boldsymbol{\beta}}_{i} - \boldsymbol{\beta}_{0i}\|_{2,n} &\leq \left(\widehat{u} + \frac{1}{c}\right) \frac{\lambda \widehat{v}_{\max}^{0} \sqrt{s}}{n \widehat{\kappa}_{\widehat{c}_{0}\widehat{\mu}_{0}}}, \\ \max_{i\in[p]} \|\widehat{\boldsymbol{\beta}}_{i} - \boldsymbol{\beta}_{0i}\|_{\ell_{1}} &\leq \left(\widehat{u} + \frac{1}{c}\right) (1 + \widehat{c}_{0}\widehat{\mu}_{0}) \frac{\lambda s \widehat{v}_{\max}^{0}}{n \widehat{\kappa}_{\widehat{c}_{0}\widehat{\mu}_{0}}^{2}} \quad and \\ \max_{i\in[p]} \|\widehat{\boldsymbol{\beta}}_{i} - \boldsymbol{\beta}_{0i}\|_{\ell_{2}} &\leq \left(\widehat{u} + \frac{1}{c}\right) \frac{\lambda \widehat{v}_{\max}^{0} \sqrt{s}}{n \widehat{\kappa}_{\widehat{c}_{0}\widehat{\mu}_{0}}^{2}}, \end{aligned}$$

where $\hat{c}_0 := (\hat{u}c+1)/(\hat{\ell}c-1)$, and $\hat{\kappa}_{\hat{c}_0\hat{\mu}_0}$ is defined via (4.2) with $C = \hat{c}_0\hat{\mu}_0$.

In order to convert the above error guarantees into convergence rate results of the form (2.11)-(2.13) for Algorithm 1, we proceed by establishing:

1. Deviation bound: $P(\lambda_n^*/n \ge c \max_{i \in [p]} \| \boldsymbol{S}_{n,i} \|_{\ell_{\infty}}) \to 1$ holds for λ_n^* given in (2.7) and $\{\boldsymbol{S}_{n,i}\}_{i \in [p]}$ in (4.6).

- 2. Ideal penalty loadings control: There are constants $\underline{v}^0, \overline{v}^0 \in (0, \infty)$ such that $P(\underline{v}^0 \leqslant \widehat{v}^0_{\min} \leqslant \widehat{v}^0_{\max} \leqslant \overline{v}^0) \to 1$. This finding also yields $P(\widehat{\mu}_0 \leqslant \overline{v}^0/\underline{v}^0) \to 1$.
- 3. Restricted eigenvalue bound: $P(\widehat{\kappa}^2_{\widehat{c}_0\widehat{\mu}_0} \ge d/2) \to 1.$
- 4. Asymptotic validity of data-driven penalty loadings: The (final) data-driven penalty loadings $\{\widehat{v}_{i,j}^{(K)}\}_{(i,j)\in[p]\times[pq]}$ obtained from Algorithm 1 are asymptotically valid in the sense of (4.7)–(4.8). This implies control of the random $\widehat{\ell}$, \widehat{u} , and thus \widehat{c}_0 .

In the following four subsections we formalize Items 1-4 in the above enumeration, from which Theorem 1 will follow.

Item 1: Deviation Bound

Recall c and γ_n given in (2.6), the penalty level λ_n^* in (2.7) and the scores $\{S_{n,i}\}_{i \in [p]}$ based on ideal penalty loadings in (4.6).

Lemma 2 (Deviation Bound). Let Assumptions 1 and 2 hold, and suppose that $(\ln p)^5 = o(n)$. Then

$$\mathbf{P}\left(\lambda_{n}^{*}/n \ge c \max_{i \in [p]} \left\| \boldsymbol{S}_{n,i} \right\|_{\ell_{\infty}} \right) \to 1.$$

Item 2: Ideal Penalty Loadings Control

The following lemma states that the ideal penalty loadings (4.4) and (4.5) are bounded from above and away from zero with probability approaching one. We obtain this control over the ideal penalty loadings (in Lemma 3) as well as over the restricted eigenvalue (in Lemma 5) by establishing a new maximal inequality (Lemma B.2) for sums of dependent sub-Weibull random variables, which may be of independent interest.

Lemma 3 (Ideal Penalty Loadings Control). Let Assumptions 1, 2.1 and 2.2 hold, and suppose that $(\ln p)^{40/(3\alpha)} = o(n)$ for α provided by Assumption 1.2. Then there are constants $\underline{v}^0, \overline{v}^0 \in (0, \infty)$, depending only on a_1, a_2, α, b_1 and b_2 , such that

$$P\left(\underline{v}^{0} \leqslant \widehat{v}_{\min}^{0} \leqslant \widehat{v}_{\max}^{0} \leqslant \overline{v}^{0}\right) \to 1,$$
(4.9)

and, thus, $P(\widehat{\mu}_0 \leq \overline{\upsilon}^0/\underline{\upsilon}^0) \to 1$.

Item 3: Restricted Eigenvalue Bound

A key ingredient in bounding the restricted eigenvalue $\hat{\kappa}^2_{\hat{c}_0\hat{\mu}_0}$ away from zero is to prove that the empirical covariance matrix

$$\widehat{\boldsymbol{\Sigma}}_{\boldsymbol{Z}} := \frac{1}{n} \mathbf{X}^{\top} \mathbf{X} = \frac{1}{n} \sum_{t=0}^{n-1} \boldsymbol{Z}_t \boldsymbol{Z}_t^{\top}$$

of the regressors $\{\boldsymbol{Z}_t\}_{t=0}^{n-1}$ is close to $\boldsymbol{\Sigma}_{\boldsymbol{Z}} = \mathrm{E}[\boldsymbol{Z}_0\boldsymbol{Z}_0^{\top}].$

Lemma 4 (Covariance Consistency). Let Assumptions 1.1, 1.2 and 2 hold, and suppose that $(\ln p)^{1+4/\alpha} = o(n)$ for α provided by Assumption 1.2. Then

$$\max_{(i,j)\in [pq]^2} |(\widehat{\boldsymbol{\Sigma}}_{\boldsymbol{Z}})_{i,j} - (\boldsymbol{\Sigma}_{\boldsymbol{Z}})_{i,j}| \lesssim_{\mathrm{P}} \sqrt{\frac{(\ln(pn))^{1+4/\alpha}}{n}} = o(1).$$

Combining the ideal penalty loading control of Lemma 3 with Lemma 4, we can bound the restricted eigenvalue away from zero.

Lemma 5 (Restricted Eigenvalue Bound). Let Assumptions 1.1, 1.2, 2 and 3 hold, suppose that $s^2(\ln(pn))^{1+4/\alpha} = o(n)$ for α provided by Assumption 1.2, and consider any collection $\{\widehat{\Upsilon}_i\}_{i\in[p]}$ of asymptotically valid penalty loadings in the sense of (4.7) holding with probability approaching one and (4.8). Then

$$P\left(\widehat{\kappa}_{\widehat{c}_{0}\widehat{\mu}_{0}}^{2} \geqslant d/2\right) \to 1,$$

where $\hat{c}_0 := (\hat{u}c+1)/(\hat{\ell}c-1)$, and $\hat{\kappa}_{\hat{c}_0\hat{\mu}_0}$ is defined via (4.2) with $C = \hat{c}_0\hat{\mu}_0$.

Item 4: Asymptotic Validity of Data-Driven Penalty Loadings

Lemma 5 applies to any asymptotically valid penalty loadings. The following result states that the data-driven penalty loadings obtained from Algorithm 1 satisfy this requirement.

Lemma 6 (Asymptotic Validity of Data-Driven Penalty Loadings). Let Assumptions 1, 2, 3 and 4 hold. Then the penalty loadings $\{\widehat{v}_{i,j}^{(K)}\}_{(i,j)\in[p]\times[pq]}$ constructed by Algorithm 1 with a fixed number of updates, $K \in \mathbb{N}_0$, are asymptotically valid.

The proof of Lemma 6 is inspired by that of Belloni et al. (2012, Lemma 11). The main idea behind the proof is to show that the penalty loadings of Algorithm 1 are close to the blocking-based ideal ones. To this end, we use our maximal inequality for sums of dependent sub-Weibull random variables (Lemma B.2).

5 Simulations

In this section we explore the finite-sample estimation error of the weighted Lasso tuned via Algorithm 1.

5.1 Data-Generating Processes

We set the number of Monte Carlo replications to 1,000 and for each design we generate samples of effective size n ranging from 100 to 2,000.⁷ We consider system sizes $p \in \{16, 32, 64, 128\}$. The largest system size is chosen to essentially match the number (127) of macroeconomic series in our empirical illustration in Section 6.

Three experimental designs are considered, all inspired by Kock and Callot (2015, Section 5). The intercepts are zero and the $\{\varepsilon_t\}_{t\in\mathbb{Z}}$ are independent centered Gaussian with a diagonal covariance matrix and .1 on the diagonal. The remaining details of the designs are as follows:

- **Design A:** The data-generating process (DGP) is a VAR(1) with $\Theta_{01} = (.5)\mathbf{I}_p$ implying a spectral radius $\rho(\Theta_{01})$ of .5. This setting is very row sparse as s = 1.
- Design B: The DGP is a VAR(4) with Θ₀₁ and Θ₀₄ block diagonal with diagonal blocks of size 4 × 4 with all entries equal to .15 and −.1, respectively. Θ₀₂ = Θ₀₃ = 0_{p×p}. Thus, s = 8 whereas the spectral radius ρ(Θ̃₀) of the companion matrix is .9.⁸
- **Design C:** The DGP is a VAR(1) with $\Theta_{01,i,j} = (-1)^{|i-j|} (.4)^{1+|i-j|}$, $i, j \in [p]$, such that the entries decay exponentially fast in magnitude as one moves away from the diagonal, yet they remain non-zero. Noting that s = p, this setting is included to investigate the effects of a violation of strict sparsity. However, Θ_{01} is a near-band matrix, which is a form of approximate sparsity. $\rho(\Theta_{01}) = .9$.

5.2 Implementation and Performance Measure

The weighted Lasso is implemented equation by equation according to Algorithm 1. We also implement the post-Lasso by running least squares on the set of variables selected by the weighted Lasso.⁹ Finally, for the sake of comparison, we implement the plain Lasso using

⁷When simulating under Design A, we initiate the process by drawing from its stationary distribution, which in this case can be solved for explicitly. When using Designs B and C, we initiate the process at all zeros (i.e. the unconditional mean) and use a burn-in of 10,000 periods.

⁸We deviate from the 5×5 blocks in Kock and Callot (2015, Experiment B, p. 333) to ensure that the block sizes are divisors of p.

⁹To be precise, our post-Lasso is tuned using a modified penalization algorithm, where refitting is done prior to constructing residuals, and the final estimates follow from refitting. See Section A.2 for details.

the BIC to choose the penalty level. Although this is a common choice in the literature, [cf., e.g., Kock and Callot (2015)], we stress that there is currently no theoretical foundation for doing so. The resulting BIC-Lasso is also implemented equation by equation.

For all procedures, the intercept is (correctly) enforced to be zero and the number of lags included is the smallest one ensuring that the model is well-specified (i.e. one in Designs A and C, but four in Design B). The number of parameters to be estimated therefore ranges from 256 in a VAR(1) of size p = 16 to 65,536 in a VAR(4) of size p = 128.

To illustrate Theorem 1, we study the maximum row-wise ℓ_2 -estimation error,

Estimation Error :=
$$\max_{i \in [p]} \left\| \widehat{\beta}_i - \beta_{0i} \right\|_{\ell_2},$$
 (5.1)

for which we report the average across the 1,000 Monte Carlo replications. In addition, Section 6 contains an empirical forecasting exercise based on macroeconomic data.

5.3 Results

The estimation errors as defined in (5.1) are depicted in Figure 5.3; the top panel contains the results for Design A, the middle one for Design B, and the bottom one for Design C. The horizontal dashed line is the error of the estimator setting all parameters equal to zero a benchmark one would hope to beat. In line with our theoretical results, the estimation error is decreasing in n for fixed p, but increasing in p for fixed n. This also holds in the approximately sparse Design C. It is worth noting that choosing the tuning parameter by the BIC often leads to a worse performance than the naïve "all zero" estimator. This is never the case for the weighted Lasso with tuning parameters chosen according to our Algorithm 1.

6 Empirical Illustration

We apply the methods from the above simulations to forecast a large set of macroeconomic variables using the Federal Reserve Economic Data (FRED) monthly data (MD) database. This database is maintained and regularly updated by the Federal Reserve Bank of St. Louis and described in detail on Michael W. McCracken's website.¹⁰ The data is pre-processed in a standard manner using the Matlab code on McCracken's website leaving us with 758 observations on p = 127 macroeconomic variables covering March 1959 through April 2022.¹¹

¹⁰https://research.stlouisfed.org/econ/mccracken/fred-databases/. See also McCracken and Ng (2016).

¹¹The data pre-processing amounts to carrying out (deterministic) stationarity inducing transformations (prepare_missing.m), then removing outliers (remove_outliers.m), and finally replacing missing values with the unconditional average of the corresponding series (as in the initialization of factors_em.m).



Figure 5.1: Estimation errors calculated according to (5.1) and averaged over the Monte Carlo replications. The dashed horizontal line indicates the error of the "all zero" estimator.

6.1 Forecasting

In each of the last 120 months (i.e. ten years) of the sample we forecast the p = 127 variables one month ahead. Specifically, we estimate VAR(q) models of orders $q \in [12]$ using a rolling estimation window of size n = 758 - 120 - 12 = 626, and create one-month-ahead outof-sample forecasts as $\hat{Y}_{t+1} := \hat{\mu} + \sum_{j=1}^{q} \hat{\Theta}_j Y_{t+1-j}$. The three methods (weighted Lasso, post-Lasso and BIC-Lasso) are implemented as in the simulations in Section 5, but we now include an unpenalized intercept in each equation to account for non-zero means of the variables.¹² For each method, each q, and each month $t \in \{639, 640, \ldots, 758\}$ to be forecast, we calculate the forecast errors $\hat{Y}_t - Y_t$. Due to the different scaling of the variables, we then calculate the inverse-variance-weighted squared forecast error (IVWSFE)

IVWSFE_t :=
$$\sum_{i=1}^{p} (\widehat{Y}_{t,i} - Y_{t,i})^2 / \widehat{\sigma}_{Y_i}^2$$
,

where $\hat{\sigma}_{Y_i}^2$ denotes the sample variance of the *i*th variable (over the entire series), and finally average over the 120 forecasts to get the mean IVWSFE (MIVWSFE).

The MIVWSFEs relative to the MIVWSFE of the VAR(1) estimated by the weighted Lasso are plotted in Figure 6.1. It is seen that the weighted Lasso tuned according to Algorithm 1 and its post-Lasso variant dominate the BIC-Lasso irrespectively of the number of lags, q, included in the model. For VAR models with more than five lags the MIVWSFE of the BIC-Lasso is up to 25% higher than that of its competitors. We also note that the MIVWSFE of the weighted Lasso and the post-Lasso hardly vary with q.¹³

References

- ADAMEK, R., S. SMEEKES, AND I. WILMS (2023): "Lasso inference for high-dimensional time series," *Journal of Econometrics*, 235, 1114–1143.
- BABII, A., E. GHYSELS, AND J. STRIAUKAS (2022): "Machine learning time series regressions with an application to nowcasting," *Journal of Business & Economic Statistics*, 40, 1094–1106.
- BASU, S. AND G. MICHAILIDIS (2015): "Regularized estimation in sparse high-dimensional time series models," *The Annals of Statistics*, 43, 1535–1567.

¹²See Section A.1 for a modification of Algorithm 1, which includes unpenalized intercepts.

¹³We also experimented with equation by equation ordinary least squares (OLS). With a lag order of one, OLS led to a MIVWSFE 40 percent higher than our VAR(1) Lasso benchmark. For higher lag orders, the implied design matrix repeatedly fell short of full rank, thus preventing a meaningful comparison.



- BELLONI, A., D. CHEN, V. CHERNOZHUKOV, AND C. HANSEN (2012): "Sparse models and methods for optimal instruments with an application to eminent domain," *Econometrica*, 80, 2369–2429.
- BICKEL, P. J., Y. RITOV, AND A. B. TSYBAKOV (2009): "Simultaneous analysis of Lasso and Dantzig selector," *The Annals of Statistics*, 1705–1732.
- CHEN, X., Q.-M. SHAO, W. B. WU, AND L. XU (2016): "Self-normalized Cramér-type moderate deviations under dependence," *The Annals of Statistics*, 44, 1593–1617.
- CHERNOZHUKOV, V., W. K. HÄRDLE, C. HUANG, AND W. WANG (2021): "LASSOdriven inference in time and space," *The Annals of Statistics*, 49, 1702–1735.
- DAVIS, R. A., P. ZANG, AND T. ZHENG (2016): "Sparse vector autoregressive modeling," Journal of Computational and Graphical Statistics, 25, 1077–1096.
- GAO, L., Q.-M. SHAO, AND J. SHI (2022): "Refined Cramér-type moderate deviation theorems for general self-normalized sums with applications to dependent random variables and winsorized mean," *The Annals of Statistics*, 50, 673–697.
- GUO, S., Y. WANG, AND Q. YAO (2016): "High-dimensional and banded vector autoregressions," *Biometrika*, 103, 889–903.

- HAN, F. AND H. LIU (2013): "Transition matrix estimation in high dimensional time series," in *International conference on machine learning*, PMLR, 172–180.
- HAN, F., H. LU, AND H. LIU (2015): "A direct estimation of high dimensional stationary vector autoregressions," *Journal of Machine Learning Research*, 16, 3115–3150.
- JAVANMARD, A. AND A. MONTANARI (2014): "Confidence intervals and hypothesis testing for high-dimensional regression," *The Journal of Machine Learning Research*, 15, 2869– 2909.
- KOCK, A. B. AND L. CALLOT (2015): "Oracle inequalities for high dimensional vector autoregressions," *Journal of Econometrics*, 186, 325–344.
- KUCHIBHOTLA, A. K. AND A. CHAKRABORTTY (2022): "Moving beyond sub-Gaussianity in high-dimensional statistics: Applications in covariance estimation and linear regression," *Information and Inference: A Journal of the IMA*, 11, 1389–1456.
- LOH, P.-L. AND M. J. WAINWRIGHT (2012): "High-dimensional regression with noisy and missing data: Provable guarantees and nonconvexity," *The Annals of Statistics*, 40, 1637–1664.
- LÜTKEPOHL, H. (2005): New introduction to multiple time series analysis, Springer Berlin, Heidelberg.
- MASINI, R. P., M. C. MEDEIROS, AND E. F. MENDES (2022): "Regularized estimation of high-dimensional vector autoregressions with weakly dependent innovations," *Journal* of *Time Series Analysis*, 43, 532–557.
- MCCRACKEN, M. W. AND S. NG (2016): "FRED-MD: A Monthly Database for Macroeconomic Research," *Journal of Business & Economic Statistics*, 34, 574–589.
- MEDEIROS, M. C. AND E. F. MENDES (2016): "*l*₁-regularization of high-dimensional timeseries models with non-Gaussian and heteroskedastic errors," *Journal of Econometrics*, 191, 255–271.
- MIAO, L., P. C. B. PHILLIPS, AND L. SU (2023): "High-dimensional VARs with common factors," *Journal of Econometrics*, 233, 155–183.
- NEGAHBAN, S. N., P. RAVIKUMAR, M. J. WAINWRIGHT, AND B. YU (2012): "A unified framework for high-dimensional analysis of *M*-estimators with decomposable regularizers," *Statistical Science*, 27, 538–557.

- SONG, S. AND P. J. BICKEL (2011): "Large vector auto regressions," arXiv preprint arXiv:1106.3915.
- TIBSHIRANI, R. (1996): "Regression shrinkage and selection via the lasso," Journal of the Royal Statistical Society. Series B (Methodological), 267–288.
- VAN DE GEER, S., P. BÜHLMANN, Y. RITOV, AND R. DEZEURE (2014): "On asymptotically optimal confidence regions and tests for high-dimensional models," *The Annals of Statistics*, 42, 1166–1202.
- VLADIMIROVA, M., S. GIRARD, H. NGUYEN, AND J. ARBEL (2020): "Sub-Weibull distributions: Generalizing sub-Gaussian and sub-Exponential properties to heavier tailed distributions," *Stat*, 9, e318.
- WONG, K. C., Z. LI, AND A. TEWARI (2020): "Lasso guarantees for β -mixing heavy-tailed time series," *The Annals of Statistics*, 48, 1124–1142.
- WU, W. B. (2005): "Nonlinear system theory: Another look at dependence," *Proceedings* of the National Academy of Sciences, 102, 14150–14154.
- WU, W. B. AND X. SHAO (2004): "Limit theorems for iterated random functions," *Journal* of Applied Probability, 41, 425–436.
- WU, W.-B. AND Y. N. WU (2016): "Performance bounds for parameter estimates of high-dimensional linear models with correlated errors," *Electronic Journal of Statistics*, 10, 352–379, publisher: Institute of Mathematical Statistics and Bernoulli Society.
- ZHANG, C.-H. AND S. S. ZHANG (2014): "Confidence intervals for low dimensional parameters in high dimensional linear models," *Journal of the Royal Statistical Society: Series* B (Statistical Methodology), 76, 217–242.
- ZHANG, X. AND G. CHENG (2018): "Gaussian approximation for high dimensional vector under physical dependence," *Bernoulli*, 24, 2640–2675.

Supplementary Appendices

Table of Contents

Α	Modified Data-Driven Penalization Algorithms	2
	A.1 Including Unpenalized Intercepts	2
	A.2 Incorporating Refitting	3
В	Auxiliary Results	5
	B.1 Functional Dependence Measures for Strictly Stationary Processes	5
	B.2 Functional Dependence Measure Calculus	5
	B.3 Maximal Inequality with Functional Dependence	6
С	Moderate Deviation with Geometric Moment Contraction	10
D	Proofs of Lemmas 1–6 and Theorem 1	12
D	Proofs of Lemmas 1–6 and Theorem 1 D.1 Proof of Master Lemma (Lemma 1)	12 12
D	Proofs of Lemmas 1–6 and Theorem 1 D.1 Proof of Master Lemma (Lemma 1) D.2 Proof of Deviation Bound (Lemma 2)	12 12 13
D	Proofs of Lemmas 1–6 and Theorem 1 D.1 Proof of Master Lemma (Lemma 1)	12 12 13 18
D	Proofs of Lemmas 1–6 and Theorem 1D.1Proof of Master Lemma (Lemma 1)D.2Proof of Deviation Bound (Lemma 2)D.3Proof of Ideal Penalty Loadings Control (Lemma 3)D.4Proof of Restricted Eigenvalue Bound (Lemma 5)	12 12 13 18 24
D	 Proofs of Lemmas 1–6 and Theorem 1 D.1 Proof of Master Lemma (Lemma 1) D.2 Proof of Deviation Bound (Lemma 2) D.3 Proof of Ideal Penalty Loadings Control (Lemma 3) D.4 Proof of Restricted Eigenvalue Bound (Lemma 5) D.5 Proof of Asymptotic Validity of Penalty Loadings from Algorithm 1 (Lemma 6) 	 12 13 18 24 26

A Modified Data-Driven Penalization Algorithms

A.1 Including Unpenalized Intercepts

Incorporating model intercepts $\mu_0 \in \mathbb{R}^p$, (2.3) becomes

$$oldsymbol{Y}_t = oldsymbol{\mu}_0 + \sum_{j=1}^q oldsymbol{\Theta}_{0j}oldsymbol{Y}_{t-j} + oldsymbol{arepsilon}_t, \quad t \in \mathbb{Z}.$$

If one is confident that intercepts belong in the model, then they should not be penalized. The resulting Lasso estimator therefore takes the form

$$\left(\widehat{\mu}_{i}(\lambda,\widehat{\boldsymbol{\Upsilon}}_{i}),\widehat{\boldsymbol{\beta}}_{i}(\lambda,\widehat{\boldsymbol{\Upsilon}}_{i})\right) \in \operatorname*{argmin}_{(\mu,\boldsymbol{\beta})\in\mathbb{R}^{1+pq}} \left\{ \frac{1}{n} \sum_{t=1}^{n} \left(Y_{t,i} - \mu - \boldsymbol{Z}_{t-1}^{\top}\boldsymbol{\beta} \right)^{2} + \frac{\lambda}{n} \|\widehat{\boldsymbol{\Upsilon}}_{i}\boldsymbol{\beta}\|_{\ell_{1}} \right\}.$$
(A.1)

The first-order condition for the intercept reveals the relation

$$\widehat{\mu}_i(\lambda, \widehat{\Upsilon}_i) = \overline{Y}_i - \overline{Z}_{-1}\widehat{\beta}_i(\lambda, \widehat{\Upsilon}_i), \qquad (A.2)$$

where

$$\overline{Y}_i := \frac{1}{n} \sum_{t=1}^n Y_{t,i} \quad \text{and} \quad \overline{Z}_{-1} := \frac{1}{n} \sum_{t=0}^{n-1} Z_t.$$
(A.3)

Concentrating out the intercept in (A.1), we see that

$$\widehat{\boldsymbol{\beta}}_{i}(\lambda,\widehat{\boldsymbol{\Upsilon}}_{i}) \in \operatorname*{argmin}_{\boldsymbol{\beta} \in \mathbb{R}^{pq}} \left\{ \frac{1}{n} \sum_{t=1}^{n} (\ddot{Y}_{t,i} - \ddot{\boldsymbol{Z}}_{t-1}^{\top} \boldsymbol{\beta})^{2} + \frac{\lambda}{n} \|\widehat{\boldsymbol{\Upsilon}}_{i} \boldsymbol{\beta}\|_{\ell_{1}} \right\},$$
(A.4)

where

$$\ddot{Y}_{t,i} := Y_{t,i} - \overline{Y}_i \text{ and } \ddot{Z}_{t-1} := Z_{t-1} - \overline{Z}_{-1}.$$
 (A.5)

The problem in (A.4) is of the form in (2.4) except that we have here replaced the outcome variable and regressors with their demeaned counterparts. The data-driven penalization algorithm now runs parallel to Algorithm 1.

Algorithm A.1 (Data-Driven Penalization with Unpenalized Intercepts). Initialize: Specify the penalty level in (A.4) as

$$\lambda_n^* := 2c\sqrt{n}\Phi^{-1}\left(1 - \gamma_n/(2p^2q)\right),$$

and specify the initial penalty loadings as

$$\widehat{v}_{i,j}^{(0)} := \sqrt{\frac{1}{n} \sum_{t=1}^{n} \ddot{Y}_{t,i}^2 \ddot{Z}_{t-1,j}^2}, \quad i \in [p], \quad j \in [pq]$$

Use λ_n^* and $\widehat{\Upsilon}_i^{(0)} := \operatorname{diag}(\widehat{v}_{i,1}^{(0)}, \dots, \widehat{v}_{i,pq}^{(0)})$ to compute a Lasso estimate $\widehat{\beta}_i^{(0)} := \widehat{\beta}_i(\lambda_n^*, \widehat{\Upsilon}_i^{(0)})$ via (A.4) for each $i \in [p]$. Store the residuals $\widehat{\widetilde{\varepsilon}}_{t,i}^{(0)} := \ddot{Y}_{t,i} - \ddot{Z}_{t-1}^{\top}\widehat{\beta}_i^{(0)}, t \in [n], i \in [p], and$ set k = 1.

Update: While $k \leq K$, specify the penalty loadings as

$$\widehat{v}_{i,j}^{(k)} := \sqrt{\frac{1}{n} \sum_{t=1}^{n} \widehat{\varepsilon}_{t,i}^{[k-1]2} \ddot{Z}_{t-1,j}^2}, \quad i \in [p], \quad j \in [pq].$$

Use λ_n^* and $\widehat{\Upsilon}_i^{(k)} := \operatorname{diag}(\widehat{v}_{i,1}^{(k)}, \dots, \widehat{v}_{i,pq}^{(k)})$ to compute a Lasso estimate as $\widehat{\beta}_i^{(k)} := \widehat{\beta}_i(\lambda_n^*, \widehat{\Upsilon}_i^{(k)})$ via (A.4) for each $i \in [p]$. Store the residuals $\widehat{\widetilde{\varepsilon}}_{t,i}^{(k)} := \ddot{Y}_{t,i} - \ddot{Z}_{t-1}^{\top}\widehat{\beta}_i^{(k)}, t \in [n], i \in [p], and$ increment $k \leftarrow k+1$.

Algorithm A.1 yields the (slope) estimates $\widehat{\beta}_i(\lambda_n^*, \widehat{\Upsilon}_i^{(K)}), i \in [p]$. Using (A.2) and (A.3), the implied intercept estimates are $\widehat{\mu}_i(\lambda_n^*, \widehat{\Upsilon}_i^{(K)}) = \overline{Y}_i - \overline{Z}_{-1}^{\top} \widehat{\beta}_i(\lambda_n^*, \widehat{\Upsilon}_i^{(K)}), i \in [p]$. Each step in Algorithm A.1 could involve refitting after Lasso selection, thus replacing the Lasso residuals with their post-Lasso counterparts—see Section A.2 for details.

A.2 Incorporating Refitting

In this section we modify Algorithm 1 to accommodate refitting after variable selection. Specifically, we consider a *post-Lasso* estimator, which satisfies

$$\widetilde{\boldsymbol{\beta}}_{i} := \widetilde{\boldsymbol{\beta}}_{i}(\lambda, \widehat{\boldsymbol{\Upsilon}}_{i}) \in \underset{\boldsymbol{\beta} \in \mathbb{R}^{pq}:}{\operatorname{supp}(\boldsymbol{\beta}) \subseteq \operatorname{supp}(\widehat{\boldsymbol{\beta}}_{i})}$$
(A.6)

which depends on the penalty level λ and the penalty loadings $\widehat{\Upsilon}_i$ through the support $\operatorname{supp}(\widehat{\beta}_i)$ of the Lasso estimator $\widehat{\beta}_i := \widehat{\beta}_i(\lambda, \widehat{\Upsilon}_i)$ in (2.4), with $\operatorname{supp}(\beta) := \{j \in [pq] : \beta_j \neq 0\}$. The support constraint allows the post-Lasso to refit coefficients of variables selected by the Lasso, while keeping zeros at zero. The data-driven penalization algorithm incorporating refitting goes as follows.

Algorithm A.2 (Data-Driven Penalization with Refitting). Initialize: Specify the penalty level in (2.4) as

$$\lambda_n^* := 2c\sqrt{n}\Phi^{-1}\left(1 - \gamma_n/(2p^2q)\right),$$

and specify the initial penalty loadings as

$$\widehat{v}_{i,j}^{(0)} := \sqrt{\frac{1}{n} \sum_{t=1}^{n} Y_{t,i}^2 Z_{t-1,j}^2}, \quad i \in [p], \quad j \in [pq].$$

Use λ_n^* and $\widehat{\Upsilon}_i^{(0)} := \operatorname{diag}(\widehat{\upsilon}_{i,1}^{(0)}, \dots, \widehat{\upsilon}_{i,pq}^{(0)})$ to compute an estimate $\widetilde{\beta}_i^{(0)} := \widetilde{\beta}_i(\lambda_n^*, \widehat{\Upsilon}_i^{(0)})$ via post-Lasso (A.6) for each $i \in [p]$. Store the residuals $\widehat{\varepsilon}_{t,i}^{(0)} := Y_{t,i} - \mathbb{Z}_{t-1}^{\top} \widetilde{\beta}_i^{(0)}, t \in [n], i \in [p]$, and set k = 1.

Update: While $k \leq K$, specify the penalty loadings as

$$\widehat{\upsilon}_{i,j}^{(k)} := \sqrt{\frac{1}{n} \sum_{t=1}^{n} \widehat{\varepsilon}_{t,i}^{(k-1)2} Z_{t-1,j}^2}, \quad i \in [p], \quad j \in [pq].$$

Use λ_n^* and $\widehat{\Upsilon}_i^{(k)} := \operatorname{diag}(\widehat{v}_{i,1}^{(k)}, \dots, \widehat{v}_{i,pq}^{(k)})$ to compute an estimate $\widetilde{\beta}_i^{(k)} := \widetilde{\beta}_i(\lambda_n^*, \widehat{\Upsilon}_i^{(k)})$ via post-Lasso (A.6) for each $i \in [p]$. Store the residuals $\widehat{\varepsilon}_{t,i}^{(k)} := Y_{t,i} - \mathbb{Z}_{t-1}^{\top} \widetilde{\beta}_i^{(k)}, t \in [n], i \in [p]$, and increment $k \leftarrow k+1$.

Algorithm A.2 could include unpenalized intercepts via an initial demeaning of variables—see Section A.1 for the necessary adjustments.

B Auxiliary Results

B.1 Functional Dependence Measures for Strictly Stationary Processes

Let $\{\varepsilon_t\}_{t\in\mathbb{Z}}$ be independent and identically distributed \mathbb{R}^{p_1} -valued random variables, and let $\{X_t\}_{t\in\mathbb{Z}}$ be \mathbb{R}^{p_2} -valued random variables having the causal representation

$$\boldsymbol{X}_t = \boldsymbol{G}(\ldots, \boldsymbol{\varepsilon}_{t-1}, \boldsymbol{\varepsilon}_t), \quad t \in \mathbb{Z},$$

where G is a measurable function of the present and past ε_t . Strict stationarity of the stochastic process $\{X_t\}_{t\in\mathbb{Z}}$ follows from G not depending on t. Let $\{\varepsilon_t^*\}_{t\in\mathbb{Z}}$ be an independent copy of $\{\varepsilon_t\}_{t\in\mathbb{Z}}$. Assuming that the i^{th} coordinate $X_{t,i} = G_i(\ldots, \varepsilon_{t-1}, \varepsilon_t)$ of X_t satisfies $E[|X_{t,i}|^r] < \infty$ for some $r \in [1, \infty)$, we define the functional (or physical) dependence measure for the coordinate process $\{X_{t,i}\}_{t\in\mathbb{Z}}$

$$\Delta_{r}^{X_{\cdot i}}(h) := \left\| X_{t,i} - G_{i}\left(\dots, \boldsymbol{\varepsilon}_{t-h-1}^{*}, \boldsymbol{\varepsilon}_{t-h}^{*}, \boldsymbol{\varepsilon}_{t-h+1}, \dots, \boldsymbol{\varepsilon}_{t} \right) \right\|_{r}, \quad h \in \mathbb{N}_{0},$$

where $\Delta_r^{X_{i}}(h)$ is interpreted as zero for h < 0. The coordinate process $\{X_{t,i}\}_{t \in \mathbb{Z}}$ is said to be a geometric moment contraction [Wu and Shao (2004)] if there are constants $b_1, b_2 \in (0, \infty)$ and $\tau \in (0, 1]$ such that

$$\Delta_r^{X_{\cdot i}}(h) \leqslant b_1 \mathrm{e}^{-b_2 h^{\tau}} \text{ for all } h \in \mathbb{N}_0.$$
(B.1)

B.2 Functional Dependence Measure Calculus

Lemma B.1. Let $\{(X_{t,1}, X_{t,2})\}_{t\in\mathbb{Z}}$, be a strictly stationary causal process with values in \mathbb{R}^2 and $\mathbb{E}[|X_{0,1}|^{2r}] < \infty$ as well as $\mathbb{E}[|X_{0,2}|^{2r}] < \infty$ for some $r \in [1, \infty)$. Then the product process $\{X_{t,1}X_{t,2}\}_{t\in\mathbb{Z}}$ is strictly stationary and causal, and its functional dependence measure satisfies

$$\Delta_{r}^{X_{\cdot 1}X_{\cdot 2}}\left(h\right) \leqslant \left\|X_{0,2}\right\|_{2r} \Delta_{2r}^{X_{\cdot 1}}\left(h\right) + \left\|X_{0,1}\right\|_{2r} \Delta_{2r}^{X_{\cdot 2}}\left(h\right), \quad h \in \mathbb{N}_{0}.$$

Proof. Since $\{(X_{t,1}, X_{t,2})\}_{t\in\mathbb{Z}}$ is a strictly stationary causal process, it has the representation

$$(X_{t,1}, X_{t,2}) = \boldsymbol{G}(\ldots, \boldsymbol{\varepsilon}_{t-1}, \boldsymbol{\varepsilon}_t),$$

for some i.i.d. process $\{\boldsymbol{\varepsilon}_t\}_{t\in\mathbb{Z}}$ and some measurable function \boldsymbol{G} . The composition of a continuous and measurable function is measurable, so the product process $\{X_{t,1}X_{t,2}\}_{t\in\mathbb{Z}}$ is both strictly stationary and causal and has finite r^{th} absolute moment by Hölder's inequality.

Let $\{\boldsymbol{\varepsilon}_t^*\}_{t\in\mathbb{Z}}$ be an independent copy of $\{\boldsymbol{\varepsilon}_t\}_{t\in\mathbb{Z}}, h\in\mathbb{N}_0$ and denote

$$(X'_{t,1},X'_{t,2}) := \boldsymbol{G}(\ldots,\boldsymbol{\varepsilon}^*_{t-h-1},\boldsymbol{\varepsilon}^*_{t-h},\boldsymbol{\varepsilon}_{t-h+1},\ldots,\boldsymbol{\varepsilon}_t).$$

Then

$$\begin{aligned} \Delta_{r}^{X_{\cdot1}X_{\cdot2}}\left(h\right) &= \left\|X_{t,1}X_{t,2} - X_{t,1}'X_{t,2} + X_{t,1}'X_{t,2} - X_{t,1}'X_{t,2}'\right\|_{r} \\ &\leqslant \left\|\left(X_{t,1} - X_{t,1}'\right)X_{t,2}\right\|_{r} + \left\|X_{t,1}'\left(X_{t,2} - X_{t,2}'\right)\right\|_{r} \\ &\leqslant \left\|X_{t,2}\right\|_{2r} \left\|X_{t,1} - X_{t,1}'\right\|_{2r} + \left\|X_{t,1}'\right\|_{2r} \left\|X_{t,2} - X_{t,2}'\right\|_{2r} \end{aligned}$$
(sub-additivity)
$$&= \left\|X_{0,2}\right\|_{2r} \Delta_{2r}^{X_{\cdot1}}\left(h\right) + \left\|X_{0,1}\right\|_{2r} \Delta_{2r}^{X_{\cdot2}}\left(h\right). \end{aligned}$$
(stationarity)

B.3 Maximal Inequality with Functional Dependence

We here establish a maximal inequality for an array of $T \in \mathbb{N}$ possibly dependent random vectors of dimension d_T possibly growing with T.

Lemma B.2. Let $\{X_t\}_{t\in\mathbb{Z}}$ be a strictly stationary causal process with each X_t a random element of \mathbb{R}^{d_T} for some $d_T \in \{2, 3, ...\}$, and $\max_{i\in[d_T]} \|X_{0,i}\|_{\psi_{\xi}} \leq K$ for some constants $\xi, K \in (0, \infty)$. Then for any $r \in [1, \infty)$, any $m_T \in [T]$ and $l_T := \lceil T/m_T \rceil$

$$\max_{i \in [d_T]} \left| \sum_{t=1}^T \left(X_{T,i} - \mathbf{E} \left[X_{0,i} \right] \right) \right| \lesssim_{\mathbf{P}} T d_T^{1/r} \Delta_{r,T} + m_T \sqrt{l_T \ln \left(d_T T \right)} + m_T (\ln(d_T T))^{\overline{\xi}}$$

as $T \to \infty$, where

$$\Delta_{r,T} := \max_{i \in [d_T]} \max_{u \in \{m_T+1,\dots,2m_T\}} \Delta_r^{X,i}(u)$$

and $\overline{\xi} := 1/\xi + 1/\min\{\xi, 1\}.$

Remark B.1. If $u \mapsto \Delta_r^{X,i}(u)$ is decreasing for all $i \in [d_T]$, one may upper bound $\Delta_{r,T}$ by $\max_{i \in [d_T]} \Delta_r^{X,i}(m_T)$.

PROOF OF LEMMA B.2. The overall strategy for proving Lemma B.2 is two-fold. First, we approximate the original (functionally) dependent process with an *m*-dependent process [see, e.g., (Zhang and Cheng, 2018)]. Suitably construed, the *m*-dependent process can next be handled using tools designed for independent observations.

To this end, write $X_t = G(\ldots, \varepsilon_{t-1}, \varepsilon_t)$, which is possible by strict stationarity and causality. Fix $m_T \in [T]$ and, thus, $l_T = \lceil T/m_T \rceil$ and abbreviate $m := m_T$, $l := l_T$,

and $d := d_T$. Define

$$\boldsymbol{Y}_k := \sum_{t=(k-1)m+1}^{\min\{km,T\}} \boldsymbol{X}_t \quad \text{and} \quad \boldsymbol{Y}_k^{(m)} := \mathrm{E}[\boldsymbol{Y}_k \mid \{\boldsymbol{\varepsilon}_t\}_{t=(k-2)m+1}^{km}], \quad k \in [l],$$

Then, per iterated expectations and the triangle inequality,

$$\max_{i \in [d]} \left| \sum_{t=1}^{T} \left(X_{t,i} - \mathbf{E} \left[X_{0,i} \right] \right) \right| = \max_{i \in [d]} \left| \sum_{k=1}^{l} \left(Y_{k,i} - \mathbf{E} \left[Y_{k,i}^{(m)} \right] \right) \right|$$

$$\leq \max_{i \in [d]} \left| \sum_{k=1}^{l} \left(Y_{k,i} - Y_{k,i}^{(m)} \right) \right| + \max_{i \in [d]} \left| \sum_{k=1}^{l} \left(Y_{k,i}^{(m)} - \mathbf{E} \left[Y_{k,i}^{(m)} \right] \right) \right| =: \mathbf{I}_{T} + \mathbf{I}\mathbf{I}_{T}.$$

We bound each of the right-hand side terms, starting with I_T . Since $\max_{i \in [n]} ||X_{0,i}||_{\psi_{\xi}} \leq K$, all absolute moments of the $X_{0,i}$ exist. Hence, for any $r \in [1, \infty)$

$$\mathbf{E}\left[\mathbf{I}_{T}\right] \leqslant \left\| \max_{i \in [d]} \left| \sum_{k=1}^{l} \left(Y_{k,i} - Y_{k,i}^{(m)} \right) \right| \right\|_{r}$$
(Jensen)

$$\leq d^{1/r} \max_{i \in [d]} \left\| \sum_{k=1}^{l} \left(Y_{k,i} - Y_{k,i}^{(m)} \right) \right\|_{r}$$
(Jensen)

$$\leq d^{1/r} \max_{i \in [d]} \sum_{k=1}^{l} \left\| Y_{k,i} - Y_{k,i}^{(m)} \right\|_{r}$$
(sub-additivity)

$$\leq d^{1/r} \max_{i \in [d]} \sum_{k=1}^{l} \sum_{t=(k-1)m+1}^{\min(km,T)} \left\| X_{t,i} - \mathbf{E} \left[X_{t,i} \mid \{ \boldsymbol{\varepsilon}_u \}_{u=(k-2)m+1}^{km} \right] \right\|_r \qquad (\text{sub-additivity})$$

$$\leq d^{1/r} \max_{i \in [d]} \sum_{k=1}^{l} \sum_{t=(k-1)m+1}^{km} \left\| X_{t,i} - \mathbf{E} \left[X_{t,i} \mid \{ \boldsymbol{\varepsilon}_u \}_{u=(k-2)m+1}^{km} \right] \right\|_r.$$
 (non-negativity)

Fix $i \in [d], k \in [l]$ and $t \in \{(k-1)m+1, \ldots, km\}$. Letting $\{\boldsymbol{\varepsilon}_t^*\}_{t \in \mathbb{Z}}$ be an independent copy of $\{\boldsymbol{\varepsilon}_t\}_{t \in \mathbb{Z}}$, we have

$$\begin{aligned} \left\| X_{t,i} - \mathbf{E} \left[X_{t,i} \mid \{ \boldsymbol{\varepsilon}_u \}_{u=(k-2)m+1}^{km} \right] \right\|_r^r \\ &= \mathbf{E} \left[\left| X_{t,i} - \mathbf{E} \left[X_{t,i} \mid \{ \boldsymbol{\varepsilon}_u \}_{u=(k-2)m+1}^t \right] \right|^r \right] \qquad (\text{causality}) \\ &= \mathbf{E} \left[\left| \mathbf{E} \left[X_{t,i} - G_i(\dots, \boldsymbol{\varepsilon}_{(k-2)m-1}^*, \boldsymbol{\varepsilon}_{(k-2)m}^*, \boldsymbol{\varepsilon}_{(k-2)m+1}, \dots, \boldsymbol{\varepsilon}_t) \mid \{ \boldsymbol{\varepsilon}_u \}_{u=-\infty}^t \right] \right|^r \right]. \qquad (\text{causality}) \end{aligned}$$

Applying Jensen's inequality (conditionally) to the right-hand side, we get

$$\begin{split} & \operatorname{E}\left[\left|\operatorname{E}\left[X_{t,i} - G_{i}(\ldots, \boldsymbol{\varepsilon}_{(k-2)m-1}^{*}, \boldsymbol{\varepsilon}_{(k-2)m}^{*}, \boldsymbol{\varepsilon}_{(k-2)m+1}, \ldots, \boldsymbol{\varepsilon}_{t}\right) \mid \{\boldsymbol{\varepsilon}_{u}\}_{u=-\infty}^{t}\right]\right|^{r}\right] \\ & \leqslant \operatorname{E}\left[\left|X_{t,i} - G_{i}(\ldots, \boldsymbol{\varepsilon}_{(k-2)m-1}^{*}, \boldsymbol{\varepsilon}_{(k-2)m}^{*}, \boldsymbol{\varepsilon}_{(k-2)m+1}, \ldots, \boldsymbol{\varepsilon}_{t})\right|^{r}\right] \\ & = \left[\Delta_{r}^{X_{\cdot,i}}(t - (k-2)m)\right]^{r}. \end{split}$$

Combining the previous two displays and adding up, by a change of variables, we see that

$$\sum_{t=(k-1)m+1}^{km} \left\| X_{t,i} - \mathbf{E} \left[X_{t,i} \mid \{ \varepsilon_u \}_{u=(k-2)m+1}^{km} \right] \right\|_r \leqslant \sum_{t=(k-1)m+1}^{km} \Delta_r^{X_{\cdot,i}} \left(t - (k-2)m \right)$$
$$= \sum_{u=m+1}^{2m} \Delta_r^{X_{\cdot,i}} \left(u \right)$$
$$\leqslant m \max_{u \in \{m+1,\dots,2m\}} \Delta_r^{X_{\cdot,i}} \left(u \right) \right).$$

Since the right-hand bound does not depend on k, it follows from our calculations that

$$\mathbf{E}[\mathbf{I}_T] \leqslant mld^{1/r} \max_{i \in [d]} \max_{u \in \{m+1,\dots,2m\}} \Delta_r^{X_{\cdot,i}}(u)) = mld^{1/r} \Delta_{r,T}.$$

By Markov's inequality and $ml \leq 2T$, we get $I_T \leq_P T d^{1/r} \Delta_{r,T}$.

To bound II_T , observe that

$$II_T \leqslant \max_{i \in [d]} \left| \sum_{k \leqslant l \text{ odd}} \left(Y_{k,i}^{(m)} - E[Y_{k,i}^{(m)}] \right) \right| + \max_{i \in [d]} \left| \sum_{k \leqslant l \text{ even}} \left(Y_{k,i}^{(m)} - E[Y_{k,i}^{(m)}] \right) \right| \\ =: \Pi_T^{\text{odd}} =: \Pi_T^{\text{even}}$$

We set up for two applications of Kuchibhotla and Chakrabortty (2022, Theorem 3.4) for the odd and even parts, respectively. We only bound II_T^{odd} ; the argument for II_T^{even} is analogous.

First, for any $r \in [1, \infty)$, $i \in [d]$, and $k \in [l]$,

$$\left\|Y_{k,i}^{(m)} - \mathbf{E}[Y_{k,i}^{(m)}]\right\|_{r} \leq 2\|Y_{k,i}^{(m)}\|_{r} \leq 2\|Y_{k,i}\|_{r} \leq 2m \|X_{0,i}\|_{r} \leq 2Kmr^{1/\xi},$$

which means that

$$\left\| m^{-1} (Y_{k,i}^{(m)} - \mathbf{E}[Y_{k,i}^{(m)}]) \right\|_r \leq 2Kr^{1/\xi}.$$

From the previous display, an application of Vladimirova et al. (2020, Theorem 1) implies

the existence of a constant $C_{\xi,K} \in (0,\infty)$, depending only on ξ and K, such that

$$\max_{i \in [d], k \in [l]} \mathbf{E} \left[\exp \left\{ \left| m^{-1} \left(Y_{k,i}^{(m)} - \mathbf{E}[Y_{k,i}^{(m)}] \right) \right|^{\xi} / C_{K,\xi}^{\xi} \right\} \right] \leqslant 2.$$

Let $l^{\text{odd}} := l_T^{\text{odd}} := |\{k \in [l] : k \text{ odd}\}|$. The l^{odd} summands $m^{-1}(\boldsymbol{Y}_k^{(m)} - \mathbb{E}[\boldsymbol{Y}_k^{(m)}])$, are independent random elements of \mathbb{R}^d . It therefore follows from Kuchibhotla and Chakrabortty (2022, Theorem 3.4) (with t there set to $\ln(dT)$) that there is a constant $C'_{K,\xi} \in (0,\infty)$, depending only on ξ and K, such that with probability at least 1 - 3/(dT),

$$\max_{i \in [d]} \left| \frac{1}{l^{\text{odd}}} \sum_{k \leq l \text{ odd}} \frac{Y_{k,i}^{(m)} - \mathbb{E}[Y_{k,i}^{(m)}]}{m} \right| \tag{B.2}$$

$$\leq 7 \sqrt{\frac{\Gamma_T \left(\ln(dT) + \ln d \right)}{l^{\text{odd}}}} + \frac{C'_{K,\xi} \ln \left(2l^{\text{odd}} \right)^{1/\xi} \left(\ln \left(dT \right) + \ln d \right)^{1/\min\{1,\xi\}}}{l^{\text{odd}}}, \tag{B.3}$$

$$\Gamma_T := \max_{i \in [d]} \frac{1}{l^{\text{odd}}} \sum_{k=1}^{l^{\text{odd}}} E\left[\left| \frac{Y_{k,i}^{(m)} - E[Y_{k,i}^{(m)}]}{m} \right|^2 \right].$$

From previous calculations, setting r = 2, we know that for all $i \in [d]$ and $k \in [l]$,

$$\mathbf{E}\left[\left|\frac{Y_{k,i}^{(m)} - \mathbf{E}[Y_{k,i}^{(m)}]}{m}\right|^{2}\right] = \left\|\frac{Y_{k,i}^{(m)} - \mathbf{E}[Y_{k,i}^{(m)}]}{m}\right\|_{2}^{2} \leqslant 2^{2}K^{2} \cdot 2^{2/\xi},$$

so $\Gamma_T \leq 2^{2+2/\xi} K^2$. Multiplying by l^{odd} and m, through simple majorizations, we see that for some constant $C''_{K,\xi} \in (0,\infty)$, depending only on ξ and K, with probability approaching one,

$$\max_{i \in [d]} \left| \sum_{k \leqslant l \text{ odd}} \left(Y_{k,i}^{(m)} - \mathbf{E}[Y_{k,i}^{(m)}] \right) \right| \leqslant C_{K,\xi}'' \left(m \sqrt{l \ln(dT)} + m \left(\ln(dT) \right)^{\overline{\xi}} \right).$$

It follows that $\operatorname{II}_T^{\operatorname{odd}} \leq_{\operatorname{P}} m\sqrt{l \ln(dT)} + m(\ln(dT))^{\overline{\xi}}$. Hence, $\operatorname{II}_T \leq_{\operatorname{P}} m\sqrt{l \ln(dT)} + m(\ln(dT))^{\overline{\xi}}$.

C Moderate Deviation with Geometric Moment Contraction

Let $\{X_t\}_{t\in\mathbb{Z}}$ be a strictly stationary causal process with elements X_t taking values in \mathbb{R} . Define the sums

$$S_{k,h} := \sum_{t=k+1}^{k+h} X_t, \quad k \in \mathbb{N}_0, \quad h \in \mathbb{N},$$

and abbreviate $S_n := S_{0,n}$. Consider the following assumptions.

Assumption C.1. $E[X_0] = 0$ and $E[X_0^4] < \infty$.

Assumption C.2. There is a constant $\omega \in (0, \infty)$, such that $\mathbb{E}[S_{k,h}^2] \ge \omega^2 h$ for all $k \in \mathbb{N}_0$ and all $h \in \mathbb{N}$.

Assumption C.3. There are constants $b_1, b_2 \in (0, \infty)$, such that $\Delta_4^{X}(h) \leq b_1 e^{-b_2 h}$ for all $h \in \mathbb{N}_0$.

Let $n \in \mathbb{N}$ and define the block size $m_n := \lfloor n^{1/5} \rfloor$ and number of blocks $l_n := \lfloor n/m_n \rfloor$, and let $H_{n,k}, k \in [l_n]$, hold the block indices as in (4.3). Define the block-normalized sum

$$T_{n} := \frac{\sum_{k=1}^{l_{n}} \sum_{t \in H_{n,k}} X_{t}}{\sqrt{\sum_{k=1}^{l_{n}} \left(\sum_{t \in H_{n,k}} X_{t}\right)^{2}}}$$

The following lemma is a consequence of Gao et al. (2022, Theorem 3.3).

Lemma C.1. Let Assumptions C.1, C.2 and C.3 hold. Then there are constants A and g_0 both in $(0, \infty)$, depending only on b_1, b_2 and ω , such that

$$x \in [0, g_0 n^{1/10}] \implies \frac{\mathrm{P}(|T_n| \ge x)}{2[1 - \Phi(x)]} \le 1 + A\left(\frac{1 + x^2}{n^{1/5}}\right).$$

Proof. If Assumptions C.1, C.2 and C.3 hold for the random process $\{X_t\}$, then they hold for $\{-X_t\}$ as well with the same constants. Applying Gao et al. (2022, Theorem 3.3) twice (with their $\alpha = 1/5$ and their $\tau = 1$), we therefore get that

$$x \in \left[0, g_0 n^{1/10}\right] \implies \max\left\{\frac{\mathcal{P}\left(T_n \ge x\right)}{1 - \Phi\left(x\right)}, \frac{\mathcal{P}\left(-T_n \ge x\right)}{1 - \Phi\left(x\right)}\right\} \leqslant 1 + A\left(\frac{1 + x^2}{n^{1/5}}\right),$$

for constants A and g_0 with the claimed dependencies. The claim then follows from $P(|T_n| \ge x) = P(T_n \ge x) + P(-T_n \ge x)$.

Next, let $\{X_t\}_{t\in\mathbb{Z}}$ be a strictly stationary causal process with elements X_t taking values in \mathbb{R}^{d_n} , where we take d_n to be at least two. Let $T_{n,j}$ denote the block-normalized sum using the j^{th} coordinate process $\{X_{t,j}\}_{t\in\mathbb{N}}$ of $\{X_t\}_{t\in\mathbb{Z}}$, such that

$$T_{n,j} = \frac{\sum_{k=1}^{l_n} \sum_{t \in H_{n,k}} X_{t,j}}{\sqrt{\sum_{k=1}^{l_n} \left(\sum_{t \in H_{n,k}} X_{t,j}\right)^2}}, \quad j \in [d_n].$$

We invoke the growth condition:

Assumption C.4. $\{\gamma_n\}_{n=1}^{\infty}$ is a non-random sequence in (0,1) satisfying $\ln(1/\gamma_n) \lesssim \ln(nd_n)$.

Lemma C.2. Let all coordinate processes $\{\{X_{t,j}\}_{i\in\mathbb{Z}}\}_{j=1}^{d_n}$ satisfy Assumptions C.1, C.2 and C.3 for the same constants b_1, b_2 and ω , let Assumption C.4 hold, and suppose that $\ln(d_n) = o(n^{1/5})$. Then there is a constant $N \in \mathbb{N}$ such that

$$n \ge N \implies \mathcal{P}\left(\max_{j \in [d_n]} |T_{n,j}| \ge \Phi^{-1}\left(1 - \frac{\gamma_n}{2d_n}\right)\right) \le 2\gamma_n.$$

Proof. Under Assumption C.4, using $d_n \ge 2$ and the Gaussian quantile bound $\Phi^{-1}(1-z) \le \sqrt{2\ln(1/z)}, z \in (0,1)$, we get

$$0 < \Phi^{-1} \left(1 - \gamma_n / (2d_n) \right) \leqslant \sqrt{2 \ln (2d_n / \gamma_n)} \leqslant C_1 \sqrt{\ln(d_n)}$$
 (C.1)

for some constant $C_1 \in (0, \infty)$. The growth condition $\ln(d_n) = o(n^{1/5})$ equips us with an $N_1 \in \mathbb{N}$ such that for the constant $g_0 \in (0, \infty)$ provided by Lemma C.1,

$$n \ge N_1 \implies \Phi^{-1} \left(1 - \gamma_n / (2d_n) \right) \le g_0 n^{1/10}.$$

For such $n \ge N_1$,

$$P\left(\max_{j\in[d_n]}|T_{n,j}| \ge \Phi^{-1}\left(1-\frac{\gamma_n}{2d_n}\right)\right)$$

$$\leqslant \sum_{j=1}^{d_n} P\left(|T_{n,j}| \ge \Phi^{-1}\left(1-\frac{\gamma_n}{2d_n}\right)\right) \qquad (\text{union bound})$$

$$\leqslant \sum_{j=1}^{d_n} 2\left[1-\Phi\left(\Phi^{-1}\left(1-\frac{\gamma_n}{2d_n}\right)\right)\right]\left[1+A\left(\frac{1+\left[\Phi^{-1}\left(1-\frac{\gamma_n}{2d_n}\right)\right]^2}{n^{1/5}}\right)\right] \qquad (\text{Lemma C.1})$$

$$\leqslant \gamma_n \left[1+A\left(\frac{1+C_1^2\ln(d_n)}{n^{1/5}}\right)\right].$$

Again using $\ln(d_n) = o(n^{1/5})$, there is a constant $N_2 \in \mathbb{N}$ such that $n \ge N_2$ implies $A[1 + C_1^2 \ln(d_n)/n^{1/5}] \le 1$. The claim now follows from setting $N = \max\{N_1, N_2\}$.

D Proofs of Lemmas 1–6 and Theorem 1

D.1 Proof of Master Lemma (Lemma 1)

PROOF OF LEMMA 1. The proof of the lemma is similar to that of Belloni et al. (2012, Lemma 6), which, in turn, builds on a strategy attributed to Bickel et al. (2009). Let $T_i := \{j \in [pq] : \beta_{0i,j} \neq 0\}$. Recall that $\widehat{\Upsilon}_i^0$ denotes the diagonal matrix diag $(\widehat{\upsilon}_{i,1}^0, \ldots, \widehat{\upsilon}_{i,pq}^0)$ of ideal penalty loadings and let $\widehat{\delta}_i := \widehat{\beta}_i - \beta_{0i}$. Expanding the quadratic part (2.5) of the objective function in (2.4), when $\lambda/n \ge c \|S_i\|_{\ell_{\infty}}$ we get

$$\begin{aligned} \widehat{Q}_{i}(\widehat{\boldsymbol{\beta}}_{i}) - \widehat{Q}_{i}(\boldsymbol{\beta}_{0i}) &= \frac{1}{n} \sum_{t=1}^{n} (\boldsymbol{Z}_{t-1}^{\top} \widehat{\boldsymbol{\delta}}_{i})^{2} - \frac{2}{n} \sum_{t=1}^{n} \varepsilon_{t,i} \boldsymbol{Z}_{t-1}^{\top} \widehat{\boldsymbol{\delta}}_{i} \\ &= \|\widehat{\boldsymbol{\delta}}_{i}\|_{2,n}^{2} - \underbrace{\left[\frac{2}{n} \sum_{t=1}^{n} (\widehat{\boldsymbol{\Upsilon}}_{i}^{0})^{-1} \boldsymbol{Z}_{t-1} \varepsilon_{t,i}\right]^{\top}}_{=\boldsymbol{S}_{n,i}^{\top}} (\widehat{\boldsymbol{\Upsilon}}_{i}^{0} \widehat{\boldsymbol{\delta}}_{i}) \\ &= \|\widehat{\boldsymbol{\delta}}_{i}\|_{2,n}^{2} - \|\boldsymbol{S}_{n,i}\|_{\ell_{\infty}} \|\widehat{\boldsymbol{\Upsilon}}_{i}^{0} \widehat{\boldsymbol{\delta}}_{i}\|_{\ell_{1}} \\ &\geq \|\widehat{\boldsymbol{\delta}}_{i}\|_{2,n}^{2} - \frac{\lambda}{cn} \|\widehat{\boldsymbol{\Upsilon}}_{i}^{0} \widehat{\boldsymbol{\delta}}_{i}\|_{\ell_{1}}, \end{aligned}$$
(Hölder)

which we can write as

$$\widehat{Q}_{i}(\widehat{\beta}_{i}) - \widehat{Q}_{i}(\beta_{0i}) \ge \|\widehat{\delta}_{i}\|_{2,n}^{2} - \frac{\lambda}{cn} \left(\|\widehat{\Upsilon}_{i}^{0}\widehat{\delta}_{iT_{i}}\|_{\ell_{1}} + \|\widehat{\Upsilon}_{i}^{0}\widehat{\delta}_{iT_{i}^{c}}\|_{\ell_{1}} \right).$$
(D.1)

Since $\widehat{\beta}_i$ is a minimizer of $\widehat{Q}_i(\cdot) + (\lambda/n) \|\widehat{\Upsilon}_i \cdot\|_{\ell_1}$,

$$\widehat{Q}_{i}(\widehat{\beta}_{i}) - \widehat{Q}_{i}(\beta_{0i}) \leqslant \frac{\lambda}{n} \left(\|\widehat{\Upsilon}_{i}\beta_{0i}\|_{\ell_{1}} - \|\widehat{\Upsilon}_{i}\widehat{\beta}_{i}\|_{\ell_{1}} \right) \leqslant \frac{\lambda}{n} \left(\|\widehat{\Upsilon}_{i}\widehat{\delta}_{iT_{i}}\|_{\ell_{1}} - \|\widehat{\Upsilon}_{i}\widehat{\delta}_{iT_{i}^{c}}\|_{\ell_{1}} \right),$$

so from the penalty loadings $\widehat{\Upsilon}_i$ satisfying (4.7), we arrive at

$$\widehat{Q}_{i}(\widehat{\boldsymbol{\beta}}_{i}) - \widehat{Q}_{i}(\boldsymbol{\beta}_{0i}) \leqslant \frac{\lambda}{n} \left(\widehat{\boldsymbol{u}} \| \widehat{\boldsymbol{\Upsilon}}_{i}^{0} \widehat{\boldsymbol{\delta}}_{iT_{i}} \|_{\ell_{1}} - \widehat{\ell} \| \widehat{\boldsymbol{\Upsilon}}_{i}^{0} \widehat{\boldsymbol{\delta}}_{iT_{i}^{c}} \|_{\ell_{1}} \right).$$
(D.2)

The two inequalities (D.1) and (D.2) combine to yield

$$\|\widehat{\boldsymbol{\delta}}_{i}\|_{2,n}^{2} + \frac{\lambda}{n} \left(\widehat{\ell} - \frac{1}{c}\right) \|\widehat{\boldsymbol{\Upsilon}}_{i}^{0}\widehat{\boldsymbol{\delta}}_{iT_{i}^{c}}\|_{\ell_{1}} \leqslant \frac{\lambda}{n} \left(\widehat{u} + \frac{1}{c}\right) \|\widehat{\boldsymbol{\Upsilon}}_{i}^{0}\widehat{\boldsymbol{\delta}}_{iT_{i}}\|_{\ell_{1}}.$$
 (D.3)

Since $1/c < \hat{\ell} \leq \hat{u}$, we deduce that

$$\widehat{v}_{\min}^{0} \|\widehat{\boldsymbol{\delta}}_{iT_{i}^{c}}\|_{\ell_{1}} \leqslant \|\widehat{\boldsymbol{\Upsilon}}_{i}^{0}\widehat{\boldsymbol{\delta}}_{iT_{i}^{c}}\|_{\ell_{1}} \leqslant \frac{\widehat{u}+1/c}{\widehat{\ell}-1/c} \|\widehat{\boldsymbol{\Upsilon}}_{i}^{0}\widehat{\boldsymbol{\delta}}_{iT_{i}}\|_{\ell_{1}} \leqslant \frac{\widehat{u}+1/c}{\widehat{\ell}-1/c} \widehat{v}_{\max}^{0} \|\widehat{\boldsymbol{\delta}}_{iT_{i}}\|_{\ell_{1}}.$$
(D.4)

It follows that

$$\|\widehat{\boldsymbol{\delta}}_{iT_i^c}\|_{\ell_1} \leqslant \frac{\widehat{u}+1/c}{\widehat{\ell}-1/c} \cdot \frac{\widehat{v}_{\max}^0}{\widehat{v}_{\min}^0} \|\widehat{\boldsymbol{\delta}}_{iT_i}\|_{\ell_1} = \widehat{c}_0 \widehat{\mu}_0 \|\widehat{\boldsymbol{\delta}}_{iT_i}\|_{\ell_1},$$

i.e. $\widehat{\delta}_i$ lies in the restricted set $\mathcal{R}_{\widehat{c}_0\widehat{\mu}_0,T_i}$ defined via (4.1) with $C = \widehat{c}_0\widehat{\mu}_0$ and $T = T_i$. From the definition (4.2) of the restricted eigenvalue we see that $\|\widehat{\delta}_i\|_{\ell_2} \leq \|\widehat{\delta}_i\|_{2,n}/\widehat{\kappa}_{\widehat{c}_0\widehat{\mu}_0}$, so by the Cauchy-Schwarz inequality and recalling that $s = \max_{i \in [p]} |T_i|$ (cf. (2.10))

$$\|\widehat{\mathbf{\Upsilon}}_{i}^{0}\widehat{\boldsymbol{\delta}}_{iT_{i}}\|_{\ell_{1}} \leqslant \widehat{v}_{\max}^{0} \|\widehat{\boldsymbol{\delta}}_{iT_{i}}\|_{\ell_{1}} \leqslant \widehat{v}_{\max}^{0} \sqrt{s} \|\widehat{\boldsymbol{\delta}}_{iT_{i}}\|_{\ell_{2}} \leqslant \widehat{v}_{\max}^{0} \sqrt{s} \|\widehat{\boldsymbol{\delta}}_{i}\|_{\ell_{2}} \leqslant \frac{\widehat{v}_{\max}^{0} \sqrt{s} \|\boldsymbol{\delta}_{i}\|_{2,n}}{\widehat{\kappa}_{\widehat{c}_{0}\widehat{\mu}_{0}}}.$$
 (D.5)

Hence, continuing our string of inequalities from (D.3), we see that

$$\|\widehat{\boldsymbol{\delta}}_{i}\|_{2,n}^{2} + \underbrace{\frac{\lambda}{n}\left(\widehat{\ell} - \frac{1}{c}\right)\|\widehat{\boldsymbol{\Upsilon}}_{i}^{0}\widehat{\boldsymbol{\delta}}_{iT_{i}^{c}}\|_{\ell_{1}}}_{\geqslant 0} \leqslant \frac{\lambda}{n}\left(\widehat{u} + \frac{1}{c}\right)\frac{\widehat{\upsilon}_{\max}^{0}\sqrt{s}\|\widehat{\boldsymbol{\delta}}_{i}\|_{2,n}}{\widehat{\kappa}_{\widehat{c}_{0}\widehat{\mu}_{0}}},$$

from which we arrive at the prediction error bound

$$\|\widehat{\boldsymbol{\delta}}_i\|_{2,n} \leqslant \left(\widehat{u} + \frac{1}{c}\right) \frac{\lambda \widehat{v}_{\max}^0 \sqrt{s}}{n\widehat{\kappa}_{\widehat{c}_0\widehat{\mu}_0}}$$

As the right-hand bound is independent of i, the uniformity in $i \in [p]$ follows. The ℓ_2 error bound then follows from the already established $\|\widehat{\delta}_i\|_{\ell_2} \leq \|\widehat{\delta}_i\|_{2,n}/\widehat{\kappa}_{\widehat{c}_0\widehat{\mu}_0}$, and the ℓ_1 bound subsequently follows from $\|\widehat{\delta}_i\|_{\ell_1} \leq (1+\widehat{c}_0\widehat{\mu}_0) \|\widehat{\delta}_{iT_i}\|_{\ell_1}$ and $\|\widehat{\delta}_{iT_i}\|_{\ell_1} \leq \sqrt{s}\|\widehat{\delta}_i\|_{\ell_2}$. \Box

D.2 Proof of Deviation Bound (Lemma 2)

We first state and prove some supporting lemmas.

Lemma D.1. If Assumptions 1.1, 1.2, 2.1 and 2.2 hold, then, for α provided by Assumption 1.2, the following statements hold.

(1) For all $i \in [pq]$,

$$||Z_{0,i}||_{\psi_{\alpha}} \leqslant \frac{a_1 b_1}{1 - e^{-b_2}}.$$

(2) For any $r \in [1, \infty)$ and any $i \in [pq]$,

$$||Z_{0,i}||_r \leq \frac{a_1 b_1 r^{1/\alpha}}{1 - e^{-b_2}}.$$

(3) For any $r \in [1, \infty)$, any $i \in [pq]$ and any $j \in [pq]$

$$\|\varepsilon_{0,i}Z_{-1,j}\|_r \leqslant \frac{a_1^2 b_1 r^{2/\alpha}}{1 - e^{-b_2}}$$

(4) For any $i \in [p]$,

$$\left\|\boldsymbol{Z}_{0}^{\top}\boldsymbol{\beta}_{0i}\right\|_{\psi_{\alpha}} \leqslant \frac{a_{1}b_{1}\mathrm{e}^{-b_{2}}}{1-\mathrm{e}^{-b_{2}}}.$$

PROOF OF LEMMA D.1. Strict stationarity (which follows from Assumptions 1.1 and 2.1) allows us to represent the companion VAR(1) in (2.2) as an infinite vector moving average $[VMA(\infty)]$,

$$\boldsymbol{Z}_{t} = \sum_{h=0}^{\infty} \widetilde{\boldsymbol{\Theta}}_{0}^{h} \widetilde{\boldsymbol{\varepsilon}}_{t-h}, \quad t \in \mathbb{Z},$$
(D.6)

where the series converges absolutely almost surely. Moreover, under Assumption 2.2 the series converges in $\|\cdot\|_{\psi_{\alpha}}$ norm. To establish Claim 1, since for $i \in [pq]$, $Z_{t-1,i} = Y_{t-j,k}$ for some $(j,k) \in [q] \times [p]$, for any α provided by Assumption 1.2, we have

$$\begin{split} \|Z_{0,i}\|_{\psi_{\alpha}} &\leqslant \max_{j \in [p]} \|Y_{0,j}\|_{\psi_{\alpha}} \qquad (\text{strict stationarity}) \\ &= \max_{j \in [p]} \left\|\sum_{h=0}^{\infty} (\widetilde{\Theta}_{0}^{h})_{j,i} \widetilde{\varepsilon}_{0-h}\right\|_{\psi_{\alpha}} \qquad (\text{VMA}(\infty) \text{ representation}) \\ &= \max_{j \in [p]} \left\|\sum_{h=0}^{\infty} (\widetilde{\Theta}_{0}^{h})_{j,1:p} \varepsilon_{0-h}\right\|_{\psi_{\alpha}} \qquad (\widetilde{\varepsilon}_{p+1:pq} \equiv \mathbf{0}_{p(q-1)\times 1}) \\ &\leqslant \max_{j \in [p]} \sum_{h=0}^{\infty} \|(\widetilde{\Theta}_{0}^{h})_{j,1:p} \varepsilon_{0-h}\|_{\psi_{\alpha}} \qquad (\text{countable sub-additivity}) \\ &\leqslant \|\varepsilon_{0}\|_{\psi_{\alpha}} \max_{j \in [p]} \sum_{h=0}^{\infty} \|(\widetilde{\Theta}_{0}^{h})_{j,1:p}\|_{\ell_{2}} \qquad (\text{sub-Weibullness and stationarity}) \\ &\leqslant a_{1} \sum_{h=0}^{\infty} b_{1} e^{-b_{2}h} = \frac{a_{1}b_{1}}{1 - e^{-b_{2}}}. \qquad (\text{Assumptions 1.2 and 2.2}) \end{split}$$

Claim 2 is immediate from Claim 1 and the definition of a sub-Weibull norm. To establish

Claim 3, note that for any $r \in [1, \infty)$, $i \in [p]$ and $j \in [pq]$, using also independence, we get

$$\|\varepsilon_{0,i}Z_{-1,j}\|_{r} = \|\varepsilon_{0,i}\|_{r} \|Z_{0,j}\|_{r} \leqslant r^{1/\alpha} \|\varepsilon_{0,i}\|_{\psi_{\alpha}} \|Z_{0,j}\|_{r} \leqslant r^{1/\alpha} \|\varepsilon_{0}\|_{\psi_{\alpha}} \|Z_{0,j}\|_{r} \leqslant a_{1}r^{1/\alpha} \|Z_{0,j}\|_{r}.$$

Claim 3 now follows from Claim 2. For Claim 4, since β_{0i}^{\top} is the i^{th} row of $\widetilde{\Theta}_0$, letting $e_i \in \mathbb{R}^{pq}$ be the i^{th} unit vector, from (D.6) we see that

$$oldsymbol{Z}_0^ opoldsymbol{eta}_{0i} = \sum_{h=0}^\infty oldsymbol{eta}_{0i}^ op \widetilde{oldsymbol{\Theta}}_0^h \widetilde{oldsymbol{arepsilon}}_{0-h} = \sum_{h=0}^\infty oldsymbol{e}_i^ op \widetilde{oldsymbol{\Theta}}_0^{h+1} \widetilde{oldsymbol{arepsilon}}_{0-h} = \sum_{h=0}^\infty (\widetilde{oldsymbol{\Theta}}_0^{h+1})_{i,1:p} oldsymbol{arepsilon}_{0-h}.$$

It follows that

$$\begin{split} \left\| \boldsymbol{Z}_{0}^{\top} \boldsymbol{\beta}_{0i} \right\|_{\psi_{\alpha}} &\leq \sum_{h=0}^{\infty} \left\| (\widetilde{\boldsymbol{\Theta}}_{0}^{h+1})_{i,1:p} \boldsymbol{\varepsilon}_{0} \right\|_{\psi_{\alpha}} \qquad \text{(countable sub-additivity and stationarity)} \\ &\leq \left\| \boldsymbol{\varepsilon}_{0} \right\|_{\psi_{\alpha}} \sum_{h=0}^{\infty} \left\| (\widetilde{\boldsymbol{\Theta}}_{0}^{h+1})_{i,1:p} \right\|_{\ell_{2}} \qquad \text{(sub-Weibullness)} \\ &\leq a_{1} \sum_{h=0}^{\infty} b_{1} \mathrm{e}^{-b_{2}(h+1)} \qquad \text{(Assumptions 1.2 and 2.2)} \\ &= \frac{a_{1} b_{1} \mathrm{e}^{-b_{2}}}{1 - \mathrm{e}^{-b_{2}}}, \end{split}$$

which holds uniformly in $i \in [p]$.

Lemma D.2. If Assumption 2.2 holds, then

$$\max_{i \in [pq]} \left\| (\widetilde{\Theta}_0^h)_{i,1:p} \right\|_{\ell_2} \leq (1+b_1) \mathrm{e}^{b_2(q-1)} \mathrm{e}^{-b_2 h} \text{ for any } h \in \mathbb{N}_0.$$

PROOF OF LEMMA D.2. For q = 1 the result is immediate from Assumption 2.2. For q > 1, by construction of the companion matrix $\tilde{\Theta}_0$, we know that

$$(\widetilde{\Theta}_{0}^{1})_{p+1:2p,1:p} = \mathbf{I}_{p} \text{ and } (\widetilde{\Theta}_{0}^{1})_{kp+1:(k+1)p,1:p} = \mathbf{0}_{p \times p}, k \in \{2, \dots, q-1\}.$$

Moreover, we have the recursion

$$(\widetilde{\Theta}_0^h)_{kp+1:(k+1)p,1:p} = (\widetilde{\Theta}_0^{h-1})_{(k-1)p+1:kp,1:p}, \quad k \in \{1, \dots, q-1\}, \quad h \in \{2, 3, \dots\}.$$

Hence, for $h \ge 1$, it follows that

$$(\widetilde{\Theta}_{0}^{h})_{\cdot,1:p} = \begin{pmatrix} (\widetilde{\Theta}_{0}^{h})_{1:p,1:p} \\ (\widetilde{\Theta}_{0}^{h-1})_{1:p,1:p} \\ (\widetilde{\Theta}_{0}^{h-2})_{1:p,1:p} \\ \vdots \\ (\widetilde{\Theta}_{0}^{h-(q-2)})_{1:p,1:p} \\ (\widetilde{\Theta}_{0}^{h-(q-1)})_{1:p,1:p} \end{pmatrix},$$

with the convention that $(\widetilde{\Theta}_0^i)_{1:p,1:p} = \mathbf{0}_{p \times p}$ for integers i < 0 and $(\widetilde{\Theta}_0^0)_{1:p,1:p} = \mathbf{I}_p$. Using Assumption 2.2, for $h \ge 1$ we therefore get

$$\begin{aligned} \max_{i \in [pq]} \left\| (\widetilde{\Theta}_{0}^{h})_{i,1:p} \right\|_{\ell_{2}} &= \max_{j \in \{h - (q-1), \dots, h\}} \max_{i \in [p]} \left\| (\widetilde{\Theta}_{0}^{j})_{i,1:p} \right\|_{\ell_{2}} \\ &\leqslant \max_{j \in \{h - (q-1), \dots, h\}} \left\{ b_{1} e^{-b_{2}j} \mathbf{1}(j > 0) + \mathbf{1}(j = 0) \right\} \\ &\leqslant \max_{j \in \{h - (q-1), \dots, h\}} \left\{ (1 + b_{1}) e^{-b_{2}j} \right\} \\ &= (1 + b_{1}) e^{b_{2}(q-1)} e^{-b_{2}h}. \end{aligned}$$

Lemma D.3. Let Assumptions 1.1, 1.2 and 2 hold. Then for any $r \in [1, \infty)$, there are constants $C_1, C_2 \in (0, \infty)$, depending only on a_1, α, b_1, b_2, q and r, such that

- (1) $\max_{i \in [pq]} \Delta_r^{Z_{\cdot,i}}(h) \leqslant C_1 e^{-b_2 h}$ for any $h \in \mathbb{N}_0$, and
- (2) $\max_{i \in [p], j \in [pq]} \Delta_r^{\varepsilon_{\cdot,i} Z_{\cdot-1,j}}(h) \leqslant C_2 \mathrm{e}^{-b_2 h} \text{ for any } h \in \mathbb{N}_0.$

PROOF OF LEMMA D.3. Fix $r \in [1, \infty)$ and $i \in [p]$. Under Assumptions 1.1, 1.2, 2.1 and 2.2 the r^{th} absolute moments of both $Z_{t-1,i}$ and $\varepsilon_{t,i}Z_{t-1,j}$ are finite (cf. Lemma D.1, Parts 2 and 3). Thus, $\Delta_r^{Z_{\cdot,i}}(h)$ and $\Delta_r^{\varepsilon_{\cdot,i}Z_{\cdot-1,j}}(h)$ are well defined for all $h \in \mathbb{N}$. Defining

$$oldsymbol{G}\left(\ldots,oldsymbol{arepsilon}_{t-1},oldsymbol{arepsilon}_{t}
ight):=\sum_{h=0}^{\infty}(\widetilde{oldsymbol{\Theta}}_{0}^{h})_{\cdot,1:p}oldsymbol{arepsilon}_{t-h}.$$

one has the representation $Z_{t,i} = G_i(\ldots, \varepsilon_{t-1}, \varepsilon_t)$ for $i \in [pq]$, cf. (D.6). Letting $\{\varepsilon_t^*\}_{t \in \mathbb{Z}}$ be

an independent copy of $\{\boldsymbol{\varepsilon}_t\}_{t\in\mathbb{Z}}$, Claim 1 follows from

$$\begin{split} \Delta_{r}^{Z,i}(h) &= \left\| \sum_{\ell=h}^{\infty} (\widetilde{\Theta}_{0}^{\ell})_{i,1:p} \left(\varepsilon_{t-\ell} - \varepsilon_{t-\ell}^{*} \right) \right\|_{r} \qquad (\text{causality}) \\ &\leqslant \sum_{\ell=h}^{\infty} \left\| (\widetilde{\Theta}_{0}^{\ell})_{i,1:p} \left(\varepsilon_{t-\ell} - \varepsilon_{t-\ell}^{*} \right) \right\|_{r} \qquad (\text{countable sub-additivity}) \\ &\leqslant \sum_{\ell=h}^{\infty} r^{1/\alpha} \left\| (\widetilde{\Theta}_{0}^{\ell})_{i,1:p} \left(\varepsilon_{t-\ell} - \varepsilon_{t-\ell}^{*} \right) \right\|_{\psi_{\alpha}} \qquad (\text{marginal sub-Weibullness}) \\ &\leqslant r^{1/\alpha} \left\| \varepsilon_{0} - \varepsilon_{0}^{*} \right\|_{\psi_{\alpha}} \sum_{\ell=h}^{\infty} \left\| (\widetilde{\Theta}_{0}^{\ell})_{i,1:p} \right\|_{\ell_{2}} \qquad (\text{joint sub-Weibullness and stationarity}) \\ &\leqslant 2a_{1}r^{1/\alpha} \sum_{\ell=h}^{\infty} (1+b_{1})e^{b_{2}(q-1)}e^{-b_{2}\ell} \qquad (\text{countable sub-additivity and Lemma D.2}) \\ &= \underbrace{\frac{2a_{1}(1+b_{1})e^{b_{2}(q-1)}r^{1/\alpha}}{1-e^{-b_{2}}}e^{-b_{2}h}. \end{split}$$

To establish Claim 2, fix $i \in [p]$ and $j \in [pq]$. On the one hand, taking h = 0 leads to

$$\begin{aligned} \Delta_{r}^{\varepsilon_{\cdot,i}Z_{\cdot-1,j}}\left(0\right) &= \left\|\varepsilon_{t,i}Z_{t-1,j} - \varepsilon_{t,i}^{*}G_{j}\left(\dots,\varepsilon_{t-2}^{*},\varepsilon_{t-1}^{*}\right)\right\|_{r} \\ &\leqslant 2 \left\|\varepsilon_{t,i}Z_{t-1,j}\right\|_{r} \qquad \text{(sub-additivity)} \\ &\leqslant \frac{2a_{1}^{2}b_{1}r^{2/\alpha}}{1 - e^{-b_{2}}} \qquad \text{(Part 3 of Lemma D.1)} \\ &=: C' = C'e^{-b_{2}h}. \end{aligned}$$

On the other hand, for $h \in \mathbb{N}$,

Claim 2 now follows from setting $C_2 := \max\{C', a_1C_1r^{1/\alpha}\}.$

Based on these supporting lemmas, we are now able to give the proof of Lemma 2. PROOF OF LEMMA 2. We set up for an application of Lemma C.2. Consider the stochastic

processes $\{\{\varepsilon_{t,i}Z_{t-1,j}\}_{t\in\mathbb{Z}}\}, (i,j) \in [p] \times [pq]$, of which there are a total of $p^2q(=d_n)$. Since the $\{Z_t\}_{t\in\mathbb{Z}}$ admit the VMA(∞) representation in (D.6), each process $\{\{\varepsilon_{t,i}Z_{t-1,j}\}_{t\in\mathbb{Z}}\}, (i,j) \in [p] \times [pq]$, is both strictly stationary and causal with elements taking values in \mathbb{R} . Setting r = 4, Part 3 of Lemma D.1 and Part 2 of Lemma D.3 respectively show that there are constants $C, C' \in (0, \infty)$, depending only on a_1, α, b_1, b_2 and q, such that both $\max_{i\in[p],j\in[pq]} \|\varepsilon_{0,i}Z_{-1,j}\|_4 \leq C$ and $\max_{i\in[p],j\in[pq]} \Delta_4^{\varepsilon_{i,i}Z_{i-1,j}}(h) \leq C' e^{-b_2 h}$ for all $h \in \mathbb{N}$. Since the $\{\varepsilon_t\}_{t\in\mathbb{Z}}$ are centered, so are the processes $\{\{\varepsilon_{t,i}Z_{t-1,j}\}_{t\in\mathbb{Z}}\}, (i,j) \in [p] \times [pq]$. It follows that Assumptions C.1 and C.3 hold (with b_1 there equal to the constant C'). To establish Assumption C.2, note that for any $i \in [p], j \in [pq], k \in \mathbb{N}_0$ and $h \in \mathbb{N}$,

$$E\left[\left(\sum_{t=k+1}^{k+h} \varepsilon_{t,i} Z_{t-1,j}\right)^2\right] = \sum_{t=k+1}^{k+h} E\left[\varepsilon_{t,i}^2 Z_{t-1,j}^2\right]$$
(centered)
$$= \sum_{t=k+1}^{k+h} E\left[\varepsilon_{t,i}^2\right] E\left[Z_{t-1,j}^2\right]$$
(independence)
$$\ge \sum_{t=k+1}^{k+h} E\left[\varepsilon_{t,i}^2\right] E\left[\varepsilon_{t-1,j}^2\right]$$
(E $\left[Z_{t-1,j}^2\right] \ge E\left[\varepsilon_{t-1,j}^2\right]$)
$$\ge h\Lambda_{\min}\left(\Sigma_{\varepsilon}\right)^2$$
$$\ge a_2^2h,$$
(Assumption 1.3)

implying that Assumption C.2 is satisfied for the constant $\omega = a_2 \in (0, \infty)$. The derived constants depend on neither *i* nor *j*. Choosing the sequence γ_n in Assumption C.4 to be the non-random sequence γ_n in (2.6), we have both $\ln(1/\gamma_n) \leq \ln(pqn)$ and $\gamma_n \to 0$. The desired conclusion therefore follows from Lemma C.2.

D.3 Proof of Ideal Penalty Loadings Control (Lemma 3)

The proof of Lemma 3 essentially follows from the following result.

Lemma D.4. Let Assumptions 1.1, 1.2, 2.1 and 2.2 hold, suppose that $(\ln p)^{40/(3\alpha)} = o(n)$ for α provided by Assumption 1.2, let $v_{i,j}^0 := (\mathbb{E}[\varepsilon_{0,i}^2 Z_{-1,j}^2])^{1/2}$ for $i \in [p]$ and $j \in [pq]$, and define

$$\widehat{\Delta}^{0} := \max_{i \in [p], j \in [pq]} \left| (\widehat{v}^{0}_{i,j})^{2} - (v^{0}_{i,j})^{2} \right|, \tag{D.7}$$

where $\widehat{v}_{i,j}^0$ is given in (4.5). Then $\widehat{\Delta}^0 \to_{\mathrm{P}} 0$.

PROOF OF LEMMA D.4. We set up for an application of Lemma B.2. Observe first that

$$\begin{split} \widehat{\Delta}^{0} &= \max_{i \in [p], j \in [pq]} \left| \frac{1}{n} \sum_{k=1}^{l_{n}} \left(\sum_{t \in H_{n,k}} \varepsilon_{t,i} Z_{t-1,j} \right)^{2} - \mathbf{E} \left[\varepsilon_{0,i}^{2} \right] \mathbf{E} \left[Z_{0,j}^{2} \right] \right| \\ &= m_{n} \max_{i \in [p], j \in [pq]} \left| \frac{1}{l_{n}} \sum_{k=1}^{l_{n}} \left(\frac{1}{m_{n}^{2}} \left(\sum_{t \in H_{n,k}} \varepsilon_{t,i} Z_{t-1,j} \right)^{2} - \frac{1}{m_{n}} \mathbf{E} \left[\varepsilon_{0,i}^{2} \right] \mathbf{E} \left[Z_{0,j}^{2} \right] \right) \right| \\ &= \frac{m_{n}}{l_{n}} \max_{i \in [p], j \in [pq]} \left| \sum_{k=1}^{l_{n}} \left(m_{n}^{-2} (V_{k,i,j}^{(n)})^{2} - \mathbf{E} \left[m_{n}^{-2} (V_{0,i,j}^{(n)})^{2} \right] \right) \right|, \end{split}$$

where we have defined

$$V_{k,i,j}^{(n)} := \sum_{t \in H_{n,k}} \varepsilon_{t,i} Z_{t-1,j}, \quad i \in [p], \quad j \in [pq], \quad k \in [l_n].$$

The (skeleton) process $\{\mathbf{V}_{k}^{(n)}\}_{k\in\mathbb{Z}} = \{V_{k,i,j}^{(n)}: i\in[p], j\in[pq]\}_{k\in\mathbb{Z}}$ inherits the strict stationary from $\{\boldsymbol{\varepsilon}_{t}\boldsymbol{Z}_{t-1}^{\top}\}_{t\in\mathbb{Z}}$.

By a change of indices, we can write $V_{k,i,j}^{(n)}$ as

$$V_{k,i,j}^{(n)} = \sum_{t=(k-1)m_n}^{km_n - 1} \varepsilon_{t+1,i} Z_{t,j}.$$
 (D.8)

Iterating on (2.2), for $t \in \{(k-1)m_n, \ldots, km_n - 1\}$ we can express

$$Z_{t,j} = (\widetilde{\Theta}_0^{t-(k-1)m_n})_{j,\cdot} \mathbf{Z}_{(k-1)m_n} + \sum_{h=0}^{t-(k-1)m_n-1} (\widetilde{\Theta}_0^h)_{j,1:p} \boldsymbol{\varepsilon}_{t-h}.$$
 (D.9)

where in (D.9) we understand the sum $\sum_{h=0}^{-1} (\widetilde{\Theta}_0^h)_{j,1:p} \varepsilon_{t-h}$ as zero. Again iterating on (2.2), we see that

$$\boldsymbol{Z}_{(k-1)m_n} = \widetilde{\boldsymbol{\Theta}}_0^{m_n} \boldsymbol{Z}_{(k-2)m_n} + \sum_{h=0}^{m_n-1} (\widetilde{\boldsymbol{\Theta}}_0^h)_{\cdot,1:p} \boldsymbol{\varepsilon}_{(k-1)m_n-h}$$
$$= \sum_{\ell=0}^{\infty} \sum_{h=\ell m_n}^{(\ell+1)m_n-1} (\widetilde{\boldsymbol{\Theta}}_0^h)_{\cdot,1:p} \boldsymbol{\varepsilon}_{(k-1)m_n-h}.$$
(D.10)

Combining (D.8)–(D.10), we arrive at

$$V_{k,i,j}^{(n)} = \sum_{t=(k-1)m_n}^{km_n-1} \varepsilon_{t+1,i} \sum_{h=0}^{t-(k-1)m_n-1} (\widetilde{\Theta}_0^h)_{j,1:p} \varepsilon_{t-h} + \left(\sum_{t=(k-1)m_n}^{km_n-1} \varepsilon_{t+1,i} (\widetilde{\Theta}_0^{t-(k-1)m_n})_{j,\cdot}\right) \left(\sum_{\ell=0}^{\infty} \sum_{h=\ell m_n}^{(\ell+1)m_n-1} (\widetilde{\Theta}_0^h)_{\cdot,1:p} \varepsilon_{(k-1)m_n-h}\right).$$
(D.11)

The right-hand side of (D.11) defines an \mathbb{R} -valued mapping $G_{i,j}^{(n)}$, which (using strict stationarity) does not depend on k. Define the matrices

$$\boldsymbol{\varepsilon}_{H_{n,k}} := \begin{bmatrix} \boldsymbol{\varepsilon}_{(k-1)m_n+1} : \cdots : \boldsymbol{\varepsilon}_{km_n} \end{bmatrix}^\top, \quad k \in \mathbb{Z},$$
 (D.12)

and gather the mappings $G_{i,j}^{(n)}, i \in [p], j \in [pq]$, into an $\mathbb{R}^{p \times pq}$ -valued mapping $\mathbf{G}^{(n)}$. Then we have expressed $\mathbf{V}_k^{(n)}$ in the form

$$\mathbf{V}_k^{(n)} = \mathbf{G}^{(n)}\left(\ldots, oldsymbol{arepsilon}_{H_{n,k-1}}, oldsymbol{arepsilon}_{H_{n,k}}
ight)$$

which shows that $\{\mathbf{V}_k^{(n)}\}_{k\in\mathbb{Z}}$ is causal.

Next, for any $i \in [p], j \in [pq], r \in [1, \infty)$ and $k \in [l_n]$,

$$\begin{split} \left\| m_n^{-2} (V_{k,i,j}^{(n)})^2 \right\|_r &= m_n^{-2} \left\| \left(\sum_{t \in H_{n,k}} \varepsilon_{t,i} Z_{t-1,j} \right)^2 \right\|_r \\ &= m_n^{-2} \left\| \sum_{t=1}^{m_n} \varepsilon_{t,i} Z_{t-1,j} \right\|_{2r}^2 \\ &\leqslant \| \varepsilon_{0,i} Z_{-1,j} \|_{2r}^2 \qquad \text{(sub-additivity and stationarity)} \\ &\leqslant Cr^{4/\alpha} \qquad \text{(Lemma D.1, Part 3)} \end{split}$$

for some constant $C \in (0, \infty)$ depending only on a_1, α, b_1 and b_2 . It follows that

$$\max_{i \in [p], j \in [pq]} \left\| m_n^{-2} (V_{k,i,j}^{(n)})^2 \right\|_{\psi_{\alpha/4}} \leqslant C,$$

showing that the sub-Weibullness requirement of Lemma B.2 is satisfied with the constants (K,ξ) there equal to $(C,\alpha/4)$. Invoking Lemma B.2 for the vectorization of $\{m_n^{-2}V_{k,i,j}^{(n)}: i \in \mathbb{C}\}$

 $[p], j \in [pq]\}_{k \in \mathbb{Z}}$ with $r = 2, T = l_n, d_T = p^2 q$ and $m_T = 1$, we see that, as $n \to \infty$,

$$\begin{aligned} \widehat{\Delta}^{0} &= \frac{m_{n}}{l_{n}} \max_{i \in [p], j \in [pq]} \left| \sum_{k=1}^{l_{n}} \left(m_{n}^{-2} (V_{k,i,j}^{(n)})^{2} - \mathbf{E} \left[m_{n}^{-2} (V_{0,i,j}^{(n)})^{2} \right] \right) \right| \\ &\lesssim_{\mathbf{P}} \frac{m_{n}}{l_{n}} \left(l_{n} \left(p^{2}q \right)^{1/2} \Delta_{2,n} + \sqrt{l_{n} \ln \left(p^{2}q l_{n} \right)} + \left(\ln \left(p^{2}q l_{n} \right) \right)^{8/\alpha} \right) \\ &= m_{n} \left(p^{2}q \right)^{1/2} \Delta_{2,n} + \frac{m_{n}}{l_{n}} \sqrt{l_{n} \ln \left(p^{2}q l_{n} \right)} + \frac{m_{n}}{l_{n}} \left(\ln \left(p^{2}q l_{n} \right) \right)^{8/\alpha} \\ &=: \mathbf{I}_{n} + \mathbf{II}_{n} + \mathbf{III}_{n} \end{aligned}$$

with

$$\Delta_{2,n} := \max_{i \in [p], j \in [pq]} \Delta_2^{m_n^{-2}(V_{\cdot,i,j}^{(n)})^2} (2) \,.$$

We handle each of the right-hand side terms in turn, starting with II_n and III_n . Recalling that $m_n = n^{1/5}$, $l_n = n^{4/5}$, and $(\ln p)^{40/(3\alpha)} = o(n)$ (Assumption 4), we have that

$$II_n \leqslant Cn^{-3/5} [n^{4/5} (\ln p + \ln n)]^{1/2} = C \sqrt{\frac{\ln p + \ln n}{n^{2/5}}} = o(1),$$

for some constant $C \in (0, \infty)$, and

$$III_n \leqslant C' n^{-3/5} \left(\ln p + \ln n \right)^{8/\alpha} = C' \left(\frac{\left(\ln p + \ln n \right)^{40/(3\alpha)}}{n} \right)^{3/5} = o(1).$$

for some constant $C' \in (0, \infty)$. It remains to show that

$$m_n p \Delta_{2,n} = o(1)$$

Specifically, we will show that

$$\Delta_{2,n} \lesssim e^{-b_2 m_n},\tag{D.13}$$

such that

$$m_n p \Delta_{2,n} \lesssim e^{-b_2 n^{1/5} + \ln(p) + \frac{1}{5} \ln(n)} = o(1).$$

To do so, note that (D.8)-(D.10) imply that

$$V_{1,i,j}^{(n)} = \sum_{t=0}^{m_n-1} \varepsilon_{t+1,i} (\widetilde{\Theta}_0^{t+m_n})_{j,\cdot} \mathbf{Z}_{-m_n} + \sum_{t=0}^{m_n-1} \varepsilon_{t+1,i} \sum_{h=0}^{m_n-1} (\widetilde{\Theta}_0^{h+t})_{j,1:p} \mathbf{\varepsilon}_{-h} + \sum_{t=0}^{m_n-1} \varepsilon_{t+1,i} \sum_{h=0}^{t-1} (\widetilde{\Theta}_0^{h})_{j,1:p} \mathbf{\varepsilon}_{t-h},$$

such that, with $\{\boldsymbol{\varepsilon}_t^*\}_{t\in\mathbb{Z}}$ an independent copy of $\{\boldsymbol{\varepsilon}_t\}_{t\in\mathbb{Z}}$, and with

$$oldsymbol{Z}^*_{-m_n} := \sum_{h=0}^\infty (\widetilde{oldsymbol{\Theta}}^h_0)_{\cdot,1:p} oldsymbol{arepsilon}^*_{-m_n-h},$$

for any $r \in [1, \infty)$,

$$\begin{split} \Delta_{r}^{V_{\cdot,i,j}^{(n)}}(2) &= \left\| G_{i,j}^{(n)} \left(\dots, \varepsilon_{H_{n,-2}}, \varepsilon_{H_{n,-1}}, \varepsilon_{H_{n,0}}, \varepsilon_{H_{n,1}} \right) - G_{i,j}^{(n)} \left(\dots, \varepsilon_{H_{n,-2}}^{*}, \varepsilon_{H_{n,-1}}^{*}, \varepsilon_{H_{n,0}}, \varepsilon_{H_{n,1}} \right) \right\|_{r} \\ &= \left\| \sum_{t=0}^{m_{n}-1} \varepsilon_{t+1,i} (\widetilde{\Theta}_{0}^{t+m_{n}})_{j,\cdot} \left(Z_{-m_{n}} - Z_{-m_{n}}^{*} \right) \right\|_{r} \\ &= \left\| \sum_{t=0}^{m_{n}-1} \varepsilon_{t+1,i} (\widetilde{\Theta}_{0}^{t+m_{n}})_{j,\cdot} \sum_{h=0}^{\infty} (\widetilde{\Theta}_{0}^{h})_{\cdot,1:p} \left(\varepsilon_{-m_{n}-h} - \varepsilon_{-m_{n}-h}^{*} \right) \right\|_{r} \\ &= \left\| \sum_{t=0}^{m_{n}-1} \sum_{h=0}^{\infty} \varepsilon_{t+1,i} (\widetilde{\Theta}_{0}^{h+t+m_{n}})_{j,1:p} \left(\varepsilon_{-m_{n}-h} - \varepsilon_{-m_{n}-h}^{*} \right) \right\|_{r} \end{split}$$

Countable sub-additivity, followed by independence and strict stationarity yield

$$\Delta_{r}^{V_{\cdot,i,j}^{(n)}}(2) \leq \max_{i \in [p]} \|\varepsilon_{0,i}\|_{r} \sum_{t=0}^{m_{n}-1} \sum_{h=0}^{\infty} \left\| (\widetilde{\Theta}_{0}^{h+t+m_{n}})_{j,1:p} \left(\varepsilon_{-m_{n}-h} - \varepsilon_{-m_{n}-h}^{*} \right) \right\|_{r}.$$

Assumption 1.2 yields $\max_{i \in [p]} \|\varepsilon_{0,i}\|_r \leq r^{1/\alpha} \|\varepsilon_0\|_{\psi_{\alpha}} \leq a_1 r^{1/\alpha}$. Moreover, using sub-Weibullness, sub-additivity, stationarity and Assumptions 1.2 and 2.2, the right-hand side summands in the previous display can be bounded as follows

$$\begin{split} \left\| (\widetilde{\Theta}_{0}^{h+t+m_{n}})_{j,1:p} \left(\boldsymbol{\varepsilon}_{-m_{n}-h} - \boldsymbol{\varepsilon}_{-m_{n}-h}^{*} \right) \right\|_{r} &\leq r^{1/\alpha} \left\| (\widetilde{\Theta}_{0}^{h+t+m_{n}})_{j,1:p} \left(\boldsymbol{\varepsilon}_{-m_{n}-h} - \boldsymbol{\varepsilon}_{-m_{n}-h}^{*} \right) \right\|_{\psi_{\alpha}} \\ &\leq 2r^{1/\alpha} \left\| \boldsymbol{\varepsilon}_{0} \right\|_{\psi_{\alpha}} \left\| (\widetilde{\Theta}_{0}^{h+t+m_{n}})_{j,1:p} \right\|_{\ell_{2}} \\ &\leq 2a_{1}b_{1}r^{1/\alpha}\mathrm{e}^{-b_{2}(h+t+m_{n})}. \end{split}$$

The previous two displays therefore combine to produce

$$\begin{split} \Delta_r^{V_{r,i,j}^{(n)}}(2) &\leqslant a_1 r^{1/\alpha} \sum_{t=0}^{m_n - 1} \sum_{h=0}^{\infty} 2a_1 b_1 r^{1/\alpha} \mathrm{e}^{-b_2(h+t+m_n)} \\ &= 2a_1^2 b_1 r^{2/\alpha} \mathrm{e}^{-b_2 m_n} \sum_{t=0}^{m_n - 1} \mathrm{e}^{-b_2 t} \sum_{h=0}^{\infty} \mathrm{e}^{-b_2 h} \\ &= 2a_1^2 b_1 r^{2/\alpha} \mathrm{e}^{-b_2 m_n} \frac{1 - \mathrm{e}^{-b_2 m_n}}{1 - \mathrm{e}^{-b_2}} \cdot \frac{1}{1 - \mathrm{e}^{-b_2}} \\ &\leqslant \frac{2a_1^2 b_1}{(1 - \mathrm{e}^{-b_2})^2} r^{2/\alpha} \mathrm{e}^{-b_2 m_n}. \end{split}$$

Furthermore, Minkowski's inequality, stationarity, and Part 3 of Lemma D.1 yield

$$\|V_{0,i,j}^{(n)}\|_{2r} \leqslant m_n \max_{i \in [p], j \in [pq]} \|\varepsilon_{0,i} Z_{-1,j}\|_{2r} \leqslant \frac{2^{2/\alpha} a_1^2 b_1}{1 - e^{-b_2}} r^{2/\alpha} m_n.$$

Hence, for any $r \in [1, \infty)$ and some constant $C \in (0, \infty)$ depending only on a_1, α, b_1 and b_2 ,

$$\Delta_{r}^{m_{n}^{-2}(V_{.,i,j}^{(n)})^{2}}(2) = m_{n}^{-2} \Delta_{r}^{(V_{.,i,j}^{(n)})^{2}}(2)$$

$$\leq 2m_{n}^{-2} \|V_{0,i,j}^{(n)}\|_{2r} \Delta_{2r}^{V_{.,i,j}^{(n)}}(2)$$

$$\leq Cr^{4/\alpha} m_{n}^{-1} e^{-b_{2}m_{n}}.$$
(Lemma B.1)

As the latter upper bound does not depend on (i, j), setting r = 2, it follows that $\Delta_{2,n} \leq_{\mathbf{P}} m_n^{-1} e^{-b_2 m_n}$, which is more than enough for the desired $\Delta_{2,n} \leq_{\mathbf{P}} e^{-b_2 m_n}$.

PROOF OF LEMMA 3. Recall that $v_{i,j}^0 = (E[\varepsilon_{0,i}^2 Z_{-1,j}^2])^{1/2}$ for $i \in [p]$ and $j \in [pq]$. Using Assumption 1, we have

$$\min_{i \in [p], j \in [pq]} (v_{i,j}^0)^2 = \min_{i \in [p], j \in [pq]} \mathbf{E}[\varepsilon_{0,i}^2] \mathbf{E}[Z_{-1,j}^2] \geqslant a_2^2 > 0,$$

so the $\{(v_{i,j}^0)^2\}_{i\in[p],j\in[pq]}$ are bounded away from zero by the constant $a_2^2 =: C$. Part 3 of Lemma D.1 (with r = 2) shows that $\{(v_{i,j}^0)^2\}_{i\in[p],j\in[pq]}$ are bounded from above by a constant $C' \in (0,\infty)$ depending only on a_1, α, b_1 and b_2 . Recalling $\widehat{\Delta}^0$ in (D.7), it follows that

$$\min_{i \in [p], j \in [pq]} (\widehat{v}_{i,j}^0)^2 \ge \min_{i \in [p], j \in [pq]} (v_{i,j}^0)^2 - \widehat{\Delta}^0 \quad \text{and} \quad \max_{i \in [p], j \in [pq]} (\widehat{v}_{i,j}^0)^2 \le \max_{i \in [p], j \in [pq]} (v_{i,j}^0)^2 + \widehat{\Delta}^0.$$

Lemma D.4 shows that $\widehat{\Delta} \to_{\mathrm{P}} 0$. Hence, (4.9) is satisfied with, e.g., $\underline{v}^0 := \sqrt{C/2}$ and $\overline{v}^0 := \sqrt{2C'}$.

D.4 Proof of Restricted Eigenvalue Bound (Lemma 5)

PROOF OF LEMMA 4. We set up for an application of Lemma B.2. Assumptions 1.1 and 2.1 yield the VMA(∞) representation (D.6), so $\{\mathbf{Z}_t\}_{t\in\mathbb{Z}}$ is strictly stationary and causal. These properties are inherited by the process $\{\mathbf{Z}_t\mathbf{Z}_t^{\top}\}_{t\in\mathbb{Z}}$ with elements taking values in $\mathbb{R}^{pq\times pq}$. Part 1 of Lemma D.1 shows that, for α provided by Assumption 1.2,

$$\max_{i \in [pq]} \|Z_{0,i}\|_{\psi_{\alpha}} \leqslant \frac{a_1 b_1}{1 - e^{-b_2}} =: C_1.$$

Combined with the equivalent definitions of sub-Weibullness in Vladimirova et al. (2020, Theorem 1) and the bound in Kuchibhotla and Chakrabortty (2022, Proposition D.2), the previous display shows that for some constant $C_2 \in (0, \infty)$, depending only on a_1, α, b_1 and b_2 ,

$$\max_{(i,j)\in [pq]^2} \|Z_{0,i}Z_{0,j}\|_{\psi_{\alpha/2}} \leqslant C_2.$$

Part 2 of Lemma D.1 shows that

$$\max_{i \in [pq]} \|Z_{0,i}\|_4 \leqslant \frac{4^{1/\alpha} a_1 b_1}{1 - e^{-b_2}} =: C_3,$$

and Part 1 of Lemma D.3 shows that there is a constant $C_4 \in (0, \infty)$, depending only on a_1, α, b_1, b_2 and q, such that

$$\max_{i \in [pq]} \Delta_4^{Z_{i,i}}(h) \leqslant C_4 \mathrm{e}^{-b_2 h}, \quad h \in \mathbb{N}_0.$$

It follows from Lemma B.1 that

$$\max_{(i,j)\in[pq]^2}\Delta_2^{Z_{\cdot,i}Z_{\cdot,j}}(h) \leqslant 2C_3C_4\mathrm{e}^{-b_2h}, \quad h\in\mathbb{N}_0.$$

From the previous display and Remark B.1, we see that for any choice of $m_n \in [n]$,

$$\max_{(i,j)\in [pq]^2} \max_{u\in\{m_n+1,\dots,2m_n\}} \Delta_2^{Z_{\cdot,i}Z_{\cdot,j}} (u) \leq 2C_3 C_4 e^{-b_2 m_n}.$$

By hypothesis we have $(\ln p)^{1+4/\alpha} = o(n)$, which implies that $\ln p = o(n)$, such that we eventually have $\lceil \ln(pn)/b_2 \rceil \in [n]$. Applying Lemma B.2 for the vectorization of $\{Z_t Z_t^{\top}\}_{t \in \mathbb{Z}}$ with $\xi = \alpha/2$, $K = C_2$, r = 2, T = n, $d_T = p^2 q^2$ and $m_T = \lceil \ln(pn)/b_2 \rceil$, we see that, as

 $n \to \infty$,

$$\max_{(i,j)\in[pq]^2} \left| (\hat{\Sigma}_{\mathbf{Z}})_{i,j} - (\hat{\Sigma}_{\mathbf{Z}})_{i,j} \right| \\
= \frac{1}{n} \max_{(i,j)\in[pq]^2} \left| \sum_{t=1}^n \left(Z_{t,i} Z_{t,j} - \mathbb{E} \left[Z_{0,i} Z_{0,j} \right] \right) \right| \\
\lesssim_{\mathrm{P}} \frac{1}{n} \left(n [(pq)^2]^{1/2} \mathrm{e}^{-b_2 m_T} + m_T \sqrt{l_T \ln((pq)^2 n)} + m_T (\ln((pq)^2 n))^{4/\alpha} \right) \\
\lesssim_{\mathrm{P}} \frac{1}{n} + \sqrt{\frac{(\ln(pn))^2}{n}} + \frac{(\ln(pn))^{1+4/\alpha}}{n} \\
\lesssim \sqrt{\frac{(\ln(pn))^{1+4/\alpha}}{n}} = o(1). \qquad (\alpha \in (0, 2])$$

PROOF OF LEMMA 5. Let $T \subseteq [pq]$ and $\boldsymbol{\delta} \in \mathbb{R}^{pq}$ satisfy $|T| \in [s]$ and $||\boldsymbol{\delta}_{T^c}||_{\ell_1} \leq \hat{c}_0 \hat{\mu}_0 ||\boldsymbol{\delta}_T||_{\ell_1}$, where we define $\hat{c}_0 = (\hat{u}c + 1)/(\hat{\ell}c - 1)$ for some random sequences $\hat{\ell}$ and \hat{u} provided by asymptotic validity (4.7)–(4.8) of the penalty loadings $\{\hat{\boldsymbol{\Upsilon}}_i\}_{i=1}^p$. It follows that $\boldsymbol{\delta} \in \mathcal{R}_{\hat{c}_0\hat{\mu}_0,T}$ and, thus,

$$\|\boldsymbol{\delta}\|_{\ell_1} \leqslant (1+\widehat{c}_0\widehat{\mu}_0)\|\boldsymbol{\delta}_T\|_{\ell_1} \leqslant (1+\widehat{c}_0\widehat{\mu}_0)\sqrt{s}\|\boldsymbol{\delta}\|_{\ell_2}.$$

Abbreviating $\widehat{\Delta} := \max_{(i,j) \in [pq]^2} |(\widehat{\Sigma}_{\mathbf{Z}})_{i,j} - (\Sigma_{\mathbf{Z}})_{i,j}|$, we then have

$$\begin{split} \boldsymbol{\delta}^{\top} \boldsymbol{\Sigma}_{\boldsymbol{Z}} \boldsymbol{\delta} &- \boldsymbol{\delta}^{\top} \widehat{\boldsymbol{\Sigma}}_{\boldsymbol{Z}} \boldsymbol{\delta} = \boldsymbol{\delta}^{\top} (\boldsymbol{\Sigma}_{\boldsymbol{Z}} - \widehat{\boldsymbol{\Sigma}}_{\boldsymbol{Z}}) \boldsymbol{\delta} \leqslant |\boldsymbol{\delta}^{\top} (\widehat{\boldsymbol{\Sigma}}_{\boldsymbol{Z}} - \boldsymbol{\Sigma}_{\boldsymbol{Z}}) \boldsymbol{\delta}| \\ & \leqslant \left\| (\widehat{\boldsymbol{\Sigma}}_{\boldsymbol{Z}} - \boldsymbol{\Sigma}_{\boldsymbol{Z}}) \boldsymbol{\delta} \right\|_{\ell_{1}} \qquad (\text{H\"{o}lder}) \\ & \leqslant \widehat{\boldsymbol{\Delta}} \| \boldsymbol{\delta} \|_{\ell_{1}}^{2} \qquad (\text{H\"{o}lder}) \\ & \leqslant (1 + \widehat{c}_{0} \widehat{\mu}_{0})^{2} s \| \boldsymbol{\delta} \|_{\ell_{2}}^{2} \widehat{\boldsymbol{\Delta}}. \end{split}$$

For $\delta \neq 0$ the previous display rearranges to

$$\frac{\boldsymbol{\delta}^{\top} \widehat{\boldsymbol{\Sigma}}_{\boldsymbol{Z}} \boldsymbol{\delta}}{\|\boldsymbol{\delta}\|_{\ell_2}^2} \geq \frac{\boldsymbol{\delta}^{\top} \boldsymbol{\Sigma}_{\boldsymbol{Z}} \boldsymbol{\delta}}{\|\boldsymbol{\delta}\|_{\ell_2}^2} - (1 + \widehat{c}_0 \widehat{\mu}_0)^2 s \widehat{\Delta}.$$

It follows that

$$\widehat{\kappa}_{\widehat{c}_{0}\widehat{\mu}_{0}}^{2} \geq \min_{\substack{T \subseteq [pq]: \\ |T| \in [s]}} \inf_{\substack{\boldsymbol{\delta} \in \mathcal{R}_{\widehat{c}_{0}\widehat{\mu}_{0},T}: \\ \boldsymbol{\delta} \neq \boldsymbol{0}}} \frac{\boldsymbol{\delta}^{\top} \boldsymbol{\Sigma}_{\boldsymbol{Z}} \boldsymbol{\delta}}{\|\boldsymbol{\delta}\|_{\ell_{2}}^{2}} - (1 + \widehat{c}_{0}\widehat{\mu}_{0})^{2} s\widehat{\Delta} \geq \Lambda_{\min}(\boldsymbol{\Sigma}_{\boldsymbol{Z}}) - (1 + \widehat{c}_{0}\widehat{\mu}_{0})^{2} s\widehat{\Delta}.$$

By Assumption 3 it thus suffices to show that $(1 + \hat{c}_0 \hat{\mu}_0)^2 s \hat{\Delta} \to_P 0$. To this end, note that asymptotic validity of penalty loadings $\{\hat{\Upsilon}_i\}_{i=1}^p$ implies $\hat{\ell} \to_P 1$ and $\hat{u} \to_P u$ for some

constant $u \in [1, \infty)$. It follows that $\widehat{c}_0 = (\widehat{u}c + 1)/(\widehat{\ell}c - 1) \to_{\mathrm{P}} (uc + 1)/(c - 1) \in [1, \infty)$. Lemma 3 establishes $\widehat{\mu}_0 \lesssim_{\mathrm{P}} 1$, and therefore $\widehat{c}_0 \widehat{\mu}_0 \lesssim_{\mathrm{P}} 1$ as well. The growth condition $s^2(\ln(pn))^{1+4/\alpha} = o(n)$ implies $(\ln(pn))^{1+4/\alpha} = o(n)$ and so by Lemma 4

$$\widehat{\Delta} \lesssim_{\mathrm{P}} \sqrt{\frac{(\ln(pn))^{1+4/\alpha}}{n}}$$

which further implies $(1 + \hat{c}_0 \hat{\mu}_0)^2 s \hat{\Delta} \rightarrow_{\mathrm{P}} 0$.

D.5 Proof of Asymptotic Validity of Penalty Loadings from Algorithm 1 (Lemma 6)

Lemma D.5 (Convergence Rates for Lasso with Asymptotically Valid Loadings). Let Assumptions 1, 2, 3 and 4 hold, and let $\{\widehat{\Upsilon}_i\}_{i=1}^p$ be any collection of asymptotically valid penalty loadings. Then Lasso estimates $\widehat{\beta}_i := \widehat{\beta}_i(\lambda_n^*, \widehat{\Upsilon}_i)$ based on the penalty level λ_n^* in (2.7) and the penalty loadings $\{\widehat{\Upsilon}_i\}_{i=1}^p$ satisfy (2.11)–(2.13).

PROOF OF LEMMA D.5. By Lemma 2 $P(\lambda_n^*/n \ge c \max_{i \in [p]} || S_{n,i} ||_{\ell_{\infty}}) \to 1$, and by Lemma 3 there exist constants $\underline{v}^0, \overline{v}^0 \in (0, \infty)$ such that $P(\underline{v}^0 \le \widehat{v}_{\min}^0 \le \widehat{v}_{\max}^0 \le \overline{v}^0) \to 1$ as well as $P(\widehat{\mu}_0 \le \overline{v}^0/\underline{v}^0) \to 1$. Since the penalty loadings $\{\widehat{\Upsilon}_i\}_{i=1}^p$ are asymptotically valid, there are non-negative random variables $\widehat{\ell} := \widehat{\ell}_n$ and $\widehat{u} := \widehat{u}_n, n \in \mathbb{N}$, such that (4.7) with probability approaching one and (4.8). For any choice of such random variables, we have the restricted eigenvalue bound $P(\widehat{\kappa}_{\widehat{c}_0\widehat{\mu}_0}^2 \ge d/2) \to 1$ (Lemma 5), where $\widehat{c}_0 := (\widehat{u}c + 1)/(\widehat{\ell}c - 1)$, and $\widehat{\kappa}_{\widehat{c}_0\widehat{\mu}_0}^2$ is defined via (4.2) with $C = \widehat{c}_0\widehat{\mu}_0$. Thus, it follows from Lemma 1 that

$$\max_{i \in [p]} \|\widehat{\boldsymbol{\beta}}_{i} - \boldsymbol{\beta}_{0i}\|_{2,n} \lesssim_{\mathrm{P}} \frac{\lambda_{n}^{*}\sqrt{s}}{n},$$
$$\max_{i \in [p]} \|\widehat{\boldsymbol{\beta}}_{i} - \boldsymbol{\beta}_{0i}\|_{\ell_{1}} \lesssim_{\mathrm{P}} \frac{\lambda_{n}^{*}s}{n} \text{ and}$$
$$\max_{i \in [p]} \|\widehat{\boldsymbol{\beta}}_{i} - \boldsymbol{\beta}_{0i}\|_{\ell_{2}} \lesssim_{\mathrm{P}} \frac{\lambda_{n}^{*}\sqrt{s}}{n}.$$

Thus, (2.11)–(2.13) follow as $\Phi^{-1}(1-z) \leq \sqrt{2\ln(1/z)}, z \in (0,1)$, implies $\lambda_n^* \lesssim \sqrt{n\ln(pn)}$.

PROOF OF LEMMA 6. The proof is divided into three steps. Step 1 establishes asymptotic validity of the *initial* penalty loadings $\{\widehat{v}_{i,j}^{(0)}\}_{(i,j)\in[p]\times[pq]}$, while Step 2 proves asymptotic validity of the *once updated* penalty loadings $\{\widehat{v}_{i,j}^{(1)}\}_{(i,j)\in[p]\times[pq]}$. Step 3 observes that the argument in Step 2 can be iterated a finite number of times to arrive at the asymptotic validity of the *K* times updated loadings $\{\widehat{v}_{i,j}^{(K)}\}_{(i,j)\in[p]\times[pq]}$ for any fixed $K \in \mathbb{N}_0$.

Step 1: To establish asymptotic validity of the *initial* penalty loadings, let

$$v_{i,j}^{(0)} := (\mathbf{E}[(\widehat{v}_{i,j}^{(0)})^2])^{1/2} = (\mathbf{E}[Y_{0,i}^2 Z_{-1,j}^2])^{1/2}, \quad (i,j) \in [p] \times [pq],$$

and

$$v_{i,j}^0 := (\mathbf{E}[(\widehat{v}_{i,j}^0)^2])^{1/2} = (\mathbf{E}[\varepsilon_{0,i}^2 Z_{-1,j}^2])^{1/2}, \quad (i,j) \in [p] \times [pq].$$

Since $\boldsymbol{\varepsilon}_0$ is centered (Assumption 1) and independent of $\boldsymbol{Z}_{-1,j}$, for any $(i,j) \in [p] \times [pq]$

$$\mathbf{E}[Y_{0,i}^2 Z_{-1,j}^2] = \mathbf{E}[(\mathbf{Z}_{-1}^\top \boldsymbol{\beta}_{0i} + \varepsilon_{0,i})^2 Z_{-1,j}^2] = \mathbf{E}[(\mathbf{Z}_{-1}^\top \boldsymbol{\beta}_{0i})^2 Z_{-1,j}^2] + \mathbf{E}[\varepsilon_{0,i}^2 Z_{-1,j}^2]$$

such that

$$(v_{i,j}^{(0)})^2 = \mathbf{E}[(\mathbf{Z}_0^{\top} \boldsymbol{\beta}_{0i})^2 Z_{0,j}^2] + (v_{i,j}^0)^2.$$
(D.14)

Parts 2 and 4 of Lemma D.1 together with the Cauchy-Schwarz inequality show that, under Assumptions 1.1, 1.2, 2.1 and 2.2 there is a constant $C_1 \in (0, \infty)$, depending only on a_1, α, b_1 and b_2 , such that

$$\max_{i \in [p], j \in [pq]} \mathbf{E}[(\boldsymbol{Z}_0^{\top} \boldsymbol{\beta}_{0i})^2 Z_{0,j}^2] \leqslant C_1.$$

It follows from (D.14) that for any $(i, j) \in [p] \times [pq]$,

$$(v_{i,j}^0)^2 \leqslant (v_{i,j}^{(0)})^2 \leqslant C_1 + (v_{i,j}^0)^2.$$
 (D.15)

Lemma 3 shows that there is a constant $\underline{v}^0 \in (0, \infty)$, depending only on a_1, a_2, α, b_1 and b_2 , such that $P(\hat{v}^0_{\min} \ge \underline{v}^0) \to 1$. To simplify notation, without loss of generality, we henceforth argue as if $P(\hat{v}^0_{\min} \ge \underline{v}^0) = 1$ for all n. Lemma D.4 shows that $\widehat{\Delta}^0$ in (D.7) satisfies

$$\widehat{\Delta}^{0} = \max_{i \in [p], j \in [pq]} \left| (\widehat{v}_{i,j}^{0})^{2} - (v_{i,j}^{0})^{2} \right| \xrightarrow{\mathrm{P}} 0.$$
(D.16)

Suppose for now that also

$$\widehat{\Delta}^{(0)} := \max_{i \in [p], j \in [pq]} \left| (\widehat{\upsilon}_{i,j}^{(0)})^2 - (\upsilon_{i,j}^{(0)})^2 \right| \xrightarrow{\mathbf{P}} 0.$$
(D.17)

(We will establish (D.17) below.) We now construct non-negative random scalars $\hat{\ell}^{(0)} := \hat{\ell}_n^{(0)}$ and $\hat{u}^{(0)} := \hat{u}_n^{(0)}, n \in \mathbb{N}$, and a constant $u^{(0)} \in [1, \infty)$ such that (4.7) holds with probability approaching one and (4.8) holds. To construct $\hat{\ell}^{(0)}$, observe that for any $(i, j) \in [p] \times [pq]$, by the triangle inequality and (D.15),

$$(\widehat{v}_{i,j}^{(0)})^2 \ge (v_{i,j}^{(0)})^2 - \widehat{\Delta}^{(0)} \ge (v_{i,j}^0)^2 - \widehat{\Delta}^{(0)} \ge (\widehat{v}_{i,j}^0)^2 - (\widehat{\Delta}^0 + \widehat{\Delta}^{(0)}).$$

On the one hand, since $0 < \underline{v}^0 \leq \hat{v}_{\min}^0 \leq \hat{v}_{i,j}^0$, we can continue this string of inequalities to get

$$(\widehat{\upsilon}_{i,j}^{(0)})^2 \geqslant \left(1 - \frac{\widehat{\Delta}^0 + \widehat{\Delta}^{(0)}}{(\underline{\upsilon}^0)^2}\right) (\widehat{\upsilon}_{i,j}^0)^2.$$

On the other hand, we know that penalty loadings are non-negative. These observations suggest

$$\widehat{\ell}^{(0)} := \sqrt{\max\left\{0, 1 - \frac{\widehat{\Delta}^0 + \widehat{\Delta}^{(0)}}{(\underline{\nu}^0)^2}\right\}}$$
(D.18)

To construct $\hat{u}^{(0)}$, observe that for any $(i, j) \in [p] \times [pq]$, by the triangle inequality and the upper bound in (D.15),

$$(\widehat{v}_{i,j}^{(0)})^2 \leqslant (v_{i,j}^{(0)})^2 + \widehat{\Delta}^{(0)} \leqslant (v_{i,j}^0)^2 + C_1 + \widehat{\Delta}^{(0)} \leqslant (\widehat{v}_{i,j}^0)^2 + C_1 + \widehat{\Delta}^0 + \widehat{\Delta}^{(0)}.$$

Again, since $0 < \underline{v}^0 \leqslant \widehat{v}_{\min}^0 \leqslant \widehat{v}_{i,j}^0$, we can continue this string of inequalities to get

$$(\widehat{\upsilon}_{i,j}^{(0)})^2 \leqslant \left(1 + \frac{C_1 + \widehat{\Delta}^0 + \widehat{\Delta}^{(0)}}{(\underline{\upsilon}^0)^2}\right) (\widehat{\upsilon}_{i,j}^0)^2.$$

The previous displays suggests

$$\widehat{u}^{(0)} := \sqrt{1 + \frac{C_1 + \widehat{\Delta}^0 + \widehat{\Delta}^{(0)}}{(\underline{v}^0)^2}} \tag{D.19}$$

By construction of (D.18) and (D.19), from (D.16) and the working hypothesis (D.17), we see that (4.7) holds with probability approaching one. We also see that $\hat{\ell}^{(0)} \to_{\rm P} 1$ and $\hat{u}^{(0)} \to_{\rm P} u^{(0)}$ for the constant $u^{(0)} := (1 + C_1/(\underline{v}^0)^2)^{1/2} \in [1, \infty)$, which depends only on a_1, a_2, α, b_1 and b_2 , showing that also (4.8) holds. Hence, as long as we can establish (D.17), the initial penalty loadings $\{\hat{v}_{i,j}^{(0)}\}_{(i,j)\in[p]\times[pq]}$ are asymptotically valid.

To this end, we set up for an application of Lemma B.2. Part 1 of Lemma D.1 shows that

$$\max_{i \in [pq]} \|Z_{0,i}\|_{\psi_{\alpha}} \leqslant \frac{a_1 b_1}{1 - e^{-b_2}} =: C_2.$$
 (D.20)

Combined with the equivalent definitions of sub-Weibullness in Vladimirova et al. (2020, Theorem 1) and the bound in Kuchibhotla and Chakrabortty (2022, Proposition D.2), the previous display shows that for some constant $C_3 \in (0, \infty)$, depending only on a_1, α, b_1 and b_2

$$\max_{(i,j)\in[p]\times[pq]} \left\| Y_{0,i}^2 Z_{-1,j}^2 \right\|_{\psi_{\alpha/4}} \leqslant C_3.$$

Setting r = 4, Part 2 of Lemma D.1 shows that

$$\max_{i \in [pq]} \|Z_{0,i}\|_4 \leqslant \frac{4^{1/\alpha} a_1 b_1}{1 - e^{-b_2}} =: C_4,$$

and Part 1 of Lemma D.3 shows that for some constant $C_5 \in (0, \infty)$, depending only on a_1, α, b_1, b_2 and q,

$$\max_{i \in [pq]} \Delta_4^{Z_{\cdot,i}}(h) \leqslant C_5 \mathrm{e}^{-b_2 h}, \quad h \in \mathbb{N}_0.$$

Repeated use of Lemma B.1 and the previous two displays imply that for some constant $C_6 \in (0, \infty)$, depending only on a_1, α, b_1, b_2 and q,

$$\max_{(i,j)\in[p]\times[pq]} \Delta_1^{Y^2_{\cdot,i}Z^2_{\cdot-1,j}}(h) \leqslant C_6 e^{-b_2 h}, \quad h \in \mathbb{N}_0.$$

By hypothesis we have $(\ln p)^{40/(3\alpha)} = o(n)$, which using $\alpha \in (0, 2]$ implies that $\ln p = o(n)$, such that eventually $\lceil 2\ln(pn)/b_2 \rceil \in [n]$. Applying Lemma B.2 for the vectorization of the strictly stationary and causal matrix process $\{[Y_{t,i}^2 Z_{t-1,j}^2]_{(i,j)\in[p]\times[pq]}\}_{t\in\mathbb{Z}}$ with $r = 1, \xi = \alpha/4, K = C_3, T = n, d_T = p^2q$ and $m_T = \lceil 2\ln(pn)/b_2 \rceil$, we see that, as $n \to \infty$,

$$\begin{split} \widehat{\Delta}^{(0)} &= \max_{(i,j) \in [p] \times [pq]} \left| (\widehat{v}_{i,j}^{(0)})^2 - (v_{i,j}^{(0)})^2 \right| \\ &= \frac{1}{n} \max_{(i,j) \in [p] \times [pq]} \left| \sum_{t=1}^n \left(Y_{t,i}^2 Z_{t-1,j}^2 - \operatorname{E}[Y_{0,i}^2 Z_{-1,j}^2] \right) \right| \\ &\lesssim_{\operatorname{P}} \frac{1}{n} \left(n p^2 q e^{-b_2 m_T} + m_T \sqrt{l_T \ln(p^2 q n)} + m_T (\ln(p^2 q n))^{8/\alpha} \right) \\ &\lesssim_{\operatorname{P}} \frac{1}{n} + \sqrt{\frac{(\ln(pn))^2}{n}} + \frac{(\ln(pn))^{1+8/\alpha}}{n} \\ &\lesssim_{\operatorname{P}} \sqrt{\frac{(\ln(pn))^{1+8/\alpha}}{n}} \\ &\lesssim_{\operatorname{P}} \sqrt{\frac{(\ln(pn))^{40/(3\alpha)}}{n}} = o(1), \end{split}$$
(D.21)

where we again use $\alpha \in (0, 2]$.

Step 2: If K = 0, then there is nothing left to show. We therefore presume that the number of updates $K \in \mathbb{N}$, and consider the once updated penalty loadings $\{\widehat{v}_{i,j}^{(1)}\}_{(i,j)\in[p]\times[pq]}$ from (2.9). As observed in Step 1, Lemma 3 shows that there is a constant $\underline{v}^0 \in (0, \infty)$, depending only on a_1, a_2, α, b_1 and b_2 , such that $P(\widehat{v}_{\min}^0 \ge \underline{v}^0) \to 1$. As in Step 1, we (without loss of generality) continue to argue as if $P(\widehat{v}_{\min}^0 \ge \underline{v}^0) = 1$ for all n. Now, the proof of Lemma

3 shows (D.16). Suppose for now that also

$$\widehat{\Delta}^{(1)} := \max_{(i,j)\in[p]\times[pq]} \left| (\widehat{v}_{i,j}^{(1)})^2 - (v_{i,j}^0)^2 \right| \xrightarrow{\mathrm{P}} 0.$$
(D.22)

(We will establish (D.22) below.) We now construct random scalars $\hat{\ell}^{(1)} := \hat{\ell}_n^{(1)}$ and $\hat{u}^{(1)} := \hat{u}_n^{(1)}, n \in \mathbb{N}$, such that (4.7) holds with probability approaching one and (4.8) holds with $\hat{u}^{(1)} \to_{\mathrm{P}} 1$. To construct $\hat{\ell}^{(1)}$, fix $(i, j) \in [p] \times [pq]$. On the one hand, by the triangle inequality and $0 < \underline{v}^0 \leq \hat{v}_{\min}^0 \leq \hat{v}_{i,j}^0$, we get

$$(\widehat{v}_{i,j}^{(1)})^2 \ge (v_{i,j}^0)^2 - \widehat{\Delta}^{(1)} \ge (\widehat{v}_{i,j}^0)^2 - (\widehat{\Delta}^0 + \widehat{\Delta}^{(1)}) \ge \left(1 - \frac{\Delta^0 + \widehat{\Delta}^{(1)}}{(\underline{v}^0)^2}\right) (\widehat{v}_{i,j}^0)^2.$$

On the other hand, penalty loadings are non-negative. These observations suggest

$$\widehat{\ell}^{(1)} := \sqrt{\max\left\{0, 1 - \frac{\widehat{\Delta}^0 + \widehat{\Delta}^{(1)}}{(\underline{\nu}^0)^2}\right\}},\tag{D.23}$$

and similar arguments lead us to

$$\widehat{u}^{(1)} := \sqrt{1 + \frac{\widehat{\Delta}^0 + \widehat{\Delta}^{(1)}}{(\underline{v}^0)^2}}.$$
(D.24)

By construction of (D.23) and (D.24), from (D.16) and the working hypothesis (D.22), we see that (4.7) holds with probability approaching one. Furthermore, $\hat{\ell}^{(1)} \rightarrow_{\rm P} 1$ and $\hat{u}^{(1)} \rightarrow_{\rm P} 1$, showing that also (4.8) holds, now with u = 1. Hence, it remains to establish (D.22).

To this end, define the (infeasible) penalty loadings

$$\widetilde{v}_{i,j}^0 := \left(\frac{1}{n} \sum_{t=1}^n \varepsilon_{t,i}^2 Z_{t-1,j}^2\right)^{1/2}, \quad (i,j) \in [p] \times [pq],$$

which correspond to the ideal penalty loadings without any blocking. To show $\widehat{\Delta}^{(1)} \to_{\mathrm{P}} 0$, by the triangle inequality, it suffices that

$$\widetilde{\Delta}^{0} := \max_{(i,j)\in[p]\times[pq]} \left| (\widetilde{v}_{i,j}^{0})^{2} - (v_{i,j}^{0})^{2} \right| \xrightarrow{\mathcal{P}} 0 \quad \text{and} \quad \widetilde{\Delta}^{(1)} := \max_{(i,j)\in[p]\times[pq]} \left| (\widehat{v}_{i,j}^{(1)})^{2} - (\widetilde{v}_{i,j}^{0})^{2} \right| \xrightarrow{\mathcal{P}} 0.$$

To show that $\widetilde{\Delta}^0 \to_{\mathcal{P}} 0$, note that

$$\widetilde{\Delta}^0 = \frac{1}{n} \max_{(i,j) \in [p] \times [pq]} \left| \sum_{t=1}^n \left(\varepsilon_{t,i}^2 Z_{t-1,j}^2 - \mathbf{E}[\varepsilon_{0,i}^2 Z_{-1,j}^2] \right) \right|.$$

By Assumption 1.2 one has $\max_{i \in [p]} \|\varepsilon_{0,i}\|_{\psi_{\alpha}} \leq a_1$. In addition, $\Delta_4^{\varepsilon_{\cdot,i}}(h) = 0$ for $h \in \mathbb{N}$ while $\Delta_4^{\varepsilon_{\cdot,i}}(0)$ is bounded. Thus, $\{\varepsilon_{t,i}\}_{t \in \mathbb{Z}}$ is a geometric moment contraction and replacing $Y_{t,i}$ with $\varepsilon_{t,i}$ in the arguments starting at (D.20) and leading to (D.21) yields $\widetilde{\Delta}^0 \to_{\mathbb{P}} 0$.

To show that $\widetilde{\Delta}^{(1)} \to_{\mathrm{P}} 0$, recall that $\widehat{\varepsilon}_{t,i}^{(0)} = Y_{t,i} - \mathbf{Z}_{t-1}^{\top} \widehat{\boldsymbol{\beta}}_{i}^{(0)} = \varepsilon_{t,i} - \mathbf{Z}_{t-1}^{\top} \widehat{\boldsymbol{\delta}}_{i}^{(0)}$, with $\widehat{\boldsymbol{\delta}}_{i}^{(0)} := \widehat{\boldsymbol{\beta}}_{i}^{(0)} - \boldsymbol{\beta}_{0i}$ being the estimation error arising from the initial penalty loadings. Hence,

$$\begin{split} \widetilde{\Delta}^{(1)} &= \max_{(i,j)\in[p]\times[pq]} \left| \frac{1}{n} \sum_{t=1}^{n} \left((\widehat{\varepsilon}_{ti}^{(0)})^2 - \varepsilon_{ti}^2 \right) Z_{t-1,j}^2 \right| \\ &= \max_{(i,j)\in[p]\times[pq]} \left| \frac{1}{n} \sum_{t=1}^{n} \left[(Z_{t-1}^\top \widehat{\delta}_i^{(0)})^2 - 2\varepsilon_{t,i} Z_{t-1}^\top \widehat{\delta}_i^{(0)} \right] Z_{t-1,j}^2 \right| \\ &\leqslant 2 \max_{(i,j)\in[p]\times[pq]} \left| \frac{1}{n} \sum_{t=1}^{n} \varepsilon_{t,i} Z_{t-1,j}^2 Z_{t-1}^\top \widehat{\delta}_i^{(0)} \right| + \max_{(i,j)\in[p]\times[pq]} \left| \frac{1}{n} \sum_{t=1}^{n} |Z_{t-1}' \widehat{\delta}_i^{(0)}|^2 Z_{t-1,j}^2 \right| \\ &= :\mathrm{In} \end{split}$$

To show $\widetilde{\Delta}^{(1)} \to_{\mathcal{P}} 0$, we argue $\mathcal{I}_n \to_{\mathcal{P}} 0$ and $\mathcal{II}_n \to_{\mathcal{P}} 0$, in turn. First,

$$I_{n} \leqslant \max_{(i,j)\in[p]\times[pq]} \sqrt{\frac{1}{n} \sum_{t=1}^{n} \varepsilon_{t,i}^{2} Z_{t-1,j}^{4}} \|\widehat{\boldsymbol{\delta}}_{i}^{(0)}\|_{2,n}}$$
(Cauchy-Schwarz)
$$\leqslant \max_{\substack{t\in[n] \ j\in[pq] \\ =:I_{n}^{(a)}}} \max_{(i,j)\in[p]\times[pq] \atop =:I_{n}^{(b)}} \widetilde{\upsilon}_{i,j}^{0} \cdot \max_{\substack{i\in[p] \\ =:I_{n}^{(c)}}} \|\widehat{\boldsymbol{\delta}}_{i}^{(0)}\|_{2,n}}.$$
(D.25)

We next handle each of the terms in (D.25), starting with $I_n^{(c)}$. From Lemma D.5 and the asymptotic validity of the initial loadings (Step 1), we know that

$$\mathbf{I}_n^{(c)} = \max_{i \in [p]} \|\widehat{\boldsymbol{\delta}}_i^{(0)}\|_{2,n} \lesssim_{\mathbf{P}} \sqrt{s \ln(pn)/n}.$$

Turning to $I_n^{(b)}$, as

$$\widetilde{\Delta}^0 = \max_{(i,j) \in [p] \times [pq]} \left| (\widetilde{v}^0_{i,j})^2 - (v^0_{i,j})^2 \right| \xrightarrow{\mathbf{P}} 0$$

and

$$\max_{(i,j)\in[p]\times[pq]} v_{i,j}^0 = \max_{(i,j)\in[p]\times[pq]} (\mathbf{E}[\varepsilon_{0,i}^2 Z_{-1,j}^2])^{1/2} \leqslant \frac{2^{1/\alpha} a_1^2 b_1}{1-\mathrm{e}^{-b_2}} < \infty,$$

we must have

$$\mathbf{I}_{n}^{(b)} = \max_{(i,j)\in[p]\times[pq]}\widetilde{\upsilon}_{i,j}^{0} \lesssim_{\mathbf{P}} 1.$$

Concerning $I_n^{(a)}$, Part 1 of Lemma D.1 yields

$$\max_{j \in [pq]} \|Z_{0,j}\|_{\psi_{\alpha}} \leqslant \frac{a_1 b_1}{1 - e^{-b_2}}$$

Thus, by Vladimirova et al. (2020, Theorem 1), there exists a constant $C_7 \in (0, \infty)$, depending only on a_1, α, b_1 and b_2 , such that

$$\mathcal{P}(|Z_{0,j}| \ge x) \le 2\mathbf{e}^{-(x/C_7)^{\alpha}}$$

holds for all $x \in (0, \infty)$ and all $j \in [pq]$. Hence, by a union bound

$$\max_{t \in [n]} \max_{j \in [pq]} |Z_{t-1,j}| \lesssim_{\mathbf{P}} (\ln(pn))^{1/\alpha},$$

so gathering our findings in (D.25), we arrive at

$$I_n \lesssim_P (\ln(pn))^{1/\alpha} \cdot \sqrt{\frac{s \ln(pn)}{n}} = \sqrt{\frac{s(\ln(pn))^{1+2/\alpha}}{n}} = o(1).$$

Our previous calculations also show that

$$\begin{aligned} \mathrm{II}_{n} &= \max_{(i,j)\in[p]\times[pq]} \left| \frac{1}{n} \sum_{t=1}^{n} |\mathbf{Z}_{t-1}^{\top} \widehat{\boldsymbol{\delta}}_{i}^{(0)}|^{2} Z_{t-1,j}^{2} \right| \leq \max_{t\in[n]} \max_{j\in[pq]} Z_{t-1,j}^{2} \cdot \max_{i\in[p]} \|\widehat{\boldsymbol{\delta}}_{i}^{(0)}\|_{2,n}^{2} \\ &\lesssim_{\mathrm{P}} (\ln(pn))^{2/\alpha} \cdot \frac{s\ln(pn)}{n} = \frac{s(\ln(pn))^{1+2/\alpha}}{n} = o\left(1\right). \end{aligned}$$

It follows that the once updated penalty loadings $\{\widehat{v}_{i,j}^{(1)}\}_{(i,j)\in[p]\times[pq]}$ are asymptotically valid.

Step 3: To finish the proof, observe that asymptotic validity of the once updated penalty loadings (Step 2) and Lemma D.5 imply

$$\max_{i \in [p]} \|\widehat{\boldsymbol{\delta}}_i^{(1)}\|_{2,n} \lesssim_{\mathbf{P}} \sqrt{s \ln(pn)/n},$$

with $\widehat{\delta}_{i}^{(1)} := \widehat{\beta}_{i}^{(1)} - \beta_{0i}, i \in [p]$, being the estimation error arising from the once updated penalty loadings. Since the rate of convergence of the resulting $\|\widehat{\delta}_{i}^{(k)}\|_{2,n}$ remains $\sqrt{s \ln(pn)/n}$ for any finite k, we can iterate on the argument given in Step 2 to obtain asymptotic validity of the penalty loadings constructed from $K \in \mathbb{N}$ updates.

D.6 Proof of Convergence Rates for Lasso with Data-Driven Penalization (Theorem 1)

PROOF OF THEOREM 1. The assumptions of the theorem suffice for asymptotic validity of the penalty loadings $\{\widehat{v}_{i,j}^{(K)}\}_{(i,j)\in[p]\times[pq]}$ generated by Algorithm 1 with any fixed number $K \in \mathbb{N}_0$ of updates, cf. Lemma 6. The rates (2.11)–(2.13) thus follow from Lemma D.5. \Box