ANICK RESOLUTION FOR THE FREE UNITARY QUANTUM GROUP

ALEXANDER MANG

ABSTRACT. A resolution P of the co-unit of the Hopf *-algebra $\mathcal{O}(U_n^+)$ of representative functions on van Daele and Wang's free unitary quantum group U_n^+ in terms of free $\mathcal{O}(U_n^+)$ -modules is computed for arbitrary n.

A different such resolution was recently found by Baraquin, Franz, Gerhold, Kula and Tobolski. While theirs has desirable properties which P lacks, P is still good enough to compute the (previously known) quantum group cohomology and comes instead with an important advantage: P can be arrived at without the clever combination of certain results potentially very particular to U_n^+ that enabled the aforementioned authors to find their resolution. Especially, P relies neither on the resolution for O_n^+ obtained by Collins, Härtel and Thom nor the one for $SL_2(q)$ found by Hadfield and Krähmer.

Rather, as shown in the present article, the recursion defining the Anick resolution of the co-unit of $\mathcal{O}(U_n^+)$ can be solved in closed form. That suggests a potential strategy for determining the cohomologies of arbitrary easy quantum groups.

This is a case study in computing the quantum group cohomology of the discrete duals of so-called easy compact quantum groups.

More precisely, the Anick resolution of the trivial right module of the algebra of representative functions on the free unitary quantum group U_n^+ is computed for arbitrary n and used to deduce the (previously known) cohomology of the dual of U_n^+ .

 \Box Which findings did the case study produce? – Section 1

 \Box What are "easy" quantum groups and U_n^+ ? – Section 6.1

 \Box What are Anick resolutions? – Section 2

- \Box Why do the results of the study hold? Sections 3–5
- \Box Why do the case study in the first place? Section 6.2
- \Box Which conclusions can be drawn from the results? Section 6.3

Date: March 12, 2024.

²⁰²⁰ Mathematics Subject Classification. 20G42.

Key words and phrases. Anick resolution, Gröbner basis, free unitary quantum group, easy quantum group, compact quantum group, quantum group cohomology.

1. Main result

The main statement of the article is greatly shortened by the use of the following abbreviations.

Notation. In the context of any set E with a specified involution *, for any $n \in \mathbb{Z}_{>0}$ and any $n \times n$ -matrix $v = (v_{j,i})_{(j,i)\in[n]^{\times 2}}$ of elements of E consider besides v also the three matrices v^{t} , v^{\star} , v^{\ddagger} defined by for any $(j,i) \in [n]^{\times 2}$,

$$v_{j,i}^{t} := v_{i,j}, \qquad v_{j,i}^{\star} := v_{r(j),r(i)}^{\star}, \qquad v_{j,i}^{\dagger} := v_{r(j),r(i)}^{\star},$$

where $r: \mathbb{Z} \to \mathbb{Z}, i \mapsto n - i + 1$ is the reflection at n.

Main result. For any field \mathbb{K} , any $n \in \mathbb{Z}_{>0}$ with $2 \leq n$ and any 2nelemental set $E = \{u_{j,i}^{\circ}, u_{j,i}^{\bullet}\}_{i,j=1}^{n}$, if $u_{j,i}^{\circ} := u_{j,i}^{\bullet}$ and $u_{j,i}^{\bullet} := u_{j,i}^{\circ}$ for any $(j,i) \in [n]^{\times 2}$, if $u := (u_{j,i}^{\circ})_{(j,i) \in [n]^{\times 2}}$, if $V := \{u, u^{t}, u^{\star}, u^{\ddagger}\}$ (which means |V| = 4), if A is the quotient of the free unital \mathbb{K} -algebra $\mathbb{K}\langle E \rangle$ generated by E by the two-sided ideal J generated by

$$R := \{ \sum_{s=1}^{n} v_{j,\mathbf{r}(s)} v_{s,\mathbf{r}(i)}^{\ddagger} - \delta_{j,i} \, 1 | \, v \in V \land (j,i) \in [n]^{\times 2} \}$$

and if $\varepsilon \colon A \to \mathbb{K}$ is the unique morphism of complex algebras with for any $v \in V$ and $(j, i) \in [n]^{\times 2}$,

$$\varepsilon(v_{j,i}+J) := \delta_{j,i}$$

which turns \mathbb{K} into a right A-module \mathbb{K}_{ε} , then the following are true: If for any $v \in V$, any $(j, i) \in [n]^{\times 2}$ and any $\ell \in \mathbb{Z}_{>0}$,

$$x_{j,i}^{v} \equiv x_{j,i}^{v,\ell} \coloneqq \begin{cases} v_{j,\mathbf{r}(i)} & |\ell = 1\\ v_{j,n}(v_{1,n}^{\dagger}v_{1,n})^{\frac{\ell-2}{2}}v_{1,\mathbf{r}(i)}^{\dagger} & |\ell \ even\\ v_{j,n}(v_{1,n}^{\dagger}v_{1,n})^{\frac{\ell-3}{2}}v_{1,n}^{\dagger}v_{1,\mathbf{r}(i)} & |\ell \neq 1 \land \ell \ odd \end{cases}$$

if for any $\ell \in \mathbb{Z}_{>0}$,

$$C_{\ell} := \{ x_{j,i}^{v,\ell} | v \in V \land (j,i) \in [n]^{\times 2} \} \subseteq \mathbb{K} \langle E \rangle$$

(which means in particular $|C_1| = 2n^2$ and $|C_2| = 4n^2 - 2$ and $|C_\ell| = 4n^2$ for any $\ell \in \mathbb{Z}_{>0}$ with $3 \leq \ell$), if for any $\ell \in \mathbb{Z}$ with $-1 \leq \ell$,

$$P_{\ell} := \begin{cases} \mathbb{K}_{\varepsilon} & |\ell = -1\\ A & |\ell = 0\\ \mathbb{K}C_{\ell} \otimes_{\mathbb{K}} A & |1 \leq \ell \end{cases}$$

(i.e., if P_{ℓ} is the free right A-module over C_{ℓ} for each $\ell \in \mathbb{Z}_{>0}$), and if $(d_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ is such that for each $\ell \in \mathbb{Z}_{\geq 0}$ the mapping d_{ℓ} is a morphism $P_{\ell} \to P_{\ell-1}$ of right A-modules and such that for any $v \in V$ and any $(j,i) \in [n]^{\times 2}$,

$$d_0(1+J) \coloneqq 1$$

and

$$d_1(x_{j,i}^v \otimes (1+J)) \coloneqq v_{j,\mathbf{r}(i)} - \delta_{j,\mathbf{r}(i)} 1 + J$$

and

$$\begin{aligned} d_2(x_{j,i}^v \otimes (1+J)) \\ &\coloneqq \sum_{s=1}^n x_{j,s}^v \otimes (v_{s,r(i)}^{\ddagger} + J) + x_{r(j),i}^{v^{\ddagger}} \otimes (1+J) \\ &\quad -\delta_{j,n}\delta_{i,n} \sum_{s=2}^n (\sum_{t=1}^n x_{t,s}^v \otimes (v_{s,r(t)}^{\ddagger} + J) + x_{s,r(s)}^{v^{\ddagger}} \otimes (1+J)) \end{aligned}$$

and for any $\ell \in \mathbb{Z}_{\geq 0}$ with $3 \leq \ell$, if for any $w \in V$ the symbol ℓw is short for w if ℓ is odd and for w^{\ddagger} if ℓ is even, then

$$\begin{split} d_{\ell}(x_{j,i}^{v}\otimes(1+J)) & \coloneqq \sum_{s=1}^{n} x_{j,s}^{v}\otimes(^{\ell}v_{s,\mathrm{r}(i)}+J) + (-1)^{\ell} x_{\mathrm{r}(j),i}^{v^{\dagger}}\otimes(1+J) \\ & + \delta_{j,n}((x_{1,1}^{v^{\dagger}}+\delta_{\ell,3}\sum_{s=2}^{n-1} x_{s,s}^{v^{\dagger}})\otimes(^{\ell}v_{\mathrm{r}(i),n}^{t}+J) \\ & - \delta_{i,n}(\sum_{s=1}^{n} x_{1,s}^{v^{\dagger}}\otimes(^{\ell}v_{s,n}^{t}+J) + (-1)^{\ell} x_{n,1}^{v^{\star}}\otimes(1+J))) \\ & + \delta_{j,1}\delta_{i,n}(-1)^{\ell}(x_{1,1}^{v^{\star}}+\delta_{\ell,3}\sum_{s=2}^{n-1} x_{s,s}^{v^{\star}})\otimes(1+J), \end{split}$$

then the complex

$$\cdots \xrightarrow{d_3} \mathbb{K}C_2 \otimes A \xrightarrow{d_2} \mathbb{K}C_1 \otimes A \xrightarrow{d_1} A \xrightarrow{d_0} \mathbb{K}_{\varepsilon}$$

of right A-modules is exact. (In other words, $(P_{\ell}, d_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ is a free resolution of \mathbb{K}_{ε} in terms of right A-modules.) In fact, it is precisely the Anick resolution of \mathbb{K}_{ε} with respect to the degreewise lexicographic extension of the total order on E defined by, for any $\{\mathbf{c}, \mathbf{d}\} \subseteq \{\circ, \bullet\}$ and any $\{(j, i), (t, s)\} \subseteq [n]^{\times 2}$ with respect to the lexicographic order on $[n]^{\times 2}$,

$$\begin{split} u^{\mathfrak{d}}_{t,s} &\leq u^{\mathfrak{c}}_{j,i} \Leftrightarrow \quad (\mathfrak{d},\mathfrak{c}) = \bullet \circ \\ & \vee (\mathfrak{d},\mathfrak{c}) = \bullet \bullet \wedge (j,i) < (t,s) \\ & \vee (\mathfrak{d},\mathfrak{c}) = \circ \circ \wedge (t,s) < (j,i). \end{split}$$

Finally, if $\mathbb{K} = \mathbb{C}$, then A is the underlying algebra and ε the co-unit of the CQG algebra $\mathcal{O}(U_n^+)$ of representative functions on the free unitary quantum group U_n^+ .

This is an inferior resolution compared to the one previously obtained by Baraquin, Franz, Gerhold, Kula and Tobolski in [BFGKT23] (see Appendix A for a quasi-isomorphism). However, it can still be used to compute the (previously known) quantum group cohomology

$$H^{\cdot}(\hat{U}_{n}^{+}) \cong \operatorname{Ext}_{A}^{\cdot}(\mathbb{C}_{\varepsilon}, \mathbb{C}_{\varepsilon}),$$

as demonstrated in Section 5.

The significance of the Main result is that it is evidence that perhaps Anick resolutions can be used to compute homological invariants of easy quantum groups. For a discussion in that respect see Section 6.

2. Reminder on Anick resolutions

Anick presented the resolutions named after him in [Ani86]. They are commonly explained within the framework of non-commutative Gröbner bases developed by Mora in [Mor85]. Bergman's diamond lemma from [Ber78] also fits in this framework. All three are recalled below.

2.1. **Tips.** For any vector space F over any field \mathbb{K} , any \mathbb{K} -basis B, any vector $v \in F$ and any basis element $b \in B$ let $b \lrcorner v$ be the coefficient of v with respect to b. Given any total order \leq on B and any $v \in F$ with $v \neq 0$, there exists a unique element of B, called the *tip* of v with respect to B and \leq and denoted by the (non-standard) symbol $\diamond v$, which is \leq -maximal among the elements $b \in B$ with the property that $b \lrcorner v \neq 0$. Moreover, for any subset J of F let $\diamond J := \{\diamond v | v \in J \land v \neq 0\}$.

A well-order is any total order with respect to which any non-empty subset has a minimal element. By a transfinite generalization of Gaussian elimination, if \leq is a well-order on B, then any K-vector subspace J induces a decomposition of F into a direct sum $F = J \oplus \text{span}(B \setminus J)$ of K-vector spaces, where $B \setminus J$ is the set complement of $\diamond J$ in B, sometimes referred to as the non-tips of J with respect to B and \leq . For any $v \in F$ the unique $b \in \text{span}(B \setminus J)$ such that $v - b \in J$ is called the normal form of v with respect to J, B and \leq .

In particular, for any $\{b, b'\} \subseteq B \setminus J$ with $b \neq b'$ the classes b + J and b' + J are then distinct. And, the set $\{b + J | b \in B \setminus J\}$ is a K-basis of the quotient K-vector space F/J of F by J. The mapping which for any $v \in F$ sends the class v + J to the normal form of v is a well-defined K-linear splitting $F/J \to F$ of the projection map $F \to F/J$.

2.2. Monoidal ideals. In the following, a monoid B is any set equipped with an associative binary operation with a neutral element. Any $b \in B$ is said to divide any $b' \in B$ if there exist $\{b_1, b_2\} \subseteq B$ with $b_1bb_2 = b'$. Equivalently, b' is called a multiple of b.

A monoidal ideal of B is any subset of B which also contains all multiples of its elements. Given any monoidal ideal I of B, there may or may not exist a subset M which generates I as a monoidal ideal and which is minimal with that property with respect to the subset inclusion partial order \subseteq . If such a set M exists, it is unique and will, for brevity, be referred to as the *core* of I, which is non-standard terminology.

2.3. Monomial orders. A well-order on a monoid B is called *admissible* if for any $\{b, b', b_1, b_2\} \subseteq B$, whenever b < b', then also $b_1b < b_1b'$ and $bb_2 < b'b_2$. If there exists at least one admissible order on B, then either B is a trivial monoid, consisting only of a single element, or B has no left or right absorbing elements. Moreover, it can be seen that then any monoidal ideal of B has a core.

Let E be any set. If $B = \langle E \rangle$ is the free monoid over E, then any admissible order on $\langle E \rangle$ is called a *monomial order* for E. One way of

obtaining monomial orders for E is to extend a given well-order on E degreewise lexicographically: That means that for any $\{m, m'\} \subseteq \mathbb{Z}_{\geq 0}$ and any $e \in E^{\times m}$ and $e' \in E^{\times m'}$ it is defined that $e_1 e_2 \dots e_m < e'_1 e'_2 \dots e'_{m'}$ if and only if either m < m' or simultaneously m = m' and there exists $j \in [m]$ such that $e_i = e'_i$ for any $i \in [j-1]$ and $e_j < e'_j$.

2.4. Gröbner bases. Let E be any set, \mathbb{K} any field, $F = \mathbb{K}\langle E \rangle$ the free \mathbb{K} -algebra over E, moreover 1 its unit, \leq any monomial order for E and J any (two-sided) ideal of $\mathbb{K}\langle E \rangle$. Then, $B = \langle E \rangle$ is a basis of $\mathbb{K}\langle E \rangle$ and, with respect to that basis and \leq , the tips $I = \diamond J$ of J form a monoidal ideal in $\langle E \rangle$. The elements of $\langle E \rangle$ are usually called monomials in E and the non-tips $\langle E \rangle \langle \diamond J$ reduced monomials. Because \leq is a well-order the monoidal ideal $\diamond J$ admits a core M, sometimes called the obstructions of J with respect to \leq .

A Gröbner basis of J with respect to \leq is any subset G of J such that $\diamond G$ generates $\diamond J$ as a monoidal ideal of $\langle E \rangle$ or, equivalently, such that $M \subseteq \diamond G$. Any Gröbner basis G is said to be minimal if $0 \notin G$ and if there are no $\{g, g'\} \subseteq G$ with $g \neq g'$ such that $\diamond g$ divides $\diamond g'$. That is the case if and only if $\diamond G = M$. Even further, any minimal Gröbner basis is called *reduced* if $\diamond g \lrcorner g = 1$ for any $g \in G$ and if there exist no $\{g, g'\} \subseteq G$ such that $g \neq g'$ and $b \lrcorner g' \neq 0$ and such that $\diamond g$ divides b. In fact, with respect to \leq there is exactly one reduced Gröbner basis of J. It is in particular a \mathbb{K} -basis of J.

2.5. **Diamond lemma.** For any $v \in \mathbb{K}\langle E \rangle$ with $v \neq 0$ a one-step reduction by v with respect to \leq is any \mathbb{K} -vector space endomorphism of $\mathbb{K}\langle E \rangle$ for which there exist $\{b_1, b_2\} \subseteq \langle E \rangle$ such that $b_1(\diamond v)b_2$ is mapped to $b_1(\diamond v - \frac{1}{\diamond v \lrcorner v}v)b_2$ and any other element of $\langle E \rangle$ to itself.

Let G be any subset of $\mathbb{K}\langle E \rangle$. A reduction by G with respect to \leq is any element of a certain submonoid of the monoid formed by the \mathbb{K} -vector space endomorphisms of $\mathbb{K}\langle E \rangle$ under composition, namely the one generated by the one-step reductions by arbitrary non-zero elements of G.

An inclusion ambiguity of G with respect to \leq is any quadruple (g, g', b_1, b_2) such that $\{g, g'\} \subseteq G$ and $\{b_1, b_2\} \subseteq \langle E \rangle$ and $g \neq 0$ and $g' \neq 0$ and $g \neq g'$ and $b_1(\diamond g)b_2 = \diamond g'$. And it is then said to resolve if there exist reductions φ_1 and φ_2 by G with respect to \leq such that $\varphi_1(\diamond g' - \frac{1}{\diamond g' \cdot g'}g') = \varphi_2(b_1(\diamond g - \frac{1}{\diamond g \cdot g}g)b_2)$. Similarly, an overlap ambiguity of G with respect to \leq is any tuple

Similarly, an overlap ambiguity of G with respect to \leq is any tuple (g_1, g_2, b_1, b_2) with $\{g_1, g_2\} \subseteq G$ and $\{b_1, b_2\} \subseteq \langle E \rangle$ and $g_1 \neq 0$ and $g_2 \neq 0$ such that $(b_1, b_2) \neq (1, 1)$, such that $(\diamond g_1)b_2 = b_1(\diamond g_2)$ and such that neither is b_2 divided by $\diamond g_2$ nor b_1 by $\diamond g_1$. It is said to resolve if there exist reductions φ_1 and φ_2 by G such that $(\varphi_1 - \frac{1}{\diamond g_1 \cup g_1}g_1)b_2) = \varphi_2(b_1(\diamond g_2 - \frac{1}{\diamond g_2 \cup g_2}g_2)).$

Proposition 2.1 (Bergman's diamond lemma). With respect to \leq , the set G is a Gröbner basis of J if and only if G generates J as an ideal and all inclusion and overlap ambiguities of G resolve.

2.6. Chains. Suppose $E \cap \diamond J = \emptyset$ and let $pC_0 := C_0 := \{1\}$ be the set containing the unit of $\mathbb{K}\langle E \rangle$ and let $pC_1 := C_1 := E$.

Suppressing the dependency on J and \leq in the notation, for any $\ell \in \mathbb{Z}_{\geq 0}$ with $2 \leq \ell$, any $m \in \mathbb{Z}_{\geq 0}$ and any $e \in E^{\times m}$ the monomial $c = e_1 e_2 \dots e_m$ is called an ℓ -pre-chain, in symbols: $c \in pC_\ell$, if and only if there exist $\alpha \in [m]^{\times (\ell-1)}$ and $\beta \in [m]^{\times (\ell-1)}$ such that either $\ell = 2$ and $1 = \alpha_1 \leq \beta_1 = m$ or $3 \leq \ell$ and $1 = \alpha_1 < \alpha_2 \leq \beta_1$ and $\alpha_{\ell-1} \leq \beta_{\ell-2} < \beta_{\ell-1} = m$ and $\alpha_{s+1} \leq \beta_s < \alpha_{s+2}$ for any $s \in [\ell-3]$ and such that $e_{\alpha_s} e_{\alpha_{s+1}} \dots e_{\beta_s} \in M$ for any $s \in [\ell-1]$.

Furthermore, c is called an ℓ -chain, an element of C_{ℓ} , if and only if there exist α and β as above such that, additionally, for any $\ell' \in [\ell]$ with $2 \leq \ell'$ and any $s \in [\beta_{\ell'-1} - 1]$ the monomial $e_1e_2 \dots e_s$ is not an element of $pC_{\ell'}$. If so, then (α, β) is unique with those extended properties and, in the present article, will be referred to as the ℓ -chain indices of c. Beware that compared to [Ani86] the numbering of $(C_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ has been shifted by 1. An ℓ -chain here would be an $(\ell - 1)$ -chain there.

2.7. Combinatorial differentials and splittings. For any $\ell \in \mathbb{Z}_{>0}$, any $m \in \mathbb{Z}_{\geq 0}$, any $e \in E^{\times m}$ and any ℓ -chain $c = e_1 e_2 \dots e_m \in C_\ell$ let $\partial_\ell(c)$ be defined as (1, c) if $\ell = 1$, as $(e_1, e_2 e_3 \dots e_m)$ if $\ell = 2$, and otherwise, if (α, β) are the chain indices of c, as $(e_1 e_2 \dots e_{\beta_{\ell-2}}, e_{\beta_{\ell-2}+1} e_{\beta_{\ell-2}+2} \dots e_m)$. It can be proved that ∂_ℓ is a mapping $C_\ell \to C_{\ell-1} \times (\langle E \rangle \backslash \diamond J)$ if $2 \leq \ell$.

Moreover, let $Z_0 := (C_0 \times (\langle E \rangle \backslash \diamond J)) \setminus \{(1,1)\}$ and for any $\ell \in \mathbb{Z}_{>0}$ define Z_ℓ to be the set of all $(c,d) \in C_\ell \times (\langle E \rangle \backslash \diamond J)$ such that, if $(a,b) = \partial_\ell(c)$, then $bd \in \diamond J$.

For any $\ell \in \mathbb{Z}_{\geq 0}$ and any $(c,d) \in Z_{\ell}$, if $\{m,t\} \subseteq \mathbb{Z}_{\geq 0}$ and $e \in E^{\times (m+t)}$ are such that $c = e_1 e_2 \dots e_m$ and $d = e_{m+1} e_{m+2} \dots e_{m+t}$, then let $\int_{\ell} (c,d) := (e_1 e_2 \dots e_{\beta_{\ell}}, e_{\beta_{\ell}+1} e_{\beta_{\ell}+2} \dots e_{m+t})$, where $\beta_{\ell} \in [m+t]$ is such that $m < \beta_{\ell}$, such that, if $\ell = 0$, then $\beta_{\ell} = 1$, such that, if $1 \leq \ell$, then $e_{\alpha_{\ell}} e_{\alpha_{\ell}+1} \dots e_{\beta_{\ell}} \in M$, where $\alpha_{\ell} \in [m]$ is given by 1 if $\ell = 1$, by min $\{i \in \mathbb{Z} \land \exists j \in \mathbb{Z} : 1 < i \leq m < j < m + t \land e_i e_{i+1} \dots e_j \in M\}$ if $\ell = 2$, and by min $\{i \in \mathbb{Z} \land \exists j \in \mathbb{Z} : \beta_{\ell-2} < i \leq m < j < m + t \land e_i e_{i+1} \dots e_j \in M\}$ if $3 \leq \ell$, where $((\alpha_1, \alpha_2, \dots, \alpha_{\ell-1}), (\beta_1, \beta_2, \dots, \beta_{\ell-1}))$ are the ℓ -chain indices of c. One proves that \int_{ℓ} is a mapping $Z_{\ell} \to C_{\ell+1} \times \langle E \rangle$.

Note that ∂_{ℓ} , Z_{ℓ} and \int_{ℓ} are left implicit in [Ani86]. They can be interpreted as the combinatorial base of, respectively, the differentials, cycles and splittings of the resolution given below.

2.8. Chain modules. Let A denote the quotient K-algebra $\mathbb{K}\langle E \rangle/J$ of $\mathbb{K}\langle E \rangle$ by J and let ε be any morphism $A \to \mathbb{K}$ of K-algebras. Then ε induces a right A-module structure \mathbb{K}_{ε} on K. Let $P_{-1} := \mathbb{K}_{\varepsilon}$ and for any $\ell \in \mathbb{Z}_{\geq 0}$ let $P_{\ell} := \mathbb{K}C_{\ell} \otimes_{\mathbb{K}} A$, where $\mathbb{K}C_{\ell}$ is the free \mathbb{K} -vector space over C_{ℓ} .

There then exists a unique total order \leq_{ℓ} on the K-basis $B_{\ell} := \{c \otimes (b+J) | (c,b) \in C_{\ell} \times (\langle E \rangle \backslash \diamond J) \}$ of P_{ℓ} such that for any $\{c,c'\} \subseteq C_{\ell}$ and any $\{b,b'\} \subseteq \langle E \rangle \backslash \diamond J$ the inequality $c \otimes b \leq_{\ell} c' \otimes b'$ holds if and only if $cb \leq c'b'$. The tips of non-zero vectors in P_{ℓ} with respect to \leq_{ℓ} will be indicated by the symbol \diamond_{ℓ} .

2.9. Anick resolution. There exist unique families $(d_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ and $(\iota_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ of mappings such that the following conditions are met. For each $\ell \in \mathbb{Z}_{\geq 0}$ the mapping d_{ℓ} is a morphism $P_{\ell} \to P_{\ell-1}$ of right A-modules and ι_{ℓ} is a K-linear map $\ker(d_{\ell}) \to P_{\ell+1}$, where $\ker(d_{\ell}) \subseteq P_{\ell}$ is the kernel of d_{ℓ} as a K-linear map. The map d_0 satisfies $d_0(1 \otimes (1+J)) = 1$ and for any $e \in E = C_1$ it holds $d_1(e \otimes (1+J)) = 1 \otimes (e - \varepsilon(e) + J)$. For each $\ell \in \mathbb{Z}_{\geq 0}$ with $2 \leq \ell$ and any $c \in C_{\ell}$, if $\partial_{\ell}(c) = (a, b)$, then $d_{\ell}(c \otimes (1+J)) = a \otimes (b+J) - \iota_{\ell-2}(d_{\ell-1}(a \otimes (b+J)))$. For any $\ell \in \mathbb{Z}_{\geq 0}$ and any $v \in \ker(d_{\ell})$ with $v \neq 0$, if $\lambda = \diamond_{\ell} v \lrcorner v$, if $c \in C_{\ell}$ and $d \in \langle E \rangle \backslash \diamond J$ are such that $\diamond_{\ell} v = c \otimes (d+J)$, in which case it can be proved that $(c, d) \in Z_{\ell}$, and if $(g, h) = \int_{\ell} (c, d)$, then $\iota_{\ell}(v) = \lambda g \otimes (h+J) + \iota_{\ell}(v - \lambda d_{\ell+1}(g \otimes (h+J)))$.

Proposition 2.2 (Anick's resolution). $(P_{\ell}, d_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ is a resolution of \mathbb{K}_{ε} in terms of free right A-modules.

3. GRÖBNER BASIS FOR U_n^+

From now on let \mathbb{K} , n, E, V, R, J, A and \leq be as in the Main result. In this section the reduced Gröbner basis of J with respect to \leq is computed.

Definition 3.1. For any $v \in V$ and any $(j, i) \in [n]^{\times 2}$ let

$$e_{j,i}^{v} := v_{j,n} v_{1,\mathbf{r}(i)}^{\ddagger} - \left(-\sum_{s=2}^{n} v_{j,\mathbf{r}(s)} v_{s,\mathbf{r}(i)}^{\ddagger} + \delta_{j,i} 1 \right) \in \mathbb{K} \langle E \rangle$$

$$f^{v} := v_{n,n} v_{1,1}^{\ddagger} - \left(\sum_{s=2}^{n} \sum_{t=1}^{n-1} v_{t,\mathbf{r}(s)} v_{s,\mathbf{r}(t)}^{\ddagger} - (n-2) 1 \right) \in \mathbb{K} \langle E \rangle.$$

and then let

$$G := \{ f^v, e^v_{j,i} | v \in V \land (j,i) \in [n]^{\times 2} \setminus \{(n,n)\} \}.$$

With these abbreviations, $R = \{e_{j,i}^v | v \in V \land (j,i) \in [n]^{\times 2}\}$. Eventually, G will be shown to be the reduced Gröbner basis.

3.1. Auxiliary results. During the computation of the Gröbner basis the following fundamental consequences of the definition of the monomial order \leq will be used frequently.

Lemma 3.2. $w \in \{v, v^{\dagger}, v^{\star}, v^{\ddagger}\}$ for any $\{v, w\} \subseteq V$.

Proof. The below tables gives the argument for each of the 4×4 cases.

v	u	u	u	u	u^{t}	u^{t}	u^{t}	u^{t}	u^{\star}	u^{\star}	u^{\star}	u^{\star}	$ u^{\ddagger} $	u^{\ddagger}	u^{\ddagger}	u^{\ddagger}	
w	u	u^{t}	u^{\star}	u^{\ddagger}	u	u^{t}	u^{\star}	u^{\ddagger}	u	u^{t}	u^{\star}	u^{\ddagger}	u	u^{t}	u^{\star}	u^{\ddagger} u^{\ddagger}	
w	v	v^{t}	v^{\star}	v^{\ddagger}	v^{t}	v	v^{\ddagger}	v^{\star}	v^{\star}	v^{\ddagger}	v	v^{t}	v^{\ddagger}	v^{\star}	v^{t}	v	
That proves the claim.																	

That proves the claim.

Insofar as case distinctions are necessary, Lemma 3.2 provides the four pairwise exclusive possibilities which need to be considered.

Lemma 3.3. For any $\{v, w\} \subseteq V$ and any $\{(j, i), (t, s)\} \subseteq [n]^{\times 2}$,

 $w_{t,s} = v_{j,i}$

if and only if one of the following exclusive conditions is met: $\square w = v \text{ and } (t, s) = (j, i).$

 $\square w = v^{t} and (t, s) = (i, j).$

In particular, $w_{t,s} \neq v_{j,i}$ whenever $w \in \{v^{\star}, v^{\ddagger}\}$.

Proof. Follows from the assumption that $E = \{u_{\ell,k}^{\circ}, u_{\ell,k}^{\bullet} | (\ell, k) \in [n]^{\times 2} \}$ has $2n^2$ elements and Lemma 3.2.

Lemma 3.4. For any $\{v, w\} \subseteq V$ and $\{(j, i), (t, s)\} \subseteq [n]^{\times 2}$,

$$w_{t,s} < v_{j,i}$$

if and only if one of the following mutually exclusive conditions is met: $\square w \in \{v^{\star}, v^{\ddagger}\} and v \in \{u, u^{t}\}.$ $\Box w = v and$

$$v \in \{u, u^{\star}\} \land (t, s) < (j, i)$$
$$v \in \{u^{t}, u^{\ddagger}\} \land (s, t) < (i, j).$$

 $\Box w = v^{t}$ and

V

$$\begin{aligned} v \in \{u, u^{\star}\} & \land \ (s, t) < (j, i) \\ \forall & v \in \{u^{\mathrm{t}}, u^{\ddagger}\} \ \land \ (t, s) < (i, j). \end{aligned}$$

In particular, $v_{j,s} < v_{j,i}$ if and only if s < i and, likewise, $v_{t,i} < v_{j,i}$ if and only if t < j.

Proof. The case $w \in \{v^*, v^{\ddagger}\}$ is clear, as then $(w, v) \in \{u^*, u^{\ddagger}\} \times \{u, u^{\ddagger}\}$. Moreover, the case $w = v^{t}$ follows from the case w = v by exchanging the roles $t \leftrightarrow s$. But also in order to prove the assertion in the remaining case w = v it can be assumed that $v \in \{u, u^*\}$. That is because, if $v \in \{u^{t}, u^{\dagger}\}$, it is possible to apply the case $v \in \{u, u^{\star}\}$ after exchanging both $t \leftrightarrow s$ and $j \leftrightarrow i$.

If v = u, then $v_{t,s} < v_{j,i}$ if and only if $u_{t,s}^{\circ} < u_{j,i}^{\circ}$, which by definition

holds precisely if and only if (t, s) < (j, i). If $v = u^*$, then $v_{t,s} < v_{j,i}$ if and only if $u^{\bullet}_{\mathbf{r}(t),\mathbf{r}(s)} < u^{\bullet}_{\mathbf{r}(j),\mathbf{r}(i)}$, which by definition is equivalent to $(\mathbf{r}(j), \mathbf{r}(i)) < (\mathbf{r}(t), \mathbf{r}(s))$. The simultaneous application of r in both components is an anti-automorphism of the lexicographic order of $[n]^{\times 2}$. Indeed, $(\mathbf{r}(j), \mathbf{r}(i)) < (\mathbf{r}(t), \mathbf{r}(s))$ means that

either r(j) < r(t) or both r(j) = r(t) and r(i) < r(s). That is true if and only if either t < j or both t = j and s < i, i.e., if and only if (t, s) < (j, i). Hence, the assertion is true.

One particular consequence of Lemma 3.4 deserves emphasis.

Lemma 3.5. For any $v \in V$, any $\{(t,s), (j,i)\} \subseteq [n]^{\times 2}$ and any $w \in \{v, v^t\},\$

$$w_{s,t}^{\ddagger} < v_{i,j}^{\ddagger} \quad \Leftrightarrow \quad w_{t,s} < v_{j,i}.$$

Proof. The claim follows by comparing the conclusion of Lemma 3.4 to the conclusion it yields when applied $(w^{\ddagger}, v^{\ddagger})$ in the role of (w, v). Because $w \in \{v, v^{\ddagger}\}$, and thus $w^{\ddagger} \in \{v^{\star}, v^{\ddagger}\} = \{v^{\ddagger}, v^{\ddagger, t}\}$ only the last two cases of Lemma 3.4 need to be considered. Obviously, w = v is equivalent to $w^{\ddagger} = v^{\ddagger}$ and, likewise, $w = v^{t}$ to $w^{\ddagger} = v^{\ddagger, t} = v^{\star}$. That is why the second case for (w, v) must be compared to the second case for $(w^{\ddagger}, v^{\ddagger})$ and, the third case for (w, v) to the third case for $(w^{\ddagger}, v^{\ddagger})$. Within each of these two cases, however, the exchange $v \leftrightarrow v^{\ddagger}$ has swapped the condition $v \in \{u, u^{t}\}$ for $v \in \{u^{\star}, u^{\ddagger}\}$ and vice versa. But that change has the same effect as exchanging the roles $t \leftrightarrow s$ and $j \leftrightarrow i$ would. Thus, the comparison between the two conclusions of Lemma 3.4 proves the claim.

3.2. Relations as non-minimal Gröbner basis. It is actually more convenient to first prove that R itself happens to be a (non-minimal) Gröbner basis. Recall that \diamond indicates tips with respect to \leq .

Lemma 3.6. For any
$$v \in V$$
 and any $(j,i) \in [n]^{\times 2}$,
 $\diamond e_{j,i}^v = v_{j,n} v_{1,r(i)}^{\ddagger}, \qquad \diamond f^v = v_{n,n} v_{1,1}^{\ddagger}.$

Proof. The claim about $e_{j,i}^v$ is equivalent to $v_{j,r(s)}v_{s,r(i)}^{\ddagger} < v_{j,n}v_{1,r(i)}^{\ddagger}$ holding for any $s \in [n]$ with 1 < s. This in turn is certainly true if already $v_{j,r(s)} < v_{j,n}$ since \leq was obtained by (degreewise) lexicographic extension. By the addendum to Lemma 3.4 the latter holds, irrespective of the value of v, since 1 < s requires r(s) < n. That proves the assertion about $e_{j,i}^v$.

What is claimed about f^v is that $v_{t,r(s)}v_{s,r(t)}^{\ddagger} < v_{n,n}v_{1,1}^{\ddagger}$ for any $\{s,t\} \subseteq [n]$ with 1 < s and t < n. As before this will follow already from $v_{t,r(s)} < v_{n,n}$. A second application of Lemma 3.4 shows that this last condition is met if and only if either $v \in \{u, u^*\}$ and (t, r(s)) < (n, n) or $v \in \{u^t, u^{\ddagger}\}$ and (r(s), t) < (n, n). Again, no matter what v is, the respective inequality is true since t < n and since r(s) < n by 1 < s. That concludes the proof.

Lemma 3.7. Any inclusion ambiguity of R is of the form

$$(e_{n,n}^{v^{t}}, e_{n,n}^{v}, 1, 1)$$

for some $v \in V$.

Proof. Let (g, g', b_1, b_2) be any inclusion ambiguity of R. There exist $\{v, w\} \subseteq V$ and $\{(j, i), (\ell, k)\} \subseteq [n]^{\times 2}$ with $g = e_{\ell,k}^w$ and $g' = e_{j,i}^v$. Because $\diamond g = \diamond e_{\ell,k}^w$ and $\diamond g' = \diamond e_{j,i}^v$ have the same degree by Lemma 3.6 the assumption $b_1(\diamond g)b_2 = \diamond g'$ requires $b_1 = b_2 = 1$ and $\diamond g = \diamond g'$. By Lemma 3.6 the latter means

$$w_{\ell,n} w_{1,\mathbf{r}(k)}^{\ddagger} = v_{j,n} v_{1,\mathbf{r}(i)}^{\ddagger}.$$

Since that requires in particular $w_{\ell,n} = v_{j,n}$ Lemma 3.3 excludes $w \in \{v^*, v^{\ddagger}\}$, which by Lemma 3.2 then requires $w \in \{v, v^{\ddagger}\}$.

Moreover, $w \neq v$. Indeed, otherwise $v_{\ell,n}v_{1,\mathbf{r}(k)}^{\ddagger} = v_{j,n}v_{1,\mathbf{r}(i)}^{\ddagger}$ would follow, which by Lemma 3.3 would demand $(\ell, n) = (j, n)$ and $(1, \mathbf{r}(k)) = (1, \mathbf{r}(i))$. That would imply $(\ell, k) = (j, i)$ and thus ultimately $e_{\ell,k}^w = e_{j,i}^v$, in contradiction to the assumption $g \neq g'$.

in contradiction to the assumption $g \neq g'$. Hence, instead, $w = v^{t}$ and thus $v_{n,\ell}v_{r(k),1}^{\dagger} = v_{j,n}v_{1,r(i)}^{\dagger}$. By Lemma 3.3 it follows that $(n,\ell) = (j,n)$ and (r(k),1) = (1,r(i)), which is to say $i = j = k = \ell = n$. Hence, $e_{\ell,k}^{w} = e_{n,n}^{v^{t}}$ and $e_{j,i}^{v} = e_{n,n}^{v}$. And this is the only inclusion ambiguity, as claimed.

Lemma 3.8. All inclusion ambiguities of R resolve.

Proof. By Lemma 3.7 it suffices to show that for any $v \in V$ and $w := v^{t}$ the terms $\diamond e_{n,n}^{v} - e_{n,n}^{v}$ and $\diamond e_{n,n}^{w} - e_{n,n}^{w}$ reduce to the same expression. Indeed, by definition,

$$e_{n,n}^v - e_{n,n}^v = -\sum_{s=2}^n v_{n,r(s)} v_{s,1}^{\ddagger} + 1 = -\sum_{s=2}^n w_{r(s),n} w_{1,s}^{\ddagger} + 1 = -\sum_{s=1}^{n-1} w_{s,n} w_{1,r(s)}^{\ddagger} + 1 = -\sum_{s=1}^{n-1} \diamond e_{s,s}^w + 1$$

where the third identity is obtained by reindexing with r. Once reduced by $e_{s,s}^w$ for each $s \in [n]$ with s < n the above takes on the form

$$\diamond e_{n,n}^v - e_{n,n}^v \rightsquigarrow -\sum_{s=1}^{n-1} (-\sum_{t=2}^n w_{s,r(t)} w_{t,r(s)}^{\dagger} + 1) + 1$$

= $\sum_{s=1}^{n-1} \sum_{t=2}^n v_{r(t),s} v_{r(s),t}^{\dagger} - (n-2)1.$ (1)

Exchanging the roles $v \leftrightarrow w$ in (1) yields

$$\begin{aligned} \diamond e_{n,n}^{w} - e_{n,n}^{w} &\rightsquigarrow \sum_{s=1}^{n-1} \sum_{t=2}^{n} w_{\mathbf{r}(t),s} w_{\mathbf{r}(s),t}^{\ddagger} - (n-2) \mathbf{1} \\ &= \sum_{s=1}^{n-1} \sum_{t=2}^{n} v_{s,\mathbf{r}(t)} v_{t,\mathbf{r}(s)}^{\ddagger} - (n-2) \mathbf{1}. \end{aligned}$$

Switching the names of the summation indices $s \leftrightarrow t$ here gives

$$\diamond e_{n,n}^w - e_{n,n}^w \rightsquigarrow \sum_{s=2}^n \sum_{t=1}^{n-1} v_{t,\mathbf{r}(s)} v_{s,\mathbf{r}(t)}^{\ddagger} - (n-2)1,$$

which after transforming the domains of the summations with **r** each amounts to

$$\diamond e_{n,n}^w - e_{n,n}^w \rightsquigarrow \sum_{s=1}^{n-1} \sum_{t=2}^n v_{\mathbf{r}(t),s} v_{\mathbf{r}(s),t}^{\ddagger} - (n-2)1.$$
 (2)

Since the right-hand sides of (1) and (2) agree, all inclusion ambiguities resolve.

Lemma 3.9. Any overlap ambiguity of R is of the form $(e_{\ell,1}^{v^{\ddagger}}, e_{1,i}^{v}, v_{\ell,n}^{\ddagger}, v_{1,r(i)}^{\ddagger})$

for some $v \in V$ and $(\ell, i) \in [n]^{\times 2}$.

Proof. Let (g_1, g_2, b_1, b_2) be any overlap ambiguity of R. Then there exist $\{v, w\} \subseteq V$ and $\{(j, i), (\ell, k)\} \subseteq [n]^{\times 2}$ with $g_1 = e^w_{\ell, k}$ and $g_2 = e^v_{j, i}$.

Moreover, necessarily, $\{b_1, b_2\} \subseteq E$ for the following reasons. If, say, b_2 had degree 2 or greater, then because $\diamond g_2 = \diamond e_{j,i}^v$ has degree 2 by Lemma 3.6 the identity $(\diamond g_1)b_2 = b_1(\diamond g_2)$ would imply that $\diamond g_2$ divides b_2 , in contradiction to the assumption. Likewise, b_1 has degree less than 2 because also $\diamond g_1 = \diamond e_{\ell,k}^w$ has degree 2 by the same lemma. Furthermore, by $(\diamond g_1)b_2 = b_1(\diamond g_2)$ the monomials b_1 and b_2 must have the same degree because $\diamond g_1$ and $\diamond g_2$ do. As it was assumed that $(b_1, b_2) \neq (1, 1)$ that only leaves the possibility that $\{b_1, b_2\} \subseteq E$.

It follows that the identity $(\diamond g_1)b_2 = b_1(\diamond g_2)$, which by Lemma 3.6 is equivalent to

$$w_{\ell,n} w_{1,\mathbf{r}(k)}^{\ddagger} b_2 = b_1 v_{j,n} v_{1,\mathbf{r}(i)}^{\ddagger},$$

requires $b_1 = w_{\ell,n}$ and $b_2 = v_{1,r(i)}^{\ddagger}$ and, most of all, $v_{j,n} = w_{1,r(k)}^{\ddagger}$. By Lemmata 3.2 and 3.3 the latter demands $w^{\ddagger} \in \{v, v^{\dagger}\}$, which is to say $w \in \{v^{\star}, v^{\ddagger}\}$. If $w = v^{\star}$ was true, it would follow that $v_{j,n} = v_{r(k),1}$ and thus (j,n) = (r(k), 1) by Lemma 3.3. Because that would violate the assumption $2 \leq n$ it must instead be the case that $w = v^{\ddagger}$ and thus $b_1 = v_{\ell,n}^{\ddagger}$ and $v_{j,n} = v_{1,r(k)}$. According to Lemma 3.3 then (j,n) = (1,r(k)), i.e., j = k = 1. Thus, $g_1 = e_{\ell,1}^{v^{\ddagger}}$ and $g_2 = e_{1,i}^{v}$. That proves the claim.

Lemma 3.10. All overlap ambiguities of R resolve.

Proof. By Lemma 3.9 it is enough to prove that for any $v \in V$ and $w := v^{\ddagger}$ and any $(\ell, i) \in [n]^{\times 2}$ the polynomials $(\diamond e_{\ell,1}^w - e_{\ell,1}^w) w_{1,\mathbf{r}(i)}$ and $w_{\ell,n}(\diamond e_{1,i}^v - e_{1,i}^v)$ have a common reduction.

On the one hand, the definitions imply

~ ~ ~

$$\begin{aligned} (\diamond e_{\ell,1}^w - e_{\ell,1}^w) w_{1,\mathbf{r}(i)} &= (-\sum_{s=2}^n w_{\ell,\mathbf{r}(s)} w_{s,n}^{\ddagger} + \delta_{\ell,1} 1) w_{1,\mathbf{r}(i)} \\ &= -\sum_{s=2}^n w_{\ell,\mathbf{r}(s)} v_{s,n} v_{1,\mathbf{r}(i)}^{\ddagger} + \delta_{\ell,1} w_{1,\mathbf{r}(i)} \\ &= -\sum_{s=2}^n w_{\ell,\mathbf{r}(s)} (\diamond e_{s,i}^v) + \delta_{\ell,1} w_{1,\mathbf{r}(i)}, \end{aligned}$$

which upon reduction by $e_{s,i}^v$ for each $s \in [n]$ with 1 < s becomes

$$(\diamond e_{\ell,1}^{w} - e_{\ell,1}^{w}) w_{1,\mathbf{r}(i)} \rightsquigarrow -\sum_{s=2}^{n} w_{\ell,\mathbf{r}(s)} (-\sum_{t=2}^{n} v_{s,\mathbf{r}(t)} v_{t,\mathbf{r}(i)}^{\ddagger} + \delta_{s,i} 1) + \delta_{\ell,1} w_{1,\mathbf{r}(i)} = \sum_{s,t=2}^{n} w_{\ell,\mathbf{r}(s)} v_{s,\mathbf{r}(t)} w_{t,\mathbf{r}(i)} - w_{\ell,\mathbf{r}(i)} + \delta_{i,1} w_{\ell,n} + \delta_{\ell,1} w_{1,\mathbf{r}(i)}, \quad (3)$$

where the last identity is due to

$$\sum_{s=2}^{n} \delta_{s,i} w_{\ell,\mathbf{r}(s)} = \sum_{s=1}^{n} \delta_{s,i} w_{\ell,\mathbf{r}(s)} - \delta_{1,i} w_{\ell,\mathbf{r}(1)} = w_{\ell,\mathbf{r}(i)} - \delta_{i,1} w_{\ell,n}.$$

On the other hand, by definition

$$w_{\ell,n}(\diamond e_{1,i}^{v} - e_{1,i}^{v}) = w_{\ell,n}(-\sum_{t=2}^{n} v_{1,r(t)}v_{t,r(i)}^{\ddagger} + \delta_{1,i}1)$$

= $-\sum_{t=2}^{n} w_{\ell,n}w_{1,r(t)}^{\ddagger}w_{t,r(i)} + \delta_{i,1}w_{\ell,n}$
= $-\sum_{t=2}^{n}(\diamond e_{\ell,t}^{w})w_{t,r(i)} + \delta_{i,1}w_{\ell,n},$

which when reduced by $e^w_{\ell,t}$ for each $t \in [n]$ with 1 < t takes on the form

$$w_{\ell,n}(\diamond e_{1,i}^{v} - e_{1,i}^{v}) \rightsquigarrow -\sum_{t=2}^{n} (-\sum_{s=2}^{n} w_{\ell,\mathbf{r}(s)} w_{s,\mathbf{r}(t)}^{\ddagger} + \delta_{\ell,t} 1) w_{t,\mathbf{r}(i)} + \delta_{i,1} w_{\ell,n} = \sum_{s,t=2}^{n} w_{\ell,\mathbf{r}(s)} v_{s,\mathbf{r}(t)} w_{t,\mathbf{r}(i)} - w_{\ell,\mathbf{r}(i)} + \delta_{\ell,1} w_{1,\mathbf{r}(i)} + \delta_{i,1} w_{\ell,n}, \quad (4)$$

where now the last step comes from

$$\sum_{t=2}^{n} \delta_{t,\ell} w_{t,\mathbf{r}(i)} = \sum_{t=1}^{n} \delta_{t,\ell} w_{t,\mathbf{r}(i)} - \delta_{1,\ell} w_{1,\mathbf{r}(i)} = w_{\ell,\mathbf{r}(i)} - \delta_{\ell,1} w_{1,\mathbf{r}(i)}.$$

Since (3) and (4) have the same right-hand sides, all overlap ambiguities resolve.

Proposition 3.11. *R* is a Gröbner basis of *J*.

Proof. Follows from Lemmata 3.8 and 3.10 by Proposition 2.1, Bergman's diamond lemma. \Box

3.3. Reduced Gröbner basis. Wth Proposition 3.11 at hand it is now not hard to prove that G is the reduced Gröbner basis.

Lemma 3.12. For any $v \in V$,

$$f^v = e^v_{n,n} - \sum_{s=1}^{n-1} e^{v^{t}}_{s,s}$$

Proof. If $w := v^{t}$, then for any $s \in [n]$ by definition

$$e_{s,s}^{w} = w_{s,n}w_{1,r(s)}^{\ddagger} - (-\sum_{t=2}^{n} w_{s,r(t)}w_{t,r(s)}^{\ddagger} + 1)$$

= $\sum_{t=1}^{n} v_{r(t),s}v_{r(s),t}^{\ddagger} - 1$
= $\sum_{t=1}^{n} v_{t,s}v_{r(s),r(t)}^{\ddagger} - 1.$

Summing up these identities thus yields

$$\sum_{s=1}^{n-1} e_{s,s}^w = \sum_{s=1}^{n-1} \sum_{t=1}^n v_{t,s} v_{\mathbf{r}(s),\mathbf{r}(t)}^{\ddagger} - (n-1)\mathbf{1}$$
$$= \sum_{s=2}^n \sum_{t=1}^n v_{t,\mathbf{r}(s)} v_{\mathbf{s},\mathbf{r}(t)}^{\ddagger} - (n-1)\mathbf{1}.$$

Hence, subtracting this equality from the definition

$$e_{n,n}^v = v_{n,n}v_{1,1}^{\ddagger} + \sum_{s=2}^n v_{n,\mathbf{r}(s)}v_{s,\mathbf{r}(n)}^{\ddagger} - 1$$

of $e_{n,n}^v$ yields

$$e_{n,n}^{v} - \sum_{s=1}^{n-1} e_{s,s}^{w} = v_{n,n} v_{1,1}^{\ddagger} - \sum_{s=2}^{n} \sum_{t=1}^{n-1} v_{t,r(s)} v_{s,r(t)}^{\ddagger} + (n-2)1 = f^{v},$$
which is what was claimed.

Proposition 3.13. G is the reduced Gröbner basis of J.

Proof. Recall that G results from R by replacing $\{e_{n,n}^v | v \in V\}$ by $\{f^v | v \in V\}$. Lemma 3.12 therefore shows that G generates the same ideal as R. Moreover, $\diamond G = \diamond R$ by Lemma 3.6. Since R is a Gröbner basis of J by Proposition 3.11, hence so is G.

It remains to prove that G is reduced, i.e., that the tip of any element of G does not divide any monomial occurring in any other element of G. In fact, for degree reasons, any such divisibility relation is already necessarily an identity. Let $\{g, g'\} \subseteq G$ be arbitrary and suppose that $\diamond g$ occurs in g'. By definition of G there then exist $\{v, w\} \subseteq V$ and $\{(j,i), (\ell, k)\} \subseteq [n]^{\times 2}$ such that $(\ell, k) \neq (n, n) \neq (j, i)$ and $g \in \{f^w, e^w_{\ell, k}\}$ and $g' \in \{f^v, e^v_{j, i}\}$.

Step 1. First, $(g,g') \neq (e_{\ell,k}^w, f^v)$ is proved indirectly. If (g,g') was $(e_{\ell,k}^w, f^v)$, there would exist $(t,s) \in [n]^{\times 2}$ such that either both 1 < s and t < n or (t,s) = (n,1) and such that $w_{\ell,n}w_{1,r(k)}^{\dagger} = v_{t,r(s)}v_{s,r(t)}^{\dagger}$. By Lemmata 3.2 and 3.3 it would follow $w \in \{v, v^t\}$.

Case 1.1. If w = v was the case, then Lemma 3.3 would demand both $(\ell, n) = (t, \mathbf{r}(s))$ and $(1, \mathbf{r}(k)) = (s, \mathbf{r}(t))$. In other words, $\ell = t = k$ and s = 1. By assumption on s that would force t = n and thus $(\ell, k) = (n, n)$ in contradiction to the assumptions on ℓ and k.

Case 1.2. And in the opposite case that $w = v^t$ the same lemma would require both $(\ell, n) = (\mathbf{r}(s), t)$ and $(1, \mathbf{r}(k)) = (\mathbf{r}(t), s)$, which is to say $\ell = \mathbf{r}(s) = k$ and t = n. Again, t = n would entail s = 1, which would yield the contradiction $(\ell, k) = (n, n)$.

Step 2. Similarly, $(g, g') \neq (f^w, e_{j,i}^v)$. Indeed, if (g, g') was $(f^w, e_{j,i}^v)$, there would exist $s \in [n]$ such that $w_{n,n}w_{1,1}^{\ddagger} = v_{j,r(s)}v_{s,r(i)}^{\ddagger}$. Lemmata 3.2 and 3.3 would imply $w \in \{v, v^t\}$ and hence (j, r(s)) = (n, n) and (s, r(i)) = (1, 1). In particular, $(j, i) \neq (n, n)$ would be violated.

Step 3. From Steps 1 and 2 it follows $(g, g') \in \{(f^w, f^v), (e^w_{\ell,k}, e^v_{j,i})\}$. Hence, there are two cases to distinguish. It is enough to prove g = g' in each of them.

Case 3.1: If $(g, g') = (f^w, f^v)$, then there need to exist $(t, s) \in [n]^{\times 2}$ such that either both 1 < s and t < n or (t, s) = (n, 1) and such that $w_{n,n}w_{1,1}^{\ddagger} = v_{t,r(s)}v_{s,r(t)}^{\ddagger}$. Lemmata 3.2 and 3.3 then force in particular $w \in \{v, v^t\}$. Because $f^{v^t} = f^v$ that proves g = g', no matter whether w = v or $w = v^t$.

Case 3.2: Alternatively, if $(g, g') = (e_{\ell,k}^w, e_{j,i}^v)$, there is $s \in [n]$ such that $w_{\ell,n}w_{1,r(k)}^{\ddagger} = v_{j,r(s)}v_{s,r(i)}^{\ddagger}$. Hence, $w \in \{v, v^t\}$ by Lemmata 3.2 and 3.3. If w was v^t , that would require both $(n, \ell) = (j, r(s))$ and (r(k), 1) = (s, r(i)) and thus in particular (j, i) = (n, n), which is false. Hence, instead, w = v and thus $(\ell, n) = (j, r(s))$ and (1, r(k)) = (s, r(i)). That implies $(\ell, k) = (j, i)$ and, in conclusion, g = g', which means that G is reduced.

4. Anick resolution for U_n^+

In addition to the assumptions from Section 3 let also ε and \mathbb{K}_{ε} be as in the Main result. In this section, the Anick resolution of \mathbb{K}_{ε} with respect to \leq as a right *A*-module is computed explicitly by solving the determining recursion equations.

4.1. Auxiliary results. Recall that \diamond indicates tips and let M be the core of $\diamond J$, i.e., the minimal generating set of the monoidal ideal $\diamond J$ in the monoid $\langle E \rangle$, as explained in Section 2.2.

The next proposition guarantees in particular that $E \cap \diamond J = \emptyset$ and thus that Anick's construction is applicable.

Proposition 4.1.
$$M = \{v_{j,n}v_{1,r(i)}^{\ddagger} | v \in V \land (j,i) \in [n]^{\times 2}\}$$

Proof. By Proposition 3.11 the set G is a reduced and thus in particular minimal Gröbner basis of J, whence $M = \diamond G$. Thus, Lemma 3.6 proves the claim.

The following simple fact will be used so often that it seems worthy of its own lemma.

Lemma 4.2. For any $v \in V$ and $\{j, i\} \subseteq [n]^{\times 2}$, $v_{j_1,i_1}v_{i_2,j_2}^{\dagger} \notin \diamond J$

if and only if

$$(i_1, i_2) \neq (n, 1) \neq (j_1, j_2).$$

Proof. The contraposition, $b := v_{j_1,i_1} v_{i_2,j_2}^{\ddagger} \in \diamond J$ if and only if $(i_1, i_2) = (n, 1)$ or $(j_1, j_2) = (n, 1)$, is what is proved below.

Sufficient condition. First, if $(i_1, i_2) = (n, 1)$, then $b = v_{j_1,n} v_{1,r(r(j_2))}^{\ddagger} = \diamond e_{j_1,r(j_2)}^v$ by Lemma 3.6 and thus $b \in \diamond J$ because $e_{j_1,r(j_2)}^v \in R$ and $R \subseteq J$. Likewise, if $(j_1, j_2) = (n, 1)$ and if $w := v^t$, then $b = v_{n,i_1} v_{i_2,1}^{\ddagger} = w_{i_1,n} w_{1,r(r(i_2))}^{\ddagger} = \diamond e_{i_1,r(i_2)}^w$ and hence $b \in \diamond J$. Necessary condition. Conversely, if $b \in \diamond J$, then there exist $a \in V$.

Necessary condition. Conversely, if $b \in \diamond J$, then there exist $a \in M$ and $\{b_1, b_2\} \subseteq \langle E \rangle$ with $b_1 a b_2 = b$. Because a has degree 2 by Proposition 4.1, the same degree as b, that demands $b_1 = b_2 = 1$ and thus a = b, which then proves that actually $b \in M$. Hence, by a second application of Proposition 4.1 there exist $w \in V$ and $(t, s) \in [n]^{\times 2}$ with

$$v_{j_1,i_1}v_{i_2,j_2}^{\dagger} = w_{t,n}v_{1,\mathbf{r}(s)}^{\dagger}.$$
(5)

By Lemmata 3.2 and 3.3 that requires $w \in \{v, v^t\}$.

Case 1. If w = v, then according to Lemma 3.3 the identity (5) demands $(j_1, i_1) = (t, n)$ and $(i_2, j_2) = (1, s)$, in particular $(i_1, i_2) = (n, 1)$.

Case 2. Alternatively, if $w = v^t$, then (5) requires $(i_1, j_1) = (t, n)$ and $(j_2, i_2) = (1, s)$ by Lemma 3.3, which implies $(j_1, j_2) = (n, 1)$ in particular. That concludes the proof.

For any $\ell \in \mathbb{Z}_{>0}$, any $v \in V$ and any $(j,i) \in [n]^{\times 2}$ let $x_{j,i}^{v,\ell}$ and ℓv be as in the Main result. It is important to note that there are non-trivial identities between the $x_{j,i}^{v,\ell}$ for different values of (v, j, i) – however, only for $\ell \leq 2$.

Lemma 4.3. For any $\{v, w\} \subseteq V$, any $\{(j, i), (t, s)\} \subseteq [n]^{\times 2}$ and any $\ell \in \mathbb{Z}_{>0}$ the identity

$$x_{t,s}^{w,\ell} = x_{j,i}^{v,\ell}$$

holds if and only if both w = v and (t, s) = (j, i) or one of the following mutually exclusive conditions is met:

 $\Box \ \ell = 1 \ and \ w = v^{\mathsf{t}} \ and \ (t,s) = (\mathsf{r}(i),\mathsf{r}(j))$ $\Box \ \ell = 2 \ and \ w = v^{\mathsf{t}} \ and \ (t,s) = (n,n) = (j,i).$

Proof. The proof requires a distinction of cases.

Case 1. If $\ell = 1$, the definitions and Lemma 3.3 imply that

$$x_{t,s}^{w,1} = w_{t,\mathbf{r}(s)} = v_{j,\mathbf{r}(i)} = x_{j,i}^{v,1}$$

is satisfied if and only if either both w = v and $(t, \mathbf{r}(s)) = (j, \mathbf{r}(i))$ or both $w = v^{t}$ and $(t, \mathbf{r}(s)) = (\mathbf{r}(i), j)$. That proves the claim in this case.

Case 2. Similarly, if $\ell = 2$, then the identity

$$x_{t,s}^{w,2} = w_{t,n} w_{1,\mathbf{r}(s)}^{\ddagger} = v_{j,n} v_{1,\mathbf{r}(i)}^{\ddagger} = x_{j,i}^{v,2}$$

holds if and only if both $w_{t,n} = v_{j,n}$ and $w_{1,r(s)}^{\ddagger} = v_{1,r(i)}^{\ddagger}$. By Lemma 3.3 that is true if and only if either w = v and (t, n) = (j, n) and (1, r(s)) = (1, r(i)) or $w = v^{\dagger}$ and (n, t) = (j, n) and (r(s), 1) = (1, r(i)). The first of these possibilities, of course, means that (w, t, s) = (v, j, i), the other that $w = v^{\dagger}$ and (t, s) = (n, n) = (j, i). Hence, the claim is true for $\ell = 2$ as well.

Case 3. If, next, $3 \leq \ell$, then what needs to be proved is that $x_{t,s}^{w,\ell}$ can only equal $x_{j,i}^{v,\ell}$ if (w,t,s) = (v,j,i). The vectors $x_{t,s}^{w,\ell}$ and $x_{j,i}^{v,\ell}$ being equal is equivalent to the conditions that either ℓ is odd and

$$w_{t,n}(w_{1,n}^{\dagger}w_{1,n})^{\frac{\ell-3}{2}}w_{1,n}^{\dagger}w_{1,\mathbf{r}(s)} = v_{j,n}(v_{1,n}^{\dagger}v_{1,n})^{\frac{\ell-3}{2}}v_{1,n}^{\dagger}v_{1,\mathbf{r}(i)}$$
(6)

or ℓ is even and

$$w_{t,n}(w_{1,n}^{\dagger}w_{1,n})^{\frac{\ell-4}{2}}w_{1,n}^{\dagger}w_{1,n}w_{1,r(s)}^{\dagger} = v_{j,n}(v_{1,n}^{\dagger}v_{1,n})^{\frac{\ell-4}{2}}v_{1,n}^{\dagger}v_{1,n}v_{1,r(i)}^{\dagger}.$$
 (7)

Any of (6) and (7) requires in particular $w_{1,n}^{\ddagger} = v_{1,n}^{\ddagger}$. By Lemma 3.3 that is only possible if either both w = v and (1, n) = (1, n) or both $w = v^{t}$ and (n, 1) = (1, n). As the second of these conclusions would violate the assumption that $2 \le n$ it must be the first one which is true.

Of course, (6) and (7) moreover demand, on the one hand, $w_{t,n} = v_{j,n}$ and, on the other hand, $w_{1,r(s)} = v_{1,r(i)}$ respectively $w_{1,r(s)}^{\dagger} = v_{1,r(i)}^{\dagger}$. According to Lemma 3.3, given that w = v and thus $w^{\dagger} = v^{\dagger}$, these identities are only satisfied if (t, n) = (j, n) and (1, r(s)) = (1, r(i)). But

that is only true if (t, s) = (j, i). Thus, the claim has been verified for all ℓ .

It will moreover be necessary to make comparisons between those vectors with respect to \leq .

Lemma 4.4. For any $\ell \in \mathbb{Z}_{>0}$, any $\{v, w\} \subseteq V$ and $\{(j, i), (t, s)\} \subseteq$ $[n]^{\times 2}$, the inequality

$$x_{t,s}^{w,\ell} < x_{j,i}^{v,\ell}$$

holds if and only if either both $w \in \{v^*, v^{\ddagger}\}$ and $v \in \{u, u^{t}\}$ or one of the following mutually exclusive conditions is met.

 $\Box \ \ell = 1$ and, alternatively, $-w = v \text{ and } v \in \{u, u^{\star}\} \text{ and } (t, i) < (j, s)$ -w = v and $v \in \{u^{t}, u^{\ddagger}\}$ and (i, t) < (s, j) $-w = v^{t}$ and $v \in \{u, u^{\star}\}$ and $(\mathbf{r}(j), t) < (s, \mathbf{r}(i))$ $-w = v^{t} \text{ and } v \in \{u^{t}, u^{t}\} \text{ and } (t, \mathbf{r}(j)) < (\mathbf{r}(i), s).$ \Box $\ell = 2$ and, alternatively, -w = v and (t, i) < (j, s). $-w = v^{t}$ and $v \in \{u, u^{\star}\}$ and j = n and either $t \neq n$ or $t = n \neq i$. $-w = v^{t}$ and $v \in \{u^{t}, u^{t}\}$ and either $t \neq n$ or $t = s = n = j \neq i$. $\square 3 < \ell$ and, alternatively, -w = v and (t, i) < (j, s). $-w = v^{t}$ and $v \in \{u, u^{\star}\}$ and j = n. $-w = v^{t}$ and $v \in \{u^{t}, u^{t}\}$ and $t \neq n$. In particular, $x_{j,s}^{v,\ell} < x_{j,i}^{v,\ell}$ if and only if i < s and, likewise, $x_{t,i}^{v,\ell} < x_{j,i}^{v,\ell}$ if and only if j < t.

Proof. No matter the value of ℓ , there exist $\{b, a\} \subseteq [n]$ and $\{q, p\} \subseteq \langle E \rangle$ such that $x_{t,s}^{w,\ell} = w_{t,b}q$ and $x_{j,i}^{v,\ell} = v_{j,a}p$. If $w \in \{v^*, v^{\ddagger}\}$, then already $w_{t,b} < v_{j,a}$ if and only if $v \in \{u, u^{\ddagger}\}$ by Lemma 3.4 and thus, as claimed, $x_{t,s}^{w,\ell} < x_{j,i}^{v,\ell}$ if and only if $v \in \{u, u^t\}$ since \leq was extended (degreewise) lexicographically. For that reason it can be assumed in the following that $w \notin \{v^*, v^{\ddagger}\}$, which by Lemma 3.2 means that $w \in \{v, v^{\dagger}\}$. It is necessary to distinguish both between different values of ℓ and between whether w is v or v^{t} .

Case 1. If $\ell = 1$, then by Lemma 3.4 the inequality

$$x_{t,s}^{w,1} = w_{t,\mathbf{r}(s)} < v_{j,\mathbf{r}(i)} = x_{j,i}^{v,1}$$

is true if and only if either w = v and $v \in \{u, u^*\}$ and $(t, \mathbf{r}(s)) < (j, \mathbf{r}(i))$ or w = v and $v \in \{u^{t}, u^{\dagger}\}$ and $(\mathbf{r}(s), t) < (\mathbf{r}(i), j)$ or $w = v^{t}$ and $v \in \{u, u^{\star}\}$ and $(\mathbf{r}(s), t) < (j, \mathbf{r}(i))$ or $w = v^{t}$ and $v \in \{u^{t}, u^{\dagger}\}$ and $(t, \mathbf{r}(s)) < (\mathbf{r}(i), j)$. Since r reverses the order of [n] that proves the claim in this case.

Case 2. For $\ell = 2$, the inequality $x_{t,s}^{w,2} = w_{t,n} w_{1,r(s)}^{\ddagger} < v_{j,n} v_{1,r(i)}^{\ddagger} = x_{j,i}^{v,2}$ is equivalent to

$$w_{t,n}w_{\mathbf{r}(s),1} < v_{j,n}v_{\mathbf{r}(i),1}$$
 (8)

by Lemma 3.5 since $w \in \{v, v^t\}$. It is necessary to distinguish further whether w is v and v^t and between different values of v.

Case 2.1. If w = v, then by Lemma 3.3 and by the addendum to Lemma 3.4 condition (8) is equivalent to the statement that either t < j or both t = j and r(s) < r(i), which is to say i < s. In other words, (8) is equivalent to (t, i) < (j, s) if w = v, as asserted.

Case 2.2. If, next, $w = v^t$ and $v \in \{u, u^*\}$, then Lemmata 3.3 and 3.4 imply that (8) holds if and only if either (n, t) < (j, n) or both (n, t) = (j, n) and (1, r(s)) < (r(i), 1). Those conditions are equivalent to either t < n = j or both t = n = j and 1 < r(i), i.e., i < n, being true. In particular, j = n is required for (8) to hold. And, if so, then (8) is valid if and only if either $t \neq n$ or both t = n and $i \neq n$.

Case 2.3. Finally, if $w = v^t$ and $v \in \{u^t, u^t\}$, then by Lemmata 3.3 and 3.4 the inequality (8) is satisfied if and only if either (t, n) < (n, j)or both (t, n) = (n, j) and $(\mathbf{r}(s), 1) < (1, \mathbf{r}(i))$. This is true if and only if either t < n or both t = n = j and $\mathbf{r}(s) = 1 < \mathbf{r}(i)$, i.e., either $t \neq n$ or $t = s = n = j \neq i$. That is what was claimed in this case.

Case 3. Lastly, for $3 \leq \ell$, the definitions imply that $x_{t,s}^{w,\ell} < x_{j,i}^{v,\ell}$ is true if and only if either ℓ is odd and

$$w_{t,n}(w_{1,n}^{\dagger}w_{1,n})^{\frac{\ell-3}{2}}w_{1,n}^{\dagger}w_{1,\mathbf{r}(s)} < v_{j,n}(v_{1,n}^{\dagger}v_{1,n})^{\frac{\ell-3}{2}}v_{1,n}^{\dagger}v_{1,\mathbf{r}(i)}$$

or ℓ is even and

$$w_{t,n}(w_{1,n}^{\ddagger}w_{1,n})^{\frac{\ell-2}{2}}w_{1,\mathbf{r}(s)}^{\ddagger} < v_{j,n}(v_{1,n}^{\ddagger}v_{1,n})^{\frac{\ell-2}{2}}v_{1,\mathbf{r}(i)}^{\ddagger}$$

Since $w \in \{v, v^t\}$ these conditions can be simplified with the help of Lemma 3.5 into the form that, if ℓ is odd,

$$w_{t,n}w_{n,1}(w_{1,n}w_{n,1})^{\frac{\ell-3}{2}}w_{1,\mathbf{r}(s)} < v_{j,n}v_{n,1}(v_{1,n}v_{n,1})^{\frac{\ell-3}{2}}v_{1,\mathbf{r}(i)}$$
(9)

and, if ℓ is even,

$$w_{t,n}w_{n,1}(w_{1,n}w_{n,1})^{\frac{\ell-4}{2}}w_{1,n}w_{\mathbf{r}(s),1} < v_{j,n}v_{n,1}(v_{1,n}v_{n,1})^{\frac{\ell-4}{2}}v_{1,n}v_{\mathbf{r}(i),1}.$$
 (10)

Again, the argument depends on the value of (w, v).

Case 3.1. If w = v, then also $w_{n,1}(w_{n,1}w_{1,n})^{\frac{\ell-3}{2}} = v_{n,1}(v_{n,1}v_{1,n})^{\frac{\ell-3}{2}}$ if ℓ is odd and, likewise, $w_{n,1}(w_{1,n}w_{n,1})^{\frac{\ell-4}{2}}w_{1,n} = v_{n,1}(v_{1,n}v_{n,1})^{\frac{\ell-4}{2}}v_{1,n}$ if ℓ is even. Hence, (9) is true if and only if $v_{t,n}v_{1,r(s)} < v_{j,n}v_{1,r(i)}$ and (10) if and only if $v_{t,n}v_{r(s),1} < v_{j,n}v_{r(i),1}$. By Lemma 3.3 and the addendum to Lemma 3.4 that is true if and only if either t < j or both t = j and r(s) < r(i), regardless of what parity ℓ has. That means $x_{t,s}^{w,\ell} < x_{j,i}^{v,\ell}$ if and only if (t,i) < (j,s), as was claimed.

Case 3.2. If $w = v^t$ and $v \in \{u, u^*\}$, then the assumption $2 \leq n$ requires (1, n) < (n, 1) and thus $w_{n,1} < v_{n,1}$ by Lemma 3.4. In consequence,

each of (9) and (10) is satisfied if and only if $w_{t,n} \leq v_{j,n}$. According to Lemmata 3.3 and 3.4 that is equivalent to the condition that $(n,t) \leq (j,n)$. Because the latter is met if and only if j = n that proves the claim in this case.

Case 3.3. In the final case that $w = v^t$ and $v \in \{u^t, u^t\}$ the fact that (n, 1) < (1, n) is false implies by Lemma 3.4 that also $w_{n,1} < v_{n,1}$ is false. Hence, any one of (9) and (10) is equivalent to $w_{t,n} < v_{j,n}$. By Lemma 3.4 that is true if and only if (t, n) < (n, j), i.e., if $t \neq n$. As this is what was claimed in this case that concludes the proof. \Box

The central identity for the recursion will be the following.

Lemma 4.5. For any $\ell \in \mathbb{Z}_{>0}$, any $v \in V$ and any $(j,i) \in [n]^{\times 2}$, if $2 \leq \ell$, then

$$x_{j,i}^{v,\ell} = x_{j,1}^{v,\ell-1} \cdot {}^{\ell}v_{1,\mathbf{r}(i)}$$
(11)

and, if $3 \leq \ell$, then

$$x_{j,i}^{v,\ell-1} = x_{j,1}^{v,\ell-2} \cdot {}^{\ell}v_{1,\mathbf{r}(i)}^{\ddagger}, \qquad (12)$$

where \cdot is the multiplication of $\mathbb{K}\langle E \rangle$.

Proof. Because ${}^{\ell}v_{1,r(i)}^{\ddagger} = {}^{\ell-1}v_{1,r(i)}$ identity (12) follows from (11), which is thus the only one that needs proving.

Case 1. If $\ell = 2$, then by definition

$$x_{j,i}^{v,2} = v_{j,n} v_{1,\mathbf{r}(i)}^{\ddagger}, \qquad x_{j,1}^{v,1} = v_{j,\mathbf{r}(1)} = v_{j,n}, \qquad {}^{\ell} v_{1,\mathbf{r}(i)} = v_{1,\mathbf{r}(i)}^{\ddagger}$$

where the last identity is due to ℓ being even. That proves (11) in this case.

Case 3. If ℓ is odd with $3 \leq \ell$, then ${}^{\ell}v_{1,\mathbf{r}(i)} = v_{1,\mathbf{r}(i)}$ as well as

$$\begin{aligned} x_{j,i}^{v,\ell} &= v_{j,n} (v_{1,n}^{\dagger} v_{1,n})^{\frac{\ell-3}{2}} v_{1,n}^{\dagger} v_{1,r(i)}, \\ x_{j,1}^{v,\ell-1} &= v_{j,n} (v_{1,n}^{\dagger} v_{1,n})^{\frac{(\ell-1)-2}{2}} v_{1,r(1)}^{\dagger} &= v_{j,n} (v_{1,n}^{\dagger} v_{1,n})^{\frac{\ell-3}{2}} v_{1,n}^{\dagger}. \end{aligned}$$

Comparing the polynomials now proves (11).

Case 3. Finally, if ℓ is even and $4 \leq \ell$, then it was defined that

$$\begin{aligned} x_{j,i}^{v,\ell} &= v_{j,n} (v_{1,n}^{\dagger} v_{1,n})^{\frac{\ell-2}{2}} v_{1,r(i)}^{\dagger} = v_{j,n} (v_{1,n}^{\dagger} v_{1,n})^{\frac{\ell-4}{2}} v_{1,n}^{\dagger} v_{1,n} v_{1,r(i)}^{\dagger}, \\ x_{j,1}^{v,\ell-1} &= v_{j,n} (v_{1,n}^{\dagger} v_{1,n})^{\frac{(\ell-1)-3}{2}} v_{1,n}^{\dagger} v_{1,r(1)} = v_{j,n} (v_{1,n}^{\dagger} v_{1,n})^{\frac{\ell-4}{2}} v_{1,n}^{\dagger} v_{1,n} \end{aligned}$$

since $\ell - 1$ is odd. Given that by definition ${}^{\ell}v_{1,r(i)} = v_{1,r(i)}^{\ddagger}$ the identity (11) is satisfied in this instance as well.

4.2. Chains, combinatorial differentials, cycles, splittings. When computing Anick's resolution, the first order of business is to determine the sets of chains and the combinatorial differentials, cycles and splittings.

4.2.1. Chains. For any $\ell \in \mathbb{Z}_{\geq 0}$ let pC_{ℓ} respectively C_{ℓ} denote the sets of ℓ -pre-chains and -chains with respect to J and \leq .

Lemma 4.6. For any $\ell \in \mathbb{Z}_{>0}$ with $2 \leq \ell$, any $v \in V$ and any $(j,i) \in [n]^{\times 2}$ any $c \in \langle E \rangle$ is equal to $x_{j,i}^{v,\ell}$ if and only if there exists $e \in E^{\times \ell}$ such that $c = e_1 e_2 \dots e_{\ell}$ and such that for any $s \in [\ell - 1]$,

$$e_{s}e_{s+1} = \begin{cases} {}^{s}v_{j,n}{}^{s}v_{1,n}^{\ddagger} & |s=1 \\ {}^{s}v_{1,n}{}^{s}v_{1,n}^{\ddagger} & |2 \leq s \leq \ell - 2 \\ {}^{s}v_{1,n}{}^{s}v_{1,r(i)}^{\ddagger} & |s=\ell-1. \end{cases}$$
(13)

Proof. Follows from the definition by distinguishing between even and odd ℓ .

Proposition 4.7. For any $\ell \in \mathbb{Z}_{>0}$,

$$\mathbf{p}C_{\ell} = C_{\ell} = \{ x_{j,i}^{v,\ell} | v \in V \land (j,i) \in [n]^{\times 2} \}.$$

Moreover, for any $v \in V$ and any $(j,i) \in [n]^{\times 2}$, if (α,β) are the chainindices of $x_{j,i}^{v,\ell}$, then $\alpha_s = s$ and $\beta_s = s + 1$ for any $s \in [\ell - 1]$.

Proof. The proof varies with ℓ . Hence, a case distinction is necessary.

Case 1. If $\ell = 1$, then $pC_1 = C_1 = E$ already by definition and the addendum is vacuously true. The only actual claim is that $E = \{x_{j,i}^{v,1} | v \in V \land (j,i) \in [n]^{\times 2}\}$. But that is clear because $x_{j,i}^{v,1} = v_{j,r(i)}$ for any $v \in V$ and $(j,i) \in [n]^{\times 2}$.

Case 2. Also in the case that $\ell = 2$ most of the claims are clear a priori. Indeed, irrespective of the particular choices of E, J and \leq , it can be seen that $pC_2 = C_2 = M$ and that the chain indices of any 2-chain c are necessarily (1, m), where m is the degree of c. Hence, the only non-trivial claim in this case is that $M = \{x_{j,i}^{v,2} | v \in V \land (j,i) \in [n]^{\times 2}\}$. But again that is obvious by Proposition 4.1 because $x_{j,i}^{v,2} = v_{j,n}v_{1,r(i)}^{\ddagger}$ for any $v \in V$ and $(j,i) \in [n]^{\times 2}$.

Case 3. The proof for the case $3 \leq \ell$ is divided into two steps. The first is to prove $pC_{\ell} \subseteq \{x_{j,i}^{v,\ell} | v \in V \land (j,i) \in [n]^{\times 2}\}$. Because $C_{\ell} \subseteq pC_{\ell}$ by definition, it is then enough to check, second, that $\{x_{j,i}^{v,\ell} | v \in V \land (j,i) \in [n]^{\times 2}\} \subseteq C_{\ell}$, in the process of which the addendum will be proved as well.

Step 3.1. For any $m \in \mathbb{Z}_{\geq 0}$ and any $e \in E^{\times 2}$, if $c = e_1 e_2 \dots e_m \in pC_{\ell}$, then there exist $\alpha \in [m]^{\times \ell - 1}$ and $\beta \in [m]^{\times \ell - 1}$ such that $1 = \alpha_1 < \alpha_2 \leq \beta_1$ and $\alpha_{\ell-1} \leq \beta_{\ell-2} < \beta_{\ell-1} = m$ and $\alpha_{s+1} \leq \beta_s < \alpha_{s+2}$ for any $s \in [\ell - 3]$ and such that $e_{\alpha_s} e_{\alpha_s+1} \dots e_{\beta_s} \in M$ for any $s \in [\ell - 1]$. Since all elements of M have degree 2 by Proposition 4.1 this last condition implies $\beta_s = \alpha_s + 1$ for each $s \in [\ell - 1]$. That implies $\alpha_s = s$ and $\beta_s = s + 1$ for each $s \in [\ell - 1]$. In particular, $m = \beta_{\ell-1} = \ell$. In fact, by Proposition 4.1 for any $s \in [\ell - 1]$ there exist $v^{(s)} \in V$ and $(j_s, i_s) \in [n]^{\times 2}$ with $e_{\alpha_s} e_{\alpha_s+1} \dots e_{\beta_s} = e_s e_{s+1} = v_{j_s,n}^{(s)} v_{1,r(i_s)}^{(s)^{\ddagger}}$.

If $v := v^{(1)}$, if $j := j_1$ and if $i := i_{\ell-1}$, then $c = x_{j,i}^{v,\ell}$ by the following argument. For each $s \in [\ell - 2]$ the equality $\beta_s = \alpha_{s+1}$ demands $v_{1,r(i_s)}^{(s)\ddagger} = e_{s+1} = v_{j_{s+1},n}^{(s+1)}$. According to Lemma 3.3 that requires $v^{(s+1)} \in$ $\{v^{(s)\star}, v^{(s)\ddagger}\}$ for each $s \in [\ell - 2]$. If $v^{(s+1)}$ was $v^{(s)\star}$ for such s, it would follow that $(1, r(i_s)) = (n, j_{s+1})$, which would violate the assumption $2 \leq n$. Hence, instead, $v^{(s+1)} = v^{(s)\ddagger}$ and $(1, r(i_s)) = (j_{s+1}, n)$, i.e., $j_{s+1} = 1 = i_s$, for each $s \in [\ell - 2]$. In particular, $v^{(s)} = {}^{s}v$ for any $s \in [\ell - 1]$. Actually, altogether that proves that c satisfies the identities (13) in Lemma 4.6 and must thus be equal to $x_{j,i}^{v,\ell}$, as claimed.

(13) in Lemma 4.6 and must thus be equal to $x_{j,i}^{v,\ell}$, as claimed. Step 3.2. For any $v \in V$ and $(j,i) \in [n]^{\times 2}$, if $\alpha_s = s$ and $\beta_s = s + 1$ for any $s \in [\ell - 1]$, then $x_{j,i}^{v,\ell}$ is an ℓ -chain with chain indices (α, β) , as explained hereafter.

Of course, $1 = \alpha_1 < \alpha_2 \leq \beta_1$ by $\alpha_2 = 2 = \beta_1$. The degree of $x_{j,i}^{v,\ell}$ is ℓ and $\alpha_{\ell-1} \leq \beta_{\ell-2} < \beta_{\ell-1} = \ell$ holds by $\alpha_{\ell-1} = \ell - 1$ and $\beta_{\ell-2} = \ell - 1$. Likewise, for any $s \in [\ell - 3]$ the definitions $\alpha_{s+1} = s + 1$ and $\beta_s = s + 1$ and $\alpha_{s+2} = s + 2$ imply $\alpha_{s+1} \leq \beta_s < \alpha_{s+2}$.

Let $e \in E^{\times \ell}$ be the unique sequence of generators with $x_{j,i}^{v,\ell} = e_1 e_2 \dots e_\ell$. By Lemma 4.6 then the identities (13) hold. According to Proposition 4.1 those then prove $e_{\alpha_s} e_{\alpha_s+1} \dots e_{\beta_s} = e_s e_{s+1} \in M$ for each $s \in [\ell - 1]$. Hence, $x_{j,i}^{v,\ell} \in pC_\ell$. Moreover, for any $\ell' \in [\ell]$ with $2 \leq \ell'$ and any $t \in [\beta_{\ell'-1} - 1]$ the

Moreover, for any $\ell' \in [\ell]$ with $2 \leq \ell'$ and any $t \in [\beta_{\ell'-1} - 1]$ the monomial $e_1e_2 \ldots e_t$ is no element of $pC_{\ell'}$ because its degree is less than $\beta_{\ell'-1} = (\ell' - 1) + 1 = \ell'$, whereas by Step 3.1 any element of $pC_{\ell'}$ is of the form $x_{b,a}^{w,\ell'}$ for some $w \in V$ and $(b,a) \in [n]^{\times 2}$ and thus of degree ℓ' .

In conclusion, $x_{j,i}^{v,\ell}$ is an ℓ -chain and (α, β) are its chain indices. That verifies the claim in the case $3 \leq \ell$ and thus completes the proof overall.

4.2.2. Combinatorial differential. For any $\ell \in \mathbb{Z}_{>0}$ let ∂_{ℓ} be the combinatorial differential associated with J and \leq from Section 2.7.

Proposition 4.8. For any $\ell \in \mathbb{Z}_{>0}$ with $2 \leq \ell$, any $v \in V$ and any $(j,i) \in [n]^{\times 2}$,

$$\partial_{\ell}(x_{j,i}^{v,\ell}) = (x_{j,1}^{v,\ell-1}, {}^{\ell}v_{1,\mathbf{r}(i)}).$$

Proof. If (α, β) are the chain indices of $x_{j,i}^{v,\ell}$, then $\alpha_s = s$ and $\beta_s = s + 1$ for any $s \in [\ell - 1]$ by the addendum to Proposition 4.7. Accordingly, if $e \in E^{\times \ell}$ is such that $x_{j,i}^{v,\ell} = e_1 e_2 \dots e_\ell$, then by definition $\partial_\ell(x_{j,i}^{v,\ell}) := (e_1 e_2 \dots e_{\beta_{\ell-2}}, e_{\beta_{\ell-2}+1} e_{\beta_{\ell-2}+2} \dots e_\ell) = (e_1 e_2 \dots e_{\ell-1}, e_\ell)$. Since $e_1 e_2 \dots e_{\ell-1} = x_{j,1}^{v,\ell-1}$ and $e_\ell = {}^\ell v_{1,r(i)}$ by (11) in Lemma 4.5 that proves the claim.

4.2.3. Combinatorial cycles. For any $\ell \in \mathbb{Z}_{\geq 0}$ let Z_{ℓ} be the set of combinatorial cycles associated with J and \leq from Section 2.7.

Proposition 4.9. For any $\ell \in \mathbb{Z}_{>0}$ with $3 \leq \ell$,

$$Z_{\ell-2} = \{ (x_{j,1}^{v,\ell-2}, {}^{\ell}v_{1,\mathbf{r}(i)}^{\ddagger}h) | v \in V \land (j,i) \in [n]^{\times 2}, h \in \langle E \rangle \backslash \diamond J \}$$

Proof. Each inclusion is proved separately.

Step 1. For any $v \in V$ and any $(j,i) \in [n]^{\times 2}$ the pair $\partial_{\ell-2}(x_{j,1}^{v,\ell-2})$ is given by $(1, x_{j,1}^{v,1})$ if $\ell = 3$ and otherwise by $(x_{j,1}^{v,\ell-3}, \ell^{-2}v_{1,\mathbf{r}(1)})$ according to Proposition 4.8. By definition, $x_{j,1}^{v,1} = v_{j,\mathbf{r}(1)} = {}^{3}v_{j,n}$ and $\ell^{-2}v_{1,\mathbf{r}(1)} = \ell v_{1,n}$. Both ${}^{3}v_{j,n}{}^{3}v_{1,\mathbf{r}(i)}^{\ddagger}$ and $\ell v_{1,n}{}^{\ell}v_{1,\mathbf{r}(i)}^{\ddagger}$ are elements of $M \subseteq \diamond J$ by Proposition 4.1. Hence, for any $h \in \langle E \rangle \backslash \diamond J$ neither ${}^{3}v_{j,n}{}^{3}v_{1,\mathbf{r}(i)}^{\ddagger}h$ nor $\ell v_{1,n}{}^{\ell}v_{1,\mathbf{r}(i)}^{\ddagger}h$ is normal, which makes $(x_{j,1}^{v,\ell-2}, \ell v_{1,\mathbf{r}(i)}^{\ddagger}h)$ an element of $Z_{\ell-2}$.

Step 2. Let $\{m,t\} \subseteq \mathbb{Z}_{\geq 0}$ and $e \in E^{\times (m+t)}$ be arbitrary with the property, that, if $c = e_1 e_2 \dots e_m$ and $d = e_{m+1} e_{m+2} \dots e_{m+t}$, then $(c,d) \in Z_{\ell-2}$. By Proposition 4.7 there then exist $v \in V$ and $(j,s) \in [V]^{\times 2}$ with $c = x_{j,s}^{v,\ell-2}$ and, in particular, $m = \ell - 2$. Consequently, $(a,b) \coloneqq \partial_{\ell-2}(x_{j,s}^{v,\ell-2})$ is given by $(1, x_{j,s}^{v,1}) = (1, {}^1v_{j,r(s)})$ if $\ell = 3$ and otherwise by $(x_{j,1}^{v,\ell-3}, {}^{\ell-2}v_{1,r(s)})$ according to Proposition 4.8.

Since $(c, d) \in Z_{\ell-2}$ then $bd \in \diamond J$. Because $b \notin \diamond J$ that requires $d \neq 1$, which is to say $1 \leq t$. Since $d \notin \diamond J$ the assumption that $bd \in \diamond J$ then demands the existence of $z \in [t]$ with $be_{m+1}e_{m+2} \dots e_{m+z} \in \diamond J$. In fact there must exist $z \in [t]$ with $be_{m+1}e_{m+2} \dots e_{m+z} \in M$ because otherwise the assumption $d \notin \diamond J$ would be violated. In fact, by nature of Mthere can exist only such z. Even more precisely, because any element of M has degree 2 by Proposition 4.1 and since b has degree 1 the index z must be 1. In conclusion, if $h := e_{m+2}e_{m+3} \dots e_{m+t}$, then, first, $h \notin \diamond J$ because already $d \notin \diamond J$ and, second, $d = e_{m+1}h$ and $be_{m+1} \in M$. Another implication of Proposition 4.1 is therefore the existence of $w \in V$ and $(q, i) \in [n]^{\times 2}$ with $be_{m+1} = w_{q,n}w_{1,r(i)}^{\ddagger}$. By Lemma 3.3 the ensuing equalities $b = {}^{3}v_{j,r(s)} = w_{q,n}$ for $\ell = 3$ and $b = {}^{\ell}v_{1,r(s)} = w_{q,n}$ for $4 \leq \ell$ demand $w \in \{{}^{\ell}v, {}^{\ell}v^{t}\}$ for any $3 \leq \ell$.

Case 2.1. If $\ell = 3$ and $w = {}^{\ell}v$, then by the same lemma, $(j, \mathbf{r}(s)) = (q, n)$. Thus, $c = x_{j,1}^{v,1}$ and $e_{m+1} = w_{1,\mathbf{r}(i)}^{\ddagger} = v_{1,\mathbf{r}(i)}^{\ddagger}$ and hence $d = e_{m+1}h = {}^{3}v_{1,\mathbf{r}(i)}^{\ddagger}h$. Ultimately, $(c, d) = (x_{j,1}^{v,1}, {}^{3}v_{1,\mathbf{r}(i)}^{\ddagger}h)$. That proves the claim in this case.

Case 2.2. In case $\ell = 3$ and $w = {}^{\ell}v^{t}$, the equality $v_{j,r(s)} = w_{q,n}$ requires (j, r(s)) = (n, q) by Lemma 3.3. With the help of Lemma 4.3 it follows $c = x_{j,s}^{v,1} = x_{n,r(q)}^{v,1} = x_{q,1}^{v^{t},1}$. Because, moreover, $e_{m+1} = w_{1,r(i)}^{\dagger} = v_{1,r(i)}^{\star}$ and thus $d = e_{m+1}h = {}^{3}v_{1,r(i)}^{\star}h$ the conclusion $(c, d) = (x_{q,1}^{v^{t},1}, {}^{3}v_{1,r(i)}^{\star}h)$ proves the claim in this instance.

Case 2.3. Finally, for $4 \leq \ell$ Lemma 3.3 and the equality ${}^{\ell}v_{1,r(s)} = w_{q,n}$ actually require $w = {}^{\ell}v$ and (1, r(s)) = (q, n) because if w was ${}^{\ell}v^{t}$ the then inevitable identity between (1, r(s)) and (n, q) would contradict

the assumption $2 \leq n$. In conclusion, s = 1 and thus $c = x_{j,1}^{v,\ell-2}$ and $e_{m+1} = w_{1,r(i)}^{\dagger} = {}^{\ell}v_{1,r(i)}^{\dagger}$, i.e., $(c,d) = (x_{j,1}^{v,\ell-2}, {}^{\ell}v_{1,r(i)}^{\dagger}h)$, which concludes the proof.

4.2.4. Combinatorial splitting. For any $\ell \in \mathbb{Z}_{\geq 0}$ let \int_{ℓ} be the combinatorial splitting associated with J and \leq from Section 2.7.

Proposition 4.10. For any $\ell \in \mathbb{Z}_{>0}$ with $3 \leq \ell$, any $v \in V$, any $(j,i) \in [n]^{\times 2}$ and any $h \in \langle E \rangle \backslash \diamond J$,

$$\int_{\ell-2} (x_{j,1}^{v,\ell-2}, {}^{\ell}v_{1,\mathbf{r}(i)}^{\ddagger}h) = (x_{j,i}^{v,\ell-1}, h).$$

Proof. With the abbreviation $m := \ell - 2$, let $t \in \mathbb{Z}_{>0}$ and $e \in [n]^{\times (m+t)}$ be such that $x_{j,1}^{v,\ell-2} = e_1e_2 \dots e_m$ and ${}^{\ell}v_{1,r(i)}^{\ddagger}h = e_{m+1}e_{m+2}\dots e_{m+t}$. Since by Proposition 4.7 the chain indices (α, β) of $x_{j,1}^{v,\ell-2}$ satisfy $\alpha_s = s$ and $\beta_s = s+1$ for any $s \in [\ell-3]$ there is exactly one $\alpha_{\ell-2} \in \mathbb{Z}_{>0}$ such that $\alpha_{\ell-2} = 1$ if $\ell = 3$, such that $1 < \alpha_{\ell-2} \le m$ if $\ell = 4$ and such that $\beta_{\ell-4} < \alpha_{\ell-2} \le m$ if $5 \le \ell$, namely, $\alpha_{\ell-2} = m$. Because any element of M has degree 2 by Proposition 4.1 the unique $\beta_{\ell-2} \in \mathbb{Z}_{>0}$ with $m < \beta_{\ell-2} \le m+t$ and $e_{\alpha_{\ell-2}}e_{\alpha_{\ell-2}+1}\dots e_{\beta_{\ell-2}} \in M$ is $\beta_{\ell-2} = \alpha_{\ell-2}+1 = m+1$. In consequence, $\int_{\ell-2}(x_{j,1}^{v,\ell-2}, {}^{\ell}v_{1,r(i)}^{\ddagger}h) = (e_1e_2\dots e_{\beta_{\ell-2}}, e_{\beta_{\ell-2}+1}e_{\beta_{\ell-2}+2}\dots e_{m+t})$ is given by $(e_1e_2\dots e_{m+1}, e_{m+2}e_{m+3}\dots e_{m+t})$. Evidently, $e_{m+2}e_{m+3}\dots e_{m+t}$ is equal to h. And (12) in Lemma 4.5 implies $e_1e_2\dots e_{m+1} = x_{j,i}^{v,\ell-1}$. Hence, the claim is true. □

4.3. Zeroth and first order. In addition to the assumptions from the previous two sections, for any $\ell \in \mathbb{Z}_{\geq 0}$ let $P_{\ell} = \mathbb{K}C_{\ell} \otimes A$ be the free right A-module over the set C_{ℓ} of ℓ -chains and \leq_{ℓ} its order, tips with respect to which are indicated by \diamond_{ℓ} , from Section 2.8 and let d_{ℓ} and ι_{ℓ} be the order- ℓ differential and splitting, respectively, of the Anick resolution from Section 2.9.

Since the set C_0 of 0-chains always has only one element, the unit 1 of $\mathbb{K}\langle E \rangle$, it is common to identify the right A-modules $\mathbb{K}C_0 \otimes A$ and $A = P_0$. This practice is adopted in the following. The differential of order 0 does not depend on E, J or \leq either. It is always given by the map d_0 from the Main theorem.

Proposition 4.11. $d_0(1 \otimes 1) = 1$.

The first-order differential of any Anick resolution is completely determined by the set E of generators and the augmentation ε . Taking into account the aforementioned identification it is precisely the mapping d_1 from Main theorem.

Proposition 4.12. For any $v \in V$ and $(j,i) \in [n]^{\times 2}$, $d_1(x_{i,i}^v \otimes 1) = v_{i,r(i)} - \delta_{j,r(i)} 1 + J.$ Hence, it is only the orders two and up which need computing. That is carried out over the next three sections.

Beware that, in order to increase legibility, for any $p \in \mathbb{K}\langle E \rangle$ the symbol p will from now on also be used for the element p + J of A.

4.4. Second order. Computing d_2 requires determining (some of) the values of the splitting ι_0 . Note that, taking into acccount the identification $\mathbb{K}C_0 \otimes A \cong A$, the order \leq_0 is simply \leq and the combinatorial splitting \int_0 becomes the mapping $\langle E \rangle \backslash (\{1\} \cup \diamond J) \to C_1 \times \langle E \rangle$ which for any $m \in \mathbb{Z}_{>0}$ and any $e \in E^{\times m}$ with $e_1 e_2 \ldots e_m \in \langle E \rangle \backslash \diamond J$ satisfies $\int_0 (e_1 e_2 \ldots e_m) = (e_1, e_2 \ldots e_m).$

For computing d_2 it is convenient to distinguish two cases.

4.4.1. Generic case. First, $d_2(x_{j,i}^v \otimes 1)$ is computed for any $v \in V$ and any $(j,i) \in [n]^{\times 2}$ with $(j,i) \neq (n,n)$. That takes n+1 steps. For any $\{s,t\} \subseteq \mathbb{Z}$ let $\zeta_{s,t} \in \mathbb{K}$ be 1 if $s \leq t$ and 0 otherwise.

Lemma 4.13. For any $m \in [n]$ and any $(j,i) \in [n]^{\times 2} \setminus \{(n,n)\},\$

$$d_{2}(x_{j,i}^{v} \otimes 1) = \sum_{s=1}^{m} x_{j,s}^{v} \otimes v_{s,r(i)}^{\ddagger}$$

$$+ \iota_{0} \Big(\sum_{s=m+1}^{n} v_{j,r(s)} v_{s,r(i)}^{\ddagger} + \zeta_{r(j),m} v_{r(j),r(i)}^{\ddagger} - \delta_{j,i} 1 \Big).$$
(14)

Proof. The claim is proved by induction over m. It is convenient, though, to first note that for any $k \in [n]$ by Proposition 4.12

$$-d_{1}(x_{j,k}^{v} \otimes v_{k,\mathbf{r}(i)}^{\dagger}) = -d_{1}(x_{j,k}^{v} \otimes 1)v_{k,\mathbf{r}(i)}^{\dagger}$$

$$= -(v_{j,\mathbf{r}(k)} - \delta_{j,\mathbf{r}(k)}1)v_{k,\mathbf{r}(i)}^{\dagger}$$

$$= -v_{j,\mathbf{r}(k)}v_{k,\mathbf{r}(i)}^{\dagger} + \delta_{\mathbf{r}(j),k}v_{\mathbf{r}(j),\mathbf{r}(i)}^{\dagger}.$$
 (15)

Induction base. For m = 1 the claim is that

$$d_{2}(x_{j,i}^{v} \otimes 1) = x_{j,1}^{v} \otimes v_{1,\mathbf{r}(i)}^{\ddagger} + \iota_{0} \left(\sum_{s=2}^{n} v_{j,\mathbf{r}(s)} v_{s,\mathbf{r}(i)}^{\ddagger} + \delta_{\mathbf{r}(j),1} v_{\mathbf{r}(j),\mathbf{r}(i)}^{\ddagger} - \delta_{j,i} 1 \right)$$
(16)

because $\zeta_{\mathbf{r}(j),1} = \delta_{\mathbf{r}(j),1}$ by $1 \leq \mathbf{r}(j)$.

Because by Proposition 4.8

$$\partial_2(x_{j,i}^{v,2}) = (x_{j,1}^{v,1}, v_{1,\mathbf{r}(i)}^{\ddagger})$$

the definition of d_2 implies

$$d_2(x_{j,i}^v \otimes 1) = x_{j,1}^v \otimes v_{1,\mathbf{r}(i)}^{\ddagger} + \iota_0(-d_1(x_{j,1}^v \otimes v_{1,\mathbf{r}(i)}^{\ddagger})).$$
(17)

Hence, it is enough to show that the vector (17) is the same as the argument of ι_0 on the right-hand side of (16).

And, indeed, specializing k to 1 in (15) gives

$$-d_1(x_{j,1}^v \otimes v_{1,\mathbf{r}(i)}^{\ddagger}) = -v_{j,n}v_{1,\mathbf{r}(i)}^{\ddagger} + \delta_{\mathbf{r}(j),1}v_{\mathbf{r}(j),\mathbf{r}(i)}^{\ddagger}.$$
 (18)

Given that $-v_{j,n}v_{1,\mathbf{r}(i)}^{\ddagger} = \sum_{s=2}^{n} v_{j,\mathbf{r}(s)}v_{s,\mathbf{r}(i)}^{\ddagger} - \delta_{j,i}\mathbf{1}$, this thus proves the claim for m = 1.

Induction step. Let (14) be true for some m with $m \leq n-1$. Then it also holds for m + 1 in place of m for the following reasons. Because $(j,i) \neq (n,n)$ and $1 \neq m+1$ the vector $v_{j,\mathbf{r}(m+1)}v_{m+1,\mathbf{r}(i)}^{\ddagger}$ is normal by Lemma 4.2. It is the tip of the argument of ι_0 in (14). Indeed, any other term in the same grouped sum is less because already $v_{j,\mathbf{r}(s)} < v_{j,\mathbf{r}(m+1)}$ for any $s \in [n]$ with m + 1 < s, i.e., r(s) < r(m + 1), by the addendum to Lemma 3.4. And all remaining terms have lower degree.

Because by definition

$$\int_0 (v_{j,\mathbf{r}(m+1)} v_{m+1,\mathbf{r}(i)}^{\ddagger}) = (x_{j,m+1}^{v,1}, v_{m+1,\mathbf{r}(i)}^{\ddagger})$$

the equality in (14) hence remains true if $x_{j,m+1}^v \otimes v_{m+1,r(i)}^{\ddagger}$ is added to the right-hand side of (14) outside of ι_0 and $-d_1(x_{i,m+1}^v \otimes v_{m+1,r(i)}^{\ddagger})$ to the inside.

That evidently changes the outside in the same way as replacing mby m + 1 would. At the same time, by (15) for k = m + 1,

$$-d_1(x_{j,m+1}^v \otimes v_{m+1,\mathbf{r}(i)}^{\dagger}) = -v_{j,\mathbf{r}(m+1)}v_{m+1,\mathbf{r}(i)}^{\dagger} + \delta_{\mathbf{r}(j),m+1}v_{\mathbf{r}(j),\mathbf{r}(i)}^{\dagger}, \quad (19)$$

Keeping in mind that $\zeta_{\mathbf{r}(j),m} + \delta_{\mathbf{r}(j),m+1} = \zeta_{\mathbf{r}(j),m+1}$ it is now clear that adding the vector in (19) to the argument of ι_0 in (14) has the same effect as replacing m by m + 1 there. Thus, the claim is true for all $m \in [n].$

Proposition 4.14. For any $v \in V$ and any $(j,i) \in [n]^{\times 2} \setminus \{(n,n)\},\$ $d_2(x_{j,i}^v \otimes 1) = \sum_{s=1}^n x_{j,s}^v \otimes v_{s,\mathbf{r}(i)}^{\ddagger} + x_{\mathbf{r}(j),i}^{v^{\ddagger}} \otimes 1.$ (20)

Proof. In the special m = n the identity (14) in Lemma 4.13 reads

$$d_2(x_{j,i}^v \otimes 1) = \sum_{s=1}^n x_{j,s}^v \otimes v_{s,\mathbf{r}(i)}^{\ddagger} + \iota_0(v_{\mathbf{r}(j),\mathbf{r}(i)}^{\ddagger} - \delta_{j,i} 1).$$
(21)

because $\zeta_{\mathbf{r}(j),n} = 1$. Since $v_{\mathbf{r}(j),\mathbf{r}(i)}^{\ddagger}$ is normal and greater than 1 it is the tip of the argument of ι_0 . Because it was defined that

$$\int_0 (v_{\mathbf{r}(j),\mathbf{r}(i)}^{\ddagger}) = (x_{\mathbf{r}(j),i}^{v^{\ddagger},1}, 1)$$

the veracity of (21) is hence not affected by adding on the right-hand side $x_{\mathbf{r}(j),i}^{v^{\dagger}} \otimes 1$ to the outside of ι_0 and $-d_1(x_{\mathbf{r}(j),i}^{v^{\dagger}} \otimes 1)$ to the inside. Since by Proposition 4.12

$$-d_1(x_{r(j),i}^{v^{\ddagger}} \otimes 1) = -v_{r(j),r(i)}^{\ddagger} + \delta_{j,i} 1$$

the outcome of that addition is precisely (20).

4.4.2. Special case. It remains to compute $d_2(x_{n,n}^v \otimes 1)$ for any $v \in V$. Actually, since $x_{n,n}^v = v_{n,n} v_{1,1}^{\dagger}$ is invariant under replacing v with v^t it is enough to consider $v \in \{u, u^{\star}\}$, which greatly simplifies the proof. This time exactly n^2 steps are necessary to solve the recursion.

24

Lemma 4.15. For any $v \in \{u, u^*\}$ and any $\{\ell, m\} \subseteq [n]$ with $2 \leq \ell$,

$$d_{2}(x_{n,n}^{v} \otimes 1) = x_{n,1}^{v} \otimes v_{1,1}^{\ddagger} - \sum_{s=2}^{n} \sum_{t=\ell}^{n-1} x_{t,s}^{v} \otimes v_{s,r(t)}^{\ddagger}$$

$$- \sum_{s=2}^{m} x_{\ell-1,s}^{v} \otimes v_{s,r(\ell-1)}^{\ddagger}$$

$$+ \iota_{0} \Big(- \sum_{s=m+1}^{n} v_{\ell-1,r(s)} v_{s,r(\ell-1)}^{\ddagger} - \sum_{s=2}^{n} \sum_{t=1}^{\ell-2} v_{t,r(s)} v_{s,r(t)}^{\ddagger}$$

$$- \sum_{s=\ell}^{n-1} v_{r(s),r(s)}^{\ddagger} - \zeta_{r(\ell-1),m} v_{r(\ell-1),r(\ell-1)}^{\ddagger} + v_{1,1}^{\ddagger}$$

$$+ (n-2) 1 \Big).$$

$$(22)$$

Proof. The claim is proved by nested induction, an outer one over $r(\ell)$ and an inner one over m.

Part 1: Outer induction base. As base case $\ell = n$ what needs to be shown is that for any $m \in [n]$,

$$d_{2}(x_{n,n}^{v} \otimes 1) = x_{n,1}^{v} \otimes v_{1,1}^{\ddagger} - \sum_{s=2}^{m} x_{n-1,s}^{v} \otimes v_{s,r(n-1)}^{\ddagger} + \iota_{0} \Big(-\sum_{s=m+1}^{n} v_{n-1,r(s)} v_{s,r(n-1)}^{\ddagger} - \sum_{s=2}^{n} \sum_{t=1}^{n-2} v_{t,r(s)} v_{s,r(t)}^{\ddagger} - \zeta_{r(n-1),m} v_{r(n-1),r(n-1)}^{\ddagger} + v_{1,1}^{\ddagger} + (n-2) 1 \Big).$$

$$(23)$$

Part 1.1: Inner induction base. In particular, as the start m = 1 it is claimed that

where the two grouped sums in the argument of ι_0 appearing in (23) have been comined into one and $\zeta_{r(n-1),1} = 0$ has been used. And, indeed, because by Proposition 4.8

$$\partial_2(x_{n,n}^{v,2}) = (x_{n,1}^{v,1}, v_{1,1}^{\ddagger})$$

and because by Proposition 4.12

$$\begin{aligned} -d_1(x_{n,1}^v \otimes 1)v_{1,1}^{\dagger} &= -(v_{n,n} - 1)v_{1,1}^{\dagger} \\ &= -v_{n,n}v_{1,1}^{\dagger} + v_{1,1}^{\dagger} \\ &= -(\sum_{s=2}^n \sum_{t=1}^{n-1} v_{t,r(s)}v_{s,r(t)}^{\dagger} - (n-2)1) + v_{1,1}^{\dagger} \end{aligned}$$

that is what the definition

$$d_2(x_{n,n}^v \otimes 1) = x_{n,1}^v \otimes v_{1,1}^{\ddagger} + \iota_0(-d_1(x_{n,1}^v \otimes v_{1,1}^{\ddagger}))$$

implies.

Part 1.2: Inner induction step. Now, suppose (23) holds for some $m \in [n]$ with $m \leq n-1$. By the ensuing argument (23) remains true if m is replaced by m+1.

By Lemma 4.2 the vector $v_{n-1,r(m+1)}v_{m+1,r(n-1)}^{\dagger}$ is normal. It is moreover the tip of the argument of ι_0 (23) as explained hereafter. For degree reasons the only monomials which could possibly be greater or equal

appear in the third line of (23). More precisely, the other summands in this line are precisely the $v_{t,r(s)}v_{s,r(t)}^{\dagger}$ for $\{t,s\} \subseteq [n]$ with $s \neq 1$ such that either t < n - 1 or both t = n - 1 and m + 1 < s, which is to say with (t, r(s)) < (n - 1, r(m + 1)). Because $v \in \{u, u^*\}$ was assumed, Lemma 3.4 proves that already $v_{t,r(s)} < v_{n-1,r(m+1)}$ for those s and t, which then ensures $v_{t,r(s)}v_{s,r(t)}^{\dagger} < v_{n-1,r(m+1)}v_{m+1,r(n-1)}^{\dagger}$.

Because by definition

$$\int_0 (v_{n-1,\mathbf{r}(m+1)} v_{m+1,\mathbf{r}(n-1)}^{\ddagger}) = (x_{n-1,m+1}^{v,1}, v_{m+1,\mathbf{r}(n-1)}^{\ddagger})$$

the identity in (23) is hence preserved if $-x_{n-1,m+1}^v \otimes v_{m+1,r(n-1)}^{\dagger}$ is added outside of ι_0 and $d_1(x_{n-1,m+1}^v \otimes v_{m+1,r(n-1)}^{\dagger})$ inside on the right-hand side.

This addition evidently transforms the exterior of ι_0 in (23) in the same way as replacing m by m + 1 would. At the same time, adding

$$d_1(x_{n-1,m+1}^v \otimes v_{m+1,r(n-1)}^{\dagger}) = (v_{n-1,r(m+1)} - \delta_{n-1,r(m+1)} 1)v_{m+1,r(n-1)}^{\dagger} = v_{n-1,r(m+1)}v_{m+1,r(n-1)}^{\dagger} - \delta_{r(n-1),m+1}v_{r(n-1),r(n-1)}^{\dagger}.$$

to the argument of ι_0 in (23) has the same effect there because $\zeta_{r(n-1),m} + \delta_{r(n-1),m+1} = \zeta_{r(\ell-1),m+1}$. In conclusion, the base case for the outer induction has been established.

Part 2: Outer induction step. Next, suppose (22) holds for some $\ell \in [n]$ with $3 \leq \ell$ and all $m \in [n]$. It remains to show that (22) is true with $\ell - 1$ in place of ℓ , i.e., that for any $m \in [n]$,

$$d_{2}(x_{n,n}^{v} \otimes 1) = x_{n,1}^{v} \otimes v_{1,1}^{\dagger} - \sum_{s=2}^{n} \sum_{t=\ell-1}^{n-1} x_{t,s}^{v} \otimes v_{s,r(t)}^{\dagger}$$

$$- \sum_{s=2}^{m} x_{\ell-2,s}^{v} \otimes v_{s,r(\ell-2)}^{\dagger}$$

$$+ \iota_{0} \left(-\sum_{s=m+1}^{n} v_{\ell-2,r(s)} v_{s,r(\ell-2)}^{\dagger} - \sum_{s=2}^{n} \sum_{t=1}^{\ell-3} v_{t,r(s)} v_{s,r(t)}^{\dagger}$$

$$- \sum_{s=\ell-1}^{n-1} v_{r(s),r(s)}^{\dagger} - \zeta_{r(\ell-2),m} v_{r(\ell-2),r(\ell-2)}^{\dagger} + v_{1,1}^{\dagger}$$

$$+ (n-2)1 \right).$$

$$(24)$$

Part 2.1: Inner induction base. As the base case m = 1 this means proving that

$$d_{2}(x_{n,n}^{v} \otimes 1) = x_{n,1}^{v} \otimes v_{1,1}^{\ddagger} - \sum_{s=2}^{n} \sum_{t=\ell-1}^{n-1} x_{t,s}^{v} \otimes v_{s,r(t)}^{\ddagger} + \iota_{0} \Big(-\sum_{s=2}^{n} \sum_{t=1}^{\ell-2} v_{t,r(s)} v_{s,r(t)}^{\ddagger} - \sum_{s=\ell-1}^{n-1} v_{r(s),r(s)}^{\ddagger} + v_{1,1}^{\ddagger} + (n-2)1 \Big).$$

$$(25)$$

where, again, two grouped sums in the argument of ι_0 in (24) have been combined into one and where it has been used that $\ell \leq n$ implies $\ell - 2 \leq n - 2$ and thus $3 \leq r(\ell - 2)$, which is why $\zeta_{r(\ell-2),1} = 0$.

And that is precisely what the special case m = n of the induction hypothesis (22) says: The two grouped sums on the exterior of ι_0 can be combined into one. The first grouped sum in the fourth line vanishes. And because, of course, $\zeta_{r(\ell-1),n} = 1$ the terms $-\sum_{s=\ell}^{n-1} v_{r(s),r(s)}^{\ddagger} - \sum_{s=\ell}^{n-1} v_{r(s),r(s)}^{\ddagger}$ $\zeta_{\mathbf{r}(\ell-1),n} v_{\mathbf{r}(\ell-1),\mathbf{r}(\ell-1)}^{\ddagger} \text{ equal } -\sum_{s=\ell-1}^{n-1} v_{\mathbf{r}(s),\mathbf{r}(s)}^{\ddagger}.$

Part 2.2: Inner induction step. Finally, let (24) be true for some $m \in [n]$ with m < n - 1. It follows the proof that (24) is still true if m is replaced by m+1.

The vector $v_{\ell-2,r(m+1)}v_{m+1,r(\ell-2)}^{\ddagger}$ is normal by Lemma 4.2. It is the tip of the argument of ι_0 in (24). Indeed, already by degree it is greater than any monomial appearing in the last two lines of (24). The remaining summands inside ι_0 are exactly all $v_{t,\mathbf{r}(s)}v_{s,\mathbf{r}(t)}^{\ddagger}$ for $\{s,t\} \subseteq [n]$ with $s \neq 1$ such that either $t < \ell - 2$ or both $t = \ell - 2$ and m + 1 < s. Since $(t, \mathbf{r}(s)) < (\ell - 2, \mathbf{r}(m+1))$ and thus $v_{t,\mathbf{r}(s)} < v_{\ell-2,\mathbf{r}(m+1)}$ by Lemma 3.4 also $v_{t,\mathbf{r}(s)}v_{s,\mathbf{r}(t)}^{\ddagger} < v_{\ell-2,\mathbf{r}(m+1)}v_{m+1,\mathbf{r}(\ell-2)}^{\ddagger}$. Consequently, given that by definition

$$\int_0 (v_{\ell-2,\mathbf{r}(m+1)} v_{m+1,\mathbf{r}(\ell-2)}^{\ddagger}) = (x_{\ell-2,m+1}^{v,1}, v_{m+1,\mathbf{r}(\ell-2)}^{\ddagger}),$$

the equation resulting from (24) by adding on the right-hand side $-x_{\ell-2,m+1}^v \otimes v_{m+1,\mathbf{r}(\ell-2)}^{\ddagger}$ outside ι_0 and $d_1(-x_{\ell-2,m+1}^v \otimes v_{m+1,\mathbf{r}(\ell-2)}^{\ddagger})$ inside is true as well.

Said addition effectively increases m by 1 in (24) outside of ι_0 . Simultaneously, by Proposition 4.12

$$d_{1}(x_{\ell-2,m+1}^{v} \otimes v_{m+1,r(\ell-2)}^{\ddagger}) = (v_{\ell-2,r(m+1)} - \delta_{\ell-2,r(m+1)} 1) v_{m+1,r(\ell-2)}^{\ddagger} = v_{\ell-2,r(m+1)} v_{m+1,r(\ell-2)}^{\ddagger} - \delta_{r(\ell-2),m+1} v_{r(\ell-2),r(\ell-2)}^{\ddagger}$$
(26)

and $\zeta_{r(\ell-2),m} + \delta_{r(m+1),\ell-2} = \zeta_{r(\ell-2),m+1}$. An inspection of the argument of ι_0 in (24) therefore proves that the addition of the term (26) has the same effect as replacing m by m+1 there. Thus, (24) is still true for m+1 instead of m, which concludes the proof of (22) for arbitrary ℓ and m.

Lemma 4.16. For any $v \in \{u, u^*\}$ and any $m \in [n]$ with $m \leq n - 1$,

$$d_{2}(x_{n,n}^{v} \otimes 1) = x_{n,1}^{v} \otimes v_{1,1}^{\ddagger} - \sum_{s=2}^{n} \sum_{t=1}^{n-1} x_{t,s}^{v} \otimes v_{s,r(t)}^{\ddagger} - \sum_{s=1}^{m} x_{r(s),s}^{v^{\ddagger}} \otimes 1 \qquad (27) + \iota_{0} \Big(-\sum_{s=m+1}^{n-1} v_{r(s),r(s)}^{\ddagger} + v_{1,1}^{\ddagger} + r(m+3) 1 \Big).$$

Proof. Once more the proof goes by induction over m.

Induction base. For m = 1 what needs to be shown is that

$$d_{2}(x_{n,n}^{v} \otimes 1) = x_{n,1}^{v} \otimes v_{1,1}^{\ddagger} - \sum_{s=2}^{n} \sum_{t=1}^{n-1} x_{t,s}^{v} \otimes v_{s,r(t)}^{\ddagger} - x_{n,1}^{v^{\ddagger}} \otimes 1 + \iota_{0} \left(-\sum_{s=2}^{n-1} v_{r(s),r(s)}^{\ddagger} + v_{1,1}^{\ddagger} + (n-3) 1 \right).$$
(28)

In the special case $\ell = 2$ and m = n the grouped sum in the third line of (22) in Lemma 4.15 can be merged with that in the second line. Moreover, all the arguments of ι_0 in the fourth line vanish. Since, moreover, $\zeta_{r(1),n} = 1$ the resulting identity then reads

$$d_2(x_{n,n}^v \otimes 1) = x_{n,1}^v \otimes v_{1,1}^{\ddagger} - \sum_{s=2}^n \sum_{t=1}^{n-1} x_{t,s}^v \otimes v_{s,r(t)}^{\ddagger} + \iota_0 \Big(-\sum_{s=1}^{n-1} v_{r(s),r(s)}^{\ddagger} + v_{1,1}^{\ddagger} + (n-2) \, 1 \Big).$$
(29)

Obviously, $v_{n,n}^{\ddagger}$ is normal and the tip in the argument of ι_0 in (29) by Lemma 3.4. Because by definition

$$\int_0 (v_{n,n}^{\ddagger}) = (x_{n,1}^{v^{\ddagger},1}, 1)$$

the veracity of (29) is thus not affected by adding $-x_{n,1}^{v^{\ddagger}} \otimes 1$ outside of ι_0 and $d_1(x_{n,1}^{v^{\ddagger}} \otimes 1)$ inside.

Since, according to Proposition 4.12,

$$d_1(x_{n,1}^{v^\ddagger} \otimes 1) = v_{n,n}^\ddagger - 1$$

this change turns (29) into (28), providing the induction base.

Induction step. Let $m \in [n]$ be such that $m \leq n-2$ and such that (27) is true. The entire argument of ι_0 in (27) is evidently normal and by Lemma 3.4 has tip $v_{r(m+1),r(m+1)}^{\ddagger}$. Because

$$\int_0 (v_{\mathbf{r}(m+1),\mathbf{r}(m+1)}^{\ddagger}) = (x_{\mathbf{r}(m+1),m+1}^{v^{\ddagger},1},1)$$

a further valid identity is produced by adding to (27) on the right-hand side $-x_{r(m+1),m+1}^{v^{\ddagger}} \otimes 1$ outside of ι_0 and $d_1(x_{r(m+1),m+1}^{v^{\ddagger}} \otimes 1)$ inside. Since Proposition 4.12 implies

$$d_1(x_{\mathbf{r}(m+1),m+1}^{v^{\ddagger}}\otimes 1) = v_{\mathbf{r}(m+1),\mathbf{r}(m+1)}^{\ddagger} - 1$$

the identity resulting from the additions is just (27) with m replaced by m+1. \square

Proposition 4.17. For any $v \in V$,

$$d_2(x_{n,n}^v \otimes 1) = x_{n,1}^v \otimes v_{1,1}^{\ddagger} - \sum_{s=2}^n \sum_{t=1}^{n-1} x_{t,s}^v \otimes v_{s,r(t)}^{\ddagger}$$
(30)
$$- \sum_{s=2}^n x_{s,r(s)}^{v^{\ddagger}} \otimes 1 + x_{1,n}^{v^{\ddagger}} \otimes 1.$$

Proof. Since Lemma 4.3 shows that (30) is invariant under exchanging the roles $v \leftrightarrow v^{t}$ it is sufficient to prove it for any $v \in \{u, u^{\star}\}$.

Instantiating (27) from Lemma 4.16 at m = n - 1 yields the identity

$$d_{2}(x_{n,n}^{v} \otimes 1) = x_{n,1}^{v} \otimes v_{1,1}^{\ddagger} - \sum_{s=2}^{n} \sum_{t=1}^{n-1} x_{t,s}^{v} \otimes v_{s,r(t)}^{\ddagger} - \sum_{s=1}^{n-1} x_{r(s),s}^{v^{\ddagger}} \otimes 1 \qquad (31)$$
$$+ \iota_{0}(v_{1,1}^{\ddagger} - 1)$$

since r(n+2) = n - (n+2) + 1 = -1. Because the tip of the argument of ι_0 in (31) is $v_{1,1}^{\ddagger}$ and because

$$\int_0 (v_{1,1}^{\ddagger}) = (x_{1,n}^{v^{\ddagger}}, 1)$$

adding on the right-hand side of (31) the vector $x_{1,n}^{v^{\ddagger}} \otimes 1$ to the exterior of ι_0 while at the same time subtracting

$$d_2(x_{1,n}^{v^{\ddagger}} \otimes 1) = v_{1,1}^{\ddagger} - 1$$

from the interior gives another true identity. As that procedure transforms (31) into (30) this proves the claim.

4.4.3. Synthesis. Finally, it will be useful at certain points to have a formula for d_2 that covers both the generic and the special case simultaneously.

Theorem 4.18. For any $v \in V$ and any $(j, i) \in [n]^{\times 2}$,

$$d_{2}(x_{j,i}^{v} \otimes 1) = \sum_{s=1}^{n} x_{j,s}^{v} \otimes v_{s,r(i)}^{\ddagger} + x_{r(j),i}^{v^{\ddagger}} \otimes 1$$

$$- \delta_{j,n} \delta_{i,n} \sum_{s=2}^{n} (\sum_{t=1}^{n} x_{t,s}^{v} \otimes v_{s,r(t)}^{\ddagger} + x_{s,r(s)}^{v^{\ddagger}} \otimes 1).$$
(32)

Proof. It is clear by Proposition 4.14 that (32) is valid if $(j, i) \neq (n, n)$. In the case (j, i) = (n, n) the sum of all degree-two terms in (32) amounts to

$$x_{n,1}^{v} \otimes v_{1,r(n)}^{\ddagger} + \sum_{s=2}^{n} x_{n,s}^{v} \otimes v_{s,r(n)}^{\ddagger} - \sum_{s=2}^{n} \sum_{t=1}^{n} x_{t,s}^{v} \otimes v_{s,r(t)}^{\ddagger}$$

As the grouped sum in the middle is canceled by the one on the right it is now evident that (32) also agrees with (30) in Proposition 4.17. \Box

4.5. Third order. While the formula for d_3 is very similar to that for d_ℓ for odd $\ell \in \mathbb{Z}_{>0}$ with $3 < \ell$, the fact that d_2 is so different from d_ℓ for even $\ell \in \mathbb{Z}_{>0}$ with $2 < \ell$, makes it necessary to treat d_3 separately.

4.5.1. Case-independent identities. It is convenient to treat all basis vectors simultaneously up to a certain point and then distinguish cases for the remainder of the proof. There will be n recursion steps before the case distinction.

Lemma 4.19. For any $v \in V$, any $(j,i) \in [n]^{\times 2}$ and any $m \in [n]$,

$$d_{3}(x_{j,i}^{v} \otimes 1) = \sum_{s=1}^{m} x_{j,s}^{v} \otimes v_{s,r(i)}$$

$$+ \iota_{1} \Big(\sum_{t=m+1}^{n} x_{j,1}^{v} \otimes v_{1,r(t)}^{\ddagger} v_{t,r(i)} - \sum_{s=2}^{n} \sum_{t=1}^{m} x_{j,s}^{v} \otimes v_{s,r(t)}^{\ddagger} v_{t,r(i)}$$

$$- \sum_{s=1}^{m} x_{r(j),s}^{v^{\ddagger}} \otimes v_{s,r(i)} - \delta_{i,1} x_{j,i}^{v} \otimes 1$$

$$+ \zeta_{n,m} \delta_{j,n} \Big(\sum_{s=2}^{n} \sum_{t=1}^{n} x_{t,s}^{v} \otimes v_{s,r(t)}^{\ddagger} v_{n,r(i)}$$

$$+ \sum_{s=2}^{n} x_{s,r(s)}^{v^{\ddagger}} \otimes v_{n,r(i)} \Big) \Big).$$

$$(33)$$

Proof. The assertion will be proved by induction over m. It is economic, though, to first note that, if using the shorthand

$$q_{j} \coloneqq \delta_{j,n} (\sum_{s=2}^{n} \sum_{t=1}^{n} x_{t,s}^{v} \otimes v_{s,r(t)}^{\ddagger} + \sum_{s=2}^{n} x_{s,r(s)}^{v^{\ddagger}} \otimes 1),$$

Theorem 4.18 implies for any $k \in [n]$,

$$-d_{2}(x_{j,k}^{v} \otimes v_{k,\mathbf{r}(i)}) = -(\sum_{s=1}^{n} x_{j,s}^{v} \otimes v_{s,\mathbf{r}(k)}^{\dagger} + x_{\mathbf{r}(j),k}^{v^{\dagger}} \otimes 1 - \delta_{n,k} q_{j})v_{k,\mathbf{r}(i)}$$
$$= -x_{j,1}^{v} \otimes v_{1,\mathbf{r}(k)}^{\dagger}v_{k,\mathbf{r}(i)} - \sum_{s=2}^{n} x_{j,s}^{v} \otimes v_{s,\mathbf{r}(k)}^{\dagger}v_{k,\mathbf{r}(i)} \quad (34)$$
$$-x_{\mathbf{r}(j),k}^{v^{\dagger}} \otimes v_{k,\mathbf{r}(i)} + \delta_{n,k} q_{j}v_{k,\mathbf{r}(i)}.$$

Induction base. Because $2 \leq n$ and thus $\zeta_{n,1} = 0$ what is claimed in the special case m = 1 is that

$$d_{3}(x_{j,i}^{v} \otimes 1) = x_{j,1}^{v} \otimes v_{1,\mathbf{r}(i)}$$

$$+ \iota_{1} \Big(\sum_{t=2}^{n} x_{j,1}^{v} \otimes v_{1,\mathbf{r}(t)}^{\dagger} v_{t,\mathbf{r}(i)} - \sum_{s=2}^{n} x_{j,s}^{v} \otimes v_{s,\mathbf{r}(1)}^{\dagger} v_{1,\mathbf{r}(i)}$$

$$- x_{\mathbf{r}(j),1}^{v^{\dagger}} \otimes v_{1,\mathbf{r}(i)} - \delta_{i,1} x_{j,i}^{v} \otimes 1 \Big).$$
(35)

Since by Proposition 4.8

$$\partial_3(x_{j,i}^{v,3}) = (x_{j,1}^{v,2}, v_{1,\mathbf{r}(i)})$$

the definition of d_3 implies

$$d_3(x_{j,i}^v \otimes 1) = x_{j,1}^v \otimes v_{1,\mathbf{r}(i)} + \iota_1(-d_2(x_{j,1}^v \otimes v_{1,\mathbf{r}(i)})).$$
(36)

Thus it is enough to prove that the arguments of ι_1 on the respective right-hand sides of (35) and (36) are equal.

Reducing $v_{1,r(1)}^{\ddagger}v_{1,r(i)}$ in (34) for k = 1 yields

$$-d_{2}(x_{j,1}^{v} \otimes v_{1,\mathbf{r}(i)}) = -x_{j,1}^{v} \otimes (-\sum_{t=2}^{n} v_{1,\mathbf{r}(t)}^{\dagger} v_{t,\mathbf{r}(i)} + \delta_{i,1} 1)$$

$$-\sum_{s=2}^{n} x_{j,s}^{v} \otimes v_{s,\mathbf{r}(1)}^{\dagger} v_{1,\mathbf{r}(i)} - x_{\mathbf{r}(j),1}^{v^{\dagger}} \otimes v_{1,\mathbf{r}(i)},$$

$$(37)$$

which proves the claimed agreement of the two arguments of ι_0 . Hence, (33) is true for m = 1.

Induction step. Let (33) hold for some $m \in [n]$ with $m \leq n - 1$. As shown hereafter, (33) remains true if m is replaced by m + 1.

The values of v, j and i be as they may, $x_{j,1}^{v} \otimes v_{1,r(m+1)}^{\ddagger} v_{m+1,r(i)}$ is normal by Lemma 4.2. It is the tip of the argument of ι_1 in the induction hypothesis (33) for the following reasons. First of all, by $\zeta_{n,m} = 0$ the

last two lines of (33) are zero. Furthermore, any vector in the fourth line is less already by degree. Hence, the only vectors against which to compare are in the third line of (33). For any $s \in [n]$ with 1 < s the fact that (j, 1) < (j, s) implies $x_{j,s}^v < x_{j,1}^v$ by Lemma 4.4 and thus $x_{j,s}^v \otimes v_{s,r(t)}^{\ddagger} v_{t,r(i)} <_1 x_{j,1}^v \otimes v_{1,r(m+1)}^{\ddagger} v_{m+1,r(i)}$ for arbitrary $t \in [n]$. Similarly, for any $t \in [n]$ with m + 1 < t the addendum to Lemma 3.4 proves $v_{1,r(t)}^{\ddagger} < v_{1,r(m+1)}^{\ddagger}$ and thus $x_{j,1}^v \otimes v_{1,r(t)}^{\ddagger} v_{t,r(i)} <_1 x_{j,1}^v \otimes v_{1,r(m+1)}^{\ddagger} v_{m+1,r(i)}$. In consequence, since by Proposition 4.10

$$\int_{1} (x_{j,1}^{v,1}, v_{1,\mathbf{r}(m+1)}^{\ddagger} v_{m+1,\mathbf{r}(i)}) = (x_{j,m+1}^{2,v}, v_{m+1,\mathbf{r}(i)})$$

the veracity of (33) is not affected by adding on the right-hand side $x_{j,m+1}^v \otimes v_{m+1,\mathbf{r}(i)}$ outside of ι_1 and $-d_2(x_{j,m+1}^v \otimes v_{m+1,\mathbf{r}(i)})$ inside. Clearly, on the exterior of ι_1 this addition has the same effect as

Clearly, on the exterior of ι_1 this addition has the same effect as replacing m by m + 1. Simultaneously, specializing (34) to k = m + 1yields

$$-d_{2}(x_{j,m+1}^{v} \otimes v_{m+1,\mathbf{r}(i)}) = -x_{j,1}^{v} \otimes v_{1,\mathbf{r}(m+1)}^{\dagger} v_{m+1,\mathbf{r}(i)} - \sum_{s=2}^{n} x_{j,s}^{v} \otimes v_{s,\mathbf{r}(m+1)}^{\dagger} v_{m+1,\mathbf{r}(i)} - x_{\mathbf{r}(j),m+1}^{v\dagger} \otimes v_{m+1,\mathbf{r}(i)} + \delta_{n,m+1} q_{j} v_{m+1,\mathbf{r}(i)}.$$
(38)

Because $\zeta_{n,m} + \delta_{n,m+1} = \delta_{n,m+1}$ it is now evident that the addition of the vector (38) changes the argument of ι_1 in (33) in the same way as increasing m by 1 would. Hence, the assertion is true for all m.

Lemma 4.20. For any $v \in V$ and any $(j, i) \in [n]^{\times 2}$,

$$d_{3}(x_{j,i}^{v} \otimes 1) = \sum_{s=1}^{n} x_{j,s}^{v} \otimes v_{s,r(i)}$$

$$+ \iota_{1} \Big(-\sum_{s=1}^{n} x_{r(j),s}^{v^{\dagger}} \otimes v_{s,r(i)} - x_{j,i}^{v} \otimes 1 \\ + \delta_{j,n} \Big(\sum_{s=2}^{n} \sum_{t=1}^{n} x_{t,s}^{v} \otimes v_{s,r(t)}^{\dagger} v_{n,r(i)} + \sum_{s=2}^{n} x_{s,r(s)}^{v^{\dagger}} \otimes v_{n,r(i)} \Big) \Big).$$
(39)

Proof. In the special case m = n the first grouped sum in the third line of (33) in Lemma 4.19 vanishes and what remains is the identity

$$d_{3}(x_{j,i}^{v} \otimes 1) = \sum_{s=1}^{n} x_{j,s}^{v} \otimes v_{s,r(i)}$$

$$+ \iota_{1} \left(-\sum_{s=2}^{n} x_{j,s}^{v} \otimes \sum_{t=1}^{n} v_{s,r(t)}^{\dagger} v_{t,r(i)} - \sum_{s=1}^{n} x_{r(j),s}^{v^{\dagger}} \otimes v_{s,r(i)} - \delta_{i,1} x_{j,i}^{v} \otimes 1 + \delta_{j,n} \left(\sum_{s=2}^{n} \sum_{t=1}^{n} x_{t,s}^{v} \otimes v_{s,r(t)}^{\dagger} v_{n,r(i)} + \sum_{s=2}^{n} x_{s,r(s)}^{v^{\dagger}} \otimes v_{n,r(i)} \right) \right).$$
(40)

And because

$$-\sum_{s=2}^{n} x_{j,s}^{v} \otimes \sum_{t=1}^{n} v_{s,r(t)}^{\ddagger} v_{t,r(i)} - \delta_{i,1} x_{j,i}^{v} \otimes 1$$

$$= -\sum_{s=2}^{n} \delta_{s,i} x_{j,s}^{v} \otimes 1 - \delta_{i,1} x_{j,1}^{v} \otimes 1 = -\sum_{s=1}^{n} \delta_{s,i} x_{j,s}^{v} \otimes 1 = -x_{j,i}^{v} \otimes 1$$

$$\text{that already proves (39).} \qquad \Box$$

From now on it makes more sense to distinguish cases, ultimately four of them.

4.5.2. *Towards the first two cases.* For the first two out of the four cases that need to be considered there is a common starting point, obtained via one recursion step.

Lemma 4.21. For any $v \in V$ and any $(j,i) \in [n]^{\times 2}$, if $j \neq n$, then $d_3(x_{j,i}^v \otimes 1) = \sum_{s=1}^n x_{j,s}^v \otimes v_{s,r(i)} - x_{r(j),i}^{v^{\ddagger}} \otimes 1 \qquad (41)$ $+ \iota_1 \left(-\sum_{s=1}^n x_{r(j),s}^{v^{\ddagger}} \otimes v_{s,r(i)} - x_{j,i}^v \otimes 1 + d_2(x_{r(j),i}^{v^{\ddagger}} \otimes 1)) \right).$

Proof. What the identity (39) in Lemma 4.20 says in the special case $j \neq n$ is that

$$d_{3}(x_{j,i}^{v} \otimes 1) = \sum_{s=1}^{n} x_{j,s}^{v} \otimes v_{s,r(i)} + \iota_{1}(-\sum_{s=1}^{n} x_{r(j),s}^{v^{\ddagger}} \otimes v_{s,r(i)} - x_{j,i}^{v} \otimes 1).$$
(42)

The argument of ι_1 in this identity is obviously normal and its tip is $x_{r(j),1}^{v^{\dagger}} \otimes v_{1,r(i)}$ because already $x_{r(j),s}^{v^{\dagger}} < x_{r(j),1}^{v^{\dagger}}$ for any $s \in [n]$ with 1 < s by Lemma 4.4.

Because by Proposition 4.10

$$\int_{1} (x_{\mathbf{r}(j),1}^{v^{\ddagger},1}, v_{1,\mathbf{r}(i)}) = (x_{\mathbf{r}(j),i}^{v^{\ddagger},2}, 1)$$

another valid identity is produced by adding to (42) on the right-hand side the term $-x_{r(j),i}^{v^{\ddagger}} \otimes 1$ on the exterior of ι_1 and $d_2(x_{r(j),i}^{v^{\ddagger}} \otimes 1)$. The resulting equation is precisely (41).

4.5.3. *First case.* The first case is very simple, requiring no further recursions.

Proposition 4.22. For any $v \in V$ and any $(j,i) \in [n]^{\times 2}$, if $j \neq n$ and $(j,i) \neq (1,n)$, then

$$d_3(x_{j,i}^v \otimes 1) = \sum_{s=1}^n x_{j,s}^v \otimes v_{s,r(i)} - x_{r(j),i}^{v^{\ddagger}} \otimes 1.$$
(43)

Proof. By Proposition 4.14 the assumption that $(j, i) \neq (1, n)$ implies

$$d_2(x_{\mathbf{r}(j),i}^{v^{\ddagger}} \otimes 1) = \sum_{s=1}^n x_{\mathbf{r}(j),s}^{v^{\ddagger}} \otimes v_{s,\mathbf{r}(i)} + x_{j,i}^v \otimes 1,$$

which is precisely the negative of the remainder of the argument of ι_1 in (41). Hence, Lemma 4.21 proves the assertion.

4.5.4. Second case. The second case is still handled quickly, in n steps.

Lemma 4.23. For any $v \in V$ and any $m \in [n]$,

$$d_{3}(x_{1,n}^{v} \otimes 1) = \sum_{s=1}^{n} x_{1,s}^{v} \otimes v_{s,1} - x_{n,n}^{v^{\ddagger}} \otimes 1 - \sum_{s=2}^{m} x_{r(s),r(s)}^{v^{\star}} \otimes 1 + \iota_{1} \left(-\sum_{s=1}^{n} \sum_{t=m+1}^{n} x_{r(t),s}^{v^{\star}} \otimes v_{s,t}^{t} - \sum_{s=m+1}^{n} x_{s,r(s)}^{v^{t}} \otimes 1 \right).$$

$$(44)$$

Proof. The proof goes by induction over m.

Induction base. For the base case m = 1 what needs to be shown is that

$$d_{3}(x_{1,n}^{v} \otimes 1) = \sum_{t=1}^{n} x_{1,s}^{v} \otimes v_{s,1} - x_{n,n}^{v^{\dagger}} \otimes 1 \qquad (45)$$
$$+ \iota_{1} \Big(-\sum_{s=1}^{n} \sum_{t=2}^{n} x_{r(t),s}^{v^{\star}} \otimes v_{s,t}^{t} - \sum_{s=2}^{n} x_{s,r(s)}^{v^{t}} \otimes 1 \Big).$$

Since by Proposition 4.17

$$d_2(x_{n,n}^{v^{\ddagger}} \otimes 1) = x_{n,1}^{v^{\ddagger}} \otimes v_{1,1} - \sum_{s=2}^n \sum_{t=1}^{n-1} x_{t,s}^{v^{\ddagger}} \otimes v_{s,r(t)} - \sum_{s=2}^n x_{s,r(s)}^v \otimes 1 + x_{1,n}^v \otimes 1$$

the conclusion (41) of Lemma 4.21 in the special case (j, i) = (1, n)reads

$$d_{3}(x_{1,n}^{v} \otimes 1) = \sum_{s=1}^{n} x_{1,s}^{v} \otimes v_{s,1} - x_{n,n}^{v^{\ddagger}} \otimes 1$$

$$+ \iota_{1} \Big(-\sum_{s=2}^{n} \sum_{t=1}^{n} x_{t,s}^{v^{\ddagger}} \otimes v_{s,r(t)} - \sum_{s=2}^{n} x_{s,r(s)}^{v} \otimes 1 \Big).$$
(46)

where two instances each of $x_{1,n}^v \otimes 1$ and $x_{n,1}^{v^{\ddagger}} \otimes v_{1,1}$ with different signs have canceled each other out and where two grouped sums have been combined into one. Hence, it is enough to rewrite the argument of ι_1 in (46) until it agrees with that in (45).

Reindexing the summation over t with r and afterwards exchanging the names $s \leftrightarrow t$ of the summation indices makes the argument of ι_1 in (46) take on the form

$$-\sum_{s=1}^{n}\sum_{t=2}^{n}x_{\mathbf{r}(s),t}^{v^{\ddagger}}\otimes v_{t,s}-\sum_{s=2}^{n}x_{s,\mathbf{r}(s)}^{v}\otimes 1$$

Recognizing that for any $\{s,t\} \subseteq [n]$ not only $v_{t,s} = v_{s,t}^{t}$ but by Lemma 4.3 also

$$x_{\mathbf{r}(s),t}^{v^{\ddagger},1} = x_{\mathbf{r}(t),s}^{v^{\star},1}$$

as well as

$$x_{s,r(s)}^{v,1} = x_{s,r(s)}^{v^{t},1}$$

now shows the claimed equality between the arguments in (46) and (45), thus proving (44) for m = 1.

Induction step. Let (44) hold for some $m \in [n]$ with $m \leq n-1$. By

the following argument (44) is also true for m + 1 in place of m. The normal vector $x_{r(m+1),1}^{v^*} \otimes v_{1,m+1}^t$ is the tip of the argument of ι_1 in (44) as explained hereafter. For reasons of degree the only terms which could possibly be greater or equal are of the form $x_{r(t),s}^{v^{\star}} \otimes v_{s,t}^{t}$ for $\{s,t\} \subseteq [n]$ such that either m+1 < t or m+1 = t and 1 < s, which is to say such that (m+1,1) < (t,s). But those conditions are actually equivalent to (1, m+1) < (s, t). Since thus both $(\mathbf{r}(t), 1) < (\mathbf{r}(m+1), s)$ and $(1, \mathbf{r}(t)) < (s, \mathbf{r}(m+1))$ for such t and s Lemma 4.4 proves that then $x_{\mathbf{r}(t),s}^{v^{\star}} < x_{\mathbf{r}(m+1),1}^{v^{\star}}$, irrespective of the value of v.

Because by Proposition 4.10

$$\int_{1} (x_{\mathbf{r}(m+1),1}^{v^{\star},1}, v_{1,m+1}^{\mathbf{t}}) = (x_{\mathbf{r}(m+1),\mathbf{r}(m+1)}^{v^{\star},2}, 1)$$

the definition of ι_1 implies that a valid identity is derived by adding to the right-hand side of (44) the term $-x_{r(m+1),r(m+1)}^{v^*} \otimes 1$ outside of ι_1 and $d_2(x_{r(m+1),r(m+1)}^{v^*} \otimes 1)$ inside of it.

Evidently, the outside of this new identity is the same as that of (44) for m + 1 instead of m. Likewise, the addition of, by Proposition 4.14,

$$d_2(x_{r(m+1),r(m+1)}^{v^*} \otimes 1) = \sum_{s=1}^n x_{r(m+1),s}^{v^*} \otimes v_{s,m+1}^{t} + x_{m+1,r(m+1)}^{v^{t}} \otimes 1.$$

modifies the argument of ι_1 in (44) in the same way as replacing m by m+1 would. That proves (44) to hold for arbitrary m.

Proposition 4.24. For any $v \in V$,

$$d_3(x_{1,n}^v \otimes 1) = \sum_{s=1}^n x_{1,s}^v \otimes v_{s,1} - x_{n,n}^{v^{\ddagger}} \otimes 1 - \sum_{s=1}^{n-1} x_{s,s}^{v^{\star}} \otimes 1.$$
(47)

Proof. The claim (47) is precisely (44) in Lemma 4.23 for m = n with $\sum_{s=2}^{n} x_{r(s),r(s)}^{v^*} \otimes 1$ reindexed.

4.5.5. Towards the last two cases. Just like for the first two cases there is a common starting point for the last two, arrived at by n-1 steps.

Lemma 4.25. For any $v \in V$, any $i \in [n]$ and any $m \in [n]$ with $m \leq n-1$,

$$\begin{aligned} d_{3}(x_{n,i}^{v} \otimes 1) \\ &= \sum_{s=1}^{n} x_{n,s}^{v} \otimes v_{s,r(i)} + \sum_{s=2}^{m} x_{r(s),r(s)}^{v^{t}} \otimes v_{r(i),n}^{t} \qquad (48) \\ &+ \iota_{1} \Big(\sum_{s=m+1}^{n} x_{r(s),1}^{v^{t}} \otimes v_{1,s}^{t} v_{r(i),n}^{t} + \sum_{s=m+1}^{n} \sum_{t=2}^{n} x_{r(s),t}^{v^{t}} \otimes v_{t,s}^{t} v_{r(i),n}^{t} \\ &+ \sum_{s=m+1}^{n} x_{s,r(s)}^{v^{\star}} \otimes v_{r(i),n}^{t} - \sum_{s=1}^{n} x_{r(s),n}^{v^{\star}} \otimes v_{r(i),s}^{t} - x_{r(i),1}^{v^{t}} \otimes 1 \Big). \end{aligned}$$

Proof. The proof is by induction over m.

Induction base. For the base case m = 1 what needs to be proved is that

$$d_{3}(x_{n,i}^{v} \otimes 1) = \sum_{s=1}^{n} x_{n,s}^{v} \otimes v_{s,r(i)}$$

$$+ \iota_{1} \Big(\sum_{s=2}^{n} \sum_{t=1}^{n} x_{r(s),t}^{v^{t}} \otimes v_{t,s}^{\star} v_{r(i),n}^{t} \\ + \sum_{s=2}^{n} x_{s,r(s)}^{v^{\star}} \otimes v_{r(i),n}^{t} - \sum_{s=1}^{n} x_{r(s),n}^{v^{\star}} \otimes v_{r(i),s}^{t} - x_{r(i),1}^{v^{t}} \otimes 1 \Big).$$
(49)

But (49) is precisely what (39) in Lemma 4.20 says in the special case j = n after the summation over t is reindexed with r and for any $\{s,t\} \subseteq [n]$ with 1 < s the identities

$$x_{\mathbf{r}(t),s}^{v,1} \otimes v_{s,t}^{\ddagger} v_{n,\mathbf{r}(i)} = x_{\mathbf{r}(s),t}^{v^{\mathsf{t}},1} \otimes v_{t,s}^{\star} v_{\mathbf{r}(i),n}^{\mathsf{t}}$$

and

$$x_{s,\mathbf{r}(s)}^{v^{\ddagger},1} \otimes v_{n,\mathbf{r}(i)} = x_{s,\mathbf{r}(s)}^{v^{\star},1} \otimes v_{\mathbf{r}(i),n}^{\mathbf{t}}$$

and

$$x_{\mathbf{r}(n),s}^{v^{\star},1} \otimes v_{s,\mathbf{r}(i)} = x_{\mathbf{r}(s),n}^{v^{\star},1} \otimes v_{\mathbf{r}(i),s}^{\mathbf{t}}$$

and

$$x_{n,i}^{v,1} = x_{\mathbf{r}(i),1}^{v^{\mathrm{t}},1},$$

each partly implied by Lemma 4.3, are taken into account.

Induction step. Let (48) be valid for some $m \in [n]$ with $m \leq n-2$. By the ensuing argument it holds for m + 1 in place of m.

Regardless of the value of *i*, the vector $x_{r(m+1),1}^{v_{t}^{t}} \otimes v_{1,m+1}^{\star} v_{r(i),n}^{t}$ is normal by Lemma 4.2 since $m + 1 \neq n$. It is moreover the tip of the argument of ι_1 in (48) for the following reasons. Any vector in the fourth line of (48) is dominated already by degree. All the remaining other vectors in the argument are of the form $x_{\mathbf{r}(s),t}^{v^{t}} \otimes v_{t,s}^{\star} v_{\mathbf{r}(i),n}^{t}$ for $\{s,t\} \subseteq [n]$ with $m+1 \leq s$ such that either $2 \leq t$ or both t=1 and m+1 < s. The latter condition can also be stated as (1, m+1) < (t, s), which is actually equivalent to (m + 1, 1) < (s, t). Because thus both $(1, \mathbf{r}(s)) < (t, \mathbf{r}(m+1))$ and $(\mathbf{r}(s), 1) < (\mathbf{r}(m+1), t)$ Lemma 4.4 proves $x_{\mathbf{r}(s),t}^{v^{t}} < x_{\mathbf{r}(m+1),1}^{v^{t}}$, which makes $x_{\mathbf{r}(m+1),1}^{v^{t}} \otimes v_{1,m+1}^{\star} v_{\mathbf{r}(i),n}^{t}$ the tip. Because moreover by Proposition 4.10

$$\int_{1} (x_{\mathbf{r}(m+1),1}^{\mathbf{v}^{t},1}, v_{1,m+1}^{\star} v_{\mathbf{r}(i),n}^{\mathbf{t}}) = (x_{\mathbf{r}(m+1),\mathbf{r}(m+1)}^{\mathbf{v}^{t},2}, v_{\mathbf{r}(i),n}^{\mathbf{t}})$$

the equality in (48) is preserved if $x_{r(m+1),r(m+1)}^{v^{t}} \otimes v_{r(i),n}^{t}$ is added on the right-hand side to the exterior of ι_1 and $-d_2(x_{r(m+1),r(m+1)}^{v^t} \otimes v_{r(i),n}^t)$ to the interior.

Evidently, outside of ι_1 this has the same effect on (48) as replacing m by m + 1 would. Because $(r(m + 1), r(m + 1)) \neq (n, n)$ by $1 \leq m$ Proposition 4.14 implies

$$-d_{2}(x_{r(m+1),r(m+1)}^{v^{v}} \otimes v_{r(i),n}^{t})) = -(\sum_{t=1}^{n} x_{r(m+1),t}^{v^{t}} \otimes v_{t,m+1}^{\star} + x_{m+1,r(m+1)}^{v^{\star}} \otimes 1)v_{r(i),n}^{t} \\ = -x_{r(m+1),1}^{v^{t}} \otimes v_{1,m+1}^{t}v_{r(i),n}^{t} - \sum_{t=2}^{n} x_{r(m+1),t}^{v^{t}} \otimes v_{t,m+1}^{t}v_{r(i),n}^{t} \tag{50}) \\ - x_{m+1,r(m+1)}^{v^{\star}} \otimes v_{r(i),n}^{t}.$$

Hence, a comparison with (48) shows that adding the term (50) to the argument of ι_1 in (48) is the same as increasing m by 1 there. That verifies the claim for any m.

Lemma 4.26. For any $v \in V$ and any $i \in [n]$,

$$d_{3}(x_{n,i}^{v} \otimes 1) = \sum_{s=1}^{n} x_{n,s}^{v} \otimes v_{s,r(i)} + \sum_{s=2}^{n-1} x_{s,s}^{v^{t}} \otimes v_{r(i),n}^{t}$$

$$+ \iota_{1}(x_{1,1}^{v^{t}} \otimes v_{1,n}^{t} v_{r(i),n}^{t} + \sum_{t=2}^{n} x_{1,t}^{v^{t}} \otimes v_{t,n}^{\star} v_{r(i),n}^{t}$$

$$+ x_{n,1}^{v^{\star}} \otimes v_{r(i),n}^{t} - \sum_{s=1}^{n} x_{1,s}^{v^{t}} \otimes v_{s,r(i)} - x_{n,i}^{v} \otimes 1).$$
(51)

Proof. Because $\sum_{s=2}^{n-1} x_{\mathbf{r}(s),\mathbf{r}(s)}^{v^{t}} \otimes v_{\mathbf{r}(i),n}^{t} = \sum_{s=2}^{n-1} x_{s,s}^{v^{t}} \otimes v_{\mathbf{r}(i),n}^{t}$ and because by Lemma 4.3

 $-\sum_{s=1}^{n} x_{r(s),n}^{v^{\star}} \otimes v_{r(i),s}^{t} - x_{r(i),1}^{v^{t}} \otimes 1 = -\sum_{s=1}^{n} x_{1,s}^{v^{\ddagger}} \otimes v_{s,r(i)} - x_{n,i}^{v} \otimes 1$

the claim is the conclusion (48) of Lemma 4.25 in the special case m = n - 1.

4.5.6. *Third case.* Again, one of the two remaining cases is quicker to deal with than the other, requiring only a single step.

Lemma 4.27. For any $v \in V$ and any $i \in [n]$, if $i \neq n$, then

$$d_{3}(x_{n,i}^{v} \otimes 1) = \sum_{s=1}^{n} x_{n,s}^{v} \otimes v_{s,r(i)} + \sum_{s=1}^{n-1} x_{s,s}^{v^{t}} \otimes v_{r(i),n}^{t} \qquad (52)$$
$$+ \iota_{1} \Big(-\sum_{s=1}^{n} x_{1,s}^{v^{\dagger}} \otimes v_{s,r(i)} - x_{n,i}^{v} \otimes 1 \Big).$$

Proof. Because $i \neq n$ the vector $x_{1,1}^{v^t} \otimes v_{1,n}^* v_{r(i),n}^t$ is normal by Lemma 4.2. It is the tip of the argument of ι_1 in (51) in Lemma 4.26 for the following reasons. Any term in the fourth line of (51) is dominated already by degree. And for any $t \in [n]$ with 1 < t, i.e., r(t) < n, the fact that both (1,1) < (1,t) and (1,1) < (t,1) demands $x_{1,t}^{v^t} < x_{1,1}^{v^t}$ by Lemma 4.4. Hence, $x_{1,1}^{v^t} \otimes v_{1,n}^* v_{r(i),n}^t$ subjugates also any other term in the third line of (51).

Given that by Proposition 4.10

$$\int_{1} (x_{1,1}^{v^{\mathrm{t},1}}, v_{1,n}^{\star} v_{\mathrm{r}(i),n}^{\mathrm{t}}) = (x_{1,1}^{v^{\mathrm{t},2}}, v_{\mathrm{r}(i),n}^{\mathrm{t}}),$$

adding $x_{1,1}^{v^{t}} \otimes v_{r(i),n}^{t}$ to the outside of ι_{1} in (51) and $-d_{2}(x_{1,1}^{v^{t}} \otimes v_{r(i),n}^{t})$ to the inside does not affect the equality.

This addition turns the outside of ι_1 in (51) into that of (52). Since Proposition 4.14 implies

$$-d_2(x_{1,1}^{v^{t}} \otimes v_{\mathbf{r}(i),n}^{t}) = -\sum_{t=1}^{n} x_{1,t}^{v^{t}} \otimes v_{t,n}^{\star} v_{\mathbf{r}(i),n}^{t} - x_{n,1}^{v^{\star}} \otimes v_{\mathbf{r}(i),n}^{t}$$
(53)

adding (53) the inside of ι_1 likewise makes the arguments of ι_1 in (51) and (52) agree. That proves the claim.

Proposition 4.28. For any $v \in V$ and any $i \in [n]$, if $i \neq n$, then

$$d_{3}(x_{n,i}^{v} \otimes 1) = \sum_{s=1}^{n} x_{n,s}^{v} \otimes v_{s,r(i)} + \sum_{s=1}^{n-1} x_{s,s}^{v^{t}} \otimes v_{r(i),n}^{t} - x_{1,i}^{v^{\dagger}} \otimes 1.$$
(54)

Proof. The argument of ι_1 on the right-hand side of (52) in Lemma 4.27 is obviously entirely normal. Its tip is $x_{1,1}^{v^{\dagger}} \otimes v_{1,\mathbf{r}(i)}$, which dominates $x_{n,i}^{v} \otimes 1$ by degree and which, according to Lemma 4.4, is greater than $x_{1,s}^{v^{\dagger}} \otimes v_{s,\mathbf{r}(i)}$ for any $s \in [n]$ with 1 < s because (1,1) < (1,s) and (1,1) < (s,1), and thus already $x_{1,s}^{v^{\dagger}} < x_{1,1}^{v^{\dagger}}$.

In consequence, since by Proposition 4.10

$$\int_{1} (x_{1,1}^{v^{\downarrow},1}, v_{1,\mathbf{r}(i)}) = (x_{1,i}^{v^{\downarrow},2}, 1)$$

the veracity of (52) is not affected by adding $-x_{1,i}^{v^{\dagger}} \otimes 1$ to the outside of ι_1 and $d_2(x_{1,i}^{v^{\dagger}} \otimes 1)$ to the inside. Because $d_2(x_{1,i}^{v^{\dagger}} \otimes 1)$ by Proposition 4.14 is precisely the negative of the argument of ι_1 in (52) that proves (54). \Box 4.5.7. Fourth case. It only remains to treat a single case, in n+2 steps. Lemma 4.29. For any $v \in V$ and any $m \in [n]$,

$$d_{3}(x_{n,n}^{v} \otimes 1) = \sum_{s=1}^{n} x_{n,s}^{v} \otimes v_{s,1} + \sum_{s=2}^{n-1} x_{s,s}^{v^{t}} \otimes v_{1,n}^{t} - \sum_{s=2}^{m} x_{1,s}^{v^{t}} \otimes v_{s,n}^{t} \qquad (55)$$

+ $\iota_{1} \left(-\sum_{s=m+1}^{n} x_{1,1}^{v^{t}} \otimes v_{1,r(s)}^{\star} v_{s,n}^{t} + \sum_{s=1}^{m} \sum_{t=2}^{n} x_{1,t}^{v^{t}} \otimes v_{t,r(s)}^{\star} v_{s,n}^{t} \right)$
+ $\sum_{s=1}^{m} x_{n,s}^{v^{\star}} \otimes v_{s,n}^{t} - \sum_{s=1}^{n} x_{1,s}^{v^{\dagger}} \otimes v_{s,1}^{t} \right).$

Proof. The claim is proved by induction over m.

Induction base. In the base case m = 1 the asserted identity (55) reads

$$d_{3}(x_{n,n}^{v} \otimes 1) = \sum_{s=1}^{n} x_{n,s}^{v} \otimes v_{s,1} + \sum_{s=2}^{n-1} x_{s,s}^{v^{t}} \otimes v_{1,n}^{t}$$
(56)
+ $\iota_{1} \Big(-\sum_{s=2}^{n} x_{1,1}^{v^{t}} \otimes v_{1,r(s)}^{\star} v_{s,n}^{t} + \sum_{t=2}^{n} x_{1,t}^{v^{t}} \otimes v_{t,n}^{\star} v_{1,n}^{t}$
+ $x_{n,1}^{v^{\star}} \otimes v_{1,n}^{t} - \sum_{s=1}^{n} x_{1,s}^{v^{\dagger}} \otimes v_{s,1} \Big).$

What (51) in Lemma 4.26 says in the special case i = n is that

$$d_{3}(x_{n,n}^{v} \otimes 1) = \sum_{s=1}^{n} x_{n,s}^{v} \otimes v_{s,1} + \sum_{s=2}^{n-1} x_{s,s}^{v^{t}} \otimes v_{1,n}^{t}$$

$$+ \iota_{1} \left(x_{1,1}^{v^{t}} \otimes v_{1,n}^{\star} v_{1,n}^{t} + \sum_{t=2}^{n} x_{1,t}^{v^{t}} \otimes v_{t,n}^{\star} v_{1,n}^{t} \right)$$

$$+ x_{n,1}^{v^{\star}} \otimes v_{1,n}^{t} - \sum_{s=1}^{n} x_{1,s}^{v^{\dagger}} \otimes v_{s,1} - x_{n,n}^{v} \otimes 1 \right).$$
(57)

Therefore, it suffices to prove that the arguments of ι_1 in (56) and (57) are equal.

Because $x_{n,n}^v = x_{1,1}^{v^t}$ by Lemma 4.3 and thus

$$\begin{aligned} x_{1,1}^{v^{t}} \otimes v_{1,n}^{\star} v_{1,n}^{t} - x_{n,n}^{v} \otimes 1 &= x_{1,1}^{v^{t}} \otimes (-\sum_{s=2}^{n} v_{1,r(s)}^{\star} v_{s,n}^{t} + 1) - x_{1,1}^{v^{t}} \otimes 1 \\ &= -\sum_{s=2}^{n} x_{1,1}^{v^{t}} \otimes v_{1,r(s)}^{\star} v_{s,n}^{t} \end{aligned}$$

that is indeed the case. Hence, (56) is true.

Induction step. Let $m \in [n]$ be arbitrary with $m \leq n-1$ and with the property that (55) is satisfied. Then (55) still holds if m is replaced by m+1 by the ensuing argument.

Since $1 \leq m$ the monomial $v_{1,r(m+1)}^{\star}v_{m+1,n}^{t}$ is normal by Lemma 4.2 and the vector $x_{1,1}^{v^{t}} \otimes v_{1,r(m+1)}^{\star}v_{m+1,n}^{t}$ is the tip of the argument of ι_{1} in (55) for the following reasons. Any vector in the fourth line of (55) is less already by degree. The only terms against which to compare therefore live in the third line and are of the form $x_{1,t}^{v^{t}} \otimes v_{t,r(s)}^{\star}v_{s,n}^{t}$ for $\{s,t\} \subseteq [n]$ with either 1 < t or both t = 1 and m + 1 < s. If 1 < t, then the fact that both (1,1) < (1,t) and (1,1) < (t,1) implies $x_{1,t}^{v^{t}} < x_{1,1}^{v^{t}}$ by Lemma 4.4 and thus $x_{1,t}^{v^{t}} \otimes v_{t,r(s)}^{\star}v_{s,n}^{t} <_{1} x_{1,1}^{v^{t}} \otimes v_{1,r(m+1)}^{\star}v_{m+1,n}^{t}$. Alternatively, if 1 = t and m + 1 < s, which is to say, r(s) < r(m + 1), then it is

the addendum to Lemma 3.4 that guarantees $v_{1,\mathbf{r}(s)}^{\star} < v_{1,\mathbf{r}(m+1)}^{\star}$ and thus $x_{1,1}^{v^{t}} \otimes v_{1,\mathbf{r}(s)}^{\star} v_{s,n}^{t} <_{1} x_{1,1}^{v^{t}} \otimes v_{1,\mathbf{r}(m+1)}^{\star} v_{m+1,n}^{t}$ as well. Hence, and because by Proposition 4.10

$$\int_{1} (x_{1,1}^{v^{t},1}, v_{1,r(m+1)}^{\star} v_{m+1,n}^{t}) = (x_{1,m+1}^{v^{t},2}, v_{m+1,n}^{t})$$

adding to the right-hand side of (55) outside of ι_1 the term $-x_{1,m+1}^{v^t} \otimes v_{m+1,n}^t$ and inside the term $d_2(x_{1,m+1}^{v^t} \otimes v_{m+1,n}^t)$ gives another true identity.

Evidently, this has the same effect on the outside of ι_1 in (55) as replacing m by m + 1 would have. Because by Proposition 4.14

$$d_{2}(x_{1,m+1}^{v^{t}} \otimes v_{m+1,n}^{t}) = \sum_{t=1}^{n} x_{1,t}^{v^{t}} \otimes v_{t,r(m+1)}^{\star} v_{m+1,n}^{t} + x_{n,m+1}^{v^{\star}} \otimes v_{m+1,n}^{t}$$
(58)

the addition of (58) likewise changes the argument of ι_1 in (55) in the same way as increasing m by 1 would. Hence, the claim is true for all m.

Lemma 4.30. For any $v \in V$,

$$d_{3}(x_{n,n}^{v} \otimes 1) = \sum_{s=1}^{n} x_{n,s}^{v} \otimes v_{s,1} + \sum_{s=2}^{n-1} x_{s,s}^{v^{t}} \otimes v_{1,n}^{t}$$

$$-\sum_{s=2}^{n} x_{1,s}^{v^{t}} \otimes v_{s,n}^{t} + x_{n,1}^{v^{\star}} \otimes 1$$

$$+ \iota_{1} \left(-\sum_{s=1}^{n} x_{1,s}^{v^{\dagger}} \otimes v_{s,1} - x_{n,n}^{v} \otimes 1 \right).$$
(59)

Proof. In the special case m = n the first grouped sum in the third line of (55) in Lemma 4.29 vanishes and, taking into account the fact that $\sum_{s=1}^{n} v_{t,r(s)}^{\star} v_{s,n}^{t} = \delta_{t,1} 1$ implies

$$\sum_{s=1}^{n} \sum_{t=2}^{n} x_{1,t}^{v^{t}} \otimes v_{t,r(s)}^{\star} v_{s,n}^{t} = 0,$$

what remains of (55) is the identity

$$\begin{aligned} d_3(x_{n,n}^v \otimes 1) \\ &= \sum_{s=1}^n x_{n,s}^v \otimes v_{s,1} + \sum_{s=2}^{n-1} x_{s,s}^{v^{\rm t}} \otimes v_{1,n}^{\rm t} - \sum_{s=2}^n x_{1,s}^{v^{\rm t}} \otimes v_{s,n}^{\rm t} \\ &+ \iota_1 \Big(\sum_{s=1}^n x_{n,s}^{v^{\star}} \otimes v_{s,n}^{\rm t} - \sum_{s=1}^n x_{1,s}^{v^{\dagger}} \otimes v_{s,1} \Big). \end{aligned}$$
(60)

The tip of the argument of ι_1 in (60) is $x_{n,1}^{v^*} \otimes v_{1,n}^{t}$. Indeed, on the one hand, for any $s \in [n]$ with 1 < s already $x_{n,s}^{v^*} < x_{n,1}^{v^*}$ by the addendum to Lemma 4.4, and thus $x_{n,s}^{v^*} \otimes v_{s,n}^{t} <_1 x_{n,1}^{v^*} \otimes v_{1,n}^{t}$. On the other hand, for any $s \in [n]$, even s = 1 by $2 \leq n$, the facts that both (1,1) = $(\mathbf{r}(n), 1) < (s, \mathbf{r}(1)) = (s, n)$ and $(1, 1) = (1, \mathbf{r}(n)) < (\mathbf{r}(1), s) = (n, s)$ imply $x_{1,s}^{v^*} < x_{n,1}^{v^*}$ by Lemma 4.4 and consequently $x_{1,s}^{v^*} \otimes v_{s,1} <_1 x_{n,1}^{v^*} \otimes v_{1,n}^{t}$.

Therefore and since by Proposition 4.10

$$\int_{1} (x_{n,1}^{v^{\star},1}, v_{1,n}^{t}) = (x_{n,1}^{v^{\star},2}, 1)$$

adding on the right-hand side of (60) the term $x_{n,1}^{v^*} \otimes 1$ to the exterior of ι_1 and $-d_2(x_{n,1}^{v^*} \otimes 1)$ to the interior preserves the equality.

Of course, this change is precisely what is necessary to turn the outside of ι_1 in (60) into that of (59). At the same time, since by Proposition 4.14

$$-d_2(x_{n,1}^{v^*} \otimes 1) = -\sum_{s=1}^n x_{n,s}^{v^*} \otimes v_{s,n}^{t} - x_{1,1}^{v^t} \otimes 1$$
(61)

and since $-x_{1,1}^{v^{t}} \otimes 1 = -x_{n,n}^{v} \otimes 1$ by Lemma 4.3 the addition of the vector (61) likewise transforms the argument of ι_{1} in (60) into that of (59), which proves the latter identity.

Proposition 4.31. For any $v \in V$,

$$d_{3}(x_{n,n}^{v} \otimes 1) = \sum_{s=1}^{n} x_{n,s}^{v} \otimes v_{s,1} + \sum_{s=2}^{n-1} x_{s,s}^{v^{t}} \otimes v_{1,n}^{t} - \sum_{s=2}^{n} x_{1,s}^{v^{t}} \otimes v_{s,n}^{t} \qquad (62) + x_{n,1}^{v^{\star}} \otimes 1 - x_{1,n}^{v^{\dagger}} \otimes 1.$$

Proof. The tip of the argument of ι_1 in (59) in Lemma 4.30 is given by $x_{1,1}^{v^{\ddagger}} \otimes v_{1,1}$ because already $x_{1,s}^{v^{\ddagger}} < x_{1,1}^{v^{\ddagger}}$ for any $s \in [n]$ with 1 < s by the addendum to Lemma 4.4 and thus $x_{1,s}^{v^{\ddagger}} \otimes v_{s,1} <_1 x_{1,1}^{v^{\ddagger}} \otimes v_{1,1}$ and because $x_{n,n}^{v} \otimes 1 <_1 x_{1,1}^{v^{\ddagger}} \otimes v_{1,1}$ for degree reasons.

In consequence, since by Proposition 4.10

$$\int_{1} (x_{1,1}^{v^{\ddagger},1}, v_{1,1}) = (x_{1,n}^{v^{\ddagger},2}, 1)$$

the validity of (59) is not affected by adding $-x_{1,n}^{v^{\ddagger}} \otimes 1$ to the outside of ι_1 on the right-hand side and $d_2(x_{1,n}^{v^{\ddagger}} \otimes 1)$ to the inside.

Obviously that makes the new outside of ι_1 in (59) agree with that in (62). And since by Proposition 4.14 the vector

$$d_2(x_{1,n}^{v^{\ddagger}} \otimes 1) = \sum_{s=1}^n x_{1,s}^{v^{\ddagger}} \otimes v_{s,1} + x_{n,n}^v \otimes 1$$

is precisely the negative of the argument of ι_1 in (59) that verifies (62) and thus concludes the proof.

4.5.8. *Synthesis.* The individual results obtained in the four cases distinguished combine into the aggregated statement from the Main result.

Theorem 4.32. For any $v \in V$ and any $(j, i) \in [n]^{\times 2}$,

$$d_{3}(x_{j,i}^{v} \otimes 1) = \sum_{s=1}^{n} x_{j,s}^{v} \otimes v_{s,r(i)} - x_{r(j),i}^{v^{\dagger}} \otimes 1 \qquad (63)$$

+ $\delta_{j,n} (\sum_{s=1}^{n-1} x_{s,s}^{v^{\dagger}} \otimes v_{r(i),n}^{t} - \delta_{i,n} (\sum_{s=1}^{n} x_{1,s}^{v^{\dagger}} \otimes v_{s,n}^{t} - x_{n,1}^{v^{\star}} \otimes 1))$
- $\delta_{j,1} \delta_{i,n} \sum_{s=1}^{n-1} x_{s,s}^{v^{\star}} \otimes 1.$

Proof. By the vanishing of the last two lines, in the case $j \neq n$ and $(j, i) \neq (1, n)$ the formula (63) collapses into (43) from Proposition 4.22.

If, still $j \neq n$ but (j, i) = (1, n), while the third line of (63) is zero, the fourth becomes $-\sum_{s=1}^{n-1} x_{s,s}^{v^*} \otimes 1$, which is why in this case (63) is precisely (47) from Proposition 4.24.

In the instance that $j = n \neq i$ the two last lines of (63) amount to nothing more than $\sum_{s=1}^{n-1} x_{s,s}^{v^{t}} \otimes v_{r(i),n}^{t}$, which proves (63) identical to (54) from Proposition 4.28 in this case.

Finally, for (j, i) = (n, n) the sum of all degree-2 terms in (63) is

$$\sum_{s=1}^{n} x_{n,s}^{v} \otimes v_{s,1} + \sum_{s=1}^{n-1} x_{s,s}^{v^{t}} \otimes v_{1,n}^{t} - \sum_{s=1}^{n} x_{1,s}^{v^{t}} \otimes v_{s,n}^{t}.$$
 (64)

Because the term $x_{1,1}^{v^{t}} \otimes v_{1,n}^{t}$ appears in both the middle and right grouped sum in (64), but with different signs, (64) actually agrees with the second line of (62) in Proposition 4.31. Now it is evident that (62) is true in all cases.

4.6. Fourth and higher orders. The fourth and higher orders of the resolution can be computed in a uniform manner.

Assumption 4.33. Let $\ell \in \mathbb{Z}_{\geq 0}$ be such that $4 \leq \ell$ and for any $v \in V$ and $(j, i) \in [n]^{\times 2}$,

$$\begin{aligned} d_{\ell-1}(x_{j,i}^{v} \otimes 1) &= \sum_{s=1}^{n} x_{j,s}^{v} \otimes^{\ell-1} v_{s,r(i)} + (-1)^{\ell-1} x_{r(j),i}^{v^{\dagger}} \otimes 1 \\ &+ \delta_{j,n}((x_{1,1}^{v^{\dagger}} + \delta_{\ell-1,3} \sum_{s=2}^{n-1} x_{s,s}^{v^{\dagger}}) \otimes^{\ell-1} v_{r(i),n}^{t} \\ &- \delta_{i,n}(\sum_{s=1}^{n} x_{1,s}^{v^{\dagger}} \otimes^{\ell-1} v_{s,n}^{t} + (-1)^{\ell-1} x_{n,1}^{v^{\star}} \otimes 1)) \\ &+ \delta_{j,1} \delta_{i,n}(-1)^{\ell-1} (x_{1,1}^{v^{\star}} + \delta_{\ell-1,3} \sum_{s=2}^{n-1} x_{s,s}^{v^{\star}}) \otimes 1. \end{aligned}$$

Note that, if $\ell = 4$, this assumption holds by Theorem 4.32.

4.6.1. Case-independent identities. It is convenient to carry out n steps of the recursion for all basis vectors simultaneously and distinguish cases afterwards.

Lemma 4.34. For any $v \in V$, any $(j,i) \in [n]^{\times 2}$ and any $m \in [n]$,

$$d_{\ell}(x_{j,i}^{v} \otimes 1) = \sum_{s=1}^{m} x_{j,s}^{v} \otimes^{\ell} v_{s,r(i)}$$

$$+ \iota_{\ell-2} \Big(\sum_{s=m+1}^{n} x_{j,1}^{v} \otimes^{\ell} v_{1,r(s)}^{\dagger} {}^{\ell} v_{s,r(i)}$$

$$- \sum_{s=2}^{n} \sum_{t=1}^{m} x_{j,s}^{v} \otimes^{\ell} v_{s,r(t)}^{\dagger} {}^{\ell} v_{t,r(i)}$$

$$+ (-1)^{\ell} \sum_{s=1}^{m} x_{r(j),s}^{v^{\dagger}} \otimes^{\ell} v_{s,r(i)} - \delta_{i,1} x_{j,1}^{v} \otimes 1$$

$$- \delta_{j,n} (x_{1,1}^{v^{\dagger}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\dagger}}) \otimes \sum_{s=1}^{m} {}^{\ell} v_{r(s),n}^{\star} {}^{\ell} v_{r(i),s}$$

$$+ \zeta_{n,m} (\delta_{j,n} (\sum_{s=1}^{n} x_{1,s}^{v^{\dagger}} \otimes^{\ell} v_{s,n}^{\star} {}^{\ell} v_{r(i),n}^{t} - (-1)^{\ell} x_{n,1}^{v^{\star}} \otimes^{\ell} v_{r(i),n}^{t})$$

$$+ \delta_{j,1} (-1)^{\ell} (x_{1,1}^{v^{\star}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\star}}) \otimes^{\ell} v_{r(i),n}^{t}) \Big).$$
(65)

Proof. The proof goes by induction over m. It is economic though to first note that for any $k \in [n]$ renaming the i in Assumption 4.33 to k and multiplying the resulting identity with $-\ell v_{k,r(i)}$ from the right

yields

$$-d_{\ell-1}(x_{j,k}^{v} \otimes^{\ell} v_{k,r(i)}) = -x_{j,1}^{v} \otimes^{\ell} v_{1,r(k)}^{\dagger} {}^{\ell} v_{k,r(i)} - \sum_{s=2}^{n} x_{j,s}^{v} \otimes^{\ell} v_{s,r(k)}^{\dagger} {}^{\ell} v_{k,r(i)}$$

$$+ (-1)^{\ell} x_{r(j),k}^{v^{\dagger}} \otimes^{\ell} v_{k,r(i)}$$

$$+ \delta_{j,n}(-(x_{1,1}^{v^{t}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{t}}) \otimes^{\ell} v_{r(k),n}^{\star} {}^{\ell} v_{r(i),k}$$

$$+ \delta_{k,n}(\sum_{s=1}^{n} x_{1,s}^{v^{t}} \otimes^{\ell} v_{s,n}^{\star} {}^{\ell} v_{r(i),n}^{t} - (-1)^{\ell} x_{n,1}^{v^{\star}} \otimes^{\ell} v_{r(i),n}^{t}))$$

$$+ \delta_{j,1} \delta_{k,n}(-1)^{\ell} (x_{1,1}^{v^{\star}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\star}}) \otimes^{\ell} v_{r(i),n}^{t},$$

$$(66)$$

where, also, the s = 1 term has been splitt off from the first grouped sum on the right-hand side.

Induction base. Since $2 \leq n$ and thus $\zeta_{n,1} = 0$, in the base case m = 1 what needs to be proved is that

$$d_{\ell}(x_{j,i}^{v} \otimes 1) = x_{j,1}^{v} \otimes^{\ell} v_{1,\mathbf{r}(i)}$$

$$+ \iota_{\ell-2} \Big(\sum_{s=2}^{n} x_{j,1}^{v} \otimes^{\ell} v_{1,\mathbf{r}(s)}^{\dagger} {}^{\ell} v_{s,\mathbf{r}(i)} - \sum_{s=2}^{n} x_{j,s}^{v} \otimes^{\ell} v_{s,\mathbf{r}(1)}^{\dagger} {}^{\ell} v_{1,\mathbf{r}(i)}$$

$$+ (-1)^{\ell} x_{\mathbf{r}(j),1}^{v^{\dagger}} \otimes^{\ell} v_{1,\mathbf{r}(i)} - \delta_{i,1} x_{j,1}^{v} \otimes 1$$

$$- \delta_{j,n} (x_{1,1}^{v^{\dagger}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\dagger}}) \otimes^{\ell} v_{\mathbf{r}(1),n}^{\star} {}^{\ell} v_{\mathbf{r}(i),1}^{t} \Big).$$

$$(67)$$

Given that by Proposition 4.8

$$\partial_{\ell}(x_{j,i}^{v,\ell}) = (x_{j,1}^{v,\ell-1}, {}^{\ell}v_{1,\mathbf{r}(i)})$$

the definition of d_{ℓ} means in this case

$$d_{\ell}(x_{j,i}^{v} \otimes 1) = x_{j,1}^{v} \otimes^{\ell} v_{1,\mathbf{r}(i)} + \iota_{\ell-2}(-d_{\ell-1}(x_{j,1}^{v} \otimes^{\ell} v_{1,\mathbf{r}(i)})).$$

Hence, it suffices to prove that $-d_{\ell-1}(x_{j,1}^v \otimes {}^{\ell}v_{1,\mathbf{r}(i)})$ agrees with the argument of $\iota_{\ell-2}$ in (67). By $2 \leq n$ the last two lines of (66) do not survive instantiation at k = 1, which thus yields

$$-d_{\ell-1}(x_{j,1}^{v} \otimes^{\ell} v_{1,\mathbf{r}(i)}) = -x_{j,1}^{v} \otimes^{\ell} v_{1,\mathbf{r}(1)}^{\ddagger} v_{1,\mathbf{r}(i)} - \sum_{s=2}^{n} x_{j,s}^{v} \otimes^{\ell} v_{s,\mathbf{r}(1)}^{\ddagger} v_{1,\mathbf{r}(i)} + (-1)^{\ell} x_{\mathbf{r}(j),1}^{v\dagger} \otimes^{\ell} v_{1,\mathbf{r}(i)} - \delta_{j,n}(x_{1,1}^{v^{t}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{t}}) \otimes^{\ell} v_{\mathbf{r}(1),n}^{\star} v_{\mathbf{r}(i),1}^{t}.$$

Since ${}^{\ell}v_{1,r(1)}^{\dagger}{}^{\ell}v_{1,r(i)} = -\sum_{s=2}^{n}{}^{\ell}v_{1,r(s)}^{\dagger}{}^{\ell}v_{s,r(i)} + \delta_{i,1} 1$ it is now visible that $-d_{\ell-1}(x_{j,1}^{v} \otimes v_{1,r(i)}^{\dagger})$ is precisely the argument of $\iota_{\ell-2}$ in (67). Hence, the claim holds for m = 1.

Induction step. Now, suppose $m \in [n]$ is such that $m \leq n-1$ and such that (65) is true. It follows the proof that (65) still holds if m is replaced by m + 1.

Since $1 \leq m$ the monomial ${}^{\ell}v_{1,r(m+1)}^{\dagger}{}^{\ell}v_{m+1,r(i)}$ is normal by Lemma 4.2 and the vector $x_{j,1}^{v} \otimes {}^{\ell}v_{1,r(m+1)}^{\dagger}{}^{\ell}v_{m+1,r(i)}$ is the tip of the argument of $\iota_{\ell-2}$ in (65) for the reasons given hereafter. It dominates the other terms

in the grouped sum where it originates because ${}^{\ell}v_{1,r(s)}^{\ddagger} < {}^{\ell}v_{1,r(m+1)}^{\ddagger}$ for any $s \in [n]$ with m + 1 < s by the addendum to Lemma 3.4. It is also greater than any of the terms in the grouped sum in the fourth line of (65) because $x_{j,t}^v < x_{j,1}^v$ for any $t \in [n]$ with 1 < t by the addendum to Lemma 4.4. Any term in the fifth line of (65) is less already for degree reasons. Also, any term in the sixth line is dominated because, if it is non-zero, then j = n and hence $x_{s,s}^{v^{\dagger}} < x_{j,n}^v$ for any $s \in [n]$ with $s \neq n$ by Lemma 4.4.

Because by Proposition 4.10

$$\int_{\ell-2} (x_{j,1}^{v,\ell-2}, {^\ell v_{1,\mathbf{r}(m+1)}^{\ddagger}}^{\ell} v_{m+1,\mathbf{r}(i)}) = (x_{j,m+1}^{v,\ell-1}, {^\ell v_{m+1,\mathbf{r}(i)}})$$

the veracity of (65) is not affected by adding on the right-hand side $x_{j,m+1}^v \otimes {}^{\ell}v_{m+1,\mathbf{r}(i)}$ outside of $\iota_{\ell-2}$ and $-d_{\ell-1}(x_{j,m+1}^v \otimes {}^{\ell}v_{m+1,\mathbf{r}(i)})$ inside. It is clear that this has the same effect on the outside of $\iota_{\ell-2}$ in (65)

It is clear that this has the same effect on the outside of $\iota_{\ell-2}$ in (65) as replacing m by m + 1 would have. Simultaneously, specializing the k in (66) to m + 1 and separating the fifth line from the fourth and combining the fifth and sixth lines shows that

$$-d_{\ell-1}(x_{j,m+1}^{v} \otimes^{\ell} v_{m+1,r(i)}) = -x_{j,1}^{v} \otimes^{\ell} v_{1,r(m+1)}^{\dagger} v_{m+1,r(i)} - \sum_{s=2}^{n} x_{j,s}^{v} \otimes^{\ell} v_{s,r(m+1)}^{\dagger} v_{m+1,r(i)} \quad (68)$$
$$+ (-1)^{\ell} x_{r(j),m+1}^{v^{\dagger}} \otimes^{\ell} v_{m+1,r(i)}$$
$$- \delta_{j,n}(x_{1,1}^{v^{\dagger}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\dagger}}) \otimes^{\ell} v_{r(m+1),n}^{\star} v_{r(i),m+1}$$
$$+ \delta_{m+1,n}(\delta_{j,n}(\sum_{s=1}^{n} x_{1,s}^{v^{\dagger}} \otimes^{\ell} v_{s,n}^{\star} v_{r(i),n}^{t} - (-1)^{\ell} x_{n,1}^{v^{\star}} \otimes^{\ell} v_{r(i),n}^{t})$$
$$+ \delta_{j,1}(-1)^{\ell}(x_{1,1}^{v^{\star}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\star}}) \otimes^{\ell} v_{r(i),n}^{t}).$$

Keeping in mind that $\zeta_{n,m} + \delta_{n,m+1} = \zeta_{n,m+1}$, a comparison now proves that adding the term (68) to the argument of $\iota_{\ell-2}$ in (65) has the same effect as replacing m by m+1 there.

Lemma 4.35. For any $v \in V$ and any $(j, i) \in [n]^{\times 2}$,

$$d_{\ell}(x_{j,i}^{v} \otimes 1) = \sum_{s=1}^{n} x_{j,s}^{v} \otimes^{\ell} v_{s,r(i)}$$

$$+ \iota_{\ell-2} ((-1)^{\ell} \sum_{s=1}^{n} x_{r(j),s}^{v^{\dagger}} \otimes^{\ell} v_{s,r(i)} - x_{j,i}^{v} \otimes 1$$

$$+ \delta_{j,n} (\sum_{s=1}^{n} x_{1,s}^{v^{\dagger}} \otimes^{\ell} v_{s,n}^{\star} \ell_{r(i),n}^{t} - (-1)^{\ell} x_{n,1}^{v^{\star}} \otimes^{\ell} v_{r(i),n}^{t}$$

$$- \delta_{i,n} (x_{1,1}^{v^{\dagger}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\dagger}}) \otimes 1)$$

$$+ \delta_{j,1} (-1)^{\ell} (x_{1,1}^{v^{\star}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\star}}) \otimes^{\ell} v_{r(i),n}^{t}).$$

$$(69)$$

Proof. The special case m = n of (65) in Lemma 4.34 amounts to the identity

$$d_{\ell}(x_{j,i}^{v} \otimes 1) = \sum_{s=1}^{n} x_{j,s}^{v} \otimes {}^{\ell} v_{s,r(i)}$$

$$+ \iota_{\ell-2} \Big(-\sum_{s=2}^{n} x_{j,s}^{v} \otimes \sum_{t=1}^{n} {}^{\ell} v_{s,r(t)}^{\dagger} {}^{\ell} v_{t,r(i)}$$

$$+ (-1)^{\ell} \sum_{s=1}^{n} x_{r(j),s}^{v\dagger} \otimes {}^{\ell} v_{s,r(i)} - \delta_{i,1} x_{j,1}^{v} \otimes 1$$

$$+ \delta_{j,n} (-(x_{1,1}^{v^{t}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{t}}) \otimes \sum_{s=1}^{n} {}^{\ell} v_{r(s),n}^{\star} {}^{\ell} v_{r(i),s}$$

$$+ \sum_{s=1}^{n} x_{1,s}^{v^{t}} \otimes {}^{\ell} v_{s,n}^{\star} v_{r(i),n}^{t} - (-1)^{\ell} x_{n,1}^{v^{\star}} \otimes {}^{\ell} v_{r(i),n}^{t})$$

$$+ \delta_{j,1} (-1)^{\ell} (x_{1,1}^{v^{\star}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\star}}) \otimes {}^{\ell} v_{r(i),n}^{t}).$$

$$(70)$$

From the fact that $\sum_{t=1}^{n} {}^{\ell} v_{s,\mathbf{r}(t)}^{\dagger} {}^{\ell} v_{t,\mathbf{r}(i)} = \delta_{s,i} 1$ for any $s \in [n]$ with $2 \leq s$ it follows that

$$-\sum_{s=2}^{n} x_{j,s}^{v} \otimes \sum_{t=1}^{n} {}^{\ell} v_{s,r(t)}^{\ddagger} {}^{\ell} v_{t,r(i)} - \delta_{i,1} x_{j,1}^{v} \otimes 1$$

= $-\sum_{s=2}^{n} \delta_{s,i} x_{j,s}^{v} \otimes 1 - \delta_{1,i} x_{j,1}^{v} \otimes 1 = -\sum_{s=1}^{n} \delta_{s,i} x_{j,s}^{v} \otimes 1 = -x_{j,i}^{v} \otimes 1.$

That proves that the third and fourth lines of (70) actually agree with the third line of (69).

Similarly, because $\sum_{s=1}^{n} {}^{\ell} v_{\mathbf{r}(s),n}^{\star} {}^{\ell} v_{\mathbf{r}(i),s}^{t} = \delta_{i,n} 1$ the fifth line of (70) is equal to the fifth line of (69). And that means the identities (70) and (69) are equivalent.

At this point it is most efficient to distinguish four cases and treat them individually.

4.6.2. *First case*. The first case is handled in only one further step.

Proposition 4.36. For any $v \in V$ and any $(j,i) \in [n]^{\times 2}$, if $j \neq n \neq i$, then

$$d_{\ell}(x_{j,i}^{v} \otimes 1) = \sum_{s=1}^{n} x_{j,s}^{v} \otimes {}^{\ell}v_{s,r(i)} + (-1)^{\ell} x_{r(j),i}^{v^{\ddagger}} \otimes 1.$$
(71)

Proof. Under the assumptions that both $j \neq n$ and $i \neq n$ the fourth and fifth lines of (69) in Lemma 4.35 are zero, which hence specializes to the statement that

$$d_{\ell}(x_{j,i}^{v} \otimes 1) = \sum_{s=1}^{n} x_{j,s}^{v} \otimes {}^{\ell} v_{s,r(i)}$$

$$+ \iota_{\ell-2} ((-1)^{\ell} \sum_{s=1}^{n} x_{r(j),s}^{v^{\dagger}} \otimes {}^{\ell} v_{s,r(i)} - x_{j,i}^{v} \otimes 1$$

$$+ \delta_{j,1} (-1)^{\ell} (x_{1,1}^{v^{\star}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\star}}) \otimes {}^{\ell} v_{r(i),n}^{t}).$$
(72)

The vector $x_{r(j),1}^{v^{\dagger}} \otimes {}^{\ell}v_{1,r(i)}$ is the tip of the argument of $\iota_{\ell-2}$ in (72) for the following reasons. The term $x_{j,i}^{v} \otimes 1$ is less already by degree. All the other summands in that line of (72) are dominated because $x_{r(j),s}^{v^{\dagger}} < x_{r(j),1}^{v^{\dagger}}$ for any $s \in [n]$ with 1 < s by the addendum to Lemma 4.4. Finally, the entire third line of (72) is less because, if it is non-zero, then j = 1, i.e., r(j) = n and hence $x_{s,s}^{v^*} < x_{r(j),1}^{v^{\dagger}}$ for any $s \in [n]$ with s < n by Lemma 4.4.

Because moreover by Proposition 4.10

$$\int_{\ell-2} (x_{\mathbf{r}(j),1}^{v^{\ddagger},\ell-2}, {}^{\ell}v_{1,\mathbf{r}(i)}) = (x_{\mathbf{r}(j),i}^{v^{\ddagger},\ell-1}, 1)$$

another valid identity is produced by adding to the right-hand side of (72) the term $(-1)^{\ell} x_{\mathbf{r}(j),i}^{v^{\ddagger}} \otimes 1$ outside of $\iota_{\ell-2}$ and $-(-1)^{\ell} d_{\ell-1}(x_{\mathbf{r}(j),i}^{v^{\ddagger},\ell-1} \otimes 1)$ inside.

That addition transforms the outside of $\iota_{\ell-1}$ in (72) into that in (71). And since by $r(j) \neq 1$ and $i \neq n$ Assumption 4.33 gives

$$-(-1)^{\ell} d_{\ell-1}(x_{\mathbf{r}(j),i}^{v^{\dagger},\ell-1} \otimes 1) = -(-1)^{\ell} \sum_{s=1}^{n} x_{\mathbf{r}(j),s}^{v^{\dagger}} \otimes {}^{\ell} v_{s,\mathbf{r}(i)} + x_{j,i}^{v} \otimes 1 -\delta_{j,1}(-1)^{\ell} (x_{1,1}^{v^{\star}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\star}}) \otimes {}^{\ell} v_{\mathbf{r}(i),n}^{\mathsf{t}}$$

the same is true of the inside of $\iota_{\ell-2}$, which proves the claim.

4.6.3. Second case. For the second case, $d_{\ell}(x_{j,i}^v \otimes 1)$ for j = n and $i \in [n]$ with $i \neq n$, it takes two more recursion steps to eliminate $\iota_{\ell-2}$ entirely.

Lemma 4.37. For any $v \in V$ and any $i \in [n]$, if $i \neq n$, then

$$d_{\ell}(x_{n,i}^{v} \otimes 1) = \sum_{s=1}^{n} x_{n,s}^{v} \otimes {}^{\ell} v_{s,\mathbf{r}(i)} + x_{1,1}^{v^{t}} \otimes {}^{\ell} v_{\mathbf{r}(i),n}^{t} + \iota_{\ell-2} \big((-1)^{\ell} \sum_{s=1}^{n} x_{1,s}^{v^{t}} \otimes {}^{\ell} v_{s,\mathbf{r}(i)} - x_{n,i}^{v} \otimes 1 \big).$$
(73)

Proof. Under the assumption $i \neq n$ instantiating Lemma 4.35 at j = nyields

$$d_{\ell}(x_{n,i}^{v} \otimes 1) = \sum_{s=1}^{n} x_{n,s}^{v} \otimes {}^{\ell}v_{s,r(i)}$$

$$+ \iota_{\ell-2} \Big(\sum_{s=1}^{n} x_{1,s}^{v^{t}} \otimes {}^{\ell}v_{s,n}^{\star} {}^{\ell}v_{r(i),n}^{t} - (-1)^{\ell} x_{n,1}^{v^{\star}} \otimes {}^{\ell}v_{r(i),n}^{t}$$

$$+ (-1)^{\ell} \sum_{s=1}^{n} x_{1,s}^{v^{\dagger}} \otimes {}^{\ell}v_{s,r(i)} - x_{n,i}^{v} \otimes 1 \Big).$$
(74)

Since $i \neq n$ the monomial ${}^{\ell}v_{1,n}^{\star}{}^{\ell}v_{r(i),n}^{t}$ is reduced by Lemma 4.2. The vector $x_{1,1}^{v^{t}} \otimes {}^{\ell}v_{1,n}^{\star} {}^{\ell}v_{r(i),n}^{t}$ is the tip of the argument of $\iota_{\ell-2}$ on the right-hand side of (74). Indeed, for degree reasons only the terms in the same grouped sum where $x_{1,1}^{v^{t}} \otimes {}^{\ell}v_{1,n}^{\star} {}^{\ell}v_{\mathbf{r}(i),n}$ comes from could possibly be greater or equal. However, for any $s \in [n]$ with 1 < s the addendum to Lemma 4.4 ensures that already $x_{1,s}^{v^t} < x_{1,1}^{v^t}$ and hence $\begin{array}{c} x_{1,s}^{v^{\mathrm{t}}} \otimes {}^{\ell} v_{s,n}^{\star} {}^{\ell} v_{\mathrm{r}(i),n}^{\mathrm{t}} <_{\ell-2} x_{1,1}^{v^{\mathrm{t}}} \otimes {}^{\ell} v_{1,n}^{\star} {}^{\ell} v_{\mathrm{r}(i),n}^{\mathrm{t}}.\\ \text{Given that by Proposition 4.10} \end{array}$

$$\int_{\ell-2} (x_{1,1}^{v^{t},\ell-2}, {}^{\ell}v_{1,n}^{\star} {}^{\ell}v_{\mathbf{r}(i),n}^{t}) = (x_{1,1}^{v^{t},\ell-1}, {}^{\ell}v_{\mathbf{r}(i),n}^{t}),$$

adding $x_{1,1}^{v^{t}} \otimes^{\ell} v_{\mathbf{r}(i),n}^{t}$ outside of $\iota_{\ell-2}$ and $-d_{\ell-1}(x_{1,1}^{v^{t}} \otimes^{\ell} v_{\mathbf{r}(i),n}^{t})$ inside turns (74) into another true identity.

This change brings the outside of $\iota_{\ell-2}$ in (74) exactly into the form of the outside in (73). Moreover, by Assumption 4.33

$$d_{\ell-1}(x_{1,1}^{v^{t}} \otimes 1) = \sum_{s=1}^{n} x_{1,s}^{v^{t}} \otimes {}^{\ell}v_{s,n}^{\star} - (-1)^{\ell} x_{n,1}^{v^{\star}} \otimes 1,$$

which is why

$$-d_{\ell-1}(x_{1,1}^{v^{t}} \otimes {}^{\ell}v^{t}_{\mathbf{r}(i),n}) = -\sum_{s=1}^{n} x_{1,s}^{v^{t}} \otimes {}^{\ell}v^{\star}_{s,n} {}^{\ell}v^{t}_{\mathbf{r}(i),n} + (-1)^{\ell} x_{n,1}^{v^{\star}} \otimes {}^{\ell}v^{t}_{\mathbf{r}(i),n}.$$
(75)

Evidently adding the vector (75) to the argument of $\iota_{\ell-2}$ in (74) produces precisely the argument of $\iota_{\ell-2}$ in (73). Thus, the claim is true.

Proposition 4.38. For any $v \in V$ and any $i \in [n]$, if $i \neq n$, then

$$d_{\ell}(x_{n,i}^{v} \otimes 1) = \sum_{s=1}^{n} x_{n,s}^{v} \otimes {}^{\ell}v_{s,\mathbf{r}(i)} + x_{1,1}^{v^{t}} \otimes {}^{\ell}v_{\mathbf{r}(i),n}^{t} + (-1)^{\ell} x_{1,i}^{v^{\ddagger}} \otimes 1.$$
(76)

Proof. The vector $x_{1,1}^{v^{\ddagger}} \otimes^{\ell} v_{1,\mathbf{r}(i)}$ is the tip of the argument of $\iota_{\ell-2}$ in (73) because $x_{1,s}^{v^{\ddagger}} < x_{1,1}^{v^{\ddagger}}$ for any $s \in [n]$ with 1 < s by the addendum to Lemma 4.4. And, of course, $x_{n,i}^v \otimes 1$ is dominated for degree reasons.

Since by Proposition 4.10

$$\int_{\ell-2} (x_{1,1}^{v^{\ddagger},\ell-2}, {}^{\ell}v_{1,\mathbf{r}(i)}) = (x_{1,i}^{v^{\ddagger},\ell-1}, 1)$$

it follows that the equality in (73) is preserved by adding to its righthand side $(-1)^{\ell} x_{1,i}^{v^{\dagger}} \otimes 1$ outside of $\iota_{\ell-2}$ and $-(-1)^{\ell} d_{\ell-1}(x_{1,i}^{v^{\dagger}} \otimes 1)$ inside.

That immediately makes the outsides of $\iota_{\ell-2}$ in (73) and (76) agree. At the same time $i \neq n$ implies according to Assumption 4.33

$$-(-1)^{\ell} d_{\ell-1}(x_{1,i}^{v^{\ddagger}} \otimes 1) = -(-1)^{\ell} \sum_{s=1}^{n} x_{1,s}^{v^{\ddagger}} \otimes {}^{\ell} v_{s,\mathbf{r}(i)} + x_{n,i}^{v} \otimes 1.$$

Hence, the argument of $\iota_{\ell-2}$ in (73) is canceled by the addition of $-(-1)^{\ell} d_{\ell-1}(x_{1,i}^{v^{\ddagger}} \otimes 1)$, which then proves (76).

4.6.4. Third case. In the third case, computing $d_{\ell}(x_{i,n}^v \otimes 1)$ for any $j \in [n]$ with $j \neq n$, two recursion steps are necessary to solve.

Lemma 4.39. For any $v \in V$ and any $j \in [n]$, if $j \neq n$, then

$$d_{\ell}(x_{j,n}^{v} \otimes 1) = \sum_{s=1}^{n} x_{j,s}^{v} \otimes^{\ell} v_{s,1} + (-1)^{\ell} x_{r(j),n}^{v^{\dagger}} \otimes 1$$

$$+ \delta_{j,1} \iota_{\ell-2} \big((-1)^{\ell} \sum_{s=1}^{n} x_{1,s}^{v^{\star}} \otimes^{\ell} v_{s,n}^{t} - x_{n,1}^{v^{\dagger}} \otimes 1 \big).$$
(77)

Proof. Specializing i to n in Lemma 4.35 and recognizing that the assumption $j \neq n$ removes the fourth and fifth lines of (69) gives

$$d_{\ell}(x_{j,n}^{v} \otimes 1) = \sum_{s=1}^{n} x_{j,s}^{v} \otimes^{\ell} v_{s,1}$$

$$+ \iota_{\ell-2} \left((-1)^{\ell} \sum_{s=1}^{n} x_{r(j),s}^{v^{\dagger}} \otimes^{\ell} v_{s,1} - x_{j,n}^{v} \otimes 1 \right)$$

$$+ \delta_{j,1} (-1)^{\ell} \left(x_{1,1}^{v^{\star}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\star}} \right) \otimes^{\ell} v_{1,n}^{t} \right).$$

$$(78)$$

The vector $x_{r(j),1}^{v^{\ddagger}} \otimes {}^{\ell}v_{1,1}$ is the tip of the argument of $\iota_{\ell-2}$ in (78). It dominates the other terms in the grouped sum from where it stems because for any $s \in [n]$ with 1 < s necessarily $x_{r(j),s}^{v^{\ddagger}} < x_{r(j),1}^{v^{\ddagger}}$ by the addendum to Lemma 4.4. And the terms in the third line of (78) are less because, if they are non-zero, then j = 1, i.e., r(j) = n and hence $x_{s,s}^{v^*} < x_{r(j),1}^{v^{\ddagger}}$ for any $s \in [n]$ with $s \neq n$ according to Lemma 4.4.

Because by Proposition 4.10

$$\int_{\ell-2} (x_{\mathbf{r}(j),1}^{v^{\dagger},\ell-2}, {}^{\ell}v_{1,1}) = (x_{\mathbf{r}(j),n}^{v^{\dagger},\ell-1}, 1)$$

a true identity is thus derived by adding $(-1)^{\ell} x_{r(j),n}^{v^{\dagger}} \otimes 1$ on the righthand side to the outside of $\iota_{\ell-2}$ in (78) and $-(-1)^{\ell} d_{\ell-2}(x_{r(j),n}^{v^{\dagger}} \otimes 1)$ to the inside.

That addition turns the outside of (78) into that of (77). Simultaneously, since $r(j) \neq 1$, by Assumption 4.33

$$-(-1)^{\ell} d_{\ell-1}(x_{r(j),n}^{v^{\dagger}} \otimes 1) = -(-1)^{\ell} \sum_{s=1}^{n} x_{r(j),s}^{v^{\dagger}} \otimes^{\ell} v_{s,1} + x_{j,n}^{v} \otimes 1 + \delta_{j,1}(-(-1)^{\ell} (x_{1,1}^{v^{\star}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\star}}) \otimes^{\ell} v_{1,n}^{t} + (-1)^{\ell} \sum_{s=1}^{n} x_{1,s}^{v^{\star}} \otimes^{\ell} v_{s,n}^{t} - x_{n,1}^{v^{t}} \otimes 1).$$
(79)

It is now evident that when the vector (79) and the argument of $\iota_{\ell-2}$ in (78) are added together the result is precisely the argument of $\iota_{\ell-2}$ in (77). And by what was said before that proves the assertion.

Proposition 4.40. For any $v \in V$ and any $j \in [n]$, if $j \neq n$, then

$$d_{\ell}(x_{j,n}^{v} \otimes 1) = \sum_{s=1}^{n} x_{j,s}^{v} \otimes {}^{\ell}v_{s,1} + (-1)^{\ell} x_{r(j),n}^{v^{\dagger}} \otimes 1 + \delta_{j,1}(-1)^{\ell} x_{1,1}^{v^{\star}} \otimes 1.$$
(80)

Proof. For any $s \in [n]$ with 1 < s the inequality $x_{1,s}^{v^*} < x_{1,1}^{v^*}$ holds by the addendum to Lemma 4.4. That makes $x_{1,1}^{v^*} \otimes {}^{\ell}v_{1,n}^{t}$ the tip of the argument of $\iota_{\ell-2}$ on the right-hand side of (77) in Lemma 4.39.

By Proposition 4.10

$$\int_{\ell-2} (x_{1,1}^{v^{\star},\ell-2}, {}^{\ell}v_{1,n}^{t}) = (x_{1,1}^{v^{\star},\ell-1}, 1).$$

Note that $\iota_{\ell-2}$ has a scalar factor of $\delta_{j,1}$ in (77). For those reasons another valid identity is obtained by adding to the right-hand side of (77) the terms $\delta_{j,1}(-1)^{\ell} x_{1,1}^{v^{\star}} \otimes 1$ outside of $\iota_{\ell-2}$ and $-(-1)^{\ell} d_{\ell-1}(x_{1,1}^{v^{\star}} \otimes 1)$ inside of it.

It is clear that this change turns the outside of $\iota_{\ell-2}$ in (77) into that in (80). And because according to Assumption 4.33

$$-(-1)^{\ell} d_{\ell-1}(x_{1,1}^{v^{\star}} \otimes 1) = -(-1)^{\ell} \sum_{s=1}^{n} x_{1,s}^{v^{\star}} \otimes {}^{\ell} v_{s,n}^{t} + x_{n,1}^{v^{t}} \otimes 1$$

the addition of $-d_{\ell-1}(x_{1,1}^{v^*} \otimes 1)$ exactly cancels out the argument of $\iota_{\ell-2}$ in (77). That verifies (80).

4.6.5. Fourth case. It remains to compute $d_{\ell}(x_{n,n}^v \otimes 1)$, which is the one requiring the most effort, another n+2 recursion steps.

Lemma 4.41. For any $v \in V$ and any $m \in [n]$,

$$d_{\ell}(x_{n,n}^{v} \otimes 1) = \sum_{s=1}^{n} x_{n,s}^{v} \otimes^{\ell} v_{s,1} - \sum_{s=2}^{m} x_{1,s}^{v^{t}} \otimes^{\ell} v_{s,n}^{t}$$

$$+ \iota_{\ell-2} \Big(-\sum_{s=m+1}^{n} x_{1,1}^{v^{t}} \otimes^{\ell} v_{1,r(s)}^{\star} \ell v_{s,n}^{t} \\
+ \sum_{s=2}^{n} \sum_{t=1}^{m} x_{1,s}^{v^{t}} \otimes^{\ell} v_{s,r(t)}^{\star} \ell v_{t,n}^{t} \\
- (-1)^{\ell} \sum_{s=1}^{m} x_{n,s}^{v^{\star}} \otimes^{\ell} v_{s,n}^{t} + (-1)^{\ell} \sum_{s=1}^{n} x_{1,s}^{v^{t}} \otimes^{\ell} v_{s,1} \\
- x_{n,n}^{v} \otimes 1 - \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{t}} \otimes 1 \\
- \zeta_{n,m} (-1)^{\ell} (x_{1,1}^{v^{t}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{t}}) \otimes^{\ell} v_{n,n}).$$

$$(81)$$

Proof. The proof goes by induction over m.

Induction base. By $\zeta_{n,1} = 0$ what needs to be proved in the base case m = 1 is that

$$d_{\ell}(x_{n,n}^{v} \otimes 1) = \sum_{s=1}^{n} x_{n,s}^{v} \otimes^{\ell} v_{s,1}$$

$$+ \iota_{\ell-2}(-\sum_{s=2}^{n} x_{1,1}^{v^{\dagger}} \otimes^{\ell} v_{1,r(s)}^{\star} \ell v_{s,n} + \sum_{s=2}^{n} x_{1,s}^{v^{\dagger}} \otimes^{\ell} v_{s,n}^{\star} v_{1,n}^{\star}$$

$$- (-1)^{\ell} x_{n,1}^{v^{\star}} \otimes^{\ell} v_{1,n}^{t} + (-1)^{\ell} \sum_{s=1}^{n} x_{1,s}^{v^{\dagger}} \otimes^{\ell} v_{s,1}$$

$$- x_{n,n}^{v} \otimes 1 - \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\dagger}} \otimes 1).$$
(82)

Specializing both i and j to n in (69) in Lemma 4.35 gives, after some rearrangement,

$$d_{\ell}(x_{n,n}^{v} \otimes 1) = \sum_{s=1}^{n} x_{n,s}^{v} \otimes {}^{\ell}v_{s,1}$$

$$+ \iota_{\ell-2} \Big(\sum_{s=1}^{n} x_{1,s}^{v^{t}} \otimes {}^{\ell}v_{s,n}^{\star} {}^{\ell}v_{1,n}^{t} - x_{1,1}^{v^{t}} \otimes 1$$

$$- (-1)^{\ell} x_{n,1}^{v^{\star}} \otimes {}^{\ell}v_{1,n}^{t} + (-1)^{\ell} \sum_{s=1}^{n} x_{1,s}^{v^{\dagger}} \otimes {}^{\ell}v_{s,1}$$

$$- x_{n,n}^{v} \otimes 1 - \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{t}} \otimes 1 \Big).$$

$$(83)$$

Since ${}^{\ell}v_{1,n}^{\star}{}^{\ell}v_{1,n}^{t} = -\sum_{t=2}^{n} {}^{\ell}v_{1,r(t)}^{\star}{}^{\ell}v_{t,n}^{t} + 1$ the terms in the argument of $\iota_{\ell-2}$ in the second line of (83) can also be written as

$$\begin{split} \sum_{s=1}^{n} x_{1,s}^{v^{\mathrm{t}}} \otimes {}^{\ell} v_{s,n}^{\star} {}^{\ell} v_{1,n}^{\mathrm{t}} - x_{1,1}^{v^{\mathrm{t}}} \otimes 1 \\ &= -\sum_{t=2}^{n} x_{1,1}^{v^{\mathrm{t}}} \otimes {}^{\ell} v_{1,r(t)}^{\star} {}^{\ell} v_{t,n}^{\mathrm{t}} + \sum_{s=2}^{n} x_{1,s}^{v^{\mathrm{t}}} \otimes {}^{\ell} v_{s,n}^{\star} v_{1,n}^{\star}, \end{split}$$

i.e., as the terms in the argument of $\iota_{\ell-2}$ in the second line of (82). Hence, the claim is true for m = 1.

Induction step. Let $m \in [n]$ be such that $m \leq n-1$ and such that (81) holds. By the argument below, (81) is then still true if m is replaced by m+1.

Since $1 \leq m$ the monomial ${}^{\ell}v_{1,r(m+1)}^{\star}{}^{\ell}v_{m+1,n}^{t}$ is normal by Lemma 4.2. The tip of the argument of $\iota_{\ell-2}$ on the right-hand side of (81) is the vector $x_{1,1}^{v^{t}} \otimes {}^{\ell}v_{1,r(m+1)}^{\star}{}^{\ell}v_{m+1,n}^{t}$ for the following reasons. It is greater than any other summand in the grouped sum where it originates because for any $s \in [n]$ with m+1 < s necessarily ${}^{\ell}v_{1,r(s)}^{\star} < {}^{\ell}v_{1,r(m+1)}^{\star}$ by Lemma 3.4

and thus $x_{1,1}^{v^{t}} \otimes {}^{\ell}v_{1,r(s)}^{\star} {}^{\ell}v_{s,n}^{t} <_{\ell-2} x_{1,1}^{v^{t}} \otimes {}^{\ell}v_{1,r(m+1)}^{\star} {}^{\ell}v_{m+1,n}^{t}$. Any term in the fourth line of (81) is dominated because according to the addendum to Lemma 4.4 already $x_{1,s}^{v^{t}} < x_{1,1}^{v^{t}}$ for any $s \in [n]$ with 1 < s. And all the remaining terms are less for degree reasons.

Because by Proposition 4.10

$$\int_{\ell-2} (x_{1,1}^{v^{\mathrm{t}},\ell-2}, {}^{\ell}v_{1,\mathrm{r}(m+1)}^{\star} {}^{\ell}v_{m+1,n}^{\mathrm{t}}) = (x_{1,m+1}^{v^{\mathrm{t}},\ell-1}, {}^{\ell}v_{m+1,n}^{\mathrm{t}}),$$

if on the right-hand side of (81) the vector $-x_{1,m+1}^{v^{t}} \otimes {}^{\ell}v_{m+1,n}^{t}$ is added outside of $\iota_{\ell-2}$ and the vector $d_{\ell-1}(x_{1,m+1}^{v^{t}} \otimes {}^{\ell}v_{m+1,n}^{t})$ inside, the equality is preserved.

Obviously, on the exterior of $\iota_{\ell-2}$ in (81) this has the same effect as replacing m by m+1 would. At the same time by $2 \le n$ Assumption 4.33 gives

$$d_{\ell-1}(x_{1,m+1}^{v^{t}} \otimes 1) = \sum_{s=1}^{n} x_{1,s}^{v^{t}} \otimes {}^{\ell}v_{s,r(m+1)}^{\star} - (-1)^{\ell} x_{n,m+1}^{v^{\star}} \otimes 1 - \delta_{m+1,n} (-1)^{\ell} (x_{1,1}^{v^{\dagger}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\dagger}}) \otimes 1.$$

Multiplying with ${}^{\ell}v_{m+1,n}^{t}$ from the right and splitting off the s = 1 term in the first line yields

$$d_{\ell-1}(x_{1,m+1}^{v^{t}} \otimes^{\ell} v_{m+1,n}^{t}) = x_{1,1}^{v^{t}} \otimes^{\ell} v_{1,r(m+1)}^{\star} {}^{\ell} v_{m+1,n}^{t} + \sum_{s=2}^{n} x_{1,s}^{v^{t}} \otimes^{\ell} v_{s,r(m+1)}^{\star} {}^{\ell} v_{m+1,n}^{t}$$

$$- (-1)^{\ell} x_{n,m+1}^{v^{\star}} \otimes^{\ell} v_{m+1,n}^{t}$$

$$- \delta_{n,m+1} (-1)^{\ell} (x_{1,1}^{v^{t}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{t}}) \otimes^{\ell} v_{n,n},$$
(84)

where it has been used that $\delta_{m+1,n} {}^{\ell} v_{m+1,n}^{t} = \delta_{n,m+1} {}^{\ell} v_{n,n}$ at the end. Because $\zeta_{n,m} + \delta_{n,m+1} = \zeta_{n,m+1}$ it is now evident that adding the vector (84) to the argument of $\iota_{\ell-2}$ changes (81) in the same way as increasing m by 1. That completes the induction.

Lemma 4.42. For any $v \in V$,

$$d_{\ell}(x_{n,n}^{v} \otimes 1) = \sum_{s=1}^{n} x_{n,s}^{v} \otimes^{\ell} v_{s,1} - \sum_{s=2}^{n} x_{1,s}^{v^{t}} \otimes^{\ell} v_{s,n}^{t} - (-1)^{\ell} x_{n,1}^{v^{\star}} \otimes 1 \qquad (85)$$

+ $\iota_{\ell-2} ((-1)^{\ell} \sum_{s=1}^{n} x_{1,s}^{v^{\dagger}} \otimes^{\ell} v_{s,1} - x_{n,n}^{v} \otimes 1$
- $(x_{1,1}^{v^{t}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{t}}) \otimes 1).$

Proof. Instatiating (81) in Lemma 4.41 at m = n renders the third and fourth lines of (81) zero. In the case of the third line that is obvious. The fourth line vanishes because

$$\sum_{s=2}^{n} x_{1,s}^{v^{t}} \otimes \sum_{t=1}^{n} {}^{\ell} v_{s,r(t)}^{\star} {}^{\ell} v_{t,n}^{t} = \sum_{s=2}^{n} \delta_{s,1} x_{1,s}^{v^{t}} \otimes 1 = 0.$$

Hence, what survives the specialization to m = n is the identity

$$d_{\ell}(x_{n,n}^{v} \otimes 1) = \sum_{s=1}^{n} x_{n,s}^{v} \otimes^{\ell} v_{s,1} - \sum_{s=2}^{n} x_{1,s}^{v^{\dagger}} \otimes^{\ell} v_{s,n}^{t} \qquad (86) + \iota_{\ell-2}(-(-1)^{\ell} \sum_{s=1}^{n} x_{n,s}^{v^{\star}} \otimes^{\ell} v_{s,n}^{t} + (-1)^{\ell} \sum_{s=1}^{n} x_{1,s}^{v^{\dagger}} \otimes^{\ell} v_{s,1} - x_{n,n}^{v} \otimes 1 - \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\dagger}} \otimes 1 - (-1)^{\ell} (x_{1,1}^{v^{\dagger}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\star}}) \otimes^{\ell} v_{n,n}).$$

The vector $x_{n,1}^{v^*} \otimes^{\ell} v_{1,n}^{t}$ is the tip of the argument of $\iota_{\ell-2}$ in (86). Indeed, any other term in the same grouped sum is less by the addendum to Lemma 4.4 because $x_{n,s}^{v^*} < x_{n,1}^{v^*}$ for any $s \in [n]$ with 1 < s. Any term of the form $x_{1,s}^{v^{\dagger}} \otimes^{\ell} v_{s,1}$ for some $s \in [n]$ is dominated because already $x_{1,s}^{v^{\dagger}} < x_{n,1}^{v^{\star}}$ by Lemma 4.4. For the same reason $x_{s,s}^{v^{\dagger}} < x_{n,1}^{v^{\star}}$ for any $s \in [n]$ with s < n. Hence, $x_{n,1}^{v^{\star}} \otimes^{\ell} v_{1,n}^{t}$ is also greater than any term in the fifth line of (86). Of course all other terms are less already for reasons of degree.

In consequence and because, according to Proposition 4.10,

$$\int_{\ell-2} (x_{n,1}^{v^{\star},\ell-2}, {}^{\ell}v_{1,n}^{t}) = (x_{n,1}^{v^{\star},\ell-1}, 1),$$

if $-(-1)^{\ell} x_{n,1}^{v^{\star}} \otimes 1$ is added to the outisde of $\iota_{\ell-2}$ and $(-1)^{\ell} d_{\ell-1}(x_{n,1}^{v^{\star}} \otimes 1)$ to the inside on the right-hand side of (86) the identity remains true.

With this change, the outside of $\iota_{\ell-2}$ in (86) becomes that in (85). And since $2 \leq n$ Assumption 4.33 implies

$$(-1)^{\ell} d_{\ell-1}(x_{n,1}^{v^{\star}} \otimes 1) = (-1)^{\ell} \sum_{s=1}^{n} x_{n,s}^{v^{\star}} \otimes {}^{\ell} v_{s,n}^{t} - x_{1,1}^{v^{t}} \otimes 1 + (-1)^{\ell} (x_{1,1}^{v^{\dagger}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\dagger}}) \otimes {}^{\ell} v_{n,n}.$$
(87)

This shows that adding the term (87) to the argument of $\iota_{\ell-2}$ in (86) transforms the latter into the argument in (85). That concludes the proof.

Proposition 4.43. For any $v \in V$,

$$d_{\ell}(x_{n,n}^{v} \otimes 1) = \sum_{s=1}^{n} x_{n,s}^{v} \otimes^{\ell} v_{s,1} - \sum_{s=2}^{n} x_{1,s}^{v^{t}} \otimes^{\ell} v_{s,n}^{t}$$

$$-(-1)^{\ell} x_{n,1}^{v^{\star}} \otimes 1 + (-1)^{\ell} x_{1,n}^{v^{\ddagger}} \otimes 1.$$
(88)

Proof. The tip of the argument of $\iota_{\ell-2}$ in (85) in Lemma 4.42 is $x_{1,1}^{v^{\ddagger}} \otimes^{\ell} v_{1,1}$. Any other term in the grouped sum in the third line of (85) is less because $x_{1,s}^{v^{\ddagger}} < x_{1,1}^{v^{\ddagger}}$ for any $s \in [n]$ with 1 < s by the addendum to Lemma 4.4. And all remaining terms are dominated for degree reasons.

Since by Proposition 4.10

$$\int_{\ell-2} (x_{1,1}^{v^{\ddagger},\ell-2},{}^{\ell}v_{1,1}) = (x_{1,n}^{v^{\ddagger},\ell-1},1)$$

it follows that the veracity of (85) is not affected by adding the vector $(-1)^{\ell} x_{1,n}^{v^{\ddagger}} \otimes 1$ to the outside of $\iota_{\ell-2}$ and $-(-1)^{\ell} d_{\ell-1}(x_{1,n}^{v^{\ddagger}} \otimes 1)$ to the inside.

This change makes the outsides of $\iota_{\ell-2}$ in (85) and in (88) agree. On the other hand, since by Assumption 4.33

$$-(-1)^{\ell} d_{\ell-1}(x_{1,n}^{v^{\ddagger}} \otimes 1) = -(-1)^{\ell} \sum_{s=1}^{n} x_{1,s}^{v^{\ddagger}} \otimes^{\ell} v_{s,1} + x_{n,n}^{v} \otimes 1$$

$$+ (x_{1,1}^{v^{\ddagger}} + \delta_{\ell,4} \sum_{s=2}^{n-1} x_{s,s}^{v^{\ddagger}}) \otimes 1$$
(89)

the argument of $\iota_{\ell-2}$ in (85) is actually the negative of the term (89). That proves (88).

4.6.6. Synthesis. The four particular results obtained in the different cases assemble into the claim from the Main result. Recall that $4 \leq \ell$.

Theorem 4.44. For any $v \in V$ and $(j,i) \in [n]^{\times 2}$,

$$d_{\ell}(x_{j,i}^{v} \otimes 1) = \sum_{s=1}^{n} x_{j,s}^{v} \otimes^{\ell} v_{s,r(i)} + (-1)^{\ell} x_{r(j),i}^{v^{\dagger}} \otimes 1 \qquad (90) + \delta_{j,n}(x_{1,1}^{v^{\dagger}} \otimes^{\ell} v_{r(i),n}^{t} - \delta_{i,n}(\sum_{s=1}^{n} x_{1,s}^{v^{\dagger}} \otimes^{\ell} v_{s,n}^{t} + (-1)^{\ell} x_{n,1}^{v^{\star}} \otimes 1)) + \delta_{j,1}\delta_{i,n}(-1)^{\ell} x_{1,1}^{v^{\star}} \otimes 1.$$

Proof. The first two lines of (90) agree with (71) from Proposition 4.36. Hence, in the case that $j \neq n$ and $i \neq n$ the claim is true.

If j = n but $i \neq n$, then the only non-zero term in the last two lines of (90) is $x_{1,1}^{v^{t}} \otimes {}^{\ell}v_{r(i),n}^{t}$. That is why (90) is identical to (76) from Proposition 4.38 in this case.

In the opposite instance that $j \neq n$ but i = n the third line of (90) vanishes and the fourth reads $\delta_{j,1}(-1)^{\ell} x_{1,1}^{v^{\star}} \otimes 1$. That is precisely the term which needs to be added on the right-hand side to turn the first two lines of (90) into (80) from Proposition 4.40.

In the final case where j = i = n the last line of (90) is zero and what remains of the right-hand side is

$$\sum_{s=1}^{n} x_{n,s}^{v} \otimes^{\ell} v_{s,1} + (-1)^{\ell} x_{1,n}^{v^{\ddagger}} \otimes 1$$

$$+ x_{1,1}^{v^{\ddagger}} \otimes^{\ell} v_{1,n}^{t} - \sum_{s=1}^{n} x_{1,s}^{v^{\ddagger}} \otimes^{\ell} v_{s,n}^{t} - (-1)^{\ell} x_{n,1}^{v^{\star}} \otimes 1$$
(91)

Because the term $x_{1,1}^{v^{t}} \otimes {}^{\ell}v_{1,n}^{t}$ also occurs in the grouped sum in the middle of the second line of (91), but with opposite sign, this proves that (90) is actually the same as (88) from Proposition 4.43 in this case. Hence, the claim holds for all (j, i).

5. Recomputing the cohomology of U_n^+

In this section it is demonstrated that the Anick resolution of the right A-module \mathbb{K}_{ε} obtained in Section 4 is good enough to compute

$$\operatorname{Ext}_{A}^{\cdot}(\mathbb{K}_{\varepsilon},\mathbb{K}_{\varepsilon}),$$

which in the case $\mathbb{K} = \mathbb{C}$ gives the quantum group cohomology $H^{\cdot}(U_n^+)$ of the discrete dual of the free unitary quantum group U_n^+ . Note that, as special cases of much further-reaching results, the orders 1 and 2

of $\operatorname{Ext}_{A}^{\cdot}(\mathbb{K}_{\varepsilon}, \mathbb{K}_{\varepsilon})$ had already been computed by Das, Franz, Kula and Skalski in [DFKS18] and the higher orders by Baraquin, Franz, Gerhold, Kula and Tobolski in [BFGKT23].

Let Hom_A be the external hom functor from the category of right Amodules to the category of \mathbb{K} -vector spaces. Since P_{ℓ} is finite-dimensional as an A-module for each $\ell \in \mathbb{Z}_{\geq 0}$ it is enough to compute the defect of the \mathbb{K} -linear map $\operatorname{Hom}_A(d_{\ell}, \mathbb{K}_{\varepsilon})$ for each $\ell \in \mathbb{Z}_{>0}$.

As in Section 4 the difference between p and p + J will be suppressed in the notation for any $p \in \mathbb{K}\langle E \rangle$.

Notation 5.1. For any $\ell \in \mathbb{Z}_{>0}$, any $g \in \text{Hom}_A(P_\ell, \mathbb{K}_{\varepsilon})$, any $v \in V$ and any $(j, i) \in [n]^{\times 2}$ let

$$g_{j,i}^{v} := g(x_{j,i}^{v,\ell} \otimes 1).$$

5.1. First order. Computing the defect of $\operatorname{Hom}_A(d_1, \mathbb{K}_{\varepsilon})$ is easy.

Lemma 5.2. For any $g \in \operatorname{Hom}_A(P_0, \mathbb{K}_{\varepsilon})$, any $v \in V$ and $(j, i) \in [n]^{\times 2}$, $(g \circ d_1)(x_{j,i}^v \otimes 1) = 0.$

Proof. Since P_0 is the regular A-module and since by Proposition 4.12

$$(g \circ d_1)(x_{j,i}^v \otimes 1) = g(v_{j,i} - \delta_{j,i} 1) = g(v_{j,i}) - \delta_{j,i} g(1)$$

the fact that $g(v_{j,i}) = \delta_{j,i} g(1)$ by definition of \mathbb{K}_{ε} proves the claim. \Box

Lemma 5.3. The defect and rank of $\operatorname{Hom}_A(d_1, \mathbb{K}_{\varepsilon})$ are 1 and 0 respectively.

Proof. The domain \mathbb{K} -vector space $\operatorname{Hom}_A(P_0, \mathbb{K}_{\varepsilon}) = \operatorname{Hom}_A(A, \mathbb{K}_{\varepsilon}) \cong \mathbb{K}$ of $\operatorname{Hom}_A(d_1, \mathbb{K}_{\varepsilon})$ is one-dimensional. Hence, the claim follows from Lemma 5.2 by the dimension formula for \mathbb{K} -linear maps. \Box

5.2. Second order. The defect and rank of $\operatorname{Hom}_A(d_2, \mathbb{K}_{\varepsilon})$ can still be computed without much effort. Recall Notation 5.1.

Lemma 5.4. For any $g \in \text{Hom}_A(P_1, \mathbb{K}_{\varepsilon})$, any $v \in V$ and $(j, i) \in [n]^{\times 2}$, $(g \circ d_2)(x_{j,i}^v \otimes 1) = g_{j,r(i)}^v + g_{r(i),j}^{v^*} - \delta_{j,n} \delta_{i,n} \sum_{s=2}^n (g_{r(s),s}^v + g_{s,r(s)}^{v^*})$. (92)

Proof. Theorem 4.18 shows that $(g \circ d_2)(x_{j,i}^v \otimes 1)$ is given by

$$\sum_{s=1}^{n} g(x_{j,s}^{v} \otimes v_{s,r(i)}^{\ddagger}) + g(x_{r(j),i}^{v^{\ddagger}} \otimes 1) - \delta_{j,n} \delta_{i,n} \sum_{s=2}^{n} (\sum_{t=1}^{n} g(x_{t,s}^{v} \otimes v_{s,r(t)}^{\ddagger}) + g(x_{s,r(s)}^{v^{\ddagger}} \otimes 1)).$$

By $g(x_{j,s}^v \otimes v_{s,\mathbf{r}(i)}^{\ddagger}) = \delta_{s,\mathbf{r}(i)} g_{j,s}^v$ and $g(x_{t,s}^v \otimes v_{s,\mathbf{r}(t)}^{\ddagger}) = \delta_{s,\mathbf{r}(t)} g_{t,s}^v$ this is equal to

$$g_{j,\mathbf{r}(i)}^{v} + g_{\mathbf{r}(j),i}^{v^{\ddagger}} - \delta_{j,n}\delta_{i,n}\sum_{s=2}^{n}(g_{\mathbf{r}(s),s}^{v} + g_{s,\mathbf{r}(s)}^{v^{\ddagger}}).$$
(93)

Since for any $s \in [n]$ Lemma 4.3 implies $x_{r(j),i}^{v^{\ddagger},1} = x_{r(i),j}^{v^{\dagger},1}$ and $x_{s,r(s)}^{v^{\ddagger},1} = x_{s,r(s)}^{v^{\dagger},1}$ and, thus,

$$g_{\mathbf{r}(j),i}^{v^{\ddagger}} = g_{\mathbf{r}(i),j}^{v^{\star}}, \qquad g_{s,\mathbf{r}(s)}^{v^{\ddagger}} = g_{s,\mathbf{r}(s)}^{v^{\star}}$$

the vector (93) is the right-hand side of (92).

Lemma 5.5. Any $g \in \text{Hom}_A(P_1, \mathbb{K}_{\varepsilon})$ satisfies $g \circ d_2 = 0$ if and only if for any $v \in V$ and any $(j, i) \in [n]^{\times 2}$,

$$g_{j,i}^{v^{\star}} = -g_{i,j}^{v}. \tag{94}$$

In fact, (94) for any $(j,i) \in [n]^{\times 2}$ is satisfied by any $v \in V$ as soon as it is by an arbitrary one $v \in V$.

Proof. Step 1: Equivalence. By Lemma 5.4 it is enough to prove that the conditions from the claim are satisfied if and only if for any $v \in V$ and any $(j,i) \in [n]^{\times 2}$,

$$g_{j,\mathbf{r}(i)}^{v} + g_{\mathbf{r}(i),j}^{v^{\star}} - \delta_{j,n} \delta_{i,n} \sum_{s=2}^{n} (g_{\mathbf{r}(s),s}^{v} + g_{s,\mathbf{r}(s)}^{v^{\star}}) = 0.$$
(95)

Obviously, if (94) holds for any $v \in V$ and $(j, i) \in [n]^{\times 2}$, then so does (95).

Conversely, suppose (95) is true for any $v \in V$ and $(j, i) \in [n]^{\times 2}$. Let $v \in V$ and $(j, i) \in [n]^{\times 2}$ be arbitrary. If $(j, i) \neq (n, 1)$, which is to say $(j, r(i)) \neq (n, n)$, condition (95) with $(v^*, j, r(i))$ in the role of (v, j, i) collapse to the demand that $g_{j,i}^{v^*} + g_{i,j}^v = 0$. Hence, all that is left to prove is that also $g_{n,1}^{v^*} = -g_{1,n}^v$. But for (v, 1, 1) in the role of (v, j, i) condition (95) reads $g_{1,n}^v + g_{n,1}^{v^*} = 0$. Hence, the claimed equivalence is true.

Step 2: Addendum. It remains to prove the addendum. Hence, suppose there exists some $w \in V$ such that, for v = w specifically, (94) holds for any $(j,i) \in [n]^{\times 2}$. By symmetry it is then true that (94) is satisfied for any $v \in \{w, w^*\}$ and $(j,i) \in [n]^{\times 2}$. Because Lemma 4.3 and the assumption imply

$$g_{j,i}^{w^{\ddagger}} = g_{\mathbf{r}(i),\mathbf{r}(j)}^{w^{\star}} = -g_{\mathbf{r}(j),\mathbf{r}(i)}^{w} = -g_{i,j}^{w^{\dagger}}$$

equation (94) is actually satisfied for any $v \in V$ and $(j, i) \in [n]^{\times 2}$. That concludes the proof.

Lemma 5.6. Both defect and rank of $\operatorname{Hom}_A(d_2, \mathbb{K}_{\varepsilon})$ are n^2 .

Proof. The domain space $\operatorname{Hom}_A(P_1, \mathbb{K}_{\varepsilon}) = \operatorname{Hom}_A(\mathbb{K}C_1 \otimes_{\mathbb{K}} A, \mathbb{K}_{\varepsilon}) \cong \mathbb{K}C_1$ of $\operatorname{Hom}_A(d_2, \mathbb{K}_{\varepsilon})$ has dimension $|C_1| = |\{x_{j,i}^{u,1}, x_{j,i}^{u^*,1}\}_{j,i=1}^n| = 2n^2$. Lemma 5.5 shows that the kernel of $\operatorname{Hom}_A(d_2, \mathbb{K}_{\varepsilon})$ is in particular free over the set $\{x_{j,i}^{u,1}\}_{j,i=1}^n$, which has n^2 many elements. Hence, the claim follows by the dimension formula. \Box

5.3. Third and higher orders. It is most economic to first treat all the maps $\operatorname{Hom}_A(d_{\ell}, \mathbb{K}_{\varepsilon})$ for $3 \leq \ell$ simultaneously up to a certain point and then distinguish whether $\ell = 3$ or not.

5.3.1. Common results for orders three and higher. The strategy remains the same as for orders 1 and 2. Only now, more effort is required.

Lemma 5.7. For any $\ell \in \mathbb{Z}_{>0}$ with $3 \leq \ell$, any $g \in \operatorname{Hom}_A(P_{\ell-1}, \mathbb{K}_{\varepsilon})$, any $v \in V$ and any $(j, i) \in [n]^{\times 2}$,

$$(g \circ d_{\ell})(x_{j,i}^{v} \otimes 1) = g_{j,r(i)}^{v} + (-1)^{\ell} g_{r(j),i}^{v^{\dagger}}$$

$$(96)$$

$$- \delta_{j,n} \delta_{i,n} \left(g_{1,n}^{v^{\dagger}} + (-1)^{\ell} g_{n,1}^{v^{\star}} \right)$$

$$+ \delta_{j,1} \delta_{i,n} (-1)^{\ell} \left(g_{1,1}^{v^{\star}} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{\star}} \right)$$

$$+ \delta_{j,n} \delta_{i,1} \left(g_{1,1}^{v^{\dagger}} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{\dagger}} \right).$$

Proof. According to Theorems 4.32 and 4.44 the vector $(g \circ d_{\ell})(x_{j,i}^v \otimes 1)$ is given by

$$\begin{split} &\sum_{s=1}^{n} g(x_{j,s}^{v} \otimes^{\ell} v_{s,r(i)}) + (-1)^{\ell} g(x_{r(j),i}^{v^{\dagger}} \otimes 1) \\ &+ \delta_{j,n} \left(g(x_{1,1}^{v^{t}} \otimes^{\ell} v_{r(i),n}^{t}) + \delta_{\ell,3} \sum_{s=2}^{n-1} g(x_{s,s}^{v^{t}} \otimes^{\ell} v_{r(i),n}^{t}) \right. \\ &- \delta_{i,n} \left(\sum_{s=1}^{n} g(x_{1,s}^{v^{t}} \otimes^{\ell} v_{s,n}^{t}) + (-1)^{\ell} g(x_{n,1}^{v^{\star}} \otimes 1) \right) \right) \\ &+ \delta_{j,1} \delta_{i,n} (-1)^{\ell} \left(g(x_{1,1}^{v^{\star}} \otimes 1) + \delta_{\ell,3} \sum_{s=2}^{n-1} g(x_{s,s}^{v^{\star}} \otimes 1) \right). \end{split}$$

Recognizing $g(x_{j,s}^{v} \otimes^{\ell} v_{s,r(i)}) = \delta_{s,r(i)} g_{j,s}^{v}$ and $g(x_{s,s}^{v^{t}} \otimes^{\ell} v_{r(i),n}^{t}) = \delta_{r(i),n} g_{s,s}^{v^{t}} = \delta_{i,1} g_{s,s}^{v^{t}}$ and $g(x_{1,s}^{v^{t}} \otimes^{\ell} v_{s,n}^{t}) = \delta_{s,n} g_{1,s}^{v^{t}}$ for any $s \in [n]$ and reordering the lines shows that the above is equal to the right-hand side of (96). \Box

Lemma 5.8. For any $\ell \in \mathbb{Z}_{>0}$ with $3 \leq \ell$, any $g \in \text{Hom}_A(P_{\ell-1}, \mathbb{K}_{\varepsilon})$ satisfies $g \circ d_{\ell} = 0$ if and only if for any $v \in V$, all in conjunction, for any $(j, i) \in [n]^{\times 2}$ with $(j, i) \neq (1, 1)$,

$$g_{j,i}^{v^{t}} = -(-1)^{\ell} g_{\mathbf{r}(j),\mathbf{r}(i)}^{v^{\star}} - \delta_{j,n} \delta_{i,n} \left(g_{1,1}^{v} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v} \right), \qquad (97)$$

$$g_{j,i}^{v^{\downarrow}} = -(-1)^{\ell} g_{\mathbf{r}(j),\mathbf{r}(i)}^{v} - \delta_{j,n} \delta_{i,n} \left(g_{1,1}^{v^{\star}} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{\star}} \right)$$
(98)

and

$$g_{1,1}^{v^{\ddagger}} = -(-1)^{\ell} g_{1,1}^{v^{\ddagger}} - (-1)^{\ell} g_{n,n}^{v} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{\star}}$$
(99)

as well as

$$g_{n,n}^{v^{\star}} = -\delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{\star}} + (-1)^{\ell} (g_{n,n}^{v} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v}).$$
(100)

In fact, (97)-(100) for any $(j,i) \in [n]^{\times 2}$ with $(j,i) \neq (1,1)$ are all satisfied by each $v \in V$ as soon as they are all satisfied by an arbitrary one $v \in V$.

Proof. The claimed equivalence and the addendum will be proved separately.

Step 1: Equivalence. By Lemma 5.7 it is enough to prove that the conditions from the claim and the demand that for any $v \in V$ and

 $(j,i)\in [n]^{\times 2},$

$$0 = g_{j,r(i)}^{v} + (-1)^{\ell} g_{r(j),i}^{v^{\dagger}}$$

$$- \delta_{j,n} \delta_{i,n} \left(g_{1,n}^{v^{\dagger}} + (-1)^{\ell} g_{n,1}^{v^{\star}} \right)$$

$$+ \delta_{j,1} \delta_{i,n} (-1)^{\ell} \left(g_{1,1}^{v^{\star}} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{\star}} \right)$$

$$+ \delta_{j,n} \delta_{i,1} \left(g_{1,1}^{v^{\dagger}} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{\dagger}} \right)$$

$$(101)$$

are equivalent. The two implications will be treated separately.

Step 1.1. First, let (97)–(100) hold for any $v \in V$ and any $(j,i) \in [n]^{\times 2}$ with $(j,i) \neq (1,1)$. For any $v \in V$ instantiating (97) at (j,i) = (1,n)produces the identity $g_{1,n}^{v^{t}} = -(-1)^{\ell} g_{n,1}^{v^{\star}}$. The latter shows that the second line of (101) vanishes for arbitrary $v \in V$ and $(j,i) \in [n]^{\times 2}$. Now, let $v \in V$ and $(j,i) \in [n]^{\times 2}$ be arbitrary. Then (101) is satisfied by (v, j, i) for the following reasons.

Case 1.1.1. If $(j, i) \neq (n, 1)$, then in addition to the second also the fourth line of (101) is zero, which means that what needs to be shown is that

$$0 = g_{j,\mathbf{r}(i)}^{v} + (-1)^{\ell} g_{\mathbf{r}(j),i}^{v^{\ddagger}} + \delta_{j,1} \delta_{i,n} (-1)^{\ell} (g_{1,1}^{v^{\star}} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{\star}}).$$

Multiplying this identity with $(-1)^{\ell}$ shows it equivalent to (98) for r(j) in the role j. Since $(r(j), i) \neq (1, 1)$ by assumption that proves (101) in this case.

Case 1.1.2. Alternatively, for (j, i) = (n, 1) condition (101) collapses to the demand that

$$0 = g_{n,n}^{v} + (-1)^{\ell} g_{1,1}^{v^{\dagger}} + (g_{1,1}^{v^{\dagger}} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{\dagger}}).$$

But this is a mere rearrangement of (97) for (v^{t}, n, n) in the role of (v, j, i) there. Hence, (101) is true in all cases. That proves one half of the asserted equivalence.

Step 1.2. Conversely, suppose (101) is true for any $v \in V$ and $(j,i) \in [n]^{\times 2}$. Then, for any $v \in V$ with $(v^{\ddagger}, 1, 1)$ in place of (v, j, i) equation (101) gives

$$0 = g_{1,n}^{v^{\ddagger}} + (-1)^{\ell} g_{n,1}^{v}, \qquad (102)$$

while instantiating (101) at (v, j, i) = (v, n, n) yields

$$0 = g_{n,1}^{v} + (-1)^{\ell} g_{1,n}^{v^{\ddagger}} - (g_{1,n}^{v^{\ddagger}} + (-1)^{\ell} g_{n,1}^{v^{\star}}).$$
(103)

For any $v \in V$ multiplying (102) with $(-1)^{\ell}$ and subtracting (103) from the resulting identity now proves $0 = g_{1,n}^{v^{t}} + (-1)^{\ell} g_{n,1}^{v^{\star}}$. Hence, the second line of (101) is actually zero no matter what (v, j, i) is.

In consequence, for any $v \in V$ and any $(j, i) \in [n]^{\times 2}$ with $(v, \mathbf{r}(j), i)$ in place of (v, j, i) assumption (101) implies

$$0 = g_{\mathbf{r}(j),\mathbf{r}(i)}^{v} + (-1)^{\ell} g_{j,i}^{v^{\dagger}}$$

$$+ \delta_{j,n} \delta_{i,n} (-1)^{\ell} (g_{1,1}^{v^{\star}} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{\star}})$$

$$+ \delta_{j,1} \delta_{i,1} (g_{1,1}^{v^{\dagger}} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{\dagger}}).$$
(104)

Multiplying (104) with $(-1)^{\ell}$ and solving for $g_{j,i}^{v^{\ddagger}}$ produces for any $v \in V$ and any $(j,i) \in [n]^{\times 2}$ the identity

$$g_{j,i}^{v^{\ddagger}} = -(-1)^{\ell} g_{r(j),r(i)}^{v}$$

$$- \delta_{j,n} \delta_{i,n} (g_{1,1}^{v^{\star}} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{\star}})$$

$$- \delta_{j,1} \delta_{i,1} (-1)^{\ell} (g_{1,1}^{v^{t}} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{t}}).$$
(105)

Evidently, for any $v \in V$ any $(j, i) \in [n]^{\times 2}$ with $(j, i) \neq (1, 1)$ there is no difference between (105) and (98), which proves the latter.

But replacing v by v^* transforms (98) into (97), which has thus been verified as well for any $v \in V$ any $(j,i) \in [n]^{\times 2}$ with $(j,i) \neq (1,1)$. It only remains to prove (99) and (100).

Applying (97) to (s, s) in the role of (j, i) for any $v \in V$ and $s \in [n]$ with $s \neq 1$ shows $g_{s,s}^{v^{t}} = -(-1)^{\ell} g_{r(s),r(s)}^{v^{\star}}$ and thus in particular for any $v \in V$,

$$\delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{t}} = -(-1)^{\ell} \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{\star}}.$$
 (106)

Inserting (106) into (105) evaluated at (j, i) = (1, 1) now proves (99) for any $v \in V$.

Multiplying (99) with $(-1)^{\ell}$ and rearranging yields

$$g_{1,1}^{v^{t}} = -g_{n,n}^{v} - (-1)^{\ell} g_{1,1}^{v^{t}} + (-1)^{\ell} \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{\star}}$$
(107)

for any $v \in V$. Replacing v by v^* in (107) thus shows for any $v \in V$

$$g_{1,1}^{v^{\ddagger}} = -g_{n,n}^{v^{\star}} - (-1)^{\ell} g_{1,1}^{v^{\ddagger}} + (-1)^{\ell} \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v}.$$
(108)

For any $v \in V$, equating the right-hand sides of (99) and (108) and then adding $(-1)^{\ell} g_{1,1}^{v^{t}}$ on both sides of the resulting equation shows

$$-(-1)^{\ell} g_{n,n}^{\nu} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{\nu^{\star}} = -g_{n,n}^{\nu^{\star}} + (-1)^{\ell} \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{\nu}, \quad (109)$$

which is equivalent to (100). Hence, the asserted equivalence of the conditions in the claim with $g \circ d_{\ell} = 0$ is true.

Step 2: Addendum. The proof of the addendum is split into two steps.

Step 2.1. For the following reasons, given any $w \in V$, if the conditions in the claim are satisfied by v = w, then they are also true for $v = w^*$. In the cases of (97) and (98) this intermediate claim is clearly true because replacing v by v^* transforms (97) into (98) and vice versa. Since (100) is equivalent to

$$g_{n,n}^{v^*} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^*} = (-1)^{\ell} (g_{n,n}^v + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^v)$$
(110)

and since multiplication with $(-1)^{\ell}$ has the same effect on (110) as replacing v by v^* identity (100) indeed holds for any $v = w^*$ as soon as it does for v = w.

As noted before, (100) being true for v = w implies that also (109) holds for v = w. Inserting (109) into the assumption (99) for v = w produces the identity (read for v = w)

$$g_{1,1}^{v^{\ddagger}} = -(-1)^{\ell} g_{1,1}^{v^{\ddagger}} - g_{n,n}^{v^{\star}} + (-1)^{\ell} \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v},$$

which on multiplication with $(-1)^{\ell}$ becomes a mere rearrangement of

$$g_{1,1}^{v^{t}} = -(-1)^{\ell} g_{1,1}^{v^{\ddagger}} - (-1)^{\ell} g_{n,n}^{v^{\star}} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v},$$

i.e., (99) for $v = w^*$. Hence, (97)–(100) for any $(j,i) \in [n]^{\times 2}$ with $(j,i) \neq (1,1)$ automatically also hold for $v = w^*$ if they do for v = w.

Step 2.2. What is left to prove after Step 2.1 is that, if the conditions in the claim are satisfied by some $v = w \in V$, then so they are by $v = w^{t}$.

It has already been noted that (97) being true for v = w and any $(j,i) \in [n]^{\times 2}$ with $(j,i) \neq (1,1)$ implies (106) for this v = w. Analogously, summing up (98) for (s,s) in place of (j,i) over all $s \in [n]$ with $s \notin \{1,n\}$ shows that (106) is also valid for $v = w^*$. In fact, since multiplying with $-(-1)^{\ell}$ has the same effect on (106) as replacing v by v^{\ddagger} it has thus been shown that (106) is true for any $v \in V$.

Instantiating (97) respectively (98) at (v, j, i) = (w, n, n) yields that

$$g_{n,n}^{v^{t}} = -(-1)^{\ell} g_{1,1}^{v^{\star}} - g_{1,1}^{v} - \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v}$$

is true for v = w and $v = w^*$. Inserting into this (106) for $v = w^{\text{t}}$ respectively $v = w^{\ddagger}$ proves

$$g_{n,n}^{v^{t}} = -(-1)^{\ell} g_{1,1}^{v^{\star}} - g_{1,1}^{v} + (-1)^{\ell} \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{\ddagger}}$$

for v = w and $v = w^*$. By multiplying with $(-1)^{\ell}$ and solving for $g_{1,1}^{v^*}$ it follows

$$g_{1,1}^{v^{\star}} = -(-1)^{\ell} g_{1,1}^{v} - (-1)^{\ell} g_{1,1}^{v^{t}} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{t}}$$

for v = w and $v = w^*$, respectively, i.e., exactly (99) for $v = w^t$ and $v = w^{\ddagger}$. Hence, (99) is true for any $v \in V$ by Step 2.1.

It has already been seen in Step 1.2 that (99) being true for any $v \in V$ implies that (100) holds for all $v \in V$, in particular for $v = w^{t}$.

Thus, all that still needs proving at this point is that (97) and (98) are true for $v = w^{t}$ and any $(j,i) \in [n]^{\times 2}$. For $(j,i) \neq (n,n)$ this is immediately clear by replacing (j,i) by $(\mathbf{r}(j),\mathbf{r}(i))$ in (98) respectively (97) and then multiplying with $-(-1)^{\ell}$.

Multiplying (99) for v = w respectively $v = w^*$ with $-(-1)^{\ell}$ and solving for $g_{n,n}^v$ shows

$$g_{n,n}^{v} = -(-1)^{\ell} g_{1,1}^{v^{\dagger}} - (g_{1,1}^{v^{\dagger}} - (-1)^{\ell} \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{\star}}),$$

whence it follows by using (106) for $v = w^*$ respectively v = w that

$$g_{n,n}^{v} = -(-1)^{\ell} g_{1,1}^{v^{\ddagger}} - (g_{1,1}^{v^{\ddagger}} + \delta_{\ell,3} \sum_{s=2}^{n-1} g_{s,s}^{v^{\ddagger}})$$

for v = w respectively $v = w^*$. Since this is precisely (97) respectively (98) for $(v, j, i) = (w^t, n, n)$, the proof is complete.

With Lemma 5.8 at hand it is time to distinguish the cases $\ell = 3$ and $4 \leq \ell$.

5.3.2. Third order. In the case of $\operatorname{Hom}_A(d_3, \mathbb{K}_{\varepsilon})$ there is still a little more work to do in order to determine the defect and rank.

Lemma 5.9. Any $g \in \text{Hom}_A(P_2, \mathbb{K}_{\varepsilon})$ satisfies $g \circ d_3 = 0$ if and only if for any $v \in V$, all in conjunction, for any $(j, i) \in [n]^{\times 2}$ with $(j, i) \notin \{(1, 1), (n, n)\},\$

$$g_{j,i}^{v^*} = g_{r(j),r(i)}^{v^*},\tag{111}$$

$$g_{j,i}^{v^{*}} = g_{\mathbf{r}(j),\mathbf{r}(i)}^{v} \tag{112}$$

and

$$g_{1,1}^{v^{\ddagger}} = g_{1,1}^{v^{\ddagger}} + g_{n,n}^{v} + \sum_{s=2}^{n-1} g_{s,s}^{v^{\star}}$$
(113)

as well as

$$g_{n,n}^{v^{\star}} = -\sum_{s=2}^{n-1} g_{s,s}^{v^{\star}} - \sum_{s=2}^{n} g_{s,s}^{v}$$
(114)

and

$$g_{1,1}^{v^{\star}} = \sum_{s=1}^{n} g_{s,s}^{v}.$$
 (115)

In fact, (111)–(115) for any $(j,i) \in [n]^{\times 2}$ with $(j,i) \notin \{(1,1), (n,n)\}$ are all satisfied by each $v \in V$ as soon as they are all satisfied by an arbitrary one $v \in V$.

Proof. Both the claimed equivalence and the addendum can be proved simultaneously by showing that for any fixed $v \in V$ the conditions (111)– (115) are satisfied for any $(j,i) \in [n]^{\times 2}$ with $(j,i) \notin \{(1,1), (n,n)\}$ if and only if the conditions (97)–(100) from Lemma 5.8 are satisfied for $\ell = 3$ and any $(j,i) \in [n]^{\times 2}$ with $(j,i) \neq (1,1)$.

Obviously, for $\ell = 3$ and $(j, i) \notin \{(1, 1), (n, n)\}$ the conditions (97) and (111) are identical, as are (98) and (112). Likewise, $\ell = 3$ provided, (99) and (113) agree, and so do (100) and (114). Thus, all that remains to be shown is that, assuming all of these, the equations (97) for (j, i) = (n, n), i.e.,

$$g_{n,n}^{v^{t}} = g_{1,1}^{v^{\star}} - \sum_{s=1}^{n-1} g_{s,s}^{v}, \qquad (116)$$

and (98) for (j, i) = (n, n), i.e.,

$$g_{n,n}^{v^{\ddagger}} = g_{1,1}^{v} - \sum_{s=1}^{n-1} g_{s,s}^{v^{\star}}, \qquad (117)$$

are together equivalent to (115).

Because $x_{n,n}^{v^{\dagger},2} = x_{n,n}^{v,2}$ and $x_{n,n}^{v^{\dagger},2} = x_{n,n}^{v^{\star},2}$ by Lemma 4.3 the assumption $\ell = 3$ requires $g_{n,n}^{v^{\dagger}} = g_{n,n}^{v}$ and $g_{n,n}^{v^{\dagger}} = g_{n,n}^{v^{\star}}$. Because the first of the latter two identities reveals that (116) and (115) are equivalent, it is clear that (116) and (117) imply (115) and that it suffices to prove that (115) requires (117).

And, indeed, by using (114) in the second and (115) in the fourth step it can be concluded that

$$g_{n,n}^{v^{*}} = g_{n,n}^{v^{*}}$$

$$= -\sum_{s=2}^{n-1} g_{s,s}^{v^{*}} - \sum_{s=2}^{n} g_{s,s}^{v}$$

$$= g_{1,1}^{v} - \sum_{s=2}^{n-1} g_{s,s}^{v^{*}} - \sum_{s=1}^{n} g_{s,s}^{v}$$

$$= g_{1,1}^{v} - \sum_{s=1}^{n-1} g_{s,s}^{v^{*}},$$

which thus concludes the proof.

Lemma 5.10. Both defect and rank of $\operatorname{Hom}_A(d_3, \mathbb{K}_{\varepsilon})$ are $2n^2 - 1$.

Proof. The domain $\operatorname{Hom}_A(P_2, \mathbb{K}_{\varepsilon}) = \operatorname{Hom}_A(\mathbb{K}C_2 \otimes_{\mathbb{K}} A, \mathbb{K}_{\varepsilon}) \cong \mathbb{K}C_2$ of $\operatorname{Hom}_A(d_3, \mathbb{K}_{\varepsilon})$ has dimension $|C_2| = 4n^2 - 2$ by Lemma 4.3. For a fixed $v \in V$, say v = u, the (addendum to) Lemma 5.9 characterizes membership of an arbitrary $g \in \operatorname{Hom}_A(P_2, \mathbb{K}_{\varepsilon})$ in the kernel of $\operatorname{Hom}_A(d_3, \mathbb{K}_{\varepsilon})$ by the equations (111)–(115) for $(j, i) \in [n]^{\times 2}$ with $(j, i) \notin \{(1, 1), (n, n)\}$.

These conditions are linearly independent for the following reasons. The variables $g_{j,i}^{v^{t}}$ and $g_{j,i}^{v^{t}}$ for $(j,i) \in [n]^{\times 2}$ with $(j,i) \notin \{(1,1), (n,n)\}$ appear only in the equations (111) and (112), which are obviously linearly independent. Hence, the question is reduced to whether (113)–(115) are linearly independent. But (113) features the variables $g_{1,1}^{v^{t}}$ and $g_{1,1}^{v^{t}}$ which appear neither in (114) nor in (115). And, likewise, (114) and (115) each have a variable, namely $g_{n,n}^{v^{\star}}$ respectively $g_{1,1}^{v^{\star}}$, which occurs in no other equation.

Since there are $(n^2 - 2) + (n^2 - 2) + 1 + 1 + 1 = 2n^2 - 1$ independent equations for $4n^2 - 2$ variables, the claim follows by the dimension formula.

5.3.3. Fourth and higher orders. For orders $\ell \in \mathbb{Z}_{\geq 0}$ with $4 \leq \ell$ the defect and rank of $\operatorname{Hom}_A(d_\ell, \mathbb{K}_{\varepsilon})$ is now easy to compute.

Lemma 5.11. Both defect and rank of $\operatorname{Hom}_A(d_\ell, \mathbb{K}_{\varepsilon})$ are $2n^2$ for each $\ell \in \mathbb{Z}_{\geq 0}$ with $4 \leq \ell$.

Proof. The domain $\operatorname{Hom}_A(P_{\ell-1}, \mathbb{K}_{\varepsilon}) = \operatorname{Hom}_A(\mathbb{K}C_{\ell-1} \otimes_{\mathbb{K}} A, \mathbb{K}_{\varepsilon}) \cong \mathbb{K}C_{\ell-1}$ of $\operatorname{Hom}_A(d_{\ell}, \mathbb{K}_{\varepsilon})$ has dimension $|C_{\ell-1}| = 4n$ by Lemma 4.3. For a fixed $v \in V$, say v = u, the (addendum to) Lemma 5.8 characterizes membership of an arbitrary $g \in \operatorname{Hom}_A(P_{\ell-1}, \mathbb{K}_{\varepsilon})$ in the kernel of $\operatorname{Hom}_A(d_{\ell}, \mathbb{K}_{\varepsilon})$ by the equations (97)–(100) for $(j, i) \in [n]^{\times 2}$ with $(j, i) \neq (1, 1)$.

These conditions are linearly independent. Indeed, for any $(j,i) \in [n]^{\times 2}$ with $(j,i) \neq (1,1)$ the variable $g_{j,i}^{v^{t}}$ appears only in the corresponding (97) and nowhere else in the system of equations. Likewise, for such

(j,i) the variable $g_{j,i}^{v^{t}}$ is found only in the respective (98) and nowhere else. Furthermore, in the form of $g_{1,1}^{v^{t}}$ and $g_{n,n}^{v^{\star}}$, each of (99) and (100) respectively exhibits a variable which appears in no other equation.

Since there are $(n^2-1)+(n^2-1)+1+1=2n^2$ independent equations for $4n^2$ variables, the claim follows by the dimension formula.

5.4. Synthesis. Of course, the result of the computation was already known in advance from [DFKS18] and [BFGKT23].

Proposition 5.12. *For any* $\ell \in \mathbb{Z}_{>0}$ *,*

$$\dim_{\mathbb{K}}(\operatorname{Ext}_{A}^{\ell}(\mathbb{K}_{\varepsilon},\mathbb{K}_{\varepsilon})) = \begin{cases} n^{2} & |\ell = 1\\ n^{2} - 1 & |\ell = 2\\ 1 & |\ell = 3\\ 0 & |4 \leq \ell. \end{cases}$$

Proof. Because $\operatorname{Ext}_{A}^{\ell}(\mathbb{K}_{\varepsilon},\mathbb{K}_{\varepsilon})$ is the quotient vector space of the kernel of $\operatorname{Hom}_{A}(d_{\ell+1},\mathbb{K}_{\varepsilon})$ by the image of $\operatorname{Hom}_{A}(d_{\ell},\mathbb{K}_{\varepsilon})$, the dimension of $\operatorname{Ext}_{A}^{\ell}(\mathbb{K}_{\varepsilon},\mathbb{K}_{\varepsilon})$ is given by the difference $d_{\ell+1} - r_{\ell}$ of the defect $d_{\ell+1}$ of $\operatorname{Hom}_{A}(d_{\ell+1},\mathbb{K}_{\varepsilon})$ and the rank r_{ℓ} of $\operatorname{Hom}_{A}(d_{\ell},\mathbb{K}_{\varepsilon})$. Because $r_{1} = 0$ and $d_{2} = r_{2} = n^{2}$ and $d_{3} = r_{3} = 2n^{2} - 1$ and $d_{\ell} = r_{\ell} = 2n^{2}$ if $4 \leq \ell$ by Lemmata 5.3, 5.6, 5.10 and 5.11 that proves the claim. \Box

6. DISCUSSION

The present article is a case study in computing homological invariants of easy quantum groups.

- \Box Reminder on easy quantum groups, invariants Section 6.1
- \Box Literature context and motivation Section 6.2
- \Box Conclusions and outlook Section 6.3
- \Box Various other comments and remarks Section 6.4

6.1. Quantum groups background. One main approach to quantizing the notion of groups was developed by Woronowicz in [Wor87; Wor91; Wor98], culminating in the definition of compact quantum groups (see Section 6.1.1). While there are no free compact quantum groups, the universal compact matrix quantum groups $U_n^+(Q)$ discovered by Van Daele and Wang in [DW96; Wan95] come close in spirit (see Section 6.1.2). A special case of those is the free unitary quantum group U_n^+ , the subject of the present article. It arises from the unitary group U_n by a process termed liberation by Banica and Speicher in [BS09], which makes U_n^+ a so-called easy quantum groups (see Section 6.1.3). The findings of this article have promising implications for the feasibility of computing the quantum group cohomology (see Section 6.1.4) of the discrete duals of easy quantum groups (see Section 6.2 and 6.3). 6.1.1. Compact quantum groups. Compact quantum groups as defined by Woronowicz in [Wor87; Wor91; Wor98] are (formally dual to) – not necessarily commutative – C^* -algebras with the same kind of extra structure as the group law of a compact group G would induce on the commutative C^* -algebra C(G) of continuous complex-valued functions over it. The exact definition is not important for the purposes of the present article. It is enough to know that there exists a full and essentially surjective-on-objects contravariant functor \mathcal{O} from the category of compact quantum groups to the category of CQG algebras introduced by Dijkhuizen and Koornwinder in [DK94].

A CQG algebra is a Hopf *-algebra which admits a positive faithful integral. And, a Hopf *-algebra is a complex Hopf algebra equipped with a conjugate-linear anti-multiplicative co-multiplicative involution, the star *. An integral h on a Hopf *-algebra A is positive and faithful if and only if $h(a^*a) > 0$ for any $a \in A$ with $a \neq 0$. In particular, by forgetting (the star and) the co-multiplication but not its co-unit ε any CQG algebra can be interpreted as an augmented algebra and comes with a natural (right) module structure \mathbb{C}_{ε} on \mathbb{C} . That is the only part of the compact quantum group structure that has appeared explicitly so far in the present article.

6.1.2. Universal compact matrix quantum groups. Particular objects of the category of compact quantum groups are Van Daele and Wang's universal compact $n \times n$ -matrix quantum groups $U_n^+(Q)$ defined in [DW96; Wan95]. The latter can be understood as limits in appropriate subcategories, which motivates the name. The indexing parameters nand Q can be any positive integer respectively any invertible $n \times n$ matrix with complex entries. The augmented algebra underlying the CQG algebra $\mathcal{O}(U_n^+(Q))$ is the universal complex algebra with $2n^2$ generators $E = \{u_{j,i}^{\circ}, u_{j,i}^{\bullet}\}_{i,j=1}^n$ subject to the relations consisting of for any $(j, i) \in [n]^{\times 2}$ the polynomials


$$\sum_{s=1}^{n} u_{j,s}^{\circ} u_{i,s}^{\bullet} - \delta_{j,i} 1, \qquad \sum_{p,s,t=1}^{n} Q_{t,s} Q_{p,i}^{-1} u_{t,j}^{\circ} u_{s,p}^{\bullet} - \delta_{j,i} 1, \\
\sum_{s=1}^{n} u_{s,j}^{\bullet} u_{s,i}^{\circ} - \delta_{j,i} 1, \qquad \sum_{p,s,t=1}^{n} Q_{j,t} Q_{s,p}^{-1} u_{t,s}^{\bullet} u_{i,p}^{\circ} - \delta_{j,i} 1$$
(118)

together with the morphism of complex algebras to \mathbb{C} with $u_{j,i}^{\circ} \mapsto \delta_{j,i}$ and $u_{j,i}^{\bullet} \mapsto \delta_{j,i}$ for any $(j,i) \in [n]^{\times 2}$. In the special case that Q is the identity $n \times n$ -matrix, $U_n^+(Q)$ is simply denoted by U_n^+ . Then, for any $(j,i) \in [n]^{\times 2}$ the second polynomials in the the two rows of (118) become $\sum_{s=1}^n u_{s,j}^{\circ} u_{s,i}^{\bullet} - \delta_{j,i} 1$ and $\sum_{s=1}^n u_{j,s}^{\circ} u_{i,s}^{\circ} - \delta_{j,i} 1$ respectively. Hence, in that case the relations are precisely R and the augmentation precisely ε from the Main result.


6.1.3. Easy quantum groups. Adding to the definition of $\mathcal{O}(U_n^+)$ the relations $u_{t,s}u_{j,i} - u_{j,i}u_{t,s}$ for any $\{(t,s), (j,i)\} \subseteq [n]^{\times 2}$ produces the CQG algebra of the unitary group U_n . Of course it must have been by the reverse process, removing those relations from the presentation

of $\mathcal{O}(U_n)$, that Van Daele and Wang arrived at the definition of U_n^+ in the first place. In [BS09], Banica and Speicher gave this method of quantization by straight-up removing rather than deforming relations the name *liberation*. With the modifications by Tarrago and Weber [TW18] Banica and Speicher's construction provides a way of exhibiting new examples of quantum subgroups of U_n^+ , called *easy*, by finding solutions to special combinatorics problems.


More precisely, if $\{ {}^{\Box}1, {}^{\Box}2, \ldots \}$ and $\{ {}_{\Box}1, {}_{\Box}2, \ldots \}$ are two disjoint copies of $\mathbb{Z}_{>0}$, the *upper* respectively *lower points*, then for any $(k, \ell) \in \mathbb{Z}_{\geq 0}$ a *partition* p from any string $\mathfrak{c} \in \{ \circ, \bullet \}^{\times k}$ of *upper colors* to any *lower colors* $\mathfrak{d} \in \{ \circ, \bullet \}^{\times \ell}$ is the set of equivalence classes, its *blocks*, of an equivalence relation on $\{ {}^{\Box}a, {}_{\Box}b | a \in [k], b \in [\ell] \}$, its *points*.



Given any partition $p: \mathfrak{c} \to \mathfrak{d}$, its *adjoint* $p^*: \mathfrak{d} \to \mathfrak{c}$ is obtained by exchanging the roles of upper and lower points, $p^* = \{\{ {}^{\Box}b | {}_{\Box}b \in \mathsf{B} \} \cup \{ {}_{\Box}a | {}^{\Box}a \in \mathsf{B} \} \}_{\mathsf{B} \in p}$.




For any $\{k_t, \ell_t\} \subseteq \mathbb{Z}_{\geq 0}$, any $\mathbf{c}_t \in \{\circ, \bullet\}^{\times k_t}$, any $\mathfrak{d}_t \in \{\circ, \bullet\}^{\times k_t}$ and any partition $p_t \colon \mathbf{c}_t \to \mathfrak{d}_t$ for each $t \in \{1, 2\}$ the *tensor product* $p_1 \otimes p_2$ is the partition $\mathbf{c}_1 \mathbf{c}_2 \to \mathfrak{d}_1 \mathfrak{d}_2$ obtained by horizontal concatenation, $p_1 \otimes p_2 \coloneqq p_1 \cup \{\{ \exists (k_1 + a) \mid a \in [k_2] \land \exists a \in \mathsf{B}\} \cup \{ \exists (\ell_1 + b) \mid b \in [\ell_2] \land \exists b \in \mathsf{B}\} \}_{\mathsf{B} \in p_2}$.




Finally, for any $\{k, \ell, m\} \subseteq \mathbb{Z}_{\geq 0}$, any $\mathfrak{c} \in \{\circ, \bullet\}^{\times k}$, $\mathfrak{d} \in \{\circ, \bullet\}^{\times \ell}$, and $\mathfrak{e} \in \{\circ, \bullet\}^{\times \mathfrak{e}}$ and any partitions $p: \mathfrak{c} \to \mathfrak{d}$ and $q: \mathfrak{d} \to \mathfrak{e}$ the composition $qp: \mathfrak{c} \to \mathfrak{e}$ of (q, p) is the vertical concatenation under forgetting of loops, $qp:=\{\mathsf{A} \in p \land \mathsf{A} \subseteq \{^{\circ}1, ^{\circ}2, \ldots\}\} \cup \{\mathsf{C} \in q \land \mathsf{C} \subseteq \{_{\circ}1, _{\circ}2, \ldots\}\} \cup \{\bigcup\{\mathsf{A} \cap \{^{\circ}1, ^{\circ}2, \ldots\} \mid \mathsf{A} \in p \land \exists j \in B: \ _{\circ}j \in \mathsf{A}\} \cup \bigcup\{\mathsf{C} \cap \{_{\circ}1, _{\circ}2, \ldots\} \mid \mathsf{C} \in q \land \exists i \in B: \ ^{\circ}i \in \mathsf{C}\}\}_{B \in s} \setminus \{\emptyset\}$, where s is the set of equivalence classes of the finest equivalence relation on $[\ell]$ coarser than the equivalence relation

with the classes $\{\{j \in [\ell] \land_{\square} j \in \mathsf{A}\}\}_{\mathsf{A} \in p} \setminus \{\emptyset\}$ and the equivalence relation with the classes $\{\{i \in [\ell] \land^{\square} i \in \mathsf{C}\}\}_{\mathsf{C} \in q} \setminus \{\emptyset\}$.



With those definitions, a *category of (two-colored) partitions* is any set of partitions which is closed under forming adjoints, tensor products and (where defined) composition and which contains the partitions



For each category of two-colored partitions there is an associated quantum subgroup of U_n^+ whose CQG algebra is obtained by adding to the definition of $\mathcal{O}(U_n^+)$ for any $\{k, \ell\} \subseteq \mathbb{Z}_{\geq 0}$, any $\mathfrak{c} \in \{\circ, \bullet\}^{\times k}$ and $\mathfrak{d} \in \{\circ, \bullet\}^{\times \ell}$, any partition $p: \mathfrak{c} \to \mathfrak{d}$ of the category and any $g \in [n]^{\times k}$ and $j \in [n]^{\times \ell}$ the relation

$$\sum_{i \in [n]^{\times \ell}} \zeta_{p, \ker(g^{\Box_{\Box}}i)} u^{\mathfrak{d}_{1}}_{j_{1}, i_{1}} u^{\mathfrak{d}_{2}}_{j_{2}, i_{2}} \dots u^{\mathfrak{d}_{\ell}}_{j_{\ell}, i_{\ell}}$$

$$- \sum_{h \in [n]^{\times k}} \zeta_{p, \ker(h^{\Box_{\Box}}j)} u^{\mathfrak{c}_{1}}_{h_{1}, g_{1}} u^{\mathfrak{c}_{2}}_{h_{2}, g_{2}} \dots u^{\mathfrak{c}_{k}}_{h_{k}, g_{k}},$$
(120)

where for any $g \in [n]^{\times k}$ and $j \in [n]^{\times \ell}$ the number $\zeta_{p,\ker(g^{\circ}_{\square}j)}$ is defined as 1 if and only if the equivalence relation embodied by p is finer than the equivalence relation with the classes $\{\{ {}^{\square}a | a \in [k] \land g_a = z\} \cup \{ {}_{\square}b | b \in [\ell] \land j_b = z\}\}_{z \in [n]} \setminus \{\emptyset\}$ and as 0 otherwise.

In particular, U_n^+ itself corresponds exactly to the smallest of all possible categories of partitions. The non-zero ones among the polynomials (120) of the partitions (119) are (up to a sign) the relations (118) (for Q being the identity).

While many other categories of partitions have been found in a collective effort by various authors (see [BS09; BCS10; Web13; RW14; RW16a; RW16b; TW18; Gro18; MW20; MW21a; MW21b; MW21c; Maa21]), the full classification is still open. And, in the vast majority of cases little to nothing is known about the easy quantum groups corresponding to these categories. Critically, it is unclear when two easy quantum groups are isomorphic, especially across different n. Homological invariants can help with making these distinctions.

6.1.4. *Quantum group cohomology*. One invariant which can possibly discriminate between at least some easy quantum groups is the quantum group cohomology of their duals (see [Bic17] for an introduction). The fact that in terminology this cohomology is attributed not to the easy

quantum groups but the discrete quantum groups (see [Van96]) arising as their generalized Pontryagin duals is immaterial. A coefficient module which can be chosen for any discrete quantum group is the complex numbers. It provides the broadest possible range for comparisons. Given any compact quantum group G, the quantum group cohomology with trivial coefficients of its discrete dual is defined as

$$H^{\cdot}(G) = \operatorname{Ext}_{\mathcal{O}(G)}^{\cdot}(\mathbb{C}_{\varepsilon}, \mathbb{C}_{\varepsilon}),$$

i.e., as the derived functor at \mathbb{C}_{ε} of the contravariant external hom functor $\operatorname{Hom}_{\mathcal{O}(G)}(\cdot, \mathbb{C}_{\varepsilon})$ from the category of (right) $\mathcal{O}(G)$ -modules to the category of complex vector spaces. Thus, in order to compute these invariants of compact quantum groups it is necessary to have a projective resolution of \mathbb{C}_{ε} in terms of right $\mathcal{O}(G)$ -modules.

6.2. Motivation. The present article provides a projective resolution of the trivial module of the CQG algebra of the easy quantum group U_n^+ and uses it to compute the cohomology of the discrete dual of U_n^+ (see Section 6.1 for definitions). Of course, before the work that led to this article had even begun, Baraquin, Franz, Gerhold, Kula and Tobolski had already found in [BFGKT23] such a resolution and thus managed to determine the third and higher orders of said cohomology after Das, Franz, Kula and Skalski had previously computed the first two orders in [DFKS18; DFKS23]. In fact, in much greater generality than the present article, the results of [BFGKT23] apply, to the cohomology of the dual of the universal compact $n \times n$ -matrix quantum group $U_n^+(Q)$ or, technically, the more general universal co-sovereign Hopf algebra $\mathcal{H}(F)$ from [Bic01] for generic F, i.e., any invertible scalar matrix F such that tr(F) = 0 if and only if $tr(F^{-1}) = 0$ and such that there exist no roots of unity q of order greater than 2 with $q^2 - \sqrt{\operatorname{tr}(F)\operatorname{tr}(F^{-1})}q + 1 = 0$. Actually, transcending the results of the present article even further, [BFGKT23] determines in addition the bi-algebra cohomology of $\mathcal{H}(F)$, also for generic F (although their results hold in greater generality by [Bic23]), and Hochschild cohomology with arbitrary one-dimensional coefficients. Finally, it must be emphasized that Baraquin, Franz, Gerhold, Kula and Tobolski's resolution is not only finite but in fact minimal, namely zero from the fourth order onwards, whereas the one presented here is infinite. Given that what the current article provides can be achieved in much greater generality with much better results, why even write it in the first place?

The article grew out of the attempt to continue the work of [Man23] and compute after the first also the second cohomology of the discrete duals of all easy quantum groups G. As demonstrated by Bichon, Franz and Gerhold in [BFG17] and then Das, Franz, Gerhold, Kula and Skalski in [DFKS18; DFKS23], at least for certain G, the standard resolution is good enough for this purpose (see Appendix B for an explanation of their method). However, after finding myself consistently failing with

this strategy, even for relatively uncomplicated G, I decided to look for smaller resolutions. An analysis of the techniques used for obtaining resolutions for the very few G for which this has been achieved, left me clueless as to how to adapt those to arbitrary G.

Researching instead general strategies for going from the presentation of an augemented algebra in terms of generators and relations to a projective resolution of the trivial module I learned about Anick resolutions (see [Ani86]) from [Mus23]. I wondered whether it was feasible to use those for my purpose. There is a theorem guaranteeing the existence of an Anick resolution for each augmented algebra and monomial order. That means, in principle, this is a resolution which could be used for any G. Of course, there is a price. Its differentials are defined recursively. Hence, with those resolutions (just like with the standard resolution) the difficulty lies in computing the cohomology from the resolution. In contrast, with the computations of quantum group cohomologies in the literature the challenge is to find a resolution in the first place and prove its exactness. Computing the cohomology from an Anick resolution would presumably get easier if the recursion could be solved, i.e., if there was a closed-form expression for the differentials. As far as I was able to ascertain, it is not known whether this is always possible. Eventually, I decided to try. Having no idea what to expect, it seemed prudent to test on the easy G for which the set of relations $\mathcal{O}(G)$ is the smallest, U_n^+ . And for U_n^+ , it turns out, there is a non-recursive formula for the differential, making the cohomology computation simple.

The present article is effectively me reporting on the outcome of that experiment. Although it computes a cohomology that is already known, it is ultimately aimed at determining new, unknown ones. While it provides an explicit resolution for a quantum group where a better one is already available, it does so by very different means. At this point in time the only easy quantum groups G the full cohomologies of whose duals have been determined are U_n^+ , the free orthogonal quantum group O_n^+ defined in [Van96] and the free symmetric quantum group S_n^+ of [Wan98] (see Appendix C for an analysis of the three computations in the literature). And, whereas it is unclear to me where to even start with the methods used in those cases, the strategy employed here is, at least in principle, immediately applicable to arbitrary easy G (see Section 6.3).

6.3. **Conclusions.** The present article is a case study, a single data point, anecdotal evidence. As such, very few rigorous inferences are possible. That is not its purpose. Rather, it is meant to generate ideas and suggest future research directions.

On the object level, it shows that for each $n \in \mathbb{Z}$ with $2 \leq n$ there exists a monomial order on the set E of generators in the standard presentation of the CQG algebra $\mathcal{O}(U_n^+)$ of the easy quantum group

 U_n^+ with respect to which the Anick resolution of the co-unit admits a closed-form, non-recursive expression. (This resolution is different from the one obtained by Baraquin, Franz, Gerhold, Kula and Tobolski in [BFGKT23].) And it shows that the Anick resolution is good enough to compute from it the (previously known) quantum group cohomology of the dual of U_n^+ in all orders.

On the meta level, the article suggests the following.

1) The most important conclusion to be drawn from these findings is that Anick resolutions might be a way to reliably compute the cohomologies of the discrete duals of easy quantum groups G. For any G the trivial $\mathcal{O}(G)$ -module has such a resolution. If a Gröbner basis of $\mathcal{O}(G)$ can be found, there is, in principle, nothing preventing the investigation of the cohomology by the method used here. The only question is whether there is a closed formula and whether the number of recursion steps needed to prove it is managable or not.

2) In particular, if one is just interested in the second cohomology, this approach might be worth considering. The alternative method employed so far (see Appendix B) uses the standard resolution, which is likely much larger than the Anick resolution already in low degrees.

3) The formulas for the differentials $(d_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ become periodic from $\ell = 4$ onwards. Additionally, the $(d_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ depend on n only in a very benign way. Probably it would have been possible to guess the formula for general n and ℓ knowing just the orders, say, $\ell \leq 6$ in the special cases $n \leq 5$. Perhaps the Anick resolutions of \mathbb{K}_{ε} for other G behave similarly. In that case it might be quicker to use a computer algebra system to compute the resolution for small ℓ and n first, then form a hypothesis for general ℓ and n and try to use the Gröbner basis to prove exactness, rather than solve the recursion.

6.4. Additional remarks. The final section offers some context for the choices made in the article and discusses potential modifications.

6.4.1. On the choice of monomial order. When working with U_n^+ or with Kac-type compact $n \times n$ -matrix quantum groups in general it is customary to consider besides the fundamental representation matrix uits transpose u^t , its conjugate \overline{u} and its adjoint u^{\dagger} . In contrast, rather than the usual $\{u, u^t, \overline{u}, u^{\dagger}\}$ the Main result (and the entire article) only references $V = \{u, u^t, u^\star, u^{\ddagger}\}$. The switch $(\overline{u}, u^{\dagger}) \rightsquigarrow (u^\star, u^{\ddagger})$ is due to the particular choice of monomial order for $E = \{u_{j,i}, u_{j,i}^*\}_{i,j=1}^n$.

the particular choice of monomial order for $E = \{u_{j,i}, u_{j,i}^*\}_{i,j=1}^n$. More precisely, it is the fact that $\{u_{j,i}\}_{i,j=1}^n = \{u_{j,i}^\circ\}_{i,j=1}^n$ was given the lexicographic order but $\{u_{j,i}^*\}_{i,j=1}^n = \{u_{j,i}^\circ\}_{i,j=1}^n$ the *opposite* of the lexicographic order, that motivates this perspective. This choice may or may not seem natural. But, importantly, equipping both $\{u_{j,i}^\circ\}_{i,j=1}^n$ and $\{u_{j,i}^\bullet\}_{i,j=1}^n$ with the lexicographic order would not have worked. **Remark 6.1.** *R* is *no* Gröbner basis of *J* with respect to the degreewise lexicographic extension of the total order on *E* defined by for any $\{c, \mathbf{d}\} \subseteq \{\circ, \bullet\}$ and any $\{(j, i), (t, s)\} \subseteq [n]^{\times 2}$,

$$\begin{split} u^{\mathfrak{d}}_{t,s} &\leq u^{\mathfrak{c}}_{j,i} \Leftrightarrow \quad (\mathfrak{d},\mathfrak{c}) = \bullet \circ \\ & \lor (\mathfrak{d},\mathfrak{c}) \in \{\bullet \bullet, \circ \circ\} \land (t,s) < (j,i). \end{split}$$

In particular, R is not a universal Gröbner basis of J.

Proof. Actually, Proposition 2.1 is only an excerpt of Bergman's diamond lemma from [Ber78]. The full version states, among other things, (although not in these words) that any orbit (in the sense of monoid actions) of the reductions by a Gröbner basis has exactly one fixed point. In particular, in order to refute that R is a Gröbner basis, it suffices to exhibit a polynomial $b \in \mathbb{K}\langle E \rangle$ and reductions φ_1 and φ_2 by R with $\varphi_1(b) \neq \varphi_2(b)$ such that both $\varphi_1(b)$ and $\varphi_2(b)$ are fixed points of the reductions by R.

If $b := u_{n,n}^{\circ} u_{n,n}^{\bullet} u_{n,n}^{\circ}$, then, on the one hand, reduction of b by $u_{n,n}^{\circ} u_{n,n}^{\bullet} + \sum_{t=1}^{n-1} u_{t,n}^{\circ} u_{t,n}^{\bullet} - 1$ produces

$$-\sum_{t=1}^{n-1} u_{t,n}^{\circ} u_{t,n}^{\bullet} u_{n,n}^{\circ} + u_{n,n}^{\circ},$$

which when reduced by $u_{t,n}^{\circ}u_{t,n}^{\bullet} + \sum_{s=1}^{n-1} u_{t,s}^{\circ}u_{t,s}^{\bullet} - 1$ for each $t \in [n-1]$ becomes

$$\sum_{s,t=1}^{n-1} u_{t,s}^{\circ} u_{t,s}^{\bullet} u_{n,n}^{\circ} - (n-2) u_{n,n}^{\circ}.$$
(121)

On the other hand, if reduced by $u_{n,n}^{\bullet}u_{n,n}^{\circ} + \sum_{t=1}^{n-1} u_{t,n}^{\bullet}u_{t,n}^{\circ} - 1$ the polynomial *b* takes on the form

$$-\sum_{t=1}^{n-1} u_{n,n}^{\circ} u_{t,n}^{\bullet} u_{t,n}^{\circ} + u_{n,n}^{\circ},$$

which after reduction by $u_{t,n}^\bullet u_{t,n}^\circ + \sum_{s=1}^{n-1} u_{t,s}^\bullet u_{t,s}^\circ - 1$ for each $t\in [n-1]$ reads

$$\sum_{s,t=1}^{n-1} u_{n,n}^{\circ} u_{t,s}^{\bullet} u_{t,s}^{\circ} - (n-2) u_{n,n}^{\circ}.$$
(122)

Since (121) and (122) are distinct and fixed under reductions by R the set R cannot be a Gröbner basis.

Of course, there may still exist monomial orders which are different from the one in the Main result and which exhibit even more desirable properties. But, if one is interested in quantum groups, an order which makes the set R encoding an essential part of what makes U_n^+ a quantum "group" seemed a reasonable choice.

6.4.2. Left modules, bi-modules. While the co-unit of $A = \mathcal{O}(U_n^+)$ is resolved as a right module in the Main result, with the obvious modifications the same method could produce a resolution in terms of left modules. However, a different question is whether there is also a version for bi-modules. While that is not directly addressed by Anick in [Ani86], there is of course the option of interpreting bi-modules over A

as, say, right modules over the enveloping algebra $A^{\mathrm{op}} \otimes A$ of A. The enveloping algebra can also be expressed as a universal algebra over \mathbb{K} . Any disjoint union $E_1 \cup E_2$ of two copies E_1 and E_2 of E could serve as a generating set. The relations $\bigcup_{k=0}^2 R_k$ would consist of one copy $R_2 = \{r((e_2)_{e_2 \in E_2}) | r \in R\}$ of $R \subseteq \mathbb{K} \langle E \rangle$ imposed on E_2 , another copy, but reversed, $R_1 = \{r^{\mathrm{op}}((e_1)_{e_1 \in E_1}) | r \in R\}$ of R for E_1 , and the demand $R_0 = \{e_1 - e_2 | (e_1, e_2) \in E_1 \times E_2\}$ that E_1 and E_2 commute. One may want to try to as monomial order for $E_1 \cup E_2$ the same order as in the Main Theorem on E_2 and the same or the opposite of that on E_1 together with the prescription that the maximum of E_1 is less than the minimum of E_2 . Probably there is a then a way of obtaining a Gröbner basis of $A^{\mathrm{op}} \otimes A$ from that of A. Perhaps a similar connection exists between the sets of chains of $A^{\mathrm{op}} \otimes A$ and A.

Be that as it may, depending on the intended application, this might not be the best approach, for example, to computing the Hochschild cohomology of A. That is because A is not just an algebra but has the structure of a Hopf algebra.

Remark 6.2. There exists a unique algebra morphism $\Delta : A \to A \otimes A$ such that for any $v \in V$ and $(j, i) \in [n]^{\times 2}$,

$$\Delta(v_{j,i}) = \sum_{s=1}^n \begin{cases} v_{j,s} \otimes v_{s,i} & | v \in \{u, u^\star\} \\ v_{s,i} \otimes v_{j,s} & | v \in \{u^t, u^\sharp\}. \end{cases}$$

 Δ turns A into a Hopf algebra with co-unit ε and with antipode S given by the unique algebra morphism $A \to A^{\text{op}}$ with for any $v \in V$ and $(j,i) \in [n]^{\times 2}$,

$$S(v_{j,i}) = v_{\mathbf{r}(j),\mathbf{r}(i)}^{\ddagger}$$

Proof. Well-knonw consequence of [DW96].

In consequence (see [Bic13, Proposition 2.1]), there are natural isomorphisms between the Hochschild cohomology of A with coefficients in a given bi-module X and $\operatorname{Tor}^{A}(\mathbb{K}_{\varepsilon}, X')$ or $\operatorname{Ext}^{i}_{A}(\mathbb{K}_{\varepsilon}, X'')$ for the left A-module X' and the right A-module X'' with actions given by




where \triangleright and \triangleleft are the left respectively right A-actions on X.

6.4.3. Yetter-Drinfeld modules. The chain complex constructed by Baraquin, Franz, Gerhold, Kula and Skalski is [BFGKT23] resolves \mathbb{K}_{ε} not just as an $\mathcal{O}(U_n^+)$ -module but in fact as a Yetter-Drinfeld module with respect to the co-action given by the unit 1 of $\mathcal{O}(U_n^+)$. It is only natural to ask whether the modules $(P_{\ell})_{\ell \in \mathbb{Z}_{>0}}$ from the Main result can

be equipped with co-actions which also make $(d_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ a resolution in co-modules.

A vector space V together with a right action \triangleleft and a right co-action γ of a Hopf algebra A with multiplication μ , co-multiplication Δ and antipode S is a right Yetter-Drinfeld module if



In particular, the Hopf algebra A itself becomes the *regular Yetter*-Drinfeld module if equipped with the co-action



More generally, given any right co-module T over A with co-action κ , the free right A-module $T \otimes A$ over T can be turned into the *free Yetter-Drinfeld module* over (T, κ) by means of the co-action




With respect to this co-action on $T \otimes A$ any morphism $d: T \otimes A \to V$ of right A-modules into any right Yetter-Drinfeld module V with coaction γ is already a morphism of co-modules if


$$\begin{array}{c}
 d & V_A \\
 T & 1 \\
 \kappa & T
 \end{array} \\
 K & T
 \end{array} =
 \begin{array}{c}
 V \gamma & A \\
 V \gamma & V \\
 d & A \\
 T & 1
 \end{array}$$
(125)

In particular, the co-unit ε of the Hopf algebra A is a morphism of co-modules with respect to the Yetter-Drinfeld co-action (123) on A. Beyond that the above proves the following about the objects of the Main result.

Remark 6.3. For any family $(\kappa_{\ell})_{\ell \in \mathbb{Z}_{>0}}$ such that for each $\ell \in \mathbb{Z}_{>0}$ the mapping κ_{ℓ} is a right co-action of A on $\mathbb{K}C_{\ell}$, such that



and such that for any $\ell \in \mathbb{Z}_{>0}$ with $2 \leq \ell$



the family $(d_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ becomes a resolution in co-modules if P_0 is considered the regular Yetter-Drinfeld module and P_{ℓ} the free Yetter-Drinfeld module over $(\mathbb{K}C_{\ell}, \kappa_{\ell})$ for any $\ell \in \mathbb{Z}_{>0}$.

Proof. As mentioned above, $d_0 = \varepsilon$ is a co-module morphism since P_0 has the co-action (123). For the same reason condition (126) is precisely what (125) means for d_1 . Likewise, (127) spells out (125) for d_ℓ for $\ell \in \mathbb{Z}_{>0}$ with $2 \leq \ell$ because both P_ℓ and $P_{\ell-1}$ are equipped with the co-action (124).

Perhaps a recursive algorithm can be developed for computing such a family $(\kappa_{\ell})_{\ell \in \mathbb{Z}_{>0}}$ like for $(d_{\ell})_{\ell \in \mathbb{Z}_{>0}}$ itself.

Appendix A. Relationship to the known resolution

This section answers the question of how the resolution from the Main result relates to the one previously found by Baraquin, Franz, Gerhold, Kula, Tobolski in [BFGKT23]. I.e., an explicit quasi-isomorphism is given. As before, for any $p \in \mathbb{K}\langle E \rangle$ the element p + J of A will be referred to as p. And this convention is extended to matrices of elements of A. On p. 20 of [BFGKT23] the following result about the universal co-sovereign Hopf algebra $\mathcal{H}(F)$ can be found verbatim.

"Theorem 5.2. Let $F = E^t E^{-1} \in \operatorname{GL}_n(\mathbb{C})$ be a generic asymmetry and $\mathcal{H} = \mathcal{H}(F)$. Then the following sequence

,

$$0 \to \begin{pmatrix} \mathcal{H} \\ \mathcal{H} \end{pmatrix} \stackrel{\Phi_3^{\mathcal{H}}}{\longrightarrow} \begin{pmatrix} M_n(\mathcal{H}) \\ M_n(\mathcal{H}) \end{pmatrix} \stackrel{\Phi_2^{\mathcal{H}}}{\longrightarrow} \begin{pmatrix} M_n(\mathcal{H}) \\ M_n(\mathcal{H}) \\ \mathcal{H} \end{pmatrix} \stackrel{\Phi_1^{\mathcal{H}}}{\longrightarrow} \begin{pmatrix} \mathcal{H} \\ \mathcal{H} \end{pmatrix} \stackrel{\varepsilon}{\longrightarrow} \mathbb{C}_{\varepsilon} \to 0$$

with the mappings

$$\Phi_{3}^{\mathcal{H}} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} -Fa + (EuE^{-t})b \\ F^{-1}va - Fb \end{pmatrix},$$

$$\Phi_{2}^{\mathcal{H}} \begin{pmatrix} A \\ B \end{pmatrix} = \begin{pmatrix} A + (E^{-1}u^{t}BE)^{t} \\ B + (v^{t}E^{-1}AE)^{t} \\ 0 \end{pmatrix},$$

$$\Phi_{1}^{\mathcal{H}} \begin{pmatrix} A \\ B \\ c \end{pmatrix} = \begin{pmatrix} \operatorname{tr}(-A + u^{t}B) + c \\ \operatorname{tr}(Ev^{t}E^{-1}A - B) - c \end{pmatrix}$$

for $a, b, c \in \mathcal{H}$, $A, B \in M_n(\mathcal{H})$, yields a projective resolution of the counit of \mathcal{H} ."

Expressed in a way closer to the language of the present article a special case of [BFGKT23, Theorem 5.2] might read as follows.

Proposition A.1. If A, ε and \mathbb{K}_{ε} are as in the Main result, if $\overline{\circ} := \bullet$ and $\overline{\bullet} := \circ$, if

$$\begin{split} C_0' &= \{z^{\circ}, z^{\bullet}\}, \\ C_1' &= \{b_{j,i}^{\circ}, b_{j,i}^{\bullet}, y\}_{i,j=1}^n, \\ C_2' &= \{a_{j,i}^{\circ}, a_{j,i}^{\bullet}\}_{i,j=1}^n, \\ C_3' &= \{x^{\circ}, x^{\bullet}\} \end{split}$$

are such that $|C'_0| = |C'_3| = 2$ and $|C'_1| = 2n^2 + 1$ and $|C'_2| = 2n^2$, if for any $\ell \in \mathbb{Z}$ with $-1 \leq \ell$,

$$P'_{\ell} := \begin{cases} \mathbb{K}_{\varepsilon} & |\ell = -1 \\ \mathbb{K}C'_{\ell} \otimes A & |0 \leq \ell \leq 3 \\ 0 & |4 \leq \ell \end{cases}$$

if $(d'_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ is such that for each $\ell \in \mathbb{Z}_{\geq 0}$ the mapping d'_{ℓ} is a morphism $P'_{\ell} \to P'_{\ell-1}$ of right A-modules and such that for any $(j, i) \in [n]^{\times 2}$ and any $\mathfrak{c} \in \{\circ, \bullet\}$,

$$\begin{aligned} d_0'(z^{\mathfrak{c}} \otimes 1) &= 1, \\ d_1'(b_{j,i}^{\mathfrak{c}} \otimes 1) &= z^{\overline{\mathfrak{c}}} \otimes u_{j,i}^{\overline{\mathfrak{c}}} - \delta_{j,i} z^{\mathfrak{c}} \otimes 1, \\ d_1'(y \otimes 1) &= (z^{\circ} - z^{\bullet}) \otimes 1, \\ d_2'(a_{j,i}^{\mathfrak{c}} \otimes 1) &= \sum_{t=1}^n b_{i,t}^{\overline{\mathfrak{c}}} \otimes u_{j,t}^{\overline{\mathfrak{c}}} + b_{j,i}^{\mathfrak{c}} \otimes 1, \\ d_3'(x^{\mathfrak{c}} \otimes 1) &= \sum_{s,t=1}^n a_{t,s}^{\overline{\mathfrak{c}}} \otimes u_{t,s}^{\overline{\mathfrak{c}}} - \sum_{s=1}^n a_{s,s}^{\mathfrak{c}} \otimes 1 \end{aligned}$$

and such that d'_{ℓ} is the zero morphism of right A-modules for each $\ell \in \mathbb{Z}_{>0}$ with $4 \leq \ell$, then the complex

$$\cdots \longrightarrow 0 \longrightarrow \mathbb{K}P'_3 \otimes A \xrightarrow{d'_3} \mathbb{K}P'_2 \otimes A \xrightarrow{d'_2} \mathbb{K}P'_1 \otimes A \xrightarrow{d'_1} \mathbb{K}P'_0 \otimes A \xrightarrow{d'_0} \mathbb{K}_{\varepsilon}$$

of right A-modules is exact, meaning that $(P'_{\ell}, d'_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ is a resolution of \mathbb{K}_{ε} in terms of free right A-modules.

It can then be checked that a chain map from the chain complex in the Main result can be defined as follows.

Proposition A.2. If $(f_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ is such that for any $\ell \in \mathbb{Z}_{\geq 0}$ the mapping f_{ℓ} is a morphism of right A-modules $P_{\ell} \to P'_{\ell}$, such that for any $(j, i) \in [n]^{\times 2}$

$$f_0(1) = \frac{1}{2} \left(z^\circ + z^\bullet \right) \otimes 1$$

and

$$f_1(x_{j,i}^u \otimes 1) = b^{\bullet}_{j,\mathbf{r}(i)} \otimes 1 - \frac{1}{2} y \otimes (u^{\circ}_{j,\mathbf{r}(i)} + \delta_{j,\mathbf{r}(i)} 1),$$

$$f_1(x_{j,i}^{u^{\star}} \otimes 1) = b^{\circ}_{\mathbf{r}(j),i} \otimes 1 + \frac{1}{2} y \otimes (u^{\bullet}_{\mathbf{r}(j),i} + \delta_{\mathbf{r}(j),i} 1)$$

and

$$f_2(x_{j,i}^u \otimes 1) = a_{i,j}^\circ \otimes 1 - \delta_{j,n} \delta_{i,n} \sum_{s=1}^{n-1} \sum_{t=1}^n a_{t,s}^\bullet \otimes u_{t,s}^\bullet,$$

$$f_2(x_{j,i}^{u^t} \otimes 1) = \sum_{s=1}^n a_{s,j}^\bullet \otimes u_{s,i}^\bullet - \delta_{j,n} \delta_{i,n} \sum_{s=1}^{n-1} a_{s,s}^\circ \otimes 1,$$

$$f_2(x_{j,i}^{u^\star} \otimes 1) = a_{r(i),r(j)}^\bullet \otimes 1 - \delta_{j,n} \delta_{i,n} \sum_{s=2}^n \sum_{t=1}^n a_{t,s}^\circ \otimes u_{t,s}^\circ,$$

$$f_2(x_{j,i}^{u^t} \otimes 1) = \sum_{s=1}^n a_{s,r(j)}^\circ \otimes u_{s,r(i)}^\circ - \delta_{j,n} \delta_{i,n} \sum_{s=2}^n a_{s,s}^\bullet \otimes 1,$$

and such that $f_{\ell} = 0$ for any $\ell \in \mathbb{Z}_{\geq 0}$ with $3 \leq \ell$, then $(f_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ is a chain map $(P_{\ell}, d_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}} x \to (P'_{\ell}, d'_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ of complexes of right A-modules.

A chain map in the converse direction is for example the following.

Proposition A.3. If $(f'_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ is such that for any $\ell \in \mathbb{Z}_{\geq 0}$ the mapping f'_{ℓ} is a morphism of right A-modules $P'_{\ell} \to P_{\ell}$, such that for any $(j,i) \in [n]^{\times 2}$

$$f'_0(z^\circ \otimes 1) = 1,$$

$$f'_0(z^\bullet \otimes 1) = 1$$

and

$$\begin{aligned} f_1'(b_{j,i}^{\circ} \otimes 1) &= x_{\mathbf{r}(j),i}^{u^*}, \\ f_1'(b_{j,i}^{\bullet} \otimes 1) &= x_{j,\mathbf{r}(i)}^{u}, \\ f_1'(y \otimes 1) &= 0 \end{aligned}$$

and

$$f_{2}'(a_{j,i}^{\circ} \otimes 1) = x_{i,j}^{u} \otimes 1 + \delta_{j,n} \delta_{i,n} \sum_{s=1}^{n-1} x_{s,s}^{u^{t}} \otimes 1,$$

$$f_{2}'(a_{j,i}^{\bullet} \otimes 1) = x_{r(i),r(j)}^{u^{\star}} \otimes 1 + \delta_{j,1} \delta_{i,1} \sum_{s=1}^{n-1} x_{s,s}^{u^{\ddagger}} \otimes 1$$

and

$$\begin{aligned} f_3'(x^{\circ} \otimes 1) &= \sum_{s=1}^n x_{\mathbf{r}(s),s}^{u^{\star}}, \\ f_3'(x^{\bullet} \otimes 1) &= \sum_{s=1}^n x_{\mathbf{r}(s),s}^{u}, \end{aligned}$$

and such that $f'_{\ell} = 0$ for any $\ell \in \mathbb{Z}_{\geq 0}$ with $4 \leq \ell$, then $(f'_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ is a chain map $(P'_{\ell}, d'_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}} \rightarrow (P_{\ell}, d_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ of complexes of right A-modules.

Composing the two chain maps in the two possible ways produces the maps below.

Proposition A.4. For any $\ell \in \mathbb{Z}_{\geq 0}$ and any $(j, i) \in [n]^{\times 2}$,

$$(f_0 \circ f'_0)(z^{\circ} \otimes 1) = (f_0 \circ f'_0)(z^{\bullet} \otimes 1) = \frac{1}{2}(z^{\circ} + z^{\bullet}) \otimes 1$$

and

$$(f_1 \circ f'_1)(b^{\circ}_{j,i} \otimes 1) = b^{\circ}_{j,i} \otimes 1 + \frac{1}{2} y \otimes (u^{\bullet}_{j,i} + \delta_{j,i}1), (f_1 \circ f'_1)(b^{\bullet}_{j,i} \otimes 1) = b^{\bullet}_{j,i} \otimes 1 - \frac{1}{2} y \otimes (u^{\circ}_{j,i} + \delta_{j,i}1), (f_1 \circ f'_1)(y \otimes 1) = 0$$

and $f_2 \circ f'_2 = \text{id}$ and $f_\ell \circ f'_\ell = 0$ for any $\ell \in \mathbb{Z}_{\geq 0}$ with $3 \leq \ell$.

Proposition A.5. For any $\ell \in \mathbb{Z}_{\geq 0}$, if $\ell \leq 1$, then $f_{\ell} \circ f'_{\ell} = \mathrm{id}$, if $\ell = 2$, then for any $(j,i) \in [n]^{\times 2}$, if $v \in \{u, u^{\star}\}$, then

$$(f_2 \circ f'_2)(x^v_{j,i} \otimes 1) = x^v_{j,i} \otimes 1 - \delta_{j,n} \delta_{i,n} (\sum_{s=2}^n \sum_{t=1}^n x^{v^*}_{s,t} \otimes v^*_{t,s} + \sum_{s=1}^{n-1} x^{v^{\ddagger}}_{s,s} \otimes (v^{\ddagger}_{n,n} - 1))$$

and, if $v \in \{u^{t}, u^{\ddagger}\}$, then

$$(f_2 \circ f'_2)(x^v_{j,i} \otimes 1)$$

$$= \sum_{s=1}^n x^{v^{\dagger}}_{\mathbf{r}(j),s} \otimes v^{\dagger}_{s,\mathbf{r}(i)} + \delta_{j,1} \sum_{s=1}^{n-1} x^{v^{\star}}_{s,s} \otimes v^{\star}_{\mathbf{r}(i),n} - \delta_{j,n} \delta_{j,n} \sum_{s=1}^{n-1} v^{t}_{s,s} \otimes 1$$
and, if $3 \leq \ell$, then $f_\ell \circ f'_\ell = 0$.

Finally, choices of chain homotopies from the compositions of the chain maps to the identity chain maps are given by the below maps.

Proposition A.6. If $(D'_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ is such that for any $\ell \in \mathbb{Z}_{\geq 0}$ the mapping D'_{ℓ} is a morphism of right A-modules $P'_{\ell-1} \to P'_{\ell}$, such that $D'_{0} = 0$, such that

$$D'_1(z^{\circ} \otimes 1) = -\frac{1}{2} y \otimes 1,$$

$$D'_1(z^{\bullet} \otimes 1) = \frac{1}{2} y \otimes 1,$$

and such $D'_{\ell} = 0$ for any $\ell \in \mathbb{Z}_{\geq 0}$ with $2 \leq \ell$, then $(D'_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ is a chain homotopy $(f_{\ell} \circ f'_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}} \to \text{id of chain maps of complexes of right A$ modules.

Proposition A.7. If $(D_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ is such that for any $\ell \in \mathbb{Z}_{\geq 0}$ the mapping D_{ℓ} is a morphism of right A-modules $P_{\ell-1} \to P_{\ell}$, such that $D'_0 = D'_1 = D'_2 = 0$ and such that for any $\ell \in \mathbb{Z}_{\geq 0}$ with $3 \leq \ell$, any $v \in V$ and any $(j, i) \in [n]^{\times 2}$, if $v \in \{u, u^*\}$, then

$$D_{\ell}(x_{j,i}^{v} \otimes 1) = (-1)^{\ell} \delta_{j,n} \delta_{i,n} \left(x_{n,1}^{v^{\star}} + \delta_{\ell,3} \sum_{s=2}^{n-1} x_{r(s),s}^{v^{\star}} \right) \otimes 1$$

and, if $v \in \{u^{t}, u^{\ddagger}\}$, then

$$D_{\ell}(x_{j,i}^{v} \otimes 1) = (-1)^{\ell} \left(x_{r(j),i}^{v^{\ddagger}} + \delta_{j,1} \delta_{i,n} x_{1,1}^{v^{\star}} \right) \otimes 1,$$

then $(D_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}}$ is a chain homotopy $(f'_{\ell} \circ f_{\ell})_{\ell \in \mathbb{Z}_{\geq 0}} \to \text{id of chain maps of complexes of right A-modules.}$

Taken together the chain maps and chain homotopies above thus constitute a quasi-isomorphism between the two chain complexes.

Appendix B. On using the standard resolution

For the interested reader this section explains the strategy used by Bichon, Franz and Gerhold in [BFG17] and Das, Franz, Kula and Skalski in [DFKS18; DFKS23] to compute the orders 1 and 2 of the quantum group cohomology of the discrete duals of certain easy quantum groups, including U_n^+ , from the standard resolution. (Those authors investigated these quantum groups not with the objective of discerning isomorphism classes of easy quantum groups. Rather, they were led by an interest in the Calabi-Yau property of [Gin06], which generalizes Poincaré duality, and the AC property of [FGT15], which guarantees the existence of an associated Schürmann triple.)

B.1. Standard resolution. Using the standard resolution in order to compute the quantum group cohomology of the dual of a compact quantum group G is equivalent to determining the *Hochschild cohomolo*gy (see [Hoc56]) of $\mathcal{O}(G)$ with coefficients in the bi-module ${}_{\varepsilon}\mathbb{C}_{\varepsilon}$ obtained by letting ε induce both actions on \mathbb{C} (see, e.g., [Bic13]). Given any bimodule X over any algebra A over any field \mathbb{K} , if for any $k \in \mathbb{Z}_{\geq 0}$ the differential ∂_{HS}^k is the linear map $\mathrm{Hom}_{\mathbb{K}}(A^{\otimes k}, X) \to \mathrm{Hom}_{\mathbb{K}}(A^{\otimes (k+1)}, X)$

iv

with for any $\psi \in \operatorname{Hom}_{\mathbb{K}}(A^{\otimes k}, X)$ and any $(a_0, a_1, \ldots, a_k) \in A^{\times k+1}$,

 $\partial_{\mathrm{HS}}^k \psi(a_0 \otimes a_1 \otimes \ldots \otimes a_k)$

$$=\sum_{i=0}^{k+1}(-1)^{i}\begin{cases}a_{0}\psi(a_{1}\otimes a_{2}\otimes\ldots\otimes a_{k})&|i=0\\\psi(a_{0}\otimes a_{1}\otimes\ldots\otimes a_{i-2}\otimes a_{i-1}a_{i}&|\\\otimes a_{i+1}\otimes a_{i+2}\otimes\ldots\otimes a_{k})&|0$$

then the k-th Hochschild cohomology $H^k_{\text{HS}}(A, X)$ of A with coefficients in X is the quotient vector space of the space

$$Z^{k}_{\mathrm{HS}}(A, X) = \{ \gamma \in \mathrm{Hom}_{\mathbb{K}}(A^{\otimes k}, X) \land \partial^{k}_{\mathrm{HS}}\gamma = 0 \}$$

of k-co-cycles by the space $B_{\rm HS}^k(A, X)$ of k-co-boundaries given by 0 if k = 0 and otherwise by

$$B^k_{\mathrm{HS}}(A, X) = \{ \partial^{k-1}_{\mathrm{HS}} \psi | \psi \in \mathrm{Hom}_{\mathbb{K}}(A^{\otimes (k-1)}, X) \},\$$

where $A^{\otimes 0}$ means \mathbb{K} and where $\operatorname{Hom}_{\mathbb{K}}$ is the internal hom functor of vector spaces. Specializing to $A = \mathcal{O}(G)$ and $X = {}_{\varepsilon}\mathbb{C}_{\varepsilon}$ gives the quantum group cohomology with trivial coefficients of the dual of G.

B.2. First order. The space $B^1_{HS}(A, X)$ is 0 then and computing the first cohomology thus boils down to determining $Z^1_{\text{HS}}(A, X)$ alone. That is possible to do for all easy quantum groups more or less simultaneously, as shown in [Man23]. The key is to use the following folk theorem (see, e.g., [KR17, Lemma 1.9]). On the direct sum $A \oplus X$ an algebra structure $A \boxplus_0 X$ with a useful property is defined by letting $(a_1, x_1)(a_2, x_2) := (a_1a_2, a_1x_2 + x_1a_2)$ for any $\{(a_1, x_1), (a_2, x_2)\} \subseteq A \oplus X$. Any $\psi \in \operatorname{Hom}_{\mathbb{K}}(A, X)$ satisfies $\partial^{1}_{\mathrm{HS}}\psi = 0$ if and only if the linear map (id, ψ) with $a \mapsto (a, \gamma(a))$ for any $a \in A$ is an algebra morphism $A \to A \boxplus_0 X$. If A is a universal algebra with generators E and relations $R \subseteq \mathbb{K}\langle E \rangle$, the map (id, ψ) is such an algebra morphism if and only if $r((e, \psi(e))_{e \in E}) = 0$ in $A \boxplus_0 X$ for any $r \in R$. That gives a characterization of $Z^{1}_{HS}(A, X)$ in terms of linear equations induced by R. Because in case $A = \mathcal{O}(G)$ the polynomials R are all of the form (120) the resulting equations can be predicted well enough to solve them in general, even for those easy quantum groups whose categories of partitions are still unknown.

B.3. Second order. Already in order 2 however, the space of coboundaries is not only non-zero but infinite-dimensional in general. Nonetheless the authors of [BFG17] and [DFKS18; DFKS23] managed to compute the cohomology. Three insights are necessary to understand their strategy. First, if any $\gamma \in Z^2_{\text{HS}}(A, X)$ is called *normalized* if $\gamma(1 \otimes 1) = 0$, then $H^2_{\text{HS}}(A, X)$ is also the quotient vector space of the normalized 2-co-cycles by the normalized 2-co-boundaries. Second, the authors of [BFG17] provided in their Lemma 5.4 (or [DFKS18, Lemma 4.1]) a key tool, which, if generalized, turns out to extend the above folk theorem: Given any normalized $\gamma \in Z^2_{\text{HS}}(A, X)$ the rule that $(a_1, x_1)(a_2, x_2) := (a_1a_2, a_1x_2 + x_1a_2 - \gamma(a_1 \otimes a_2))$ for any $\{(a_1, x_1), (a_2, x_2)\} \subseteq A \oplus X$ also defines an algebra structure $A \boxplus_{\gamma} X$ on $A \oplus X$. Now, for any $\psi \in \operatorname{Hom}_{\mathbb{K}}(A, X)$ such that $\partial^{1}_{\operatorname{HS}}\psi$ is normalized (which means $\psi(1) = 0$) the map (id, ψ) is an algebra morphism $A \to A \boxplus_{\gamma} X$ if and only if $\partial^1_{\mathrm{HS}} \psi = \gamma$. Thus, if A is a universal algebra with generators E and relations R any normalized $\gamma \in Z^2_{HS}(A, X)$ is a co-boundary if and only if there exists $\psi \in \operatorname{Hom}_{\mathbb{K}}(A, X)$ such that $\psi(1) = 0$ and $r((e, \psi(e))_{e \in E}) = 0$ in $A \boxplus_{\gamma} X$ for any $r \in R$. In fact, it can be seen that any such ψ is already uniquely determined by its values on E (generalizing [DFKS18, pp.23–25, after (\Leftarrow)]). Solving these equations characterizing when a normalized 2-co-cycle is a 2-coboundary can be understood as finding a vector space V and what Das, Franz, Kula and Skalski call a *defect map*, a linear map Δ from the space of normalized 2-co-cycles into V whose kernel is the space of normalized 2-co-boundaries (see the proof of [DFKS18, Theorem 4.6]). Computing $H^2_{\rm HS}(A, X)$ is then equivalent to determining the image of Δ . The third part of the strategy employed in [BFG17] and [DFKS18] consists in an ansatz for achieving just that. In the case where X is additionally equipped with an algebra structure compatible with the Aactions, for any $(\eta, \theta) \in Z^1_{\mathrm{HS}}(A, X)^{\times 2}$ a normalized 2-co-cycle is given by the so-called *cup product* $\eta \cup \theta$ defined by $a_1 \otimes a_2 \mapsto \eta(a_1)\theta(a_2)$ for any $\{a_1, a_2\} \subseteq A$. If any normalized 2-co-cycle is co-homologous to a linear combination of cup products, the image of Δ coincides with the span of the images of cup products under Δ . In the case $A = \mathcal{O}(G)$ and $X = {}_{\varepsilon}\mathbb{C}_{\varepsilon}$ for the G investigated in [BFG17] and [DFKS18], in particular for $G = U_n^+$, this turns out to be true (see [DFKS18, p. 26]).

B.4. Additional tools for the second order. This strategy succeeds also for certain other easy quantum groups G than the ones from [BFG17] and [DFKS18], as demonstrated by [Wen20]. When trying to carry it out for all easy G simultaneously, it may be useful to extend the strategy by two additional parts. First, the folk theorem already used twice can be employed a third time to address the question whether for a given G any normalized $\gamma \in Z^2_{HS}(A, X)$ is co-homologous to a linear combination of cup products. It namely implies that this is the case if and only if there exist $\psi \in \operatorname{Hom}_{\mathbb{K}}(A, X)$ with $\psi(1) = 0$ and $m \in \mathbb{Z}_{\geq 0}$ and $\{\eta_i, \theta_i\}_{i=1}^m \subseteq Z^1_{\mathrm{HS}}(A, X)$ such that, if $\varphi := \gamma - \sum_{i=1}^m \eta_i \cup \theta_i$, then (id, ψ) is an algebra morphism $A \to A \boxplus_{\varphi} X$. As before, if A is generated by E subject to relations R, this means one has to check whether $r((e, \psi(e))_{e \in E}) = 0$ in $A \boxplus_{\varphi} X$ for any $r \in R$. Solving those conditions likely requires a lot of knowledge about $Z^2_{\text{HS}}(A, X)$. Therefore, the second piece one might want to add to the strategy from [BFG17] and [DFKS18] is the following. The space $\{\psi \in \operatorname{Hom}_{\mathbb{K}}(A, X) \land \psi(1) = 0\}$ can be turned into an A-bi-module A|X via the left action \triangleright and right action

⊲ defined by $(a \triangleright \psi)(b) := a\psi(b)$ and $(\psi \triangleleft a)(b) := \psi(ba) - \psi(b)a$ for any element ψ and any $\{a, b\} \subseteq A$. By restricting the natural isomorphism Λ : Hom_K($A \otimes A, X$) → Hom_K($A, Hom_{K}(A, X)$) induced by the tensor hom adjunction it can be seen that the space of normalized elements of $Z^{2}_{HS}(A, X)$ is isomorphic to $Z^{1}_{HS}(A, A|X)$ (see [Hoc56]). Hence, the folk theorem can be applied a fourth and last time to conclude that for any normalized element γ of $Z^{2}_{HS}(A, X)$ the map (id, $\Lambda(\gamma)$) is an algebra morphism $A \to A \boxplus_0 A | X$. For A universal with generators Eand relations R that means $r((e, \Lambda(\gamma))_{e \in E}) = 0$ in $A \boxplus_0 A | X$ for any element r of the ideal generated by R.

APPENDIX C. OTHER METHODS

At this point in time the only easy quantum groups the cohomologies of whose duals have been determined are O_n^+ , S_n^+ and U_n^+ . For the interested reader this section summarizes how this was achieved in each case and what would be required in order to adapt the strategies to other easy quantum groups.

C.1. Free orthogonal quantum group O_n^+ . The free orthogonal quantum group O_n^+ introduced by Wang and Van Daele in [DW96] is easy by [BS09]. Its associated category of partitions is generated by the partitions (119) and



The cohomology of the dual of O_n^+ was computed by Collins, Härtel and Thom in [CHT09] directly from a projective resolution of the co-unit of $\mathcal{O}(O_n^+)$. And the latter was a hard-won achievement. According to Bemerkung 3.1.1 in Härtel's (German-language) PhD thesis [Här08] it was found by "computer-assisted guessing, using, besides lots of custom-made software, the applications Plural, Gap and Magma" (my translation).

In other words, it would presumably be difficult to reproduce this remarkable feat for each of the many easy quantum groups whose cohomologies want to be computed. At least it seems quite unclear how and where to start.


An alternative way to come up with the same resolution was later found by Bichon in [Bic13]. In fact, he provided a free resolution in terms of Yetter-Drinfeld modules of the trivial module over $\mathcal{B}(E)$, the symmetry hopf algebra of a non-degenerate bi-linear form introduced by Dubois-Violette and Launer in [DL90]. The key argument is to use monoidal equivalence of co-module categories between $\mathcal{B}(E)$ and $\mathcal{O}(SL_2(q))$ for specific q. That allows one to transport over a noncommutative Koszul resolution of the trivial $\mathcal{O}(SL_2(q))$ -module found

ALEXANDER MANG

by Hadfield and Krähmer in [HK05]. The monoidal equivalence used was discovered by Bichon in [Bic03]. It is obtained by constructing an explicit Hopf bi-Galois extension between $\mathcal{O}(SL_2(q))$ and $\mathcal{B}(E)$ for specific q, which suffices according to a theorem by Schauenburg from [Sch96]. If one is just interested in $\mathcal{O}(O_n^+)$, i.e., trivial E, one can also use the equivalence Banica constructed in [Ban96] by determining the simple co-modules of $\mathcal{O}(O_n^+)$. In particular, both proofs depend on the detailed understanding of $SL_2(q)$ that is available.

Adapting this strategy to easy quantum groups G other than O_n^+ thus has three prerequisites. First, a resolution of the trivial module over some hopf algebra \mathcal{A} as well as knowledge of, second, \mathcal{A} , and, third, $\mathcal{O}(G)$ sufficient to construct a monoidal equivalence or Hopf bi-Galois object. Unfortunately, besides the presentation of $\mathcal{O}(G)$ in terms of generators and relations granted by the associated category of partitions, nothing is known about $\mathcal{O}(G)$ in almost all cases G. Moreover, there is no reason why their co-module categories are necessary equivalent to a known category of co-modules of some \mathcal{A} or that, if so, a resolution is available of the trivial \mathcal{A} -module.

C.2. Free symmetric quantum group S_n^+ . In [BS09] it is recognized that the free symmetric quantum group S_n^+ or quantum permutation group defined by Wang in [Wan98] is easy, namely corresponds to the category of partitions generated by the partitions (119) and



The cohomology of the dual of S_n^+ was determined in [BFG17] by Bichon, Franz and Gerhold. Key to their proof is to establish that $\mathcal{O}(S_n^+)$ is Calabi-Yau of dimension 3 in the sense of [Gin06], provided $n \leq 4$. In fact, this is proved in greater generality for the quantum symmetry algebras of semi-simple measured algebras of finite dimension greater than 3 with normalizable tracial measure. And the proof of that crucially relies on a monoidal equivalence of co-module categories between the quantum symmetry algebra and $\mathcal{O}(PSL_2(q))$ proved by Mrozinski in [Mro15]. That in turn required detailed knowledge of the simple co-modules of both the quantum symmetry algebra and $\mathcal{O}(SO_3(q))$ for at least certain q.

Knowing that $\mathcal{O}(S_n^+)$ is Calabi-Yau enables Bichon, Franz and Gerhold to conclude that the cohomological dimension is 3. Hence, they only need to compute the cohomology up to the third order. The orders 1 and 2 are computed directly from the standard resolution. For the third order they use a duality between cohomology and homology implied by the Calabi-Yau property. In consequence, rather than compute the cohomology in the third order, it suffices to determine the homology in the first. This again is achieved via the standard resolution. Bichon, Franz and Gerhold's method thus strongly relies on a particular property of S_n^+ . How many further easy quantum groups G can be regarded as quantum symmetry groups of measured algebras is unclear. Of course, even if G is not a quantum symmetry group, it could still be Calabi-Yau. Though, verifying that might then require a further generalization of Mrozinski's result. Such an extension seems likely to demand significant knowledge about the simple co-modules of $\mathcal{O}(G)$. Sadly, though, besides a combinatorial approach to computing partial fusion rules in [FW16], figuring out the simple co-modules of $\mathcal{O}(G)$ for general easy G remains an open problem.

C.3. Free unitary quantum group U_n^+ . As was already mentioned, in [BFGKT23], Baraquin, Franz, Gerhold, Kula and Tobolski not only compute the cohomology of the dual of the easy quantum group U_n^+ . They give a free resolution of $\mathcal{H}(F)$ for generic F and determine its Hochschild cohomology with one-dimensional coefficients and its bialgebra cohomology.

Their proof builds on the finite free resolution of the co-unit of $\mathcal{B}(E)$ found by Bichon in [Bic13]. Baraquin, Franz, Gerhold, Kula and Tobolski prove that, if $F = E^{t}E$, then $\mathcal{H}(F)$ appears as a Hopf subalgebra of the co-product $\mathcal{B}(E) * \mathbb{CZ}_2$ of $\mathcal{B}(E)$ and the group algebra of the cyclic group \mathbb{Z}_2 of order two. Namely, $\mathcal{H}(F)$ is precisely the socalled glued free product $\mathcal{B}(E) * \mathbb{CZ}_2$ in the sense of [TW16] of $\mathcal{B}(E)$ and \mathbb{CZ}_2 . Already in [Ban97] Banica had observed (although not in those terms) that $\mathcal{O}(U_n^+)$ is the glued free product $\mathcal{O}(O_n^+) * \mathbb{CZ}$ of $\mathcal{O}(O_n^+)$ and the group algebra of the integers. Gromada later refined this result by narrowing \mathbb{Z} to \mathbb{Z}_2 in [Gro22]. It is also his study of the glued product that when combined with the monoidal equivalence between the comodules of $\mathcal{B}(E)$ and $\mathcal{O}(SL_2(q))$ from [Bic03] is the basis of the proof that $\mathcal{H}(F) \cong \mathcal{B}(E) * \mathbb{CZ}_2$.

The authors of [BFGKT23] then combine these two ingredients, the resolution for $\mathcal{B}(E)$ and the isomorphism $\mathcal{H}(F) \cong \mathcal{B}(E) \cong \mathbb{C}\mathbb{Z}_2$ as follows. They prove that a projective resolution of the trivial module over the co-product of two Hopf algebras can be obtained as the direct sum of projective resolutions of the co-factors. In that way they construct from the projective resolution for $\mathcal{B}(E)$ and the known one for $\mathbb{C}\mathbb{Z}_2$ a resolution for $\mathcal{B}(E) * \mathbb{C}\mathbb{Z}_2$. The fact that co-semi-simple Hopf algebras are faithfully flat over Hopf subalgebras, proved by Chirvasitu in [Chi14], enables them to apply the restriction of scalars functor to turn that free resolution for $\mathcal{B}(E) * \mathbb{C}\mathbb{Z}_2$ into a projective resolution for $\mathcal{B}(E) \cong \mathbb{C}\mathbb{Z}_2$ and thus for $\mathcal{H}(F)$. In fact, it turns out that the resolution so obtained consists of free modules which can be further upgraded to free Yetter-Drinfeld modules. Finally, by using the theory of Yetter-Drinfeld modules Baraquin, Franz, Gerhold, Kula and Tobolski manage to exchange the premise that $F = E^{t}E$ for the assumption that F is generic.

If taken as a template for easy G other than U_n^+ , the strategy would be to express $\mathcal{O}(G)$ as a Hopf subalgebra of $\mathcal{O}(H)$ for some compact quantum group H for which a projective resolution is available and then applying the restriction of scalars functors. More narrowly, $\mathcal{O}(H)$ could be a co-product of two Hopf algebras with known resolutions and $\mathcal{O}(G)$ could be a glued free product. In particular, such a presentation exists if the degree of reflection (in the sense of [Gro18; Gro22]) of the category associated with G is non-zero. Because the majority of already all so-called non-hyperoctahedral categories of partitions, which make up "three quarters" of all categories, have degree of reflection zero by [MW21c], a more general product construction would likely be in order for arbitrary G. The issue is that establishing such a subalgebra relationship in the first place requires detailed knowledge of both comodule categories. And, again, for general G, with very few exceptions, that is not available.

In summary, for any $G \in \{O_n^+, S_n^+, U_n^+\}$ the known resolution of the trivial $\mathcal{O}(G)$ -module either was guessed with computer assistance, is the standard resolution (in low degrees) or was found by establishing a monoidal equivalence of co-modules between $\mathcal{O}(G)$ and a Hopf algebra with known resolution. Of course, it was never an objective of any of the authors to compute the quantum group cohomology of the duals of arbitrary easy quantum groups G. If, however, one is interested in that, there would be significant difficulties to overcome trying to adapt these methods, most importantly, the lack of knowledge of the simple co-modules of $\mathcal{O}(G)$.

References

[Ani86]	David J Anick. "On the homology of associative algebras".
	In: Transactions of the American Mathematical Society
	296.2 (1986), pp. 641–659. DOI: 10.2307/2000383.
[Ban96]	Teodor Banica. "Théorie des représentations du groupe
	quantique compact libre $O(n)$ ". In: Comptes rendus de
	l'Académie des sciences. Série 1, Mathématique 322.3
	(1996), pp. 241-244. URL: https://arxiv.org/abs/
	math/9806063.
[Ban97]	Teodor Banica. "Le groupe quantique compact libre $U(n)$ ".
	In: Communications in mathematical physics 190.1 (1997),
	рр. 143–172. DOI: 10.1007/s002200050237.
[BCS10]	Teodor Banica, Stephen Curran, and Roland Speicher.
	"Classification results for easy quantum groups". In: Pa-
	cific journal of mathematics 247.1 (2010), pp. 1–26. DOI:
	10.2140/pjm.2010.247.1.

- [Ber78] George M Bergman. "The diamond lemma for ring theory". In: *Advances in mathematics* 29.2 (1978), pp. 178– 218. DOI: 10.1016/0001-8708(78)90010-5.
- [BFG17] Julien Bichon, Uwe Franz, and Malte Gerhold. "Homological properties of quantum permutation algebras". In: New York J. Math 23 (2017), pp. 1671–1695. URL: https: //www.emis.de/journals/NYJM/nyjm/j/2017/23-74p.pdf.
- [BFGKT23] Isabelle Baraquin, Uwe Franz, Malte Gerhold, Anna Kula, and Mariusz Tobolski. Free resolutions for free unitary quantum groups and universal cosovereign Hopf algebras. 2023. DOI: 10.48550/arXiv.2309.07767.
- [Bic01] Julien Bichon. "Cosovereign Hopf algebras". In: Journal of Pure and Applied Algebra 157.2 (2001), pp. 121–133. DOI: https://doi.org/10.1016/S0022-4049(00)00024-4.
- [Bic03] Julien Bichon. "The Representation Category of the Quantum Group of a Non-degenerate Bilinear Form". In: *Communications in Algebra* 31.10 (2003), pp. 4831–4851. DOI: 10.1081/AGB-120023135.
- [Bic13] Julien Bichon. "Hochschild homology of Hopf algebras and free Yetter–Drinfeld resolutions of the counit". In: *Compositio Mathematica* 149.4 (2013), pp. 658–678. DOI: 10.1112/S0010437X12000656.
- [Bic17] Julien Bichon. "Homological invariants for discrete quantum groups". In: Conf. Topological Quantum Groups and Harmonic Analysis. 2017, pp. 15–19. URL: https://lmbp. uca.fr/~bichon/Seoul.pdf.
- [Bic23] Julien Bichon. Bialgebra cohomology and exact sequences. 2023. DOI: https://doi.org/10.48550/arXiv.2309. 10434.
- [BS09] Teodor Banica and Roland Speicher. "Liberation of orthogonal Lie groups". In: Advances in Mathematics 222.4 (2009), pp. 1461–1501. DOI: 10.1016/j.aim.2009.06. 009.
- [Chi14] Alexandru Chirvasitu. "Cosemisimple Hopf algebras are faithfully flat over Hopf subalgebras". In: Algebra & Number Theory 8.5 (2014), pp. 1179–1199. DOI: 10.2140/ant. 2014.8.1179.
- [CHT09] Benoît Collins, Johannes Härtel, and Andreas Thom. "Homology of free quantum groups". In: Comptes Rendus Mathematique 347.5-6 (2009), pp. 271–276. DOI: 10.1016/ j.crma.2009.01.021.

- [DFKS18] Biswarup Das, Uwe Franz, Anna Kula, and Adam Skalski.
 "Lévy-Khintchine decompositions for generating functionals on algebras associated to universal compact quantum groups". In: Infinite Dimensional Analysis, Quantum Probability and Related Topics 21.03 (2018), p. 1850017. DOI: 10.1142/S0219025718500170.
- [DFKS23] Biswarup Das, Uwe Franz, Anna Kula, and Adam Skalski.
 "Second cohomology groups of the Hopf*-algebras associated to universal unitary quantum groups". In: Annales de l'Institut Fourier 73.2 (2023), pp. 479–509. DOI: 10.5802/aif.3527.
- [DK94] Mathijs S Dijkhuizen and Tom H Koornwinder. "CQG algebras: a direct algebraic approach to compact quantum groups". In: Letters in Mathematical Physics 32.4 (1994), pp. 315–330. DOI: 10.1007/BF00761142.
- [DL90] Michel Dubois-Violette and Guy Launer. "The quantum group of a non-degenerate bilinear form". In: *Physics Letters B* 245.2 (1990), pp. 175–177. DOI: 10.1016/0370–2693(90)90129–T.
- [DW96] Alfons van Daele and Shuzhou Wang. "Universal quantum groups". In: *International Journal of Mathematics* 7.02 (1996), pp. 255–263. DOI: 10.1142/S0129167X96000153.
- [FGT15] Uwe Franz, Malte Gerhold, and Andreas Thom. "On the Lévy-Khinchin decomposition of generating functionals". In: Communications on Stochastic Analysis 9.4 (2015), p. 6. DOI: 10.31390/cosa.9.4.06.
- [FW16] Amaury Freslon and Moritz Weber. "On the representation theory of partition (easy) quantum groups". In: Journal für die reine und angewandte Mathematik (Crelles Journal) 2016.720 (2016), pp. 155–197. DOI: 10.1515/ crelle-2014-0049.
- [Gin06] Victor Ginzburg. "Calabi-yau algebras". In: arXiv preprint math/0612139 (2006). DOI: 10.48550/ARXIV.MATH/ 0612139.
- [Gro18] Daniel Gromada. "Classification of globally colorized categories of partitions". In: Infinite Dimensional Analysis, Quantum Probability and Related Topics 21.04 (2018), p. 1850029. DOI: 10.1142/S0219025718500297.
- [Gro22] Daniel Gromada. "Gluing compact matrix quantum groups". In: Algebras and Representation Theory 25.1 (2022), pp. 53– 88. DOI: 10.1007/s10468-020-10010-0.
- [Här08] Johannes Härtel. "Reduktionssysteme zur Berechnung einer Auflösung der orthogonalen freien Quantengruppen $A_O(n)$ ". PhD thesis. Georg-August-Universität Göttingen, 2008. URL: 10.53846/goediss-2505.

xii

- [HK05] Tom Hadfield and Ulrich Krähmer. "Twisted Homology of Quantum SL(2)". In: *K*-Theory 4.34 (2005), pp. 327–360. DOI: 10.1007/s10977-005-3118-2.
- [Hoc56] Gerald Hochschild. "Relative homological algebra". In: Transactions of the American Mathematical Society 82.1 (1956), pp. 246–269. DOI: 10.2307/1992988.
- [KR17] David Kyed and Sven Raum. "On the L2-Betti numbers of universal quantum groups". In: Mathematische Annalen 369 (2017), pp. 957–975. DOI: 10.1007/s00208-017-1531-5.
- [Maa21] Laura Maaßen. "Representation categories of compact matrix quantum groups". PhD thesis. RWTH Aachen, 2021. URL: http://publications.rwth-aachen.de/ record/822228/files/822228.pdf.
- [Man23] Alexander Mang. First cohomology with trivial coefficients of all unitary easy quantum group duals. 2023. DOI: 10. 48550/arXiv.2305.12494.
- [Mor85] Ferdinando Mora. "Gröbner bases for non-commutative polynomial rings". In: International Conference on Applied Algebra, Algebraic Algorithms, and Error-Correcting Codes. Springer. 1985, pp. 353–362. DOI: 10.1007/3-540-16776-5_740.
- [Mro15] Colin Mrozinski. "Quantum automorphism groups and SO(3)-deformations". In: Journal of Pure and Applied Algebra 219.1 (2015), pp. 1–32. DOI: 10.1016/j.jpaa. 2014.04.006.
- [Mus23] Adya Musson-Leymarie. On Anick resolution: from the original setting to the language of non-commutative Groebner bases. 2023. DOI: 10.48550/arXiv.2307.06144.
- [MW20] Alexander Mang and Moritz Weber. "Categories of twocolored pair partitions part I: categories indexed by cyclic groups". In: *The Ramanujan Journal* 53 (2020), pp. 181– 208. DOI: 10.1007/s11139-019-00149-w.
- [MW21a] Alexander Mang and Moritz Weber. "Categories of twocolored pair partitions Part II: Categories indexed by semigroups". In: *Journal of Combinatorial Theory, Series A* 180 (2021), p. 105409. DOI: 10.1016/j.jcta.2021. 105409.
- [MW21b] Alexander Mang and Moritz Weber. "Non-hyperoctahedral categories of two-colored partitions part I: new categories". In: *Journal of Algebraic Combinatorics* 54 (2021), pp. 475–513. DOI: 10.1007/s10801-020-00998-5.
- [MW21c] Alexander Mang and Moritz Weber. "Non-hyperoctahedral Categories of Two-Colored Partitions Part II: All

Possible Parameter Values". In: *Applied Categorical Structures* 29 (2021), pp. 951–982. DOI: 10.1007/s10485-021-09641-1.

- [RW14] Sven Raum and Moritz Weber. "The combinatorics of an algebraic class of easy quantum groups". In: Infinite Dimensional Analysis, Quantum Probability and Related Topics 17.03 (2014), p. 1450016. DOI: 10.1142/ S0219025714500167.
- [RW16a] Sven Raum and Moritz Weber. "Easy quantum groups and quantum subgroups of a semi-direct product quantum group". In: *Journal of Noncommutative Geometry* 9.4 (2016), pp. 1261–1293. DOI: 10.4171/jncg/223.
- [RW16b] Sven Raum and Moritz Weber. "The full classification of orthogonal easy quantum groups". In: Communications in Mathematical Physics 341.3 (2016), pp. 751–779. DOI: 10.1007/s00220-015-2537-z.
- [Sch96] Peter Schauenburg. "Hopf bigalois extensions". In: *Communications in Algebra* 24.12 (1996), pp. 3797–3825. DOI: 10.1080/00927879608825788.
- [TW16] Pierre Tarrago and Moritz Weber. "Unitary Easy Quantum Groups: The Free Case and the Group Case". In: *International Mathematics Research Notices* 2017.18 (Aug. 2016), pp. 5710–5750. DOI: 10.1093/imrn/rnw185.
- [TW18] Pierre Tarrago and Moritz Weber. "The classification of tensor categories of two-colored noncrossing partitions".
 In: Journal of Combinatorial Theory, Series A 154 (2018), pp. 464–506. DOI: 10.1016/j.jcta.2017.09.003.
- [Van96] Alfons Van Daele. "Discrete quantum groups". In: *Journal* of Algebra 180.2 (1996), pp. 431–444. DOI: 10.1006/jabr. 1996.0075.
- [Wan95] Shuzhou Wang. "Free products of compact quantum groups". In: *Communications in Mathematical Physics* 167.3 (1995), pp. 671–692. DOI: 10.1007/BF02101540.
- [Wan98] S Wang. "Quantum Symmetry Groups of Finite Spaces".
 In: Commun. Math. Phys. 195.math. OA/9807091 (1998),
 pp. 195–211. DOI: 10.1007/s002200050385.
- [Web13] Moritz Weber. "On the classification of easy quantum groups". In: Advances in Mathematics 245 (2013), pp. 500–533. DOI: 10.1016/j.aim.2013.06.019.
- [Wen20] Alexander Wendel. "Hochschild cohomology of free easy quantum groups". Bachelor's Thesis. Saarland University, Apr. 2020.
- [Wor87] Stanisław L Woronowicz. "Compact matrix pseudogroups". In: Communications in Mathematical Physics 111.4 (1987), pp. 613–665. DOI: 10.1007/BF01219077.

[Wor91]	SL Woronowicz. "A remark on compact matrix quantum
	groups". In: letters in mathematical physics 21.1 (1991),
	рр. 35–39. DOI: 10.1007/ВF01221411.
[Wor98]	Stanisław I. Woronowicz "Compact quantum groups" In:

[Wor98] Stanisław L Woronowicz. "Compact quantum groups". In: Symétries quantiques (Les Houches, 1995) 845.884 (1998), p. 98. URL: https://www.impan.pl/~pmh/teach/ intro3/CQG3.pdf.

HAMBURG UNIVERSITY, MATHEMATICS DEPARTMENT, BUNDESSTRASSE 55, ROOM 316, 20146 HAMBURG, GERMANY, ALEXANDER.MANG@UNI-HAMBURG.DE Email address: alex@alexandermang.net URL: https://alexandermang.net