

THE BERGE-FÜREDI CONJECTURE ON THE CHROMATIC INDEX OF HYPERGRAPHS WITH LARGE HYPEREDGES

ALAIN BRETTO, ALAIN FAISANT, AND FRANÇOIS HENNECART

ABSTRACT. This paper is concerned with two conjectures which are intimately related. The first is a generalization to hypergraphs of Vizing's Theorem on the chromatic index of a graph and the second is the well-known conjecture of Erdős, Faber and Lovász which deals with the problem of coloring a family of cliques intersecting in at most one vertex. We are led to study a special class of uniform and linear hypergraphs for which a number of properties are established.

1. Introduction

In this paper, we address the problem of coloring hyperedges of a hypergraph \mathcal{H} , in the case where \mathcal{H} is loopless and linear [2, 3]. Two main conjectures have been proposed. The first one (the last chronologically) is a conjecture of several authors which is a generalization of Vizing's Theorem on the coloration of edges in a graph (cf. [28]). We postpone to Section 2 all the required definitions and notations.

Theorem 1.1 (VIZING'S THEOREM, 1964). *The chromatic index of any simple graph Γ satisfies*

$$q(\Gamma) \leq \Delta(\Gamma) + 1.$$

One can wonder if this result can be generalized to hypergraphs. Actually, this theorem suggested the following conjecture.

Conjecture 1.2 (GENERALIZED VIZING'S THEOREM). *Every linear hypergraph \mathcal{H} without loop verifies*

$$(1) \quad q(\mathcal{H}) \leq \Delta([\mathcal{H}]_2) + 1,$$

where $\Delta([\mathcal{H}]_2)$ is the maximum degree of the 2-section $[\mathcal{H}]_2$ of \mathcal{H} .

This conjecture was independently proposed, around 1985 by several authors, including Berge (cf. [3]), Füredi (cf. [16] where the conjecture is proven for intersecting hypergraphs) and Meyniel (unpublished).

The second one is closely related to the first and deals on the coloring of vertices of a graph formed by a families of n cliques each having n vertices and intersecting in at most one vertex.

Conjecture 1.3 (ERDŐS-FABER-LOVÁSZ). *Every graph Γ formed by a family of n cliques each having n vertices and intersecting in at most one vertex has chromatic number verifying $\chi(\Gamma) \leq n$.*

Date: March 12, 2024.

This conjecture can be implemented in hypergraph's language by considering cliques as hyperedges. Transposing it to the dual gives the following statement.

Conjecture 1.4 (ERDŐS-FABER-LOVÁSZ). *Every linear hypergraph \mathcal{H} without loop and having n vertices verifies $q(\mathcal{H}) \leq n$.*

It is easy to see that Conjecture 1.2 implies Conjecture 1.4 since $\Delta([\mathcal{H}]_2) \leq n-1$ when \mathcal{H} is linear. There are very few results on these conjectures, especially on Conjecture 1.2 (see [16, 3]).

In this article we solve the Berge-Füredi Conjecture 1.2 whenever the antirank is bounded from below by the square root of the number of vertices. We first provide a criterion ensuring that \mathcal{H} satisfies the desired bound (1) and use it to show our main result in regards to Conjecture 1.2 (cf. Theorem 1.6).

Theorem 1.5. *Let $\mathcal{H} = (V, E)$ be a linear hypergraph such that*

$$(2) \quad \forall e \in E, \exists x \in e \text{ such that } \deg_{\mathcal{H}}(x) \leq |e|.$$

Then

$$(3) \quad q(\mathcal{H}) \leq \Delta([\mathcal{H}]_2) + 1.$$

Note that if $\Delta(\mathcal{H}) \leq \text{ar}(\mathcal{H})$ then (2) is satisfied, whence $q(\mathcal{H}) \leq \Delta([\mathcal{H}]_2) + 1$ by Theorem 1.5. Thus the validity of Conjecture 1.2 depends actually on the remaining case $\Delta(\mathcal{H}) > \text{ar}(\mathcal{H})$.

In [24] the author shows that $q(\mathcal{H}) \leq |V|$ whenever $\text{ar}(\mathcal{H}) > |V|^{1/2}$. In [29] it is proven that it remains true under the weaker condition $\text{ar}(\mathcal{H}) \geq |V|^{1/2}$. We improve this result by showing the following statement.

Theorem 1.6. *Let $\mathcal{H} = (V, E)$ be a linear hypergraph such that $\text{ar}(\mathcal{H}) \geq |V|^{1/2}$.*

Then

$$q(\mathcal{H}) \leq \Delta([\mathcal{H}]_2) + 1.$$

The special case in Theorem 1.6 where $\text{ar}(\mathcal{H}) = |V|^{1/2}$ highlights an attractive class, denoted \mathcal{H}_k , of linear and k -uniform hypergraphs which will be studied separately (see beginning of Section 4 for detailed definition). We will obtain various remarkable properties in Theorem 3.6 and Section 4 (see Theorems 4.3 and 4.4) which will raise up several questions. This is the core of this paper.

2. Definitions, notations and general properties

Throughout this article, we shall use the notation $|S|$ for the cardinality of a finite set S . In what follows $\mathcal{H} = (V, E)$ denotes a hypergraph where $V(\mathcal{H}) = V$ is the set of vertices and $E(\mathcal{H}) = E$ is the set of hyperedges.

- \mathcal{H} is said to be *linear* if $|e \cap e'| \leq 1$ for any pair of distinct hyperedges $e, e' \in E$.
- \mathcal{H} is said to be *k -uniform* if $|e| = k$ for any $e \in E$.
- We denote by $\text{ar}(\mathcal{H})$ the *antirank* of \mathcal{H} , namely the minimum cardinality of hyperedges of \mathcal{H} . We have $\text{ar}(\mathcal{H}) \geq 2$ if and only if \mathcal{H} has no loop. If \mathcal{H} has no hyperedge then $\text{ar}(\mathcal{H})$ is fixed to be equal to ∞ . The *rank* of \mathcal{H} is defined by $r(\mathcal{H}) := \max\{|e|, e \in E\}$.
- For $x \in V$ let $\mathcal{H}(x) = \{e \in E : e \ni x\}$ denote the *star centered at x* .
- For $e \in E$, $\mathcal{H} \setminus e$ denotes the partial hypergraph $(V, E \setminus \{e\})$.

- The *degree of a vertex* x in \mathcal{H} is defined by $\deg_{\mathcal{H}}(x) = |\{e \in E : x \in e\}|$. The *minimum degree* is denoted by $\delta(\mathcal{H})$ and the *maximum degree* by $\Delta(\mathcal{H})$. We have the following identity

$$(4) \quad \sum_{x \in V} \deg_{\mathcal{H}}(x) = \sum_{e \in E} |e|,$$

which implies

$$(5) \quad |E| \leq \frac{|V|\Delta(\mathcal{H})}{\ar(\mathcal{H})}.$$

When \mathcal{H} is linear and if $\delta(\mathcal{H}) \geq 2$ and $\ar(\mathcal{H}) \geq 2$, we have the uniform upper bounds

$$(6) \quad \forall x \in V, \quad \deg_{\mathcal{H}}(x) \leq \frac{|V| - 1}{\ar(\mathcal{H}) - 1},$$

$$\forall e \in E, \quad |e| \leq \frac{|E| - 1}{\delta(\mathcal{H}) - 1}.$$

- \mathcal{H} is said *d-regular* if $\deg_{\mathcal{H}}(x) = d$ for any $x \in V$.
- The *degree of a hyperedge* e in \mathcal{H} is given by

$$d_{\mathcal{H}}(e) = |\{a \in E \setminus \{e\} : a \cap e \neq \emptyset\}|.$$

Note that this definition is different from that sometimes given in the literature (see for instance [4]). We have the following bounds:

$$(7) \quad d_{\mathcal{H}}(e) \leq |E| - 1 \quad \text{and} \quad \sum_{e \in E} d_{\mathcal{H}}(e) \leq |E|(|E| - 1).$$

Moreover when \mathcal{H} is linear we can develop $d_{\mathcal{H}}(e)$ as

$$d_{\mathcal{H}}(e) = \sum_{x \in e} \sum_{\substack{a \in E \setminus \{e\} \\ a \cap e = \{x\}}} 1 = \sum_{x \in e} \sum_{\substack{a \in E \setminus \{e\} \\ a \ni x}} 1$$

thus

$$(8) \quad d_{\mathcal{H}}(e) = \sum_{x \in e} (\deg_{\mathcal{H}}(x) - 1) \quad \text{and} \quad \sum_{e \in E} d_{\mathcal{H}}(e) = \sum_{x \in V} (\deg_{\mathcal{H}}(x) - 1) \deg_{\mathcal{H}}(x).$$

- The *2-section* of \mathcal{H} is the simple graph, denoted by $[\mathcal{H}]_2$, whose vertices are those of \mathcal{H} and for which two distinct vertices form an edge if and only if they both belong to a common hyperedge e of \mathcal{H} . For any $x \in V$ we have

$$(9) \quad \deg_{[\mathcal{H}]_2}(x) = \sum_{\substack{e \in E \\ e \ni x}} (|e| - 1),$$

giving the following lower bound on the maximum degree in $[\mathcal{H}]_2$

$$(10) \quad \Delta([\mathcal{H}]_2) \geq (\ar(\mathcal{H}) - 1) \Delta(\mathcal{H})$$

and, when \mathcal{H} is linear, the following identity

$$(11) \quad \sum_{x \in V} \deg_{[\mathcal{H}]_2}(x) = \sum_{e \in E} (|e|^2 - |e|).$$

- The *line-graph* of \mathcal{H} denoted by $L(\mathcal{H})$ is the simple graph whose vertices are hyperedges of \mathcal{H} and such that there is an edge between $e, e' \in E$, $e \neq e'$, if $e \cap e' \neq \emptyset$. We have $\deg_{L(\mathcal{H})}(e) = d_{\mathcal{H}}(e)$ for any $e \in E$.
- Let A be the incidence matrix of the hypergraph \mathcal{H} supposed without any isolated vertex and A^t be its transpose. The hypergraph whose incidence matrix is A^t is called the *dual* hypergraph of \mathcal{H} and denoted by $\mathcal{H}^* = (V^*, E^*)$. Clearly \mathcal{H}^* is without isolated vertex and $(\mathcal{H}^*)^* = \mathcal{H}$. We may notice that the second identity in (8) is related to (11) by duality. Note also that $\delta(\mathcal{H}) = \text{ar}(\mathcal{H}^*)$.
- A *k-coloring of hyperedges or hyperedge k-coloring* of a hypergraph \mathcal{H} is the assignment of one color from the set $\{1, 2, 3, \dots, k\}$ to every hyperedge of \mathcal{H} in such a way that no two intersecting hyperedges have the same color.
- The *chromatic index* of a hypergraph \mathcal{H} , denoted by $q(\mathcal{H})$, is the least integer k such that there exists a hyperedge k -coloring of \mathcal{H} . It is not difficult to see that (cf. [4])

$$\delta(\mathcal{H}) \leq \Delta(\mathcal{H}) \leq \Delta_0(\mathcal{H}) \leq q(\mathcal{H}),$$

where $\Delta_0(\mathcal{H})$ denotes the cardinality of the largest subset S of E such that $e, e' \in S \implies e \cap e' \neq \emptyset$.

- A hyperedge $e \in E$ is said *critical* in \mathcal{H} if $q(\mathcal{H} \setminus e) = q(\mathcal{H}) - 1$. \mathcal{H} is itself said *critical* if all its hyperedges are critical.
- Let Γ be a simple graph. A *k-coloring* of the vertices of Γ is an assignment of one color from the set $\{1, 2, 3, \dots, k\}$ to every vertex of the graph such that no two adjacent vertices have the same color. The smallest k such that Γ has a k -coloring, denoted by $\chi(\Gamma)$, is called the *chromatic number* of Γ . It is known (see for instance [6, Corollary 7.1.7, p. 235]) that $\chi(\Gamma) \leq \Delta(\Gamma) + 1$, thus we have the upper bound

$$(12) \quad q(\mathcal{H}) = \chi(L(\mathcal{H})) \leq \Delta(L(\mathcal{H})) + 1 \leq r(\mathcal{H})(\Delta(\mathcal{H}) - 1) + 1.$$

Hence if \mathcal{H} is k -uniform, that is $r(\mathcal{H}) = \text{ar}(\mathcal{H}) = k$, then by (10) the bound (3) holds whenever $\text{ar}(\mathcal{H}) \geq \Delta(\mathcal{H})$. By Theorem 1.5 and the remark that follows, the uniformity of \mathcal{H} is actually not required.

- The *strong chromatic number* of a hypergraph \mathcal{H} is the chromatic number of its 2-section. We denote it by $\chi(\mathcal{H})$ and we have $\chi(\mathcal{H}) = \chi([\mathcal{H}]_2)$. Notice that the (*weak*) *chromatic number* of a hypergraph \mathcal{H} (not used in this article) is the least integer k for which there exists a vertex k -coloring of \mathcal{H} such that any hyperedge of \mathcal{H} is not monochromatic.
- In a graph or a hypergraph, we say that a color i is *incident to* a vertex x if there is a (hyper-)edge with color i which contains x .
- Two linear hypergraphs $\mathcal{H} = (V, E)$ and $\mathcal{H}' = (V', E')$ without loop are said to be *isomorphic* if there exists a bijection f from V onto V' such that

$$\{x_1, \dots, x_t\} \in E \iff \{f(x_1), \dots, f(x_t)\} \in E'.$$

In this case it is clear that $q(\mathcal{H}) = q(\mathcal{H}')$.

- \mathcal{H} is said to be a *finite affine plane* if there exists $k \geq 2$ such that
 - $|V| = k^2$ and $\forall x \in V$, $\deg_{\mathcal{H}}(x) = k + 1$;
 - $|E| = k^2 + k$ and $\forall e \in E$, $|e| = k$;
 - $\forall x, y \in V$, $x \neq y$, there exists a unique hyperedge e such that $\{x, y\} \subset e$;

– $\forall a \in E$ and $x \in V \setminus a$, there exists a unique hyperedge e such that $x \in e$ and $e \cap a = \emptyset$.

Hence \mathcal{H} is a linear k -uniform and $(k + 1)$ -regular hypergraph. Moreover E can be partitioned into $k + 1$ sets of k disjoint hyperedges. This implies that $q(\mathcal{H}) = k + 1$.

When k is a prime power, let denote by \mathbb{F}_k the Galois field with cardinality k . By identifying lines with hyperedges the *field plane* \mathbb{F}_k^2 is a finite affine plane and denoted by \mathcal{A}_k .

For a same prime power k , it may exist several non-isomorphic finite affine planes (for instance when $k = 9$) with k^2 vertices. When k is not a prime power it is conjectured that there is no finite affine plane with k^2 vertices (known as the Prime Power Conjecture).

3. Proofs and remarks

3.1. When all hyperedges are large. The following result shows that the number hyperedges in a linear hypergraph hugely depends on the antirank in a anti-proportional way. Before we notice that (5) and (10) give together $|E| \leq \frac{n}{k^2-k} \Delta([\mathcal{H}]_2)$ (with the notation of the next lemma) without any restriction on $2 \leq k \leq n$.

Lemma 3.1. *Let $\mathcal{H} = (V, E)$ be a linear hypergraph with n vertices and $k = \text{ar}(\mathcal{H})$ be its antirank. If $k^2 > n$ then*

$$(13) \quad |E| \leq \frac{n(k-1)}{k^2-n},$$

and

$$(14) \quad \Delta(\mathcal{H}) \leq k.$$

Proof. We may assume that

$$(15) \quad |E| \geq \frac{n}{2k}$$

since otherwise (13) is plainly satisfied whenever $k^2 > n$. From (8), Cauchy inequality and (4), we infer

$$\sum_{e \in E} (d_{\mathcal{H}}(e) + |e|) = \sum_{x \in V} \text{deg}_{\mathcal{H}}(x)^2 \geq \frac{1}{n} \left(\sum_{x \in V} \text{deg}_{\mathcal{H}}(x) \right)^2 = \frac{1}{n} \left(\sum_{e \in E} |e| \right)^2$$

Thus

$$\sum_{e \in E} d_{\mathcal{H}}(e) \geq n(\lambda^2 - \lambda)$$

where $\lambda = \lambda(\mathcal{H}) = \frac{1}{n} \sum_{e \in E} |e|$. Since $\lambda \geq \frac{k|E|}{n}$, we have $\lambda \geq 1/2$ by (15), thus $\sum_{e \in E} d_{\mathcal{H}}(e) \geq k|E| \left(\frac{k|E|}{n} - 1 \right)$. We obtain by (7)

$$|E|(|E| - 1) \geq k|E| \left(\frac{k|E|}{n} - 1 \right)$$

giving

$$\left(\frac{k^2}{n} - 1 \right) |E| \leq k - 1.$$

This yields (13). Bound (14) follows from (6). \square

As a direct consequence of (13) we get by (10)

$$q(\mathcal{H}) \leq |E| \leq \frac{n}{k^2 - n} \times \frac{\Delta([\mathcal{H}]_2)}{\Delta(\mathcal{H})}$$

whenever $k^2 > n$. If in addition $n = o(k^2)$ then $q(\mathcal{H}) = o\left(\frac{\Delta([\mathcal{H}]_2)}{\Delta(\mathcal{H})}\right)$ as $n \rightarrow \infty$ (see also §3.3). Note also that (14) implies (2). So when $k^2 > n$, (3) will follow from Theorem 1.5. In order to deduce Theorem 1.6 from it, we essentially need to consider the equality case $n = k^2$ (see §3.3).

3.2. Proof of Theorem 1.5. We first show a simple but useful lemma related to critical hyperedges.

Lemma 3.2. *Let $\mathcal{H} = (V, E)$ be a hypergraph without loop and $e \in E$ be a critical hyperedge, that is $q(\mathcal{H} \setminus e) = q(\mathcal{H}) - 1$. Then $q(\mathcal{H}) - 1 \leq d_{\mathcal{H}}(e)$.*

Proof. Assume by contradiction that $q(\mathcal{H}) - 2 \geq d_{\mathcal{H}}(e)$. Let $k = q(\mathcal{H}) - 1$ and $C = \{c_1, \dots, c_k\}$ be the set of colors which are necessary to color the hyperedges of $\mathcal{H} \setminus e$. In order to color \mathcal{H} it remains to fix a permitted color for e . Since e intersects $d_{\mathcal{H}}(e) \leq q(\mathcal{H}) - 2$ hyperedges, there exists $c_i \in C$ which is not used for coloring those hyperedges. One then assigns the color c_i to e and we get $q(\mathcal{H}) \leq k$, a contradiction. \square

We continue with two easy properties in regard to Condition (2).

Lemma 3.3. *Let $\mathcal{H} = (V, E)$ be a hypergraph satisfying (2). Then*

$$(16) \quad \forall e \in E \text{ such that } |e| = \text{ar}(\mathcal{H}), \exists x \in e \text{ such that } \deg_{\mathcal{H}}(x) \leq \text{ar}(\mathcal{H}),$$

and for any $e' \in E$ the partial hypergraph $\mathcal{H}' = \mathcal{H} \setminus e' = (V, E \setminus \{e'\})$ also satisfies (2).

Proof. The implication (2) \implies (16) is plain.

Let $e \in E \setminus \{e'\}$. Since \mathcal{H} satisfies (2), there exists $x \in V$ such that $\deg_{\mathcal{H}}(x) \leq |e|$. We easily conclude since $\deg_{\mathcal{H}'}(x) \leq \deg_{\mathcal{H}}(x)$. \square

Remark 3.4. We stress that fact that Condition (2) is hereditary, namely

$$\mathcal{H} \text{ satisfies (2)} \implies \mathcal{H} \setminus e' \text{ satisfies (2)},$$

but it is no longer the case with Condition (16). This hereditary property will play a central role along the induction argument in the proof of Theorem 1.5.

Proof of Theorem 1.5. We argue by induction on the number of hyperedges.

If $|E| = 1$, clearly $q(\mathcal{H}) = 1 \leq \Delta([\mathcal{H}]_2) + 1$ and we are done.

Let $m \geq 1$ and $\mathcal{H} = (V, E)$ with $|E| = m + 1$ hyperedges.

Let e_0 be such that $|e_0| = \text{ar}(\mathcal{H})$ and $x_0 \in e_0$ such that $\deg_{\mathcal{H}}(x_0) \leq |e_0| = \text{ar}(\mathcal{H})$. Let $\mathcal{H}_0 = \mathcal{H} \setminus e_0$. Clearly \mathcal{H}_0 has m hyperedges and satisfies (2) since $\deg_{\mathcal{H}_0}(x) \leq \deg_{\mathcal{H}}(x)$. Therefore we may apply our induction hypothesis to \mathcal{H}_0 . We distinguish two cases.

- If $q(\mathcal{H}_0) = q(\mathcal{H}) - 1$ then by Lemma 3.2 and (10) we get

$$\begin{aligned}
q(\mathcal{H}) - 1 \leq d_{\mathcal{H}}(e_0) &= \sum_{\substack{x \in e_0 \\ x \neq x_0}} (\deg_{\mathcal{H}}(x) - 1) + \deg_{\mathcal{H}}(x_0) - 1 \\
&\leq (|e_0| - 1) \left(\max_{x \in e_0} \deg_{\mathcal{H}}(x) - 1 \right) + |e_0| - 1 \\
&= (\text{ar}(\mathcal{H}) - 1) \max_{x \in e_0} \deg_{\mathcal{H}}(x) \\
&\leq (\text{ar}(\mathcal{H}) - 1) \Delta(\mathcal{H}) \\
&\leq \Delta([\mathcal{H}]_2).
\end{aligned}$$

- Otherwise $q(\mathcal{H}_0) = q(\mathcal{H})$ and by induction hypothesis we get

$$q(\mathcal{H}) = q(\mathcal{H}_0) \leq \Delta([\mathcal{H}_0]_2) + 1 \leq \Delta([\mathcal{H}]_2) + 1.$$

This finishes the proof of Theorem 1.5. \square

3.3. Proof of Theorem 1.6. If $\mathcal{H} = (V, E)$ satisfies (2) then by Theorem 1.5 we infer the desired result. Assume the contrary, that is there exists $e_0 \in E$ such that

$$(17) \quad \forall x \in e_0 \quad \deg_{\mathcal{H}}(x) \geq |e_0| + 1 \geq \text{ar}(\mathcal{H}) + 1 = k + 1$$

where

$$(18) \quad k = \text{ar}(\mathcal{H}).$$

Our assumption on the antirank of \mathcal{H} can be rewritten as

$$(19) \quad k \geq |V|^{1/2}.$$

We shall use the notation $S = \{a \in E : a \cap e_0 = \emptyset\}$ the set of hyperedges which are *parallel* to e_0 and let $s = |S|$. Then

$$(20) \quad |E| = d_{\mathcal{H}}(e_0) + 1 + s.$$

Step 1: let $x \in e_0$. By (17), (18) and (19) we have following inequalities:

$$\begin{aligned}
k^2 \geq |V| &\geq \deg_{[\mathcal{H}]_2}(x) + 1 = \sum_{\substack{e \in E \\ e \ni x}} (|e| - 1) + 1 = \sum_{\substack{e \in E \setminus \{e_0\} \\ e \ni x}} (|e| - 1) + |e_0| \\
&\geq (\deg_{\mathcal{H}}(x) - 1)(k - 1) + k \geq k(k - 1) + k = k^2.
\end{aligned}$$

This implies that the above inequalities are all equalities. This gives

- $|V| = k^2$,
- $\deg_{[\mathcal{H}]_2}(x) = k^2 - 1$,
- $\deg_{\mathcal{H}}(x) = k + 1$,

We infer

$$(21) \quad d_{\mathcal{H}}(e_0) = \sum_{x \in e_0} (\deg_{\mathcal{H}}(x) - 1) = k^2,$$

and

$$(22) \quad |E| \geq d_{\mathcal{H}}(e_0) + 1 = k^2 + 1.$$

Step 2: we prove that \mathcal{H} is k -uniform and that $\forall y \in V \setminus e_0, \deg_{\mathcal{H}}(y) \geq k$.

- $|e_0| = k$ by (17), since $\deg_{\mathcal{H}}(x) = k + 1$ for $x \in e_0$.
- Let $a \in E \setminus (S \cup \{e_0\})$. Then $a \cap e_0 = \{x\}$ for some vertex $x \in e_0$.
As in Step 1, we may again write

$$k^2 - 1 = \deg_{[H]_2}(x) = \sum_{\substack{e \in E \\ e \ni x}} (|e| - 1) = \sum_{\substack{e \in E \setminus \{a\} \\ e \ni x}} (|e| - 1) + |a| - 1$$

thus

$$k^2 - 1 \geq (\deg_{\mathcal{H}}(x) - 1)(k - 1) + |a| - 1 = k(k - 1) + |a| - 1.$$

This implies $|a| \leq k$ yielding $|a| = k$ by (18).

- Let $e_0 = \{x_0, x_1, \dots, x_{k-1}\}$. Then for any $0 \leq i \leq k - 1$, we have $\deg_{\mathcal{H}}(x_i) = k + 1$. We let $\mathcal{H}(x_i) = \{e_0, e_{i,1}, \dots, e_{i,k}\}$ be the star centered at x_i , $0 \leq i \leq k - 1$. From above $|e_{i,j}| = k$, $1 \leq j \leq k$, thus for any $0 \leq i \leq k - 1$, we have

$$\left| e_0 \cup \bigcup_{j=1}^k e_{i,j} \right| = \sum_{j=1}^k (|e_{i,j}| - 1) + |e_0| = k(k - 1) + k = k^2 = |V|,$$

giving

$$(23) \quad V = e_0 \cup \bigcup_{j=1}^k e_{i,j}, \quad 0 \leq i \leq k - 1.$$

This implies that any vertex y in $V \setminus \{e_0\}$ is adjacent to x_i for any $0 \leq i \leq k - 1$. Hence for such y there exist distinct hyperedges a_i , $0 \leq i \leq k - 1$, such that both x_i and y are in a_i . This gives $\deg_{\mathcal{H}}(y) \geq k$.

- Let $a \in S$. We see by (23) that any vertex of a is in some $e_{0,j} \setminus \{x_0\}$, $1 \leq j \leq k$. Since \mathcal{H} is linear, we have $|a \cap e_{0,j}| \leq 1$, $1 \leq j \leq k$. We obtain $|a| \leq k$ and finally $|a| = k$ by (18).

We infer that \mathcal{H} is k -uniform.

Step 3: let $a, b \in S$, $a \neq b$. If $y \in a \cap b$, then y is adjacent to each x_i , $0 \leq i \leq k - 1$ through k many hyperedges. Therefore considering in addition a and b , it follows that y is contained in $k + 2$ many hyperedges, a contradiction. Hence $a \cap b = \emptyset$. In other words two hyperedges parallel to e_0 are indeed *parallel*.

Since \mathcal{H} is k -uniform we have $k^2 = |V| \geq |e_0| + \sum_{a \in S} |a| = k + ks$ hence $0 \leq s \leq k - 1$. By (20) and (21) we infer

$$|E| = d_{\mathcal{H}}(e_0) + 1 + s = k^2 + s + 1.$$

Step 4: the partial hypergraph $\tilde{\mathcal{H}} = (V, E \setminus S)$ belongs to the class \mathcal{H}_k of all hypergraphs $\tilde{\mathcal{H}} = (\tilde{V}, \tilde{E})$ satisfying

- $\tilde{\mathcal{H}}$ is linear and k -uniform,
- $|\tilde{V}| = k^2$,
- $|\tilde{E}| = k^2 + 1$,
- and there exists $e_0 \in \tilde{E}$ such that
 - $\forall x \in e_0$, $\deg_{\tilde{\mathcal{H}}}(x) = k + 1$,
 - $\forall x \in V \setminus \{e_0\}$, $\deg_{\tilde{\mathcal{H}}}(x) \geq k$.

If we have a hyperedge coloring of $\tilde{\mathcal{H}}$, then assigning to all hyperedges in S the same color already assigned to e_0 provides a hyperedge coloring of \mathcal{H} . Thus $q(\mathcal{H}) = q(\tilde{\mathcal{H}})$. It remains to show that $q(\tilde{\mathcal{H}}) \leq \Delta([\mathcal{H}]_2) + 1$. This will follow from Theorem 3.6 iii) below. Note that it is possible to get this bound by a shorter argument but instead we make the choice to study more widely the class \mathcal{H}_k .

Before stating and proving Theorem 3.6 we outline *geometric* properties for hypergraphs of \mathcal{H}_k .

Proposition 3.5. *Let $k \geq 2$, $\mathcal{H} = (V, E) \in \mathcal{H}_k$, $e_0 \in E$ satisfy (17) and $e \in E \setminus \{e_0\}$. Then*

- i) *We have $d_{\mathcal{H}}(e_0) = k^2$ and for any $a \in E$, $a \cap e_0 \neq \emptyset$. In other words there is no hyperedge parallel to e_0 .*
- ii) *There exists $x_0 \in e_0$ such that $e \cap e_0 = \{x_0\}$.*
- iii) *For any $x \in e_0 \setminus \{x_0\}$ there exists a unique hyperedge $a \in \mathcal{H}(x) \setminus \{e_0\}$ such that $a \cap e = \emptyset$.*
- iv) *Assume $k \geq 3$. It may exist $x \in V \setminus (e_0 \cup e)$ and distinct hyperedges a, a' such that $a \cap e = a' \cap e = \emptyset$ and $a \cap a' = \{x\}$.*
- v) *It is not true in general that $a \cap e = a' \cap e = \emptyset \implies a \cap a' = \emptyset$.*

Proof. i) By (21) and since $|E| = k^2 + 1$, all hyperedges intersect e_0 .

ii) follows from i).

iii) Existence: otherwise for any hyperedge $a \neq e_0$ and containing x , we have $a \cap e = \{y_a\}$ for some vertex $y_a \in e \setminus \{x_0\}$. Since \mathcal{H} is linear the mapping $a \in \mathcal{H}(x) \setminus \{e_0\} \mapsto y_a \in e \setminus \{x_0\}$ is injective. But $|\mathcal{H}(x)| = \deg_{\mathcal{H}}(x) = k + 1$, hence $|e| \geq k + 1$, a contradiction.

Uniqueness: let a satisfy the desired conclusion. Then $a \cap e_0 = \{x\}$ and $a \cap e = \emptyset$. By ii), each of the $k - 1$ many vertices of $e \setminus \{x_0\}$ lies in exactly one of the $k - 1$ many hyperedges of $\mathcal{H}(x) \setminus \{e_0, a\}$. This gives a mapping $y \in e \setminus \{x_0\} \mapsto a_y \in \mathcal{H}(x) \setminus \{e_0, a\}$ which is injective by linearity of \mathcal{H} . Since $|e \setminus \{x_0\}| = k - 1 = |\mathcal{H}(x) \setminus \{e_0, a\}|$ it is a bijection. It follows that for any $a' \in \mathcal{H}(x) \setminus \{e_0, a\}$, there exists $y \in e \setminus \{x_0\}$ such that $a' \cap e = \{y\}$.

iv) and v) will follow from the construction of \mathcal{H}' in iv) of Proposition 4.2. \square

Theorem 3.6. *For any $k \geq 2$ and any $\mathcal{H} \in \mathcal{H}_k$, we have*

- i) $\Delta([\mathcal{H}]_2) = k^2 - 1$ and $\delta([\mathcal{H}]_2) = k^2 - k$;
- ii) $k + 1 \leq q(\mathcal{H}) \leq 1 + k \lceil \frac{k}{2} \rceil$;
- iii) $q(\mathcal{H}) \leq \Delta([\mathcal{H}]_2)$.

Proof. i) Let $x_i \in e_0$, then x_i is adjacent to any other vertex of V . Thus $\deg_{[\mathcal{H}]_2}(x_i) = k^2 - 1$. If $x \in V \setminus e_0$ then x belongs to k different hyperedges thus admits $k^2 - k$ adjacent vertices. We infer $\Delta([\mathcal{H}]_2) = k^2 - 1$ and $\delta([\mathcal{H}]_2) = k^2 - k$.

ii) We fix c_0 the color of e_0 , and for each even $0 \leq i \leq k - 1$ and any $0 \leq j \leq k$ we color $e_{i,j}$ with $c_{i,j}$. We need exactly

$$1 + \sum_{\substack{0 \leq i \leq k-1 \\ i \text{ even}}} k = 1 + k \left\lceil \frac{k}{2} \right\rceil$$

many different colors. Our aim is to show that they are sufficiently numerous to provide a hyperedge coloring of \mathcal{H} . For this it remains to color all hyperedges $e_{i,h}$ where $1 \leq i \leq k-1$ is odd and $1 \leq h \leq k$.

We fix an odd integer $1 \leq i \leq k-1$.

- For each $1 \leq j \leq k$ there exists a unique $1 \leq h = h(j) \leq k$ such that $e_{i-1,j} \cap e_{i,h} = \emptyset$: it suffices to apply iii) of Proposition 3.5 with $e = e_{i-1,j}$ and $x = x_i$.

We thus may define the map

$$f: \begin{array}{ccc} \{e \in E \setminus \{e_0\} : x_{i-1} \in e\} & \longrightarrow & \{e \in E \setminus \{e_0\} : x_i \in e\} \\ e_{i-1,j} & \longmapsto & e_{i,h(j)} \end{array}$$

- f is injective. Otherwise it would exist $1 \leq j \neq j' \leq k$ such that $h(j) = h(j')$. Hence

$$e_{i,h(j)} \setminus \{x_i\} \subset \bigsqcup_{\substack{1 \leq t \leq k \\ t \neq j, j'}} e_{i-1,t} \setminus \{x_{i-1}\}.$$

Since $|e_{i,h(j)} \setminus \{x_i\}| = k-1$ and $|(e_{i,h(j)} \setminus \{x_i\}) \cap e_{i-1,t} \setminus \{x_{i-1}\}| \leq 1$ for any $1 \leq t \neq j, j' \leq k$ we deduce from the *pigeon hole principle* that $|e_{i,h(j)} \cap e_{i-1,t}| \geq 2$ for some t , a contradiction to the linearity of \mathcal{H} .

- Thus f is bijective, that is for any $1 \leq h \leq k$ there exists $1 \leq j \leq k$ such that $e_{i,h} \cap e_{i-1,j} = \emptyset$. This shows that hyperedge $e_{i,h}$ can be colored with $c_{i-1,j}$.

Statement iii) follows from i) and ii). \square

Remarks 3.7. i) There exist several sharp results for the chromatic index of linear k -uniform and d -regular hypergraphs \mathcal{H} yielding in particular the asymptotic upper bound $q(\mathcal{H}) \leq (1+o(1))d$ when $d \rightarrow \infty$, where the implied function in $o(1)$ may depend on k (see [19, Section 2] for a rich survey on this topic). However they do not apply when $d = k$.

- ii) The size of the hyperedge coloring of $\mathcal{H} \in \mathcal{H}_k$ given in Theorem 3.6 is certainly not optimal. For instance when $k = 3$ we shall see in Proposition 4.2 that $q(\mathcal{H}) \leq 5$ while the algorithm provided by Theorem 3.6 yields $q(\mathcal{H}) \leq 7$. It could be interesting to improve the upper bound in ii) and to give the order of magnitude of $\max\{q(\mathcal{H}), \mathcal{H} \in \mathcal{H}_k\}$ in terms of k . See Proposition 4.2 for additional properties.

4. Studying the class \mathcal{H}_k

4.1. General facts. For any integer $k \geq 2$, we denote by \mathcal{H}_k the set hypergraphs $\mathcal{H} = (V, E)$ such that

- (C0) \mathcal{H} is linear and k -uniform,
- (C1) $|V| = k^2$,
- (C2) $|E| = k^2 + 1$,

and there exists a hyperedge e_0 such that

- (C3) $\forall x \in e_0, \deg_{\mathcal{H}}(x) = k+1$,
- (C4) $\forall x \in V \setminus e_0, \deg_{\mathcal{H}}(x) \geq k$.

We start with easy consequences of the definition.

Proposition 4.1. *Let $\mathcal{H} = (V, E)$ satisfy Conditions (C0)-(C4). Then*

- i) $\forall x \in V \setminus e_0, \deg_{\mathcal{H}}(x) = k,$
- ii) $\forall e \in E \setminus \{e_0\}, d_{\mathcal{H}}(e) = k^2 - k + 1.$

Proof. i) By (4) and (C2), we have $\sum_{x \in V} \deg_{\mathcal{H}}(x) = (k^2 + 1)k$, hence by (C3) and (C4)

$$\begin{aligned} k^3 + k &= (k + 1)k + (k^2 - k)k \leq (k + 1)k + (k^2 - k) \min_{x \in V \setminus e_0} \deg_{\mathcal{H}}(x) \\ &\leq (k + 1)k + \sum_{x \in V \setminus e_0} \deg_{\mathcal{H}}(x) = (k^2 + 1)k, \end{aligned}$$

since there are $(k^2 - k)$ vertices not belonging to e_0 . It follows that $\forall x \in V \setminus e_0, \deg_{\mathcal{H}}(x) = k$.

ii) For any $e \in E \setminus \{e_0\}$, we get from from the first equation in (8), (C3) and i)

$$d_{\mathcal{H}}(e) = \sum_{x \in e \cap e_0} k + \sum_{x \in e \setminus e_0} (k - 1) = k + (k - 1)^2 = k^2 - k + 1$$

since $|e \cap e_0| = 1$ and $|e \setminus e_0| = k - 1$. □

Proposition 4.2. *For any prime power k there exists a hypergraph denoted by $\widehat{\mathcal{A}}_k$ in the class \mathcal{H}_k such that*

- i) $q(\widehat{\mathcal{A}}_k) = k + 1,$
- ii) $\widehat{\mathcal{A}}_k$ is not maximal (for inclusion) in the class of all linear and k -uniform hypergraphs with k^2 vertices,
- iii) $\widehat{\mathcal{A}}_k$ contains only one critical hyperedge, namely the unique hyperedge e_0 satisfying (C3) and (C4).

Proof. The field plane $\mathcal{A}_k = \mathbb{F}_k^2$ over \mathbb{F}_k provides a $(k + 1)$ -regular and k -uniform linear hypergraph with k^2 vertices when considering the $k^2 + k$ lines as its hyperedges. By removing any $k - 1$ non intersecting hyperedges (namely parallel lines) from \mathcal{A}_k we obtain a hypergraph $\widehat{\mathcal{A}}_k$ of \mathcal{H}_k .

i) Lines of \mathcal{A}_k can be partitioned according to their slope. For each slope, any point appears only once on some line of \mathcal{A}_k . This shows that all the lines having a common slope can be colored with the same color. Since there are $k + 1$ distinct slopes in \mathcal{A}_k , including the vertical one, we can conclude.

ii) This follows from the construction of $\widehat{\mathcal{A}}_k$.

iii) The unique line without any parallel line in $\widehat{\mathcal{A}}_k$ is obviously critical, while removing any other line does not decrease the chromatic index. □

4.2. The cases $k = 2, 3.$

Theorem 4.3. *We have:*

- i) \mathcal{H}_2 contains just the graph $\widehat{\mathcal{A}}_2$ up to isomorphism. It satisfies $q(\widehat{\mathcal{A}}_2) = 3$ and $\text{Aut}(\widehat{\mathcal{A}}_2) \simeq \mathfrak{S}_2 \times \mathfrak{S}_2$, the Klein four-group.
- ii) \mathcal{H}_3 possesses exactly two non isomorphic elements $\widehat{\mathcal{A}}_3$ and \mathcal{H}'_3 which satisfy $q(\widehat{\mathcal{A}}_3) = 4$ and $q(\mathcal{H}'_3) = 5$.
- iii) $\text{Aut}(\widehat{\mathcal{A}}_3) = \mathfrak{S}_3 \rtimes \mathfrak{S}_3$ and $\text{Aut}(\mathcal{H}'_3) \simeq \mathfrak{S}_2 \times \mathfrak{S}_3 = D_6$, the dihedral group of degree 6 and order 12.

Proof. i) For $k = 2$, we see easily that any hypergraph \mathcal{H} in \mathcal{H}_2 is obtained by removing a single edge from the complete simple graph on four vertices K_4 , so that $\mathcal{H} = \widehat{\mathcal{A}}_2$ and $q(\mathcal{H}) = 3$ by Proposition 4.2. Clearly any automorphism of \mathcal{H} is allowed to permute both vertices of degree 2 in one side, and independently both vertices of degree 3 on the other. Hence the result.

ii) Let $(V, E) \in \mathcal{H}_3$. We suppose that $V = \mathbb{F}_3^2$ where we identify the vertex (x, y) with the integer $3x + y$ whose development in base 3 is xy . We may assume that the *special* hyperedge e_0 is $\{0, 1, 2\}$ simply denoted 012. The 3 other hyperedges containing 0 can be set to 036, 048 and 057 with the same short-hand notation. Note that at this stage, the vertices 3, 6 can be permuted, likewise for 4, 8 and for 5, 7. These 4 hyperedges are lines in the field plane \mathbb{F}_3^2 . Now the 3 hyperedges containing 1 and different from e_0 are

$$13u, 16v, 1xw \quad \text{where } \{u, v, w, x\} = \{4, 5, 7, 8\}.$$

We must have $x = 4$ or 8 since $\{w, x\} \neq \{5, 7\}$ by the facts that hyperedge 057 is already defined and \mathcal{H} is linear. Switching if necessary 4 and 8, we may assume that $x = 4$. This implies $w \in \{5, 7\}$. Switching if necessary 5 and 7 we may set $w = 7$. Since 3 and 6 are also permutable we may set $u = 8$ and $v = 5$. Up to isomorphism, hyperedges

$$012, 036, 048, 057, 147, 138, 156$$

are in E . We may observe on the following figure that they are lines in \mathbb{F}_3^2 .




FIGURE 1. A partial hypergraph of \mathcal{H} with 7 lines *immersed* in the field plane \mathcal{A}_3 .

Let

$$2xw, 23u, 26v \quad \text{where } u \in \{4, 5, 7\}, v \in \{4, 7, 8\}, \{w, x\} = \{\{4, 5\}, \{5, 8\}, \{7, 8\}\},$$

be the 3 remaining hyperedges. At least one of them is a line in the plane. Indeed, otherwise we would have

$$u \in \{4, 5\}, v \in \{7, 8\}, \{w, x\} = \{\{4, 5\}, \{7, 8\}\},$$

a contradiction since u, v, w, x are distinct.

Applying if necessary a linear transformation with matrix $\begin{pmatrix} \alpha & 0 \\ \beta & 1 \end{pmatrix}$ for which the vertices 0, 1, 2 are fixed, we may assume that 258 is in E . We get $\{u, v\} = \{4, 7\}$.

- First case: if the set of hyperedges is

$$E = \{012, 036, 048, 057, 147, 138, 156, 258, 237, 246\},$$

all the hyperedges are lines in the plane. We thus obtain the hypergraph $\mathcal{H} = \widehat{\mathcal{A}}_3$.




FIGURE 2. The hypergraph $\widehat{\mathcal{A}}_3$.

- Second case: if the set of hyperedges is

$$E' = \{012, 036, 048, 057, 147, 138, 156, 258, 234, 276\},$$

we let $\mathcal{H}'_3 = (V, E')$.




FIGURE 3. The hypergraph \mathcal{H}'_3 .

We infer that up to isomorphism the class \mathcal{H}_3 reduces to $\{\widehat{\mathcal{A}}_3, \mathcal{H}'_3\}$.

We have $q(\mathcal{H}'_3) \leq 5$ since

$$0 \curvearrowright 012; \quad 1 \curvearrowright 036, 147, 258; \quad 2 \curvearrowright 048, 156; \quad 3 \curvearrowright 057, 234; \quad 4 \curvearrowright 138, 276;$$

provides a hyperedge 5-coloring of \mathcal{H}'_3 . Now suppose by contradiction that 4 colors are sufficient. Without loss of generality, we may assume that we have made the following partial assignment of colors:

$$0 \curvearrowright 012; \quad 1 \curvearrowright 036, 147; \quad 2 \curvearrowright 048, 156; \quad 3 \curvearrowright 057, 138.$$

We see that hyperedge 234 then necessarily requires an additional color, contradiction. Hence $q(\mathcal{H}'_3) = 5$. Since $q(\widehat{\mathcal{A}}_3) = 4$ by i), \mathcal{H}'_3 is not isomorphic to $\widehat{\mathcal{A}}_3$. We conclude that $|\mathcal{H}'_3| = 2$.

Note also that \mathcal{H}'_3 is maximal in the class of 3-uniform linear hypergraphs: adding a new hyperedge yields a non linear hypergraph. In contrast $\widehat{\mathcal{A}}_3 = (V, E)$ can be completed with two additional lines, namely 345, 678. This shows by a different way that \mathcal{H}'_3 and $\widehat{\mathcal{A}}_3$ are non isomorphic.

iii) We use notation of ii) and only sketch the proof. Since e_0 contains all the vertices of degree $k + 1$ in \mathcal{H} , the subset e_0 of V is fixed by any automorphism φ of \mathcal{H} . Denoting by $\varphi|_{e_0}$ the restriction of φ to e_0 , we get in both case $\mathcal{H} = \widehat{\mathcal{A}}_3, \mathcal{H}'_3$, the homomorphism

$$g : \begin{array}{ccc} \text{Aut}(\mathcal{H}) & \longrightarrow & \mathfrak{S}_3 \\ \varphi & \longmapsto & \varphi|_{e_0} \end{array}$$

- When $\mathcal{H} = \widehat{\mathcal{A}}_3$ we have

$$\ker g = \langle p, t \rangle \simeq \mathfrak{S}_3 \quad \text{where} \quad p = (4 \ 3 \ 5)(6 \ 7 \ 8) \text{ and } t = (4 \ 6)(5 \ 7)(3 \ 8)$$

with usual notation for permutation running on the vertices $0, 1, \dots, 8$. We check easily that both permutations $u = (0 \ 1)(4 \ 5)(6 \ 8)$ and $v = (0 \ 1 \ 2)(4 \ 3 \ 5)$ are automorphisms of $\widehat{\mathcal{A}}_3$. Since $g(u) = (0 \ 1)$ and $g(v) = (0 \ 1 \ 2)$, g is surjective and $|\text{Aut}(\widehat{\mathcal{A}}_3)| = 36$.

Let $K = \langle u, v \rangle$. Since $u^2 = v^3 = \text{id}$ and $u \circ v = v^2 \circ u$, we have $K \simeq \mathfrak{S}_3$. Moreover $K \cap \ker g = \{\text{id}\}$ thus finally $\text{Aut}(\widehat{\mathcal{A}}_3) = K \cdot \ker g$. Since $t \circ v \neq v \circ t$, this is a semi direct product, hence $\text{Aut}(\widehat{\mathcal{A}}_3) \simeq \mathfrak{S}_3 \rtimes \mathfrak{S}_3$.

- When $\mathcal{H} = \mathcal{H}'_3$ we have

$$\ker g = \langle q \rangle \simeq \mathfrak{S}_2 \quad \text{where} \quad q = (3 \ 6)(4 \ 7)(5 \ 8).$$

Both permutations $r = (0 \ 1)(3 \ 4)(6 \ 7)$ and $s = (0 \ 1 \ 2)(4 \ 3 \ 5 \ 7)$ are automorphisms of \mathcal{H}'_3 . Since $g(r) = (0 \ 1)$ and $g(s) = (0 \ 1 \ 2)$, g is surjective and $|\text{Aut}(\mathcal{H}'_3)| = 12$.

Let $K' = \langle u', v' \rangle$ where $u' = (0 \ 1 \ 2)(4 \ 8 \ 3)(5 \ 6 \ 7)$ and $v' = (0 \ 1)(3 \ 4)(6 \ 7)$ are automorphisms of \mathcal{H}'_3 . Since $u'^3 = v'^2 = \text{id}$ and $v' \circ u' = u'^2 \circ v'$, we have $K' \simeq \mathfrak{S}_3$. Moreover $K' \cap \ker g = \{\text{id}\}$ thus finally $\text{Aut}(\mathcal{H}'_3) = K' \cdot \ker g \simeq \mathfrak{S}_3 \times \mathfrak{S}_2 = D_6$ since clearly the product is direct. \square

4.3. The general case.

Theorem 4.4. *For any prime power $k \geq 3$, there exists a hypergraph $\mathcal{H}' = \mathcal{H}'_k$ in \mathcal{H}_k such that*

- $q(\mathcal{H}') = 2k - 1$ and any hyperedge $(2k - 1)$ -coloring of \mathcal{H}' is unique up to an affine permutation on the vertices,
- \mathcal{H}' contains exactly k many critical hyperedges,
- \mathcal{H}' is maximal (for inclusion) in the class of all linear and k -uniform hypergraphs with k^2 vertices,
- \mathcal{H}' and $\widehat{\mathcal{A}}_k$ are non isomorphic.

Proof. In the field plane \mathcal{A}_k let x_0 be an arbitrary vertex and $\mathcal{H}(x_0) = \{e_0, e_1, \dots, e_k\}$ be the star centered at x_0 . Let $e_1 = \{x_0, \dots, x_{k-1}\}$ and a_1, \dots, a_{k-1} be the parallel lines to e_0 with $a_i \cap e_1 = \{x_i\}$, $1 \leq i \leq k-1$. We apply to \mathcal{A}_k the following changes on its set of hyperedges:

- e_2, \dots, e_k are removed,
- for any $1 \leq i \leq k-1$, a_i is replaced by the hyperedge $a'_i = \{x_0\} \cup (a_i \setminus \{x_i\})$,
- e_1 and all other hyperedges, namely those intersecting $e_0 \setminus \{x_0\}$ including e_0 itself, are kept unchanged.




FIGURE 4. From \mathcal{A}_k to \mathcal{H}' . Vertices of e_0 and e_1 are labelled so that the lines $\langle x_i, y_i \rangle$ ($1 \leq i \leq k-1$) are parallel in \mathcal{A}_k and \mathcal{H}' .

We get a k -uniform hypergraph $\mathcal{H}' = (V', E')$ such that $|V'| = |V(\mathcal{A}_k)| = k^2$, $|E'| = |E(\mathcal{A}_k)| - (k-1) = k^2 + 1$ and $\deg_{\mathcal{H}'}(x) = \deg_{\mathcal{A}_k}(x) = k+1$ for any $x \in e_0$. Moreover $V' \setminus e_0 = \{x_{i,j}, 1 \leq i \leq k, 1 \leq j \leq k-1\}$ where $\{x_{i,j}\} = e_i \cap a_j$. It follows that $\deg_{\mathcal{H}'}(x_{i,j}) = \deg_{\mathcal{A}_k}(x_{i,j}) - 1$. Finally $\mathcal{H}' \in \mathcal{H}_k$.

i) We now estimate $q(\mathcal{H}')$.

- We first fix a hyperedge coloring $\mathbf{c} : \mathcal{H}' \rightarrow \mathbb{N}$ of \mathcal{H}' . Hyperedges in the star $\mathcal{H}'(x_0) = \{e_0, a'_1, \dots, a'_{k-1}, e_1\}$ are respectively colored by $k+1$ many distinct colors in the set $\mathcal{C} = \{c_0, c_1, \dots, c_{k-1}, c_k\}$. Now the star $\mathcal{H}'(x_1)$ is composed by e_1 and the lines $\langle x_1, y_i \rangle$ ($1 \leq i \leq k-1$) of \mathcal{A}_k passing thru x_1 and y_i respectively where we have set $e_0 = \{x_0, y_1, \dots, y_{k-1}\}$. Vertices of e_0 are labelled so that the hyperedges passing thru x_i, y_i ($1 \leq i \leq k-1$) are parallel. We denote by c'_i ($1 \leq i \leq k-1$) the $k-1$ many distinct colors that are assigned to those lines $\langle x_1, y_i \rangle$ ($1 \leq i \leq k-1$). For any $2 \leq j \leq k-1$, each line $\langle x_1, y_i \rangle$ intersects a_j in \mathcal{A}_k , thus intersects a'_j in \mathcal{H}' since the intersection vertex cannot be x_j by linearity of \mathcal{H}' . Thus $c'_i \notin \mathcal{C} \setminus \{c_1\}$ for any $1 \leq i \leq k-1$. Since there

is at most one line $\langle x_1, y_i \rangle$ ($1 \leq i \leq k-1$) that are colored by c_1 , we infer $q(\mathcal{H}') \geq k+1+k-2=2k-1$.

Note that if none of the lines $\langle x_1, y_i \rangle$ ($1 \leq i \leq k-1$) is colored by c_1 , then the given hyperedge coloring \mathbf{c} has at least $2k$ many colors. We use this fact in order to get an optimal coloring of \mathcal{H}' in what follows.

- We proceed as follows. We color as before $\mathcal{H}'(x_0)$ by $k+1$ many distinct colors c_0, \dots, c_k . We observed that $k-2$ among the $k-1$ lines $\langle x_1, y_i \rangle$ ($1 \leq i \leq k-1$) require new colors c_{k+1}, \dots, c_{2k-2} . We have to show that we can color the remaining hyperedges of \mathcal{H}' without any additional color.

Let e be a not yet colored hyperedge of \mathcal{H}' .

– if e is the last line $\langle x_1, y_{i_1} \rangle$ passing thru x_1 , then we must color it with c_1 ; we denote by $\lambda \in \mathbb{F}_k \setminus \{0\}$ its slope in the field plane \mathcal{A}_k ;

– if $e \in \{e'_2, \dots, e'_{k-1}\}$ is a parallel line to e_1 , then we must color it with c_k since it intersects $e_0, a'_i, \langle x_1, y_i \rangle$ ($1 \leq i \leq k-1$);

– for each $2 \leq j \leq k-1$, the unique parallel $e = \langle x_j, y_{i_j} \rangle$ to $\langle x_1, y_{i_1} \rangle$ needs to be colored with c_j since it intersects e_0, e_1, a'_i ($i \neq j$) and $\langle x_1, y_i \rangle$ ($i \neq i_1$);

– for any $2 \leq j \leq k-1$ and $1 \leq i \leq k-1$, $i \neq i_j$, the line $e = \langle x_j, y_i \rangle$ can only be colored with the same color than that of its unique parallel in $\mathcal{H}'(x_1)$.

By this way only hyperedges of \mathcal{H}' that are parallel lines in \mathcal{A}_k may have the same color. Indeed by construction, each color c_{k+i-1} ($1 \leq i \leq k-1$) is only assigned to disjoint hyperedges of \mathcal{H}' which are parallel lines in \mathcal{A}_k . Color c_0 is exclusive to e_0 . Finally color c_j ($1 \leq j \leq k-1$) is only assigned to both hyperedges a'_j and $\langle x_j, y_{i_j} \rangle$ which are non intersecting.

We thus have a hyperedge $(2k-1)$ -coloring of \mathcal{H}' . Hence $q(\mathcal{H}') = 2k-1$.

- We have shown in fact that hyperedge $(2k-1)$ -coloring of \mathcal{H}' are parametrized by the slope $\lambda \in \mathbb{F}_k \setminus \{0\}$ of the parallel lines $\langle x_j, y_{i_j} \rangle$ ($1 \leq j \leq k-1$). Conversely different slopes λ yield distinct coloring when vertices and hyperedges of \mathcal{H}' are fixed beforehand. Hence if we ignore color labels we infer that \mathcal{H}' admits exactly $k-1$ many hyperedge $(2k-1)$ -coloring $\mathbf{c}_{\mathcal{H}', \lambda}$, $\lambda \in \mathbb{F}_k \setminus \{0\}$.

Precisely each slope $\lambda \in \mathbb{F}_k \setminus \{0\}$ is associated to some admissible one to one mapping $j \mapsto i_j$ from $\{1, \dots, k-1\}$ on itself. It yields a permutation σ on the y_j 's and the e'_j 's respectively $\sigma : y_j \mapsto y_{i_j}$, $e'_j \mapsto e'_{i_j}$ which extends to a one to one mapping from the vertices of e'_j onto those of $\sigma(e'_j)$ ($1 \leq j \leq k-1$) by sending $a'_i \cap e'_j$ on $a'_i \cap \sigma(e'_j)$ ($1 \leq i \leq k-1$), see Figure 4. Since \mathcal{H}' is immersed in the field plane $\mathcal{A}_k = \mathbb{F}_k^2$ we may consider that $e_0 = \{0\} \times \mathbb{F}_k$ and $e_1 = \mathbb{F}_k \times \{0\}$. It follows σ is an affine transformation of the field plane, namely the *dilation from axis e_1 towards axis e_0 by a factor of λ* . We have $\sigma(\mathcal{H}') = \mathcal{H}'$.

Indeed

– σ permutes the e'_j 's,

– $\sigma(e_0) = e_0$, $\sigma(e_1) = e_1$, $\sigma(a'_i) = a'_i$ ($2 \leq i \leq k-1$),

– $\sigma(\langle x_i, y_j \rangle) = \langle x_i, \sigma(y_j) \rangle$ ($1 \leq i, j \leq k-1$).

It is thus an automorphism of \mathcal{H}' . Finally

$$\mathbf{c}_{\mathcal{H}', \lambda}(e) = \mathbf{c}_{\mathcal{H}', 1}(\sigma^{-1}(e)), \quad \text{for any } e \in E(\mathcal{H}').$$

This shows that for any coloring \mathbf{c} of \mathcal{H}' there exists an *affine* automorphism of \mathcal{H}' such that $\mathbf{c}(e) = \mathbf{c}_{\mathcal{H}', 1}(\sigma^{-1}(e))$ for any $e \in E(\mathcal{H}')$.

ii) Here we show that only $e_0, a'_1, \dots, a'_{k-1}$ are critical hyperedges in \mathcal{H}' .

- By the coloring process implemented above, a'_j ($2 \leq j \leq k-1$) is the unique hyperedge of \mathcal{H}' with color c_j . By symmetry on the vertices x_i ($2 \leq i \leq k-1$), there exists a hyperedge coloring of \mathcal{H}' such that a'_1 has an exclusive color c_1 . Hence any a'_j ($1 \leq j \leq k$) is critical. It is clear that e_0 is also critical since it intersects all other hyperedges.
- Any parallel of e_1 , including e_1 itself, intersects all the hyperedges of \mathcal{H}' which are not parallel to e_1 . Removing one of them does not affect the number of required colors. Thus none of them is critical.
- In the first part of the proof of i), we notice that just coloring $e_0, e_1, a'_i, \langle x_1, y_i \rangle$ ($1 \leq i \leq k-1$) already required $2k-1$ colors. We infer that none of the lines $\langle x_j, y_i \rangle$ is critical, $2 \leq j \leq k-1$ and $1 \leq i \leq k-1$. By symmetry of \mathcal{H}' with regard to the vertices x_j , this is also true for the lines $\langle x_1, y_i \rangle$, $1 \leq i \leq k-1$.

iii) Let \mathcal{H}'' be a k -uniform linear hypergraph with k^2 vertices such that $E(\mathcal{H}') \subset E(\mathcal{H}'')$. Let $e \in E(\mathcal{H}'')$. Set $e'_1 = e_1$ and denote by e'_2, \dots, e'_k its parallel lines in \mathcal{A}_k (see Figure 4). These hyperedges are also in \mathcal{H}' hence form a partition of $V(\mathcal{H}')$. We thus have $e \cap e'_i = \{z_i\}$ for distinct vertices z_i , $1 \leq i \leq k$, and $e = \{z_1, \dots, z_k\}$.

- If $z_1 = x_0$ then z_2 is not in $e'_1 = e_1$ and does belong to some a'_j . Since $x_0 \in a'_j \in E(\mathcal{H}') \subset E(\mathcal{H}'')$ and \mathcal{H}'' is linear, we infer $e = a'_j \in E(\mathcal{H}')$.
- Assume that $z_1 = x_j$ for some $1 \leq j \leq k-1$. Since $k \geq 3$ and $|a'_j \cap e| \leq 1$, there exists $2 \leq i \leq k$ such that $z_i \notin a'_j$. This implies that z_1 and z_i lie on two distinct proper parallels to e_0 in \mathcal{A}_k . We then see that the line $\langle z_1, z_i \rangle$ containing both z_1 and z_i in \mathcal{A}_k is distinct from the a_t 's, $1 \leq t \leq k-1$, and from the e_t 's, $0 \leq t \leq k$. Thus it is a hyperedge of \mathcal{H}' as well of \mathcal{H}'' . By linearity of \mathcal{H}'' we conclude that $e = \langle z_1, z_i \rangle \in E(\mathcal{H}')$.

iv) In view of Proposition 4.2 the result follows equivalently from i), ii) or iii). \square

4.4. Remarks and questions.

- As a generalization of iii) of Theorem 4.3 and an extension of iv) of Theorem 4.4, it should be interesting to determine and to compare the groups of automorphisms of $\widehat{\mathcal{A}}_k$ and \mathcal{H}'_k .
- In Proposition 4.2 we investigated the case when k is a prime power and we proved that \mathcal{H}_k is not empty and that $|\mathcal{H}_k| \geq 2$. Estimating $|\mathcal{H}_k|$ seems to be a hard problem.
- When k is not a prime power it is not obvious neither known that \mathcal{H}_k is non-empty. In contrast with the prime power case we should mention that it is conjectured (known as the prime power Conjecture) that there is no affine plane with k^2 points. For instance for $k=6$ this is related to the 36 officers problem posed by Euler and to the existence of Graeco-Latin squares of order 6: there is no affine plane of order 6 (see [27, 26]). However one can ask the question whether or not \mathcal{H}_6 is empty.
- Our construction of \mathcal{H}'_k the Theorem 4.4 starts from the affine plane \mathcal{A}_k . We may ask the question whether or not all hypergraphs of \mathcal{H}_k can be derived in a certain way from \mathcal{A}_k . However, as also noticed in [9], we need to avoid hypergraphs in which parallel is an equivalence relation.

5. Further results

5.1. Some consequences. We first derive the following result establishing Conjecture 1.4 for hypergraphs with small maximum degree.

Theorem 5.1. *Let $\mathcal{H} = (V, E)$ be a linear hypergraph with n vertices and without loop such that $\Delta(\mathcal{H}) \leq \sqrt{n} + 1$. Then $q(\mathcal{H}) \leq n$.*

Proof. Let $\mathcal{H}' = (V, E')$ be a critical partial hypergraph of \mathcal{H} such that $q(\mathcal{H}') = q(\mathcal{H})$.

- If $\text{ar}(\mathcal{H}') \geq \sqrt{n}$ we infer from [24] or from Theorem 1.6 that $q(\mathcal{H}) = q(\mathcal{H}') \leq n$.
- If $\text{ar}(\mathcal{H}') < \sqrt{n}$ then we take $e \in E'$ such that $|e| = \text{ar}(\mathcal{H}')$. By Lemma 3.2 and (8) we get

$$\begin{aligned} q(\mathcal{H}) - 1 &= q(\mathcal{H}') - 1 \leq d_{\mathcal{H}'}(e) = \sum_{x \in e} (\deg_{\mathcal{H}'}(x) - 1) \\ &\leq |e|(\Delta(\mathcal{H}') - 1) \leq \text{ar}(\mathcal{H}')(\Delta(\mathcal{H}) - 1) < n. \end{aligned}$$

Hence the result. □

Note that by (12) we have $q(\mathcal{H}) \leq n$ whenever $r(\mathcal{H})(\Delta(\mathcal{H}) - 1) < n$. Thus by letting $u = \Delta(\mathcal{H}) - \sqrt{n} - 1$ with $0 < u \leq \sqrt{n} - 2$ we see that Conjecture 1.4 holds under the additional strong hypothesis on the rank $r(\mathcal{H}) \leq \sqrt{n} - u$.

The second statement extends the validity of Theorem 1.6 when \mathcal{H} is a uniform linear hypergraph. The extension is moderate but allows us to drop down the threshold $\sqrt{|V|}$ for the antirank.

Theorem 5.2. *Let $k \geq 2$ and $\mathcal{H} = (V, E)$ be a k -uniform linear hypergraph with n vertices satisfying $n \leq k^2 + k - 2$. Then $q(\mathcal{H}) \leq \Delta([\mathcal{H}]_2) + 1$.*

Proof. By Theorem 1.6 we may assume that $n \geq k^2$. We set $u = n - k^2$. Hence

$$(24) \quad 0 \leq u \leq k - 2.$$

Since any partial hypergraph of \mathcal{H} satisfies theorem's hypotheses, we may assume that \mathcal{H} is critical. We distinguish two cases

- We first assume that there exists $e_0 \in E$ and $x_0 \in e_0$ such that $\deg_{\mathcal{H}}(x_0) \leq |e_0| = k$. By Lemma 3.2 and (8) we get

$$\begin{aligned} q(\mathcal{H}) - 1 &\leq d_{\mathcal{H}}(e_0) = \sum_{x \in e_0} (\deg_{\mathcal{H}}(x) - 1) \\ &= \sum_{x \in e_0 \setminus \{x_0\}} \deg_{\mathcal{H}}(x) - |e_0| + \deg_{\mathcal{H}}(x_0) \\ &\leq (|e_0| - 1)\Delta(\mathcal{H}) = (k - 1)\Delta(\mathcal{H}) = \Delta([\mathcal{H}]_2). \end{aligned}$$

- We now assume that $\forall x \in V \deg_{\mathcal{H}}(x) \geq k + 1$. By (6) we get

$$\forall x \in V \quad \deg_{\mathcal{H}}(x) \leq \frac{k^2 + k - 3}{k - 1} < k + 2.$$

hence \mathcal{H} is $(k + 1)$ -regular. By (4) we thus infer

$$|E| = \frac{(k + 1)n}{k} = k^2 + k + u + \frac{u}{k},$$

implying $u = 0$ by (24) since $|E|$ is an integer. Whence $|V| = k^2$ and $|E| = k^2 + k$. Our aim is to show that \mathcal{H} is a finite affine plane yielding $q(\mathcal{H}) = k + 1$.

For any $x \in V$ we have $\deg_{[\mathcal{H}]_2}(x) = \sum_{e \in \mathcal{H}(x)} (|e| - 1) = k^2 - 1 = |V| - 1$ hence any pair of distinct vertices (x, y) in \mathcal{H} are adjacent. Denote by $e_{x,y}$ the unique hyperedge containing both x and y .

Let $e \in E$. Then by (8) we have $d_{\mathcal{H}}(e) = \sum_{x \in e} (\deg_{\mathcal{H}}(x) - 1) = k^2$. Since $|E| = k^2 + k$ and $d_{\mathcal{H}}(e) = k^2$, there are $k - 1$ many hyperedges a_1, \dots, a_{k-1} such that $a_i \cap e = \emptyset$, $1 \leq i \leq k - 1$. If there were $y \in a_i \cap a_j$ with $i \neq j$, then $\mathcal{H}(y)$ would contain $k + 2$ distinct hyperedges: a_i, a_j , and $e_{x,y}$, $x \in e$, a contradiction to $|\mathcal{H}(y)| = \deg_{\mathcal{H}}(y) = k + 1$. Hence $a_i \cap a_j = \emptyset$ if $i \neq j$. This gives

$$\left| e \cup \bigcup_{i=1}^{k-1} a_i \right| = |e| + \sum_{i=1}^{k-1} |a_i| = k^2 = |V|,$$

thus any $x \in V \setminus e$ belongs to exactly one of the hyperedges a_i , $1 \leq i \leq k - 1$, that is a parallel to e . We conclude that \mathcal{H} is a finite affine plane and that $q(\mathcal{H}) = k + 1$. \square

5.2. A generalization of Theorem 1.6. Let $\mathcal{H} = (V, E)$ be a hypergraph with n vertices. Then we can split E as the disjoint union $E = E' \sqcup E''_- \sqcup E''_+$ where

$$\begin{aligned} E' &= \{e \in E : |e| \geq \sqrt{n}\}, \\ E''_- &= \{e \in E \setminus E' : \exists x_0 \in e \text{ such that } \deg_{\mathcal{H}}(x_0) \leq |e|\}, \\ E''_+ &= \{e \in E \setminus E' : \forall x \in e, \deg_{\mathcal{H}}(x) > |e|\}. \end{aligned}$$

Theorems 1.5 and 1.6 can be combined in order to obtain the following slight extension of both results. It shows that the bound (1) is true when $E''_+ = \emptyset$.

Theorem 5.3. *Let $\mathcal{H} = (V, E)$ be a linear hypergraph with n vertices and without loop such that E can be partitioned as $E = E' \sqcup E''$ in such a way that the partial hypergraphs $\mathcal{H}' = (V, E')$ and $\mathcal{H}'' = (V, E'')$ satisfy the following properties:*

- i) $\text{ar}(\mathcal{H}') \geq \sqrt{n}$,
- ii) $\forall e \in E'', |e| < \sqrt{n}$,
- iii) $\forall e \in E'', \exists x_0 \in e$ such that $\deg_{\mathcal{H}}(x_0) \leq |e|$.

Then $q(\mathcal{H}) \leq \Delta([\mathcal{H}]_2) + 1$.

Proof. We argue by induction on $|E''|$. It runs as in the proof of Theorem 1.5.

When $E'' = \emptyset$ then $\mathcal{H} = \mathcal{H}'$ and the result follows directly from Theorem 1.6.

When $E'' = \{e_0, e_1, \dots, e_m\}$ then the partial hypergraph

$$\mathcal{H}_0 := \mathcal{H} \setminus e_0 = (V, E' \sqcup (E'' \setminus \{e_0\}))$$

satisfies conditions i)-iii) and the induction hypothesis applies to \mathcal{H}_0 . We even can assume that $|e_0| = \text{ar}(\mathcal{H})$ since $|e_i| < \sqrt{n} \leq \text{ar}(\mathcal{H}')$ for all $0 \leq i \leq m$. We distinguish two cases.

- If $q(\mathcal{H}) = q(\mathcal{H}_0)$, then by applying induction hypothesis to \mathcal{H}_0 we infer

$$q(\mathcal{H}) = q(\mathcal{H}_0) \leq \Delta([\mathcal{H}_0]_2) + 1 \leq \Delta([\mathcal{H}]_2) + 1.$$

- If $q(\mathcal{H}) = q(\mathcal{H}_0) + 1$ then we fix $x_0 \in e_0$ such that $\deg_{\mathcal{H}}(x_0) \leq |e_0|$ and we get by Lemma 3.2

$$\begin{aligned} q(\mathcal{H}) &\leq d_{\mathcal{H}}(e_0) + 1 = \sum_{\substack{x \in e_0 \\ x \neq x_0}} (\deg_{\mathcal{H}}(x) - 1) + \deg_{\mathcal{H}}(x_0) \\ &\leq (|e_0| - 1) \max_{x \in e} \deg_{\mathcal{H}}(x) - |e_0| + 1 + \deg_{\mathcal{H}}(x_0) \\ &\leq \Delta([\mathcal{H}]_2) + 1, \end{aligned}$$

since $|e_0| = \text{ar}(\mathcal{H})$.

This completes the induction. \square

Remarks 5.4. i) Before applying this theorem it should be convenient to remove all isolated vertices and all but one vertices of degree one in each hyperedge so that the resulting hypergraph still verifies conditions i)-iii). Indeed this will not change the chromatic index but may improve on the upper bound.

- ii) Let $\mathcal{H} = (V, E)$ and assume that $|E''_+| = r \geq 1$. The partial hypergraph $\tilde{\mathcal{H}} = (V, E \setminus E''_+)$ satisfies Theorem 5.3. Letting $\mathcal{H}''_+ = (V, E''_+)$ we infer

$$(25) \quad q(\mathcal{H}) \leq q(\tilde{\mathcal{H}}) + q(\mathcal{H}''_+) \leq \Delta([\tilde{\mathcal{H}}]_2) + 1 + r \leq \Delta([\mathcal{H}]_2) + 1 + r.$$

Note that r can be very large, as large as the total number of hyperedges. If $\mathcal{H} = (V, E)$ is a k -uniform and $(k+1)$ -regular linear hypergraph with k^2 vertices, then $|E| = k(k+1)$ by (4). Adding an isolated vertex to \mathcal{H} gives a hypergraph $\hat{\mathcal{H}} = (\hat{V}, \hat{E})$ in which $\hat{E} = \hat{E}''_+$. In this case the bound (25) is trivial.

- iii) In fact the induction argument in the proof of Theorem 5.3 applies as well on the set of hyperedges

$$E_- := \{e \in E : \exists x_0 \in e \text{ such that } \deg_{\mathcal{H}}(x_0) \leq |e|\}.$$

Hence if the partial hypergraph $(V, E \setminus E_-)$ satisfies Berge-Füredi bound (1) then (V, E) also does. It follows that Conjecture 1.2 is true in general if it holds for every hypergraph (V, E) such that E_- is empty.

5.3. Consequences for the Erdős-Faber-Lovász Conjecture. We noticed earlier that Conjecture 1.2 implies Conjecture 1.4, since $\Delta([\mathcal{H}]_2) \leq |V| - 1$ for any hypergraph $\mathcal{H} = (V, E)$. Therefore we have the following corollaries.

Corollary 5.5. *Let \mathcal{H} be a linear hypergraph with n vertices and satisfying (2). Then $q(\mathcal{H}) \leq n$.*

Corollary 5.6. *Let \mathcal{H} be a linear hypergraph with n vertices and satisfying $\text{ar}(\mathcal{H}) \geq n^{1/2}$. Then $q(\mathcal{H}) \leq n$.*

Corollary 5.7. *Let $\mathcal{H} = (V, E' \sqcup E'')$ be a linear hypergraph with n vertices and satisfying hypotheses of Theorem 5.3. Then $q(\mathcal{H}) \leq n$.*

REFERENCES

- [1] G. Araujo-Pardo and A. Vázquez-Ávila, A note on Erdős-Faber-Lovász Conjecture and edge coloring of complete graphs, *Ars Combin.* **129** (2016), 287–298.
- [2] C. Berge, Motivations and history of some of my conjectures, *Discrete Mathematics* **165-166** (1997), 61–70.

- [3] C. Berge, On the chromatic index of a linear hypergraph and the Chvátal Conjecture, *Annals of the New York Academy of Sciences* **555** (1989), 40–44.
- [4] A. Bretto, Hypergraph Theory: An Introduction, *Mathematical Engineering*, Springer, 2013.
- [5] A. Bretto, A. Faisant and F. Hennecart, On hyperedge coloring of weakly triangulated hypergraphs and well ordered hypergraphs, *Discrete Mathematics* **343** (2020), #112059.
- [6] A. Bretto, A. Faisant and F. Hennecart, Elements of Graph Theory: From Basic Concepts to Modern Developments, *European Mathematical Society*, 2022.
- [7] C. J. Colbourn and M. J. Colbourn, The chromatic index of cyclic Steiner 2-designs, *Intern. J. Math. and Math. Sci.* **5**(4) (1982), 823–825.
- [8] O. Couture, Unsolved Problems in Mathematics, First Edition, 2012 *Published by: White Word Publications 4735/22 Prakashdeep Bldg, Ansari Road, Darya Ganj, Delhi - 110002*
- [9] S. Dow, A completion problem for finite affine planes, *Combinatorica* **6** (1986), 321–325.
- [10] T. Dvoraák, Chromatic index of hypergraphs and Shannon’s Theorem, *Eur. J. Comb.* **21** (2000), 585–591.
- [11] P. Erdős, Problems and results in graph theory and combinatorial analysis, Graph theory and related topics *Proc. Conf., Univ. Waterloo, Waterloo, Ont., 1977*, 153–163, Academic Press, New York-London, 1979.
- [12] P. Erdős, On the combinatorial problems which I would most like to see solved, *Combinatorica* **1** (1982), 25–42.
- [13] P. Erdős, Some of my Favorite Problems in Number Theory, Combinatorics and Geometry, <http://www.revistas.usp.br/resenhasimeusp/article/viewFile/74798/78366>.
- [14] V. Faber, The Erdős-Faber-Lovász conjecture: the uniform regular case, *J. Comb.* **1**(2) (2010), 113–120.
- [15] V. Faber and D. G. Harris, Edge-coloring linear hypergraphs with medium-sized edges. *Random Struct Alg.* **55** (2019), 153–159.
- [16] Z. Füredi, The chromatic index of simple hypergraphs, *Graphs and Combin.* **2**(1) (1986), 89–92.
- [17] L. Haddad and C. Tardif, A clone-theoretic formulation of the Erdős-Faber-Lovász conjecture, *Discussiones Mathematicae Graph Theory* **24** (2004), 545–549.
- [18] N. Hindman, On a conjecture of Erdős, Faber and Lovász about n -colorings, *Canad. J. Math.* **33** (1981), 563–570.
- [19] D. Y. Kang, T. Kelly, D. Kühn, A. Methuku and D. Osthus, Graph and hypergraph colouring via nibble methods: A survey, *arXiv:2106.13733*, 2021.
- [20] J. Kahn, Coloring nearly-disjoint hypergraphs with $q(H) \leq n + o(n)$ colors, *J. Combin. Theory Ser. B* **59**(1) (1992), 31–39.
- [21] J. Khan, Asymptotics of hypergraph matching, covering and coloring problems. *Proceedings of the International Congress of Mathematicians, Zürich, Switzerland 1994*, Birkhäuser Verlag, Basel, Switzerland, 1353–1362, 1995.
- [22] H. Klein and M. Margraf, A remark on the conjecture of Erdős, Faber and Lovász, *Journal of Geometry* **88** (2008), 116–119.
- [23] D. Romero and A. Sanchez-Arroyo, Advances on the Erdős-Faber-Lovász conjecture. In *Grimmet, Geoffrey, McDiarmid, Colin, Combinatorics, Complexity, and Chance: A Tribute to Dominic Welsh*, Oxford Lecture Series in Mathematics and Its Applications, Oxford University Press, 285–298, 2007.
- [24] A. Sanchez-Arroyo, The Erdős-Faber-Lovász conjecture for dense hypergraphs, *Discrete Mathematics* **308** (2008), 991–992.
- [25] P. D. Seymour, Packing nearly-disjoint sets, *Combinatorica* **2**(1) (1982), 91–97.
- [26] D. R. Stinson, A short proof of the nonexistence of a pair of orthogonal Latin squares of order six, *J. Combin. Theory Ser. A* **36** (1984), 373–376.
- [27] G. Tarry, Le problème des 36 officiers, *Compte Rendu de l’Association Française pour l’Avancement des Sciences Naturelles* **1** (1900), 122–123; **2** (1901), 170–203.
- [28] V. Vizing, On an estimate of the chromatic class of a p -graph (in Russian), *Diskret. Analiz.* **3** (1964), 25–30.
- [29] Z. Whang, The Erdős-Faber-Lovász Conjecture for gap-restricted hypergraphs, *Engineering* **16** (2024), 47–59.

[30] <http://www.math.illinois.edu/~dwest/regs/efl.html>

[31] List of unsolved problems in mathematics. <https://en.wikipedia.org/>

NORMANDIEUNICAEN, GREYC CNRS-UMR 6072, CAEN, FRANCE

Email address: `alain.bretto@info.unicaen.fr`

UNIVERSITÉ JEAN MONNET, ICJ UMR5208, CNRS, ECOLE CENTRALE DE LYON, INSA LYON,
UNIVERSITE CLAUDE BERNARD LYON 1, 42023 SAINT-ETIENNE, FRANCE

Email address: `faisant@univ-st-etienne.fr`, `francois.hennecart@univ-st-etienne.fr`