

LINEAR INDEPENDENCE FOR $C_\ell^{(1)}$ BY USING $C_{2\ell}^{(1)}$

MIRKO PRIMC AND GORAN TRUPČEVIĆ

ABSTRACT. In this note we prove linear independence of the combinatorial spanning set for standard $C_\ell^{(1)}$ -module $L(k\Lambda_0)$ by establishing a connection with the combinatorial basis of Feigin-Stoyanovsky's type subspace $W(k\Lambda_0)$ of $C_{2\ell}^{(1)}$ -module $L(k\Lambda_0)$. It should be noted that the proof of linear independence for the basis of $W(k\Lambda_0)$ is obtained by using simple currents and intertwining operators in the vertex operator algebra $L(k\Lambda_0)$.

1. INTRODUCTION

Let \mathfrak{g} be a simple Lie algebra of type C_ℓ , $\ell \geq 2$, and

$$\hat{\mathfrak{g}} = \coprod_{j \in \mathbb{Z}} \mathfrak{g} \otimes t^j + \mathbb{C}c$$

the associated affine Lie algebra of type $C_\ell^{(1)}$. For a basis B_ℓ of \mathfrak{g} (see Subsection 2.4) we have the basis $\bar{B}_\ell \cup \{c\}$ of $\hat{\mathfrak{g}}$, where

$$\bar{B}_\ell = \{x(n) \mid x \in B_\ell, n \in \mathbb{Z}\}$$

and, as usual, $b(n) = b \otimes t^n$. In [PŠ2] it is conjectured that the spanning set consisting of monomial vectors

$$(1.1) \quad u(\pi)v_{k\Lambda_0} = \left(\prod_{b \in B_\ell, n < 0} b(n)^{m_b(n)} \right) v_{k\Lambda_0},$$

where the exponents $\pi = (m_b(n) \mid b \in B_\ell, n < 0)$ are $L_{C_\ell^{(1)}}(k\Lambda_0)$ -admissible, is a basis of the vacuum standard $\hat{\mathfrak{g}}$ -module $L_{C_\ell^{(1)}}(k\Lambda_0)$ of the given level k , with a highest weight vector $v_{k\Lambda_0}$.

In this note we prove this conjecture by using the construction [BPT] of a basis of Feigin-Stoyanovsky's subspace $W_{C_\ell^{(1)}}(k\Lambda_0) \subset L_{C_\ell^{(1)}}(k\Lambda_0)$, consisting, for certain $B_\ell^1 \subset B_\ell$, of monomial vectors

$$w(\pi)v_{k\Lambda_0} = \left(\prod_{b \in B_\ell^1, n < 0} b(n)^{m_b(n)} \right) v_{k\Lambda_0},$$

where the exponents $\pi = (m_b(n) \mid b \in B_\ell^1, n < 0)$ are $W_{C_\ell^{(1)}}(k\Lambda_0)$ -admissible. As noted in [P3], there is a bijection

$$B_\ell \longleftrightarrow B_{2\ell}^1$$

such that the corresponding combinatorial conditions of $L_{C_\ell^{(1)}}(k\Lambda_0)$ -admissible π and $W_{C_{2\ell}^{(1)}}(k\Lambda_0)$ -admissible π coincide (so from here on we use only the term $(k\Lambda_0)$ -admissible).

So the key idea for the proof of linear independence of the set of monomial vectors (1.1) is to embed Lie algebra \mathfrak{g} of type C_ℓ into \mathfrak{g}' of type $C_{2\ell}$ together with its standard module

$$L_{C_\ell^{(1)}}(k\Lambda_0) \subset L_{C_{2\ell}^{(1)}}(k\Lambda_0) \supset W_{C_{2\ell}^{(1)}}(k\Lambda_0),$$

and then, by using inner derivations $T_{\underline{1}}, \dots, T_{\underline{\ell}}$ of \mathfrak{g}' , connect the sets B_ℓ and $B_{2\ell}^1$ and monomials $u(\pi)$ and $w(\pi)$ in the universal enveloping algebra $U(\widehat{\mathfrak{g}'})$:

$$T_{\underline{1}}^{m_{\underline{1}}(\pi)} \dots T_{\underline{\ell}}^{m_{\underline{\ell}}(\pi)}: u(\pi) \rightarrow w(\pi).$$

For $k = 1$ linear independence of the set of monomial vectors (1.1) is proved in [PŠ1], [DK] and [R].

It should be noted that the proof of linear independence for the basis of $W(k\Lambda_0)$ in [BPT] is obtained by using simple currents and intertwining operators for the vertex operator algebra $L_{C_\ell^{(1)}}(k\Lambda_0)$. This method was used in [CLM1] and [CLM2] to find Rogers-Ramanujan and Rogers-Selberg recursions for the characters of Feigin-Stoyanovsky's subspaces of standard $A_1^{(1)}$ -modules, and it seems that this approach is very well-suited for inductive arguments and recursions (cf. [B, J1, J2, P1, P2, T1, T2]). However, it is not clear whether this approach can yield a proof of linear independence of bases for all $L_{C_\ell^{(1)}}(\Lambda)$ which are implicitly conjectured in [CMPP]. J. Dousse and I. Konan proved in [DK] unspecialized character formula for all level 1 standard modules $L_{C_\ell^{(1)}}(\Lambda)$ by using crystal base theory, and M. Russel proved in [R] Andrews-Bressoud type identities by solving recursions which involved all level 1 principally specialized characters of $L_{C_\ell^{(1)}}(\Lambda)$, but not only them(!?). Groebner basis approach is used in [PŠ1] in $k = 1$ case, but for higher levels this most general approach gets to be very difficult (cf. [PŠ3]).

In this note we are interested in vacuum standard modules for affine Lie algebra of type $C_\ell^{(1)}$, $\ell \geq 2$, but most of our considerations hold as well for $\ell = 1$ when $C_1^{(1)} \cong A_1^{(1)}$. In particular, we have Theorem 2.1 for $l = 1$, which was proved by different methods in [MP] and [FKLMM]. Our proof is, in a way, inspired by E. Feigin's degeneration procedure in [F].

2. STANDARD MODULES FOR AFFINE LIE ALGEBRAS

2.1. Affine Lie algebras. Let \mathfrak{g} be a simple complex Lie algebra, \mathfrak{h} a Cartan subalgebra, R the corresponding root system and θ the maximal root with respect to some fixed basis of the root system. For each root α we fix a root vector x_α in \mathfrak{g} . Via a symmetric invariant bilinear form $\langle \cdot, \cdot \rangle$ on \mathfrak{g} we identify \mathfrak{h} and \mathfrak{h}^* and we assume that $\langle \theta, \theta \rangle = 2$. Set

$$\widehat{\mathfrak{g}} = \prod_{j \in \mathbb{Z}} \mathfrak{g} \otimes t^j + \mathbb{C}c, \quad \widetilde{\mathfrak{g}} = \widehat{\mathfrak{g}} + \mathbb{C}d.$$

Then $\widetilde{\mathfrak{g}}$ is the associated untwisted affine Kac-Moody Lie algebra (cf. [K]) with the commutator

$$[x(i), y(j)] = [x, y](i+j) + i\delta_{i+j,0}(x, y)c.$$

Here, as usual, $x(i) = x \otimes t^i$ for $x \in \mathfrak{g}$ and $i \in \mathbb{Z}$, c is the canonical central element, and $[d, x(i)] = ix(i)$. We identify \mathfrak{g} and $\mathfrak{g} \otimes 1$. In the later sections, we will often

use the notation x_n for $x(n)$. Let B be a basis of \mathfrak{g} consisting of root vectors and elements of \mathfrak{h} . Set

$$\bar{B} = \{x(n) \mid x \in B, n \in \mathbb{Z}\}.$$

Then $\bar{B} \cup \{c\}$ is a basis of $\hat{\mathfrak{g}}$. A given linear order \leq on B we extend it to a linear order on \bar{B} by

$$(2.1) \quad x(n) < y(m) \quad \text{iff} \quad n < m \text{ or } n = m \text{ and } x < y.$$

2.2. \mathbb{Z} -gradings of \mathfrak{g} by miniscule coweights. Let R^\vee be the dual root system of R and fix a minuscule coweight $\omega \in P(R^\vee)$. Set $\Gamma = \{\alpha \in R \mid \omega(\alpha) = 1\}$ and

$$\mathfrak{g}_0 = \mathfrak{h} + \sum_{\omega(\alpha)=0} \mathfrak{g}_\alpha, \quad \mathfrak{g}_{\pm 1} = \sum_{\alpha \in \pm \Gamma} \mathfrak{g}_\alpha, \quad \hat{\mathfrak{g}}_0 = \mathfrak{g}_0 \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}c, \quad \hat{\mathfrak{g}}_{\pm 1} = \mathfrak{g}_{\pm 1} \otimes \mathbb{C}[t, t^{-1}].$$

Then on \mathfrak{g} and $\hat{\mathfrak{g}}$ we have \mathbb{Z} -gradings

$$\mathfrak{g} = \mathfrak{g}_{-1} + \mathfrak{g}_0 + \mathfrak{g}_1 \quad \text{and} \quad \hat{\mathfrak{g}} = \hat{\mathfrak{g}}_{-1} + \hat{\mathfrak{g}}_0 + \hat{\mathfrak{g}}_1.$$

Let $\mathfrak{h}'_0 \subset \mathfrak{h}$ be a Cartan subalgebra of $\mathfrak{g}'_0 = [\mathfrak{g}_0, \mathfrak{g}_0]$. Note that $\mathfrak{g}_0 = \mathfrak{g}'_0 + \mathbb{C}\omega$ is a reductive Lie algebra, $\mathfrak{g}_{\pm 1}$ are commutative subalgebras of \mathfrak{g} and \mathfrak{g}_0 -modules, and that $\hat{\mathfrak{g}}_{\pm 1}$ are commutative subalgebras of $\hat{\mathfrak{g}}$ and $\hat{\mathfrak{g}}_0$ -modules.

2.3. Standard modules and Feigin-Stoyanovsky's type subspaces. As usual, we denote by $\Lambda_0, \dots, \Lambda_\ell$ the fundamental weights of $\tilde{\mathfrak{g}}$. For a given integral dominant weight $\Lambda = k_0\Lambda_0 + \dots + k_\ell\Lambda_\ell$ we denote by $L(\Lambda)$ or $L_{\tilde{\mathfrak{g}}}(\Lambda)$ the standard (i.e. integral highest weight) $\tilde{\mathfrak{g}}$ -module with the highest weight Λ , by v_Λ a fixed highest weight vector and by $k = \Lambda(c)$ the level of the module (cf. [K]). In this note we shall be interested mostly in the vacuum standard modules $L(k\Lambda_0)$.

For a standard $\tilde{\mathfrak{g}}$ -module $L(\Lambda)$ and a given \mathbb{Z} -grading on $\hat{\mathfrak{g}}$ as above, we define the corresponding Feigin-Stoyanovsky's type subspace $W(\Lambda)$ or $W_{\tilde{\mathfrak{g}}}(\Lambda)$ as

$$W(\Lambda) = U(\hat{\mathfrak{g}}_1)v_\Lambda \subset L(\Lambda).$$

Note that $U(\hat{\mathfrak{g}}_1) \cong S(\hat{\mathfrak{g}}_1)$ since $\hat{\mathfrak{g}}_1$ is commutative.

2.4. A basis of $\hat{\mathfrak{g}}$ of the type $C_\ell^{(1)}$. Let \mathfrak{g} be a simple Lie algebra of the type C_ℓ and $\mathfrak{g} = \mathfrak{n}_- + \mathfrak{h} + \mathfrak{n}_+$ a Cartan decomposition. We identify the root system

$$R = \{\pm\epsilon_i \pm \epsilon_j \mid i, j = 1, \dots, \ell\} \setminus \{0\},$$

where $\{\epsilon_i \mid i = 1, \dots, \ell\}$ is the canonical basis in \mathbb{R}^ℓ . For $1 \leq a, b \leq \ell$ fix root vectors in \mathfrak{g} denoted as

- ab for the root $\epsilon_a + \epsilon_b$, $a \leq b$,
- \underline{ab} for the root $-\epsilon_a - \epsilon_b$, $a \geq b$,
- $a\bar{b}$ for the root $\epsilon_a - \epsilon_b$, $a \neq b$,
- $a\underline{a} = h_a$ (a simple coroot in \mathfrak{h}).

These vectors form a weight basis $B_\ell = B$ of \mathfrak{g} . Set

$$(2.2) \quad 1 > \dots > \ell > \underline{\ell} > \dots > \underline{1}.$$

We write B_ℓ in a triangular scheme with the element $ab \in B_\ell$ in a -th column and b -th row. For $\ell = 3$ we have B_3

11
 12 22
 13 23 33
 13 23 33 33
 12 22 32 32 22
11 21 31 31 21 11

Note that 11 is the root vector for the highest root $\theta = 2\epsilon_1$ of \mathfrak{g} .


We write $\bar{B}_\ell = \bar{B}$ for the affine Lie algebra $\hat{\mathfrak{g}}$ of the type $C_\ell^{(1)}$. In order to describe the combinatorial monomial basis of $L(k\Lambda_0)$ we need a subset $\bar{B}_\ell^{<0} \subset \bar{B}$,

$$\bar{B}_\ell^{<0} = B_\ell \otimes t^{-1} \cup B_\ell \otimes t^{-2} \cup B_\ell \otimes t^{-3} \cup \dots,$$

written as the array composed of a sequence of triangles as in [T3]. For $\ell = 2$ we have the array $\bar{B}_2^{<0}$

11₋₁ 12₋₁ 12₋₁ 11₋₁ 11₋₂
 22₋₁ 22₋₁ 21₋₁ 12₋₂ 22₋₂
 22₋₁ 21₋₁ 12₋₂ 22₋₂ 22₋₂
 11₋₁ 11₋₂ 21₋₂ 21₋₂ 11₋₂
 11₋₃ 12₋₃ 12₋₃ 11₋₃ 11₋₄
 ⋮ ⋮ ⋮ ⋮ ⋮

where $ab_{-n} = ab(-n) = ab \otimes t^{-n}$. We shall write this array of root vectors $\bar{B}_\ell^{<0}$ rotated for $\pi/4$, denoting only the position \circ of an element $x \in \bar{B}_\ell^{<0}$:



Of course, it is easy to see that the second element in the top row is $x = 22_{-2}$. A downward path \mathcal{Z} in the array $\bar{B}_\ell^{<0}$ is a sequence (or a subset) with an element in the top row followed by an adjacent element in the second row, and so on all the way to an element in the bottom row. An example of a downward path is denoted with the dashed line.

For $\ell > 2$ the array $\bar{B}_\ell^{<0}$ is composed of a sequence of triangles so that for triangles $B_\ell \otimes t^{-n}$ and $B_\ell \otimes t^{-n-1}$ the adjacent sides are

$$(\underline{11}_{-n}, \dots, \underline{\ell 1}_{-n}, \underline{\ell 1}_{-n}, \dots, \underline{11}_{-n}) \quad \text{and} \quad (11_{-n-1}, \dots, 1\ell_{-n-1}, 1\ell_{-n-1}, \dots, 1\underline{1}_{-n-1})$$

and the weight difference between the corresponding points is given by

$$\text{wt}(1a_{-n-1}) = -\alpha_0 + \text{wt}(a\underline{1}_{-n}),$$

where $\alpha_0 = -\theta + \delta$ is the simple root and δ is the imaginary root for the affine Lie algebra $\hat{\mathfrak{g}}$. The array of root vectors $\bar{B}_\ell^{<0}$ rotated for $\pi/4$ has $2\ell + 1$ rows and through each point in the first row there is $2^{2\ell}$ downward paths.

2.5. Combinatorial spanning set of $C_\ell^{(1)}$ -modules $L(k\Lambda_0)$. By using (2.2) we define a lexicographical order \leq on B_ℓ and we extend it to \bar{B}_ℓ by (2.1). A monomial

$$(2.3) \quad \pi = \prod_{b(j) \in \bar{B}_\ell^{<0}} b(j)^{m_{b(j)}} \in S(\tilde{\mathfrak{g}})$$

can be interpreted as a colored partition π : for $m_{b(j)} > 0$ we say that $b(j)$ is a part of degree $|b(j)| = j$ and color $b \in B_\ell$ which appears in the partition $m_{b(j)}$ times. We define the degree and length of a colored partition π as

$$|\pi| = \sum_{b(j) \in \bar{B}_\ell^{<0}} j \cdot m_{b(j)}, \quad \ell(\pi) = \sum_{b(j) \in \bar{B}_\ell^{<0}} m_{b(j)}.$$

For a positive integer k we say that a monomial (colored partition) π is $k\Lambda_0$ -admissible if for every downward path \mathcal{Z} in the array $\bar{B}_\ell^{<0}$ we have

$$\sum_{b(j) \in \mathcal{Z}} m_{b(j)} \leq k.$$

For a monomial π of length s ,

$$\pi = x_1 x_2 \dots x_s, \quad x_1 \leq x_2 \leq \dots \leq x_s,$$

we define the corresponding monomial in the enveloping algebra

$$(2.4) \quad u(\pi) = x_1 x_2 \dots x_s \in U(\tilde{\mathfrak{g}})$$

and the corresponding monomial vector in the standard module

$$u(\pi)v_{k\Lambda_0} \in L(k\Lambda_0).$$

It is proved in [PŠ2] that the set of monomial vectors $u(\pi)v_{k\Lambda_0}$ with $k\Lambda_0$ -admissible π is a spanning set of $L(k\Lambda_0)$. Moreover, in [PŠ1] it is proved that this is a basis for $k = 1$ and it is conjectured that this spanning set is a basis for all integer $k > 1$. The main result of this paper is

Theorem 2.1. *For a positive integer k the set of monomial vectors $u(\pi)v_{k\Lambda_0}$ with $k\Lambda_0$ -admissible π is a basis of $L_{C_\ell^{(1)}}(k\Lambda_0)$.*

We prove linear independence of the spanning set by establishing a connection with the combinatorial basis of Feigin-Stoyanovsky's type subspace $W(k\Lambda_0)$ of $C_{2\ell}^{(1)}$ -module $L(k\Lambda_0)$ constructed in [BPT].

2.6. Combinatorial bases of Feigin-Stoyanovsky type subspaces for $C_{2\ell}^{(1)}$. For \mathfrak{g} of the type C_ℓ there is (up to conjugation) only one \mathbb{Z} -grading

$$\mathfrak{g} = \mathfrak{g}_{-1} + \mathfrak{g}_0 + \mathfrak{g}_1$$

with $\mathfrak{h} \subset \mathfrak{g}_0$ and we choose the basis

$$B_\ell^1 = \{ij \mid 1 \leq i \leq j \leq \ell\}$$

for \mathfrak{g}_1 organized in a triangular scheme. For $\ell = 3$ we have the elements of B_3^1 in the upper shaded triangle:

11
 12 22
 13 23 33
 13 23 33 33
 12 22 32 32 22
 11 21 31 31 21 11

From now on, for fixed $\ell \geq 2$, we consider $B_{2\ell}^1$. For $\ell = 3$ we have a basis B_6^1 of \mathfrak{g}_1 for \mathfrak{g} of the type C_6 :

11
 12 22
 13 23 33
 14 24 34 44
 15 25 35 45 55
 16 26 36 46 56 66

In order to describe the combinatorial monomial basis of Feigin-Stoyanovsky's subspace $W(k\Lambda_0)$ for $C_{2\ell}^{(1)}$ we need a subset $\bar{B}_{2\ell}^{1,<0} \subset \bar{B}_{2\ell}$,

$$\bar{B}_{2\ell}^{1,<0} = B_{2\ell}^1 \otimes t^{-1} \cup B_{2\ell}^1 \otimes t^{-2} \cup B_{2\ell}^1 \otimes t^{-3} \cup \dots,$$

written as the array composed of a sequence of triangles. For $\ell = 2$ we have the array $\bar{B}_4^{1,<0}$

11₋₁ 12₋₁ 13₋₁ 14₋₁ 11₋₂
 22₋₁ 23₋₁ 24₋₁ 12₋₂ 22₋₂
 33₋₁ 34₋₁ 13₋₂ 23₋₂ 33₋₂
 44₋₁ 14₋₂ 24₋₂ 34₋₂ 44₋₂
 11₋₃ 12₋₃ 13₋₃ 14₋₃ 11₋₄
 ∴ ∴ ∴ ∴ ∴

where $ij_{-n} = ij(-n) = ij \otimes t^{-n}$. We shall write this array of root vectors $\bar{B}_4^{1,<0}$ rotated for $\pi/4$, denoting only the position \circ of an element $x \in \bar{B}_4^{1,<0}$:

∘ ∘ ∘
 ∘ ∘ ∘ ∘
 ∘ ∘ ∘ ∘ ∘
 ∘ ∘ ∘ ∘ ∘
 ∘ ∘ ∘ ∘ ∘

The array $\bar{B}_{2\ell}^{1,<0}$ rotated for $\pi/4$ has $2\ell + 1$ rows. A downward path \mathcal{Z} in the array $\bar{B}_{2\ell}^{1,<0}$ is defined as before, and through each point in the first row there is $2^{2\ell}$ downward paths.

Like before, a monomial

$$\pi = \prod_{b(j) \in \bar{B}_{2\ell}^{1,<0}} b(j)^{m_{b(j)}} \in S(\tilde{\mathfrak{g}}_1)$$

FIGURE 1. Bases B_6 , shaded B_3 and boxed B_6^1

11
12 22
13 23 33
14 24 34 44
15 25 35 45 55
16 26 36 46 56 66
16 26 36 46 56 66 66
15 25 35 45 55 65 65 55
14 24 34 44 54 64 64 54 44
13 23 33 43 53 63 63 53 43 33
12 22 32 42 52 62 62 52 42 32 22
11 21 31 41 51 61 61 51 41 31 21 11

can be interpreted as a colored partition π with colors $b \in B_{2\ell}^1$. Formally the definition of $k\Lambda_0$ -admissible colored partitions π on the array $\bar{B}_{2\ell}^{1, <0}$ is the same as before.

Note that the Lie algebra \mathfrak{g}_1 is commutative, so the enveloping algebra is commutative, that is $U(\mathfrak{g}_1) \cong S(\mathfrak{g}_1)$. It is proved in [BPT] that the set of monomial vectors $u(\pi)v_{k\Lambda_0}$ with $k\Lambda_0$ -admissible π is a basis set of the Feigin-Stoyanovsky's subspace $W(k\Lambda_0)$.

3. THE PROOF OF LINEAR INDEPENDENCE

Let \mathfrak{g}' be a simple Lie algebra of the type $C_{2\ell}$. For any subset $\{i_1, \dots, i_\ell\} \subset \{1, \dots, 2\ell\}$, a subalgebra generated by root vectors

$$\{ab \mid a, b \in \{i_1, \dots, i_\ell, \underline{i}_\ell, \dots, \underline{i}_1\}, a \geq b\}$$


is a Lie algebra of the type C_ℓ . This induces inclusions of the corresponding affine Lie algebra $C_\ell^{(1)}$ and its vacuum standard module

$$L_{C_\ell}(k\Lambda_0) \subset L_{C_{2\ell}}(k\Lambda_0).$$

In the remaining considerations, we fix a subset $\{1, \dots, \ell\} \subset \{1, \dots, 2\ell\}$ and the corresponding subalgebra $\mathfrak{g} \subset \mathfrak{g}'$; for $\ell = 3$ we see on Figure 1 the shaded basis $B_3 \subset B_6$ and the boxed basis $B_6^1 \subset B_6$. We consider the basis $\bar{B}_\ell^{<0}$ as a subset of $\bar{B}_{2\ell}^{<0}$ and similarly for monomials in $\bar{B}_\ell^{<0}$ and $\bar{B}_{2\ell}^{<0}$ and the corresponding monomial vectors. The basic idea for the proof of linear independence is to shift colors of monomials in $\bar{B}_\ell^{<0}$ to $\bar{B}_{2\ell}^{1, <0}$ (see Figure 2). The same combinatorial description of the spanning set of the vacuum standard module $L_{C_\ell^{(1)}}(k\Lambda_0)$ from [PŠ2] and of the basis of Feigin-Stoyanovsky's subspace $W_{C_{2\ell}^{(1)}}(k\Lambda_0)$ from [BPT] gives linear independence of the first set.

To shift colors from B_ℓ to $B_{2\ell}^1$, or from \bar{B}_ℓ to $\bar{B}_{2\ell}^1$, we use the adjoint action $T_\ell, \dots, T_{\underline{1}}$ by suitable elements of \mathfrak{g}' . Note that for $b \in B_{2\ell}$ and $n < 0$ we have

$$(3.1) \quad T_{\underline{a}}(b_n) = (T_{\underline{a}}b)_n.$$

FIGURE 2. Shifting colors from B_3 to B_6^1 

Example 3.1. For $\ell = 3$ we first move the shaded elements of B_3 in $\underline{3}$ -row to 4-row in B_6^1 with $T_3 = \text{ad}(34)$ (see Figure 2):

$$\underline{33} \mapsto \underline{43}, \quad \underline{32} \mapsto \underline{42}, \quad \underline{31} \mapsto \underline{41} \quad \text{and} \quad \underline{13} \mapsto 14, \quad \underline{23} \mapsto 24, \quad \underline{33} \mapsto 34, \quad \underline{43} \mapsto 44,$$

and $T_3 = \text{ad}(34)$ is zero on all other shaded elements together with newly shaded $\underline{42}$ and $\underline{41}$. Now we have shaded 4-row in B_6^1 and last two rows with shaded 4×2 rectangle at the beginning and 2×2 triangle at the end. Next we move the shaded elements in $\underline{2}$ -row to 5-row in B_6^1 with $T_2 = \text{ad}(25)$:

$$\underline{22} \mapsto \underline{52}, \quad \underline{21} \mapsto \underline{51} \quad \text{and} \quad \underline{12} \mapsto 15, \quad \underline{22} \mapsto 25, \quad \underline{32} \mapsto 35, \quad \underline{42} \mapsto 45, \quad \underline{52} \mapsto 55$$

and $T_2 = \text{ad}(25)$ is zero on all other shaded elements together with newly shaded $\underline{51}$. Now we have shaded 5-row in B_6^1 and one last row with shaded 5×1 “rectangle” at the beginning and 1×1 “triangle” at the end. Finally we move the shaded elements in $\underline{1}$ -row to 6-row with $T_1 = \text{ad}(16)$:

$$\underline{11} \mapsto \underline{61} \quad \text{and} \quad \underline{11} \mapsto 16, \quad \underline{21} \mapsto 26, \quad \underline{31} \mapsto 36, \quad \underline{41} \mapsto 46, \quad \underline{51} \mapsto 56, \quad \underline{61} \mapsto 66.$$

In general, $B_{2\ell}$ is the triangle with vertices $11, \underline{11}, \underline{11}$; $B_{2\ell}^1$ is the triangle with vertices $11, 1(2\ell), (2\ell)(2\ell)$ —we think of it as boxed elements; and B_ℓ is the union of two $\ell \times \ell$ triangles with vertices $11, 1\ell, \ell\ell$ and $\ell\ell, \underline{\ell 1}, \underline{11}$ and one $\ell \times \ell$ square with vertices $\ell\ell, 1\ell, \underline{11}, \underline{\ell 1}$ —we think of it as shaded elements.

For $1 \leq a \leq \ell$ set

$$(3.2) \quad T_a = \text{ad}(t_a): \mathfrak{g}' \rightarrow \mathfrak{g}', \quad t_a = a(2\ell - a + 1) \in B_{2\ell} \subset \mathfrak{g}'.$$

We use inner derivations T_a to shift in steps colors in B_ℓ into colors in $B_{2\ell}^1$, row by row. First we apply $T_\ell = \text{ad}(\ell(\ell + 1))$ and move the vertical catheti of the lower triangle $[\underline{\ell\ell}, \underline{\ell 1}]$ onto the vertical segment $[(\ell + 1)\underline{\ell}, (\ell + 1)\underline{1}]$, forming a new shaded $\ell \times (\ell + 1)$ rectangle. Then we move with $T_\ell = \text{ad}(\ell(\ell + 1))$ the upper side of newly formed shaded rectangle $[\underline{1\ell}, (\ell + 1)\underline{\ell}]$ into the $(\ell + 1)$ -row $[1(\ell + 1), (\ell + 1)(\ell + 1)]$ of boxed triangle $B_{2\ell}^1$. The result is the shaded union of $(\ell + 1) \times (\ell + 1)$ triangle on the top, $(\ell + 1) \times (\ell - 1)$ rectangle at the right angle and $(\ell - 1) \times (\ell - 1)$ triangle on the right.

In $r < \ell$ steps we get the shaded union of $(\ell + r) \times (\ell + r)$ triangle on the top, $(\ell + r) \times (\ell - r)$ rectangle at the right angle and $(\ell - r) \times (\ell - r)$ triangle on the right, so we can proceed by applying $T_{\ell-r} = \text{ad}((\ell - r)(\ell + r + 1))$.

Our intuitive considerations above we can rephrase with formulas:

Lemma 3.2.

$$\begin{aligned} T_a \underline{ab}_n &\in \mathbb{C}^\times (2\ell - a + 1) \underline{b}_n, \\ T_a \underline{ca}_n &\in \mathbb{C}^\times c(2\ell - a + 1)_n, \end{aligned}$$

for $1 \leq b \leq a \leq \ell$, $1 \leq c \leq 2\ell - a + 1$. In the first case, there is a shift from the \underline{a} -column to the $(2\ell - a + 1)$ -column, while in the second case, there is a shift from the \underline{a} -row to the $(2\ell - a + 1)$ -row. Moreover

$$\begin{aligned} T_{\underline{a}}^2 \underline{ca}_n &= 0, \\ T_{\underline{a}}^2 \underline{ab}_n &= 0 \text{ for } b < a, \\ T_{\underline{a}}^3 \underline{aa}_n &= 0, \\ T_{\underline{a}}^2 \underline{bc}_n &= 0 \text{ for } b, c \notin \{\underline{a}, \underline{2\ell - a + 1}\}. \end{aligned}$$

Since an inner derivation on $\widehat{\mathfrak{g}}$ extends to a derivation on $U(\widehat{\mathfrak{g}})$, we have:

Lemma 3.3. *For a positive integer m , $1 \leq b \leq a \leq \ell$ and $1 \leq c \leq 2\ell - a + 1$, we have*

$$\begin{aligned} T_{\underline{a}}^{2m} (\underline{aa})^m &\in \mathbb{C}^\times ((2\ell - a + 1)(2\ell - a + 1))^m, \\ T_{\underline{a}}^{2m+1} (\underline{aa})^m &= 0, \\ T_{\underline{a}}^m (\underline{ab})^m &\in \mathbb{C}^\times ((2\ell - a + 1)\underline{b})^m, \\ T_{\underline{a}}^{m+1} (\underline{ab})^m &= 0 \\ T_{\underline{a}}^m (\underline{ca})^m &\in \mathbb{C}^\times (c(2\ell - a + 1))^m, \\ T_{\underline{a}}^{m+1} (\underline{ca})^m &= 0. \end{aligned}$$

Proof. By Lemma 3.2 we have

$$T_{\underline{a}}^2 (\underline{aa}) \in \mathbb{C}^\times (2\ell - a + 1)(2\ell - a + 1) \quad \text{and} \quad T_{\underline{a}}^3 (\underline{aa}) = 0.$$

The monomial $(\underline{aa})^m$ has m factors \underline{aa} , and if we act with the derivation $T_{\underline{a}}$ on each of this factors twice, we get something proportional to $((2\ell - a + 1)(2\ell - a + 1))^m$, and if we act on some factor more than two times we get 0. So the first two statements follow from the pigeonhole principle. For other four statements we argue similarly. \square

For a monomial π with variables in $\bar{B}_\ell^{<0}$ given by (2.3), and $1 \leq a \leq \ell$, define

$$\begin{aligned} m_{\underline{a}}(\pi) &= \sum_{b=1}^{\ell} \sum_{n < 0} m_{b\underline{a}}(n) + \sum_{b=1}^{a-1} \sum_{n < 0} m_{\underline{a}b}(n) + 2 \sum_{n < 0} m_{\underline{aa}}(n) + \sum_{b=a+1}^{\ell} \sum_{n < 0} m_{\underline{ba}}(n), \\ M(\pi) &= \sum_{a=1}^{\ell} m_{\underline{a}}(\pi). \end{aligned}$$

For $x \in \mathfrak{g}' \subset \widehat{\mathfrak{g}}$ we have the inner derivation $T = \text{ad } x$ on $U(\widehat{\mathfrak{g}})$. Since $xv_{k\Lambda_0} = 0$, for $u \in U(\widehat{\mathfrak{g}})$ we have $x(uv_{k\Lambda_0}) = [x, u]v_{k\Lambda_0} = (Tu)v_{k\Lambda_0}$. In particular, for $T_{\underline{a}}$ and $t_{\underline{a}}$ defined by (3.2) we have

$$t_{\underline{a}}(uv_{k\Lambda_0}) = (T_{\underline{a}}u)v_{k\Lambda_0}.$$

Let

$$\begin{aligned} t(\pi) &= t_{\underline{1}}^{m_{\underline{1}}(\pi)} t_{\underline{2}}^{m_{\underline{2}}(\pi)} \dots t_{\underline{\ell}}^{m_{\underline{\ell}}(\pi)} \\ &= 1(2\ell)^{m_{\underline{1}}(\pi)} 2(2\ell - 1)^{m_{\underline{2}}(\pi)} \dots \ell(\ell + 1)^{m_{\underline{\ell}}(\pi)}, \\ T(\pi) &= T_{\underline{1}}^{m_{\underline{1}}(\pi)} T_{\underline{2}}^{m_{\underline{2}}(\pi)} \dots T_{\underline{\ell}}^{m_{\underline{\ell}}(\pi)}. \end{aligned}$$

Lemma 3.4. *For a monomial π given by (2.3) we have*

$$t(\pi)(u(\pi)v_{k\Lambda_0}) = (T(\pi)u(\pi))v_{k\Lambda_0} = w(\pi')v_{k\Lambda_0},$$

where $w(\pi')$ is a monomial in $\bar{B}_{2\ell}^{1, <0} \subset \bar{B}_{2\ell}^{<0}$ such that

$$\begin{aligned} m_{ab(n)}(\pi') &= m_{ab(n)}(\pi), \\ m_{a(2\ell-b+1)(n)}(\pi') &= m_{\underline{a}\underline{b}(n)}(\pi), \\ m_{(2\ell-a+1)(2\ell-b+1)(n)}(\pi') &= m_{\underline{a}\underline{b}(n)}(\pi), \\ m_{\underline{a}\underline{b}(n)}(\pi') &= 0, \\ m_{\underline{a}\underline{b}(n)}(\pi') &= 0, \end{aligned}$$

for $1 \leq a, b \leq \ell$.

Proof. Assume that the monomial $u(\pi)$ defined by (2.4) has at least one factor $b(j)^{m_{b(j)}}$, $m_{b(j)} > 0$, with color b from the $\underline{\ell}$ -row of B_ℓ . To shift colors in the monomial $u(\pi)$, first act with $T_{\underline{\ell}}^{m_{\underline{\ell}}(\pi)} = \ell(\ell+1)^{m_{\underline{\ell}}(\pi)}$. This shifts colors from $\underline{\ell}$ -column to $(\ell+1)$ -column, and from $\underline{\ell}$ -row to $(\ell+1)$ -row. Monomial may have factors

$$1\underline{\ell}(n)^{m_{1\underline{\ell}(n)}}, 2\underline{\ell}(n)^{m_{2\underline{\ell}(n)}}, \dots, \underline{\ell}\underline{\ell}(n)^{m_{\underline{\ell}\underline{\ell}(n)}}, \underline{\ell}\underline{\ell}(n)^{m_{\underline{\ell}\underline{\ell}(n)}}$$

with colors in ℓ -row for several values of $n < 0$, but because of (3.1) $T_{\underline{\ell}}$ acts only on colors of these factors. Arguing as in the proof of Lemma 3.3, we see we need to act

$$m' = \sum_{n < 0} \sum_{b=1}^{\ell} m_{b\underline{\ell}(n)}$$

times with $T_{\underline{\ell}}$ to move all factors with colors $1\underline{\ell}, \dots, \underline{\ell}\underline{\ell}$ to factors with colors $1(\ell+1), \dots, \ell(\ell+1)$, and acting with $T_{\underline{\ell}}^{m'+1}$ will produce 0. Schematically we write

$$T_{\underline{\ell}}^{m'} : 1\underline{\ell}(n)^{m_{1\underline{\ell}(n)}}, \dots, \underline{\ell}\underline{\ell}(n)^{m_{\underline{\ell}\underline{\ell}(n)}}, \forall n \rightsquigarrow 1(\ell+1)(n)^{m_{1\underline{\ell}(n)}}, \dots, \ell(\ell+1)(n)^{m_{\underline{\ell}\underline{\ell}(n)}}, \forall n.$$

We also see that we need to act

$$m'' = 2 \sum_{n < 0} m_{\underline{\ell}\underline{\ell}(n)}$$

times with $T_{\underline{\ell}}$ to move all factors with color $\underline{\ell}\underline{\ell}$ to factors with colors $(\ell+1)(\ell+1)$, and acting with $T_{\underline{\ell}}^{m''+1}$ will produce 0. Schematically we write

$$T_{\underline{\ell}}^{m''} : \underline{\ell}\underline{\ell}(n)^{m_{\underline{\ell}\underline{\ell}(n)}}, \forall n \rightsquigarrow (\ell+1)(\ell+1)(n)^{m_{1\underline{\ell}(n)}}, \forall n.$$

So acting with $T_{\underline{\ell}}^{m'+m''}$ we move colors from $\underline{\ell}$ -row to $(\ell+1)$ row. But regardless whether there is or there is not a factor in $u(\pi)$ with color in $\underline{\ell}$ -row of B_ℓ , we must not forget to move the $\underline{\ell}$ -column below $\underline{\ell}\underline{\ell}$ to $(\ell+1)$ -column, for which we need to act

$$m''' = \sum_{n < 0} \sum_{b=1}^{\ell-1} m_{\underline{\ell}b}(n)$$

times with $T_{\underline{\ell}}$. Schematically we write

$$T_{\underline{\ell}}^{m'''} : \underline{\ell}b(n)^{m_{\underline{\ell}b(n)}}, b < \ell, \forall n \rightsquigarrow (\ell+1)\underline{b}(n)^{m_{\underline{\ell}b(n)}}, b < \ell, \forall n.$$

After that we proceed acting with $T_{\underline{a}}^{m_{\underline{a}}}$ for $a = \ell-1, \dots, 2, 1$, and arguing in a similar way. \square

The proof of linear independence. Let

$$(3.3) \quad \sum_{\pi \in S} C_\pi u(\pi) v_{k\Lambda_0} = 0,$$

for some nonempty set S of $k\Lambda_0$ -admissible monomials in $\bar{B}_\ell^{<0}$. Let $M = \max_{\pi \in S} M(\pi)$.

Let $\pi_0 \in S$ be such that $M(\pi_0) = M$. Acting by $t(\pi_0)$ on (3.3) one gets

$$\sum_{\pi \in S} C_\pi (T(\pi_0)u(\pi)) v_{k\Lambda_0} = 0.$$

By using the pigeonhole principle, and an argument similar to the proof of Lemma 3.3, we see that

$$T(\pi_0)u(\pi) = \begin{cases} C'_\pi w(\pi'), & C'_\pi \in \mathbb{C}^\times \text{ if } t(\pi) = t(\pi_0), \\ 0 & \text{otherwise.} \end{cases}$$

Hence,

$$\sum_{\pi \in S'} C'_\pi C_\pi w(\pi') v_{k\Lambda_0} = 0,$$

for some nonempty subset $S' \subset S$. This is a relation of linear dependence between monomials from $\bar{B}_{2\ell}^{1, <0}$. Since these are, again, $k\Lambda_0$ -admissible, and hence linearly independent, it follows that $C_\pi = 0$ for $\pi \in S'$. In particular, $C_{\pi_0} = 0$. By induction one gets $C_\pi = 0$ for all $\pi \in S$, i.e. the set of monomial vectors $u(\pi)v_{k\Lambda_0}$ with $k\Lambda_0$ -admissible π is linearly independent, and hence a basis of $L(k\Lambda_0)$. \square

ACKNOWLEDGEMENT

This work has been supported in part by Croatian Science Foundation under the project IP-2022-10-9006.

REFERENCES

- [B] I. Baranović, *Combinatorial bases of Feigin-Stoyanovsky's type subspaces of level 2 standard modules for $D_4^{(1)}$* , Comm. Algebra **39** (2011), 1007–1051.
- [BPT] I. Baranović, M. Primc and G. Trupčević, *Bases of Feigin-Stoyanovsky's type subspaces for $C_\ell^{(1)}$* , The Ramanujan Journal **45** (2018), 265–289.
- [CLM1] S. Capparelli, J. Lepowsky and A. Milas, *The Rogers-Ramanujan recursion and intertwining operators*, Comm. Contemporary Math. **5** (2003), 947–966.
- [CLM2] S. Capparelli, J. Lepowsky and A. Milas, *The Rogers-Selberg recursions, the Gordon-Andrews identities and intertwining operators*, Ramanujan J. **12** (2006), 379–397.
- [CMPP] S. Capparelli, A. Meurman, A. Primc and M. Primc *New partition identities from $C_\ell^{(1)}$ -modules*, Glasnik Matematički **57** (2022), 161–184.
- [DK] J. Dousse and I. Konan, *Characters of level 1 standard modules of $C_n^{(1)}$ as generating functions for generalised partitions*. arXiv:2212.12728 [math.CO], 2022.
- [FKLMM] B. Feigin, R. Kedem, S. Loktev, T. Miwa and E. Mukhin, *Combinatorics of the $\widehat{\mathfrak{sl}}_2$ spaces of coinvariants*, Transformation Groups **6** (2001), 25–52.
- [FS] A. V. Stoyanovsky and B. L. Feigin, *Functional models of the representations of current algebras, and semi-infinite Schubert cells*, (Russian) Funktsional. Anal. i Prilozhen. **28** (1994), no. 1, 68–90, 96; translation in Funct. Anal. Appl. **28** (1994), no. 1, 55–72; preprint B. Feigin and A. Stoyanovsky, *Quasi-particles models for the representations of Lie algebras and geometry of flag manifold*, hep-th/9308079, RIMS 942.
- [F] E. Feigin, *The PBW filtration*, Represent. Theory **13** (2009), 165–181.
- [H] J. E. Humphreys, *Introduction to Lie Algebras and Representation Theory*, Springer, New York, 1972.
- [J1] M. Jerković, *Recurrence relations for characters of affine Lie algebra $A_\ell^{(1)}$* , J. Pure Appl. Algebra **213** (2009), 913–926.

- [J2] M. Jerković, *Recurrences and characters of Feigin-Stoyanovsky's type subspaces*, Vertex operator algebras and related areas, 113–123, *Contemp. Math.* **497**, Amer. Math. Soc., Providence, 2009.
- [K] V.G. Kac, *Infinite-dimensional Lie algebras*, 3rd ed. Cambridge University Press, Cambridge, 1990.
- [MP] A. Meurman and M. Primc, *Annihilating fields of standard modules of $sl(2, \mathbb{C})$ and combinatorial identities*, *Memoirs of the Amer. Math. Soc.* **137**, No. 652 (1999).
- [P1] M. Primc, *(k, r) -admissible configurations and intertwining operators*, *Contemp. Math.* **442** (2007), 425–434.
- [P2] M. Primc, *Combinatorial bases of modules for affine Lie algebra $B_2^{(1)}$* , *Cent. Eur. J. Math.* **11** (2013), 197–225.
- [P3] M. Primc, *Some combinatorial coincidences for standard representations of affine Lie algebras*, in the book D. Adamović, P. Papi, *Affine, Vertex and W -algebras*, Springer INdAM Series (SINDAMS, Vol. **37**) (2019), 203–218.
- [PŠ1] M. Primc and T. Šikić, *Combinatorial bases of basic modules for affine Lie algebras $C_n^{(1)}$* , *J. Math. Phys.*, **57** (2016), 1–19.
- [PŠ2] M. Primc and T. Šikić, *Leading terms of relations for standard modules of affine Lie algebras $C_n^{(1)}$* , *Ramanujan J.* **48** (2019), 509–543.
- [PŠ3] M. Primc and T. Šikić, *Combinatorial relations for level 2 standard $C_n^{(1)}$ -modules*, *J. Math. Phys.* **64** (2023); doi: 10.1063/5.0145719.
- [R] M. C. Russell, *A companion to the Andrews-Gordon and Andrews-Bressoud identities, and recent conjectures of Capparelli, Meurman, Primc, and Primc*, arXiv:2306.16251.
- [T1] G. Trupčević, *Combinatorial bases of Feigin-Stoyanovsky's type subspaces of level 1 standard $\tilde{sl}(\ell + 1, \mathbb{C})$ -modules*, *Comm. Algebra* **38** (2010), 3913–3940.
- [T2] G. Trupčević, *Combinatorial bases of Feigin-Stoyanovsky's type subspaces of higher-level standard $\tilde{sl}(\ell + 1, \mathbb{C})$ -modules*, *J. Algebra* **322** (2009), 3744–3774.
- [T3] G. Trupčević, *Bases of standard modules for affine Lie algebras of type $C_\ell^{(1)}$* , *Communications in Algebra* **46 (8)** (2018), 3663–3673.

(M. Primc) FACULTY OF SCIENCE, UNIVERSITY OF ZAGREB, ZAGREB, CROATIA
Email address: primc@math.hr

(G. Trupčević) FACULTY OF TEACHER EDUCATION, UNIVERSITY OF ZAGREB, ZAGREB, CROATIA
Email address: goran.trupcevic@ufzg.hr