

Sheaves of (∞, ∞) -categories

Zach Goldthorpe

March 2024

Abstract

We provide a functorial presentation of the $(\infty, 1)$ -category of sheaves of (n, r) -categories for all $-2 \leq n \leq \infty$ and $0 \leq r \leq n + 2$ based on complete Segal space objects. In this definition, the equivalences of sheaves of (∞, ∞) -categories are defined inductively, so we also provide a localisation at the coinductive equivalences to define the $(\infty, 1)$ -category of sheaves of ω -categories as well.

We prove that the $(\infty, 1)$ -category of sheaves of (∞, ∞) -categories, and the $(\infty, 1)$ -category of sheaves of ω -categories both define distributors over the underlying topos of sheaves of spaces. Moreover, we show that these distributors define terminal and initial fixed points with respect to the construction of complete Segal space objects in a distributor.

We conclude with a sheafification result: the category of sheaves of (n, r) -categories over a site \mathcal{C} can be presented as a strongly reflective localisation of $\mathbf{Fun}(\mathcal{C}^{\text{op}}, \mathbf{Cat}_{(n,r)})$, where the localisation functor preserves fibre products over $\mathbf{Sh}(\mathcal{C})$, with the analogous result also holding for sheaves of ω -categories.

Contents

1	Introduction	1
1.1	Organisation of paper	3
1.2	Notation and terminology	3
2	Sheaves of higher categories	4
2.1	Polysimplicial sheaves	4
2.2	Higher category objects in sheaves	6
2.3	Sheaves of (∞, ∞) -categories and of ω -categories	8
3	Distributors and complete Segal spaces	11
3.1	Preliminaries	11
3.2	The ∞ -categories of distributors	12
3.3	Functoriality of complete Segal spaces	14
3.4	Distributors of sheaves of higher categories	16
3.5	Geometric morphisms and sheafification	18
	References	21

1 Introduction

A higher category is a generalisation of an ordinary category consisting of objects, 1-morphisms between objects, 2-morphisms between 1-morphisms, 3-morphisms between 2-morphisms, *ad infinitum*. For $-2 \leq n \leq \infty$ and $0 \leq r \leq n + 2$, an (n, r) -category is a higher category where

- parallel k -morphisms are equivalent for $k > n$, and
- k -morphisms are invertible for $k > r$.

For $r < \infty$, the notion of (n, r) -category is unambiguous up to an action of $(\mathbb{Z}/2)^r$ by [BSP21]. However, there are several inequivalent notions of (∞, ∞) -category, due to nuances in the concept of invertibility in a purely infinite-dimensional setting.

We have a tower of inclusions

$$\mathbf{Cat}_{(\infty,0)} \subseteq \mathbf{Cat}_{(\infty,1)} \subseteq \mathbf{Cat}_{(\infty,2)} \subseteq \dots$$

where each inclusion $\mathbf{Cat}_{(\infty,r)} \subseteq \mathbf{Cat}_{(\infty,r+1)}$ admits a left adjoint $\pi_{\leq r}$ and a right adjoint $\kappa_{\leq r}$. For an $(\infty, r+1)$ -category \mathcal{C} , we obtain the (∞, r) -category $\pi_{\leq r}\mathcal{C}$ by formally inverting the $(r+1)$ -morphisms of \mathcal{C} . On the other hand, $\kappa_{\leq r}\mathcal{C}$ is just the maximal sub- (∞, r) -category of \mathcal{C} , obtained by discarding the non-invertible $(r+1)$ -morphisms. From these adjoints, we obtain two natural constructions of the $(\infty, 1)$ -category of (∞, ∞) -categories:

$$\begin{aligned} \mathbf{Cat}_{(\infty,\infty)} &:= \varprojlim \left(\dots \rightarrow \mathbf{Cat}_{(\infty,2)} \xrightarrow{\kappa_{\leq 1}} \mathbf{Cat}_{(\infty,1)} \xrightarrow{\kappa_{\leq 0}} \mathbf{Cat}_{(\infty,0)} \right) \\ \mathbf{Cat}_{\omega} &:= \varprojlim \left(\dots \rightarrow \mathbf{Cat}_{(\infty,2)} \xrightarrow{\pi_{\leq 1}} \mathbf{Cat}_{(\infty,1)} \xrightarrow{\pi_{\leq 0}} \mathbf{Cat}_{(\infty,0)} \right) \end{aligned}$$

Roughly speaking, $\mathbf{Cat}_{(\infty,\infty)}$ consists of higher categories wherein equivalences are defined inductively from the identity k -morphisms, whereas \mathbf{Cat}_{ω} consists of higher categories wherein equivalences are defined coinductively; see [Gol23, Remark 3.0.5].

There are several approaches to defining higher categories, but they are largely guided by one of the two principles:

- An $(n+1, r+1)$ -category is a category that is suitably *enriched* in (n, r) -categories; that is, an $(n+1, r+1)$ -category is a suitable category \mathcal{C} such that $\mathrm{Hom}_{\mathcal{C}}(x, y)$ is an (n, r) -category for every $x, y \in \mathcal{C}$.
- An $(n+1, r+1)$ -category is a suitable category object *internal* to (n, r) -categories; that is, an $(n+1, r+1)$ -category consists of an (n, r) -category \mathcal{C}_0 of objects, and an (n, r) -category \mathcal{C}_1 of morphisms, such that $(\mathcal{C}_0, \mathcal{C}_1)$ satisfies a suitable analogue of the axioms of category theory.

Although enrichment provides a more concise inductive description of higher categories— $(n+1, r+1)$ -categories should be precisely the categories enriched in (n, r) -categories—a suitably general theory of enrichment proved too difficult to develop without already establishing a sufficiently general theory of higher categories.

On the other hand, a suitable theory of internalisation was more readily available using the tools from abstract homotopy theory. The caveat is that a general category object \mathcal{C}_{\bullet} in $\mathbf{Cat}_{(n,r)}$ has more structure than that of an $(n+1, r+1)$ -category. Between objects in \mathcal{C}_0 , there are two distinct notions of a morphism between them: there are the objects of \mathcal{C}_1 , and there are also the morphisms of \mathcal{C}_0 .

One way to avoid this caveat is to assert that \mathcal{C}_0 is a set; that is, \mathcal{C}_0 is a $(0, 0)$ -category. We can then think of \mathcal{C}_0 as the underlying set of objects of the $(n+1, r+1)$ -category. This leads to the notion of higher *Segal categories*, which is developed in detail, for instance, in [Sim11]. However, the “underlying set” of objects in an ordinary category is not invariant under equivalence, and this complicates the notion of equivalence between higher Segal categories. A functor $\mathcal{C}_{\bullet} \rightarrow \mathcal{D}_{\bullet}$ of $(n+1, r+1)$ -categories should be an equivalence precisely if it is fully faithful on the (n, r) -categories of morphisms, and essentially surjective on the objects. Since essential surjectivity cannot be tested on the underlying functor $\mathcal{C}_0 \rightarrow \mathcal{D}_0$, it must be defined by extracting the higher equivalences from the categorical structure of \mathcal{C}_{\bullet} and \mathcal{D}_{\bullet} .

Alternatively, we can instead assert that \mathcal{C}_0 is an ∞ -groupoid; that is, all k -morphisms in \mathcal{C}_0 are invertible. Then, \mathcal{C}_0 serves as the underlying n -groupoid of \mathcal{C}_{\bullet} , which is an invariant of \mathcal{C}_{\bullet} under equivalence. By the Homotopy Hypothesis, we can therefore view \mathcal{C}_0 as a space. This leads to the notion of higher *Segal spaces*. However, this means that the objects of \mathcal{C}_{\bullet} have two distinct notions of equivalence:

- Path-connectedness in the space \mathcal{C}_0 , and

- Isomorphism from the categorical structure of \mathcal{C}_\bullet .

Therefore, we restrict to *complete Segal spaces*, which are those Segal spaces for which these two notions of equivalence coincide. Although the model for $(n+1, r+1)$ -categories becomes more technical, requiring the completeness condition, the resulting equivalences are easier to define: a functor $\mathcal{C}_\bullet \rightarrow \mathcal{D}_\bullet$ of complete Segal spaces modelling $(n+1, r+1)$ -categories is an equivalence precisely if $\mathcal{C}_0 \rightarrow \mathcal{D}_0$ is a homotopy equivalence (giving essential surjectivity), and $\mathcal{C}_1 \rightarrow \mathcal{D}_1$ is an equivalence of (n, r) -categories (giving fully faithfulness).

Complete Segal spaces were first defined by Rezk in [Rez01]. Barwick then defined r -fold iterated complete Segal spaces as a model for (∞, r) -categories in [Bar05] (see [BSP21, §14]). Lurie expanded this construction and axiomatised its iterative nature in [Lur09b, §1], providing a means to construct complete Segal space objects in any *distributor*.

Intuitively, a distributor \mathcal{Y} is a locally presentable $(\infty, 1)$ -category with a choice of full subcategory $\mathcal{X} \subseteq \mathcal{Y}$ consisting of the “spaces.” Then, the $(\infty, 1)$ -category $\mathbf{CSS}_{\mathcal{X}}(\mathcal{Y})$ of complete Segal space object in \mathcal{Y} consists of categories \mathcal{C}_\bullet internal to \mathcal{Y} such that $\mathcal{C}_0 \in \mathcal{X}$ (i.e., \mathcal{C}_0 is “space-like”), and \mathcal{C}_\bullet is complete. The axioms of a distributor imply that the subcategory \mathcal{X} of “spaces” is always an $(\infty, 1)$ -topos; that is, an $(\infty, 1)$ -category of sheaves of spaces. In particular, r -fold complete Segal spaces over \mathcal{X} define a model for sheaves of (∞, r) -categories.

In this paper, we use the language of complete Segal spaces to define a presentation of the $(\infty, 1)$ -category of sheaves of (n, r) -categories or ω -categories over an $(\infty, 1)$ -topos \mathcal{X} as a localisation of r -fold simplicial objects in \mathcal{X} . We then show that both the $(\infty, 1)$ -categories of sheaves of (∞, ∞) -categories, and of ω -categories, form distributors over \mathcal{X} . In fact, both categories are completely characterised by $\mathbf{CSS}_{\mathcal{X}}$. More precisely, we construct an $(\infty, 1)$ -category $\mathbf{Dist}_{\mathcal{X}}^R$ of distributors over \mathcal{X} and show that sheaves of (∞, ∞) -categories and of ω -categories satisfy universal properties with respect to the construction of complete Segal space objects:

Theorem A (Theorems 3.4.3 and 3.4.5). *The $(\infty, 1)$ -category of sheaves of (∞, ∞) -categories is the terminal object in the $(\infty, 1)$ -category $\mathbf{Fix}_{\mathbf{CSS}}^R$ of distributors \mathcal{Y} over \mathcal{X} with an equivalence $\mathcal{Y} \xrightarrow{\sim} \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y})$. On the other hand, the $(\infty, 1)$ -category of sheaves of ω -categories is the initial object in $\mathbf{Fix}_{\mathbf{CSS}}^R$.*

1.1 Organisation of paper

Section 2 focuses on an explicit presentation of sheaves of higher categories. In Section 2.1, we define polysimplicial sheaves to serve as the underlying structure of sheaves of higher categories. We then provide an explicit definition of sheaves of (n, r) -categories for $-2 \leq n \leq \infty$ and $0 \leq r \leq n+2$ in terms of polysimplicial sheaves in Section 2.2. In Section 2.3, we prove that sheaves of (∞, ∞) -categories can be constructed as a suitable limit of sheaves of (∞, r) -categories, analogous to the construction of $\mathbf{Cat}_{(\infty, \infty)}$. We then define sheaves of ω -categories, and characterise those sheaves of (∞, ∞) -categories that are sheaves of ω -categories in terms of truncation towers in Corollary 2.3.5.

In Section 3, we compare the constructions of Section 2 to the construction of complete Segal spaces in distributors. We recall the preliminary theory of distributors and complete Segal spaces in Section 3.1. Then, in Section 3.2, we construct the $(\infty, 1)$ -category $\mathbf{Dist}_{\mathcal{X}}^R$, and prove that it has all small limits and colimits. We show that the construction of complete Segal spaces is functorial over $\mathbf{Dist}_{\mathcal{X}}^R$ in Section 3.3, and moreover that it preserves certain limits and colimits. In Section 3.4, we show that the constructions in Section 2 can be expressed in terms of complete Segal space objects in distributors, and prove Theorem A. Then, we briefly explore the functoriality of the construction of sheaves of higher categories in Section 3.5, and moreover prove that presheaves of higher categories admit a relatively left exact sheafification functor.

1.2 Notation and terminology

In this paper, unless otherwise specified, unprefix categorical notions refer to their $(\infty, 1)$ -categorical analogues: “categories” are $(\infty, 1)$ -categories, “topoi” are $(\infty, 1)$ -topoi, and “(co)limits” are $(\infty, 1)$ -(co)limits.

Let $\mathbf{\Delta}$ denote the simplex category, and let \mathcal{S} denote the category of homotopy types. Given a small category \mathcal{C} , let $\mathcal{P}(\mathcal{C}) := \mathbf{Fun}(\mathcal{C}^{\text{op}}, \mathcal{S})$ denote the category of presheaves on \mathcal{C} .

The categories $\mathbf{Cat}_{(n,r)}$ for $-2 \leq n \leq \infty$ and $0 \leq r \leq n$ and \mathbf{Cat}_ω refer to the categories of (small) higher categories defined in [Gol23, Definitions 3.0.1 and 3.0.4]. On the other hand, let $\widehat{\mathbf{Cat}}_\infty$ denote the huge category of large categories.

For a regular cardinal λ , say that a functor is λ -*accessible* if it preserves λ -filtered colimits. In particular, say that a functor is *accessible* if it is ω -accessible. A category \mathcal{C} is said to be a λ -*accessible localisation* of \mathcal{D} if \mathcal{C} embeds fully faithfully into \mathcal{D} via a λ -accessible right adjoint, and the localisation is moreover said to be *left exact* if the corresponding left adjoint is left exact.

For a regular cardinal λ , say that a category is *locally λ -presentable* if it is a λ -accessible localisation of a presheaf category (which are called “ λ -compactly generated” categories in [Lur09a]), and say that a category is *locally presentable* if it is locally λ -presentable for some $\lambda \gg 0$ (which are called “presentable” categories in [Lur09a]). Denote by \mathbf{Pres}_∞^L or \mathbf{Pres}_∞^R the huge categories of locally presentable categories, and left or right adjoints, respectively.

Say that a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ *creates limits* if any $p : K^\triangleleft \rightarrow \mathcal{C}$ is a limit cone if and only if $F \circ p$ is a limit cone. The concept of creating colimits is entirely dual.

2 Sheaves of higher categories

Throughout this section, fix a topos \mathcal{X} , presented as a left exact accessible localisation of $\mathcal{P}(\mathcal{C})$ for some small category \mathcal{C} , where the localisation functor is denoted $(-)^{\#} : \mathcal{P}(\mathcal{C}) \rightarrow \mathcal{X}$. The goal of this section is to explicitly construct the category $\mathbf{Sh}_{(n,r)}(\mathcal{C})$ of sheaves of (n,r) -categories over \mathcal{C} , where $-2 \leq n \leq \infty$ and $0 \leq r \leq n + 2$, as well as the category $\mathbf{Sh}_\omega(\mathcal{C})$ of sheaves of ω -categories.

2.1 Polysimplicial sheaves

Definition 2.1.1. Define the *polysimplex category* Σ_∞ as the following subcategory of $\Delta^{\times\infty}$.

- The objects of Σ_∞ are infinite sequences $[\vec{k}] = (k_0, k_1, k_2, \dots)$ of simplices such that there exists some $r \geq 0$ such that $k_n = 0$ if and only if $n \geq r$. Call r the *dimension* of $[\vec{k}]$, denoted $\dim[\vec{k}] := r$.
- The morphisms $\vec{\phi} : [\vec{k}] \rightarrow [\vec{\ell}]$ are sequences of morphisms $\phi_n : [k_n] \rightarrow [\ell_n]$ in Δ such that there exists some $r \geq 0$ such that $\phi_n = 0$ if and only if $n \geq r$.

Denote by Σ_r the full subcategory of Σ_∞ spanned by polysimplices of dimension $\leq r$.

Remark 2.1.2. The category Σ_r is equivalent to the quotient category Θ^r of $\Delta^{\times r}$ defined in [Sim00, §2]. We choose different notation so as to not confuse the polysimplex category with Joyal’s disk category.

Definition 2.1.3. For a polysimplex $[\vec{k}]$ and $r \geq 0$, define the polysimplices

$$\begin{aligned} [\vec{k}]_{<r} &:= (k_0, k_1, k_2, \dots, k_{r-1}, 0, \dots) \\ [\vec{k}]_{\geq r} &:= (k_r, k_{r+1}, k_{r+2}, \dots) \end{aligned}$$

These constructions extend to define endofunctors on Σ_∞ . Note that $[-]_{<r}$ restricts to a right adjoint to the inclusion $\Sigma_r \subseteq \Sigma_\infty$.

Given two polysimplices $[\vec{k}]$ and $[\vec{\ell}]$, let $[\vec{k}, \vec{\ell}]$ denote their concatenation; that is, the unique polysimplex such that $[\vec{k}, \vec{\ell}]_{<r} = [\vec{k}]$ and $[\vec{k}, \vec{\ell}]_{\geq r} = [\vec{\ell}]$, where $r = \dim[\vec{k}]$.

Definition 2.1.4. A *polysimplicial sheaf* over $(\mathcal{C}, (-)^{\#})$ is a polysimplicial object in \mathcal{X} ; that is, a functor $\Sigma_\infty^{\text{op}} \rightarrow \mathcal{X}$.

For a polysimplicial sheaf $X : \Sigma_\infty^{\text{op}} \rightarrow \mathcal{X}$ and a polysimplex $[\vec{k}]$, let $\Delta[\vec{k}, X] : \Sigma_\infty^{\text{op}} \rightarrow \mathcal{X}$ denote the polysimplicial sheaf given by the left Kan extension of X along the functor $[\vec{k}, -] : \Sigma_\infty \rightarrow \Sigma_\infty$. By definition, it follows for all polysimplicial sheaves X' that we have a natural equivalence

$$\text{Map}(\Delta[\vec{k}, X], X') \simeq \text{Map}(X, X'_{\vec{k}, \bullet})$$

We are also interested in a couple of special cases:

- For a polysimplex $[\vec{k}]$ and object $U \in \mathcal{C}$, define the U -local \vec{k} -polysimplex to be $\Delta_U[\vec{k}] := \Delta[\vec{k}, h_U^\#]$, which corepresents the functor $\mathbf{Fun}(\Sigma_\infty^{\text{op}}, \mathcal{X}) \rightarrow \mathcal{S}$ mapping $X_\bullet \mapsto X_{\vec{k}}(U)$.
- Define the *suspension* of a polysimplicial sheaf X to be $\Sigma X := \Delta[1, X]$, which is left adjoint to the *desuspension* $\Omega X' := X'_{1, \bullet}$.

Remark 2.1.5. The functor $[0, -] : \Sigma_\infty \rightarrow \Sigma_\infty$ is constant on the terminal object $[0] \in \Sigma_\infty$. In particular, the polysimplicial sheaf $\Delta[0, X]$ is the terminal polysimplicial sheaf, so we may denote this object by $\Delta[0]$.

On the other hand, the U -local 0-polysimplex $\Delta_U[0]$ is *not* the terminal object; rather, it is the corepresenting object for the functor mapping $X_\bullet \mapsto X_0(U)$. This is precisely the representable sheaf $\Delta_U[0] \simeq h_U^\#$.

Remark 2.1.6. Since $\mathbf{Fun}(\Sigma_\infty^{\text{op}}, \mathcal{X})$ is a localisation of $\mathcal{P}(\Sigma_\infty \times \mathcal{C})$, every polysimplicial sheaf is a canonical colimit of the local polysimplices $\Delta_U[\vec{k}]$.

Remark 2.1.7. Left Kan extension along $\Sigma_r \subseteq \Sigma_\infty$ defines a fully faithful functor $\mathbf{Fun}(\Sigma_r^{\text{op}}, \mathcal{X}) \subseteq \mathbf{Fun}(\Sigma_\infty^{\text{op}}, \mathcal{X})$. We may refer to objects in the essential image of this inclusion as r -polysimplicial sheaves.

Explicitly, we view $X : \Sigma_r^{\text{op}} \rightarrow \mathcal{X}$ as an r -polysimplicial sheaf by taking $X_{\vec{k}} := X_{\vec{k} < r}$ for every $[\vec{k}] \in \Sigma_\infty$.

We now establish some useful preliminary results before studying higher category objects in \mathcal{X} .

Definition 2.1.8. For a polysimplex $[\vec{k}]$ and an object $U \in \mathcal{C}$, define the U -local \vec{k} -spine to be the pushout

$$\text{Sp}_U[\vec{k}] := \underbrace{\Delta_U[1, \vec{k}_{\geq 1}] \sqcup_{\Delta[0]} \cdots \sqcup_{\Delta[0]} \Delta_U[1, \vec{k}_{\geq 1}]}_{k_0}$$

in $\mathbf{Fun}(\Sigma_\infty^{\text{op}}, \mathcal{X})$, induced by the inert maps $[0], [1] \hookrightarrow [k_0]$ in Δ . In particular, the U -local \vec{k} -spine admits a canonical inclusion $\text{Sp}_U[\vec{k}] \subseteq \Delta_U[\vec{k}]$.

Recall from [Lur09a, Definition 5.5.4.5] that a class W of morphisms in a category \mathcal{D} is *strongly saturated* if it is closed under pushouts along arbitrary morphisms of \mathcal{D} , it is closed under colimits in $\mathbf{Fun}([1], \mathcal{D})$, and it satisfies 2-out-of-3.

Lemma 2.1.9. For $0 \leq r \leq \infty$, let $X : \Sigma_r^{\text{op}} \rightarrow \mathcal{X}$. If $[\ell] = [\ell^1] \sqcup_{[0]} [\ell^2]$ in Δ , then the inclusion

$$\Delta[\ell^1, X] \sqcup_{\Delta[0]} \Delta[\ell^2, X] \hookrightarrow \Delta[\ell, X]$$

lies in the strongly saturated class of morphisms generated by the local spine inclusions in $\mathbf{Fun}(\Sigma_{r+1}^{\text{op}}, \mathcal{X})$.

Proof. By definition, the strongly saturated class of morphisms generated by the local spine inclusions is closed under colimits in $\mathbf{Fun}([1], \mathbf{Fun}(\Sigma_{r+1}^{\text{op}}, \mathcal{X}))$. Therefore, Remark 2.1.6 allows us to reduce to the case where $X \simeq \Delta_U[\vec{k}]$. In this situation, consider the composite

$$\text{Sp}_U[\ell, \vec{k}] \xrightarrow{\sim} \text{Sp}_U[\ell^1, \vec{k}] \sqcup_{\Delta[0]} \text{Sp}_U[\ell^2, \vec{k}] \xrightarrow{i} \Delta_U[\ell^1, \vec{k}] \sqcup_{\Delta[0]} \Delta_U[\ell^2, \vec{k}] \xrightarrow{j} \Delta_U[\ell, \vec{k}]$$

The map j is precisely the inclusion $\Delta[\ell^1, X] \sqcup_{\Delta[0]} \Delta[\ell^2, X] \hookrightarrow \Delta[\ell, X]$. Note that the composite is a local spine inclusion, and i is a pushout of local spine inclusions. Therefore, j is in the strongly saturated class by 2-out-of-3. \square

Proposition 2.1.10. For $0 \leq r \leq \infty$, let W be any class of morphisms in $\mathbf{Fun}(\Sigma_r^{\text{op}}, \mathcal{X})$. Then, the following classes generate the same strongly saturated class of morphisms in $\mathbf{Fun}(\Sigma_{r+1}^{\text{op}}, \mathcal{X})$:

- (1) $\{\Delta[\ell^1, X] \sqcup_{\Delta[0]} \Delta[\ell^2, X] \hookrightarrow \Delta[\ell, X] : [\ell] = [\ell^1] \sqcup_{[0]} [\ell^2]; X : \Sigma_r^{\text{op}} \rightarrow \mathcal{X}\} \cup \{\Delta[\ell, f] : | : f \in W; \ell > 0\}$
- (2) $\{\text{Sp}_U[\vec{k}] \hookrightarrow \Delta_U[\vec{k}] : | : [\vec{k}] \in \Sigma_{r+1}; U \in \mathcal{C}\} \cup \{\Sigma f : | : f \in W\}$

Proof. Let W_1 and W_2 denote the strongly saturated classes of morphisms generated by (1) and (2), respectively, then clearly $W_2 \subseteq W_1$.

Conversely, any $\Delta[\ell^1, X] \sqcup_{\Delta[0]} \Delta[\ell^2, X] \hookrightarrow \Delta[\ell, X]$ lies in W_2 by Lemma 2.1.9. For a polysimplicial sheaf X_\bullet , let

$$\mathrm{Sp}[\ell, X] := \underbrace{\Sigma X \sqcup_{\Delta[0]} \cdots \sqcup_{\Delta[0]} \Sigma X}_\ell \hookrightarrow \Delta[\ell, X]$$

then this inclusion also lies in W_2 .

For $f : X_\bullet \rightarrow X'_\bullet$ in W , the induced map $\mathrm{Sp}[\ell, X] \rightarrow \mathrm{Sp}[\ell, X']$ is a pushout of elements of W_2 and is thus an element of W_2 as well. From the commutative square

$$\begin{array}{ccc} \mathrm{Sp}[\ell, X] & \xrightarrow{\in W_2} & \Delta[\ell, X] \\ \in W_2 \downarrow & & \downarrow \Delta[\ell, f] \\ \mathrm{Sp}[\ell, X'] & \xrightarrow{\in W_2} & \Delta[\ell, X'] \end{array}$$

it follows from 2-out-of-3 that $\Delta[\ell, f] \in W_2$, as desired. \square

2.2 Higher category objects in sheaves

Definition 2.2.1. For $U \in \mathcal{C}$, left Kan extension of $\Delta_U[-] : \mathbf{\Delta} \rightarrow \mathbf{Fun}(\Sigma_r^{\mathrm{op}}, \mathcal{X})$ for any $0 \leq r \leq \infty$ along the Yoneda embedding $\mathbf{\Delta} \hookrightarrow \mathcal{P}(\mathbf{\Delta})$ induces a U -local realisation functor $|-|_U^\# : \mathcal{P}(\mathbf{\Delta}) \rightarrow \mathbf{Fun}(\Sigma_r^{\mathrm{op}}, \mathcal{X})$. By definition, this realisation is left adjoint to the functor $\mathbf{Fun}(\Sigma_r^{\mathrm{op}}, \mathcal{X}) \rightarrow \mathcal{P}(\mathbf{\Delta})$ sending $X_\bullet \mapsto X_\bullet(U)$.

Definition 2.2.2. Let $\mathbb{I} : \mathbf{\Delta}^{\mathrm{op}} \rightarrow \mathbf{Set}$ denote the nerve of the walking isomorphism. Explicitly, we have $\mathbb{I}_n = \{\perp, \top\}^{\times(n+1)}$ for all $n \geq 0$, where the face maps are given by projections, and the degeneracy maps are given by diagonal inclusions. On the other hand, let $\mathbb{S}^m := \partial\Delta[m+1]$ denote the simplicial m -sphere for $m \geq -1$.

For $-2 \leq n \leq \infty$ and $0 \leq r \leq n+2$, let $W_{(n,r)}$ denote the strongly saturated class of morphisms in $\mathbf{Fun}(\Sigma_\infty^{\mathrm{op}}, \mathcal{X})$ generated by:

- (W1) $\Sigma^m \mathrm{Sp}_U[\vec{k}] \hookrightarrow \Sigma^m \Delta_U[\vec{k}]$ for $U \in \mathcal{C}$, $m \geq 0$, and $[\vec{k}] \in \Sigma_\infty$,
- (W2) $\Sigma^m |\mathbb{I}|_U^\# \rightarrow \Sigma^m \Delta_U[0]$ for $U \in \mathcal{C}$, and $m \geq 0$,
- (W3) $\Sigma^m (\mathbb{S}^{k-m} \times \Delta_U[0]) \rightarrow \Sigma^m \Delta_U[0]$ for $U \in \mathcal{C}$, and finite $m \geq r$ and $k > n$, and
- (W4) $\Sigma^m \Delta_U[1] \rightarrow \Sigma^m \Delta_U[0]$ for $U \in \mathcal{C}$, and finite $m \geq r$.

Call a polysimplicial sheaf a *sheaf of (n, r) -categories* on \mathcal{C} if it is $W_{(n,r)}$ -local. Denote by $\mathbf{Sh}_{(n,r)}(\mathcal{C})$ the full subcategory of $\mathbf{Fun}(\Sigma_\infty^{\mathrm{op}}, \mathcal{X})$ spanned by the sheaves of (n, r) -categories.

Remark 2.2.3. Following the proof of [Lur09b, Proposition 1.5.4], suppose \mathcal{M} is a left proper combinatorial simplicial model category such that its underlying ∞ -category is a topos \mathcal{X} . For $-2 \leq n \leq \infty$ and $0 \leq r \leq n+2$, we can endow $\mathbf{Fun}(\Sigma_\infty^{\mathrm{op}}, \mathcal{M})$ with a simplicial model structure such that

- (C) A morphism $f : X_\bullet \rightarrow X'_\bullet$ is a cofibration precisely if the map $X_{\vec{k}} \rightarrow X'_{\vec{k}}$ is a cofibration in \mathcal{M} for every $[\vec{k}] \in \Sigma_\infty$,
- (W) pointwise weak equivalences are among the weak equivalences in $\mathbf{Fun}(\Sigma_\infty^{\mathrm{op}}, \mathcal{M})$, and
- (F) An object X is fibrant if and only if it is injectively fibrant, and the induced map $N(\Sigma_\infty^{\mathrm{op}})^{\mathrm{op}} \rightarrow N(X)$ in the underlying ∞ -category $\mathbf{Fun}(\Sigma_\infty^{\mathrm{op}}, \mathcal{X})$ is a sheaf of (n, r) -categories.

In fact, by Lemma 2.2.8 below, we can restrict to a Quillen equivalent model structure on the subcategory $\mathbf{Fun}(\Sigma_r^{\mathrm{op}}, \mathcal{M})$.

Remark 2.2.4. We can generalise Definition 2.2.1 to any $X : \Sigma_r^{\text{op}} \rightarrow \mathcal{X}$ by taking the left Kan extension of $\Delta[-, X] : \Delta \rightarrow \mathbf{Fun}(\Sigma_{r+1}^{\text{op}}, \mathcal{X})$ along the Yoneda embedding $\Delta \hookrightarrow \mathcal{P}(\Delta)$. This defines an action $(-) \odot X : \mathcal{P}(\Delta) \rightarrow \mathbf{Fun}(\Sigma_{r+1}^{\text{op}}, \mathcal{X})$.

By Remark 2.1.6, we have for any simplicial space S that locality with respect to $\Sigma^m | S|_U^\# \rightarrow \Sigma^m \Delta_U[0]$ for all $U \in \mathcal{C}$ is equivalent to locality with respect to $\Sigma^m(S \odot X) \rightarrow \Sigma^m X$ for all polysimplicial sheaves X . In particular, combined with Proposition 2.1.10, we see that $W_{(n,r)}$ is independent of the choice of site \mathcal{C} . Therefore, we may also denote the category of sheaves of (n, r) -categories on (a site of) \mathcal{X} by $\mathbf{Sh}_{(n,r)}(\mathcal{X})$ to avoid specifying a site of definition.

Below, we provide a recursive characterisation of sheaves of (n, r) -categories.

Definition 2.2.5. For $0 \leq r \leq \infty$, the canonical inclusion $\Sigma_{r+1} \subseteq \Delta \times \Sigma_r$ admits a left adjoint given by the mapping

$$([n], [\vec{k}]) \mapsto \begin{cases} [0], & n = 0 \\ [n, \vec{k}], & n > 0 \end{cases}$$

which induces a fully faithful inclusion $\mathbf{Fun}(\Sigma_{r+1}^{\text{op}}, \mathcal{X}) \hookrightarrow \mathbf{Fun}(\Delta^{\text{op}}, \mathbf{Fun}(\Sigma_r^{\text{op}}, \mathcal{X}))$.

Proposition 2.2.6. For $-2 \leq n \leq \infty$ and $0 \leq r \leq n + 2$, the essential image of the fully faithful inclusion

$$\mathbf{Sh}_{(n+1, r+1)}(\mathcal{C}) \subseteq \mathbf{Fun}(\Sigma_\infty^{\text{op}}, \mathcal{X}) \hookrightarrow \mathbf{Fun}(\Delta^{\text{op}}, \mathbf{Fun}(\Sigma_\infty^{\text{op}}, \mathcal{X}))$$

consists of those $F : \Delta^{\text{op}} \rightarrow \mathbf{Fun}(\Sigma_\infty^{\text{op}}, \mathcal{X})$ such that

- (S1) $F_0 : \Sigma_\infty^{\text{op}} \rightarrow \mathcal{X}$ is essentially constant,
- (S2) F_1 is a sheaf of (n, r) -categories over \mathcal{C} ,
- (S3) F satisfies the Segal condition: for every $k \geq 2$, the canonical map

$$F_k \rightarrow \underbrace{F_1 \times_{F_0} \cdots \times_{F_0} F_1}_k$$

is an equivalence in $\mathbf{Fun}(\Sigma_\infty^{\text{op}}, \mathcal{X})$, and

- (S4) F is Rezk-complete: it is local with respect to $|\mathbb{I}|_U^\# \rightarrow \Delta_U[0]$ for every $U \in \mathcal{C}$.

Remark 2.2.7. If F satisfies (S1), (S2), and (S3) and $n = \infty$, then F_k is a sheaf of (∞, r) -categories over \mathcal{C} for every $k \geq 0$. In particular, we can characterise $\mathbf{Sh}_{(\infty, r+1)}(\mathcal{C})$ as the full subcategory of $\mathbf{Fun}(\Delta^{\text{op}}, \mathbf{Sh}_{(\infty, r)}(\mathcal{C}))$ spanned by those $F : \Delta^{\text{op}} \rightarrow \mathbf{Sh}_{(\infty, r)}(\mathcal{C})$ such that

- F_0 is essentially constant,
- F satisfies the Segal condition, and
- F is Rezk-complete.

Note that if n is finite, then F_0 is a sheaf of $(n + 1)$ -groupoids, and therefore F cannot be identified with a functor $\Delta^{\text{op}} \rightarrow \mathbf{Sh}_{(n,r)}(\mathcal{C})$.

Proof of Proposition 2.2.6. A functor $F : \Delta^{\text{op}} \times \Sigma_\infty^{\text{op}} \rightarrow \mathcal{X}$ factors through the left adjoint $\Delta \times \Sigma_\infty \rightarrow \Sigma_\infty$ of Definition 2.2.5 if and only if F satisfies (S1). Assuming F satisfies (S1), we therefore tacitly identify F with its underlying functor $F : \Sigma_\infty^{\text{op}} \rightarrow \mathcal{X}$.

By definition, (S2) is equivalent to asserting that F is local with respect to $\Sigma(W_{(n,r)})$; that is, F is local with respect to:

- (W1') $\Sigma^{m+1} \text{Sp}_U[\vec{k}] \hookrightarrow \Sigma^{m+1} \Delta_U[\vec{k}]$ for $U \in \mathcal{C}$, $m \geq 0$, and $[\vec{k}] \in \Sigma_\infty$,
- (W2') $\Sigma^{m+1} |\mathbb{I}|_U^\# \rightarrow \Sigma^{m+1} \Delta_U[0]$ for $U \in \mathcal{C}$, and $m \geq 0$,

(W3') $\Sigma^{m+1}(\mathbb{S}^{k-m} \times \Delta_U[0]) \rightarrow \Sigma^{m+1}\Delta_U[0]$ for $U \in \mathcal{C}$, and finite $m \geq r$ and $k > n$, and

(W4') $\Sigma^{m+1}\Delta_U[1] \rightarrow \Sigma^{m+1}\Delta_U[0]$ for $U \in \mathcal{C}$, and finite $m \geq r$.

all of which are a subset of the generators for $W_{(n+1,r+1)}$. In particular, F is the image of a sheaf of $(n+1, r+1)$ -categories if and only if F satisfies (S1) and (S2), and is moreover local with respect to

- $\mathrm{Sp}_U[\vec{k}] \hookrightarrow \Delta_U[\vec{k}]$ for $[\vec{k}] \in \Sigma_r$ and $U \in \mathcal{C}$, and
- $|\mathbb{I}_U^\# \rightarrow \Delta_U[0]$ for $U \in \mathcal{C}$.

Locality with respect to $|\mathbb{I}_U^\# \rightarrow \Delta_U[0]$ is precisely condition (S4). It remains to show that (S3) is equivalent to locality with respect to the remaining spine inclusions.

The Segal condition (S3) is equivalent to locality with respect to $\mathrm{Sp}[k, X] \hookrightarrow \Delta[k, X]$ for every $k \geq 2$ and all polysimplicial sheaves X . By Remark 2.1.6, it suffices to assert locality with respect to $\mathrm{Sp}_U[\vec{k}] \hookrightarrow \Delta_U[\vec{k}]$ for $U \in \mathcal{C}$ and all $[\vec{k}] \in \Sigma_\infty$. Assuming (S2), locality with respect to $\Sigma^{m+1}\Delta_U[1] \rightarrow \Sigma^{m+1}\Delta_U[0]$ for all finite $m \geq r$ implies that the Segal condition is equivalent to locality with respect to $\mathrm{Sp}_U[\vec{k}] \hookrightarrow \Delta_U[\vec{k}]$ for $U \in \mathcal{C}$ and $[\vec{k}] \in \Sigma_r$, as desired. \square

Lemma 2.2.8. *For $0 \leq r < \infty$, the category $\mathbf{Sh}_{(\infty,r)}(\mathcal{C})$ is precisely the full subcategory of $\mathbf{Sh}_{(\infty,\infty)}(\mathcal{C})$ spanned by those $X : \Sigma_\infty^{\mathrm{op}} \rightarrow \mathcal{X}$ that factor through $[-]_{<r} : \Sigma_\infty^{\mathrm{op}} \rightarrow \Sigma_r^{\mathrm{op}}$.*

Proof. We prove the case where $r = 0$; the rest follow by induction with Proposition 2.2.6. When $r = 0$, we are showing that $X : \Sigma_\infty^{\mathrm{op}} \rightarrow \mathcal{X}$ is a sheaf of $(\infty, 0)$ -categories precisely if it is essentially constant.

Note that $W_{(\infty,0)}$ is generated by $\Sigma^m \mathrm{Sp}_U[\vec{k}] \hookrightarrow \Sigma^m \Delta_U[\vec{k}]$ and $\Sigma^m \Delta_U[1] \rightarrow \Sigma^m \Delta_U[0]$ for $U \in \mathcal{C}$, $m \geq 0$, and $[\vec{k}] \in \Sigma_\infty$. By induction with Proposition 2.1.10, $W_{(\infty,0)}$ therefore contains the morphisms $\Delta_U[\vec{k}, 1] \rightarrow \Delta_U[\vec{k}]$ for all $[\vec{k}] \in \Sigma_\infty$, as well as the morphisms

$$\underbrace{\Delta_U[\vec{k}, 1] \sqcup_{\Delta_U[\vec{k}]} \cdots \sqcup_{\Delta_U[\vec{k}]} \Delta_U[\vec{k}, 1]}_{\ell} \simeq \Delta[\vec{k}, \mathrm{Sp}_U[\ell]] \rightarrow \Delta_U[\vec{k}, \ell]$$

for all $[\vec{k}] \in \Sigma_\infty$ and $\ell \geq 0$. This implies that $W_{(\infty,0)}$ contains all maps of the form $\Delta_U[\vec{k}, \ell] \rightarrow \Delta_U[\vec{k}]$, implying that X is $W_{(\infty,0)}$ -local if and only if X is essentially constant, as desired. \square

Proposition 2.2.9. *For $-2 \leq n \leq \infty$ and $0 \leq r < n+2$, the fully faithful inclusion $\mathbf{Sh}_{(n,r)}(\mathcal{C}) \subseteq \mathbf{Sh}_{(n,r+1)}(\mathcal{C})$ admits a right adjoint $\kappa_{\leq r}$.*

Proof. First suppose $n = \infty$. The right adjoint to the inclusion $\mathbf{Fun}(\Sigma_r^{\mathrm{op}}, \mathcal{X}) \rightarrow \mathbf{Fun}(\Sigma_{r+1}^{\mathrm{op}}, \mathcal{X})$ is given by restriction to $\Sigma_r \subseteq \Sigma_{r+1}$ (see Remark 2.1.7). By Lemma 2.2.8, restriction descends to a right adjoint $\kappa_{\leq r} : \mathbf{Sh}_{(\infty,r+1)}(\mathcal{C}) \rightarrow \mathbf{Sh}_{(\infty,r)}(\mathcal{C})$. The general case follows from observing that the above right adjoint descends further to a right adjoint $\mathbf{Sh}_{(n,r+1)}(\mathcal{C}) \rightarrow \mathbf{Sh}_{(n,r)}(\mathcal{C})$ for $n < \infty$. \square

Remark 2.2.10. On the other hand, it follows by definition that every inclusion $\mathbf{Sh}_{(n,r)}(\mathcal{C}) \subseteq \mathbf{Sh}_{(n',r')}(\mathcal{C})$ with $n \leq n'$ and $r \leq r'$ admits a left adjoint $\pi_{\leq r}$.

2.3 Sheaves of (∞, ∞) -categories and of ω -categories

Proposition 2.3.1. *The category $\mathbf{Sh}_{(\infty,\infty)}(\mathcal{C})$ is equivalent to the limit*

$$\varprojlim \left(\cdots \rightarrow \mathbf{Sh}_{(\infty,2)}(\mathcal{C}) \xrightarrow{\kappa_{\leq 1}} \mathbf{Sh}_{(\infty,1)}(\mathcal{C}) \xrightarrow{\kappa_{\leq 0}} \mathbf{Sh}_{(\infty,0)}(\mathcal{C}) \right)$$

in $\widehat{\mathbf{Cat}}_\infty$.

Proof. Since $\Sigma_\infty \simeq \varinjlim_r \Sigma_r$, we have an equivalence $\mathbf{Fun}(\Sigma_\infty^{\mathrm{op}}, \mathcal{X}) \simeq \varprojlim_r \mathbf{Fun}(\Sigma_r^{\mathrm{op}}, \mathcal{X})$. This allows us to identify the limit $\varprojlim_r \mathbf{Sh}_{(\infty,r)}(\mathcal{C})$ of right adjoints with the full subcategory of $\mathbf{Fun}(\Sigma_\infty^{\mathrm{op}}, \mathcal{X})$ on those $X : \Sigma_\infty^{\mathrm{op}} \rightarrow \mathcal{X}$ that restrict on Σ_r to a sheaf of (∞, r) -categories for every $0 \leq r < \infty$, but these are precisely the sheaves of (∞, ∞) -categories. \square

One may also be interested in the sheaves obtained as as the analogous limit of left adjoints:

Definition 2.3.2. As in Remark 2.2.10, the inclusions $\mathbf{Sh}_{(\infty,r)}(\mathcal{C}) \hookrightarrow \mathbf{Sh}_{(\infty,r+1)}(\mathcal{C})$ admit left adjoints $\pi_{\leq r}$. Define the category of *sheaves of ω -categories* over \mathcal{C} to be the analogous limit

$$\mathbf{Sh}_\omega(\mathcal{C}) := \varprojlim \left(\cdots \rightarrow \mathbf{Sh}_{(\infty,2)}(\mathcal{C}) \xrightarrow{\pi_{\leq 1}} \mathbf{Sh}_{(\infty,1)}(\mathcal{C}) \xrightarrow{\pi_{\leq 0}} \mathbf{Sh}_{(\infty,0)}(\mathcal{C}) \right)$$

in $\widehat{\mathbf{Cat}}_\infty$.

Remark 2.3.3. As $\mathbf{Sh}_\omega(\mathcal{C})$ is given as a limit of left adjoints between locally presentable categories, it is locally presentable. Moreover, we have by [Lur09a, Proposition 5.5.3.13 and Corollary 5.5.3.4] that it can be computed equivalently as the colimit

$$\mathbf{Sh}_\omega(\mathcal{C}) \simeq \varinjlim^R (\mathbf{Sh}_{(\infty,0)}(\mathcal{C}) \subseteq \mathbf{Sh}_{(\infty,1)}(\mathcal{C}) \subseteq \mathbf{Sh}_{(\infty,2)}(\mathcal{C}) \subseteq \cdots)$$

in \mathbf{Pres}_∞^R .

To understand the relationship between $\mathbf{Sh}_\omega(\mathcal{C})$ and $\mathbf{Sh}_{(\infty,\infty)}$, we have the following more general result about colimits in \mathbf{Pres}_∞^R .

Lemma 2.3.4. *Fix a locally presentable category \mathcal{B} , and let $(\mathcal{A}_k \subseteq \mathcal{B})_{k \in K}$ be a filtered diagram of locally presentable full subcategories of \mathcal{B} such that each inclusion $\mathcal{A}_k \subseteq \mathcal{B}$ admits a left adjoint $\pi_k : \mathcal{B} \rightarrow \mathcal{A}_k$. Then, the following categories in $(\widehat{\mathbf{Cat}}_\infty)_{/\mathcal{B}}$ are equivalent:*

- (i) *The colimit $\varinjlim_k^R \mathcal{A}_k$, computed in \mathbf{Pres}_∞^R , with its canonical map $\varinjlim_k^R \mathcal{A}_k \rightarrow \mathcal{B}$,*
- (ii) *The closure $\overline{\mathcal{A}}$ of $\mathcal{A} := \bigcup_k \mathcal{A}_k$ in \mathcal{B} under limits, with the inclusion $\overline{\mathcal{A}} \subseteq \mathcal{B}$.*
- (iii) *The full subcategory $\pi_\omega \mathcal{B}$ of \mathcal{B} spanned by all $x \in \mathcal{B}$ such that the canonical map $x \rightarrow \pi_\omega x := \varprojlim_k \pi_k x$ is an equivalence.*

In particular, the canonical map $\varinjlim_k^R \mathcal{A}_k \rightarrow \mathcal{B}$ is fully faithful, and its left adjoint is given by sending $x \in \mathcal{B}$ to $\pi_\omega x = \varprojlim_k \pi_k x$.

Proof. We first establish the equivalence between (ii) and (iii). By definition, there is certainly a fully faithful inclusion $\pi_\omega \mathcal{B} \subseteq \overline{\mathcal{A}}$. Suppose $x \in \overline{\mathcal{A}}$; that is, let $x \in \mathcal{B}$ be some limit of objects in $\mathcal{A} = \bigcup_k \mathcal{A}_k$. Then, it is also a canonical limit of objects in \mathcal{A} ; that is, the canonical map

$$x \rightarrow \varprojlim (\mathcal{A}_{x/} \rightarrow \mathcal{B})$$

is an equivalence in \mathcal{B} . The above map factors through $x \rightarrow \pi_\omega x$ via the map $\varprojlim_k \pi_k x \rightarrow \varprojlim (\mathcal{A}_{x/} \rightarrow \mathcal{B})$ induced by the functor $\pi_\bullet x : K^{\text{op}} \rightarrow \mathcal{A}_{x/}$ that sends $k \mapsto \pi_k x$. It therefore suffices to show that this functor $\pi_\bullet x : K^{\text{op}} \rightarrow \mathcal{A}_{x/}$ is co-initial (that is, that $(\pi_\bullet x)^{\text{op}} : K \rightarrow (\mathcal{A}_{x/})^{\text{op}}$ is cofinal in the sense of [Lur09a, §4.1]), as this would prove that the induced map $\varprojlim_k \pi_k x \rightarrow \varprojlim (\mathcal{A}_{x/} \rightarrow \mathcal{B})$ is an equivalence, and thus show that $x \in \pi_\omega \mathcal{B}$ by 2-out-of-3.

By Quillen's Theorem A for ∞ -categories [Lur09a, Theorem 4.1.3.1], $\pi_\bullet x$ is co-initial if and only if for every $(x \rightarrow y) \in \mathcal{A}_{x/}$, the category $K^{\text{op}} \times_{\mathcal{A}_{x/}} \mathcal{A}_{x//y}$ is weakly contractible. Note that $K^{\text{op}} \times_{\mathcal{A}_{x/}} \mathcal{A}_{x//y}$ is the category of factorisations of $x \rightarrow y$ through some $x \rightarrow \pi_\ell x$.

Fix $(x \rightarrow y) \in \mathcal{A}_{x/}$. Then, $y \in \mathcal{A}_k$ for some $k \in K$. Let $\chi : J \rightarrow K^{\text{op}} \times_{\mathcal{A}_{x/}} \mathcal{A}_{x//y}$ be an arbitrary diagram, then each $j \in J$ is sent through χ to some factorisation $x \rightarrow \pi_{\ell_j} x \rightarrow y$ of $x \rightarrow y$. Since K^{op} is cofiltered, we can extend the diagram $J \rightarrow K^{\text{op}}$ to a cone $J^\triangleleft \rightarrow K^{\text{op}}$, where the cone point maps to a vertex ℓ with an arrow $\ell \rightarrow \ell_j$ in K^{op} for each $j \in J$. We can choose the cone so that we also have an arrow $\ell \rightarrow k$ in K^{op} .

The arrows $\ell \rightarrow \ell_j$ ensure that $\pi_{\ell_j} x \in \mathcal{A}_\ell$ for each $j \in K$. Our choice of ℓ ensures also that $y \in \mathcal{A}_\ell$. Therefore, χ corestricts to a map $J \rightarrow (K^{\text{op}})_{\ell/} \times_{(\mathcal{A}_\ell)_{x/}} (\mathcal{A}_\ell)_{x//y}$. Since $x \rightarrow \pi_\ell x$ is initial in $(\mathcal{A}_\ell)_{x/}$ from being

left adjoint to the inclusion $\mathcal{A}_\ell \subseteq \mathcal{B}$, it follows by [Lur09a, Proposition 1.2.13.8] that we have a lift

$$\left\{ \begin{array}{ccc} & (\mathcal{A}_\ell)_{x//y} & \\ \nearrow J^\triangleleft & \downarrow & \\ & (\mathcal{A}_\ell)_{x/} & \end{array} \right\} \iff \left\{ \begin{array}{ccc} & ((\mathcal{A}_\ell)_{x//y})_{\pi_\ell x/} & \\ \nearrow J & \downarrow & \\ & ((\mathcal{A}_\ell)_{x/})_{\pi_\ell x/} & \end{array} \right\}$$

which induces a lift

$$\begin{array}{ccccc} J & \longrightarrow & (K^{\text{op}})_{\ell/} \times_{(\mathcal{A}_\ell)_{x/}} (\mathcal{A}_\ell)_{x//y} & \longrightarrow & (\mathcal{A}_\ell)_{x//y} \\ \downarrow & \nearrow & \downarrow & \lrcorner & \downarrow \\ J^\triangleleft & \longrightarrow & (K^{\text{op}})_{\ell/} & \longrightarrow & (\mathcal{A}_\ell)_{x/} \end{array}$$

This shows that $K^{\text{op}} \times_{\mathcal{A}_{x/}} \mathcal{A}_{x//y}$ satisfies the right lifting property against all inclusions $J \hookrightarrow J^\triangleleft$, which implies that $K^{\text{op}} \times_{\mathcal{A}_{x/}} \mathcal{A}_{x//y}$ is weakly contractible, as desired. This establishes the equivalence between (ii) and (iii).

Choose a regular cardinal $\lambda \gg |K|$ such that each $\mathcal{A}_k \subseteq \mathcal{B}$ is closed under λ -filtered colimits, and such that λ -filtered colimits commute with λ -small limits in \mathcal{B} . It then follows that $\pi_\omega \mathcal{B}$ is closed under λ -filtered colimits, since λ -filtered colimits commute with the $|K|$ -small limit construction of $\pi_\omega x = \varprojlim_k \pi_k x$. As $\pi_\omega \mathcal{B} \simeq \overline{\mathcal{A}}$ is closed under λ -filtered colimits and arbitrary limits, it then follows from the reflection theorem [RS22, Theorem 6.2] that $\pi_\omega \mathcal{B}$ is locally presentable.

Moreover, $\pi_\omega : \mathcal{B} \rightarrow \pi_\omega \mathcal{B}$ defines a left adjoint to the inclusion. Indeed, the adjoint functor theorem ensures that the inclusion $\pi_\omega \mathcal{B} \subseteq \mathcal{B}$ admits a left adjoint L . Since the inclusion $\mathcal{A}_k \subseteq \mathcal{B}$ factors through $\overline{\mathcal{A}} \simeq \pi_\omega \mathcal{B}$, it follows from taking left adjoints to these inclusions that we have a natural equivalence $\pi_k L \simeq \pi_k$ for every $k \in K$. Taking the limit over k , this implies $\pi_\omega L \simeq \pi_\omega$. On the other hand, $Lx \in \pi_\omega \mathcal{B}$ by definition, meaning that $Lx \rightarrow \pi_\omega Lx \simeq \pi_\omega x$ is an equivalence for every $x \in \mathcal{B}$, showing that $L \simeq \pi_\omega$.

It remains to show that the inclusions $\mathcal{A}_k \subseteq \overline{\mathcal{A}}$ exhibit $\overline{\mathcal{A}}$ as the colimit $\varinjlim_k^R \mathcal{A}_k$ in \mathbf{Pres}_∞^R . For an arbitrary locally presentable category \mathcal{E} , consider the fibre product

$$\begin{array}{ccc} \mathbf{Fun}^{(R)}(\mathcal{A}, \mathcal{E}) & \xrightarrow{\sim} & \varprojlim_k \mathbf{Fun}^R(\mathcal{A}_k, \mathcal{E}) \\ \downarrow & \lrcorner & \downarrow \\ \mathbf{Fun}(\mathcal{A}, \mathcal{E}) & \xrightarrow{\sim} & \varprojlim_k \mathbf{Fun}(\mathcal{A}_k, \mathcal{E}) \end{array}$$

where the limits are computed in $\widehat{\mathbf{Cat}}_\infty$; that is, $\mathbf{Fun}^{(R)}(\mathcal{A}, \mathcal{E})$ is the full subcategory of $\mathbf{Fun}(\mathcal{A}, \mathcal{E})$ spanned by those functors $F : \mathcal{A} \rightarrow \mathcal{E}$ whose restrictions $F| : \mathcal{A}_k \rightarrow \mathcal{E}$ are right adjoints for every $k \in K$. Since \mathcal{E} has all limits, we have from [Lur09a, Lemma 4.3.2.13 and Corollary 4.3.2.16] a fully faithful right Kan extension functor $\mathbf{Fun}^{(R)}(\mathcal{A}, \mathcal{E}) \hookrightarrow \mathbf{Fun}(\overline{\mathcal{A}}, \mathcal{E})$ right adjoint to restriction along $\mathcal{A} \subseteq \overline{\mathcal{A}}$.

By definition, $\overline{\mathcal{A}}$ is the limit closure of \mathcal{A} in \mathcal{B} , so the right Kan extension \overline{F} of any functor $F \in \mathbf{Fun}^{(R)}(\mathcal{A}, \mathcal{E})$ preserves arbitrary limits. Moreover, if we choose a regular cardinal $\lambda \gg |K|$ such that $F| : \mathcal{A}_k \rightarrow \mathcal{E}$ preserves λ -filtered colimits for each $k \in K$, and λ -filtered colimits commute with λ -small limits in \mathcal{E} , then the corresponding right Kan extension $\overline{F} : \overline{\mathcal{A}} \rightarrow \mathcal{E}$ preserves λ -filtered colimits as well, since $\overline{\mathcal{A}} \simeq \pi_\omega \mathcal{B}$ ensures that every object of $\overline{\mathcal{A}}$ is a $|K|$ -small limit of objects in \mathcal{A} . In summary, the right Kan extension of any functor $F \in \mathbf{Fun}^{(R)}(\mathcal{A}, \mathcal{E})$ is necessarily a right adjoint.

Conversely, the limit-preservation of right adjoints ensures that every right adjoint $\overline{\mathcal{A}} \rightarrow \mathcal{E}$ arises in this way. Therefore, taking right Kan extension induces an equivalence of categories

$$\mathbf{Fun}^R(\varinjlim_k^R \mathcal{A}_k, \mathcal{E}) \simeq \varprojlim_k \mathbf{Fun}^R(\mathcal{A}_k, \mathcal{E}) \simeq \mathbf{Fun}^{(R)}(\mathcal{A}, \mathcal{E}) \simeq \mathbf{Fun}^R(\overline{\mathcal{A}}, \mathcal{E})$$

Since $\mathcal{E} \in \mathbf{Pres}_\infty^R$ is arbitrary, it follows that $\overline{\mathcal{A}} \simeq \varinjlim_k^R \mathcal{A}_k$, as desired. \square

Corollary 2.3.5. Let $\pi_{\leq r} : \mathbf{Sh}_{(\infty, \infty)}(\mathcal{C}) \rightarrow \mathbf{Sh}_{(\infty, r)}(\mathcal{C})$ denote the left adjoints to inclusion for $0 \leq r < \infty$. For $X_{\bullet} \in \mathbf{Sh}_{(\infty, \infty)}(\mathcal{C})$, define

$$\pi_{\leq \omega} X := \varprojlim_r \pi_{\leq r} X$$

Then, $\mathbf{Sh}_{\omega}(\mathcal{C})$ is equivalent to the full subcategory of $\mathbf{Sh}_{(\infty, \infty)}(\mathcal{C})$ spanned by those sheaves X_{\bullet} for which $X \rightarrow \pi_{\leq \omega} X$ is an equivalence. Moreover, $\pi_{\leq \omega}$ defines a left adjoint to the inclusion $\mathbf{Sh}_{\omega}(\mathcal{C}) \subseteq \mathbf{Sh}_{(\infty, \infty)}(\mathcal{C})$.

3 Distributors and complete Segal spaces

In this section, we relate the categories $\mathbf{Sh}_{(n, r)}(\mathcal{C})$ and $\mathbf{Sh}_{\omega}(\mathcal{C})$ constructed in Section 2 to the general theory of complete Segal space objects in a distributor.

3.1 Preliminaries

We recall the necessary theory from [Lur09b] below.

Definition 3.1.1. [Lur09b, Definition 1.2.1] Fix a category \mathcal{Y} and a full subcategory \mathcal{X} . Say that \mathcal{Y} is an \mathcal{X} -distributor if the following hold:

- (D1) \mathcal{X} and \mathcal{Y} are locally presentable,
- (D2) The inclusion $\mathcal{X} \subseteq \mathcal{Y}$ admits both a left adjoint $\pi_{\mathcal{X}}$ called \mathcal{X} -truncation, and a right adjoint $\kappa_{\mathcal{X}}$ called \mathcal{X} -core,
- (D3) For all $y \rightarrow x$ in \mathcal{Y} with $x \in \mathcal{X}$, the pullback functor $\mathcal{X}/_x \rightarrow \mathcal{Y}/_y$ preserves colimits, and
- (D4) The functor $\chi : \mathcal{X} \rightarrow (\widehat{\mathbf{Cat}}_{\infty})^{\text{op}}, x \mapsto \mathcal{Y}/_x$, which sends morphisms in \mathcal{X} to pullback functors, preserves limits.

Remark 3.1.2. [Lur09b, Example 1.2.3 and Remark 1.2.6] If \mathcal{Y} is an \mathcal{X} -distributor, then \mathcal{X} is necessarily a topos. Conversely, every topos \mathcal{X} is a distributor relative to itself.

Definition 3.1.3. For \mathcal{X} a finitely complete category, let $\mathbf{Cat}(\mathcal{X})$ denote the full subcategory of $\mathbf{Fun}(\Delta^{\text{op}}, \mathcal{X})$ spanned by those $X : \Delta^{\text{op}} \rightarrow \mathcal{X}$ satisfying the *Segal condition*; that is, the canonical maps

$$X_n \rightarrow X_1 \times_{X_0} \cdots \times_{X_0} X_1$$

are equivalences for every $n \geq 0$.

Further, define $\mathbf{Grpd}(\mathcal{X})$ to be the full subcategory of $\mathbf{Cat}(\mathcal{X})$ spanned by those X_{\bullet} such that whenever $[n] = S \cup S'$ with $S \cap S' = \{i\}$, the induced map

$$X_n \rightarrow X_S \times_{X_{\{i\}}} X_{S'}$$

is an equivalence.

Remark 3.1.4. If \mathcal{X} is locally presentable, then so are $\mathbf{Cat}(\mathcal{X})$ and $\mathbf{Grpd}(\mathcal{X})$. Indeed, $\mathbf{Cat}(\mathcal{S})$ is the strongly reflective localisation of $\mathcal{P}(\Delta)$ at the maps $\Delta[1] \sqcup_{\Delta[0]} \cdots \sqcup_{\Delta[0]} \Delta[1] \hookrightarrow \Delta[n]$ for $n \geq 0$, and $\mathbf{Grpd}(\mathcal{S})$ is similarly a strongly reflective localisation of $\mathbf{Cat}(\mathcal{S})$. As every locally presentable category is a strongly reflective localisation of a presheaf category, it follows likewise that $\mathbf{Cat}(\mathcal{X})$ is a strongly reflective localisation of $\mathbf{Fun}(\Delta^{\text{op}}, \mathcal{X})$, and $\mathbf{Grpd}(\mathcal{X})$ is a strongly reflective localisation of $\mathbf{Cat}(\mathcal{X})$.

Definition 3.1.5. Fix a topos \mathcal{X} and an \mathcal{X} -distributor \mathcal{Y} . Then, define $\mathbf{Cat}_{\mathcal{X}}(\mathcal{Y})$ to be the full subcategory of $\mathbf{Cat}(\mathcal{Y})$ spanned by those $Y \in \mathbf{Cat}(\mathcal{Y})$ such that $Y_0 \in \mathcal{X}$.

Remark 3.1.6. As $\mathbf{Cat}_{\mathcal{X}}(\mathcal{Y}) \simeq \mathbf{Cat}(\mathcal{Y}) \times_{\mathcal{X}} \mathcal{Y}$ is a fibre product of right adjoints between locally presentable categories, $\mathbf{Cat}_{\mathcal{X}}(\mathcal{Y})$ is also locally presentable, and the inclusion $\mathbf{Cat}_{\mathcal{X}}(\mathcal{Y}) \subseteq \mathbf{Cat}(\mathcal{Y})$ is a right adjoint.

Remark 3.1.7. The category $\mathbf{Cat}_X(\mathcal{Y})$ is denoted by $\mathbf{SS}_X(\mathcal{Y})$ in [Lur09b, Definition 1.2.7], and is referred to as the category of *Segal space objects* in \mathcal{Y} .

Lemma 3.1.8. [Lur09b, Proposition 1.1.14] *If \mathcal{X} is finitely complete, then the inclusion $\mathbf{Grpd}(\mathcal{X}) \subseteq \mathbf{Cat}(\mathcal{X})$ admits a right adjoint $X_\bullet \mapsto X_\bullet^\sim$.*

Proof sketch. If $\mathcal{X} \simeq \mathcal{S}$, then we construct the right adjoint explicitly. For $X_\bullet \in \mathbf{Cat}(\mathcal{S})$, define the $\mathrm{Ho}(\mathcal{S})$ -enriched category hX where its objects are the underlying points of X_0 , and the mapping space $\mathrm{Hom}_{hX}(x, y)$ is the homotopy class of the fibre product $\{x\} \times_{X_0} X_1 \times_{X_0} \{y\}$ in $\mathrm{Ho}(\mathcal{S})$. Then, we define X_n^\sim to be the full subspace of X_n spanned by those cells $\sigma \in X_n$ for which each $d_i\sigma$ descends to an isomorphism in hX .

If $\mathcal{X} \simeq \mathcal{P}(\mathcal{C})$, then $\mathbf{Cat}(\mathcal{X}) \simeq \mathbf{Fun}(\mathcal{C}^{\mathrm{op}}, \mathbf{Cat}(\mathcal{S}))$ and $\mathbf{Grpd}(\mathcal{X}) \simeq \mathbf{Fun}(\mathcal{C}^{\mathrm{op}}, \mathbf{Grpd}(\mathcal{S}))$, so the right adjoint in this case is given pointwise by the right adjoint for \mathcal{S} .

For a general \mathcal{X} , let $X_\bullet \in \mathbf{Cat}(\mathcal{X})$, and consider the object $jX_\bullet \in \mathbf{Cat}(\mathcal{P}(\mathcal{X}))$ obtained by pointwise post-composition with the Yoneda embedding $j : \mathcal{X} \hookrightarrow \mathcal{P}(\mathcal{X})$. Then, the reflection $(jX_\bullet)^\sim \in \mathbf{Grpd}(\mathcal{P}(\mathcal{X}))$ turns out to be pointwise representable, allowing it to descend to an object $X_\bullet^\sim \in \mathbf{Grpd}(\mathcal{X})$. \square

Corollary 3.1.9. [Lur09b, Notation 1.2.9] *For an \mathcal{X} -distributor \mathcal{Y} , the inclusion $\mathbf{Grpd}(\mathcal{X}) \subseteq \mathbf{Cat}_X(\mathcal{Y})$ admits a right adjoint $\mathrm{Gp} : \mathbf{Cat}_X(\mathcal{Y}) \rightarrow \mathbf{Grpd}(\mathcal{X})$.*

Remark 3.1.10. Explicitly, $\mathrm{Gp}_\bullet Y \simeq (\kappa_{X,*} Y_\bullet)^\sim$, where $\kappa_{X,*}$ applies the core pointwise. By the proof sketch of Lemma 3.1.8, the canonical map $\mathrm{Gp}_0 Y \rightarrow Y_0$ is an equivalence for all $Y_\bullet \in \mathbf{Cat}_X(\mathcal{Y})$.

Definition 3.1.11. Fix an \mathcal{X} -distributor \mathcal{Y} . Say that a morphism $f : Y_\bullet \rightarrow Y'_\bullet$ in $\mathbf{Cat}_X(\mathcal{Y})$ is a *Segal equivalence* if the following conditions hold:

(E1) f is *fully faithful*, in that

$$\begin{array}{ccc} Y_1 & \longrightarrow & Y'_1 \\ \downarrow & & \downarrow \\ Y_0 \times Y_0 & \longrightarrow & Y'_0 \times Y'_0 \end{array}$$

is a pullback square, and

(E2) f is *essentially surjective* in that the induced map $|\mathrm{Gp}_\bullet Y| \rightarrow |\mathrm{Gp}_\bullet Y'|$ is an equivalence in \mathcal{X} .

Say that $Y_\bullet \in \mathbf{Cat}_X(\mathcal{Y})$ is a *complete Segal space* if it is local with respect to the Segal equivalences. Denote by $\mathbf{CSS}_X(\mathcal{Y})$ the full subcategory of $\mathbf{Cat}_X(\mathcal{Y})$ spanned by the complete Segal spaces.

Remark 3.1.12. By [Lur09b, Theorem 1.2.13], $Y_\bullet \in \mathbf{Cat}_X(\mathcal{Y})$ is a complete Segal space if and only if $\mathrm{Gp}_\bullet Y$ is essentially constant; that is, $\mathrm{Gp}_\bullet Y$ lies in the essential image of the diagonal $\mathcal{X} \subseteq \mathbf{Grpd}(\mathcal{X})$.

Proposition 3.1.13. [Lur09b, Proposition 1.3.2] *If \mathcal{Y} is an \mathcal{X} -distributor, then the fully faithful diagonal functor $\mathcal{X} \subseteq \mathbf{CSS}_X(\mathcal{Y})$ exhibits $\mathbf{CSS}_X(\mathcal{Y})$ as an \mathcal{X} -distributor as well.*

Remark 3.1.14. The truncation functor $\pi_X : \mathbf{CSS}_X(\mathcal{Y}) \rightarrow \mathcal{X}$, being left adjoint to the diagonal, is given by geometric realisation $\pi_X Y \simeq |Y_\bullet|$. On the other hand, the core functor $\kappa_X : \mathbf{CSS}_X(\mathcal{Y}) \rightarrow \mathcal{X}$ is given by $\kappa_X Y \simeq Y_0$.

3.2 The ∞ -categories of distributors

Definition 3.2.1. Fix a topos \mathcal{X} . Then, define \mathbf{Dist}_X^L to be the subcategory of $(\widehat{\mathbf{Cat}_\infty})_{\mathcal{X}}$ where the objects are \mathcal{X} -distributors $\mathcal{X} \subseteq \mathcal{Y}$, and a functor $\psi : \mathcal{Z} \rightarrow \mathcal{Y}$ under \mathcal{X} is a morphism of \mathbf{Dist}_X^L if and only if

(L1) ψ preserves colimits, and

(L2) ψ preserves truncation, in that $\pi_X \psi \simeq \pi_X$.

On the other hand, define \mathbf{Dist}_X^R to be the subcategory of $(\widehat{\mathbf{Cat}_\infty})_{\mathcal{X}}$ where the objects are again the \mathcal{X} -distributors $\mathcal{X} \subseteq \mathcal{Y}$, and a functor $\phi : \mathcal{Y} \rightarrow \mathcal{Z}$ under \mathcal{X} is a morphism of \mathbf{Dist}_X^R if and only if

(R1) ϕ preserves limits and λ -filtered colimits for some $\lambda \gg 0$, and

(R2) ϕ preserves cores, in that $\kappa_{\mathcal{X}}\phi \simeq \kappa_{\mathcal{X}}$.

Remark 3.2.2. Since \mathcal{X} -distributors are necessarily locally presentable, (L1) is equivalent by [Lur09a, Corollary 5.5.2.9] to asserting that ψ admits a right adjoint ϕ . As truncation is left adjoint to the inclusion of \mathcal{X} , (L2) is then equivalent to asserting that the right adjoint ϕ lies under \mathcal{X} ; that is, $\phi|_{\mathcal{X}} \simeq \text{id}_{\mathcal{X}}$.

Dually, (R1) is likewise equivalent to asserting that ϕ admits a left adjoint ψ . As core is right adjoint to the inclusion of \mathcal{X} , (R2) is equivalent to asserting that the left adjoint ψ lies under \mathcal{X} ; that is $\psi|_{\mathcal{X}} \simeq \text{id}_{\mathcal{X}}$. This demonstrates that the two categories of Definition 3.2.1 are formally dual: $\mathbf{Dist}_{\mathcal{X}}^L \simeq (\mathbf{Dist}_{\mathcal{X}}^R)^{\text{op}}$.

Proposition 3.2.3. *For any topos \mathcal{X} , the functor $\mathbf{Dist}_{\mathcal{X}}^R \rightarrow (\widehat{\mathbf{Cat}}_{\infty})_{/\mathcal{X}}$ sending an \mathcal{X} -distributor \mathcal{Y} to its core $\kappa_{\mathcal{X}} : \mathcal{Y} \rightarrow \mathcal{X}$ creates all small limits.*

Proof. Let $\mathcal{Y}_{\bullet} : K \rightarrow \mathbf{Dist}_{\mathcal{X}}^R$ be a small diagram indexed by a simplicial set K . The limit of the diagram $K \rightarrow (\widehat{\mathbf{Cat}}_{\infty})_{/\mathcal{X}}$ can be computed as the limit of the corresponding diagram $K^{\triangleright} \rightarrow \widehat{\mathbf{Cat}}_{\infty}$ that sends the cocone point of K^{\triangleright} to \mathcal{X} via core maps $\kappa_{\mathcal{X}} : \mathcal{Y}_k \rightarrow \mathcal{X}$. By [Lur09a, Theorem 5.5.3.18], we can compute this limit instead as a limit of the diagram $\tilde{\mathcal{Y}}_{\bullet} : K^{\triangleright} \rightarrow \mathbf{Pres}_{\infty}^R$.

Let $\mathcal{Y} := \varprojlim_k \tilde{\mathcal{Y}}_k$, then \mathcal{Y} is locally presentable, and the projection map $\kappa_{\mathcal{X}} : \mathcal{Y} \rightarrow \mathcal{X}$ is a right adjoint. The left adjoint coincides with the functor $\mathcal{X} \rightarrow \mathcal{Y} \simeq \varprojlim_k \tilde{\mathcal{Y}}_k$ in $\widehat{\mathbf{Cat}}_{\infty}$ induced by the inclusions $\mathcal{X} \subseteq \mathcal{Y}_k$ for $k \in K^{\triangleright}$. In particular, since these inclusions are fully faithful right adjoints, the same is true for the left adjoint $\mathcal{X} \rightarrow \mathcal{Y}$. This proves that \mathcal{Y} satisfies (D1) and (D2).

Let $y \rightarrow x$ in \mathcal{Y} with $x \in \mathcal{X}$. Note that $\mathcal{Y}_{/y} \simeq \varprojlim_k (\tilde{\mathcal{Y}}_k)_{/y_k}$, where y_k is the image of y under the projection $\mathcal{Y} \rightarrow \mathcal{Y}_k$, and the map $\mathcal{X}_{/x} \rightarrow \mathcal{Y}_{/y}$ is the canonical map induced by the functors $\mathcal{X}_{/x} \rightarrow (\tilde{\mathcal{Y}}_k)_{/y_k}$ for each $k \in K^{\triangleright}$. Since each \mathcal{Y}_k is an \mathcal{X} -distributor, the functor $\mathcal{X}_{/x} \rightarrow (\tilde{\mathcal{Y}}_k)_{/y_k}$ is a left adjoint for every $k \in K^{\triangleright}$. By [Lur09a, Proposition 5.5.3.13], $\mathcal{X}_{/x} \rightarrow \mathcal{Y}_{/y}$ must be a left adjoint as well, establishing (D3).

For $k \in K^{\triangleright}$, let $\chi^k : \mathcal{X} \rightarrow (\widehat{\mathbf{Cat}}_{\infty})^{\text{op}}$ denote the functor mapping $x \in \mathcal{X}$ to the category $(\tilde{\mathcal{Y}}_k)_{/x}$. Since each χ^k preserves limits from $\tilde{\mathcal{Y}}_k$ being an \mathcal{X} -distributor, the same is true for the induced functor $\bar{\chi} : \mathcal{X} \rightarrow \mathbf{Fun}(K^{\triangleright}, (\widehat{\mathbf{Cat}}_{\infty})^{\text{op}})$ that sends each $x \in \mathcal{X}$ to the functor mapping $k \mapsto (\tilde{\mathcal{Y}}_k)_{/x}$. Now, the functor $\chi : \mathcal{X} \rightarrow (\widehat{\mathbf{Cat}}_{\infty})^{\text{op}}$ sending $x \mapsto \mathcal{Y}_{/x} \simeq \varprojlim_k (\tilde{\mathcal{Y}}_k)_{/x}$ is precisely the composite of limit-preserving functors

$$\mathcal{X} \xrightarrow{\bar{\chi}} \mathbf{Fun}(K^{\triangleright}, (\widehat{\mathbf{Cat}}_{\infty})^{\text{op}}) \xrightarrow{\varprojlim} (\widehat{\mathbf{Cat}}_{\infty})^{\text{op}}$$

and thus preserves limits, establishing (D4).

Therefore, the limit \mathcal{Y} is an \mathcal{X} -distributor, and the canonical projections $\mathcal{Y} \rightarrow \mathcal{Y}_k$ are core-preserving right adjoints. Suppose we have a cone from an \mathcal{X} -distributor \mathcal{Z} to \mathcal{Y}_{\bullet} in $\mathbf{Dist}_{\mathcal{X}}^R$. This induces a cone from \mathcal{Z} to $\tilde{\mathcal{Y}}_{\bullet}$ in \mathbf{Pres}_{∞}^R , inducing an essentially unique right adjoint $\mathcal{Z} \rightarrow \mathcal{Y}$. Moreover, since the functors $\mathcal{Z} \rightarrow \mathcal{Y}_k$ lie under \mathcal{X} , the same is true for $\mathcal{Z} \rightarrow \mathcal{Y}$, ensuring that the canonical functor $\mathcal{Z} \rightarrow \mathcal{Y}$ is a morphism of $\mathbf{Dist}_{\mathcal{X}}^R$, proving that \mathcal{Y} is indeed a limit of \mathcal{Y}_{\bullet} in $\mathbf{Dist}_{\mathcal{X}}^R$. \square

Proposition 3.2.4. *For any topos \mathcal{X} , the functor $\mathbf{Dist}_{\mathcal{X}}^L \rightarrow (\widehat{\mathbf{Cat}}_{\infty})_{/\mathcal{X}}$ sending an \mathcal{X} -distributor \mathcal{Y} to its truncation $\pi_{\mathcal{X}} : \mathcal{Y} \rightarrow \mathcal{X}$ creates all small limits.*

Proof. Let $\mathcal{Y}_{\bullet} : K \rightarrow \mathbf{Dist}_{\mathcal{X}}^L$ be a small diagram indexed by a simplicial set K . The limit of the diagram $K \rightarrow (\widehat{\mathbf{Cat}}_{\infty})_{/\mathcal{X}}$ can be computed as a limit of the corresponding diagram $K^{\triangleright} \rightarrow \widehat{\mathbf{Cat}}_{\infty}$ that sends the cocone point of K^{\triangleright} to \mathcal{X} via truncation maps $\pi_{\mathcal{X}} : \mathcal{Y}_k \rightarrow \mathcal{X}$. By [Lur09a, Proposition 5.5.3.13], we can compute this limit instead as a limit of the diagram $\tilde{\mathcal{Y}}_{\bullet} : K^{\triangleright} \rightarrow \mathbf{Pres}_{\infty}^L$.

Let $\mathcal{Y} := \varprojlim_k \tilde{\mathcal{Y}}_k$, then \mathcal{Y} is locally presentable, and the projection map $\pi_{\mathcal{X}} : \mathcal{Y} \rightarrow \mathcal{X}$ is a left adjoint. The right adjoint coincides with the functor $\mathcal{X} \rightarrow \mathcal{Y} \simeq \varprojlim_k \tilde{\mathcal{Y}}_k$ in $\widehat{\mathbf{Cat}}_{\infty}$ induced by the inclusions $\mathcal{X} \subseteq \mathcal{Y}_k$ for $k \in K^{\triangleright}$. In particular, since these inclusions are fully faithful left adjoints, the same is true for the right adjoint $\mathcal{X} \rightarrow \mathcal{Y}$. This proves that \mathcal{Y} satisfies (D1) and (D2). Moreover, that \mathcal{Y} satisfies (D3) and (D4) follows the argument in Proposition 3.2.3 verbatim.

Therefore, the limit \mathcal{Y} is an \mathcal{X} -distributor, and the canonical projections $\mathcal{Y} \rightarrow \mathcal{Y}_k$ are truncation-preserving left adjoints. Suppose we have a cone from an \mathcal{X} -distributor \mathcal{Z} to \mathcal{Y}_\bullet in $\mathbf{Dist}_{\mathcal{X}}^L$. This induces a cone from \mathcal{Z} to $\tilde{\mathcal{Y}}_\bullet$ in \mathbf{Pres}_{∞}^L , inducing an essentially unique left adjoint $\mathcal{Z} \rightarrow \mathcal{Y}$. Moreover, since the functors $\mathcal{Z} \rightarrow \mathcal{Y}_k$ lie under \mathcal{X} , the same is true for $\mathcal{Z} \rightarrow \mathcal{Y}$, ensuring that the canonical functor $\mathcal{Z} \rightarrow \mathcal{Y}$ is a morphism of $\mathbf{Dist}_{\mathcal{X}}^L$, proving that \mathcal{Y} is indeed a limit of \mathcal{Y}_\bullet in $\mathbf{Dist}_{\mathcal{X}}^L$. \square

Remark 3.2.5. From the duality in Remark 3.2.2, $\mathbf{Dist}_{\mathcal{X}}^R$ has all small limits and colimits (and likewise for $\mathbf{Dist}_{\mathcal{X}}^L$). By [Gol23, Lemma 2.3.11], the forgetful functor $\mathbf{Dist}_{\mathcal{X}}^R \rightarrow \mathbf{Pres}_{\infty}^R$ creates all small limits indexed by a weakly contractible simplicial set; that is, weakly contractible limits of \mathcal{X} -distributors can be computed on their underlying categories. On the other hand, the topos \mathcal{X} viewed as a distributor over itself is simultaneously the initial and terminal object in $\mathbf{Dist}_{\mathcal{X}}^R$.

3.3 Functoriality of complete Segal spaces

Definition 3.3.1. Let \mathcal{Y} be a category with full subcategory $\mathcal{X} \subseteq \mathcal{Y}$. Say that a functor $\phi : \mathcal{Y} \rightarrow \mathcal{Z}$ *preserves fibre products over \mathcal{X}* if for every fibre product $y^1 \times_x y^2$ in \mathcal{Y} with $x \in \mathcal{X}$, the induced map $\phi(y^1 \times_x y^2) \rightarrow \phi(y^1) \times_{\phi(x)} \phi(y^2)$ is an equivalence in \mathcal{Z} .

If \mathcal{X} has finite limits, and $\mathcal{X} \subseteq \mathcal{Y}$ preserves these limits, then say that a functor $\phi : \mathcal{Y} \rightarrow \mathcal{Z}$ is *relatively left exact* (relative to \mathcal{X}) if $\phi|_{\mathcal{X}} : \mathcal{X} \rightarrow \mathcal{Z}$ is left exact, and ϕ preserves fibre products over \mathcal{X} .

The functorial nature of the construction of complete Segal spaces can be summarised as follows:

Lemma 3.3.2. *Let $g_* : \mathcal{X} \rightarrow \mathcal{X}'$ be a geometric morphism of topoi. Fix an \mathcal{X} -distributor \mathcal{Y} and an \mathcal{X}' -distributor \mathcal{Y}' , and let $\phi : \mathcal{Y} \rightarrow \mathcal{Y}'$ be a functor such that*

- ϕ extends g_* , in that $\phi|_{\mathcal{X}} \simeq g_*$,
- ϕ commutes with cores, in that $\phi \kappa_{\mathcal{X}} \simeq \kappa_{\mathcal{X}'} \phi$, and
- ϕ preserves fibre products over \mathcal{X} .

Then, $\phi_ : \mathbf{Fun}(\Delta^{\text{op}}, \mathcal{Y}) \rightarrow \mathbf{Fun}(\Delta^{\text{op}}, \mathcal{Y}')$ restricts to a functor $\phi_* : \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}) \rightarrow \mathbf{CSS}_{\mathcal{X}'}(\mathcal{Y}')$.*

Remark 3.3.3. In the situation of Lemma 3.3.2, $\phi_* : \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}) \rightarrow \mathbf{CSS}_{\mathcal{X}'}(\mathcal{Y}')$ acts pointwise on the underlying functors. Since limits of complete Segal spaces are computed pointwise, this ensures that ϕ_* preserves fibre products over \mathcal{X} . Likewise, Remark 3.1.14 ensures that ϕ_* commutes with cores as well. Therefore, Lemma 3.3.2 combined with Proposition 3.1.13 implies an *endofunctorial* nature of the construction of complete Segal spaces.

Proof of Lemma 3.3.2. Since ϕ is relatively left exact, it follows that ϕ_* restricts to a functor $\mathbf{Cat}_{\mathcal{X}}(\mathcal{Y}) \rightarrow \mathbf{Cat}_{\mathcal{X}'}(\mathcal{Y}')$. Similarly, the left exactness of the left adjoint $g^* \dashv g_*$ induces a commutative square of left adjoints

$$\begin{array}{ccc} \mathbf{Grpd}(\mathcal{X}') & \hookrightarrow & \mathbf{Cat}(\mathcal{X}') \\ g^* \downarrow & & \downarrow g^* \\ \mathbf{Grpd}(\mathcal{X}) & \hookrightarrow & \mathbf{Cat}(\mathcal{X}) \end{array}$$

Taking right adjoints implies for any $X_\bullet \in \mathbf{Cat}(\mathcal{X})$ that $(g_* X_\bullet)^\sim \simeq g_*(X_\bullet^\sim)$. It therefore follows for any $Y_\bullet \in \mathbf{Cat}_{\mathcal{X}}(\mathcal{Y})$ that

$$\begin{aligned} \mathbf{Gp}_\bullet(\phi_* Y) &\simeq ((\kappa_{\mathcal{X}'} \circ \phi)_* Y)^\sim && \text{(Remark 3.1.10)} \\ &\simeq ((\phi \circ \kappa_{\mathcal{X}})_* Y)^\sim && (\phi \text{ commutes with cores}) \\ &\simeq ((g_* \circ \kappa_{\mathcal{X}})_* Y)^\sim && (\phi \text{ extends } g_*) \\ &\simeq g_*(\kappa_{\mathcal{X},*} Y)^\sim \simeq g_*(\mathbf{Gp}_\bullet Y) && \text{(Remark 3.1.10)} \end{aligned}$$

In particular, if Y_\bullet is a complete Segal space, then $\mathbf{Gp}_\bullet(\phi_* Y) \simeq g_* \mathbf{Gp}_\bullet Y$ is essentially constant, ensuring that $\phi_* Y_\bullet$ is a complete Segal space by Remark 3.1.12. \square

We make the functoriality of the construction of complete Segal spaces precise by realising the construction as a functor on the categories of distributors defined in Section 3.2. For this, we need a preliminary lemma:

Lemma 3.3.4. *Fix a topos \mathcal{X} , and let \mathcal{Y} be an \mathcal{X} -distributor. Choose $\lambda \gg 0$ such that \mathcal{Y} is locally λ -presentable and the core functor $\kappa_{\mathcal{X}} : \mathcal{Y} \rightarrow \mathcal{X}$ is λ -accessible. Then,*

- (i) $\mathbf{CSS}_{\mathcal{X}}(\mathcal{Y})$ is stable under λ -filtered colimits in $\mathbf{Fun}(\Delta^{\text{op}}, \mathcal{Y})$.
- (ii) λ -filtered colimits in $\mathbf{CSS}_{\mathcal{X}}(\mathcal{Y})$ commute with λ -small limits.
- (iii) The core functor $\kappa_{\mathcal{X}} : \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}) \rightarrow \mathcal{X}$ preserves λ -filtered colimits.

Proof. This follows the proof of [Lur09b, Proposition 1.2.29] *mutatis mutandis*, noting that λ -filtered colimits commute with λ -small limits in any locally λ -presentable category. \square

Proposition 3.3.5. *For a topos \mathcal{X} , the construction of complete Segal spaces on \mathcal{X} -distributors defines a functor $\mathbf{CSS}_{\mathcal{X}} : \mathbf{Dist}_{\mathcal{X}}^R \rightarrow \mathbf{Dist}_{\mathcal{X}}^R$, where the functor $\phi_* : \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}) \rightarrow \mathbf{CSS}_{\mathcal{X}}(\mathcal{Z})$ induced by any $\phi : \mathcal{Y} \rightarrow \mathcal{Z}$ in $\mathbf{Dist}_{\mathcal{X}}^R$ acts pointwise.*

Proof. Let $\phi : \mathcal{Y} \rightarrow \mathcal{Z}$ be a morphism of $\mathbf{Dist}_{\mathcal{X}}^R$. Then, Lemma 3.3.2 implies that pointwise application of ϕ defines a functor $\phi_* : \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}) \rightarrow \mathbf{CSS}_{\mathcal{X}}(\mathcal{Z})$. Moreover, this functor preserves cores and small limits.

Choose $\lambda \gg 0$ so that $\phi : \mathcal{Y} \rightarrow \mathcal{Z}$ is a λ -accessible functor between locally λ -presentable categories. Then, Lemma 3.3.4 implies that λ -filtered colimits in $\mathbf{CSS}_{\mathcal{X}}(\mathcal{Y})$ and $\mathbf{CSS}_{\mathcal{X}}(\mathcal{Z})$ are computed pointwise, and are thus preserved by ϕ_* . Therefore, ϕ_* is indeed a morphism of $\mathbf{Dist}_{\mathcal{X}}^R$. \square

Remark 3.3.6. By Remark 3.2.2, the construction of complete Segal spaces on \mathcal{X} -distributors also defines an endofunctor on $\mathbf{Dist}_{\mathcal{X}}^L$. However, the functor $\psi_! : \mathbf{CSS}_{\mathcal{X}}(\mathcal{Z}) \rightarrow \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y})$ induced by $\psi : \mathcal{Z} \rightarrow \mathcal{Y}$ does not act pointwise; rather, if $\text{Seg}_{\mathcal{X}} : \mathbf{Fun}(\Delta^{\text{op}}, \mathcal{Y}) \rightarrow \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y})$ denotes a left adjoint to the inclusion, then $\psi_!$ is given on $Z_{\bullet} \in \mathbf{CSS}_{\mathcal{X}}(\mathcal{Z})$ by $(\psi_! Z)_{\bullet} \simeq \text{Seg}_{\mathcal{X}}(\psi_* Z_{\bullet})$.

Theorem 3.3.7. *For a topos \mathcal{X} , the functor $\mathbf{CSS}_{\mathcal{X}} : \mathbf{Dist}_{\mathcal{X}}^R \rightarrow \mathbf{Dist}_{\mathcal{X}}^R$ preserves weakly contractible limits.*

Proof. Let $\mathcal{Y}_{\bullet} : K \rightarrow \mathbf{Dist}_{\mathcal{X}}^R$ be a diagram indexed by a weakly contractible simplicial set K , and let $\mathcal{Y} := \varprojlim_k \mathcal{Y}_k$, with canonical projection maps $\varpi_k : \mathcal{Y} \rightarrow \mathcal{Y}_k$. Consider the commutative square

$$\begin{array}{ccc} \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}) & \longrightarrow & \varprojlim_k \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}_k) \\ \downarrow & & \downarrow \\ \mathbf{Fun}(\Delta^{\text{op}}, \mathcal{Y}) & \xrightarrow{\sim} & \varprojlim_k \mathbf{Fun}(\Delta^{\text{op}}, \mathcal{Y}_k) \end{array}$$

where the limits are computed in $\widehat{\mathbf{Cat}}_{\infty}$. By Remark 3.2.5, the limit $\mathcal{Y} = \varprojlim_k \mathcal{Y}_k$ can be computed on underlying categories in $\widehat{\mathbf{Cat}}_{\infty}$, implying that the bottom arrow in the above diagram is an equivalence. Remark 3.2.5 also implies that it suffices to show that the top arrow in the above diagram is an equivalence as well. As all other functors are fully faithful, the top arrow is fully faithful. Therefore, it remains to show that the top arrow is essentially surjective.

From the commutative diagram, we can identify $\varprojlim_k \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}_k)$ with the full subcategory of $\mathbf{Fun}(\Delta^{\text{op}}, \mathcal{Y})$ spanned by those $Y : \Delta^{\text{op}} \rightarrow \mathcal{Y}$ for which $\varpi_k Y : \Delta^{\text{op}} \rightarrow \mathcal{Y}_k$ lies in $\mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}_k)$ for every $k \in K$. It suffices to show that any such functor $Y : \Delta^{\text{op}} \rightarrow \mathcal{Y}$ is already a complete Segal space in \mathcal{Y} .

Certainly $Y_0 \in \mathcal{X}$, since $\varpi_k Y_0 \in \mathcal{X}$ for all $k \in K$. Since ϖ_k is a morphism of $\mathbf{Dist}_{\mathcal{X}}^R$, it preserves limits. In particular, for $n \geq 0$, the canonical map

$$Y_n \rightarrow \underbrace{Y_1 \times_{Y_0} \cdots \times_{Y_0} Y_1}_n$$

in \mathcal{Y} descends via ϖ_k to an equivalence in \mathcal{Y}_k for every $k \in K$, and is therefore also an equivalence. In particular, $Y_\bullet \in \mathbf{Cat}_{\mathcal{X}}(\mathcal{Y})$.

As K is weakly contractible, it is nonempty, so fix some $k_0 \in K$. Since ϖ_{k_0} preserves cores and limits, the proof of Lemma 3.3.2 implies that $\mathbf{Gp}_\bullet(\varpi_{k_0,*}Y_\bullet) \simeq \mathbf{Gp}_\bullet Y$. Since $\mathbf{Gp}_\bullet(\varpi_{k_0,*}Y_\bullet)$ is essentially constant, this proves that Y_\bullet is a complete Segal space, as desired. \square

An analogue to Theorem 3.3.7 for colimits is likely untrue in general, so we settle for a result that suffices for our intended application.

Theorem 3.3.8. *Fix a topos \mathcal{X} . For any \mathcal{X} -distributor \mathcal{Z} and filtered diagram $(\mathcal{Y}_k \subseteq \mathcal{Z})_{k \in K}$ of inclusions of \mathcal{X} -distributors in $\mathbf{Dist}_{\mathcal{X}}^R$, the canonical functor $\varinjlim_k^R \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}_k) \rightarrow \mathbf{CSS}_{\mathcal{X}}(\varinjlim_k^R \mathcal{Y}_k)$ is an equivalence of \mathcal{X} -distributors, where the colimits are computed in $\mathbf{Dist}_{\mathcal{X}}^R$.*

Proof. By Remark 3.2.5 and [Lur09a, Theorem 5.5.3.18], the forgetful functor $\mathbf{Dist}_{\mathcal{X}}^R \rightarrow \mathbf{Pres}_{\infty}^R$ creates weakly contractible colimits. For $k \in K$, let $\pi_k : \mathcal{Z} \rightarrow \mathcal{Y}_k$ denote a left adjoint to the corresponding inclusion. Then, define \mathcal{Y} to be the full subcategory of \mathcal{Z} spanned by those $z \in \mathcal{Z}$ such that $z \rightarrow \pi_\omega z := \varprojlim_k \pi_k z$ is an equivalence. By Lemma 2.3.4, we have that $\mathcal{Y} \simeq \varinjlim_k^R \mathcal{Y}_k$, and π_ω defines a left adjoint to the inclusion $\mathcal{Y} \subseteq \mathcal{Z}$.

The inclusions $\mathcal{Y}_k \subseteq \mathcal{Z}$ and $\mathcal{Y} \subseteq \mathcal{Z}$ induce fully faithful right adjoint functors $\mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}_k) \subseteq \mathbf{CSS}_{\mathcal{X}}(\mathcal{Z})$ and $\mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}) \subseteq \mathbf{CSS}_{\mathcal{X}}(\mathcal{Z})$, respectively, since the action on complete Segal spaces is pointwise. For $k \in K$, let $\pi_{k,!} : \mathbf{CSS}_{\mathcal{X}}(\mathcal{Z}) \rightarrow \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}_k)$ denote the left adjoint to the corresponding inclusion. Then, Lemma 2.3.4 implies that $\varinjlim_k^R \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}_k)$ can be identified with the full subcategory of $\mathbf{CSS}_{\mathcal{X}}(\mathcal{Z})$ spanned by those $Y_\bullet \in \mathbf{CSS}_{\mathcal{X}}(\mathcal{Z})$ for which $Y \rightarrow \pi_{\omega,!} Y := \varprojlim_k \pi_{k,!} Y$ is an equivalence, and $\pi_{\omega,!} : \mathbf{CSS}_{\mathcal{X}}(\mathcal{Z}) \rightarrow \varinjlim_k^R \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}_k)$ is left adjoint to the inclusion.

We have a commutative triangle

$$\begin{array}{ccc} \varinjlim_k^R \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}_k) & \xrightarrow{\quad} & \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}) \\ & \searrow & \swarrow \\ & \mathbf{CSS}_{\mathcal{X}}(\mathcal{Z}) & \end{array}$$

implying that the canonical functor $\varinjlim_k^R \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}_k) \rightarrow \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y})$ is fully faithful. It remains to prove that the functor is essentially surjective.

By definition, $\mathbf{CSS}_{\mathcal{X}}(\mathcal{Y})$ consists of those $Y_\bullet \in \mathbf{CSS}_{\mathcal{X}}(\mathcal{Z})$ for which $Y_n \in \mathcal{Y}$ for every $n \geq 0$; that is, $Y_n \rightarrow \pi_\omega Y_n$ is an equivalence for every $n \geq 0$. In other words, $\mathbf{CSS}_{\mathcal{X}}(\mathcal{Y})$ is the full subcategory of $\mathbf{CSS}_{\mathcal{X}}(\mathcal{Z})$ spanned by those Y_\bullet for which $Y_\bullet \rightarrow \pi_{\omega,*} Y_\bullet$ is an equivalence of functors $\mathbf{\Delta}^{\text{op}} \rightarrow \mathcal{Z}$.

For $Y_\bullet \in \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y})$, it therefore follows that the map $Y \rightarrow \pi_{\omega,!} Y$ factors as the chain of equivalences

$$\begin{aligned} Y &\simeq \text{Seg}_{\mathcal{X}}(Y) && (Y \text{ is a complete Segal space}) \\ &\simeq \text{Seg}_{\mathcal{X}}(\pi_{\omega,*} Y) && (Y \simeq \pi_{\omega,*} Y \text{ from the previous paragraph}) \\ &\simeq \pi_{\omega,!} Y && (\text{Remark 3.3.6}) \end{aligned}$$

and is therefore an equivalence, proving that Y_\bullet lies in the essential image of $\varinjlim_k^R \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}_k)$, as desired. \square

3.4 Distributors of sheaves of higher categories

As in Section 2, fix a topos \mathcal{X} , presented as a left exact accessible localisation of $\mathcal{P}(\mathcal{C})$ for some small category \mathcal{C} , where the localisation functor is denoted $(-)^{\#} : \mathcal{P}(\mathcal{C}) \rightarrow \mathcal{X}$. In this section, we aim to use the language of distributors and complete Segal spaces to further study the categories of sheaves of (n, r) -categories over \mathcal{C} , particularly when $n = \infty$.

Lemma 3.4.1. *Let \mathcal{Y} be an \mathcal{X} -distributor. Then, $Y_\bullet \in \mathbf{Cat}_{\mathcal{X}}(\mathcal{Y})$ is a complete Segal space if and only if it is local with respect to $|\mathbb{I}|_U^{\#} \rightarrow \Delta_U[0]$ for every $U \in \mathcal{C}$.*

Proof. Since $\mathbf{Gp}_\bullet Y \in \mathbf{Grpd}(\mathcal{X})$ satisfies the Segal condition, it is essentially constant if and only if the map $\mathbf{Gp}_1 Y \rightarrow \mathbf{Gp}_0 Y$ is an equivalence, which is equivalent to asserting that the underlying functor $\mathbf{Gp}_\bullet Y : \Delta^{\text{op}} \rightarrow \mathcal{X}$ is local with respect to $\Delta_U[1] \rightarrow \Delta_U[0]$ for every $U \in \mathcal{C}$. Now, note that

$$\begin{aligned}
\text{Hom}_{\mathbf{Fun}(\Delta^{\text{op}}, \mathcal{X})}(\Delta_U[1], \mathbf{Gp}_\bullet Y) &\simeq \text{Hom}_{\mathcal{P}(\Delta)}(\Delta[1], \mathbf{Gp}_\bullet Y(U)) && \text{(Definition 2.1.4)} \\
&\simeq \text{Hom}_{\mathbf{Cat}(\mathcal{S})}(\Delta[1], \mathbf{Gp}_\bullet Y(U)) && \text{(both } \Delta[1] \text{ and } \mathbf{Gp}_\bullet Y(U) \text{ satisfy the Segal condition)} \\
&\simeq \text{Hom}_{\mathbf{Grpd}(\mathcal{S})}(\mathbb{I}, \mathbf{Gp}_\bullet Y(U)) && (\mathbb{I} \text{ is the groupoidification of } \Delta[1]) \\
&\simeq \text{Hom}_{\mathcal{P}(\Delta)}(\mathbb{I}, \mathbf{Gp}_\bullet Y(U)) && (\mathbf{Grpd}(\mathcal{S}) \subseteq \mathcal{P}(\Delta) \text{ is fully faithful)} \\
&\simeq \text{Hom}_{\mathbf{Fun}(\Delta^{\text{op}}, \mathcal{X})}(|\mathbb{I}|_U^\#, \mathbf{Gp}_\bullet Y) && \text{(Definition 2.2.1)} \\
&\simeq \text{Hom}_{\mathbf{Grpd}(\mathcal{X})}(|\mathbb{I}|_U^\#, \mathbf{Gp}_\bullet Y) && \text{(both } |\mathbb{I}|_U^\# \text{ and } \mathbf{Gp}_\bullet Y \text{ are groupoid objects in } \mathcal{X}) \\
&\simeq \text{Hom}_{\mathbf{Cat}_{\mathcal{X}}(y)}(|\mathbb{I}|_U^\#, Y_\bullet) && \text{(Corollary 3.1.9)}
\end{aligned}$$

and similarly for $\Delta_U[0]$. Therefore, $\mathbf{Gp}_\bullet Y$ is essentially constant if and only if Y_\bullet is local with respect to $|\mathbb{I}|_U^\# \rightarrow \Delta_U[0]$ for every $U \in \mathcal{C}$, as desired. \square

Corollary 3.4.2. *The category $\mathbf{Sh}_{(\infty, r)}(\mathcal{C})$ is an \mathcal{X} -distributor for every $0 \leq r < \infty$. Moreover, we have a canonical equivalence $\mathbf{Sh}_{(\infty, r+1)}(\mathcal{C}) \simeq \mathbf{CSS}_{\mathcal{X}}(\mathbf{Sh}_{(\infty, r)}(\mathcal{C}))$.*

Proof. This is clear if $r = 0$, since $\mathbf{Sh}_{(\infty, 0)}(\mathcal{C}) \simeq \mathcal{X}$. The result for $r > 0$ follows by induction from combining Lemma 3.4.1 with Proposition 2.2.6. \square

That iterated application of $\mathbf{CSS}_{\mathcal{X}}$ to \mathcal{X} yields sheaves of (∞, r) -categories was already suggested in [Lur09b, Variant 1.3.8]. We are more interested in the case $r = \infty$. From the inductive formula in Corollary 3.4.2, we might expect that $\mathbf{Sh}_{(\infty, \infty)}(\mathcal{C})$ is an \mathcal{X} -distributor with a canonical equivalence $\mathbf{Sh}_{(\infty, \infty)}(\mathcal{C}) \simeq \mathbf{CSS}_{\mathcal{X}}(\mathbf{Sh}_{(\infty, \infty)}(\mathcal{C}))$. In fact, more is true:

Theorem 3.4.3. *Let $\mathbf{Fix}_{\mathbf{CSS}}^R$ denote the category whose objects are \mathcal{X} -distributors \mathcal{Y} with an equivalence $\mathcal{Y} \xrightarrow{\sim} \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y})$, and whose morphisms are the morphisms of $\mathbf{Dist}_{\mathcal{X}}^R$ that commute with these equivalences. Then, the terminal object of $\mathbf{Fix}_{\mathbf{CSS}}^R$ is equivalent to $\mathbf{Sh}_{(\infty, \infty)}(\mathcal{C})$.*

Proof. By Proposition 2.3.1 and Corollary 3.4.2, we can compute $\mathbf{Sh}_{(\infty, \infty)}(\mathcal{C})$ as the limit

$$\mathbf{Sh}_{(\infty, \infty)}(\mathcal{C}) \simeq \varprojlim \left(\cdots \rightarrow \mathbf{CSS}_{\mathcal{X}}^3(\mathcal{X}) \xrightarrow{\kappa_{\mathcal{X}, *}} \mathbf{CSS}_{\mathcal{X}}^2(\mathcal{X}) \xrightarrow{\kappa_{\mathcal{X}, *}} \mathbf{CSS}_{\mathcal{X}}(\mathcal{X}) \xrightarrow{\kappa_{\mathcal{X}}} \mathcal{X} \right)$$

By Remark 3.2.5, this limit can be lifted to a limit in $\mathbf{Dist}_{\mathcal{X}}^R$, proving that $\mathbf{Sh}_{(\infty, \infty)}(\mathcal{C})$ is indeed an \mathcal{X} -distributor. Moreover, Theorem 3.3.7 proves that the induced functor $\mathbf{CSS}_{\mathcal{X}}(\mathbf{Sh}_{(\infty, \infty)}(\mathcal{C})) \rightarrow \mathbf{Sh}_{(\infty, \infty)}(\mathcal{C})$ is an equivalence. Since \mathcal{X} is the terminal object in $\mathbf{Dist}_{\mathcal{X}}^R$, the theorem now follows from [Gol23, Corollaries 2.2.9 and 2.3.3]. \square

Remark 3.4.4. The crux of the above proof is the observation that $\mathbf{Sh}_{(\infty, \infty)}(\mathcal{C})$ can be realised as an instance of Adámek's construction. In particular, this shows that $\mathbf{Sh}_{(\infty, \infty)}(\mathcal{C})$ has a stronger universal property: it is the terminal $\mathbf{CSS}_{\mathcal{X}}$ -coalgebra in $\mathbf{Dist}_{\mathcal{X}}^R$.

For completeness, we can also describe the initial fixed point:

Theorem 3.4.5. *The initial object in the category $\mathbf{Fix}_{\mathbf{CSS}}^R$ defined in Theorem 3.4.3 is equivalent to the category $\mathbf{Sh}_\omega(\mathcal{C})$ of Definition 2.3.2.*

Proof. By Remark 2.3.3, Corollary 3.4.2, and Remark 3.2.5, $\mathbf{Sh}_\omega(\mathcal{C})$ is equivalent to the colimit

$$\mathbf{Sh}_\omega(\mathcal{C}) \simeq \varinjlim^R (\mathcal{X} \subseteq \mathbf{CSS}_{\mathcal{X}}(\mathcal{X}) \subseteq \mathbf{CSS}_{\mathcal{X}}^2(\mathcal{X}) \subseteq \mathbf{CSS}_{\mathcal{X}}^3(\mathcal{X}) \subseteq \cdots)$$

in \mathbf{Dist}_X^R . Since each $\mathbf{CSS}_X^r(\mathcal{X}) \simeq \mathbf{Sh}_{(\infty,r)}(\mathcal{C})$ embeds fully faithfully into $\mathbf{Sh}_{(\infty,\infty)}(\mathcal{C})$ via a morphism of \mathbf{Dist}_X^R , it follows by Theorem 3.3.8 that the induced functor $\mathbf{Sh}_\omega(\mathcal{C}) \rightarrow \mathbf{CSS}_X(\mathbf{Sh}_\omega(\mathcal{C}))$ is an equivalence. As \mathcal{X} is the initial object in \mathbf{Dist}_X^R , the theorem now follows from [Gol23, Corollaries 2.2.9 and 2.3.3]. \square

Remark 3.4.6. Let $\mathcal{C} \simeq *$, so that $\mathcal{X} \simeq \mathcal{S}$. Then, \mathcal{S} -distributors are also called *absolute distributors*. Haugseng proved in [Hau15, Theorem 7.18] that there is an equivalence $\mathcal{Y}\mathbf{Cat} \simeq \mathbf{CSS}_\mathcal{S}(\mathcal{Y})$ between \mathcal{Y} -enriched categories and complete Segal spaces in \mathcal{Y} for every absolute distributor \mathcal{Y} .

Following [Gol23, Definition 3.0.1 and Remark 3.0.2], this implies that we have an equivalence of categories $\mathbf{Sh}_{(n,r)}(*) \simeq \mathbf{Cat}_{(n,r)}$ for all $-2 \leq n \leq \infty$ and finite $0 \leq r \leq n$. Then, taking appropriate limits following [Gol23, Definition 3.0.4], we moreover have equivalences $\mathbf{Sh}_{(\infty,\infty)}(*) \simeq \mathbf{Cat}_{(\infty,\infty)}$ and $\mathbf{Sh}_\omega(*) \simeq \mathbf{Cat}_\omega$.

In particular, we see that both $\mathbf{Cat}_{(\infty,\infty)}$ and \mathbf{Cat}_ω are absolute distributors, and by [Gol23, Theorem 3.3.12] are simultaneously universal fixed points with respect to enrichment and the construction of complete Segal spaces.

3.5 Geometric morphisms and sheafification

In this section, we prove comparison results between categories of sheaves. We prove in Corollary 3.5.5 that the construction $\mathbf{Sh}_{(n,r)}(-)$ or $\mathbf{Sh}_\omega(-)$ is functorial with respect to geometric morphisms, sending geometric morphisms to right adjoints with relatively left exact left adjoints. We likewise prove that the truncation functors $\pi_{\leq r} : \mathbf{Sh}_{(\infty,r')}(\mathcal{C}) \rightarrow \mathbf{Sh}_{(\infty,r)}(\mathcal{C})$ are relatively left exact in Proposition 3.5.8 and Corollary 3.5.9. Finally, we show in Theorem 3.5.10 that $\mathbf{Sh}_{(n,r)}(\mathcal{C})$ is a relatively left exact accessible localisation of $\mathbf{Fun}(\mathcal{C}^{\text{op}}, \mathbf{Cat}_{(n,r)})$, and likewise $\mathbf{Sh}_\omega(\mathcal{C})$ is a relatively left exact accessible localisation of $\mathbf{Fun}(\mathcal{C}^{\text{op}}, \mathbf{Cat}_\omega)$.

Throughout the section, fix a topos \mathcal{X} , exhibited as a left exact accessible localisation of $\mathcal{P}(\mathcal{C})$ for some small category \mathcal{C} throughout this section.

Lemma 3.5.1. *Suppose K is sifted in the sense of [Lur09a, Definition 5.5.8.1]. Then, for all topoi \mathcal{X} , the functor $\varinjlim : \mathbf{Fun}(K, \mathcal{X}) \rightarrow \mathcal{X}$ preserves fibre products over \mathcal{X} , where \mathcal{X} is viewed as a full subcategory of $\mathbf{Fun}(K, \mathcal{X})$ spanned by the essentially constant functors.*

Proof. Suppose $F \rightarrow x \leftarrow G$ is a cospan of functors $K \rightarrow \mathcal{X}$, where $x \in \mathcal{X}$ is viewed as a constant functor. Since K is sifted, the diagonal $K \rightarrow K \times K$ is cofinal. Therefore,

$$\begin{aligned} \left(\varinjlim_{k \in K} F(k) \right) \times_x \left(\varinjlim_{\ell \in K} G(\ell) \right) &\simeq \varinjlim_{(k,\ell) \in K \times K} \left(F(k) \times_x G(\ell) \right) && \text{(colimits are universal in } \mathcal{X} \text{)} \\ &\simeq \varinjlim_{j \in K} \left(F(j) \times_x G(j) \right) && \text{(} K \rightarrow K \times K \text{ is cofinal)} \end{aligned}$$

showing that $\varinjlim : \mathbf{Fun}(K, \mathcal{X}) \rightarrow \mathcal{X}$ preserves the fibre product $F \times_x G$. \square

Lemma 3.5.2. *Fix an \mathcal{X} -distributor \mathcal{Y} . Suppose a morphism $Y_\bullet \rightarrow Z_\bullet$ in $\mathbf{Cat}_\mathcal{X}(\mathcal{Y})$ is a Segal equivalence. Then, for any cospan $Z_\bullet \rightarrow x \leftarrow Z'_\bullet$ in $\mathbf{Cat}_\mathcal{X}(\mathcal{Y})$ with $x \in \mathcal{X}$, the induced map $Y \times_x Z' \rightarrow Z \times_x Z'$ is a Segal equivalence.*

Proof. The functor $Y \times_x Z' \rightarrow Z \times_x Z'$ is certainly fully faithful in the sense of (E1), since limits commute with limits. We need to show that the map is also essentially surjective in the sense of (E2); that is, $|\mathbf{Gp}_\bullet(Y \times_x Z')| \rightarrow |\mathbf{Gp}_\bullet(Z \times_x Z')|$ is an equivalence in \mathcal{X} .

Since \mathbf{Gp}_\bullet is a right adjoint that restricts to the identity on \mathcal{X} , we have $\mathbf{Gp}_\bullet(Y \times_x Z') \simeq (\mathbf{Gp}_\bullet Y) \times_x (\mathbf{Gp}_\bullet Z')$ and likewise for $Z \times_x Z'$. On the other hand, since Δ^{op} is sifted by [Lur09a, Lemma 5.5.8.4], geometric realisation also preserves fibre products over \mathcal{X} by Lemma 3.5.1.

Therefore, $Y \times_x Z' \rightarrow Z \times_x Z'$ satisfies (E2) if and only if the map $|\mathbf{Gp}_\bullet Y| \times_x |\mathbf{Gp}_\bullet Z'| \rightarrow |\mathbf{Gp}_\bullet Z| \times_x |\mathbf{Gp}_\bullet Z'|$ is an equivalence, which follows from the fact that this is the base change of the map $|\mathbf{Gp}_\bullet Y| \rightarrow |\mathbf{Gp}_\bullet Z|$ along $|\mathbf{Gp}_\bullet Z'| \rightarrow x$, which is an equivalence since $Y \rightarrow Z$ is a Segal equivalence. \square

Proposition 3.5.3. *For an \mathcal{X} -distributor \mathcal{Y} , the left adjoint $\text{Seg}_{\mathcal{X}} : \mathbf{Cat}_{\mathcal{X}}(\mathcal{Y}) \rightarrow \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y})$ to inclusion preserves fibre products over \mathcal{X} .*

Proof. Suppose $Y \rightarrow x \leftarrow Y'$ is a cospan in $\mathbf{Cat}_{\mathcal{X}}(\mathcal{Y})$, where $x \in \mathcal{X}$. Since $(\text{Seg}_{\mathcal{X}} Y) \times_x (\text{Seg}_{\mathcal{X}} Y')$ is a limit of complete Segal spaces, it is also a complete Segal space. Therefore, it suffices to show that the map $Y \times_x Y' \rightarrow (\text{Seg}_{\mathcal{X}} Y) \times_x (\text{Seg}_{\mathcal{X}} Y')$ is a Segal equivalence, as it would then induce an equivalence $\text{Seg}_{\mathcal{X}}(Y \times_x Y') \rightarrow (\text{Seg}_{\mathcal{X}} Y) \times_x (\text{Seg}_{\mathcal{X}} Y')$.

Since $Y \times_x Y' \rightarrow (\text{Seg}_{\mathcal{X}} Y) \times_x (\text{Seg}_{\mathcal{X}} Y')$ factors as

$$Y \times_x Y' \rightarrow (\text{Seg}_{\mathcal{X}} Y) \times_x Y' \rightarrow (\text{Seg}_{\mathcal{X}} Y) \times_x (\text{Seg}_{\mathcal{X}} Y')$$

the fact that it is a Segal equivalence follows from Lemma 3.5.2, since $Y \rightarrow \text{Seg}_{\mathcal{X}} Y$ and $Y' \rightarrow \text{Seg}_{\mathcal{X}} Y'$ are Segal equivalences. \square

Lemma 3.5.4. *In the situation of Lemma 3.3.2, suppose $\phi : \mathcal{Y} \rightarrow \mathcal{Y}'$ admits a left adjoint ψ . Then, the induced functor $\phi_* : \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}) \rightarrow \mathbf{CSS}_{\mathcal{X}'}(\mathcal{Y}')$ admits a left adjoint $\psi_!$. If moreover $\psi : \mathcal{Y}' \rightarrow \mathcal{Y}$ preserves fibre products over \mathcal{X}' , then the same is true for $\psi_!$.*

Proof. Choose $\lambda \gg 0$ so that $\phi : \mathcal{Y} \rightarrow \mathcal{Z}$ is a λ -accessible functor between locally λ -presentable categories. Then, Lemma 3.3.4 implies that λ -filtered colimits in $\mathbf{CSS}_{\mathcal{X}}(\mathcal{Y})$ and $\mathbf{CSS}_{\mathcal{X}}(\mathcal{Z})$ are computed pointwise, and are thus preserved by ϕ_* . In particular, [Lur09a, Corollary 5.5.2.9] implies that ϕ_* admits a left adjoint $\psi_!$.

If ψ preserves fibre products over \mathcal{X}' , then the induced functor $\psi_* : \mathbf{Cat}_{\mathcal{X}'}(\mathcal{Y}') \rightarrow \mathbf{Cat}_{\mathcal{X}}(\mathcal{Y})$ is left adjoint to $\phi_* : \mathbf{Cat}_{\mathcal{X}}(\mathcal{Y}) \rightarrow \mathbf{Cat}_{\mathcal{X}'}(\mathcal{Y}')$. Taking left adjoints of the functors in the commutative square

$$\begin{array}{ccc} \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}) & \xrightarrow{\phi_*} & \mathbf{CSS}_{\mathcal{X}'}(\mathcal{Y}') \\ \downarrow & & \downarrow \\ \mathbf{Cat}_{\mathcal{X}}(\mathcal{Y}) & \xrightarrow[\phi_*]{} & \mathbf{Cat}_{\mathcal{X}'}(\mathcal{Y}') \end{array}$$

yields the commutative square in the diagram

$$\begin{array}{ccccc} \mathbf{CSS}_{\mathcal{X}'}(\mathcal{Y}) & \longleftarrow & \mathbf{Cat}_{\mathcal{X}'}(\mathcal{Y}') & \xrightarrow{\psi_*} & \mathbf{Cat}_{\mathcal{X}}(\mathcal{Y}) \\ & \searrow \text{id} & \downarrow \text{Seg}_{\mathcal{X}'} & & \downarrow \text{Seg}_{\mathcal{X}} \\ & & \mathbf{CSS}_{\mathcal{X}'}(\mathcal{Y}') & \xrightarrow[\psi_!]{=} & \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}) \end{array}$$

where the vertical functors preserve fibre products over \mathcal{X}' by Proposition 3.5.3, and the inclusion preserves all limits from being a right adjoint. Therefore, $\psi_!$ preserves fibre products over \mathcal{X}' as well. \square

Corollary 3.5.5. *Given a geometric morphism $g_* : \mathbf{Sh}(\mathcal{C}) \rightarrow \mathbf{Sh}(\mathcal{C}')$ of topoi of sheaves, we have for all $-2 \leq n \leq \infty$ and $0 \leq r \leq n$ an adjunction*

$$g^* : \mathbf{Sh}_{(n,r)}(\mathcal{C}') \rightleftarrows \mathbf{Sh}_{(n,r)}(\mathcal{C}) : g_*$$

and similarly an adjunction

$$g^* : \mathbf{Sh}_{\omega}(\mathcal{C}') \rightleftarrows \mathbf{Sh}_{\omega}(\mathcal{C}) : g_*$$

where each g_* acts pointwise, and the corresponding left adjoints g^* are relatively left exact.

Lemma 3.5.6. *Let K be a filtered simplicial set, and $(\mathcal{A}^k \rightarrow \mathcal{B}^k)_{k \in K}$ a diagram in $\mathbf{Fun}(\Delta[1], \mathbf{Cat}_{\infty})$ consisting of cofinal functors. Then, the colimit of this diagram is again a cofinal functor.*

Proof. Let $\mathcal{A} \rightarrow \mathcal{B}$ denote the colimit of the diagram of cofinal functors. By Quillen's Theorem A [Lur09a, Theorem 4.1.3.1], we show for any $B \in \mathcal{B}$ that $\mathcal{A}_{B/} := \mathcal{A} \times_{\mathcal{B}} \mathcal{B}_{B/}$ is weakly contractible.

Choose $k \in K$ so that $B \in \mathcal{B}^k$; that is, so that B is in the essential image of the coprojection $\mathcal{B}^k \rightarrow \mathcal{B}$. Since K is filtered, the canonical functor $K_{k/} \rightarrow K$ is cofinal. Therefore,

$$\begin{aligned}
\mathcal{A} \times_{\mathcal{B}} \mathcal{B}_{B/} &\simeq \left(\varinjlim_{\ell} \mathcal{A}^{\ell} \right) \times \left(\varinjlim_{\ell} \mathcal{B}^{\ell} \right)_{B/} \\
&\simeq \left(\varinjlim_{k \rightarrow \ell} \mathcal{A}^{\ell} \right) \times \left(\varinjlim_{k \rightarrow \ell} \mathcal{B}^{\ell} \right)_{B/} && \text{(cofinality of } K_{k/} \rightarrow K) \\
&\simeq \left(\varinjlim_{k \rightarrow \ell} \mathcal{A}^{\ell} \right) \times \left(\varinjlim_{k \rightarrow \ell} \mathcal{B}_{B/}^{\ell} \right) && (B \in \mathcal{B}^k) \\
&\simeq \varinjlim_{k \rightarrow \ell} \left(\mathcal{A}^{\ell} \times_{\mathcal{B}^{\ell}} \mathcal{B}_{B/}^{\ell} \right) && \text{(filtered colimits are left exact in } \mathbf{Cat}_{\infty})
\end{aligned}$$

By assumption, every $\mathcal{A}^{\ell} \times_{\mathcal{B}^{\ell}} \mathcal{B}_{B/}^{\ell}$ is weakly contractible for every $k \rightarrow \ell$, implying the same for $\mathcal{A} \times_{\mathcal{B}} \mathcal{B}_{B/}$, as desired. \square

Corollary 3.5.7. *Sifted categories are closed under filtered colimits in \mathbf{Cat}_{∞} .*

Proposition 3.5.8. *For $0 \leq r < r' \leq \infty$, the functor $\pi_{\leq r} : \mathbf{Sh}_{(\infty, r')}(\mathcal{C}) \rightarrow \mathbf{Sh}_{(\infty, r)}(\mathcal{C})$ is relatively left exact.*

Proof. We prove the proposition by induction on r , so suppose first that $r = 0$.

Let Δ_{∞} denote the full subcategory of $\Delta^{\times \infty}$ spanned by those $[\vec{k}] = (k_0, k_1, k_2, \dots)$ such that $k_n = 0$ for all $n \gg 0$. For $0 \leq r' \leq \infty$, denote by $\Delta_{r'}$ the full subcategory of Δ_{∞} spanned by those $[\vec{k}]$ such that $k_n = 0$ for all $n \geq r'$. Then, we have a fully faithful inclusion $\Sigma_{r'} \subseteq \Delta_{r'}$ for every $0 \leq r' \leq \infty$.

The functor $\pi_{\leq 0} : \mathbf{Sh}_{(\infty, r')}(\mathcal{C}) \rightarrow \mathbf{Sh}_{(\infty, 0)}(\mathcal{C}) \simeq \mathcal{X}$ is left adjoint to the diagonal $\mathcal{X} \subseteq \mathbf{Sh}_{(\infty, r')}(\mathcal{C})$. The composite of right adjoints

$$\mathcal{X} \subseteq \mathbf{Sh}_{(\infty, r')}(\mathcal{C}) \subseteq \mathbf{Fun}(\Sigma_{r'}^{\text{op}}, \mathcal{X}) \subseteq \mathbf{Fun}(\Delta_{r'}^{\text{op}}, \mathcal{X})$$

where the rightmost inclusion is right Kan extension along $\Sigma_{r'} \subseteq \Delta_{r'}$ is precisely the diagonal inclusion, which implies that $\pi_{\leq 0}$ is the restriction of the left adjoint $\varinjlim : \mathbf{Fun}(\Delta_{r'}^{\text{op}}, \mathcal{X}) \rightarrow \mathcal{X}$ to the diagonal. By Lemma 3.5.1, it suffices to prove that $\Delta_{r'}^{\text{op}}$ is sifted for every $0 \leq r' \leq \infty$.

For $r' < \infty$, note that $\Delta_{r'} \simeq \Delta^{\times r'}$. Since Δ^{op} is sifted by [Lur09a, Lemma 5.5.8.4], and [Lur09a, Corollary 4.1.1.13 and Proposition 4.1.1.3(2)] imply that sifted categories are closed under finite products, it follows that $\Delta_{r'}^{\text{op}}$ is sifted for r' finite. Finally, since $\Delta_{\infty} \simeq \varinjlim (\Delta_0 \subseteq \Delta_1 \subseteq \Delta_2 \subseteq \dots)$, it follows from Corollary 3.5.7 that $\Delta_{\infty}^{\text{op}}$ is sifted as well.

Now, let $r > 0$, and suppose all $\pi_{\leq (r-1)}$ preserve fibre products over \mathcal{X} . Then,

$$\pi_{\leq (r-1), *}: \mathbf{Fun}(\Delta^{\text{op}}, \mathbf{Sh}_{(\infty, r'-1)}(\mathcal{C})) \rightarrow \mathbf{Fun}(\Delta^{\text{op}}, \mathbf{Sh}_{(\infty, r-1)}(\mathcal{C}))$$

restricts to a functor $\mathbf{Cat}_{\mathcal{X}}(\mathbf{Sh}_{(\infty, r'-1)}(\mathcal{C})) \rightarrow \mathbf{Cat}_{\mathcal{X}}(\mathbf{Sh}_{(\infty, r-1)}(\mathcal{C}))$ that also preserves fibre products over \mathcal{X} .

As $\pi_{\leq r} \simeq \pi_{\leq (r-1), !}$ is the image of $\pi_{\leq (r-1)}$ under $\mathbf{CSS}_{\mathcal{X}} : \mathbf{Dist}_{\mathcal{X}}^L \rightarrow \mathbf{Dist}_{\mathcal{X}}^L$, and $\pi_{\leq (r-1)}$ preserves fibre products over \mathcal{X} by assumption, it follows by Lemma 3.5.4 that $\pi_{\leq r}$ preserves fibre products over \mathcal{X} as well. \square

Corollary 3.5.9. *The localisation functor $\pi_{\leq \omega} : \mathbf{Sh}_{(\infty, \infty)}(\mathcal{C}) \rightarrow \mathbf{Sh}_{\omega}(\mathcal{C})$ is relatively left exact.*

The remainder of this section is dedicated to proving the following result:

Theorem 3.5.10 (Sheafification). *For $-2 \leq n \leq \infty$ and $0 \leq r \leq n + 2$, the category $\mathbf{Sh}_{(n, r)}(\mathcal{C})$ includes fully faithfully into $\mathbf{Fun}(\mathcal{C}^{\text{op}}, \mathbf{Cat}_{(n, r)})$, and this inclusion admits a relatively left exact left adjoint.*

Likewise, $\mathbf{Sh}_{\omega}(\mathcal{C})$ includes fully faithfully into $\mathbf{Fun}(\mathcal{C}^{\text{op}}, \mathbf{Cat}_{\omega})$, and the inclusion admits a relatively left exact left adjoint.

Proof. Follows from combining Proposition 3.5.12 below with Lemma 3.5.4. \square

Lemma 3.5.11. *Let \mathcal{Y} be an \mathcal{X} -distributor. For every small category \mathcal{K} , the category $\mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathcal{Y})$ is a distributor over $\mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathcal{X})$. Moreover, $\mathbf{CSS}_{\mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathcal{X})}(\mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathcal{Y})) \simeq \mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}))$.*

Proof. $\mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathcal{X})$ and $\mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathcal{Y})$ are certainly locally presentable, verifying (D1). The inclusion $\mathcal{X} \subseteq \mathcal{Y}$ preserves small limits and colimits, and limits and colimits in functor categories are computed pointwise, which proves (D2). By [Lur09b, Corollary 1.2.5], $\mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathcal{Y})$ is then a distributor over $\mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathcal{X})$ if and only if the following holds:

- (*) For any simplicial set J and natural transformation $\bar{\alpha} : \bar{p} \Rightarrow \bar{q} : J^{\triangleright} \rightarrow \mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathcal{Y})$ such that \bar{q} is a colimit diagram in $\mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathcal{X})$ and the naturality squares of $\alpha = \bar{\alpha}|_K$ are pullback squares, then \bar{p} is a colimit diagram in $\mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathcal{Y})$ if and only if the naturality squares of $\bar{\alpha}$ are pullback squares.

This condition can be checked pointwise, and [Lur09b, Corollary 1.2.5] implies that the analogous results hold for $\mathcal{X} \subseteq \mathcal{Y}$. Therefore, $\mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathcal{Y})$ is then a distributor over $\mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathcal{X})$, as desired.

Since $\mathbf{Fun}(\Delta^{\mathrm{op}}, \mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathcal{Y})) \simeq \mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathbf{Fun}(\Delta^{\mathrm{op}}, \mathcal{Y}))$ and fibre products are computed pointwise in functor categories, we have $\mathbf{Cat}_{\mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathcal{X})}(\mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathcal{Y})) \simeq \mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathbf{Cat}_{\mathcal{X}}(\mathcal{Y}))$. Finally, since the conditions for a transformation to be a Segal equivalence can be checked pointwise, this equivalence descends to an equivalence $\mathbf{CSS}_{\mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathcal{X})}(\mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathcal{Y})) \simeq \mathbf{Fun}(\mathcal{K}^{\mathrm{op}}, \mathbf{CSS}_{\mathcal{X}}(\mathcal{Y}))$. \square

Proposition 3.5.12. *Let $-2 \leq n \leq \infty$ and $0 \leq r \leq n + 2$. If $\mathcal{X} \simeq \mathcal{P}(\mathcal{C})$, then we have an equivalence $\mathbf{Sh}_{(n,r)}(\mathcal{C}) \simeq \mathbf{Fun}(\mathcal{C}^{\mathrm{op}}, \mathbf{Cat}_{(n,r)})$. Similarly, $\mathbf{Sh}_{\omega}(\mathcal{C}) \simeq \mathbf{Fun}(\mathcal{C}^{\mathrm{op}}, \mathbf{Cat}_{\omega})$.*

Proof. Suppose first that $n = \infty$. If $r = 0$, then there is nothing to prove, as $\mathbf{Sh}_{(\infty,0)}(\mathcal{C}) \simeq \mathcal{P}(\mathcal{C})$ and $\mathbf{Cat}_{(\infty,0)} \simeq \mathcal{S}$. Then, iteratively applying Lemma 3.5.11 proves the proposition if r is finite. For $r = \infty$, we have

$$\begin{aligned} \mathbf{Sh}_{(\infty,\infty)}(\mathcal{C}) &\simeq \varprojlim \left(\cdots \rightarrow \mathbf{Sh}_{(\infty,2)}(\mathcal{C}) \xrightarrow{\kappa_{\leq 1}} \mathbf{Sh}_{(\infty,1)}(\mathcal{C}) \xrightarrow{\kappa_{\leq 0}} \mathbf{Sh}_{(\infty,0)}(\mathcal{C}) \right) \\ &\simeq \varprojlim \left(\cdots \rightarrow \mathbf{Fun}(\mathcal{C}^{\mathrm{op}}, \mathbf{Cat}_{(\infty,2)}) \xrightarrow{\kappa_{\leq 1,*}} \mathbf{Fun}(\mathcal{C}^{\mathrm{op}}, \mathbf{Cat}_{(\infty,1)}) \xrightarrow{\kappa_{\leq 0,*}} \mathbf{Fun}(\mathcal{C}^{\mathrm{op}}, \mathbf{Cat}_{(\infty,0)}) \right) \\ &\simeq \mathbf{Fun} \left(\mathcal{C}^{\mathrm{op}}, \varprojlim \left(\cdots \rightarrow \mathbf{Cat}_{(\infty,2)} \xrightarrow{\kappa_{\leq 2}} \mathbf{Cat}_{(\infty,1)} \xrightarrow{\kappa_{\leq 1}} \mathbf{Cat}_{(\infty,0)} \right) \right) \\ &\simeq \mathbf{Fun}(\mathcal{C}^{\mathrm{op}}, \mathbf{Cat}_{(\infty,\infty)}) \end{aligned}$$

Similarly,

$$\begin{aligned} \mathbf{Sh}_{\omega}(\mathcal{C}) &\simeq \varprojlim \left(\cdots \rightarrow \mathbf{Sh}_{(\infty,2)}(\mathcal{C}) \xrightarrow{\pi_{\leq 1}} \mathbf{Sh}_{(\infty,1)}(\mathcal{C}) \xrightarrow{\pi_{\leq 0}} \mathbf{Sh}_{(\infty,0)}(\mathcal{C}) \right) \\ &\simeq \varprojlim \left(\cdots \rightarrow \mathbf{Fun}(\mathcal{C}^{\mathrm{op}}, \mathbf{Cat}_{(\infty,2)}) \xrightarrow{\pi_{\leq 1,*}} \mathbf{Fun}(\mathcal{C}^{\mathrm{op}}, \mathbf{Cat}_{(\infty,1)}) \xrightarrow{\pi_{\leq 0,*}} \mathbf{Fun}(\mathcal{C}^{\mathrm{op}}, \mathbf{Cat}_{(\infty,0)}) \right) \\ &\simeq \mathbf{Fun} \left(\mathcal{C}^{\mathrm{op}}, \varprojlim \left(\cdots \rightarrow \mathbf{Cat}_{(\infty,2)} \xrightarrow{\pi_{\leq 2}} \mathbf{Cat}_{(\infty,1)} \xrightarrow{\pi_{\leq 1}} \mathbf{Cat}_{(\infty,0)} \right) \right) \\ &\simeq \mathbf{Fun}(\mathcal{C}^{\mathrm{op}}, \mathbf{Cat}_{\omega}) \end{aligned}$$

Now suppose $n < \infty$. Note that $\mathbf{Sh}_{(n,r)}(\mathcal{C})$ is the localisation of $\mathbf{Sh}_{(\infty,r)}(\mathcal{C}) \simeq \mathbf{Fun}(\mathcal{C}^{\mathrm{op}}, \mathbf{Cat}_{(\infty,r)})$ at the maps $\Sigma^m(\mathbb{S}^{k-m} \times \Delta_U[0]) \rightarrow \Sigma^m \Delta_U[0]$ for $m \geq r$, $k > n$, and $U \in \mathcal{C}$. A functor $F : \mathcal{C}^{\mathrm{op}} \rightarrow \mathbf{Cat}_{(\infty,r)}$ is local with respect to $\Sigma^m(\mathbb{S}^{k-m} \times \Delta_U[0]) \rightarrow \Sigma^m \Delta_U[0]$ if and only if $F(U) \in \mathbf{Cat}_{(\infty,r)}$ is local with respect to $\Sigma^m \mathbb{S}^{k-m} \rightarrow \Sigma^m *$. By [GH15, Proposition 6.1.7], it follows that F lies in $\mathbf{Sh}_{(n,r)}(\mathcal{C})$ if and only if $F(U) \in \mathbf{Cat}_{(n,r)}$ for every $U \in \mathcal{C}$, as desired. \square

References

- [Bar05] Clark Barwick. $(\infty, n)\mathbf{Cat}$ as a closed model category. PhD thesis, University of Pennsylvania, 2005. Unpublished.

- [BSP21] Clark Barwick and Christopher Schommer-Pries. On the unicity of the theory of higher categories. *Journal of the American Mathematical Society*, 34:1011–1058, 2021. arXiv:1112.0040v6. doi:10.1090/jams/972.
- [GH15] David Gepner and Rune Haugseng. Enriched ∞ -categories via non-symmetric ∞ -operads. *Advances in Mathematics*, 279:575–716, 2015. arXiv:1312.3178v4. doi:10.1016/j.aim.2015.02.007.
- [Gol23] Z. Goldthorpe. Homotopy theories of (∞, ∞) -categories as universal fixed points with respect to weak enrichment. *International Mathematics Research Notices*, 2023. arXiv:2307.00442. doi:10.1093/imrn/rnad196.
- [Hau15] Rune Haugseng. Rectification of enriched ∞ -categories. *Algebraic & Geometric Topology*, 15(4):1931–1982, 2015. arXiv:1312.3881v4. doi:10.2140/agt.2015.15.1931.
- [Lur09a] Jacob Lurie. *Higher Topos Theory*, volume 170 of *the Annals of Mathematical Studies*. Princeton University Press, 2009. arXiv:math/0608040v4.
- [Lur09b] Jacob Lurie. $(\infty, 2)$ -categories and the Goodwillie calculus I, 2009. Preprint. arXiv:0905.0462v2.
- [Rez01] Charles Rezk. A model for the homotopy theory of homotopy theory. *Transactions of the American Mathematical Society*, 353:973–1007, 2001. arXiv:9811037v3. doi:10.1090/S0002-9947-00-02653-2.
- [RS22] Shaul Ragimov and Tomer Schlank. The ∞ -categorical reflection theorem and applications, 2022. Preprint. arXiv:2207.09244v1.
- [Sim00] Carlos Simpson. A closed model structure for n -categories, internal *Hom*, n -stacks and generalized Seifert-Van Kampen, 2000. Preprint. arXiv:9704006v2.
- [Sim11] Carlos Simpson. *Homotopy Theory of Higher Categories*, volume 19 of *New Mathematical Monographs*. Cambridge University Press, 2011. hal-00449826. doi:10.1017/CB09780511978111.