

Minimal Fractional Topological Insulator in half-filled conjugate moiré Chern bands

Chao-Ming Jian¹ and Cenke Xu²

¹*Department of Physics, Cornell University, Ithaca, New York 14853, USA*

²*Department of Physics, University of California, Santa Barbara, CA 93106*

(Dated: March 13, 2024)

We propose a “minimal” fractional topological insulator (mFTI), motivated by the recent experimental report on the signatures of FTI at total filling factor $\nu_{\text{tot}} = 3$ in a transition metal dichalcogenide (TMD) moiré system. The observed FTI at $\nu_{\text{tot}} = 3$ is likely given by a topological state living in a pair of half-filled conjugate Chern bands with Chern number $C = \pm 1$ on top of another pair of fully-filled conjugate Chern bands. We propose the mFTI as a strong candidate topological state in the half-filled conjugate Chern bands. The mFTI is characterized by the following features: (1) It is a fully gapped topological order with 32 Abelian anyons (including the electron); (2) the minimally-charged anyon carries electric charge $e^* = e/2$, together with the spin-Hall conductivity, implying that the mFTI’s robust edge state remains gapless whenever time-reversal symmetry and charge conversation are present; (3) the mFTI is “minimal” in the sense that it has the smallest total quantum dimension (a metric for the topological order’s complexity) within all the topological orders that can potentially be realized at the same electron filling of the system and with the same Hall transports; (4) the mFTI is the common descendant of multiple valley-decoupled “product topological orders” with larger quantum dimensions. It is also a symmetry-enriched topological order promoted from multiple symmetry-protected topological states, after gauging part of their symmetries.

I. INTRODUCTION

The recent discovery of fractional Chern insulator^{1,2} and fractional quantum anomalous Hall insulator at zero magnetic field³⁻⁷ have led to a new excitement on the strongly correlated states of matter in the moiré systems. Especially, the time-reversal-invariant fractional topological insulator (FTI) reported most recently in transition metal dichalcogenide (TMD) homobilayer moiré heterostructure⁸, twisted bilayer MoTe₂ to be more precise, represents a potentially new topological state of matter which could be fundamentally different from the previously known fractional quantum Hall state. In this TMD moiré system with spin-valley locking, the electron bands relevant to the experimental observations include two bands with Chern number $C = +1$ at valley-1 (spin-up), and two bands with opposite Chern number $C = -1$ at valley-2 (spin-down).⁹ The observed FTI signal appears at the total filling $\nu_{\text{tot}} = 3$. If the system preserves the time-reversal symmetry, and one ignores the fully occupied lower bands in each valley, the most relevant bands are a pair of time-reversal conjugate Chern bands with $C = \pm 1$, each with electron filling factor $\nu = 1/2$. The experiment⁸ on twisted bilayer MoTe₂ observed a series of appealing evidence that supports the existence of a certain type of time-reversal-invariant FTI with fractionalized helical edge states and spin-Hall conductivity $\sigma^{\text{sh}} = 1/2$ (when we only consider the contribution from the half-filled bands). The experimental evidence includes the nonlocal transport signal, the quantization of the edge conductance at fractional filling, and the suppression of the edge conductance when the time-reversal symmetry is explicitly broken by an in-plane magnetic field, analogous to the quantum spin Hall insulator¹⁰. It is worth noting that the relevant twisted bilayer MoTe₂

system exhibits an S_z spin quantum number conservation in addition to the charge conservation and time-reversal symmetry.⁸

The experimental facts have not yet completely pinned down the very nature of the reported FTI. A possibility is that the FTI is simply a product of a pair of time-reversal conjugate topological orders (TO) in each valley, like some of the FTIs considered theoretically in the past^{11,12} (For a review of previous theoretical discussion of FTI, please refer to Ref. 13. We also refer to the seminal work Ref. 14 for general properties of generic FTIs). In the context of the TMD moiré system, the Chern band at valley-1 (valley-2) with Chern number $C = 1$ ($C = -1$) can be viewed as a Landau level due to the real space pseudo magnetic field^{15,16} (as illustrated in the upper panel of Fig. 1). For a half-filled Landau level, the candidate topological order of electrons could be the nonabelian Pfaffian state¹⁷⁻¹⁹, anti-Pfaffian state²⁰, PH-Pfaffian state²¹, the Abelian $U(1)_8$ state²², or the “331” state^{23,24}. Hence, the FTI could be the product of one of the TOs mentioned above in one valley, and its time-reversal conjugate in the other valley. In such product TOs, the TOs from the two valleys essentially decouple.

In this work, we explore what kind of FTI is the “minimal” one, in the sense that it gives the desired experimental signals (especially the spin-Hall conductivity) and also has the minimal total quantum dimension (which is, in simple terms, a metric for the topological order’s complexity). We take charge conversation, spin- S_z conservation, and time-reversal symmetry into account in the search for the minimal FTI. It turns out that the minimal FTI (mFTI) is *NOT* a simple product of a pair of conjugate TOs of electrons from the two valleys, though it can be viewed as the common descendant of multiple “product TOs” via anyon condensations. Our mFTI also has




FIG. 1. Mutual flux attachment construction of the mFTI. Upper panel: The pair of conjugate Chern bands with Chern number $C = \pm 1$ are both half-filled. The green and yellow arrows indicate their contribution to the effective magnetic field seen by the inter-valley Cooper pair and the effective spin-magnetic field seen by the inter-valley excitons. In total, the Cooper pair sees no net magnetic field while the exciton sees a spin flux $\Phi^s = 2\Phi_0$ per moire unit cell. Lower panel: The inter-valley Cooper pair (green ball) is attached with flux $-4\Phi_0$ seen by the spin-1 inter-valley exciton (yellow ball), forming a composite boson. Hence, the exciton sees a total zero magnetic field when each valley has a filling factor $\nu = 1/2$. The Cooper pair also sees the exciton as a $-4\Phi_0$ flux. This flux attachment turns the exciton into another composite boson. $e_{1,2}$ and $h_{1,2}$ stand for the electrons and holes in the two valleys.

a spin-Hall conductivity $\sigma^{\text{sh}} = 1/2$, which is consistent with the experimental observation once the fully filled bands are included. This spin-Hall conductivity is also half of the value for the elementary non-interacting quantum spin-Hall insulators^{25,26}. The minimally charged anyon of the mFTI carries an electric charge $e^* = e/2$. In contrast, all the product TOs with the same response must have a minimal charge equal to or smaller than $e/4$. Additionally, according to the criterion established in Ref. 14, our mFTI has robust gapless edge states that remain protected by time-reversal symmetry and charge conservation even when S_z is not conserved.

II. CONSTRUCTION OF THE MFTI

The mFTI may be perceived and constructed in various ways. Let us first explore the most intuitive flux-attachment picture of the mFTI, which is based on the kinematics of both the inter-valley Cooper pair, and also the inter-valley exciton. In the TMD system, the motion of electrons in each valley is governed by a real-space pseudo magnetic field. Hence, the physics of a Chern band with Chern number $C = +1$ can be viewed as a Landau level in a magnetic field^{15,16}. More precisely, an electron in valley-1(2) will see magnetic flux $\pm\Phi_0 = 2\pi$

in each moiré unit cell. If one performs the standard flux attachment to electrons (independently for each valley), it is most natural to construct the conjugate composite Fermi liquid state discussed recently^{27–30}, which is a compressible state in the bulk. However, there is another alternative flux attachment which naturally leads to an incompressible FTI.

Let's start by considering the bosonic objects of the system: the inter-valley Cooper pairs and inter-valley excitons. A spin singlet inter-valley Cooper pair (whose annihilation operator is labeled as b^c) carrying charge- $2e$ sees zero net magnetic fields from the two valleys. An inter-valley exciton (labeled as b^s) carrying zero charges and spin-1 sees “spin” flux $\Phi^s = 2\Phi_0$ in each moiré unit cell. b^s should be viewed as a bosonic rotor. It can have both positive and negative density fluctuation but has zero average density when the system is time-reversal invariant. At half-filling of the conjugate pair of Chern bands, the density of b^c is at $1/2$ particle per moiré unit cell. It is energetically favorable for each b^c to be attached with flux $-4\Phi_0$ seen by b^s , such that b^s sees zero total spin flux. Likewise, b^c will view each b^s as $-4\Phi_0$ flux. Here, we note that since b^s has zero average density, b^c still sees zero total flux. A schematic illustration of the flux attachment is shown in Fig. 1. After the mutual flux attachment, the “composite bosons”, formed by b^c and b^s bound with fluxes, can condense. The condensate of the composite bosons is an incompressible state without any spontaneous symmetry breaking due to the attached gauge fluxes.

The state constructed using the mutual flux attachment picture described above is described by the following Chern-Simons theory:

$$\mathcal{L}_{\text{CS}} = \frac{4i}{2\pi} a^s da^c + \frac{i}{2\pi} A^s da^s + \frac{2i}{2\pi} A^e da^c. \quad (1)$$

Here $a^s da^c$ is a short-handed notation for $\epsilon_{\mu\nu\rho} a_\mu^s \partial_\nu a_\rho^c$. a_μ^s and a_μ^c are the “dual gauge field” of the current of b^s and b^c respectively: $J_\mu^{c,s} = \frac{1}{2\pi} \epsilon_{\mu\nu\rho} \partial_\nu a_\rho^{c,s}$. A_μ^e and A_μ^s are the background charge and “spin gauge fields”: $A_\mu^e = (A_{1,\mu} + A_{2,\mu})/2$, $A_\mu^s = A_{1,\mu} - A_{2,\mu}$. Here $A_{1,2}$ are introduced as the background gauge fields of valley-1 (spin-up) and valley-2 (spin-down) electrons. They arise from the conservation of electron number in each valley/spin species. If we integrate out the dynamical gauge fields a^c and a^s from Eq. (1), the response theory for this state is

$$\mathcal{L}_{\text{res}} = \frac{1}{2} \frac{i}{2\pi} A^e dA^s = \frac{1}{2} \frac{i}{4\pi} A_1 dA_1 - \frac{1}{2} \frac{i}{4\pi} A_2 dA_2, \quad (2)$$

indicating the desired spin-Hall conductivity $\sigma^{\text{sh}} = 1/2$. Also, there are no “diagonal” Hall responses for the electric charge and S_z quantum numbers separately, which is consistent with time-reversal symmetry.

The mutual Chern-Simons theory Eq. (1) implies that the mFTI is actually a \mathbb{Z}_4 TO “enriched” by two $U(1)$ symmetries and time-reversal symmetry \mathcal{T} , i.e. it is a type of symmetry-enriched topological orders. Within

the \mathbb{Z}_4 TO, the “minimally-charged” anyon \mathbf{e} carries physical electric charge $e/2$, and the “minimal-spin” anyon \mathbf{m} carries a physical spin quantum number $S_z = \hbar/4$. Both of these anyons are self-bosons, but they see each other as a $\Phi_0/4$ flux, i.e. they have mutual statistical angle $\pi/2$. The \mathbf{em} bound state carries an electric charge and a spin quantum number $(e/2, \hbar/4)$, and it is an anyon with self-statistical angle $\pi/2$, i.e. its topological spin is $\theta_{\mathbf{em}} = e^{i\pi/2} = i$. This \mathbb{Z}_4 TO also preserves the time-reversal symmetry \mathcal{T} , which keeps the anyon \mathbf{e} invariant and maps the anyon \mathbf{m} to its anti-particle \mathbf{m}^3 . Curiously, \mathbf{m}^2 is a Kramers doublet under time reversal with $\mathcal{T}^2 = -1$. This result can be seen using both the edge theory in Sec. III and the general arguments of Ref. 14, which we will elaborate more in Sec. IV.

This \mathbb{Z}_4 TO have 16 anyons modulo the electrons (and holes), which can be represented as composite states $\mathbf{e}^p \mathbf{m}^q$ with $p, q = 0, \dots, 3$. However, since this \mathbb{Z}_4 TO is constructed with bosonic objects b^c and b^s , the background gapped electrons and holes were not yet taken into account. Hence, there exist another 16 anyons, which are bound states of $\mathbf{e}^p \mathbf{m}^q$ and the electron. Note that the bound state of $\mathbf{e}^p \mathbf{m}^q$ and the electron has a different topological spin from $\mathbf{e}^p \mathbf{m}^q$. Here, we’ve effectively included the electron as a “transparent” Abelian anyon (that braids trivially with all other anyons), which is a convenient technical choice for showing the minimality of this \mathbb{Z}_4 TO later. With the electrons and holes included, the \mathbb{Z}_4 topological order has in total 32 Abelian anyons, resulting in a squared total quantum dimension $\mathcal{D}^2 = 32$. In Sec. IV, we will prove that $\mathcal{D}^2 = 32$ is the minimal squared total quantum dimension compatible with the given Hall transports associated with the electric charge and spin quantum number S_z .

III. EXPERIMENTAL SIGNATURES

Following the standard derivation of the edge states from the bulk CS theory^{31,32}, we obtain the Lagrangian for the $1d$ edge based on the bulk CS theory Eq. 1

$$\mathcal{L}_{\text{edge}} = \frac{4}{4\pi} i \partial_\tau \phi^c \partial_x \phi^s - \frac{4}{4\pi} i \partial_\tau \phi^s \partial_x \phi^c + \dots, \quad (3)$$

where the “...” is the kinetic energy that will be written in a different basis below. $\partial_x \phi^c$ and $\partial_x \phi^s$ are respectively the density of the Cooper pair b^c and spin-1 exciton b^s at the boundary. We have the identification $e^{i4\phi^c} \sim b^c$ and $e^{i4\phi^s} \sim b^s$. Under the time-reversal symmetry \mathcal{T} , the Cooper pair b^c we introduced is invariant, while the spin-1 exciton operator b^s shares the same behavior as the lowering operator of a spin-1/2 object, i.e. $\mathcal{T} b^s \mathcal{T}^{-1} = -b^{s\dagger}$. Therefore, the time-reversal action on the edge theory is given by $\mathcal{T} : \phi^c \rightarrow -\phi^c, \phi^s \rightarrow \phi^s + \pi/4$.³³ The anyon \mathbf{m}^2 is identified with the field $e^{i2\phi_s}$, which implies that \mathbf{m}^2 is a Kramers doublet under time reversal with $\mathcal{T}^2 = -1$.

If the spin S_z conservation is broken but time-reversal symmetry is still preserved, an extra term $\cos(8\phi_s)$ is allowed in the edge theory Eq. 3. When this term is relevant, it will lead to two-fold degenerate ground states at the edge, which are characterized by nonzero expectation values of gauge invariant and time-reversal odd operator $\cos(4\phi_s)$, or $\sin(4\phi_s)$. Hence, the edge theory remains gapless unless there is either spontaneous or explicit time-reversal symmetry breaking (assuming charge is conserved). A different way to understand the stability of the gapless edge using only the bulk data will be given below.

By recombining ϕ^s and ϕ^c into $\phi_{1,2} = \phi^c \pm \phi^s$, we obtain a pair of counter-propagating (or helical) modes

$$\mathcal{L}_{\text{edge}} = \frac{2}{4\pi} i \partial_\tau \phi_1 \partial_x \phi_1 - \frac{2}{4\pi} i \partial_\tau \phi_2 \partial_x \phi_2 - V_{IJ} \partial_x \phi_I \partial_x \phi_J. \quad (4)$$

The density $\partial_x \phi_1$ (or $\partial_x \phi_2$) carries charge e and spin $\hbar/2$ (or $-\hbar/2$). We have added the kinetic energy with the velocity matrix V_{IJ} in Eq. (4) which arises from the non-topological part of the system. If V_{IJ} is proportional to Pauli matrix σ^z , the left and right counter-propagating modes do not interact with each other, and this edge theory would lead to a fractionally quantized $1d$ conductance per edge:

$$G = \frac{1}{2} \frac{e^2}{h}. \quad (5)$$

In a two-terminal measurement, the two-terminal conductance receives contributions from two edges connecting the two leads. At the total filling factor $\nu_{\text{tot}} = 3$, there are two completely filled bands with $C = \pm 1$ in addition to the mFTI state in the half-filled conjugate Chern bands. Therefore, the total two-terminal conductance should be $G = 3e^2/h$, which was observed experimentally in the twisted bilayer MoTe_2 at $\nu_{\text{tot}} = 3$.⁸

To give a full analysis of the $1d$ edge conductance with counter-propagating fractionalized modes, one needs careful analysis of the physics in the $1d$ channel as well as in the metallic lead and the contact^{34–36}. We expect the full analysis to be more involved than the edge state of the quantum spin-Hall insulator without fractionalization^{37,38}, and we defer the systematic study to the future.

Our mFTI is essentially a bosonic symmetry-enriched TO in the sense that although the system is made of correlated electrons/holes, a single electron/hole is always gapped in the bulk and the boundary. Hence, there is always a single particle gap in the system, despite the existence of the gapless charge modes at the edge, contrary to most fractional quantum Hall states. This effect is analogous to the bosonic SPT state proposed to be realized in the bilayer graphene³⁹.

The smallest electric charge carried by the anyons of the mFTI is $e^* = e/2$. The odd value of $\sigma^{\text{sh}}/e^* = 1$ in the mFTI guarantees the gaplessness of the edge whenever time-reversal symmetry and charge conservation are present (even without the S_z conservation).¹⁴ In contrast, using similar arguments as Ref. 14, one can show

that, given the spin-Hall conductivity $\sigma^{\text{sh}} = 1/2$ (and the vanishing “diagonal” Hall responses associated with the charge and spin quantum numbers), any TO that is a product of two decoupled TOs from the two valleys must have excitations with fractional electric charges equal to or smaller than $e/4$. Therefore, the mFTI and the product TOs from the two valleys can be experimentally distinguished by probing the electric charge fractionalization. Examples of such probes include the shot noise of quantum point contacts⁴⁰, local capacitance probe of localized charge states⁴¹, and Coulomb oscillations in antidots⁴².

IV. BOOTSTRAPPING THE MINIMAL FTI

In this section, we bootstrap the minimal topological order (TO) allowed by the Hall responses under the charge and the S_z -spin $U(1)$ symmetries and show that $\mathcal{D}^2 = 32$ is the minimal squared total quantum dimension compatible with a spin-Hall conductivity $\sigma^{\text{sh}} = 1/2$, zero electric Hall responses, and zero Hall response with respect to the spin S_z quantum number. The vanishing of the diagonal Hall responses associated with either the electric charge or S_z is required by the time-reversal symmetry.

For technical convenience, we prefer to re-organize the charge and spin $U(1)$ symmetries into $U(1)_{\uparrow} \times U(1)_{\downarrow}$, where $U(1)_{\uparrow, \downarrow}$ are associated with the charge conversation within each of the valleys 1 and 2 (which are locked to electron spins \uparrow and \downarrow). The Hall responses we focus on in this paper are equivalent to the combination of (1) $U(1)_{\uparrow}$ -Hall conductivity $\sigma_1^{\text{h}} = 1/2$, (2) $U(1)_{\downarrow}$ -Hall conductivity $\sigma_2^{\text{h}} = -1/2$, and (3) zero mixed $U(1)_{\uparrow} \times U(1)_{\downarrow}$ response. For an anyon x , we denote its fractional charge under $U(1)_{\uparrow} \times U(1)_{\downarrow}$ as $q_x = (q_{x,1}, q_{x,2})$, which is defined modulo the charge of local bosonic objects, i.e. $q_x \sim q_x + (n, m)$ for $n, m \in \mathbb{Z}$ with $m + n$ even. The fractional electric charge and S_z -spin of the anyon x is then given by $(q_{x,1} + q_{x,2})e$ and $(q_{x,1} - q_{x,2})\hbar/2$. In the TO, the electrons and holes are all identified as the transparent fermionic particle f with $q_f = (\pm 1, 0) \sim (0, \pm 1)$. Two such f particles fuse into a trivial anyon $\mathbf{1}$, i.e. $f \times f = \mathbf{1}$.

To bootstrap the minimal TO, we start by considering the anyons $v_{1,2}$ created by the adiabatic insertion of the 2π flux of $U(1)_{\uparrow, \downarrow}$. They must be Abelian anyons⁴³. Their fractional charges are given by the Hall conductivities $\sigma_{1,2}^{\text{h}}$: $q_{v_1} = (1/2, 0)$ and $q_{v_2} = (0, -1/2)$. Their topological spins are $\theta_{v_1} = e^{i\pi\sigma_1^{\text{h}}} = i$ and $\theta_{v_2} = e^{i\pi\sigma_2^{\text{h}}} = -i$, which are the consequences of the Aharonov-Bohm (AB) phase between charge and flux. Based on the AB effect, the mutual statistics between v_1 and v_2 must be trivial because v_1 (v_2) does not carry any fractional $U(1)_{\downarrow}$ -charge ($U(1)_{\uparrow}$ -charge). The anyons $v_{1,2}^2$ are associated with the 4π flux of $U(1)_{\uparrow, \downarrow}$. They are both self-bosons, i.e. $\theta_{v_1^2} = \theta_{v_1}^4 = 1$ and $\theta_{v_2^2} = \theta_{v_2}^4 = 1$. They have charges $q_{v_1^2} = (1, 0)$ and $q_{v_2^2} = (0, -1)$, which is non-trivial com-

pared to local bosonic objects. Hence, v_1^2 and v_2^2 are non-trivial anyons. Since $\theta_{v_1^2} = \theta_{v_2^2}$ and $q_{v_1^2} \sim q_{v_2^2}$, there are two scenarios: (1) v_1^2 and v_2^2 belongs to the same anyon type, and (2) v_1^2 and v_2^2 are different anyons.

For the second scenario with $v_1^2 \neq v_2^2$, there are at least 32 different anyons of the form $v_1^n v_2^m f^k$ with $n, m = 0, 1, 2, 3$ and $k = 0, 1$. They can be distinguished from each other based on their topological spins, fractional charges, and the assumption that $v_1^2 \neq v_2^2$. Moreover, based on the AB-phase, v_2^2 and v_1^2 braids trivially with any of the 32 anyons listed above. For a TO to be “complete”⁴⁴, there must be some extra anyons that braid non-trivially v_1^2 and v_2^2 . Consequently, the squared total quantum dimension must be larger than 32 in this scenario. Therefore, to search for the minimal TO, we only need to focus on the first scenario.

For the first scenario with $v_1^2 = v_2^2$, we show below that the minimal squared total quantum dimension is exactly $\mathcal{D}^2 = 32$. Let s denote the minimal (positive) integer s such that $v_1^{2s} = v_2^{2s}$ equals the trivial anyon $\mathbf{1}$. Given that $v_1^2 = v_2^2$ is non-trivial, $s \geq 2$. In the spirit of bootstrapping the minimal TO, we should focus on the case with $s = 2$. Similar arguments as below, when applied to the cases with $s \geq 3$, will lead to a squared total quantum dimension larger than 32.

From now on, we focus on the first scenario with $s = 2$. A list of 8 distinct Abelian anyons purely consists of $2\pi\mathbb{Z}$ fluxes of $U(1)_{\uparrow, \downarrow}$ is given by

$$V = \{\mathbf{1}, v_1, v_1^2, v_1^3, v_2, v_2^3, v_1 v_2, v_1^3 v_2\}. \quad (6)$$

By including the transparent fermion f , one finds 16 different Abelian anyons $\{\mathbf{1}, f\} \times V$. They can be distinguished by their topological spins and fractional charges. Note that v_1^2 braids trivially with all anyons in $\{\mathbf{1}, f\} \times V$. Therefore, there must be an extra anyon, called it y , that braids non-trivially with $v_1^2 = v_2^2$. However, it must braid trivially with $v_1^4 = v_2^4 = \mathbf{1}$. Using the AB-phase interpretation of the braiding with v_1^n and v_2^m , we conclude that the fractional charge of y must be $q_y = (1/4 + \mathbb{Z}, 1/4 + \mathbb{Z})$. Without loss of generality, we can pick

$$q_y = (1/4, 1/4). \quad (7)$$

For other choices of q_y , we can find another anyon with the fractional charge $(1/4, 1/4)$ by fusing y with one of the Abelian anyons in $\{\mathbf{1}, f\} \times V$. From the AB effect, we can obtain mutual statistics between y and $v_1^n v_2^m f^k$:

$$M_{y, v_1^n v_2^m f^k} = e^{i\frac{2\pi}{4}(n+m)}. \quad (8)$$

Therefore, the topological spins of the anyons $y v_1^n v_2^m f^k$ are all related to the topological spin of y via

$$\theta_{y v_1^n v_2^m f^k} = \theta_y \times (i)^{n^2} (-i)^{m^2} e^{i\frac{2\pi}{4}(n+m)} (-1)^k. \quad (9)$$

Using this result of the topological spins (and fractional charges), we can examine the set of anyons $\{\mathbf{1}, y\} \times \{\mathbf{1}, f\} \times V$. This set may have redundancy because y

and yv_1^2f share the same topological spin and fractional charge. As a result, there are two types, type-I and type-II, of minimal TOs.

For *type-I* minimal TOs, y and yv_1^2f are distinct. Consequently, there are 32 different anyons in

$$\mathcal{C}_I = \{1, y\} \times \{1, f\} \times V. \quad (10)$$

For minimality, \mathcal{C}_I should be the entire set of anyons, and each anyon should be Abelian, which yields a squared total quantum dimension $\mathcal{D}^2 = 32$. The \mathbb{Z}_4 FTI constructed in Sec. II provides an example of such type-I minimal TOs. v_1 , v_2 , and y correspond to the anyons \mathbf{em} and $\mathbf{e}^3\mathbf{m}$, \mathbf{e} respectively. Note that the \mathbb{Z}_4 FTI not only has the desired Hall responses but also preserves the time-reversal symmetry. Moreover, \mathbf{m}^2 , which corresponds to v_1v_2 , is the anyon generated by the insertion of a 2π flux in each valley. Its time-reversal conjugate should be associated with a -2π flux in each valley, which still belongs to the same anyon type \mathbf{m}^2 given that $v_1^2v_2^2 \sim 1$. Knowing that \mathbf{m}^2 is related to its time-reversal partner by a 4π flux in each valley and that the spin Hall conductivity is $\sigma^{\text{sh}} = 1/2$, \mathbf{m}^2 must be a Kramers doublet under time reversal¹⁴, which is consistent with the analysis in Sec. III. This statement have used the fact that the system is ultimately built out of the electron and holes that are Kramers doublets.

For *type-II* minimal TOs, y and yv_1^2f are identical. In other words, the fusion of y and v_1^2f , a self-fermion, yields y . Consequently, y is a non-Abelian anyon whose quantum dimension of y is at least $\sqrt{2}$. Now, there are 24 distinct anyons

$$\mathcal{C}_{II} = (\{1, f\} \times V) \cup (y \times V). \quad (11)$$

For minimality, \mathcal{C}_{II} should be the entire set of anyons, and the quantum dimension of each anyon in $y \times V$ should be exactly $\sqrt{2}$, which again yields a squared total quantum dimension of $\mathcal{D}^2 = 32$. Note that all the anyons in $y \times V$ share the same quantum dimension because of the Abelian nature of anyons in V . Such type-II minimal TOs must be non-Abelian. In the next section, we briefly discuss examples of such type-II minimal TO. It turns out that the examples we found all break the time-reversed symmetry. At the same time, they also require non-trivial couplings between the two valleys.

We remark that the \mathbb{Z}_4 FTI is a promising candidate for the mFTI not only because it has the minimal total quantum dimension compatible with the desired Hall responses. The \mathbb{Z}_4 FTI preserves time-reversal symmetry, while all other examples of minimal TOs we found turn out to break the time-reversal symmetry.

V. ALTERNATIVE CONSTRUCTIONS OF THE MFTI

A. Descending from larger topological orders




FIG. 2. The mFTI we propose is the common descendant of multiple product TOs with larger quantum dimensions. “Pf” stands for Pfaffian. It can also be viewed as a symmetry-enriched TO promoted from multiple bosonic SPTs by “gauging” the discrete subgroup of the symmetries.

In this section, we present other constructions of the same \mathbb{Z}_4 mFTI. The mFTI can be constructed through a “top-down” approach by descending from “larger” TOs, such as a product of two separate TOs in each valley. The simplest example of such TO, which can be realized at the $\nu = 1/2$ filling of both valleys, is the $U(1)_8 \times U(1)_{-8}$ state. The $U(1)_8$ TO realized in a half-filled Landau level can be viewed as a $\nu_b = 1/8$ bosonic Laughlin state for Cooper pairs²², described by a level-8 Chern-Simons theory for a dynamical $U(1)$ gauge field a , whose flux is dual to the density of the *intra-valley* Cooper pair. The anyons of the $U(1)_8 \times U(1)_{-8}$ state correspond to the gauge charges (n_1, n_2) of the two gauge fields, whose topological spins are given by $e^{i\pi(n_1^2 - n_2^2)/8}$. This anyon also carries electric charge $(n_1 - n_2)e/4$, and spin $S_z = (n_1 + n_2)\hbar/8$.

The anyon $(4, 4)$ is obviously a self-boson, hence it can condense. The anyon $(4, 4)$ can be created by adiabatically inserting 4π fluxes of $U(1)_\uparrow$ and $U(1)_\downarrow$ in both valleys. It carries spin $S_z = \hbar$. To avoid spontaneously breaking the spin S_z symmetry in the condensate, we bind the $(4, 4)$ anyon with the physical inter-valley exciton b^s with $S_z = -\hbar$ to neutralize its S_z spin. The mutual braiding statistical between the neutralized bosonic anyon $(4, 4)$ and a general anyon (n_1, n_2) is $e^{i\pi(n_1 - n_2)}$. Hence, the condensate of $(4, 4)$ would confine all the anyons whose n_1 and n_2 have opposite parity. The condensate also identifies anyons (n_1, n_2) and $(n_1 + 4, n_2 + 4)$. Consequently, there are, in total, 16 anyons left (before including the electrons/holes), the same as the mFTI. Using the notation of Sec. IV, the anyon v_1 and v_2 correspond to $(2, 0)$, and $(0, 2)$. The one of the most elementary deconfined anyons is $(1, 1)$, which is a self-boson with electric charge 0 and spin $S_z = \hbar/4$. Hence, $(1, 1)$ is identified with the anyon \mathbf{m} of the mFTI. Similarly, the anyon $(1, -1)$ can be identified as anyon \mathbf{e} of the mFTI. The anyons $(1, 1)$ and $(1, -1)$ also have the desired mutual braiding statistics $e^{i\pi/2}$. At this point, we can confirm that the condensate of the neutralized anyon $(4, 4)$ is the mFTI with \mathbb{Z}_4 TO. As an alternative approach, how the $U(1)_8 \times U(1)_{-8}$ TO descends, via the anyon conden-

sation of (4,4), to the desired mFTI can also be shown explicitly using the K -matrix formalism.

In fact, one can show that if we start with the other obvious product TOs, such as the Pfaffian \times Pfaffian, $331 \times \overline{331}$, anti-Pfaffian \times anti-Pfaffian, or PH-Pfaffian \times PH-Pfaffian, there is always a self-bosonic anyon created by inserting 4π fluxes in both valleys. Then, condensing this anyon after neutralizing its S_z spin will always lead to the same mFTI state.

As an aside, we can perform a similar “descending” construction starting from a product TO with *different* Abelian TOs in the two valleys, such as $331 \times U(1)_{-8}$. A similar anyon condensation in $331 \times U(1)_{-8}$ results in a type-I minimal TO that breaks time-reversal symmetry (manifested by its net chiral central charge $c = 1$). Hence, the so-constructed type-I minimal TO is not a candidate for mFTI even though they share the same Hall responses with respect to the electric charge and the S_z quantum number.

B. “Promoting” SPT states

As we remarked earlier, the mFTI is a symmetry-enriched topological (SET) order. The subject of SET has attracted enormous theoretical interest in the past decade^{45–48}. One very general way to construct nontrivial SET states is by “gauging” a part of the discrete symmetries of symmetry-protected topological (SPT) states^{49,50}, i.e. by “promoting” a part of the discrete global symmetries to local gauge invariance, and coupling the state to the gauge fields accordingly.

In our current case, the mFTI can be constructed through the same procedure. The SPT state we start with could be constructed by the inter-valley exciton operator b^s , and “one quarter” of the inter-valley Cooper pair b^c . In other words, we formally fractionalize b^c into four bosonic partons each carrying charge- $e/2$: $b^c = (\tilde{b}^c)^4$. Then, \tilde{b}^c must coupled to a Z_4 gauge field. Now, we construct a minimal level-1 bosonic SPT state between b^s and \tilde{b}^c ⁵¹, which can be described in various formulations, including the Chern-Simons theory⁵², or the nonlinear sigma model^{51,53}. The most convenient formulation for our current purpose is the Chern-Simons theory developed in Ref. 52. The Chern-Simons theory for many $(2+1)d$ bosonic SPT states takes a universal form:

$$\mathcal{L}_{\text{spt},1} = \frac{i}{2\pi} \tilde{a}^c da^s + \frac{i}{2\pi} A^s da^s + \frac{1}{2} \frac{i}{2\pi} A^e d\tilde{a}^c, \quad (12)$$

where \tilde{a}_μ^c and a_μ^s are the dual of the currents of \tilde{b}^c and b^s . Now, we need to gauge this SPT by coupling \tilde{b}^c to a Z_4 gauge field, which can be captured by adding the following terms

$$\mathcal{L}_{g,1} = \frac{i}{2\pi} c_1 d\tilde{a}^c + \frac{4i}{2\pi} c_1 dc_2. \quad (13)$$

The gauge fields $c_{1,\mu}$ and $c_{2,\mu}$ are introduced as auxiliary dynamical gauge fields that describe the Z_4 gauge

field through the last mutual Chern-Simons term. The coupling $\frac{i}{2\pi} c_1 d\tilde{a}^c$ is the minimal coupling between the Z_4 gauge field and the current of \tilde{b}^c . Combining the Lagrangians of Eq. (12) and Eq. (13), the Chern-Simons couplings of all the dynamical gauge fields $a_I = (\tilde{a}^c, a^s, c_1, c_2)$ can be organized into a K -matrix theory

$$\mathcal{L}_{\text{cs},1} = \frac{i}{4\pi} K_1^{IJ} a_I da_J, \quad K_1 = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 4 \\ 0 & 0 & 4 & 0 \end{pmatrix}. \quad (14)$$

The charge-1 particle of the gauge field c_1 is the minimal \mathbf{e} anyon of the Z_4 gauge field, it carries a 2π flux of \tilde{a}^c , which carries physical electric charge $e/2$, as expected. The charge-1 particle of gauge field c_2 carries $\pi/2$ flux of c_1 , which is equivalent to $\pi/2$ flux of a^s through the equation of motion of the gauge fields. The $\pi/2$ flux of a^s carries S_z -spin $\hbar/4$, hence the charge-1 of c_2 corresponds to the anyon \mathbf{m} of our mFTI.

To make a direct connection to the original CS theory Eq. (1), we note that the 2×2 block of the K -matrix associated with (\tilde{a}^c, a^s) has determinant -1 . Hence, \tilde{a}^c and a^s can be integrated out safely without compromising the nature of the TO. After integrating out (\tilde{a}^c, a^s) , and shifting $c_1 + \frac{1}{2} A^e \rightarrow c_1$, the theory Eq. 14 returns to the form of Eq. (1), except now (c_1, c_2) plays the role of (a^s, a^c) .

Another way of constructing the same mFTI state is by introducing the “half-partons” for both b^c and b^s : $b^c \sim (\tilde{b}^c)^2$, and $b^s \sim (\tilde{b}^s)^2$. Now, \tilde{b}^c and \tilde{b}^s carry electric charge e and quantum number $S_z = \hbar/2$, respectively. And, they are both coupled to their own Z_2 gauge fields. We can make \tilde{b}^c and \tilde{b}^s form a level-1 bosonic SPT state. The entire theory now reads

$$\begin{aligned} \mathcal{L}_{\text{cs},2} = & \frac{i}{2\pi} \tilde{a}^c d\tilde{a}^s + \frac{i}{2\pi} (A^e + c_1) d\tilde{a}^c + \frac{2i}{2\pi} c_1 dc_2 \\ & + \frac{i}{2\pi} \left(\frac{1}{2} A^s + c'_1\right) d\tilde{a}^s + \frac{2i}{2\pi} c'_1 dc'_2. \end{aligned} \quad (15)$$

Again, \tilde{a}^c and \tilde{a}^s are dual to the currents of \tilde{b}^c and \tilde{b}^s respectively, and they can be integrated out safely. After integrating out \tilde{a}^c and \tilde{a}^s , the $\mathcal{L}_{\text{cs},2}$ reduces to a CS theory for (c_1, c_2, c'_1, c'_2) with K -matrix

$$K_2 = \begin{pmatrix} 0 & 2 & 1 & 0 \\ 2 & 0 & 0 & 0 \\ 1 & 0 & 0 & 2 \\ 0 & 0 & 2 & 0 \end{pmatrix} \simeq K'_2 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 4 \\ 0 & 0 & 4 & 0 \end{pmatrix}. \quad (16)$$

One can show that the K_2 matrix is equivalent to the K'_2 -matrix up to an $\text{SL}(4, \mathbb{Z})$ transformation, and K'_2 again describes a Z_4 gauge field. The last two components of gauge fields of the K'_2 -matrix couple to the external gauge field A^e and A^s in the same way as a^c and a^s in Eq. (1).

C. Example of type-II minimal TO

The non-Abelian “type-II minimal TO” discussed in Sec. IV can be obtained as descendants of the product TOs (with larger total quantum dimensions). As an example, We start from the product TO Pfaffian $\times U(1)_{-8}$. The anyons of the Pfaffian TO is a subset of the topological order Ising $\times U(1)_8$. Here, “Ising” stands for the Ising TO, which has three anyons $\{\mathbb{1}, \sigma, \psi\}$ with $\mathbb{1}$ the trivial anyon, σ the Ising anyon with quantum dimension $\sqrt{2}$, and ψ the self fermion. Their topological spins are $\theta_{\mathbb{1}} = 1$, $\theta_{\sigma} = e^{i2\pi/16}$, and $\theta_{\psi} = -1$ respectively. For the $U(1)_8 \times U(1)_{-8}$ sector of Pfaffian $\times U(1)_{-8}$, we can still use the charge (n_1, n_2) of the gauge fields of the $U(1)_8 \times U(1)_{-8}$ Chern-Simons theory as a part of the anyon label. The contribution to the topological spins from the gauge charge (n_1, n_2) is $\theta_{(n_1, n_2)} = e^{i2\pi(n_1^2 - n_2^2)/16}$. The full set of anyons of Pfaffian $\times U(1)_{-8}$ is given by

$$\begin{aligned} (\mathbb{1}, n_1, n_2) & \text{ with } n_1 = 0, 2, 4, 6 \text{ and } n_2 = 0, 1, \dots, 7, \\ (\psi, n_1, n_2) & \text{ with } n_1 = 0, 2, 4, 6 \text{ and } n_2 = 0, 1, \dots, 7, \\ (\sigma, n_1, n_2) & \text{ with } n_1 = 1, 3, 5, 7 \text{ and } n_2 = 0, 1, \dots, 7, \end{aligned} \quad (17)$$

where $(\psi, 4, 0)$ is identified as the transparent fermion f , i.e. the electron or the hole. The electric charge and S_z quantum numbers of these anyons are given by $(n_1 - n_2)e/4$ and $(n_1 + n_2)\hbar/8$. Note that $(\mathbb{1}, 4, 4)$ is self-boson without electric charge. Its S_z quantum number can be neutralized by adding a local boson, i.e. an intervalley exciton.

By condensing the neutralized version of $(\mathbb{1}, 4, 4)$ in Pfaffian $\times U(1)_{-8}$, we obtain a type-II minimal TO with 24 deconfined anyons. Using the notation of Sec. IV, the 16 Abelian anyons in $\{1, f\} \times V$ are given by

$$(\mathbb{1}, n_1, n_2) \text{ and } (\psi, n_1, n_2) \quad \text{with } n_{1,2} = 0, 2, 4, 6 \quad (18)$$

with the identification $(\mathbb{1}/\psi, n_1, n_2) \sim (\mathbb{1}/\psi, n_1 + 4, n_2 + 4)$.⁵⁴ In particular, we can identify $v_1 = (\mathbb{1}, 2, 0)$ and $v_2 = (\mathbb{1}, 0, 2)$. The 8 non-Abelian quantum-dimension- $\sqrt{2}$ anyons are given by

$$(\sigma, n_1, n_2) \quad \text{with } n_{1,2} = 1, 3, 5, 7 \quad (19)$$

with the identification $(\sigma, n_1, n_2) \sim (\sigma, n_1 + 4, n_2 + 4)$. In particular, we have $y = (\sigma, 1, 7)$.

This non-Abelian TO, described by the condensate of $(\mathbb{1}, 4, 4)$, is one of the type-II minimal TO that can be

realized in a pair of half-filled conjugate Chern bands. It has the same Hall responses with respect to the electric charge and the S_z quantum number as the mFTI. However, this state breaks the time-reversal symmetry because it has the same chiral central charge $c = 1/2$ as the original Pfaffian $\times U(1)_{-8}$ TO. Consequently, this type-II minimal TO also has a quantized thermal-Hall response.

More examples of type-II minimal TOs can be constructed as the descendants of other combinations of a non-Abelian TO (which could be the Pfaffian TO, the anti-Pfaffian TO, the PH-Pfaffian TO, and their conjugates) in one valley and an Abelian TO (which could be $U(1)_8$, the 331 state, and their conjugates) in the other valley. The so-constructed type-II minimal TOs all break time-reversal symmetry.

VI. SUMMARY

Motivated by a recent experiment⁸, in this work, we proposed a minimal fractional topological insulator that can potentially be realized in a pair of conjugate Chern bands, both with electron filling $\nu = 1/2$. The mFTI we proposed has the minimal total quantum dimension amongst all the topological orders that can be potentially realized in the same setting (especially with the same experimentally observed fractionally quantized transport signal). We caution that a more thorough analysis, similar to that for the $\nu = 2/3$ fractional quantum state^{34–36}, is required to understand the transport quantization. We found that the minimally-charged anyon in mFTI has fractional charge $e^* = e/2$, which is different from all product TOs. The gapless edge state of this mFTI is protected by charge conservation and time-reversal symmetry even when S_z is not conserved. We also showed that the mFTI is the common descendant of various product TOs of the two valleys, which have larger quantum dimensions. The mFTI is also the common symmetry-enriched TO that can be promoted from different bosonic SPT states by gauging the discrete subgroup of their symmetries.

The authors thank Leon Balents and Matthew Fisher for very helpful discussions. C.X. is supported by the Simons foundation through the Simons investigator program. C.-M.J. is supported by a faculty startup grant at Cornell University. While finishing this paper, we became aware of another independent work⁵⁵ studying the edge state of FTIs that are a product of conjugate TOs from two valleys, which, as we discussed, should have a larger quantum dimension than the mFTI constructed in this paper.

¹ E. M. Spanton, A. A. Zibrov, H. Zhou, T. Taniguchi, K. Watanabe, M. P. Zaletel, and A. F. Young, Science

360, 62 (2018).

² Y. Xie, A. T. Pierce, J. M. Park, D. E. Parker, E. Khalaf,

- P. Ledwith, Y. Cao, S. H. Lee, S. Chen, P. R. Forrester, K. Watanabe, T. Taniguchi, A. Vishwanath, P. Jarillo-Herrero, and A. Yacoby, *Nature (London)* **600**, 439 (2021), arXiv:2107.10854 [cond-mat.mes-hall].
- ³ J. Cai, E. Anderson, C. Wang, X. Zhang, X. Liu, W. Holtzmann, Y. Zhang, F. Fan, T. Taniguchi, K. Watanabe, Y. Ran, T. Cao, L. Fu, D. Xiao, W. Yao, and X. Xu, *Nature* **622**, 63 (2023).
- ⁴ H. Park, J. Cai, E. Anderson, Y. Zhang, J. Zhu, X. Liu, C. Wang, W. Holtzmann, C. Hu, Z. Liu, T. Taniguchi, K. Watanabe, J.-H. Chu, T. Cao, L. Fu, W. Yao, C.-Z. Chang, D. Cobden, D. Xiao, and X. Xu, *Nature* **622**, 74 (2023).
- ⁵ Y. Zeng, Z. Xia, K. Kang, J. Zhu, P. Knapp, C. Vaswani, K. Watanabe, T. Taniguchi, K. F. Mak, and J. Shan, “Integer and fractional chern insulators in twisted bilayer mote2,” (2023), arXiv:2305.00973 [cond-mat.mes-hall].
- ⁶ F. Xu, Z. Sun, T. Jia, C. Liu, C. Xu, C. Li, Y. Gu, K. Watanabe, T. Taniguchi, B. Tong, *et al.*, arXiv preprint arXiv:2308.06177 (2023).
- ⁷ Z. Lu, T. Han, Y. Yao, A. P. Reddy, J. Yang, J. Seo, K. Watanabe, T. Taniguchi, L. Fu, and L. Ju, “Fractional quantum anomalous hall effect in a graphene moire superlattice,” (2023), arXiv:2309.17436 [cond-mat.mes-hall].
- ⁸ K. Kang, B. Shen, Y. Qiu, K. Watanabe, T. Taniguchi, J. Shan, and K. F. Mak, “Observation of the fractional quantum spin hall effect in moiré mote2,” (2024), arXiv:2402.03294 [cond-mat.mes-hall].
- ⁹ F. Wu, T. Lovorn, E. Tutuc, I. Martin, and A. H. MacDonald, *Phys. Rev. Lett.* **122**, 086402 (2019).
- ¹⁰ M. König, S. Wiedmann, C. Brüne, A. Roth, H. Buhmann, L. W. Molenkamp, X.-L. Qi, and S.-C. Zhang, *Science* **318**, 766 (2007), <https://www.science.org/doi/pdf/10.1126/science.1148047>.
- ¹¹ N. H. Lindner, E. Berg, G. Refael, and A. Stern, *Phys. Rev. X* **2**, 041002 (2012).
- ¹² M. Cheng, *Phys. Rev. B* **86**, 195126 (2012).
- ¹³ A. Stern, *Annual Review of Condensed Matter Physics* **7**, 349–368 (2016).
- ¹⁴ M. Levin and A. Stern, *Phys. Rev. Lett.* **103**, 196803 (2009).
- ¹⁵ N. Morales-Durán, N. Wei, and A. H. MacDonald, “Magic angles and fractional chern insulators in twisted homobilayer tmds,” (2023), arXiv:2308.03143 [cond-mat.str-el].
- ¹⁶ N. Paul, Y. Zhang, and L. Fu, *Science Advances* **9** (2023), 10.1126/sciadv.abn1401.
- ¹⁷ G. Moore and N. Read, *Nuclear Physics B* **360**, 362 (1991).
- ¹⁸ M. Greiter, X.-G. Wen, and F. Wilczek, *Phys. Rev. Lett.* **66**, 3205 (1991).
- ¹⁹ X.-G. Wen, *Phys. Rev. Lett.* **70**, 355 (1993).
- ²⁰ M. Levin, B. I. Halperin, and B. Rosenow, *Physical Review Letters* **99** (2007), 10.1103/physrevlett.99.236806.
- ²¹ D. T. Son, *Phys. Rev. X* **5**, 031027 (2015).
- ²² M. Greiter, X. Wen, and F. Wilczek, *Nuclear Physics B* **374**, 567 (1992).
- ²³ X. G. Wen and A. Zee, *Phys. Rev. B* **46**, 2290 (1992).
- ²⁴ B. J. Overbosch and X.-G. Wen, arXiv e-prints, arXiv:0804.2087 (2008), arXiv:0804.2087 [cond-mat.mes-hall].
- ²⁵ C. L. Kane and E. J. Mele, *Phys. Rev. Lett.* **95**, 226801 (2005).
- ²⁶ B. A. Bernevig and S.-C. Zhang, *Phys. Rev. Lett.* **96**, 106802 (2006).
- ²⁷ Y.-H. Zhang, “Composite fermion insulator in opposite-fields quantum hall bilayers,” (2018), arXiv:1810.03600 [cond-mat.str-el].
- ²⁸ N. Myerson-Jain, C.-M. Jian, and C. Xu, “The conjugate composite fermi liquid,” (2023), arXiv:2311.16250 [cond-mat.str-el].
- ²⁹ Y.-H. Zhang, “Vortex spin liquid with neutral fermi surface and fractional quantum spin hall effect at odd integer filling of moiré chern band,” (2024), arXiv:2402.05112 [cond-mat.str-el].
- ³⁰ Z. D. Shi, H. Goldman, Z. Dong, and T. Senthil, “Excitonic quantum criticality: from bilayer graphene to narrow chern bands,” (2024), arXiv:2402.12436 [cond-mat.str-el].
- ³¹ X. G. Wen, *Phys. Rev. Lett.* **64**, 2206 (1990).
- ³² X. G. Wen, *Phys. Rev. B* **43**, 11025 (1991).
- ³³ In principle, one can consider a more general time-reversal action with $\phi^c \rightarrow -\phi^c + \alpha_0$ for some finite constant α_0 . However, this constant α_0 can be absorbed by redefining the time-reversal symmetry as the original one followed by an extra charge U(1) rotation. The redefined time-reversal action remains to be a symmetry action of order 2. We caution that one cannot redefine the time-reversal symmetry by combining it with arbitrary spin S_z rotations because the resulting action is generically not an order-2 action anymore.
- ³⁴ C. L. Kane, M. P. A. Fisher, and J. Polchinski, *Phys. Rev. Lett.* **72**, 4129 (1994).
- ³⁵ C. L. Kane and M. P. A. Fisher, *Phys. Rev. B* **51**, 13449 (1995).
- ³⁶ C. L. Kane and M. P. A. Fisher, *Phys. Rev. B* **52**, 17393 (1995).
- ³⁷ C. Xu and J. E. Moore, *Phys. Rev. B* **73**, 045322 (2006).
- ³⁸ C. Wu, B. A. Bernevig, and S.-C. Zhang, *Phys. Rev. Lett.* **96**, 106401 (2006).
- ³⁹ Z. Bi, R. Zhang, Y.-Z. You, A. Young, L. Balents, C.-X. Liu, and C. Xu, *Phys. Rev. Lett.* **118**, 126801 (2017).
- ⁴⁰ L. Saminadayar, D. C. Glatli, Y. Jin, and B. Etienne, *Phys. Rev. Lett.* **79**, 2526 (1997).
- ⁴¹ J. Martin, S. Ilani, B. Verdene, J. Smet, V. Umansky, D. Mahalu, D. Schuh, G. Abstreiter, and A. Yacoby, *Science* **305**, 980 (2004), <https://www.science.org/doi/pdf/10.1126/science.1099950>.
- ⁴² A. Kou, C. M. Marcus, L. N. Pfeiffer, and K. W. West, *Phys. Rev. Lett.* **108**, 256803 (2012).
- ⁴³ M. Cheng, M. Zaletel, M. Barkeshli, A. Vishwanath, and P. Bonderson, *Phys. Rev. X* **6**, 041068 (2016).
- ⁴⁴ A. Kitaev, *Annals of Physics* **321**, 2 (2006), arXiv:cond-mat/0506438 [cond-mat.mes-hall].
- ⁴⁵ A. M. Essin and M. Hermele, *Phys. Rev. B* **87**, 104406 (2013).
- ⁴⁶ A. Mesaros and Y. Ran, *Phys. Rev. B* **87**, 155115 (2013).
- ⁴⁷ L.-Y. Hung and X.-G. Wen, *Phys. Rev. B* **87**, 165107 (2013).
- ⁴⁸ Y.-M. Lu and A. Vishwanath, *Phys. Rev. B* **93**, 155121 (2016).
- ⁴⁹ X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, *Phys. Rev. B* **87**, 155114 (2013).
- ⁵⁰ X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, *Science* **338**, 1604 (2012).
- ⁵¹ T. Senthil and M. Levin, *Phys. Rev. Lett.* **110**, 046801 (2013).
- ⁵² Y.-M. Lu and A. Vishwanath, *Phys. Rev. B* **86**, 125119 (2012).
- ⁵³ C. Xu and T. Senthil, *Phys. Rev. B* **87**, 174412 (2013).

⁵⁴ $n_{1,2}$ should be viewed as integers mod 8.

⁵⁵ J. May-Mann, A. Stern, and T. Devakul, “Theory of

half-integer fractional quantum spin hall insulator edges,” (2024), arXiv:2403.03964 [cond-mat.mes-hall].