From the Fokker-Planck equation to a contact Hamiltonian system

Shin-itiro GOTO

Center for Mathematical Science and Artificial Intelligence, Chubu University, 1200 Matsumoto-cho, Kasugai, Aichi 487-8501, Japan

March 13, 2024

Abstract

The Fokker-Planck equation is one of the fundamental equations in nonequilibrium statistical mechanics, and this equation is known to be derived from the Wasserstein gradient flow equation with a free energy. This gradient flow equation describes relaxation processes and is formulated on Riemannian manifolds. Meanwhile contact Hamiltonian systems are also known to describe relaxation processes. Hence a relation between these two equations is expected to be clarified, which gives a solid foundation in geometric statistical mechanics. In this paper a class of contact Hamiltonian systems is derived from a class of the Fokker-Planck equations on Riemannian manifolds. In the course of the derivation, the Fokker-Planck equation is shown to be written as a diffusion equation with a weighted Laplacian without any approximation, which enables to employ a theory of eigenvalue problems.

Contents

1	Introduction	1
2	From the Fokker-Planck equation to a diffusion equation2.1The Fokker-Planck equation as a Wasserstein gradient flow2.2Diffusion equation with a weighted Laplacian2.3Spectrum associated with the Fokker-Planck equation	3 3 5 10
3	From the diffusion equation to a contact Hamiltonian system3.1Contact geometry3.2Derivation of a contact Hamiltonian system3.3Dynamics depending on externally applied fields3.4Example	 13 14 21 24
4	Discussions and Conclusion	27
A	Appendix A.1 A simplified proof for Theorem 2.2 A.2 Various weighted Laplacians A.3 An expression of the Fokker-Planck equation A.4 Several choices of z	 29 29 29 31 33

1 Introduction

Nonequilibrium statistical mechanics is a developing branch of physics, and studies relations between microscopic dynamics and macroscopic nonequilibrium thermodynamic phenomena [26]. In theoretical physics, considerable activity is being devoted to establish a reliable theory, where such a theory is expected to provide mathematical methodologies for analyzing nonequilibrium phenomena. To this end various fundamental equations have been proposed and analyzed, where such fundamental equations are the Fokker-Planck equation, master equations, and so on [44]. Their applications include not only to physical sciences, but also to mathematical engineering [36, 33]. In addition, by applying these equations to toy models, explicit mathematical expressions of thermodynamic quantities have been obtained, where such toy models include the Glauber equation [12, 6], Brownian motion [23], and gas systems [44]. Recalling that some of these developments lie in the applications of well-established mathematics, such as stochastic partial differential equations [35], dynamical systems theory [38], and geometry [8, 11], one expects that further developments of this research field will lie in the application of advanced mathematics.

Among various mathematical formulations that have been proposed so far, one class relies on geometric methods. Examples include Wasserstein geometry [43] and contact geometry [31], where Wasserstein geometry studies Wasserstein spaces for probability density functions [42], and contact geometry is an odd-dimensional cousin of symplectic geometry [4]. Contact geometry has been applied to thermodynamics and statistical mechanics [32, 2, 40, 21, 19, 20], since it is a differential geometry endowed with an appropriate 1-form. This 1-form describes the laws of thermodynamics. As mentioned above, since there are several theories, any comparison among them is needed to establish a solid foundation of geometric nonequilibrium statistical mechanics. As a first step, a Wasserstein gradient flow was compared with a contact Hamiltonian one for a particular class of models [7], where relaxation was investigated. Since relaxation process is a simple nonequilibrium phenomenon and this links nonequilibrium states and equilibrium states, the clarification of how to express relaxation processes should be a meaningful target to advance our understanding of nonequilibrium systems. Beyond this pioneering study, more general approaches should be explored. One of the advantages of involving Wasserstein geometry is that techniques developed in Riemannian geometry can be employed, where Riemannian geometry has been applied to various physical problems [1]. This advantage is expected to be used efficiently in future explorations.

In this paper a class of contact Hamiltonian equation as ODEs (Ordinary Differential Equations) is derived from the Fokker-Planck equation by means of Riemannian geometry, where this Fokker-Planck equation is a PDE (Partial Differential Equation). This PDE is the Wasserstein gradient flow of a free energy. Here a set of macroscopic thermodynamic variables is represented as a point of a contact manifold, where these variables are defined by integrating some functions with the weight of a solution of the Fokker-Planck equation. The most principal step for deriving a contact Hamiltonian system is to put fast time-scales appeared in a solution of the Wasserstein gradient equation zero. Then main theorems in this paper are informally summarized as follows:

Claim 1.1. (Informal version of Theorem 2.2). A diffusion equation with a weighted Laplacian is derived from the Fokker-Planck equation without any approximation.

Claim 1.2. (Informal version of Theorem 3.8). The Fokker-Planck equation is equivalent to a contact Hamiltonian system when fast time-scales in the diffusion equation are neglected.

In addressing Claim 1.1, a weighted Laplacian is introduced in this paper. It should be emphasized that there are several weighted Laplacians in the literature, and they are sometimes called the Witten Laplacians. On the one hand, as summarized in [22], although one existing Witten Laplacian is closely related to the Fokker-Planck equation, there is no exact one-to-one correspondence between a heat equation and the Fokker-Planck equation, where heat equation is a synonym of diffusion equation. On the other hand, the weighted Laplacian in this paper is shown to be equivalent to another existing self-adjoint operator acting on a function [41] (see Section A.3 of this paper). This existing operator is shown to be a Witten Laplacian, and yields another form of the Fokker-Planck equation.

The rest of this paper is organized as follows. In Section 2, how the Fokker-Planck equation is derived from the Wasserstein gradient flow is summarized. Then a diffusion equation with a weighted Laplacian is derived without any approximation. Since this diffusion equation exactly corresponds to the Fokker-Planck equation, several properties of the introduced Laplacian are clarified here and employed for later sections. In Section 3, after contact geometry is briefly summarized, the contact Hamiltonian flow is derived with the use of the information about a spectrum of the weighted Laplacian. In addition, an explicit example of how the theory is applied is shown. Finally, Section 4 summarizes the present study, and possible future directions are presented.

2 From the Fokker-Planck equation to a diffusion equation

As shown in [25], the Fokker-Planck equation can be derived from a variational principle with a free-energy functional in the context of Wasserstein geometry.

In this section after recalling how to derive the Fokker-Planck equation, it is shown that this equation induces a diffusion equation with a weighted Laplacian.

2.1 The Fokker-Planck equation as a Wasserstein gradient flow

In this subsection necessary notations in Riemannian geometry are fixed, and then how the Fokker-Planck equation is derived is summarized.

Let (\mathcal{M}, q) be an *m*-dimensional Riemannian manifold with q being a Riemannian metric tensor field, and $\star 1$ a canonical volume-form on \mathcal{M} . The boundary of \mathcal{M} is denoted by $\partial \mathcal{M}$. An interpretation of \mathcal{M} in this paper is phase space of a microscopic dynamical system. An example of \mathcal{M} is a symplectic manifold, and q is a compatible metric with respect to an endowed symplectic form on \mathcal{M} . In the discussion below the manifold need not be symplectic. The spaces of vector fields and k-forms on \mathcal{M} are denoted by $\Gamma T \mathcal{M}$ and $\Gamma \Lambda^k \mathcal{M}$, respectively. Since 0-forms are identified with functions, $\Gamma \Lambda^0 \mathcal{M}$ denotes the space of functions on \mathcal{M} . The Hodge map is introduced and denoted by $\star : \Gamma \Lambda^k \mathcal{M} \to \Gamma \Lambda^{m-k} \mathcal{M}, (k = 0, 1, \dots, m)$. Given \star , its inverse is denoted by \star^{-1} . Let Id be the identical operator: $\Gamma \Lambda^k \mathcal{M} \to \Gamma \Lambda^k \mathcal{M}, (k = 0, 1, ..., m)$. Vector fields in a patch on \mathcal{M} can explicitly be written in coordinates, and coordinate expressions give systems of ordinary differential equations (ODEs). Since ODEs describe continuous-time dynamical systems, vector fields are often identified with dynamical systems in this paper. Then $d: \Gamma \Lambda^k \mathcal{M} \to \Gamma \Lambda^{k+1} \mathcal{M}, (k = 0, 1, ..., m)$ is the exterior derivative and i_X is the interior product with respect to $X \in \Gamma T \mathcal{M}$. After defining an inner product for forms, the so-called co-derivative d^{\dagger} , is introduced as in the standard way. This d^{\dagger} is often expressed in the literature as δ [5], but δ is not used in this paper. To clearly distinguish nationally the adjoint of d and the Hodge map \star , d^{*} is not used in this paper, where d^{*} is often employed to denote the adjoint of d. Some of the details of d^{\dagger} are summarized in this section so that a cousin d^{\dagger}_{G} is analogously introduced in this paper. Throughout the paper integrals over \mathcal{M} are assumed to commute with $\partial/\partial t$, where $t \in \mathbb{R}$ denotes time.

To describe the Fokker-Planck equation on \mathcal{M} , let h be a function on \mathcal{M} called a microscopic Hamiltonian, β denote the inverse temperature with the Boltzmann constant $k_{\rm B}$ being unity. Throughout this paper $\beta > 0$ is constant in time and over \mathcal{M} . The space of other externally applied fields \mathbb{R}^n will be introduced in later sections. It is known that the Fokker-Planck equation can be derived as a Wasserstein gradient flow associated with a free energy on the Euclidean spaces [25]. Some extension to Riemannian manifolds can be established [29]. To recall this derivation and to fix notations, in the following, the Fokker-Planck equation on \mathcal{M} for $\rho_t \in \Gamma \Lambda^0 \mathcal{M}$, $(\forall t \in \mathbb{R})$

$$\frac{\partial}{\partial t}\rho_t = -\frac{1}{\beta} \,\mathrm{d}^\dagger \,\mathrm{d}\,\rho_t - \mathrm{d}^\dagger\,\rho_t \,\mathrm{d}\,h,\tag{1}$$

is derived as a Wasserstein gradient flow. The ρ_t is interpreted as a time-dependent distribution function. Hence ρ_t is demanded to be normalized:

$$\int_{\mathcal{M}} \rho_t \star 1 = 1 \qquad \forall t \in \mathbb{R}.$$
(2)

For a simple case where $\mathcal{M} = \mathbb{R}$ and $g = dx \otimes dx$ with $x \in \mathbb{R}$, a coordinate expression of (1) will be given in (39). To derive (1), preliminary formulae are shown below. Let d^{\dagger} be the co-derivative that is

$$d^{\dagger}: \Gamma \Lambda^{k} \mathcal{M} \to \Gamma \Lambda^{m-1} \mathcal{M}, \qquad k = 0, 1, \dots, m$$
$$\alpha \quad \mapsto \quad d^{\dagger} \alpha,$$

with [5]

$$d^{\dagger} \alpha := (-1)^k \star^{-1} d \star \alpha, \qquad k = 0, 1, \dots, m.$$

From this definition it follows that $d^{\dagger} \Phi = 0$ for any function Φ . To see that d^{\dagger} is an adjoint operator of d, introduce the inner-product

$$\langle , \rangle : \Gamma \Lambda^k \mathcal{M} \times \Gamma \Lambda^k \mathcal{M} \quad \to \quad \mathbb{R}, \qquad k = 0, 1, \dots, m$$
$$(\alpha^{\mathrm{I}}, \alpha^{\mathrm{II}}) \quad \mapsto \quad \langle \alpha^{\mathrm{I}}, \alpha^{\mathrm{II}} \rangle,$$

with

$$\langle \alpha^{\mathrm{I}}, \alpha^{\mathrm{II}} \rangle := \int_{\mathcal{M}} \alpha^{\mathrm{I}} \wedge \star \alpha^{\mathrm{II}}.$$
 (3)

For instance, for any $\alpha \in \Gamma \Lambda^1 \mathcal{M}$ and $\Phi \in \Gamma \Lambda^0 \mathcal{M}$ with boundary values on $\partial \mathcal{M}$, it follows that

$$\left\langle d^{\dagger} \alpha, \Phi \right\rangle = \int_{\mathcal{M}} (d^{\dagger} \alpha) \Phi \star 1 = -\int_{\mathcal{M}} (\star^{-1} d \star \alpha) \Phi \star 1 = -\int_{\mathcal{M}} \Phi(d \star \alpha)$$
$$= -\int_{\mathcal{M}} d(\Phi \star \alpha) + \int_{\mathcal{M}} d\Phi \wedge \star d\alpha = -[\Phi \star \alpha]_{\partial \mathcal{M}} + \langle d\Phi, \alpha \rangle$$
$$= \langle \alpha, d\Phi \rangle,$$
(4)

where the boundary term on $\partial \mathcal{M}$ has been assumed to vanish. The operator div is defined as

$$\operatorname{div} := -\operatorname{d}^{\dagger}.$$

In addition the corresponding potential $\phi_{\dot{\rho}_t} \in \Gamma \Lambda^0 \mathcal{M}$ is introduced such that the continuity equation holds [1]:

$$\dot{\rho}_t = -\operatorname{div}(\rho_t \,\mathrm{d}\,\phi_{\dot{\rho}_t}) = \mathrm{d}^\dagger \,\rho_t \,\mathrm{d}\,\phi_{\dot{\rho}_t}, \quad \forall \ t \in \mathbb{R},$$
(5)

where the abbreviation has been introduced:

$$\dot{\rho}_t := \frac{\partial}{\partial t} \rho_t.$$

Using these introduced operators and integrals, one can derive (1) as follows. Recall that \mathcal{M} is phase space of a microscopic dynamical system, and that g is given on \mathcal{M} as a Riemannian metric. In addition, $h \in \Gamma \Lambda^0 \mathcal{M}$ has been given as a Hamiltonian. Let the free energy F be written as

$$F = \beta^{-1}S - H,$$

$$S = -\langle \rho_t, \ln \rho_t \rangle = -\int_{\mathcal{M}} \rho_t \ln \rho_t \star 1,$$

$$H = \langle h, \rho_t \rangle = \int_{\mathcal{M}} h\rho_t \star 1.$$
(6)

Then from (5), (4), and

$$\int_{\mathcal{M}} \dot{\rho}_t \star 1 = 0,$$

it follows that

$$\dot{S} = -\int_{\mathcal{M}} \dot{\rho}_t (\ln \rho_t + 1) \star 1 = -\int_{\mathcal{M}} \dot{\rho}_t \ln \rho_t \star 1 = -\left\langle \operatorname{d}^{\dagger} \rho_t \operatorname{d} \phi_{\dot{\rho}_t} , \ln \rho_t \right\rangle$$
$$= -\left\langle \rho_t \operatorname{d} \phi_{\dot{\rho}_t} , \operatorname{d} \ln \rho_t \right\rangle = -\left\langle \operatorname{d} \phi_{\dot{\rho}_t} , \rho_t \operatorname{d} \ln \rho_t \right\rangle = -\left\langle \phi_{\dot{\rho}_t} , \operatorname{d}^{\dagger} \rho_t \operatorname{d} \ln \rho_t \right\rangle.$$

To reduce this equation further, by employing

$$\rho_t \operatorname{d} \ln \rho_t = \rho_t \frac{1}{\rho_t} \operatorname{d} \rho_t = \operatorname{d} \rho_t,$$

one has that

$$\dot{S} = -\left\langle \phi_{\dot{\rho}_t} , \, \mathrm{d}^{\dagger} \, \mathrm{d} \, \rho_t \right\rangle = -\left\langle \, \mathrm{d}^{\dagger} \, \mathrm{d} \, \rho_t , \, \phi_{\dot{\rho}_t} \right\rangle.$$

In addition, simple calculations yield

$$\dot{H} = \int_{\mathcal{M}} h\dot{\rho}_t \star 1 = \int_{\mathcal{M}} h(\mathrm{d}^{\dagger} \rho_t \,\mathrm{d} \,\phi_{\dot{\rho}_t}) \star 1 = \left\langle h, \,\mathrm{d}^{\dagger} \rho_t \,\mathrm{d} \,\phi_{\dot{\rho}_t} \right\rangle$$
$$= \left\langle \rho_t \,\mathrm{d} \,h, \,\mathrm{d} \,\phi_{\dot{\rho}_t} \right\rangle = \left\langle \,\mathrm{d}^{\dagger} \,\rho_t \,\mathrm{d} \,h, \,\phi_{\dot{\rho}_t} \right\rangle.$$

By defining $\operatorname{grad}_W F \in \Gamma \Lambda^0 \mathcal{M}$ such that

$$\dot{F} = \langle \operatorname{grad}_W F, \phi_{\dot{\rho}_t} \rangle,$$

one identifies

$$\operatorname{grad}_W F = -\beta^{-1} \operatorname{d}^{\dagger} \operatorname{d} \rho_t - \operatorname{d}^{\dagger} \rho_t \operatorname{d} h.$$

Thus the equation called the gradient flow equation,

$$\frac{\partial}{\partial t}\rho_t = \operatorname{grad}_W F,$$

is the Fokker-Planck equation, (1). See [7] for the case of discrete phase space.

For a stationary solution $\rho_{\rm G}$ that satisfies $\dot{\rho}_{\rm G} = 0$ for (1), introduce the partition function $Z_{\rm G}$ such that

$$\int_{\mathcal{M}} \rho_{\rm G} \star 1 = 1. \tag{7}$$

The stationary solution satisfies

$$\mathrm{d}^{\dagger} \left(\beta^{-1} \,\mathrm{d} \,\rho_{\mathrm{G}} + \rho_{\mathrm{G}} \,\mathrm{d} \,h \right) = 0,$$

and it is the Gibbs distribution function:

$$\rho_{\rm G} = Z_{\rm G}^{-1} \exp(-\beta h),\tag{8}$$

where the stationary property can be verified from (8) with

$$\beta^{-1} d \rho_{\rm G} = -Z_{\rm G}^{-1} \exp(-\beta h) d h = -\rho_{\rm G} d h.$$
(9)

2.2 Diffusion equation with a weighted Laplacian

In this subsection a diffusion equation with a weighted Laplacian is derived from the Fokker-Planck equation, where no approximation is made. The main steps for this derivation are to employ a change of variables and to define an appropriate weighted Laplacian. After deriving the diffusion equation, its basic properties are shown.

To derive a version of the diffusion equation in an appropriate variable from the Fokker-Planck equation, introduce Φ_t such that

$$\rho_t = \rho_{\rm G} \Phi_t, \tag{10}$$

where $\rho_{\rm G}$ has been defined in (8). To write the equation for Φ_t in terms of a self-adjoint operator with respect to an inner product, several operators and an inner product are defined below. First, let $\star_{\rm G}$ and its inverse be given by

$$\star_{\mathrm{G}}: \Gamma \Lambda^{k} \mathcal{M} \to \Gamma \Lambda^{m-k} \mathcal{M} \quad \text{and} \quad \star_{\mathrm{G}}^{-1}: \Gamma \Lambda^{m-k} \mathcal{M} \to \Gamma \Lambda^{k} \mathcal{M}, \qquad k = 0, 1, \dots, m$$

with

$$\star_{\rm G} := \rho_{\rm G} \star, \text{ and } \star_{\rm G}^{-1} = \rho_{\rm G}^{-1} \star^{-1},$$

for $\rho_{\rm G} \neq 0$. Notice that

$$\star_{\mathbf{G}}^{-1} \star_{\mathbf{G}} = \star_{\mathbf{G}} \star_{\mathbf{G}}^{-1} = \mathrm{Id}$$

In addition, let d_G^{\dagger} be the map such that

$$\begin{aligned} \mathbf{d}_{\mathbf{G}}^{\dagger} &: \Gamma \Lambda^{k} \mathcal{M} &\to \Gamma \Lambda^{m-1} \mathcal{M}, \qquad k = 0, 1, \dots, m \\ \alpha &\mapsto \mathbf{d}_{\mathbf{G}}^{\dagger} \alpha, \end{aligned}$$

with

$$\mathbf{d}_{\mathbf{G}}^{\dagger} \alpha := (-1)^{k} \star_{\mathbf{G}}^{-1} \mathbf{d} \star_{\mathbf{G}} \alpha, \qquad k = 0, 1, \dots, m$$

From this definition it follows that $d_G^{\dagger} \Phi = 0$ for any function Φ . Thus,

$$d_{\rm G}^{\dagger} = \rho_{\rm G}^{-1} d^{\dagger} \rho_{\rm G}, \qquad \text{or equivalently} \qquad d^{\dagger} = \rho_{\rm G} d_{\rm G}^{\dagger} \rho_{\rm G}^{-1}, \tag{11}$$

from which $d_G^{\dagger} d_G^{\dagger} = 0$. This d_G^{\dagger} is called the *co-derivative associated with* ρ_G in this paper. Introduce the *weighted Laplacian* Δ_G acting on functions on \mathcal{M}

$$\Delta_{\mathbf{G}} := \mathbf{d}_{\mathbf{G}}^{\dagger} \, \mathbf{d},\tag{12}$$

which reduces to the standard Laplacian on \mathcal{M} when $\rho_{\rm G} = 1$. To discuss properties of the weighted Laplacian $\triangle_{\mathbf{G}}$, introduce

$$\begin{array}{cccc} \langle \;,\; \rangle_{\mathbf{G}} : \Gamma \Lambda^{k} \mathcal{M} \times \Gamma \Lambda^{k} \mathcal{M} \; \to \; \mathbb{R} \\ & (\alpha^{\mathbf{I}}, \alpha^{\mathbf{II}}) \; \mapsto \; \left\langle \; \alpha^{\mathbf{I}}, \; \alpha^{\mathbf{II}} \right\rangle_{\mathbf{G}} \end{array}$$

with

$$\langle \alpha^{\mathrm{I}}, \alpha^{\mathrm{II}} \rangle_{\mathrm{G}} = \int_{\mathcal{M}} (\alpha^{\mathrm{I}} \wedge \star \alpha^{\mathrm{II}}) \rho_{\mathrm{G}}, \quad \forall \alpha^{\mathrm{I}}, \alpha^{\mathrm{II}} \in \Gamma \Lambda^{k} \mathcal{M}, \quad k = 0, 1, \dots, m.$$
 (13)

For the case of k = 0, this definition is written as

$$\left\langle \Phi^{\mathrm{I}}, \Phi^{\mathrm{II}} \right\rangle_{\mathrm{G}} = \int_{\mathcal{M}} \Phi^{\mathrm{I}} \Phi^{\mathrm{II}} \rho_{\mathrm{G}} \star 1, \quad \forall \ \Phi^{\mathrm{I}}, \Phi^{\mathrm{II}} \in \Gamma \Lambda^{0} \mathcal{M}.$$

Then $\langle \ , \ \rangle_G, \ d_G^{\dagger}, \ \triangle_G$ have the following properties. These properties are employed below without mentioning explicitly.

Lemma 2.1. 1.

$$\left\langle \alpha^{\mathrm{I}}, \alpha^{\mathrm{II}} \right\rangle_{\mathrm{G}} = \left\langle \alpha^{\mathrm{II}}, \alpha^{\mathrm{I}} \right\rangle_{\mathrm{G}}, \qquad \forall \alpha^{\mathrm{I}}, \alpha^{\mathrm{II}} \in \Gamma \Lambda^{k} \mathcal{M}, \quad k = 0, 1, \dots, m.$$

2. (Adjoint of d_{G}^{\dagger} is d). When the following boundary condition is satisfied:

$$\Phi \,\mathrm{d} \star \rho_{\mathrm{G}} \alpha |_{\partial \mathcal{M}} = 0,$$

it follows that

$$\left\langle \, \mathrm{d}_{\mathrm{G}}^{\dagger} \, \alpha \,, \, \Phi \, \right\rangle_{\mathrm{G}} = \left\langle \, \alpha \,, \, \mathrm{d} \, \Phi \, \right\rangle_{\mathrm{G}}, \qquad \forall \, \alpha \in \Gamma \Lambda^{1} \mathcal{M}, \, \, \Phi \in \Gamma \Lambda^{0} \mathcal{M}.$$

3. (Adjoint of d is d_{G}^{\dagger}). When the following boundary condition is satisfied:

$$\alpha^{\mathrm{I}} \wedge \star \rho_{\mathrm{G}} \alpha^{\mathrm{II}}|_{\partial \mathcal{M}} = 0$$

it follows that

$$\left\langle \mathrm{d}\,\alpha^{\mathrm{I}}\,,\,\alpha^{\mathrm{II}}\,\right\rangle_{\mathrm{G}} = \left\langle \,\alpha^{\mathrm{I}}\,,\,\mathrm{d}_{\mathrm{G}}^{\dagger}\,\alpha^{\mathrm{II}}\,\right\rangle_{\mathrm{G}},\qquad\forall\alpha^{\mathrm{I}}\in\Gamma\Lambda^{k}\mathcal{M},\quad\alpha^{\mathrm{II}}\in\Gamma\Lambda^{k+1}\mathcal{M},\quad k=0,1,\ldots,m.$$

4. (Decomposition of \triangle_{G} into d). When the boundary condition is satisfied:

$$\Phi^{\mathrm{II}} \mathrm{d} \star \rho_{\mathrm{G}} \mathrm{d} \Phi^{\mathrm{I}}|_{\partial \mathcal{M}} = 0,$$

it follows that

$$\left\langle \bigtriangleup_{G} \Phi^{I}, \Phi^{II} \right\rangle_{G} = \left\langle d \Phi^{I}, d \Phi^{II} \right\rangle_{G}, \qquad \forall \Phi^{I}, \Phi^{II} \in \Gamma \Lambda^{0} \mathcal{M}.$$

5. (Self-adjoint property of $\triangle_{\rm G}$). When the boundary conditions are satisfied:

$$\Phi^{\mathrm{II}} \,\mathrm{d} \star \rho_{\mathrm{G}} \,\mathrm{d} \,\Phi^{\mathrm{I}}|_{\partial \mathcal{M}} = 0, \quad \mathrm{and} \quad \Phi^{\mathrm{I}} \,\mathrm{d} \star \rho_{\mathrm{G}} \,\mathrm{d} \,\Phi^{\mathrm{II}}|_{\partial \mathcal{M}} = 0,$$

it follows that

$$\left\langle \bigtriangleup_{\mathrm{G}} \Phi^{\mathrm{I}}, \, \Phi^{\mathrm{II}} \right\rangle_{\mathrm{G}} = \left\langle \, \Phi^{\mathrm{I}}, \, \bigtriangleup_{\mathrm{G}} \Phi^{\mathrm{II}} \, \right\rangle_{\mathrm{G}}, \qquad \forall \, \Phi^{\mathrm{I}}, \Phi^{\mathrm{II}} \in \Gamma \Lambda^{0} \mathcal{M}.$$

Proof. 1. The statement follows by $\alpha^{I} \wedge \star \alpha^{II} = \alpha^{II} \wedge \star \alpha^{I}$ for all $\alpha^{I}, \alpha^{II} \in \Gamma \Lambda^{k} \mathcal{M}, k = 0, 1, \dots, m$.

2. Since the co-derivative associated with $\rho_{\rm G}$ acting on $\alpha \in \Gamma \Lambda^1 \mathcal{M}$ is

$$\mathbf{d}_{\mathbf{G}}^{\dagger} \,\alpha = -\rho_{\mathbf{G}}^{-1} \star^{-1} \mathbf{d} \star \rho_{\mathbf{G}} \alpha,$$

it follows that

$$\left\langle d_{\mathrm{G}}^{\dagger} \alpha, \Phi \right\rangle_{\mathrm{G}} = -\int_{\mathcal{M}} (\rho_{\mathrm{G}}^{-1} \star^{-1} \mathrm{d} \star \rho_{\mathrm{G}} \alpha) \wedge \star \Phi \rho_{\mathrm{G}} = -\int_{\mathcal{M}} \Phi (\mathrm{d} \star \rho_{\mathrm{G}} \alpha)$$
$$= -\int_{\mathcal{M}} \mathrm{d}(\Phi \star \rho_{\mathrm{G}} \alpha) + \int_{\mathcal{M}} \mathrm{d} \Phi \wedge \star \rho_{\mathrm{G}} \alpha = -\Phi \, \mathrm{d} \star \rho_{\mathrm{G}} \alpha|_{\partial \mathcal{M}} + \int_{\mathcal{M}} \mathrm{d} \Phi \wedge \star \rho_{\mathrm{G}} \alpha$$
$$= 0 + \left\langle \mathrm{d} \Phi, \alpha \right\rangle_{\mathrm{G}} = \left\langle \alpha, \mathrm{d} \Phi \right\rangle_{\mathrm{G}}.$$

3. It follows that

$$\begin{split} \left\langle \mathrm{d}\,\alpha^{\mathrm{I}}\,,\,\alpha^{\mathrm{II}}\,\right\rangle_{\mathrm{G}} &= \int_{\mathcal{M}} \mathrm{d}\,\alpha^{\mathrm{I}} \wedge \star \rho_{\mathrm{G}} \alpha^{\mathrm{II}} = \int_{\mathcal{M}} \mathrm{d}(\alpha^{\mathrm{I}} \wedge \star \rho_{\mathrm{G}} \alpha^{\mathrm{II}}) - (-1)^{k} \int_{\mathcal{M}} \alpha^{\mathrm{I}} \wedge \mathrm{d} \star \rho_{\mathrm{G}} \alpha^{\mathrm{II}} \\ &= 0 - (-1)^{k} \int_{\mathcal{M}} \alpha^{\mathrm{I}} \wedge \mathrm{d} \star \rho_{\mathrm{G}} \alpha^{\mathrm{II}} = (-1)^{k+1} \int_{\mathcal{M}} \alpha^{\mathrm{I}} \wedge \star (\rho_{\mathrm{G}} \rho_{\mathrm{G}}^{-1}) \star^{-1} \mathrm{d} \star \rho_{\mathrm{G}} \alpha^{\mathrm{II}} \\ &= (-1)^{k+1} \int_{\mathcal{M}} \alpha^{\mathrm{I}} \wedge \star \rho_{\mathrm{G}} (\rho_{\mathrm{G}}^{-1} \star^{-1} \mathrm{d} \star \rho_{\mathrm{G}}) \alpha^{\mathrm{II}} \\ &= (-1)^{k+1} \int_{\mathcal{M}} \alpha^{\mathrm{I}} \wedge \star \rho_{\mathrm{G}} (-1)^{k+1} \mathrm{d}_{\mathrm{G}}^{\dagger} \alpha^{\mathrm{II}} \\ &= \int_{\mathcal{M}} \alpha^{\mathrm{I}} \wedge \star \rho_{\mathrm{G}} \mathrm{d}_{\mathrm{G}}^{\dagger} \alpha^{\mathrm{II}} = \left\langle \alpha^{\mathrm{I}}\,,\,\mathrm{d}_{\mathrm{G}}^{\dagger} \alpha^{\mathrm{II}} \right\rangle_{\mathrm{G}}. \end{split}$$

4. It follows from item 1 that

$$\left\langle \bigtriangleup_{G} \Phi^{I} , \Phi^{II} \right\rangle_{G} = \left\langle d^{\dagger}_{G} d \Phi^{I} , \Phi^{II} \right\rangle_{G} = \left\langle d \Phi^{I} , d \Phi^{II} \right\rangle_{G}.$$

5. Combining the proved equations, one has

$$\left\langle \bigtriangleup_{G} \Phi^{I}, \, \Phi^{II} \right\rangle_{G} = \left\langle \, d^{\dagger}_{G} \, d \, \Phi^{I}, \, \Phi^{II} \right\rangle_{G} = \left\langle \, d \, \Phi^{I}, \, d \, \Phi^{II} \right\rangle_{G} = \left\langle \, \Phi^{I}, \, d^{\dagger} \, d \, \Phi^{II} \right\rangle_{G} = \left\langle \, \Phi^{I}, \, \bigtriangleup_{G} \Phi^{II} \right\rangle_{G}.$$

In Lemma 2.1, several boundary conditions have been imposed. These conditions are satisfied on closed manifolds due to $\partial \mathcal{M} = \emptyset$. For non-closed manifolds, such as \mathbb{R}^m , these conditions are not strong, because for the case that h is divergent sufficiently fast at the boundary, $\rho_{\rm G} = Z_{\rm G}^{-1} \exp(-\beta h)$ in (8) makes the boundary terms to vanish. In what follows all of these boundary conditions are assumed to be satisfied.

In terms of the weighted Laplacian $\triangle_{\rm G}$ and the change of variables (10), it is shown below, as the main theorem in this subsection, that the Fokker-Planck equation is equivalent to the diffusion equation with the weighted Laplacian.

Theorem 2.2. (Diffusion equation from the Fokker-Planck equation). The function Φ_t in (10) satisfies the modified diffusion equation:

$$\frac{\partial}{\partial t}\Phi_t = -\beta^{-1}\triangle_{\mathcal{G}}\Phi_t,\tag{14}$$

where $\rho_t = \rho_G \Phi_t$ solves the Fokker-Planck equation (1) with $\rho_G \neq 0$ on $\mathcal{M} \setminus \partial \mathcal{M}$, and $\Delta_G = d_G^{\dagger} d$ defined in (12) is the weighted Laplacian.

Proof. By integrating $\dot{\rho}_t \zeta$ over \mathcal{M} with ζ being an arbitrary function, and employing (1), one has

$$\frac{\partial}{\partial t} \langle \rho_t, \zeta \rangle = -\beta^{-1} \left\langle d^{\dagger} d \rho_t, \zeta \right\rangle - \left\langle d^{\dagger} \rho_t d h, \zeta \right\rangle.$$
(15)

After expressing (15) in terms of $\langle , \rangle_{\rm G}$ in (13), and recalling that ζ is arbitrary, one completes the proof. The details of calculations are as follows.

The left hand side of (15) is written as

$$\frac{\partial}{\partial t} \left\langle \rho_t , \zeta \right\rangle = \frac{\partial}{\partial t} \left\langle \Phi_t , \zeta \right\rangle_{\rm G}$$

The right hand side of (15) is written as follows. First, using (11) and (9), one has

$$\begin{aligned} -\beta^{-1} \left\langle d^{\dagger} d \rho_{t}, \zeta \right\rangle &= -\beta^{-1} \left\langle \rho_{\mathrm{G}} d^{\dagger}_{\mathrm{G}} \rho_{\mathrm{G}}^{-1} d \rho_{t}, \zeta \right\rangle = -\beta^{-1} \left\langle d^{\dagger}_{\mathrm{G}} \rho_{\mathrm{G}}^{-1} (d \rho_{t}), \zeta \right\rangle_{\mathrm{G}} \\ &= -\beta^{-1} \left\langle d^{\dagger}_{\mathrm{G}} \rho_{\mathrm{G}}^{-1} (\rho_{\mathrm{G}} d \Phi_{t} + \Phi_{t} d \rho_{\mathrm{G}}), \zeta \right\rangle_{\mathrm{G}} \\ &= -\beta^{-1} \left\langle d^{\dagger}_{\mathrm{G}} (d \Phi_{t} + \rho_{\mathrm{G}}^{-1} \Phi_{t} d \rho_{\mathrm{G}}), \zeta \right\rangle_{\mathrm{G}} \\ &= -\beta^{-1} \left\langle \Delta_{\mathrm{G}} \Phi_{t} + d^{\dagger}_{\mathrm{G}} \rho_{\mathrm{G}}^{-1} \Phi_{t} d \rho_{\mathrm{G}}, \zeta \right\rangle_{\mathrm{G}}, \end{aligned}$$

and

$$\begin{aligned} -\left\langle \,\mathrm{d}^{\dagger}\,\rho_{t}\,\mathrm{d}\,h\,,\,\zeta\,\right\rangle &= -\left\langle\,\rho_{\mathrm{G}}^{-1}\,\mathrm{d}^{\dagger}(\rho_{\mathrm{G}}\rho_{\mathrm{G}}^{-1})\rho_{t}\,\mathrm{d}\,h\,,\,\zeta\,\right\rangle_{\mathrm{G}} = -\left\langle\,\mathrm{d}_{\mathrm{G}}^{\dagger}\,\rho_{\mathrm{G}}^{-1}(\rho_{\mathrm{G}}\Phi_{t})\,\mathrm{d}\,h\,,\,\zeta\,\right\rangle_{\mathrm{G}} \\ &= -\left\langle\,\mathrm{d}_{\mathrm{G}}^{\dagger}\,\Phi_{t}\,\mathrm{d}\,h\,,\,\zeta\,\right\rangle_{\mathrm{G}} = \beta^{-1}\left\langle\,\mathrm{d}_{\mathrm{G}}^{\dagger}\,\rho_{\mathrm{G}}^{-1}\Phi_{t}\,\mathrm{d}\,\rho_{\mathrm{G}}\,,\,\zeta\,\right\rangle_{\mathrm{G}}.\end{aligned}$$

Hence, by summing the two obtained equations, the right hand side of (15) is expressed as

$$-\beta^{-1}\langle \Delta_{\mathbf{G}}\Phi_t, \zeta \rangle_{\mathbf{G}}.$$

Since the right hand side equals the left hand side, one has

$$\frac{\partial}{\partial t} \left\langle \Phi_t , \zeta \right\rangle_{\mathcal{G}} = -\beta^{-1} \left\langle \bigtriangleup_{\mathcal{G}} \Phi_t , \zeta \right\rangle_{\mathcal{G}}.$$

Since ζ is arbitrary, one has (14).

Another derivation of (14) is shown in Section A.1. The operator \triangle_{G} is related to the so-called Witten Laplacians, where the Witten Laplacians are widely discussed in several contexts. To avoid confusion, in this paper, a weighted Laplacian associated with an inner product is called the *Witten Laplacian*, when a weight is not identical. Hence several Witten Laplacians can be defined.

Remark 2.3. There are several studies on how to rewrite the Fokker-Planck equation in the literature. In this Remark some of them are briefly discussed.

In Theorem 2.2 the change of variables, (10), is employed. This sort of change of variables was also employed in [16] for deriving a diffusion equation from the continuous-time master equation without approximation. In addition, some changes of variables are discussed in Sections 5 and 6 of [36]. In particular in Section 6.3 of [36], a transformation is presented in which a method of eigenfunction expansions can be applied to the Fokker-Planck equation, and one operator in [36] is not self-adjoint (or equivalently, that operator is non-hermitian). Meanwhile the present operator $\Delta_{\rm G}$ is self-adjoint as shown in Lemma 2.1, and hence a general theory of self-adjoint operators on manifolds can fully be applied to the study of the Fokker-Planck equation. In the literature another self-adjoint operator acting on $\Phi_t = \rho_G^{-1} \rho_t$ has been known [41, 28]. To briefly discuss such an operator, let $L_{\beta,h}$ be the operator such that $-L_{\beta,h}\Phi_t$ is the right hand side of (14) (see Propositions A.2 and A.3 in Section A.3 of this paper for properties of $L_{\beta,h}$). It is unclear how to generalize $L_{\beta,h}$ so that it can be applied to a k-form with $k \ge 1$. An advantageous point of $\Delta_{\rm G}$ is that it can be generalized so that its action to k-forms is defined, $\Gamma \Lambda^k \mathcal{M} \to \Gamma \Lambda^k \mathcal{M}, k = 0, \dots, m$. This generalization should be $d_{G}^{\dagger} d + d d_{G}^{\dagger}$, by following the definition of the standard Laplacian for k-forms [5]. Another advantageous point is as follows. As shown in Proposition A.3, $\triangle_{\rm G}$ is equal to a version of the Witten Laplacian studied in [10] up to some constant. Hence existing theorems about the Witten Laplacian can be applied to the Fokker-Planck equation.

In the rest of this subsection properties of Φ_t and (14) in Theorem 2.2 are shown.

In theorem 2.2, the diffusion equation is derived. Conversely, as shown below, the diffusion equation with some conditions yield the Fokker-Planck equation.

Proposition 2.4. Let ρ_0 be an equilibrium distribution function associated with a function h such that

 $\dot{\rho}_0 = 0$, and $d \rho_0 = -\beta \rho_0 d h$

In addition, let $\triangle_0 = d_0^{\dagger} d$ be the weighted Laplacian with $d_0^{\dagger} = \rho_0^{-1} d \rho_0$. Then the diffusion equation associated with \triangle_0 ,

$$\frac{\partial}{\partial t}\Phi_t = -\beta^{-1}\triangle_0\Phi_t \tag{16}$$

induces the Fokker-Planck equation.

Proof. Let $\rho_t = \Phi_t \rho_0$. Multiplying both sides of (16) by ρ_0 , one has

$$\frac{\partial}{\partial t}\rho_t = -\beta^{-1}\rho_0 \operatorname{d}_0^{\dagger} \operatorname{d} \Phi_t = -\beta^{-1}\rho_0(\rho_0^{-1} \operatorname{d}^{\dagger} \rho_0) \operatorname{d} \Phi_t = -\beta^{-1} \operatorname{d}^{\dagger} \rho_0 \operatorname{d} \Phi_t.$$

Since $d \rho_t = \rho_0 d \Phi_t + \Phi_t d \rho_0$ and $d \rho_0 = -\beta \rho_0 d h$, one has

$$\frac{\partial}{\partial t}\rho_t = -\beta^{-1} \operatorname{d}^{\dagger}(\operatorname{d} \rho_t - \Phi_t \operatorname{d} \rho_0) = -\beta^{-1} \operatorname{d}^{\dagger} \operatorname{d} \rho_t - \operatorname{d}^{\dagger} \rho_t \operatorname{d} h,$$

which is the Fokker-Planck equation, (1).

There are several properties of Φ_t as shown below.

Lemma 2.5.

$$\langle \Phi_t, 1 \rangle_{\mathcal{G}} = \langle 1, 1 \rangle_{\mathcal{G}} = 1, \text{ and } \left\langle \dot{\Phi}_t, 1 \right\rangle_{\mathcal{G}} = 0.$$

Proof. Straightforward calculations show that

$$\langle \Phi_t, 1 \rangle_{\mathcal{G}} = \int_{\mathcal{M}} \Phi_t \rho_{\mathcal{G}} \star 1 = \int_{\mathcal{M}} \rho_t \star 1 = 1, \text{ and } \langle 1, 1 \rangle_{\mathcal{G}} = \int_{\mathcal{M}} \rho_{\mathcal{G}} \star 1 = 1,$$

due to (2) and (7). In addition, it follows that

$$\left\langle \dot{\Phi}_{t}, 1 \right\rangle_{\mathbf{G}} = \left\langle -\beta^{-1} \triangle_{\mathbf{G}} \Phi, 1 \right\rangle_{\mathbf{G}} = -\beta^{-1} \left\langle \Phi, \Delta_{\mathbf{G}} 1 \right\rangle_{\mathbf{G}} = -\beta^{-1} \left\langle \Phi, 0 \right\rangle_{\mathbf{G}} = 0,$$

or when $\partial/\partial t$ and integral over \mathcal{M} commutes,

$$\left\langle \dot{\Phi}_t, 1 \right\rangle_{\mathcal{G}} = \frac{\mathrm{d}}{\mathrm{d}\,t} \left\langle \Phi_t, 1 \right\rangle_{\mathcal{G}} = \frac{\mathrm{d}}{\mathrm{d}\,t} \mathbf{1} = 0.$$

To show that $\Phi_t = 1$ is stable under (14), one can construct a Lyapunov function, where the state $\Phi_t = 1$ is equivalent to $\rho_t = \rho_G \Phi_t$. To show this property of Φ_t , let Υ_t be the function of t defined as

$$\Upsilon_t := \frac{\beta}{2} \int_{\mathcal{M}} \mathrm{d}\,\Phi_t \wedge \rho_{\mathrm{G}} \star \mathrm{d}\,\Phi_t = \frac{\beta}{2} \left\langle \,\mathrm{d}\,\Phi_t \,,\,\mathrm{d}\,\Phi_t \,\right\rangle_{\mathrm{G}}, \quad \forall t \in \mathbb{R}.$$

Then the following statements show that $\rho_{\rm G}$ is attractive under the time-development.

Proposition 2.6. The function Υ_t is a Lyapunov function for (14), and $\Phi_t = \text{const.}$ is asymptotically stable. **Proof.** It immediately follows that $\Upsilon_t \ge 0, \forall t \in \mathbb{R}$. The derivative of Υ_t with respect to t is calculated with (14) to be

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}t} \Upsilon_t &= \beta \left\langle \mathrm{d} \dot{\Phi}_t \,, \, \mathrm{d} \Phi_t \right\rangle_{\mathrm{G}} = \beta \left\langle -\beta^{-1} \, \mathrm{d} \bigtriangleup_{\mathrm{G}} \Phi_t \,, \, \mathrm{d} \Phi_t \right\rangle_{\mathrm{G}} = - \left\langle \bigtriangleup_{\mathrm{G}} \Phi_t \,, \, \mathrm{d}_{\mathrm{G}}^{\dagger} \, \mathrm{d} \Phi_t \right\rangle_{\mathrm{G}} \\ &= - \left\langle \bigtriangleup_{\mathrm{G}} \Phi_t \,, \, \bigtriangleup_{\mathrm{G}} \Phi_t \right\rangle_{\mathrm{G}} \le 0, \end{aligned}$$

for all $t \in \mathbb{R}$. Combining $\Upsilon_t \ge 0$ and $\dot{\Upsilon}_t \le 0$, one concludes that Υ_t is a Lyapunov function. The equality holds when $\triangle_{\mathbf{G}} \Phi_t = 0$. Notice for c being a constant that $\triangle_{\mathbf{G}} c = \mathbf{d}_{\mathbf{G}}^{\dagger} \mathbf{d} c = \mathbf{d}_{\mathbf{G}}^{\dagger} \mathbf{0} = 0$. Applying the Lyapunov theorem to this inequality, one completes the proof.

From Proposition 2.6, the following holds:

Corollary 2.7.

$$\lim_{t \to \infty} \Phi_t = 1, \quad \text{and} \quad \lim_{t \to \infty} \rho_t = \rho_G.$$

Proof. By Proposition 2.6, $\Phi_t \to c$ as $t \to \infty$, where c is constant, one has that

$$\lim_{t \to \infty} \rho_t = \rho_{\rm G} \lim_{t \to \infty} \Phi_t = \rho_{\rm G} c.$$

This and the normalization conditions, (2) and (7), yield

$$\int_{\mathcal{M}} \lim_{t \to \infty} \rho_t \star 1 = c \int_{\mathcal{M}} \rho_{\mathcal{G}} \star 1 = 1$$

from which c = 1. This completes the prove.

2.3 Spectrum associated with the Fokker-Planck equation

In Section 3, ODEs will be derived by reducing the diffusion equation (14), where (14) has been derived from the Fokker-Planck equation. To this end several preliminary arguments are necessary, and they are discussed in this subsection. Recall how the standard diffusion equation associated with the unweighted Laplacian reduces to a set of ODEs. The strategy for reducing the modified diffusion equation is to modify this reduction for the standard diffusion equation. Then, one recognizes that the spectrum of the self-adjoint operator $\Delta_{\rm G}$ plays a central role in this reduction associated with the Fokker-Planck equation. Hence in this subsection attention is focused on the eigenvalue problem of $\Delta_{\rm G}$ first. Then its basic properties and applications are shown.

Let ϕ_s be an eigenfunction of $\triangle_{\mathbf{G}}$ labeled by s,

$$\triangle_{\mathbf{G}}\phi_s = \lambda_s\phi_s, \qquad s = 0, 1, 2, \dots, \tag{17}$$

where λ_s is an eigenvalue, and all the elements of $\{\phi_s\}$ are assumed to be orthonormal,

$$\langle \phi_s , \phi_{s'} \rangle_{\mathcal{G}} = \delta_{ss'}, \qquad s, s' = 0, 1, 2, \dots,$$
 (18)

with $\delta_{ss'}$ being the Kronecker delta, giving unity if s = s' and 0 otherwise. In addition all the elements of $\{\phi_s\}$ are assumed to be non-identically vanishing.

The following simple statement plays one of the central roles in the later sections.

Lemma 2.8. In (17), every eigenvalue λ_s is equal or greater than zero.

Proof. From (17), it follows for each s that

$$\lambda_s \left\langle \phi_s \,,\, \phi_s \right\rangle_{\mathcal{G}} = \left\langle \lambda_s \phi_s \,,\, \phi_s \right\rangle_{\mathcal{G}} = \left\langle \bigtriangleup_{\mathcal{G}} \phi_s \,,\, \phi_s \right\rangle_{\mathcal{G}} = \left\langle \operatorname{d}_{\mathcal{G}}^{\dagger} \operatorname{d} \phi_s \,,\, \phi_s \right\rangle_{\mathcal{G}} = \left\langle \operatorname{d} \phi_s \,,\, \operatorname{d} \phi_s \right\rangle_{\mathcal{G}}.$$

Combining this,

 $\left< \, \phi_s \, , \, \phi_s \, \right>_{\rm G} = 1, \qquad {\rm and} \qquad \left< \, {\rm d} \, \phi_s \, , \, {\rm d} \, \phi_s \, \right>_{\rm G} \ge 0,$

one has that $\lambda_s \geq 0$.

In this paper ϕ_s is called the *s*-th mode, and the following ordering is assumed:

$$\lambda_0 = 0 \le \lambda_1 \le \lambda_2 \le \cdots.$$

The following is a basic statement for the eigenvalue problem.

Proposition 2.9. In (17), the 0-th mode always exists, and it is given by

$$\phi_0 = 1.$$

Proof. The 0-th mode ϕ_0 is obtained from $\triangle_G \phi_0 = 0$ as $\phi_0 = 1$, where $\langle \phi_0, \phi_0 \rangle_G = \langle 1, 1 \rangle_G = 1$ is satisfied as required in (18).

To obtain ODEs that are reduced from the Fokker-Planck equation, expand Φ_t in terms of $\{\phi_s\}$ as

$$\Phi_t = \sum_{s=0}^{\infty} a_t^s \phi_s, \qquad \phi_s \in \Gamma \Lambda^0 \mathcal{M}$$
⁽¹⁹⁾

where

$$a^{s}: \mathbb{R} \to \mathbb{R} \qquad s = 0, 1, \dots,$$
$$t \mapsto a^{s}_{t}.$$

In this paper (19) is called the *eigenfunction expansion* of the Fokker-Planck equation or that of Φ_t . As shown below, this expansion yields the dynamical system for a_t^s , and its time-scale is $\beta^{-1}\lambda_s$ for each s.

Proposition 2.10. For each s, a_t^s satisfies the ODE

$$\frac{\mathrm{d}}{\mathrm{d}\,t}a_t^s = -\beta^{-1}\lambda_s a_t^s$$

whose solution is given by

$$a_t^s = a_0^s \exp(-\beta^{-1}\lambda_s t)$$

Proof. To derive the equation of a_t^s , take the inner product of (14) and ϕ_s with a fixed s:

$$\left\langle \phi_s , \dot{\Phi}_t \right\rangle_{\mathbf{G}} = \left\langle \phi_s , -\beta^{-1} \triangle_{\mathbf{G}} \Phi_t \right\rangle_{\mathbf{G}}.$$

The left and right hand sides of this are

$$\left\langle \phi_s \,, \, \frac{\partial \Phi_t}{\partial t} \right\rangle_{\mathbf{G}} = \sum_{s'} \frac{\mathrm{d} \, a_t^{s'}}{\mathrm{d} \, t} \left\langle \phi_s \,, \, \phi_{s'} \right\rangle_{\mathbf{G}} = \frac{\mathrm{d} \, a_t^s}{\mathrm{d} \, t},$$

$$\left\langle \phi_s \,, \, -\beta^{-1} \triangle_{\mathbf{G}} \Phi_t \right\rangle_{\mathbf{G}} = -\beta^{-1} \sum_{s'} \lambda_{s'} a_t^{s'} \left\langle \phi_s \,, \, \phi_{s'} \right\rangle_{\mathbf{G}} = -\beta^{-1} \lambda_s a_t^s,$$

respectively. Since they are equal, one has that

$$\frac{\mathrm{d}}{\mathrm{d}t}a_t^s = -\beta^{-1}\lambda_s a_t^s, \qquad \forall \ s$$

whose solution is immediately obtained. This yields the desired expression of the solution.

Remark 2.11. The 0-th mode is interpreted as the equilibrium state. To see this, consider the special case where there is only the 0-th mode, that is, $\Phi_t = a_t^0 \phi_0$ with $\phi_0 = 1$. Then, it follows from Proposition 2.10 that $\Phi_t = a_0^0$, where a_0^0 is constant. From (10), it follows that $\rho_t = \rho_G \Phi_t = a_0^0 \rho_G$. This and normalization conditions, (2) and (7), yield $a_0^0 = 1$. Hence this special case consisting only of the 0-th mode yields $\rho_t = \rho_G$, and the 0-th mode expresses the equilibrium distribution function ρ_G .

Remark 2.12. It follows from Proposition 2.10 with (19) that Φ_t has the countable time-scales $\{\beta^{-1}\lambda_s\}$:

$$\Phi_t = \sum_{s=0}^{\infty} a_0^s \exp(-\beta^{-1}\lambda_s)\phi_s.$$

In addition, since $\rho_t = \rho_G \Phi_t$ in (10), ρ_t also has the countable time-scales.

There are several applications of the eigenfunction expansion of the Fokker-Planck equation. In what follows one of them is considered. Recall that the Fokker-Planck equation describes the time-evolution of the probability distribution function, and hence one can define expectation value (or expected value) with respect to the solution to the Fokker-Planck equation by integrating a function over possible states with a weight. In this paper these expectation values are assumed to be identified with nonequilibrium thermodynamic variables.

To write the expectation values in terms of a geometric language, let B_1 be a function on \mathcal{M} . The expectation value of B_1 at t is expressed in terms of \mathbb{E}_t , where

$$\mathbb{E}_{t}: \Gamma \Lambda^{0} \mathcal{M} \to \mathbb{R}$$

$$B_{1} \mapsto \mathbb{E}_{t}[B_{1}]$$

$$\mathbb{E}_{t}[B_{1}] := \int_{\mathcal{M}} B_{1} \rho_{t} \star 1.$$
(20)

with

This time-dependent quantity can be written in terms of the inner-product:

$$\mathbb{E}_t[B_1] = \int_{\mathcal{M}} B_1 \rho_t \star 1 = \int_{\mathcal{M}} B_1 \Phi_t \rho_{\mathrm{G}} \star 1 = \langle B_1, \Phi_t \rangle_{\mathrm{G}}$$

If B_1 does not depend on t, then the following holds:

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}_t[B_1] = \int_{\mathcal{M}} B_1 \rho_{\mathrm{G}} \dot{\Phi}_t \star 1 = -\beta^{-1} \langle B_1, \, \triangle_{\mathrm{G}} \Phi_t \, \rangle_{\mathrm{G}} = -\beta^{-1} \langle \triangle_{\mathrm{G}} B_1, \, \Phi_t \, \rangle_{\mathrm{G}}$$

Remark 2.13. When $B_1 = 1$, the normalization condition for ρ_t is reconstructed in the sense that

$$\frac{\mathrm{d}}{\mathrm{d}t} \int_{\mathcal{M}} \rho_t \star 1 = \frac{\mathrm{d}}{\mathrm{d}t} \mathbb{E}_t[1] = -\beta^{-1} \left\langle \bigtriangleup_{\mathrm{G}} 1, \Phi_t \right\rangle_{\mathrm{G}} = 0,$$

due to $riangle_G 1 = d_G^{\dagger} d 1 = d_G^{\dagger} 0 = 0.$

There are several formulae regarding the expansion of B_1 in terms of eigenfunctions. To show these, expand $B_1 \in \Gamma \Lambda^0 \mathcal{M}$ in terms of $\{\phi_s\}$:

$$B_1 = \sum_{s'=0}^{\infty} b_{s'} \phi_{s'}, \quad \text{where} \quad b_s = \langle \phi_s \,, \, B_1 \, \rangle_{\mathcal{G}} \, \in \mathbb{R}, \qquad \forall \ s$$

This and

$$\rho_t = \rho_{\rm G} \Phi_t = \rho_{\rm G} \sum_{s=0}^{\infty} a_t^s \phi_s,$$

yield

$$\mathbb{E}_t[B_1] = \int_{\mathcal{M}} B_1 \rho_t \star 1 = \sum_{s'} b_{s'} \int_{\mathcal{M}} \phi_{s'} \sum_s a_t^s \phi_s \rho_G \star 1$$
$$= \sum_s \sum_{s'} b_{s'} a_t^s \langle \phi_{s'}, \phi_s \rangle_G = \sum_s b_s a_t^s.$$

and if $b_s = 0$ for all s, then it follows that

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}_t[B_1] = -\beta^{-1} \left\langle \bigtriangleup_{\mathrm{G}} B_1, \Phi_t \right\rangle_{\mathrm{G}} = -\beta^{-1} \sum_s \lambda_s b_s a_t^s$$

In addition, one has

$$\mathbb{E}_t[\phi_s] = \int_{\mathcal{M}} \phi_s \rho_t \star 1 = \sum_{s'} \langle \phi_{s'}, \phi_s \rangle_{\mathcal{G}} a_t^{s'} = a_t^s, \quad \text{and} \quad \frac{\mathrm{d}}{\mathrm{d}\,t} \mathbb{E}_t[\phi_s] = -\beta^{-1} \lambda_s a_t^s = -\beta^{-1} \lambda_s \mathbb{E}_t[\phi_s].$$

The first equation above and (19) yield

$$\Phi_t = \sum_s (\mathbb{E}_t[\phi_s])\phi_s.$$

Before closing this section, it should be mentioned that, in the literature, there are several operators that are analogous to $d_{\rm G}^{\dagger} d$ [22]. To compare the approach in this paper with the existing approaches, several calculations are shown in appendix of this paper (see Section A.2). In addition in the case where the standard exterior derivative and co-derivative d and d^{\dagger} are only employed, one can also have a PDE for $\Phi_t = \rho_{\rm G}^{-1} \rho_t$, as shown in Section A.3.

3 From the diffusion equation to a contact Hamiltonian system

In Section 2, a diffusion equation associated with a weighted Laplacian has been derived. In this section, this diffusion equation is shown to induce a contact Hamiltonian system. In contact geometry, contact Hamiltonian systems are defined on contact manifolds and employed to describe thermodynamics, where contact geometry is an odd-dimensional analogue of symplectic geometry [4, 30]. After recalling contact geometry briefly and fix our notation, it is shown how the diffusion equation induces a contact Hamiltonian system.

3.1 Contact geometry

In this subsection necessary background of contact geometry is summarized.

A contact manifold is a pair $(\mathcal{N}, \ker \alpha)$, where \mathcal{N} is an odd-dimensional manifold and $\ker \alpha = \{X \in \Gamma T\mathcal{N} | \alpha(X) = 0\}$ with α being a contact form. The interpretation of \mathcal{N} in this section is thermodynamic phase space, whose coordinates represent macroscopic thermodynamic variables. Let $\dim \mathcal{N} = 2n + 1$, and denote by **d** the exterior derivative, $\mathbf{d} : \Gamma \Lambda^k \mathcal{N} \to \Gamma \Lambda^{k+1} \mathcal{N}$, $(k = 0, 1, \dots, 2n + 1)$. A contact form α on \mathcal{N}

is a 1-form that the form $\alpha \wedge \mathbf{d} \alpha \wedge \cdots \wedge \mathbf{d} \alpha \in \Gamma \Lambda^{2n+1} \mathcal{N}$ does not vanish anywhere. There are canonical coordinates (p, q, z) with $p = (p_1, \ldots, p_n)$ and $q = (q^1, \ldots, q^n)$ such that the contact form is written as

$$\alpha = \mathbf{d} \, z - \sum_{j=1}^n p_j \, \mathbf{d} \, q^j.$$

As an example, the pair consisting of $T^*\mathbb{R}^n \times \mathbb{R}$ and ker α is a contact manifold, where $T^*\mathbb{R}^n$ is the cotangent bundle of \mathbb{R}^n , q denotes a point of \mathbb{R}^n , p coordinates of $T^*_q\mathbb{R}^n$, z coordinate of \mathbb{R} , and α is given above. Given a function \mathcal{H} on \mathcal{N} , a vector field $X_{\mathcal{H}}$ satisfying

$$i_{X_{\mathcal{H}}} \mathbf{d} \alpha = -\mathbf{d} \mathcal{H} + (R\mathcal{H})\alpha, \text{ and } i_{X_{\mathcal{H}}} \alpha = \mathcal{H}$$

is called a *contact Hamiltonian vector field*, where \mathcal{H} is called a *contact Hamiltonian*, R is called the *Reeb* vector field that is the unique vector field satisfying $i_R \alpha = 1$ and $i_R \mathbf{d} \alpha = 0$. In the canonical coordinates R and $X_{\mathcal{H}}$ are written as

$$R = \frac{\partial}{\partial z}, \qquad X_{\mathcal{H}} = \dot{z}\frac{\partial}{\partial z} + \sum_{j=1}^{n} \left(\dot{p}_{j}\frac{\partial}{\partial p_{j}} + \dot{q}^{j}\frac{\partial}{\partial q^{j}} \right),$$

where

$$\dot{q}^{j} = -\frac{\partial \mathcal{H}}{\partial p_{j}}, \quad \dot{p}_{j} = \frac{\partial \mathcal{H}}{\partial q^{j}} + p_{j}\frac{\partial \mathcal{H}}{\partial z}, \quad \dot{z} = \mathcal{H} - \sum_{j=1}^{n} p_{j}\frac{\partial \mathcal{H}}{\partial p_{j}}, \qquad j = 1, \dots, n.$$
 (21)

Remark 3.1. To relate (21) with a dynamical system, \dot{t} in the left hand side is identified with d/dt. If \mathcal{H} is written as $\mathcal{H} = \gamma_1 \mathcal{H}^{(0)}$ with some function $\mathcal{H}^{(0)}$ and constant $\gamma_1 > 0$, then γ_1 gives a natural time-scale. To see this, let $t_1 := \gamma_1 t$. Then t_1 can be recognized as a scaled time with a time scale γ_1 . If introducing t_1 , then γ_1 does not explicitly appear in the dynamical system (21).

In the applications of contact geometry to thermodynamics, α is employed to express the laws of thermodynamics, and $X_{\mathcal{H}}$ preserves ker α . Equilibrium thermodynamic phase space is modeled by a class of Legendrian submanifolds [32, 2]. The Legendrian submanifold \mathcal{L} is a submanifold, where dim $\mathcal{L} = n$ and the pullback of α to \mathcal{L} vanishes. One example of \mathcal{L} is expressed with a function ψ of q in the canonical coordinates as

$$\mathcal{L}_{\psi} := \bigcup_{q \in \mathbb{R}} \mathcal{L}_{\psi}(q), \ \subset T^* \mathbb{R}^n \times \mathbb{R}, \quad \text{with} \quad \mathcal{L}_{\psi}(q) := \left\{ \left. (p, q, z) \right| \ p_j = \frac{\partial \psi}{\partial q^j}(q), \quad z = \psi(q), \quad j = 1, \dots, n \right\}.$$

There are several approaches to express nonequilibrium thermodynamic processes by means of contact geometry, [13, 6, 18].

3.2 Derivation of a contact Hamiltonian system

To derive a contact Hamiltonian system from the Wasserstein gradient flow or equivalently the Fokker-Planck equation, thermodynamic variables are needed to be defined, since contact Hamiltonian equations are equations for thermodynamic variables. In general, thermodynamic variables form pairs, and they consist of intensive and extensive parameters [3]. In this paper they are called *primal and conjugate (or dual) variables*, where primal variables often express externally applied fields, and the conjugate or dual variables express expectation values. An example of this pair is an applied magnetic field and the magnetization for an Ising spin system, where the magnetization is obtained by integrating a function over a state space with the weight of a distribution function. As is known in equilibrium thermodynamics, given primal variables, thermodynamic conjugate variables are obtained by differentiation of a moment generating function with respect to the primal variables (see [27, 39] for applications of moment generating functions in statistical mechanics). In this paper, this known approach for equilibrium systems is extended to that for nonequilibrium systems. In this subsection, after introducing time-dependent thermodynamic variables and a time-dependent moment generating function, it is shown how a contact Hamiltonian equation is derived from the Fokker-Planck equation. The principal step in this derivation, from a PDE to a system of ODEs, is to integrate functions over \mathcal{M} with a weight. In calculating various time-dependent quantities, the Fokker-Planck equation (1) is identified with the diffusion equation (14) via the change of variables (10).

To describe a set of primal variables q and its conjugate ones, let

$$q = (q^1, q^2, \dots, q^n) \in \mathbb{R}^n, \quad B = (B_1, B_2, \dots, B_n) \text{ with } B_j \in \Gamma \Lambda^0 \mathcal{M}, \quad j = 1, \dots, n$$

and

$$q \cdot B := \sum_{j=1}^{n} q^{j} B_{j} \in \Gamma \Lambda^{0} \mathcal{M}.$$

The time-dependent moment generating function,

$$M: \mathbb{R} \times \mathbb{R}^n \quad \to \quad \mathbb{R},$$
$$(t,q) \quad \mapsto \quad M_t(q)$$

is defined as

$$M_t(q) := \mathbb{E}_t[\exp(q \cdot B)] = \int_{\mathcal{M}} \rho_{\mathcal{G}} \Phi_t \exp(q \cdot B) \star 1 = \langle \Phi_t, \exp(q \cdot B) \rangle_{\mathcal{G}}.$$
 (22)

This generating function is employed to obtain expectation values. By assuming the commutability between $\partial/\partial q^j$ and the integral over \mathcal{M} , one has from (22) that

$$\frac{\partial}{\partial q^j} M_t(q) = \mathbb{E}_t[B_j \exp(q \cdot B)] = \int_{\mathcal{M}} \rho_{\mathcal{G}} \Phi_t B_j \exp(q \cdot B) \star 1 = \langle \Phi_t, B_j \exp(q \cdot B) \rangle_{\mathcal{G}},$$
(23)

where Φ_t has been given in (10) and this Φ_t is independent of $q \cdot B$ but depends on h. An interpretation of q^j and

$$\mathbb{E}_t[B_j \exp(q \cdot B)]$$
 or $\mathbb{E}_t[B_j]$

for each j is a pair of thermodynamic primal and time-dependent conjugate variables.

Since time-evolution is governed by the Fokker-Planck equation, asymptotic limits of (22) and (23) are obtained with Corollary 2.7 as

$$\lim_{t \to \infty} M_t(q) = \langle \exp(q \cdot B), 1 \rangle_{\mathcal{G}} = \psi_{\mathcal{G}}(q),$$
$$\lim_{t \to \infty} \frac{\partial}{\partial q^j} M_t(q) = \langle B_j \exp(q \cdot B), 1 \rangle_{\mathcal{G}} = \frac{\partial \psi_{\mathcal{G}}}{\partial q^j}(q), \qquad j = 1, \dots, n$$

where we have introduced $\psi_{\mathrm{G}}: \mathbb{R}^n \to \mathbb{R}$ with

$$\psi_{\mathcal{G}}(q) := \langle \exp(q \cdot B), 1 \rangle_{\mathcal{G}} = \int_{\mathcal{M}} \exp(q \cdot B) \rho_{\mathcal{G}} \star 1.$$
(24)

In addition, the asymptotic limit of $M_t(q)$ is expressed as

$$\lim_{t \to \infty} M_t(q) = \frac{1}{Z_G} \int_{\mathcal{M}} e^{-\beta \tilde{h}} \star 1, \qquad \tilde{h} := h - \beta^{-1} q \cdot B$$
$$= \frac{Z_G(q)}{Z_G(0)}, \qquad Z_G(q) := \int_{\mathcal{M}} e^{-\beta h + q \cdot B} \star 1$$

where an interpretation of \tilde{h} is the sum of a microscopic Hamiltonian and an energy due to an externally applied field $-\beta^{-1}q \cdot B$, where $\tilde{h} = \ln(\rho_{\rm G} \exp(q \cdot B)) + \text{const.}$ To see this interpretation with n = 1, consider a system consisting of interacting Ising spins as an example. Then h describes spin-spin interactions, $\beta^{-1}q$ a constant expressing an externally applied magnetic filed, and B denotes magnetization, the mean of Ising spins. Analogously, $\ln(\rho_t \exp(q \cdot B))$, the log of the integrand of (22) at t, is possibly interpreted as the sum of a microscopic Hamiltonian and an energy due to an externally applied field up to an additional constant. **Remark 3.2.** The function $\psi_{\rm G}$ is convex, since for any $q \in \mathbb{R}^n$ the Hessian matrix $(\partial^2 \psi_{\rm G}/\partial q^j \partial q^k)$ is positive semi-definite:

$$\sum_{j=1}^{n} \sum_{k=1}^{n} q^{j} \frac{\partial^{2} \psi_{\mathrm{G}}}{\partial q^{j} \partial q^{k}} q^{k} = \sum_{j=1}^{n} \sum_{k=1}^{n} q^{j} q^{k} \int_{\mathcal{M}} B_{j} B_{k} \, \mathrm{e}^{q \cdot B} \, \rho_{\mathrm{G}} \star 1 = \int_{\mathcal{M}} (q \cdot B)^{2} \, \mathrm{e}^{q \cdot B} \, \rho_{\mathrm{G}} \star 1 \ge 0.$$

Under these preliminary definitions, attention can now be focused on a contact geometric description of a system, where this system is derived from the Fokker-Planck equation. Discussions below are divided into 2 cases. One is full-time scale dynamics, and the other is slowest-time scale dynamics. In the latter, a contact Hamiltonian system is derived.

Full times-scale dynamics: a non-closed system

In the following several variables are introduced so that a contact geometric description of the Fokker-Planck equation is established. Meanwhile the derivation of a contact Hamiltonian is not given here.

Introduce $z(t;q), p_j(t;q)$ with $q = (q^1, q^2, \dots, q^n)$ as

$$z(t;q) := M_t(q) = \langle \Phi_t, \exp(q \cdot B) \rangle_{\mathrm{G}},$$

$$p_j(t;q) := \frac{\partial}{\partial q^j} M_t(q) = \langle \Phi_t, B_j \exp(q \cdot B) \rangle_{\mathrm{G}}, \qquad j = 1, \dots, n$$

These variables z and p contain all the time-scales $\{\beta^{-1}\lambda_s\}$ as discussed in Remark 2.12, therefore its corresponding dynamics is called *full time-scale dynamics* in this paper. As is shown below, this dynamics on a contact manifold is not written as a simple contact Hamiltonian system. Meanwhile making an approximation discussed later allows to write it as a simple contact Hamiltonian system, where this approximation scheme is viewed as a projection of a function onto a subspace spanned by a set of lower eigenfunctions.

Let (p,q,z) be coordinates of $T^*\mathbb{R}^n \times \mathbb{R}$, and let α be the 1-form on $T^*\mathbb{R}^n \times \mathbb{R}$ as

$$\alpha = \mathbf{d} \, z - \sum_{j=1}^n p_j \, \mathbf{d} \, q^j.$$

Then $(T^*\mathbb{R}^n \times \mathbb{R}, \ker \alpha)$ is a contact manifold. Points on the Legendrian submanifold expressed in coordinates,

$$\mathcal{L}_{\psi_{\mathrm{G}}} := \bigcup_{q \in \mathbb{R}} \mathcal{L}_{\psi_{\mathrm{G}}}(q), \quad \mathcal{L}_{\psi_{\mathrm{G}}}(q) := \left\{ \left. (p, q, z) \right| z = \psi_{\mathrm{G}}(q), \quad p_{j} = \frac{\partial \psi_{\mathrm{G}}}{\partial q^{j}}(q), \quad j = 1, \dots, n \right\}, \tag{25}$$

are interpreted as points of the equilibrium state. Consider dynamical behavior of z(t;q) on the z-axis, and that of p(t;q) on the *p*-axis. A natural question is whether or not the long-time limit of the point (p(t;q),q,z(t;q)) is on the Legendrian submanifold. The answer is as follows.

Proposition 3.3. The long-time limit of the point (p(t;q),q,z(t;q)) is on the Legendrian submanifold $\mathcal{L}_{\psi_{\mathrm{G}}}(q)$, where dynamics of z(t;q) and p(t;q) obeys the diffusion equation (14) associated with the Fokker-Planck equation.

Proof. The long-time limits of z and p are obtained by using $\Phi_t \to 1$, $(t \to \infty)$ shown in Corollary 2.7 as

$$\lim_{t \to \infty} z(t;q) = \langle \exp(q \cdot B), 1 \rangle_{\mathcal{G}} = \psi_{\mathcal{G}}(q),$$
$$\lim_{t \to \infty} p_j(t;q) = \langle B_j \exp(q \cdot B), 1 \rangle_{\mathcal{G}} = \frac{\partial}{\partial q^j} \langle \exp(q \cdot B), 1 \rangle_{\mathcal{G}} = \frac{\partial \psi_{\mathcal{G}}}{\partial q^j}(q), \qquad j = 1, \dots, n.$$

Hence the point in the limit $t \to \infty$ is on the Legendrian submanifold $\mathcal{L}_{\psi_{\mathrm{G}}}(q)$.

Remark 3.4. In many cases, q = 0 is substituted after evaluating p_j , and it is useful to expand z and p in terms of q. They are calculated to be

$$\lim_{t \to \infty} z(t;q) = \psi_{\mathcal{G}}(q) = 1 + \sum_{j=1}^{n} q^{j} \langle B_{j}, 1 \rangle_{\mathcal{G}} + \mathcal{O}(q^{2}),$$
$$\lim_{t \to \infty} p_{j}(t;q) = \frac{\partial \psi_{\mathcal{G}}}{\partial q^{j}} = \frac{\partial}{\partial q^{j}} \left\langle 1 + (q \cdot B) + \frac{(q \cdot B)^{2}}{2!} + \mathcal{O}(q^{3}), 1 \right\rangle_{\mathcal{G}}$$
$$= \langle B_{j}, 1 \rangle_{\mathcal{G}} + \sum_{k=1}^{n} q^{k} \langle B_{j}, B_{k} \rangle_{\mathcal{G}} + \mathcal{O}(q^{2}).$$

After clarifying the long-time limit, the next point of discussion is dynamical behavior of z and p governed by the Fokker-Planck equation. First, observe that

$$\frac{\mathrm{d}}{\mathrm{d}t}z = \left\langle \dot{\Phi}_t, \exp(q \cdot B) \right\rangle_{\mathrm{G}} = -\beta^{-1} \left\langle \bigtriangleup_{\mathrm{G}} \Phi_t, \exp(q \cdot B) \right\rangle_{\mathrm{G}} = -\beta^{-1} \left\langle \Phi_t, \bigtriangleup_{\mathrm{G}} \exp(q \cdot B) \right\rangle_{\mathrm{G}}, \\ \frac{\mathrm{d}}{\mathrm{d}t}p_j = \left\langle \dot{\Phi}_t, B^j \exp(q \cdot B) \right\rangle_{\mathrm{G}} = -\beta^{-1} \left\langle \Phi_t, \bigtriangleup_{\mathrm{G}} (B^j \exp(q \cdot B)) \right\rangle_{\mathrm{G}}, \quad j = 1, \dots, n.$$

In general, the right hand sides of the equations above are not written in terms of z and p. In this sense, they are not closed.

Slowest times-scale dynamics: a contact Hamiltonian system

To obtain a closed dynamical system on $T_q^* \mathbb{R}^n \times \mathbb{R}$, some approximation is introduced below. This approximation is to neglect the fast-time scales, and leads to a contact Hamiltonian system. More precisely, this approximation scheme is based on a projection of Φ_t onto the subspace spanned by ϕ_0 and ϕ_1 . This projection of Φ_t then induces the corresponding projections of z and p.

Before addressing this approximation, recall that the 0-th mode exists as shown in Proposition 2.9. Then the next point of discussion is the 1st mode. The following 2 cases for the 1st mode are considered separately. However the conclusion on one side is equal to a conclusion on the other.

• Case I: There is no degeneracy:

$$0 = \lambda_0 < \lambda_1 < \lambda_2 < \cdots . \tag{26}$$

In the case of nodegeneracy, consider the following formal expansion:

$$\Phi_{t} = \sum_{s=0}^{\infty} a_{t}^{s} \phi_{s} = \sum_{s=0}^{\infty} a_{0}^{s} \exp(-\beta^{-1}\lambda_{s}t)\phi_{s}$$

$$= a_{t}^{0} \phi_{0} + a_{t}^{1} \phi_{1} + \dots = a_{0}^{0} \phi_{0} e^{-\beta^{-1}\lambda_{0}t} + a_{0}^{1} \phi_{1} e^{-\beta^{-1}\lambda_{1}t} + \dots$$

$$= \Phi_{\infty} + a_{0}^{1} \phi_{1} e^{-\beta^{-1}\lambda_{1}t} + \dots$$
(27)

Note that the terms $a_0^s \phi_s$, (s = 1, 2, ...) are written from (27) as

$$a_{0}^{1}\phi_{1} = \lim_{t \to \infty} e^{\beta^{-1}\lambda_{1}t} \left(\Phi_{t} - \Phi_{\infty}\right), \quad a_{0}^{2}\phi_{2} = \lim_{t \to \infty} e^{\beta^{-1}\lambda_{2}t} \left(\Phi_{t} - \Phi_{\infty} - a_{0}^{1}\phi_{1} e^{-\beta^{-1}\lambda_{1}t}\right), \quad \cdots$$

Then, differentiating Φ_t in (27) with respect to t and recalling $\Phi_{\infty} = 1$ from Corollary 2.7, one has

$$\dot{\Phi}_t = -\beta^{-1}\lambda_1 a_0^1 \phi_1 e^{-\beta^{-1}\lambda_1 t} + \cdots$$
$$= -\beta^{-1}\lambda_1 (\Phi_t - 1) + \cdots .$$

Let us define the equation associated with the approximation by this neglect:

Definition 3.5. Dynamics generated by

$$\overline{\Phi}_t(\xi) = 1 + a_t^1 \phi_1(\xi), \quad \forall \xi \in \mathcal{M}, \ \forall t \in \mathbb{R}$$
(28)

that satisfies the PDE

$$\frac{\partial}{\partial t}\overline{\Phi}_t = -\beta^{-1}\lambda_1(\overline{\Phi}_t - 1) \tag{29}$$

is called the *slowest time-scale dynamics*, its solution $\overline{\Phi}_t$ is called the *slowest time-scale mode*, and the equation the *slowest time-scale mode equation*, in this paper.

Notice that (28) can be viewed as a projection of Φ_t onto the subspace spanned by ϕ_0 and ϕ_1 , and can also be viewed as a truncation of the eigenvalue function expansion.

• Case II: There is degeneracy of the form,

$$0 = \lambda_0 < \lambda_1 = \lambda_2 = \dots = \lambda_\nu < \lambda_{\nu+1} < \dots$$

Analogous to (28), let

$$\overline{\Phi}_t^{\mathrm{dgt}} = 1 + (a_0^1 \phi_1 + \dots + a_0^{\nu} \phi_{\nu}) \exp(-\beta^{-1} \lambda_1 t).$$

Then this $\overline{\Phi}_t^{\mathrm{dgt}}$ satisfies

$$\frac{\partial}{\partial t}\overline{\Phi}_t^{\rm dgt} = -\beta^{-1}\lambda_1 \left(\overline{\Phi}_t^{\rm dgt} - 1\right). \tag{30}$$

Since (30) in Case II can be identical to (29) in Case I, these 2 cases are not distinguished below. The following is a remarkable property of (28).

Proposition 3.6. The slowest mode (28) satisfies the modified diffusion equation (14).

Proof. Differentiation of (28) yields

$$\begin{aligned} \frac{\partial}{\partial t} \overline{\Phi}_t &= -\beta^{-1} \lambda_1 a_t^1 \phi_1 = -\beta^{-1} \lambda_1 \left(\overline{\Phi}_t - 1 \right), \\ \Delta_{\mathcal{G}} \overline{\Phi}_t &= a_t^1 \Delta_{\mathcal{G}} \phi_1 = \lambda_1 a_t^1 \phi_1 = \lambda_1 \left(\overline{\Phi}_t - 1 \right). \end{aligned}$$

Combining these equations, one has that

$$\frac{\partial}{\partial t}\overline{\Phi}_t = -\beta^{-1} \triangle_{\mathbf{G}}\overline{\Phi}_t,$$

which is (14).

Proposition 3.6 shows that the approximation, or the projection, (28) is adequate in the sense that full time-scale mode $\overline{\Phi}_t$ and slowest time-scale mode $\overline{\Phi}_t$ obey the same PDE. Hence, this adequate approximation preserves, for example, Lie symmetries of differential equations [34].

It is straightforward to verify by $\beta^{-1}\lambda_1 > 0$ that

$$\overline{\Phi}_t = 1 + (\overline{\Phi}_0 - 1) \exp(-\beta^{-1}\lambda_1 t)$$

= $1 + a_0^1 \exp(-\beta^{-1}\lambda_1 t)\phi_1,$
$$\lim_{t \to \infty} \overline{\Phi}_t = 1.$$
 (31)

To compare the asymptotic behavior of Φ_t and $\overline{\Phi}_t$, the following can be employed:

$$|\Phi_t(\xi) - \overline{\Phi}_t(\xi)| = |a_0^2 \phi_2(\xi) e^{-\beta^{-1}\lambda_2 t} + \cdots|, \quad \forall \xi \in \mathcal{M}.$$

This yields

$$\lim_{t \to \infty} \left| \Phi_t(\xi) - \overline{\Phi}_t(\xi) \right| = 0, \quad \forall \ \xi \in \mathcal{M}$$

To see the approximation in the original variable, introduce

$$\overline{\rho}_t = \rho_{\rm G} \overline{\Phi}_t. \tag{32}$$

Then this $\overline{\rho}_t$ satisfies the PDE

$$\frac{\partial}{\partial t}\overline{\rho}_t = -\beta^{-1}\lambda_1 \left(\overline{\rho}_t - \rho_{\rm G}\right)$$

which is derived by differentiating both sides of (32) with respect to t. In addition, $\overline{\rho}_t$ is a relevant approximation in the sense that the asymptotic limits coincide:

$$\lim_{t \to \infty} |\rho_t(\xi) - \overline{\rho}_t(\xi)| = \rho_{\mathcal{G}}(\xi) \lim_{t \to \infty} |\Phi_t(\xi) - \overline{\Phi}_t(\xi)| = 0, \quad \forall \xi \in \mathcal{M}.$$

When neglecting higher order (or fast time-scale) modes, i.e., $\phi_s = 0$ is substituted for $s \ge 2$, as shown below, a closed system for analogues of z and p is obtained. To obtain such a closed system of ODEs, introduce the slowest time-scale moment generating function

$$\overline{M} : \mathbb{R} \times \mathbb{R}^n \quad \to \quad \mathbb{R}$$
$$(t,q) \quad \mapsto \quad \overline{M}_t(q)$$

with

$$\overline{M}_t(q) := \left\langle \overline{\Phi}_t \,, \, \exp(q \cdot B) \right\rangle_{\mathcal{G}}$$

and introduce the dynamical variables \overline{z} and \overline{p} with $\overline{p} = (\overline{p}_1, \dots, \overline{p}_n)$ such that

$$\overline{z}(t;q) := \overline{M}_t(q) = \left\langle \overline{\Phi}_t \,, \, \exp(q \cdot B) \right\rangle_{\mathcal{G}},\tag{33}$$

$$\overline{p}_{j}(t;q) := \frac{\partial}{\partial q^{j}} \overline{M}_{t}(q) = \left\langle \overline{\Phi}_{t}, B_{j} \exp(q \cdot B) \right\rangle_{\mathrm{G}}, \qquad j = 1, \dots, n.$$
(34)

Consider dynamical behavior of $\overline{z}(t;q)$ on the z-axis, and that of $\overline{p}(t;q)$ on the p-axis. One then finds that there are similarity between z(t;q) and $\overline{z}(t;q)$, and between p(t;q) and $\overline{p}(t;q)$. These are described as follows.

Proposition 3.7. The long-time limit of the point $(\overline{p}(t;q), q, \overline{z}(t;q))$ is on the Legendrian submanifold $\mathcal{L}_{\psi_{\mathrm{G}}}(q)$ of the contact manifold, where dynamics of $\overline{z}(t;q)$ and $\overline{p}(t;q)$ obeys the slowest time-scale mode equation (29).

Proof. A proof is similar to that of Proposition 3.3.

The main theorem in this paper is as follows. With the variables \overline{z} and \overline{p} , one has a contact Hamiltonian system.

Theorem 3.8. (Approximate Fokker-Planck equation and contact Hamiltonian system). The time-development of \overline{z} , \overline{p} and q is identified with a contact Hamiltonian system on $T^*\mathbb{R}^n \times \mathbb{R}$, where dynamics of \overline{z} and \overline{p} is governed by the slowest time-scale dynamics.

Proof. Substituting the slowest time-scale mode (29) into the time-derivative of \overline{z} and \overline{p} , one has the closed dynamical system:

$$\frac{\mathrm{d}}{\mathrm{d}t}\overline{z} = -\beta^{-1}\lambda_1 \left\langle \overline{\Phi}_t - 1, \exp(q \cdot B) \right\rangle_{\mathrm{G}} = \beta^{-1}\lambda_1(\psi_{\mathrm{G}}(q) - \overline{z})$$
$$\frac{\mathrm{d}}{\mathrm{d}t}\overline{p}_j = -\beta^{-1}\lambda_1 \left\langle \overline{\Phi}_t - 1, B^j \exp(q \cdot B) \right\rangle_{\mathrm{G}} = \beta^{-1}\lambda_1 \left(\frac{\partial\psi_{\mathrm{G}}}{\partial q^j} - \overline{p}_j \right), \quad j = 1, 2, \dots, n$$

To imitate this dynamics of \overline{z} and \overline{p} by means of contact geometry, one introduces a dynamical system on a contact manifold. More specifically, a contact Hamiltonian system is introduced as follows. Choose \mathcal{H} as a contact Hamiltonian to be [13, 6]

$$\mathcal{H}(\overline{p}, q, \overline{z}) = \gamma_1(\psi_{\rm G}(q) - \overline{z}),\tag{35}$$

with $\gamma_1 \neq 0$ being some constant. From the coordinate expression of the contact Hamiltonian vector field (21), the corresponding contact Hamiltonian system is expressed as

$$\dot{q}^{j} = 0, \quad \dot{\overline{p}}_{j} = \gamma_{1} \left(\frac{\partial \psi_{\mathrm{G}}}{\partial q^{j}} - \overline{p}_{j} \right), \quad \dot{\overline{z}} = \gamma_{1} (\psi_{\mathrm{G}}(q) - \overline{z}), \qquad j = 1, \dots, n.$$

By identifying

 $\gamma_1 = \beta^{-1} \lambda_1,$

one recognizes that the Fokker-Planck equation with the slowest time-scale is exactly same as the contact Hamiltonian system. $\hfill \Box$

Integral curves of the contact Hamiltonian flow in Theorem 3.8 are explicitly written in coordinates as

$$q_j(t) = q_j(0), \qquad \overline{p}_j(t;q) = \frac{\partial \psi_{\rm G}}{\partial q^j}(q) + \left(\frac{\partial \psi_{\rm G}}{\partial q^j}(q) - \overline{p}_j(0;q)\right) {\rm e}^{-\gamma_1 t}, \quad j = 1, \dots, n$$

and

$$\overline{z}(t;q) = \psi(q) + (\psi_{\mathrm{G}}(q) - \overline{z}(0;q)) \,\mathrm{e}^{-\gamma_{1}t}$$

As can be seen from this explicit expression, γ_1 is the natural time-scale (see also Remark 3.1). From the expression of $\overline{p}_j(t;q)$, one can express an approximated expectation value of B_1 , as well as the exact expectation value that has been defined in (20). To express the approximated one for a given $B_1 \in \Gamma \Lambda^1 \mathcal{M}$ at t, introduce

$$\overline{\mathbb{E}}_t : \Gamma \Lambda^1 \mathcal{M} \to \mathbb{R}
B_1 \mapsto \overline{\mathbb{E}}_t [B_1]$$

as

$$\overline{\mathbb{E}}_t[B_1] := \int_{\mathcal{M}} B_1 \overline{\rho}_t \star 1 = \int_{\mathcal{M}} B_1 \rho_G \overline{\Phi}_t \star 1.$$
(36)

From (36) with $\overline{p}_{j}(t;q)$ in (34), one has

$$\overline{\mathbb{E}}_t[B_j] = \overline{p}_j(t;0) = \frac{\partial \psi_{\mathrm{G}}}{\partial q^j}(0) + \left(\frac{\partial \psi_{\mathrm{G}}}{\partial q^j}(0) - \overline{p}_j(0;0)\right) \mathrm{e}^{-\gamma_1 t}, \quad j = 1, \dots, n.$$

In addition, from (33) and (36), $\overline{z}(t;q)$ is interpreted as the approximated moment generating function:

$$\overline{\mathbb{E}}_t \left[\exp(q \cdot B) \right] = \overline{z}(t;q).$$

Remark 3.9. By Theorem 3.8, a procedure is obtained, in which, for a given pair $(h, q) \in \Gamma \Lambda^0 \mathcal{M} \times \mathbb{R}^n$, a contact Hamiltonian $\mathcal{H} \in \Gamma \Lambda^0(T^*\mathbb{R}^n \times \mathbb{R})$ is obtained systematically. This \mathcal{H} is given by

$$\mathcal{H}(\overline{p}, q, \overline{z}) = \beta^{-1} \lambda_1(\psi_{\mathrm{G}}(q) - \overline{z}),$$

where $\psi_{\rm G}(q)$ has been given by (24) as the integral involving *h* over $\mathcal{M}, \overline{z}$ and \overline{p} have been given by (33) and (34) respectively, and λ_1 has been the 1st eigenvalue of $\Delta_{\rm G}$.

Remark 3.10. Given a function $B_1 \in \Gamma \Lambda^0 \mathcal{M}$ that does not depend on t, consider $\mathbb{E}_t[B_1]$ in (20) and $\overline{\mathbb{E}}_t[B_1]$ in (36), where they are unrelated to a contact Hamiltonian system in general. Then from Corollary 2.7 and (31), it follows that the asymptotic limits of these variables coincide:

$$\lim_{t \to \infty} \mathbb{E}_t[B_1] = \lim_{t \to \infty} \overline{\mathbb{E}}_t[B_1] = \langle 1, B_1 \rangle_{\mathrm{G}}.$$

In this subsection the contact geometric description of the Fokker-Planck equation has been discussed, where the variable \overline{z} is chosen to be the moment generating function, and the equilibrium distribution function does not depend on q and B. Meanwhile in the literature dynamics of the minus of a free-energy F, (6), is often considered [13, 18, 7]. To compare the present choice of z with the existing one, the case with z = -F is considered in Section A.4. Note that in [15], another choice of z was studied.

3.3 Dynamics depending on externally applied fields

In Section 3.2, $\Phi_t = \rho_{\rm G}^{-1}\rho_t$ given by (10) is employed to show the relation between the Fokker-Planck equation and a contact Hamiltonian system, where ρ_t obeys (1). The asymptotic limit of ρ_t is the stationary distribution function $\rho_{\rm G}$ given by (8), and $\rho_{\rm G}$ is independent of $q \cdot B$. Here, recall that q is a set of primal variables and and the integrals of B yield the conjugate thermodynamic variables, as mentioned in Section 3.2. This B is interpreted as a set of externally applied fields. Meanwhile, one can consider the case that the stationary distribution depends on $(q \cdot B)$, and it is discussed in this subsection.

Consider the case that the stationary distribution depends on externally applied fields. For this case, one introduces

$$\begin{split} \widetilde{h} &:= h - \beta^{-1}(q \cdot B), \quad \in \Gamma \Lambda^0(\mathcal{M} \times \mathbb{R}^n), \\ \widetilde{\rho}_{\mathrm{G}} &:= \widetilde{Z}_{\mathrm{G}}^{-1} \exp(-\beta \widetilde{h}) = \widetilde{Z}_{\mathrm{G}}^{-1} \exp(-\beta h + q \cdot B), \quad \in \Gamma \Lambda^0(\mathcal{M} \times \mathbb{R}^n), \end{split}$$

where \widetilde{Z}_{G} depends on β and q. The principal role of \widetilde{Z}_{G} is to normalize $\widetilde{\rho}_{G}$ such that

$$\int_{\mathcal{M}} \widetilde{\rho}_{\mathrm{G}} \star 1 = 1.$$

Note that the integral is over \mathcal{M} , not $\mathcal{M} \times \mathbb{R}^n$. In addition, $\star 1$ is a volume form on \mathcal{M} , not on $\mathcal{M} \times \mathbb{R}^n$. The relation,

$$\beta^{-1} \operatorname{d} \widetilde{\rho}_{\mathrm{G}} = -\widetilde{\rho}_{\mathrm{G}} \operatorname{d} h,$$

holds and is used to derive a diffusion equation from the Fokker-Planck equation. The Fokker-Planck equation for $\tilde{\rho}_t \in \Gamma \Lambda^0(\mathcal{M} \times \mathbb{R}^n)$ is

$$\frac{\partial}{\partial t}\widetilde{\rho}_t = -\beta^{-1} d^{\dagger} d\widetilde{\rho}_t - d^{\dagger} \widetilde{\rho}_t d\widetilde{h}$$
$$= -\beta^{-1} d^{\dagger} d\widetilde{\rho}_t - d^{\dagger} \widetilde{\rho}_t (dh - \beta^{-1}q \cdot dB)$$

Notice that the time-evolution of $\tilde{\rho}_t$ depends on q and B, which is dissimilar to the time-evolution of ρ_t .

To derive the corresponding diffusion equation, introduce Φ_t such that

$$\widetilde{\rho}_t = \widetilde{\rho}_{\rm G} \Phi_t$$

and the inner product

$$\left\langle \, \widetilde{\alpha}^{\mathrm{I}} \, , \, \widetilde{\alpha}^{\mathrm{II}} \, \right\rangle_{\widetilde{G}} = \int_{\mathcal{M}} \left(\widetilde{\alpha}^{\mathrm{I}} \wedge \star \widetilde{\alpha}^{\mathrm{II}} \right) \widetilde{\rho}_{\mathrm{G}}$$

for $\widetilde{\alpha}^{\mathrm{I}}, \widetilde{\alpha}^{\mathrm{II}} \in \Gamma \Lambda^{k}(\mathcal{M} \times \mathbb{R}^{n}), k = 0, 1, \dots, m$, such that

$$i_{\partial/\partial q^j} \widetilde{\alpha}^{\mathrm{I}} = i_{\partial/\partial q^j} \widetilde{\alpha}^{\mathrm{II}} = 0, \qquad j = 1, \dots, n.$$

In addition, the weighted Laplacian can be defined as

$$\triangle_{\widetilde{\mathbf{G}}} := \mathbf{d}_{\widetilde{\mathbf{G}}}^{\dagger} \, \mathbf{d},$$

with $d_{\widetilde{G}}^{\dagger} \alpha := \widetilde{\rho}_{G}^{-1} d^{\dagger} \widetilde{\rho}_{G} \alpha$ for all $\alpha \in \Gamma \Lambda^{k}(\mathcal{M} \times \mathbb{R}^{n}), k = 0, 1, \dots, m$. It is then natural to consider the eigenvalue problem:

$$\Delta_{\widetilde{\mathbf{G}}}\phi_s = \lambda_s\phi_s, \qquad s = 0, 1, \dots,$$

and the formal expansion:

$$\widetilde{\Phi}_t = \sum_{s=0}^{\infty} \widetilde{a}_t^s \widetilde{\phi}_s, \qquad \widetilde{\phi}_s \in \Gamma \Lambda^0(\mathcal{M} \times \mathbb{R}^n),$$

with \tilde{a}_t^s being a function $t \mapsto \tilde{a}_t^s$, $\forall s$. Note that, unlike λ_s , $\tilde{\lambda}_s$ depends on q for each s in general. Normalizations are imposed such that

$$\left\langle \widetilde{\phi}_{s}, \widetilde{\phi}_{s'} \right\rangle_{\widetilde{\mathbf{G}}} = \delta_{ss'}. \qquad s, s' = 0, 1, \dots.$$

In addition, one can show that $\lambda_s \geq 0$ for all s by arguing as in Lemma 2.8.

Repeating the derivation of the diffusion equation for Φ_t (see Theorem 2.2), one derives

$$\frac{\partial}{\partial t}\widetilde{\Phi}_t = -\beta^{-1} \triangle_{\widetilde{\mathbf{G}}}\widetilde{\Phi}_t.$$

To see the asymptotic limit of $\tilde{\rho}_t$, let

$$\widetilde{\Upsilon}_t := \frac{\beta}{2} \int_{\mathcal{M}} \mathrm{d}\, \widetilde{\Phi}_t \wedge \widetilde{\rho}_{\mathrm{G}} \star \mathrm{d}\, \widetilde{\Phi}_t.$$

Then applying a similar argument in Proposition 2.6, one has that $\widetilde{\Upsilon}_t$ is a Lyapunov function. By following the same argument as in Corollary 2.7, the asymptotic limit of $\widetilde{\rho}_t$ is $\widetilde{\rho}_G$:

$$\lim_{t \to \infty} \widetilde{\Phi}_t(\widetilde{\xi}) = 1, \quad \text{and} \quad \lim_{t \to \infty} \widetilde{\rho}_t(\widetilde{\xi}) = \widetilde{\rho}_{\mathrm{G}}(\widetilde{\xi}), \quad \forall \widetilde{\xi} \in \mathcal{M} \times \mathbb{R}^n$$

To derive a contact Hamiltonian system, define the slowest time-scale mode $\tilde{\Phi}_t$ as

$$\widetilde{\Phi}_t^{(1)} := 1 + \widetilde{a}_t^1 \widetilde{\phi}_1.$$

This $\widetilde{\Phi}_t^{(1)}$ satisfies

$$\frac{\partial}{\partial t}\widetilde{\Phi}_t^{(1)} = -\beta^{-1}\widetilde{\lambda}_1\left(\widetilde{\Phi}_t^{(1)} - 1\right).$$

Dynamics generated by this PDE is called, by abuse of terminology, the *slowest mode dynamics* in this paper. In addition $\widetilde{\Phi}_t^{(1)}$ satisfies

$$\Delta_{\widetilde{\mathbf{G}}}\widetilde{\Phi}_t^{(1)} = \widetilde{\lambda}_1 \left(\widetilde{\Phi}_t^{(1)} - 1 \right),$$

and the diffusion equation:

$$\frac{\partial}{\partial t}\widetilde{\Phi}_t^{(1)} = -\beta^{-1} \triangle_{\widetilde{\mathbf{G}}} \widetilde{\Phi}_t^{(1)}$$

Since $\widetilde{\lambda}_1 > 0$, this $\widetilde{\Phi}_t^{(1)}$ satisfies

$$\lim_{t \to \infty} \widetilde{\Phi}_t^{(1)}(\widetilde{\xi}) = 1, \quad \forall \widetilde{\xi} \in \mathcal{M} \times \mathbb{R}^n$$

To construct a contact Hamiltonian system, introduce

$$\widetilde{z}^{(1)}(t;q) := \int_{\mathcal{M}} \rho_{\mathrm{G}} \widetilde{\Phi}_{t}^{(1)} \exp(q \cdot B) \star 1 = \left\langle \widetilde{\Phi}_{t}^{(1)}, \exp(q \cdot B) \right\rangle_{\mathrm{G}}, \tag{37}$$

$$\widetilde{p}_{j}^{(1)}(t;q) := \int_{\mathcal{M}} \rho_{\mathrm{G}} \widetilde{\Phi}_{t}^{(1)} B_{j} \exp(q \cdot B) \star 1 = \left\langle \widetilde{\Phi}_{t}^{(1)}, B_{j} \exp(q \cdot B) \right\rangle_{\mathrm{G}}, \quad j = 1, \dots, n.$$
(38)

Note that $\rho_{\rm G} \exp(q \cdot B) = \widetilde{Z}_G \widetilde{\rho}_{\rm G} (\neq \widetilde{\rho}_{\rm G})$ and that the employed inner product has been $\langle , \rangle_{\rm G}$ in (13), not $\langle , \rangle_{\rm \widetilde{G}}$. The reason why $\langle , \rangle_{\rm \widetilde{G}}$ is not employed is that the quantity $\langle B_i, B_j \rangle_{\rm \widetilde{G}}$ depends on t even for time-independent functions B_i and B_j . This t-dependence yields

$$\frac{\mathrm{d}}{\mathrm{d}\,t}\left\langle \,\widetilde{\Phi}_{t}^{(1)} \,,\, B_{j} \,\right\rangle_{\widetilde{\mathrm{G}}} \neq \left\langle \,\dot{\widetilde{\Phi}}_{t}^{(1)} \,,\, B_{j} \,\right\rangle_{\widetilde{\mathrm{G}}} \,, \quad j=1,\ldots,n,$$

which may not lead to a closed equation.

Consider dynamical behavior of $\tilde{z}^{(1)}$ on the z-axis, and that of $\tilde{p}^{(1)}$ on the p-axis. Regarding the timeevolution of (37) and (38), one has the following.

Proposition 3.11. (Approximate Fokker-Planck equation depending on externally applied field). The timedevelopment of $\tilde{z}^{(1)}$ and $\tilde{p}_j^{(1)}$, (j = 1, ..., n) is described on $T^*\mathbb{R}^n \times \mathbb{R}$, where dynamics of $\tilde{z}^{(1)}$ and $\tilde{p}_j^{(1)}$ is governed by the slowest time-scale dynamics. Their asymptotic limits are on the Legendrian submanifold generated by $\psi_{\rm G}$.

Proof. From (37) and (38), one has that

$$\frac{\mathrm{d}}{\mathrm{d}t}\widetilde{z}^{(1)} = \left\langle \dot{\widetilde{\Phi}}_{t}^{(1)}, \exp(q \cdot B) \right\rangle_{\mathrm{G}} = -\beta^{-1}\widetilde{\lambda}_{1} \left\langle \widetilde{\Phi}_{t}^{(1)} - 1, \exp(q \cdot B) \right\rangle_{\mathrm{G}} = \beta^{-1}\widetilde{\lambda}_{1} \left(\psi_{\mathrm{G}}(q) - \widetilde{z}^{(1)} \right),$$

$$\frac{\mathrm{d}}{\mathrm{d}t}\widetilde{p}_{j}^{(1)} = \left\langle \dot{\widetilde{\Phi}}_{t}^{(1)}, B_{j}\exp(q \cdot B) \right\rangle_{\mathrm{G}} = -\beta^{-1}\widetilde{\lambda}_{1} \left\langle \widetilde{\Phi}_{t}^{(1)} - 1, B_{j}\exp(q \cdot B) \right\rangle_{\mathrm{G}}$$

$$= \beta^{-1}\widetilde{\lambda}_{1} \left(\frac{\partial\psi_{\mathrm{G}}}{\partial q^{j}}(q) - \widetilde{p}_{j}^{(1)} \right), \qquad j = 1, \dots, n,$$

where $\psi_{\rm G}$ has been defined in (24). Then it follows from (37) and (38) that

$$\lim_{t \to \infty} \tilde{z}^{(1)}(t;q) = \langle 1, \exp(q \cdot B) \rangle_{\mathrm{G}} = \psi_{\mathrm{G}}(q),$$
$$\lim_{t \to \infty} \tilde{p}^{(1)}_{j}(t;q) = \langle 1, B_{j} \exp(q \cdot B) \rangle_{\mathrm{G}} = \frac{\partial \psi_{\mathrm{G}}}{\partial q^{j}}(q), \quad j = 1, \dots, n$$

This shows that, for a fixed q, the asymptotic limit of a point is on a point of the Legendrian submanifold $\mathcal{L}_{\psi_{G}}(q)$.

Notice that the obtained system of ODEs in Proposition 3.11 is different to that in Theorem 3.8, since the eigenvalue $\tilde{\lambda}_1$ depends on q in general. To see this difference, consider the contact Hamiltonian

$$\widetilde{\mathcal{H}}^{(1)}(\widetilde{p}^{(1)}, q, \widetilde{z}^{(1)}) = \beta^{-1} \widetilde{\lambda}_1(q) \left(\psi_{\mathrm{G}}(q) - \widetilde{z}^{(1)} \right),$$

which is analogous to but different to \mathcal{H} in (35). From (21), one has the contact Hamiltonian system:

$$\dot{\tilde{z}}^{(1)} = \tilde{\mathcal{H}}^{(1)} = \beta^{-1} \tilde{\lambda}_1(q) \left(\psi_{\mathrm{G}}(q) - \tilde{z}^{(1)} \right), \dot{q}^j = 0, \quad \dot{\tilde{p}}_j^{(1)} = \beta^{-1} \tilde{\lambda}_1 \left(\frac{\partial \psi_{\mathrm{G}}}{\partial q^j} - \tilde{p}_j^{(1)} \right) + \beta^{-1} \frac{\partial \tilde{\lambda}_1}{\partial q^j} \left(\psi_{\mathrm{G}}(q) - \tilde{z}^{(1)} \right), \quad j = 1, \dots, n.$$

The differences between $\dot{\tilde{p}}_{j}^{(1)}$ and $\dot{\bar{p}}_{j}$ are

$$\dot{\widetilde{p}}_{j}^{(1)} - \dot{\overline{p}}_{j} = \beta^{-1} \frac{\partial \widetilde{\lambda}_{1}}{\partial q^{j}} \left(\psi_{\mathrm{G}}(q) - \widetilde{z}^{(1)} \right), \quad j = 1, \dots, n,$$

which vanish in the asymptotic limit, $t \to \infty$. From this discussion associated with Theorem 3.11, one notices the following.

Remark 3.12. If $\partial \tilde{\lambda}_1 / \partial q^j = 0$ for all j, then the time-development of $\tilde{z}^{(1)}$ in (37) and $\tilde{p}^{(1)}$ in (38) obeys a contact Hamiltonian system.

3.4 Example

In this subsection the case of $\mathcal{M} = \mathbb{R}$ is considered as an example. After discussing basic properties for this case, a Hamiltonian h is chosen and focused, and its corresponding contact Hamiltonian \mathcal{H} and its variables are explicitly obtained. This analysis demonstrates how the general theory of this paper is applied.

Let x be the coordinate of \mathbb{R} , where x also denotes a point of \mathbb{R} . In addition let the Riemannian metric and its canonical volume-form to be

$$g = \mathrm{d} \, x \otimes \mathrm{d} \, x, \qquad \star 1 = \mathrm{d} \, x$$

The equilibrium state is totally characterized by the Gibbs distribution function in (8). Then the next target is to analyze nonequilibrium states with the Fokker-Planck equation in the present framework. From straightforward calculations, the Fokker-Planck equation (1) is expressed in coordinates as

$$\frac{\partial}{\partial t}\rho_t = \beta^{-1} \frac{\partial^2}{\partial x^2} \rho_t + \frac{\partial}{\partial x} \left(\rho_t \frac{\partial h}{\partial x} \right),$$
$$\frac{\partial}{\partial t}\rho_t = \frac{\partial}{\partial x} \left(\beta^{-1} \frac{\partial}{\partial x} + \frac{\partial h}{\partial x} \right) \rho_t.$$
(39)

The point of departure is to discuss the eigenvalues of $\triangle_{\rm G}$. The eigenvalue problem $\triangle_{\rm G}\phi_s = \lambda_s\phi_s$ for an *s*-th mode is equivalent to

$$-\rho_{\rm G}^{-1} \star^{-1} \mathrm{d} \star \rho_{\rm G} \mathrm{d} \phi_s = \lambda_s \phi_s,$$

which is,

that is

$$d \star \rho_{\rm G} d \phi_s = -\lambda_s \rho_{\rm G} \phi_s \star 1.$$

This reduces further. To ease of notation, ' denotes $\partial/\partial x$, with the use of

$$\rho'_{\rm G} = -\beta h' \rho_{\rm G}, \quad \text{and} \quad \star \,\mathrm{d}\, x = 1,$$

one calculates

$$d \star \rho_{G} d \phi_{s} = d(\star \rho_{G} \phi'_{s} d x) = d(\rho_{G} \phi'_{s}) = (\rho_{G} \phi''_{s} + \rho'_{G} \phi'_{s}) d x$$
$$= (\rho_{G} \phi''_{s} - \beta h' \rho_{G} \phi'_{s}) d x,$$
$$\lambda_{s} \rho_{G} \phi_{s} \star 1 = \lambda_{s} \rho_{G} \phi_{s} d x.$$

Hence the eigenvalue problem is equivalent to

$$\phi_s'' - \beta h' \phi_s' + \lambda_s \phi_s = 0. \tag{40}$$

This equation is a special case of the Strum-Liouville equation, and various theorems are known [45]. For s = 0 with $\lambda_0 = 0$, (40) reduces to

$$\phi_0'' = \beta h' \phi_0'$$

Its solution is found to be

$$\phi_0(x) = 1$$

where $\langle \phi_0, \phi_0 \rangle_G = 1$ is satisfied so that (18) holds for s = s' = 0. The modes ϕ_s with $s \ge 1$ should satisfy $\langle \phi_s, \phi_0 \rangle_G = 0$.

On this manifold, we consider the Hamiltonian on \mathbb{R} ,

$$h(x) = \frac{x^2}{2\mu}.\tag{41}$$

Physically this h is the total energy of a free particle with $\mu > 0$ being mass. In addition, x is interpreted as momentum of the free particle, the metric g measures the norm of the momentum for a unit mass particle, and (41) can be written as

$$h(x) = \frac{1}{2\mu}g\left(\mathfrak{p}^{\sharp}, \mathfrak{p}^{\sharp}\right), \quad \mathfrak{p} = x \,\mathrm{d}\, x = g(\mathfrak{p}^{\sharp}, -) \in T^{*}_{\mathfrak{q}}Q, \quad \mathfrak{p}^{\sharp} = x\frac{\partial}{\partial x} \in T_{\mathfrak{q}}Q, \quad \mathfrak{q} \in Q,$$

with Q being a 1-dimensional manifold. For this Hamiltonian the Gibbs distribution function (8) is

$$\rho_{\rm G}(x) = Z_{\rm G}^{-1} \exp\left(-\beta \frac{x^2}{2\mu}\right), \quad \text{with} \quad Z_{\rm G} = \sqrt{\frac{2\pi\mu}{\beta}},$$

the weighted Laplacian $\triangle_{\mathbf{G}}$ acting on a function f on \mathbb{R} is

$$\Delta_{\mathbf{G}}f = -\rho_{\mathbf{G}}^{-1} \star^{-1} \mathrm{d} \star \rho_{\mathbf{G}} \mathrm{d} f = -\mathrm{e}^{\beta x^{2}/(2\mu)} \frac{\partial}{\partial x} \left(\mathrm{e}^{-\beta x^{2}/(2\mu)} \frac{\partial f}{\partial x} \right) = -\frac{\partial^{2} f}{\partial x^{2}} + \beta \mu^{-1} x \frac{\partial f}{\partial x}$$

and the eigenvalue equation is written as

$$-\phi_s'' + \beta \mu^{-1} x \phi_s' = \lambda_s \phi_s, \quad s = 0, 1, \dots$$

This equation is the Hermite differential equation up to some coefficients, and the solutions for lower eigenvalue labels are

$$\lambda_0 = 0, \quad \phi_0(x) = 1,$$

 $\lambda_1 = \beta \mu^{-1}, \quad \phi_1(x) = \phi_1(1) x, \quad \cdots,$

where $\phi_1(1)$ is the normalization constant. Note that $\langle \phi_0, \phi_0 \rangle_G = 1$ and $\langle \phi_1, \phi_0 \rangle_G = 0$ are satisfied, and $\phi_1(1)$ is determined by $\langle \phi_1, \phi_1 \rangle_G = 1$. The explicit form of $\phi_1(1)$ is obtained with

$$\int_{-\infty}^{\infty} x^2 \,\mathrm{e}^{-\beta x^2/(2\mu)} \,\mathrm{d}\, x = \frac{1}{2} \sqrt{\pi \frac{(2\mu)^3}{\beta^3}},$$

as

$$\phi_1(1) = \left(\frac{\beta^3}{2\pi\mu^3}\right)^{1/4}.$$

The slowest time-scale mode equation (29) is expressed for this model with $\lambda_1 = \beta \mu^{-1}$ as

$$\frac{\partial}{\partial t}\overline{\Phi}_t = -\mu^{-1}(\overline{\Phi}_t - 1).$$

The slowest time-scale mode is then

$$\overline{\Phi}_t(x) = 1 + c x \exp(-\mu^{-1}t)$$
, where $c := \overline{\Phi}_0(1) - 1 = a_0^0 \phi_1(1)$.

This $\overline{\Phi}_t$ satisfies the modified diffusion equation, as stated in Proposition 3.6 and verified from

$$\frac{\partial}{\partial t}\overline{\Phi}_t = -\mu^{-1}c \, x \, \mathrm{e}^{-\mu^{-1}t} \quad \text{and} \quad -\beta^{-1} \triangle_{\mathrm{G}}\overline{\Phi}_t = -\beta^{-1}c \, \mathrm{e}^{-\mu^{-1}t} \left(-\frac{\partial^2 x}{\partial x^2} + \beta \mu^{-1} x \frac{\partial x}{\partial x} \right).$$

In addition, from (32) with this $\overline{\Phi}_t$, one has the approximate distribution function $\overline{\rho}_t = \rho_G \overline{\Phi}_t$:

$$\overline{\rho}_t(x) = Z_{\mathcal{G}}^{-1} \exp(-\beta x^2/(2\mu)) \left[1 + c x \exp\left(-\mu^{-1} t\right)\right],$$

which is a truncation of the exact distribution function $\rho_t(x) = \rho_G \Phi_t(x)$:

$$\rho_t(x) = Z_{\rm G}^{-1} \exp(-\beta x^2/(2\mu)) \left[1 + c x \exp\left(-\mu^{-1} t\right) + \phi_2(x) \exp(-\beta^{-1} \lambda_2 t) + \cdots\right].$$

This $\overline{\rho}_t$ captures the asymmetry $\overline{\rho}_t(-x) - \overline{\rho}_t(x) \neq 0$ at finite t with $x \neq 0$, and shows relaxation $\overline{\rho}_t(-x) - \overline{\rho}_t(x) \rightarrow 0$, $t \rightarrow \infty$. These properties also hold for ρ_t . For a set of some functions $B = (B_1, \ldots, B_n)$ and $q = (q^1, \ldots, q^n) \in \mathbb{R}^n$, let

$$\psi_{\mathcal{G}}(q) = \langle \exp(q \cdot B), 1 \rangle_{\mathcal{G}} = \int_{\mathbb{R}} \exp(q \cdot B(x)) \rho_{\mathcal{G}}(x) \, \mathrm{d} x$$
$$= Z_{\mathcal{G}}^{-1} \int_{\mathbb{R}} \exp\left(-\beta \frac{x^2}{2\mu} + q \cdot B(x)\right) \, \mathrm{d} x.$$

and

$$\overline{z}(t;q) = \langle \overline{\Phi}_t, \exp(q \cdot B) \rangle_{\mathcal{G}} = Z_{\mathcal{G}}^{-1} \int_{\mathbb{R}} \overline{\Phi}_t \exp(-\beta x^2/(2\mu) + q \cdot B) \, \mathrm{d} x,$$

$$\overline{p}_j(t;q) = \langle \overline{\Phi}_t, B_j \exp(q \cdot B) \rangle_{\mathcal{G}}$$

$$= Z_{\mathcal{G}}^{-1} \int_{\mathbb{R}} \overline{\Phi}_t B_j(x) \exp(-\beta x^2/(2\mu) + q \cdot B) \, \mathrm{d} x, \quad j = 1, \dots, n.$$

By Theorem 3.8, the dynamical system for $(\overline{p}, q, \overline{z})$ is written with the contact Hamiltonian

$$\mathcal{H}(\overline{p}, q, \overline{z}) = \mu^{-1}(\psi_{\mathrm{G}}(q) - \overline{z}).$$

In particular choose n = 1 and B = x as a simple case. In this case, one has

$$\psi_{\mathcal{G}}(q) = Z_{\mathcal{G}}^{-1} \int_{-\infty}^{\infty} \exp\left(-\beta \frac{x^2}{2\mu} + qx\right) dx$$
$$= Z_{\mathcal{G}}^{-1} \exp\left(\frac{\mu}{2\beta}q^2\right) \int_{-\infty}^{\infty} \exp\left(-\frac{\beta}{2\mu}(x-\beta^{-1}\mu q)^2\right) dx = \exp\left(\frac{\mu}{2\beta}q^2\right).$$

The asymptotic limit of a point $(\overline{p}, q, \overline{z})$ on $T^*\mathbb{R} \times \mathbb{R}$ is a point on the Legendre submanifold generated by this $\psi_{\mathbf{G}}$:

$$\mathcal{L}_{\psi_{\mathrm{G}}} = \left\{ (\overline{p}, q, \overline{z}) \mid \overline{p} = \frac{\mu}{\beta} q \exp\left(\frac{\mu}{2\beta} q^2\right), \ \overline{z} = \exp\left(\frac{\mu}{2\beta} q^2\right) \right\},$$

Then the approximated expectation value of $B_1 = x$ at t, defined in (36), is calculated as

$$\begin{aligned} \overline{\mathbb{E}}_{t}[x] &= \overline{p}(t;0) = Z_{\mathrm{G}}^{-1} \int_{-\infty}^{\infty} \overline{\Phi}_{t} x \exp(-\beta x^{2}/(2\mu)) \, \mathrm{d} \, x \\ &= \sqrt{\frac{\beta}{2\pi\mu}} \int_{-\infty}^{\infty} \left(1 + cx \,\mathrm{e}^{-\mu^{-1}t}\right) x \,\mathrm{e}^{-\beta x^{2}/(2\mu)} \, \mathrm{d} \, x = c \sqrt{\frac{\beta}{2\pi\mu}} \,\mathrm{e}^{-\mu^{-1}t} \int_{-\infty}^{\infty} x^{2} \,\mathrm{e}^{-\beta x^{2}/(2\mu)} \, \mathrm{d} \, x \\ &= c \mu \beta^{-1} \,\mathrm{e}^{-\mu^{-1}t}, \end{aligned}$$

where $c = a_0^0 \phi_1(1)$ is a constant involving the initial point for the contact Hamiltonian system. The physical interpretation of $\overline{\mathbb{E}}_t[x]$ is, by recalling (41), the average of the momentum x. To obtain more accurate expressions, fast time-scales with $\lambda_2, \lambda_3, \ldots$ should be taken into account.

Consider the case where dynamics involves $q \cdot B$ as discussed in Section 3.3. The most critical parameter is the 1st eigenvalue of $\triangle_{\tilde{G}}$. For the case that n = 1, B = x, and

$$\widetilde{h}(x) = \frac{x^2}{2\mu} - \beta^{-1}qx,$$

one starts with

$$\widetilde{\rho}_{\rm G} = \widetilde{Z}_{\rm G}^{-1} \exp\left(-\beta h + qx\right) = \widetilde{Z}_{\rm G}^{-1} \exp\left(-\frac{\beta}{2\mu}x^2 + qx\right),$$

where the partition function is calculated so that $\tilde{\rho}_{\rm G}$ is normalized as

$$\widetilde{Z}_{\rm G} = \int_{-\infty}^{\infty} \exp\left(-\frac{\beta}{2\mu}x^2 + qx\right) dx = \exp\left(\frac{\mu}{2\beta}q^2\right) \sqrt{\frac{2\pi\mu}{\beta}}.$$

The eigenvalue equation $d_{\widetilde{G}}^{\dagger} d \widetilde{\phi}_s = \widetilde{\lambda}_s \phi_s$, $s = 0, 1, \ldots$, is written in coordinates as

$$-\widetilde{\phi}_{s}^{\prime\prime} + (\beta\mu^{-1}x - q)\widetilde{\phi}_{s}^{\prime} = \widetilde{\lambda}_{s}\widetilde{\phi}_{s}, \qquad s = 0, 1, \dots$$

whose solutions for lower eigenvalue labels are

$$\lambda_0 = 0, \qquad \phi_0(x) = 1$$

$$\widetilde{\lambda}_1 = \beta \mu^{-1}, \qquad \widetilde{\phi}_1(x) = \widetilde{\phi}_1(1) \left(x - \beta^{-1} \mu q \right), \quad \cdots .$$

with $\tilde{\phi}_1(1)$ being a normalization constant. From Remark 3.12 with $\partial \tilde{\lambda}_1 / \partial q = 0$, it follows that the dynamical system for $\tilde{z}^{(1)}$ and $\tilde{p}^{(1)}$ with $B_1(x) = x$ is a contact Hamiltonian system on $T^*\mathbb{R} \times \mathbb{R}$.

4 Discussions and Conclusion

In this paper, it has been shown how the Fokker-Planck equation yields a class of contact Hamiltonian systems. A crucial point in this derivation has been to derive a diffusion equation and to employ several properties of a weighted Laplacian, where the diffusion equation with the weighted Laplacian have been derived from the Fokker-Planck equation on a Riemannian manifold. This Laplacian is self-adjoint with respect to an introduced inner product for forms, and the slowest time-scale eigenfunctions and eigenvalues have been employed to construct closed dynamical equations for expectation values. This set of dynamical equations has been shown to be a contact Hamiltonian vector field.

In the following contact geometric equilibrium and nonequilibrium statistical mechanics are briefly summarized, so that the significance of the claims in this paper is illustrated.

• Equilibrium statistical mechanics provides links between phase space of microscopic dynamical systems and equilibrium thermodynamic phase space. To show this schematically, let \mathcal{M} be phase space of a microscopic dynamical system, x be a set of coordinates of \mathcal{M} , \mathbb{R}^n be the space for externally applied field, $\mathcal{N} = T^*\mathbb{R}^n \times \mathbb{R}$ be thermodynamic phase space, and $\mathcal{L} \subset \mathcal{N}$ be a Legendrian submanifold expressing a set of equilibrium states. Since the space of Hamiltonians is $\Gamma\Lambda^0\mathcal{M}$, one of the roles of equilibrium statistical mechanics is summarized as the diagram,

$$\Gamma\Lambda^0 \mathcal{M} \times \mathbb{R}^n \xrightarrow[\text{statistical mechanics}]{\text{equilibrium}} \mathcal{L} \subset \mathcal{N}, \tag{42}$$

where $A \xrightarrow{C} B$ means that C provides B from A. To express (42) in coordinates, let h be a Hamiltonian, $q \in \mathbb{R}^n$, and \mathcal{L}_{ψ_0} be the Legendrian submanifold generated by some function ψ_0 . Then (42) is expressed in coordinates as

$$(h(x),q) \xrightarrow[\text{statistical mechanics}]{\text{equilibrium}} \mathcal{L}_{\psi_0}(q).$$

$$(43)$$

The diagram (43) is a standard treatment of contact geometric thermodynamics [31].

• Nonequilibrium statistical mechanics provides links between microscopic dynamical systems and nonequilibrium thermodynamic systems. When restricting attention to derivations of macroscopic thermodynamic dynamics from microscopic systems, by identifying dynamical systems with vector fields on manifolds, the role of nonequilibrium statistical mechanics is summarized as the diagram,

$$\Gamma T \mathcal{M} \times \mathbb{R}^n \xrightarrow[\text{statistical mechanics}]{\text{nonequilibrium}} \Gamma T \mathcal{N}.$$
(44)

For the case that \mathcal{M} is a symplectic manifold, a Hamiltonian on \mathcal{M} determines a vector field uniquely, and a contact Hamiltonian on \mathcal{N} determines a vector field uniquely. Note that, by recalling the use of a Hamiltonian h in equilibrium statistical mechanics, this h as an element of $\Gamma \Lambda^0 \mathcal{M}$ need not induce a vector field. When a Hamiltonian and external fields are enough to specify microscopic dynamics, (44) can be expressed as

$$\Gamma\Lambda^{0}\mathcal{M} \times \mathbb{R}^{n} \xrightarrow{\text{nonequilibrium}} \Gamma\Lambda^{0}\mathcal{N}.$$
(45)

In coordinates, when a pair (h, q) is given, (45) is expressed with x for \mathcal{M} as

$$(h(x),q) \xrightarrow[\text{statistical mechanics}]{\text{non-modulibrium}} \mathcal{H}(\overline{p},q,\overline{z}),$$
(46)

where $\mathcal{H} \in \Gamma \Lambda^0 \mathcal{N}$ is a contact Hamiltonian and $(\overline{p}, q, \overline{z})$ is a point of \mathcal{N} .

In this paper, one explicit realization of (46) has been proposed, as has been summarized as Theorem 3.8 with Remark 3.9. One recognizes that one open problem in contact geometric thermodynamics had been to systematically determine a contact Hamiltonian \mathcal{H} from a given h. A solution to this problem has been given in this paper.

There remain unsolved problems that have not been addressed in this paper. They include the following.

- Obtain a contact Hamiltonian from a given dynamical spin system, where phase space of the spin system is discrete. For instance, consider the Glauber system that is a well-studied toy model in nonequilibrium statistical mechanics. Since its dynamics is based on the master equation, and is defined on discrete phase space, the reduction method presented in this paper is inapplicable. An approach for this discrete system is expected to be established by combining the study in Section 3 of this paper and techniques developed in [16], where in [16], a discrete diffusion equation was shown to be derived from a master equation.
- Establish a perturbation theory for Section 3 of this paper, so that the faster time-scale modes of the weighted Laplacian are incorporated order by order.
- Establish a method to obtain contact Hamiltonians from nonlinear Fokker-Planck equations [9], since a natural extension of the linear Fokker-Planck equation is a nonlinear one.
- Extend the existing supersymmetric method for solving the Fokker-Planck equation by means of the present method [37], since the present method may include the so-called supersymmetric method that can be applied to the Fokker-Planck equation.
- Explore a relation between the present study and information geometry [24]. This is because information geometry is compatible with nonequilibrium statistical mechanics [15, 14], clearer relations are expected to be found with the present approach.
- Show how symmetries associated with \mathcal{M} are reflected in \mathcal{N} . In particular for the case when \mathcal{M} is a symplectic manifold, how symplectic reductions in \mathcal{M} are reflected in \mathcal{N} should be clarified [4, 30].
- Extend the present theory to describe systems with phase transitions [17, 18], where singularities are developed on free-energy curves.
- Explore how symplectic topology and contact topology [30] can be applied to the present theory.

By addressing these together with the present study, it is expected that a relevant and sophisticated geometric methodology will be established for dealing with nonequilibrium phenomena.

Data availability statement

No new data were created or analyzed in this study.

Acknowledgments

The author would like to thank Leonid Polterovich for fruitful discussions, comments on this study, and for useful references. These discussions and thoughtful comments helped improve the paper.

ORCID ID

Shin-itiro Goto https://orcid.org/0000-0002-5249-1054

A Appendix

A.1 A simplified proof for Theorem 2.2

The following is another proof for Theorem 2.2.

Proof. Using

$$d \rho_t = \Phi_t d \rho_G + \rho_G d \Phi_t = \Phi_t (-\beta \rho_G d h) + \rho_G d \Phi_t$$

$$\rho_t d h = \rho_G \Phi_t d h,$$

$$\beta^{-1} d \rho_t + \rho_t d h = \beta^{-1} \rho_G d \Phi_t,$$

$$d^{\dagger} (\beta^{-1} d \rho_t + \rho_t d h) = \beta^{-1} d^{\dagger} (\rho_G d \Phi_t),$$

and substituting $\rho_t = \rho_G \Phi_t$ into the Fokker-Planck equation, (1), one obtains that

$$\rho_{\rm G} \frac{\partial}{\partial t} \Phi_t = -\beta^{-1} \,\mathrm{d}^{\dagger} (\rho_{\rm G} \,\mathrm{d}\,\Phi_t).$$

To reduce this equation further, the identity $(d^{\dagger} \rho_{\rm G})\alpha = (\rho_{\rm G} d_{\rm G}^{\dagger})\alpha$, for all $\alpha \in \Gamma \Lambda^k \mathcal{M}, k = 0, 1, \dots, m$ is applied. This holds due to

$$d^{\dagger} \rho_{\rm G} \alpha = \rho_{\rm G} \rho_{\rm G}^{-1} d^{\dagger} \rho_{\rm G} \alpha = \rho_{\rm G} d_{\rm G}^{\dagger} \alpha,$$

where (11) has been employed. Applying the identity with $\alpha = d \Phi_t$, one has that

$$\frac{\partial}{\partial t} \Phi_t = -\beta^{-1} \operatorname{d}_{\mathrm{G}}^{\dagger} \operatorname{d} \Phi_t.$$

This is identical to the modified diffusion equation (14) with the diffusion constant β^{-1} due to $\Delta_{\rm G} = d_{\rm G}^{\dagger} d$.

A.2 Various weighted Laplacians

The operator $\triangle_{\rm G} = d_{\rm G}^{\dagger} d$ in (12), with $d_{\rm G}^{\dagger} = \rho_{\rm G}^{-1} d^{\dagger} \rho_{\rm G}$ is closely related to the so-called Witten Laplacian, and there are several variants of the Witten Laplacian. In this subsection some of them are briefly discussed.

1. Firstly, define d_W as

$$d_{\mathbf{W}}: \Gamma \Lambda^{k} \mathcal{M} \to \Gamma \Lambda^{k+1} \mathcal{M}, \quad k = 0, 1, \dots, m$$
$$\alpha \mapsto d_{\mathbf{W}} \alpha$$

with

$$d_{\rm W} := e^{\beta h} \, \mathrm{d} \, \mathrm{e}^{-\beta h} \,. \tag{47}$$

For example, the action of (47) to a function Φ is

$$d_{W} \Phi = e^{\beta h} d(e^{-\beta h} \Phi) = d \Phi - \beta \Phi d h, \quad \forall \Phi \in \Gamma \Lambda^{0} \mathcal{M},$$

and $d_W d_W \Phi = 0$ can be verified. The corresponding adjoint operator of (47) with respect to \langle , \rangle_G in (13) is denoted by d_W^{\dagger} . Note that if another inner product was chosen, such as the standard one \langle , \rangle in (3), then the following argument would be significantly changed. The adjoint operator of d_W with respect to \langle , \rangle_G is shown simply as

$$\mathbf{d}_{\mathbf{W}}^{\dagger} = \mathbf{d}^{\dagger} \,. \tag{48}$$

To verify (48), let $\alpha^{I} \in \Gamma \Lambda^{k-1} \mathcal{M}$ and $\alpha^{II} \in \Gamma \Lambda^{k} \mathcal{M}$. Assume that the boundary term vanish. Then it is verified by

$$\begin{split} \left\langle \, d_W \, \alpha^I \,, \, \alpha^{II} \, \right\rangle_G &= \int_{\mathcal{M}} (e^{\beta h} \, d \, e^{-\beta h} \, \alpha^I) \wedge \star e^{-\beta h} \, \alpha^{II} = \int_{\mathcal{M}} d(e^{-\beta h} \, \alpha^I) \wedge \star \alpha^{II} \\ &= e^{-\beta h} \, \alpha^I \wedge \star \alpha^{II}|_{\partial \mathcal{M}} + (-1)^k \int_{\mathcal{M}} e^{-\beta h} \, \alpha^I \wedge \star (\star^{-1} \, d \star \alpha^{II}) \\ &= \int_{\mathcal{M}} \alpha^I \wedge \star e^{-\beta h} \, d^\dagger \, \alpha^{II} = \left\langle \, \alpha^I \,, \, d^\dagger \, \alpha^{II} \, \right\rangle_G. \end{split}$$

By extending the standard procedure to define the Laplacian on a Riemannian manifold, one defines $\triangle_W : \Gamma \Lambda^0 \mathcal{M} \to \Gamma \Lambda^0 \mathcal{M}$. This \triangle_W acting on a function is defined as $\triangle_W := d_W^{\dagger} d_W$, which is

$$\triangle_{\mathrm{W}} = \mathrm{d}^{\dagger} \, \mathrm{d}_{\mathrm{W}},$$

It can be shown that eigenvalues of this Witten Laplacian are equal or greater than 0. Both $\triangle_{\rm G}$ and $\triangle_{\rm W}$ can be extended for k-forms if needed.

The explicit form of the Witten Laplacian \triangle_{W} acting on a function ρ is

$$\Delta_{\mathrm{W}}\rho = \mathrm{d}^{\dagger}\,\mathrm{d}_{\mathrm{W}}\,\rho = \mathrm{d}^{\dagger}\,\mathrm{d}\,\rho - \beta\,\mathrm{d}^{\dagger}(\rho\,\mathrm{d}\,h).$$

There is a sign difference between the equation $\dot{\rho} = -\beta^{-1} \Delta_{\rm W} \rho$ and the Fokker-Planck equation (1).

2. Secondly, define

$$D: \Gamma \Lambda^{k} \mathcal{M} \to \Gamma \Lambda^{k+1} \mathcal{M}, \qquad k = 0, 1, \dots, m$$

$$\alpha \mapsto D \alpha$$

with

$$\mathbf{D} := \mathbf{e}^{-\beta h} \, \mathbf{d} \, \mathbf{e}^{\beta h} \,. \tag{49}$$

This is also employed in the literature. The adjoint of D with respect to the standard or unweighted inner product \langle , \rangle in (3) is obtained. Write this adjoint of (49) as D[†], and it is

$$\mathbf{D}^{\dagger} = \mathbf{e}^{\beta h} \, \mathbf{d}^{\dagger} \, \mathbf{e}^{-\beta h} \,. \tag{50}$$

When the boundary term vanishes, D^{\dagger} in (50) is verified to be the adjoint of D for $\alpha^{I} \in \Gamma \Lambda^{k-1} \mathcal{M}$ and $\alpha^{II} \in \Gamma \Lambda^{k} \mathcal{M}$ as

$$\begin{split} \left\langle \operatorname{D} \alpha^{\mathrm{I}}, \, \alpha^{\mathrm{II}} \right\rangle &= \int_{\mathcal{M}} (\mathrm{e}^{-\beta h} \, \mathrm{d} \, \mathrm{e}^{\beta h} \, \alpha^{\mathrm{I}}) \wedge \star \alpha^{\mathrm{II}} = \int_{\mathcal{M}} \mathrm{d} (\mathrm{e}^{\beta h} \, \alpha^{\mathrm{I}}) \wedge \mathrm{e}^{-\beta h} \star \alpha^{\mathrm{II}} \\ &= \int_{\mathcal{M}} \mathrm{d} (\mathrm{e}^{\beta h} \, \alpha^{\mathrm{I}} \wedge \mathrm{e}^{-\beta h} \star \alpha^{\mathrm{II}}) + (-1)^{k} \int_{\mathcal{M}} \mathrm{e}^{\beta h} \, \alpha^{\mathrm{I}} \wedge \star (\star^{-1} \, \mathrm{d} \star \mathrm{e}^{-\beta h} \, \alpha^{\mathrm{II}}) \\ &= \alpha^{\mathrm{I}} \wedge \star \alpha^{\mathrm{II}} |_{\partial \mathcal{M}} + (-1)^{k} \int_{\mathcal{M}} \alpha^{\mathrm{I}} \wedge \star (\mathrm{e}^{\beta h} \star^{-1} \, \mathrm{d} \star \mathrm{e}^{-\beta h} \, \alpha^{\mathrm{II}}) \\ &= \left\langle \alpha^{\mathrm{I}}, \, \mathrm{e}^{\beta h} \, \mathrm{d}^{\dagger} \, \mathrm{e}^{-\beta h} \, \alpha^{\mathrm{II}} \right\rangle. \end{split}$$

There are several ways to express D and D^{\dagger} , including

$$\mathbf{D} = \mathbf{d} + \beta \, \mathbf{d} \, h \wedge, \qquad \mathbf{D}^{\dagger} = \mathbf{d}^{\dagger} + (-1)^m \beta \imath_{(\mathbf{d} h)^{\sharp}},$$

where the following isomorphism has been introduced:

$$\begin{aligned} {}^{\sharp}: \Gamma \Lambda^1 \mathcal{M} & \to & \Gamma T \mathcal{M} \\ \alpha & \mapsto & \alpha^{\sharp}, \end{aligned}$$

with $\alpha^{\sharp} \in \Gamma T \mathcal{M}$ being such that $\alpha = g(\alpha^{\sharp}, -) \in \Gamma \Lambda^1 \mathcal{M}$. The action of D^{\dagger} to a k-form α is calculated from (50) as

$$D^{\dagger} \alpha = e^{\beta h} (-1)^{k} \star^{-1} d(e^{-\beta h} \star \alpha) = (-1)^{k} \star^{-1} e^{\beta h} [e^{-\beta h} (-\beta d h) \wedge \star \alpha + e^{-\beta h} d \star \alpha]$$

= $(-1)^{k} \star^{-1} [(-\beta d h) \wedge \star \alpha + d \star \alpha] = (d^{\dagger} \alpha) + (-1)^{k+1} \beta \star^{-1} (d h \wedge \star \alpha).$

The action of $D^{\dagger}D$ to a function is calculated from

$$D\Phi = d\Phi + \beta \Phi dh, \quad \forall \Phi \in \Gamma \Lambda^0 \mathcal{M}$$

as

$$\begin{split} \mathrm{D}^{\dagger}\,\mathrm{D}\,\Phi &= \mathrm{D}^{\dagger}(\mathrm{d}\,\Phi + \beta\Phi\,\mathrm{d}\,h) = \mathrm{d}^{\dagger}(\mathrm{d}\,\Phi + \beta\Phi\,\mathrm{d}\,h) + \beta\star^{-1}\left[\mathrm{d}\,h\wedge\star(\mathrm{d}\,\Phi + \beta\Phi\,\mathrm{d}\,h)\right] \\ &= \mathrm{d}^{\dagger}\,\mathrm{d}\,\Phi + \beta\,\mathrm{d}^{\dagger}(\Phi\,\mathrm{d}\,h) + \beta\star^{-1}\left(\mathrm{d}\,h\wedge\star\mathrm{d}\,\Phi\right) + \beta^{2}\Phi\star^{-1}\left(\mathrm{d}\,h\wedge\star\mathrm{d}\,h\right) \\ &= \mathrm{d}^{\dagger}\,\mathrm{d}\,\Phi - \beta\star^{-1}\left(\mathrm{d}\,h\wedge\star\mathrm{d}\,\Phi + \Phi\,\mathrm{d}\star\mathrm{d}\,h\right) + \beta\star^{-1}\left(\mathrm{d}\,h\wedge\star\mathrm{d}\,\Phi\right) + \beta^{2}\Phi\star^{-1}\left(\mathrm{d}\,h\wedge\star\mathrm{d}\,h\right) \\ &= \mathrm{d}^{\dagger}\,\mathrm{d}\,\Phi - \beta\Phi\star^{-1}\,\mathrm{d}\star\mathrm{d}\,h + \beta^{2}\Phi\star^{-1}\left(\mathrm{d}\,h\wedge\star\mathrm{d}\,h\right) \\ &= \mathrm{d}^{\dagger}\,\mathrm{d}\,\Phi - \beta\Phi\,\mathrm{d}^{\dagger}\,\mathrm{d}\,h + \beta^{2}\Phi\,\mathrm{d}^{-1}\left(\mathrm{d}\,h\wedge\star\mathrm{d}\,h\right) \\ &= \mathrm{d}^{\dagger}\,\mathrm{d}\,\Phi + \beta\Phi\,\mathrm{d}^{\dagger}\,\mathrm{d}\,h + \beta^{2}\Phi\,g((\mathrm{d}\,h)^{\sharp}, (\mathrm{d}\,h)^{\sharp}), \end{split}$$

where the following have been employed:

$$d^{\dagger} \Phi dh = -\star^{-1} d(\Phi \star dh) = -\star^{-1} (d\Phi \wedge \star dh + \Phi d \star dh) = -\star^{-1} (dh \wedge \star d\Phi + \Phi d \star dh),$$

and

$$\star^{-1}(\mathrm{d}\,h\wedge\star\mathrm{d}\,h) = g((\mathrm{d}\,h)^{\sharp}, (\mathrm{d}\,h)^{\sharp}).$$

Note that it is unclear how $D^{\dagger}\,D$ is exactly corresponding to the Fokker-Planck equation.

A.3 An expression of the Fokker-Planck equation

There are several expressions of (14). In this section one of them is shown. Specifically, a PDE for $\Phi_t = \rho_{\rm G}^{-1} \rho_t$ without use of $\Delta_{\rm G}$ is shown below.

Proposition A.1. The Fokker-Planck equation (1) can be written in terms of Φ_t and $\Delta := d^{\dagger} d$ as

$$\frac{\partial}{\partial t}\Phi_t = -\beta^{-1} \Delta \Phi_t - g((\mathrm{d}\,\Phi_t)^\sharp, (\mathrm{d}\,h)^\sharp),\tag{51}$$

with $\alpha^{\sharp} \in \Gamma T \mathcal{M}$ being such that $\alpha = g(\alpha^{\sharp}, -) \in \Gamma \Lambda^{1} \mathcal{M}$.

Proof. The derivation of (51) is to substitute the assumed form $\rho_t = \rho_G \Phi_t$ into the Fokker-Planck equation (1). The details of calculations are as follows.

Since $\rho_{\rm G} = \exp(-\beta h + c)$ with $c = -\ln Z_{\rm G}$ and $\rho_t = \rho_{\rm G} \Phi_t$ in (8) and (10), one has

$$\mathrm{d}\,\rho_t = \mathrm{e}^{-\beta h + c} (\mathrm{d}\,\Phi_t - \beta \Phi_t \,\mathrm{d}\,h),$$

and then

$$d^{\dagger}(\rho_t d h) = -\star^{-1} d(\rho_t \star d h)$$

= $-\star^{-1} (d \rho_t \wedge \star d h + \rho_t d \star d h)$
= $-\star^{-1} [e^{-\beta h + c} (d \Phi_t - \beta \Phi_t d h) \wedge \star d h + e^{-\beta h + c} \Phi_t d \star d h]$
= $-e^{-\beta h + c} \star^{-1} [(d \Phi_t - \beta \Phi_t d h) \wedge \star d h + \Phi_t d \star d h],$

and

$$\begin{split} \beta^{-1} d^{\dagger} d\rho_t &= -\beta^{-1} \star^{-1} d \star (d\rho_t) = -\beta^{-1} \star^{-1} d[e^{-\beta h + c} \star (d\Phi_t - \beta\Phi_t dh)] \\ &= -\beta^{-1} \star^{-1} \left[-\beta e^{-\beta h + c} dh \wedge \star (d\Phi_t - \beta\Phi_t dh) + e^{-\beta h + c} d \star (d\Phi_t - \beta\Phi_t dh) \right] \\ &= e^{-\beta h + c} \star^{-1} [dh \wedge \star (d\Phi_t - \beta\Phi_t dh) - \beta^{-1} d \star (d\Phi_t - \beta\Phi_t dh)]. \end{split}$$

Summing the 2 terms above, one has that

$$\begin{split} \beta^{-1} d^{\dagger} d\rho_t + d^{\dagger}(\rho_t dh) &= -\beta^{-1} e^{-\beta h + c} \star^{-1} d \star (d\Phi_t - \beta\Phi_t dh) - e^{-\beta h + c} \Phi_t \star^{-1} d \star dh \\ e^{\beta h - c} \left(\beta^{-1} d^{\dagger} d\rho_t + d^{\dagger}(\rho_t dh) \right) &= \beta^{-1} d^{\dagger} (d\Phi_t - \beta\Phi_t dh) + \Phi_t d^{\dagger} dh \\ &= \beta^{-1} d^{\dagger} d\Phi_t - d^{\dagger} \Phi_t dh + \Phi_t d^{\dagger} dh \\ &= \beta^{-1} d^{\dagger} d\Phi_t + \star^{-1} (d\Phi_t \wedge \star dh) \,, \end{split}$$

where in the last line

$$d^{\dagger} (\Phi_t d h) = -\star^{-1} d (\Phi_t \star d h) = -\star^{-1} (d \Phi_t \wedge \star d h + \Phi_t d \star d h) = -\star^{-1} (d \Phi_t \wedge \star d h) + \Phi_t d^{\dagger} d h,$$

has been used. Since

$$\star^{-1}(\mathrm{d}\,\Phi_t\wedge\star\mathrm{d}\,h) = g((\mathrm{d}\,\Phi_t)^{\sharp},(\mathrm{d}\,h)^{\sharp}),$$

the Fokker-Planck equation (1) can be written in terms of Φ_t and $\Delta := d^{\dagger} d$ as (51).

This derived PDE for Φ_t , (51), is the PDE discussed in Section 2 of [28], up to some constant and the convention of the sign for the Laplacian. In [28], the transform $\rho_t \mapsto \Phi_t = \rho_{\rm G}^{-1} \rho_t$ is called the *ground state transform*. In addition to [28], several studies of (51) exist in the literature [41].

To see a property of the right hand side of (51), define

$$\begin{array}{rcl} \mathcal{L}_{\beta,h}:\Gamma\Lambda^{0}\mathcal{M}&\to&\Gamma\Lambda^{0}\mathcal{M}\\ &\Phi&\mapsto&\mathcal{L}_{\beta,h}\,\Phi, \end{array}$$

with

$$\mathcal{L}_{\beta,h} \Phi := \beta^{-1} \operatorname{d}^{\dagger} \operatorname{d} \Phi + \star^{-1} (\operatorname{d} \Phi \wedge \star \operatorname{d} h), \quad \forall \Phi \in \Gamma \Lambda^{0} \mathcal{M},$$
(52)

so that (51) is written as

$$\frac{\partial}{\partial t}\Phi_t = -\operatorname{L}_{\beta,h}\Phi_t.$$

The following is known in [41], and its proof is written with the notation employed in this paper as follows.

Proposition A.2. When the boundary term vanishes, this $L_{\beta,h}$ is self-adjoint with respect to $\langle , \rangle_{G}, L_{\beta,h}^{\dagger} = L_{\beta,h}$: $\langle L_{\beta,h} \Phi^{I}, \Phi^{II} \rangle_{G} = \langle \Phi^{I}, L_{\beta,h} \Phi^{II} \rangle_{G}, \qquad \forall \Phi^{I}, \Phi^{II} \in \Gamma \Lambda^{0} \mathcal{M},$

$$\left\langle \mathbf{L}_{\beta,h} \Psi \right\rangle_{\mathbf{G}} = \left\langle \Psi \right\rangle_{\mathbf{G}}, \quad \forall \Psi , \Psi$$

in addition, the following holds:

$$\left\langle L_{\beta,h} \Phi^{I}, \Phi^{II} \right\rangle_{G} = \beta^{-1} \left\langle d \Phi^{I}, d \Phi^{II} \right\rangle_{G}, \qquad \forall \Phi^{I}, \Phi^{II} \in \Gamma \Lambda^{0} \mathcal{M}.$$

Proof. They are proved as follows. The main part of the proof is to show $\langle L_{\beta,h} \Phi^{I}, \Phi^{II} \rangle_{G} = \beta^{-1} \langle d \Phi^{I}, d \Phi^{II} \rangle_{G}$. After showing this, by exchanging Φ^{I} and Φ^{II} , the proof is completed.

First, one calculates $\langle L_{\beta,h} \Phi^{I}, \Phi^{II} \rangle_{G}$ straightforwardly. From the identity $d(e^{-\beta h+c} \Phi^{II}) = e^{-\beta h+c} (d \Phi^{II} - \beta \Phi^{II} dh)$, it follows that

$$\begin{split} \beta^{-1} \left\langle \operatorname{d}^{\dagger} \operatorname{d} \Phi^{\mathrm{I}}, \, \Phi^{\mathrm{II}} \right\rangle_{\mathrm{G}} &= -\beta^{-1} \int_{\mathcal{M}} (\mathrm{e}^{-\beta h} \, \Phi^{\mathrm{II}}) \operatorname{d} \star \operatorname{d} \Phi^{\mathrm{I}} \\ &= \beta^{-1} \int_{\mathcal{M}} \operatorname{d} (\mathrm{e}^{-\beta h + c} \, \Phi^{\mathrm{II}}) \wedge \star \operatorname{d} \Phi^{\mathrm{I}} - \beta^{-1} \operatorname{e}^{-\beta h + c} \Phi^{\mathrm{II}} \star \operatorname{d} \Phi^{\mathrm{I}}|_{\partial \mathcal{M}} \\ &= \beta^{-1} \int_{\mathcal{M}} \operatorname{d} (\mathrm{e}^{-\beta h + c} \, \Phi^{\mathrm{II}}) \wedge \star \operatorname{d} \Phi^{\mathrm{I}} \\ &= \beta^{-1} \left\langle \operatorname{d} \Phi^{\mathrm{II}}, \, \operatorname{d} \Phi^{\mathrm{I}} \right\rangle_{\mathrm{G}} - \left\langle \Phi^{\mathrm{II}} \operatorname{d} h, \, \operatorname{d} \Phi^{\mathrm{I}} \right\rangle_{\mathrm{G}}, \end{split}$$

and

$$\left\langle \star^{-1} (\mathrm{d} \, \Phi^{\mathrm{I}} \wedge \star \mathrm{d} \, h) \,, \, \Phi^{\mathrm{II}} \right\rangle_{\mathrm{G}} = \int_{\mathcal{M}} \mathrm{e}^{-\beta h + c} \, \Phi^{\mathrm{II}} (\mathrm{d} \, \Phi^{\mathrm{I}} \wedge \star \mathrm{d} \, h) = \int_{\mathcal{M}} \mathrm{d} \, \Phi^{\mathrm{I}} \wedge \star \mathrm{e}^{-\beta h + c} \, \Phi^{\mathrm{II}} \, \mathrm{d} \, h \\ = \left\langle \Phi^{\mathrm{II}} \, \mathrm{d} \, h \,, \, \mathrm{d} \, \Phi^{\mathrm{I}} \right\rangle_{\mathrm{G}} \,.$$

Summing these 2 obtained equations, one has that

$$\left\langle \mathbf{L}_{\beta,h} \Phi^{\mathrm{I}}, \Phi^{\mathrm{II}} \right\rangle = \beta^{-1} \left\langle \mathrm{d} \Phi^{\mathrm{II}}, \mathrm{d} \Phi^{\mathrm{I}} \right\rangle_{\mathrm{G}}$$

Exchanging Φ^{I} and Φ^{II} in the equation above, one has $\langle L_{\beta,h} \Phi^{I}, \Phi^{II} \rangle_{G} = \langle \Phi^{I}, L_{\beta,h} \Phi^{II} \rangle_{G}$.

From Proposition A.2, the eigenvalues of $L_{\beta,h}$ are greater or equal to zero, which can be proved by a similar manner as in the proof of Lemma 2.8.

The relation between $L_{\beta,h}$ and \triangle_{G} is given as follows.

Proposition A.3.

$$\triangle_{\mathbf{G}} \Phi = \beta \operatorname{L}_{\beta,h} \Phi, \qquad \forall \Phi \in \Gamma \Lambda^0 \mathcal{M}.$$

Proof. By straightforward calculations with (12) and (52), one has

$$\begin{split} \triangle_{\mathbf{G}} \Phi &= \mathrm{d}_{\mathbf{G}}^{\dagger} \mathrm{d} \Phi = -\mathrm{e}^{\beta h - c} \star^{-1} \mathrm{d} \left(\mathrm{e}^{-\beta h + c} \star \mathrm{d} \Phi \right) \\ &= -\mathrm{e}^{\beta h - c} \star^{-1} \mathrm{e}^{-\beta h + c} \left(-\beta \, \mathrm{d} \, h \wedge \star \mathrm{d} \, \Phi + \mathrm{d} \star \mathrm{d} \, \Phi \right) \\ &= \beta \star^{-1} \left(\mathrm{d} \, \Phi \wedge \star \mathrm{d} \, h \right) - \star^{-1} \mathrm{d} \star \mathrm{d} \, \Phi \\ &= \beta g \left((\mathrm{d} \, \Phi)^{\sharp}, (\mathrm{d} \, h)^{\sharp} \right) + \mathrm{d}^{\dagger} \, \mathrm{d} \, \Phi \\ &= \beta \, \mathrm{L}_{\beta, h} \, \Phi, \end{split}$$

which is the desired equality.

In the case of compact manifolds, the lower bound of the non-trivial first eigenvalue of $L_{\beta,h}$ was estimated in [10]. Since the eigenvalues of Δ_G coincide with the ones of $\beta L_{\beta,h}$ due to Proposition A.3, this estimate and its related studies are expected to clarify a general property of relaxation processes that take place on compact manifolds.

A.4 Several choices of z

In the literature [6, 18], the case with z = -F is studied, where F is given in (6). In this subsection, for the case of z = -F, it is shown that a construction of a closed dynamical system for \overline{z} is possible, where \overline{z} obeys the slowest time-scale dynamics. In addition $z = \langle \Phi_t, h \rangle_{\rm G}$ is also considered, where h is the microscopic Hamiltonian. By recalling (6), this z is interpreted as internal energy, z = H.

Choose z to be -F, that is expressed in terms of \langle , \rangle_{G} :

$$z := -F = \beta^{-1} \langle \Phi_t, \ln(\rho_{\rm G} \Phi_t) \rangle_{\rm G} + \langle \Phi_t, h \rangle_{\rm G}.$$

With the slowest time-scale dynamics developed in Section 3.2, introduce

$$\overline{z}(t) := \beta^{-1} \left\langle \overline{\Phi}_t \,, \, \ln(\rho_{\rm G} \overline{\Phi}_t) \right\rangle_{\rm G} + \left\langle \overline{\Phi}_t \,, \, h \right\rangle_{\rm G},$$

where $\overline{\Phi}_t$ obeys (28). Notice from Remark 3.10 that

$$\lim_{t \to \infty} z(t) = \lim_{t \to \infty} \overline{z}(t) = \overline{z}(\infty),$$

where

$$\overline{z}(\infty) := \beta^{-1} \langle 1, \ln \rho_{\rm G} \rangle_{\rm G} + \langle 1, h \rangle_{\rm G}$$

To derive an ODE for \overline{z} , with the use of (29), differentiation of \overline{z} with respect to t is calculated as

$$\begin{split} \frac{\mathrm{d}}{\mathrm{d}t}\overline{z} &= \beta^{-1} \left[-\beta^{-1}\lambda_1 \left\langle \overline{\Phi}_t - 1, \ln(\rho_{\mathrm{G}}\overline{\Phi}_t) \right\rangle_{\mathrm{G}} - \beta^{-1}\lambda_1 \left\langle \overline{\Phi}_t - 1, 1 \right\rangle_{\mathrm{G}} \right] - \beta^{-1}\lambda_1 \left\langle \overline{\Phi}_t - 1, h \right\rangle_{\mathrm{G}} \\ &= -\beta^{-1}\lambda_1 \left[\beta^{-1} \left\langle \overline{\Phi}_t, \ln(\rho_{\mathrm{G}}\overline{\Phi}_t) \right\rangle_{\mathrm{G}} + \left\langle \overline{\Phi}_t, h \right\rangle_{\mathrm{G}} - \beta^{-1} \left\langle 1, \ln(\rho_{\mathrm{G}}\overline{\Phi}_t) \right\rangle_{\mathrm{G}} \right] \\ &+ \beta^{-1}\lambda_1 \left\langle 1, h \right\rangle_{\mathrm{G}} - \beta^{-2}\lambda_1 \left\langle \overline{\Phi}_t - 1, 1 \right\rangle_{\mathrm{G}} \\ &= -\beta^{-1}\lambda_1 \left[\overline{z} - \beta^{-1} \left\langle 1, \ln(\rho_{\mathrm{G}}\overline{\Phi}_t) \right\rangle_{\mathrm{G}} \right] + \beta^{-1}\lambda_1 \left\langle 1, h \right\rangle_{\mathrm{G}} - \beta^{-2}\lambda_1 \left\langle \overline{\Phi}_t, 1 \right\rangle_{\mathrm{G}} \\ &= -\beta^{-1}\lambda_1 \left[\overline{z} - \beta^{-1} \left\langle 1, \ln\rho_{\mathrm{G}} \right\rangle_{\mathrm{G}} \right] + \beta^{-1}\lambda_1 \left\langle 1, h \right\rangle_{\mathrm{G}} \\ &= -\beta^{-1}\lambda_1 \left(\overline{z} - \overline{z}(\infty) \right). \end{split}$$

The right hand side in the obtained equation is written in terms of \overline{z} only, and in this sense this equation is a closed system.

As another example, choose $z := \langle \Phi_t, h \rangle_{\mathcal{G}}$, and let

$$\overline{z} := \left\langle \overline{\Phi}_t \,, \, h \right\rangle_{\mathcal{G}} = \overline{\mathbb{E}}_t[h].$$

In this case, one has the closed dynamical system for \overline{z} :

$$\frac{\mathrm{d}}{\mathrm{d}t}\overline{z} = -\beta^{-1}\lambda_1 \left(\overline{z} - \langle 1, h \rangle_{\mathrm{G}}\right).$$

The asymptotic limit of \overline{z} is obtained as $\overline{z} \to \langle 1, h \rangle_{\rm G}, t \to \infty$, which is

$$\lim_{t \to \infty} \overline{E}_t[h] = \int_{\mathcal{M}} h\rho_{\rm G} \star 1.$$

This can be compared with full time-scale dynamics as follows (see also Remark 3.10). Without introducing a contact manifold, the asymptotic limit of full time-scale dynamics is written for $z = \langle \Phi_t, h \rangle_G$, as $z \to \langle 1, h \rangle_G$, $t \to \infty$ from Corollary 2.7. Thus the asymptotic limit of z is the same as that of \overline{z} .

References

- Abraham R, Marsden JE, and Ratiu T 1991 Manifolds, Tensor Analysis, and Applications 2nd Edition, Springer
- [2] Bravetti A 2018 Contact geometry and thermodynamics, Int. J. Geom. Meth. Mod. Phys. 1940003 (51pp)
- [3] Callen HB 1985 Thermodynamics and an introduction to thermostatistics 2nd Eds John Willy and Sons

- [4] Cannas da Silva A 2008 Lectures on Symplectic Geometry Springer
- [5] Choquet-Bruhat Y, DeWitt-Mortte C, and Dillard-Bleick M 1997 Analysis, Manifolds and Physics Part I:Basics Elsevier
- [6] Entov M, and Polterovich L 2023 Contact topology and non-equilibrium thermodynamics Nonlinearity 36 3349-3375
- [7] Entov M, Polterovich L, and Ryzhik L 2023 Geometric aspects of a spin chain arXiv:2311.14205
- [8] Ezra GS 2002 Geometric approach to response theory in non-Hamiltonian systems J. Chem. Math. 32 339-360
- [9] Frank TD 2005 Nonlinear Fokker-Planck equations: fundamentals and applications Springer
- [10] Futaki A, Li Haizhong , and Li Xiang-Dong 2013 On the first eigenvalue of the Witten–Laplacian and the diameter of compact shrinking solitons Ann. Glob. Anal. Geom. 44 105–114
- [11] Gay-Balmaz F, Yoshimura H 2018 Dirac structures in nonequilibrium thermodynamics, J. Math. Phys. 59 012701 (29pp)
- [12] Glauber RJ 1963 Time-dependent statistics of the Ising model J. Math. Phys. 4, 294–307
- [13] Goto S 2015 Legendre submanifolds in contact manifolds as attractors and geometric nonequilibrium thermodynamics, J. Math. Phys. 56 073301 (30pp)
- [14] Goto S 2016 Contact geometric descriptions of vector fields on dually flat spaces and their applications in electric circuit models and nonequilibrium statistical mechanics, J. Math. Phys. 57 102702 (40pp)
- [15] Goto S and H Hino H 2020 Information and contact geometric description of expectation variables exactly derived from master equations, *Phys. Scr.* 95 015207 (14pp)
- [16] Goto S and Hino H 2020 Diffusion equations from master equations A discrete geometric approach J. Math. Phys. 61 113301 (27pp)
- [17] Goto S 2022 Nonequilibrium thermodynamic process with hysteresis and metastable states— A contact Hamiltonian with unstable and stable segments of a Legendre submanifold J. Math. Phys. 63 053302 (25pp)
- [18] Goto S, Lerer S, and Polterovich L 2023 Contact geometric approach to Glauber dynamics near a cusp and its limitation, J. Phys. A 56 125001 (16pp)
- [19] Grmela M 2014 Contact geometry of mesoscopic thermodynamics and dynamics Entropy 16 1652–1686
- [20] Gromov D, and Caines PE 2015 Stability of composite thermodynamic systems with interconnection constraints IET Control Theory & Applications 9 1629-1636
- [21] Haslach HW Jr 1997 Geometric structure of the non-equilibrium thermodynamics of homogeneous systems Rep. Math. Phys. 39 147–62
- [22] Helffer B, and Nier F 2005 Hypoelliptic Estimates and Spectral Theory for Fokker-Planck Operators and Witten Laplacians Springer
- [23] Hsu EP 2002 Stochastic Analysis on Manifolds, American Mathematical Society
- [24] Ito S 2023 Geometric thermodynamics for the Fokker–Planck equation: stochastic thermodynamic links between information geometry and optimal transport *Information geometry* 7 S441–S483

- [25] Jordan R, Kinderlehrer D, and Otto F 1997 Free energy and the Fokker-Planck equation Physica D 107 265-271
- [26] Kubo R, Toda M, and Hashitsume N 1991 Statistical Physics II Springer
- [27] Kubo R 1962 Generalized Cumulant Expansion Method J.Phys.Soc.Jpn 17 1100-1120
- [28] Li XD 2011 Perelman's W-entropy for the Fokker-Planck equation over complete Riemannian manifold, Bull. Sci. Math. 135 871-882
- [29] Lotto J 2008 Some Geometric Calculations on Wasserstein Space, Commun. Math. Phys. 277 423-437
- [30] McDuff D, and Salamon D 2016 Introduction to Symplectic topology 3rd ed, Oxford University Press
- [31] Mrugala R, Nulton JD, Schön JC, and Salamon P 1990 Statistical approach to the geometric structure of thermodynamics, *Phys. Rev. A* 41 3156-3160
- [32] Mrugala R 2000 Geometrical Methods in Thermodynamics. In: Sieniutycz, S., De Vos, A. (eds) Thermodynamics of Energy Conversion and Transport Springer
- [33] Nishimori H 2001 Statistical Physics of Spin Glasses and Information Processes Oxford University Press
- [34] Olver PJ 1986 Applications of Lie Groups to Differential Equations Springer
- [35] Pavliotis GA 2014 Stochastic Processes and Applications Springer
- [36] Risken H 1989 The Fokker-Planck Equation Second Edition Springer
- [37] Rosu HC 1997 Supersymmetric Fokker-Planck strict isospectrality Phys. Rev. E 56 2269-2271
- [38] Ruelle D 1999 Smooth Dynamics and New Theoretical Ideas in Nonequilibrium Statistical Mechanics J. Stat. Phys. 95, 393-468
- [39] Salazar DSP 2023 Moment generating function bound from detailed fluctuation theorem arxiv:2302.02998
- [40] van der Schaft A, and Maschke B 2018 Geometry of Thermodynamic Processes Entropy 20 925 (23pp)
- [41] Markowich PA, and Villani C 2000 On the trend to equilibrium for the Fokker-Planck equation: an interplay between physics and functional analysis *Mat. Contemp* 19 1-29
- [42] Villani C 2008 Optimal transport, old and new Springer
- [43] Vu TV, and Saito K 2023 Thermodynamic Unification of Optimal Transport: Thermodynamic Uncertainty Relation, Minimum Dissipation, and Thermodynamic Speed Limits Phys. Rev. X 13 011013 (45pp)
- [44] Zwanzig R 2001 Nonequilibrium statistical mechanics Oxford University Press
- [45] Zetti A 2005 Sturm-Liouville Theory American Mathematical Society