

# Spatially oscillating correlation functions in $(2 + 1)$ -dimensional four-fermion models: The mixing of scalar and vector modes at finite density

Marc Winstel\*

*Institut für Theoretische Physik, Goethe-Universität,  
Max-von-Laue-Straße 1, D-60438 Frankfurt am Main, Germany.*

(Dated: March 13, 2024)

In this work, we demonstrate that the mixing of scalar and vector condensates produces spatially oscillating, but exponentially damped correlation functions in fermionic theories at finite density and temperature. We find a regime exhibiting this oscillatory behavior in a Gross-Neveu-type model that also features vector interactions within the mean-field approximation. The existence of this regime aligns with expectations based on symmetry arguments, that are also applicable to QCD at finite baryon density. We compute the phase diagram including both homogeneous phases and regions with spatially oscillating, exponentially damped correlation functions at finite temperature and chemical potential for different strengths of the vector coupling. Furthermore, we find that inhomogeneous condensates are disfavored compared to homogeneous ones akin to previous findings without vector interactions [1]. We show that our results are valid for a broad class of  $(2 + 1)$ -dimensional models with local four-fermion interactions.

Keywords: Scalar-vector mixing, stability analysis, quantum pion liquid, oscillating meson propagators, inhomogeneous phases, phase diagram, mean-field

## CONTENTS

I. Introduction	2
II. Fermionic model with scalar-vector mixing	3
A. Mean-field approximation	4
III. Methods for computing the phase diagram in the presence of mixing	4
A. Homogeneous condensates	4
B. Renormalization and parameter fixing	5
C. The Hessian matrix and bosonic two-point functions in the presence of mixing	6
D. Structure of the Hessian and its physical consequences	7
IV. Phase diagram in the presence of mixing	8
A. Homogeneous condensation	8
B. Stability analysis with mixing	11
1. Symmetric phase	11
2. Complex conjugate eigenvalues for $q = 0$	11
3. Complex-conjugate eigenvalue pairs emerging at $q \neq 0$	15
V. Implications for general four-fermion models in $(2 + 1)$ dimensions	17
VI. Conclusions	18
VII. Acknowledgments	19
A. Conventions for the Wick rotation	19
B. Formulas for the stability analysis with mixing	20
1. Finite temperature expressions for non-vanishing $q$	20
2. Finite temperature expressions for $q = 0$	22
References	23

---

\* [winstel@itp.uni-frankfurt.de](mailto:winstel@itp.uni-frankfurt.de)

## I. INTRODUCTION

Quantum field theories (QFTs) with fermionic four-point interactions – so-called four-fermion (FF) models – and Yukawa models are often used as qualitative descriptions of fermionic matter in various branches of physics, mostly high-energy and condensed matter physics [2–36]. An important application is the phenomenological low-energy description of chiral symmetry breaking in Quantum Chromodynamics (QCD) at finite temperature  $T$  and chemical potential  $\mu$ . Although originally designed as models for the strong interaction between nucleons, a common example for QCD-inspired models are the Nambu-Jona-Lasinio (NJL) and the quark-meson (QM) model, which describe the interaction of fermions through the exchange of light mesons, such as, e.g., pions [37–39]. When formulated in the chiral limit, i.e., without a bare mass term for the fermions, these models typically feature an homogeneous broken phase (HBP) with a chiral condensate  $\langle\bar{\psi}\psi\rangle \neq 0$  acting as a dynamically generated mass term in the vacuum. Above a critical temperature, a second-order phase transition from the HBP to the symmetry-restored phase (SP) occurs where chiral symmetry is restored – in consistency with the chiral cross-over of QCD with physical quark masses<sup>1</sup> [41]. The effective model description is of particular relevance for phenomenological predictions at non-vanishing density, since QCD suffers from complex Boltzmann weights appearing in the partition function at  $\mu \neq 0$  [42]. The complex weights in the partition function significantly complicate first-principle calculations, e.g, using lattice field theory. Therefore, the development of techniques with access to the QCD phase diagram in the  $(\mu, T)$  plane is an active field of research [43–50].

Thus, except for regions of the QCD phase diagram at  $\mu = 0$  and small  $\mu/T$ , theorists have relied on effective models for the study of the phase diagram of strongly-interacting matter at intermediate densities and temperatures, see, e.g., Refs. [2, 4]. Model studies are often carried out in the mean-field approximation where bosonic quantum fluctuations are suppressed – in order to reduce the complexity of the calculations. With respect to the chiral phase transition, computations assuming homogeneous ground states typically discuss the scenario of a first-order phase transition including a critical end point (CEP) at non-vanishing  $\mu$  – in consistency with investigations of QCD using functional methods [43, 44]. When allowing for inhomogeneous condensates in these model calculations the first order phase transition region is replaced by the alternative scenario of a so-called inhomogeneous phase (IP) – a phase where the chiral condensate is a function of the spatial coordinates, i.e,  $\langle\bar{\psi}\psi\rangle = f(\mathbf{x})$ , and translational symmetry is spontaneously broken (see Ref. [51] for a review). A phenomenon expected to be closely related to the IP is the so-called moat regime – where a negative bosonic wave-function renormalization is obtained. Accordingly, a modified dispersion relation with a minimal energy at a non-vanishing momentum is obtained [52], which is the reason that the moat regime is also called a precursor phenomenon of an IP, see also Ref. [53] for an example of a  $(1+1)$ -dimensional model featuring a moat regime which is present in large parts of the IP and the SP. In general, spatially oscillatory behavior of quantities that are related to particles with the moat dispersion relation should be favored within a moat regime. Scenarios, which are alternatives to translational symmetry breaking in an IP, include a liquid crystal-like behavior [54–59], where correlations are oscillatory and of quasi-long range order with polynomial suppression, and a so-called ‘quantum pion liquid’ with oscillatory, but exponentially suppressed two-point functions [60–63]. Both of these scenarios are related to some sort of disordering of the IP through bosonic quantum fluctuations. The observation of the moat regime in a recent Functional Renormalization Group study [44] as well as a few other, albeit limited and exploratory studies of IPs in QCD [64, 65] suggest that IPs or, in general, regimes with spatially oscillatory behavior are relevant in QCD at finite density.

Inhomogeneous phases are very common in  $(1+1)$ -dimensional models at least within the mean-field approximation [53, 66–72]. However, in recent literature there is an on-going discussion whether IPs persist when allowing for bosonic quantum fluctuations [73–79]. In  $(3+1)$  dimensions, IPs have been observed in various models over the last decades [80–86]. However, recent studies [87, 88] show an inherit dependence of the IP on the regulator value and the choice of the regularization scheme in the NJL model caused by the chemical potentials and momenta of the inhomogeneous condensates being in the order of the regulator. Thus, one can argue that there is no predictive power of the NJL model results with respect to IPs<sup>2</sup>, as long as there is no good argument that an effective theory for QCD at finite density could behave as a low-dimensional or strongly-regulated NJL model [89, 90]. In contrast, the moat regime seems to be a stable feature of the NJL model [88] – as will be reported on in an upcoming publication following Ref. [88].

Due to the non-renormalizability of FF models in  $(3+1)$  dimensions and the problems arising from this non-renormalizability as well as due to the application to condensed matter systems, it is common to study these models in  $(2+1)$  dimensions [91–101], where FF models are renormalizable. In a preceding work [1], we have shown the absence

<sup>1</sup> In the case of formulating these models with a non-vanishing bare quark mass, the chiral symmetry is also only approximate and a crossover is observed instead of a second-order phase transition. The explicit symmetry breaking can then be compensated at high temperature and chemical potential in a super restoration phase boundary, as found in Ref. [40].

<sup>2</sup> It is probably necessary to explore the regularization dependence of the IP in models where the regulators can be tuned to higher values than in the NJL – such as, e.g., the QM model.

of IPs and moat regimes in a variety of FF models and Yukawa models by analyzing the stability of homogeneous ground states. These results are in consistency with previous findings that minimize the effective action of the  $(2+1)$ -dimensional Gross-Neveu (GN) model on the lattice [102–105] and generalizes this result to the whole class of models with Lorentz-scalar interaction channels.

In this work, we expand on the analysis in Ref. [1] by including vector interactions  $\sim (\bar{\psi}\gamma_\nu\psi)^2$ . In contrast to nuclear matter, where the Walecka-model – featuring interactions with vector mesons – is commonly used, the inclusion of vector interactions is often not part of effective model calculations of QCD matter except for a few studies [106–110]. However, the fundamental QCD action gives rise to Yukawa-type interactions between fermions and vector mesons, such as the  $\omega$  meson, arising from a resonance of the FF interaction in the corresponding vector channels [111–113]. At non-vanishing  $\mu$ , the temporal component of the  $\omega$  meson couples directly to the density  $\langle\psi^\dagger\psi\rangle$  via the Yukawa interaction  $\sim\psi^\dagger\omega_0\psi$ . Thus, this interaction is expected to play an important role in finite density QCD.

In the present work, we focus on the effects of mixing between scalar and vector condensates on the  $(\mu, T)$  phase diagram by analyzing the stability of homogeneous ground states. For this sake a FF model including both scalar and vector interactions is defined in Section II. The renormalization of the FF interaction and the methods for the computation of homogeneous condensates are briefly described in Section III A and Section III B. The stability analysis including the computation of the Hessian matrix in field space – featuring mixing of different condensates – is presented in Section III C. Also, we discuss further implications of the analysis regarding oscillatory correlations functions in Section III D and its relation to symmetries realized in finite density QCD. In Section IV A, we present original results on the homogeneous phase diagram of the theory. The appearance of oscillatory correlation functions in a quantum pion liquid regime as well as the absence of IPs are discussed in Section IV B. In Section V, we show that our results are generic for all models with local FF interactions based on a similar argumentation as in Ref. [1] before concluding in Section VI.

## II. FERMIONIC MODEL WITH SCALAR-VECTOR MIXING

In order to study mixing induced by the interplay of a repulsive vector and an attractive scalar interaction, we study the action

$$\mathcal{S}_{\text{mix}}[\bar{\psi}, \psi] = \int_0^\beta d\tau \int d^2x \left\{ \bar{\psi} (\not{\partial} + \gamma_3\mu) \psi - \left[ \frac{\lambda_S}{2N} (\bar{\psi}\psi)^2 + \frac{\lambda_V}{2N} \left( (\bar{\psi}i\gamma_3\psi)^2 + (\bar{\psi}i\vec{\gamma}\psi)^2 \right) \right] \right\}, \quad (1)$$

where  $\psi$  contains  $N$  four-component spinors, describing a generic fermion field without a bare mass,  $\mu$  is the chemical potential and  $\gamma_\nu = (\gamma_3, \vec{\gamma})$  are the  $4 \times 4$  Dirac matrices<sup>3</sup>, where  $\vec{\gamma} = (\gamma_1, \gamma_2)$ . The spacetime integration goes over a  $(2+1)$ -dimensional Euclidean spacetime volume, where the inverse temperature determines the temporal extent  $\beta = 1/T$ . Our conventions for the Wick rotation are given in Appendix A.

Introducing auxiliary bosonic fields in an inverse, shifted Gaussian integration yields the partially bosonized model

$$\mathcal{S}[\bar{\psi}, \psi, \sigma, \omega_\nu] = \int d^3x \left[ \bar{\psi} Q \psi + \frac{\omega_\nu\omega_\nu}{2\lambda_V} + \frac{\sigma^2}{2\lambda_S} \right], \quad Z = \int \prod_{\phi=\{\bar{\psi}, \psi, \sigma, \omega_\nu\}} \mathcal{D}\phi e^{-S[\bar{\psi}, \psi, \sigma, \omega_\nu]} \quad (2)$$

with the Dirac operator

$$Q[\sigma, \omega_\nu] = \gamma_\nu\partial_\nu + \gamma_3\mu + \sigma + i\gamma_\nu\omega_\nu. \quad (3)$$

The scalar field  $\sigma$  and the vector field  $\omega_\nu = (\omega_3, \vec{\omega})$  are linked to fermionic expectation values via

$$\langle\sigma\rangle(x) = -\frac{\lambda_S}{N} \langle\bar{\psi}\psi\rangle(x), \quad \langle\omega_3\rangle(x) = -\frac{\lambda_V}{N} i \langle\bar{\psi}\gamma_3\psi\rangle(x), \quad \langle\vec{\omega}\rangle(x) = -\frac{\lambda_V}{N} i \langle\bar{\psi}\vec{\gamma}\psi\rangle(x). \quad (4)$$

Assuming that the theories' invariance under rotations of the spatial coordinates  $\mathbf{x} = (x_1, x_2)$  remains intact, only condensation of the scalar field  $\sigma$  and the temporal component of the vector field  $\omega_3$  is allowed. Thus, this assumption implies that  $\langle\vec{\omega}\rangle = 0$ . In order to study IPs, a stability analysis about homogeneous condensates is applied, which also respects the invariance of the theory under spatial rotations, and no assumptions on the functional form of  $\sigma(\mathbf{x})$  or  $\omega_\nu(\mathbf{x})$  are made. A non-vanishing expectation value of  $\sigma$  indicates the breaking of the discrete chiral symmetry of this model [91, 103, 104], since it acts as a dynamically generated mass term for the fermions. This mass term is

<sup>3</sup> These fulfill the Clifford algebra with the Euclidean metric  $\text{diag}(1, 1, 1)$ .

parity-even, while there is also the possibility for a parity-odd mass term in  $(2+1)$  dimensions [1, 114]. Thus, we often refer to  $\langle\sigma\rangle$  as the chiral condensate, since it is directly linked through Eq. (4) to  $\langle\bar{\psi}\psi\rangle$ . Also, we note that in our conventions (compare Appendix A) the quark number density is given by  $n(x) = -\langle\bar{\psi}\gamma_3\psi\rangle(x)/N$  and, thus,  $\omega_3$  is required to be a purely imaginary field with  $\text{Im}\omega_3(x) = \lambda_V n(x)/N$  in order for the baryon density to be real-valued. This is directly linked to the repulsive nature of the Yukawa interaction between quarks and the vector meson  $\omega_3$  and the corresponding FF interaction.

### A. Mean-field approximation

For the remainder of this work, we use the so-called mean-field approximation, which in this context means the suppression of bosonic quantum fluctuations in the partition function (2). In the case of FF models such as Eq. (1), this suppression can directly be obtained by taking the limit of the parameter  $N \rightarrow \infty$ . Since the fermion fields appear only as bilinears in Eq. (2), one can integrate their fluctuations out. The obtained effective action<sup>4</sup>

$$\frac{1}{N}\mathcal{S}_{\text{eff}}[\sigma, \omega_\nu] = \int d^3x \left[ \frac{\omega_\nu\omega_\nu}{2\lambda_V} + \frac{\sigma^2}{2\lambda_S} \right] - \text{Tr} \ln Q \quad (5)$$

is then a functional of the bosonic fields only. In the mean-field approximation, observables can be computed by evaluating them on the global minimum of the effective action<sup>5</sup>, denoted by  $\sigma = \bar{\Sigma}$  and  $\omega_\nu = \bar{\Omega}_\nu$ . Since the interactions do not mix between the  $N$  fermion fields, the effective action is proportional to  $N$  and the minimization of  $\mathcal{S}_{\text{eff}}$  is independent of  $N$ . Hence, the obtained phase diagram would be identical when using the mean-field method as a semi-classical approximation of the QFT with any integer number of  $N$ . The mean-field approximation has turned out to be a decent starting point for investigations of phase diagrams, since ordered phases as, e.g., an HBP or an IP, are typically weakened through bosonic quantum fluctuations, see, e.g., Refs. [73–77, 100]. Thus, if an HBP or an IP does not exist in the mean-field approximation, it is very likely that those phases also do not exist in the full QFT.

## III. METHODS FOR COMPUTING THE PHASE DIAGRAM IN THE PRESENCE OF MIXING

In this chapter, we present methods for computing the phase diagram of the FF model with mixing, see Eq. (1).

### A. Homogeneous condensates

As a starting point for the investigation of the phase structure of the model (1), the thermodynamic ground state with the assumption of homogeneous condensates  $\sigma = \bar{\sigma}$ ,  $\omega_3 = \bar{\omega}_3$  is determined. We assume that the spatial components of the vector field respect the invariance of the partition function (2) under spatial rotations, i.e.  $\vec{\omega} = 0$ . With this assumption one can compute the homogeneous effective potential at fixed  $\mu$  and  $T$

$$\bar{U}^{(\mu, T)}(\bar{\sigma}, \bar{\omega}_3) = \frac{1}{N} \frac{\mathcal{S}_{\text{eff}}[\bar{\sigma}, \bar{\omega}_3]}{\beta V} \quad (6)$$

using standard techniques of thermal field theory. Inspecting Eq. (2) under the assumption of  $\omega_3 = \bar{\omega}_3$ , it is convenient to absorb the vector condensate into an effective chemical potential  $\bar{\mu} = \mu + i\bar{\omega} = \mu - \lambda_V \bar{n}$ , where  $\bar{n}$  is the homogeneous quark number density per fermion species, see the discussion below the Ward identity (4). The homogeneous effective potential is given by

$$\bar{U}^{(\mu, T)}(\bar{\sigma}, \bar{\omega}_3) = \frac{\bar{\omega}_3^2}{2\lambda_V} + \frac{\bar{\sigma}^2}{2\lambda_S} - \frac{N_\gamma}{2} \int \frac{d^2p}{(2\pi)^2} \left\{ E + \frac{1}{\beta} \left[ \ln \left( 1 + e^{-\beta(E+\bar{\mu})} \right) + \ln \left( 1 + e^{-\beta(E-\bar{\mu})} \right) \right] \right\}, \quad (7)$$

where the last two terms are identical to the effective potential of the GN model at chemical potential  $\bar{\mu}$  and temperature  $T$ , as, e.g., studied in Refs. [53, 91, 92, 102, 103]. In Eq. (7), summation over fermionic Matsubara frequencies

<sup>4</sup> In the mean-field approximation, the effective action is directly proportional to the quantum effective action, usually defined as the Legendre transform with respect to of the logarithm of the partition function.

<sup>5</sup> In the case of multiple, degenerate global minima, which are linked through a symmetry transformation, one has to formally introduce a small symmetry breaking parameter  $h$  and extrapolate to a vanishing  $h$ . In the mean-field approximation, this can be directly implemented by picking one of the degenerate minima. In the case of obtaining a point in the phase diagram with a first order phase transition, where this issue can lead to ambiguities, we will refrain from evaluating any observables depending on the minimum of the effective action. We refer to Ref. [1] for a similar discussion.

was already performed and  $E = \sqrt{\vec{p}^2 + \bar{\sigma}^2}$ , where  $\vec{p}$  are the spatial momenta. The dimensionality of the spinors is denoted by  $N_\gamma = 4$ .

The computation of homogeneous condensates is standard in the literature, see, e.g., Refs. [51, 53, 66, 91, 115]. In order to determine the extrema of the effective potential (7) with respect to  $\bar{\sigma}$  and  $\bar{\omega}_3$ , we use the so-called gap equations

$$\frac{\partial \bar{U}^{(\mu, T)}}{\partial \bar{\sigma}} = 0 \leftrightarrow \frac{\bar{\sigma}}{\lambda_S} - \bar{\sigma} \frac{N_\gamma}{2} \int \frac{d^2 p}{(2\pi)^2} \frac{1 - n_F(E) - n_{\bar{F}}(E)}{E} \equiv \bar{\sigma} \left( \frac{1}{\lambda_S} - \ell_1(\mu, T, \bar{\sigma}, \bar{\omega}_3) \right) = 0 \quad (8)$$

and

$$\frac{\partial \bar{U}^{(\mu, T)}}{\partial \bar{\omega}} = 0 \leftrightarrow \frac{\bar{\omega}_3}{\lambda_V} = \frac{N_\gamma}{2} i \int \frac{d^2 p}{(2\pi)^2} [n_F(E) - n_{\bar{F}}(E)] \equiv i\bar{n}(\mu, T, \bar{\sigma}, \bar{\omega}_3), \quad (9)$$

where we defined the Fermi distributions  $n_F(x) = (1 + e^{\beta(x - \bar{\mu})})^{-1}$  and  $n_{\bar{F}}(x) = (1 + e^{\beta(x + \bar{\mu})})^{-1}$ . Solving this set of coupled equations, one computes all extrema of  $\bar{U}$  with respect to  $\bar{\sigma}$  and  $\bar{\omega}_3$  and determines the global minimum, denoted as  $(\bar{\Sigma}(\mu, T), \bar{\Omega}_3(\mu, T))$ , respectively, by inserting the all found extrema back into the effective potential (7).

The absorption of  $\bar{\omega}_3$  into a chemical potential  $\bar{\mu}$  can be useful for the interpretation of results as well as for computational purposes. As there is no way to distinct the dynamical contribution to the chemical potential through the condensation of the vector field  $\bar{\Omega}_3$  experimentally,  $\bar{\mu}$  is also the phenomenologically relevant quantity. Thus, one could argue to use  $\bar{\mu}$  and  $T$  as external parameters of the model and, instead, treat the effective potential as a function of  $\bar{\sigma}$  and  $\mu$ , such as done in Refs. [106, 107]. This is also advantageous for the minimization of the effective potential Eq. (7), since one can directly solve Eq. (8) at fixed  $\bar{\mu}$  and  $T$ . Inserting the solution  $\bar{\Sigma}$  into Eq. (9), it is possible to directly compute  $\bar{\Omega}_3$  and, thus, the chemical potential from  $\mu = \bar{\mu} - i\bar{\Omega}_3$ . However, this procedure leads to ambiguities in the  $(\mu, T)$  phase diagram, as multiple  $(\bar{\Sigma}, \bar{\mu})$  pairs can belong to the same chemical potential  $\mu$ . Then, one would need to iterate through all computed tuples  $(\bar{\sigma}, \bar{\mu})$  to get the correct global minimum of  $\bar{U}$  for a certain  $(\mu, T)$  and draw a correct picture of the phase diagram. Thus, in our numerical setup we keep  $\bar{\sigma}$  and  $\bar{\omega}_3$  as variables.

## B. Renormalization and parameter fixing

As can be seen from Eq. (8), the computation of homogeneous condensates in the model (1) involves the evaluation of integrals with linear ultra-violet (UV) divergences that require regularization. As originally demonstrated in Refs. [92, 116], FF interactions are renormalizable order by order in  $1/N$  in  $(2+1)$  dimensions. Thus, we can remove the divergences by imposing a renormalization condition. By fixing the dynamically generated fermion mass in the vacuum  $\bar{\Sigma}(\mu = 0, T = 0) = \bar{\Sigma}_0$ , one can absorb the divergence using the coupling  $\lambda_S$  through the gap equation (8) in the vacuum. Using a sharp UV cutoff  $\Lambda$  for the spatial loop momenta, we determine

$$\ell_1(0, 0, \bar{\sigma}, 0) = \frac{N_\gamma}{4\pi} (\Lambda - |\bar{\sigma}|) \quad (10)$$

such that

$$\frac{1}{\lambda_S} = \frac{N_\gamma}{4\pi} (\Lambda - |\bar{\Sigma}_0|). \quad (11)$$

Note that we used that  $\bar{\omega}_3 = 0$  in the vacuum, as can be read off directly from Eq. (9). Using this procedure and sending  $\Lambda \rightarrow \infty$ , we obtain

$$L_1(\mu, T, \bar{\sigma}, \bar{\omega}_3) \equiv \frac{1}{\lambda_S} - \ell_1(\mu, T, \bar{\sigma}, \bar{\omega}_3) = N_\gamma \left[ \frac{|\bar{\sigma}| - |\bar{\Sigma}_0|}{4\pi} + \int \frac{d^2 p}{(2\pi)^2} \frac{n_F(E) + n_{\bar{F}}(E)}{2E} \right], \quad (12)$$

which is finite and is all that is required to render the observables of interest in this work finite. Thus, the chiral condensate in the vacuum  $\bar{\Sigma}_0$  will be used as the physical scale to construct dimensionless ratios for all other quantities in this work.

Since the gap equation for  $\bar{\omega}_3$  (9) vanishes in the vacuum,  $\lambda_V$  can only be fixed by working at  $\mu \neq 0$ , e.g., by imposing a value for  $\bar{\omega}_3$  at a non-vanishing  $\mu$  and  $T = 0$ . Alternatively, we could tune  $\lambda_V$  to the inverse mass of the physical isoscalar vector meson, as is common in phenomenological models, see, e.g., Ref. [110]. Such a treatment would, however, be inconsistent compared to the treatment of  $\lambda_S$ , which cannot be tuned to a phenomenological value for  $m_\sigma$  because of the renormalization procedure, where  $1/\lambda_S \sim \Lambda$ . Since we consider the action (1) as a toy model for the mixing of scalar and vector modes, we prefer not to apply a phenomenologically motivated parameter setting. Instead, we do not fix  $\lambda_V$  but treat it as a free parameter to study its influence on the phase diagram of the theory similar to, e.g., Refs. [106, 107].

### C. The Hessian matrix and bosonic two-point functions in the presence of mixing

In order to compute the phase structure of the theory, the ground state  $(\sigma, \omega_\nu) = (\Sigma(\mathbf{x}), \Omega_\nu(\mathbf{x}))$  needs to be determined. Without specifying a particular ansatz for the dependence of the condensates on the spatial coordinates, this is a difficult functional minimization problem, that was so far not consistently solved in the literature. However, an efficient strategy to test whether, e.g., an IP is favored compared to homogeneous ground states  $\bar{\Phi}_j$  is to perform a stability analysis, where the homogeneous condensates are perturbed by arbitrary functions of infinitesimal amplitude  $\delta\phi_j(\mathbf{x})$ . References [53, 78] give an in-depth discussion of the method and its advantages and drawbacks on the example of a model with an analytical solution for  $\Sigma(\mathbf{x})$  in the whole  $(\mu, T)$  plane. Additional implications of the analysis, that are not discussed in Refs. [53, 78], will be subject of Section III D.

The main idea of the method is to determine whether the non-vanishing leading-order coefficients in an expansion with respect to the inhomogeneous perturbation is negative and, thus, leads to a lower free energy. In this case there is an instability of homogeneous condensates  $\bar{\Phi}_j$  (or linear combinations of the condensates) towards an IP. The first non-vanishing contribution when perturbing around solutions of the gap Eqs. (8) and (9) is of second order in the bosonic fields, since the first order contribution is proportional to the gap Eqs. [53]. In the case of multiple bosonic fields, the second order contribution is given by a Hessian matrix  $H_{\phi_j\phi_k}$ , which needs to be diagonalized in field space by finding its eigenvalues  $\Gamma_{\varphi_j}^{(2)}$  and eigenvectors  $\varphi_j$ , where, in general, one can find  $\varphi_j \neq \phi_j$ . In Ref. [1] multiple examples of the diagonalization procedure are given. The eigenvalues  $\Gamma_{\varphi_j}^{(2)}$  are called the bosonic two-point functions<sup>6</sup> and are typically expressed as functions of the bosonic momentum  $q$  corresponding to the respective Fourier mode of  $\delta\varphi_j$ <sup>7</sup>. The stability analysis is equivalent to a saddle point approximation of the partition function (2) with respect to the bosonic fields. Investigations of this type have recently been used a lot in the literature, see, e.g., Refs. [1, 85, 89, 90, 103, 104, 117].

For a detailed derivation of the bosonic two-point functions in models with multiple FF or Yukawa-type interactions we refer to appendix A of Ref. [1]. The focus of this Sec. is on the important differences in the analysis compared to the one in Ref. [1] caused by the included vector interactions. As discussed in the aforementioned Refs., the Hessian matrix can be derived by considering all second-order functional derivative terms with respect to the bosonic fields  $\phi_j$ . One then has to find a basis transformation mapping  $\phi_j$  on a field basis  $\varphi_j$  diagonalizing  $H$  such that the eigenvalues of  $H$  are the two-point functions  $\Gamma_{\varphi_j}^{(2)}(q)$ . With Lorentz-scalar interaction, one usually finds  $\varphi_j = \phi_j$ . This is not the case, when studying a model with vector interactions, such as Eq. (2) with  $\vec{\phi} = (\sigma, \omega_\nu)$ . Then, not only  $\varphi_j \neq \phi_j$  but also the eigenvectors of  $H$  in field space have to be determined for every value of  $q$ , i.e.  $\varphi_j(q) = \sum_k c_{j,k}(q)\phi_k$ .

In a model with a repulsive vector interaction, as mediated by  $\omega_3$ , the situation gets even more complicated. As discussed above, the homogeneous condensate  $\bar{\omega}_3 \sim i\bar{n}$  is purely imaginary. In contrast to the real-valued condensate  $\bar{\sigma}$  being analyzed by studying its stability with respect to real valued perturbations  $\delta\sigma$ , one has to treat fluctuations  $\delta\omega_3$  about  $\bar{\omega}_3$  in the complex plane. As recently discussed in Ref. [110], fluctuations must be in the direction of the steepest descent of the effective action corresponding to the stable Lefshetz thimble. The analysis of fluctuations about the homogeneous ground state is equivalent to saddle point approximation of the path integral, which in turn is only well-defined when the stable Lefshetz thimble is used [118]. Inspecting the effective action Eq. (2) under these aspects reveals that one has to consider real-valued fluctuations  $\delta\omega_3$  about the purely imaginary  $\bar{\omega}_3$ . This is demonstrated in a similar, NJL-type model with vector interactions in  $(3+1)$  dimensions in Ref. [119]. In Figure 5 therein, the Lefshetz thimble in the field space of the temporal vector component is depicted.

The resulting Hessian for model (2) is given by

$$H_{\phi_j\phi_k}(q) = \delta_{\phi_j, \phi_k} \left( \frac{\delta_{\phi_j, \sigma}}{\lambda_S} + \frac{1 - \delta_{\phi_j, \sigma}}{\lambda_V} \right) + \frac{1}{\beta} \sum_n \int \frac{d^2p}{(2\pi)^2} \text{tr} [S(\nu_n, \vec{p} + \vec{q}) c_j S(\nu_n, \vec{p}) c_k], \quad (13)$$

where we define the propagator of a free fermion with mass  $\bar{\Sigma}$  at chemical potential  $\bar{\mu} = \mu + i\bar{\Omega}_3$  and temperature  $T$  as

$$S = \left( \hat{Q}[\bar{\Sigma}, \bar{\Omega}_3 \delta_{\nu,3}] \right)^{-1}, \quad (14)$$

where  $\hat{Q}$  denotes the Fourier transform of  $Q[\sigma, \omega_\nu]$ , compare Eq. (3). Also,  $\nu_n = 2\pi(n - \frac{1}{2})T$  are fermionic Matsubara frequencies,  $\vec{q} = (q, 0)$ , where the spatial momentum integration can be chosen such that  $q$  is aligned along one

<sup>6</sup> In the mean-field approximation, this quantity is precisely the two-point-one-particle-irreducible vertex function, as it is often defined in QFT textbooks.

<sup>7</sup> The two-point function does not depend on the direction of the vector  $\vec{q}$ , because the computation involves the integration over loop-momenta, which can always be rotated such that the resulting integrals only depend on the absolute value of  $q$  respecting rotational invariance.

coordinate axis and we defined the bare vertices  $\vec{c} = (1, -\gamma_3, i\gamma_1, i\gamma_2)^T$ . Note that purely imaginary perturbations about  $\bar{\omega}_3$  would correspond to a vertex  $i\gamma_3$  instead of  $-\gamma_3$ . From Eq. (13), the cyclic property of the trace gives  $H_{\phi_j\phi_k} = H_{\phi_k\phi_j}$ . Formulas for the evaluation of  $H_{\phi_j\phi_k}(q)$  can be found in Appendix B. One finds  $H_{\sigma\omega_\nu} \propto \bar{\Sigma}$ , such mixing between the scalar and the vector mode is only relevant within the HBP. In practical computations, we compute  $H_{\phi_j\phi_k}(q)$  for a fixed  $q$  and diagonalize the resulting matrix numerically using *Python3* with various libraries [120–122].

#### D. Structure of the Hessian and its physical consequences

In the following, we will analyze the structure of the Hessian (13). A similar discussion can be found in the recent work [110], where a Polyakov-loop QM model with vector mesons has been studied using a static stability analysis, i.e., the Hessian was studied at  $q = 0$  only. Since  $\omega_\nu$  are components of a vector field, they pick up a sign under charge conjugation, i.e.,  $\mathcal{C}\omega_\nu\mathcal{C}^{-1} = -\omega_\nu$  (see Ref. [123] for a construction of the charge conjugation operation on four-dimensional spinors in  $(2+1)$  Euclidean spacetime dimensions). This breaking of charge conjugation symmetry is expected at finite density due to the excess of particles over anti-particles. Since  $H_{\sigma\omega_3} = H_{\omega_3\sigma}$  is non-vanishing at  $\mu \neq 0$  and purely imaginary, we conclude that the Hessian is non-Hermitian when  $\mathcal{C}$ -symmetry is broken. On the level of homogeneous condensation the breaking of  $\mathcal{C}$ -symmetry is realized through  $\bar{\omega}_3 \neq 0$  at  $\mu \neq 0$ , as will be discussed in Section IV A. However, the homogeneous condensates as well as the Hessian are still invariant under the combined operation of the linear  $\mathcal{C}$  transformation and anti-linear transformation of complex conjugation  $\mathcal{K}$ . A remaining invariance under a combined anti-linear  $\mathcal{CK}$  operation while the charge conjugation symmetry is broken at finite density can also be found in QCD itself as well as other QCD-inspired theories, such as Polyakov-NJL models [110, 124, 125].

Due to the invariance under the  $\mathcal{CK}$  operation the Hessian obeys the relation  $H = AH^*A$ , where  $*$  denotes complex conjugation and  $A = \text{diag}(1, -1, 1, 1)$ . It directly follows that  $H(q)$  possesses the same set of eigenvalues  $\lambda_j$  as  $H^*(q)$  and, consequently, the eigenvalues are either real-valued or come in complex-conjugate pairs. In the case of real-valued  $\lambda_j(q)$ , their interpretation as bosonic two-point functions  $\Gamma_{\varphi_j}^{(2)}(q)$  has been extensively discussed in the previous section and also in Ref. [1]. However, it has recently been demonstrated that the competition between repulsive and attractive interactions can induce complex-conjugate eigenvalue pairs in the stability analysis based on the study of mixing in Euclidean field theories with  $\mathcal{PT}$ -type symmetries<sup>8</sup> [60, 62, 63]. The existence of complex-conjugate eigenvalue pairs leads to bosonic correlation functions with spatial sine-like modulations in addition to the ordinary exponential decay and is directly related to the invariance under a combination of an linear and an anti-linear symmetry operation, such as the  $\mathcal{CK}$  operation [124].

The arguments for the existence of these oscillating bosonic correlation functions is briefly recapitulated here. As discussed in Ref. [60] and Ref. [62] for the static Hessian  $H(q = 0)$ , the inverse propagator matrix of dynamical bosonic fields is given by  $q^2 + H(0)$ , where the  $q^2$ -term comes from an a priori kinetic term as included in Refs. [60, 62, 63]. Then, one can directly classify the behavior of propagators by just inspecting the eigenvalues  $\Gamma_{\varphi_j}^{(2)}(0)$  of the static Hessian  $H(0)$ . The classification stems from the fact that roots of the expression  $q^2 + H(0)$  correspond to poles of the corresponding propagators of their dynamical bosonic fields. Ordinary, stable homogeneous phases with exponentially decaying propagators yield Hessian matrices with positive eigenvalues. An even number of negative eigenvalues corresponds to a stable ground state with respect to homogeneous perturbations but unstable with respect to inhomogeneous ones. Based on this analysis, one can obtain indications about IPs without including the full-momentum dependence of  $H$ . For an uneven number of negative eigenvalues in the static Hessian, the system is completely unstable against both homogeneous and inhomogeneous perturbations corresponding to a set of field values, which are not a stable minimum of the system. The complex-conjugate eigenvalues give rise to what is defined in Ref. [63] as modulated exponential decay, where one finds spatial oscillations with a momentum scale related to the imaginary part of the eigenvalue of  $H(0)$ . The real part of the eigenvalue is then a scale for the exponential decay, as it is with ordinary screening poles. This behavior [58] is also observed when an inhomogeneous ground condensate, such as, e.g., the chiral spiral, is destabilized through the fluctuations of Goldstone modes of  $O(N)$  symmetry breaking [61]. This analysis can be seen as an extension of the common stability analysis discussed before. However, it has, to our knowledge, so far only been performed with static modes [134].

In contrast to the previous works [63, 134], we consider auxiliary fields and, thus, do not a priori include a kinetic term for the bosonic fields. Thus, the bosonic two-point functions  $\Gamma_{\varphi_j}^{(2)}(q)$  are just the inverse of the bosonic correlation

<sup>8</sup> Here,  $\mathcal{PT}$ -type symmetry means symmetry under an linear symmetry operation  $\mathcal{P}$  and an arbitrary anti-linear symmetry operation  $\mathcal{T}$ . Field theories with  $\mathcal{PT}$ -type symmetries typically describe theories, that are not invariant under either  $\mathcal{P}$  or  $\mathcal{T}$  separately, but show invariance under their combined operation.  $\mathcal{PT}$ -type symmetries are also widely studied in quantum mechanics, optics and condensed matter for their unique properties [126–130]. Also, they are also known to be useful in resolving sign problems in lattice field theory simulations, see, e.g., Refs. [60, 131–133].

functions  $G_{\varphi_j}$  for the fields  $\varphi_j$ . However, the low-momentum expansion of the bosonic two-point functions results in

$$G_{\varphi_j}^{-1} \approx Z_{\varphi_j} q^2 + \Gamma_{\varphi_j}^{(2)}(0) \quad (15)$$

generating a kinetic term in the quantum effective action, where we defined the bosonic wave-function renormalization

$$Z_{\varphi_j} = \frac{1}{2} \left. \frac{d^2 \Gamma_{\varphi_j}^{(2)}}{dq^2} \right|_{q=0} \quad (16)$$

as the coefficient of the kinetic term generated by the fermion loop. Thus, one can perform a classification similar to the one just described by finding the roots of Eq. (15). It is important to note that  $Z_{\varphi_j}$  can have a small<sup>9</sup>, but non-vanishing imaginary part. As one can see from Eq. (15), both  $\Gamma_{\varphi_j}^{(2)}(0)$  and  $Z_{\varphi_j}$  being real-valued leads to purely imaginary propagator poles such that one obtains the usual exponential decay of  $G_{\varphi_j}$ . However, a non-vanishing real part of the obtained roots yields the above described oscillatory behavior of the propagator.

The analysis of the momentum-dependent Hessian  $H(q)$  is, however, more elaborate. In Section IV we will obtain the existence of regimes with complex-conjugate bosonic two-point functions  $\Gamma_{\varphi_j}^{(2)}(q)$  at both  $q = 0$  and  $q \neq 0$ , but also regimes in the  $(\mu, T)$  phase diagram of the FF model (2), where  $\Gamma_{\varphi_j}^{(2)}(q = 0) \in \mathbb{R}$  but  $\Gamma_{\varphi_j}^{(2)}(q \neq 0) \in \mathbb{C}$ . To our knowledge, the interpretation of  $H(q)$  developing complex-conjugate eigenvalue pairs at  $q \neq 0$  while the eigenvalues at  $q = 0$  are real-valued is unclear. Especially, the low-momentum expansion of the two-point function is not meaningful at non-vanishing momenta as one computes the Hessian by expanding about the homogeneous field values  $(\sigma, \omega_\nu) = (\bar{\Sigma}, \bar{\Omega}_3 \delta_{\nu,3})$ . Statements with respect to the existence of an IPs can, however, be made as discussed above, as long as the bosonic two-point functions are real-valued.

#### IV. PHASE DIAGRAM IN THE PRESENCE OF MIXING

We proceed by presenting our results for the phase diagram. Thereby, the vector coupling  $\lambda_V$  is treated as a free parameter and is varied to study varying strengths of the vector coupling. In order to define a scale for the strength of the scalar coupling  $\lambda_S^R$  used for comparison to the used values of the vector coupling  $\lambda_V$ , we define  $\lambda_S^R = 1/\Gamma_\sigma^{(2)}(q = 0)$  in the vacuum. Since the Hessian Eq. (13) is diagonal in the vacuum, this yields  $\lambda_S^R = \pi$ , where the result for  $\Gamma_\sigma^{(2)}$  in the vacuum can be found by computing Eq. (B2) in the limit of zero temperature and chemical potential, see Appendix A of Ref. [89] for the explicit expression. Thus, we consider  $\lambda_V \bar{\Sigma}_0 \in [0, \pi]$ . Note that all dimensionful quantities in plots are expressed in units of the chiral condensate in the vacuum  $\bar{\Sigma}_0$ .

##### A. Homogeneous condensation

As the homogeneous ground states are the input for the stability analysis, we first compute the phase diagram of model (1) when assuming homogeneous condensation  $(\sigma, \omega_\nu) = (\bar{\sigma}, \bar{\omega}_3 \delta_{\nu,3})$ . Minimizing the effective potential Eq. (7) with respect to  $\bar{\sigma}$  and  $\bar{\omega}_3$  as described in Section III A, we determine the thermodynamic ground state  $(\bar{\sigma}, \bar{\omega}_3) = (\bar{\Sigma}, \bar{\Omega}_3)$ . We distinct between the SP, where  $\bar{\Sigma} = 0$  and chiral symmetry is restored, and the HBP, where chiral symmetry is spontaneously broken by a non-vanishing  $\bar{\Sigma}$ . By studying the gap Eq. (8) and the effective potential (7) in Section III A, one obtains that the phase boundary between HBP and SP in the  $(\bar{\mu}, T)$  of the FF model defined in Eq. (1) is identical to the one of the  $(2+1)$ -dimensional GN model in the  $(\mu, T)$  plane, as first determined in Ref. [91]. The analytic expression for the critical chemical potential  $\mu_c$  of the GN model is given by

$$\mu_c(T) = T \text{arcosh} \left( 0.5 e^{\bar{\Sigma}_0/T} - 1 \right). \quad (17)$$

We use this analytical result as a cross-check for our numerical results at  $\lambda_V \neq 0$ , by computing  $\bar{\mu}_c = \mu_c + i\bar{\Omega}_3$  at the obtained phase boundary  $\mu_c(\lambda_V, T)$  and comparing the result to the case without vector coupling  $\mu_c(0.0, T)$ , as given by the right hand side of Eq. (17). The phase boundary in the  $(2+1)$ -dimensional GN model is of second order, except for the point  $(\mu/\bar{\Sigma}_0 = 1.0, T/\bar{\Sigma}_0 = 0.0)$  where one obtains an effective potential, which is flat for  $\bar{\sigma} \in [0.0, 1.0]$ .

In Fig. 1, we plot the phase boundary lines of the model (1) for different values of  $\lambda_V \bar{\Sigma}_0 \in [0.0, \pi]$ . Similar to findings with NJL-type models featuring vector interactions [106, 108, 109], an enlargement of the HBP is observed

<sup>9</sup> By small, it is meant that the imaginary part is typically at least two orders of magnitude smaller than the real part.






FIG. 1. Phase boundary lines between the HBP and the SP in the  $(\mu, T)$  space for 5 different values of the vector coupling  $\lambda_V$ . The  $\lambda_V = 0.0$  phase boundary corresponds to the known analytic solution of the GN model in  $(2+1)$  dimensions [91, 103].

when increasing the vector coupling. The extent of the HBP in the  $(\mu, T)$  plane grows monotonically with  $\lambda_V$ . For  $\lambda_V \bar{\Sigma}_0/\pi = 0.1, 0.3$ , the critical chemical potential, compared to the GN model result, increases when the temperature is decreased. Increasing  $\lambda_V$  further, leads to a change in this behavior, as mostly evident for  $\lambda_V \bar{\Sigma}_0 = \pi$  (visualized by the blue dot-dashed line). At this value of the vector coupling, the difference in the critical chemical potential compared to the GN model result, i.e.,  $\mu_c(\lambda_V, T) - \mu_c(0.0, T)$ , is larger for, e.g.,  $T/\bar{\Sigma}_0 = 0.4$  than for  $T/\bar{\Sigma}_0 = 0.0$ . This results in a back-bending shape of the transition line, which reminds of a spinodal line in typical NJL or GN model phase diagrams [115]. In this case, however, the left spinodal corresponds directly to the phase boundary line, since the phase transition is of second order and the order parameter  $\bar{\Sigma}(\mu, T)$  goes continuously to zero when crossing the boundary line from the HBP to the SP.

The order of the phase transition becomes evident when studying the behavior of the order parameter  $\bar{\Sigma}(\mu, T)$  as plotted in Fig. 2, where the value of  $\bar{\Omega}_3(\mu, T)$  for  $\lambda_V \bar{\Sigma}_0/\pi = 0.1, 1.0$  is visualized in a triangulated contour color map. For all studied values of the vector coupling, one observes a continuous decrease of the chiral condensate when increasing the temperature starting within the HBP. At  $T = 0$ , we obtain  $\bar{\Sigma}(\mu, T = 0) = \bar{\Sigma}_0, \forall \mu \leq \bar{\Sigma}_0$  in consistency with the Silver Blaze property. When further increasing the chemical potential above  $\bar{\Sigma}_0$ , one again obtains a continuous decrease of the chiral condensate from  $\bar{\Sigma}(\mu/\bar{\Sigma}_0 = 1, T = 0) = \bar{\Sigma}_0$  to  $\bar{\Sigma}(\mu = \mu_c, T = 0) = 0.0$  at the phase transition to the SP. This can be explained in context of the flatness of the effective potential of the GN model at  $(\mu/\bar{\Sigma}_0 = 1, T = 0)$ . This flatness as a function of  $\bar{\sigma}$  is also present in the effective potential at  $\lambda_V \neq 0$  at  $(\mu/\bar{\Sigma}_0 = 1, T = 0)$ . However, the additional contribution due to the non-vanishing  $\bar{\omega}_3$  causes solutions with higher densities to be favored, compare Eq. (7). The coupling of the gap Eqs. for  $\bar{\Sigma}$  and  $\bar{\Omega}_3$ , Eq. (8) and Eq. (9) respectively, leads then to a simultaneous decrease of  $\bar{\Sigma}$  and a continuous increase of  $\bar{\Omega}_3$  when increasing  $\mu$  for all chemical potentials  $\mu \in [\bar{\Sigma}_0, \mu_c]$ . For all values of  $\lambda_V$ , the chemical potentials in this interval correspond to  $\bar{\mu} = 1.0$ , since this is the only point at zero temperature where the gap Eq. (8) allows for solutions  $\bar{\Sigma}(\mu, T = 0)$  other than  $\bar{\Sigma}/\bar{\Sigma}_0 = 1.0$  or  $\bar{\Sigma}/\bar{\Sigma}_0 = 0.0$ . Due to the coupling of the gap Eqs. for  $\bar{\Sigma}$  and  $\bar{\Omega}_3$ , the solution of the gap Eq. for  $\bar{\Omega}_3(\mu, T = 0)$  can directly be read of the plot for this range of chemical potentials as it is given by  $\bar{\Omega}_3 = \mu - \bar{\mu}_c(T = 0) = \mu - \bar{\Sigma}_0$ . The density is given by  $\bar{N} = -i\bar{\Omega}_3/\lambda_V$  due to the gap Eq. (9) or, equivalently, the Ward identity (4) for  $\omega_3$ .

In the GN model, the density jumps from 0 to  $\bar{\Sigma}_0^2/(2\pi)$  when crossing the phase transition towards the HBP at zero temperature<sup>10</sup> [135]. This jump becomes a continuous transition for all  $\lambda_V > 0.0$  and  $\bar{N}/\bar{\Sigma}_0^2 = 1/(2\pi)$  is reached in the models with vector interaction directly at  $\mu = \mu_c$ . All other values of  $\bar{N}(\mu, T = 0)/\bar{\Sigma}_0^2 \in [0.0, 1/(2\pi))$  are obtained when continuously increasing the chemical potential from  $\mu/\bar{\Sigma}_0 > 1.0$ , whereas  $\bar{\Sigma}(\mu, T = 0)$  continuously decreases as discussed above. This continuous decrease for  $\lambda_V > 0.0$  is enforced by the additional term  $\propto -\bar{n}^2$  in the effective potential (7). The gap equation Eq. (9) allows a non-vanishing density only at  $\mu/\bar{\Sigma}_0 > 1.0$  and  $\bar{\Sigma}/\bar{\Sigma}_0 < 1.0$  such that at  $\mu/\bar{\Sigma}_0 = 1.0$  only the zero density solution is allowed. This is another clear indication that the phase boundary for  $\lambda_V > 0.0$  is of second order, also at  $T = 0$ .

At non-vanishing temperatures the relation between the chemical potential and the value of  $\bar{\Omega}_3$  becomes non-trivial and has to be determined via the numerical solution of the gap Eqs. (8), (9) and finding the global minimum of

<sup>10</sup> Note that there is a factor of 2 difference in the definition of the density between this work and Ref. [135].




FIG. 2. Contour color maps in the  $(\mu, T)$  plane for the value of the chiral condensate  $\bar{\Sigma}(\mu, T)$  for **(left)**  $\lambda_V \bar{\Sigma}_0 / \pi = 0.1$ , **(right)**  $\lambda_V \bar{\Sigma}_0 / \pi = 1.0$ . The green lines represent the second order phase boundary of the  $(2+1)$ -dimensional GN model, while the blue lines correspond to the phase boundary of the model (1) for the respective value of  $\lambda_V$ . Continuous data for the contour plots is obtained using triangulation provided by *Matplotlib* in *Python3* [120, 136]. Note that the plot range in the  $\mu$  and the  $T$  axis differs from plot to plot in order to make the contour lines visible.




FIG. 3. Contour color maps in the  $(\mu, T)$  plane for the value of the density  $\bar{N}(\mu, T)\pi = -i\bar{\Omega}_3(\mu, T)\pi/\lambda_V$  for **(left)**  $\lambda_V \bar{\Sigma}_0 / \pi = 0.1$ , **(right)**  $\lambda_V \bar{\Sigma}_0 / \pi = 1.0$ . The green lines represent the second order phase boundary of the  $(2+1)$ -dimensional GN model, while the blue lines correspond to the phase boundary of the model (1) for the respective value of  $\lambda_V$ . Continuous data for the contour plots is obtained using triangulation provided by *Matplotlib* in *Python3* [120, 136]. Note that the plot range in the  $\mu$  axis, the  $T$  axis and the discrete color bar differs from plot to plot in order to make the contour lines visible. Also, the discrete contour levels are not necessarily linearly distributed.

the effective potential (7). Fig. 3 depicts the density as a triangulated contour color map in the  $(T, \mu)$  plane for  $\lambda_V \bar{\Sigma}_0 / \pi = 0.1, 0.5, 1.0$ , respectively. Again, at  $T = 0$  we observe the Silver Blaze property as the density is zero for  $\mu / \bar{\Sigma}_0 \in [0.0, 1.0]$ . For non-vanishing temperature, this no longer holds and one observes the onset of the density for  $\mu / \bar{\Sigma}_0 < 1.0$ . In general, the density at fixed chemical potential grows monotonically with the temperature and vice versa. Comparing the phase diagrams for different vector couplings, one generally observes that the density is smaller for larger  $\lambda_V$  when comparing at fixed values of  $T$  and  $\mu$ .

Note that, in contrast to NJL model results [106], one does never obtain a first-order phase transition in the  $(\mu, T)$  plane. This is also relevant in the context of an IP, that typically covers the region of a first order phase transition between the HBP and SP that is present when restricting to homogeneous phases. It is worth to note that there exists a first order phase transition at  $\bar{\mu} / \bar{\Sigma}_0 = 1.0$  and  $T / \bar{\Sigma}_0 = 0$  for any value of  $\lambda_V$ . This is caused by the effective potential of the GN model being flat within  $\bar{\sigma} = [0.0, \bar{\Sigma}_0]$ , but the additional contribution of  $\bar{\Omega}_3$  favors solutions of the gap Eqs. with a high density leading to a jump of the global minimum from  $(\bar{\Sigma} / \bar{\Sigma}_0, \bar{\Omega}_3 / \bar{\Sigma}_0) = (1.0, 0.0)$  at  $(\bar{\mu} / \bar{\Sigma}_0 = 1.0 - \epsilon, T / \bar{\Sigma}_0 = 0.0)$  to  $(\bar{\Sigma} / \bar{\Sigma}_0, \bar{\Omega}_3 / \bar{\Sigma}_0) = (0.0, i\lambda_V \bar{\Sigma}_0 / (2\pi))$  at  $(\bar{\mu} = 1.0, T = 0.0)$ , where  $\epsilon$  can be

infinitesimally small. This demonstrates that one has to be very careful about drawing  $(\mu, T)$  phase diagrams, when using  $\bar{\mu}$  as an external parameter in the computation and the chemical potential  $\mu$  as a variable instead. As discussed in Section III A, the same value of  $(\bar{\mu}, T)$  can correspond to multiple points in the  $(\mu, T)$  plane or vice versa such that one has to always compare the corresponding values of the effective potential (7).

## B. Stability analysis with mixing

### 1. Symmetric phase

Within the SP the Hessian  $H_{\phi_j \phi_k}$  is diagonal for the original field basis, i.e.,  $\vec{\varphi} = \vec{\phi} = (\sigma, \omega_\nu)$ , since all non-vanishing off-diagonal elements are proportional to  $\bar{\Sigma}$ , see Eq. (13) and the formulas in Appendix B. In this case, the analysis is similar to the one described in Section III C. The eigenvalues/diagonal elements of  $H$  are the respective bosonic two-point functions  $\Gamma_{\varphi_j}^{(2)}$  of the fields  $(\sigma, \omega_\nu)$ . Thus, they can be analyzed to study (the absence of) instabilities towards the IP and the existence of the moat regime, similar to the analysis in Ref. [1] without vector interactions.

In the left plot of Fig. 4, the bosonic two-point functions  $\Gamma_{\varphi_j}^{(2)}$  are plotted as functions of the momentum  $q$  of the perturbation  $\delta\varphi_j$  for  $\lambda_V \bar{\Sigma}_0/\pi = 0.1$  and  $(\mu/\bar{\Sigma}_0, T/\bar{\Sigma}_0) = (1.051, 0.238)$ . This point in the phase diagram lies directly on the second order phase boundary between the SP and HBP (compare Fig. 1). One of the two-point functions is zero at  $q = 0$  and corresponds to the order parameter  $\sigma$  undergoing the phase transition. Since the two-point functions  $\Gamma_{\varphi_j}^{(2)}(q)$  can be interpreted as the curvature of the effective action (5) in the direction of  $\varphi_j(q)$ , it is expected that this curvature goes to zero for  $q = 0$  and  $\varphi_j = \sigma$  at the second-order homogeneous phase transition. The field  $\omega_\nu$  is not an order parameter and, thus, the corresponding bosonic two-point function do not show signals of the phase transition.

We always obtain that the two-point functions  $\Gamma_{\phi_j}^{(2)}(q)$  are monotonically increasing functions as can directly be seen for  $T = 0$  by taking the zero temperature limit for all diagonal elements of the Hessian using the formulas in Appendix B and setting  $\bar{\Sigma} = 0$ . An example for the bosonic two-point functions within the SP is plotted on the right side of Fig. 4. In the plot, we used the largest of the studied vector couplings  $\lambda_V \bar{\Sigma}_0/\pi = 1.0$ . A non-monotonic behavior of the two-point function is never observed within in the SP, which we studied for a large range of chemical potentials and temperatures. For  $\Gamma_\sigma^{(2)}$ , this result was already presented in Ref. [1], while in this work we also studied the two-point functions  $\Gamma_{\omega_\nu}^{(2)}(q)$ , that correspond to the included vector interactions. In the SP, however, the analysis yields the same conclusion as the one in Ref. [1]. We do neither observe an instability towards an IP nor a moat regime for all studied vector couplings. Together with the argument that all observed IPs feature a second order phase boundary towards the SP – which would be detected by the stability analysis [53] – have so far never been observed, we consider the absence of such an instability a strong indication for the non-existence of an IP within this model.

In the GN model there still exists a degeneracy between inhomogeneous condensates and homogeneous phases at zero temperature as found with a particular ansatz function [135]. This is also consistent with the stability analysis, where the bosonic two-point function  $\Gamma_\sigma^{(2)}(q)$  is flat and vanishes for a certain interval in  $q$  is found at the point  $(\mu/\bar{\Sigma}_0 = 1.0, T/\bar{\Sigma}_0 = 0)$  [1] – the same point in the phase diagram where also the homogeneous potential is flat. This flatness of the bosonic two-point function also occurs at the critical chemical potential  $\mu_c$  at any value of  $\lambda_V$  at  $T = 0.0$ , again indicating a similar degeneracy between the SP and the IP as before. This property could already be guessed from the right plot in Fig. 4, which is still at finite, but low enough temperature such that the two-point function  $\Gamma_\sigma^{(2)}$  is almost flat for small  $q$ . We expect, however, that a degenerate condensate would not be given by the one-dimensional kink ansatz from Ref. [135] (see Figure 5.4 therein), since its density  $\bar{n}$  remains homogeneous and is smaller than  $\bar{n}/\bar{\Sigma}_0^2 = 1/(2\pi)$ , which is the density corresponding to  $\bar{\Sigma} = 0.0$  when solving the gap Eqs. (8), (9) at  $(\mu/\bar{\Sigma}_0 = 1.0, T/\bar{\Sigma}_0 = 0)$ . Thus, the homogeneous solution  $(\bar{\Sigma}/\bar{\Sigma}_0 = 0.0, \bar{\Omega}_3/\bar{\Sigma}_0 = i\lambda_V \bar{\Sigma}_0/(2\pi))$  is expected to be favored over the ansatz in Ref. [135], as the bosonized action favors solution of the gap Eqs. with higher densities, compare Eq. (2) using  $\bar{\Omega}_3 = i\lambda_V \bar{n}$ .

### 2. Complex conjugate eigenvalues for $q = 0$

Within the HBP, one obtains mixing between  $\sigma$  and  $\omega_3$  when inspecting the static Hessian  $H(q = 0)$ , since  $H_{\sigma\omega_3}(q = 0) \neq 0$ . When considering  $q \neq 0$ , there can also occur mixing involving the spatial components of  $\omega_\nu$ , see the discussion in Section III C and Appendix B for the Hessian matrix elements. We start by focusing on the static Hessian  $H(q = 0)$ . To study these static mixing effects, the Hessian (13) can be considered only for  $\phi_j, \phi_k \in \{\sigma, \omega_3\}$ . Perturbations about the spatial components of the vector field are not interesting in this case, since the static Hessian is diagonal anyway with respect to  $\omega_1$  and  $\omega_2$ .




FIG. 4. Bosonic two-point functions  $\Gamma_{\varphi_j}^{(2)}(q)$  as functions of the momentum of the perturbation  $q$ . **(left)** At the homogeneous second order phase transition for  $\lambda_V \bar{\Sigma}_0/\pi = 0.1$ . **(right)** Within the SP for  $\lambda_V \bar{\Sigma}_0/\pi = 1.0$ . Note the different plot ranges for the  $y$  axis in the left and right panel.

At any temperature and  $\mu \neq 0$  within the HBP, one obtains mixing between  $\sigma$  and  $\omega_3$  such that the physical basis  $\varphi_j \neq \phi_j$ . As extensively discussed in Section III D, this mixing can lead to complex-conjugate eigenvalue pairs of the static Hessian  $H(0)$  depending on the parameters  $\mu$  and  $T$ . An example of this phenomenon is shown in Fig. 5, where the real and imaginary eigenvalues of the Hessian matrix  $H_{\phi_j \phi_k}(q)$  with  $\phi_j, \phi_k \in \{\sigma, \omega_3\}$  are plotted for  $(\mu/\bar{\Sigma}_0 = 1.03, T/\bar{\Sigma}_0 = 0.05)$  and  $\lambda_V \bar{\Sigma}_0/\pi = 1.0$ . The eigenvectors  $\varphi_a$  and  $\varphi_b$  are given by  $q$ -dependent linear combinations of  $\sigma$  and  $\omega_3$ , i.e.,  $\varphi_j(q) = c_j(q)\sigma + d_j(q)\omega_3(q)$ . The non-vanishing imaginary part of the eigenvalues for  $q = 0$  decreases as a function of  $q$  in this analysis such that real-valued eigenvalues are obtained for some relatively small  $q/\bar{\Sigma}_0 \approx 0.5$ . Then, also the degeneracy  $\text{Re} \Gamma_{\varphi_a}^{(2)} = \text{Re} \Gamma_{\varphi_b}^{(2)}$  is no longer enforced by  $\mathcal{CK}$  invariance (see the discussion in Section III D) resulting in an apparently non-analytic behavior of both two-point functions at  $q/\bar{\Sigma}_0 \approx 0.5$  and  $q/\bar{\Sigma}_0 \approx 3.0$ . It is interesting to note that whenever the real parts of  $\Gamma_{\varphi_{a/b}}^{(2)}(q)$  are equal to each other one also observes a non-vanishing imaginary part of  $\Gamma_{\varphi_a}^{(2)}(q) = -\Gamma_{\varphi_b}^{(2)}(q)$ . This is a result of the invariance under the  $\mathcal{CK}$  and the fact that both eigenvectors always need to fulfill  $\varphi_a \neq \varphi_b$ . In the static case with  $q = 0$ , the complex-conjugate eigenvalue pairs leads to bosonic propagators that are sinusoidal modulated alongside the usual exponential decay, see the discussions in [60, 63, 124] and Section III D. This behavior is clearly different from the usually obtained exponential decay and is caused by propagator poles developing non-vanishing real and imaginary parts. Note that the behavior of the two-point functions in Fig. 5 can change at  $q \neq 0$  when including perturbations about  $\omega_1$  and  $\omega_2$ , as discussed above. Thus, Fig. 5 should only be understood as an example for the effects of mixing and not as a full solution of the momentum dependence of the Hessian matrix  $H(q)$  of the full model (1). It would be the full solution of the momentum dependence when only interactions proportional to  $\sigma$  and  $\omega_3$  would be studied.

To characterize the regime with spatial oscillatory behavior of propagators in the  $(\mu, T)$  plane, we use the maximal imaginary part of the eigenvalues at  $q = 0$  given by

$$k_0 = \max_{\varphi_j} \left( \text{Im} \Gamma_{\varphi_j}^{(2)}(q = 0) \right). \quad (18)$$

This is an important scale for the momentum of the sinusoidal oscillation of the bosonic propagator, as a non-vanishing  $k_0$  induces a non-vanishing real part of the propagator poles, see the discussion in Section III D. In Fig. 6, we plot  $k_0$  in the  $(\mu, T)$  plane using a color code for  $\lambda_V \bar{\Sigma}_0/\pi \in \{0.6, 0.8, 1.0\}$ . A region with  $k_0 \neq 0$  for chemical potentials  $\mu/\bar{\Sigma}_0 > 1.0$  and rather small temperatures is observed for all three vector couplings. We note that  $\vec{\varphi}(q = 0) = (z_1\sigma + z_2\omega_3, z_2^*\sigma + z_1^*\omega_3, \omega_1, \omega_2)$  with complex-valued coefficients  $z_1$  and  $z_2$  for all studied points in the phase diagram and all studied vector couplings  $\lambda_V$ . The extent of the region with complex-conjugate eigenvalue pairs both in  $\mu$  and in  $T$  direction strongly depends on the value of the vector coupling. For  $\lambda_V \bar{\Sigma}_0/\pi = 0.6$ , this region's extent is significantly smaller than for  $\lambda_V \bar{\Sigma}_0/\pi = 1.0$ . For all studied vector couplings, the width in the  $\mu$  direction of the regime with oscillating propagator behavior decreases for larger temperatures, until  $k_0$  goes to zero. This is caused by thermal fluctuations suppressing the oscillatory behavior in the propagators, as can be seen from  $k_0$  decreasing monotonically with the temperature at fixed  $\mu$ . Such a behavior is typical for regimes with spatial oscillations, because the thermal fluctuations tend to destroy ordering in general [54, 61, 73, 76]. For all three vector couplings,  $k_0$  jumps




FIG. 5. The real and imaginary part of the bosonic two-point functions  $\Gamma_{\varphi_j}^{(2)}(q)$  are plotted as functions of the momentum of the perturbation  $q$ . The bosonic two-point functions are obtained as eigenvalues of  $H_{\phi_j, \phi_k}$  with  $\phi_j, \phi_k \in \{\sigma, \omega_3\}$ . Note that only  $\text{Im} \Gamma_{\varphi_a}^{(2)}(q)$  is plotted, since  $\text{Im} \Gamma_{\varphi_a}^{(2)} = -\text{Im} \Gamma_{\varphi_b}^{(2)}$ .

from zero to a non-vanishing value when crossing  $\mu/\bar{\Sigma}_0 > 1.0$  at zero temperature in consistency with the Silver Blaze property. This can also be derived from the formulas in Appendix B for the off-diagonal elements. Investigations of  $k_0$  as show that  $k_0(\mu, T)$  seems to have a rapid but continuous onset from zero when increasing  $\mu$  from the left of the regime with spatial oscillations at any  $T \neq 0$ . In the typical literature, the transition from  $k_0 = 0$  to  $k_0 \neq 0$  is also called disorder line [61] – as it is not a phase transition in the typical sense, but marks a distinctly different behavior of propagators.

In Fig. 7, we plot  $k_0$  at zero temperature as a function of  $\lambda_V \bar{\Sigma}_0$  and  $\mu/\bar{\Sigma}_0$  and observe the above described non-analytic behavior at all  $\lambda_V \bar{\Sigma}_0 > 0.5\pi$ . Precisely at  $\lambda_{V,c} \bar{\Sigma}_0 = 0.5\pi$ , however, there is a continuous onset of  $k_0$ . For  $\lambda_V \bar{\Sigma}_0 < 0.5\pi$ , we do not find  $k_0 \neq 0$  at all. Since the width of the region with  $k_0 \neq 0$  in the  $\mu$  direction is largest at  $T = 0$  for all studied vector couplings, we expect that one finds  $k_0 = 0$  for all vector couplings lower than  $\lambda_{V,c}$  both at zero and non-zero temperature. Thus, we expect there is no region with complex-conjugate eigenvalues appearing in the whole  $(\mu, T)$  plane for  $\lambda_V \bar{\Sigma}_0 < 0.5\pi$ . From Fig. 7 it becomes clear that the extent of the region with complex-conjugate eigenvalues of  $H(0)$  grows with increasing vector coupling when  $\lambda_V > \lambda_{V,c}$ . Also, the value of  $k_0$  grows monotonically when increasing  $\lambda_V$  from any value larger than  $\lambda_{V,c}$ . Both observations show that the increase of the vector coupling increases the mixing effects, which is expected since an increase of  $\lambda_V$  lowers the difference  $H_{\sigma\sigma}(0) - H_{\omega_3\omega_3}(0)$ . This difference between the two diagonal elements needs to be smaller than the product of the off-diagonal elements such that the eigenvalues of the corresponding  $2 \times 2$  block are complex-conjugate eigenvalue pairs. Thus, the increase of mixing effects also amounts to a growth of the extent of the regime with spatially oscillating propagators and the maximal obtained values of  $k_0$ .

Although  $k_0$  is a useful observable to quantify the appearance of complex-conjugate eigenvalues, it is not the unique scale determining the oscillation of the propagator  $G_{\varphi_j}$ . Instead, the scales for the oscillation and the exponential decay of  $G_{\varphi_j}$  are given by the respective real and imaginary parts of the roots  $q_{\varphi_j}^{1,2}$  of Eq. (15). These roots are given by

$$q_{\varphi_j}^{1,2} = \pm i \sqrt{\frac{\Gamma_{\varphi_j}^{(2)}(0)}{Z_{\varphi_j}}}. \quad (19)$$

In order to compute these roots numerically, one can use the fact that  $\varphi_j(q)$  has a weak dependence on  $q$  around  $q = 0$ . In practice, the computation of  $Z_{\varphi_j}$  involves the discrete differentiation of the eigenvalues of  $H$  (compare Eq. (15) and Eq. (16)). Thus, its computation has to be performed by carefully taking into account the discretization error in  $q$  and the change of basis  $\varphi_j(q)$  in this area. Also, the imaginary part of  $Z_{\varphi_j}$  is very small such that one also encounters problems with under-flowing of double precision. In test runs of this evaluation the maximum value for  $\text{Im} Z_{\varphi_j}$  encountered was on the order of  $10^{-3}$ , while  $\text{Re} Z_{\varphi_j}$  is typically of order  $10^{-1}$ . However, the problems involving the numerical evaluation of  $Z$  make a precise determination of the poles using Eq. (19) impractical with respect to the study of the whole phase diagram.




FIG. 6. Contour color maps in the  $(\mu, T)$  plane for the value of the maximal imaginary part of the eigenvalues at  $q = 0$ , denoted by  $k_0$ , for **(top left)**  $\lambda_V \bar{\Sigma}_0/\pi = 0.6$ , **(top right)**  $\lambda_V \bar{\Sigma}_0/\pi = 0.8$ , **(bottom)**  $\lambda_V \bar{\Sigma}_0/\pi = 1.0$ . The green lines represent the second order phase boundary of the  $(2 + 1)$ -dimensional GN model, while the blue lines correspond to the phase boundary of the model (1) for the respective value of  $\lambda_V$ . Continuous data for the contour plots is obtained using triangulation provided by *Matplotlib* in *Python3* [120, 136]. Note that the plot range in the  $\mu$  axis, the  $T$  axis and the discrete color bar differs from plot to plot in order to make the contour lines visible.

Hence, we decided to use<sup>11</sup>

$$\theta = \arg \left( \sqrt{\frac{\Gamma_\chi^{(2)}(0)}{\operatorname{Re} Z_\chi}} \right) = \frac{1}{2} \arccos \left( \frac{\operatorname{Re} \Gamma_\chi^{(2)}(0)}{|\Gamma_\chi^{(2)}(0)|} \right), \quad \chi = \arg \max_{\varphi_j} \left( \operatorname{Im} \Gamma_{\varphi_j}^{(2)}(q=0) \right), \quad (20)$$

in order to compare the scale of oscillation in comparison to the exponential decay. Eq. (20) gives the complex argument  $\theta$  of  $q^{1,2}$  when setting the imaginary part of  $Z_{\varphi_j}$  to zero. As argued above, the error of this approximation should be rather small. Note that the numerical computation of  $\theta$  using Eq. (20) does require an evaluation of  $Z_{\varphi_j}$ .

We find  $r = \tan \theta = \operatorname{Re} q_\chi^{1,2} / \operatorname{Im} q_\chi^{1,2}$  as the ratio of the oscillation frequency  $\operatorname{Re} q_\chi^{1,2}$  and the decay rate  $\operatorname{Im} q_\chi^{1,2}$ . In Fig. 8, we plot  $r$  for two different values  $\lambda_V \bar{\Sigma}_0 = 0.6\pi$  and  $\lambda_V \bar{\Sigma}_0 = \pi$ . The maximal value of  $r$  obtained is 0.58 for  $\lambda_V \bar{\Sigma}_0 = \pi$  such that frequency of the oscillation is larger than half of the exponential decay rate at this point in the phase diagram. Similar to the behavior of  $k_0$ , we find that for smaller vector couplings lower values of  $r$  are obtained in general. This is expected, since mixing effects will not be as drastic for lower vector couplings, see the discussion above. For  $\lambda_V \bar{\Sigma}_0 = 0.6\pi$ , the maximal value of  $r$  is roughly 0.3. In general, the obtained contour lines for  $r$  are very similar to the ones of  $k_0$ , see Fig. 6. The figure demonstrates that for large parts of the regime with oscillatory behavior the wavelengths are in the order of the inverse of the exponential decay rate.

<sup>11</sup> Note that by definition of  $\Gamma_\chi^{(2)}$  its imaginary part is always positive such that the determination of the argument in Eq. (20) is always valid.




FIG. 7. Contour color maps in the  $(\lambda_V, \mu)$  plane for the value of the maximal imaginary part of the eigenvalues at  $q = 0$ , denoted by  $k_0$ , at zero temperature. Continuous data for the contour plots is obtained using triangulation provided by *Matplotlib* in *Python3* [120, 136].




FIG. 8. Contour color maps in the  $(\mu, T)$  plane for the ratio  $r$  between the frequency of the spatial oscillation and the exponential decay rate of the exponential decay of the propagator. **(left)**  $\lambda_V \bar{\Sigma}_0/\pi = 0.6$ , **(right)**  $\lambda_V \bar{\Sigma}_0/\pi = 1.0$ . The green lines represent the second order phase boundary of the  $(2 + 1)$ -dimensional GN model, while the blue lines correspond to the phase boundary of the model (1) for the respective value of  $\lambda_V$ . Continuous data for the contour plots is obtained using triangulation provided by *Matplotlib* in *Python3* [120, 136]. Note that the plot range in the  $\mu$  axis, the  $T$  axis and the discrete color bar differs from plot to plot in order to make the contour lines visible.

### 3. Complex-conjugate eigenvalue pairs emerging at $q \neq 0$

Already in Fig. 5, where only mixing of  $\sigma$  and  $\omega_3$  was taken into account, one can observe the appearance of complex-conjugate eigenvalues of the Hessian at  $q \neq 0$ . However, considering a  $4 \times 4$  Hessian matrix  $H_{\phi_j \phi_k}(q)$  with  $\phi_j, \phi_k \in \{\sigma, \omega_\nu\}$  yields a more involved mixing pattern, since also  $\omega_1$  contributes to mixing effects with  $\sigma$  when studying  $q \neq 0$ . The other vector component  $\omega_2$  is not mixing with  $\sigma$  (but with the other components of  $\omega_\nu$ ), since we choose  $\mathbf{q} = (q, 0)$  in Section III C. The roles of  $\omega_1$  and  $\omega_2$  are exchanged if we choose  $\mathbf{q}$  to be aligned with the  $x_2$  axis. Since our whole analysis is invariant under spatial rotations, the eigenvalues are expected to be independent of the chosen spatial direction  $\mathbf{q}/|\mathbf{q}|$ .

The more complicated mixing pattern is depicted in Fig. 9. From the plot one obtains that  $H(0)$  has real eigenvalues, but then develops complex-conjugate eigenvalue pairs at some value of  $q = q_B$ , where we define

$$q_B = \min_{q \in C} q, \quad C = \{q \in [0, \infty) | \text{Im} \Gamma_{\phi_j}^{(2)}(q) \neq 0\}. \quad (21)$$




FIG. 9. The real and imaginary part of the bosonic two-point functions  $\Gamma_{\varphi_j}^{(2)}(q)$  as functions of the momentum of the perturbation  $q$ . The bosonic two-point functions are obtained as eigenvalues of  $\hat{H}_{\phi_j, \phi_k}$  with  $\phi_j, \phi_k \in \{\sigma, \omega_\nu\}$ . Note that the non-vanishing imaginary part does not necessarily belong to similar eigenvectors  $\varphi_j(q)$  and complex-valued two-point functions appear as complex-conjugate pairs.

In this case, one obtains that only two of the four eigenvalues have non-vanishing imaginary parts for fixed  $q$ , while the other two are real-valued. As can also be seen from Fig. 9, complex-conjugate eigenvalue pairs can be obtained for multiple intervals in  $q$ . This leads to the rather complicated behavior of two-point functions with the real-parts of different eigenvalues becoming degenerate depending on the value of  $q$ . Since complex-conjugate eigenvalue pairs can occur for all of the eigenvalues  $\Gamma_{\varphi_j}^{(2)}$  with  $j \in \{1, 2, 3, 4\}$ , the yellow line in the plot always only describes the appearance of the maximal imaginary part in any of those eigenvalues. Since the eigenvectors of  $H$  can be strongly  $q$ -dependent especially for large  $q$ , one might even argue that the association of the eigenvalues using functions  $\Gamma_{\varphi_j}^{(2)}(q)$  is not very insightful. Nevertheless, Fig. 9 certainly demonstrates the involved mixing effects between scalar and vector modes. Also it shows that there is certainly non-monotonic behavior of the real parts of the two-point functions such that one cannot exclude the appearance of moat regime. However, also the bosonic wave-function renormalization  $Z_{\varphi_j}$  can become complex, see Eq. (16), and might not be a decent criterion for a moat regime where  $\arg \min_q \text{Re} \Gamma_{\varphi_j}^{(2)}(q) \neq 0$ . This is caused by non-analytic behavior of  $\Gamma_{\varphi_j}^{(2)}(q)$  around the regions with non-vanishing imaginary parts leading also to non-monotonic behavior. For these reasons we refrain from studying the moat regime within the HBP and focus on the complex-valued eigenvalues instead.

In the left plot of Fig. 10, we plot a color map in the  $(T, \mu)$  plane encoding  $q_B$  for  $\lambda_V \bar{\Sigma}_0 / \pi$ . In the right plot of this figure we visualize

$$k_{\max} = \max_{j,q} \left( \text{Im} \Gamma_{\varphi_j}^{(2)}(q) \right) \quad (22)$$

also using a color code. Fig. 10 demonstrates that complex-conjugate eigenvalues appear in large parts of the HBP except for a rather small region at small temperatures and chemical potentials. For some parts of this region the obtained imaginary parts are rather small. In consistency with the Silver Blaze property,  $k_{\max} = 0$  for  $T = 0$  and  $\mu / \bar{\Sigma}_0 < 1.0$ . For rather small chemical potentials,  $k_{\max} / \bar{\Sigma}_0$  is of the order of  $10^{-3}$  and  $q_B / \bar{\Sigma}_0 \gg 1$ . Accordingly, the interpretation in terms of oscillating propagators as in the static case is not possible as a low-momentum expansion is not meaningful. However, if existent, any oscillating effects in this region should be negligible anyhow given that any relevant imaginary part should be smaller than  $k_{\max} / \bar{\Sigma}_0 \sim 10^{-3}$ . For larger chemical potentials, one obtains  $q_B / \bar{\Sigma}_0 < 1.0$  making a low-momentum expansion of the inverse propagator more sensible. We note that due to the computational demands of computing multiple momentum integrals for the determination of the matrix entries of  $H(q)$  and its diagonalization, the resolution in  $q$  for the computation of the data Fig. 10 was chosen as  $\Delta q = 0.2$ . The rather coarse resolution in  $q$  results in inaccuracies in the determination of  $q_B$  and  $k_{\max}$ , because the intervals, where complex-valued eigenvalues occur, can be smaller than  $\Delta q$ . This is evident for some data points around the homogeneous phase boundary as well as small  $\mu$  and  $T / \bar{\Sigma}_0 \in [0.5, 0.7]$ , where  $q_B$  appears to have jumps when changing  $\mu$  or  $T$ . At these data points, it is likely that for some intervals in  $q$ , which are smaller than  $\Delta q$ , complex-valued  $\Gamma_{\varphi_j}^{(2)}(q \neq 0)$  appear, that were missed such that the correct  $q_B$  differs from the depicted data point. Anyhow, the






FIG. 10. Contour color maps in the  $(\mu, T)$  plane for  $\lambda_V \bar{\Sigma}_0/\pi = 1.0$  encoding **(left)**  $q_B$  (see Eq. (21)) **(right)**  $k_{\max}$  (see Eq. (22)). The green lines represent the second order phase boundary of the  $(2+1)$ -dimensional GN model, while the blue lines correspond to the phase boundary of the model (1). Note that the plot range in the discrete color bar differs from plot to plot. The color bar for  $q_B$  is cut off at  $q_B/\bar{\Sigma}_0 = 4.0$  such that the behavior of lower  $q_B$  could be visualized accurately. The discretization of  $q$  is given by  $\Delta q = 0.2$  resulting in discretization errors in the computation of  $q_B$  and  $k_{\max}$ .

value of  $k_{\max}$  is likely to be small anyhow within these intervals. Still, it is unclear whether complex-conjugate eigenvalues appearing at  $q \neq 0$  have any consequence on the behavior of propagators.

Overall, to the best of our knowledge there is no clear physical explanation for the appearance of complex-conjugate two-point function pairs appearing at  $q \neq 0$  in the literature. Also, this work marks the first observation of this phenomenon in the literature – again to the knowledge of the author. For small  $q_B$ , a similar interpretation as in the static case of correlation functions with sinusoidal modulations might be meaningful, again performing a low-momentum expansion of the inverse propagators  $G_{\varphi_j}^{-1}$ , compare Eq. (15) around some non-vanishing  $q$  and determining propagator poles similar to Eq. (19). However, in the regions with larger  $q_B/\bar{\Sigma}_0 > 1.0$  such an expansion is certainly not sensible. An inversion and four-transformation of the obtained  $\Gamma_{\varphi_j}^{(2)}$  to compute  $G_{\varphi_j}(x, y)$  within these regions yields an ordinary exponential decay in the HBP. This is expected, since  $k_{\max}$  is rather small compared to the real-part of the two-point functions in these parameter regions (see Fig. 10).

## V. IMPLICATIONS FOR GENERAL FOUR-FERMION MODELS IN $(2+1)$ DIMENSIONS

In this section, we will argue that the previous analysis and, consequently, the phase diagram of the simpler model (1) is identical to  $(2+1)$ -dimensional FF models with all kind of local FF interactions. The general prototype of these models is defined by the action

$$\mathcal{S}_{\text{FF}}[\bar{\psi}, \psi] = \int_0^\beta d\tau \int d^2x \left\{ \bar{\psi} (\not{\partial} + \gamma_3 \mu) \psi - \sum_{j=1}^{16} \left[ \frac{\lambda_S}{2N} (\bar{\psi} c_j \psi)^2 + \frac{\lambda_V}{2N} \left( (\bar{\psi} i c_j \gamma_3 \psi)^2 + (\bar{\psi} i c_j \vec{\tau} \psi)^2 \right) \right] \right\}, \quad (23)$$

where  $\psi$  now contains  $2N$  four-component spinors (due to an additional "isospin" degree of freedom compared to the fields in Eq. (1)). The interaction vertices  $c_{j,\nu}$  are  $8 \times 8$  matrices in isospin and spin space and elements of

$$C = (c_j)_{j=1, \dots, 16} \equiv (1, i\gamma_4, i\gamma_5, \gamma_{45}, \vec{\tau}, i\vec{\tau}\gamma_4, i\vec{\tau}\gamma_5, \vec{\tau}\gamma_{45}), \quad (24)$$

where  $\vec{\tau}$  is the vector of Pauli-matrices acting on the isospin degrees of freedom. The matrices  $\gamma_4$  and  $\gamma_5$  anti-commute with the  $\gamma_\nu$ , while  $\gamma_{45} \equiv i\gamma_4\gamma_5$ , consequently, commutes with the  $\gamma_\nu$ . This model is invariant under a global  $U(4N)$  chiral symmetry (see appendix A of Ref. [1] for details).

When neglecting vector interactions by setting  $\lambda_V = 0$ , this model has already been studied with respect to IPs and the moat regime in Ref. [1]. Also therein, it was shown that the FF model (23) can easily be generalized to a Yukawa model by including local self-interactions and a kinetic term for the bosonic fields (after bosonization similar to Eq. (2)) without altering the above analysis, as was also done in Ref. [1]. We refrain from doing this generalization

to a Yukawa model for the action (23). This would further complicate the analysis through the medium induced mixing of vector and scalar modes, which would also affect the purely bosonic terms.

For the FF model (23) and any FF model with a subset of its interaction channels, we will now demonstrate that their Hessian is block-diagonal with  $4 \times 4$  blocks that are identical to the  $4 \times 4$  Hessian matrix (13) of the simpler model Eq. (1). Therefore, Eq. (23) is bosonized, as before, with auxiliary bosonic fields  $(\phi_a, v_{a,\nu})$ . These fulfill the Ward identities

$$\langle \phi_a \rangle = -\frac{\lambda_S}{N} \langle \bar{\psi} c_a \psi \rangle, \quad \langle v_{a,\nu} \rangle = -i \frac{\lambda_V}{N} \langle \bar{\psi} c_a \gamma_\nu \psi \rangle. \quad (25)$$

Similar to the discussion in Section II, one can assume that  $\bar{v}_{a,j} = 0$  for  $j = 1, 2$ , leaving invariance under spatial rotation intact, where  $\bar{v}_{a,j}$  are the homogeneous condensates of the vector fields. Then, the derivation of the Hessian is similar to the one described in Ref. [1] and in Section III C. Through the  $U(4N)$  global symmetry transformations, one can choose the values of  $\phi_a = \bar{\sigma} \delta_{a,0}$  such that there is no mixing between those fields, i.e.,  $H_{\phi_a \phi_b} = 0$  for  $a \neq b$ . A similar transformation can be made for the fields  $v_{a,\nu}$ , such that  $H_{v_{a,\nu} v_{b,\nu}} = 0$  for  $a \neq b$ . With inspection of the trace in Eq. (13), one can then already infer that  $H_{\phi_a, \nu v_{b,\nu}} = 0$  for  $a \neq b$ . Thus, one obtains mixing only for each respective  $4 \times 4$  block in  $H$  with  $a = b$ , i.e., only  $H_{\phi_a, \nu v_{a,\nu}}$ ,  $H_{\phi_a, \nu \phi_a, \nu}$  and  $H_{v_{a,\nu} v_{a,\nu}}$  can be non-vanishing. Again, inspecting the trace in Eq. (13) and using the exchange properties of  $c_j \in C$  with each other and the  $\gamma_\nu$ , one can derive that the matrix elements of these respective  $4 \times 4$  blocks in  $H$  are identical to the  $4 \times 4$  Hessian matrix of Eq. (1) given by Eq. (13), respectively. Thus, the Hessian of Eq. (23) is block-diagonal consisting of 16 matrices of size  $4 \times 4$ , which are identical to the Hessian of the model studied in the previous sections. Consequently, also the phase diagram is identical to the one of Eq. (1) with the enhancement of the HBP when increasing  $\lambda_V$ , the absence of an IP and the existence of regimes with complex-conjugate eigenvalues of  $H$ .

With this analysis and the results in Ref. [89], we have demonstrated the absence of an IP in  $(2+1)$ -dimensional FF models with all kind of local interaction terms. Instead, we found a regime with spatially oscillating, but exponentially damped bosonic correlation functions when mixing between scalar and vector modes is allowed and the vector coupling exceeds a certain value of  $\lambda_{V,c} \bar{\Sigma}_0 = 0.5\pi$ , see, e.g., Fig. 7 and Fig. 8 as well as the discussion in Section IV B 2.

## VI. CONCLUSIONS

In this work, we analyzed the stability of homogeneous ground states for a  $(2+1)$ -dimensional FF model (1) with both scalar and vector interactions with particular emphasis on the effects of mixing between scalar and vector modes. The analysis was performed in the mean-field approximation, i.e., neglecting bosonic quantum fluctuations. Also, we showed that our findings hold for a more general FF model (23) consisting of all relevant interaction channels in  $(2+1)$  dimensions. We have shown the stability of homogeneous ground states against inhomogeneous perturbations and argued that this is strong evidence for the absence of an IP in all  $(2+1)$ -dimensional fermionic theories with local FF interactions. Instead, a regime with spatially oscillating, but exponentially damped mesonic correlation functions has been detected through the appearance of complex-conjugate eigenvalues in the Hessian matrix for static perturbations  $H(q=0)$ . This regime is often also termed ‘quantum pion liquid’ or ‘quantum spin liquid’ [61] – in analogy to condensed matter literature – and is directly related to the invariance of our model under  $\mathcal{CK}$  transformation at  $\mu \neq 0$  [60, 63]. We also find regions with complex-conjugate eigenvalues in the Hessian matrix for perturbations with a non-vanishing momentum, i.e.,  $H(q=q_B \neq 0)$ , while  $H(0)$  has real-valued eigenvalues. So far, the implications of this phenomenon are not discussed in the literature. However, we argued that the value of the obtained imaginary parts are so small that the mesonic correlation functions will follow an ordinary exponential decay within these regions. The whole phase diagram of the FF model (1) was presented for different values of the vector coupling including the regimes with ‘quantum pion liquid’ behavior – which are located at low  $T$  and intermediate  $\mu$  within the HBP. The typical ratio of the frequency of the oscillation and the exponential decay rate of the mesonic propagators is such that the oscillatory behavior could have an effect on related observables.

The finding of the ‘quantum pion liquid’ is to the best of our knowledge the first finding of this regime generated by the interplay of a attractive, scalar and a repulsive, vector FF interaction – apart from the report of this phenomenon in Ref. [110], where it was not the main focus of the study. Since the ‘quantum pion liquid’ is closely related to the invariance under the  $\mathcal{CK}$  operation [124], we think this regime could also be of relevance in the phase diagram of QCD at non-vanishing densities. However, this work should be understood as a first, qualitative investigation of the underlying mechanism through mixing of scalar and vector modes. The used  $(2+1)$ -dimensional models are too simplistic to make quantitative predictions for the phase diagram of QCD. We also want to note that the authors of Ref. [124] also report non-vanishing complex-conjugate eigenvalue pairs appearing in their static Hessian matrix (compare Figs. 17-21) in a PNJL model. However, these non-vanishing imaginary parts are generated by an analysis of Polyakov-loop effects and the mixing contributions stemming from fermionic point-like interactions are not included.

In QCD, the contributions of both the gluon effects and of the mixing between the chiral condensate and vector mesons will play an important role and might lead to the existence of a ‘quantum pion liquid’ regime at non-vanishing densities.

There are several possibilities to extend on the present work. A straightforward extension would be to study a  $(3 + 1)$ -dimensional QM model including Polyakov-loop effects as well as vector meson interactions as in Ref. [110]. This allows to study the mixing of Polyakov-loop, chiral condensate and vector mesons in the same model. Also, one could make phenomenologically more relevant predictions with respect to the parameter regions, where the ‘quantum pion liquid’ should appear. Moreover, it would certainly be interesting to study mixing effects beyond the mean-field approximation. Thereby, it is interesting to note that already in Refs. [97, 137] oscillating mesonic correlation functions have been measured in a  $(2 + 1)$ -dimensional GN model, i.e., without including vector interactions, at finite number of fermion flavors  $N$  using lattice field theory. In Ref. [1], we argued that these findings are reminiscent from an IP that exists within the mean-field approximation at finite lattice spacing [102, 103]. However, as discussed in Refs. [73, 138], the naive and, following a similar argumentation, also the staggered fermion discretization yields off-diagonal interaction terms generated by the doublers, that contain  $\gamma$  matrices. Thus, it is up to speculation whether the oscillating mesonic correlation functions observed in Refs. [97, 137] might as well be generated by mixing effects as discussed in the present work. When going beyond the mean-field approximation using a model that directly contains diagonal vector interactions as Eq. (1), complex weights appear in the path integral [119, 139]. This likely complicates using  $1/N$  expansion techniques [140, 141] or other analytical approaches like non-abelian bosonization [77, 79]. Thus, one would have to either rely on Lefschetz thimble approaches [119, 142] or Functional Renormalization Group techniques [44].

Certainly, regimes where mesonic correlation functions are oscillatory will have effects on, e.g., the propagation of pions in heavy-ion-collision experiments. Regarding experimental observables, direct consequences of spatially oscillatory regimes have to be worked out. Using the most regime dispersion relation, this has been recently done using Hambury, Brown and Twist interferometry [134, 143] leading to signal peaks in two-particle correlation functions. To study consequences for experimental observables in the ‘quantum pion liquid’ regime one would first have to work out the phenomenologically relevant scales of the oscillation and exponential decay, e.g., using a Polyakov-loop QM model as described above.

## VII. ACKNOWLEDGMENTS

I thank M. Buballa, T. Motta, Z. Nussinov, L. Pannullo, R. Pisarski, F. Rennecke, D. H. Rischke, S. Schindler, M. Ogilvie, M. Wagner for fruitful discussions related to this work. Especially, I thank F. Rennecke, S. Schindler and M. Ogilvie for help with the interpretation of complex-conjugate eigenvalues appearing in the Hessian matrix. Furthermore, I like to thank A. Koenigstein, L. Pannullo, R. Pisarski, S. Schindler and M. Wagner for helpful comments on this manuscript. I acknowledge the support of the *Deutsche Forschungsgemeinschaft* (DFG, German Research Foundation) through the collaborative research center trans-regio CRC-TR 211 ‘‘Strong-interaction matter under extreme conditions’’ – project number 315477589 – TRR 211. I acknowledge the support of the *Helmholtz Graduate School for Hadron and Ion Research*.

All numerical results in this work were obtained using *Python3* [120] with various libraries [121, 122]. The plots were designed using the *Matplotlib* package [136].

### Appendix A: Conventions for the Wick rotation

In  $(2 + 1)$ -dimensional spacetime with the metric  $\eta_{\mu\nu} = \text{diag}(1, -1, -1)$  we demonstrate the Wick rotation using the action and partition function

$$\mathcal{S}_{M,\text{mix}}[\bar{\psi}, \psi] = \int dx_0 \int d^2x \left\{ \bar{\psi} (i\gamma_M^\nu \partial_\nu) \psi + \left[ \frac{\lambda_S}{2N} (\bar{\psi} \psi)^2 + \frac{\lambda_V}{2N} (\bar{\psi} i\gamma_M^\nu \psi) (\bar{\psi} i\gamma_{M,\nu} \psi) \right] \right\}, \quad Z = \int \mathcal{D}\bar{\psi} \mathcal{D}\psi e^{i\mathcal{S}_{M,\text{mix}}[\bar{\psi}, \psi]} \quad (\text{A1})$$

with most quantities as defined below Eq. (1). The Gamma matrices  $\gamma_M^\nu$  fulfill  $\{\gamma_M^\mu, \gamma_M^\nu\} = 2\eta_{\mu\nu}$ , as usual. For the Wick rotation we use the conventions

$$\tau = ix_0, \quad \gamma_3 = \gamma_M^0, \quad \gamma_i = i\gamma_M^i, \quad i \in \{1, 2\}, \quad (\text{A2})$$

such that one obtains  $(\gamma_\nu)^2 = 1$ . This changes the vector FF interaction term in the following way

$$(\bar{\psi} i\gamma_M^\nu \psi) (\bar{\psi} i\gamma_{M,\nu} \psi) = (\bar{\psi} i\gamma_M^0 \psi)^2 - (\bar{\psi} i\gamma_M^i \psi)^2 = (\bar{\psi} i\gamma_3 \psi)^2 + \bar{\psi} (i\gamma_i \psi)^2 \quad (\text{A3})$$

where the additional factor  $(-i)^2 = -1$ . With the other standard changes of the action and the definition  $\mathcal{S}_{M,\text{mix}} = i\mathcal{S}_{M,\text{mix}}$  one obtains the Euclidean action Eq. (1).

In these conventions the density  $n$  is given by

$$n = \frac{1}{N} \frac{d \ln Z}{d\mu} = -\frac{\langle \bar{\psi} \gamma_3 \psi \rangle}{N}, \quad (\text{A4})$$

such that the Ward identity for  $\omega_3$  (4) follows. Note that these conventions and the Ward identity may differ from recent works, such as Ref. [119].

## Appendix B: Formulas for the stability analysis with mixing

In this section, we collect formulas needed for computation of the Hessian matrix elements  $H_{\phi_j, \phi_k} = H_{\phi_k, \phi_j}$  with  $\vec{\phi} = (\sigma, \omega_\nu)$ , compare Eq. (13). Inserting

$$S(\nu_n, \mathbf{p}) = \frac{-i\vec{p} + \bar{\Sigma}}{\tilde{p}^2 + \bar{\Sigma}^2}, \quad \tilde{p} = (\nu_n - i\bar{\mu}, \mathbf{p})^T \quad (\text{B1})$$

and the respective vertex  $\vec{c} = (1, -\gamma_3, i\gamma_1, i\gamma_2)$  for  $\vec{\phi} = (\sigma, \omega_3, \omega_1, \omega_2)$  in Eq. (13), respectively, one obtains for the diagonal elements

$$H_{\sigma\sigma} = \frac{1}{\lambda_S} + \frac{N_\gamma}{\beta} \int \frac{d^2 p}{(2\pi)^2} \sum_{n=-\infty}^{\infty} \frac{-\tilde{p}^2 - \mathbf{p}\mathbf{q} + \bar{\Sigma}^2}{(\nu_n - i\bar{\mu})^2 + (\mathbf{p} + \mathbf{q})^2 + \bar{\Sigma}^2} \frac{1}{(\nu_n - i\bar{\mu})^2 + \mathbf{p}^2 + \bar{\Sigma}^2}, \quad (\text{B2})$$

$$H_{\omega_\nu \omega_\nu} = \frac{1}{\lambda_S} + (1 - 2\delta_{\nu,3}) \frac{N_\gamma}{\beta} \int \frac{d^2 p}{(2\pi)^2} \sum_{n=-\infty}^{\infty} \frac{(2\delta_{\nu,\alpha} - 1)\tilde{p}_\alpha \tilde{p}_\alpha + (2\delta_{\nu,1} - 1)p_1 q - \bar{\Sigma}^2}{(\nu_n - i\bar{\mu})^2 + (\mathbf{p} + \mathbf{q})^2 + \bar{\Sigma}^2} \frac{1}{(\nu_n - i\bar{\mu})^2 + \mathbf{p}^2 + \bar{\Sigma}^2}, \quad (\text{B3})$$

where we choose the angle integration such that  $q$  lies on the  $x_1$  axis. The off-diagonal elements are given by

$$H_{\sigma\omega_\nu} = (\delta_{\nu,3}(-i-1) + 1) \bar{\Sigma} \frac{N_\gamma}{\beta} \int \frac{d^2 p}{(2\pi)^2} \sum_{n=-\infty}^{\infty} \frac{2\tilde{p}_\nu + \delta_{\nu,1} p_1 q}{(\nu_n - i\bar{\mu})^2 + (\mathbf{p} + \mathbf{q})^2 + \bar{\Sigma}^2} \frac{1}{(\nu_n - i\bar{\mu})^2 + \mathbf{p}^2 + \bar{\Sigma}^2}, \quad (\text{B4})$$

$$H_{\omega_1 \omega_2} = \frac{N_\gamma}{\beta} \int \frac{d^2 p}{(2\pi)^2} \sum_{n=-\infty}^{\infty} \frac{2p_1 p_2 + q p_2}{(\nu_n - i\bar{\mu})^2 + (\mathbf{p} + \mathbf{q})^2 + \bar{\Sigma}^2} \frac{1}{(\nu_n - i\bar{\mu})^2 + \mathbf{p}^2 + \bar{\Sigma}^2}, \quad (\text{B5})$$

$$H_{\omega_3 \omega_j} = -\frac{N_\gamma}{\beta} \int \frac{d^2 p}{(2\pi)^2} \sum_{n=-\infty}^{\infty} \frac{2\tilde{p}_3 p_j + \delta_{j,1} \tilde{p}_3 q}{(\nu_n - i\bar{\mu})^2 + (\mathbf{p} + \mathbf{q})^2 + \bar{\Sigma}^2} \frac{1}{(\nu_n - i\bar{\mu})^2 + \mathbf{p}^2 + \bar{\Sigma}^2}, \quad j \in \{1, 2\}. \quad (\text{B6})$$

$$(\text{B7})$$

Note that in consistency with the repulsive nature of the Yukawa interaction  $\bar{\psi} \omega_3 \psi$ , the Hessian matrix elements with only one  $\omega_3$  index are purely imaginary.

### 1. Finite temperature expressions for non-vanishing $\mathbf{q}$

In order to evaluate the Hessian at finite temperature and chemical potential, we perform the Matsubara summation. For some of the above expression, one needs to do some additional manipulations in order to perform the summation in an easier way or to circumvent problems in the contour integration stemming from  $\tilde{p}_3^2$  terms in the numerator. Equation (B2) is exactly the two-point function of the GN model, and was calculated multiple times, e.g., in Refs. [89, 90, 103], see Appendix B of Ref. [1] for the evaluation of the two-point function in various limits of  $\bar{\Sigma}$ ,  $T$  and  $q$ . The other entries, however, have to our knowledge not been computed. Using some manipulation of the numerator of Eq. (B3) for  $\nu = 3$  as well as rewriting the denominator in a partial fraction (see Eqs. (4.14 – 16) in Ref. [3] for the general idea) one finds that

$$H_{\omega_3 \omega_3} = \frac{1}{\lambda_V} - \ell_1(\mu, T, \bar{\Sigma}, \bar{\Omega}_3) + L_{2,+}(\bar{\mu}, T, \bar{\Sigma}, q) + \ell_3(\mu, T, \bar{\Sigma}, q), \quad (\text{B8})$$

where  $\ell_1$  is known from the GN model and appears in the gap equation (8),

$$L_{2,+}(\bar{\mu}, T, \bar{\Sigma}, q) = \frac{1}{2} \int \frac{d^2 p}{(2\pi)^2} \sum_{n=-\infty}^{\infty} \frac{N_\gamma}{\beta} \frac{(q^2 + 4\bar{\Sigma}^2)}{(\nu_n - i\bar{\mu})^2 + \mathbf{p}^2 + \bar{\Sigma}^2} \frac{1}{(\nu_n - i\bar{\mu})^2 + (\mathbf{p} + \mathbf{q})^2 + \bar{\Sigma}^2} \equiv \frac{1}{2} (q^2 + 4\bar{\Sigma}^2) \ell_2(\mu, T, \bar{\Sigma}, q) \quad (\text{B9})$$

is (up to a factor of 2) the momentum dependence of the two-point function in the GN model (compare Eq. (13) of Ref. [1]) and the new contribution

$$\ell_3(\bar{\mu}, T, \bar{\Sigma}, q) = \frac{2N_\gamma}{\beta} \int \frac{d^2 p}{(2\pi)^2} \sum_{n=-\infty}^{\infty} \frac{\mathbf{p}^2 + \mathbf{p}\mathbf{q}}{(\nu_n - i\bar{\mu})^2 + \mathbf{p}^2 + \bar{\Sigma}^2} \frac{1}{(\nu_n - i\bar{\mu})^2 + (\mathbf{p} + \mathbf{q})^2 + \bar{\Sigma}^2}. \quad (\text{B10})$$

The Matsubara summation can be performed as usual by analytic continuation and a contour integral. Then, one obtains

$$\ell_3(\bar{\mu}, T, \bar{\Sigma}, q) = 2N_\gamma \int \frac{d^2 p}{(2\pi)^2} \frac{(p^2 + \mathbf{p}\mathbf{q})}{q^2 + 2\mathbf{p}\mathbf{q}} \left[ \frac{1}{2E} (1 - n_F(E) - n_{\bar{F}}(E)) - \frac{1}{2E_q} (1 - n_F(E_q) - n_{\bar{F}}(E_q)) \right] \quad (\text{B11})$$

with  $E$ ,  $n_F(x)$ ,  $n_{\bar{F}}(x)$  as defined in Section III A and  $E_q = \sqrt{(p+q)^2 + \bar{\Sigma}^2}$ . This expression can be split up between  $p_1 < q/2$  and  $p_1 > q/2$ , manipulated and evaluated using a Cauchy-Principal value (similar to how  $L_2$  can be evaluated) and amounts to

$$\ell_3(\bar{\mu}, T, \bar{\Sigma}, q) = \frac{N_\gamma}{(4\pi)} \left\{ \int_0^{q/2} dp \frac{p}{E} \frac{p^2 - q^2/2}{q\sqrt{q^2 - 4p^2}} [1 - n_F(E) - n_{\bar{F}}(E)] + \int_0^\infty \frac{p}{E} [1 - n_F(E) - n_{\bar{F}}(E)] \right\}, \quad (\text{B12})$$

where the latter term is linearly divergent and identical to  $\ell_1$ . Thus, the divergences of  $\ell_1$  and  $\ell_3$  exactly cancel each other out such that Eq. (B8) is finite. We are aware that these divergences occur because of splitting up the numerator of Eq. (B3) for  $\nu = 3$ , as described above. However, this procedure is easier than dealing with the  $(\nu_n - i\bar{\mu})^2$  term in the numerator, since the standard method of analytic continuation and contour integration cannot be done as usual, since the integrand does not fall off fast enough.

In turn, the other diagonal elements are rather straight forward

$$H_{\omega_j \omega_j} = \frac{1}{\lambda_V} - \ell_1 + \frac{1}{2} q^2 \ell_2 + \frac{2N_\gamma}{\beta} \int \frac{d^2 p}{(2\pi)^2} \sum_{n=-\infty}^{\infty} \frac{p_j^2 + \delta_{j,1} p_1 q}{(\nu_n - i\bar{\mu})^2 + \mathbf{p}^2 + \bar{\Sigma}^2} \frac{1}{(\nu_n - i\bar{\mu})^2 + (\mathbf{p} + \mathbf{q})^2 + \bar{\Sigma}^2}. \quad (\text{B13})$$

Splitting up the numerator and treating the remaining integrals with standard techniques (shifts, inversions of the integration variable) allow to identify already known integral structures for  $j = 1$  and one obtains

$$H_{\omega_1 \omega_1} = \frac{1}{\lambda_V} - \ell_1 + \frac{1}{2} q^2 \ell_2(\mu, T, \bar{\Sigma}, q) + \ell_1 + q^2 \ell_2(\mu, T, \bar{\Sigma}, q) = \frac{1}{\lambda_V} + \frac{3}{2} q^2 \ell_2(\mu, T, \bar{\Sigma}, q). \quad (\text{B14})$$

Also, one finds for  $j = 2$

$$H_{\omega_2 \omega_2} = \frac{1}{\lambda_V} + \frac{N_\gamma}{2\pi q} \int_0^{q/2} \frac{p^3}{E\sqrt{q^2 - 4p^2}} [1 - n_F(E) - n_{\bar{F}}(E)], \quad (\text{B15})$$

which is again finite. Note that the differences in the expression for  $H_{\omega_1 \omega_1}$  and  $H_{\omega_2 \omega_2}$  come from the choice on  $\mathbf{q} = (q, 0)$ . The matrix elements  $H_{\omega_2 \omega_2}$  and  $H_{\omega_1 \omega_1}$  would be exchanged if we would have chosen  $\mathbf{q} = (0, q)$ , as expected since the analysis is invariant under spatial rotations. The limits of zero temperature in the above expressions is rather straightforward.

The mixing between  $\sigma$  and  $\omega_3$  is given by

$$H_{\sigma \omega_3} = -i\bar{\Sigma} \frac{2N_\gamma}{\beta} \int \frac{d^2 p}{(2\pi)^2} \sum_{n=-\infty}^{\infty} \frac{\tilde{p}_3}{(\nu_n - i\bar{\mu})^2 + \mathbf{p}^2 + \bar{\Sigma}^2} \frac{1}{(\nu_n - i\bar{\mu})^2 + (\mathbf{p} + \mathbf{q})^2 + \bar{\Sigma}^2}. \quad (\text{B16})$$

This expression can be evaluated using a contour integral, as the analytic continuation of the integrand is well-behaved when closing the contour at infinity. The result yields

$$H_{\sigma \omega_3}(q) = i\bar{\Sigma} N_\gamma \int \frac{d^2 p}{(2\pi)^2} \frac{1}{\mathbf{q}^2 + 2\mathbf{p}\mathbf{q}} [(n_F(E) - n_{\bar{F}}(E)) - (n_F(E_q) - n_{\bar{F}}(E_q))] \quad (\text{B17})$$

$$= -i\bar{\Sigma}N_\gamma \int_0^{q/2} \frac{dp}{2\pi} \frac{p}{q\sqrt{q^2 - 4p^2}} [n_F(E) - n_{\bar{F}}(E)] \equiv -i\bar{\Sigma}\ell_4(\mu, T, \bar{\Sigma}, q),$$

where one can directly see that this expression vanishes in the SP and for  $\mu = 0$ .

Also, one obtains

$$H_{\sigma\omega_1} = 2\bar{\Sigma}|q|\ell_2(\mu, T, \bar{\Sigma}, q) \quad (\text{B18})$$

and  $H_{\sigma\omega_2} = 0$ , where the term with  $j = 2$  vanishes using symmetry arguments in the angle integration<sup>12</sup> (compare Eq. (B4) for  $\nu = 2$ ). Again, the two matrix elements  $H_{\sigma\omega_1}$  and  $H_{\sigma\omega_2}$  would be exchanged, if we would setup the momentum integration such that  $\mathbf{q}$  would be aligned with the  $p_2$  axis. In a similar way as for  $H_{\sigma\omega_2}$ , we obtain that  $H_{\omega_1\omega_2} = H_{\omega_3\omega_2} = 0$ . Lastly, one obtains

$$H_{\omega_3\omega_1} = 0 \quad (\text{B19})$$

through splitting up the integral in Eq. (B6) into two integrals, where each integral contains one of the summands in the numerator. Both integrals turn to be proportional to  $\ell_4$ , but with different signs and prefactors such that both contributions cancel each other.

## 2. Finite temperature expressions for $q = 0$

For the  $q = 0$  expressions (and even more limits) of  $H_{\sigma\sigma}$  and all integrals related to it, we refer again to Ref. [87]. Note that several of the integrals in the beginning of Appendix B vanish, when the limit of  $q$  going to zero is taken. More precisely, we find

$$H_{\sigma\omega_j}(q = 0) = 0, \quad H_{\omega_3\omega_j}(q = 0) = 0, \quad j \in \{1, 2\}, \quad (\text{B20})$$

$$H_{\omega_1\omega_2}(q = 0) = 0, \quad (\text{B21})$$

as can be seen from going into polar coordinates after Matsubara summation and performing the angle integration, respectively. The Hessian (13) becomes diagonal w.r.t. the  $2 \times 2$  block with  $\phi_j, \phi_k \in \{\omega_1, \omega_2\}$ . Thus, complex-conjugate eigenvalue pairs can only be generated by mixing of  $\sigma$  and  $\omega_3$  at  $q = 0$ , allowing to study this phenomenon by only taking the respective  $2 \times 2$  block in Eq. (13). For the rest of the diagonal elements, we find

$$H_{\omega_j\omega_j}(q = 0) = \frac{1}{\lambda_V} - \ell_1(\mu, T, \bar{\Sigma}) + \ell_3(\mu, T, \bar{\Sigma}, q = 0) \quad (\text{B22})$$

$$\begin{aligned} &= \frac{1}{\lambda_V} - \ell_1 + \frac{N_\gamma}{8\pi} \int_0^\infty dp \frac{p^3}{E^3} [1 - (1 + \beta E)(n_F(E) + n_{\bar{F}}(E)) + \beta E(n_F^2(E) + n_{\bar{F}}^2(E))] \\ &= \frac{1}{\lambda_V} + \frac{N_\gamma}{8\pi} \int_{|\bar{\Sigma}|}^\infty dE \left\{ \beta E [n_F^2(E) + n_{\bar{F}}^2(E) - n_F(E) - n_{\bar{F}}(E)] + \right. \\ &\quad \left. - \frac{\bar{\Sigma}^2}{E^2} [1 - (1 + \beta E)(n_F(E) + n_{\bar{F}}(E)) + \beta E(n_F^2(E) + n_{\bar{F}}^2(E))] \right\}, \end{aligned} \quad (\text{B23})$$

which is finite due to the suppression of high momenta in the first summand of the integrand, and

$$H_{\omega_3\omega_3}(q = 0) = \frac{1}{\lambda_V} - \ell_1(\mu, T, \bar{\Sigma}) + 2\bar{\Sigma}^2\ell_2(\mu, T, \bar{\Sigma}, q = 0) + \ell_3(\mu, T, \bar{\Sigma}, q = 0), \quad (\text{B24})$$

which is identical to  $H_{\omega_j\omega_j}$  up to an additional contribution of  $2\bar{\Sigma}^2\ell_2$ . The diagonal element  $H_{\sigma\sigma}$  is given by the bosonic two-point function  $\Gamma_\sigma^{(2)}(q)$  in the  $(2 + 1)$ -dimensional GN model and studied in this limit in Refs. [1, 103]. For the remaining off-diagonal element we find

$$H_{\sigma\omega_3} = -i\bar{\Sigma} \frac{N_\gamma}{\pi} \int_0^\infty dE \left\{ \beta E [n_F^2(E) - n_{\bar{F}}^2(E)] - [n_F(E) - n_{\bar{F}}(E)] \right\}. \quad (\text{B25})$$

<sup>12</sup> At first glance, this seems to be a trivial integration, but one still needs to perform a Cauchy principal value before using symmetry arguments.

- 
- [1] L. Pannullo and M. Winstel, Absence of inhomogeneous chiral phases in (2+1)-dimensional four-fermion and Yukawa models, *Phys. Rev. D* **108**, 036011 (2023), [arXiv:2305.09444 \[hep-ph\]](#).
- [2] M. Asakawa and K. Yazaki, Chiral Restoration at Finite Density and Temperature, *Nucl. Phys. A* **504**, 668 (1989).
- [3] S. P. Klevansky, The Nambu–Jona-Lasinio model of quantum chromodynamics, *Reviews of Modern Physics* **64**, 649 (1992).
- [4] O. Scavenius, A. Mocsy, I. N. Mishustin, and D. H. Rischke, Chiral phase transition within effective models with constituent quarks, *Phys. Rev. C* **64**, 045202 (2001), [arXiv:nucl-th/0007030](#).
- [5] S. Hands and D. N. Walters, Evidence for BCS diquark condensation in the (3+1)-d lattice NJL model, *Phys. Lett. B* **548**, 196 (2002), [arXiv:hep-lat/0209140](#).
- [6] M. Sadzikowski, Coexistence of pion condensation and color superconductivity in two flavor quark matter, *Phys. Lett. B* **553**, 45 (2003), [arXiv:hep-ph/0210065](#).
- [7] S. Hands and D. N. Walters, Numerical portrait of a relativistic BCS gapped superfluid, *Phys. Rev. D* **69**, 076011 (2004), [arXiv:hep-lat/0401018](#).
- [8] B.-J. Schaefer and J. Wambach, The Phase diagram of the quark meson model, *Nucl. Phys. A* **757**, 479 (2005), [arXiv:nucl-th/0403039](#).
- [9] J. E. Drut and T. A. Lahde, Is graphene in vacuum an insulator?, *Phys. Rev. Lett.* **102**, 026802 (2009), [arXiv:0807.0834 \[cond-mat.str-el\]](#).
- [10] B. Hiller, A. A. Osipov, A. H. Blin, and J. da Providencia, Effects of quark interactions on dynamical chiral symmetry breaking by a magnetic field, *SIGMA* **4**, 024 (2008), [arXiv:0802.3193 \[hep-ph\]](#).
- [11] R. Gatto and M. Ruggieri, Quark Matter in a Strong Magnetic Background, *Lect. Notes Phys.* **871**, 87 (2013), [arXiv:1207.3190 \[hep-ph\]](#).
- [12] D. Roscher, J. Braun, and J. E. Drut, Inhomogeneous phases in one-dimensional mass- and spin-imbalanced Fermi gases, *Phys. Rev. A* **89**, 063609 (2014), [arXiv:1311.0179 \[cond-mat.quant-gas\]](#).
- [13] K. G. Klimenko, R. N. Zhokhov, and V. C. Zhukovsky, Superconductivity phenomenon induced by external in-plane magnetic field in (2+1)-dimensional Gross-Neveu type model, *Mod. Phys. Lett. A* **28**, 1350096 (2013), [arXiv:1211.0148 \[hep-th\]](#).
- [14] V. V. Braguta, S. N. Valgushev, A. A. Nikolaev, M. I. Polikarpov, and M. V. Ulybyshev, Interaction of static charges in graphene within Monte-Carlo simulation, *Phys. Rev. B* **89**, 195401 (2014), [arXiv:1306.2544 \[cond-mat.str-el\]](#).
- [15] J. O. Andersen, W. R. Naylor, and A. Tranberg, Phase diagram of QCD in a magnetic field: A review, *Rev. Mod. Phys.* **88**, 025001 (2016), [arXiv:1411.7176 \[hep-ph\]](#).
- [16] S. Hands, W. Armour, and C. Strouthos, Graphene as a Lattice Field Theory, *PoS CPOD2014*, 016 (2015), [arXiv:1501.01895 \[cond-mat.str-el\]](#).
- [17] J. A. Gracey, Loop calculations for models of graphene, *PoS LL2018*, 070 (2018), [arXiv:1808.05030 \[hep-th\]](#).
- [18] J. A. Gracey, Large  $N$  critical exponents for the chiral Heisenberg Gross-Neveu universality class, *Phys. Rev. D* **97**, 105009 (2018), [arXiv:1801.01320 \[hep-th\]](#).
- [19] T. C. Lang and A. M. Läuchli, Quantum Monte Carlo Simulation of the Chiral Heisenberg Gross-Neveu-Yukawa Phase Transition with a Single Dirac Cone, *Phys. Rev. Lett.* **123**, 137602 (2019), [arXiv:1808.01230 \[cond-mat.str-el\]](#).
- [20] R. L. S. Farias, D. C. Duarte, G. a. Krein, and R. O. Ramos, Thermodynamics of quark matter with a chiral imbalance, *Phys. Rev. D* **94**, 074011 (2016), [arXiv:1604.04518 \[hep-ph\]](#).
- [21] T. G. Khunjua, K. G. Klimenko, R. N. Zhokhov, and V. C. Zhukovsky, Inhomogeneous charged pion condensation in chiral asymmetric dense quark matter in the framework of NJL<sub>2</sub> model, *Phys. Rev. D* **95**, 105010 (2017), [arXiv:1704.01477 \[hep-ph\]](#).
- [22] B. Knorr, Critical chiral Heisenberg model with the functional renormalization group, *Phys. Rev. B* **97**, 075129 (2018), [arXiv:1708.06200 \[cond-mat.str-el\]](#).
- [23] R.-A. Tripolt, B.-J. Schaefer, L. von Smekal, and J. Wambach, The low-temperature behavior of the quark-meson model, *Physical Review D* **97**, 034022 (2018), [arXiv:1709.05991 \[hep-ph, physics.hep-th, physics.nucl-th\]](#).
- [24] V. V. Braguta, M. I. Katsnelson, A. Y. Kotov, and A. M. Trunin, Catalysis of Dynamical Chiral Symmetry Breaking by Chiral Chemical Potential in Dirac semimetals, *Phys. Rev. B* **100**, 085117 (2019), [arXiv:1904.07003 \[cond-mat.str-el\]](#).
- [25] E. J. Ferrer and V. de la Incera, Absence of Landau-Peierls Instability in the Magnetic Dual Chiral Density Wave Phase of Dense QCD, *Phys. Rev. D* **102**, 014010 (2020), [arXiv:1902.06810 \[nucl-th\]](#).
- [26] P. Lakaschus, M. Buballa, and D. H. Rischke, Competition of inhomogeneous chiral phases and two-flavor color superconductivity in the NJL model, *Phys. Rev. D* **103**, 034030 (2021), [arXiv:2012.07520 \[hep-ph\]](#).
- [27] B. S. Lopes, S. S. Avancini, A. Bandyopadhyay, D. C. Duarte, and R. L. S. Farias, Hot QCD at finite isospin density: Confronting the SU(3) Nambu–Jona-Lasinio model with recent lattice data, *Phys. Rev. D* **103**, 076023 (2021), [arXiv:2102.02844 \[hep-ph\]](#).
- [28] F. Attanasio, L. Rammelmüller, J. E. Drut, and J. Braun, Pairing patterns in polarized unitary Fermi gases above the superfluid transition, *Phys. Rev. A* **105**, 063317 (2022), [arXiv:2112.07309 \[cond-mat.quant-gas\]](#).
- [29] W. Gyory and V. de la Incera, Phase transitions and resilience of the magnetic dual chiral density wave phase at finite temperature and density, *Phys. Rev. D* **106**, 016011 (2022), [arXiv:2203.14209 \[nucl-th\]](#).
- [30] F. Attanasio, J. P. Klinger, and J. M. Pawłowski, Low energy effective theories on the lattice with coloured noise, *PoS LATTICE2022*, 290 (2023), [arXiv:2212.12354 \[hep-lat\]](#).

- [31] F. Attanasio, M. Bauer, R. Kapust, and J. M. Pawłowski, Harmonically trapped fermions in one dimension: A finite-temperature lattice Monte Carlo study, *Phys. Rev. A* **109**, 033305 (2024), [arXiv:2309.06766 \[cond-mat.quant-gas\]](#).
- [32] E. J. Ferrer, W. Gyory, and V. de la Incera, Thermal phonon fluctuations and stability of the magnetic dual chiral density wave phase in dense QCD, (2023), [arXiv:2307.05621 \[nucl-th\]](#).
- [33] A. Ayala, A. Bandyopadhyay, R. L. S. Farias, L. A. Hernández, and J. L. Hernández, QCD equation of state at finite isospin density from the linear sigma model with quarks: The cold case, (2023), [arXiv:2301.13633 \[hep-ph\]](#).
- [34] A. Podo and L. Santoni, Fermions at finite density in the path integral approach, (2023), [arXiv:2312.14753 \[hep-th\]](#).
- [35] S. Pitsinigos and A. Schmitt, Chiral crossover versus chiral density wave in dense nuclear matter, *Phys. Rev. D* **109**, 014024 (2024), [arXiv:2309.01603 \[nucl-th\]](#).
- [36] S. M. A. Tabatabaee Mehr, Chiral symmetry breaking and phase diagram of dual chiral density wave in a rotating quark matter, *Phys. Rev. D* **108**, 094042 (2023), [arXiv:2306.11753 \[nucl-th\]](#).
- [37] Y. Nambu and G. Jona-Lasinio, Dynamical model of elementary particles based on an analogy with superconductivity. i, *Phys. Rev.* **122**, 345 (1961).
- [38] Y. Nambu and G. Jona-Lasinio, Dynamical model of elementary particles based on an analogy with superconductivity. ii, *Phys. Rev.* **124**, 246 (1961).
- [39] M. Gell-Mann and M. Levy, The axial vector current in beta decay, *Nuovo Cim.* **16**, 705 (1960).
- [40] T. Inagaki, D. Kimura, and H. Shimoji, Super Restoration of Chiral Symmetry in Massive Four-Fermion Interaction Models, (2023), [arXiv:2306.00470 \[hep-ph\]](#).
- [41] S. Borsanyi, Z. Fodor, C. Hoelbling, S. D. Katz, S. Krieg, C. Ratti, and K. K. Szabo (Wuppertal-Budapest), Is there still any  $T_c$  mystery in lattice QCD? Results with physical masses in the continuum limit III, *JHEP* **09**, 073, [arXiv:1005.3508 \[hep-lat\]](#).
- [42] P. de Forcrand, Simulating QCD at finite density, *PoS LAT2009*, 010 (2009), [arXiv:1005.0539 \[hep-lat\]](#).
- [43] C. S. Fischer, QCD at finite temperature and chemical potential from Dyson–Schwinger equations, *Prog. Part. Nucl. Phys.* **105**, 1 (2019), [arXiv:1810.12938 \[hep-ph\]](#).
- [44] W.-j. Fu, J. M. Pawłowski, and F. Rennecke, QCD phase structure at finite temperature and density, *Phys. Rev. D* **101**, 054032 (2020), [arXiv:1909.02991 \[hep-ph\]](#).
- [45] J. Bernhardt, C. S. Fischer, and P. Isserstedt, Finite-volume effects in baryon number fluctuations around the QCD critical endpoint, *Phys. Lett. B* **841**, 137908 (2023), [arXiv:2208.01981 \[hep-ph\]](#).
- [46] T. F. Motta, M. Buballa, and C. S. Fischer, Towards a Stability Analysis of Inhomogeneous Phases in the QCD Phase Diagram, *Acta Phys. Polon. Supp.* **16**, 8 (2023).
- [47] J. Bernhardt and C. S. Fischer, QCD phase transitions in the light quark chiral limit, *Phys. Rev. D* **108**, 114018 (2023), [arXiv:2309.06737 \[hep-ph\]](#).
- [48] F. Ihssen, J. M. Pawłowski, F. R. Sattler, and N. Wink, Towards quantitative precision for QCD at large densities, (2023), [arXiv:2309.07335 \[hep-th\]](#).
- [49] T. F. Motta, J. Bernhardt, M. Buballa, and C. S. Fischer, Toward a stability analysis of inhomogeneous phases in QCD, *Phys. Rev. D* **108**, 114019 (2023), [arXiv:2306.09749 \[hep-ph\]](#).
- [50] N. Zorbach, J. Stoll, and J. Braun, Optimization and Stabilization of Functional Renormalization Group Flows, (2024), [arXiv:2401.12854 \[hep-ph\]](#).
- [51] M. Buballa and S. Carignano, Inhomogeneous chiral condensates, *Prog. Part. Nucl. Phys.* **81**, 39 (2015), [arXiv:1406.1367 \[hep-ph\]](#).
- [52] R. D. Pisarski and F. Rennecke, Signatures of Moat Regimes in Heavy-Ion Collisions, *Phys. Rev. Lett.* **127**, 152302 (2021), [arXiv:2103.06890 \[hep-ph\]](#).
- [53] A. Koenigstein, L. Pannullo, S. Rechenberger, M. J. Steil, and M. Winstel, Detecting inhomogeneous chiral condensation from the bosonic two-point function in the (1 + 1)-dimensional Gross–Neveu model in the mean-field approximation\*, *J. Phys. A* **55**, 375402 (2022), [arXiv:2112.07024 \[hep-ph\]](#).
- [54] K. Kolehmainen and G. Baym, PION CONDENSATION AT FINITE TEMPERATURE. 2. SIMPLE MODELS INCLUDING THERMAL EXCITATIONS OF THE PION FIELD, *Nucl. Phys. A* **382**, 528 (1982).
- [55] E. H. Fradkin, *Field Theories of Condensed Matter Physics*, Vol. 82 (Cambridge Univ. Press, Cambridge, UK, 2013).
- [56] Y. Hidaka, K. Kamikado, T. Kanazawa, and T. Noumi, Phonons, pions and quasi-long-range order in spatially modulated chiral condensates, *Phys. Rev. D* **92**, 034003 (2015), [arXiv:1505.00848 \[hep-ph\]](#).
- [57] T.-G. Lee, E. Nakano, Y. Tsue, T. Tatsumi, and B. Friman, Landau-Peierls instability in a Fulde-Ferrell type inhomogeneous chiral condensed phase, *Phys. Rev. D* **92**, 034024 (2015), [arXiv:1504.03185 \[hep-ph\]](#).
- [58] O. Akerlund, P. de Forcrand, and T. Rindlisbacher, Oscillating propagators in heavy-dense QCD, *JHEP* **10**, 055, [arXiv:1602.02925 \[hep-lat\]](#).
- [59] R. D. Pisarski, V. V. Skokov, and A. M. Tsvetik, Fluctuations in cool quark matter and the phase diagram of Quantum Chromodynamics, *Phys. Rev. D* **99**, 074025 (2019), [arXiv:1801.08156 \[hep-ph\]](#).
- [60] M. A. Schindler, S. T. Schindler, L. Medina, and M. C. Ogilvie, Universality of Pattern Formation, *Phys. Rev. D* **102**, 114510 (2020), [arXiv:1906.07288 \[hep-lat\]](#).
- [61] R. D. Pisarski, A. M. Tsvetik, and S. Valgushev, How transverse thermal fluctuations disorder a condensate of chiral spirals into a quantum spin liquid, *Phys. Rev. D* **102**, 016015 (2020), [arXiv:2005.10259 \[hep-ph\]](#).
- [62] M. A. Schindler, S. T. Schindler, and M. C. Ogilvie,  $\mathcal{PT}$  symmetry, pattern formation, and finite-density QCD [10.1088/1742-6596/2038/1/012022](#) (2021), [arXiv:2106.07092 \[hep-lat\]](#).
- [63] M. A. Schindler, S. T. Schindler, and M. C. Ogilvie, Finite-density QCD,  $\mathcal{PT}$  symmetry, and exotic phases, *PoS LATTICE2021*, 555 (2022), [arXiv:2110.07761 \[hep-lat\]](#).



- [64] D. V. Deryagin, D. Y. Grigoriev, and V. A. Rubakov, Standing wave ground state in high density, zero temperature QCD at large  $N_c$ , *Int. J. Mod. Phys. A* **7**, 659 (1992).
- [65] D. Müller, M. Buballa, and J. Wambach, Dyson-Schwinger study of chiral density waves in QCD, *Phys. Lett. B* **727**, 240 (2013), [arXiv:1308.4303 \[hep-ph\]](#).
- [66] M. Thies and K. Urlichs, Revised phase diagram of the Gross-Neveu model, *Phys. Rev. D* **67**, 125015 (2003), [arXiv:hep-th/0302092](#).
- [67] M. Thies, From relativistic quantum fields to condensed matter and back again: Updating the Gross-Neveu phase diagram, *J. Phys. A* **39**, 12707 (2006), [arXiv:hep-th/0601049](#).
- [68] M. Thies, Chiral spiral in the presence of chiral imbalance, *Physical Review D* **98**, 096019 (2018), [arXiv:1810.03921](#).
- [69] M. Thies, Phase structure of the 1+1 dimensional Nambu–Jona-Lasinio model with isospin, *Physical Review D* **101**, 014010 (2020), [arXiv:1911.11439](#).
- [70] M. Thies, First-order phase boundaries of the massive 1+1 dimensional Nambu–Jona-Lasinio model with isospin, *Physical Review D* **101**, 074013 (2020), [arXiv:2002.01190](#).
- [71] M. Thies, Duality study of the chiral Heisenberg–Gross-Neveu model in 1+1 dimensions, *Phys. Rev. D* **102**, 096006 (2020), [arXiv:2008.13119 \[hep-th\]](#).
- [72] M. Thies, Gross-Neveu model with  $O(2)_L \times O(2)_R$  chiral symmetry: Duality with Zakharov-Mikhailov model and large N solution, *Phys. Rev. D* **107**, 076024 (2023), [arXiv:2302.07660 \[hep-th\]](#).
- [73] J. Lenz, L. Pannullo, M. Wagner, B. Wellegehausen, and A. Wipf, Inhomogeneous phases in the Gross-Neveu model in 1+1 dimensions at finite number of flavors, *Phys. Rev. D* **101**, 094512 (2020), [arXiv:2004.00295 \[hep-lat\]](#).
- [74] J. J. Lenz, M. Mandl, and A. Wipf, Inhomogeneities in the two-flavor chiral Gross-Neveu model, *Phys. Rev. D* **105**, 034512 (2022), [arXiv:2109.05525 \[hep-lat\]](#).
- [75] J. J. Lenz and M. Mandl, Remnants of large- $N_f$  inhomogeneities in the 2-flavor chiral Gross-Neveu model, *PoS LATTICE2021*, 415 (2022), [arXiv:2110.12757 \[hep-lat\]](#).
- [76] J. Stoll, N. Zorbach, A. Koenigstein, M. J. Steil, and S. Rechenberger, Bosonic fluctuations in the (1 + 1)-dimensional Gross-Neveu(-Yukawa) model at varying  $\mu$  and  $T$  and finite  $N$ , (2021), [arXiv:2108.10616 \[hep-ph\]](#).
- [77] R. Ciccone, L. Di Pietro, and M. Serone, Inhomogeneous Phase of the Chiral Gross-Neveu Model, *Phys. Rev. Lett.* **129**, 071603 (2022), [arXiv:2203.07451 \[hep-th\]](#).
- [78] A. Koenigstein, *Non-perturbative aspects of (low-dimensional) quantum field theories*, Ph.D. thesis, Goethe U., Frankfurt (main) (2023).
- [79] R. Ciccone, L. Di Pietro, and M. Serone, Anomalies and Persistent Order in the Chiral Gross-Neveu model, (2023), [arXiv:2312.13756 \[hep-th\]](#).
- [80] M. Kutschera, W. Broniowski, and A. Kotlorz, Quark matter with pion condensate in an effective chiral model, *Nucl. Phys. A* **516**, 566 (1990).
- [81] M. Kutschera, W. Broniowski, and A. Kotlorz, Quark matter with pion condensate in an effective chiral model, *Nucl. Phys. A* **525**, 585C (1991).
- [82] D. Nickel, Inhomogeneous phases in the Nambu-Jona-Lasino and quark-meson model, *Phys. Rev. D* **80**, 074025 (2009), [arXiv:0906.5295 \[hep-ph\]](#).
- [83] S. Carignano, M. Buballa, and B.-J. Schaefer, Inhomogeneous phases in the quark-meson model with vacuum fluctuations, *Phys. Rev. D* **90**, 014033 (2014), [arXiv:1404.0057 \[hep-ph\]](#).
- [84] M. Buballa and S. Carignano, Inhomogeneous chiral phases away from the chiral limit, *Phys. Lett. B* **791**, 361 (2019), [arXiv:1809.10066 \[hep-ph\]](#).
- [85] S. Carignano and M. Buballa, Inhomogeneous chiral condensates in three-flavor quark matter, *Phys. Rev. D* **101**, 014026 (2020), [arXiv:1910.03604 \[hep-ph\]](#).
- [86] M. Buballa, S. Carignano, and L. Kurth, Inhomogeneous phases in the quark-meson model with explicit chiral-symmetry breaking, *Eur. Phys. J. ST* **229**, 3371 (2020), [arXiv:2006.02133 \[hep-ph\]](#).
- [87] L. Pannullo, M. Wagner, and M. Winstel, Inhomogeneous phases in the 3+1-dimensional Nambu-Jona-Lasinio model and their dependence on the regularization scheme, *PoS LATTICE2022*, 156 (2023), [arXiv:2212.05783 \[hep-lat\]](#).
- [88] L. Pannullo, *Inhomogeneous Phases and the Moat Regime in Nambu-Jona-Lasinio-Type Models*, Ph.D. thesis, Goethe U., Frankfurt (main) (2023).
- [89] L. Pannullo, Inhomogeneous condensation in the Gross-Neveu model in noninteger spatial dimensions  $1 \leq d < 3$ , *Phys. Rev. D* **108**, 036022 (2023), [arXiv:2306.16290 \[hep-ph\]](#).
- [90] A. Koenigstein and L. Pannullo, Inhomogeneous condensation in the Gross-Neveu model in noninteger spatial dimensions  $1 \leq d < 3$ . II. Nonzero temperature and chemical potential, (2023), [arXiv:2312.04904 \[hep-ph\]](#).
- [91] K. G. Klimenko, Phase Structure of Generalized Gross-Neveu Models, *Z. Phys. C* **37**, 457 (1988).
- [92] B. Rosenstein, B. J. Warr, and S. H. Park, The Four Fermi Theory Is Renormalizable in (2+1)-Dimensions, *Phys. Rev. Lett.* **62**, 1433 (1989).
- [93] S. Hands, A. Kocic, and J. B. Kogut, The Four Fermi model in three-dimensions at nonzero density and temperature, *Nucl. Phys. B* **390**, 355 (1993), [arXiv:hep-lat/9206024](#).
- [94] T. Inagaki, T. Kouno, and T. Muta, Phase structure of four fermion theories at finite temperature and chemical potential in arbitrary dimensions, *Int. J. Mod. Phys. A* **10**, 2241 (1995), [arXiv:hep-ph/9409413](#).
- [95] L. Del Debbio and S. Hands, Monte Carlo simulation of the three-dimensional Thirring model, *Phys. Lett. B* **373**, 171 (1996), [arXiv:hep-lat/9512013](#).
- [96] T. Appelquist and M. Schwetz, The (2+1)-dimensional NJL model at finite temperature, *Phys. Lett. B* **491**, 367 (2000), [arXiv:hep-ph/0007284](#).

- [97] S. Hands, J. B. Kogut, C. G. Strouthos, and T. N. Tran, Fermi surface phenomena in the (2+1)-d four Fermi model, *Phys. Rev. D* **68**, 016005 (2003), arXiv:hep-lat/0302021.
- [98] D. Ebert, K. G. Klimenko, P. B. Kolmakov, and V. C. Zhukovsky, Phase transitions in hexagonal, graphene-like lattice sheets and nanotubes under the influence of external conditions, *Annals Phys.* **371**, 254 (2016), arXiv:1509.08093 [cond-mat.mes-hall].
- [99] M. Mandl, J. J. Lenz, and A. Wipf, Magnetic catalysis in the 1-flavor Gross-Neveu model in 2 + 1 dimensions, *PoS LATTICE2022*, 152 (2023), arXiv:2211.10333 [hep-lat].
- [100] J. J. Lenz, M. Mandl, and A. Wipf, Magnetic catalysis in the (2+1)-dimensional Gross-Neveu model, *Phys. Rev. D* **107**, 094505 (2023), arXiv:2302.05279 [hep-lat].
- [101] S. Hands, B. Lucini, and S. Morrison, Critical behavior in the dense planar NJL model, *Phys. Rev. Lett.* **86**, 753 (2001), arXiv:hep-lat/0008027.
- [102] R. Narayanan, Phase diagram of the large  $N$  Gross-Neveu model in a finite periodic box, *Phys. Rev. D* **101**, 096001 (2020), arXiv:2001.09200 [hep-th].
- [103] M. Buballa, L. Kurth, M. Wagner, and M. Winstel, Regulator dependence of inhomogeneous phases in the (2+1)-dimensional Gross-Neveu model, *Phys. Rev. D* **103**, 034503 (2021), arXiv:2012.09588 [hep-lat].
- [104] L. Pannullo, M. Wagner, and M. Winstel, Inhomogeneous Phases in the Chirally Imbalanced 2 + 1-Dimensional Gross-Neveu Model and Their Absence in the Continuum Limit, *Symmetry* **14**, 265 (2022), arXiv:2112.11183 [hep-lat].
- [105] M. Winstel and L. Pannullo, Stability of homogeneous chiral phases against inhomogeneous perturbations in 2+1 dimensions, in *39th International Symposium on Lattice Field Theory* (2022) arXiv:2211.04414 [hep-ph].
- [106] C. Sasaki, B. Friman, and K. Redlich, Quark Number Fluctuations in a Chiral Model at Finite Baryon Chemical Potential, *Phys. Rev. D* **75**, 054026 (2007), arXiv:hep-ph/0611143.
- [107] K. Redlich, B. Friman, and C. Sasaki, QCD phase diagram and charge fluctuations, *J. Phys. G* **34**, S437 (2007), arXiv:hep-ph/0702296.
- [108] S. Carignano, D. Nickel, and M. Buballa, Influence of vector interaction and Polyakov loop dynamics on inhomogeneous chiral symmetry breaking phases, *Phys. Rev. D* **82**, 054009 (2010), arXiv:1007.1397 [hep-ph].
- [109] S. Carignano, M. Schramm, and M. Buballa, Influence of vector interactions on the favored shape of inhomogeneous chiral condensates, *Phys. Rev. D* **98**, 014033 (2018), arXiv:1805.06203 [hep-ph].
- [110] M. Haensch, F. Rennecke, and L. von Smekal, Medium Induced Mixing and Critical Modes in QCD, (2023), arXiv:2308.16244 [hep-ph].
- [111] J. Braun, L. Fister, J. M. Pawłowski, and F. Rennecke, From Quarks and Gluons to Hadrons: Chiral Symmetry Breaking in Dynamical QCD, *Phys. Rev. D* **94**, 034016 (2016), arXiv:1412.1045 [hep-ph].
- [112] F. Rennecke, Vacuum structure of vector mesons in QCD, *Phys. Rev. D* **92**, 076012 (2015), arXiv:1504.03585 [hep-ph].
- [113] K. Fukushima, J. M. Pawłowski, and N. Strodthoff, Emergent hadrons and diquarks, *Annals Phys.* **446**, 169106 (2022), arXiv:2103.01129 [hep-ph].
- [114] R. D. Pisarski, Chiral Symmetry Breaking in Three-Dimensional Electrodynamics, *Phys. Rev. D* **29**, 2423 (1984).
- [115] M. Buballa, NJL model analysis of quark matter at large density, *Phys. Rept.* **407**, 205 (2005), arXiv:hep-ph/0402234.
- [116] G. Gat, A. Kovner, and B. Rosenstein, Chiral phase transitions in  $d = 3$  and renormalizability of four Fermi interactions, *Nucl. Phys. B* **385**, 76 (1992).
- [117] E. Nakano and T. Tatsumi, Chiral symmetry and density wave in quark matter, *Phys. Rev. D* **71**, 114006 (2005), arXiv:hep-ph/0411350.
- [118] A. Alexandru, G. Basar, P. F. Bedaque, and N. C. Warrington, Complex paths around the sign problem, *Rev. Mod. Phys.* **94**, 015006 (2022), arXiv:2007.05436 [hep-lat].
- [119] Y. Mori, K. Kashiwa, and A. Ohnishi, Lefschetz thimbles in fermionic effective models with repulsive vector-field, *Phys. Lett. B* **781**, 688 (2018), arXiv:1705.03646 [hep-lat].
- [120] G. Van Rossum and F. L. Drake, *Python 3 Reference Manual* (CreateSpace, Scotts Valley, CA, 2009).
- [121] P. Virtanen *et al.*, SciPy 1.0—Fundamental Algorithms for Scientific Computing in Python, *Nature Meth.* **17**, 261 (2020), arXiv:1907.10121 [cs.MS].
- [122] C. R. Harris, K. J. Millman, S. J. van der Walt, R. Gommers, P. Virtanen, D. Cournapeau, E. Wieser, J. Taylor, S. Berg, N. J. Smith, R. Kern, M. Picus, S. Hoyer, M. H. van Kerkwijk, M. Brett, A. Haldane, J. F. del Río, M. Wiebe, P. Peterson, P. Gérard-Marchant, K. Sheppard, T. Reddy, W. Weckesser, H. Abbasi, C. Gohlke, and T. E. Oliphant, Array programming with numpy, *Nature* **585**, 357–362 (2020).
- [123] D. D. Scherer and H. Gies, Renormalization Group Study of Magnetic Catalysis in the 3d Gross-Neveu Model, *Phys. Rev. B* **85**, 195417 (2012), arXiv:1201.3746 [cond-mat.str-el].
- [124] H. Nishimura, M. C. Ogilvie, and K. Pangaeni, Complex Saddle Points and Disorder Lines in QCD at finite temperature and density, *Phys. Rev. D* **91**, 054004 (2015), arXiv:1411.4959 [hep-ph].
- [125] A. Felski, A. Beygi, and S. P. Klevansky, Non-Hermitian extension of the Nambu–Jona-Lasinio model in 3+1 and 1+1 dimensions, *Phys. Rev. D* **101**, 116001 (2020), arXiv:2004.04011 [hep-ph].
- [126] C. M. Bender and S. Boettcher, Real spectra in nonHermitian Hamiltonians having PT symmetry, *Phys. Rev. Lett.* **80**, 5243 (1998), arXiv:physics/9712001.
- [127] R. El-Ganainy, K. G. Makris, M. Khajavikhan, Z. H. Musslimani, S. Rotter, and D. N. Christodoulides, Non-Hermitian physics and PT symmetry, *Nature Phys.* **14**, 11 (2018).
- [128] M.-A. Miri and A. Alù, Exceptional points in optics and photonics, *Science* **363**, eaar7709 (2019).
- [129] Y. Ashida, Z. Gong, and M. Ueda, Non-Hermitian physics, *Adv. Phys.* **69**, 249 (2021), arXiv:2006.01837 [cond-mat.mes-hall].

- [130] C. M. Bender and D. W. Hook, PT-symmetric quantum mechanics, (2023), [arXiv:2312.17386 \[quant-ph\]](#).
- [131] P. N. Meisinger and M. C. Ogilvie, PT Symmetry in Classical and Quantum Statistical Mechanics, *Phil. Trans. Roy. Soc. Lond. A* **371**, 20120058 (2013), [arXiv:1208.5077 \[math-ph\]](#).
- [132] M. C. Ogilvie and L. Medina, Simulation of Scalar Field Theories with Complex Actions, *PoS LATTICE2018*, 157 (2018), [arXiv:1811.11112 \[hep-lat\]](#).
- [133] M. C. Ogilvie, M. A. Schindler, and S. T. Schindler, Finite-density QCD,  $\mathcal{PT}$  symmetry, and dual algorithms, *PoS LATTICE2021*, 417 (2022), [arXiv:2110.14009 \[hep-lat\]](#).
- [134] F. Rennecke, R. D. Pisarski, and D. H. Rischke, Particle interferometry in a moat regime, *Phys. Rev. D* **107**, 116011 (2023), [arXiv:2301.11484 \[hep-ph\]](#).
- [135] K. Urlichs, *Baryons and baryonic matter in four-fermion interaction models*, Ph.D. thesis, Universität Erlangen-Nuremberg (2007).
- [136] J. D. Hunter, Matplotlib: A 2d graphics environment, *Computing in Science & Engineering* **9**, 90 (2007).
- [137] C. G. Strouthos, Mesons at finite baryon density in (2+1)-dimension, *Prog. Theor. Phys. Suppl.* **153**, 69 (2004), [arXiv:hep-lat/0308021](#).
- [138] Y. Cohen, S. Elitzur, and E. Rabinovici, A MONTE CARLO STUDY OF THE GROSS-NEVEU MODEL, *Nucl. Phys. B* **220**, 102 (1983).
- [139] A. Alexandru, G. Basar, P. F. Bedaque, G. W. Ridgway, and N. C. Warrington, Monte Carlo calculations of the finite density Thirring model, *Phys. Rev. D* **95**, 014502 (2017), [arXiv:1609.01730 \[hep-lat\]](#).
- [140] K. G. Klimenko, Three-dimensional (anti-psi psi)\*\*2 model and optimized expansion, *Mod. Phys. Lett. A* **9**, 1767 (1994).
- [141] J.-L. Kneur, M. B. Pinto, R. O. Ramos, and E. Staudt, Updating the phase diagram of the Gross-Neveu model in 2+1 dimensions, *Phys. Lett. B* **657**, 136 (2007), [arXiv:0705.0673 \[hep-ph\]](#).
- [142] M. Cristoforetti, F. Di Renzo, and L. Scorzato (AuroraScience), New approach to the sign problem in quantum field theories: High density QCD on a Lefschetz thimble, *Phys. Rev. D* **86**, 074506 (2012), [arXiv:1205.3996 \[hep-lat\]](#).
- [143] K. Fukushima, Y. Hidaka, K. Inoue, K. Shigaki, and Y. Yamaguchi, HBT signature for clustered substructures probing primordial inhomogeneity in hot and dense QCD matter, (2023), [arXiv:2306.17619 \[hep-ph\]](#).