Effects of diffusion and advection on predator prey dynamics in an advective patchy environment

Qi Wang *

Abstract

In this paper, we consider a specialist predator-prey patchy model over the closed stream network. We study the dynamics and the asymptotic profiles of positive steady states according to the mortality rate of the specialist predators, advection and diffusion rates. We verify that the specialist predators can successfully invade as long as the mortality rate is sufficiently small. On the other hand, the impacts of diffusion and advection on the asymptotic profiles of positive steady states and on the concentration of the species are given.

Keywords: Patchy environment; Predator-prey system; Global dynamics; Asymptotic profiles Mathematics Subject Classification (2010): 34D05, 34D23, 37C75, 92D25, 92D40

1 Introduction

It is well-known that whether the individuals survive or not in a predominantly unidirectional flow environments (such as streams, rivers, lakes or oceans) depends on both the environment and the growth of interacting species. Sometimes, despite the flow induced washout, the species can always persist in their habitats for many generations. This phenomenon is so-called "drift paradox" [19,20,23]. To explain this paradox, Speirs and Gurney [29] proposed a reaction-diffusion-advection equation and indicated that the persistence of single species is possible when the flow speed is slow relative to the diffusion and the stream is long enough. Intuitively, the predominantly unidirectional flow will carry individuals to the downstream end or drive individuals out of the system, which may be crowded or hostile. However, random dispersal may drive them to some favorable locations in the upstream [7]. Therefore, one can see the joint impacts of both undirectional and directed dispersal rates on the population dynamics of the species are usually complicated and have attracted the attention of many researchers [6,9,11,17,18,20,29].

In real ecological advective environments, there are various inter-specific relationships including competition, predation. The two-species reaction-diffusion-advection competition models have been widely studied in [4, 10, 11, 13-15, 21, 30-32, 35-38]. Among them, some results illustrate the conditions under which the species u or v is stable/unstable. Various results on the global dynamics of the two-species reaction-diffusion-advection competition models are also presented.

Regarding the dynamics of two species competition models in a river network, the authors in [1-3,7,8] investigated a two-species Lotka-Volterra competition patchy model over a inland stream. Let $n \ge 3$, $\boldsymbol{u} = (u_1, u_2, \dots, u_n)$ and $\boldsymbol{v} = (v_1, v_2, \dots, v_n)$ be the population densities of two competing species, respectively, where u_i and v_i are the densities in patch *i*. Suppose that the dispersal patterns of the individuals and the configuration of the patches are shown in Fig.??

^{*}College of Science, University of Shanghai for Science and Technology, Shanghai 200093, P.R. China. Email: qwang@usst.edu.cn.

The competition patchy model over the stream network in Fig. ?? in [1-3, 7, 8] is:

$$\begin{cases} \frac{\mathrm{d}u_i}{\mathrm{d}t} = \sum_{j=1}^n (d_1 D_{ij} + q_1 Q_{ij}) u_j + r_i u_i (1 - \frac{u_i + v_i}{k_i}), & i \in \{1, 2, \cdots, n\}, \ t > 0, \\ \frac{\mathrm{d}v_i}{\mathrm{d}t} = \sum_{j=1}^n (d_2 D_{ij} + q_2 Q_{ij}) v_j + r_i v_i (1 - \frac{u_i + v_i}{k_i}), & i \in \{1, 2, \cdots, n\}, \ t > 0, \\ u(0) = u_0 \ge \neq 0, \ v(0) = v_0 \ge \neq 0, \end{cases}$$
(1.1)

where d_1, d_2 are random movement rates; q_1, q_2 represent directed movement rates; the matrices $D = (D_{ij})$ and $Q = (Q_{ij})$ are the random movement pattern and directed drift pattern of individuals, respectively, where

$$D = \begin{pmatrix} -1 & 1 & 0 & \cdots & \cdots & 0\\ 1 & -2 & 1 & \cdots & \cdots & 0\\ 0 & \ddots & \ddots & \ddots & & \\ \vdots & & \ddots & \ddots & \ddots & \\ 0 & \cdots & \cdots & 1 & -2 & 1\\ 0 & \cdots & \cdots & 1 & -1 \end{pmatrix}, Q = \begin{pmatrix} -1 & 0 & 0 & \cdots & \cdots & 0\\ 1 & -1 & 0 & \cdots & \cdots & 0\\ 0 & \ddots & \ddots & \ddots & \\ \vdots & & \ddots & \ddots & \\ 0 & \cdots & \cdots & 1 & -1 & 0\\ 0 & \cdots & \cdots & 1 & 0 \end{pmatrix}.$$
(1.2)

Recently, Chen, Liu and Wu [2] considered (1.1) with three patches. They studied if one species is treated as a resident species, whether the other species can invade or not under some assumptions on the carrying capacity.

Predation is another common phenomenon in advective environments. For instance, herbivorous zooplankton and phytoplankton in water columns. Compared with the competitive systems, up to now, the studies of the evolution of dispersal in predator-prey systems under advective environments are relatively few. How do the prey persist in advective environments and avoid predations successfully? How do the predators invade the prey and coexist in advective environments? To better understand these issues, Hilker and Lewis [5] proposed the following predator-prey system in advective environments.

$$\begin{cases} N_t = d_1 N_{xx} - q N_x + f(N, P) N, & x \in (0, L), t > 0, \\ P_t = d_2 P_{xx} - \tau q P_x + g(N, P) P, & x \in (0, L), t > 0. \end{cases}$$
(1.3)

Here N(x,t) and P(x,t) are the population densities of the prey and predator species at time t and location x, respectively. d_1, d_2 are dispersal rates, q denotes the effective advection rate of the prey, L is the domain length. The function f(N, P) is given by

$$f(N,P) = r_1(x)(1 - \frac{N}{K_1(x)}) - aP,$$
(1.4)

and the function g(N, P) is given by

$$g(N, P) = \beta aN - \gamma \text{ or } g(N, P) = r_2(x)(1 - \frac{P}{K_2(x)}) + \beta aN,$$
 (1.5)

where $r_i(x)(i = 1, 2)$ and $K_i(x)(i = 1, 2)$ account for the intrinsic growth rate and the carrying capacity of the prey and predators at location x, respectively, a is the predation rate, β is the trophic conversion efficiency, and γ is the mortality rate of the predators.

By numerical simulations and traveling wave speed approximations, Hilker and Lewis achieved the coexistence, extinction of both species and survival of one species. Based on [5], some mathematicians studied the dynamics of the predator-prey model in open advective environments further [12, 25, 26, 34]. Furthermore, Wang et al. in [27, 33] investigated the dynamics of some predator-prey models in closed advective environments and obtained some significant outcomes.

Motivated by [2, 27], in this paper we let $n \ge 3$, $N = (N_1, N_2, \dots, N_n)$, $P = (P_1, P_2, \dots, P_n)$ and consider the following specialist predator-prev patchy model over the closed stream network in Fig.??:

$$\begin{cases} \frac{\mathrm{d}N_i}{\mathrm{d}t} = \sum_{j=1}^n (d_1 D_{ij} + q Q_{ij}) N_j + r_i N_i (1 - \frac{N_i}{k_i}) - a N_i P_i, & i \in \{1, 2, \cdots, n\}, \ t > 0, \\ \frac{\mathrm{d}P_i}{\mathrm{d}t} = \sum_{j=1}^n (d_2 D_{ij} + \tau q Q_{ij}) P_j + P_i (\beta a N_i - \gamma), & i \in \{1, 2, \cdots, n\}, \ t > 0, \\ \mathbf{N}(0) \ge \neq \mathbf{0}, \ \mathbf{P}(0) \ge \neq \mathbf{0}, \end{cases}$$
(1.6)

where N_i and P_i are the densities of the prey and predator in patch *i* respectively. Here $r_i (i \in \{1, 2, \dots, n\})$ and $k_i (i \in \{1, 2, \dots, n\})$ account for the intrinsic growth rate and the carrying capacity of the prey in patch *i*, respectively. The meanings of a, β, γ are the same as them in (1.5).

The goal of this paper is to study the dynamics of system (1.6), and investigate the similarities or differences between (1.6) and the specialist predator-prey model in [27].

For convenience, we set $\mathcal{A} = \{1, 2, \dots, n\}, \mathcal{B} = \{2, \dots, n-1\}, \mathcal{C} = \{1, \dots, n-1\}, \mathbf{r} = (r_1, r_2, \dots, r_n), \mathbf{r} + b = (r_1 + b, r_2 + b, \dots, r_n + b), r_{max} = \max_{i \in \mathcal{A}} r_i, r_{min} = \min_{i \in \mathcal{A}} r_i.$ We write $\mathbf{r} \ge \mathbf{0} (\mathbf{r} \gg \mathbf{0})$ if $r_i \ge 0 (r_i > 0)$ for all $i \in \mathcal{A}$, and $\mathbf{r} > \mathbf{0}$ if $\mathbf{r} \ge \mathbf{0}$ and $\mathbf{r} \neq \mathbf{0} := (0, \dots, 0).$

At the end of this section, we make some assumptions throughout this paper:

$$\mathbf{r}, \mathbf{k} \gg \mathbf{0}, d_1, d_2, a, \beta, \tau > 0, q \ge 0$$
 and there exist $i \ne j$ such that $k_i \ne k_j$. (H)

The rest of this paper is organized as follows. In Section 2, we state the main results. In Section 3, we provide some preliminary results. In Section 4, we show our main results.

2 Main Results

Before describing precisely the outcome of our paper, we first recall some existing results on the dynamics of the following single-species model:

$$\begin{cases} \frac{\mathrm{d}N_i}{\mathrm{d}t} = \sum_{j=1}^n (d_1 D_{ij} + q Q_{ij}) N_j + r_i N_i (1 - \frac{N_i}{k_i}), & i \in \mathcal{A}, \ t > 0, \\ \mathbf{N}(0) > \mathbf{0}. \end{cases}$$
(2.1)

It is known in [2, Lemma 2.2](see also [16]) that system (2.1) admits a unique positive steady state, denoted by $\boldsymbol{\theta} = (\theta_1, \dots, \theta_n) \gg 0$, which is globally asymptotically stable for any $d_1 > 0$ and $q \ge 0$.

We are now in a position to state our main results.

Theorem 2.1. Suppose (**H**) hold. Let $(\mathbf{N}(t), \mathbf{P}(t))$ be the solution of (1.6). There exists a critical curve $\gamma = \gamma^*(q, d_1, d_2) \in (0, +\infty)$ continuously dependent on q, d_1, d_2 such that:

(1) if $\gamma > \gamma^*$, then $(\mathbf{N}(t), \mathbf{P}(t))$ converges uniformly to $(\boldsymbol{\theta}, \mathbf{0})$ as $t \to +\infty$;

(2) if $\gamma < \gamma^*$, then system (1.6) is uniformly persistent in the sense that there exists $\eta > 0$ such that the solution $(\mathbf{N}(t), \mathbf{P}(t))$ satisfies $\liminf_{t \to +\infty} (N(t))_{min} \ge \eta$, $\liminf_{t \to +\infty} (P(t))_{min} \ge \eta$. Moreover, system (1.6) admits a positive steady state.

Remark 2.1. Theorem 2.1 indicates that the dynamical behavior of system (1.6) is depicted completely by the critical curve $\gamma = \gamma^*(q, d_1, d_2)$. More precisely, $\gamma = \gamma^*(q, d_1, d_2)$ classifies the dynamical behavior of system (1.6) into two scenarios: (i) coexistence; and (ii) persistence of prey only. Biologically, a large death rate of the predators makes it difficult to invade, while the predators can invade successfully only when the mortality rate of the predators is less than the critical mortality rate γ^* .

Theorem 2.2. Suppose (**H**) hold. The threshold value $\gamma^*(q, d_1, d_2)$ satisfies the following properties: (1) $\lim_{a \to +\infty} \gamma^*(q, d_1, d_2) = \beta a k_n$ for $d_1, d_2 > 0$;

(2)
$$\lim_{d_1 \to +\infty} \gamma^*(q, d_1, d_2) = \frac{\beta a \sum_{i=1}^n r_i}{\sum_{i=1}^n \frac{r_i}{k_i}} \text{ for } d_2 > 0, \ q \ge 0,$$

(3) if $q \ge \max\{r_{max} + 2\sqrt{d_1r_{max}}, 2r_{max}\}$, then $\lim_{d_1 \to 0^+} \gamma^*(q, d_1, d_2) = \mu(k_n)$ for $d_2 > 0$, q > 0, where $\mu(k_n) > 0$ is the principal eigenvalue of the following system

$$\begin{cases} -(d_2 + \tau q)\vartheta_1 + d_2\vartheta_2 = \mu(k_n)\vartheta_1, \\ (d_2 + \tau q)\vartheta_{j-1} + (-2d_2 - \tau q)\vartheta_j + d_2\vartheta_{j+1} = \mu(k_n)\vartheta_j, \ j \in \mathcal{B}, \\ (d_2 + \tau q)\vartheta_{n-1} - d_2\vartheta_n + \beta ak_n\vartheta_n = \mu(k_n)\vartheta_n, \end{cases}$$
(2.2)

and $\mu(k_n)$ is strictly increasing with respect to k_n ;

(4) $\lim_{d_1\to 0^+} \gamma^*(0, d_1, d_2) = \lambda_1(d_2, 0, \beta a \mathbf{k}) > 0$ for $d_2 > 0$, where $\lambda_1(d_2, 0, \beta a \mathbf{k})$ is defined in subsection 3.1:

 $(5) \lim_{d_2 \to 0^+} \gamma^*(q, d_1, d_2) = \max\{\beta a\theta_1 - \tau q, \cdots, \beta a\theta_{n-1} - \tau q, \beta a\theta_n\}, and \lim_{d_2 \to +\infty} \gamma^*(q, d_1, d_2) = \frac{\sum_{i=1}^n \beta a\theta_i}{n}$ for $d_1 > 0, q \ge 0$.

Remark 2.2. It is easy to see that $\mu(0) = 0$.

Theorem 2.2 (1), (2) and (5) indicate that in closed advective patchy environments $\gamma^*(q, d_1, d_2)$ has a positive lower bound for any $q \ge 0$ and $d_1, d_2 > 0$. This implies that if the prey's diffusion rate $d_1 > 0$ is fixed, then the specialist predators can invade successfully as long as their death rate is suitably small. For a small diffusion rate d_1 of the prey, (3) and (4) show that the specialist predators can always invade successfully in both non-advective and advective environments. Moreover, given sufficiently small death rate of the specialist predators, when the carrying capacity of the prey at patch n increases, it is more convenient for the predators to invade. This phenomenon is different from the one in [27], where they verified that it is difficult for the specialist predators to invade in advective environments, when d_1 is small enough.

Next, we investigate the asymptotic profiles of the positive steady state (N^*, P^*) .

Theorem 2.3. Suppose (**H**) hold, and $0 < \gamma < \beta a k_n$. Let $(\mathbf{N}^*, \mathbf{P}^*)$ be a positive steady state of (1.6). Then it satisfies $(N_i^*, P_i^*) \to (0, 0)$ uniformly for $i \in \mathcal{C}$ as $q \to +\infty$, and $(N_n^*, P_n^*) \to (\frac{\gamma}{\beta a}, \frac{1}{a}(r_n - \frac{r_n}{k_n}\frac{\gamma}{\beta a}))$ as $q \to +\infty$.

Theorem 2.4. Suppose (**H**) hold, $0 < \gamma < \beta a k_{min}$. Let $(\mathbf{N}^*, \mathbf{P}^*)$ be a positive steady state of (1.6). Then it satisfies

$$(N_i^*, P_i^*) \to (\frac{\gamma}{\beta a}, (1 + \frac{\tau q}{d_2})^{i-1} \frac{\sum_{j=1}^n r_j (1 - \frac{\gamma}{\beta a k_j})}{\frac{d_2 a}{\tau q} [(1 + \frac{\tau q}{d_2})^n - 1]})$$
(2.3)

uniformly for $i \in \mathcal{A}$ as $d_1 \to +\infty$.

Theorem 2.5. Suppose (**H**) hold, $0 < \gamma < \gamma^*(q, d_1, d_2)$, and $q > \max\left\{r_{max} + 2\sqrt{d_1r_{max}}, \frac{2\beta aN_n^*}{\tau}\right\}$. Let $(\mathbf{N}^*, \mathbf{P}^*)$ be a positive steady state of (1.6). Then it satisfies $(N_i^*, P_n^*) \to (\theta_i, 0)$ as $d_2 \to 0^+$. Moreover,

$$\max_{i \in \mathcal{C}} \left(P_i^* - P_n^* \left(\frac{d_1}{d_2 + \tau q} \right)^{n-i} \right) \to 0 \text{ as } d_2 \to 0^+.$$
(2.4)

Theorem 2.6. Suppose (**H**) hold, $0 < \gamma < \frac{\beta a \sum_{i}^{n} \theta_{i}}{n}$. Let $(\mathbf{N}^{*}, \mathbf{P}^{*})$ be a positive steady state of (1.6). Then it satisfies $(N_{i}^{*}, P_{n}^{*}) \rightarrow (\hat{\theta}_{i}, z)$ as $d_{2} \rightarrow +\infty$, where $(\hat{\theta}_{i}, z)$ satisfies

$$\begin{cases} \sum_{j=1}^{n} (d_1 D_{ij} + q Q_{ij}) \hat{\theta}_j + \hat{\theta}_i (r_i - \frac{r_i}{k_i} \hat{\theta}_i - az) = 0, \quad i \in \mathcal{A}, \\ \beta a \sum_{i=1}^{n} \hat{\theta}_i = \gamma n, \qquad \qquad i \in \mathcal{A}. \end{cases}$$

$$(2.5)$$

Theorem 2.3 implies that sufficiently large advection will affect the distribution of species in the network. More precisely, two species will coexist and concentrate at the patch n if the advection rate is large enough. Theorems 2.4 and 2.6 show that two species can coexist for large diffusion rates. Theorem 2.2(5) and 2.5 illustrate the successful invasion of the specialist predators once diffusion rate d_2 is enough small, but their population density tends to zero as the diffusion rate tends to zero. This phenomenon is partially due to the lack of other food sources of the specialist predators except the prey. Another reason is that the specialist predators cannot catch up with the prey when their diffusion rate is sufficiently small.

3 Preliminaries

In this section, we aim to exhibit some fundamental results, which will be utilized in later sections.

3.1 The linear eigenvalue problem

We first consider an auxiliary linear eigenvalue problem.

$$\sum_{j=1}^{n} (dD_{ij} + qQ_{ij})\phi_j + h_i\phi_i = \lambda\phi_i, \quad i \in \mathcal{A},$$
(3.1)

where $d > 0, q \ge 0$. It is well-known from [1,2] that problem (3.1) admits a unique principal eigenvalue $\lambda_1(d, q, \mathbf{h}) = s(dD + qQ + diag(h_i))$ (which is the spectral bound of $dD + qQ + diag(h_i)$) associated with a nonnegative eigenvector $\boldsymbol{\phi} = (\phi_1, \dots, \phi_n) > 0$, since the matrix $dD + qQ + diag(h_i)$ is irreducible and essentially nonnegative for any $d > 0, q \ge 0$, where D and Q are defined by (1.2).

Lemma 3.1. Let D and Q be defined in (1.2), d, q > 0, and $\lambda_1(d, q, h)$ be the principal eigenvalue of (3.1). Suppose that $h \gg 0$ and there exist $i \neq j$ such that $h_i \neq h_j$. Then, we have the following:

(i) for each fixed d > 0, $\lambda_1(d, q, h)$ is strictly decreasing with respect to q in $[0, +\infty)$ when $h_1 \ge \cdots \ge h_n$ with at least one strict inequality, and $\lambda_1(d, q, h)$ is strictly increasing with respect to q when $h_1 \le \cdots \le h_n$ with at least one strict inequality;

(*ii*) $\lim_{q \to +\infty} \lambda_1(d, q, h) = h_n;$

(*iii*)
$$\lim_{d\to 0} \lambda_1(d, q, \mathbf{h}) = \max\{h_1 - q, \cdots, h_{n-1} - q, h_n\}$$
 and $\lim_{d\to +\infty} \lambda_1(d, q, \mathbf{h}) = \frac{\sum_{i=1}^n h_i}{n};$
(*iv*) if $\mathbf{h} > \mathbf{g}$, then $\lambda_1(d, q, \mathbf{h}) \ge \lambda_1(d, q, \mathbf{g})$, and the equality holds only if $\mathbf{h} = \mathbf{g}$.

Proof: The proofs of (i)-(iii) are standard and we refer to [1] for the proofs. We now show the statement (iv). Indeed, we can rewrite (3.1) as the following forms:

$$\begin{cases} -(d+q)\phi_1 + d\phi_2 + h_1\phi_1 = \lambda_1\phi_1, \\ (d+q)\phi_{j-1} + (-2d-q)\phi_j + d\phi_{j+1} + h_j\phi_j = \lambda_1\phi_j, \ j \in \mathcal{B}, \\ (d+q)\phi_{n-1} - d\phi_n + h_n\phi_n = \lambda_1\phi_n. \end{cases}$$
(3.2)

Differentiating (3.2) with respect to h_1 , we have

$$\begin{cases} -(d+q)\frac{\partial\phi_1}{\partial h_1} + d\frac{\partial\phi_2}{\partial h_1} + h_1\frac{\partial\phi_1}{\partial h_1} + \phi_1 = \frac{\partial\lambda_1}{\partial h_1}\phi_1 + \lambda_1\frac{\partial\phi_1}{\partial h_1}, \\ (d+q)\frac{\partial\phi_{j-1}}{\partial h_1} + (-2d-q)\frac{\partial\phi_j}{\partial h_1} + d\frac{\partial\phi_{j+1}}{\partial h_1} + h_j\frac{\partial\phi_j}{\partial h_1} = \frac{\partial\lambda_1}{\partial h_1}\phi_j + \lambda_1\frac{\partial\phi_j}{\partial h_1}, \ j \in \mathcal{B}, \\ (d+q)\frac{\partial\phi_{n-1}}{\partial h_1} - d\frac{\partial\phi_n}{\partial h_1} + h_n\frac{\partial\phi_n}{\partial h_1} = \frac{\partial\lambda_1}{\partial h_1}\phi_n + \lambda_1\frac{\partial\phi_n}{\partial h_1}. \end{cases}$$
(3.3)

Multiplying the first(second, third) equation of (3.3) by $\phi_1(\phi_j, \phi_n)$ and the first(second, third) equation of (3.2) by $\frac{\partial \phi_1}{\partial h_1} (\frac{\partial \phi_j}{\partial h_1}, \frac{\partial \phi_n}{\partial h_1})$, and subtracting each other, it then follows that

$$\begin{cases} d(\frac{\partial\phi_2}{\partial h_1}\phi_1 - \phi_2\frac{\partial\phi_1}{\partial h_1}) + \phi_1^2 = \frac{\partial\lambda_1}{\partial h_1}\phi_1^2, \\ (d+q)(\frac{\partial\phi_{j-1}}{\partial h_1}\phi_j - \phi_{j-1}\frac{\partial\phi_j}{\partial h_1}) + d(\frac{\partial\phi_{j+1}}{\partial h_1}\phi_j - \phi_{j+1}\frac{\partial\phi_j}{\partial h_1}) = \frac{\partial\lambda_1}{\partial h_1}\phi_j^2, \ j \in \mathcal{B}, \\ (d+q)(\frac{\partial\phi_{n-1}}{\partial h_1}\phi_n - \phi_{n-1}\frac{\partial\phi_n}{\partial h_1}) = \frac{\partial\lambda_1}{\partial h_1}\phi_n^2. \end{cases}$$
(3.4)

Let $b_i = \left(\frac{d}{d+q}\right)^{i-1}$. Then we have that $b_i d = b_{i+1}(d+q)$. This implies that

$$b_{1}d(\frac{\partial\phi_{2}}{\partial h_{1}}\phi_{1}-\phi_{2}\frac{\partial\phi_{1}}{\partial h_{1}}\phi_{1}')+b_{2}(d+q)(\frac{\partial\phi_{1}}{\partial h_{1}}\phi_{2}-\phi_{1}\frac{\partial\phi_{2}}{\partial h_{1}})$$

$$+b_{2}d(\frac{\partial\phi_{3}}{\partial h_{1}}\phi_{2}-\phi_{3}\frac{\partial\phi_{2}}{\partial h_{1}})+b_{3}(d+q)(\frac{\partial\phi_{2}}{\partial h_{1}}\phi_{3}-\phi_{2}\frac{\partial\phi_{3}}{\partial h_{1}})$$

$$+b_{3}d(\frac{\partial\phi_{4}}{\partial h_{1}}\phi_{3}-\phi_{4}\frac{\partial\phi_{3}}{\partial h_{1}})+b_{4}(d+q)(\frac{\partial\phi_{3}}{\partial h_{1}}\phi_{4}-\phi_{3}\frac{\partial\phi_{4}}{\partial h_{1}})$$

$$+\cdots$$

$$+b_{n-1}d(\frac{\partial\phi_{n}}{\partial h_{1}}\phi_{n-1}-\phi_{n}\frac{\partial\phi_{n-1}}{\partial h_{1}})+b_{n}(d+q)(\frac{\partial\phi_{n-1}}{\partial h_{1}}\phi_{n}-\phi_{n-1}\frac{\partial\phi_{n}}{\partial h_{1}})$$

$$=0.$$
(3.5)

Then by (3.4) and (3.5), there holds

$$\frac{\partial \lambda_1}{\partial h_1} (b_1 \phi_1^2 + \dots + b_n \phi_n^2) = b_1 \phi_1^2 > 0.$$
(3.6)

Therefore, $\frac{\partial \lambda_1}{\partial h_1} > 0$. Similarly, one can have that $\frac{\partial \lambda_1}{\partial h_i} > 0$ for all $i \in \mathcal{A}$. The proof is completed. \Box

Lemma 3.2. Let D and Q be defined in (1.2), $d > 0, q \ge 0$, and $\lambda_1(d, q, h)$ be the principal eigenvalue of (3.1). Suppose that $h \gg 0$. If $h_1 = h_2 = \cdots = h_n := h_0$, then $\lambda_1(d, q, h) = h_0$ for all $d > 0, q \ge 0$.

Indeed this lemma can be deduced directly, we omit the proof here.

3.2 Properties of the positive steady state of single species model

Recall that for $d_1 > 0$ and q > 0, (2.1) possesses a unique positive steady state $\boldsymbol{\theta} = (\theta_1, \dots, \theta_n) \gg 0$, which satisfies

$$\sum_{l=1}^{n} (d_1 D_{il} + q Q_{il})\theta_l + r_i \theta_i (1 - \frac{\theta_i}{k_i}) = 0, \quad i \in \mathcal{A},$$

$$(3.7)$$

i.e.

$$\begin{cases} -(d_1+q)\theta_1 + d_1\theta_2 + r_1\theta_1(1-\frac{\theta_1}{k_1}) = 0, \\ (d_1+q)\theta_{j-1} + (-2d_1-q)\theta_j + d_1\theta_{j+1} + r_j\theta_j(1-\frac{\theta_j}{k_j}) = 0, \ j \in \mathcal{B}, \\ (d_1+q)\theta_{n-1} - d_1\theta_n + r_n\theta_n(1-\frac{\theta_n}{k_n}) = 0. \end{cases}$$
(3.8)

Then we collect some useful results for $\boldsymbol{\theta}$.

Lemma 3.3. Suppose
$$d_1 > 0, q \ge 0$$
. Then we have
(i) $\frac{\min_{i \in \mathcal{A}} b_i k_i}{b_i} \le \theta_i \le \frac{\max_{i \in \mathcal{A}} b_i k_i}{b_i}$, where $b_i = (\frac{d}{d+q})^{i-1}$;
(ii) if $k_1 \le k_2 \le \cdots \le k_n$, then $\theta_1 \le \theta_2 \le \cdots \le \theta_n$;

(iii) if $q \ge r_{max} + 2\sqrt{d_1 r_{max}}$, then θ_i is strictly increasing with respect to *i*; (iv) if $q \ge \max\{r_{max} + 2\sqrt{d_1 r_{max}}, 2r_{max}\}$, then for $i \in C$

$$\theta_n \left(\frac{d_1 + q + \frac{2r_{max}}{k_{min}}\theta_n}{d_1}\right)^{i-n} < \theta_i < \theta_n \left(\frac{d_1 + q - 2r_{max}}{d_1}\right)^{i-n}.$$
(3.9)

Moreover,

$$\lim_{q \to +\infty} \theta_n = \lim_{d_1 \to 0} \theta_n = k_n, \tag{3.10}$$

and

$$\lim_{q \to +\infty} \max_{i \in \mathcal{A}} (\theta_i - \theta_n (\frac{d_1}{d_1 + q})^{n-i}) = \lim_{d_1 \to 0} \max_{i \in \mathcal{A}} (\theta_i - \theta_n (\frac{d_1}{d_1 + q})^{n-i}) = 0;$$
(3.11)

$$(v) \lim_{d_1 \to +\infty} \theta_i = \frac{\sum_{i=1}^n r_i}{\sum_{i=1}^n \frac{r_i}{k_i}};$$

(vi) for $d_1 > 0$ small, if q = 0, then $\max_{i \in \mathcal{A}} |\theta_i - k_i| < C_* d_1$ where $C_* = \frac{2 \max_{i \in \mathcal{A}} (\sum_{l=1} D_i k_l)}{r_{min}}$

Proof: (i) Let $b_i = \left(\frac{d_1}{d_1+q}\right)^{i-1}$ and $w_i = b_i \theta_i (i \in \mathcal{A})$. We see that w_i satisfies

$$\begin{cases} (d_1+q)(w_2-w_1)+r_1(1-\frac{w_1}{b_1k_1})w_1=0,\\ d_1w_{j-1}+(-2d_1-q)w_j+(d_1+q)w_{j+1}+r_j(1-\frac{w_j}{b_jk_j})w_j=0, \ j\in\mathcal{B},\\ d_1(w_{n-1}-w_n)+r_n(1-\frac{w_n}{b_nk_n})w_n=0. \end{cases}$$
(3.12)

If $w_1 = \max_{i \in \mathcal{A}} w_i$, then $w_2 - w_1 \leq 0$, which implies from the first equation of (3.12) that $w_i \leq w_1 \leq b_1 k_1$. Similarly, if $w_n = \max_{i \in \mathcal{A}} w_i$, then $w_i \leq w_n \leq b_n k_n$. If $w_{i_0} = \max_i w_i$ for some $i_0 \in \mathcal{B}$, then $w_{i_0-1} - 2w_{i_0} + w_{i_0+1} \leq 0$. The second equation of (3.12) then yields that $w_i \leq w_{i_0} \leq b_{i_0} k_{i_0}$. Therefore, $w_i \leq \max_{i \in \mathcal{A}} b_i k_i$ and $\theta_i \leq \frac{\max_{i \in \mathcal{A}} b_i k_i}{b_i}$. Another inequality $\frac{\min_{i \in \mathcal{A}} b_i k_i}{b_i} \leq \theta_i$ can be obtained similarly. (ii) Set $a_l = \frac{\theta_{l+1} - \theta_l}{\theta_l} (l \in \mathcal{C})$. Then following equations can be deduced directly by (3.8).

$$\begin{cases} d_1 a_1 - q + r_1 (1 - \frac{\theta_1}{k_1}) = 0, \\ d_1 \left[(a_j - a_{j-1})\theta_j + a_{j-1}^2 \theta_{j-1} \right] - q a_{j-1} \theta_{j-1} + r_j \theta_j (1 - \frac{\theta_j}{k_j}) = 0, \ j \in \mathcal{B}, \\ q - d_1 a_{n-1} + r_n \frac{\theta_n}{\theta_{n-1}} (1 - \frac{\theta_n}{k_n}) = 0. \end{cases}$$
(3.13)

Suppose that $a_1 \leq 0$. By the first equation in (3.13), we conclude that $k_1 > \theta_1$. Then $k_2 \geq k_1 > \theta_1 \geq \theta_2$. This implies that $1 - \frac{\theta_2}{k_2} > 0$ and further $(a_2 - a_1)\theta_2 + a_1^2\theta_1 < 0$ by the second equation of (3.13) with j = 2. Thus $a_2 < a_1 \leq 0$, which implies that $\theta_3 < \theta_2 < k_2 \leq k_3$, $1 - \frac{\theta_3}{k_3} > 0$. By the second equation of (3.13) with j = 3, we arrive at $a_3 < a_2$. Thus, by induction, it follows that

$$a_{n-1} < \dots < a_1 \le 0,$$
 (3.14)

which indicates that

$$\theta_n < \theta_{n-1} < \dots < \theta_2 \le \theta_1 < k_1 \le \dots \le k_n.$$
(3.15)

However, the third equation of (3.13) yields that

$$d_1 a_{n-1} = q + r_n \frac{\theta_n}{\theta_{n-1}} (1 - \frac{\theta_n}{k_n}) > 0,$$
(3.16)

a contradiction to (3.14). Therefore, $a_1 > 0$.

Suppose that $a_2 \leq 0$. Using the second equation in (3.8) with j = 2 (which is equivalent to $r_2\theta_2(1 - \frac{\theta_2}{k_2}) + d_1a_2\theta_2 - (d_1 + q)a_1\theta_1 = 0$), we arrive at $k_3 \geq k_2 > \theta_2 \geq \theta_3$. It then follows from the second equation in (3.13) with j = 3 that $a_3 < a_2 \leq 0$. Thus $\theta_4 < \theta_3 \leq \theta_2 < k_2 \leq k_3 \leq k_4$. By induction, there holds

$$a_{n-1} < \dots < a_2 \le 0,$$
 (3.17)

and furthermore $\theta_n < \cdots < \theta_3 \leq \theta_2 < k_n$, which contradicts to the third equation in (3.13). Hence $a_2 > 0$. By the similar methods as above, we reach that $\theta_n > \cdots > \theta_1$.

(iii) By a transformation $v_i = (\frac{d_1}{d_1+q})^{\frac{i-1}{2}}\theta_i$, we can get that

$$\begin{cases} \sqrt{d_1(d_1+q)}(v_2-v_1) + (r_1(1-\frac{\theta_1}{k_1}) - d_1 - q + \sqrt{d_1(d_1+q)})v_1 = 0, \\ \sqrt{d_1(d_1+q)}(v_{j-1} - 2v_j + v_{j+1}) + (r_j(1-\frac{\theta_j}{k_j}) - 2d_1 - q + 2\sqrt{d_1(d_1+q)})v_j = 0, \ j \in \mathcal{B}, \\ (d_1+q)\theta_{n-1} - d_1\theta_n + r_n\theta_n(1-\frac{\theta_n}{k_n}) = 0. \end{cases}$$
(3.18)

Since $q \ge r_{max} + 2\sqrt{d_1 r_{max}}$, we see from the first and second equations of (3.18) that $v_1 < v_2 < \cdots < v_n$. Therefore, $\theta_1 < \theta_2 < \cdots < \theta_n$.

(iv) Set
$$\Theta_i = \theta_n (\frac{d_1 + q - 2r_{max}}{d_1})^{i-n}$$
. By direct calculations, we obtain from $q \ge 2r_{max}$ that

$$\sum_{l=1}^{n} (d_{1}D_{1l} + qQ_{1l})\Theta_{l} + r_{1}\Theta_{1}(1 - \frac{\theta_{1}}{k_{1}}) = -(d_{1} + q)\Theta_{1} + d_{1}\Theta_{2} + r_{1}\Theta_{1}(1 - \frac{\theta_{1}}{k_{1}})$$

$$= \theta_{n}[(-d_{1} - q)(\frac{d_{1} + q - 2r_{max}}{d_{1}})^{1-n} + d_{1}(\frac{d_{1} + q - 2r_{max}}{d_{1}})^{2-n} + r_{1}(\frac{d_{1} + q - 2r_{max}}{d_{1}})^{1-n}(1 - \frac{\theta_{1}}{k_{1}})]$$

$$< \theta_{n}(\frac{d_{1} + q - 2r_{max}}{d_{1}})^{1-n}(-d_{1} - q + d_{1}(\frac{d_{1} + q - 2r_{max}}{d_{1}}) + r_{max}) \leq 0,$$
(3.19)

and

$$\begin{split} \sum_{l=1}^{n} (d_{1}D_{jl} + qQ_{jl})\Theta_{l} + r_{j}\Theta_{j}(1 - \frac{\theta_{j}}{k_{j}}) &= (d_{1} + q)\Theta_{j-1} + (-2d_{1} - q)\Theta_{j} + d_{1}\Theta_{j+1} + r_{j}\Theta_{j}(1 - \frac{\theta_{j}}{k_{j}}) \\ &= \theta_{n}(\frac{d_{1} + q - 2r_{max}}{d_{1}})^{j-n-1}[\frac{q - 2r_{max}}{d_{1}}(d_{1}(\frac{d_{1} + q - 2r_{max}}{d_{1}}) - d_{1} - q) \\ &+ r_{1}(\frac{d_{1} + q - 2r_{max}}{d_{1}}) - r_{1}(\frac{d_{1} + q - 2r_{max}}{d_{1}})\frac{\theta_{j}}{k_{j}}], \\ &< \theta_{n}(\frac{d_{1} + q - 2r_{max}}{d_{1}})^{j-n-1}[d_{1} + q + (r_{max} - 2d_{1} - q)(\frac{d_{1} + q - 2r_{max}}{d_{1}}) \\ &+ d_{1}(\frac{d_{1} + q - 2r_{max}}{d_{1}})^{2}] \leq 0, \ j \in \mathcal{B}. \end{split}$$

$$(3.20)$$

Now, set $u_i = \Theta_i - \theta_i$ satisfying

$$\sum_{l=1}^{n} (d_1 D_{jl} + q Q_{jl}) u_l + r_j u_j (1 - \frac{\theta_j}{k_j}) < 0, \ j \in \mathcal{C}.$$
(3.21)

Clearly, $u_n = 0$. By a further transformation $u_i = z_i \left(\frac{d_1}{d_1 + q}\right)^{\frac{1-i}{2}}$, we get that

$$\begin{cases} \sqrt{d_1(d_1+q)}(z_2-z_1) + (r_1(1-\frac{\theta_1}{k_1}) - d_1 - q + \sqrt{d_1(d_1+q)})z_1 < 0, \\ \sqrt{d_1(d_1+q)}(z_{j-1} - 2z_j + z_{j+1}) + (r_j(1-\frac{\theta_j}{k_j}) - 2d_1 - q + 2\sqrt{d_1(d_1+q)})z_j < 0, \ j \in \mathcal{B}. \end{cases}$$
(3.22)

If there exists $i_0 \in \mathcal{B}$ such that $0 \ge z_{i_0} = \min_{i \in \mathcal{A}} z_i$, then $(z_{i_0-1}-2z_{i_0}+z_{i_0+1}) \ge 0$. Since $q \ge r_{max}+2\sqrt{d_1r_{max}}$ leads to $r_{i_0}(1-\frac{\theta_{i_0}}{k_{i_0}})-2d_1-q+2\sqrt{d_1(d_1+q)} < 0$, then there holds

$$\sqrt{d_1(d_1+q)}(z_{i_0-1}-2z_{i_0}+z_{i_0+1}) + \left(r_{i_0}(1-\frac{\theta_{i_0}}{k_{i_0}})-2d_1-q+2\sqrt{d_1(d_1+q)}\right)z_{i_0} \ge 0,$$
(3.23)

which is a contradiction. Hence, $z_i > \min\{0, z_1\}$ for $i \in C$. If $z_1 < 0$, then $z_2 > z_1$. By $q \ge r_{max} + 2\sqrt{d_1 r_{max}}$, there holds

$$\sqrt{d_1(d_1+q)}(z_2-z_1) + (r_1(1-\frac{\theta_1}{k_1}) - d_1 - q + \sqrt{d_1(d_1+q)})z_1 > 0,$$
(3.24)

which contradicts to (3.22). Hence $z_1 \ge 0$, and therefore $z_i > 0$, that is, $\theta_i < \Theta_i$ for $i \in C$. This proves the second inequality in (3.9).

Next, set
$$\vartheta_i = \theta_n \left(\frac{d_1 + q + \frac{2r_{max}}{k_{min}}\theta_n}{d_1}\right)^{i-n}$$
. Then we have from $q \ge r_{max} + 2\sqrt{d_1 r_{max}}$ that

$$\sum_{l=1}^n (d_1 D_{il} + qQ_{il})\vartheta_i + r_i\vartheta_i(1 - \frac{\theta_i}{k_i}) > \sum_{l=1}^n (d_1 D_{il} + qQ_{il})\vartheta_i - r_{max}\vartheta_i\frac{\theta_n}{k_{min}} > 0.$$
(3.25)

By the same arguments, the first inequality in (3.9) also holds.

A direct summation of the equation of θ_i gives

$$\sum_{i=1}^{n} r_i \theta_i = \sum_{i=1}^{n} r_i \frac{\theta_i^2}{k_i}.$$
(3.26)

It then follows from (3.9) that

$$\theta_n^2 \min_{i \in \mathcal{A}} \frac{r_i}{k_i} \sum_{i=1}^n \left(\frac{d_1 + q + \frac{2r_{max}}{k_{min}}\theta_n}{d_1}\right)^{2i-2n} \le r_{max}\theta_n \sum_{i=1}^n \left(\frac{d_1 + q - 2r_{max}}{d_1}\right)^{i-n},\tag{3.27}$$

which implies that

$$\min_{i \in \mathcal{A}} \frac{r_i}{k_i} \theta_n \frac{1 - (\frac{d_1}{d_1 + q + \frac{2r_{max}}{k_{min}}})^{2n}}{1 - (\frac{d_1}{d_1 + q + \frac{2r_{max}}{k_{min}}})^2} < \frac{r_{max}}{1 - \frac{d_1}{d_1 + q - 2r_{max}}}.$$
(3.28)

Obviously, for sufficiently large q or sufficiently small d_1 , we have that

$$1 - \left(\frac{d_1}{d_1 + q + \frac{2r_{max}}{k_{min}}\theta_n}\right)^{2n} \ge \frac{1}{2}, \ 1 - \left(\frac{d_1}{d_1 + q + \frac{2r_{max}}{k_{min}}\theta_n}\right)^2 > 0, \ \text{and} \ 1 - \frac{d_1}{d_1 + q - 2r_{max}} \ge \frac{1}{2}.$$
 (3.29)

Hence, we can deduce that

$$\min_{i \in \mathcal{A}} \frac{r_i}{k_i} \theta_n < 4r_{max}, \text{ for } q \gg 1 \text{ or } d_1 \ll 1.$$
(3.30)

Next, we show the desired limit. Dividing (3.26) by θ_n , we then observe that

$$\theta_n \sum_{i=1}^n \frac{r_i}{k_i} (\frac{\theta_i}{\theta_n})^2 = \sum_{i=1}^n r_i \frac{\theta_i}{\theta_n}.$$
(3.31)

Since

$$\left(\frac{d_1 + q + \frac{2r_{max}}{k_{min}}\theta_n}{d_1}\right)^{i-n} \le \frac{\theta_i}{\theta_n} \le \left(\frac{d_1 + q - 2r_{max}}{d_1}\right)^{i-n},\tag{3.32}$$

then we see that

$$\lim_{q \to +\infty} \frac{\theta_i}{\theta_n} = \lim_{d_1 \to 0} \frac{\theta_i}{\theta_n} = 0, \ \lim_{q \to +\infty} \frac{\theta_i}{\theta_n} = \lim_{d_1 \to 0} (\frac{\theta_i}{\theta_n})^2 = 0, \text{ for } i \in \mathcal{C},$$
(3.33)

which, together with (3.30), allows us to obtain the limit (3.10).

In view of (3.9), we reach that

$$\theta_n ((\frac{d_1}{d_1 + q + \frac{2r_{max}}{k_{min}}}\theta_n)^{n-i} - (\frac{d_1}{d_1 + q})^{n-i}) \le \theta_i - \theta_n (\frac{d_1}{d_1 + q})^{n-i} \le \theta_n ((\frac{d_1}{d_1 + q - 2r_{max}})^{n-i} - (\frac{d_1}{d_1 + q})^{n-i}).$$

$$(3.34)$$

Therefore, the limit (3.11) is correct.

(v) By statement (i), we have that θ_i is uniformly bounded for $i \in \mathcal{A}$ as $d_1 \to +\infty$. Dividing the system (3.8) by d_1 leads to

$$\begin{cases} \theta_2 - \theta_1 - \frac{q}{d_1} \theta_1 + \frac{r_1}{d_1} \theta_1 (1 - \frac{\theta_1}{k_1}) = 0, \\ (1 + \frac{q}{d_1}) \theta_{j-1} + (-2 - \frac{q}{d_1}) \theta_j + \theta_{j+1} + \frac{r_j}{d_1} \theta_j (1 - \frac{\theta_j}{k_j}) = 0, \ j \in \mathcal{B}, \\ (1 + \frac{q}{d_1}) \theta_{n-1} - \theta_n + \frac{r_n}{d_1} \theta_n (1 - \frac{\theta_n}{k_n}) = 0. \end{cases}$$
(3.35)

Then we have that

$$\theta_1 = \theta_2 = \dots = \theta_n := \rho \ge 0, \tag{3.36}$$

as $d_1 \to +\infty$. By applying the statement (i) again, one can get that $\rho > 0$. Adding (3.35) up, we get

$$\sum_{i=1}^{n} r_i \theta_i (1 - \frac{\theta_i}{k_i}) = 0, \qquad (3.37)$$

which means that $\lim_{d_1 \to +\infty} \theta_i = \rho = \frac{\sum_{i=1}^n r_i}{\sum_{i=1}^n \frac{r_i}{k_i}}$.

(vi) The idea of the proof of this statement is to get a sequence of appropriate upper and lower solutions in terms of d_1 . For each $d_1 > 0$, letting $\bar{\theta}_i = k_i + C_* d_1$, we compute

$$\sum_{l=1}^{n} (d_1 D_{il} \bar{\theta}_l) + \frac{r_i}{k_i} \bar{\theta}_i (k_i - \bar{\theta}_i) = d_1 (\sum_{l=1}^{n} D_{il} k_l - C_* r_i (1 + C_* \frac{d_1}{k_i}))$$

$$\leq d_1 (\sum_{l=1}^{n} D_{il} k_l - C_* r_i) \leq 0,$$
(3.38)

i.e. $\bar{\boldsymbol{\theta}}$ is an upper solution.

Setting $\underline{\theta}_i = k_i - C_* d_1$, we see that for $d_1 > 0$ small, $\underline{\theta}_i > 0$. Then we compute, for $d_1 > 0$ small

$$\sum_{l=1}^{n} (d_1 D_{il} \underline{\theta}_l) + \frac{r_i}{k_i} \underline{\theta}_i (k_i - \underline{\theta}_i) = d_1 (\sum_{l=1}^{n} D_{il} k_l + C_* r_i - C_*^2 \frac{d_1 r_i}{k_i})$$

$$= d_1 (\sum_{l=1}^{n} D_{il} k_l + \frac{1}{2} C_* r_i + C_* r_i (\frac{1}{2} - C_* \frac{d_1}{k_i})) \ge d_1 (\sum_{l=1}^{n} D_{il} k_l + \frac{1}{2} C_* r_i) \ge 0,$$
(3.39)

i.e. $\underline{\theta}$ is a lower solution. Clearly, $\overline{\theta}_i > \underline{\theta}_i$. Therefore, by the similar arguments in the proof of [2, Lemma 2.2](see also [16]), we have that $\overline{\theta}_i > \theta_i > \underline{\theta}_i$, and statement (vi) is correct. This ends the proof.

3.3 Maximum Principle

In this subsection, we shall introduce some important maximum/comparison principles.

Lemma 3.4 (Strong maximum principle I). Let w satisfies

$$\begin{cases} -(d+q)w_1 + (d+q)w_2 + c_1w_1 \ge 0, \\ dw_{j-1} + (-2d-q)w_j + (d+q)w_{j+1} + c_jw_j \ge 0, \ j \in \mathcal{B}, \\ dw_{n-1} - dw_n + c_nw_n \ge 0, \end{cases}$$
(3.40)

where $c_i \in \mathbb{R}$. If $w \leq 0$, then either $w \ll 0$ or w = 0.

Proof: Set $\Sigma = \{i \in \mathcal{A} : w_i = 0\}$. We need to show $\Sigma = \{1, 2, \dots, n\}$ if $\Sigma \neq \emptyset$. If $w_1 = 0$, then we may find from the first equation of (3.40) and $w \leq 0$ that $w_2 = 0$. Furthermore, the second and the third equations of (3.40) imply that $w_3 = \cdots = w_n = 0$. Similarly, we arrive at w = 0 provided that $w_n = 0$. If now $w_1 < 0, w_2 = 0$, by $w_3 \leq 0 = w_2$, we then have from the second equation of (3.40) that

$$0 \le dw_1 + (-2d - q)w_2 + (d + q)w_3 + c_2w_2 = d(w_1 - w_2) + (d + q)(w_3 - w_2) < 0,$$
(3.41)

which is a contradiction. This implies that w_1 can not be negative if $w_2 = 0$. That is, $w_2 = 0$ yields that $w_1 = w_2 = 0$. By the similar method, one can get that w = 0 provided that there exists a $i_0 \in \mathcal{A}$ such that $w_{i_0} = 0$.

Lemma 3.5 (Strong maximum principle II). Let $c_i \in \mathbb{R}$, u satisfy

$$\sum_{j=1}^{n} (dD_{ij} + qQ_{ij})u_j + c_i u_i \le 0, \quad i \in \mathcal{A},$$
(3.42)

where the inequality above is strict for some $j_0 \in A$. Suppose that $\lambda_1(d, q, c) < 0$. Then there holds $u \gg 0$.

Proof: Indeed, after setting $w_i = (\frac{d}{d+q})^{i-1}u_i$, (3.42) is equivalent to

$$\begin{cases} -(d+q)w_1 + (d+q)w_2 + c_1w_1 \le 0, \\ dw_{j-1} + (-2d-q)w_j + (d+q)w_{j+1} + c_jw_j \le 0, \ j \in \mathcal{B}, \\ dw_{n-1} - dw_n + c_nw_n \le 0. \end{cases}$$
(3.43)

We shall verify this lemma in several steps.

Step 1. Let $\phi > 0$ be the non-negative eigenvector corresponding to $\lambda_1(d, q, c)$. Then $\phi \gg 0$. Clearly, it follows from (3.1) that $\bar{\phi}_i := (\frac{d}{d+q})^{i-1} \phi_i$ satisfies

$$\begin{cases} -(d+q)\bar{\phi}_{1} + (d+q)\bar{\phi}_{2} + c_{1}\bar{\phi}_{1} = \lambda_{1}(d,q,c)\bar{\phi}_{1} \leq 0, \\ d\bar{\phi}_{j-1} + (-2d-q)\bar{\phi}_{j} + (d+q)\bar{\phi}_{j+1} + c_{j}\bar{\phi}_{j} = \lambda_{1}(d,q,c)\bar{\phi}_{j} \leq 0, \ j \in \mathcal{B}, \\ d\bar{\phi}_{n-1} - d\bar{\phi}_{n} + c_{n}\bar{\phi}_{n} = \lambda_{1}(d,q,c)\bar{\phi}_{n} \leq 0, \end{cases}$$
(3.44)

and the inequality above is strict for some $j_0 \in \mathcal{A}$. Furthermore, Lemma 3.4 yields that $\bar{\phi} \gg 0$, and hence $\phi \gg 0$.

Step 2. $\boldsymbol{w} \gg \boldsymbol{0}$. Suppose that there exists a $i_0 \in \mathcal{A}$ such that $w_{i_0} = \min_{i \in \mathcal{A}} w_i < 0$. Then we can find some $\xi > 0$ such that $\boldsymbol{w} + \xi \bar{\boldsymbol{\phi}} \ge \boldsymbol{0}$. In fact, we can define $\xi_0 = \inf\{\xi > 0 : w_i + \xi \bar{\boldsymbol{\phi}}_i \ge 0 \ \forall i \in \mathcal{A}\} > 0$. Therefore $\boldsymbol{w} + \xi_0 \bar{\boldsymbol{\phi}} \ge \boldsymbol{0}$. Note that $\boldsymbol{w} + \xi_0 \bar{\boldsymbol{\phi}}$ satisfies

$$\begin{cases} -(d+q)(w_{1}+\xi_{0}\bar{\phi}_{1})+(d+q)(w_{2}+\xi_{0}\bar{\phi}_{2})+c_{1}(w_{1}+\xi_{0}\bar{\phi}_{1})\leq 0,\\ d(w_{j-1}+\xi_{0}\bar{\phi}_{j-1})+(-2d-q)(w_{j}+\xi_{0}\bar{\phi}_{j})+(d+q)(w_{j+1}+\xi_{0}\bar{\phi}_{j+1})+c_{j}(w_{j}+\xi_{0}\bar{\phi}_{j})\leq 0, \ j\in\mathcal{B},\\ d(w_{n-1}+\xi_{0}\bar{\phi}_{n-1})-d(w_{n}+\xi_{0}\bar{\phi}_{n})+c_{n}(w_{n}+\xi_{0}\bar{\phi}_{n})\leq 0, \end{cases}$$

$$(3.45)$$

and the inequality above is also strict for some $j_0 \in \mathcal{A}$. This implies that $\boldsymbol{w} + \xi_0 \bar{\boldsymbol{\phi}} = \boldsymbol{0}$ can not happen. We than conclude from Lemma 3.4 that $\boldsymbol{w} + \xi_0 \bar{\boldsymbol{\phi}} \gg \boldsymbol{0}$, which contradicts the definition of ξ_0 . Therefore, $\boldsymbol{w} \geq \boldsymbol{0}$. Combined with Lemma 3.4 again, we reach that $\boldsymbol{w} \gg \boldsymbol{0}$. That is, $\boldsymbol{u} \gg \boldsymbol{0}$. **Lemma 3.6** (Comparison principle). Let w(t) satisfy

$$\begin{cases} \frac{\mathrm{d}w_1}{\mathrm{d}t} \ge -(d+q)w_1 + (d+q)w_2 + c_1w_1, \ 0 < t < T, \\ \frac{\mathrm{d}w_j}{\mathrm{d}t} \ge dw_{j-1} + (-2d-q)w_j + (d+q)w_{j+1} + c_iw_i, \ j \in \mathcal{B}, \ 0 < t < T, \\ \frac{\mathrm{d}w_n}{\mathrm{d}t} \ge dw_{n-1} - dw_n + c_nw_n, \ 0 < t < T, \\ w(0) > \mathbf{0}, \end{cases}$$
(3.46)

where $c_i \in \mathbb{R}$. Then $\boldsymbol{w}(t) \geq \boldsymbol{0}$.

Proof: Consider the case where $c_i \leq 0$ for every $i \in A$, and the strict inequality

$$\frac{\mathrm{d}w_i}{\mathrm{d}t} > \sum_{j=1}^n (dD_{ij} + qQ_{ij})w_j + c_i w_i \tag{3.47}$$

firstly. If there exists a point $(i_0, t_0) \in \mathcal{A} \times (0, T]$ such that $w_{i_0}(t_0) = \inf_{(i,t) \in \mathcal{A} \times (0,T)} w_i(t) < 0$, (3.47) then yields that

$$0 \ge \frac{\mathrm{d}w_{i_0}}{\mathrm{d}t} > \sum_{j=1}^n (dD_{i_0j} + qQ_{i_0j})w_j + c_{i_0}w_{i_0} \ge 0,$$
(3.48)

which is a contradiction. This contradiction shows that $w(t) \ge 0$.

For the case

$$\frac{\mathrm{d}w_i}{\mathrm{d}t} \ge \sum_{j=1}^n (dD_{ij} + qQ_{ij})w_j + c_i w_i, \tag{3.49}$$

write $w_i^{\varepsilon}(t) = w_i(t) + \varepsilon t$ where $\varepsilon > 0$. Then $\frac{\mathrm{d}w_i^{\varepsilon}}{\mathrm{d}t} > \sum_{j=1}^n (dD_{ij} + qQ_{ij})w_j^{\varepsilon} + c_i w_i^{\varepsilon}$ and so $w_i^{\varepsilon} \ge 0$. Let $\varepsilon \to 0$ to find $w_i \ge 0$. This proves the theorem for the case $c_i \le 0$. For arbitrary c_i we choose $b_i \ge c_i$ and define

to find $w_i \ge 0$. This proves the theorem for the case $c_i \le 0$. For arbitrary c_i , we choose $b_i \ge c_i$ and define $v_i(t) = e^{-b_i t} w_i(t)$. It is easy to see from (3.46) that $v_i(t)$ satisfies

$$\begin{cases} \frac{\mathrm{d}v_1}{\mathrm{d}t} \ge -(d+q)v_1 + (d+q)v_2 + (c_1 - b_1)v_1, \ 0 < t < T, \\ \frac{\mathrm{d}v_j}{\mathrm{d}t} \ge dv_{j-1} + (-2d-q)v_j + (d+q)v_{j+1} + (c_i - b_i)v_i, \ j \in \mathcal{B}, \ 0 < t < T, \\ \frac{\mathrm{d}v_n}{\mathrm{d}t} \ge dv_{n-1} - dv_n + (c_n - b_n)v_n, \ 0 < t < T, \\ \mathbf{v}(0) > \mathbf{0}. \end{cases}$$
(3.50)

Since $c_i - b_i \leq 0$, the above conclusion for $c_i \leq 0$ implies that $v \geq 0$. The conclusion $w(t) \geq 0$ follows from $w_i(t) = e^{b_i t} v_i(t)$.

Lemma 3.7 (Strong maximum principle III). Let w(t) satisfy

$$\begin{cases} \frac{\mathrm{d}w_1}{\mathrm{d}t} \leq -(d+q)w_1 + (d+q)w_2 + c_1w_1, \ 0 < t < T, \\ \frac{\mathrm{d}w_j}{\mathrm{d}t} \leq dw_{j-1} + (-2d-q)w_j + (d+q)w_{j+1} + c_iw_i, \ j \in \mathcal{B}, \ 0 < t < T, \\ \frac{\mathrm{d}w_n}{\mathrm{d}t} \leq dw_{n-1} - dw_n + c_nw_n, \ 0 < t < T, \\ \boldsymbol{w}(0) < \mathbf{0}, \end{cases}$$
(3.51)

where $c_i \in \mathbb{R}$. If $w_{i_0}(t_0) = \max_{i \in \mathcal{A}, t > 0} w_i(t) = M \ge 0$ for some $t_0 > 0, i_0 \in \mathcal{A}$, then $w_i(t) \equiv M$ for all $t < t_0$.

Proof: Without loss of generality, we may assume that $c_i \leq 0$, otherwise one can set $v_i(t) = e^{-b_i t} w_i(t)$ for $b_i \geq c_i$ and consider $v_i(t)$. Suppose by contradiction that there exists $(i_1, t_1) \in \mathcal{A} \times (0, t_0)$ such that $w_{i_1}(t_1) \leq M_1 < M$. Set $u_i(t) = M - w_i(t) - (M - M_1)he^{-\alpha(t-t_1)}$, where h is a small positive constant. We then have that

$$\sum_{j=1}^{n} (dD_{ij} + qQ_{ij})u_j + c_i u_i - \frac{\mathrm{d}u_i}{\mathrm{d}t} \le -e^{\alpha(t-t_1)}(M - M_1)(\alpha h + c_i h + \sum_{j=1}^{n} (dD_{ij} + qQ_{ij})h) < 0, \quad (3.52)$$

provided $\alpha \gg 1$. Since $u_i(t_1) = M - w_i(t_1) - (M - M_1)h \ge (1 - h)(M - M_i) > 0$ for $0 < h \ll 1$. It then follows from Lemma 3.6 that $u(t) \ge 0$. Furthermore, $w_i(t) \le M - (M - M_1)he^{-\alpha(t-t_1)} < M$ for $t > t_1$. This implies that $w_{i_0}(t_0) < M$, which is a contradiction. The proof of this lemma is hence finished. \Box

4 Proof of the main results

This section is devoted to the proofs of our main results.

4.1 Dynamic classification in the $q - \gamma$ plane

Clearly, the steady states of (1.6) consists of the trivial steady state (0, 0), the semi-trivial steady state $(\theta, 0)$ and the positive steady state (N^*, P^*) .

Lemma 4.1. Suppose that (\mathbf{H}) hold. The trivial steady state $(\mathbf{0}, \mathbf{0})$ of (1.6) is always unstable.

Proof: The linearization eigenvalue problem of(1.6) at (0,0) is

$$\begin{cases} \sum_{j=1}^{n} (d_1 D_{ij} + q Q_{ij}) \phi_j + r_i \phi_i = \lambda \phi_i, \ i \in \mathcal{A}, \\ \sum_{j=1}^{n} (d_2 D_{ij} + \tau q Q_{ij}) \psi_j - \gamma \psi_i = \lambda \psi_i \ i \in \mathcal{A}. \end{cases}$$

$$(4.1)$$

Note by $r \gg 0$ that $\lambda_1(d_1, q, r) > 0$. We conclude that (0, 0) is always unstable.

Lemma 4.2. Suppose that (**H**) hold. There exists a unique continuous critical curve $\Gamma : \gamma = \gamma^*(q, d_1, d_2)$ such that $(\theta, \mathbf{0})$ is locally asymptotically stable if $\gamma > \gamma^*(q, d_1, d_2)$ and unstable if $0 < \gamma < \gamma^*(q, d_1, d_2)$.

Proof: The linearization eigenvalue problem of (1.6) at $(\theta, 0)$ is

$$\begin{cases} \sum_{j=1}^{n} (d_1 D_{ij} + q Q_{ij}) \phi_j + r_i (1 - \frac{2\theta_i}{k_i}) \phi_i - a \theta_i \psi_i = \mu \phi_i, \ i \in \mathcal{A}, \\ \sum_{j=1}^{n} (d_2 D_{ij} + \tau q Q_{ij}) \psi_j + (\beta a \theta_i - \gamma) \psi_i = \mu \psi_i \ i \in \mathcal{A}. \end{cases}$$

$$(4.2)$$

Clearly, the local stability of $(\boldsymbol{\theta}, \mathbf{0})$ is determined by the sign of the principal eigenvalue of $L\psi_i = \sum_{j=1}^{n} (d_2 D_{ij} + \tau q Q_{ij})\psi_j + (\beta a \theta_i - \gamma)\psi_i$; namely, $(\boldsymbol{\theta}, \mathbf{0})$ is locally asymptotically stable if $\lambda_1(d_2, \tau q, \beta a \boldsymbol{\theta} - \gamma) < 0$ and unstable if $\lambda_1(d_2, \tau q, \beta a \boldsymbol{\theta} - \gamma) > 0$. Furthermore, we have

$$\lambda_1(d_2, \tau q, \beta a \boldsymbol{\theta} - \gamma) = \lambda_1(d_2, \tau q, \beta a \boldsymbol{\theta}) - \gamma.$$
(4.3)

Therefore, we let $\gamma^*(q, d_1, d_2) = \lambda_1(d_2, \tau q, \beta a \theta)$. Then it follows from Lemma 3.1(iv) that

$$\begin{cases} \lambda_1(d_2, \tau q, \beta a \boldsymbol{\theta} - \gamma) > 0 \text{ if } 0 < \gamma < \gamma^* \\ \lambda_1(d_2, \tau q, \beta a \boldsymbol{\theta} - \gamma) = 0 \text{ if } \gamma = \gamma^* \\ \lambda_1(d_2, \tau q, \beta a \boldsymbol{\theta} - \gamma) < 0 \text{ if } \gamma > \gamma^*. \end{cases}$$

$$(4.4)$$

Proof of Theorem 2.1: We shall split the proof into several steps. Step 1. Let $U_i(t) = (\frac{d_1}{d_1+q})^{i-1}N_i(t)$ and $V_i(t) = (\frac{d_2}{d_2+\tau q})^{i-1}P_i(t)$, where $(\mathbf{N}(t), \mathbf{P}(t))$ is the solution of (1.6). Then $(\mathbf{U}(t), \mathbf{V}(t))$ satisfies

$$\frac{dU_{1}}{dt} = -(d_{1}+q)U_{1} + (d_{1}+q)U_{2} + r_{1}U_{1}(1-\frac{N_{1}}{k_{1}}) - aU_{1}P_{1},
\frac{dU_{j}}{dt} = d_{1}U_{j-1} + (-2d_{1}-q)U_{j} + (d_{1}+q)U_{j+1} + r_{j}U_{j}(1-\frac{N_{j}}{k_{j}}) - aU_{j}P_{j}, \ j \in \mathcal{B},
\frac{dU_{n}}{dt} = d_{1}U_{n-1} - d_{1}U_{n} + r_{n}U_{n}(1-\frac{N_{n}}{k_{n}}) - aU_{n}P_{n},
\frac{dV_{1}}{dt} = -(d_{2}+\tau q)V_{1} + (d_{2}+\tau q)V_{2} + V_{1}(\beta aN_{1}-\gamma),
\frac{dV_{j}}{dt} = d_{2}V_{j-1} + (-2d_{2}-\tau q)V_{j} + (d_{2}+\tau q)V_{j+1} + V_{j}(\beta aN_{j}-\gamma), \ j \in \mathcal{B},
\frac{dV_{n}}{dt} = d_{2}V_{n-1} - d_{2}V_{n} + V_{n}(\beta aN_{n}-\gamma),
U(0) = (1, \frac{d_{1}}{d_{1}+q}, \cdots, (\frac{d_{1}}{d_{1}+q})^{n-1}) \cdot \mathbf{N}(0) > \mathbf{0},
V(0) = (1, \frac{d_{2}}{d_{2}+\tau q}, \cdots, (\frac{d_{2}}{d_{2}+\tau q})^{n-1}) \cdot \mathbf{P}(0) > \mathbf{0}.$$
(4.5)

Clearly, Lemma 3.7 leads to $U, V \gg 0$ since U(0), V(0) > 0, and then $N, P \gg 0$. Note that

$$\begin{cases} \frac{dU_1}{dt} \le -(d_1+q)U_1 + (d_1+q)U_2 + r_1U_1(1-\frac{U_1}{k_1}), \\ \frac{dU_j}{dt} \le d_1U_{j-1} + (-2d_1-q)U_j + (d_1+q)U_{j+1} + r_jU_j(1-\frac{U_j}{k_j}(\frac{d_1+q}{d_1})^{j-1}), \ j \in \mathcal{B}, \\ \frac{dU_n}{dt} \le d_1U_{n-1} - d_1U_n + r_nU_n(1-\frac{U_n}{k_n}(\frac{d_1+q}{d_1})^{n-1}). \end{cases}$$

$$(4.6)$$

Let $C_1 = \max_{i \in \mathcal{A}} \{k_i, U_i(0)\}$, and $\mathcal{U}_i = U_i - C_1$. It then follows that \mathcal{U}_i satisfies $\mathcal{U}_i(0) \leq 0$ and

$$\begin{cases} \frac{\mathrm{d}\mathcal{U}_{1}}{\mathrm{d}t} \leq -(d_{1}+q)\mathcal{U}_{1} + (d_{1}+q)\mathcal{U}_{2} + r_{1}C_{1}(-\frac{\mathcal{U}_{1}}{k_{1}}), \\ \frac{\mathrm{d}\mathcal{U}_{j}}{\mathrm{d}t} \leq d_{1}\mathcal{U}_{j-1} + (-2d_{1}-q)\mathcal{U}_{j} + (d_{1}+q)\mathcal{U}_{j+1} + r_{j}C_{1}(-\frac{\mathcal{U}_{j}}{k_{j}}(\frac{d_{1}+q}{d_{1}})^{j-1}), \ j \in \mathcal{B}, \\ \frac{\mathrm{d}\mathcal{U}_{n}}{\mathrm{d}t} \leq d_{1}\mathcal{U}_{n-1} - d_{1}\mathcal{U}_{n} + r_{n}C_{1}(-\frac{\mathcal{U}_{n}}{k_{n}}(\frac{d_{1}+q}{d_{1}})^{n-1}). \end{cases}$$
(4.7)

From the above inequalities, one can get that $\mathcal{U}_i \leq 0$ for $i \in \mathcal{A}$ and $t \geq 0$, and hence $U_i \leq C_1$, $0 < N_i \leq C_1 (\frac{d_1+q}{d_1})^{n-1}$ for t > 0.

$$\begin{aligned} Step \ 2. \text{ Next, we verify } V_i \text{ and } P_i \text{ are bounded uniformly for } i \in \mathcal{A}, t > 0. \text{ We rewrite system } (4.5) \text{ as} \\ \begin{cases} \frac{dU_1}{dt} = -(d_1 + q)U_1 + (d_1 + q)U_2 + r_1U_1(1 - \frac{N_1}{k_1}) - aU_1P_1, \\ (\frac{d_1 + q}{d_1})^{j-1}\frac{dU_j}{dt} = (d_1U_{j-1} + (-2d_1 - q)U_j + (d_1 + q)U_{j+1} + r_jU_j(1 - \frac{N_j}{k_j}) - aU_jP_j)(\frac{d_1 + q}{d_1})^{j-1}, \ j \in \mathcal{B} \\ (\frac{d_1 + q}{d_1})^{n-1}\frac{dU_n}{dt} = (d_1U_{n-1} - d_1U_n + r_nU_n(1 - \frac{U_n}{k_n}) - aU_nP_n)(\frac{d_1 + q}{d_1})^{n-1}, \\ \frac{dV_1}{dt} = -(d_2 + \tau q)V_1 + (d_2 + \tau q)V_2 + V_1(\beta aN_1 - \gamma), \\ (\frac{d_2 + \tau q}{d_2})^{j-1}\frac{dV_j}{dt} = (d_2V_{j-1} + (-2d_2 - \tau q)V_j + (d_2 + \tau q)V_{j+1} + V_j(\beta aN_j - \gamma))(\frac{d_2 + \tau q}{d_2})^{j-1}, \ j \in \mathcal{B}, \\ (\frac{d_2 + \tau q}{d_2})^{n-1}\frac{dV_n}{dt} = (d_2V_{n-1} - d_2V_n + V_n(\beta aN_n - \gamma))(\frac{d_2 + \tau q}{d_2})^{n-1}, \\ U(0) > \mathbf{0}, \\ V(0) > \mathbf{0}. \end{aligned}$$

Let $w(t) = \sum_{i=1}^{n} (\beta (\frac{d_1+q}{d_1})^{i-1} U_i + (\frac{d_2+\tau q}{d_2})^{i-1} V_i)$. Adding all the equations in (4.8) up yields that $w'(t) + \gamma w(x) \le \beta (r_{max}+\gamma) \sum_{i=1}^{n} (\frac{d_1+q}{d_1})^{i-1} U_i \le \beta (r_{max}+\gamma) C_1 n (\frac{d_1+q}{d_1})^{n-1}$. (4.9)

By Gronwall inequality, we have

$$w(t) \le w(0)e^{-\gamma t} + \frac{\beta(r_{max} + \gamma)C_1 n(\frac{d_1 + q}{d_1})^{n-1}}{\gamma} (1 - e^{-\gamma t}), \tag{4.10}$$

which implies that w(t) is bounded for all t > 0. Then one can get that $0 < V_i \le (\frac{d_2 + \tau q}{d_2})^{i-1}V_i < w(t)$ is also bounded uniformly. Hence $P_i(t)$ is bounded uniformly and the statement (i) holds.

Step 3. Consider the case $\gamma > \gamma^*$.

Step 3.1. Let f(t) be the solution of

$$\begin{cases} \frac{\mathrm{d}f_1}{\mathrm{d}t} = -(d_1+q)f_1 + (d_1+q)f_2 + r_1f_1(1-\frac{f_1}{k_1}), \\ \frac{\mathrm{d}f_j}{\mathrm{d}t} = d_1f_{j-1} + (-2d_1-q)f_j + (d_1+q)f_{j+1} + r_jf_j(1-\frac{f_j}{k_j}(\frac{d_1+q}{d_1})^{j-1}), \ j \in \mathcal{B}, \\ \frac{\mathrm{d}f_n}{\mathrm{d}t} = d_1f_{n-1} - d_1f_n + r_nf_n(1-\frac{f_n}{k_n}(\frac{d_1+q}{d_1})^{n-1}), \\ f(0) = (1, \frac{d_1}{d_1+q}, \cdots, (\frac{d_1}{d_1+q})^{n-1}) \cdot \mathbf{N}(0) > \mathbf{0}. \end{cases}$$

$$(4.11)$$

The comparison principle (Lemma 3.6) and (4.6) yield that $U_i \leq f_i$. Observe from [2, Lemma 2.2] that $\lim_{t \to +\infty} f_i = (\frac{d_1}{d_1 + q})^{i-1} \theta_i$. This implies that

$$\limsup_{t \to +\infty} U_i \le \left(\frac{d_1}{d_1 + q}\right)^{i-1} \theta_i \text{ for all } i \in \mathcal{A}.$$
(4.12)

Then for any $\varepsilon > 0$, there exists $T_1 > 0$ such that $N_i(t) < \theta_i + \varepsilon$ for all $t \ge T_1$. Let g(t) be the solution of

$$\begin{cases} \frac{\mathrm{d}g_1}{\mathrm{d}t} = -(d_2 + \tau q)g_1 + (d_2 + \tau q)g_2 + g_1(\beta a(\theta_1 + \varepsilon) - \gamma), \\ \frac{\mathrm{d}g_j}{\mathrm{d}t} = d_2g_{j-1} + (-2d_2 - \tau q)g_j + (d_2 + \tau q)g_{j+1} + g_j(\beta a(\theta_j + \varepsilon) - \gamma), \ j \in \mathcal{B}, \\ \frac{\mathrm{d}g_n}{\mathrm{d}t} = d_2g_{n-1} - d_2g_n + g_n(\beta a(\theta_n + \varepsilon) - \gamma), \\ g(T_1) = \mathbf{V}(T_1). \end{cases}$$
(4.13)

Clearly, one can get that $V_i(t) \leq g_i(t)$ for all $t \geq T_1$ by using Lemma 3.6. Since $\lambda_1(d_2, \tau q, \beta a \theta - \gamma) < 0$ when $\gamma > \gamma^*$, there exists $\varepsilon > 0$ small enough such that $\lambda_1(d_2, \tau q, \beta a(\theta + \varepsilon) - \gamma) < 0$. Since it is easy to see that $C(\frac{d_2}{d_2 + \tau q})^{i-1} e^{\lambda_1(d_2, \tau q, \beta a(\theta + \varepsilon) - \gamma)(t - T_1)} \phi$ is a super solution of (4.13) for *C* large enough, where ϕ is the principal eigenfunction of the eigenvalue problem (3.2) with respect to $\lambda_1(d_2, \tau q, \beta a(\theta + \varepsilon) - \gamma)$. Lemma 3.6 then yields that

$$0 \le \limsup_{t \to +\infty} V_i(t) \le \lim_{t \to +\infty} C\phi_i e^{\lambda_1(d_2, \tau q, \beta a(\boldsymbol{\theta} + \boldsymbol{\varepsilon}) - \gamma)(t - T_1)} = 0.$$
(4.14)

Step 3.2. Since $\lim_{t \to +\infty} V_i(t) = 0$, therefore, $\lim_{t \to +\infty} P_i(t) = 0$ and for any $\varepsilon > 0$, there exists $T_2 > T_1$ such that $P_i(t) \le \varepsilon$ for $t \ge T_2$, which leads to

$$\begin{cases} \frac{\mathrm{d}U_1}{\mathrm{d}t} \ge -(d_1+q)U_1 + (d_1+q)U_2 + r_1U_1(1-\frac{N_1}{k_1}) - aU_1\varepsilon, \\ \frac{\mathrm{d}U_j}{\mathrm{d}t} \ge d_1U_{j-1} + (-2d_1-q)U_j + (d_1+q)U_{j+1} + r_jU_j(1-\frac{N_j}{k_j}) - aU_j\varepsilon, \ j \in \mathcal{B}, \\ \frac{\mathrm{d}U_n}{\mathrm{d}t} \ge d_1U_{n-1} - d_1U_n + r_nU_n(1-\frac{N_n}{k_n}) - aU_n\varepsilon. \end{cases}$$

$$(4.15)$$

Let $f^{\varepsilon}(t)$ be the solution of

$$\begin{cases} \frac{\mathrm{d}f_{1}^{\varepsilon}}{\mathrm{d}t} = -(d_{1}+q)f_{1}^{\varepsilon} + (d_{1}+q)f_{2}^{\varepsilon} + r_{1}f_{1}^{\varepsilon}(1-\frac{f_{1}^{\varepsilon}}{k_{1}}) - af_{1}^{\varepsilon}\varepsilon, \\ \frac{\mathrm{d}f_{j}^{\varepsilon}}{\mathrm{d}t} = d_{1}f_{j-1}^{\varepsilon} + (-2d_{1}-q)f_{j}^{\varepsilon} + (d_{1}+q)f_{j+1}^{\varepsilon} + r_{j}f_{j}^{\varepsilon}(1-\frac{N_{j}}{k_{j}}\frac{d_{1}+q}{d_{1}})^{j-1}) - af_{j}^{\varepsilon}\varepsilon, \ j \in \mathcal{B}, \\ \frac{\mathrm{d}f_{n}^{\varepsilon}}{\mathrm{d}t} = d_{1}f_{n-1}^{\varepsilon} - d_{1}f_{n}^{\varepsilon} + r_{n}f_{n}^{\varepsilon}(1-\frac{N_{n}}{k_{n}}\frac{d_{1}+q}{d_{1}})^{n-1}) - af_{n}^{\varepsilon}\varepsilon, \\ f(T_{2}) = U(T_{2}). \end{cases}$$

$$(4.16)$$

Lemma 3.6 implies that $U_i(t) \ge f_i^{\varepsilon}(t)$ for all $t \ge T_2$. By applying [2, Lemma 2.2] and choosing $\varepsilon > 0$ sufficiently small, we have that $\lim_{t\to+\infty} f_i^{\varepsilon}(t) = \theta_i^{\varepsilon} (\frac{d_1}{d_1+q})^{i-1}$ uniformly for $i \in \mathcal{A}$, where $\theta_i^{\varepsilon} (\frac{d_1}{d_1+q})^{i-1}$ is the unique positive steady state solution of (4.16). Furthermore, by virtue of Lemma 3.3(i) and (4.16), we can obtain that $0 \le \theta_i^{\varepsilon} \le (\frac{d_1+q}{d_1})^{n-1} \max_{i\in\mathcal{A}} k_i (\frac{d_1}{d_1+q})^{i-1}$. Note that we can deduce that $\lim_{\varepsilon\to 0} \theta_i^{\varepsilon} = \theta_i$. That is

$$\liminf_{t \to +\infty} U_i(t) \ge \theta_i (\frac{d_1}{d_1 + q})^{i-1}.$$
(4.17)

Combining with (4.12), we get that

$$\lim_{t \to +\infty} U_i(t) = \theta_i (\frac{d_1}{d_1 + q})^{i-1}, \quad \lim_{t \to +\infty} N_i(t) = \theta_i.$$
(4.18)

Hence the statement (ii) holds.

Step 4. Consider the case $\gamma < \gamma^*$.

Step 4.1. System (1.6) is uniformly persistent.

In order to prove the uniform persistence of system (1.6)(or the equivalent system (4.5)) in the case where $\gamma < \gamma^*$, let $\Theta(t)$ be the solution semiflow generated by system (4.5) on the state space \mathbb{P} , where

$$\mathbb{P} = \{ (\boldsymbol{U}, \boldsymbol{V}) \in \mathbb{R}^n \times \mathbb{R}^n : \boldsymbol{U} > \boldsymbol{0}, \boldsymbol{V} > \boldsymbol{0} \}.$$
(4.19)

Define

$$\mathbb{P}_0 = \{ (\boldsymbol{U}, \boldsymbol{V}) \in \mathbb{P} : \boldsymbol{U} \neq \boldsymbol{0} \text{ and } \boldsymbol{V} \neq \boldsymbol{0} \},$$
(4.20)

and $\partial \mathbb{P}_0 = \mathbb{P} \setminus \mathbb{P}_0$. Let

$$M_{\partial} = \{ (\boldsymbol{U}(0), \boldsymbol{V}(0)) \in \partial \mathbb{P}_0 : \Theta(t)(\boldsymbol{U}(0), \boldsymbol{V}(0)) \in \partial \mathbb{P}_0, \forall t \ge 0 \},$$

$$(4.21)$$

and $\omega((\boldsymbol{U}(0), \boldsymbol{V}(0)))$ be the omega limit set of the forward orbit $\delta^+((\boldsymbol{U}(0), \boldsymbol{V}(0))) = \{\Theta(t)(\boldsymbol{U}(0), \boldsymbol{V}(0)):$ $t \geq 0$ }. By Lemma 3.7, we can conclude that \mathbb{P}_0 is open in \mathbb{P} and forward invariant under the dynamics generated by system (4.5), and $\partial \mathbb{P}_0$ contains steady state points (0,0), $(\theta,0)$. We claim that

$$\bigcup_{(\boldsymbol{U}(0),\boldsymbol{V}(0))\in M_{\partial}} \omega((\boldsymbol{U}(0),\boldsymbol{V}(0))) \in \{(\boldsymbol{0},\boldsymbol{0}), (\boldsymbol{\theta},\boldsymbol{0})\}.$$
(4.22)

In fact, for any $(U(0), V(0)) \in M_{\partial}$, we have $\Theta(t)(U(0), V(0)) \in \partial \mathbb{P}_0, \forall t \geq 0$. This implies that U(t; (U(0), V(0))) = 0 or V(t; (U(0), V(0))) = 0 for $t \ge 0$. If U(t; (U(0), V(0))) = 0 for $t \ge 0$, then $\boldsymbol{P}(t; (\boldsymbol{N}(0), \boldsymbol{P}(0)))$ satisfies

$$\begin{cases} \frac{\mathrm{d}P_1}{\mathrm{d}t} = -(d_2 + \tau q)P_1 + d_2P_2 - \gamma P_1, \\ \frac{\mathrm{d}P_j}{\mathrm{d}t} = (d_2 + \tau q)P_{j-1} + (-2d_2 - \tau q)P_j + d_2P_{j+1} - \gamma P_j, \ j \in \mathcal{B}, \\ \frac{\mathrm{d}P_n}{\mathrm{d}t} = (d_2 + \tau q)P_{n-1} - d_2P_n - \gamma P_n, \\ P(0) > \mathbf{0}. \end{cases}$$

$$(4.23)$$

It follows that $\frac{\mathrm{d}(e^{\gamma t} \sum_{i=1}^{n} P_i)}{\mathrm{d}t} = 0$, which hence implies that $0 \leq \lim_{t \to +\infty} P_i = 0$. In the case where $U(\tau_0; (U(0), V(0))) \neq \mathbf{0}$ for some $\tau_0 > 0$, we have $U(t; (U(0), V(0))) > \mathbf{0}$ for all $t > \tau_0$ by Lemma 3.7, which implies that V(t; (U(0), V(0))) = 0 for all $t > \tau_0$. Thus N(t; (N(0), P(0))) is the solution of (2.1). Then we have that either $\lim_{t \to +\infty} N_i(t) = 0$, or $\lim_{t \to +\infty} N_i(t) = \theta_i$. Hence, our claim holds. We next claim that $(\mathbf{0}, \mathbf{0}), (\boldsymbol{\theta}, \mathbf{0})$ are uniform weak repellers in the sense that

$$\limsup_{t \to +\infty} \|\Theta(t)(\boldsymbol{U}(0), \boldsymbol{V}(0)) - (\boldsymbol{0}, \boldsymbol{0})\| \ge \delta_1 \text{ for all } (\boldsymbol{U}(0), \boldsymbol{V}(0)) \in \mathbb{P}_0,$$
(4.24)

and

$$\limsup_{t \to +\infty} \|\Theta(t)(\boldsymbol{U}(0), \boldsymbol{V}(0)) - (\boldsymbol{\theta}, \boldsymbol{0})\| \ge \delta_2 \text{ for all } (\boldsymbol{U}(0), \boldsymbol{V}(0)) \in \mathbb{P}_0.$$
(4.25)

Suppose by contradiction that (4.24) is not true. Then for any $\delta > 0$, there exists $(U(0), V(0)) \in \mathbb{P}_0$ such that

$$\limsup_{t \to +\infty} \|\Theta(t)(\boldsymbol{U}(0), \boldsymbol{V}(0)) - (\boldsymbol{0}, \boldsymbol{0})\| < \delta.$$
(4.26)

Therefore, there exists $t_0 > 0$ such that for $t \ge t_0$, we have that

$$\|N(t, (N(0), P(0)))\| < \delta, \|P(t, (N(0), P(0)))\| < \delta.$$
(4.27)

Consequently, it follows from the equation for N that

$$\frac{\mathrm{d}N_i}{\mathrm{d}t} \ge \sum_{j=1}^n (d_1 D_{ij} + q Q_{ij}) N_j + N_i (r_i - \frac{r_i \delta}{k_i} - a\delta), \quad i \in \mathcal{A}, \ t \ge t_0.$$

$$(4.28)$$

In view of $(U(0), V(0)) \in \mathbb{P}_0$, by Lemma 3.7, we have $U_i(t_0) > 0$, and then $N_i(t_0) > 0$. Thus there exists an $\alpha_0 > 0$ such that $N_i(t_0) \ge \alpha_0 \phi_{1i}^{\delta}$, where ϕ_1^{δ} is the principal eigenfunction corresponding to the principal eigenvalue $\lambda_1(d_1, q, \mathbf{r} - \frac{\mathbf{r}}{\mathbf{k}}\delta - \mathbf{a}\delta)$ of (3.2). Let $\underline{N}_i(t) = \alpha_0 e^{\lambda_1(d_1, q, \mathbf{r} - \frac{\mathbf{r}}{\mathbf{k}}\delta - \mathbf{a}\delta)(t-t_0)}\phi_{1i}^{\delta}$. Then $\underline{N}_i(t)$ satisfies

$$\begin{cases} \frac{\mathrm{d}N_i}{\mathrm{d}t} = \sum_{j=1}^n (d_1 D_{ij} + q Q_{ij}) \underline{N}_j + \underline{N}_i (r_i - \frac{r_i \delta}{k_i} - a\delta), & i \in \mathcal{A}, \ t > 0, \\ \underline{N}(t_0) = \alpha_0 \boldsymbol{\phi}_1^{\boldsymbol{\delta}}. \end{cases}$$
(4.29)

The comparison principle (Lemma 3.6) yields that

$$N_i(t) \ge \underline{N}(t) \text{ for all } t \ge t_0. \tag{4.30}$$

Since $\lambda_1(d_1, q, \mathbf{r}) > 0$, by continuity, we can choose $\delta > 0$ small enough such that $\lambda_1(d_1, q, \mathbf{r} - \frac{\mathbf{r}}{\mathbf{k}}\delta - \mathbf{a}\delta) > 0$, which implies $\lim_{t \to +\infty} N_i(t, (\mathbf{N}(0), \mathbf{P}(0))) = +\infty$. This is a contradiction to $\|\mathbf{N}(t, (\mathbf{N}(0), \mathbf{P}(0)))\| < \delta$ for $t > t_0$. Hence, we conclude that $(\mathbf{0}, \mathbf{0})$ is a uniform weak repeller.

Suppose by contradiction that (4.25) is not true. Then for any $\delta > 0$, there exists $(\mathbf{N}(0), \mathbf{P}(0)) \in \mathbb{P}_0$ such that $\limsup_{t \to +\infty} \|\Theta(t)(\mathbf{N}(0), \mathbf{P}(0)) - (\boldsymbol{\theta}, \mathbf{0})\| < \delta$. Therefore, there exists $t_1 > 0$ such that for $t \ge t_1$, we have

$$\|N(t, (N(0), P(0))) - \theta\| < \delta, \|P(t, (N(0), P(0)))\| < \delta.$$
(4.31)

Therefore, it follows from the equation for \boldsymbol{P} that

$$\frac{\mathrm{d}P_i}{\mathrm{d}t} \ge \sum_{j=1}^n (d_2 D_{ij} + \tau q Q_{ij}) P_j + P_i (\beta a(\theta_i - \delta) - \gamma), \quad i \in \mathcal{A}, \ t \ge t_1.$$

$$(4.32)$$

In view of $(\mathbf{N}(0), \mathbf{P}(0)) \in \mathbb{P}_0$, by Lemma 3.7, we have $P_i(t_1) > 0$. Let ψ_1^{δ} be the principal eigenfunction with respect to $\lambda_1(d_2, \tau q, \beta a(\boldsymbol{\theta} - \boldsymbol{\delta}) - \boldsymbol{\gamma})$. Thus there exists an $\alpha_1 > 0$ such that $P_i(t_1) \geq \alpha_1 \psi_{1i}^{\delta}$. Let $\underline{P}_i(t) = \alpha_1 e^{\lambda_1(d_2, \tau q, \beta a(\boldsymbol{\theta} - \boldsymbol{\delta}) - \boldsymbol{\gamma})(t-t_1)} \psi_{1i}^{\delta}$. Then $\underline{P}_i(t)$ satisfies

$$\begin{cases} \frac{\mathrm{d}\underline{P}_{i}}{\mathrm{d}t} = \sum_{j=1}^{n} (d_{2}D_{ij} + \tau qQ_{ij})\underline{P}_{j} + \underline{P}_{i}(\beta a(\theta_{i} - \delta) - \gamma), & i \in \mathcal{A}, t \geq t_{1}, \\ \underline{P}_{i}(t_{1}) = \alpha_{1}\psi_{1i}^{\delta}. \end{cases}$$
(4.33)

It follows from Lemma 3.6 that $P_i(t) \geq \underline{P}_i(t)$ for all $t \geq t_1$. On the other hand, since $0 < \gamma < \gamma^*$, we have that $\lambda_1(d_2, \tau q, \beta a \theta - \gamma) > 0$ by (4.4). By continuity, we can choose $\delta > 0$ small enough such that $\lambda_1(d_2, \tau q, \beta a(\theta - \delta) - \gamma) > 0$, which implies $\lim_{t \to +\infty} P_i(t, (\mathbf{N}(0), \mathbf{P}(0))) = +\infty$. This is a contradiction to $\|\mathbf{P}(t, (\mathbf{N}(0), \mathbf{P}(0)))\| < \delta$ for $t > t_1$. Hence, we conclude that $(\theta, \mathbf{0})$ is a uniform weak repeller.

 $P(t, (N(0), P(0))) \| < \delta$ for $t > t_1$. Hence, we conclude that $(\delta, 0)$ is a uniform weak result.

We define now a continuous function $\mathcal{D}: \mathbb{P} \to [0,\infty)$ by

$$\mathcal{D}((\boldsymbol{N},\boldsymbol{P})) = \min\{\min_{i\in\mathcal{A}}N_i, \min_{i\in\mathcal{A}}P_i\} \text{ for any } (\boldsymbol{N},\boldsymbol{P})\in\mathbb{P}.$$
(4.34)

It follows from Lemma 3.7 that $\mathcal{D}^{-1}(0,\infty) \subset \mathbb{P}_0$ and D satisfies if $\mathcal{D}((N, \mathbf{P})) > 0$ or $(N, \mathbf{P}) \in \mathbb{P}_0$ with $\mathcal{D}((N, \mathbf{P})) = 0$, then $D(\Theta(t)(N, \mathbf{P})) > 0$, $\forall t > 0$. That is, \mathcal{D} is a generalized distance function for the semiflow $\Theta(t) : \mathbb{P} \to \mathbb{P}$ (see [28]). It follows from Step 2 that $\Theta(t)$ is point dissipative on \mathbb{P} . Moreover, it is easy to see from the Arzela-Ascoli theorem that $\Theta(t) : \mathbb{P} \to \mathbb{P}$ is compact for any t > 0. By [22, Theorem 2.6], $\Theta(t) : \mathbb{P} \to \mathbb{P}$, $t \ge 0$ admits a global compact attractor that attracts each bounded set in \mathbb{P} . Since $(\mathbf{0}, \mathbf{0})$ and $(\mathbf{\theta}, 0)$ are uniform weak repellers, we conclude that $(\mathbf{0}, \mathbf{0})$ and $(\mathbf{\theta}, 0)$ are isolated invariant sets in \mathbb{P} , and

$$W^{S}\{(\mathbf{0},\mathbf{0})\} \cap \mathcal{D}^{-1}(0,\infty) = \emptyset, W^{S}\{(\boldsymbol{\theta},0)\} \cap \mathcal{D}^{-1}(0,\infty) = \emptyset,$$
(4.35)

where $W^S\{(\mathbf{0},\mathbf{0})\}, W^S\{(\boldsymbol{\theta},0)\}$ are the stable sets of $(\mathbf{0},\mathbf{0}), (\boldsymbol{\theta},0)$, respectively (see [28]). Hence, there are no subsets of $\{(\mathbf{0},\mathbf{0})\} \cup \{(\boldsymbol{\theta},0)\}$ form a cycle in $\partial \mathbb{P}_0$. By [28, Theorem 3], there exists $\eta > 0$ such that for any $(N(0), P(0)) \in \mathbb{P}_0$,

$$\min_{(\boldsymbol{N},\boldsymbol{P})\in\omega((\boldsymbol{N}(0),\boldsymbol{P}(0)))} \mathcal{D}((\boldsymbol{N},\boldsymbol{P})) > \eta.$$
(4.36)

This implies that for any $(\mathbf{N}, \mathbf{P}) \in \mathbb{P}_0$, $\liminf_{t \to +\infty} N_i(t) \ge \eta$ and $\liminf_{t \to +\infty} P_i(t) \ge \eta$. That is, system (1.6) with initial conditions $(\mathbf{N}(0), \mathbf{P}(0)) \in \mathbb{P}_0$ is uniformly persistent.

Step 4.2. System (1.6) has a steady state solution. It follows from [22, Theorem 3.7 and Remark 3.10] that $\Theta(t) : \mathbb{P}_0 \to \mathbb{P}_0$ admits a global attractor A_0 . Then [22, Theorem 4.7] yields that $\Theta(t)$ admits at least one steady-state solution $(\mathbf{N}^*, \mathbf{P}^*) \in \mathbb{P}_0$. Furthermore, we deduce that $\mathbf{N}^*, \mathbf{P}^* \gg \mathbf{0}$ by Lemma 3.4. Thus, system (1.6) admits at least one positive steady state solution $(\mathbf{N}^*, \mathbf{P}^*)$. That is, $(\mathbf{N}^*, \mathbf{P}^*)$ satisfies

$$\begin{cases} \sum_{j=1}^{n} (d_1 D_{ij} + q Q_{ij}) N_j^* + r_i N_i^* (1 - \frac{N_i^*}{k_i}) - a N_i^* P_i^* = 0, \quad i \in \mathcal{A}, \\ \sum_{j=1}^{n} (d_2 D_{ij} + \tau q Q_{ij}) P_j^* + P_i^* (\beta a N_i^* - \gamma) = 0, \quad i \in \mathcal{A}. \end{cases}$$

$$(4.37)$$

The proof of this Theorem is hence completed.

Proof of Theorem 2.2: (1) Note that $\gamma^*(q, d_1, d_2) = \lambda_1(d_2, \tau q, \beta a \theta)$. By means of Lemma 3.1(ii) and Lemma 3.3(iv), it follows that $\lim_{a \to +\infty} \gamma^*(q, d_1, d_2) = \beta a k_n$.

(2) Since by Lemma 3.3(v), there holds $\lim_{d_1 \to +\infty} \theta_i = \frac{\sum_{i=1}^n r_i}{\sum_{i=1}^n \frac{r_i}{k_i}}$, it then indicates from Lemma 3.2 that

$$\lim_{q \to +\infty} \gamma^*(q, d_1, d_2) = \frac{\beta a \sum_{i=1}^n r_i}{\sum_{i=1}^n \frac{r_i}{k_i}}.$$
(4.38)

(3) Note from Lemma 3.3(iv) that $\lim_{d_1\to 0^+} (\theta_1, \theta_2, \cdots, \theta_n) = (0, \cdots, 0, k_n)$. Hence one has $\lim_{d_1\to 0^+} \gamma^*(q, d_1, d_2) = \mu(k_n)$, where $\mu(k_n)$ is the principal eigenvalue of (2.2). Clearly, $\mu(0) = 0$ and $\mu(k_n)$ is strictly increasing with respect to k_n .

(4) Since $\gamma^*(0, d_1, d_2) = \lambda_1(d_2, 0, \theta|_{q=0})$, by Lemma 3.3(vi), there holds

$$\lim_{d_1 \to 0^+} \gamma^*(0, d_1, d_2) = \lambda_1(d_2, 0, \beta a \mathbf{k}).$$
(4.39)

(5) If $q \ge 0$, by applying Lemma 3.1(iii), one can have that

$$\lim_{d_2 \to 0^+} \gamma^*(q, d_1, d_2) = \max\{\beta a\theta_1 - \tau q, \cdots, \beta a\theta_{n-1} - \tau q, \beta a\theta_n\}, \text{ and } \lim_{d_2 \to +\infty} \gamma^*(q, d_1, d_2) = \frac{\sum_{i=1}^n \beta a\theta_i}{n}.$$
(4.40)

The proof of this Theorem is therefore completed.

4.2 Asymptotic profiles of positive steady states

This subsection is devoted to studying the influence of diffusion and advection on the asymptotic profiles of the positive steady state of system (1.6). Therefore, we focus on the steady states system (4.37). Let $U_i = (\frac{d_1}{d_1+q})^{i-1}N_i$ and $V_i = (\frac{d_2}{d_2+\tau q})^{i-1}P_i$. Then (U, V) satisfies $\begin{cases}
-(d_1+q)U_1 + (d_1+q)U_2 + r_1U_1(1-\frac{N_1}{k_1}) - aU_1P_1 = 0, \\
d_1U_{j-1} + (-2d_1-q)U_j + (d_1+q)U_{j+1} + r_jU_j(1-\frac{N_j}{k_j}) - aU_jP_j = 0, \ j \in \mathcal{B}, \\
d_1U_{n-1} - d_1U_n + r_nU_n(1-\frac{N_n}{k_n}) - aU_nP_n = 0, \\
-(d_2+\tau q)V_1 + (d_2+\tau q)V_2 + V_1(\beta aN_1 - \gamma) = 0, \\
d_2V_{j-1} + (-2d_2-\tau q)V_j + (d_2+\tau q)V_{j+1} + V_j(\beta aN_j - \gamma) = 0, \ j \in \mathcal{B}, \\
d_2V_{n-1} - d_2V_n + V_n(\beta aN_n - \gamma) = 0.
\end{cases}$ (4.41)

Lemma 4.3. Suppose (**H**) hold. Let (N, P) be a nonnegative solution of (4.37) with $(N, P) \neq (0, 0)$. Then $0 \ll N \ll \theta$ and $\gamma < \gamma^*(q, d_1, d_2)$.

Proof: Note that, (U, V) is a nonnegative solution of (4.41) with $(U, V) \neq (0, 0)$. Lemma 3.4 yields that $U \gg 0$ and $V \gg 0$. Hence, $N \gg 0$ and $P \gg 0$. From the equations of U in (4.41), we have that

$$\begin{cases} 0 = -(d_1 + q)U_1 + (d_1 + q)U_2 + r_1U_1(1 - \frac{N_1}{k_1}) - aU_1P_1 < -(d_1 + q)U_1 + (d_1 + q)U_2 + r_1U_1(1 - \frac{N_1}{k_1}), \\ 0 = d_1U_{j-1} + (-2d_1 - q)U_j + (d_1 + q)U_{j+1} + r_jU_j(1 - \frac{N_j}{k_j}) - aU_jP_j \\ < d_1U_{j-1} + (-2d_1 - q)U_j + (d_1 + q)U_{j+1} + r_jU_j(1 - \frac{N_j}{k_j}), \ j \in \mathcal{B}, \\ 0 = d_1U_{n-1} - d_1U_n + r_nU_n(1 - \frac{N_n}{k_n}) < d_1U_{n-1} - d_1U_n + r_nU_n(1 - \frac{N_n}{k_n}). \end{cases}$$

$$(4.42)$$

Since $\boldsymbol{\theta}$ is the unique positive solution to (3.7), then $\boldsymbol{w} := \left(\theta_1, \left(\frac{d_1}{d_1+q}\right)\theta_2, \cdots, \left(\frac{d_1}{d_1+q}\right)^{n-1}\theta_n\right)$ is the unique positive solution to

$$\begin{cases} (d_1+q)(w_2-w_1)+r_1(1-\frac{w_1}{k_1})w_1=0,\\ d_1w_{j-1}+(-2d_1-q)w_j+(d_1+q)w_{j+1}+r_j(1-(\frac{d_1+q}{d_1})^{j-1}\frac{w_j}{k_j})w_j=0, \ j\in\mathcal{B},\\ d_1(w_{n-1}-w_n)+r_n(1-(\frac{d_1+q}{d_1})^{n-1}\frac{w_n}{k_n})w_n=0. \end{cases}$$
(4.43)

By the similar arguments in the proof of [2, Lemma 2.2] (see also [16]), one can conclude that $\mathbf{0} \ll \mathbf{U} \ll \mathbf{w}$, which implies $0 \ll N \ll \theta$. Furthermore, from the equations of V in (4.41) and Lemma 3.1(iv), we have

$$\gamma = \lambda_1(d_2, \tau q, \beta a \mathbf{N}) < \lambda_1(d_2, \tau q, \beta a \boldsymbol{\theta}) = \gamma^*(q, d_1, d_2).$$
(4.44)

This completes the proof.

Lemma 4.4. Suppose (**H**) hold. Let (N, P) be a nonnegative solution of (4.37) with $(N, P) \neq (0, 0)$. We have the following results:

(1) if
$$q \ge r_{max} + 2\sqrt{d_1 r_{max}}$$
, there holds $0 < N_1 < \dots < N_n$;
(2) if $q \ge \max\left\{r_{max} + 2\sqrt{d_1 r_{max}}, \frac{\beta a N_n + 2\sqrt{\beta a d_2 N_n}}{\tau}\right\}$, there holds $0 < P_1 < \dots < P_n$.

Proof: (1) It follows from Lemma 4.3 that $N \gg 0$ and $P \gg 0$. Now set $\alpha_i = \left(\frac{d_1}{d_1+q}\right)^{\frac{i-1}{2}}, \xi_i = \alpha_i N_i$ and $\zeta_i = \alpha_i P_i$. Direct calculations yield

$$\begin{cases} \sqrt{d_1(d_1+q)}(\xi_2-\xi_1) + \left[r_1(1-\frac{N_1}{k_1}) - d_1 - q + \sqrt{d_1(d_1+q)} - \frac{a\zeta_1}{\alpha_1}\right]\xi_1 = 0, \\ \sqrt{d_1(d_1+q)}(\xi_{j-1} - 2\xi_j + \xi_{j+1}) + \left[r_j(1-\frac{N_j}{k_j}) - 2d_1 - q + 2\sqrt{d_1(d_1+q)} - \frac{a\zeta_j}{\alpha_j}\right]\xi_j = 0, \ j \in \mathcal{B}. \end{cases}$$

$$(4.45)$$

If $q \geq r_{max} + 2\sqrt{d_1 r_{max}}$, then

$$\begin{cases} r_1(1 - \frac{N_1}{k_1}) - d_1 - q + \sqrt{d_1(d_1 + q)} - \frac{a\zeta_1}{\alpha_1} < 0, \\ r_j(1 - \frac{N_j}{k_j}) - 2d_1 - q + \sqrt{d_1(d_1 + q)} - \frac{a\zeta_j}{\alpha_j} < 0, \ j \in \mathcal{B}. \end{cases}$$
(4.46)

Thus, we get from (4.45) that $0 < \xi_1 < \xi_2 < \cdots < \xi_n$. This indicates that $0 < N_1 < \cdots < N_n$. (2) Set $a_i = \left(\frac{d_2}{d_2 + \tau q}\right)^{\frac{i-1}{2}}$, $\xi_i = a_i N_i$ and $\zeta_i = a_i P_i$. Then we have

$$\left(\sqrt{d_2(d_2 + \tau q)}(\zeta_2 - \zeta_1) + \left[\beta a N_1 - \gamma - d_2 - \tau q + \sqrt{d_2(d_2 + \tau q)} \right] \zeta_1 = 0, \\ \sqrt{d_2(d_2 + \tau q)}(\zeta_{j-1} - 2\zeta_j + \zeta_{j+1}) + \left[\beta a N_j - \gamma - 2d_2 - \tau q + \sqrt{d_2(d_2 + \tau q)} \right] \zeta_j = 0, \ j \in \mathcal{B}.$$
(4.47)

By means of (1), we get that

$$\begin{cases} \beta a N_1 - \gamma - d_2 - \tau q + \sqrt{d_2(d_2 + \tau q)} < \beta a N_n - d_2 - \tau q + \sqrt{d_2(d_2 + \tau q)} \le 0, \\ \beta a N_i - \gamma - 2d_2 - \tau q + \sqrt{d_2(d_2 + \tau q)} < \beta a N_n - 2d_2 - \tau q + \sqrt{d_2(d_2 + \tau q)} \le 0, \ j \in \mathcal{B}. \end{cases}$$
(4.48)

Here $q \ge \max\left\{r_{max} + 2\sqrt{d_1r_{max}}, \frac{\beta aN_n + 2\sqrt{\beta ad_2N_n}}{0 < P_1 < \cdots < P_n}\right\}$ is used. Hence, we get from (4.47) that $0 < \zeta_1 < \zeta_2 < \cdots < \zeta_n$. This indicates that $0 < P_1 < \cdots < P_n$. The proof of this lemma is therefore completed. \Box

Lemma 4.5. Suppose (**H**) hold. Let (N, P) be a nonnegative solution of (4.37) with $(N, P) \neq (0, 0)$. Then the following results hold:

(1) for fixed $d_1, d_2 > 0$, P_i is uniformly bounded for $i \in \mathcal{A}$ as $q \to +\infty$;

(2) for fixed $d_1 > 0$ and $q > \max\left\{r_{max} + 2\sqrt{d_1r_{max}}, \frac{\beta a N_n}{\tau}\right\}$, P_i is uniformly bounded for $i \in \mathcal{A}$ as $d_2 \rightarrow 0^+;$

(3) for fixed $d_2 > 0$ and $q \ge 0$, P_i is uniformly bounded for $i \in \mathcal{A}$ as $d_1 \to +\infty$; (4) for fixed $d_1 > 0$ and $q \ge 0$, P_i is uniformly bounded for $i \in \mathcal{A}$ as $d_2 \to +\infty$.

Proof: By a similar argument as in the proof of Lemma 4.3, we have $N \gg 0$ and $P \gg 0$.

(1) According to Lemma 4.4(2), we only need to show P_n is uniformly bounded as $q \to +\infty$. If this is not true, then by passing to a sequence if necessary, we may assume $P_n \to +\infty$ as $q \to +\infty$. Set $\tilde{P}_i = \frac{P_i}{P_r}$. Then

$$\begin{cases} \sum_{j=1}^{n} (d_1 D_{ij} + q Q_{ij}) N_j + r_i N_i (1 - \frac{N_i}{k_i}) - a N_i \tilde{P}_i P_n = 0, & i \in \mathcal{A}, \\ \sum_{j=1}^{n} (d_2 D_{ij} + \tau q Q_{ij}) \tilde{P}_j + \tilde{P}_i (\beta a N_i - \gamma) = 0, & i \in \mathcal{A}. \end{cases}$$
(4.49)

Since $k_i \leq k_{max}$, $0 < P_i \leq 1$ and $0 < N_i < \theta_i$ is bounded uniformly for $q \to +\infty$ by lemma 3.3 and Lemma 4.3, passing to a sequence if necessary yields that $k_i \to k_i^* \ge 0$, $\tilde{P}_i \to P_i^* \le 1$ and $N_i \to N_i^* \ge 0$. In particular, $P_n^* = 1$, since $\tilde{P}_n = 1$. By dividing the second equation of (4.49) by q, it follows that

$$\begin{cases} (-\frac{d_2}{q} - \tau)\tilde{P}_1 + \frac{d_2}{q}\tilde{P}_2 + \frac{\beta aN_1 - \gamma}{q}\tilde{P}_1 = 0, \\ (\frac{d_2}{q} + \tau)\tilde{P}_{j-1} + (-\frac{2d_2}{q} - \tau)\tilde{P}_j + \frac{d_2}{q}\tilde{P}_{j+1} + \frac{\beta aN_j - \gamma}{q}\tilde{P}_j = 0, j \in \mathcal{B}. \end{cases}$$
(4.50)

Consequently, as $q \to +\infty$, $\tilde{P}_1, \tilde{P}_2, \cdots, \tilde{P}_{n-1} \to 0$. Noting that $\sum_{i=1}^n \sum_{j=1}^n (d_2 D_{ij} + \tau q Q_{ij}) \tilde{P}_j = 0$, we deduce that

$$\sum_{i=1}^{n} \tilde{P}_i(\beta a N_i - \gamma) = 0.$$

$$(4.51)$$

A simple calculation gives $\beta a N_n^* - \gamma = 0$, which shows $N_n^* = \frac{\gamma}{\beta a}$.

Combining the first equation of (4.49) with $\sum_{i=1}^{n} \sum_{j=1}^{n} (d_1 D_{ij} + q Q_{ij}) N_j = 0$ yields that

$$aP_n \sum_{i=1}^n N_i \tilde{P}_i = \sum_{i=1}^n N_i r_i (1 - \frac{N_i}{k_i}), \qquad (4.52)$$

which implies that

$$+\infty \leftarrow aP_n = \frac{\sum_{i=1}^{n} N_i r_i (1 - \frac{N_i}{k_i})}{\sum_{i=1}^{n} N_i \tilde{P}_i}.$$
(4.53)

This is impossible. Hence P_n is uniformly bounded as $q \to +\infty$.

(2) Given $q > \max\left\{r_{max} + 2\sqrt{d_1r_{max}}, \frac{\beta a N_n}{\tau}\right\}$, by the similar arguments as in the proof of Lemma 4.4, we can deduce that for $d_2 \ll 1$, $0 < P_1 < \cdots < P_n$. Hence, we need only to show P_n is bounded (NM) P_n is bounded by the similar arguments as in the proof of Lemma 4.4, we can deduce that for $d_2 \ll 1$, $0 < P_1 < \cdots < P_n$. Hence, we need only to show P_n is bounded to be a similar argument of $P_1 < \cdots < P_n$. uniformly for $d_2 \to 0^+$. Suppose that there exist a sequence d_2^m and positive solutions (N^m, P^m) to

(4.37) such that $\lim_{m \to +\infty} d_2^m = 0$ and $P_n^m \to +\infty$ as $m \to +\infty$. Set $\tilde{P}_i^m = \frac{P_i^m}{P_n^m} \le 1$. Then we reach that

$$\begin{cases} (-d_2^m - \tau q)\tilde{P}_1^m + d_2\tilde{P}_2^m + (\beta aN_1 - \gamma)\tilde{P}_1^m = 0, \\ (d_2^m + \tau q)\tilde{P}_{j-1}^m + (-2d_2^m - \tau q)\tilde{P}_j^m + d_2^m\tilde{P}_{j+1}^m + (\beta aN_j - \gamma)\tilde{P}_j^m = 0, j \in \mathcal{B}, \\ (d_2^m + \tau q)\tilde{P}_{n-1}^m - d_2^m\tilde{P}_n^m + (\beta aN_n - \gamma)\tilde{P}_n^m = 0. \end{cases}$$
(4.54)

Since N_i is bounded for all $d_2 > 0$, we may assume that $\tilde{P}_i^m \to P_i^* \in [0,1]$ and $N_i^m \to N_i^* \ge 0$ as $m \to +\infty$. Letting $m \to +\infty$ yields

$$\begin{cases} -\tau q P_1^* + (\beta a N_1^* - \gamma) P_1^* = 0, \\ \tau q P_{j-1}^* - \tau q P_j^* + (\beta a N_j^* - \gamma) P_j^* = 0, \ j \in \mathcal{B}, \\ \tau q P_{n-1}^* + (\beta a N_n^* - \gamma) P_n^* = 0, \end{cases}$$

$$(4.55)$$

from which $P_1^* = 0$ or $\beta a N_1^* = \gamma + \tau q > 0$ follows. On the other hand, by dividing the equation of N_i^m by P_n^m and letting $m \to +\infty$, one can deduce that $N_1^* P_1^* = 0$. Hence, $P_1^* = 0$. By the similar methods above, it further follows that $P_1^* = \cdots = P_{n-1}^* = 0$, $P_n^* = 1$. Therefore, a contradiction happens according to the same arguments of deriving (4.53). Hence, P_n is bounded for $d_2 \to 0^+$.

(3) Note that N_i is bounded uniformly for $d_1 \to +\infty$. Suppose that P_i is unbounded for $i \in \mathcal{A}$ as $d_1 \to +\infty$. There exist a sequence d_1^m and positive solutions $(\mathbf{N}^m, \mathbf{P}^m)$ to (4.37) such that $d_1^m \to +\infty$ and $P_{i_0}^m := \max_{i \in \mathcal{A}} P_i^m \to +\infty$ as $m \to +\infty$. Let $\hat{P}_i^m = \frac{P_i^m}{P_{i_0}^m} \leq 1$. Then we can deduce that there exists a convergent subsequence of (N_i^m, \hat{P}_i^m) (still denoted by (N_i^m, \hat{P}_i^m)) such that $\hat{P}_i^m \to \hat{P}_i \in [0, 1]$, $N_i^m \to N_i \geq 0$ as $m \to +\infty$. Clearly, $1 = \hat{P}_{i_0}^m \to \hat{P}_{i_0} > 0$ as $m \to +\infty$.

Similar to the proof of (2), we also have $\hat{P} > 0$ and

$$\lambda_1(d_1^m, q, r_i(1 - \frac{N_i^m}{k_i}) - a\hat{P}_i^m P_{i_0}^m) = 0.$$
(4.56)

By Lemma 3.1(iii) again, we have $\lambda_1(d_1^m, q, r_i(1 - \frac{N_i^m}{k_i}) - a\hat{P}_i^m P_{i_0}^m) < 0$ as $m \to +\infty$, a contradiction occurs. Therefore, P_i is bounded for $i \in \mathcal{A}$ as $d_1 \to +\infty$.

(4) Suppose now that P_i is unbounded for $i \in \mathcal{A}$ as $d_2 \to +\infty$. There exist a sequence d_2^m and positive solutions $(\mathbf{N}^m, \mathbf{P}^m)$ to (4.37) such that $d_2^m \to +\infty$ and $P_{i_0}^m := \max_{i \in \mathcal{A}} P_i^m \to +\infty$ as $m \to +\infty$. Let $\hat{P}_i^m = \frac{P_i^m}{P_{i_0}^m} \leq 1$. Then we can deduce that there exists a convergent subsequence of (N_i^m, \hat{P}_i^m) (still denoted by (N_i^m, \hat{P}_i^m)) satisfying $\hat{P}_i^m \to \hat{P}_i \in [0, 1], N_i^m \to N_i \geq 0$ as $m \to +\infty$. Clearly, $1 = \hat{P}_{i_0}^m \to \hat{P}_{i_0} > 0$ as $m \to +\infty$.

Dividing the equation of \hat{P}_i^m by d_2^m and letting $m \to +\infty$ yield that $\hat{P}_1 = \cdots = \hat{P}_n = \hat{P}_{i_0} = 1$. Therefore, a contradictions occurs according the same arguments of deriving (4.53). P_i is hence bounded for $d_2 \to +\infty$.

The proof of this lemma is thus complete.

It follows from Theorem 2.1(2) that (1.6) has a positive steady state (N^*, P^*) when $0 < \gamma < \gamma^*(q, d_1, d_2)$. In the following, we shall investigate the positive steady state (N^*, P^*) .

Lemma 4.6. Suppose (**H**) hold, $0 < \gamma < \gamma^*(q, d_1, d_2)$. Then for $i \in C$,

$$\left(\frac{d_1 + q + 2(aP_n^* + \max_{i \in \mathcal{A}} \frac{r_i}{k_i} N_n^*)}{d_1} \right)^{i-n} N_n^* < N_i^* < \left(\frac{d_1 + q - 2r_{max}}{d_1} \right)^{i-n} N_n^*,$$

if $q \ge \max\{d_1 + 2r_{max}, r_{max} + 2\sqrt{d_1 r_{max}}\},$
$$\left(\frac{d_2 + \tau q + 2\gamma}{d_2} \right)^{i-n} P_n^* < P_i^* < \left(\frac{d_2 + \tau q - 2\beta a N_n^*}{d_2} \right)^{i-n} P_n^*, \text{ if } q \ge \max\{\frac{d_2 + 2\beta a N_n^*}{\tau}, \frac{\beta a N_n^* + 2\sqrt{\beta a N_n^* d_2}}{\tau}\}.$$
(4.57)

$$\begin{aligned} \mathbf{Proof:} \quad & \text{Set } \alpha = \frac{d_1 + q - 2r_{max}}{d_1}, \ \delta = \frac{d_1 + q + 2(aP_n^* + \max_{i \in \mathcal{A}} \frac{r_i}{k_i} N_n^*)}{d_1}, \ A = \frac{d_2 + \tau q - 2\beta a N_n^*}{d_2}, \ B = \frac{d_2 + \tau q + 2\gamma}{d_2}, \ \alpha_i = \alpha^{i-1}, \ \delta_i = \delta^{i-1}, \ A_i = A^{i-1}, \ B_i = B^{i-1}. \ \text{Let } \overline{N_i^*} = N_n^* \frac{\alpha_i}{\alpha_n}, \ \underline{N_i^*} = N_n^* \frac{\delta_i}{\delta_n}, \ \overline{P_i^*} = P_n^* \frac{A_i}{A_n}, \\ \underline{P_i^*} = P_n^* \frac{B_i}{B_n}. \ \text{Letting } u_i = (\frac{d_1}{d_1 + q})^{\frac{i-1}{2}} (\overline{N_i^*} - N_i^*), \ \text{it is obvious that} \\ \begin{cases} \sqrt{d_1(d_1 + q)}(u_2 - u_1) + \left[r_1(1 - \frac{N_1^*}{k_1}) - d_1 - q + \sqrt{d_1(d_1 + q)} - aP_1^*\right] u_1 < 0, \\ \sqrt{d_1(d_1 + q)}(u_{j-1} - 2u_j + u_{j+1}) + \left[r_j(1 - \frac{N_j^*}{k_j}) - 2d_1 - q + 2\sqrt{d_1(d_1 + q)} - aP_j^*\right] u_j < 0, \ j \in \mathcal{B}, \end{cases} \end{aligned}$$

for $q \ge \max\{d_1 + 2r_{max}, r_{max} + 2\sqrt{d_1r_{max}}\}$, some similar methods in obtaining (3.9) give rise to $u_i > 0$ and thus $\overline{N}_i^* > N_i^*$ for $i \in \mathcal{C}$.

By means of the similar arguments above, combining the equations of $\underline{N}_i^*, \overline{P}_i^*, \underline{P}_i^*$ with

$$q \ge \max\{d_1, \frac{d_2 + 2\beta ak_n}{\tau}, \frac{\beta ak_n + 2\sqrt{\beta ak_n d_2}}{\tau}, \frac{d_2 - 2\gamma}{\tau}\}$$

$$(4.59)$$

yields that $N_i^* > \underline{N}_i^*, \overline{P}_i^* > P_i^* > \underline{P}_i^*$ for $i \in \mathcal{C}$. Hence, we finish the proof of this lemma.

Proof of Theorem 2.3: Note that $0 < \gamma < \beta a k_n$ and $\lim_{q \to +\infty} \gamma^*(q, d_1, d_2) = \beta a k_n$ guarantee that there exists a sufficiently large positive constant Q such that $0 < \gamma < \gamma^*(q, d_1, d_2)$ when q > Q. This implies that a positive steady state of system (1.6) exists for all large q (see Theorem 2.1(2)).

Notice from Lemmas 4.3 and 4.5(1) that N_n^* and P_n^* are uniformly bounded as $q \to +\infty$. By choosing a subsequence if necessary, we may assume that $(N_n^*, P_n^*) \to (p_1, p_2)$ as $q \to +\infty$ for some constants p_1 and p_2 . It follows from (4.57) that

$$\frac{N_i^*}{N_n^*} \to 0, \frac{P_i^*}{P_n^*} \to 0, \text{ for } i \in \mathcal{C}, q \to +\infty.$$
(4.60)

Applying (4.37), we get

Furthermore, we reach that

$$\begin{cases} 0 = r_n - \frac{r_n}{k_n} N_n^* - a P_n^* + \sum_{i=1}^{n-1} \frac{N_i^*}{N_n^*} (r_i - \frac{r_i}{k_i} \frac{N_i^*}{N_n^*} N_n^* - a \frac{P_i^*}{P_n^*} P_n^*) \to r_n - \frac{r_n}{k_n} p_1 - a p_2, \\ 0 = \beta a N_n^* - \gamma + \sum_{i=1}^{n-1} \frac{P_i^*}{P_n^*} (\beta a \frac{N_i^*}{N_n^*} N_n^* - \gamma) \to \beta a p_1 - \gamma, \end{cases}$$
(4.62)

by letting $q \to +\infty$. Therefore, we obtain that

$$(p_1, p_2) = \left(\frac{\gamma}{\beta a}, \frac{1}{a}\left(r_n - \frac{r_n}{k_n}\frac{\gamma}{\beta a}\right)\right).$$
(4.63)

According to (4.57), we have

$$\left[\left(\frac{d_1}{d_1 + q + 2(aP_n^* + \max_{i \in \mathcal{A}} \frac{r_i}{k_i} N_n^*)} \right)^{n-i} - \left(\frac{d_1}{d_1 + q} \right)^{n-i} \right] N_n^* < N_i^* - N_n^* \left(\frac{d_1}{d_1 + q} \right)^{n-i} < \left[\left(\frac{d_1}{d_1 + q - 2r_{max}} \right)^{n-i} - \left(\frac{d_1}{d_1 + q} \right)^{n-i} \right] N_n^*,$$

$$(4.64)$$

which implies

$$\max_{i \in \mathcal{A}} \left(N_i^* - N_n^* \left(\frac{d_1}{d_1 + q} \right)^{n-i} \right) \to 0 \text{ as } q \to +\infty.$$
(4.65)

Similarly, it can be proved by (4.57) that

$$\max_{i \in \mathcal{A}} \left(P_i^* - P_n^* \left(\frac{d_2}{d_2 + \tau q} \right)^{n-i} \right) \to 0 \text{ as } q \to +\infty.$$
(4.66)

That is,

$$\max\{\max_{i\in\mathcal{A}}\left(N_{i}^{*}-N_{n}^{*}(\frac{d_{1}}{d_{1}+q})^{n-i}\right),\max_{i\in\mathcal{A}}\left(P_{i}^{*}-P_{n}^{*}(\frac{d_{2}}{d_{2}+\tau q})^{n-i}\right)\}\to 0 \text{ as } q\to+\infty,\tag{4.67}$$

which implies that $(N_i^*, P_i^*) \to (0, 0)$ uniformly for $i \in \mathcal{C}$ as $q \to +\infty$. In conclusion, we complete the proof of Theorem 2.3.

Proof of Theorem 2.4: Note that $0 < \gamma < \beta a k_{min}$ and $\lim_{d_1 \to +\infty} \gamma^*(q, d_1, d_2) = \frac{\beta a \sum_{i=1}^n r_i}{\sum_{i=1}^n \frac{r_i}{k_i}} > \beta a k_{min}$ guarantee that there exists a sufficiently large positive constant D_1 such that $0 < \gamma < \gamma^*(q, d_1, d_2)$ when $d_1 > D_1$. This implies that a positive steady state of system (1.6) exists for all large d_1 (see Theorem 2.1(2)).

It follows from Lemmas 4.3, 3.3(v) and 4.5(3) that N_i^* and P_i^* are uniformly bounded for $i \in \mathcal{A}$ as $d_1 \to +\infty$. We may assume by passing to a subsequence if necessary that $P_i^* \to \dot{P}_i$ as $d_1 \to +\infty$. Dividing the equation of N_i^* in (4.37) by d_1 and taking $d_1 \to +\infty$ lead to $N_i^* \to s \ge 0$ for all i. Let $\max_{i \in \mathcal{A}} P_i^* = P_{i_0(d_1)}^*$ and $\tilde{P}_i := \frac{P_i^*}{P_{i_0(d_1)}^*}$. Then it follows that $\tilde{P}_i \to \frac{\dot{P}_i}{P_{i_0(\infty)}^*} \ge 0$ for $d_1 \to +\infty$. Clearly, \tilde{P}_i satisfies

$$\sum_{j=1}^{n} (d_2 D_{ij} + \tau q Q_{ij}) \tilde{P}_j + \tilde{P}_i (\beta a N_i^* - \gamma) = 0.$$
(4.68)

After taking $d_1 \to +\infty$, we arrive at

$$\sum_{j=1}^{n} (d_2 D_{ij} + \tau q Q_{ij}) \frac{\dot{P}_j}{P_{i_0(\infty)}^*} + \frac{\dot{P}_i}{P_{i_0(\infty)}^*} (\beta as - \gamma) = 0.$$
(4.69)

Adding (4.69) up with respect to *i* yields

$$\sum_{i=1}^{n} \frac{\dot{P}_{i}}{P_{i_{0}(\infty)}^{*}} (\beta as - \gamma) = 0.$$
(4.70)

The above equality then implies that $s = \frac{\gamma}{\beta a}$. Letting $d_1 \to +\infty$ in the equation of P_i^* in (4.37), we see that

$$\sum_{j=1}^{n} (d_2 D_{ij} + \tau q Q_{ij}) \dot{P}_j = 0, \qquad (4.71)$$

which implies that $\dot{P}_i = (1 + \frac{\tau q}{d_2})^{i-1} \dot{P}_1$.

After letting $d_1 \to +\infty$ in the equation of N_i^* in (4.37) and adding them up respect to *i* from 1 to *n*, we can see that

$$\sum_{i=1}^{n} r_i (1 - \frac{\gamma}{\beta a k_i}) = a \dot{P}_1 \sum_{i=1}^{n} (1 + \frac{\tau q}{d_2})^{i-1}.$$
(4.72)

That is,
$$\dot{P}_1 = (1 + \frac{\tau q}{d_2})^{i-1} \frac{\sum_{j=1}^n r_j (1 - \frac{r_j}{\beta a k_j})}{\frac{d_2 a}{\tau q} [(1 + \frac{\tau q}{d_2})^n - 1]}$$
. Then (2.3) is obtained.

Proof of Theorem 2.5: Note that $0 < \gamma < \gamma^*(q, d_1, d_2)$. This implies that a positive steady state of system (1.6) exists(see Theorem 2.1(2)).

Note that N_i^* is uniformly bounded as $d_2 \to 0^+$ (see Lemma 4.3). Since $q > \max\{r_{max} + 2\sqrt{d_1 r_{max}}, \frac{2\beta a N_n^*}{\tau}\}$, by Lemma 4.5(2) and (4.57), we have that P_n^* is uniformly bounded and

$$\left(\frac{d_2 + \tau q + 2\gamma}{d_2}\right)^{i-n} P_n^* < P_i^* < \left(\frac{d_2 + \tau q - 2\beta a N_n^*}{d_2}\right)^{i-n} P_n^*$$
(4.73)

for $d_2 \to 0^+$. Therefore,

$$\max_{i} \left(P_{i}^{*} - P_{n}^{*} \left(\frac{d_{1}}{d_{2} + \tau q} \right)^{n-i} \right) \to 0 \text{ as } d_{2} \to 0^{+},$$
(4.74)

and we may suppose $(N_i^*, P_n^*) \to (N_i^0, l)$ as $d_2 \to 0^+$ by passing to a subsequence if necessary. Here $N_i^0 \ge 0, l \ge 0$.

Set $\tilde{P}_i = \frac{P_i^*}{P_n^*}$. Then \tilde{P}_i satisfies

$$\sum_{j=1}^{n} (d_2 D_{ij} + \tau q Q_{ij}) \tilde{P}_j + \tilde{P}_i (\beta a N_i^* - \gamma) = 0.$$
(4.75)

If l > 0, then we must have $\tilde{P}_i \to 0 (i \in C)$ as $d_2 \to 0^+$. Combined with (4.75), we obtain that l = 0. Setting $d_2 \to 0^+$ in the equation of N_i^* yields $N_i^0 = \theta_i$. The proof of Theorem 2.5 is thus complete. \Box

At last, before giving the proof of Theorem 2.6, we introduce the following lemma.

Lemma 4.7. Suppose (**H**) hold, $0 < \gamma < \frac{\beta a \sum_{i=1}^{n} \theta_i}{n}$ and $y \ge 0$. The system

$$\begin{cases} \sum_{j=1}^{n} (d_1 D_{ij} + q Q_{ij}) Z_j + Z_i (r_i - \frac{r_i}{k_i} Z_i - ay) = 0, & i \in \mathcal{A}, \\ \beta a \sum_{i=1}^{n} Z_i = \gamma n, & i \in \mathcal{A}. \end{cases}$$
(4.76)

admits a unique positive solution $(Z_i(\tilde{y}), \tilde{y})$.

Proof: Let $y^* = \frac{\lambda_1(d_1, q, r)}{a} > 0$. Clearly, it follows from [2, Lemma 2.2](see also [16]) that for $y \ge y^*$, $\mathbf{Z} = \mathbf{0}$ is the unique solution of the first equation of (4.76), and for $0 \le y < y^*$, the first equation of (4.76) admits a unique positive solution, denoted by $Z_i(y)$. Now choosing $0 \le y^1 < y^2 < y^*$, we obtain

$$\sum_{j=1}^{n} (d_1 D_{ij} + q Q_{ij}) Z_j + Z_i (r_i - \frac{r_i}{k_i} Z_i - ay^2) < \sum_{j=1}^{n} (d_1 D_{ij} + q Q_{ij}) Z_j + Z_i (r_i - \frac{r_i}{k_i} Z_i - ay^1).$$
(4.77)

By applying some similar arguments in the proof of [2, Lemma 2.2] (see also [16]), we have $0 < Z_i(y^2) < Z_i(y^1)$ which yields $Z_i(y)$ is strictly decreasing with respect to y in $[0, y^*]$. Thus, $F(y) = \sum_{i=1}^n Z_i(y) - \frac{\gamma n}{\beta a}$ is strictly decreasing with respect to y in $[0, y^*]$. Since $F(0) = \sum_{i=1}^n Z_i(0) - \frac{\gamma n}{\beta a} = \sum_{i=1}^n \theta_i - \frac{\gamma n}{\beta a} > 0$,

 $F(y^*) = -\frac{\gamma n}{\beta a} < 0$, we can deduce from the monotonicity of F(y) that there exists a unique $\tilde{y} \in (0, y^*)$ such that $F(\tilde{y}) = 0$. After substituting \tilde{y} into the first equation of (4.76), we deduce that (4.76) admits a unique positive solution $(Z_i(\tilde{y}), \tilde{y})$.

Proof of Theorem 2.6: Note that $0 < \gamma < \frac{\beta a \sum_{i=1}^{n} \theta_i}{n}$ and $\lim_{d_2 \to +\infty} \gamma^*(q, d_1, d_2) = \frac{\beta a \sum_{i=1}^{n} \theta_i}{n}$ guarantee that there exists a sufficiently large positive constant D_2 such that $0 < \gamma < \gamma^*(q, d_1, d_2)$ when $d_2 > D_2$. This implies that a positive steady state of system (1.6) exists for all large d_2 (see Theorem 2.1(2)).

It can be shown similarly to Lemma 4.3 and 4.5(4) that N_i^*, P_i^* are uniformly bounded as $d_2 \to +\infty$. By using similar arguments as in the proof of Theorem 2.4, we may assume that $(N_i^*, P_i^*) \to (\hat{\theta}_i, z)$ as $d_2 \to +\infty$ by passing to a subsequence if necessary, where $(\hat{\theta}_i, z)$ satisfies

$$\begin{cases} \sum_{j=1}^{n} (d_1 D_{ij} + q Q_{ij}) \hat{\theta}_j + \hat{\theta}_i (r_i - \frac{r_i}{k_i} \hat{\theta}_i - az) = 0, \quad i \in \mathcal{A}, \\ \sum_{i=1}^{n} (\beta a \hat{\theta}_i - \gamma) = 0, \qquad i \in \mathcal{A}. \end{cases}$$
(4.78)

By Lemma 4.7, it follows that $(\hat{\theta}_i, z)$ exists and is unique.

5 Discussion

In this paper, we consider the specialist predator-prey model (1.6) in a closed advective patchy environment. We obtain threshold dynamics in terms of the mortality rate of the specialist predators. In Theorem 2.1, we prove that the specialist predators will die out if their mortality rate exceeds a critical value γ^* and persist if the mortality rate is less than γ^* . In Theorem 2.1, we also prove that, no matter how large advection and diffusion rates are, the specialist predators can invade successfully as long as they maintain a small mortality rate.

We also investigate the influence of advection and diffusion on the asymptotic profiles of the positive steady state solutions of systems (1.6) and find that diffusion and advection can dramatically affect the spatial distribution of species. In Theorems 2.3, we demonstrate that two species will coexist and concentrate at the patch n for a large positive advection. This implies that the advection does not eliminate coexistence of species but significantly affect the distribution of species. The intuitive biological explanation is that predators always keep pace with the prey, allowing them to successfully invade and coexist with the prey at the patch n as flow speed increases. In Theorems 2.4 and 2.6, we show that, if the diffusion rate of the prey (resp. the predator) is large, the species will coexist.

If the prey has a small diffusion rate, in Theorems 2.1 and 2.2, we prove that the specialist predators can always invade successfully as long as the mortality rate is suitably small.

If the specialist predators have a small diffusion rate, they can successfully invade but with a small population density which tends to zero as the diffusion rate goes to zero. The biological explanation is that the specialist predators cannot catch up with the prey if they do not move fast (see Theorem 2.2(5) and 2.5).

We end this section by proposing several interesting problems that deserve further consideration. The first one is that whether there is the uniqueness of positive steady state solutions of system (1.6). Note that the arguments of deriving the uniqueness as in Step 3 of [26, Theorem 3.1] or the proof of [24, Lemma 3.3, Theorem 3.4] do not work in our model (4.37) here. For our model, a probable method to obtain the uniqueness is to verify the invertible of the coefficient matrix. However, due to the complexity of the coefficient matrix, it is to calculate the determinant or to consider the uniqueness. Another problem concerns the dynamics of the specialist predator-prey model (1.6) in open advective patchy environments. Also, how does the dynamics change for the generalist predator-prey patchy system ? We leave these challenging problems for future investigation.

References

- Chen, S., Liu, J., Wu, Y., Invasion analysis of a two-species Lotka-Volterra competition model in an advective patchy environment, *Stud. Appl. Math.* 149 (2022) 762-797.
- [2] Chen, S., Liu, J., Wu, Y., On the impact of spatial heterogeneity and drift rate in a three-patch two-species lotka-volterra competition model over a stream, Z. Angew. Math. Phys. (2023) 74:117.

- [3] Chen, S., Shi, J., Shuai, Z., Wu, Y., Global dynamics of a Lotka-Volterra competition patch model, *Nonlinearity* 35(2) (2022) 817-842.
- [4] Ge, Q., Tang, D., Global dynamics of two-species lotka-volterra competition diffusion-advection system with general carrying capacities and intrinsic growth rates, J. Dyn. Differ. Equ. (2022).
- [5] Hilker, F. M., Lewis, M. A., Predator-prey systems in streams and rivers, *Theor. Ecol.* 3 (2010) 175-193.
- [6] Huang, Q.-H., Jin, Y., Lewis, M.A., R₀ analysis of a Benthic-drift model for a stream population, SIAM J. Appl. Dyn. Syst. 15(1) (2016) 287-321.
- [7] Jiang, H., Lam, K.-Y., Lou, Y., Are two-patch models sufficient? The evolution of dispersal and topology of river network modules. *Bull. Math. Biol.* 82(10) (2020) 131 42.
- [8] Jiang, H., Lam, K.-Y., Lou, Y., Three-patch models for the evolution of dispersal in advective environments: varying drift and network topology, *Bull. Math. Biol.* 83(10) (2021) 109 46.
- [9] Jin, Y., Lewis, M.A., Seasonal influences on population spread and persistence in streams: critical domain size, SIAM J. Appl. Math. 71(4) (2011) 1241-1262.
- [10] Lam, K.Y., Lou, Y., Lutscher, F., Evolution of dispersal in closed advective environments, J. Biol. Dyn. 9(1) (2015) 188-212.
- [11] Lou, Y., Lutscher, F., Evolution of dispersal in open advective environments, J. Math. Biol. 69(6-7) (2014) 1319-1342.
- [12] Lou, Y., Nie, H., Global dynamics of a generalist predator-prey model in open advective environments, J. Math. Biol. 84 (2022) 46.
- [13] Lou, Y., Nie, H., Wang, Y. Coexistence and bistability of a competition model in open advective environments, *Math. Biosci.* 306 (2018) 10-19.
- [14] Lou, Y., Xiao, D.-M., Zhou, P., Qualitative analysis for a Lotka–Volterra competition system in advective homogeneous environment, *Discrete Contin. Dyn. Syst.* 36(2) (2016) 953-969.
- [15] Lou, Y., Zhou, P., Evolution of dispersal in advective homogeneous environment: the effect of boundary conditions, J. Differ. Equ. 259(1) (2015) 141-171.
- [16] Lu, Z.Y., Takeuchi, Y., Global asymptotic behavior in single-species discrete diffusion systems, J. Math. Biol. 32(1) (1993) 67-77.
- [17] Lutscher, F., Lewis, M.A., McCauley, E., Effects of heterogeneity on spread and persistence in rivers, Bull. Math. Biol. 68(8) (2006) 2129-2160.
- [18] Lutscher, F., McCauley, E., Lewis, M.A., Spatial patterns and coexistence mechanisms in systems with unidirectional flow, *Theor. Popul. Biol.* 71(3) (2007) 267-277.
- [19] Lutscher, F., Nisbet, R. M., Pachepsky, E., Population persistence in the face of advection, *Theor. Ecol.* 3 (2010) 271-284.
- [20] Lutscher, F. Pachepsky, E., Lewis, M. A., The effect of dispersal patterns on stream populations, SIAM J. Appl. Math. 65 (2005) 1305-1327.
- [21] Ma, L., Tang, D., Evolution of dispersal in advective homogeneous environments, Discrete Contin. Dyn. Syst. 40(10) (2020) 5815-5830.
- [22] Magal, P., Zhao, X.Q., Global attractors and steady states for uniformly persistent dynamical systems, SIAM J. Math. Anal. 37 (2005) 251-275.
- [23] Müller, K., The colonization cycle of freshwater insects, *Oecologia* 52 (1982) 202-207.

- [24] Nie, H., Hsu, S. B., Wu, J. H., Coexistence solutions of a competition model with two species in a water column, *Discrete Contin. Dyn. Syst. Ser. B* 20 (2015) 2691-2714.
- [25] Nie, H., Liu, C.R., Wang, Z. G., Global dynamics of an ecosystem in open advective environments, Int. J. Bifurc. Chaos Appl. Sci. Eng. 31 (2021) 2150087.
- [26] Nie, H., Wang, B., Wu, J. H., Invasion analysis on a predator-prey system in open advective environments, J. Math. Biol. 81 (2020) 1429-1463.
- [27] Nie, H., Xin, S. X., Shu, H. Y., Effects of diffusion and advection on predator-prey dynamics in closed environments, J. Differ. Equ. 367 (2023) 290-331.
- [28] Smith, H.L., Zhao, X.Q., Robust persistence for semidynamical systems, Nonlinear Anal. 47 (2001) 6169-6179.
- [29] Speirs, D. C., Gurney, W. S. C., Population persistence in rivers and estuaries, *Ecology* 82 (2001) 1219-1237.
- [30] Tang, D., Zhou, P., On a Lotka-Volterra competition-diffusion-advection system: homogeneity vs heterogeneity, J. Differ. Equ. 268 (2020) 1570-1599.
- [31] Vasilyeva, O., Lutscher, F., Population dynamics in rivers: analysis of steady states, Can. Appl. Math. Q. 18(4) (2010) 439-469.
- [32] Vasilyeva, O., Lutscher, F., How flow speed alters competitive outcome in advective environments, Bull. Math. Biol. 74(12) (2012) 2935-2958.
- [33] Wang, J. F., Nie, H., Invasion dynamics of a predator-prey system in closed advective environments, J. Differ. Equ. 318 (2022) 298-322.
- [34] Xin, S., Li, L., Nie, H., The effect of advection on a predator-prey model in open advective environments, *Commun. Nonlinear Sci. Numer. Simul.* 113:Paper No. 106567 17 (2022).
- [35] Yan, X., Nie, H., Zhou, P., On a competition-diffusion-advection system from river ecology: mathematical analysis and numerical study, SIAM J. Appl. Dyn. Syst. 21(1) (2022) 438-469.
- [36] Zhou, P., Xiao, D.M., Global dynamics of a classical Lotka-Volterra competition-diffusionadvection system, J. Funct. Anal. 275 (2018) 356-380.
- [37] Zhao, X.-Q., Zhou, P., On a Lotka-Volterra competition model: the effects of advection and spatial variation, *Calc. Var. Partial Differ. Equ.* 55(4) (2016) 73.
- [38] Zhou, P., On a Lotka-Volterra competition system: diffusion vs advection, Calc. Var. Partial Differ. Equ. 55(6) (2016) 137.