

Frobenius numbers associated with Diophantine triples of $x^2 + y^2 = z^r$ (extended version)

Takao Komatsu

Department of Mathematical Sciences, School of Science
Zhejiang Sci-Tech University
Hangzhou 310018 China
komatsu@zstu.edu.cn

Neha Gupta

Department of Mathematics, School of Natural Sciences
Shiv Nadar Institute of Eminence
Gautam Buddha Nagar - 201314 India
neha.gupta@snu.edu.in

Manoj Upreti

Department of Mathematics, School of Natural Sciences
Shiv Nadar Institute of Eminence
Gautam Buddha Nagar - 201314 India
mu506@snu.edu.in

MR Subject Classifications: Primary 11D07, 11D25; Secondary 05A15, 11D04,
20M14

Abstract

We give an explicit formula for the p -Frobenius number of triples associated with Diophantine equations $x^2 + y^2 = z^r$, that is, the largest positive integer that can only be represented in p ways by combining the three integers of the solutions of Diophantine equations $x^2 + y^2 = z^r$. When $r = 2$, the Frobenius number has already been given.

Keywords: Frobenius problem, Diophantine equations, Pythagorean triples, Apéry set,

1 Introduction

Diophantine equations are a fundamental part and one of the oldest branches of number theory. The main study is of polynomial equations or systems

of equations, in particular, in integers. Though there are many aspects and applications, Diophantine equations are used to characterize certain problems in Diophantine approximations. In [7, 8], we computed upper and lower bounds for the approximation of hyperbolic functions at points $1/s$ ($s = 1, 2, \dots$) by rationals x/y , such that x , y and z form Pythagorean triples. In [9, 5], we considered Diophantine approximations x/y to values ξ of hyperbolic functions, where (x, y, z) is the solution of more Diophantine equations, including $x^2 + y^2 = z^4$.

For integer $k \geq 2$, consider a set of positive integers $A = \{a_1, \dots, a_k\}$ with $\gcd(A) = \gcd(a_1, \dots, a_k) = 1$. To find the number of non-negative integral representations x_1, x_2, \dots, x_k , denoted by $d(n; A) = d(n; a_1, a_2, \dots, a_k)$, to $a_1x_1 + a_2x_2 + \dots + a_kx_k = n$ for a given positive integer n is one of the most important and interesting topics. This number is often called the *denumerant* and is equal to the coefficient of x^n in $1/(1-x^{a_1})(1-x^{a_2})\dots(1-x^{a_k})$ ([33]). Sylvester [32] and Cayley [4] showed that $d(n; a_1, a_2, \dots, a_k)$ can be expressed as the sum of a polynomial in n of degree $k-1$ and a periodic function of period $a_1a_2\dots a_k$. For two variables, a formula for $d(n; a_1, a_2)$ is obtained in [36]. For three variables in the pairwise coprime case $d(n; a_1, a_2, a_3)$, in [13], the periodic function part is expressed in terms of trigonometric functions.

For a non-negative integer p , define S_p and G_p by

$$S_p(A) = \{n \in \mathbb{N}_0 \mid d(n; A) > p\}$$

and

$$G_p(A) = \{n \in \mathbb{N}_0 \mid d(n; A) \leq p\}$$

respectively, satisfying $S_p \cup G_p = \mathbb{N}_0$, which is the set of non-negative integers. The set S_p is called *p-numerical semigroup* because $S(A) = S_0(A)$ is a usual numerical semigroup. G_p is the set of *p-gaps*. Define $g_p(A)$, $n_p(A)$ and $s_p(A)$ by

$$g_p(A) = \max_{n \in G_p(A)} n, \quad n_p(A) = \sum_{n \in G_p(A)} 1, \quad s_p(A) = \sum_{n \in G_p(A)} n,$$

respectively, and are called the *p-Frobenius number*, the *p-Sylvester number* (or *p-genus*) and the *p-Sylvester sum*, respectively. When $p = 0$, $g(A) = g_0(A)$, $n(A) = n_0(A)$ and $s(A) = s_0(A)$ are the original Frobenius number, Sylvester number (or genus) and Sylvester sum, respectively. Finding such values is one of the crucial matters in the Diophantine problem of Frobenius. More detailed descriptions of the *p-numerical semigroups* and their symmetric properties can be found in [23].

The Frobenius problem (also known as the Coin Exchange Problem or Postage Stamp Problem or Chicken McNugget Problem) has a long history and is one of the popular problems that has attracted the attention of experts as well as amateurs. For two variables $A = \{a, b\}$, it is known that

$$g(a, b) = (a - 1)(b - 1) - 1 \quad \text{and} \quad n(a, b) = \frac{(a - 1)(b - 1)}{2}$$

([33, 34]). For three or more variables, the Frobenius number cannot be given by any set of closed formulas, which can be reduced to a finite set of certain polynomials ([6]). For three variables, various algorithms have been devised for finding the Frobenius number. For example, in [28], the Frobenius number is uniquely determined by six positive integers that are the solution to a system of three polynomial equations. In [11], a general algorithm is given by using a 3×3 matrix. Nevertheless, explicit closed formulas have been found only for some special cases, including arithmetic, geometric, Mersenne, repunits and triangular (see [26, 29, 30] and references therein). We are interested in finding explicit closed forms, which is one of the most crucial matters in Frobenius problem. Our method has an advantage in terms of visually grasping the elements of the Apéry set, and is more useful to get more related values, including genus (Sylvester number), Sylvester sum [37], weighted power Sylvester sum [14, 24, 25] and so on.

We are interested in finding a closed or explicit form for the p -Frobenius number, which is more difficult when $p > 0$. For three or more variables, no concrete examples had been found until recently, when the first author succeeded in giving the p -Frobenius number as a closed-form expression for the triangular number triplet ([15]), for repunits ([16]), Fibonacci triplet ([21]), Jacobsthal triplets ([18, 19]) and arithmetic triplets ([22]).

We give an explicit formula for the p -Frobenius number of triples associated with Diophantine equations, that is the largest positive integer that can only be represented in p ways by combining the three integers of the solutions of Diophantine equations. When $p = 0$, it is the original Frobenius number in the famous Diophantine problem of Frobenius. We also obtain closed forms for the number of positive integers, and the largest positive integer that can be represented in only p ways by combining the three integers of the Diophantine triple.

In this paper, we study the numerical semigroup of the triples (x, y, z) , satisfying the Diophantine equation $x^2 + y^2 = z^r$ ($r \geq 2$). When $r = 2$, the Frobenius number of the Pythagorean triple is given in [12]. The Frobenius number of a little modified triple is studied in [20].

2 Preliminaries

We introduce the p -Apéry set (see [1]) below in order to obtain the formulas for $g_p(A)$, $n_p(A)$ and $s_p(A)$. Without loss of generality, we assume that $a_1 = \min(A)$.

Definition 1. Let p be a nonnegative integer. For a set of positive integers $A = \{a_1, a_2, \dots, a_\kappa\}$ with $\gcd(A) = 1$ and $a_1 = \min(A)$ we denote by

$$\text{Ap}_p(A) = \text{Ap}_p(a_1, a_2, \dots, a_\kappa) = \{m_0^{(p)}, m_1^{(p)}, \dots, m_{a_1-1}^{(p)}\},$$

the p -Apéry set of A , where each positive integer $m_i^{(p)}$ ($0 \leq i \leq a_1 - 1$) satisfies the conditions:

$$(i) m_i^{(p)} \equiv i \pmod{a_1}, \quad (ii) m_i^{(p)} \in S_p(A), \quad (iii) m_i^{(p)} - a_1 \notin S_p(A)$$

Note that $m_0^{(0)}$ is defined to be 0.

It follows that for each p ,

$$\text{Ap}_p(A) \equiv \{0, 1, \dots, a_1 - 1\} \pmod{a_1}.$$

When $k \geq 3$, it is hard to find any explicit form of $g_p(A)$ as well as $n_p(A)$ and $s_p(A)$. Nevertheless, the following convenient formulas are known (for a more general case, see [17]). Though finding $m_j^{(p)}$ is enough hard in general, we can obtain it for some special sequences (a_1, a_2, \dots, a_k) .

Lemma 1. Let k and p be integers with $k \geq 2$ and $p \geq 0$. Assume that $\gcd(a_1, a_2, \dots, a_k) = 1$. We have

$$g_p(a_1, a_2, \dots, a_k) = \left(\max_{0 \leq j \leq a_1-1} m_j^{(p)} \right) - a_1, \quad (1)$$

$$n_p(a_1, a_2, \dots, a_k) = \frac{1}{a_1} \sum_{j=0}^{a_1-1} m_j^{(p)} - \frac{a_1 - 1}{2}, \quad (2)$$

$$s_p(a_1, a_2, \dots, a_k) = \frac{1}{2a_1} \sum_{j=0}^{a_1-1} (m_j^{(p)})^2 - \frac{1}{2} \sum_{j=0}^{a_1-1} m_j^{(p)} + \frac{a_1^2 - 1}{12}. \quad (3)$$

Remark. When $p = 0$, the formulas (1), (2) and (3) reduce to the formulas by Brauer and Shockley [2], Selmer [31], and Tripathi [37], respectively:

$$g(a_1, a_2, \dots, a_k) = \left(\max_{1 \leq j \leq a_1-1} m_j \right) - a_1,$$

$$n(a_1, a_2, \dots, a_k) = \frac{1}{a_1} \sum_{j=1}^{a_1-1} m_j - \frac{a_1 - 1}{2},$$

$$s(a_1, a_2, \dots, a_k) = \frac{1}{2a_1} \sum_{j=1}^{a_1-1} (m_j)^2 - \frac{1}{2} \sum_{j=1}^{a_1-1} m_j + \frac{a_1^2 - 1}{12},$$

where $m_j = m_j^{(0)}$ ($1 \leq j \leq a_1 - 1$) with $m_0 = m_0^{(0)} = 0$. More general formulas using Bernoulli numbers can be seen in [14].

3 $x^2 + y^2 = z^3$

For the solution of the Diophantine equation $x^2 + y^2 = z^r$, we obtain the parameterization

$$x = \sum_{k=0}^{\lfloor r/2 \rfloor} (-1)^k \binom{r}{2k} s^{r-2k} t^{2k},$$

$$y = \sum_{k=0}^{\lfloor (r-1)/2 \rfloor} (-1)^k \binom{r}{2k+1} s^{r-2k-1} t^{2k+1},$$

$$z = s^2 + t^2,$$

where s and t are of opposite parity with $\gcd(s, t) = 1$.

The case where $r = 2$ has already been discussed in [12, 20]. Namely,

$$g_0(s^2 - t^2, 2st, s^2 + t^2) = (s - 1)(s^2 - t^2) + (s - 1)(2st) - (s^2 + t^2).$$

Let $r = 3$. The triple of the Diophantine equation $x^2 + y^2 = z^3$ is parameterized as

$$(x, y, z) = (s(s^2 - 3t^2), t(3s^2 - t^2), s^2 + t^2).$$

For convenience, we put

$$\mathbf{x} := s(s^2 - 3t^2), \quad \mathbf{y} := t(3s^2 - t^2), \quad \mathbf{z} := s^2 + t^2. \quad (4)$$

Since $\mathbf{x}, \mathbf{y}, \mathbf{z} > 0$ and $\gcd(\mathbf{x}, \mathbf{y}, \mathbf{z}) = 1$, we see that $s > \sqrt{3}t$, $\gcd(s, t) = 1$ and $s \not\equiv t \pmod{2}$.

When $\mathbf{x} > \mathbf{z}$, the Frobenius number of this triple is given as follows.


Theorem 1.

$$g_0(s(s^2 - 3t^2), t(3s^2 - t^2), s^2 + t^2) = \begin{cases} (s-1)s(s^2 - 3t^2) + (s-1)t(3s^2 - t^2) - (s^2 + t^2) \\ \quad \text{if } s > (1 + \sqrt{2})t; \\ (2s+t-1)s(s^2 - 3t^2) + (t-1)t(3s^2 - t^2) - (s^2 + t^2) \\ \quad \text{if } 2t < s < (1 + \sqrt{2})t; \\ (2s+t-1)s(s^2 - 3t^2) + (s-t-1)t(3s^2 - t^2) - (s^2 + t^2) \\ \quad \text{if } (2 + \sqrt{13})t/3 < s < 2t; \\ (5s+3t-1)s(s^2 - 3t^2) + (2t-s-1)t(3s^2 - t^2) - (s^2 + t^2) \\ \quad \text{if } 1.8139t < s < (2 + \sqrt{13})t/3; \\ (2s+t-1)s(s^2 - 3t^2) + (2s-3t-1)t(3s^2 - t^2) - (s^2 + t^2) \\ \quad \text{if } (3 + \sqrt{34})t/5 < s < 1.8139t; \\ (7s+4t-1)s(s^2 - 3t^2) + (2t-s-1)t(3s^2 - t^2) - (s^2 + t^2) \\ \quad \text{if } \sqrt{3}t < s < (3 + \sqrt{34})t/5. \end{cases}$$

Here, $1.8139\dots$ is the positive real root of $3x^4 - 7x^3 + 6x^2 - 3x - 5 = 0$.

Remark. When $\mathbf{x} < \mathbf{z}$, that is, $s\sqrt{(s-1)/(3s+1)} < t < s/\sqrt{3}$, there is no uniform pattern for the Frobenius number. We need a separate discussion for each case. See below for details.

3.1 The case where $\sqrt{3}t < s < (2 + \sqrt{3})t$



If $\sqrt{3}t < s < (2 + \sqrt{3})t$, then $0 < \mathbf{x} < \mathbf{y}$. Hence, $\mathbf{x} < \mathbf{z} < \mathbf{y}$ or $\mathbf{z} < \mathbf{x} < \mathbf{y}$.

First, let $\mathbf{z} < \mathbf{x} < \mathbf{y}$. Since $(2s+t)\mathbf{x} + (2t-s)\mathbf{y} = 2(s^2 - st - t^2)\mathbf{z}$ with $s^2 - st - t^2 > (2 - \sqrt{3})t^2$,

$$(2s+t)\mathbf{x} + (2t-s)\mathbf{y} \equiv \mathbf{z} \quad \text{and} \quad (2s+t)\mathbf{x} + (2t-s)\mathbf{y} > 0. \quad (5)$$

- The case where $(1 + \sqrt{2})t < s < (2 + \sqrt{3})t$

Then the elements of the (0-)Apéry set are given as in Table 1, where each point (X, Y) corresponds to the expression $X\mathbf{x} + Y\mathbf{y}$ and the area of the (0-)Apéry set is equal to $\mathbf{z} = s^2 + t^2$.

$(0,0)$	\cdots	$(s-1,0)$	$(s,0)$	\cdots	\cdots	$(s+t-1,0)$
\vdots		\vdots	\vdots			\vdots
$(0,t-1)$	\cdots	$(s-1,t-1)$	$(s,t-1)$	\cdots	\cdots	$(s+t-1,t-1)$
$(0,t)$	\cdots	$(s-1,t)$				
\vdots		\vdots				
\vdots		\vdots				
$(0,s-1)$	\cdots	$(s-1,s-1)$				

Table 1: $\text{Ap}_0(\mathbf{x}, \mathbf{y}, \mathbf{z})$ when $(1 + \sqrt{2})t < s < (2 + \sqrt{3})t$

Since $(1 + \sqrt{2})t < s < (2 + \sqrt{3})t$, we see that $(s + t)\mathbf{x} > (s - t)\mathbf{y}$. By $(s + t)\mathbf{x} \equiv (s - t)\mathbf{y} \pmod{\mathbf{z}}$ and by $s\mathbf{x} \equiv -t\mathbf{y} \pmod{\mathbf{z}}$ the sequence $\{\ell\mathbf{x} \pmod{\mathbf{z}}\}_{\ell=0}^{\mathbf{z}-1}$ is given by

$$\begin{aligned}
& (0,0), (1,0), \dots, (s+t-1,0), (0,s-t), (1,s-t), \dots, (s-1,s-t), \\
& (0,s-2t), (1,s-2t), \dots, (s+t-1,s-2t), \\
& (0,2s-3t), (1,2s-3t), \dots, (s-1,2s-3t), \\
& (0,2s-4t), (1,2s-4t), \dots, (s-1,2s-4t), \\
& (0,2s-5t), (1,2s-5t), \dots, (s+t-1,2s-5t) \\
& (0,3s-6t), (1,3s-6t), \dots, (s-1,3s-6t), \\
& (0,3s-7t), (1,3s-7t), \dots, (s-1,3s-7t) \\
& \dots, (s-1, st - (s-1)t). \tag{6}
\end{aligned}$$

Namely, after $(s-1, st - (s-1)t)$, the next point adding $\mathbf{x} \pmod{\mathbf{z}}$ is $(0,0)$. Note that the typical patterns in the sequence (6) are shown as follows: if

$$(0, k_1s - k_2t), (1, k_1s - k_2t), \dots, (s+t-1, k_1s - k_2t), (0, (k_1+1)s - (k_2+1)t)$$

and if $k_1s - k_2t \geq t$

$$(0, k_1s - k_2t), (1, k_1s - k_2t), \dots, (s-1, k_1s - k_2t), (0, k_1s - (k_2+1)t).$$

Since $\gcd(s, t) = 1$, all the points inside of the area in Table 1 appear in the sequence (6) just once. Since $\gcd(\mathbf{x}, \mathbf{z}) = 1$, the sequence $\{\ell\mathbf{x} \pmod{\mathbf{z}}\}_{\ell=0}^{\mathbf{z}-1}$ is equivalent to the sequence $\{\ell \pmod{\mathbf{z}}\}_{\ell=0}^{\mathbf{z}-1}$.

Comparing the elements at $(s+t-1, t-1)$ and $(s-1, s-1)$, taking possible maximal values, we find that the element at $(s-1, s-1)$ is the largest in the Apéry set because

$$(s-1)\mathbf{x} + (s-1)\mathbf{y} - ((s+t-1)\mathbf{x} + (t-1)\mathbf{y}) = t(s^2(2s-3) + (2s+t)t^2) > 0.$$

By Lemma 1 (1), we have

$$\begin{aligned} g(\mathbf{x}, \mathbf{y}, \mathbf{z}) &= (s-1)\mathbf{x} + (s-1)\mathbf{y} - \mathbf{z} \\ &= (s-1)(s-t)(s^2 + 4st + t^2) - (s^2 + t^2). \end{aligned}$$

- The case where $2t < s < (\sqrt{2} + 1)t$

Then $(s+t)\mathbf{x} \equiv (s-t)\mathbf{y} \pmod{\mathbf{z}}$ but $(s+t)\mathbf{x} < (s-t)\mathbf{y}$. Nevertheless, by $(2s+t)\mathbf{x} - (s-2t)\mathbf{y} = 2(s^2 - st - t^2)\mathbf{z} > 0$, we have $(2s+t)\mathbf{x} \equiv (s-2t)\mathbf{y} \pmod{\mathbf{z}}$ and $(2s+t)\mathbf{x} > (s-2t)\mathbf{y}$. For example, $(s, t) = (9, 4)$ satisfies this condition, so $(x, y, z) = (297, 908, 97)$. By a similar way, we know that all the elements of the (0-)Apéry set are given as in Table 2.

$(0, 0)$	\cdots	$(s-1, 0)$	$(s, 0)$	\cdots	\cdots	$(2s+t-1, 0)$
\vdots		\vdots	\vdots			\vdots
$(0, t-1)$	\cdots	$(s-1, t-1)$	$(s, t-1)$	\cdots	\cdots	$(2s+t-1, t-1)$
$(0, t)$	\cdots	$(s-1, t)$	(s, t)	\cdots	\cdots	$(2s+t-1, t)$
\vdots		\vdots	\vdots			\vdots
\vdots		\vdots	\vdots			\vdots
$(0, s-t-1)$	\cdots	$(s-1, s-t-1)$	$(s, s-t-1)$	\cdots	\cdots	$(2s+t-1, s-t-1)$

Table 2: $\text{Ap}_0(\mathbf{x}, \mathbf{y}, \mathbf{z})$ when $2t < s < (\sqrt{2} + 1)t$

Compare the elements at $(2s+t-1, t-1)$ and $(s-1, s-t-1)$, which take possible maximal values. Since the real roots of $-x^4 + 2x^3 - 3x^2 + 2x + 2 = 0$ are -0.4909 and 1.4909 , together with $s > 2t$, we see that

$$(s-1)\mathbf{x} + (s-t-1)\mathbf{y} - ((2s+t-1)\mathbf{x} + (t-1)\mathbf{y}) = -s^4 + 2s^3t - 3s^2t^2 + 2st^3 + 2t^4 < 0,$$

we find that the element at $(2s+t-1, t-1)$ is the largest in the Apéry set. By Lemma 1 (1), we have

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (2s+t-1)\mathbf{x} + (t-1)\mathbf{y} - \mathbf{z}.$$

- The case where $(2 + \sqrt{13})t/3 < s < 2t$

For example, $(s, t) = (27, 14)$ satisfies this condition, so $(x, y, z) = (3807, 27874, 925)$.

By $(2 + \sqrt{13})t/3 < s$, we have $(3s+2t)\mathbf{x} - (2s-3t)\mathbf{y} = (3s^2 - 4st - 3t^2)\mathbf{z} > 0$. So, $(3s+2t)\mathbf{x} \equiv (2s-3t)\mathbf{y} \pmod{\mathbf{z}}$ and $(3s+2t)\mathbf{x} > (2s-3t)\mathbf{y}$. Together with (5), all the elements of the (0-)Apéry set are given as in Table 3.

Compare the elements at $(3s+2t-1, 2t-s-1)$ and $(2s+t-1, s-t-1)$, which take the possible maximal values. We find that the element at

$(0, 0)$	\cdots	$(2s+t-1, 0)$	$(2s+t, 0)$	\cdots	\cdots	$(3s+2t-1, 0)$
\vdots		\vdots	\vdots			\vdots
$(0, 2t-s-1)$	\cdots	$(2s+t-1, 2t-s-1)$	$(2s+t, 2t-s-1)$	\cdots	\cdots	$(3s+2t-1, 2t-s-1)$
$(0, 2t-s)$	\cdots	$(2s+t-1, 2t-s)$				
\vdots		\vdots				
\vdots		\vdots				
$(0, s-t-1)$	\cdots	$(2s+t-1, s-t-1)$				

Table 3: $\text{Ap}_0(\mathbf{x}, \mathbf{y}, \mathbf{z})$ when $(2 + \sqrt{13})t/3 < s < 2t$

$(s-1, s-1)$ is the largest in the Apéry set because $1.8685 = (2 + \sqrt{13})t/3 < s < 2t$ and

$$\begin{aligned} & (2s+t-1)\mathbf{x} + (s-t-1)\mathbf{y} - ((3s+2t-1)\mathbf{x} + (2t-s-1)\mathbf{y}) \\ &= -(s^4 - 5s^3t + 6s^2t^2 - st^3 - 3t^4) > 0 \end{aligned}$$

for $-0.5268t < s < 3.3968t$. By Lemma 1 (1), we have

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (2s+t-1)\mathbf{x} + (s-t-1)\mathbf{y} - \mathbf{z}.$$

- The case where $(3 + \sqrt{34})t/5 < s < (2 + \sqrt{13})t/3$

For example, $(s, t) = (24, 13)$ satisfies this condition, so $(x, y, z) = (1656, 20267, 745)$.

In this case, $(3s+2t)\mathbf{x} \equiv (2s-3t)\mathbf{y} \pmod{\mathbf{z}}$ but $(3s+2t)\mathbf{x} < (2s-3t)\mathbf{y}$. Nevertheless, by $(3 + \sqrt{34})t/5 < s$ we have $(5s+3t)\mathbf{x} - (3s-5t)\mathbf{y} = (5s^2 - 6st - 5t^2)\mathbf{z} > 0$. So, $(5s+3t)\mathbf{x} \equiv (3s-5t)\mathbf{y} \pmod{\mathbf{z}}$ and $(5s+3t)\mathbf{x} > (3s-5t)\mathbf{y}$. Together with (5), all the elements of the (0-)Apéry set are given as in Table 4.

$(0, 0)$	\cdots	$(2s+t-1, 0)$	$(2s+t, 0)$	\cdots	\cdots	$(5s+3t-1, 0)$
\vdots		\vdots	\vdots			\vdots
$(0, 2t-s-1)$	\cdots	$(2s+t-1, 2t-s-1)$	$(2s+t, 2t-s-1)$	\cdots	\cdots	$(5s+3t-1, 2t-s-1)$
$(0, 2t-s)$	\cdots	$(2s+t-1, 2t-s)$				
\vdots		\vdots				
\vdots		\vdots				
$(0, 2s-3t-1)$	\cdots	$(2s+t-1, 2s-3t-1)$				

Table 4: $\text{Ap}_0(\mathbf{x}, \mathbf{y}, \mathbf{z})$ when $(3 + \sqrt{34})t/5 < s < (2 + \sqrt{13})t/3$

Comparing the elements at $(5s+3t-1, 2t-s-1)$ and $(2s+t-1, 2s-3t-1)$, taking possible maximal values, we find that there are two possibilities. By Lemma 1 (1), if

$$(2s+t-1)\mathbf{x} + (2s-3t-1)\mathbf{y} - ((5s+3t-1)\mathbf{x} + (2t-s-1)\mathbf{y})$$

$$= -3s^4 + 7s^3t - 6s^2t^2 + 3st^3 + 5t^4 > 0,$$

which is equivalent to $-0.5553t < s < 1.8139t$, namely if $1.7661t = (3 + \sqrt{34})t/5 < s < 1.8139t$, then

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (2s + t - 1)\mathbf{x} + (2s - 3t - 1)\mathbf{y} - \mathbf{z}.$$

Otherwise, namely, if $1.8139t < s < (2 + \sqrt{13})t/3 = 1.8685t$

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (5s + 3t - 1)\mathbf{x} + (2t - s - 1)\mathbf{y} - \mathbf{z}.$$

- The case where $\sqrt{3}t < s < (3 + \sqrt{34})t/5$

For example, $(s, t) = (44, 25)$ satisfies this condition, so $(x, y, z) = (2684, 129575, 2561)$.

In this case, $(5s + 3t)\mathbf{x} \equiv (3s - 5t)\mathbf{y} \pmod{\mathbf{z}}$ but $(5s + 3t)\mathbf{x} < (3s - 5t)\mathbf{y}$. Nevertheless, by $s > \sqrt{3}t = 1.732t > (4 + \sqrt{65})t/7 = 1.723t$, we have $(7s + 4t)\mathbf{x} - (4s - 7t)\mathbf{y} = (7s^2 - 8st - 7t^2)\mathbf{z} > 0$. So, $(7s + 4t)\mathbf{x} \equiv (4s - 7t)\mathbf{y} \pmod{\mathbf{z}}$ and $(7s + 4t)\mathbf{x} > (4s - 7t)\mathbf{y}$. Together with (5), all the elements of the (0-)Apéry set are given as in Table 5.

$(0, 0)$	\cdots	$(2s + t - 1, 0)$	$(2s + t, 0)$	\cdots	\cdots	$(7s + 4t - 1, 0)$
\vdots		\vdots	\vdots			\vdots
$(0, 2t - s - 1)$	\cdots	$(2s + t - 1, 2t - s - 1)$	$(2s + t, 2t - s - 1)$	\cdots	\cdots	$(7s + 4t - 1, 2t - s - 1)$
$(0, 2t - s)$	\cdots	$(2s + t - 1, 2t - s)$				
\vdots		\vdots				
\vdots		\vdots				
$(0, 3s - 5t - 1)$	\cdots	$(2s + t - 1, 3s - 5t - 1)$				

Table 5: $\text{Ap}_0(\mathbf{x}, \mathbf{y}, \mathbf{z})$ when $\sqrt{3}t < s < (3 + \sqrt{34})t/5$

Comparing the elements at $(7s + 4t - 1, 2t - s - 1)$ and $(2s + t - 1, 3s - 5t - 1)$, taking possible maximal values, we find that the element at $(7s + 4t - 1, 2t - s - 1)$ is the largest in the Apéry set because by $\sqrt{3}t < s < (3 + \sqrt{34})t/5$

$$\begin{aligned} & (2s + t - 1)\mathbf{x} + (3s - 5t - 1)\mathbf{y} - ((7s + 4t - 1)\mathbf{x} + (2t - s - 1)\mathbf{y}) \\ &= -5s^4 + 9s^3t - 6s^2t^2 + 5st^3 + 7t^4 < 0. \end{aligned}$$

Note that $-5x^4 + 9x^3 - 6x^2 + 5x + 7 = 0$ has the real roots at 0.5702 and 1.71692 and $s > \sqrt{3}t = 1.732t$. By Lemma 1 (1), we have

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (7s + 4t - 1)\mathbf{x} + (2t - s - 1)\mathbf{y} - \mathbf{z}.$$

3.2 The case where $\mathbf{x} < \mathbf{z} < \mathbf{y}$

If $\mathbf{x} < \mathbf{z} < \mathbf{y}$, then by $s(s^2 - 3t^2) < \mathbf{z}$ and $s > \sqrt{3}t$, we have

$$s\sqrt{\frac{s-1}{3s+1}} < t < \frac{s}{\sqrt{3}}.$$

As both s and t are positive integers, we can find

$$(s, t) = (2, 1), (7, 4), (9, 5), (11, 6), (14, 8), (16, 9), (21, 12), (23, 13), (26, 15), \\ (28, 16), (30, 17), (33, 19), (35, 20), (37, 21), (40, 23), \dots$$

With additional conditions $\gcd(s, t) = 1$ and $s \not\equiv t \pmod{2}$, the suitable pairs are given by

$$(s, t) = (2, 1), (7, 4), (11, 6), (16, 9), (26, 15), (30, 17), (40, 23), \dots$$

Notice that the area of the (0-)Apéry set is equal to $\mathbf{x} = s(s^2 - 3t^2)$.

In this case, there is no uniform pattern. We have to calculate the Frobenius number for each concrete case. For example, when $(s, t) = (2, 1), (7, 4), (26, 15)$, the area of the Apéry set is given as in Table 6, where each point (Z, Y) corresponds to the expression $Z\mathbf{z} + Y\mathbf{y}$ and the area of the (0-)Apéry set is equal to $\mathbf{x} = s(s^2 - 3t^2)$.

$$\boxed{(0, 0) \quad \cdots \quad (z_0, 0)}$$

Table 6: $\text{Ap}_0(\mathbf{x}, \mathbf{y}, \mathbf{z})$ when $(s, t) = (2, 1), (7, 4), (26, 15), \dots$

Here, when $(s, t) = (2, 1), (7, 4), (26, 15)$, $A = 2, 7, 26$, respectively. Then the Frobenius number is given by

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = z_0(s^2 + t^2) - s(s^2 - 3t^2).$$

When $(s, t) = (11, 6), (16, 9), (30, 17), (40, 23)$, the area of the Apéry set is given as in Table 7.

$$\boxed{\begin{array}{cccc} (0, 0) & \cdots & \cdots & \cdots \\ & \cdots & \cdots & \cdots (z_1, y_1) \\ & \cdots & \cdots & \\ & \cdots & (z_2, y_2) & \end{array}}$$

Table 7: $\text{Ap}_0(\mathbf{x}, \mathbf{y}, \mathbf{z})$ when $(s, t) = (11, 6), (16, 9), (30, 17), (40, 23), \dots$

Here,

$$\begin{aligned}
(s, t) = (11, 6) &\implies (z_1, y_1) = (37, 2), (z_2, y_2) = (28, 3); \\
(s, t) = (16, 9) &\implies (z_1, y_1) = (61, 2), (z_2, y_2) = (10, 4); \\
(s, t) = (30, 17) &\implies (z_1, y_1) = (201, 0), (z_2, y_2) = (196, 4); \\
(s, t) = (40, 23) &\implies (z_1, y_1) = (136, 2), (z_2, y_2) = (108, 3)
\end{aligned}$$

As seen, there seems to be no regularity. So, one has to find the values of (z_p, y_p) one by one.

When $(s, t) = (11, 6), (30, 17), (40, 23)$, the element at (z_1, y_1) is bigger than the element at (z_2, y_2) . Then the Frobenius number is given by

$$g(\mathbf{x}, \mathbf{y}, z) = z_1(s^2 + t^2) + y_1t(3s^2 - t^2) - s(s^2 - 3t^2).$$

When $(s, t) = (16, 9)$, the element at (z_2, y_2) is bigger than the element at (z_1, y_1) . Then the Frobenius number is given by

$$g(\mathbf{x}, \mathbf{y}, z) = z_2(s^2 + t^2) + y_2t(3s^2 - t^2) - s(s^2 - 3t^2).$$

3.3 The case $s > (2 + \sqrt{3})t$

If $s > (2 + \sqrt{3})t$, then $\mathbf{z} < \mathbf{y} < \mathbf{x}$. The elements of the (0-)Apéry set are given as in Table 8, where each point (Y, X) corresponds to the expression $Y\mathbf{y} + X\mathbf{x}$ and the area of the (0-)Apéry set is equal to $\mathbf{z} = s^2 + t^2$.

$(0, 0)$...	$(t-1, 0)$	$(t, 0)$	$(s-1, 0)$
\vdots		\vdots	\vdots			\vdots
$(0, s-1)$...	$(t-1, s-1)$	$(t, s-1)$	$(s-1, s-1)$
$(0, s)$...	$(t-1, s)$				
\vdots		\vdots				
\vdots		\vdots				
$(0, s+t-1)$...	$(t-1, s+t-1)$				

Table 8: $\text{Ap}_0(\mathbf{x}, \mathbf{y}, \mathbf{z})$ when $s > (2 + \sqrt{3})t$

By $s\mathbf{y} - t\mathbf{x} = st\mathbf{z}$, we get $s\mathbf{y} \equiv t\mathbf{x} \pmod{\mathbf{z}}$ and $s\mathbf{y} > t\mathbf{x}$. By using an additional relation $t\mathbf{y} + s\mathbf{x} = (s-t)(s+t)\mathbf{z}$, the sequence $\{\ell\mathbf{y} \pmod{\mathbf{z}}\}_{\ell=0}^{\mathbf{z}-1}$ is given by

$$\begin{aligned}
(Y, 0) & \quad (Y = 0, 1, \dots, s-1), (Y, t), \quad (Y = 0, 1, \dots, s-1), \\
(Y, 2t) & \quad (Y = 0, 1, \dots, s-1), \dots, \\
(Y, \lfloor s/t \rfloor t) & \quad (Y = 0, 1, \dots, s-1),
\end{aligned}$$

$$\begin{aligned}
& (Y, (\lfloor s/t \rfloor + 1)t) \quad (Y = 0, 1, \dots, t-1), \\
& (Y, (\lfloor s/t \rfloor + 1)t - s) \quad (Y = 0, 1, \dots, s-1), \\
& (Y, (\lfloor s/t \rfloor + 2)t - s) \quad (Y = 0, 1, \dots, s-1), \dots, \\
& (Y, (2 \lfloor s/t \rfloor + 1)t - s) \quad (Y = 0, 1, \dots, t-1), \\
& (Y, (2 \lfloor s/t \rfloor + 1)t - 2s) \quad (Y = 0, 1, \dots, s-1), \dots, \\
& (Y, (3 \lfloor s/t \rfloor + 1)t - 2s) \quad (Y = 0, 1, \dots, t-1), \\
& \dots, \\
& (Y, s) \quad (Y = 0, 1, \dots, t-1). \tag{7}
\end{aligned}$$

Note that $\gcd(s, t) = 1$. Namely, the next point of $(t-1, s)$ by adding $\mathbf{y} \pmod{\mathbf{z}}$ is $(0, 0)$. As $\gcd(\mathbf{y}, \mathbf{z}) = 1$, the sequence $\{\ell \mathbf{y} \pmod{\mathbf{z}}\}_{\ell=0}^{\mathbf{z}-1}$ matches the sequence $\{\ell \pmod{\mathbf{z}}\}_{\ell=0}^{\mathbf{z}-1}$.

Since $s > (2 + \sqrt{3})t$, we get $(s-1)\mathbf{y} + (s-1)\mathbf{x} > (t-1)\mathbf{y} + (s+t-1)\mathbf{x}$. Hence, by Lemma 1 (1), we have

$$g_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (s-1)\mathbf{y} + (s-1)\mathbf{x} - \mathbf{z}.$$

4 The case for $r = 4$

Let $r = 4$. The triple of the Diophantine equation $x^2 + y^2 = z^4$ is parameterized as

$$\begin{aligned}
(x, y, z) &= (s^4 - 6s^2t^2 + t^4, 4st(s^2 - t^2), s^2 + t^2) \\
&= ((s^2 + 2st - t^2)(s^2 - 2st - t^2), 4st(s-t)(s+t), s^2 + t^2).
\end{aligned}$$

For convenience, we put

$$\mathbf{x} := (s^2 + 2st - t^2)(s^2 - 2st - t^2), \quad \mathbf{y} := 4st(s-t)(s+t), \quad \mathbf{z} := s^2 + t^2. \tag{8}$$

Since $\mathbf{x}, \mathbf{y}, \mathbf{z} > 0$ and $\gcd(\mathbf{x}, \mathbf{y}, \mathbf{z}) = 1$, we see that $s > (1 + \sqrt{2})t$, $\gcd(s, t) = 1$ and $s \not\equiv t \pmod{2}$.

Then the Frobenius number of this triple is given as follows.

Theorem 2. *If $s^2 + t^2 < \min\{s^4 - 6s^2t^2 + t^4, 4st(s^2 - t^2)\}$, then*

$$g_0(s^4 - 6s^2t^2 + t^4, 4st(s^2 - t^2), s^2 + t^2)$$

$$= \begin{cases} (s-1)(s^4 - 6s^2t^2 + t^4) + 4st(s-1)(s^2 - t^2) - (s^2 + t^2) \\ \quad \text{if } s > (2 + \sqrt{3})t; \\ (2s + t - 1)(s^4 - 6s^2t^2 + t^4) + 4st(t-1)(s^2 - t^2) - (s^2 + t^2) \\ \quad \text{if } 2.8350t < s < (2 + \sqrt{3})t; \\ (s-1)(s^4 - 6s^2t^2 + t^4) + 4st(s-t-1)(s^2 - t^2) - (s^2 + t^2) \\ \quad \text{if } 2.5855t < s < 2.8350t; \\ (5s + 2t - 1)(s^4 - 6s^2t^2 + t^4) + 4st(3s + t - 1)(s^2 - t^2) - (s^2 + t^2) \\ \quad \text{if } (1 + \sqrt{2})t < s < 2.5855t. \end{cases}$$

Here, $2.8350\dots$ is the larger real root of $x^4 - 4x^3 + 6x^2 - 8x + 1 = 0$.
 $2.5855\dots$ is the largest real root of $2x^3 - 3x^2 - 6x + 1 = 0$.

4.1 The case where $s > \left(1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right)t$

If \mathbf{z} is not the least, for example, if $\mathbf{z} > \mathbf{x}$, then

$$\sqrt{\frac{1 + 6s^2 - \sqrt{1 + 16s^2 + 32s^4}}{2}} < t < \sqrt{\frac{1 + 6s^2 + \sqrt{1 + 16s^2 + 32s^4}}{2}}.$$

However, any integer pair (s, t) , satisfying

$$\sqrt{\frac{1 + 6s^2 - \sqrt{1 + 16s^2 + 32s^4}}{2}} < t < \frac{s}{1 + \sqrt{2}},$$

has not been found for $s < 10^7$. Even if it exists, the situation is too complicated. It is similar for the case $\mathbf{z} > \mathbf{y}$. Hence, we assume that $\mathbf{z} < \mathbf{y} < \mathbf{x}$ or $\mathbf{z} < \mathbf{x} < \mathbf{y}$.

Since $\mathbf{x} - \mathbf{y} = s^4 - 4s^3t - 6s^2t^2 + 4st^3 + t^4$ and the real roots of the equation $x^4 - 4x^3 - 6x^2 + 4x + 1 = 0$ are given by

$$\begin{aligned} 1 - \sqrt{2} - \sqrt{2(2 - \sqrt{2})} & (= -1.4966), & 1 - \sqrt{2} + \sqrt{2(2 - \sqrt{2})} & (= 0.6681), \\ 1 + \sqrt{2} - \sqrt{2(2 + \sqrt{2})} & (= -0.1989), & 1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})} & (= 5.0273), \end{aligned}$$

we have $\mathbf{x} > \mathbf{y}$ if $s > \left(1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right)t$; $\mathbf{x} < \mathbf{y}$ if $(1 + \sqrt{2})t < s < \left(1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right)t$.

- The case where $s > \left(1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right)t$

In this case, $\mathbf{z} < \mathbf{y} < \mathbf{x}$. Since $s\mathbf{y} - t\mathbf{x} = t(3s^2 - t^2)\mathbf{z} > 0$, we have $s\mathbf{y} \equiv t\mathbf{x} \pmod{\mathbf{z}}$ and $s\mathbf{y} > t\mathbf{x}$. Hence, the arrangement of the elements of the Apéry set is the same as in Table 8, where the position (Y, X) corresponds to the expression $Y\mathbf{y} + X\mathbf{x}$. By using an additional relation $t\mathbf{y} + s\mathbf{x} = s(s^2 - 3t^2)\mathbf{z}$, the sequence $\{\ell\mathbf{y} \pmod{\mathbf{z}}\}_{\ell=0}^{\mathbf{z}-1}$ is given as in (7). As $\gcd(\mathbf{y}, \mathbf{z}) = 1$, the sequence $\{\ell\mathbf{y} \pmod{\mathbf{z}}\}_{\ell=0}^{\mathbf{z}-1}$ matches the sequence $\{\ell \pmod{\mathbf{z}}\}_{\ell=0}^{\mathbf{z}-1}$.

Finally, compare the two candidates $(s-1, s-1)$ and $(t-1, s+t-1)$ to take the largest value. Since

$$(s-1)\mathbf{y} + (s-1)\mathbf{x} - ((t-1)\mathbf{y} + (s+t-1)\mathbf{x}) = t(s^3(3s-4t) + 2s^2t^2 + t^3(4s-t)) > 0,$$

the element at $(s-1, s-1)$ is the largest in the Apéry set. Therefore, by Lemma 1 (1), we have

$$g_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (s-1)\mathbf{y} + (s-1)\mathbf{x} - \mathbf{z}.$$

4.2 The case where $(1 + \sqrt{2})t < s < \left(1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right)t$

In this case, $\mathbf{z} < \mathbf{x} < \mathbf{y}$.

- The case where $(2 + \sqrt{3})t < s < \left(1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right)t$

Since $(s+t)\mathbf{x} - (s-t)\mathbf{y} = (s+t)(s^2 - 4st + t^2)\mathbf{z} > 0$, we have $(s+t)\mathbf{x} \equiv (s-t)\mathbf{y} \pmod{\mathbf{z}}$ and $(s+t)\mathbf{x} > (s-t)\mathbf{y}$. Notice that the real roots of $x^2 - 4x + 1 = 0$ are $2 - \sqrt{3}$ and $2 + \sqrt{3} = 3.7320$. Hence, the arrangement of the elements of the Apéry set is the same as in Table 1, where the position (X, Y) corresponds to the expression $X\mathbf{x} + Y\mathbf{y}$. By using an additional relation $s\mathbf{x} + t\mathbf{y} = s(s^2 - 3t^2)\mathbf{z}$, the sequence $\{\ell\mathbf{y} \pmod{\mathbf{z}}\}_{\ell=0}^{\mathbf{z}-1}$ is given as in (6). As $\gcd(\mathbf{y}, \mathbf{z}) = 1$, the sequence $\{\ell\mathbf{y} \pmod{\mathbf{z}}\}_{\ell=0}^{\mathbf{z}-1}$ matches the sequence $\{\ell \pmod{\mathbf{z}}\}_{\ell=0}^{\mathbf{z}-1}$. Since

$$(s-1)\mathbf{x} + (s-1)\mathbf{y} - ((s+t-1)\mathbf{x} + (t-1)\mathbf{y}) = t(s^3(3s-4t) + 2s^2t^2 + t^3(4s-t)) > 0,$$

the element at $(s-1, s-1)$ is the largest in the Apéry set. Therefore, we have

$$g_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (s-1)\mathbf{x} + (s-1)\mathbf{y} - \mathbf{z}.$$

- The case where $2.5855t < s < (2 + \sqrt{3})t$

In this case, $(s+t)\mathbf{x} \equiv (s-t)\mathbf{y} \pmod{\mathbf{z}}$ but $(s+t)\mathbf{x} < (s-t)\mathbf{y}$. Nevertheless,

by $(2s+t)\mathbf{x} - (s-2t)\mathbf{y} = (2s^3 - 3s^2t - 6st^2 + t^3)\mathbf{z} > 0$ we have $(2s+t)\mathbf{x} \equiv (s-2t)\mathbf{y} \pmod{\mathbf{z}}$ and $(2s+t)\mathbf{x} > (s-2t)\mathbf{y}$. Notice that the roots of the equation $2x^3 - 3x^2 - 6x + 1 = 0$ are -1.2413 , 0.1557 and 2.5855 . Since

$$(2s+t-1)\mathbf{x} + (t-1)\mathbf{y} - ((s-1)\mathbf{x} + (s-t-1)\mathbf{y}) = (s+t)(s^4 - 4s^3t + 6s^2t^2 - 8st^3 + t^4)$$

and the real roots of $x^4 - 4x^3 + 6x^2 - 8x + 1 = 0$ are 0.1380 and 2.8350 , there are two possibilities. See Table 2. If $2.8350t < s < (2+\sqrt{3})t$, the element at $(2s+t-1, t-1)$ is the largest in the Apéry set. If $2.5855t < s < 2.8350t$, the element at $(s-1, s-t-1)$ is the largest. Therefore, if $2.8350t < s < (2+\sqrt{3})t$, we have

$$g_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (2s+t-1)\mathbf{x} + (t-1)\mathbf{y} - \mathbf{z}.$$

If $2.5855t < s < 2.8350t$ (for example, $(s, t) = (14, 5), (17, 6), (18, 7), \dots$), we have

$$g_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (s-1)\mathbf{x} + (s-t-1)\mathbf{y} - \mathbf{z}.$$

- The case where $(1+\sqrt{2})t < s < 2.5855t$

In this case, $(2s+t)\mathbf{x} \equiv (s-2t)\mathbf{y} \pmod{\mathbf{z}}$ but $(2s+t)\mathbf{x} < (s-2t)\mathbf{y}$. Nevertheless, by $(5s+2t)\mathbf{x} - (2s-5t)\mathbf{y} = (5s^3 - 6s^2t - 15st^2 + 2t^3)\mathbf{z} > 0$ we have $(5s+2t)\mathbf{x} \equiv (2s-5t)\mathbf{y} \pmod{\mathbf{z}}$ and $(5s+2t)\mathbf{x} > (2s-5t)\mathbf{y}$.

Notice that the roots of $5x^3 - 6x^2 - 15x + 2 = 0$ are -1.3142 , 0.1275 and 2.3867 , and $1 + \sqrt{2} > 2.3867$

$(0, 0)$	\dots	$(3s+t-1, 0)$	$(3s+t, 0)$	\dots	\dots	$(5s+2t-1, 0)$
\vdots		\vdots	\vdots			\vdots
$(0, 3t-s-1)$	\dots	$(3s+t-1, 3t-s-1)$	$(3s+t, 3t-s-1)$	\dots	\dots	$(5s+2t-1, 3t-s-1)$
$(0, 3t-s)$	\dots	$(3s+t-1, 3t-s)$				
\vdots		\vdots				
\vdots		\vdots				
$(0, s-2t-1)$	\dots	$(3s+t-1, s-2t-1)$				

Table 9: $\text{Ap}_0(\mathbf{x}, \mathbf{y}, \mathbf{z})$ when $2.3867t < s < 2.5855t$

Since

$$\begin{aligned} & (5s+2t-1)\mathbf{x} + (3t-s-1)\mathbf{y} - ((3s+t-1)\mathbf{x} + (s-2t-1)\mathbf{y}) \\ &= (s+t)(s^4 - 4s^3t + 6s^2t^2 - 8st^3 + t^4) > 0, \end{aligned}$$

the element at $(5s+2t-1, s-1)$ is the largest in the Apéry set. Notice that the real roots of $2x^5 - 7x^4 + 8x^3 + 2x^2 - 18x + 1 = 0$ are -1.0842 , 0.0559 and 2.2938 . Therefore, by Lemma 1 (1), we have

$$g_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (5s+2t-1)\mathbf{y} + (3s+t-1)\mathbf{x} - \mathbf{z}.$$

5 The case for $r = 5$

Let $r = 5$. The triple of the Diophantine equation $x^2 + y^2 = z^5$ is parameterized as

$$(x, y, z) = (s(s^4 - 10s^2t^2 + 5t^4), t(5s^4 - 10s^2t^2 + t^4), s^2 + t^2). \quad (9)$$

For convenience, we put

$$\mathbf{x} := s(s^4 - 10s^2t^2 + 5t^4), \quad \mathbf{y} := t(5s^4 - 10s^2t^2 + t^4), \quad \mathbf{z} := s^2 + t^2.$$

Since $\mathbf{x}, \mathbf{y}, \mathbf{z} > 0$ and $\gcd(\mathbf{x}, \mathbf{y}, \mathbf{z}) = 1$, we see that $\gcd(s, t) = 1$, $s \not\equiv t \pmod{2}$, and

$$0 < s < \sqrt{\frac{5 - 2\sqrt{5}}{5}}t = 0.3249t \quad \text{or} \quad s > \sqrt{5 + 2\sqrt{5}}t = 3.0776t.$$

Notice that 0.3249 is the smaller positive root of $5x^4 - 10x^2 + 1 = 0$ and 3.0776 is the larger positive root of $x^4 - 10x^2 + 5 = 0$.

Then the Frobenius number of this triple is given as follows.

Theorem 3.

$$g_0(s(s^4 - 10s^2t^2 + 5t^4), t(5s^4 - 10s^2t^2 + t^4), s^2 + t^2)$$

$$= \left\{ \begin{array}{l}
(s-1)\mathbf{x} + (s-1)\mathbf{y} - \mathbf{z} \\
\quad \text{if } s > \left(1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right) t \\
\quad \text{or } \left(1 + \sqrt{5} - \sqrt{5 + 2\sqrt{5}}\right) t < s < \left(-1 - \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right) t; \\
(2s + t - 1)\mathbf{x} + (t - 1)\mathbf{y} - \mathbf{z} \\
\quad \text{if } 4.1894t < s < \left(1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right) t; \\
(s-1)\mathbf{x} + (s-t-1)\mathbf{y} - \mathbf{z} \\
\quad \text{if } \left(1 + \sqrt{5} + \sqrt{2(5 + \sqrt{5})}\right) t/2 < s < 4.1894t; \\
(s-1)\mathbf{x} + (s-2t-1)\mathbf{y} - \mathbf{z} \\
\quad \text{if } \left(1 + \sqrt{10} + \sqrt{2(10 + \sqrt{10})}\right) t/3 < s < 3.1098t, \\
(3s + t - 1)\mathbf{x} + (t - 1)\mathbf{y} - \mathbf{z} \\
\quad \text{if } 3.1098t < s < \left(1 + \sqrt{5} + \sqrt{2(5 + \sqrt{5})}\right) t/2, \\
(s^2 + t^2 - 1)\mathbf{x} - \mathbf{z} \\
\quad \text{if } \sqrt{5 + 2\sqrt{5}}t < s < \left(1 + \sqrt{10} + \sqrt{2(10 + \sqrt{10})}\right) t/3 \\
\quad \text{or } \left(-1 - \sqrt{5} + \sqrt{2(5 + \sqrt{5})}\right) t/2 < s < \sqrt{1 - 2/\sqrt{5}}t,
\end{array} \right.$$

$$g_0(s(s^4 - 10s^2t^2 + 5t^4), t(5s^4 - 10s^2t^2 + t^4), s^2 + t^2)$$

$$= \left\{ \begin{array}{l}
(s-1)\mathbf{x} + (s+2t-1)\mathbf{y} - \mathbf{z} \\
\quad \text{if } \left(-1 - \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right) t < s < 0.2386t, \\
(t-s-1)\mathbf{x} + (t-1)\mathbf{y} - \mathbf{z} \\
\quad \text{if } 0.2386t < s < \left(-1 - \sqrt{5} + \sqrt{2(5 + \sqrt{5})}\right) t/2, \\
(s-1)\mathbf{x} + (s+3t-1)\mathbf{y} - \mathbf{z} \\
\quad \text{if } \left(-1 - \sqrt{5} + \sqrt{2(5 + \sqrt{5})}\right) t/2 < s < 0.3215t, \\
(t-2s-1)\mathbf{x} + (t-1)\mathbf{y} - \mathbf{z} \\
\quad \text{if } 0.3215t < s < \left(-1 + \sqrt{10} + \sqrt{2(10 - \sqrt{10})}\right) t/3, \\
(t-1)\mathbf{x} + (t-1)\mathbf{y} - \mathbf{z} \\
\quad \text{if } 0 < s < \left(1 + \sqrt{5} - \sqrt{5 + 2\sqrt{5}}\right) t.
\end{array} \right.$$

Here, 4.1894... is the larger positive root of $x^6 - 4x^5 - 15x^2 + 4x + 2 = 0$.
3.1098 is the larger positive root of $2x^6 - 4x^5 - 5x^4 - 20x^2 + 4x + 3 = 0$.
0.2386 is the smaller positive root of $2x^6 + 4x^5 - 15x^4 - 4x + 1 = 0$. 0.3215
is the smaller positive root of $3x^6 + 4x^5 - 20x^4 - 5x^2 - 4x + 2 = 0$.

5.1 The case where $s > \left(1 + \sqrt{5} + \sqrt{5 + 2\sqrt{5}}\right) t$

We assume that $\mathbf{z} < \mathbf{y} < \mathbf{x}$ or $\mathbf{z} < \mathbf{x} < \mathbf{y}$. Since

$$\mathbf{x} - \mathbf{y} = s^5 - 5s^4t - 10s^3t^2 + 10s^2t^3 + 5st^4 - t^5$$

and the real roots of the equation $x^5 - 5x^4 - 10x^3 + 10x^2 + 5x - 1 = 0$ are given by

$$\begin{aligned} 1, \quad 1 - \sqrt{5} - \sqrt{5 - 2\sqrt{5}} &= -1.9626, \quad 1 - \sqrt{5} + \sqrt{5 - 2\sqrt{5}} = -0.5095, \\ 1 + \sqrt{5} - \sqrt{5 + 2\sqrt{5}} &= 0.1583, \quad 1 + \sqrt{5} + \sqrt{5 + 2\sqrt{5}} = 6.3137, \end{aligned}$$

we obtain that $\mathbf{z} < \mathbf{y} < \mathbf{x}$ if and only if

$$0.1583t = \left(1 + \sqrt{5} - \sqrt{5 + 2\sqrt{5}}\right) t < s < \sqrt{\frac{5 - 2\sqrt{5}}{5}} t = 0.3249t$$

or

$$s > \left(1 + \sqrt{5} + \sqrt{5 + 2\sqrt{5}}\right) t = 6.3137t;$$

$\mathbf{z} < \mathbf{x} < \mathbf{y}$ if and only if

$$0 < s < \left(1 + \sqrt{5} - \sqrt{5 + 2\sqrt{5}}\right) t = 0.1583t$$

or

$$3.0776t = \sqrt{5 + 2\sqrt{5}} t < s < \left(1 + \sqrt{5} + \sqrt{5 + 2\sqrt{5}}\right) t = 6.3137t.$$

- The case where $s > \left(1 + \sqrt{5} + \sqrt{5 + 2\sqrt{5}}\right) t$

In this case, $\mathbf{z} < \mathbf{y} < \mathbf{x}$. Since $s\mathbf{y} - t\mathbf{x} = st(s-t)(s+t)\mathbf{z} > 0$, we get $s\mathbf{y} \equiv t\mathbf{x} \pmod{\mathbf{z}}$ and $s\mathbf{y} > t\mathbf{x}$.

In addition, from $t\mathbf{y} + s\mathbf{x} = (s^2 + 2st - t^2)(s^2 - 2st - t^2)\mathbf{z} > 0$, we have $t\mathbf{y} \equiv (s + t - j)\mathbf{x} \pmod{\mathbf{z}}$ and $t\mathbf{y} > (s + t - j)\mathbf{x}$ ($j = 1, 2, \dots, t$). Hence, the arrangement of the elements of the Apéry set is the same as in Table 8,

where the position (Y, X) corresponds to the expression $Y\mathbf{y} + X\mathbf{x}$. Hence, the sequence $\{\ell\mathbf{y} \pmod{\mathbf{z}}\}_{\ell=0}^{\mathbf{z}-1}$ is given as in (7). As $\gcd(\mathbf{y}, \mathbf{z}) = 1$, the sequence $\{\ell\mathbf{y} \pmod{\mathbf{z}}\}_{\ell=0}^{\mathbf{z}-1}$ matches the sequence $\{\ell \pmod{\mathbf{z}}\}_{\ell=0}^{\mathbf{z}-1}$.

Finally, compare the two candidates $(s-1, s-1)$ and $(t-1, s+t-1)$ to take the largest value. Since

$$(s-1)\mathbf{y} + (s-1)\mathbf{x} - ((t-1)\mathbf{y} + (s+t-1)\mathbf{x}) = t(4s^5 - 5s^4t + 10s^2t^3 - 4st^4 - t^5) > 0,$$

the element at $(s-1, s-1)$ is the largest in the Apéry set. Notice that the largest real root of the equation $4x^5 - 5x^4 + 10x^2 - 4x - 1 = 0$ is 0.621267, which is less than $1 + \sqrt{5} + \sqrt{5 + 2\sqrt{5}}$. Therefore, by Lemma 1 (1), we have

$$g_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (s-1)\mathbf{y} + (s-1)\mathbf{x} - \mathbf{z}.$$

5.2 The case where $\sqrt{5 + 2\sqrt{5}}t < s < \left(1 + \sqrt{5} + \sqrt{5 + 2\sqrt{5}}\right)t$

In this case, $\mathbf{z} < \mathbf{x} < \mathbf{y}$.

- The case where $\left(1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right)t < s < \left(1 + \sqrt{5} + \sqrt{5 + 2\sqrt{5}}\right)t$

For example, $(s, t) = (21, 4)$ satisfies this case. Since $(s+t)\mathbf{x} - (s-t)\mathbf{y} = (s^4 - 4s^3t - 6s^2t^2 + 4st^3 + t^4)\mathbf{z}$ and the positive real roots of $x^4 - 4x^3 - 6x^2 + 4x + 1 = 0$ are

$$1 - \sqrt{2} + \sqrt{2(2 - \sqrt{2})} = 0.6681 \quad \text{and} \quad 1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})} = 5.02734,$$

if $\left(1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right)t < s < \left(1 + \sqrt{5} + \sqrt{5 + 2\sqrt{5}}\right)t$, then $(s+t)\mathbf{x} \equiv (s-t)\mathbf{y} \pmod{\mathbf{z}}$ and $(s+t)\mathbf{x} > (s-t)\mathbf{y}$. Then the area of the elements of the Apéry set is the same as in Table 2, where the position (X, Y) corresponds to the expression $X\mathbf{x} + Y\mathbf{y}$.

Compare the elements at $(s-1, s-1)$ and $(s+t-1, t-1)$. We see that

$$(s-1)\mathbf{x} + (s-1)\mathbf{y} - ((s+t-1)\mathbf{x} + (t-1)\mathbf{y}) = t(4s^5 - 5s^4t + 10s^2t^3 - 4st^4 - t^5) > 0$$

because only the positive real root of $4x^5 - 5x^4 + 10x^2 - 4x - 1 = 0$ is 0.6212, which is less than $1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})} = 5.0273$. Then the element at $(s-1, s-1)$ is the largest in the Apéry set, and we have

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (s-1)\mathbf{x} + (s-1)\mathbf{y} - \mathbf{z}.$$

- The case where $\left(1 + \sqrt{5} + \sqrt{2(5 + \sqrt{5})}\right)t/2 < s < \left(1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right)t$

If $3.5201t < s < 5.0273t$, then $(s+t)\mathbf{x} \equiv (s-t)\mathbf{y} \pmod{\mathbf{z}}$ but $(s+t)\mathbf{x} < (s-t)\mathbf{y}$. Nevertheless, $(2s+t)\mathbf{x} - (s-2t)\mathbf{y} = 2(s^4 - 2s^3t - 6s^2t^2 + 2st^3 + t^4) > 0$ because the real positive roots of $x^4 - 2x^3 - 6x^2 + 2x + 1 = 0$ are

$$\frac{1 - \sqrt{5} + \sqrt{2(5 - \sqrt{5})}}{2} = 0.5575 \quad \text{and} \quad \frac{1 + \sqrt{5} + \sqrt{2(5 + \sqrt{5})}}{2} = 3.5201.$$

Then, $(2s+t)\mathbf{x} \equiv (s-2t)\mathbf{y} \pmod{\mathbf{z}}$ and $(2s+t)\mathbf{x} > (s-2t)\mathbf{y}$. Hence, the arrangement of all the elements of the (0-)Apéry set are the same as in Table 2.

Compare the elements at $(2s+t-1, t-1)$ and $(s-1, s-t-1)$. Since $(2s+t-1)\mathbf{x} + (t-1)\mathbf{y} - ((s-1)\mathbf{x} + (s-t-1)\mathbf{y}) = s^6 - 4s^5t - 15s^2t^4 + 4st^5 + 2t^6$ and the positive real roots of $x^6 - 4x^5 - 15x^2 + 4x + 2 = 0$ are 0.5114 and 4.1894, if

$$4.1894t < s < \left(1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right)t = 6.3137t$$

(for example, $(s, t) = (15, 4)$), then the element at $(2s+t-1, t-1)$ is the largest in the Apéry set, and we have

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (2s+t-1)\mathbf{x} + (t-1)\mathbf{y} - \mathbf{z}.$$

If

$$3.0776t = \sqrt{5 + 2\sqrt{5}}t < s < 4.1894t$$

(for example, $(s, t) = (17, 4)$), then the element at $(2s+t-1, t-1)$ is the largest, and we have

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (s-1)\mathbf{x} + (s-t-1)\mathbf{y} - \mathbf{z}.$$

- The case where $\left(1 + \sqrt{10} + \sqrt{2(10 + \sqrt{10})}\right)t/3 < s < \left(1 + \sqrt{5} + \sqrt{2(5 + \sqrt{5})}\right)t/2$

If $3.0976t < s < 3.5201t$, then $(2s+t)\mathbf{x} \equiv (s-2t)\mathbf{y} \pmod{\mathbf{z}}$ but $(2s+t)\mathbf{x} < (s-2t)\mathbf{y}$. Nevertheless, $(3s+t)\mathbf{x} - (s-3t)\mathbf{y} = (3s^4 - 4s^3t - 18s^2t^2 + 4st^3 + 3t^4)\mathbf{z} > 0$ because the positive real roots of $3x^4 - 4x^3 - 18x^2 + 4x + 3 = 0$ are

$$\frac{1 - \sqrt{10} + \sqrt{2(10 - \sqrt{10})}}{3} = 0.5119$$

and

$$\frac{1 + \sqrt{10} + \sqrt{2(10 + \sqrt{10})}}{3} = 3.0976.$$

Hence, $(3s + t)\mathbf{x} \equiv (s - 3t)\mathbf{y} \pmod{\mathbf{z}}$ and $(3s + t)\mathbf{x} > (s - 3t)\mathbf{y}$.

$(0, 0)$	\dots	\dots	\dots
	\dots	\dots	$\dots (3s + t - 1, t - 1)$
	\dots	\dots	
\dots	$(s - 1, s - 2t - 1)$		

Table 10: $\text{Ap}_0(\mathbf{x}, \mathbf{y}, \mathbf{z})$ when $3.0976t < s < 3.5201t$

Comparing the possible elements (see Table 10),

$$\begin{aligned} (3s + t - 1)\mathbf{x} + (t - 1)\mathbf{y} - ((s - 1)\mathbf{x} + (s - 2t - 1)\mathbf{y}) \\ = 2s^6 - 4s^5t - 5s^4t^2 - 20s^2t^4 + 4st^5 + 3t^6. \end{aligned}$$

The positive real roots of $2x^6 - 4x^5 - 5x^4 - 20x^2 + 4x + 3 = 0$ are 0.4782 and 3.1098. Hence, if $3.0976t < s < 3.1098t$ (for example, $(s, t) = (31, 10), (87, 28)$), then $(s - 1, s - 2t - 1)$ is the largest and

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (s - 1)\mathbf{x} + (s - 2t - 1)\mathbf{y} - \mathbf{z}.$$

If $3.1098t < s < 3.5201t$ (for example, $(s, t) = (10, 3), (13, 4)$), then $(3s + t - 1, t - 1)$ is the largest and

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (3s + t - 1)\mathbf{x} + (t - 1)\mathbf{y} - \mathbf{z}.$$

- The case where $\sqrt{5 + 2\sqrt{5}}t < s < \left(1 + \sqrt{10} + \sqrt{2(10 + \sqrt{10})}\right)t/3$

If $3.0776t < s < 3.0976t$, then $(3s + t)\mathbf{x} \equiv (s - 3t)\mathbf{y} \pmod{\mathbf{z}}$ but $(3s + t)\mathbf{x} < (s - 3t)\mathbf{y}$. For example, $(34, 11), (37, 12)$ are in this case. The largest element in the Apéry set is $(s^2 + t^2 - 1, 0)$. The pattern is similar to that in Table 6. Note that $(4s + t)\mathbf{x} \equiv (s - 4t)\mathbf{y} \pmod{\mathbf{z}}$ but $s - 4t < 0$. Hence,

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (s^2 + t^2 - 1)\mathbf{x} - \mathbf{z}.$$

5.3 The case where $\left(1 + \sqrt{5} - \sqrt{5 + 2\sqrt{5}}\right)t < s < \sqrt{1 - 2/\sqrt{5}}t$

In this case, $\mathbf{x} > \mathbf{y}$. As $0.1583t < s < 0.3249t$, we get $(s, t) = (1, 4), (2, 7), (2, 9), (3, 10), (2, 11), (2, 13), (3, 14), (4, 15), (3, 16), (5, 16), (4, 17), (5, 18), (4, 19), \dots$

- The case where $\left(1 + \sqrt{5} - \sqrt{5 + 2\sqrt{5}}\right)t < s < \left(-1 - \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right)t$

If $0.1583t < s < 0.1989t$, then $(s, t) = (2, 11), (2, 13), (3, 16), \dots$. Since $(s+t)\mathbf{y} - (t-s)\mathbf{x} = (s^4 + 4s^3t - 6s^2t^2 - 4st^3 + t^4)\mathbf{z}$ and the positive real roots of $x^4 + 4x^3 - 6x^2 - 4x + 1 = 0$ are

$$-1 - \sqrt{2} + \sqrt{2(2 + \sqrt{2})} = 0.1989 \quad \text{and} \quad -1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})} = 1.4966,$$

we get $(s+t)\mathbf{y} \equiv (t-s)\mathbf{x} \pmod{\mathbf{z}}$ and $(s+t)\mathbf{y} > (t-s)\mathbf{x}$. Then the elements of the (0-)Apéry set are given as in Table 11, where each point (Y, X) corresponds to the expression $Y\mathbf{y} + X\mathbf{x}$. The situation is similar to that in Table 1.

(0, 0)
 (s + t - 1, s - 1)
	
	...	(t - 1, t - 1)	

Table 11: $\text{Ap}_0(\mathbf{x}, \mathbf{y}, \mathbf{z})$ when $0.1583t < s < 0.1989t$

We find that the element at $(t-1, t-1)$ is the largest in the Apéry set because

$$(t-1)\mathbf{y} + (t-1)\mathbf{x} - ((s+t-1)\mathbf{y} + (s-1)\mathbf{x}) = -s(s^5 + 4s^4t - 10s^3t^2 + 5st^4 - 4t^5) > 0.$$

Note that only the positive real root of $x^5 + 4x^4 - 10x^3 + 5x - 4 = 0$ is 1.6096, which is larger than 0.1989. Hence,

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (t-1)\mathbf{y} + (t-1)\mathbf{x} - \mathbf{z}.$$

- The case where $-1 - \sqrt{2} + \sqrt{2(2 + \sqrt{2})}t < s < \left(-1 - \sqrt{5} + \sqrt{2(5 + \sqrt{5})}\right)t/2$

If $0.1989t < s < 0.2840t$, then $(s, t) = (2, 9), (3, 14), (4, 15), (18, 5), \dots$. In this case, $(s+t)\mathbf{y} \equiv (t-s)\mathbf{x} \pmod{\mathbf{z}}$ but $(s+t)\mathbf{y} < (t-s)\mathbf{x}$. Nevertheless, by $(s+2t)\mathbf{y} - (t-2s)\mathbf{x} = 2(s^4 + 2s^3t - 6s^2t^2 - 2st^3 + t^4)\mathbf{z}$ and the positive real roots of $x^4 + 2x^3 - 6x^2 - 2x + 1 = 0$ are

$$\frac{-1 - \sqrt{5} + \sqrt{2(5 + \sqrt{5})}}{2} = 0.2840 \quad \text{and} \quad \frac{-1 + \sqrt{5} + \sqrt{2(5 - \sqrt{5})}}{2} = 1.7936,$$

we get $(s+2t)\mathbf{y} \equiv (t-2s)\mathbf{x} \pmod{\mathbf{z}}$ and $(s+2t)\mathbf{y} > (t-2s)\mathbf{x}$. Comparing the possible largest values of the Apéry set,

$$(t-s-1)\mathbf{y} + (t-1)\mathbf{x} - ((s+2t-1)\mathbf{y} + (s-1)\mathbf{x}) = -(2s^6 + 4s^5t - 15s^4t^2 - 4st^5 + t^6)$$

(0, 0)
 (s + 2t - 1, s - 1)
	
	...	(t - 1, t - s - 1)	

Table 12: $\text{Ap}_0(\mathbf{x}, \mathbf{y}, \mathbf{z})$ when $0.1989t < s < 0.2840t$

and the positive real roots of $2x^6 + 4x^5 - 15x^4 - 4x + 1 = 0$ are 0.2386 and 1.9552.

See Table 12. Hence, if $0.1989t < s < 0.2386t$ (for example, $(s, t) = (1, 4), (2, 9), (3, 14), \dots$), then $(s + 2t - 1, s - 1)$ takes the largest element and

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (s + 2t - 1)\mathbf{y} + (s - 1)\mathbf{x} - \mathbf{z}.$$

If $0.2386t < s < 0.2840t$ (for example, $(s, t) = (1, 4), (4, 15), (18, 5), \dots$), then $(t - 1, t - s - 1)$ takes the largest element and

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (t - 1)\mathbf{y} + (t - s - 1)\mathbf{x} - \mathbf{z}.$$

- The case where $\left(-1 - \sqrt{5} + \sqrt{2(5 + \sqrt{5})}\right) t/2 < s < \left(-1 + \sqrt{10} + \sqrt{2(10 - \sqrt{10})}\right) t/3$

If $0.2840t < s < 0.3228t$, then $(s, t) = (4, 13), (5, 16), (6, 19), (10, 31), \dots$. In this case, $(s + 2t)\mathbf{y} \equiv (t - 2s)\mathbf{xz}$ but $(s + 2t)\mathbf{y} < (t - 2s)\mathbf{x}$. Nevertheless, by $(s + 3t)\mathbf{y} - (t - 3s)\mathbf{x} = (3s^4 + 4s^3t - 18s^2t^2 - 4st^3 + 3t^4)\mathbf{z}$ and the positive real roots of $3x^4 + 4x^3 - 18x^2 - 4x + 3 = 0$ are

$$\frac{-1 - \sqrt{10} + \sqrt{2(10 + \sqrt{10})}}{3} = 1.9534$$

and

$$\frac{-1 + \sqrt{10} + \sqrt{2(10 - \sqrt{10})}}{3} = 0.3228,$$

we get $(s + 3t)\mathbf{y} \equiv (t - 3s)\mathbf{x} \pmod{\mathbf{z}}$ and $(s + 3t)\mathbf{y} > (t - 3s)\mathbf{x}$. Comparing the possible largest values of the Apéry set,

$$\begin{aligned} (t - 1)\mathbf{y} + (t - 2s - 1)\mathbf{x} - ((s + 3t - 1)\mathbf{y} + (s - 1)\mathbf{x}) \\ = -(3s^6 + 4s^5t - 20s^4t^2 - 5s^2t^4 - 4st^5 + 2t^6) \end{aligned}$$

and the positive real roots of $3x^6 + 4x^5 - 20x^4 - 5x^2 - 4x + 2 = 0$ are 0.3215 and 2.0907.

$(0, 0)$	\cdots	\cdots	\cdots
	\cdots	\cdots	$\cdots (s + 3t - 1, s - 1)$
	\cdots	\cdots	
	$\cdots (t - 1, t - 2s - 1)$		

Table 13: $\text{Ap}_0(\mathbf{x}, \mathbf{y}, \mathbf{z})$ when $0.2840t < s < 0.3228t$

See Table 13. Hence, if $0.2840t < s < 0.3215t$ (for example, $(s, t) = (4, 13), (5, 16), (6, 19)$), then $(s + 3t - 1, s - 1)$ takes the largest element and

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (s + 3t - 1)\mathbf{y} + (s - 1)\mathbf{x} - \mathbf{z}.$$

If $0.3215t < s < 0.3228t$ (for example, $(s, t) = (10, 31)$), then $(t - 1, t - 2s - 1)$ takes the largest element and

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (t - 1)\mathbf{y} + (t - 2s - 1)\mathbf{x} - \mathbf{z}.$$

- The case where $\left(-1 + \sqrt{10} + \sqrt{2(10 - \sqrt{10})}\right)t/3 < s < \sqrt{1 - 2/\sqrt{5}}t$

If $0.3228t < s < 0.3249t$, then $(s, t) = (11, 34), (12, 37), \dots$. In this case, $(s^2 + t^2 - 1, 0)$ takes the largest in the Apéry set (see Table 14) and

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (s^2 + t^2 - 1)\mathbf{y} - \mathbf{z}.$$

$(0, 0)$	\cdots	$(s^2 + t^2 - 1, 0)$
----------	----------	----------------------

Table 14: $\text{Ap}_0(\mathbf{x}, \mathbf{y}, \mathbf{z})$ when $0.3228t < s < 0.3249t$

5.4 The case where $0 < s < \left(1 + \sqrt{5} - \sqrt{5 + 2\sqrt{5}}\right)t$

In this case, $\mathbf{x} < \mathbf{y}$. Since $0 < s < 0.1538tt$, we can get $(s, t) = (1, 8 + 2j)$ ($j = 0, 1, \dots$), $(2, 15 + 2j)$ ($j = 0, 1, \dots$), $(3, 20), (3, 22), (3, 26), \dots$. Since $t\mathbf{x} - s\mathbf{y} = 4st(t - s)(s + t)\mathbf{z}$, we have $t\mathbf{x} \equiv s\mathbf{y} \pmod{\mathbf{z}}$ and $t\mathbf{x} > s\mathbf{y}$.

Then the elements of the (0-)Apéry set are given as in Table 15, where each point (X, Y) corresponds to the expression $X\mathbf{x} + Y\mathbf{y}$. The situation is similar to that in Table 15.

Comparing the possible elements, we have

$$(t - 1)\mathbf{x} + (t - 1)\mathbf{y} - ((s - 1)\mathbf{x} + (s + t - 1)\mathbf{y}) = -s(s^5 + 4s^4t - 10s^3t^2 + 5st^4 - 4t^5) > 0.$$

Notice that only the positive real root of $x^5 + 4x^4 - 10x^3 + 5x - 4 = 0$ is 1.6096. Then $(t - 1, t - 1)$ takes the largest element and

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (t - 1)\mathbf{x} + (t - 1)\mathbf{y} - \mathbf{z}.$$

(0, 0)
 (t - 1, t - 1)
	
	...	(s - 1, s + t - 1)	

Table 15: $\text{Ap}_0(\mathbf{x}, \mathbf{y}, \mathbf{z})$ when $0 < s < (1 + \sqrt{5} - \sqrt{5 + 2\sqrt{5}})t$

6 Sylvester number (genus)

6.1 The case where $r = 3$

Remember that for convenience, we use $(\mathbf{x}, \mathbf{y}, \mathbf{z})$ as in (4).

Theorem 4.

$$\begin{aligned}
& n_0(s(s^2 - 3t^2), t(3s^2 - t^2), s^2 + t^2) \\
&= \begin{cases} \frac{1}{2}((s-1)\mathbf{x} + (t-1)\mathbf{y} + 3st(s^2 - st - t^2) + (s^2 - 1)(t^2 - 1)) \\ \quad \text{if } s > (1 + \sqrt{2})t; \\ \frac{1}{2}((s-1)\mathbf{x} + (t-1)\mathbf{y} + st(5s^2 - 7st - 5t^2) + (s^2 - 1)(t^2 - 1)) \\ \quad \text{if } 2t < s < (1 + \sqrt{2})t; \\ \frac{1}{2}((s+1)\mathbf{x} - (5t-1)\mathbf{y} - st(11s^2 - 21st - 15t^2) + (s^2 - 1)(t^2 - 1)) \\ \quad \text{if } (2 + \sqrt{13})t/3 < s < 2t; \\ \frac{1}{2}((13s+1)\mathbf{x} - (17t-1)\mathbf{y} - st(45s^2 - 93st - 49t^2) + (s^2 - 1)(t^2 - 1)) \\ \quad \text{if } (3 + \sqrt{34})t/5 < s < (2 + \sqrt{13})t/3; \\ \frac{1}{2}((33s+1)\mathbf{x} - (37t-1)\mathbf{y} - st(99s^2 - 209st - 103t^2) + (s^2 - 1)(t^2 - 1)) \\ \quad \text{if } \sqrt{3}t < s < (3 + \sqrt{34})t/5. \end{cases}
\end{aligned}$$

- The case where $(1 + \sqrt{2})t < s < (2 + \sqrt{3})t$

By using Table 1, we have

$$\begin{aligned}
\sum_{w \in \text{Ap}(\mathbf{x}, \mathbf{y}, \mathbf{z})} w &= \sum_{X=0}^{s+t-1} \sum_{Y=0}^{t-1} (X\mathbf{x} + Y\mathbf{y}) + \sum_{X=0}^{s-1} \sum_{Y=t}^{s-1} (X\mathbf{x} + Y\mathbf{y}) \\
&= \frac{\mathbf{z}}{2}(-s^3 + s^4 - 3s^2t + 3s^3t + 3st^2 - 2s^2t^2 + t^3 - 3st^3 - t^4).
\end{aligned}$$

By Lemma 1 (2),

$$n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) = \frac{1}{\mathbf{z}} \sum_{w \in \text{Ap}(\mathbf{x}, \mathbf{y}, \mathbf{z})} w - \frac{\mathbf{z} - 1}{2}$$

$$= \frac{1}{2}((s-1)\mathbf{x} + (t-1)\mathbf{y} + 3st(s^2 - st - t^2) + (s^2 - 1)(t^2 - 1)).$$

- The case where $2t < s < (1 + \sqrt{2})t$

$$\begin{aligned} n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) \\ = \frac{1}{2}((s-1)\mathbf{x} + (t-1)\mathbf{y} + st(5s^2 - 7st - 5t^2) + (s^2 - 1)(t^2 - 1)). \end{aligned}$$

- The case where $(2 + \sqrt{13})t/3 < s < 2t$

$$\begin{aligned} n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) \\ = \frac{1}{2}((s+1)\mathbf{x} - (5t-1)\mathbf{y} - st(11s^2 - 21st - 15t^2) + (s^2 - 1)(t^2 - 1)). \end{aligned}$$

- The case where $(3 + \sqrt{34})t/5 < s < (2 + \sqrt{13})t/3$

$$\begin{aligned} n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) \\ = \frac{1}{2}((13s+1)\mathbf{x} - (17t-1)\mathbf{y} - st(45s^2 - 93st - 49t^2) + (s^2 - 1)(t^2 - 1)). \end{aligned}$$

- The case where $\sqrt{3}t < s < (3 + \sqrt{34})t/5$

$$\begin{aligned} n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) \\ = \frac{1}{2}((33s+1)\mathbf{x} - (37t-1)\mathbf{y} - st(99s^2 - 209st - 103t^2) + (s^2 - 1)(t^2 - 1)). \end{aligned}$$

6.1.1 The case where $\mathbf{x} < \mathbf{z} < \mathbf{y}$

If the Apéry set is arranged as in Table 6, then by

$$\sum_{w \in \text{Ap}_0(A)} w = \sum_{Z=0}^{z_0} Z\mathbf{z} = \frac{z_0(z_0+1)}{2}\mathbf{z},$$

where $z_0 + 1 = \mathbf{x}$, we have

$$n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) = \frac{1}{\mathbf{x}} \sum_{w \in \text{Ap}(\mathbf{x}, \mathbf{y}, \mathbf{z})} w - \frac{\mathbf{x}-1}{2}$$

$$= \frac{(\mathbf{x} - 1)(\mathbf{z} - 1)}{2}.$$

If the Apéry set is arranged as in Table 7, then by

$$\sum_{w \in \overline{\text{Ap}}_0(A)} w = \sum_{Z=0}^{z_1} \sum_{Y=0}^{y_1} (Z\mathbf{z} + Y\mathbf{y}) + \sum_{Z=0}^{z_2} \sum_{Y=y_1+1}^{y_2} (Z\mathbf{z} + Y\mathbf{y}),$$

we can calculate $n_0(\mathbf{x}, \mathbf{y}, \mathbf{z})$ individually.

6.1.2 The case $s > (2 + \sqrt{3})t$

By using Table 8, we have

$$\begin{aligned} \sum_{w \in \overline{\text{Ap}}(\mathbf{x}, \mathbf{y}, \mathbf{z})} w &= \sum_{Y=0}^{s-1} \sum_{X=0}^{s-1} (X\mathbf{x} + Y\mathbf{y}) + \sum_{Y=0}^{t-1} \sum_{X=s}^{s+t-1} (X\mathbf{x} + Y\mathbf{y}) \\ &= \frac{\mathbf{z}}{2}(-s^3 + s^4 - 3s^2t + 3s^3t + 3st^2 - 2s^2t^2 + t^3 - 3st^3 - t^4). \end{aligned}$$

By Lemma 1 (2),

$$\begin{aligned} n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) &= \frac{1}{\mathbf{z}} \sum_{w \in \overline{\text{Ap}}(\mathbf{x}, \mathbf{y}, \mathbf{z})} w - \frac{\mathbf{z} - 1}{2} \\ &= \frac{1}{2}((s-1)\mathbf{x} + (t-1)\mathbf{y} + 3st(s^2 - st - t^2) + (s^2 - 1)(t^2 - 1)). \end{aligned}$$

6.2 The case for $r = 4$

Remember that for convenience, we use $(\mathbf{x}, \mathbf{y}, \mathbf{z})$ as in (8).

Theorem 5. *If $s^2 + t^2 < \min\{s^4 - 6s^2t^2 + t^4, 4st(s^2 - t^2)\}$, then*

$$\begin{aligned} &n_0(s^4 - 6s^2t^2 + t^4, 4st(s^2 - t^2), s^2 + t^2) \\ &= \begin{cases} \frac{1}{2}((s+t-1)\mathbf{x} - \mathbf{y} + st(3(s^3 - t^3) + st(s-t-1)) \\ \quad + (s^2 - 1)(t^2 - 1)) & \text{if } s > (2 + \sqrt{3})t; \\ \frac{1}{2}((s+t-1)\mathbf{x} - \mathbf{y} + st(5s^3 - t^3 - st(5s+7t+1)) \\ \quad + (s^2 - 1)(t^2 - 1)) & \text{if } 2.5855t < s < (2 + \sqrt{3})t; \\ \frac{1}{2}((7s-7t+1)\mathbf{x} + \mathbf{y} + st(-35s^3 + 17t^3 + st(53s+65t-1)) \\ \quad + (s^2 - 1)(t^2 - 1)) & \text{if } (1 + \sqrt{2})t < s < 2.5855t. \end{cases} \end{aligned}$$

Here, $2.8350\dots$ is the larger real root of $x^4 - 4x^3 + 6x^2 - 8x + 1 = 0$.

6.2.1 The case where $s > \left(1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right)t$

By using Table 8, we have

$$\begin{aligned} \sum_{w \in \text{Ap}(\mathbf{x}, \mathbf{y}, \mathbf{z})} w &= \sum_{Y=0}^{s-1} \sum_{X=0}^{s-1} (Y\mathbf{y} + X\mathbf{x}) + \sum_{Y=0}^{t-1} \sum_{X=s}^{s+t-1} (Y\mathbf{y} + X\mathbf{x}) \\ &= \frac{\mathbf{z}}{2} (-s^4 + s^5 - 4s^3t + 4s^4t + 6s^2t^2 - 5s^3t^2 + 4st^3 - 7s^2t^3 - t^4 - 2st^4 + t^5). \end{aligned}$$

By Lemma 1 (2),

$$\begin{aligned} n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) &= \frac{1}{\mathbf{z}} \sum_{w \in \text{Ap}(\mathbf{x}, \mathbf{y}, \mathbf{z})} w - \frac{\mathbf{z} - 1}{2} \\ &= \frac{1}{2} ((s + t - 1)\mathbf{x} - \mathbf{y} + st(3(s^3 - t^3) + st(s - t - 1)) + (s^2 - 1)(t^2 - 1)). \end{aligned}$$

6.2.2 The case where $(1 + \sqrt{2})t < s < \left(1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right)t$

- The case where $(2 + \sqrt{3})t < s < \left(1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right)t$

By using Table 1, we have

$$\begin{aligned} \sum_{w \in \text{Ap}(\mathbf{x}, \mathbf{y}, \mathbf{z})} w &= \sum_{X=0}^{s+t-1} \sum_{Y=0}^{t-1} (X\mathbf{x} + Y\mathbf{y}) + \sum_{X=0}^{s-1} \sum_{Y=t}^{s-1} (X\mathbf{x} + Y\mathbf{y}) \\ &= \frac{\mathbf{z}}{2} ((-s^4 + s^5 - 4s^3t + 4s^4t + 6s^2t^2 - 5s^3t^2 + 4st^3 \\ &\quad - 7s^2t^3 - t^4 - 2st^4 + t^5)). \end{aligned}$$

Hence,

$$\begin{aligned} n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) &= \frac{1}{\mathbf{z}} \sum_{w \in \text{Ap}(\mathbf{x}, \mathbf{y}, \mathbf{z})} w - \frac{\mathbf{z} - 1}{2} \\ &= \frac{1}{2} ((s + t - 1)\mathbf{x} - \mathbf{y} + st(3(s^3 - t^3) + st(s - t - 1)) + (s^2 - 1)(t^2 - 1)). \end{aligned}$$

- The case where $2.5855t < s < (2 + \sqrt{3})t$

By Table 2, we have

$$n_0(\mathbf{x}, \mathbf{y}, \mathbf{z})$$

$$= \frac{1}{2}((s+t-1)\mathbf{x} - \mathbf{y} + st(5s^3 - t^3 - st(5s+7t+1)) + (s^2-1)(t^2-1)).$$

- The case where $(1 + \sqrt{2})t < s < 2.5855t$

By Table 9, we have

$$\begin{aligned} n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) &= \frac{1}{2}((7s-7t+1)\mathbf{x} + \mathbf{y} + st(-35s^3 + 17t^3 + st(53s+65t-1)) \\ &\quad + (s^2-1)(t^2-1)). \end{aligned}$$

6.3 The case for $r = 5$

Remember that for convenience, we use $(\mathbf{x}, \mathbf{y}, \mathbf{z})$ as in (9).

Theorem 6.

$$\begin{aligned} &n_0(s(s^4 - 10s^2t^2 + 5t^4), t(5s^4 - 10s^2t^2 + t^4), s^2 + t^2) \\ &= \left\{ \begin{array}{l} \frac{1}{2}((s-1)\mathbf{x} + (t-1)\mathbf{y} + st(5(s^4+t^4) - st(4s^2+14st-4t^2+1)) \\ \quad + (s^2-1)(t^2-1)) \quad \text{if } s > \left(1 + \sqrt{2} + \sqrt{2(2+\sqrt{2})}\right)t; \\ \frac{1}{2}((s-1)\mathbf{x} + (t-1)\mathbf{y} + st(7(s^4+t^4) - st(12s^2+26st-12t^2+1)) \\ \quad + (s^2-1)(t^2-1)) \\ \quad \text{if } \left(1 + \sqrt{5} + \sqrt{2(5+\sqrt{5})}\right)t/2 < s < \left(1 + \sqrt{2} + \sqrt{2(2+\sqrt{2})}\right)t; \\ \frac{1}{2}((s-1)\mathbf{x} + (t-1)\mathbf{y} + st(11(s^4+t^4) - st(20s^2+50st-20t^2+1)) \\ \quad + (s^2-1)(t^2-1)) \\ \quad \text{if } \left(1 + \sqrt{10} + \sqrt{2(10+\sqrt{10})}\right)t/3 < s < \left(1 + \sqrt{5} + \sqrt{2(5+\sqrt{5})}\right)t/2; \\ \frac{(x-1)(z-1)}{2} \\ \quad \text{if } \sqrt{5+2\sqrt{5}}t < s < \left(1 + \sqrt{10} + \sqrt{2(10+\sqrt{10})}\right)t/3; \\ \frac{1}{2}((s-1)\mathbf{x} + (t-1)\mathbf{y} + st(5(s^4+t^4) + st(4s^2-14st-4t^2-1)) \\ \quad + (s^2-1)(t^2-1)) \\ \quad \text{if } \left(1 + \sqrt{5} - \sqrt{5+2\sqrt{5}}\right)t < s < \left(-1 - \sqrt{2} + \sqrt{2(2+\sqrt{2})}\right)t; \end{array} \right. \end{aligned}$$

$$n_0(s(s^4 - 10s^2t^2 + 5t^4), t(5s^4 - 10s^2t^2 + t^4), s^2 + t^2)$$

$$= \left\{ \begin{array}{l} \frac{1}{2}((s-1)\mathbf{x} + (t-1)\mathbf{y} + st(7(s^4 + t^4) + st(12s^2 - 26st - 12t^2 - 1)) \\ \quad + (s^2 - 1)(t^2 - 1)) \\ \quad \text{if } -1 - \sqrt{2} + \sqrt{2(2 + \sqrt{2})}t < s < \left(-1 - \sqrt{5} + \sqrt{2(5 + \sqrt{5})}\right)t/2; \\ \frac{1}{2}((s-1)\mathbf{x} + (t-1)\mathbf{y} + st(11(s^4 + t^4) + st(20s^2 - 50st - 120t^2 - 1)) \\ \quad + (s^2 - 1)(t^2 - 1)) \\ \quad \text{if } \left(-1 - \sqrt{5} + \sqrt{2(5 + \sqrt{5})}\right)t/2 < s < \left(-1 + \sqrt{10} + \sqrt{2(10 - \sqrt{10})}\right)t/3; \\ \frac{(y-1)(z-1)}{2} \\ \quad \text{if } \left(-1 + \sqrt{10} + \sqrt{2(10 - \sqrt{10})}\right)t/3 < s < \sqrt{1 - 2/\sqrt{5}}t; \\ \frac{1}{2}((s-1)\mathbf{x} + (t-1)\mathbf{y} + st(5(s^4 + t^4) + st(4s^2 - 14st - 4t^2 + 1)) \\ \quad + (s^2 - 1)(t^2 - 1)) \quad \text{if } 0 < s < \left(1 + \sqrt{5} - \sqrt{5 + 2\sqrt{5}}\right)t. \end{array} \right.$$

6.3.1 The case where $s > \left(1 + \sqrt{5} + \sqrt{5 + 2\sqrt{5}}\right)t$

By using Table 8, we have

$$\begin{aligned} \sum_{w \in \text{Ap}(\mathbf{x}, \mathbf{y}, \mathbf{z})} w &= \sum_{Y=0}^{s-1} \sum_{X=0}^{s-1} (Y\mathbf{y} + X\mathbf{x}) + \sum_{Y=0}^{t-1} \sum_{X=s}^{s+t-1} (Y\mathbf{y} + X\mathbf{x}) \\ &= \frac{\mathbf{z}}{2}(-s^5 + s^6 - 5s^4t + 5s^5t + 10s^3t^2 - 9s^4t^2 + 10s^2t^3 \\ &\quad - 14s^3t^3 - 5st^4 - s^2t^4 - t^5 + 5st^5 + t^6). \end{aligned}$$

By Lemma 1 (2),

$$\begin{aligned} n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) &= \frac{1}{\mathbf{z}} \sum_{w \in \text{Ap}(\mathbf{x}, \mathbf{y}, \mathbf{z})} w - \frac{\mathbf{z} - 1}{2} \\ &= \frac{1}{2}((s-1)\mathbf{x} + (t-1)\mathbf{y} + st(5(s^4 + t^4) - st(4s^2 + 14st - 4t^2 + 1)) \\ &\quad + (s^2 - 1)(t^2 - 1)). \end{aligned}$$

6.3.2 The case where $\sqrt{5 + 2\sqrt{5}}t < s < \left(1 + \sqrt{5} + \sqrt{5 + 2\sqrt{5}}\right)t$

- The case where $\left(1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right)t < s < \left(1 + \sqrt{5} + \sqrt{5 + 2\sqrt{5}}\right)t$

By using Table 1, we have

$$\begin{aligned}
\sum_{w \in \text{Ap}(\mathbf{x}, \mathbf{y}, \mathbf{z})} w &= \sum_{X=0}^{s+t-1} \sum_{Y=0}^{t-1} (X\mathbf{x} + Y\mathbf{y}) + \sum_{X=0}^{s-1} \sum_{Y=t}^{s-1} (X\mathbf{x} + Y\mathbf{y}) \\
&= \frac{\mathbf{z}}{2} (-s^5 + s^6 - 5s^4t + 5s^5t + 10s^3t^2 - 9s^4t^2 + 10s^2t^3 \\
&\quad - 14s^3t^3 - 5st^4 - s^2t^4 - t^5 + 5st^5 + t^6).
\end{aligned}$$

By Lemma 1 (2),

$$\begin{aligned}
n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) &= \frac{1}{\mathbf{z}} \sum_{w \in \text{Ap}(\mathbf{x}, \mathbf{y}, \mathbf{z})} w - \frac{\mathbf{z} - 1}{2} \\
&= \frac{1}{2} ((s-1)\mathbf{x} + (t-1)\mathbf{y} + st(5(s^4 + t^4) - st(4s^2 + 14st - 4t^2 + 1)) \\
&\quad + (s^2 - 1)(t^2 - 1)).
\end{aligned}$$

- The case where $\left(1 + \sqrt{5} + \sqrt{2(5 + \sqrt{5})}\right) t/2 < s < \left(1 + \sqrt{2} + \sqrt{2(2 + \sqrt{2})}\right) t$
By using Table 2,

$$\begin{aligned}
n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) &= \frac{1}{2} ((s-1)\mathbf{x} + (t-1)\mathbf{y} + st(7(s^4 + t^4) - st(12s^2 + 26st - 12t^2 + 1)) \\
&\quad + (s^2 - 1)(t^2 - 1)).
\end{aligned}$$

- The case where $\left(1 + \sqrt{10} + \sqrt{2(10 + \sqrt{10})}\right) t/3 < s < \left(1 + \sqrt{5} + \sqrt{2(5 + \sqrt{5})}\right) t/2$
By using Table 10,

$$\begin{aligned}
n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) &= \frac{1}{2} ((s-1)\mathbf{x} + (t-1)\mathbf{y} + st(11(s^4 + t^4) - st(20s^2 + 50st - 20t^2 + 1)) \\
&\quad + (s^2 - 1)(t^2 - 1)).
\end{aligned}$$

- The case where $\sqrt{5 + 2\sqrt{5}} t < s < \left(1 + \sqrt{10} + \sqrt{2(10 + \sqrt{10})}\right) t/3$
By using Table 6,

$$n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) = \frac{(\mathbf{x} - 1)(\mathbf{z} - 1)}{2}.$$

6.3.3 The case where $(1 + \sqrt{5} - \sqrt{5 + 2\sqrt{5}})t < s < \sqrt{1 - 2/\sqrt{5}}t$

- The case where $(1 + \sqrt{5} - \sqrt{5 + 2\sqrt{5}})t < s < (-1 - \sqrt{2} + \sqrt{2(2 + \sqrt{2})})t$

By using Table 11, we have

$$\begin{aligned} \sum_{w \in \text{Ap}(\mathbf{x}, \mathbf{y}, \mathbf{z})} w &= \sum_{Y=0}^{s-1} \sum_{X=0}^{s-1} (Y\mathbf{y} + X\mathbf{x}) + \sum_{Y=0}^{t-1} \sum_{X=s}^{s+t-1} (Y\mathbf{y} + X\mathbf{x}) \\ &= \frac{\mathbf{z}}{2} (-s^5 + s^6 - 5s^4t + 5s^5t + 10s^3t^2 - s^4t^2 + 10s^2t^3 \\ &\quad - 14s^3t^3 - 5st^4 - 9s^2t^4 - t^5 + 5st^5 + t^6). \end{aligned}$$

By Lemma 1 (2),

$$\begin{aligned} n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) &= \frac{1}{\mathbf{z}} \sum_{w \in \text{Ap}(\mathbf{x}, \mathbf{y}, \mathbf{z})} w - \frac{\mathbf{z} - 1}{2} \\ &= \frac{1}{2} ((s-1)\mathbf{x} + (t-1)\mathbf{y} + st(5(s^4 + t^4) + st(4s^2 - 14st - 4t^2 - 1)) \\ &\quad + (s^2 - 1)(t^2 - 1)). \end{aligned}$$

- The case where $-1 - \sqrt{2} + \sqrt{2(2 + \sqrt{2})}t < s < (-1 - \sqrt{5} + \sqrt{2(5 + \sqrt{5})})t/2$

By using Table 12, we have

$$\begin{aligned} n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) &= \frac{1}{2} ((s-1)\mathbf{x} + (t-1)\mathbf{y} + st(7(s^4 + t^4) + st(12s^2 - 26st - 12t^2 - 1)) \\ &\quad + (s^2 - 1)(t^2 - 1)). \end{aligned}$$

- The case where $(-1 - \sqrt{5} + \sqrt{2(5 + \sqrt{5})})t/2 < s < (-1 + \sqrt{10} + \sqrt{2(10 - \sqrt{10})})t/3$

By using Table 13, we have

$$\begin{aligned} n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) &= \frac{1}{2} ((s-1)\mathbf{x} + (t-1)\mathbf{y} + st(11(s^4 + t^4) + st(20s^2 - 50st - 120t^2 - 1)) \\ &\quad + (s^2 - 1)(t^2 - 1)). \end{aligned}$$

- The case where $\left(-1 + \sqrt{10} + \sqrt{2(10 - \sqrt{10})}\right) t/3 < s < \sqrt{1 - 2/\sqrt{5}} t$
By using Table 14, we have

$$n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) = \frac{(\mathbf{y} - 1)(\mathbf{z} - 1)}{2}.$$

6.3.4 The case where $0 < s < \left(1 + \sqrt{5} - \sqrt{5 + 2\sqrt{5}}\right) t$

By using Table 15, we have

$$\begin{aligned} \sum_{w \in \text{Ap}(\mathbf{x}, \mathbf{y}, \mathbf{z})} w &= \sum_{X=0}^{s+t-1} \sum_{Y=0}^{t-1} (X\mathbf{x} + Y\mathbf{y}) + \sum_{X=0}^{s-1} \sum_{Y=t}^{s-1} (X\mathbf{x} + Y\mathbf{y}) \\ &= \frac{\mathbf{z}}{2} (-s^5 + s^6 - 5s^4t + 5s^5t + 10s^3t^2 - s^4t^2 + 10s^2t^3 \\ &\quad - 14s^3t^3 - 5st^4 - 9s^2t^4 - t^5 + 5st^5 + t^6). \end{aligned}$$

By Lemma 1 (2),

$$\begin{aligned} n_0(\mathbf{x}, \mathbf{y}, \mathbf{z}) &= \frac{1}{\mathbf{z}} \sum_{w \in \text{Ap}(\mathbf{x}, \mathbf{y}, \mathbf{z})} w - \frac{\mathbf{z} - 1}{2} \\ &= \frac{1}{2} ((s-1)\mathbf{x} + (t-1)\mathbf{y} + st(5(s^4 + t^4) + st(4s^2 - 14st - 4t^2 + 1)) \\ &\quad + (s^2 - 1)(t^2 - 1)). \end{aligned}$$

7 p -numerical semigroup

When $p > 0$, the formulas of p -Frobenius numbers and p -Sylvester number may be obtained though there are many different situations. For example, consider the case where $(1 + \sqrt{2})t < s < (2 + \sqrt{3})t$ on the Diophantine equation for $r = 3$. As $(s+t)\mathbf{x} \equiv (s-t)\mathbf{y} \pmod{\mathbf{z}}$, with respect to the area of the elements of $\text{Ap}_0(A)$ shown in Table 1 (taken as Range 1 (R1) in the longitudinal direction with the side-length $s+t$), the area of the elements of the first $(s-t)$ row moves in the form of filling the lower gap (modulo equal) as $0_1 \implies 1_1$. The portion protruding from the Range 1 is filled further downward from the left in order as $0_2 \implies 1_2$. The remaining regions except the first $(s-t)$ row of the elements of $\text{Ap}_0(A)$ are shifted (modulo equal) to the upper right side of Range 1 as Range 2 (R2) as $0_3 \implies 1_3$. It is as shown in Table 16. Note that $s-t > t$.

R1		R2
0 ₁	0 ₂	1 ₃ (D)
0 ₃	1 ₁	
	(C)	
	(B)	
1 ₂	(A)	

Table 16: $\text{Ap}_1(\mathbf{x}, \mathbf{y}, \mathbf{z})$ when $(1 + \sqrt{2})t < s < (2 + \sqrt{3})t$

Here, each point (X, Y) corresponds to the expression $X\mathbf{x} + Y\mathbf{y}$ and the area of the 1-Apéry set is also equal to $\mathbf{z} = s^2 + t^2$.

From Table 16, there are four candidates to take the largest value:

$$\begin{aligned} \textcircled{A} & : (s - t - 1, 2s - t - 1), & \textcircled{B} & : (s - 1, 2s + t - 1), \\ \textcircled{C} & : (s + t - 1, 2s - 1), & \textcircled{D} & : (2s + t - 1, t - 1). \end{aligned}$$

Since $s > (1 + \sqrt{2})t$, the value at \textcircled{A} is bigger than those at \textcircled{B} and \textcircled{C} . Because $(2s + t - 1)\mathbf{x} + (t - 1)\mathbf{y} - ((s - t - 1)\mathbf{x} + (2s - t - 1)\mathbf{y}) = s^4 - 4s^3t + 3s^2t^2 - 4st^3 - 2t^4$ and the only real root of $x^4 - 4x^3 + 3x^2 - 4x - 2 = 0$ is $3.5163\dots$. Therefore, if $2.4142t = (1 + \sqrt{2})t < s < 3.5163t$ then the value at \textcircled{A} is the largest and

$$g_1(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (s - t - 1)\mathbf{x} + (2s - t - 1)\mathbf{y} - \mathbf{z}.$$

If $3.5163t < s < (2 + \sqrt{3})t = 3.7320t$ (for example, $(s, t) = (18, 5), (29, 8)$) then the value at \textcircled{D} is the largest and

$$g_1(\mathbf{x}, \mathbf{y}, \mathbf{z}) = (2s + t - 1)\mathbf{x} + (t - 1)\mathbf{y} - \mathbf{z}.$$

For example, for $(s, t) = (18, 5)$, $g_1(\mathbf{x}, \mathbf{y}, \mathbf{z}) = 197871$. For $(s, t) = (29, 8)$, $g_1(\mathbf{x}, \mathbf{y}, \mathbf{z}) = 1360164$ as desired.

8 Final comments

When $r \geq 6$, we can also obtain the Frobenius numbers of the triple from $x^2 + y^2 = z^r$. However, we need to discuss the cases for each concrete value of r .

References

- [1] R. Apéry, *Sur les branches superlinéaires des courbes algébriques*, C. R. Acad. Sci. Paris **222** (1946), 1198–1200.
- [2] A. Brauer and B. M. Shockley, *On a problem of Frobenius*, J. Reine. Angew. Math. **211** (1962), 215–220.
- [3] T. C. Brown and P. J. Shiue, *A remark related to the Frobenius problem*, Fibonacci Quart. **31** (1993), 32–36.
- [4] A. Cayley, *On a problem of double partitions*, Philos. Mag. **XX** (1860), 337–341.
- [5] T. Chaichana, T. Komatsu and V. Laohakosol, *On convergents of certain values of hyperbolic functions formed from Diophantine equations*, Tokyo J. Math. **36** (2013), No. 1, 239–251.
- [6] F. Curtis, *On formulas for the Frobenius number of a numerical semigroup*, Math. Scand. **67** (1990), 190–192.
- [7] C. Elsner, T. Komatsu and I. Shiokawa, *On convergents formed by Pythagorean numbers*, Katsurada, Masanori (ed.) et al., Diophantine analysis and related fields 2006. In honor of Iekata Shiokawa. Proceedings of the conference, Keio University, Yokohama, Japan, March 7–10, 2006. Yokohama: Keio University, Department of Mathematics. Seminar on Mathematical Sciences 35, 59–76 (2006).
- [8] C. Elsner, T. Komatsu and I. Shiokawa, *Approximation of values of hypergeometric functions by restricted rationals*, J. Theor. Nombres Bordx. **19** (2007), No. 2, 393–404.
- [9] C. Elsner, T. Komatsu and I. Shiokawa, *On convergents formed from Diophantine equations*, Glas. Mat., III. Ser. **44** (2009), No. 2, 267–284.
- [10] E. Federico Elizeche and A. Tripathi, *On numerical semigroups generated by primitive Pythagorean triplets*, Integers **20** (2020), Paper No. A75, 12 pp.
- [11] L. G. Fel, *Frobenius problem for semigroups $S(d_1, d_2, d_3)$* , Funct. Anal. Other Math. **1** (2006), no. 2, 119–157.
- [12] B. K. Gil, J.-W. Han, T. H. Kim, R. H. Koo, B. W. Lee, J. Lee, K. S. Nam, H. W. Park and P.-S. Park, *Frobenius numbers of Pythagorean triples*, Int. J. Number Theory **11** (2015), no. 2, 613–619.

- [13] T. Komatsu, *On the number of solutions of the Diophantine equation of Frobenius—General case*, Math. Commun. **8** (2003), 195–206.
- [14] T. Komatsu, *Sylvester power and weighted sums on the Frobenius set in arithmetic progression*, Discrete Appl. Math. **315** (2022), 110–126.
- [15] T. Komatsu, *The Frobenius number for sequences of triangular numbers associated with number of solutions*, Ann. Comb. **26** (2022) 757–779.
- [16] T. Komatsu, *The Frobenius number associated with the number of representations for sequences of repunits*, C. R. Math., Acad. Sci. Paris **361** (2023), 73–89. <https://doi.org/10.5802/crmath.394>
- [17] T. Komatsu, *On the determination of p -Frobenius and related numbers using the p -Apéry set*, Rev. R. Acad. Cienc. Exactas Fis. Nat., Ser. A Mat., RACSAM **118** (2024), Article 58, 17 pages. DOI: 10.1007/s13398-024-01556-5
- [18] T. Komatsu and C. Pita-Ruiz, *The Frobenius number for Jacobsthal triples associated with number of solutions*, Axioms **12** (2023), no.2, Article 98, 18 pages. <https://doi.org/10.3390/axioms12020098>
- [19] T. Komatsu, S. Laishram and P. Punyani, *p -numerical semigroups of generalized Fibonacci triples*, Symmetry **15** (2023), no.4, Article 852, 13 pages. <https://doi.org/10.3390/sym15040852>
- [20] T. Komatsu and B. Sury, *p -numerical semigroups of Pythagorean triples*, preprint (2023).
- [21] T. Komatsu and H. Ying, *The p -Frobenius and p -Sylvester numbers for Fibonacci and Lucas triplets*, Math. Biosci. Eng. **20** (2023), No.2, 3455–3481. doi:10.3934/mbe.2023162
- [22] T. Komatsu and H. Ying, *The p -numerical semigroup of the triple of arithmetic progressions*, Symmetry **15** (2023), No.7, Article 1328, 17 p. <https://doi.org/10.3390/sym15071328>
- [23] T. Komatsu and H. Ying, *p -numerical semigroups with p -symmetric properties*, J. Algebra Appl. (online ready). <https://doi.org/10.1142/S0219498824502165>
- [24] T. Komatsu and Y. Zhang, *Weighted Sylvester sums on the Frobenius set*, Irish Math. Soc. Bull. **87** (2021), 21–29.

- [25] T. Komatsu and Y. Zhang, *Weighted Sylvester sums on the Frobenius set in more variables*, Kyushu J. Math. **76** (2022), 163–175.
- [26] A. M. Robles-Pérez and J. C. Rosales, *The Frobenius number for sequences of triangular and tetrahedral numbers*, J. Number Theory **186** (2018), 473–492.
- [27] Ø. J. Rødseth, *A note on Brown and Shiue’s paper on a remark related to the Frobenius problem*, Fibonacci Quart. **32** (1994), 407–408.
- [28] J. C. Rosales and P. A. Garcia-Sanchez, *Numerical semigroups with embedding dimension three*, Arch. Math. (Basel) **83** (2004), no. 6, 488–496.
- [29] J. C. Rosales, M. B. Branco and D. Torráo, *The Frobenius problem for Thabit numerical semigroups*, J. Number Theory **155** (2015), 85–99.
- [30] J. C. Rosales, M. B. Branco and D. Torráo, *The Frobenius problem for Mersenne numerical semigroups*, Math. Z. **286** (2017), 741–749.
- [31] E. S. Selmer, *On the linear diophantine problem of Frobenius*, J. Reine Angew. Math. **293/294** (1977), 1–17.
- [32] J. J. Sylvester, *On the partition of numbers*, Quart. J. Pure Appl. Math. **1** (1857), 141–152.
- [33] J. J. Sylvester, *On subinvariants, i.e. semi-invariants to binary quantics of an unlimited order*, Am. J. Math. **5** (1882), 119–136.
- [34] J. J. Sylvester, *Mathematical questions with their solutions*, Educational Times **41** (1884), 21.
- [35] J. J. Tattersall, *Elementary number theory in nine chapters*, Second edition. Cambridge University Press, Cambridge, 2005.
- [36] A. Tripathi, *The number of solutions to $ax + by = n$* , Fibonacci Quart. **38** (2000), 290–293.
- [37] A. Tripathi, *On sums of positive integers that are not of the form $ax + by$* , Amer. Math. Monthly **115** (2008), 363–364.