Vortices and Factorization

Igor Loutsenko and Oksana Yermolayeva

Laboratoire de Physique Mathématique, CRM, Université de Montréal

Abstract

We review applications of factorization methods to the problem of finding stationary point vortex patterns in two-dimensional fluid mechanics. Then we present a new class of patterns related to periodic analogs of Schrodinger operators from the "even" bi-spectral family. We also show that patterns related to rational and soliton solutions of the KdV hierarchy constitute complete solution of the problem for certain classes of vortex systems.

1 Introduction, Stationary Vortex Configurations

In fluid mechanics, factorization methods found their applications in the theory of two-dimensional flows in two opposite limits: the infinite-viscosity limit of the free-boundary flows (see e.g. [24], [25] and references therein) and in the zero-viscosity limit. Here we review the latter, namely applications to the problem of finding classes of two-dimensional stationary vortex configurations in inviscid 2d flows.

Systems of point vortices are weak solutions of the two-dimensional Euler equation. Work on their classification started more than a century and a half ago and has been developed substantially over the past several decades (for general overview see e.g. [3, 4, 14, 28, 29] and references therein).

Motion of two-dimensional inviscid and incompressible liquid is described by the system of PDEs

$$\frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v}\nabla)\mathbf{v} = -\frac{\nabla P}{\rho}, \quad \rho = \text{const}, \tag{1}$$

$$\nabla \mathbf{v} = 0, \tag{2}$$

where $\mathbf{v} = \mathbf{v}(x, y, t)$, $\mathbf{v} = (u, w)$ is the two-dimensional velocity of the flow, and P = P(x, y, t)stands for the pressure. Equation (1) is the Euler's equation, while (2) is the continuity equation. In two dimensions, the vorticity, which is a curl of velocity, has a single component ω :

$$\omega = \frac{\partial w}{\partial x} - \frac{\partial u}{\partial y}.$$

Taking curl of (1) we get

$$\frac{d\omega}{dt} = \frac{\partial\omega}{\partial t} + (v\nabla)\omega = 0.$$

In other words, the full derivative of the vorticity along the flow is zero. Then it follows that the circulation $\int_{\Omega(t)} \omega dx dy = \oint_{\partial\Omega(t)} u dx + w dy$ along the boundary of any evolving with the flow domain $\Omega(t)$ does not change with time. As a consequence for the system of point vortices

$$\omega = \sum_{i=1}^{N} \mathcal{Q}_i \delta(x - x_i(t)) \delta(y - y_i(t))$$

the strength (i.e. circulation) of each vortex Q_i is time independent and speed of *i*th vortex $(dx_i/dt, dy_i/dt)$ equals velocity of the flow averaged over a small circular domain centered at (x_i, y_i) . Since vorticity is a curl of velocity, for the system of point vortices we have

$$u - iw = U - iW + \frac{1}{2\pi i} \sum_{i=1}^{N} \frac{Q_i}{z - z_i(t)},$$
(3)

where we used complex notations z = x + iy. In (3) the full velocity field $\mathbf{v} = (u, w)$ is a sum of irrotational (i.e. zero-curl) background flow $\mathbf{V} = (U, W)$ and the flow created by superposition of N point vortices. By (2, 3) \mathbf{V} is a zero-divergence field. Since it is irrotational, it is also a gradient of a potential and as a consequence the potential is a harmonic function. In sections 1-8 we will consider unrestricted flows in the whole plane with velocities bounded at infinity. The only harmonic functions satisfying above conditions are linear functions and the background flow has constant velocity (uniform background flow). Using (3), we find the average velocity of the flow in a small disc centered at $z = z_i$, thus, obtaining the following equations of motion of vortices in the uniform background flow

$$2\pi i \frac{d\bar{z}_i}{dt} = k + \sum_{j=1, j \neq i}^N \frac{\mathcal{Q}_j}{z_i - z_j}, \quad i = 1 \dots N,$$
(4)

where k is a complex constant.

Classification of the stationary patterns of vortices is then reduced to the study of solution of system of N algebraic equations

$$k + \sum_{j=1, j \neq i}^{N} \frac{\mathcal{Q}_j}{z_i - z_j} = 0, \quad i = 1 \dots N.$$
(5)

This system also has the 2d electrostatic interpretation: (5) is nothing, but an equilibrium condition of N point electric charges Q_1, \ldots, Q_N interacting pair-wise through 2d Coulomb (logarithmic) potential and placed in the homogeneous electric field \bar{k} . The equilibrium configuration is a critical point of the electrostatic energy

$$E = k \sum_{i} \mathcal{Q}_{i} z_{i} + \sum_{i < j} \mathcal{Q}_{j} \mathcal{Q}_{j} \log(z_{i} - z_{j}) + \text{c.c.},$$

where "c.c." stands for the complex conjugate. Note that equilibrium configurations are special in the sense that system (5) does not have solution for generic parameters k and Q_i s. For instance, two-vortex (N = 2) equilibrium exists only if $k \neq 0$ and $Q_1 = -Q_2$. The stationary patterns with $k \neq 0$ are called translating equilibria (or translating configurations/patterns) because in the reference frame where the fluid rests at infinity, the whole pattern translates with constant velocity (Im k, Re k)/(2π). Below we mainly deal with equilibria without background flow, i.e. k = 0 equilibrium configurations ("static configurations/patterns"):

$$\sum_{j=1,j\neq i}^{N} \frac{\mathcal{Q}_j}{z_i - z_j} = 0, \quad i = 1 \dots N.$$
(6)

Equilibrium patterns in non-uniform background flows will be considered in section 9. Sections 10 and 11 are devoted to periodic configurations (vortex streets).

2 Vortices, Polynomials and Rational Primitives

Let us consider static equilibria of N = l + m charges (or vortices) of two species. Without lost of generality we can set values of first l charges to -1 and those of remaining m charges to Λ :

$$Q_i = \begin{cases} -1, & i = 1 \dots l \\ \Lambda, & i = l + 1 \dots l + m \end{cases}$$

Introducing polynomials in z

$$p(z) = \prod_{i=1}^{l} (z - z_i), \quad q(z) = \prod_{i=1}^{m} (z - z_{l+i}), \tag{7}$$

we first consider a situation where all charges have distinct positions, so p and q do not have multiple or common roots. One can check that the static equilibrium condition (6) for this system of two species is equivalent to the following bi-linear differential equation for the above polynomials

$$p''q - 2\Lambda p'q' + \Lambda^2 pq'' = 0, (8)$$

where prime denotes differentiation wrt z. The equivalence can be established by the partial fraction decomposition of the ratio of (8) and pq.

Note, that (8) holds not only when p and q do not have common or multiple roots, but also in special cases when such roots exist. It trivially follows from the partial fraction decomposition of the ratio of (8) and pq that if a common root exists, then it cannot be the simple one in both p and q: Multiplicities M and M' of this root, in p and q respectively, are constrained by the following relation:

$$(M - \Lambda M')^2 = M + \Lambda^2 M'.$$

In situation when a common root emerges as a limit of a generic solution, when M vortices of strength -1 and M' vortices of strength Λ collapse into a single vortex of strength $\Lambda M' - M$, the limiting vortex patterns are called the collapse configurations. There are also exist common root solutions that are not collapse solutions. For example, Demina and Kudryashov [18] presented a family of such solutions using the fact that if a pair p, q solves (8) for some Λ , then another pair $pq^{\Lambda(\Lambda-1)/2}$, $q^{\Lambda(\Lambda+1)/2}$, that corresponds to the same vortex pattern as the first one, satisfies

$$p''q - 2p'q' + pq'' = 0. (9)$$

The above equation, which is the $\Lambda = 1$ specification of (8), is called the Tkachenko equation [35] in literature on the vortex patterns¹. Its generic (i.e. no-common root) solutions describe equilibria of two species of vortices (charges) of equal magnitude but opposite sign. Bartman [9] identified the Tkachenko equation with the recurrence relation (16) for the Adler-Moser polynomials. As follows from works by Burchnall and Chaundy [7] (see the next Section) any generic polynomial solution of the Tkachenko equation is a pair of the Adler-Moser polynomials [1]

$$p(z) = P_{n-1}(z), \quad q(z) = P_n(z).$$
 (10)

In this way we get complete classification of static configuration of system of vortices (charges) of strengths ± 1 . The degree of the *n*th Adler-Moser polynomial equals n(n + 1)/2. Therefore, for this system equilibrium is only possible if the number of negative and positive charges equals l = n(n-1)/2 and m = n(n+1)/2 respectively. The equilibrium coordinates of charges z_i of each specie are roots of consecutive Adler-Moser polynomials. The *n*th Adler-Moser polynomial P_n is also a function of *n* free parameters, and the space of all possible equilibria of $N = l + m = n^2$ charges is a *n*-dimensional complex subspace of the n^2 dimensional complex space.

Details on Adler-Moser polynomials will be given in the following sections. Here, we note that these polynomials were first found by Burchnall and Chaundy [7], later re-discovered in context of the Hadamard problem [21] by Lagnese and Stellmacher, and then by Adler and Moser as polynomial τ -functions corresponding to rational solutions of the KdV hierarchy of integrable PDEs [1]. Burchnall and Chaundy [7] built these polynomials as solutions of the Tkachenko equation (9) by considering it as a linear second order ODE for p, where coefficients of ODE are determined by some polynomial $P_n = q$. Then the requirement that its two linearly independent solutions $p = P_{n-1}$ and $p = P_{n+1}$ are both polynomial is equivalent to rationality in z of two indefinite integrals (rational primitives)

$$\int \frac{p^2(z)}{q^2(z)} dz, \quad \int \frac{q^2(z)}{p^2(z)} dz$$

It turned out that, apart from the Adler-Moser case there is another case with $\Lambda = 2$ when there exists an infinite chain of polynomial solutions (8) that can be built with help of rational primitives [9, 22].

3 $\Lambda = 1$ and $\Lambda = 2$ Chains: Main Sequences

Let us return to the generic Λ case, and consider the bi-linear equation (8) as a second order linear ODE for $p = p_{n-1}(z)$ with coefficients of the equation being defined by $q = q_n(z)$. By elementary methods we find that the second linearly independent solution $p = p_n(z)$ equals

$$p_n(z) = C p_{n-1}(z) \int \frac{q_n(z)^{2\Lambda}}{p_{n-1}(z)^2} dz$$

In order to build a recursive chain we require that both p_{n-1} and p_n be polynomial in z, so the primitive

$$\int \frac{q^{2\Lambda}}{p^2} dz \tag{11}$$

¹Eq. (9) first appeared in work of Burchnall and Chaundy [7] in different context.

must be rational. Alternatively, considering (8) as an ODE for q, we come to the requirement of rationality of

$$\int \frac{p^{2/\Lambda}}{q^2} dz.$$
(12)

It is clear that pair of the above rationality conditions requires that both 2Λ and $2/\Lambda$ be integer, which is possible only if

$$\Lambda \in \left\{\frac{1}{2}, 1, 2\right\}.$$
(13)

The case $\Lambda = 1$ has already been mentioned and corresponds to the Adler-Moser polynomials. Since equation (8) is invariant under the involution $\Lambda \leftrightarrow 1/\Lambda$, $p \leftrightarrow q$, the cases $\Lambda = 1/2$ and $\Lambda = 2$ are equivalent, and, without loss of generality, both can be represented by the $\Lambda = 2$ case.

Now, let us take p and q that do not have common/multiple roots and factorize p(z) as $p(z) = (z - z_i)\tilde{p}(z)$, where $z_i, i \leq l$ is a root of p (see definition (7)). Rationality of primitive (11) implies an absence of a simple pole at $z = z_i$, which is equivalent to the condition

$$\frac{\partial}{\partial z} \left(\frac{q(z)^{2\Lambda}}{\tilde{p}(z)^2} \right)_{z=z_i} = 0.$$

Since $\tilde{p}(z_i) = p'(z_i)$ and $\tilde{p}'(z_i) = p''(z_i)/2$, we get the following equation

$$p''(z_i)q(z_i) - 2\Lambda p'(z_i)q'(z_i) = 0$$

which is clearly satisfied by solutions of (8). Similarly, rationality of (12) also follows from (8). It is worth mentioning that the converse statement also holds, i.e. the bilinear equation (8) follows from rationality of (11) and (12) (for details see [22]).

Thus, once a pair of polynomials p, q without multiple and common roots satisfying (8,13) is found, one can build an infinite chain of polynomial solutions of (8) with the help of recursive procedure for p_n and q_n mentioned at the beginning of this section.

In more detail: Since $p = z^{l} + \ldots$ and $q = z^{m} + \ldots$ from the highest symbol of (8) we get the diophantine equation connecting l and m

• $\Lambda = 1$ case: $(l - m)^2 = l + m$,

•
$$\Lambda = 2$$
 case: $(l - 2m)^2 = l + 4m$.

Its solutions are

$$l = l_i, \quad m = m_i, \qquad \text{or} \qquad l = l_i, \quad m = m_{i+1}, \quad i \in \mathbb{Z}, \tag{14}$$

where

• $\Lambda = 1$ (Adler-Moser) case: l and m are consecutive triangular numbers

$$l_i = i(2i+1), \quad m_i = i(2i-1).$$
 (15)

In terms of standard notations for the Adler-Moser polynomials P_n , we have $p_i = P_{2i}$ and $q_i = P_{2i+1}$, and (9) writes as

$$P_n''P_{n-1} - 2P_n'P_{n-1}' + P_nP_{n-1}'' = 0, \quad \deg(P_n) = n(n+1)/2.$$
(16)

• $\Lambda = 2$ case:

$$l_i = i(3i+2), \quad m_i = \frac{i(3i-1)}{2}.$$
 (17)

Now, consider bilinear relation (8) with $p = p_i$ and $q = q_i$ as a linear second-order ODE with solution q_i . Its second linearly independent solution $q = q_{i+1}$ equals

$$q_{i+1} = \left(\frac{2l_i}{\Lambda} - 2m_i + 1\right) q_i \int \frac{p_i^{2/\Lambda}}{q_i^2} dz.$$
(18)

Similarly, considering p_{i-1} and p_i as linearly independent solutions, we get

$$p_i = (2\Lambda m_i - 2l_{i-1} + 1) p_{i-1} \int \frac{q_i^{2\Lambda}}{p_{i-1}^2} dz.$$
(19)

By freedom in choosing linearly independent solutions of (8) one can also write analogs of (18) and (19) for decreasing indices

$$q_{i} = \left(\frac{2l_{i}}{\Lambda} - 2m_{i+1} + 1\right)q_{i+1} \int \frac{p_{i}^{2/\Lambda}}{q_{i+1}^{2}} dz, \quad p_{i-1} = \left(2\Lambda m_{i} - 2l_{i} + 1\right)p_{i} \int \frac{q_{i}^{2\Lambda}}{p_{i}^{2}} dz, \quad (20)$$

and one can generate p_i , and q_i iteratively in either directions starting at some *i*.

Rewriting (18,19) or (20) in the differential form, we get the first-order differential recurrence relations

$$\begin{aligned} q'_{i+1}q_i - q_{i+1}q'_i &= \left(\frac{2l_i}{\Lambda} - 2m_i + 1\right) p_i^{2/\Lambda}, \\ p'_i p_{i-1} - p_i p'_{i-1} &= \left(2\Lambda m_i - 2l_{i-1} + 1\right) q_i^{2\Lambda} \end{aligned}$$
(21)

In the case $\Lambda = 1$, the above recurrence relations are the three-term relations for the Adler-Moser polynomials P_n with $q_i = P_{2i}$ and $p_i = P_{2i+1}$:

$$P'_{i+1}P_{i-1} - P'_{i-1}P_{i+1} = (2i+1)P_i^2.$$
(22)

Examples of several first Adler-Moser polynomials are

$$P_0 = 1, \quad P_1 = z, \quad P_2 = z^3 + t_1, \quad P_3 = z^6 + 5t_1z^3 + t_2z - 5t_1^2, \quad \dots,$$
 (23)

where t_i are free parameters related to the KdV "times" [1] (we set parameter related to shift of z to zero).

In the $\Lambda = 2$ case we get

$$\begin{aligned} q'_{i+1}q_i - q_{i+1}q'_i &= (3i+1)p_i, \\ p'_ip_{i-1} - p_ip'_{i-1} &= (6i-1)q_i^4, \end{aligned}$$

where we used (17). In difference from the Adler-Moser case, there are two distinct branches of polynomials both starting at $p_0 = 1$ and $q_0 = 1$. One branch with $i \ge 0$ going in the positive direction

$$q_{0} = 1 \qquad p_{0} = 1 q_{1} = z \qquad p_{1} = z^{5} + t_{1} p_{2} = z^{16} + \frac{44}{7}s_{2}z^{12} - 32t_{1}z^{11} + 22s_{2}^{2}z^{8} - \frac{2112}{7}t_{1}s_{2}z^{7} + 1408t_{1}^{2}z^{6} + t_{2}z^{5} - 44s_{2}^{3}z^{4} + 352t_{1}s_{2}^{2}z^{3} - 1408s_{2}t_{1}^{2}z^{2} + 2816t_{1}^{3}z + t_{2}t_{1} - \frac{11}{5}s_{2}^{4}$$

$$(24)$$

etc, and another branch with $i \leq 0$ going in the negative direction

$$p_{0} = 1 \qquad q_{0} = 1 p_{-1} = z \qquad q_{0} = 1 q_{-1} = z^{2} + s_{-1} p_{-2} = z^{8} + \frac{28}{5}s_{-1}z^{6} + 14s_{-1}^{2}z^{4} \qquad q_{-2} = z^{7} + 7s_{-1}z^{5} + 35s_{-1}^{2}z^{3} + s_{-2}z^{2} + 28s_{-1}^{3}z^{2} + t_{-2}z - 7s_{-1}^{4} \qquad -35s_{-1}^{3}z + s_{-1}s_{-2} - \frac{5}{2}t_{-2}$$
(25)

etc, where t_i and s_i stand for arbitrary complex constants ² emerging in course of integrations in (18,19) or (20).

Summarizing, we emphasise again that, for $\Lambda = 1$ and $\Lambda = 2$, any no-common root solutions of the bi-linear equation (8) can be obtained by the recursive procedure described in this section. Therefore, the main sequences (23) or (25,24) determine all possible static patterns of a finite number of vortices with $Q_i \in \{-1, 1\}$ or $Q_i \in \{-1, 2\}$ respectively.

4 Darboux Transformations and Factorization Chains

The above results can be alternatively obtained with the help of factorization methods. The approach relying on these methods is more general than that used in the previous section and allows to obtain new stationary configurations. It involves theory of one-dimensional Schrödinger operator ("quantum Hamiltonian") with potential u(z):

$$H = -\partial_z^2 + u(z).$$

Let us now take the operator

$$H - \eta = -\partial_x^2 + u(z) - \eta, \qquad (26)$$

and factorize into the product of two first-order operators

$$H - \eta = A^* A,\tag{27}$$

where³

$$A^* = -\varkappa^{-1}\partial_z \varkappa = -\partial_z - \varkappa'/\varkappa, \quad A = \varkappa \partial_z \varkappa^{-1} = \partial_z - \varkappa'/\varkappa.$$
(28)

The function $\varkappa = \varkappa(z)$ is an arbitrary element of kernel of $H - \eta$, i.e. \varkappa is any eigenfunction of H corresponding to the eigenvalue η

$$H\varkappa = \eta\varkappa.$$

It trivially follows from (27) that for any λ

$$A(H - \lambda) = (\hat{H} - \lambda)A, \tag{29}$$

²Without loss of generality we omit a constant corresponding to a shift of z, so that $q_1 = p_{-1} = z$.

³Note that we do not use the special symbol \circ for composition of operators and instead e.g. $A^* \circ A$ or $\varkappa \circ \partial_z \circ \varkappa^{-1}$ etc we simply write A^*A or $\varkappa \partial_z \varkappa^{-1}$ etc.

where the new Schrödinger operator \hat{H} is obtained from H by permutation of factors A^* and A in (27):

$$\hat{H} = AA^* + \eta = -\partial_x^2 + \hat{u}, \quad \hat{u} = u - 2(\log \varkappa)''.$$
 (30)

Now, let us take an eigenfunction ψ corresponding to the eigenvalue λ of the old Schrödinger operator, i.e. $H\psi = \lambda\psi$. Then, from (29) it easy to see that the following transform of ψ :

$$\hat{\psi} = A\psi = \psi' - \frac{\varkappa'}{\varkappa}\psi, \tag{31}$$

is an eigenfunction of the new operator \hat{H} corresponding to the same eigenvalue λ , i.e.

$$\hat{H}\hat{\psi} = \lambda\hat{\psi}.$$

In other words, we can obtain a new operator \hat{H} and its eigenfunctions $\hat{\psi}(x;\lambda)$ from the old operator H and its eigenfunctions $\psi(x;\lambda)$ corresponding to the same eigenvalues λ . New operator and eigenfunctions are obtained with the help of an eigenfunction $\varkappa = \psi(x;\eta)$ of the old operator H. The transformation $H \to \hat{H}, \psi \to \hat{\psi}$ given by (30,31) is called the Darboux Transformation.

When $\lambda \neq \eta$ the transform (31) of the two dimensional kernel of $H - \lambda$ is also twodimensional. However, when $\lambda = \eta$, the transformation (31) annihilates the eigenfunction \varkappa of H, i.e. $A\varkappa = 0$: It maps the two-dimensional kernel $C_1\varkappa + C_2\varkappa \int \frac{dz}{\varkappa^2}$ of $H - \eta$ to the one dimensional sub-space

$$A\left[C_1\varkappa + C_2\varkappa \int \frac{dz}{\varkappa^2}\right] = \frac{C_2}{\varkappa}.$$

From the above, we see that function $1/\varkappa$ is an eigenfunction of the new Schrödinger operator corresponding to the eigenvalue η :

$$\hat{H}\varkappa^{-1} = \eta\varkappa^{-1}$$

Then, the general solution $\hat{\varkappa}$ of $\hat{H}\hat{\varkappa} = \eta\hat{\varkappa}$ is

$$\hat{\varkappa} = \frac{C}{\varkappa} \int \varkappa^2 dz, \qquad (32)$$

where the primitive of \varkappa^2 includes an arbitrary constant of integration. This extends the Darboux transformation of an eigenfunction (31) to the case $\lambda = \eta$. Transformations (32) are usually considered for $\eta = 0$ and are called the Darboux transformations at the zero energy level.

Darboux transformations can be iterated. Sequence of iterated transformations is presented by the factorization chain:

$$H_0 = A_0^* A_0 + \eta_1 \to H_1 = A_0 A_0^* + \eta_1 = A_1^* A_1 + \eta_2 \to H_2 = A_1 A_1^* + \eta_2 = A_2^* A_2 + \eta_3 \to \dots$$
(33)

When all $\eta_i = 0$, (33) becomes the chain of transformations at the zero energy level⁴

$$H_0 = A_0^* A_0 \to H_1 = A_0 A_0^* = A_1^* A_1 \to H_2 = A_1 A_1^* = A_2^* A_2 \to \dots,$$
(34)

⁴It would be more correctly to call it the chain of the Darboux transformations *associated with* zero energy level, rather than *at* the zero energy level, since eigenfunctions at all energy levels are transformed: Those with zero eigenvalue transform according to (32) and the rest according to (31).

where

$$A_i = \varkappa_i \partial_z \varkappa_i^{-1}, \quad A_i^* = -\varkappa_i^{-1} \partial_z \varkappa_i.$$

According to (32) and (30)

$$\varkappa_{i+1} = \frac{C_i}{\varkappa_i} \int \varkappa_i^2 dz, \quad u_{i+1} = u_i - 2(\log \varkappa_i)'', \quad H_i \varkappa_i = 0,$$
(35)

where C_i are arbitrary constants. Using (35), we can built iteratively the chain (34). Note, that at each step of the chain a free parameter, which is a constant of integration, appears in (35).

It trivially follows from (33) that H_0 and H_n are related by the intertwining operator T_n ,

$$H_n T_n = T_n H_0, \quad T_n = A_{n-1} A_{n-2} \cdots A_0.$$
 (36)

The function $T_n\psi(x;\lambda)$, where ψ is an eigenfunction of H_0 , ia an eigenfunction of H_n . Due to the Crum therem [10], the intertwining operator as well as potential u_n can be explicitly written down in terms of eigenfunctions of the initial hamiltonian H_0 . In more detail, the theorem states that the composition of n Darboux transformations, corresponding to the factorization chain (33), transforms the eigenfunction ψ , corresponding to the eigenvalue λ of $H_0 = -\partial_x^2 + u_0$ to the eigenfunction $\hat{\psi}$ of $H_n = -\partial_x^2 + u_n$, corresponding to the same eigenvalue, as the ratio of two wronskians

$$\hat{\psi} = \frac{\mathcal{W}[\psi_1, \psi_2, \dots, \psi_n, \psi]}{\mathcal{W}[\psi_1, \psi_2, \dots, \psi_n]}$$
(37)

while the potential is transformed in the following way

$$u_n = u_0 - 2 \left(\log \mathcal{W}[\psi_1, \psi_2, \dots, \psi_n] \right)''.$$
(38)

Here ψ_i are *n* eigenfunctions of the initial Schrodinger operator H_0 corresponding to the distinct eigenvalues $\eta_1, \eta_2, \ldots, \eta_n$, i.e. $H_0\psi_i = \eta_i\psi_i$, $1 \le i \le n$, called "seed functions", and \mathcal{W} stands for the Wronskian determinant. From (37), we get the Wronskian representation of the intertwining operator⁵

$$T_n[\cdot] = \frac{\mathcal{W}[\psi_1, \psi_2, \dots, \psi_n, \cdot]}{\mathcal{W}[\psi_1, \psi_2, \dots, \psi_n]}.$$
(39)

In the particular case of confluent eigenvalues, when the spacing between η_i vanishes, i.e. all η_i tend to one value η and we have the chain (34), the Crum theorem degenerates to

$$\varkappa_n = \frac{\mathcal{W}[\varkappa, \partial_\eta \varkappa, \dots, \partial_\eta^{n-1} \varkappa, \partial_\eta^n \varkappa]}{\mathcal{W}[\varkappa, \partial_\eta \varkappa, \dots, \partial_\eta^{n-1} \varkappa]}, \quad u_n = u_0 - 2\left(\log \mathcal{W}[\varkappa, \partial_\eta \varkappa, \dots, \partial_\eta^{n-1} \varkappa]\right)'', \quad (40)$$

where $\varkappa = \varkappa(z;\eta)$ is an eigenfunction of H_0 corresponding to the eigenvalue η , and \varkappa_n is an eigenfunction of H_n with the same eigenvalue. Here, the intertwining operator degenerates to

$$T_n[\,\cdot\,] = \frac{\mathcal{W}[\varkappa, \partial_\eta \varkappa, \dots, \partial_\eta^{n-1} \varkappa, \,\cdot\,]}{\mathcal{W}[\varkappa, \partial_\eta \varkappa, \dots, \partial_n^{n-1} \varkappa]}.$$

⁵Simplest proof of the Crum theorem uses the fact that, according to (31,36), T_n annihilates n seed functions ψ_i , $i = 1, \ldots, n$, i.e. kernel of T_n is spanned by these seed functions. Then, since $T_n = \partial_z^n + \ldots$, we immediately come to (39).

Eq. (40) is useful for representing the Adler-Moser polynomials in the Wronskian form. Indeed, let us take derivatives of the Schrödinger equation $H_0 \varkappa(z, \eta) = \eta \varkappa(z, \eta), H_0 = -\partial_z^2 + u_0(z)$ wrt η and then set $\eta = 0$. In this way, we get the recurrence relation for derivatives $\partial_{\eta}^n \varkappa$ at zero energy level:

$$H_0\partial_\eta^n \varkappa|_{\eta=0} = n\partial_\eta^{n-1} \varkappa|_{\eta=0}, \quad H_0 \varkappa|_{\eta=0} = 0.$$

Thus, (40) can be rewritten as

$$\varkappa_n = \frac{\tau_n}{\tau_{n-1}}, \quad \tau_n = \mathcal{W}[\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_n], \quad u_n = u_0 - 2(\log \tau_{n-1})'', \quad \tau_0 = 1,$$
(41)

where the η -independent functions $\mathcal{X}_n(z)$ are defined by the recurrence relation

$$H_0 \mathcal{X}_n = c_n \mathcal{X}_{n-1}, \quad H_0 \mathcal{X}_1 = 0, \quad H_0 = -\partial_z^2 + u_0(z).$$
 (42)

In eq. (42), c_n are arbitrary constants. At each iteration in (42) two integration constants appear, but only one of them is essential. The intertwining operator writes in terms of \mathcal{X}_i as

$$T_n[\cdot] = \frac{\mathcal{W}[\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_{n-1}, \cdot]}{\mathcal{W}[\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_{n-1}]}.$$
(43)

When we start the factorization chain (34) with the free Schrödinger operator $H_0 = -\partial_z^2$ (i.e. with $u_0 = 0$), from (42), we get

$$\mathcal{X}_{n+1}''(z) = (2n+1)\mathcal{X}_n(z), \quad \mathcal{X}_1 = z.$$
 (44)

Here⁶, the function \mathcal{X}_n is a polynomial of degree 2n-1 in z. Therefore τ_n in (41) is a polynomial of degree n(n + 1)/2. Substituting $\varkappa_n = \tau_n/\tau_{n-1}$ (see(41)) into (35), we obtain recurrence relation for τ_n which coincides with that (22) for the Adler-Moser polynomials P_n . Taking into account that degrees of τ_n and P_n are the same, we come to the conclusion that $\tau_n = P_n$. Thus, we get the Wronskian representation for the Adler-Moser polynomials

$$P_n = \mathcal{W}[\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_n],$$

where \mathcal{X}_n are defined by (44).

5 Darboux Transformation and Equilibrium Configurations

As we mentioned above, sequences of the equilibrium configurations can be produced with help of Darboux transformations. To built them, we need to introduce the factorizable function corresponding to a static configuration (6)

$$\psi(z) = C \prod_{i=1}^{N} (z - z_i)^{\mathcal{Q}_i},$$
(45)

⁶In (44) we set $c_n = 2n + 1$. For such choice of normalization constants, τ_n is monic in z.

where C is a non-zero constant and Q_i are not necessarily integers.

Now, we will show that if the Darboux transform $\hat{\psi}$ of ψ has a similar form, i.e.

$$\hat{\psi}(z) = \hat{C} \prod_{i=1}^{\hat{N}} (z - \hat{z}_i)^{\hat{\mathcal{Q}}_i}, \tag{46}$$

then $\hat{\psi}$ also corresponds to a static configuration. For this we introduce the Schrödinger operator H whose kernel contains ψ , i.e.

$$H\psi = (-\partial_z^2 + u)\psi = 0.$$

Then, using (45), we obtain

$$u = \frac{\psi''}{\psi} = (\log \psi)'' + ((\log \psi)')^2 = -\sum_i \frac{Q_i}{(z - z_i)^2} + \sum_{i,j} \frac{Q_i Q_j}{(z - z_i)(z - z_j)}.$$

Applying the partial fraction decompositions in the last sum we get

$$u = \sum_{i} \frac{\mathcal{Q}_i(\mathcal{Q}_i - 1)}{(z - z_i)^2} + 2\sum_{i} \frac{\mathcal{Q}_i}{(z - z_i)} \sum_{j \neq i} \frac{\mathcal{Q}_j}{(z_i - z_j)}.$$

Due to the equilibrium condition (6) the last term in the above equation vanishes and potential is a sum of the second order poles

$$u = \sum_{i} \frac{Q_i(Q_i - 1)}{(z - z_i)^2},$$
(47)

i.e. the equilibrium condition is equivalent to absence of the first-order poles in the potential.

Now, to prove that the Darboux transform $\hat{\psi}$ of ψ of the form (46) also correspond to an equilibrium configuration, i.e.

$$\sum_{j=1,j\neq i}^{\hat{N}} \frac{\hat{\mathcal{Q}}_j}{\hat{z}_i - \hat{z}_j} = 0, \quad i = 1\dots\hat{N},$$

we have to show that potential \hat{u} of the Darboux transformed Schrödinger operator \hat{H} is also a sum of the second order poles. Indeed, the Darboux transformed potential equals (see (32))

$$\hat{u} = u - 2(\log \psi)'',$$

and from (45) and (47) we obtain

$$\hat{u} = \sum_{i} \frac{\mathcal{Q}_i(\mathcal{Q}_i + 1)}{(z - z_i)^2},\tag{48}$$

i.e. \hat{u} is a sum of second order poles and therefore $\hat{\psi}$ of the form (46) corresponds to a static configuration.

Note, that since

$$\hat{u} = \sum_{i} \frac{\hat{\mathcal{Q}}_i(\hat{\mathcal{Q}}_i - 1)}{(z - \hat{z}_i)^2},$$

from (48), we see that the Darboux transformation acts on charges either of two ways

$$\hat{\mathcal{Q}} = -\mathcal{Q}, \quad \hat{\mathcal{Q}} = \mathcal{Q} + 1,$$
(49)

i.e. it either (i) inverts a charge or (ii) increments a charge by 1. In particular, it can create new charges with $Q_i = 1$ and annihilate charges with $Q_i = -1$. In the case of the Adler-Moser polynomials in generic configurations, it annihilates negative charges, inverts positive charges, and creates positive charges at new positions.

6 Configurations Related to Bi-spectral Systems

In section 3 we showed that the static configurations of the system consisting of charges of values ± 1 (i.e. when $\Lambda = 1$) correspond to zeros of the Adler-Moser polynomials P_n and P_{n+1} . In section 4 these polynomials were obtained with the help of sequence of the Darboux transformations (34) at zero energy level. Corresponding sequence of H_i is started from the free Schrodinger operator $H_0 = -\partial_z^2$. This sequence is known to constitute the "odd" family of bi-spectral operators⁷ [19]. There exists another sequence of the Darboux transforms consisting of "even" family, of bi-spectral operators [19]. It starts with

$$H_0 = -\partial_z^2 - \frac{1}{4z^2}.$$
 (50)

In both "even" and "odd" cases we present elements of the sequence of Darboux-transforms of kernels of H_i

$$\psi_0 \to \psi_1 \to \psi_2 \to \dots, \quad H_i \psi_i = 0$$

as the ratios

$$\psi_i = P_{i+1}/P_i, \quad P_0 = 1.$$
 (51)

According to section 4 (see eq. (32)) the Darboux transform ψ_{i+1} of ψ_i is

$$\psi_{i+1} = \frac{C_i}{\psi_i} \int \psi_i^2 dz.$$
(52)

From this and (51) we get the differential recurrent relation for P_i

$$P'_{i+1}P_{i-1} - P'_{i-1}P_{i+1} = C_i P_i^2.$$
(53)

In the "odd", i.e. the Adler-Moser case $P_0 = 1$, $P_1 = z$ (see eq. (22)). In the "even" case $H_0[z^{1/2}] = 0$ (i.e. $\psi_0 = z^{1/2}$ see (50)) and, according to (51)

$$P_0 = 1, \quad P_1 = z^{1/2}.$$
 (54)

⁷The differential operators H in z is bi-spectral if there exists a differential operator B in k and a common eigenfunction, such that $H\psi(z,k) = f(k)\psi(z,k)$, $B\psi(z,k) = g(z)\psi(z,k)$. For the even family of Schrödinger operators, the common eigenspace of H and B is two-dimensional. It is one-dimensional in the odd case.

Note that, for convenience, we set the normalization constants C_i in (53) to values at which $P_i(z)$ are monic⁸. For "even" family, from (53,54) we get $P_2 = z^2 + t_1$. So, using (51), we obtain the sequence of length 2 depending on one parameter t_1 :

$$\psi_0 = z^{1/2}, \quad \psi_1 = \frac{z^2 + t_1}{z^{1/2}}.$$
 (55)

Another application of the recurrence relation (53) with $t_1 \neq 0$ would produce a logarithmic term, so to obtain ψ_2 and ψ_3 of the form (45) we restrict t_1 to be zero and pick a new integration constant t_2 :

$$P_3 = z^{9/2} + t_2 z^{1/2}, \quad P_4 = z^8 + 6t_2 z^4 + t_3 z^2 - 3t_2^2.$$
(56)

Now, we get the sequence of length 4:

$$\psi_0 = z^{1/2}, \quad \psi_1 = z^{3/2}, \quad \psi_2 = \frac{z^4 + t_2}{z^{3/2}}, \quad \psi_3 = \frac{z^8 + 6t_2z^4 + t_3z^2 - 3t_2^2}{z^{9/2} + t_2z^{1/2}}$$

depending on two parameters t_2, t_3 .

Next, to eliminate logarithmic terms in P_5 and P_6 , one has to set both $t_1 = t_2 = 0$ and we get a sequence $\psi_0, \psi_1, \psi_2, \psi_3, \psi_4, \psi_5$ (of length 6) depending on three parameters t_3, t_4 and t_5 . And so forth.

Sequences of the even family have finite lengths (i.e. they terminate) due to the presence of charge of the third specie at the origin: At the beginning of sequence, charge 1/2 is placed at z = 0. Then, according to (49), the following transformations are produced with this charge: It is incremented by 1 during each of the m - 1 first Darboux transformations. It is inverted by the *m*th transformation and becomes equal to -m + 1/2. Then, again, it is incremented during the next m-1 steps, until it reaches value -1/2. This charge would create a logarithmic singularity in (52) at the next step, so sequence terminates.

Closing this section we note that similarly to the odd family, the wronskian representation of ψ_i exists also for the even family. It will be given in section 11 (eqs. (87) and (88)).

7 KWCC Transformation and Terminating Configurations

Let us now return to the sequence of transformations of rational functions (whose primitives are also rational functions) corresponding to 9 iterations (18,19)

$$\cdots \to \frac{q_i^{2\Lambda}}{p_{i-1}^2} \to \frac{p_i^{2/\Lambda}}{q_i^2} \to \frac{q_{i+1}^{2\Lambda}}{p_i^2} \to \frac{p_{i+1}^{2/\Lambda}}{q_{i+1}^2} \to \dots$$
(57)

Denoting function $\frac{q_i^{2\Lambda}}{p_{i-1}^2}$ from the above chain as ψ^2 and the next function $\frac{p_i^{2/\Lambda}}{q_i^2}$ as $\tilde{\psi}^2$, from (19) we get the relationships between them

$$\psi \to \tilde{\psi}, \quad \tilde{\psi}(z) = C \left(\frac{1}{\psi(z)} \int \psi(z)^2 dz\right)^{\gamma},$$
(58)

⁸In the even case, this differs from the original normalization of [19], where same $C_i = 2i + 1$ were chosen for both families.

⁹Or corresponding to the chain of solutions $\cdots \rightarrow p_{i-1}, q_i \rightarrow p_i, q_i \rightarrow p_i, q_{i+1} \rightarrow p_{i+1}, q_{i+1} \rightarrow \dots$ of (8).

where, for this step of the chain, $\gamma = 1/\Lambda$ and C stands for a non-zero constant.

Next transformation in the sequence (57) relates $\psi^2 = \frac{p_i^{2/\Lambda}}{q_i^2}$ and $\tilde{\psi}^2 = \frac{q_{i+1}^{2\Lambda}}{p_i^2}$. Using (18) we also arrive to transformation (58), this time with $\gamma = \Lambda$.

Therefore, the chain (57) can be built by the iterative applications of following transformations

$$\psi_{i+1} = C_i \left(\frac{1}{\psi_i(z)} \int \psi_i(z)^2 dz\right)^{\gamma_{i+1}},\tag{59}$$

where

$$\gamma_{i+1} = 1/\gamma_i,\tag{60}$$

and

$$\gamma_i \in \{1/2, 1, 2\}.$$
 (61)

The transformation (58) was presented by Krishnamurthy, Wheeler, Crowdy and Constantin (KWCC) in [20].

The KWCC transformation (58) is a composition of two mappings

- (i) The Darboux transformation at zero energy level $\hat{\psi} = \frac{C}{\psi} \int \psi^2 dz$ (see (32)) and
- (ii) Exponentiation $\tilde{\psi} = \hat{\psi}^{\gamma}$.

According to section 5, the Darboux transformation is a mapping between static configurations, provided the reslut of transformation $\hat{\psi}$ is factorizable, i.e. has a form (46). The exponentiation corresponds to scaling of all charges $\hat{Q}_i \to \gamma \hat{Q}_i$ and is also a mapping between static configurations. Therefore, when $\int \psi^2 dz$ is rational, the KWCC transformation is also such a mapping. Rationality of primitive of ψ_i^2 imposes restrictions on γ_i and on constants of integration.

From the above and (49) it follows that KWCC transformation (58) acts on charges either of two ways

$$\tilde{\mathcal{Q}} = -\gamma \mathcal{Q}, \quad \tilde{\mathcal{Q}} = \gamma \left(\mathcal{Q} + 1 \right).$$
 (62)

Obviously, the KWCC transformations (59,60,61) generate all classes of configurations considered above, i.e. those related to the odd and even bi-spectral families as well as to the $\Lambda = 2$ case.

To rewrite the sequence of transforms (59), (60) in terms of polynomials, we present ψ_i as

$$\psi_i = \frac{\tau_i^{\gamma_i}}{\tau_{i-1}},$$

where τ_i is a polynomial of an integer or half-integer degree d_i . Here, "polynomial of half-integer degree" in z stands for a polynomial in z times $z^{1/2}$. Then from (59) and (60) we get

$$\tau_{i+1} = (2\gamma_i d_i - 2d_{i-1} + 1)\tau_{i-1} \int \frac{\tau_i^{2\gamma_i}}{\tau_{i-1}^2} dz,$$

where

$$d_{i+1} = 2\gamma_i d_i - d_{i-1} + 1, \quad \gamma_{i+1} = 1/\gamma_i, \quad d_0 = 0.$$
(63)

In the differential form these relations write as

$$\tau_{i+1}'\tau_{i-1} - \tau_{i+1}\tau_{i-1}' = (2\gamma_i d_i - 2d_{i-1} + 1)\tau_i^{2\gamma_i}, \quad \tau_i(z) = z^{d_i} + \dots, \quad \tau_0 = 1.$$
(64)

These are actually relations (21), but now d_i are not necessarily degrees (15) or (17) of the main sequences. This is because (63, 64) are derived from the KWCC chain, rather than from the bi-linear equation (8). Because of this fact, any polynomial solution of (64) that starts from a static configuration (e.g. from a single vortex) also corresponds to a sequence of static configurations.

Apart from infinite main sequences of the $\Lambda = 1$ and $\Lambda = 2$ cases, recurrence relations (63,64) allow to find sequences of configurations of finite lengths, i.e. terminating configurations. The "even" bi-spectral family of section 6, i.e. solutions of (63,64) corresponding to $\gamma_i = 1$, $d_1 = 1/2$, is a family of terminating configurations for $\Lambda = 1$.

As an example of family of terminating configurations for $\Lambda = 2$, we can take the case $\gamma_0 = 2$, $d_1 = 2$, i.e. $\tau_1 = z^2$. Here are several first terminating sequences of τ_i :

$$\begin{array}{c}1,z^2,z^3+t_1\\1,z^2,z^3,z^2(z^9+t_2),z^9+t_3z^3-2t_2\\1,z^2,z^3,z^{11},z^3(z^6+t_3),z^2(z^{24}+\frac{20}{3}z^{18}t_3+30z^{12}t_3^2+z^9t_4-20z^6t_3^3-\frac{5}{3}t_3^4),\\z^{18}+15z^{12}t_3+z^9t_5-45z^6t_3^2+\frac{2t_3t_5-3t_4}{2}z^3+5t_3^3\end{array}$$

Similarly to the even bi-spectral family, here, a charge of the third specie undergoes transformations (62) at z = 0 (this time with $\gamma \in \{1/2, 2\}$) until the logarithmic singularity is encountered.

One can compute terminating sequences by solving (64) for τ_{i+1} as for a monic polynomial with unknown coefficients. Compatibility of resulting system of algebraic equations (which is linear in coefficients of τ_{i+1}) imposes restrictions on free parameters of τ_i . For sequences related to the even bi-spectral family there exists a wronskian representation (see section 11).

We recall that the main sequences exhaust all possible no-common root polynomial solutions of (8) as well as the collapse solutions for $\Lambda = 1$ and $\Lambda = 2$ cases. Solutions of (64) whose degrees are not in the main sequences do not satisfy bi-linear equations (8). They satisfy its generalization.

In more detail: Let us first consider sequence of the Darboux transforms at zero energy level that starts from $H_0 = -\partial_z^2 + u_0$, with non-zero $u_0 \neq 0$. At the *n*th step of the chain we have

$$u_n = u_0 - 2(\log \tau_n)'', \quad \varkappa_n = \frac{\tau_{n+1}}{\tau_n}, \quad \tau_0 = 1, \quad H_n \varkappa_n = 0,$$

Therefore

$$H_n \varkappa_n = \left(-\partial_z^2 + u_0 - 2(\log \tau_n)''\right) \left[\frac{\tau_{n+1}}{\tau_n}\right] = 0,$$

or equivalently

$$\tau_{n+1}''\tau_n - 2\tau_{n+1}'\tau_n' + \tau_{n+1}\tau_n'' - u_0\tau_n\tau_{n+1} = 0.$$
(65)

This is a generalization of the Tkachenko equation (9) that includes cases of non-integer d_i . For the even bi-spectral family, the initial potential equals to $u_0 = a(a-1)/z^2$, where a is a half-integer. Then, substituting $\tau_n = p$, $\tau_{n+1} = z^a q$ into (65), we get

$$p''q - 2p'q' + pq'' + \frac{2a}{z}(q'p - p'q) = 0.$$

In fact, this generalization is a particular case of the tri-linear equation for three species of vortices with $Q_i \in \{-1, 1, a\}$, where a single vortex of the third specie of strength a is placed at the origin z = 0 [17, 30]. In general, for multiple species we have the multi-linear equation (see e.g. [23, 29])

$$\sum_{i} \Lambda_i^2 \frac{\mathcal{P}_i''}{\mathcal{P}_i} + 2 \sum_{i < j} \Lambda_i \Lambda_j \frac{\mathcal{P}_i'}{\mathcal{P}_i} \frac{\mathcal{P}_j'}{\mathcal{P}_j} = 0,$$

where sums run over all species, Λ_i is a strength of the *i*th specie and \mathcal{P}_i is polynomial whose roots are coordinates of vortices of this specie. In the case of configurations related to the even bi-spectral family $\mathcal{P}_1 = p$, $\mathcal{P}_2 = q$, $\mathcal{P}_3 = z$. In similar situation, when $\mathcal{Q}_i \in \{-1, \Lambda, \Lambda a\}$, we have the following generalization of (8)

$$p''q - 2\Lambda p'q' + \Lambda^2 pq'' + \frac{2a\Lambda}{z}(\Lambda q'p - p'q) = 0.$$

Exploring this equation, O'Neil and Cox-Steib [30] presented the families of terminating configurations.

It is worth mentioning that all families of equilibrium patterns obtained above are related to the following chain of transformations that starts from the free Schrödinger operator

$$A_{-1}A_{-1}^* \xrightarrow{\gamma_0} A_0^*A_0 \to A_0A_0^* \xrightarrow{\gamma_1} A_1^*A_1 \to \dots, \quad \gamma_i \in \{1/2, 1, 2\},$$

where $A_{-1} = -A_{-1}^* = \partial_z$, and $\xrightarrow{\gamma_i}$ stand for exponentiations of kernels. For example, for the even bi-spectral family $\gamma_0 = 1/2$ and $\gamma_i = 1$, i > 0. Chains of this type are generalizations of (34).

8 Translating Configurations and Baker-Akhieser Functions

Translating configuration of vortices are solutions of (8). To generate them we consider the following eigenfunction

$$\psi(k,z) = C \prod_{i=1}^{N} (z - z_i(k))^{\mathcal{Q}_i} e^{kz},$$
(66)

of the Schrodinger operator

$$H\psi = (-\partial_z^2 + u)\psi = -k^2\psi.$$

The condition that potential $u = \psi''/\psi - k^2$ does not have simple poles (i.e. it is of form (47)) is equivalent to the equilibrium conditions (8) for a translating configuration. This can be shown by arguments similar to that used for the static configurations in Section 5.

In the case of the factorization chain (34) starting with the free Schrodinger operator, there exists an eigenfunction, called the "rational Baker-Akhieser function"¹⁰, which has the form (66). It is an eigenfunction of H_j that is obtained by the action of an intertwining

¹⁰In theory of integrable hierarchies of non-linear PDEs, the Baker-Akhieser function is a solution of an infinite system of linear differential equations. Associated integrable hierarchy arises as a condition of compatibility of the system. Here we deal with the Baker-Akhieser function of the KdV hierarchy.

operator with rational coefficients on the eigenfunction e^{kz} of the free Schrödinger operator, i.e. $\Psi = C(k)T_j[e^{kz}]$. Since the z-independent factor C(k) does not play role in our considerations, we will use the eigenfunction

$$\psi_j = T_j[e^{kz}] = \frac{q_j}{p_j}e^{kz},\tag{67}$$

where p_j and q_j are polynomials in z. As we have shown in section 6, the corresponding potential u_j , obtained by j Darboux transformations from $u_0 = 0$, is of the form (47). Since u_j does not contain simple poles, eigenfunction (67) corresponds to a translating configuration. It will be shown in section 12 that the class of such configurations contains all possible translating patterns consisting of finite number of vortices with $Q_i = \pm 1$.

A constructive way to write down ψ_j is to use the Wronskian representation of T_j , given by (43, 44). Then, from (67) we get

$$\psi_j = \frac{\mathcal{W}\left[\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_j, e^{kz}\right]}{\mathcal{W}\left[\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_j\right]}, \quad \mathcal{X}_1 = z, \quad \mathcal{X}_j'' = (2j+1)\mathcal{X}_{j-1},$$

and from (67) it follows that

$$q_j = e^{-kz} \mathcal{W} \left[\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_j, e^{kz} \right], \quad p_j = \mathcal{W} \left[\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_j \right] = P_j, \tag{68}$$

where P_j is the *j*th Adler-Moser polynomial. When $k \neq 0$, degree of $q_j(z)$ equals to that of $p_j(z)$. In the electrostatic interpretation (see section 1), this means that the total charge has to be zero $\sum_i Q_i = 0$ in order the system to be at rest in the homogeneous electric field $k \neq 0$. In the $k \to 0$ limit

$$\psi_j(k=0,z) = \text{const}\frac{P_{j-1}}{P_j},$$

and we get the static equilibrium with non-zero total charge. In other words, in the $k \to 0$ limit a part of positive charges moves to infinity.

9 Equilibria in Non-Homogenous Background Fields

Let us assume that the Schrödinger operator $H = -\partial_z^2 + u(z)$ has the following "quasi-rational" eigendunction

$$\psi(z) = C \prod_{i=1}^{N} (z - z_i)^{\mathcal{Q}_i} e^{\Phi(z)}, \quad H\psi = \lambda \psi,$$
(69)

where $\Phi(z)$ is a polynomial. "Quasi-rational" here means that all Q_i are integers. Requirement that potential $u = \frac{\psi''}{\psi} + \lambda$ does not have simple poles results in the system of algebraic equations

$$\Phi'(z_i) + \sum_{i=1, i \neq j}^{N} \frac{\mathcal{Q}_j}{z_i - z_j} = 0, \quad i = 1 \dots N.$$
(70)

Equations (70) are conditions of equilibrium of charges in an external harmonic electric field¹¹ or equilibrium of vortices in a background flow, see section 1. Function $\Phi(z)$ is a potential of

¹¹Such conditions were first presented by Stieltjes [33, 34] for roots of orthogonal polynomials.

the external field/background flow. Provided (70) holds,

$$u(z) = \sum_{i} \frac{\mathcal{Q}_{i}(\mathcal{Q}_{i}-1)}{(z-z_{i})^{2}} + \Phi''(z) + \Phi'(z)^{2} + 2\sum_{i} \mathcal{Q}_{i} \frac{\Phi'(z) - \Phi'(z_{i})}{z-z_{i}} - \lambda.$$

If there exists another quasi-rational eigenfunction of H which is of the form similar to (69)

$$\tilde{\psi}(z) = C \prod_{i=1}^{\tilde{N}} (z - \tilde{z}_i)^{\tilde{\mathcal{Q}}_i} e^{\Phi(z)}, \quad H\tilde{\psi} = \eta \tilde{\psi},$$

it can be taken as a "seed" for the Darboux transformation. Then the transform $\hat{\psi}$ of ψ

$$\hat{\psi} = \psi' - \frac{\psi'}{\tilde{\psi}}\psi, \quad \hat{H}\hat{\psi} = (-\partial_z^2 + \hat{u})\hat{\psi} = \lambda\hat{\psi}$$
(71)

will be also quasi-rational of the form similar to (69), i.e.

$$\hat{\psi}(z) = C \prod_{i=1}^{\hat{N}} (z - \hat{z}_i)^{\hat{\mathcal{Q}}_i} e^{\Phi(z)}$$

Since the transform of potential

$$\hat{u} = u - 2(\log \tilde{\psi})''$$

does not have first-order poles, $\hat{\psi}$ corresponds to a new equilibrium configuration in the same external field. For a set of eigenfunctions of H ("seed functions") of the form

$$\varkappa_i = R_i(z)e^{\Phi(z)}, \quad H\varkappa_i = \lambda_i\psi_i,$$

where $R_i(z)$ are rational in z, we can apply several Darboux transformations resulting in various equilibrium configurations. According to the Darboux-Crum theorem (37,38), the results of compositions of n transformations can be written in the Wronskian form

$$\psi^{(n)}(z) = \frac{\mathcal{W}[\varkappa_{k_1}, \varkappa_{k_2}, \dots, \varkappa_{k_{n-1}}, \varkappa_{k_n}]}{\mathcal{W}[\varkappa_{k_1}, \varkappa_{k_2}, \dots, \varkappa_{k_{n-1}}]}.$$
(72)

For example, for harmonic oscillator $u = z^2/2$, $\Phi(z) = -z^2/2$ and R_n are Hermite polynomials. In this case, transforms (72) correspond to various equilibria of charges in the external quadrupole field or vortex patterns in vicinity of the stagnation point of the flow encountering a solid boundary (for details and other examples, see e.g. [14, 23]). Note, that in difference from the systems considered in the previous sections, configurations corresponding to Hermite polynomials do not depend non-trivially on continuous parameters and are rather parametrised by sequences of distinct integers $k_1 < k_2 < \cdots < k_n$.

10 Equilibria of Vortex Streets - Odd Family

A periodic vortex street is a vortex pattern consisting of an infinite number of vortices situated periodically along some direction. A vortex street can be equivalently considered as a configuration of N vortices (or charges) on a cylinder. In what follows by the terms "vortex street", "periodic configuration" or "periodic pattern" we mean configuration of a finite number of charges on cylinder. Without loss of generality we can set the translation period to π . Since $\sum_{n=-\infty}^{\infty} \frac{1}{z+\pi n} = \cot z$, the stationarity condition (5) for the translating (with a speed $i\bar{k}/(2\pi)$) vortex street becomes

$$k + \sum_{j=1, j \neq i}^{N} \mathcal{Q}_j \cot(z_i - z_j) = 0, \quad i = 1 \dots N,$$
(73)

and for static periodic patterns (see (6)) we have

$$\sum_{j=1, j \neq i}^{N} \mathcal{Q}_j \cot(z_i - z_j) = 0, \quad i = 1 \dots N.$$
(74)

Similarly to vortex patterns on the plane that are related to the rational solutions of the KdV hierarchy or the odd bi-spectral family, the trigonometric soliton solutions of the hierarchy are related to the patterns on a cylinder [16, 23]. The latter can be shown by methods similar to those used in sections 5, 8.

In more detail, we introduce the trigonometric-rational analog of (45) of the following form

$$\psi(z) = C \prod_{i=1}^{N} \sin^{\mathcal{Q}_i} (z - z_i), \tag{75}$$

where Q_i , z_i correspond to the periodic static configuration (74). The "trigonometric-rational" here means that all Q_i are integers. Trigonometric-rational are special cases of the trigonometricfactorizable functions, where Q_i are real. Function (75) is an eigenfunction of the Schrödinger operator

$$H\psi = \lambda\psi, \quad H = -\partial_z^2 + u(z)$$

with the potential

$$u(z) = \frac{\psi''(z)}{\psi(z)} + \lambda.$$

Taking into account the identity

$$\cot(z - z_i)\cot(z - z_j) = -1 + (\cot(z - z_i) - \cot(z - z_j))\cot(z_i - z_j),$$

similarly to the rational case, we come to the conclusion that the static equilibrium condition is equivalent to absence of first-order poles in the potential and

$$u = \sum_{i=1}^{N} \frac{\mathcal{Q}_i(\mathcal{Q}_i - 1)}{\sin^2(z - z_i)} + \text{const.}$$

$$(76)$$

A family of the trigonometric-rational configurations can be generated by the chain of Darboux transformations (33) with $\eta_i = k_i^2$, where $0 < k_1 < k_2 < \cdots < k_i < \ldots$ is a sequence of positive integers. The chain starts from the free Schrödinger operator $H_0 = -\partial_z^2$ and it is built with the help of "seed functions" that are its periodic eigenfunctions

$$\varkappa_{k_i}(z) = \sin(k_i z + \zeta_i). \tag{77}$$

Then, according to the Crum theorem (37,38), at *n*th step of the chain we have

$$\psi^{(n)}(z) = \frac{\tau_{n+1}(z)}{\tau_n(z)}, \quad \tau_n = \mathcal{W}\left[\varkappa_{k_1}, \varkappa_{k_2}, \dots, \varkappa_{k_n}\right], \quad \tau_0 = 1.$$
(78)

Obviously, $\tau_n(z)$ are trigonometric polynomials, $\psi^{(n)}(z)$ are trigonometric-rational functions and corresponding potentials

$$u_n(z) = -2(\log \tau_n(z))''$$
(79)

are of the form (76) and do not have simple poles. Thus, eigenfunctions (78) correspond to the static vortex street configurations. Note that potentials (79) are periodic *n*-soliton solutions of the KdV hierarchy and τ_n are its *n*-soliton τ -functions (see e.g. [26] and references therein). We will demonstrate in section 12 that the set of these configurations actually contains all static vortex street patterns with $Q_i = \pm 1$.

For translating configurations (see (73)) related to the soliton solutions we can apply metod similar to that used in section 8. By analogy with quasi-rational (66), we now consider the trigonometric quasi-rational eigenfunctions

$$\psi(k,z) = C \prod_{i=1}^{N} \sin(z - z_i(k))^{\mathcal{Q}_i} e^{kz}, \quad \mathcal{Q}_i \in \mathbb{Z}.$$
(80)

According to the Darboux-Crum theorem (37), eigenfunctions, obtained by a sequence of the Darboux transformation from the formal eigenfunction e^{kz} of the free Schrödinger operator, equal

$$\psi^{(n)}(k,z) = \frac{\mathcal{W}\left[\varkappa_{k_1}, \varkappa_{k_2}, \dots, \varkappa_{k_n}, e^{kz}\right]}{\mathcal{W}\left[\varkappa_{k_1}, \varkappa_{k_2}, \dots, \varkappa_{k_n}\right]},\tag{81}$$

with \varkappa_{k_i} given by (77). Obviously $\psi^{(n)}(k, z)$ are of the form (80) and therefore correspond to the translating vortex street configurations. Since each \varkappa_{k_i} (77) depends on a free parameter, the translating and static configurations related to soliton solutions depend on n free parameters as well as on the sequence of integers $0 < k_1 < k_2 \cdots < k_n$. Up to a z-independent common factor C(k), (81) is the trigonometric Baker-Akhieser function of the KdV hierarchy¹².

The class of configurations determined by (81,77) contains all possible translating vortex street patterns with $Q_i = \pm 1$, as will be shown in section 12.

11 Equilibria of Vortex Streets - Even Family

In the previous section we considered vortex street configurations related to the trigonometric analogs of the odd family of bi-spectral potentials or soliton solutions of the KdV hierarchy. Here we consider trigonometric analogs of the even family (see section 6), i.e. examples of periodic configurations where not all Q_i are integers. For this, let us start our chain of Darboux transformations with the trigonometric Pochl-Teller potential

$$u_0 = \frac{a(a-1)}{\sin^2 z} + \frac{b(b-1)}{\cos^2 z}$$
(82)

 $^{^{12}}$ It is also the "angular part" of the Baker-Akhieser function of 2d quantum algebraically integrable systems, see e.g. [5],[12] and references therein.

with arbitrary real a and b. The simplest configuration corresponds to the eigenfunction $\sin^a z \cos^b z$ and consists of two static vortices of the strengths $Q_1 = a$, $Q_2 = b$ situated at the diametrically opposite points of the π -periodic cylinder, i.e. at $z_1 = 0$, $z_2 = \pi/2$.

Note that when a = b, or a = 1 - b, the potential (82) can be rewritten as $u_0 = 4a(a - 1)/\sin^2 2z$. By re-scaling $2z \to z$ in the Schrödinger operator, we again come to the special b = 0 case of the π -periodic potential (82).

For integers a and b, (82) is a soliton solution of KdV at specific "times" (corresponding parameters k_i and ζ_i in (78, 77) can be found e.g. in refs [6] or [15])¹³. In this case, u_0 and sequences of its Darboux transforms belong to the odd family considered in the previous section.

Due to invariance of the potential (82) wrt involutions $a \to 1-a$, $b \to 1-b$, when $a \pm b$ is not integer (generic *a* and *b*), the Schrödinger operator $H_0 = -\partial_z^2 + u_0$ has four distinct types of factorizable eigenfunctions $\varkappa_i(z)$, $H_0\varkappa_i = \lambda_i\varkappa_i$ of the form

$${}_{2}F_{1}(-i, i + a + b, 1/2 + a; \sin^{2} z) \sin^{a} z \cos^{b} z, \qquad \lambda_{i} = (2i + a + b)^{2}, \\ {}_{2}F_{1}(-i, i + 1 - a + b, 3/2 - a \sin^{2} z) \sin^{1-a} z \cos^{b} z, \qquad \lambda_{i} = (2i + 1 - a + b)^{2}, \\ {}_{2}F_{1}(-i, i + a + 1 - b, 1/2 + a; \sin^{2} z) \sin^{a} z \cos^{1-b} z, \qquad \lambda_{i} = (2i + 1 + a - b)^{2}, \\ {}_{2}F_{1}(-i, i + 2 - a - b, 3/2 - a; \sin^{2} z) \sin^{1-a} z \cos^{1-b} z, \qquad \lambda_{i} = (2i + 2 - a - b)^{2}, \end{cases}$$

$$(83)$$

where i = 0, 1, 2, ... and $_2F_1$ stands for the Gauss hypergeometric function. The first parameter of the $_2F_1$ in (83) is a non-positive integer -i, and the hypergeometric factors in (83) are polynomials in $\sin^2 z$. These factorizable eigenfunctions are trigonometric-rational up to a factor $\sin^{\pm a} z \cos^{\pm b} z$. For generic *a* and *b* all eigenvalues λ_i are distinct and all eigenfunctions in (83) are linearly independent.

From (83), with the help of the Wronskian formula (78), we can construct eigenfunctions of Darboux transforms $H_n = -\partial_z^2 + u_0 - 2(\log \tau_n)''$ of H_0 . These eigenfunctions are factorizable, and therefore correspond to static vortex street configurations. Note that, for generic a and b, only one of two linearly independent solutions of the Schrödinger equation $H_0 \varkappa_i = \lambda_i \varkappa_i$ is factorizable. Therefore, in difference from the soliton-related configurations (78,77), configurations generated by (83) do not depend on free parameters others than a and b. Similarly to systems considered in section 9, configurations with generic a and b are parametrized only by sequences of distinct integers $0 < k_1 < k_2 < \cdots < k_n$.

Even family of periodic solutions emerges when a and b become half-integers, such that

$$a = l + 1/2, \quad b = m + 1/2, \quad l, m \in \mathbb{N}.$$

In this case, eigenvalues λ_i in (83) overlap, and some of them become degenerate, with twodimensional eigenspaces¹⁴. Now, there are two types (degenerate and non-degenerate) of eigenfunctions $\varkappa_i(z)$:

$$P_i^{(-l,-m)}(\cos 2z,\zeta_i)\sin^{1/2-l}z\cos^{1/2-m}z, \quad \lambda_i = (2i+1-l-m)^2, \quad \frac{l+m}{2} \le i < l+m \\ P_i^{(l,m)}(\cos 2z)\sin^{l+1/2}z\cos^{m+1/2}z, \qquad \lambda_i = (2i+1+l+m)^2, \quad i \ge 0$$
(84)

¹³In general, sequence of *n* Darboux transformations of the chain (33), where $A_i = \partial_z - (a+i) \cot z + (b+i) \tan z$, $i = 0, 1, \ldots$, results in the shift of parameters $a \to a + n$, $b \to b + n$ in the potential (82).

¹⁴Degeneration occurs also for integer a and b. There, number of degenerate states, as we saw in the previous section, is infinite.

which are both trigonometric-rational up to the common factor $(\sin z \cos z)^{1/2}$. The first type, $P_i^{(-l,-m)}(x,\zeta)$ is a polynomial in x which depending also on a free parameter ζ . Such polynomials are called para-Jacobi Polynomials [11], [8]. Solutions of the second type involve ordinary Jacobi polynomials $P_i^{(l,m)}$ and are non-degenerate. In difference from the latter, the para-Jacobi polynomials are not orthogonal.

Thus, the even family of the static vortex street configurations parametrized by sequences of continuous free parameters as well as sequences of integers can be constructed with the help of (78) and (84) for half-integer a, b.

The simplest non-trivial example of a member of the even family corresponds to l = m = 1. Here a = b = 3/2, $P_1^{(-1,-1)}(\cos 2z, \zeta_1) = \cos 2z + 1 + \zeta_1$ and from (84) we get

$$\psi_1 = \varkappa_1 = \frac{\cos 2z + 1 + \zeta_1}{(\sin z \cos z)^{1/2}}.$$

Note that by the scaling $z \to \epsilon z$, $\psi_1 \to -\psi_1/(2\epsilon^{3/2})$ and change of parameter $\zeta_1 = -2(1+\epsilon^2 t_1)$, in the $\epsilon \to 0$ limit, we obtain $\psi_1 = (z^2 + t_1)/z^{1/2}$. This is ψ_1 (see (55)) of the even bi-spectral family of section 6.

In general, the terminating sequences of section 6 are the rational (times $z^{1/2}$) limits of the trigonometric sequences of the case a = b = m + 1/2

$$\psi_m = \mathcal{W}[\varkappa_m], \quad \psi_{m+1} = \frac{\mathcal{W}[\varkappa_m, \varkappa_{m+1}]}{\mathcal{W}[\varkappa_m]}, \dots, \psi_{2m-1} = \frac{\mathcal{W}[\varkappa_m, \varkappa_{m+1}, \dots, \varkappa_{2m-2}, \varkappa_{2m-1}]}{\mathcal{W}[\varkappa_m, \varkappa_{m+1}, \dots, \varkappa_{2m-2}]}, \quad (85)$$

where

$$\varkappa_{m+i} = (\sin z \cos z)^{1/2-m} P_{m+i}^{(-m,-m)}(\cos 2z, \zeta_{m+i}), \quad i = 0, 1, \dots, m-1$$
(86)

Here, polynomials $P_n^{(-m,-m)}$ are the special l = m case of the para-Jacobi polynomials ("para-Gegenbauer" polynomials)¹⁵

$$P_n^{(-m,-m)}(x,\zeta) = \frac{(-2)^n (n-m)! n!}{(2n-2m)!} \sum_{k=0}^{n-m} \frac{(-1)^{n-k} (2n-2m-k)!}{k! (n-m-k)! (n-k)!} \left(\frac{1+x}{2}\right)^{n-k} + \zeta \frac{(-2)^n (2n-2m+1)! (2m-n-1)!}{(n-m)!} \sum_{2(n-m)+1}^{n} \frac{(-1)^{n-k} (k-n+m-1)!}{k! (k+2m-2n-1)! (n-k)!} \left(\frac{1+x}{2}\right)^{n-k}$$

When we re-scale $z \to \epsilon z$, the eigenvalues scale as $\lambda_i \to \epsilon^2 \lambda_i$, and in the rational limit we have confluent eigenvalues, all going to $\lambda_i \to \eta = 0$. According to section 4 (see eqs. (41), (42)) in this limit the Wronskian representation of the Darboux transforms becomes

$$\psi_{m+i} = \frac{\tau_{i+1}}{\tau_i}, \quad \tau_i = \mathcal{W}[\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_i], \quad \tau_0 = 1,$$
(87)

where now

$$H_0\mathcal{X}_1(z) = 0, \quad H_0\mathcal{X}_{i+1}(z) = c_i\mathcal{X}_i(z), \quad i = 1\dots m-1$$

Here $H_0 = -\partial_z^2 + \frac{m^2 - 1/4}{z^2}$ is the rational limit of the trigonometric Schrödinger operator (82) with a = b = m + 1/2, obtained by the scaling $z \to \epsilon z$, $H_0 \to \epsilon^{-2} H_0$. In other words, $\mathcal{X}_i(z)$ are Laurent polynomials in $z^{1/2}$ that solve the chain

$$\mathcal{X}_{1} = z^{1/2-m}(z^{2m} + t_{m}), \quad -\mathcal{X}_{i+1}'' + \frac{m^{2} - 1/4}{z^{2}}\mathcal{X}_{i+1} = c_{i}\mathcal{X}_{i}, \quad i = 1, 2, \dots, m-1,$$
(88)

¹⁵They satisfy equation $((1-x^2)\partial_x^2 + 2(m-1)x\partial_x + n(n-2m+1))P_n^{(-m,-m)}(x,\zeta) = 0$ which reduces to the standard hypergeometric form by the change of variable w = (1-x)/2.

where c_i are arbitrary constants. Although two constants of integration appear at each step of the chain, only one of them is essential. The chain terminates at the (m-1)th step, since the logarithmic term appears at the *m*th step. Function \mathcal{X}_i is the $\epsilon \to 0$ limit of a linear combinations of *i* first eigenfunctions (86):

$$\epsilon^{3/2-2i-m}\varkappa_m(\epsilon z,\zeta_m),\quad \epsilon^{3/2-2i-m}\varkappa_{m+1}(\epsilon z,\zeta_{m+1}),\ldots \epsilon^{3/2-2i-m}\varkappa_{m+i-1}(\epsilon z,\zeta_{m+i}),$$

where the (j-1)th free parameter ζ_{m+j} is a linear combination of

1, $\epsilon^{2m} t_m$, $\epsilon^{2m+2} t_{m+1}$, $\epsilon^{2m+4} t_{m+1}$, $\ldots \epsilon^{2(m+j-1)} t_{m+j}$.

Coefficients of linear combinations are constants determined by the condition of existence of the limit.

For example, in m = 2 case, corresponding to the terminating sequence (56), we have

$$\begin{aligned} \mathcal{X}_1 &= \lim_{\epsilon \to 0} \epsilon^{-5/2} \varkappa_2(\epsilon z, 2 - 2\epsilon^4 t_2), \\ \mathcal{X}_2 &= \lim_{\epsilon \to 0} \epsilon^{-9/2} \left(\varkappa_3 \left(\epsilon z, -\frac{1}{2} - \frac{3}{2}\epsilon^4 t_2 - \frac{1}{2}\epsilon^6 t_3 \right) - 3\varkappa_2(\epsilon z, 2 - 2\epsilon^4 t_2) \right). \end{aligned}$$

Concluding this section we note that, in the trigonometric case, sequences of the static periodic configurations (85) can be continued with the help of solutions involving the ordinary Jacobi polynomials (see (84)). In other words, in difference from the rational limit, trigonometric sequences are not terminating. However, similarly to the rational case, these sequences depend only on finite numbers of free continuous parameters.

12 Time-Dependent Darboux Transformations, Calogero-Moser Systems and Locus Configurations

As we saw in section 3, all possible static configurations of finite number of vortices of strengths $Q_i = \pm 1$ are given by roots of the Adler-Moser polynomials. In the present section we will demonstrate that sequences of soliton τ -functions (77,78) define all possible vortex street patterns with $Q_i = \pm 1$. To do this, we will introduce dynamical systems that are completely integrable in the $\Lambda = 1$ case, and whose fixed points coincide with static and translating configurations of the point vortices. These systems are related to Darboux transforms of the time-dependent free Schrodinger (or heat) equation.

In more detail, we consider the imaginary-time Schrodinger equation with time-dependent potential

$$\frac{\partial \psi(z,t)}{\partial t} = H\psi(z,t), \quad H = -\partial_z^2 + u(z,t).$$
(89)

Here, H can be presented in the form

$$H = A^* A + (\log \varkappa)_t \,, \tag{90}$$

where subscript t stands for time derivative. In eq. (90), $\varkappa = \varkappa(z,t)$ is one of solutions of (89). The first-order differential operators A, A^* are of the same form (28) as those involved

into the time-independent Darboux transformation. Permuting factors in (90), we get the new Schrödinger operator

$$\hat{H} = AA^* + (\log \varkappa)_t.$$

Multiplying (90) by A from the left, we see that $AH = \hat{H}A - A_t$, and therefore $\hat{\psi} = A\psi$ solves the new equation

$$\frac{\partial \psi}{\partial t} = \hat{H}\hat{\psi}, \quad \hat{H} = -\partial_z^2 + \hat{u}(z,t).$$

The new potential $\hat{u}(x,t)$ and solution $\hat{\psi}$ are given by the same formulae as in the case of the time-independent Darboux transformations, i.e.

$$\hat{\psi} = A\psi = \psi' - (\log \varkappa)'\psi, \quad \hat{u} = u - 2(\log \varkappa)''.$$
(91)

Now, we take the free Schrödinger operator $H_0 = -\partial_z^2$. The corresponding imaginary-time Schrödinger (or inverse heat) equation admits polynomial solutions p(z, t)

$$p_t = -p'', \quad p = \prod_{i=1}^l (z - z_i(t)),$$

where $z_i(t)$ obey the dynamical equations

$$\frac{dz_i}{dt} = \sum_{j=1, j \neq i}^l \frac{2}{z_i - z_j}, \quad i = 1 \dots l$$

Using p as a seed function, $\varkappa = p$ in (91), we Darboux transform another polynomial solution $\psi, \psi_t = -\psi''$ into the rational solution

$$\hat{\psi} = \frac{q}{p}, \quad q = \psi' p - p' \psi. \tag{92}$$

Then, the Schrödinger operator acquires the potential $\hat{u} = -2(\log p)''$, and the new equation $\hat{H}\hat{\psi} = \hat{\psi}_t$ writes, in terms of p and q, as

 $\left(-\partial_z^2 - 2(\log p)''\right)[q/p] = \left[q/p\right]_t,$

or equivalently

$$qp_t - pq_t = q''p - 2p'q' + p''q.$$
(93)

Thus, we incorporated dynamics to the Tkachenko equation (9). It is not difficult to see that further iterations of Darboux transformations will lead to equations of the same form for consecutive transforms of polynomials.

Equation (93) results in the following dynamical system for roots of p and q, denoted as z_1, \ldots, z_l and z_{l+1}, \ldots, z_{l+m} respectively:

$$\frac{dz_i}{dt} = 2\sum_{j=1, j \neq i}^{N} \frac{Q_j}{z_i - z_j}, \quad i = 1...N, \quad N = l + m,$$
(94)

where negative and positive charges of equal magnitude $Q_i = \pm 1$ are attached to roots of q and p correspondingly. If instead of polynomials p and ψ in (92) we take pe^{kz-k^2t} and ψe^{kz-k^2t} , we will get the Galilean transform of (94)

$$\frac{dz_i}{dt} = 2k + 2\sum_{j=1, j \neq i}^{N} \frac{Q_j}{z_i - z_j}.$$
(95)

Although this system seems to resemble that of moving vortices (4), in fact, it is quite different: In contrast to (4), the system (95) contains dz_i/dt rather than complex conjugates $d\bar{z}_i/dt$. As a consequence, a systems of N vortices is a hamiltonian system of dimension 2N, while system of N particles (95) has dimension N and is not hamiltonian. Eq. (95) has several (quite artificial) physical interpretations. For example, we can interpret it as system of sinks/sources moving with the flow of the 2d ideal fluid. This and other interpretations make sense only when all z_i move along the same line (e.g. all z_i are real), except cases when all particles move with equal velocities, i.e. at fixed points of (95). These fixed points coincide with translating and static configurations (5,6) of systems of vortices.

Dynamics can be also incorporated into the bi-linear equation (8) for arbitrary Λ as

$$qp_t - \Lambda pq_t = q''p - 2\Lambda p'q' + \Lambda^2 p''q.$$

The corresponding system has the form (94), where now $Q_i = -1$ for roots of p, and $Q_i = \Lambda$ for roots of q.

System (94), with arbitrary Q_i , can be embedded into a hamiltonian system of newtonian particles interacting pairwise through inverse square potentials: Taking time derivative of eq. (94), then eliminating first order time derivatives by substituting them from (94), we get

$$\frac{d^2 z_i}{dt^2} = -4 \sum_{j=1, j \neq i}^N \frac{\mathcal{Q}_j(\mathcal{Q}_i + \mathcal{Q}_j)}{(z_i - z_j)^3}$$
(96)

(for details see e.g. [23] or [32] and references therein). For $\Lambda = 1$, the sum of two charges vanishes $Q_i + Q_j = 0$, when *i* and *j* belong to different species, and (96) decouples into two non-interacting sub-systems

$$\frac{d^2 z_i}{dt^2} = -\sum_{j: j \neq i} \frac{8}{(z_i - z_j)^3},$$

one consisting of charges $Q_i = -1$ at z_1, \ldots, z_l and another with $Q_i = 1$ at z_{l+1}, \ldots, z_{l+m} . These are completely integrable Calogero-Moser systems [27].

There is another case of decoupling, this time for three species of symmetrically situated particles: One particle of strength a at the origin z = 0, 2l particles of strength -1 at $\pm z_1, \ldots, \pm z_l$ and 2m particles of strength 1 at $\pm z_{l+1}, \ldots, \pm z_{l+m}$. Here, due to the symmetry, we have reduction from 2(l+m) + 1 to l+m degrees of freedom and decoupling into the two BC-type Calogero-Moser systems

$$\frac{d^2 z_i}{dt^2} = -\frac{(2a\pm 1)^2}{z_i^3} - \sum_{j:j\neq i} \left(\frac{8}{(z_i-z_j)^3} + \frac{8}{(z_i+z_j)^3}\right)$$

of l and m particles respectively (for review on Calogero-Moser systems related to different root systems see e.g. [31]). This case corresponds to the Darboux-transformed imaginary-time Schrödinger equation with

$$H_0 = -\partial_z^2 + \frac{a(a-1)}{z^2}.$$

Initial equation $H_0\psi = \psi_t$ has quasi-polynomial solutions $\psi = z^a \prod_{i=1}^l (z^2 - z_i(t)^2)$, and the Darboux transforms built from these solutions are quasi-rational. When *a* is half-integer, fixed points of system (94) are configurations of even bi-spectral family from section 6.

Let us now return to the main topic of this section and consider equilibrium configurations of vortices. For these configurations $d^2z_i/dt^2 = 0$. Then form (96) it follows that the equilibrium conditions (6) imply the following N locus conditions

$$\sum_{j=1, j\neq i}^{N} \frac{\mathcal{Q}_j(\mathcal{Q}_i + \mathcal{Q}_j)}{(z_i - z_j)^3} = 0, \quad i = 1 \dots N,$$

i.e. the locus conditions are necessary conditions of equilibrium. In the case of two species with $Q_i = \pm 1$, locus conditions decouple, and we have the separate set of locus conditions¹⁶ for each specie

$$\sum_{j:j\neq i} \frac{1}{(z_i - z_j)^3} = 0,$$
(97)

l conditions for the first and *m* conditions for the second specie respectively. It turned out $[2]^{17}$ that set of solutions of (97) is not empty only if the number of particles the locus is formed of is a triangular number n(n + 1)/2. Then, it was shown [1] that for this number, roots of the Adler-Moser polynomials P_n constitute complete solution of (97). As we saw in section 3, in systems of vortices in static equilibrium l = n(n - 1)/2 and m = n(n + 1)/2 particles form two loci that are roots of two consecutive Adler-Moser polynomials $P_{n-1}(z, t_1, \ldots, t_{n-2})$ and $P_n(z, t_1, \ldots, t_{n-1})$.

Conditions (97) also must hold for each specie of translating configurations (5) with $Q_i = \pm 1$. We recall that for translating configurations deg $p = \deg q$, i.e. l = m (see section 8). This means that when $k \neq 0$, p and q are Adler-Moser polynomials of the same degree, but with different sets of parameters: If $p = P_n(z, t_1, \ldots, t_{n-1})$, then $q = P_n(z + s_0, s_1, \ldots, s_{n-1})$, where s_0, \ldots, s_{n-1} are functions of t_1, \ldots, t_{n-1} and k.

We also recall that, according to section 8, the rational Baker-Akhieser function $\psi = C(k)(q/p)e^{kz}$ determines translating configuration. It is an eigenfunction of a Schrodinger operator that is obtained by a sequence of Darboux transformations from free Schrodiger operator, i.e. $H\psi = k^2\psi$, where $H = -\partial_z^2 - 2(\log p)''$. From this we get the bilinear relation for p and q:

$$p''q - 2p'q' + pq'' + 2k(q'p - p'q) = 0.$$
(98)

This relation coincides with the Tkachenko equation for translating configurations. Set of its solutions contains all translating patterns with $Q_i = \pm 1$. The Baker-Akhieser function exists

¹⁶It is worth mentioning, that the theory of algebraically integrable systems also considers loci with multiple roots, where more general locus conditions are imposed [13, 12].

¹⁷Conditions (97) first appeared in this famous paper by Airault, McKean and Moser, who introduced the term "locus".

for any locus of p and is unique for a given locus (see e.g. [13, 12] and references therein). Since the locus conditions are necessary conditions of an equilibrium, we come to the conclusion that all possible translating configurations consisting of a finite number of vortices with $Q_i = \pm 1$ are determined by the Baker-Akhieser functions and are given by (68).

In the trigonometric (periodic) case, for each of two species of opposite charges the locus conditions write as

$$\sum_{j: j \neq i} \frac{\cos(z_i - z_j)}{\sin^3(z_i - z_j)} = 0.$$

It is known (see e.g. [13, 12] and references therein) that all trigonometric locus configurations are given by roots of soliton τ -functions of the KdV hierarchy. Therefore any pair p and q that does not have common roots and solves the periodic Tkachenko equation

$$p''q - 2p'q' + pq'' + (l-m)^2 pq = 0, \quad p = \prod_{i=1}^{l} \sin(z-z_i), \quad q = \prod_{i=1}^{m} \sin(z-z_{l+i})$$
(99)

consists of soliton τ -functions. These functions are obtained with the help of factorization chain (33) starting from the free Schrödinger operator. In this chain of transformations

$$H_n \psi^{(n)} = \eta_{n+1} \psi^{(n)}, \quad H_n = -\partial_z^2 - 2(\log \tau_n)'', \quad \psi^{(n)} = \frac{\tau_{n+1}}{\tau_n}, \quad \tau_0 = 1,$$

i.e.

$$\left(-\partial_z^2 - 2(\log \tau_n)''\right) \left[\frac{\tau_{n+1}}{\tau_n}\right] = \eta_{n+1} \frac{\tau_{n+1}}{\tau_n}.$$
(100)

Since degree of the leading term of the Laurent expansion of τ_{n+1}/τ_n in exp(iz) equals $d_{n+1}-d_n$, where $d_n = \deg \tau_n$, from (100) we conclude that $\eta_{n+1} = (d_{n+1} - d_n)^2$. Then, rewriting (100) in the bi-linear form, we get the Darboux chain for τ_n :

$$\tau_{n+1}''\tau_n - 2\tau_{n+1}'\tau_n' + \tau_{n+1}\tau_n'' + (d_{n+1} - d_n)^2\tau_{n+1}\tau_n = 0, \quad d_n = \deg \tau_n, \quad \tau_0 = 1.$$
(101)

The periodic analog of the Tkachenko equation (99) is of the same form as the Darboux chain equation (101). Then, taking the above fact and the fact that p and q are necessarily soliton τ -functions (as follows from locus conditions) into account, we come to the conclusion that p and q are soliton τ -functions related by Darboux transformation. Therefore, all possible static street patterns consisting of vortices of two species of opposite strengths are given by (77,78).

For translating streets we use arguments similar to those of the rational case and come to the conclusion that the complete set of translating street patterns with $Q_i = \pm 1$ is determined by (77,81). It is worth mentioning that from the soliton theory it follows that here the relationship between p and q holds

$$q = p(z, \sigma_1, \dots, \sigma_n), \quad p = p(z, \zeta_1, \dots, \zeta_n), \quad (k + ik_j)e^{2i\sigma_j} = (k - ik_j)e^{2i\zeta_j},$$

where $p = \tau_n$ is given by (77,78). Without going into much details, we note that the above relationship can be derived using the Sato formula for the Baker-Akhieser function of the KP hierarchy and the correspondence between Hirota and Wronskian representations of the *n*-soliton τ -functions.

Closing this section, we note that, in contrast to the cases considered above, the $\Lambda = 2$ case is not exceptional as no decoupling into sub-systems of different species takes place there.

13 Conclusions and Open Problems

In the present paper we reviewed developments in application of factorization methods to the theory of point vortex patterns that took place during several last decades. In the $\Lambda = 1$ case, these development have connections with the theory of KdV hierarchy of integrable PDEs and the bi-spectral problem. Some connections with integrable hierarchies have been also found in the $\Lambda = 2$ case [17]. However, this case seems not to have as many exceptional properties: Absence of trigonometric counterparts of the main $\Lambda = 2$ sequences (24,25) makes us think that this should be so.

In more detail: We took the trigonometric analog of the bilinear equation (8)

$$\tau_{n-1}''\tau_n - 2\gamma_n\tau_{n-1}'\tau_n' + \gamma_n^2\tau_{n-1}\tau_n'' + (d_{n-1} - \gamma_n d_n)^2\tau_{n-1}\tau_n = 0,$$
(102)

where $d_n = \deg \tau_n$, $\gamma_{n-1}\gamma_n = 1$, $\gamma_n \in {\Lambda^{-1}, \Lambda}$. In the $\Lambda = 1$ case, sequence of its solutions, which is the trigonometric analog of the main sequence (Adler-Moser polynomials), is given by (77,78) with $k_n = n$. Here, the highest symbol of the *n*th Adler-Moser polynomial and its trigonometric counterpart is of degree n(n + 1)/2, i.e. in the trigonometric case $\tau_n = \xi^{n(n+1)/2} + \alpha_1 \xi^{n(n+1)/2-2} + \cdots + \alpha_n \xi^{-n(n+1)/2}$, where $\xi = \exp(iz)$. For both types of polynomials a new free parameter appears at each step of the factorization chain.

However, when $\Lambda = 2$, non-terminating solutions of the chain (102) with $\tau_n = \xi^{D_n} + \ldots$, are (up to multiplication by constants and shift of z) only $\sin^{D_n} z$, where D_n are degrees of the main sequences $(24,25)^{18}$. We searched for other configurations imposing condition that τ_i do not have common or multiple roots, i.e. for configurations of two distinct species of vortices. By direct solution of (102), we found only short terminating sequences, for example

$$\tau_0 = \xi - \frac{1}{\xi}, \tau_1 = \xi^6 - 4\xi^4 + 5\xi^2 + \frac{5t_1}{\xi^2} - \frac{4t_1}{\xi^4} + \frac{t_1}{\xi^6}, \tau_2 = \xi^5 - 5\xi^3 - t_2\xi + \frac{t_2}{\xi} - \frac{5t_1}{\xi^3} + \frac{t_1}{\xi^5},$$

where $\gamma_0 = 2$. The above example can be continued one more step if we impose condition $t_1 = t_2^2/5$. Then we will get another terminating sequence with $\tau_1 = \tau_1(z, t_2), \tau_2 = \tau_2(z, t_2)$ and $\tau_3 = \tau_3(z, t_2, t_3)$, where deg $\tau_3 = 14$.

Without going into much detail we just note that in such type of sequences τ_1 is a solution of an ODE that is reducible to the hypergeometric form. Then we can apply KWCC transformations (59) to $\psi_i = \tau_i^{\gamma_i}/\tau_{i-1}$, i = 1, 2, to continue the sequence. The transformation can be applied at most twice. Terminating configurations obtained in this way can depend on no more than two non-trivial free parameters (they depend on two parameters when deg $\tau_1 = 6, 8, 10, \ldots$). Here, a natural question arises: If the above procedure generates all possible static vortex streets configurations for two species of vortices with $Q_i \in \{-1, 2\}$?

Concerning the $\Lambda = 2$ case, we also note that methods displayed in this review are inapplicable to both the translating streets and to the translating patterns composed of finite numbers of vortices when $Q_i \in \{-1, 2\}$.

Regarding the static patterns formed by finite number of vortices, we may ask the question: Whether the main sequences (23), (24), (25) for the $\Lambda = 1$ and $\Lambda = 2$ cases are the only possible non-trivial non-terminating sequences of the KWCC transforms. Are there non-trivial sequences (59) with γ_i different from (60,61) that do not terminate?

¹⁸We recall that, according to (63), for the main sequences (24,25) $D_{i+1} - 2\gamma_i D_i + D_{i-1} = 1$, $i \in \mathbb{Z}$ and $D_0 = D_1 = 0$, $\gamma_0 = 2$.

Finally, classification of all stationary vortex patterns and understanding the role of factorization methods in this classification are ultimate questions to address in this subject.

Concluding this review, we would like to mention related problem from the theory of algebraically integrable systems, because our interest in vortex patterns stems from that theory: In the case of odd family of periodic configurations, it is convenient to rewrite the bi-linear chain (101) in terms of homogenous polynomials in two variables, replacing $\tau_n(z) = \prod_{i=1}^{d_n} \sin(z - z_i)$ with

$$\tau_n(X, Y) = R^{d_n} \tau_n(z) = \prod_{i=1}^{d_n} (Y \cos z_i - X \sin z_i),$$

where

$$X = R\cos z, \quad Y = R\sin z.$$

In the new variables, the chain (101) takes the following form

$$\tau_n \Delta \tau_{n+1} - 2(\nabla \tau_{n+1} \cdot \nabla \tau_n) + \tau_{n+1} \Delta \tau_n = 0, \quad \tau_0 = 1,$$
(103)

where Δ and ∇ stand for two-dimensional laplacian and gradient respectively. This chain is called harmonic [5], because at the first step $\Delta \tau_1 = 0$, i.e. τ_1 is a harmonic function, and $\psi_n = \tau_{n+1}/\tau_n$ are natural generalizations of harmonic functions¹⁹. The harmonic chain can be generalized to any number of dimensions: Non-terminating solutions of (103) are τ -functions for potentials of algebraically integrable Schrodinger operators (see e.g. [5], [12]). One can view (103) as a generalization of the Darboux chain (33), that starts from the free Schrodinger operator²⁰, to any number of dimensions. In one dimension, the Adler-Moser polynomials constitute complete set of non-terminating solutions of the chain. In two dimensions, the soliton related solutions exhaust all non-terminating solutions in the class of homogeneous polynomials: Family of two-dimensional solutions consists of an infinite number of branches labelled by $0 < k_1 < k_2 < \ldots$, rather than a single sequence in one dimension. Complete classification of non-terminating solutions is an open hard problem (see e.g. [5, 12])²¹.

References

- Adler M., Moser J., On a class of Polynomials connected with the Korteweg-de-Vries Equation, Comm. Math. Phys. 1978. V. 61. P. 1-30.
- [2] Airault.H., McKean H., Moser J., Rational and elliptic solutions of the KdV equation and related many body problem, Commun. Pure Appl. Math. 1977. V. 30. P. 95-148.
- [3] Aref H., Point vortex dynamics: a classical mathematics playground. J. Math. Phys. 48, (2007) 065401. (doi:10.1063/1.2425103)

¹⁹For example, ψ_n satisfy quadrature identities in "harmonic" quadrature domains (see e.g. [24, 25] and references therein).

²⁰Chain that starts from $u_0 \neq 0$ has the form $\tau_n \Delta \tau_{n+1} - 2(\nabla \tau_{n+1} \cdot \nabla \tau_n) + \tau_{n+1} \Delta \tau_n - u_0 \tau_n \tau_{n+1} = 0$, $\tau_0 = 1$. ²¹It is conjectured that families listed in [12] constitute complete solution of the problem.

- [4] Aref H., Vortices and polynomials, Fluid Dyn. Res. 39, 5–23. (2007) (doi:10.1016/ j.fluiddyn.2006.04.004)
- [5] Yu. Berest, Huygens' Principle and the Bispectral Problem, CRM Proceedings and Lecture Notes, volume 14, American Mathematical Society, Providence, RI, pp. 11-30, https://doi.org/10.1090/crmp/014/02
- [6] Berest, Y., Loutsenko, I., Huygens' Principle in Minkowski Spaces and Soliton Solutions of the Korteweg-de Vries Equation, Comm. Math. Phys. 190, 113-132 (1997)
- [7] Burchnall J.L., Chaundy T.W., A set of differential equations which can be solved by polynomials, Proc. Lonlon. Math. Soc. 1929-1930. V. 30. P. 401-414.
- [8] B. Bagchi, Y. Grandati, C. Quesne, Rational extensions of the trigonometric Darboux-Pöschl-Teller potential based on para-Jacobi polynomials, J. Math. Phys. 56, 062103 (2015)
- [9] Bartman A.B., A new interpretation of the Adler-Moser KdV polynomials: interaction of vortices, In Nonlinear and turbulent processes in physics, vol. 3 (ed. RZ Sagdeev), (1984) pp. 1175–1181. New York, NY: Harwood Academic.
- [10] M.M. Crum, Associated Sturm-Liouville systems, Quart. J. Math. Oxford 6 (1955), 121– 127.
- [11] F. Calogero and G. Yi, Can the general solution of the second-order ODE characterizing Jacobi polynomials be polynomial?, J. Phys. A 45, 095206 (2012)
- [12] Chalych O., Algebro-geometric Schrodinger operators in many dimensions, Phil. Trans. R. Soc. A (2008) 366, 947–971
- [13] Chalykh O., Feigin M., Veselov A., Multidimensional Baker-Akhiezer Functions and Huygens' Principle., Comm Math Phys 206, 533-566 (1999)
- [14] Clarkson PA. Vortices and polynomials, Stud. Appl. Math. 123, 37–62. (2009) (doi:10.1111/j.1467-9590.2009.00446.x)
- [15] P. Gaillard and V.B. Matveev, Wronskian and Casorati determinant representations for Darboux-Pöschl-Teller potentials and their difference extensions, J. Phys. A: Math. Theor. 2009, 42 404009
- [16] Adrian D. Hemery and Alexander P. Veselov, Periodic Vortex Streets and Complex Monodromy, SIGMA 10 (2014), 114
- [17] Demina, M. and Kudryashov, N., Point Vortices and Polynomials of the Sawada Kotera and Kaup – Kupershmidt Equations, Regul. Chaotic Dyn., 2011, vol. 16, no. 6, pp. 562–576.
- [18] Demina, M. and Kudryashov, N., Vortices and Polynomials: Non-Uniqueness of the Adler -Moser Polynomials for the Tkachenko Equation, J. Phys. A, 2012, vol. 45, no. 19, 195205, 12 pp.

- [19] J.J. Duistermaat and F.A. Grünbaum, Differential equations in the spectral parameters, Commun. Math. Phys. 103 (1986), 177–240.
- [20] Krishnamurthy Vikas S., Wheeler Miles H., Crowdy Darren G. and Constantin Adrian, A transformation between stationary point vortex equilibria, Proc. R. Soc. A. (2020) 4762020031020200310
- [21] J.E. Lagnese and K.L. Stellmacher, A method of generating classes of Huygens' operators, J. Math. Mech. 17 (1967), 461-472
- [22] Igor Loutsenko, Equilibrium of charges and differential equations solved by polynomials, 2004 J. Phys. A: Math. Gen. 37 1309
- [23] Loutsenko, I., Integrable Dynamics of Charges Related to the Bilinear Hypergeometric Equation, Commun. Math. Phys. 242, 251–275 (2003). https://doi.org/10.1007/s00220-003-0944-z
- [24] I. Loutsenko, The variable coefficient Hele-Shaw problem, Integrability and quadrature identities, Comm. Math.Phys 268, (2006), no.2, 465-479
- [25] Igor Loutsenko and Oksana Yermolayeva, On integrability and exact solvability in deterministic and stochastic Laplacian growth, Math. Model. Nat. Phenom. 15 (2020) 3
- [26] V. B. Matveev and M. A. Salle, Darboux Transformations and Solitons, Springer-Verlag, Berlin, 1991
- [27] J. Moser, Three integrable Hamiltonian systems connected with isospectral deformations, Adv. Math. 16 (1975), 197-220
- [28] O'Neil KA., Stationary configurations of point vortices, Trans. Am.Math. Soc. 302, 383–425 (1987). (doi:10.2307/2000850)
- [29] O'Neil KA., Minimal polynomial systems for point vortex equilibria, Phys. D 219, 69–79 (2006). (doi:10.1016/j.physd.2006.05.012)
- [30] O'Neil, K.A., Cox-Steib, N., Generalized Adler-Moser and Loutsenko polynomials for point vortex equilibria, Regul. Chaot. Dyn. 19, 523–532 (2014), https://doi.org/10.1134/S1560354714050013
- [31] M. A. Olshanetsky and A. M. Perelomov, Classical integrable finite-dimensional systems related to Lie algebras, Phys. Rep. 71 (1981), 313-400
- [32] Robert L. Sachs, On the integrable variant of the boussinesq system: Painlevé property, rational solutions, a related many-body system, and equivalence with the AKNS hierarchy, Physica D, Vol. 30, Issues 1–2, February–March 1988, Pages 1-27
- [33] T.J. Stieltjes, Sur certains polynômes qui vérifient une équation différentielle linéaire du second ordre et sur la théorie des fonctions de Lamé, Acta Math. 8 (1885) 321–326; Oeuvres Complètes 1, 434–439

- [34] G. Szegö, Orthogonal Polynomials, American Mathematical Society, 1975
- [35] Tkachenko V.K., Studying the Violations of the Superfluidity of Helium in Broad Capillaries under the Influence of Heat Flow, 1964 Unpublished thesis, Institute of Physical Problems, Moscow