

Capturing the Variability of the Nocturnal Boundary Layer through Localized Perturbation Modeling

Amandine Kaiser,^a Nikki Vercauteren,^{a,b} Sebastian Krumscheid,^c

^a *Department of Geosciences, University of Oslo, Oslo, Norway*

^b *Institut für Geophysik und Meteorologie, Universität zu Köln, Cologne, Germany*

^c *Department of Mathematics, Karlsruhe Institute of Technology, Karlsruhe, Germany*

Corresponding author: Amandine Kaiser, amandine.kaiser@geo.uio.no

ABSTRACT: A single-column model is used to investigate regime transitions within the stable atmospheric boundary layer, focusing on the role of small-scale fluctuations in wind and temperature dynamics and of turbulence intermittency as triggers for these transitions. Previous studies revealed abrupt near-surface temperature inversion transitions within a limited wind speed range. However, representing these transitions in numerical weather prediction (NWP) and climate models is a known difficulty. To shed light on boundary layer processes that explain these abrupt transitions, the Ekman layer height and its correlation with regime shifts are analyzed. A sensitivity study is performed with several types of perturbations of the wind and temperature tendencies, as well as with the inclusion of intermittent turbulent mixing through a stochastic stability function, to quantify the effect of small fluctuations of the dynamics on regime transitions. The combined results for all tested perturbation types indicate that small-scale phenomena can drive persistent regime transitions from very to weakly stable regimes, but for the opposite direction, no evidence of persistent regime transitions was found. The inclusion of intermittency prevents the model from getting trapped in the very stable regime, thus preventing the so-called "runaway cooling", an issue for commonly used short-tail stability functions. The findings suggest that using stochastic parameterizations of boundary layer processes, either through stochastically perturbed tendencies or parameters, is an effective approach to represent sharp transitions in the boundary layer regimes and is, therefore, a promising avenue to improve the representation of stable boundary layers in NWP and climate models.

1. Introduction

The stably stratified atmospheric boundary layer (SBL) is commonly classified into two stability regimes. The first one is a weakly stable regime, which corresponds to strong wind, weak stratification, and relatively continuous turbulence in both space and time, while the second one is very stable and is characterized by low wind, strong stratification, and weak or intermittent turbulence (Mahrt 2014). In the very stable regime, turbulence is not strong enough to mix the whole layer, which leads to the decoupling of the lower from the higher levels (Derbyshire 1999; Van de Wiel et al. 2012; Acevedo et al. 2015). As a result, the potential temperature difference between the surface and a higher level is large. In contrast, in the weakly stable regime, turbulence is sufficiently strong to keep the layers coupled, and therefore, the potential temperature difference is low (Vignon et al. 2017).

A conceptual model for near-surface temperature inversions showed that through feedbacks between the surface thermal processes and turbulent fluxes, the relationship between the temperature inversion strength and wind speed can be non-monotonic and exhibit a Z-shape dependency (Van de Wiel et al. 2017). For low wind speeds, the Van de Wiel et al. (2017) model has only one solution with a strong inversion. In contrast, for high wind speeds, the inversion strength is small. For intermediate wind speeds, when the thermal coupling between the surface and the atmosphere is weak, three inversions of strength ranging from strong to weak are possible solutions of the conceptual model, leading to the Z-shape dependency of the inversion on the wind speed. This non-linear dependency is commonly interpreted as a sign that the SBL is bistable, i.e., two coexisting stable equilibria exist for a fixed wind speed forcing, separated by an unstable equilibrium. This Z form has, for example, been observed empirically at Dome C, Antarctica, by Vignon et al. (2017), and the existence of an unstable equilibrium has been hypothesized to explain the scatter found in the observational data in a certain range of intermediate wind speeds (Van de Wiel et al. 2017). In principle, bistability should be analyzed using a fixed forcing parameter. However, the previous studies used a reference height as a so-called velocity crossing level, where the wind was empirically found to remain rather constant throughout a night (van Hooijdonk et al. 2015). To analyse the impact of this reference height on the observed nonlinearity, we study the growth of the Ekman layer under different forcing scenarios. By studying the growth of the

Ekman layer, the goal is to detect inherent nonlinearity in the dynamics of the SBL. In particular, field analyses highlighted the existence of a wind threshold where turbulence intensity increases rapidly (Sun et al. 2012), and such rapid increase could lead to a nonlinear growth of the layer. The Ekman layer height growth is hence chosen as a proxy for the SBL dynamics and its dependence of the forcing wind speed is analysed.

Transitions between the two SBL regimes are frequently observed in SBL dynamics worldwide (Abraham and Monahan 2019b). However, the representation of regime structures and transitions in numerical weather prediction (NWP) and climate models is inadequate, primarily due to coarse resolution (both vertical and horizontal) and an imperfect understanding of the various physical processes governing the SBL (Holtslag et al. 2013; Sandu et al. 2013). A typical solution to include turbulent mixing due to unresolved processes is to enhance the mixing (Sandu et al. 2013), but this solution blurs the sharp transitions (Baas et al. 2017). The lack of understanding of the SBL has so far prevented the emergence of a better solution. An imperfect understanding of processes leads to model uncertainties, and in fact numerous approaches have been proposed to incorporate model uncertainty in NWP. A particularly attractive solution is to use stochastic parameterization (Palmer 2001; Christensen 2020). Stochastic parameterizations recognize the importance of model uncertainty to accurately represent the mean state, and in particular they have the potential to induce regime transitions in cases where a system exhibits multiple coexisting equilibria, making their incorporation crucial for a more precise representation of the mean state of the system (Berner et al. 2017). Stochastic parameterization approaches are divided into randomized tendencies and stochastic parameterizations. The European Center for Medium-Range Weather Forecasts (ECMWF) uses, for example, a scheme that stochastically perturbs tendencies. The perturbations are added to the model a posteriori, i.e., after it has been tuned in the deterministic mode (Palmer 2017). Adding the stochasticity after the tuning has the potential to detune it. Alternatively, a priori perturbation of uncertain parameters can be used. In this case the model is tuned with the stochastic scheme included. They are called stochastic parameterization models and do not have the detuning issues as the perturbed tendency ones. In this work, the impact of perturbed tendencies and parameterization on regime transitions will be analyzed.

In a previous investigation conducted by Kaiser et al. (2024), a modified version of the conceptual model proposed by Van de Wiel et al. (2017) was employed as a representative model to examine the sensitivity of the polar stable boundary layer (SBL) to small-scale fluctuations and explore the influence of model uncertainty on the average state of the boundary layer. The findings of the aforementioned study demonstrated that random perturbations can induce transitions in the SBL regimes and impact the timing of these transitions. The model by Van de Wiel et al. (2017) is a strongly idealized model with no spatial dimension, and hence does not allow for spatially localized perturbations. The current study extends the research by Kaiser et al. (2024) using a single-column model with spatial and time dimensions. The added spatial dimension allows for more degrees of freedom for the nonlinear growth of initially small perturbations, and thereby provides a greater opportunity to capture the real-world complexities and variability of the SBL. An example where spatially localized phenomena are non-linearly amplified is given by Lan et al. (2022). The authors showed that small-scale wind profile distortions can cause very to weakly stable regime transitions. Therefore, this study analyzes the effect of perturbed wind dynamics in the single-column model. Scenarios where the perturbations are added above or below the Ekman layer are studied separately. Sun et al. (2012) showed that turbulence intermittency can occur when top-down turbulent events diffuse into an environment characterized by weak turbulence. Hence, top-down mixing events can be important feedback processes for recoupling the SBL. The direct numerical simulations of Donda et al. (2015) support the addition of perturbations to the model, as they showed that transient perturbations can be sufficient to recover a turbulent regime, i.e., for transitions from very to weakly stable regimes. Intermittent turbulence events are another possible set of processes that can trigger transitions (Mahrt 2014). These events have been observed to play a significant role in turbulent transport, particularly in highly stable conditions (Acevedo and Fitzjarrald 2003). Hence, we study the effect of including intermittent events through a stochastic stability correction on regime transitions.

Vertical mixing associated with surface heterogeneity, gravity waves, or mesoscale variability is commonly not represented in NWP models (Sandu et al. 2013). One approach to enhancing the mixing properties of the atmospheric boundary layer in NWPs is using long-tail stability functions. The issue with these types of stability functions is that they produce too much mixing

compared to observations (Chechin et al. 2019), resulting in a suppression of the backfolding of the before mentioned Z curve when plotting the temperature inversion over wind speed (Van de Wiel et al. 2017; Baas et al. 2017). On the other hand, short-tail stability functions, which suppress turbulence in the very stable case, are more consistent with turbulence theory (Chechin et al. 2019), but they can cause "runaway cooling" (Mahrt 1998; Steeneveld 2014; Kähnert 2022). One suggestion on how to circumvent the issues with stability functions was made by Maroneze et al. (2023). There, the authors suggest an adaption of a classic atmospheric boundary layer parameterization, which does not use a prescribed stability function. Instead, a turbulent heat flux prognostic equation ensures the stratification dependence on other characteristics of the mean and turbulent flows. While this approach is promising, it relies on additional equations in NWP. However, transitions from very to weakly stable regimes do not seem to be determined by internal or external state variables in a predictable manner (Abraham and Monahan 2019b). This suggests that parameterizations that facilitate such transitions in weather and climate models may need to explicitly incorporate stochastic features. Moreover, including stochasticity is one way to account for unresolved variability due to coarse resolution that could be highly relevant for triggering large-scale transitions. Therefore, in the present work, we suggest using a short-tail stability function with added stochasticity to prevent the model from getting "stuck" in the very stable regime. A particular useful example of such a randomized stability function was developed by Boyko and Vercauteren (2023). Abraham et al. (2019) created a stochastic model featuring transition probabilities dependent on stratification, while Ramsey and Monahan (2022) formulated a data-driven stochastic differential equation model to capture the variability of temperature inversion.

To study transitions in the SBL and specifically analyse the impact of localized perturbations on transitions, the paper will answer the following three questions:

1. Are the sharp regime transitions also expressed as a nonlinear growth of an Ekman layer?
2. What types of perturbation of the dynamics are particularly relevant triggers for regime transitions?

3. Can stochastic modeling be used to represent the impact of small-scale fluctuations effectively in NWP and climate models?

Questions two and three address where models and parameterizations must be improved to represent regime transitions better, which is a particularly relevant consideration for NWP.

In Section 2, the single-column model is introduced and described in detail. As part of this section, the Ekman layer height and its relation to the geostrophic wind and regime transitions is analyzed. Following that, in Section 3, two approaches are used to explore the impact of small-scale perturbations on regime transitions and identify the most impactful variables. First, in section 3a, a sensitivity analysis of the single-column model to small-scale fluctuations in the temperature and wind dynamics for different perturbation sizes is presented. In Section 3a part 1 the perturbations are defined, in Section 3a part 2, the impact of the perturbation size is quantified, and in Section 3a part 3, four different stability scenarios are analyzed in detail. Then, in Section 3b, the Ekman layer height growth is studied for the perturbed model. Lastly, Section 3c studies the impact of a randomized stability function. For this, the stability function, which is used in the rest of the paper, is replaced with the stochastic one, which was defined by Boyko and Vercauteren (2023).

2. Single-column model

a. Model description

To study which quantities can potentially trigger regime transitions in NWP and climate models, an idealized single-column model (SCM) is used, to which perturbations will be added. This model is an adapted version of a classic Reynolds-averaged Navier–Stokes 1.5-order closure model for the atmospheric boundary layer (Stull 1988). It is briefly introduced below, but a more detailed

description is given in Boyko and Vercauteren (2024). The idealized SBL is defined as:

$$\frac{\partial u}{\partial t} = f_c \cdot (v - v_G) + \frac{\partial}{\partial z} \left(K_m(\phi) \frac{\partial u}{\partial z} \right) - N_u \quad (1)$$

$$\frac{\partial v}{\partial t} = f_c \cdot (u_G - u) + \frac{\partial}{\partial z} \left(K_m(\phi) \frac{\partial v}{\partial z} \right) - N_v \quad (2)$$

$$\frac{\partial \theta}{\partial t} = \frac{\partial}{\partial z} \left(K_h(\phi) \frac{\partial \theta}{\partial z} \right) \quad (3)$$

$$\frac{\partial k}{\partial t} = \frac{\partial}{\partial z} \left(K_m(\phi) \frac{\partial k}{\partial z} \right) + K_m(\phi) \left(\left(\frac{\partial u}{\partial z} \right)^2 + \left(\frac{\partial v}{\partial z} \right)^2 \right) - \frac{g}{\theta_0} K_h(\phi) \frac{\partial \theta}{\partial z} - \frac{(\alpha_\epsilon k)^{3/2}}{l_m(\phi)} \quad (4)$$

$$\frac{d\theta_g}{dt} = \frac{1}{c_g} (R_n - H_0) - \kappa_m (\theta_g - \theta_m), \quad t > 0, \quad z = 0 \quad (5)$$

where u and v are the Reynolds averaged horizontal wind components, θ the Reynolds averaged potential temperature, k the Reynolds averaged turbulent kinetic energy (TKE), and θ_g the surface temperature whose evolution is represented through a surface energy balance and modeled using a force-restore method (Stull 1988; Garratt 1994; Acevedo et al. 2021). The thermal capacity of the soil per unit area is given by c_g , $H_0 = \rho c_p \overline{w'\theta'_0}$ is the surface sensible heat flux, where ρ is the air density, c_p is the specific heat of air at constant pressure and θ_0 is a reference temperature. The parameter $\kappa_m = 1.18\omega$ is the soil heat transfer coefficient, with ω being the Earth's angular frequency. The net radiation is given by R_n , and θ_m is the soil temperature below the surface. Additionally, u_G and v_G are the geostrophic wind components, and f_c is the Coriolis parameter. Moreover, g is the gravitational force, and α_ϵ is a modeling constant set to 0.1. To damp inertial oscillations, the relaxation terms

$$N_u = \frac{u - u_G}{\tau_r}, \quad N_v = \frac{v - v_G}{\tau_r}$$

are subtracted from the momentum equations (1) and (2). They nudge the solution towards the geostrophic wind. To ensure only a mild nudging, τ_r is set to five hours. Lastly, the exchange coefficients for momentum K_m and heat K_h are defined as

$$K_m = \alpha l_m \sqrt{k}, \quad K_h = \frac{K_m}{Pr}$$

with a Prandtl number, Pr , set to 1. The mixing length is defined as $l_m = \frac{\kappa z}{\phi(Ri) + \frac{\kappa z}{\lambda}}$ where ϕ is a stability correction function and $\lambda = 2.7 \cdot 10^{-4} \frac{u_G}{|f_c|}$ (Rodrigo and Anderson 2013) and $\kappa = 0.41$ is the von Karman constant. The gradient Richardson number, Ri , is defined as

$$Ri = \frac{\frac{g}{\theta_0} \frac{\partial \theta}{\partial z}}{\left(\frac{\partial u}{\partial z}\right)^2 + \left(\frac{\partial v}{\partial z}\right)^2}. \quad (6)$$

Two different stability functions, ϕ , are considered. The first is

$$\phi = 1 + 12Ri, \quad (7)$$

a short tail stability function, taken from Delage (1997). The second stability function is a data-driven stochastic stability correction initially suggested by Boyko and Vercauteren (2023) and will be used to model random fluctuations in the turbulent mixing within the model. It is randomized in time and discussed in detail in section 3c.

The model equations (1) to (4) are first solved in time, t , with an implicit Euler algorithm and then in space, z , with a Finite Element method. For this, the Python package FEniCS (<https://fenicsproject.org/>) is used. Equation (5) is solved using an explicit Euler algorithm. The same values as in Boyko and Vercauteren (2024) are used for all parameters, unless stated otherwise. All parameter values are summarized in Table 1 for convenience. Further details about the model and how it is numerically approximated can be found in Boyko and Vercauteren (2024).

b. Initial and boundary conditions

The initial and boundary conditions are similar to the ones in Boyko and Vercauteren (2024). They are repeated here for completeness. These initial conditions are used to initialize the non-perturbed runs. The perturbed runs (section 3) are initialized with the steady state, which is defined in section 2c, of the non-perturbed runs.

The initial condition for u is a logarithmic profile $u(z, t_0) = \frac{u^*}{\kappa} \ln\left(\frac{z}{z_0}\right)$ where $u_{init}^* = \sqrt{0.5c_f u_G^2}$ is the initial friction velocity and $c_f = 4 \cdot 10^{-3}$ is a tuning parameter to ensure the wind velocity is close to u_G at the upper boundary. The other wind velocity component, v , is constant and equal to zero

TABLE 1. Default parameters for the model (eqs. (1)–(5)).

Description	Parameter	Value	Unit
Time step size	Δt	10	s
Number of z grid points	N_z	100	-
Roughness length	z_0	0.044	m
Roughness length for heat	z_{0h}	$z_0 \cdot 0.1$	m
Domain height	H	300	m
Latitude	l	40	°
Coriolis parameter	f_c	$2 \cdot 7.27 \cdot 10^{-5} \cdot \sin(l\pi/180)$	°s ⁻¹
Geostrophic wind	u_G	-	ms ⁻¹
Geostrophic wind	v_G	0	ms ⁻¹
Relaxation time scale	τ_r	$3600 \cdot 5$	s
Gravitational acceleration	g	9.81	ms ⁻¹
Reference potential temperature	θ_0	300	K
Dissipation constant	α_ϵ	0.1	-
Angular earth velocity	ω	$2\pi/(24 \cdot 3600)$	°s ⁻¹
Thermal capacity of soil per unit area	c_g	$0.95 \cdot (1.45 \cdot 3.58 \cdot 10^6/2/\omega)^{0.5}$	Jm ⁻² K ⁻¹
Net radiation	R_n	-30	Wm ⁻²
Air density	ρ	1.225	kgm ⁻³
Air specific heat capacity	c_p	1005	Jkg ⁻¹ K ⁻¹
Restoring temperature	θ_m	300	K

as an initial condition. The initial condition for θ is given by

$$\theta(z, t_0) = \begin{cases} \theta_0 & \text{for } z \leq z_c \\ \Gamma(z - z_c) + \theta_0 & \text{otherwise,} \end{cases}$$

where $z_c = 200$ m and $\Gamma = 0.01$ is the atmospheric lapse rate. The initial profile for the TKE, k , is defined as

$$\begin{aligned}
 k(z, t_0) &= a(t_0) \ln(z + z_0) + b(t_0) \text{ with} \\
 a(t) &= \frac{k(H, t) - k(z_0, t)}{\ln(H) - \ln(z_0)}, \\
 b(t) &= k(z_0, t) - a \ln(z_0), \\
 k(z_0, t_0) &= \frac{u_{z_0}^{*2}}{\sqrt{0.087}}, \\
 k(H, t_0) &= 0, \\
 u_{z_0}^* &= \frac{\kappa}{\ln(z_1/z_0)} \sqrt{u(z_1)^2 + v(z_1)^2},
 \end{aligned}$$

where H is the domain height. The initial condition for the surface temperature, θ_g , is equal to θ_0 and for the wind speed, v , it is equal to 0.

The boundary conditions at the lower boundary, i.e., $z = z_0$, are Dirichlet boundary conditions and are given by

$$\begin{aligned}
 u(z_0, t) &= 0, \\
 v(z_0, t) &= 0, \\
 \theta(z_0, t) &= \theta_g(t), \text{ and} \\
 k(z_0, t) &= k(z_0, t_0).
 \end{aligned}$$

The upper boundary condition, $z = H$, for v is zero. For all other variables, the upper boundary condition is a Neumann condition. The gradient of θ equals the lapse rate, and all other gradients are set to zero.

c. Steady state analysis

The system is run into a steady state before analyzing the impact of transient phenomena. This state is assumed to be reached when the solutions of equations (1) – (4) are nearly constant from this state on for at least one hour. To be precise, to calculate the steady state for a variable X , i.e., $u, v, \Delta\theta$ or k , at a fixed height z_{ss} , a rolling 5-hour mean of X is calculated. This is denoted as

$\overline{X_{t_{ss}-5h}}$. A steady state is reached at the time point t_{ss} when

$$|\overline{X_{t_{ss}-5h}} - X_{t_{ss}}| \leq 0.05.$$

This means at the steady state, the variable deviates at most by 0.05 to the average of the previous 5 hours for a fixed height, z_{ss} , (see as an example figure 1 a)). The first ten simulation hours are considered the initializing period and are therefore discarded for this analysis. The length of the rolling mean window is chosen purely for modeling purposes. The maximal difference between the rolling mean of the variables is set to a fixed value for simplicity. It is the smallest value for which, for all investigated u_G , a steady state is found within a 90 h simulation. As the steady state is only determined to initialize the perturbed runs with the values of all four model variables at this state, the thresholds to compute the steady state are of minimal importance as long as the system no longer changes significantly at this state. As we compare the inversion strength for a range of geostrophic wind forcings in the following section, we choose one fixed height for all s_G to determine the steady state. This height is the same at which the effect of perturbations on regime transitions is studied in Section 3. A fixed height is chosen to make the results comparable to the ones of other studies (e.g., Abraham and Monahan (2019a)).

It shall be noted that the geostrophic wind speed $s_G = \sqrt{u_G^2 + v_G^2}$ equals u_G as v_G is set to zero for all simulations. The height z_{ss} is chosen to be equal to 20 m for all simulations.

To analyze the sensitivity to perturbations, we use the model's values at the steady state as initial conditions for the perturbed runs.

d. Stability regimes

In the literature, the before mentioned backfolding when plotting $\Delta\theta_{z_{ss}} = \theta_{z_{ss}} - \theta_{z_0}$ over a wind speed $s_{z_{ss}}$ at a fixed height z_{ss} is commonly interpreted as a sign of bistability (Van de Wiel et al. 2017; Vignon et al. 2017). This implies that there are forcing scenarios for which the system's resilience to perturbations is weaker, which might facilitate transitions after the perturbation in these cases. Van de Wiel et al. (2017) showed that the strength of the surface coupling decides on the stability of the SBL. When plotting the steady state of $\Delta\theta_{z_{ss}}$ over the wind speed $s_{z_{ss}}$, with $z_{ss} = 20$ m, which is defined as $s_{z_{ss}} = \sqrt{u_{z_{ss}}^2 + v_{z_{ss}}^2}$ for a range of geostrophic wind speeds, s_G , (see figure 2) the graph shows the same backfolding and following the arguments made by




FIG. 1. Plots of u , $\Delta\theta$, v and TKE at $z_{ss} = 20$ m for $u_G = 1.0$ ms^{-1} . The steady-state period, which is the steady state plus one hour, is shaded in red.

Van de Wiel et al. (2017) this indicates that the solution of model (1)–(5) is bistable with the before mentioned parameters. If the model is bistable, it should have two stable equilibrium solutions and one unstable one for specific wind speeds. Based on this plot, low wind speeds and high-temperature differences characterize the first stable regime; the other one has high wind speeds and low-temperature differences. The first regime is therefore very stable, and the second is weakly stable. The potential bistable region, i.e., the region for where for the same $u_{z_{ss}}$ both stable states are a possible solution, is shaded in red in figure 2. If not specified differently $\Delta\theta$ and u refer to $\Delta\theta_{ss}$ and u_{ss} . The ss index is dropped for notational convenience. Commonly, a phenomenon where the system undergoes a sudden qualitative change triggered by specific threshold conditions of a relevant parameter, i.e., a bifurcation, is studied with regard to a fixed parameter, not one of the model variables like u that is dependent on height and time. Therefore, the gray line in figure 2 shows the same values for $\Delta\theta$ just plotted over s_G instead of s . For every simulation, only s_G is changed, and in particular, the initial values are defined as in Section 2b and not perturbed. Hence, this graph cannot display any backfolding. Therefore, to investigate the bistability of the model further, it is investigated if there is an inherent nonlinearity in the boundary layer dynamics

by analyzing the growth of the Ekman layer for increasing wind forcing. The Ekman layer height is the height where s equals s_G . The Ekman layer height for all considered s_G is plotted over time in figure 3 a). The first 10 hours are considered the initialization period and are therefore not displayed. The step-like structure of the lines can be explained by the non-equidistant grid for the space dimension, z . Plot b) shows how the Ekman layer height spread over time evolves with increasing s_G . The red horizontal dashed line shows the height upper height for $\Delta\theta$, i.e., $z_{ss} = 20$ m. The gray boxes represent all $\Delta\theta$ values obtained over the 90 hours of simulation time without the initializing period for every s_G forcing. The steady-state plus the following 12 hours are colored according to s_G in plots a) and b). Plot b) shows a nearly linear relationship between the Ekman layer height and s_G . For s_G less than 1.8 ms^{-1} , the steady state is below z_{ss} . This correlates with the start of the potential weakly stable regime in the bifurcation diagram (figure 2). The observed backfolding in figure 2 and the described phenomena in the literature can hence be explained by the fact that the temperature difference is studied for a fixed height, which is below the Ekman layer height for low geostrophic winds and above it for higher wind speeds. Therefore, this backfolding could be an artifact of the fixed height at which the observations were made and not the result of a sudden rapid shift in the dynamics or an actual sign of bistability. These results suggest that large-scale forcing, like a change in geostrophic wind, is a main driver for regime transitions in the SBL, i.e., bifurcation-driven transitions. Nonetheless, localized mixing due to small-scale fluctuations and intermittent events can be an additional factor that causes regime transitions, as uncertainty in the parameterization may increase the coupling (shift to weakly stable) or decrease it (shift to very stable). These types of transitions are called noise-induced transitions. As an example, local small-scale wind speed uptake increases shearing, leading to an increase in mixing and resulting in an increase in the thickness of the SBL and further coupling (weakly stable regime) of the boundary layer. Moreover, a burst of turbulence may non-linearly amplify mixing, leading again to further coupling.

3. Sensitivity analysis

To explore the impact of small-scale perturbations on regime transitions and identify the relevant variables' small-scale fluctuations, two approaches are used. First, the wind and temperature tendencies are perturbed to represent localized acceleration/deceleration of the wind or cold/warm




FIG. 2. Plot of the steady state of $\Delta\theta$ over s at a height of 20 meters, depicting its variation over different values of forcing s_G (color-coded). Additionally, the gray dots show the steady state of $\Delta\theta$ plotted over s_G . The dashed lines mark the mean $\Delta\theta$ value for each s_G plotted over s and s_G . The transition region, highlighted in red, is accompanied by a red box marking the corresponding s_G values in the colorbar.

air advection. This is achieved by introducing deterministic localized bursts of varying strength and size to the wind and temperature dynamics (see Section 3a). In part 2 of Section 3a, the influence of the perturbation size is quantified, and in part 3, four different stability scenarios are analyzed in detail. Opting for a deterministic perturbation facilitates a more straightforward quantification of its effects than a stochastic perturbation. Moreover, computational costs are significantly lower, as a stochastic perturbation would require ensemble runs to quantify its effects. Nevertheless, exploring the effects of stochastically perturbed tendencies is an important avenue for future research. Second, the parameterization of turbulence is perturbed to study the influence of turbulence intermittency. This is realized using a stochastic stability function to incorporate localized turbulent bursts (see Section c).




FIG. 3. a): Ekman layer height plotted over time. Every grey line is a simulation with a different geostrophic wind forcing. The colored sections of the lines mark the location of the steady states and the following hour. The colors indicate the value of the geostrophic wind for that particular simulation. b): Plots of the Ekman layer height for all considered s_G . The grey boxes show all obtained height values for the whole simulation time. Again, the colored sections are the values of the Ekman layer height at the steady state and the following hour, with the colors indicating the s_G value for that particular simulation. The red dashed line in the plot marks $z_{ss} = 20$ m.

a. Perturbed model tendencies

1) PERTURBATION DEFINITION

To perturb the dynamics of the wind and temperature locally in space and time, a 2D Gaussian function is used. This function is defined as

$$p(t, z) = r \cdot \exp\left(-\left[\frac{(t-t_c)^2}{2t_s^2} + \frac{(z-z_c)^2}{2z_s^2}\right]\right). \quad (8)$$

where r is the amplitude or maximum perturbation strength, t_c and z_c are the centers of the perturbations pulse in time and space, and t_s and z_s control the time and space spread of the perturbation. The unit of r depends on which variable is perturbed. If u is perturbed, the unit of r

is ms^{-2} , while for a perturbation of θ , it is Ks^{-1} . The perturbations are localized bursts, limited in height and time and small in magnitude. They fade out over a short time span. Figure 4 shows an example of such a perturbation.

To analyze the influence of small-scale changes in the temperature dynamics and analyze if the localized perturbations can trigger regime transitions, equation (3) is perturbed as follows

$$\begin{aligned}\frac{\partial \tilde{\theta}}{\partial t} &= \frac{\partial}{\partial z} \left(K_h(\phi) \frac{\partial \theta}{\partial z} \right) \pm p(t, z) \\ &= \frac{\partial \theta}{\partial t} \pm p(t, z).\end{aligned}\tag{9}$$

The wind speed equation is perturbed similarly to analyze the sensitivity of the model to small-scale changes in the wind dynamics, i.e.,

$$\begin{aligned}\frac{\partial \tilde{u}}{\partial t} &= f_c \cdot (v - v_G) + \frac{\partial}{\partial z} \left(K_m(\phi) \frac{\partial u}{\partial z} \right) - N_u \pm p(t, z) \\ &= \frac{\partial u}{\partial t} \pm p(t, z).\end{aligned}\tag{10}$$

If the simulation starts in a very stable regime a negative perturbation is added to θ and a positive one to u to induce transitions. Similarly, if it starts in the weakly stable regime, a positive perturbation is added to θ and a negative one to u .

2) EFFECT OF THE PERTURBATION SIZE

The perturbation is characterized by four parameters that control its location and spread in space and time. Different values for the time and space spread are investigated to study the effect of the perturbation size. The center of the perturbation is fixed at 20 m and 0.5 h to allow easy comparison. This height is chosen as the steady state was determined at this height. By choosing 0.5 h as the perturbation center, a range of time spreads can be tested without the perturbation extending beyond the initial condition. The maximal time duration of the perturbation which is considered is 33 min ($t_s = 500$ s), and the maximal space extension is 40 m ($z_s = 10$ m), which is in the realm of submesoscale motions (Mahrt 2014; Vercauteren et al. 2016; Boyko and Vercauteren 2021). The smallest extension considered is 7 min ($t_s = 100$ s) in time and 4 m ($z_s = 1$ m) in space. Figure 5 shows all considered perturbations, and figure 6 shows the results of the corresponding




FIG. 4. Example of a positive perturbation (eq. (8)) with maximal perturbation strength, r , equal to 0.01, height spread $z_s = 10$ m and time spread $t_s = 200$ s.

sensitivity analysis. The sensitivity analysis determines the minimal perturbation strength for every combination of geostrophic wind, perturbation type, and size for which at least one transition takes place in the 12 h simulation time. Here, perturbation strength refers to the relation between r and the variance of the corresponding variable (u or θ) for the first 15 minutes of the simulation. This period is chosen as the simulations are not perturbed during this period for all perturbation sizes. The range of tested r values and the variance for all heights are normalized with a min-max normalization. The perturbation strength is then defined as $r_n / \text{var}(X_{n,20m}) \cdot 100$ where r_n is the normalized r range and X_n is either the normalized u or θ . For this study, u_G values between 1.0 and 2.5 ms^{-1} and r values between 0 and 0.019 were tested. For the smallest perturbation, no simulation included transitions. Therefore, no smaller perturbations were considered for this study. As expected, negative perturbations added to the temperature dynamics cause regime transitions if the simulations are started in the very stable regime ($u_G \leq 1.7 \text{ ms}^{-1}$) while a positive perturbation causes transitions for simulation started in the weakly stable regime ($u_G \geq 1.8 \text{ ms}^{-1}$) for nearly all perturbation sizes. This agrees with physical intuition as the inversion strength increases if warm air is advected over a colder surface, while cold air advection decreases the inversion strength if the advection is close to the surface. A higher perturbation strength would most likely yield the

expected results in the few cases where this does not seem to hold. However, for the parameters used in this study, a perturbation time spread of 4 m is insufficient to induce transitions. The results of the perturbed temperature dynamics show a noticeable difference when t_s is changed, whereas a change in z_s does not have as pronounced an effect. Unsurprisingly, increasing t_s leads to a decrease in the minimum perturbation strength required to cause transitions.

Analogously, positive perturbations added to the wind dynamics cause regime transitions if the simulations were started in the very stable regime ($u_G \leq 1.7 \text{ ms}^{-1}$) while a negative perturbation causes transitions for simulation started in the weakly stable regime ($u_G \geq 1.8 \text{ ms}^{-1}$) if the perturbation was large enough. The decrease in stability after an increase in wind speed can be explained by the fact that an increase in wind speed results in more shearing and, thereby, more mixing, ultimately leading to further coupling. For the here used parameters, a total perturbation time spread of 400 s, i.e., ≈ 7 min, or a total height spread of 4 m, is insufficient to get sufficient transitions related to perturbed wind dynamics. Contrary to the temperature dynamics for the perturbed wind dynamics, there is a noticeable difference between the results if z_s or t_s is changed. Unsurprisingly, if t_s increases, the minimal perturbation strength to cause transitions decreases. The same holds for an increase in z_s , just less drastic. Surprisingly, there are transitions from very to weakly stable for a negative wind perturbation. This can be explained by a wind direction change and is discussed in detail in the following section.

3) EFFECT OF PERTURBATION IN DETAIL

In this section, the focus is on four different stability scenarios:

- S1. $u_G = 1.0 \text{ ms}^{-1}$: very stable regime,
- S2. $u_G = 1.7 \text{ ms}^{-1}$: transition wind speed of very stable regime,
- S3. $u_G = 1.8 \text{ ms}^{-1}$: transition wind speed of weakly stable regime, and
- S4. $u_G = 2.5 \text{ ms}^{-1}$: weakly stable regime.

Figure 7 shows the impact of the temperature perturbation on the potential temperature, θ , the wind speed S , the wind direction γ and the inversion strength $\Delta\theta$. Each row corresponds to one of the four scenarios. The perturbation strength is chosen according to the sensitivity analysis performed in the previous section. The negative and positive perturbations are visible in the plots of θ , while




FIG. 5. This figure gives an overview of all considered perturbations. In each column, the height spread, z_s , is constant, while in every row, the time spread, t_s , is constant. That means from left to right, the spread in height increases, and from top to bottom, the time spread increases. The strength of the perturbation is indicated by color. The maximal perturbation strength is equal to 0.001 for all plotted perturbations.

the wind speed and direction do not change significantly. Therefore, the temperature perturbation impacted mainly the temperature and effectively represents the advection of cold or warm air. The plots for the temperature inversion strength (right column in figure 7) show that only for the second scenario is there a tendency for the system to stay in the new regime after the transition. For all other scenarios, while the transition occurs rapidly after the perturbation, the system tends to return to its original state for all tested perturbation strengths. This holds especially for scenario 4, where the return to the weakly stable state occurs even for the strongest perturbation after less than 7 hours.

Figure 8 is the same figure as figure 7 but with wind velocity perturbation instead of temperature perturbation. For the positive perturbation, i.e., scenarios 1 and 2, the wind speed plots look as expected. At the center of the perturbation, the wind speed is the highest and then decreases in the space and time dimension. The wind direction changes slightly but not drastically. Therefore, in these two cases, adding a perturbation to u causes mainly a perturbation in the wind speed.




FIG. 6. This figure gives an overview of the results of the sensitivity analysis for all considered perturbations (see figure 5). In each column, the height spread, z_s , is constant, while in every row, the time spread, t_s , is constant. That means from left to right, the spread in height increases, and from top to bottom, the time spread increases. The plots show the minimal perturbation strength for which that simulation includes a transition. The perturbation strength is here given in relation to the variance of the simulation’s first 15 minutes, which are not perturbed.

Contrarily, the negative perturbations significantly impact the wind direction. In fact, at the center of the perturbation, the wind turns with nearly 180° . This also explains why the wind speed increases at the center of the perturbation, even though a negative perturbation was added. The wind velocity component u decreases so much that it gets negative, which is equivalent to a wind direction change of 180° . The reversal of the wind direction explains the phenomena discovered in the previous section: a negative wind perturbation causes very to weakly stable transitions for some perturbations. This is not the intended behavior, and several approaches to prevent this have been tried. But all approaches lead to strong numerical instabilities. Hence, the results obtained with a negative wind perturbation must be treated with some reservation. Nonetheless, the temperature inversion strength plots show increased stability after the perturbation and non-persistent transitions into the very stable regime ($\Delta\theta > 5\text{ K}$).

In conclusion, the combined results for both perturbation types indicate that small-scale phenomena can drive persistent regime transitions from very to weakly stable regimes. In contrast, the phenomena tested here are insufficient to cause persistent weakly to very stable regime transitions. Therefore, it is especially relevant to account for small-scale fluctuations in the wind and temperature dynamics in weather and climate models to achieve accurate regime transition statistics for very to weakly stable transitions.

b. Changes in the system's stability

After studying the Ekman layer height in Section 2d for the purely deterministic model, we now study the Ekman layer height of a randomized version of the model to analyze if there is an inherent nonlinearity in the boundary layer dynamics in the sense that the growth of the Ekman layer becomes nonlinear for a certain range of wind speed. For this, the model with additive noise added to the temperature dynamics (eq. (9)) is initialized at the steady state of the deterministic model and run for 12 hours. The same perturbation as in Section 3a is used. The perturbation is a small, localized burst of small magnitude, which is added in the first hour of the simulation and then fades out in that hour. The center of the perturbation is at 20 m. The results are shown in figure 9. For plot a) to c), a negative temperature perturbation is added, equating to cold air's advection. As the perturbation is added at a fixed height for geostrophic winds less than 1.8 ms^{-1} , the center of the perturbation is above the Ekman layer height, while for higher wind speeds, it is below. Figure 9 c) shows a plot of the temperature profiles for the perturbed solution for $u_G = 1.6 \text{ ms}^{-1}$. The different lines correspond to different simulation time points. As $u_G < 1.8 \text{ ms}^{-1}$, the center of the perturbation is above the height of the Ekman layer. As a result of the perturbation, the temperature decreases at roughly 20 m, which is the center of the perturbation and higher than the Ekman layer at the simulation's start. Hence, the perturbation causes cooling above the Ekman layer, creating a strong vertical temperature gradient between the inversion layer and the cooled layer above. This causes an increase in mixing and, thereby, an increase in the Ekman layer height until a new equilibrium state is reached. In plot c), the temperature profiles for $u_G = 1.9 \text{ ms}^{-1}$ are shown. Here, the center of the perturbation is below the height of the Ekman layer. The negative perturbation causes a capping inversion to form between 20 m and 40 m, partially below the Ekman layer height, effectively reducing the height of the mixed layer and thus reducing the Ekman layer

height. But as a counter-movement, a convective state emerges in the first 20 m, which allows the mixing to take up more space again and thereby increases the Ekman layer height again. As the model nonlinearly amplifies this process, the Ekman layer height rises above the original height from the start of the simulation and equilibrates in this state.

For plots d) to f), a positive temperature perturbation is added, equating to warm air's advection. Again, for geostrophic winds less than 1.8 ms^{-1} , the perturbation is added above the Ekman layer height, while for higher wind speeds, the perturbation is added below. As shown in plot e) where $u_G = 1.6 \text{ ms}^{-1}$ When the perturbation is added above the Ekman layer and for low geostrophic winds, it does not affect the Ekman layer height. While the perturbation causes the temperature to increase at roughly 20 m, this is above the Ekman layer height, and hence, the inversion layer is only strengthened, and no change in mixing occurs, which would result in a change in the Ekman layer. In contrast, for $u_G = 1.9 \text{ ms}^{-1}$, the perturbation center is below the Ekman layer height as seen in plot f). Hence, even though the perturbation has the same effect as for $u_G = 1.6 \text{ ms}^{-1}$, namely the increase in temperature, the strengthening of the inversion further reduces mixing, resulting in a reduction of the Ekman layer height until it starts to return to its original state.

As a low Ekman layer correlates with a strong temperature inversion, i.e., the very stable regime, and a higher one with a weak temperature inversion, i.e., the weakly stable regime, the results above support the observation from the previous section that small-scale perturbations have a long-lasting impact for very to weakly stable transitions. This results from the top-down mixing event recoupling the previously decoupled low SBL to get a much deeper SBL. For weakly to very stable transitions, small-scale perturbations were added at a height relatively close to the surface as commonly observed in field studies (Sun et al. 2012; Lan et al. 2022), but they only have a transient impact.

c. Perturbed parameterization

As the effect of perturbed tendencies was studied in the previous section (Sect. 3) in this section, the focus lies on the effect of a perturbed parameterization. This is achieved by using a randomized stability function. The set of stability functions proposed in the literature is large. While developed with the goal to be universal, many slightly different functions have been derived based on field studies (examples are given, e.g., in Rodrigo and Anderson (2013)). They introduce

different types of model errors or uncertainty. This section uses a random stability function, which was initially developed in Boyko and Vercauteren (2023) to represent model uncertainty in the turbulence closure. The authors sought to capture unsteady mixing, potentially linked to turbulence intermittency and unresolved fluid motions. Their strategy involves incorporating stochastic parameterization by introducing randomness into the stability correction utilized in the traditional Monin–Obhukov similarity theory. They developed a data-driven stability correction function using a statistical model-based clustering technique (Boyko et al. 2022). The resulting stochastic stability function incorporates random perturbations to address localized intermittent turbulence bursts. This approach considers stochastic mixing effects from unresolved processes. The stochastic stability function is briefly introduced here, but more details are given in Boyko and Vercauteren (2023). The function is defined as

$$\rho(t, Ri) = \phi(Ri)\text{sig}(z) + \phi_s(t, Ri)(1 - \text{sig}(z)) \quad (11)$$

with $\phi(Ri) = 1 + 12Ri$ being the deterministic short-tail stability function, which is defined in section 2a. The function $\text{sig}(z) = \frac{1}{1 + \exp(-k_c(z - z_m))}$ is the logistic sigmoid function where z_m is the sigmoid's midpoint, and $k_c = 0.1$ controls the sharpness of the transition from ϕ to ϕ_s at height z_m . The stochastic part of equation (11), $\phi_s(t, Ri)$, is given as the solution to the stochastic differential equation

$$d\phi_s(t, Ri) = [1 - \Lambda(Ri)\phi_s - \Upsilon(Ri)\phi_s^2]dt + \Sigma(Ri)\phi_s dW(t), \quad \phi_s(0) = \phi_{s0}. \quad (12)$$

where $W(t)$ is a Wiener process and the scaling functions, $\Lambda, \Upsilon, \Sigma$, are defined as

$$\Lambda(Ri) = 9.3212 \tanh[0.9088 \log_{10}(Ri) - 0.0738] + 8.3220 \quad (13)$$

$$\Upsilon(Ri) = 10^{0.4294 \log_{10}(Ri) + 0.1749} \quad (14)$$

$$\Sigma(Ri) = 10^{0.8069 \tanh[0.6044 \log_{10}(Ri) - 0.8368] + \sigma_s}. \quad (15)$$

To study the impact of using the randomized stability function, the same model as in section 3a (i.e., equations (1) to (5)) is used. The noise intensity is independent of the stratification controlled by σ_s . Hence, this parameter is chosen for the sensitivity study. Figure 10 shows examples of the

stochastic stability function for four different σ_s . The plots show that a decrease in σ_s results in a decrease in the variability of ϕ . Due to the stochasticity in the definition of ϕ , the stability function creates temporally stronger mixing events (see, e.g., plot a)). The study by Boyko and Vercauteren (2023) analyzes results with three different values for σ_s . Here, the same values are used for the sensitivity study, and a high-noise scenario is added. The following four noise scenarios are studied:

N1. $\sigma_s = 1$: high noise level,

N2. $\sigma_s = 0$: medium noise level, estimated based on FLOSS II dataset (Boyko and Vercauteren 2023),

N3. $\sigma_s = -0.07$: medium noise level, adjusted noise level for FLOSS II dataset (Boyko and Vercauteren 2023) and

N4. $\sigma_s = -1$: suppressed noise level.

In addition, the same stability scenarios as in section 3a are used:

S1. $u_G = 1.0 \text{ ms}^{-1}$: very stable regime,

S2. $u_G = 1.7 \text{ ms}^{-1}$: transition wind speed of very stable regime,

S3. $u_G = 1.8 \text{ ms}^{-1}$: transition wind speed of weakly stable regime, and

S4. $u_G = 2.5 \text{ ms}^{-1}$: weakly stable regime.

For each noise and stability scenario, 200 simulations are run for 12 hours. The results are shown in figure 11. The columns in figure 11 are the different noise scenarios, while the rows correspond to the stability scenarios. In the high noise scenario (left column), $\Delta\theta$ displays a high variability, while for the scenario with the suppressed noise level (right column), the different simulations of $\Delta\theta$ are nearly equal to the mean over all simulations (red line).

The bifurcation diagrams and the plots of $\Delta\theta$ for all scenarios show that there are transitions from the very to the weakly stable regime only for the high-noise scenario. For scenarios S2. N2., S2. N3., and S2. N4., the regime threshold is barely crossed by the end of simulation time. Hence, they are not counted as full regime transitions. For none of the scenarios, there are weakly to very stable regime transitions.

In conclusion, the results indicate that the stochastic stability function prevents the model from getting "stuck" in the very stable regime, thereby preventing the so-called "runaway cooling". However, this stability function does not facilitate weakly to very stable regime transitions.

4. Summary and Conclusion

This study expands on prior research on the relevance of small-scale perturbations for SBL regime transitions. We used a 1.5 closure single-column model for the atmospheric boundary layer used in Boyko and Vercauteren (2024) to quantify the impact of perturbed tendencies and stochastic parameterizations. In particular, we studied the relevance of small-scale fluctuations of wind and temperature dynamics and intermittency for regime transitions. In addition, we addressed how they can be added to NWP and climate models. The fluctuations of the wind and temperature dynamics were included in the model by adding a 2D Gaussian function to either the function for the wind or the temperature evolution over time. Different perturbation strengths and sizes were tested in a sensitivity study. The results for both perturbation types indicate that small-scale phenomena can drive persistent regime transitions from very to weakly stable regimes. In contrast, the phenomena tested here are insufficient to cause persistent weakly to very stable regime transitions. Therefore, it is especially relevant to account for small-scale fluctuations in the wind and temperature dynamics in weather and climate models to achieve accurate regime transition statistics for very to weakly stable transitions. For the temperature dynamic perturbations, the time spread of the perturbation has a noticeable impact, while the height spread has less of an impact. For the perturbed wind dynamics, both dimensions are relevant. In addition to the impact of the perturbations on regime transitions, we studied their effect on the dynamics of the SBL by studying the growth of the Ekman layer after a perturbation for different forcing scenarios. The results agreed with the previous findings that small-scale perturbations have a long-lasting impact for very to weakly stable transitions while for weakly to very stable they only have a transient impact. Finally, to address model errors related to the Monin-Obukhov Similarity Theory, we investigated the influence of the stochastic stability function, as initially defined by Boyko and Vercauteren (2023). The parameters of the stability correction were estimated based on observational data. Analogously, this methodology can be extended to Numerical Weather Prediction models or Earth system models to integrate localized turbulent bursts. Incorporating such transient events is of substantial scientific importance,

given their potential to induce regime transitions within the stable boundary layer. Furthermore, utilizing a randomized stability function offers an alternative avenue to mitigate issues associated with the long-tail stability function by effectively reducing excessive mixing. Simultaneously, a randomized stability function could prevent the model from becoming trapped in a very stable state when employing a short-tail stability function. Our results show that the stochastic stability function prevents the model from getting trapped in the very stable regime, thus preventing the so-called "runaway cooling". However, this stability function does not facilitate weakly to very stable regime transitions.

In conclusion, the here-tested perturbations and randomizations are a promising avenue to circumvent the need for excessive mixing through a long-tail stability function while also preventing runaway cooling in NWP. The results suggest that small-scale fluctuations are most relevant for very to weakly stable regime transitions, which complements findings from Abraham and Monahan (2019b) demonstrating that large-scale forcing is the main driver for weakly to very stable regime transitions, but less for very to weakly stable transitions. Future work will entail making the perturbations of the wind and temperature dynamics stochastic and dependent on the stratification level and model resolution to make this approach applicable to NWP.




FIG. 7. Examples of simulations with perturbation added to the temperature dynamics. The plots show the simulation output of the temperature (first column), wind speed (second column), and wind direction (third column) over space and time. The right column shows the temperature inversion strength plots for a range of perturbation strengths, r , indicated by color. The parameters for the perturbation size are $z_s = 5$ m and $t_s = 300$ s. The geostrophic wind and the perturbation strength are the same in every row. They are given in the title of each plot. Every row corresponds to one of the four stability scenarios (S1. - S4.). In the first/last row, the simulation is initialized in the very/weakly stable regime outside the bifurcation region. In the second/third, the simulation starts in the very/weakly stable regime but at the transition wind speed (see figure 2).




FIG. 8. Examples of simulations with perturbation added to the wind dynamics. The plots show the simulation output of the temperature (first column), wind speed (second column), and wind direction (third column) over space and time. The right column shows the temperature inversion strength plots for a range of perturbation strengths, r , indicated by color. The parameters for the perturbation size are $z_s = 10$ m and $t_s = 300$ s. The geostrophic wind and the perturbation strength are the same in every row. They are given in the title of each plot. Every row corresponds to one of the four stability scenarios (S1. – S4.). In the first/last row, the simulation is initialized in the very/weakly stable regime outside the bifurcation region. In the second/third, the simulation starts in the very/weakly stable regime but at the transition wind speed (see figure 2).




FIG. 9. First row: Results with negatively perturbed temperature dynamics. Second row: positively perturbed temperature dynamics. First column: Ekman layer height plotted over time. Every line is a simulation with a different geostrophic wind forcing, indicated by the color. Second column: Temperature profiles for $u_G = 1.6 \text{ ms}^{-1}$ and six different simulation time points (marked as vertical dotted lines in column one). Third column: Same as second column, just for $u_G = 1.9 \text{ ms}^{-1}$.




FIG. 10. Examples of stochastic stability function with varying noise strength: a) $\sigma_s = 1$, b) $\sigma_s = 0$, c) $\sigma_s = -0.07$ and d) $\sigma_s = -1$. Each plot shows 100 examples of ϕ and $u_G = 1.0 \text{ ms}^{-1}$ for all of them.




FIG. 11. Results of the model (i.e., eq. (1) to (5)) with a randomized stability function (eq. (11)) for the four different noise scenarios N1. - N4. (columns), and four different stability scenarios S1. - S4. (rows 2-4). Each combination of scenarios was run 200 times. The first row displays bifurcation plots for each noise scenario. The colors indicate the value of the geostrophic wind, and the blue dots and the blue line correspond to the results with a short-tail stability function and no added noise. Similarly, the red dots and the red line correspond to the results with a long-tail stability function and no noise. Rows 2-4 are plots of 200 simulations of $\Delta\theta$ over time, t , for all noise and stability scenarios. The green line is the mean for each scenario over all 200 simulations.

Acknowledgments. This project has received funding from the European Union’s Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement number 945371. The computations were performed on resources provided by Sigma2 - the National Infrastructure for High-Performance Computing and Data Storage in Norway.

Data availability statement. The source code to reproduce all results can be found here: https://github.com/BoundaryLayerVercauteren/abl_scm_perturbation_study.

References

- Abraham, C., A. M. Holdsworth, and A. H. Monahan, 2019: A prototype stochastic parameterization of regime behaviour in the stably stratified atmospheric boundary layer. *Nonlinear Processes in Geophysics*, **26** (4), 401–427, <https://doi.org/10.5194/npg-26-401-2019>.
- Abraham, C., and A. H. Monahan, 2019a: Climatological Features of the Weakly and Very Stably Stratified Nocturnal Boundary Layers. Part I: State Variables Containing Information about Regime Occupation. *Journal of the Atmospheric Sciences*, **76** (11), 3455–3484, <https://doi.org/10.1175/JAS-D-18-0261.1>.
- Abraham, C., and A. H. Monahan, 2019b: Climatological Features of the Weakly and Very Stably Stratified Nocturnal Boundary Layers. Part III: The Structure of Meteorological State Variables in Persistent Regime Nights and across Regime Transitions. *Journal of the Atmospheric Sciences*, **76** (11), 3505–3527, <https://doi.org/10.1175/JAS-D-18-0274.1>.
- Acevedo, O. C., F. D. Costa, R. Maroneze, A. D. Carvalho, F. S. Puhales, and P. E. S. Oliveira, 2021: External controls on the transition between stable boundary-layer turbulence regimes. *Quarterly Journal of the Royal Meteorological Society*, **147** (737), 2335–2351, <https://doi.org/10.1002/qj.4027>.
- Acevedo, O. C., and D. R. Fitzjarrald, 2003: In the Core of the Night-Effects of Intermittent Mixing on a Horizontally Heterogeneous Surface. *Boundary-Layer Meteorology*, **106** (1), 1–33, <https://doi.org/10.1023/A:1020824109575>.
- Acevedo, O. C., L. Mahrt, F. S. Puhales, F. D. Costa, L. E. Medeiros, and G. A. Degrazia, 2015: Contrasting structures between the decoupled and coupled states of the stable boundary layer.

- Quarterly Journal of the Royal Meteorological Society*, **142 (695)**, 693–702, <https://doi.org/10.1002/qj.2693>.
- Baas, P., B. J. H. Van de Wiel, S. J. A. van der Linden, and F. C. Bosveld, 2017: From Near-Neutral to Strongly Stratified: Adequately Modelling the Clear-Sky Nocturnal Boundary Layer at Cabauw. *Boundary-Layer Meteorology*, **166 (2)**, 217–238, <https://doi.org/10.1007/s10546-017-0304-8>.
- Berner, J., and Coauthors, 2017: Stochastic Parameterization: Toward a New View of Weather and Climate Models. *Bulletin of the American Meteorological Society*, **98 (3)**, 565–588, <https://doi.org/10.1175/BAMS-D-15-00268.1>.
- Boyko, V., S. Krumscheid, and N. Vercauteren, 2022: Statistical Learning of Nonlinear Stochastic Differential Equations from Nonstationary Time Series using Variational Clustering. *Multiscale Modeling & Simulation*, 1251–1283, <https://doi.org/10.1137/21M1403989>.
- Boyko, V., and N. Vercauteren, 2021: Multiscale Shear Forcing of Turbulence in the Nocturnal Boundary Layer: A Statistical Analysis. *Boundary-Layer Meteorology*, **179 (1)**, 43–72, <https://doi.org/10.1007/s10546-020-00583-0>.
- Boyko, V., and N. Vercauteren, 2023: A stochastic stability equation for unsteady turbulence in the stable boundary layer. *Quarterly Journal of the Royal Meteorological Society*, **149 (755)**, 2125–2145, <https://doi.org/10.1002/qj.4498>.
- Boyko, V., and N. Vercauteren, 2024: Simulating the unsteady stable boundary layer with a stochastic stability equation. *Journal of Geophysical Research: Atmospheres*, **129**, <https://doi.org/10.1029/2023JD039370>.
- Chechin, D. G., I. A. Makhotina, C. Lüpkes, and A. P. Makshtas, 2019: Effect of Wind Speed and Leads on Clear-Sky Cooling over Arctic Sea Ice during Polar Night. *Journal of the Atmospheric Sciences*, **76 (8)**, 2481–2503, <https://doi.org/10.1175/JAS-D-18-0277.1>.
- Christensen, H. M., 2020: Constraining stochastic parametrisation schemes using high-resolution simulations. *Quarterly Journal of the Royal Meteorological Society*, **146 (727)**, 938–962, <https://doi.org/10.1002/qj.3717>.

- Delage, Y., 1997: Parameterising Sub-Grid Scale Vertical Transport in Atmospheric Models Under Statically Stable Conditions. *Boundary-Layer Meteorology*, **82** (1), 23–48, <https://doi.org/10.1023/A:1000132524077>.
- Derbyshire, S., 1999: Boundary-layer decoupling over cold surfaces as a physical boundary-instability. *Boundary-Layer Meteorology*, **90** (2), 297–325.
- Donda, J. M. M., I. G. S. van Hooijdonk, A. F. Moene, H. J. J. Jonker, G. J. F. van Heijst, H. J. H. Clercx, and B. J. H. Van de Wiel, 2015: Collapse of turbulence in stably stratified channel flow: a transient phenomenon. *Quarterly Journal of the Royal Meteorological Society*, **141** (691), 2137–2147, <https://doi.org/10.1002/qj.2511>.
- Garratt, J. R., 1994: Review: the atmospheric boundary layer. *Earth-Science Reviews*, **37** (1), 89–134, [https://doi.org/10.1016/0012-8252\(94\)90026-4](https://doi.org/10.1016/0012-8252(94)90026-4).
- Holtslag, A., and Coauthors, 2013: Stable atmospheric boundary layers and diurnal cycles: challenges for weather and climate models. *Bulletin of the American Meteorological Society*, **94** (11), 1691–1706, <https://doi.org/10.1175/BAMS-D-11-00187.1>.
- Kaiser, A., N. Vercauteren, and S. Krumscheid, 2024: Sensitivity of the polar boundary layer to transient phenomena. *Nonlinear Processes in Geophysics*, **31** (1), 45–60, <https://doi.org/10.5194/npg-31-45-2024>.
- Kähnert, M., 2022: Advancing the Capabilities of Numerical Weather Prediction - On the Utility of Individual Tendency Output. Doctoral thesis, The University of Bergen, URL <https://bora.uib.no/bora-xmlui/handle/11250/3030505>, accepted: 2022-11-07T14:32:07Z ISBN: 9788230845776.
- Lan, C., H. Liu, G. G. Katul, D. Li, and D. Finn, 2022: Turbulence Structures in the Very Stable Boundary Layer Under the Influence of Wind Profile Distortion. *Journal of Geophysical Research: Atmospheres*, **127** (20), e2022JD036565, <https://doi.org/10.1029/2022JD036565>.
- Mahrt, L., 1998: Stratified Atmospheric Boundary Layers and Breakdown of Models. *Theoretical and computational fluid dynamics*, **11** (3-4), 263–279, <https://doi.org/10.1007/s001620050093>.
- Mahrt, L., 2014: Stably Stratified Atmospheric Boundary Layers. *Annual Review of Fluid Mechanics*, **46** (1), 23–45, <https://doi.org/10.1146/annurev-fluid-010313-141354>.

- Maroneze, R., F. D. Costa, O. C. Acevedo, L. E. Medeiros, F. S. Puhales, V. Anabor, and L. Mortarini, 2023: A New Stable Boundary Layer Parameterization for Numerical Weather Prediction Models: A Heat Flux Budget Approach. *Boundary-Layer Meteorology*, **188** (2), 209–228, <https://doi.org/10.1007/s10546-023-00810-4>.
- Palmer, T., 2017: The primacy of doubt: Evolution of numerical weather prediction from determinism to probability. *Journal of Advances in Modeling Earth Systems*, **9** (2), <https://doi.org/10.1002/2017MS000999>.
- Palmer, T. N., 2001: A nonlinear dynamical perspective on model error: A proposal for non-local stochastic-dynamic parametrization in weather and climate prediction models. *Quarterly Journal of the Royal Meteorological Society*, **127** (572), 279–304, <https://doi.org/10.1002/qj.49712757202>.
- Ramsey, E., and A. H. Monahan, 2022: Empirical Low-Dimensional Dynamics of Atmospheric Stable Boundary Layer Temperature Inversions. *Journal of the Atmospheric Sciences*, **79** (7), 1965–1984, <https://doi.org/10.1175/JAS-D-21-0205.1>.
- Rodrigo, J. S., and P. S. Anderson, 2013: Investigation of the Stable Atmospheric Boundary Layer at Halley Antarctica. *Boundary-Layer Meteorology*, **148** (3), 517–539, <https://doi.org/10.1007/s10546-013-9831-0>.
- Sandu, I., A. C. M. Beljaars, P. Bechtold, T. Mauritsen, and G. Balsamo, 2013: Why is it so difficult to represent stably stratified conditions in numerical weather prediction (NWP) models? *Journal of Advances in Modeling Earth Systems*, **5** (2), 117–133, <https://doi.org/10.1002/jame.20013>.
- Steenefeld, G.-J., 2014: Current challenges in understanding and forecasting stable boundary layers over land and ice. 1–6, <https://doi.org/10.3389/fenvs.2014.00041/abstract>.
- Stull, R. B., 1988: *An Introduction to Boundary Layer Meteorology*. Springer Netherlands, <https://doi.org/10.1007/978-94-009-3027-8>, URL <http://link.springer.com/10.1007/978-94-009-3027-8>.
- Sun, J., L. Mahrt, R. M. Banta, and Y. L. Pichugina, 2012: Turbulence Regimes and Turbulence Intermittency in the Stable Boundary Layer during CASES-99. *Journal of the Atmospheric Sciences*, **69** (1), 338–351, <https://doi.org/10.1175/JAS-D-11-082.1>.

- Van de Wiel, B. J. H., A. F. Moene, H. J. J. Jonker, P. Baas, S. Basu, J. M. M. Donda, J. Sun, and A. Holtslag, 2012: The Minimum Wind Speed for Sustainable Turbulence in the Nocturnal Boundary Layer. *Journal of Atmospheric Sciences*, **69** (11), 3116–3127, <https://doi.org/10.1175/JAS-D-12-0107.1>.
- Van de Wiel, B. J. H., and Coauthors, 2017: Regime transitions in near-surface temperature inversions: A conceptual model. *Journal of the Atmospheric Sciences*, **74** (4), 1057–1073, <https://doi.org/10.1175/JAS-D-16-0180.1>.
- van Hooijdonk, I. G. S., J. M. M. Donda, H. J. H. Clercx, F. C. Bosveld, and B. J. H. Van de Wiel, 2015: Shear Capacity as Prognostic for Nocturnal Boundary Layer Regimes. *Journal of Atmospheric Sciences*, **72** (4), 1518–1532, <https://doi.org/10.1175/JAS-D-14-0140.1>.
- Vercauteren, N., L. Mahrt, and R. Klein, 2016: Investigation of interactions between scales of motion in the stable boundary layer. *Quarterly Journal of the Royal Meteorological Society*, **142** (699), 2424–2433, <https://doi.org/10.1002/qj.2835>.
- Vignon, E., and Coauthors, 2017: Stable boundary-layer regimes at Dome C, Antarctica: observation and analysis. *Quarterly Journal of the Royal Meteorological Society*, **143** (704), 1241–1253, <https://doi.org/10.1002/qj.2998>.