

# Presenting the topological stratified homotopy hypothesis

Lukas Waas

March 2024

## Abstract

This article is concerned with three different homotopy theories of stratified spaces: The one defined by Douteau and Henriques, the one defined by Haine, and the one defined by Nand-Lal. One of the central questions concerning these theories has been how precisely they connect with geometric and topological examples of stratified spaces, such as piecewise linear pseudomanifolds, Whitney stratified spaces, or more recently Ayala, Francis and Tanaka's conically smooth stratified spaces. More precisely, so far, it has been an open question whether there exist (semi)model structures on stratified topological spaces that present these theories, in which such relevant examples of stratified spaces are bifibrant. Here, we prove an affirmative answer to this question. As a consequence, we obtain a model categorical interpretation of a stratified homotopy hypothesis. Specifically, we show that Lurie's stratified singular simplicial set functor induces a Quillen equivalence between the semimodel category of stratified topological spaces presenting Nand-Lal's homotopy theory of stratified spaces and the left Bousfield localization of the Joyal model structure that corresponds to such  $\infty$ -categories in which every endomorphism is an isomorphism. We then perform a detailed investigation of bifibrant objects in these model structures of stratified spaces, proving a series of detection criteria and illuminating the relationship to Quinn's homotopically stratified spaces.

## 1 Introduction

Conceptually speaking, Grothendieck's homotopy hypothesis refers to the following statement:

Assigning to a topological space its  $\infty$ -groupoid of paths induces an equivalence between the homotopy theory of spaces (more precisely CW complexes) and the homotopy theory of  $\infty$ -groupoids.

Whether this statement is regarded as a theorem or a conjecture, of course, strongly relies on the precise model of  $\infty$ -categories (and thus of  $\infty$ -groupoids) one has in mind. Nowadays, the following result due to Quillen is often taken as a formal interpretation of the homotopy hypothesis:

**Theorem HH.** [Qui67] *The geometric realization and singular simplicial set adjunction*

$$|-| : \mathbf{sSet} \rightleftarrows \mathbf{Top} : \text{Sing}$$

*induces a Quillen equivalence between the Quillen model structure on topological spaces and the Kan-Quillen model structure on simplicial sets.*

That this is a formal interpretation of the homotopy hypothesis can be argued as follows.

1. The Quillen equivalence induces an equivalence

$$\mathbf{Top}[W^{-1}] \simeq \mathbf{Kan}[H_k^{-1}]$$

between topological spaces localized at weak homotopy equivalences and Kan complexes localized at homotopy equivalences of Kan complexes.

2. Quasi-categories (as introduced by Joyal and popularized by Lurie in [Lur09]) have proven to be an extremely powerful and flexible model for the theory of  $(\infty, 1)$ -categories ( $\infty$ -categories henceforth). Kan complexes are the  $\infty$ -groupoid internal to quasi-categories, and it follows from the existence of the Kan-Quillen and the Joyal model structure that the right-hand side thus defines the homotopy theory of  $\infty$ -groupoids  $\mathbf{Grpd}_\infty$ . The singular simplicial set then provides a model for the  $\infty$ -groupoid of paths in this interpretation of  $(\infty, 1)$ -categories. This justifies the usage of Kan complexes as a model for  $\infty$ -groupoids.
3. It follows from the existence of the Quillen model structure on  $\mathbf{Top}$ , or more classically an argument involving Whitehead's theorem, that the inclusion of CW complexes into topological spaces,  $\mathbf{CW} \rightarrow \mathbf{Top}$  induces an equivalence  $\mathbf{CW}[H^{-1}] \simeq \mathbf{Top}[W^{-1}]$ , where  $H$  is the class of homotopy equivalences. Since most spaces of geometric interest at least have the homotopy type of a CW complex (see, for example, [Mil59]), it follows that if one is interested in studying the homotopy types of such classical spaces, one may perform such an investigation in  $\mathbf{Top}[W^{-1}]$ . Thus, the latter can rightfully be called the homotopy theory of spaces.

Combining these insights, one obtains an equivalence

$$\mathbf{Spaces} = \mathbf{CW}[H^{-1}] \simeq \mathbf{Top}[W^{-1}] \simeq \mathbf{Kan}[H_k^{-1}] = \mathbf{Grpd}_\infty,$$

as asserted in the homotopy hypothesis.

One of the central assertions of [AFR19] is a smooth stratified version of the homotopy hypothesis. Recall that, roughly speaking and in the broadest sense, a stratified space is a topological space together with a decomposition into disjoint pieces, the so-called strata, and a stratified map is a continuous map between such objects, which has the property that the image of each stratum in the source is completely contained in a stratum in the target. Stratifications of topological spaces often arise naturally when investigating spaces with singularities, by decomposing a singular space into manifold pieces (see, for example, [Whi65; Mat12; Mat73; Tho69]) and in these scenarios the set of strata tends to naturally inherit the structure of a poset from the topological closure relation. The homotopy theory of such stratified spaces, using homotopies that also preserve the strata, was first investigated in much detail by Quinn in [Qui88]. Quinn put particular emphasis on a class of stratified spaces with excellent homotopical properties, the so-called homotopically stratified spaces (called homotopically stratified sets in [Qui88]), proving, among other results, a stratified version of the H-cobordism theorem for manifold analogues of these objects. Following a more differential topological framework, which they developed in [AFT17], in [AFR19] Ayala, Francis, and Rosenblyum developed a homotopy theory of stratified spaces within the context of differential topology. One of their central assertions was the following statement:

Let  $\mathbf{Strat}_{C^\infty}$  denote the category of conically smooth stratified spaces (with conically smooth stratified maps), and  $H_s$  the class of stratified homotopy equivalences (in the conically smooth sense). Then there is a fully faithful embedding

$$\mathbf{Strat}^{C^\infty}[H_s^{-1}] \hookrightarrow \mathbf{Cat}_\infty$$

into the homotopy theory of  $\infty$ -categories<sup>1</sup>.

---

<sup>1</sup>At the current point in time, it appears that [AFR19] is missing a definition of what exactly the category of conically smooth stratified spaces they consider is. Note that one cannot simply use the category of conically smooth stratified spaces of [AFT17], since the latter does not contain the stratified simplices. Without a definition of the stratified smooth category accommodating such spaces with corners (and the development of the necessary theory), it is currently not possible to verify the truth of the smooth stratified homotopy hypothesis.

Furthermore, they conjectured a topological analogue of this result. This relies on the exit path construction of McPherson, Treuman, Woolf, and Lurie ([Woo09; Tre09; Lur17]) associating to a stratified space an  $\infty$ -category in which morphisms are given by paths which either remain within one stratum, or start in one stratum and immediately exit into a higher one.

**Conjecture 1.0.1.** [AFR19] Topological exit paths define a fully faithful functor

$$\mathbf{Exit}: \mathbf{Strat} \rightarrow \mathbf{Cat}_\infty$$

from a homotopy theory of topological stratified spaces  $\mathbf{Strat}$  into  $\infty$ -categories.

Again, just as in the case of the classical homotopy hypothesis, any answer to this conjecture must first provide a formal interpretation of the homotopy theories in question. In this case, the difficulty lies with the left-hand side, i.e. the question after a homotopy theory of topological stratified spaces. In recent years, three of these theories have been independently suggested (not all with the intent to tackle the topological stratified homotopy hypothesis). In the following  $\mathbf{Strat}$  will always denote the category of all poset stratified spaces, with stratified maps between them (see, for example, [DW22] for an overview). One way of defining such a homotopy theory is, of course, to specify a category of topologically stratified spaces, in these cases, the category of all poset stratified spaces with stratified maps, and then to localize the latter at an appropriate class of stratified weak equivalences.

1. In [Dou21c; Hen] Douteau and Henriques independently followed an insight of [Mil13], concerning Quinn's homotopically stratified spaces (Henriques covering the case of a fixed poset, Douteau also the global case). Stratum preserving homotopy equivalences between homotopically stratified spaces are characterized as precisely such maps that induce homotopy equivalence on the strata and on the so-called  $[p < q]$ -homotopy links (where  $p < q$  are elements of the stratifying poset) which are given by the spaces of exitpaths, starting in the  $p$ -stratum and immediately exiting into the  $q$ -stratum. Since for general poset-stratified spaces pairwise homotopy links are not enough to even guarantee that the underlying map is a weak equivalence of topological spaces, [Dou21c; Hen] additionally considered so-called generalized homotopy links, obtained by replacing the stratified interval by a stratified simplex. Then a weak equivalence of stratified spaces is defined to be a stratified map that induces an isomorphism on stratifying posets and weak equivalences on all generalized homotopy links (which include the strata). We call such stratified maps **poset preserving diagrammatic equivalences** and denote the resulting  $\infty$ -category obtained by localizing poset-preserving diagrammatic equivalences by  $\mathbf{Strat}^{\mathfrak{d}, \mathfrak{p}}$ . If one restricts to spaces over a fixed poset  $P$  and stratum preserving maps (that is, such stratified maps which are given by the identity on the poset level), then taking generalized homotopy links induces an equivalence of  $\infty$ -categories

$$\mathcal{H}o\mathbf{Link}: \mathbf{Strat}_P^{\mathfrak{d}} \simeq \mathbf{Fun}((\mathbf{sd}P)^{\text{op}}, \mathbf{Spaces})$$

where  $\mathbf{sd}P$  is the category whose objects are finite increasing sequences  $[p_0 < \dots < p_n]$  in  $P$  (so-called regular flags) and whose morphisms are inclusions of subsequences.

2. In [Hai23], Haine followed the insight that the categories of exit paths associated to a classical stratified space (for example, conically stratified space or homotopically stratified space) always comes together with a conservative functor into the stratification poset. This is simply due to the fact that restricted to each stratum the construction produces the classical  $\infty$ -groupoid of paths. Furthermore, the fact that exitpaths define an  $\infty$ -category at all implies that for such stratified spaces the natural maps from the generalized homotopy links into the homotopy pullbacks of pairwise homotopy links

$$\mathcal{H}o\mathbf{Link}_{[p_0 < \dots < p_n]}(\mathcal{X}) \rightarrow \mathcal{H}o\mathbf{Link}_{p_0 < p_1}(\mathcal{X}) \times_{X_{p_1}}^H \cdots \times_{X_{p_{n-1}}}^H \mathcal{H}o\mathbf{Link}_{p_{n-1} < p_n}(\mathcal{X})$$

are weak homotopy equivalences. Hence, only certain diagrams  $D \in \mathbf{Fun}((\mathrm{sd}P)^{\mathrm{op}}, \mathbf{Spaces})$  - for which the corresponding morphism is an isomorphism in the  $\infty$ -category of spaces - may arise as the homotopy-link diagram of classical examples of stratified spaces. Diagrams that satisfy this property are called **décollages** (see [Hai23; BGH18]). The full  $\infty$ -subcategory of décollages,  $\mathbf{Déc}_P$ , turns out to be a (left) localization of  $\mathbf{Fun}((\mathrm{sd}P)^{\mathrm{op}}, \mathbf{Spaces})$  (i.e. a reflective subcategory). [Hai23] then defines his class of weak equivalences of stratified spaces over a fixed poset  $P$  as the class of such stratified maps which map into isomorphisms under

$$\mathbf{Strat}_P \xrightarrow{\mathcal{H}\mathrm{oLink}} \mathbf{Fun}((\mathrm{sd}P)^{\mathrm{op}}, \mathbf{Spaces}) \rightarrow \mathbf{Déc}_P,$$

which makes Haine's theory a left localization of Douteau and Henriques'. One may then extend this class of weak equivalences to the case of varying poset  $\mathbf{Strat}$ , by requiring a weak equivalence to induce isomorphisms on stratifying posets and induce an equivalence over fixed posets under the induced identifications. We call such stratified maps **poset preserving categorical equivalences** (for reasons which will become apparent later) and denote the resulting homotopy theory by  $\mathbf{Strat}^{\mathrm{c},P}$ .

3. Finally, in [Nan19], Nand-Lal took the approach of directly transferring the weak equivalences along Lurie's stratified singular simplicial set construction, which provides a model for the  $\infty$ -category of exit paths associated to a stratified space<sup>2</sup>:

$$\mathrm{Sing}_s: \mathbf{Strat} \rightarrow \mathbf{sSet}.$$

More precisely, a stratified map  $\mathcal{X} \rightarrow \mathcal{Y}$  is called a **categorical equivalence**, if the induced simplicial map  $\mathrm{Sing}_s(\mathcal{X}) \rightarrow \mathrm{Sing}_s(\mathcal{Y})$  is a Joyal equivalence (also called categorical equivalences; see [Lur09]). One major difference compared to the previous two theories is that weak equivalences in this setting are not required to preserve the stratification poset. In this sense, the stratifying posets are only essential to a stratified homotopy type insofar as they determine what paths are exit paths (what simplices are exit-simplices). We denote the induced homotopy theory by  $\mathbf{Strat}^{\mathrm{c}}$ .

The first answer to Conjecture 1.0.1 was given in [Hai23], where Haine used a result of Douteau ([Dou21c, Thm. 3]) to prove an equivalence of  $\infty$ -categories:

$$\mathbf{Strat}_P^{\mathrm{c}} \simeq \mathbf{AStrat}_P,$$

for  $P$  a poset, with  $\mathbf{AStrat}_P$  the  $\infty$ -category of conservative functors with target  $P$ , so-called abstract stratified homotopy types. The equivalence is constructed by passing through the equivalence of  $\mathbf{Strat}_P^{\mathrm{c}}$  with  $\mathbf{Déc}_P$ , and then, in turn, constructing an equivalence of the latter with a model for  $\mathbf{AStrat}_P$  given in the language of complete segal spaces. In this sense, the equivalence does not directly pass through Lurie's  $\mathrm{Sing}_s$  construction (from a 1-categorical perspective at least), and it is a priori unclear whether weak equivalences in  $\mathbf{Strat}_P^{\mathrm{c}}$  can be characterized through the latter. From the equivalence  $\mathbf{Strat}_P^{\mathrm{c}} \simeq \mathbf{AStrat}_P$  one may furthermore derive the existence of a class of weak equivalences  $W_c$ , such that there is a global equivalence

$$\mathbf{Strat}[W_c^{-1}] \simeq \mathbf{Lay}_\infty$$

between  $\mathbf{Strat}^{\mathrm{c},P}$  and the fully subcategory of the  $\infty$ -category of small infinity categories  $\mathbf{Lay}_\infty$ , given by such  $\infty$ -categories which have the property that every endomorphism is an isomorphism, so-called *layered* infinity categories. This provided one answer to Conjecture 1.0.1. At the same time, if we aim to replicate the well-understood and tractable scenario of the classical homotopy hypothesis, further investigation is needed.

<sup>2</sup>Strictly speaking, Nand Lal works with a full subcategory of  $\mathbf{Strat}$ , but from a homotopy theoretic perspective this difference turns out to be irrelevant (see Proposition A.0.2)

**Question 1.0.2.** Namely, we want to answer the following questions:

- Q(1) Can the equivalence be presented through Lurie’s singular simplicial set (Exit path) construction <sup>3</sup> and is  $W_c$  the class of categorical equivalences?
- Q(2) Without additional structure,  $\infty$ -categorical localizations of 1-categories are generally fairly untractable objects. What can we say about  $\mathbf{Strat}^{c,p}$  from a stratified topological perspective? For example, can we express  $\mathbf{Strat}^c$  as a subcategory of stratified spaces  $\mathbf{C}$  localized at stratified homotopy equivalences, analogously to the situation of topological spaces and CW complexes? Such a category  $\mathbf{C}$  should contain as many stratified spaces of classical interest as possible, to allow us to investigate the stratified homotopy-theoretic properties of geometrically interesting examples through the language of  $\infty$ -categories.
- Q(3) In the same line of questioning as in the previous question: The category  $\mathbf{Strat}$  admits a naive notion of mapping space, with points given by stratified maps and paths given by stratified homotopies. For what stratified spaces can we expect the mapping space  $\mathbf{Strat}^c(\mathcal{X}, \mathcal{Y})$  to have the homotopy type of this naive mapping space, or similarly, which classical examples of stratified spaces are contained in  $\mathbf{C}$ ?
- Q(4) In a geometric scenario, what is the difference between the diagrammatic and categorical versions of stratified homotopy theory?
- Q(5) How can  $\infty$ -categorical construction, such as colimits in a quasi-category, i.e., homotopy colimits, be interpreted in terms of the 1-category  $\mathbf{Strat}$ . For example, when is a pushout diagram of stratified spaces homotopy pushout in one of the homotopy theories of stratified spaces above?
- Q(6) The sets of stratified maps can even be equipped with the structure of a stratified space, with each stratum corresponding to a map of the underlying posets (see [Nan19]). In [Hug99], such stratified (decomposed) mapping spaces and their relations with stratified notions of fibrations were studied. The central result states a stratified exit path version of the classical path-space fibration. Can we replicate this result in the stratified homotopy theories described above using methods of modern abstract homotopy theory?
- Q(7) What is the relationship of the homotopy theories described above with the homotopy theory for conically smooth stratified spaces described in [AFR19]?

Classically, such questions concerning the relationship of a homotopy theory and a 1-category from which it is obtained by localization are answered through the language of model categories. Thus, we may at least partially rephrase the questions above:

**Question 1.0.3.** Do the classes of weak equivalences defining the homotopy theories  $\mathbf{Strat}^{d,p}$ ,  $\mathbf{Strat}^{c,p}$ ,  $\mathbf{Strat}^c$  extend to model structures on  $\mathbf{Strat}$  in which stratified spaces of classical interest (for example, Whitney stratified spaces or, more recently, conically smooth stratified spaces) are bifibrant. Supposing an affirmative answer, what are the properties of this model category, such as admitting a simplicial structure, Cartesian closedness, cofibrant generation? Can the equivalence  $\mathbf{Strat}^c \simeq \mathcal{Lay}_\infty$  be presented in terms of a Quillen equivalence, using Lurie’s stratified singular simplicial set construction?

In [Dou21c; Hen] model structures on  $\mathbf{Strat}$  presenting  $\mathbf{Strat}^{d,p}$  were defined. However, these model structures fail the criterion of having convenient cofibrant-fibrant objects: Not even the

---

<sup>3</sup>In [Hai23] this is claimed. We note that the proof given there ends up proving a somewhat different statement. The confusion arises from the usage of multiple different definitions of complete Segal space. See Remark D.0.1.

cone on a manifold is a cofibrant object. In [Nan19], the author defined a model structure for the subcategory of such stratified spaces,  $\mathcal{X}$ , for which  $\text{Sing}_s(\mathcal{X})$  is a quasi-category and used it to prove a Whitehead theorem for stratified spaces. The existence of a (semi)model structure as above was left open as a conjecture ([Nan19, p. 8.4.1]). While this makes, for example, piecewise linear pseudo-manifolds bifibrant, a full model category (in particular fibrant replacement) is needed to present the stratified homotopy hypothesis. Finally, in [DW22], we show that such model structures cannot exist (see Proposition 4.0.1, for a detailed proof adapted to the setting of this paper).

## 1.1 Content of this article

The goal of this paper is to prove that we can answer Question 1.0.3 affirmatively if we allow ourselves to generalize to (left) semimodel categories (see, for example, [Bar10; WB23]). We note that in practice these usually turn out to be just as powerful as model categories. In particular, the existence of semimodel structures allows us to obtain answers to Question 1.0.2. Building on the work of [Hai23], [Dou21c], and [Dou21b], we have built the foundation to prove the existence of these semimodel structures in the two preceding papers: [Waa24b] and [Waa24a]. What remains to do is to combine these results as follows:

1. In [Dou21b] and [Hai23] Douteau and Haine both respectively built simplicial counterparts to their stratified homotopy theories for the case of a fixed posets  $P$ . In [Waa24a], we extend these models to model structures for stratified simplicial sets,  $\mathbf{sStrat}$ , with varying stratification poset. Doing so, we present simplicial model categories presenting the homotopy theories  $\mathbf{Strat}^{c,p}$ ,  $\mathbf{Strat}^{d,p}$ ,  $\mathbf{Strat}^c$  as well as a fourth theory  $\mathbf{Strat}^d$ , which is the non-poset-preserving analogue to  $\mathbf{Strat}^{d,p}$ . We recall these results in Section 3.2.
2. We want to construct model structures for  $\mathbf{Strat}^{c,p}$ ,  $\mathbf{Strat}^{d,p}$ ,  $\mathbf{Strat}^c$  by transferring the model structures on the simplicial side along the stratified singular simplicial set and realization adjunction. To this end, we need to prove that the latter functor  $\mathbf{Strat} \rightarrow \mathbf{sStrat}$  creates weak equivalences for the respective theories. This is proven in Section 3.3, in fact, even more we show that the functor induces an equivalence of relative categories (Theorem 3.3.1), which follows from [DW22, Thm. 5.1].
3. Then, in Section 4, we transfer the model structures from the combinatorial framework to the topological one. To this end, we prove a transfer lemma for semimodel categories Lemma 4.1.6. Applying this result requires a deeper understanding of the interaction of the homotopy link functor with topological stratified constructions, such as pushouts along inclusions of cell complexes. This is achieved through [Waa24b, Thm. B], in which we prove the existence of certain regular neighborhoods for multistrata interactions in stratified cell complexes. In many ways, these are the topological (as opposed to smooth) analogues of the Unzip construction of [AFT17]: They provide the necessary glue to connect the geometry/topology of stratified spaces with their homotopy theory.

We may then combine these results to obtain the following result, which (using standard results on (semi)model categories) ultimately answers Questions Q(2), Q(5) and Q(6) (see, particularly, Section 4.4).

**Theorem A** (Theorems 3.3.1, 4.2.7 and 4.2.10 and Corollaries 4.2.3 and 4.3.3). *The category of stratified topological spaces  $\mathbf{Strat}$  admits the structure of three different simplicial, cofibrantly generated, Cartesian closed semimodel categories  $\mathbf{Strat}^{d,p}$ ,  $\mathbf{Strat}^{c,p}$  and  $\mathbf{Strat}^c$ , with weak equivalences given respectively by the poset preserving diagrammatic, poset preserving categorical and*

categorical equivalences. They are right-transferred along the adjunction

$$|-|_s: \mathbf{sStrat} \rightleftarrows \mathbf{Strat}: \text{Sing}_s$$

from their respective counterpart on  $\mathbf{sStrat}$ , and make the adjunction a Quillen equivalence, creating weak equivalences in both directions.

As a corollary of this result and [Waa24a, Theorem 3.3.6], we obtain the following version of the stratified homotopy hypothesis, providing answers to Question Q(1):

**Theorem B** (Theorem 4.4.4). *Mapping a simplicial set to its stratified realization (as in [Nan19]), and conversely mapping a stratified space  $\mathcal{X}$  to the underlying simplicial set of  $\text{Sing}_s \mathcal{X}$ , induces a Quillen equivalence of (semi)model categories*

$$\mathbf{sSet}^{\triangleright} \xrightleftharpoons{\cong} \mathbf{Strat}^c,$$

which creates weak equivalences in both directions.

Here,  $\mathbf{sSet}^{\triangleright}$  is the left Bousfield localization of the Joyal model structure on  $\mathbf{sSet}$  corresponding to layered  $\infty$ -categories.

$\mathbf{Strat}^{c,p}$  is obtained from  $\mathbf{Strat}^{\triangleright,p}$  in terms of a left Bousfield localization at stratified inner horn inclusions (providing a first step towards an answer to Question Q(4)), and  $\mathbf{Strat}^c$  is in turn obtained from  $\mathbf{Strat}^{c,p}$  in terms of a right Bousfield localization (Theorem 3.3.1).

To obtain a similarly convenient situation to the setup of the classical homotopy hypothesis, an answer to Question Q(3) remains to be obtained. We do so in Section 5. It is an immediate consequence of the way that the model structures are constructed via transfer from stratified simplicial sets that any stratified space which admits a piecewise linear structure (or, more generally, a cell structure) that is compatible with the stratification is cofibrant in the semimodel categories presenting  $\mathbf{Strat}^{\triangleright,p}$  and  $\mathbf{Strat}^{c,p}$ ,  $\mathbf{Strat}^{\triangleright,p}$  and  $\mathbf{Strat}^{c,p}$ . Hence, for example, by [Gor78], all Whitney stratified spaces are cofibrant. Even more, we show Proposition 5.2.10, from which it follows that any stratified space (over a finite poset) whose strata are manifolds that additionally admit a stratified notion of mapping cylinder neighborhood is cofibrant. In particular, assuming the correctness of a result of [AFR19], this implies that conically smooth stratified spaces are cofibrant. The semimodel category  $\mathbf{Strat}^c$ , turns out to be a right Bousfield localization of  $\mathbf{Strat}^{c,p}$ , whose cofibrant objects are precisely cofibrant objects in  $\mathbf{Strat}^{c,p}$  that fulfill a weak version of the classical frontier condition and have connected strata (Proposition 5.4.5). Conceptually, these are the stratified spaces (cofibrant in  $\mathbf{Strat}^{c,p}$ ) in which the stratification poset structure arises entirely in terms of the topological relations of the strata.

Faced with several different model structures and homotopy theories for stratified spaces, the obvious question between the relationship and differences between these theories, in particular in application to geometric examples, arises. We have already illustrated above that the passage from  $\mathbf{Strat}^{c,p}$  to  $\mathbf{Strat}^c$  essentially amounts to requiring that the poset structure is intrinsic to the topology of the space. The difference between  $\mathbf{Strat}^{c,p}$  and  $\mathbf{Strat}^{\triangleright,p}$  (that is, answering Question Q(4)) is more subtle. As the cofibrant objects in the categorical and the diagrammatic semimodel categories agree, the difference between the resulting homotopy theories must lie in the conditions for fibrancy. It is a result of [Lur17] that conically smooth stratified spaces are fibrant in  $\mathbf{Strat}^c$ , and hence also in  $\mathbf{Strat}^{c,p}$  and  $\mathbf{Strat}^{\triangleright,p}$ . In [Nan19], it was shown that the same holds for Quinn's homotopically stratified spaces ([Qui88]). This already covers almost any classically relevant example of stratified spaces. Thus, it appears that at least in a geometric scenario there is not much difference between the two homotopy theories  $\mathbf{Strat}^{c,p}$  and  $\mathbf{Strat}^{\triangleright,p}$  at all. In fact, we make this result rigorous in terms of the following answer to Questions Q(3) and Q(4).

**Theorem C.** *Proposition 5.1.4 and Theorem 5.1.6 Let  $\mathcal{X} \in \mathbf{Strat}$  be a metrizable stratified space. Then the following conditions are equivalent:*

- (i)  $\mathcal{X}$  is fibrant in  $\mathbf{Strat}^c$ ;
- (ii)  $\mathcal{X}$  is fibrant in  $\mathbf{Strat}^{c,p}$ ;
- (iii)  $\mathcal{X}$  is fibrant in  $\mathbf{Strat}^{0,p}$ ;
- (iv) For any pair of strata  $[p < q]$ , the starting point evaluation map  $\mathcal{H}\text{olink}_{p < q}(\mathcal{X}) \rightarrow \mathcal{X}_p$  is a Serre fibration.

*In particular, when restricted to metrizable stratified spaces, the homotopy theories defined by  $\mathbf{Strat}^{c,p}$  and  $\mathbf{Strat}^{0,p}$  (in terms of simplicial categories of bifibrant objects) agree.*

Similarly to the case of the classical homotopy hypothesis, it can be useful to restrict to a class of particularly convenient stratified spaces that mimic some of the properties of CW complexes. This is handled by Corollary 5.4.10, which in particular states that the bifibrant stratified spaces in  $\mathbf{Strat}^c$  are precisely the retracts of the so-called CFF stratified spaces, i.e., the stratified spaces  $\mathcal{X} \in \mathbf{Strat}$  fulfilling:

1.  $\text{Sing}_s \mathcal{X}$  is a quasi-category.
2.  $\mathcal{X}$  admits a stratified cell structure.
3.  $\mathcal{X}$  has non-empty connected strata, and the stratification of  $\mathcal{X}$  is induced by the frontier condition.

It follows that we may rephrase Theorem B as the following statement:

**Theorem B'.** *Denote by  $\mathbf{CFF}$  the full subcategory of  $\mathbf{Strat}$  given by CFF stratified spaces and let  $H_s$  be the class of stratified homotopy equivalences in  $\mathbf{CFF}$ . Lurie's exit path construction ([Lur17]) induces an equivalence of  $\infty$ -categories*

$$\text{Exit: } \mathbf{CFF}[H_s^{-1}] \xrightarrow{\simeq} \mathcal{Lay}_\infty$$

*where  $\mathcal{Lay}_\infty$  denotes the  $\infty$ -category of small layered  $\infty$ -categories.*

Finally, let us comment on the relationship of the homotopy theories investigated in this article with the homotopy theory of conically smooth stratified spaces investigated in [AFR19] (Question Q(7)). Supposing the existence of stratified mapping cylinder neighborhoods (tubular neighborhoods), as asserted in [AFT17, Prop. 8.2.3], it follows from Proposition 5.2.10, Proposition 5.4.5 and Theorem C that conically smooth stratified spaces are bifibrant in  $\mathbf{Strat}^c$ . Note that it follows from this result and Theorem B that the following two statements are equivalent:

1. The exit path construction induces a fully faithful embedding from the  $\infty$ -category of conically smooth stratified spaces into  $\infty$ -categories (as asserted in [AFR19]).
2. For any pair of conically smooth stratified spaces  $\mathcal{X}, \mathcal{Y}$ , the natural map

$$\mathbf{Strat}^{c,\infty}(\mathcal{X}, \mathcal{Y}) \rightarrow \mathbf{Strat}(\mathcal{X}, \mathcal{Y})$$

from the mapping space of conically smooth maps (defined in [AFR19]) into the mapping space of continuous stratified maps is a homotopy equivalence of Kan complexes.

In this sense, the smooth version of the stratified homotopy hypothesis as it is stated above is equivalent to a stratified smooth approximation theorem, as conjectured in [AFT17, Conjecture 1.5.1].



## 2 Preliminaries

In this section, we introduce and recall some of the necessary language and notation, especially for 1-categorical aspects of stratified spaces and higher homotopical frameworks.

### 2.1 Models for $(\infty, 1)$ -categories

This paper is concerned with investigating several homotopy theories of stratified spaces. By a *homotopy theory* we mean an  $(\infty, 1)$ -category, not restricting to a specific model for  $(\infty, 1)$ -categories. For the sake of readability, we will always drop the 1 and use the term  $\infty$ -category and homotopy theory synonymously. For our purposes, it is going to be extremely useful to have several different models for  $\infty$ -categories available. Specifically, we are going to use the following models.

- Notation 2.1.1.**
1. The relative categories of [BK12b], given by pairs  $(\mathbf{C}, W)$  of a 1-category  $\mathbf{C}$  with a wide subcategory  $W \subset \mathbf{C}$ . Names for relative categories will always begin with an italic letter, that is, they will be of the form: **Name**.
  2. Categories enriched over simplicial sets, also called simplicial categories (see [DK80c; Ber07a]): Names for simplicial categories will always be underlined, that is, written in the form Name.
  3. Quasi-categories in the sense of [Lur09]: Names for quasi-categories will always be bold and begin with a calligraphic letter, i.e. they will be of the form:  $\mathcal{N}\mathbf{ame}$ .

If  $\mathbf{C}$  is an  $\infty$ -category (in particular a 1-category), the mapping spaces between objects  $X, Y \in \mathbf{C}$  will be denoted by  $\mathbf{C}(X, Y)$ . In the case of simplicial categories, these have an explicit model in the obvious way. For relative categories, they are given by the hammock localization of [DK80a; BK12a]. For quasi-categories, we use any of the equivalent models of mapping spaces of [Lur09].

**Notation 2.1.2.** To study  $\infty$ -categories through some underlying 1-category which they are a localization of, we are going to make use of the language of (semi)model categories (see [Hir09] for model categories and [Bar10; WB23] for a good overview of semimodel categories). We will usually denote ordinary categories in bold letters in the form **Name**). The model structure will always be marked by adding some additional ornamentation - in the form Name<sup>5</sup> or **Name**<sup>5</sup> - to the name of the underlying simplicial or 1-category.

**Notation 2.1.3.** Often we are going to pass between different models of a homotopy theory. These passages will always follow the following ruleset for nomenclature:

1. Starting with a relative category **Name** or simplicial category Name, the underlying 1-category will be denoted **Name**. Similarly, if Name<sup>5</sup> is a simplicial model category, then we denote its underlying 1-model category by **Name**<sup>5</sup>.
2. Starting with a model category **Name**<sup>5</sup>, we denote by  $\mathbf{Name}^5$  the relative category obtained by its underlying 1-category, together with the wide subcategory of weak equivalences.
3. Starting with a relative category **Name** =  $(\mathbf{Name}, W)$ , we denote by  $\mathcal{N}\mathbf{ame}$  the quasi-category **Name** $[W^{-1}]$  obtained by taking the nerve of the underlying 1-category of **Name**, and then localizing at  $W$ .
4. In particular, following this language, if we start with a model category **Name**<sup>5</sup>, then the relative category **Name**<sup>†</sup> and the quasi-category  $\mathcal{N}\mathbf{ame}^5$  model the same  $\infty$ -category.

5. If we start with a simplicial model category  $\underline{\mathbf{Name}}^s$ , then we denote by  $\underline{\mathbf{Name}}^{s,o}$  the full simplicial subcategory of bifibrant objects. Following our notation, under the Quillen equivalence between simplicial categories and quasi-categories of [Ber07b],  $\underline{\mathbf{Name}}^{s,o}$  models the same  $\infty$ -category as  $\mathcal{N}\mathbf{ame}^s$  (this follows from [DK80b, Prop. 4.8] together with [Hin16, Prop. 1.2.1]).

For the sake of completeness, one could of course also introduce notation to pass from relative to simplicial categories, etc., but we will not make use of this change of model in this article, and thus do not take this extra step.

## 2.2 Poset stratified spaces

Next, let us introduce the basic objects of study: Topological spaces which are stratified over a partially ordered set (poset). We begin with notation for the world of posets.

**Notation 2.2.1.** We are going to use the following terminology and notation for partially ordered sets, drawn partially from [Dou21a] and [Hai23]:

- We denote by  $\mathbf{Pos}$  the category of partially ordered sets, with morphisms given by order-preserving maps.
- We denote by  $\Delta$  the full subcategory of  $\mathbf{Pos}$  given by the finite, nonempty, linearly ordered posets of the form  $[n] := \{0, \dots, n\}$ .
- Given  $P \in \mathbf{Pos}$ , we denote by  $\Delta_P$  the slice category  $\Delta/P$ . That is, objects are given by arrows  $[n] \rightarrow P$  in  $\mathbf{Pos}$ ,  $n \in \mathbb{N}$ , and morphisms are given by commutative triangles.
- We denote by  $\text{sd}P$  the **subdivision of  $P$** , given by the full subcategory of  $\Delta_P$  of such arrows  $[n] \rightarrow P$ , which are injective.
- The objects of  $\Delta_P$  are called **flags of  $P$** . We represent them by strings  $[p_0 \leq \dots \leq p_n]$ , of  $p_i \in P$ .
- Objects of  $\text{sd}P$  are called regular **flags of  $P$** . We represent them by strings  $[p_0 < \dots < p_n]$ , of  $p_i \in P$ .

**Notation 2.2.2.**  $\mathbf{Top}$  is going to denote either of the following three categories of topological spaces.

1. The category of **all topological spaces**, which we will also refer to as **general** topological spaces.
2. The category of compactly generated topological spaces, i.e. such spaces which have the final topology with respect to compact Hausdorff spaces (see, for example, [Rez17]).
3. The category of  $\Delta$ -generated topological spaces, i.e. such spaces which have the final topology with respect to realizations of simplices, or equivalently just with respect to the unit interval (compare [Dug03; Gau21]).

We denote by  $\mathbf{sSet}$  the simplicial category of simplicial sets, i.e. the category of set-valued presheaves on  $\Delta^{\text{op}}$ , equipped with the canonical simplicial structure induced by the product (see [Lur09] for all of the standard notation used for simplicial sets). We denote by  $|-|: \mathbf{sSet} \rightarrow \mathbf{Top}$  the realization functor of simplicial sets and by  $\text{Sing}: \mathbf{Top} \rightarrow \mathbf{sSet}$  its right adjoint.  $\mathbf{Top}$  naturally carries the structure of a simplicial category, tensored and powered over  $\mathbf{sSet}$ , induced by left Kan extension of the construction

$$T \otimes \Delta^n := T \times |\Delta^n|.$$

We denote the resulting simplicial category by **Top**. Furthermore, we will always consider Top to be equipped with the Quillen model structure [Qui67], which makes  $|-| \dashv \text{Sing}$  a simplicial Quillen equivalence, between **Top** and **sSet** (with the latter equipped with the Kan-Quillen model structure), which creates all weak equivalences in both directions.

**Remark 2.2.3.** Note that one commonly only defines the simplicial structure for compactly or  $\Delta$ -generated spaces. However, this is mostly due to the fact that for general topological spaces  $T$  and infinite simplicial sets  $K$ , the tensor  $T \otimes K$  does not agree with the inner product  $T \times |K|$ . Instead, it is given by a colimit of products of  $T$  with the simplices of  $K$ . Similarly, the power  $T^K$  is not given by an internal mapping space (which does not even necessarily exist for arbitrary  $K$ ) but by the limit of the mapping spaces with source given by simplices of  $K$ , which are equipped with the compact open topology. We want to emphasize that for the resulting homotopy theory the choice in underlying set-theoretic assumptions on topological spaces is immaterial.

For the remainder of this subsection, we fix some category of topological spaces **Top** as in Notation 2.2.2.

**Notation 2.2.4.** Having fixed a category of topological spaces **Top**, we then use the following notation for stratified topological spaces (all of these constructions already appear in [Dou21c] among other places). The notation and language is mostly analogous with the language for the simplicial framework.

- We think of the 1-category **Pos** as naturally embedded in **Top**, via the Alexandrov topology functor  $\mathcal{A}: \mathbf{Pos} \rightarrow \mathbf{Top}$ , equipping a poset  $P$  with the topology where the closed sets are given by the downward closed sets. By abuse of notation, we usually just write  $P$ , for the Alexandrov space corresponding to the poset  $P$  (compare [DW22, Def. 2.2]).
- For  $P \in \mathbf{Pos}$ , we denote by  $\mathbf{Strat}_P$  the slice category  $\mathbf{Top}_{/P}$ . We treat  $\mathbf{Strat}_P$  as a simplicial category, denoted  $\mathbf{Strat}_P$ , with the structure inherited from Top (see [DW22, Recol. 2.13] for a detailed definition in the  $\Delta$ -generated case).
- Objects of  $\mathbf{Strat}_P$  are called  **$P$ -stratified spaces**. They are given by a tuple  $(\mathcal{X}, s: X \rightarrow P)$ . In the literature, a  $P$ -stratified space  $(\mathcal{X}, s: X \rightarrow P)$  is often simply referred to by its underlying space  $X$ , omitting the so-called **stratification**  $s: X \rightarrow P$ . We are not going to adopt this notation here and will generally use calligraphic letters for stratified spaces and stick to the notational convention  $\mathcal{X} = (X, s_{\mathcal{X}})$  to refer to the underlying space and stratification.
- Morphisms in  $\mathbf{Strat}_P$  are called **stratum preserving maps**.
- Given a map of posets  $f: Q \rightarrow P$  and  $\mathcal{X} \in \mathbf{Strat}_P$ , we denote by  $f^*\mathcal{X} \in \mathbf{Strat}_Q$  the stratified space  $X \times_f P \rightarrow Q$ . We are mostly concerned with the case where  $f$  is given by the inclusion of a singleton  $\{p\}$ , of a subset  $\{q \sim p \mid q \in P\}$ , for  $p \in P$  and  $\sim$  some relation on the partially ordered set  $P$  (such as  $\leq$ ), or more generally, a subposet  $Q \subset P$ . We then write  $\mathcal{X}_p$  (or, respectively,  $\mathcal{X}_{\sim p}$ ,  $\mathcal{X}_Q$ ) instead of  $f^*\mathcal{X}$ . The spaces  $\mathcal{X}_p$ , for  $p \in P$  are called the **strata of  $\mathcal{X}$** .
- For  $f: Q \rightarrow P$  in **Pos**, we denote by  $f_!$  the left adjoint to the simplicial functor  $f^*: \mathbf{Strat}_Q \rightarrow \mathbf{Strat}_P$ .  $f_!$  is given on objects by  $(s_{\mathcal{X}}: X \rightarrow Q) \mapsto (f \circ s_{\mathcal{X}}: X \rightarrow P)$ .
- We denote by **Strat** the category of all stratified simplicial spaces, given by the pullback of 1-categories  $\mathbf{Top}^{[1]} \times_{\text{ev}_1} \mathbf{Pos}$ , under the Alexandrov Topology functor  $\mathcal{A}: \mathbf{Pos} \hookrightarrow \mathbf{Top}$ . In other words, objects of **Strat** are given by maps  $X \rightarrow P$ , for some  $P \in \mathbf{Pos}$ , and morphisms are given by commutative squares (see also [DW22, Def. 2.11]). Morphisms are called **stratified maps**.

- Given  $\mathcal{X} \in \mathbf{Strat}$ , we are going to use the notations  $\mathcal{X} = (X, s_{\mathcal{X}}, P_{\mathcal{X}})$  to refer, respectively, to the underlying space, the stratification, and the poset and proceed analogously for morphisms.
- $\mathbf{Strat}$  is equipped with the structure of a simplicial category (tensoring and powered over  $\mathbf{sSet}$ ),  $\mathbf{Strat}$ , with the simplicial structure induced by  $\mathcal{X} \otimes \Delta^n = (X \times \Delta^n \rightarrow X \rightarrow P_{\mathcal{X}})$ . Note that since  $\mathcal{X} \otimes -$  preserves colimits, this means that  $\mathcal{X} \otimes \emptyset$  is not stratified over  $P$ , but instead over the empty poset. Simplicial homotopies, that is, homotopies with respect to the cylinder  $- \otimes \Delta^1$  in  $\mathbf{Strat}$ , are called **stratified homotopies**. Simplicial homotopy equivalences are called **stratified homotopy equivalences**.
- The forgetful functor  $\mathbf{Strat} \rightarrow \mathbf{Top}$ ,  $\mathcal{X} \mapsto X$ , has a right adjoint. It is given by mapping  $T \in \mathbf{Top}$  to the trivially stratified space  $(T \rightarrow [0])$ . By abuse of notation, we will often write  $T$  to refer to the trivially stratified space associated to a space  $T$ .

**Remark 2.2.5.** Both  $\mathbf{Strat}$  and  $\mathbf{Strat}_P$ , for  $P \in \mathbf{Pos}$ , are bicomplete categories (see, for example, [Dou21c]). Limits and colimits in  $\mathbf{Top}$  are simply given by the limits and colimits in a slice category. Colimits in  $\mathbf{Strat}$  are computed by taking the colimit both on the space and on the poset level. Limits in  $\mathbf{Strat}$  are computed by taking the limit on the space and poset level, and then pulling back the map  $\varprojlim_i X_i \rightarrow \varprojlim_i \mathcal{A}(P_i)$  along the natural comparison map  $\mathcal{A}(\varprojlim_i P_i) \rightarrow \varprojlim_i \mathcal{A}(P_i)$ , which is an isomorphism for finite diagrams.

**Construction 2.2.6.** If our choice of  $\mathbf{Top}$  is Cartesian closed (that is, if we are in the compactly generated or  $\Delta$ -generated case) then  $\mathbf{Strat}$  is also a Cartesian closed category (see also [Nan19], for the slightly different setting of surjectively stratified spaces). We use the notation  $X^Y$  to refer to exponential in a Cartesian closed category. To be able to distinguish exponential objects in posets from those in topological spaces, we use the notation  $\mathcal{A}(P)$  to denote the Alexandrov space associated to a poset  $P$ . Given a stratified space  $\mathcal{X} \in \mathbf{Strat}$ , the right adjoint  $-^{\mathcal{X}}$  to the functor  $- \times \mathcal{X}: \mathbf{Strat} \rightarrow \mathbf{Strat}$  is constructed as follows. Given  $\mathcal{Z}$  in  $\mathbf{Strat}$ , consider a pullback square in  $\mathbf{Top}$

$$\begin{array}{ccc}
\mathcal{A}(P_{\mathcal{Z}}^{P_{\mathcal{X}}}) \times_{\mathcal{A}(P_{\mathcal{Z}})^{P_{\mathcal{X}}}} Z^{\mathcal{X}} & \longrightarrow & Z^{\mathcal{X}} \\
\downarrow & & \downarrow \\
\mathcal{A}(P_{\mathcal{Z}}^{P_{\mathcal{X}}}) & \longrightarrow & \mathcal{A}(P_{\mathcal{Z}})^{\mathcal{A}(P_{\mathcal{X}})} \longrightarrow \mathcal{A}(P_{\mathcal{Z}})^{\mathcal{X}},
\end{array} \tag{1}$$

where  $P_{\mathcal{Z}}^{P_{\mathcal{X}}}$  is the poset obtained by equipping  $\mathbf{Pos}(P_{\mathcal{X}}, P_{\mathcal{Z}})$  with the poset structure given by

$$f \leq g : \iff f(p) \leq g(p), \forall p \in P_{\mathcal{X}},$$

which defines the exponential object  $P_{\mathcal{Z}}^{P_{\mathcal{X}}}$  in  $\mathbf{Pos}$ . The lower left horizontal map between the resulting space equipped with the Alexandrov topology into the mapping space  $\mathcal{A}(P_{\mathcal{Z}})^{\mathcal{A}(P_{\mathcal{X}})}$  is always continuous, but generally only a homeomorphism if  $P_{\mathcal{X}}$  and  $P_{\mathcal{Z}}$  are finite ([May16, Cor. 2.2.11]). The stratified mapping space  $Z^{\mathcal{X}}$  is defined by the left vertical of this pullback diagram<sup>4</sup>. This construction (and its functoriality in stratified maps) induces the right adjoint  $-^{\mathcal{X}}: \mathbf{Strat} \rightarrow \mathbf{Strat}$  to  $- \times \mathcal{X}$ . In particular, we may treat  $\mathbf{Strat}$  as a symmetric monoidal category (with monoidal structure induced by the product) enriched over itself.

<sup>4</sup>Note that the underlying set of  $\mathcal{A}(P_{\mathcal{Z}}^{P_{\mathcal{X}}}) \times_{\mathcal{A}(P_{\mathcal{Z}})^{P_{\mathcal{X}}}} Z^{\mathcal{X}}$  is in natural bijection with  $\mathbf{Strat}(\mathcal{X}, \mathcal{Z})$ , which shows that we can interpret this construction as a choice of topology on  $\mathbf{Strat}(\mathcal{X}, \mathcal{Z})$ , which will generally be finer than the initial topology inherited from  $Z^{\mathcal{X}}$ .

## 2.3 Stratified simplicial sets and stratified realization

The approach to constructing and investigating homotopy theories of stratified spaces that we take in this article is to prove that a lot of homotopy theory can be transferred from the combinatorial world. Let us quickly recall some notation and terminology concerning stratified simplicial sets, as introduced and investigated in [Dou21a; Hai23]. We have surveyed and expanded the results on homotopy theories of stratified simplicial sets in [Waa24a], to which we refer for more details.

**Notation 2.3.1.** We use the following terminology and notation for (stratified) simplicial sets, drawn partially from [Dou21a] and [Hai23]:

- When we treat **sSet** as a model category, this will generally be with respect to the Kan-Quillen model structure (see [Qui67]), unless otherwise noted. When we use Joyal’s model structure for quasi-categories ([JT08]) instead, we will denote this model category by **sSet**<sup>J</sup>.
- We think of **Pos** as being naturally embedded in **sSet**, via the nerve functor (compare [Hai23]). By abuse of notation, we just write  $P$ , for the simplicial set given by the nerve of  $P \in \mathbf{Pos}$ .
- For  $P \in \mathbf{Pos}$ , we denote by **sStrat** <sub>$P$</sub>  the slice category **sSet**<sub>/ $P$</sub> , which is equivalently given by the category of set-valued presheaves on  $\Delta_P$ . We treat **sStrat** <sub>$P$</sub>  as a simplicial category, denoted **sStrat** <sub>$P$</sub> , with the structure inherited from **sSet** (see [DW22, Recol. 2.21.]). Objects of **sStrat** <sub>$P$</sub>  are called  **$P$ -stratified simplicial sets**.
- Most of the remaining language and notation we are going to use for stratified simplicial sets can be copied mutatis mutandis from the topological setting in Section 2.2. See [Waa24a, Subsec. 1.1], for a detailed list.
- The forgetful functor **sStrat**  $\rightarrow$  **sSet**,  $\mathcal{X} \mapsto X$  will be denoted  $\mathcal{F}$ , has a right adjoint and a left adjoint. The left adjoint is given by left Kan extending the functor on simplices:  $\Delta^n \mapsto \{\Delta^n \xrightarrow{1_{\Delta^n}} \Delta^n = [n]\}$ . We denote it by  $\mathcal{L}: \mathbf{sSet} \rightarrow \mathbf{sStrat}$ . The right adjoint is given by mapping  $K \in \mathbf{sSet}$  to the trivially stratified simplicial set  $\{K \rightarrow [0]\}$ . By abuse of notation, we will often write  $K$  to refer to the trivially stratified simplicial set associated to a simplicial set  $K$ .

**Notation 2.3.2.** We are going to need some additional notation for flags and stratified simplices.

- For a flag  $\mathcal{J} = [p_0 \leq \dots \leq p_n] \in \Delta_P$ , we write  $\Delta^{\mathcal{J}}$  for the image of  $\mathcal{J}$  in **sStrat** <sub>$P$</sub>  under the Yoneda embedding  $\Delta_P \hookrightarrow \mathbf{sStrat}_P$ . Equivalently,  $\Delta^{\mathcal{J}}$  is given by the unique simplicial map  $\Delta^n \rightarrow P$  mapping  $i \mapsto p_i$ .  $\Delta^{\mathcal{J}}$  is called the **stratified simplex associated to  $\mathcal{I}$** .
- Given a stratified simplex  $\Delta^{\mathcal{J}}$ , for  $\mathcal{J} = [p_0 \leq \dots \leq p_n]$ , we write  $\partial\Delta^{\mathcal{J}}$  for its **stratified boundary**, given by the composition  $\partial\Delta^n \rightarrow \Delta^n \rightarrow P$ .
- Furthermore, for  $0 \leq k \leq n$ , we write  $\Lambda_k^{\mathcal{J}} \subset \Delta^{\mathcal{J}}$ , for the stratified subsimplicial set given by the composition  $\Lambda_k^n \rightarrow \Delta^n \rightarrow P$  (we use the horn notation as in [Lur09]). The stratum preserving map  $\Lambda_k^{\mathcal{J}} \hookrightarrow \Delta^{\mathcal{I}}$  is called the **stratified horn inclusion associated to  $\mathcal{J}$  and  $k$** . The inclusion  $\Lambda_k^{\mathcal{J}} \hookrightarrow \Delta^{\mathcal{I}}$  is called **admissible**, if  $p_k = p_{k+1}$  or  $p_k = p_{k-1}$ . The inclusion  $\Lambda_k^{\mathcal{J}} \hookrightarrow \Delta^{\mathcal{I}}$  is called **inner** if  $0 < k < n$ .
- It will also be convenient to have a concise notation for the images of simplices, horns, and boundaries under  $\mathcal{L}: \mathbf{sSet} \rightarrow \mathbf{sStrat}$ . These are denoted by replacing the exponent  $n \in \mathbb{N}$ , by the poset  $[n]$ . That is, we write  $\Delta^{[n]} := \mathcal{L}(\Delta^n)$ ,  $\partial\Delta^{[n]} := \mathcal{L}(\partial\Delta^n)$ ,  $\Lambda_k^{[n]} := \mathcal{L}(\Lambda_k^n)$ , for  $0 \leq k \leq n$ .

**Recollection 2.3.3.** [Dou21a] For fixed  $P \in \mathbf{Pos}$ , the two simplicial categories  $\mathbf{Strat}_P$  and  $\mathbf{sStrat}_P$  are connected through a singular simplicial set, realization style adjunction, denoted

$$|-|_s: \mathbf{sStrat}_P \rightleftarrows \mathbf{Strat}_P: \text{Sing}_s.$$

The left adjoint is constructed by mapping a stratified simplex  $\Delta^{\mathcal{J}} \rightarrow P$ , with  $\mathcal{J} = [p_0, \leq \dots, \leq p_n]$ , to the stratified space

$$\begin{aligned} |\Delta^n| &\rightarrow P \\ x &\mapsto \sup\{p_i \in \mathcal{J} \mid x_i > 0\}, \end{aligned}$$

where we consider  $|\Delta^n|$  as embedded in  $\mathbb{R}^{n+1} \cong \mathbb{R}^{\mathcal{J}}$ . If we consider  $\mathbf{sStrat}_P$  as the category of set-valued presheaves on  $\Delta_P$ , then by the logic of a nerve and realization functor,  $\text{Sing}_s \mathcal{X}$  is hence given the stratified simplicial set

$$\text{Sing}_s \mathcal{X}(\mathcal{J}) = \mathbf{Strat}_P(|\Delta^{\mathcal{J}}|_s, \mathcal{X})$$

with the obvious structure morphisms. The adjunction  $|-|_s \dashv \text{Sing}_s$  is simplicial.

**Recollection 2.3.4.** [DW22, Recl. 2.23] The two functors  $|-|_s$  and  $\text{Sing}_s$  are compatible with functoriality in the poset, in the sense that for any morphism  $f: P \rightarrow Q$  there are natural isomorphisms

$$\begin{aligned} |-|_s f! &\cong f! |-|_s \\ \text{Sing}_s f^* &\cong f^* \text{Sing}_s. \end{aligned}$$

It follows, that the adjunctions

$$|-|_s: \mathbf{sStrat}_P \rightleftarrows \mathbf{Strat}_P: \text{Sing}_s,$$

extend to a global adjunction

$$|-|_s: \mathbf{sStrat} \rightleftarrows \mathbf{Strat}: \text{Sing}_s,$$

denoted the same, by a slight abuse of notation. Specifically, the maps of simplicial mapping spaces are given by applying the fixed poset versions of  $|-|_s \dashv \text{Sing}_s$  component-wise, under the identifications

$$\begin{aligned} \mathbf{Strat}(\mathcal{S}, \mathcal{T}) &\cong \bigsqcup_{f \in \mathbf{Pos}(P_{\mathcal{S}}, P_{\mathcal{T}})} \mathbf{Strat}_{P_{\mathcal{X}}}(\mathcal{S}, f^* \mathcal{T}) \cong \bigsqcup_{f \in \mathbf{Pos}(P_{\mathcal{S}}, P_{\mathcal{T}})} \mathbf{Strat}_{P_{\mathcal{T}}}(f! \mathcal{S}, \mathcal{T}); \\ \mathbf{sStrat}(\mathcal{X}, \mathcal{Y}) &\cong \bigsqcup_{f \in \mathbf{Pos}(P_{\mathcal{X}}, P_{\mathcal{Y}})} \mathbf{sStrat}_{P_{\mathcal{X}}}(\mathcal{X}, f^* \mathcal{Y}) \cong \bigsqcup_{f \in \mathbf{Pos}(P_{\mathcal{X}}, P_{\mathcal{Y}})} \mathbf{sStrat}_{P_{\mathcal{Y}}}(f! \mathcal{X}, \mathcal{Y}). \end{aligned}$$

### 3 Homotopy theories of stratified objects

Equipping  $\mathbf{Strat}$  with a simplicial structure and powers (even an enrichment over Kan complexes) allows us to study stratified spaces from a homotopy-theoretic perspective. It follows from [DK87, §2.5, 2.6] that the  $\infty$ -category presented by  $\mathbf{Strat}$  is equivalently given by localizing  $\mathbf{Strat}$  at stratified homotopy equivalences  $H_s^5$ . It is, however, a general paradigm in abstract homotopy

<sup>5</sup>We make now use of this statement, so we omit a proof, but this formally follows from the fact that the powering of  $\mathbf{Strat}$  under  $\mathbf{sSet}$  can be used to define an equivalence of relative categories between  $(\mathbf{Strat}, H_s)$  and the flattening of  $\mathbf{Strat}$ .

theory that it is often more fruitful to consider a class of weak equivalences larger than the homotopy equivalences with respect to the simplicial structure (for example, weak homotopy equivalences of topological spaces, instead of homotopy equivalences). We may encode this information in terms of a relative category given by equipping  $\mathbf{Strat}$  with the additional data of a class of weak equivalences  $W$ . The  $\infty$ -categorical localization  $\mathbf{Strat}[W^{-1}]$  then defines a homotopy theory of stratified spaces. In the presence of sufficient extra structure, this process of localizing a larger class of maps is often equivalent to restricting the simplicial category to a subclass of particularly nice (i.e., bifibrant) objects (formally, this is [GH05, Prop. 1.1.10] or, more classically, [DK80b, §7]). For example, if one is interested in studying topological manifolds, by Whitehead’s theorem (and the fact that topological manifolds have the homotopy type of CW complexes), it is often perfectly sufficient to do so in the setting of weak homotopy equivalences. Following this perspective from classical homotopy theory, the question of a good notion of weak equivalence of stratified spaces and hence after a convenient homotopy theory of (poset) stratified spaces, arose.

### 3.1 Homotopy theories of stratified topological spaces

The question after a convenient homotopy theory of stratified spaces was already investigated in [Qui88] by Quinn, who took the approach of restricting the class of objects, rather than increasing the class of weak equivalences. Quinn followed the insight that in geometric scenarios a stratified space  $\mathcal{W}$ , with finitely many strata and minimal stratum  $p$ , can be decomposed into a diagram

$$\mathcal{W}' \leftarrow \mathcal{L} \rightarrow \mathcal{W}_p$$

with  $\mathcal{X} \rightarrow \mathcal{W}_p$  a fiber bundle with fiber stratified over  $P_{>p}$ , and  $\mathcal{W}' \leftarrow \mathcal{L}$  a stratum preserving boundary inclusion over  $P_{>p}$ . If we inductively repeat this process with  $\mathcal{W}_{>p}$  and  $\mathcal{L}$ , we ultimately end up with a diagram of spaces indexed over the regular flags in  $\mathrm{sd}P$ , that is, over  $\mathrm{sd}P^{\mathrm{op}}$ . In a less geometric scenario, we may not have access to the geometric link  $\mathcal{L}$  and its further decomposition into smaller pieces. However, we may still consider a homotopy theoretic analogue.

**Recollection 3.1.1.** Let  $\mathcal{X} \in \mathbf{Strat}_P$ , and  $\mathcal{I} \in \mathrm{sd}P$ . The  $\mathcal{I}$ -th homotopy link of  $\mathcal{X}$ ,  $\mathcal{H}o\mathrm{Link}_{\mathcal{I}}(\mathcal{X}) \in \mathbf{Top}$ , is the topological space obtained by equipping  $\mathbf{Strat}_P(|\Delta^{\mathcal{I}}|_s, \mathcal{X})$  with the compact open topology (or the respective Kelleyfication thereof). If  $\mathcal{I} = \{p\}$ , then  $\mathcal{H}o\mathrm{Link}_{\mathcal{I}}(\mathcal{X}) = \mathcal{X}_p$  and if  $\mathcal{I} = [p < q]$ , then  $\mathcal{H}o\mathrm{Link}_{\mathcal{I}}(\mathcal{X})$  is the space of paths which start in  $\mathcal{X}_p$  and immediately exit into the  $q$ -stratum. This construction is functorial, both in  $\mathcal{I}$  and  $\mathcal{X}$ , inducing a right-adjoint, simplicial functor

$$\mathcal{H}o\mathrm{Link}: \mathbf{Strat}_P \rightarrow \mathbf{Fun}((\mathrm{sd}P)^{\mathrm{op}}, \mathbf{Top}).$$

It will often be more convenient to present  $\mathcal{H}o\mathrm{Link}_{\mathcal{I}}\mathcal{X}$  as a simplicial set. By abuse of notation, we will also denote by  $\mathcal{H}o\mathrm{Link}$  the composition

$$\mathcal{H}o\mathrm{Link}: \mathbf{Strat}_P \rightarrow \mathbf{Fun}((\mathrm{sd}P)^{\mathrm{op}}, \mathbf{Top}) \xrightarrow{\mathrm{Sing}_*} \mathbf{Fun}((\mathrm{sd}P)^{\mathrm{op}}, \mathbf{sSet}).$$

If  $\mathcal{W}$  is a Whitney stratified space with two strata, then the diagram  $\mathcal{H}o\mathrm{Link}_{\mathcal{I}}(\mathcal{W})$  is weakly equivalent to the geometric decomposition diagram illustrated above.

Quinn studied a class of (metrizable) stratified spaces, for which the natural evaluation maps  $\mathcal{H}o\mathrm{Link}_{p < q}\mathcal{X} \rightarrow \mathcal{X}_p$  are Hurewicz fibrations and which additionally fulfill a cofibrancy condition for inclusions of strata (see [Qui88] for details), so-called homotopically stratified spaces (also called homotopically stratified sets). In this type of framework he showed, for example, a stratified analogue of the h-cobordism theorem. In [Mil13], Miller proved a Whitehead-style theorem for homotopically stratified spaces: A stratum preserving map between the latter is a stratified

homotopy equivalence if and only if it induces homotopy equivalences on all pairwise homotopy links and strata. Inspired by this, both Douteau and Henriques (see [Dou21c; Hen]) independently suggested the following class of weak equivalences for stratified spaces.

**Recollection 3.1.2.** [Dou21c; DW22] We call a stratum preserving map  $\mathcal{X} \rightarrow \mathcal{Y} \in \mathbf{Strat}_P$  a **diagrammatic equivalence**, if and only if it induces weak homotopy equivalences on all generalized homotopy links  $\mathcal{H}oLink_{\mathcal{I}}$ ,  $\mathcal{I} \in sdP$ . We denote by  $\mathbf{Strat}_P^{\mathfrak{d}}$ , the relative category obtained by equipping  $\mathbf{Strat}_P$  with the class of diagrammatic equivalences. It follows from [Dou21c, Thm. 3] that the homotopy link functor

$$\mathcal{H}oLink: \mathbf{Strat}_P \rightarrow \mathbf{Fun}((sdP)^{op}, \mathbf{Top})$$

induces an equivalence of quasi-categories

$$\mathbf{Strat}_P^{\mathfrak{d}} \simeq \mathbf{Fun}((sdP)^{op}, \mathbf{Top})[\{\text{pointwise weak equivalences}\}^{-1}] \simeq \mathbf{Fun}((sdP)^{op}, \mathbf{Spaces}).$$

This result can be extremely useful insofar, as it allows one to study stratified spaces in terms of presheaves on a fairly simple category, which is a homotopy theoretical setting that is well understood. At first glance, if one takes the perspective of Miller’s Whitehead theorem for homotopically stratified sets, it may seem surprising that generalized homotopy links of flags  $\mathcal{I}$  of dimension greater equal to 2 are part of the data detecting weak equivalences. Roughly speaking, higher homotopy links cannot be ignored, as even though they are not necessary to detect stratified homotopy equivalences between two homotopically stratified spaces, they nevertheless provide obstructions for such maps to exist at all. Compare this to the situation of equivalences of categories: Whether a functor is an equivalence of categories may be detected in terms of objects and hom-sets, but whether a functor exists at all, requires us to consider higher-dimensional data, namely the composition laws in the categories involved. If one has a version of a stratified homotopy hypothesis, then this analogy can even be made rigorous.

There is, however, another way of homotopically approximating the framework of homotopically stratified spaces. It follows from [Lur17, A.5] or [Nan19, Thm. 8.1.2.6.] that the homotopy link diagrams arising from classical examples of stratified spaces usually have the following property:

**Recollection 3.1.3.** [Hai23; Waa24a] A diagram  $D \in \mathbf{Fun}((sdP)^{op}, \mathbf{Spaces})$  is called a **décollage**, if for every regular flag  $\mathcal{I} = [p_0 < \dots < p_n] \in sdP$ , the following canonical morphism

$$D(\mathcal{I}) \rightarrow D(p_0 < p_1) \times_{D(p_1)} \dots \times_{D(p_{n-1})} D(p_{n-1} < p_n).$$

is a weak homotopy equivalence (that is, an isomorphism in the  $\infty$ -category of spaces). If  $D$  is presented by a commutative diagram in topological spaces or simplicial sets, this condition is equivalent to the natural map

$$D(\mathcal{I}) \rightarrow D(p_0 < p_1) \times_{D(p_1)}^H \dots \times_{D(p_{n-1})}^H D(p_{n-1} < p_n).$$

into the iterative homotopy pullback being a weak homotopy equivalence. In particular, if the homotopy link  $\mathcal{H}oLink_{\mathcal{I}}(\mathcal{X})$  of a stratified space  $\mathcal{X}$  is a décollage, then, roughly speaking, exit paths in  $\mathcal{X}$  admit compositions which are unique up to higher coherence. The homotopy theory of such décollages can be constructed in terms of a left Bousfield localization of  $\mathbf{Diag}_P := \mathbf{Fun}((sdP)^{op}, \mathbf{sSet})$ , equipped with the injective model structure, by localizing at such weak equivalences, which are local with respect to décollages (see [Waa24a, Subsec. 2.2]). In other words,  $f: E \rightarrow E'$  is a weak equivalence in the model structure that presents décollages, if and only



if for every injectively fibrant diagram  $D \in \underline{\mathbf{Diag}}_P$  that has the décollage property the induced map of mapping spaces

$$\underline{\mathbf{Diag}}_P(E', D) \rightarrow \underline{\mathbf{Diag}}_P(E, D)$$

is a weak homotopy equivalence. We denote this left-Bousfield localization of the injective model structure on  $\underline{\mathbf{Diag}}_P$  by  $\underline{\mathbf{Diag}}_P^{\text{d  }}$ . We say that a stratum preserving map  $f: \mathcal{X} \rightarrow \mathcal{Y} \in \mathbf{Strat}_P$  is a **categorical equivalence**, if and only if the induced morphism  $\mathcal{H}o\text{Link}_{\mathcal{I}}(\mathcal{X}) \rightarrow \mathcal{H}o\text{Link}_{\mathcal{I}}(\mathcal{Y})$  is a weak equivalence in  $\text{Fun}((\text{sd}P)^{\text{op}}, \mathbf{sSet})^{\text{d  }}$ , and denote the corresponding relative category by  $\mathbf{Strat}_P^{\text{c}}$ . Categorical weak equivalences are precisely the weak equivalences of stratified spaces suggested by Peter Haine in [Hai23]. It follows by construction and [Dou21c, Thm. 3] that the induced functor of quasi-categories

$$\mathcal{H}o\text{Link}_{\mathcal{I}}: \mathbf{Strat}_P^{\text{c}} \rightarrow \mathbf{Diag}_P^{\text{d  }}$$

is an equivalence. That is, if we localize categorical equivalences in  $\mathbf{Strat}_P$ , we obtain the homotopy theory of décollages. [BGH18; Hai23, Thm. 1.1.7] states that  $\mathbf{Diag}_P^{\text{d  }}$  is in turn equivalent to the homotopy theory of quasi-categories with a conservative functor in  $P$ , so-called **abstract stratified homotopy types**. This already shows that  $\mathbf{Strat}_P^{\text{c}}$  fulfills a version of a stratified homotopy hypothesis ([Hai23]).

However, in [Hai23] it was not yet known whether the equivalence between abstract stratified homotopy types and  $\mathbf{Strat}_P^{\text{c}}$  could be constructed on the nose through the stratified singular simplicial set construction. In fact, it was not known whether any categorical equivalence even induced a categorical equivalence on stratified singular simplicial sets. We are going to prove that this is indeed the case (see Theorem 3.3.1). Before we do so, let us generalize from the setting of stratum preserving maps to flexible posets. There are two apparent ways to generalize to the case of a flexible poset. The first essentially amounts to first requiring an isomorphism on the poset level, and then a weak equivalence under the resulting of homotopy theories over fixed posets:

**Recollection 3.1.4.** Let  $f: \mathcal{X} \rightarrow \mathcal{Y} \in \mathbf{Strat}$ . We say  $f$  is a **poset preserving diagrammatic (categorical) equivalence** if the underlying map of posets  $P(f): P_{\mathcal{X}} \rightarrow P_{\mathcal{Y}}$  is an isomorphism and the induced stratum preserving map  $P(f)!: \mathcal{X} \rightarrow \mathcal{Y}$  is a diagrammatic (categorical) equivalence. We denote the resulting relative categories on  $\mathbf{Strat}$  by  $\mathbf{Strat}^{\text{d}, P}$  and  $\mathbf{Strat}^{P, \text{c}}$ . This is how weak equivalence in  $\mathbf{Strat}$  are defined respectively in [Dou21a; Hai23].

This definition of homotopy theories on  $\mathbf{Strat}$  has the advantage that by making use of the Grothendieck bifibration  $\mathbf{Strat} \rightarrow \mathbf{Pos}$ , most questions about the global homotopy theories may be reduced to questions of the fibers. From a conceptual point of view, however, it has the side effect that the resulting homotopy theory contains a lot of highly pathological stratified spaces. Namely, both  $\mathbf{Strat}^{\text{d}, P}$  and  $\mathbf{Strat}^{P, \text{c}}$  contain a fully faithful copy of  $\mathbf{Pos}$ , given by empty stratified spaces. If we are looking to obtain a theory which embeds fully faithfully into  $\mathbf{Cat}_{\infty}$ , i.e. fulfills a homotopy hypothesis closer to the classical one, then the stratification poset should generally not be a homotopy invariant, but merely a gadget which specifies the allowable paths in a stratified space. This can be achieved by following the other possible approach to generalize the definitions in Recollections 3.1.2 and 3.1.3. Namely, one can transfer weak equivalences along a functor defined on stratified maps:

**Construction 3.1.5.** *The extended homotopy link  $\hat{\mathcal{H}}o\text{Link}_n(\mathcal{X})$  of a stratified space  $\mathcal{X} \in \mathbf{Strat}$ , is the bisimplicial set given by*

$$n \mapsto \underline{\mathbf{Strat}}([\Delta^{[n]}]_s, \mathcal{X}),$$

with the obvious functoriality in  $n$ . Note that

$$\begin{aligned} \underline{\mathbf{Strat}}(|\Delta^{[n]}|_s, \mathcal{X}) &= \bigsqcup_{\mathcal{I} \in N(P)_n} \underline{\mathbf{Strat}}_{P_{\mathcal{X}}}(|\Delta^{\mathcal{I}}|_s, \mathcal{X}) \\ &= \bigsqcup_{\mathcal{I} \in \text{sd}P_{\mathcal{X}}} \mathcal{H}\text{olink}_{\mathcal{I}}(\mathcal{X}) \sqcup \bigsqcup_{\mathcal{J} \in N(P_{\mathcal{X}})_n, \text{degenerate}} \underline{\mathbf{Strat}}_{P_{\mathcal{X}}}(|\Delta^{\mathcal{J}}|_s, \mathcal{X}). \end{aligned}$$

In particular,  $\hat{\mathcal{H}}\text{olink}_n(\mathcal{X})$  contains the data of all homotopy links of  $\mathcal{X}$ . Furthermore, whenever  $\mathcal{J}$  is a flag that degenerates from a nondegenerate flag  $\mathcal{I}$ , then the degeneracy map induces weak equivalences  $\mathcal{H}\text{olink}_{\mathcal{I}}(\mathcal{X}) \rightarrow \underline{\mathbf{Strat}}_{P_{\mathcal{X}}}(|\Delta^{\mathcal{J}}|_s, \mathcal{X})$ . In this sense, if we remove homotopically redundant data, then  $\hat{\mathcal{H}}\text{olink}_n(\mathcal{X})$  stores exactly the data of all homotopy links of  $\mathcal{X}$ . At the same time, allowing for degenerate simplices in homotopy links makes  $\hat{\mathcal{H}}\text{olink}_n$  functorial in stratified maps, not only stratum preserving ones.

**Definition 3.1.6.** A stratified map  $f: \mathcal{X} \rightarrow \mathcal{Y}$  is called a **diagrammatic equivalence**, if it induces weak homotopy equivalences of simplicial sets on all extended homotopy links  $\hat{\mathcal{H}}\text{olink}_n(\mathcal{X}) \rightarrow \hat{\mathcal{H}}\text{olink}_n(\mathcal{Y})$ . We denote by  $\mathbf{Strat}^{\text{d}}$  the relative category given by  $\mathbf{Strat}$  together with diagrammatic equivalences.

Finally, let us consider the categorical analogue of  $\mathbf{Strat}^{\text{d}}$ .

**Recollection 3.1.7.** In [Nan19], Nand-Lal defined a stratified map  $f: \mathcal{X} \rightarrow \mathcal{Y}$  to be a **categorical equivalence**, if and only if the underlying simplicial map of

$$\text{Sing}_s(f): \text{Sing}_s(\mathcal{X}) \rightarrow \text{Sing}_s(\mathcal{Y})$$

is a categorical equivalence of simplicial sets. We denote by  $\mathbf{Strat}^{\text{c}}$  the relative category given by  $\mathbf{Strat}$  together with categorical equivalences.

As we have already illustrated in the introduction, defining a homotopy theory of stratified spaces in terms of an  $\infty$ -categorical localization, especially in terms of maps which are themselves obtained through a localization, comes with an immediate series of questions. To just name a few: Are the stratum preserving categorical equivalences detected by  $\text{Sing}_s$ ? Do we have any intrinsic description of the mapping spaces in the  $\infty$ -categorical localizations? More precisely, how do they relate to the classical mapping spaces of stratified maps given by the simplicial structure on  $\mathbf{Strat}$ ? In the same vein of questioning, if we restrict ourselves to classical examples of stratified space, do we obtain the same homotopy theory as is investigated in [Qui88]? And if this is the case, how come [Qui88] has no need for higher homotopy links in his definition of a good class of stratified spaces? The classical way of making localizations of 1-categories more tractable is, of course, the theory of model categories. In [Dou21c], the relative categories  $\mathbf{Strat}_p^{\text{d}}$  and  $\mathbf{Strat}^{\text{d}, \text{p}}$ , were extended to model structures. However, these model structures have only limited expressive power when it comes to investigating classical examples of stratified spaces, since these are almost never bifibrant (for example, no stratified cone on a manifold is ever bifibrant). As we have already illustrated in the introduction, the goal of this work is to construct (semi)model structures for the homotopy theories defined in this section that make as many of such classical examples bifibrant.

## 3.2 Combinatorial models for stratified homotopy theory

One general strategy of constructing model categories is to first construct a model structure in a framework where this can easily be done, i.e., a category of presheaves or a similarly convenient setting, and then to transfer the theory along some adjunction. In this section, we will discuss simplicial analogues to the homotopy theories defined in the last section. We have investigated these theories in detail in [Waa24a], connecting the theories of [Hai23; Dou21a] and constructing combinatorial models for  $\mathbf{Strat}^{\text{c}}$  and  $\mathbf{Strat}^{\text{d}}$ . Let us begin with the case of  $\mathbf{Strat}_p^{\text{d}}$ .

**Recollection 3.2.1.** [Dou21a; DW22] The **Douteau-Henriques model structure** on  $\underline{\mathbf{sStrat}}_P$ , defined first in [Dou21a], is the Cisinski model structure (see [Cis19, Thm. 2.4.19]) induced by the simplicial cylinder  $X \mapsto X \otimes \Delta^1$ , with the empty set of anodyne extensions. This defines a combinatorial, cofibrant, simplicial model structure on  $\underline{\mathbf{sStrat}}_P$ , which may be characterized as the minimal model structure (in the sense of the smallest possible class of weak equivalences) in which the cofibrations are the monomorphisms and all stratified simplicial homotopy equivalences are weak equivalences. We denote the resulting simplicial model category  $\underline{\mathbf{sStrat}}_P^{\mathfrak{d}}$ .  $\underline{\mathbf{sStrat}}_P^{\mathfrak{d}}$  is cofibrantly generated by the classes of stratified boundary inclusions and admissible horn inclusions. It follows purely abstractly that weak equivalences between stratified simplicial sets  $\mathcal{X}, \mathcal{Y}$  which have the horn filling property with respect to admissible horn inclusions are precisely such stratum preserving simplicial maps  $\mathcal{X} \rightarrow \mathcal{Y}$ , for which the induced simplicial map of the simplicial homotopy links

$$\mathrm{HoLink}_{\mathcal{I}}(\mathcal{X}) := \underline{\mathbf{sStrat}}_P(\Delta^{\mathcal{I}}, \mathcal{X}) \rightarrow \underline{\mathbf{sStrat}}_P(\Delta^{\mathcal{I}}, \mathcal{Y}) =: \mathrm{HoLink}_{\mathcal{I}}(\mathcal{Y})$$

is a weak homotopy equivalence, for all  $\mathcal{I} \in \mathrm{sd}P$ . More surprisingly, in [DW22] it was shown that this detection criterion holds for all  $\mathcal{X}, \mathcal{Y}$  and that no fibrancy assumptions are necessary. Furthermore, in [DW22] it was also shown that mapping  $\mathcal{X} \in \underline{\mathbf{sStrat}}$  to the simplicial presheaf on  $\mathrm{sd}P$  given by  $\mathcal{I} \mapsto \underline{\mathbf{Strat}}(\Delta^{\mathcal{I}}, \mathcal{X})$  induces a Quillen-equivalence

$$\mathbf{Diag}_P \stackrel{\cong}{\simeq} \underline{\mathbf{sStrat}}_P^{\mathfrak{d}}: \mathrm{HoLink}$$

where the left hand side denotes the category of simplicial presheaves,  $\mathbf{Fun}((\mathrm{sd}P)^{\mathrm{op}}, \mathbf{sSet})$  equipped with the injective (or projective) model structure. Even more, this Quillen-equivalence creates weak equivalences in both directions. In particular,  $\underline{\mathbf{sStrat}}_P^{\mathfrak{d}}$  presents the  $\infty$ -category of space-valued diagrams indexed over  $\mathrm{sd}P^{\mathrm{op}}$ .

**Recollection 3.2.2.** [Hai23] The **Joyal-Kan model structure** on  $\underline{\mathbf{sStrat}}_P$  is obtained by left Bousfield localizing  $\underline{\mathbf{sStrat}}_P^{\mathfrak{d}}$  at the class of stratified inner horn inclusions. Fibrant objects are precisely the stratified simplicial sets  $\mathcal{X}$ , for which the underlying simplicial set  $X$  is a quasi-category and  $s_{\mathcal{X}}: X \rightarrow P_{\mathcal{X}}$  is a conservative functor. It follows that  $\underline{\mathbf{sStrat}}_P^{\mathfrak{c}}$  presents the  $\infty$ -category of conservative functors from a quasi-category into  $P$ , also called **abstract stratified homotopy types** over  $P$ . We can explicitly present the equivalence between décollages and abstract stratified homotopy types of [Hai23] in terms of a Quillen equivalence. In [Waa24a, Subsec. 2.2], we have shown that the adjunction

$$\mathbf{Fun}((\mathrm{sd}P)^{\mathrm{op}}, \mathbf{sSet})^{\mathrm{dé}} \rightleftarrows \underline{\mathbf{sStrat}}_P^{\mathfrak{c}}: \mathrm{HoLink}$$

induces a Quillen equivalence between  $\underline{\mathbf{sStrat}}_P^{\mathfrak{c}}$  and the model structure for décollages, which creates weak equivalences in both directions. This lifts the equivalence between abstract stratified homotopy types and décollages already proven in [BGH18] to the level of model categories.

**Recollection 3.2.3.** [Waa24a] Both the Douteau-Henriques as well as the Joyal-Kan model structures admit a global analogue on  $\underline{\mathbf{sStrat}}$  (named accordingly) obtained by gluing the model structures together along the Grothendieck bifibration  $\underline{\mathbf{sStrat}} \rightarrow P$ , using [CM20, Thm 4.4]. We denote by  $\underline{\mathbf{sStrat}}^{\mathfrak{d}, P}$  and  $\underline{\mathbf{sStrat}}^{\mathfrak{c}, P}$  the simplicial model categories with underlying category  $\underline{\mathbf{sStrat}}$ , defined by applying [CM20, Thm 4.4] to the forgetful functor

$$P: \underline{\mathbf{sStrat}} \rightarrow \mathbf{Pos},$$

with the fiberwise model structures given by  $\underline{\mathbf{sStrat}}_P^{\mathfrak{d}}$  and  $\underline{\mathbf{sStrat}}_P^{\mathfrak{c}}$ , for  $P \in \mathbf{Pos}$ , respectively. In more detail, a stratified simplicial map  $f: \mathcal{X} \rightarrow \mathcal{Y}$  is a cofibration, if and only if the induced

map  $P(f): \mathcal{X} \rightarrow \mathcal{Y}$  is a monomorphism, and weak equivalences are precisely such maps for which  $P(f)$  is an isomorphism, and  $P(f): \mathcal{X} \rightarrow \mathcal{Y}$  is a diagrammatic (or, respectively, categorical) equivalence.  $\mathbf{sStrat}^{\mathfrak{d},\mathfrak{p}}$  and  $\mathbf{sStrat}^{\mathfrak{c},\mathfrak{p}}$  are cofibrant combinatorial model categories. A set of generating cofibrations for both model categories is given by the set of stratified boundary inclusions  $\{\partial\Delta^{[n]} \hookrightarrow \Delta^{[n]} \mid n \in \mathbb{N}\}$ , together with the two morphisms

$$\begin{array}{ccc} \emptyset & \longrightarrow & \emptyset \\ \downarrow & & \downarrow \\ \emptyset & \longrightarrow & [0], \end{array} \quad \begin{array}{ccc} \emptyset & \longrightarrow & \emptyset \\ \downarrow & & \downarrow \\ [0] \sqcup [0] & \hookrightarrow & [1]. \end{array} \quad (2)$$

Furthermore, in  $\mathbf{sStrat}^{\mathfrak{d},\mathfrak{p}}$  acyclic cofibrations are generated by the stratified admissible horn inclusions

$$\begin{array}{ccc} \Lambda_k^n & \longrightarrow & \Delta^n \\ & \searrow & \swarrow \\ & [m] & \end{array}, \quad (3)$$

for  $n, m \in \mathbb{N}$ .

Generally, the poset of a stratified simplicial set may contain a lot of redundant elements and relations that do not reflect in the simplicial set itself. For example,  $\mathbf{sStrat}_P^{\mathfrak{c}}$  does not present a subcategory of  $\mathbf{Cat}_\infty$ , but instead an  $\infty$ -category of categories together with a conservative functor into a poset, and contains a fully faithful copy of the category of posets, given by mapping a poset  $P$  to the functor  $\emptyset \rightarrow P$ . If we are looking to obtain a version of a stratified homotopy hypothesis, referring only to layered  $\infty$ -categories without a choice of additional data, a further (right) localization is necessary. This uses the fact that any layered  $\infty$ -category naturally comes with a functor to the poset of its isomorphism classes, with relations induced by morphisms between the latter.

**Recollection 3.2.4.** [Waa24a, Subsec. 3.2] Given a stratified simplicial set  $\mathcal{X} \in \mathbf{Strat}$ , its refined poset  $P_{\mathcal{X}^\tau}$  is the poset generated with elements the vertices of  $\mathcal{X}$ , and a relation  $x \leq y$ , if and only if there is a path of 1-simplices

$$x = x_0 \leftrightarrow x_1 \leftrightarrow x_2 \leftrightarrow \dots \leftrightarrow x_n = y$$

where only simplices that are contained within a stratum of  $\mathcal{X}$  are allowed to point in direction of  $x$ . In other words,  $P_{\mathcal{X}^\tau}$  is the poset whose elements are the path component of strata of  $\mathcal{X}$  and whose relations are generated by exit paths in  $\mathcal{X}$ . The stratification map of  $\mathcal{X}$ ,  $s_{\mathcal{X}}: X \rightarrow P_{\mathcal{X}}$ , factors through  $P_{\mathcal{X}^\tau}$ , inducing a new stratification of  $X$ . The resulting stratified simplicial set is called **the refinement of  $\mathcal{X}$**  and denoted  $\mathcal{X}^\tau$ , and comes with a natural stratified simplicial map  $\mathcal{X}^\tau \rightarrow \mathcal{X}$ . We denote by  $\mathbf{sStrat}^{\mathfrak{d}}$  ( $\mathbf{sStrat}^{\mathfrak{c}}$ ) the two (simplicial) right Bousfield localizations of respectively  $\mathbf{sStrat}^{\mathfrak{d},\mathfrak{p}}$  and  $\mathbf{sStrat}^{\mathfrak{c},\mathfrak{p}}$  obtained by localizing the refinement maps (the existence of these was proven in [Waa24a, Thm. 3.2.18]). They are respectively called the **diagrammatic and categorical** model structure on  $\mathbf{sStrat}$ . Both form a combinatorial model category, with cofibrations generated by the class of stratified boundary inclusions  $\{\partial\Delta^{[n]} \hookrightarrow \Delta^{[n]} \mid n \in \mathbb{N}\}$ , together with the boundary inclusion  $\partial\Delta^{[1]} \rightarrow \Delta^{[1]}$ , into the trivially stratified simplex. The fibrant objects of  $\mathbf{sStrat}^{\mathfrak{c}}$ , i.e., the refined abstract stratified homotopy types are precisely what [BGH18] call the **0-connected abstract stratified homotopy types** - such abstract stratified homotopy types which have 0-connected strata and pairwise homotopy links. Forgetting the underlying stratification poset induces an equivalence

$$\mathcal{AStrat}^\tau \simeq \mathcal{Lay}_\infty$$

between the the homotopy theory of refined abstract stratified homotopy types  $\mathcal{A}\mathbf{Strat}^{\mathfrak{r}}$  and the homotopy theory of layered  $\infty$ -categories  $\mathcal{L}\mathbf{ay}_{\infty}$  (see [BGH18] and [Waa24a, Prop. 3.1.8 and Prop. 3.3.6]).

### 3.3 Equivalences of homotopy theories of stratified objects

Now that we have introduced both topological as well as simplicial homotopy theories of stratified spaces, the obvious question concerning the relation between the two arises. More precisely, what are the homotopy-theoretic properties of the adjunctions  $|-|_s \dashv \mathbf{Sing}_s$ ? We prove that each homotopy theory of stratified topological spaces is, in fact, equivalent to its simplicial counterpart, with the equivalence induced through the adjunctions  $|-|_s \dashv \mathbf{Sing}_s$ .

**Theorem 3.3.1** (Partially already found in [DW22; Hai23]). *Let  $\mathbf{R}$  be any of the relative categories of topological stratified spaces of Section 3.1 and let  $\mathbf{sR}$  be its simplicial counterpart from Section 3.2. Then, the adjunction  $|-|_s \dashv \mathbf{Sing}_s$  preserves all weak equivalences in both directions (i.e. induces functors of relative categories), and has unit and counit a weak equivalence (in other words, it is a strict homotopy equivalence of relative categories in the sense of [BK12b]). In particular, the adjunctions induce equivalences of quasi-categories*

$$\begin{array}{ll} \mathbf{sSet}_P^{\mathfrak{d}} \simeq \mathbf{Strat}_P^{\mathfrak{d}}; & \mathbf{sSet}_P \simeq \mathbf{Strat}_P^{\mathfrak{c}}; \\ \mathbf{sStrat}^{\mathfrak{d},\mathfrak{p}} \simeq \mathbf{Strat}^{\mathfrak{d},\mathfrak{p}}; & \mathbf{sStrat}^{\mathfrak{c},\mathfrak{p}} \simeq \mathbf{Strat}^{\mathfrak{c},\mathfrak{p}}; \\ \mathbf{sStrat}^{\mathfrak{d}} \simeq \mathbf{Strat}^{\mathfrak{d}}; & \mathbf{sStrat}^{\mathfrak{c}} \simeq \mathbf{Strat}^{\mathfrak{c}}. \end{array}$$

Before we give a proof, let us remark on the history of these results.

**Remark 3.3.2.** The existence of an equivalence of quasi-categories  $\mathbf{sStrat}^{\mathfrak{d},\mathfrak{p}} \simeq \mathbf{Strat}^{\mathfrak{d},\mathfrak{p}}$  was first shown in [Dou21b], in terms of a Quillen equivalence. [Hai23] used the equivalence between abstract stratified homotopy types  $\mathbf{Strat}_P^{\mathfrak{d}} \simeq \mathbf{Diag}_P^{\mathfrak{d}\acute{e}}$  and décollages proven in [BGH18], as well as [Dou21c, Thm. 3], to subsume that  $\mathbf{Strat}_P^{\mathfrak{d}}$  was equivalent to a left localization of  $\mathbf{Strat}_P^{\mathfrak{d}}$ . If one carefully follows the argument in [Hai23], this localization turns out to be precisely  $\mathbf{Strat}_P^{\mathfrak{c}}$ . However, the resulting equivalence  $\mathbf{Strat}_P^{\mathfrak{c}} \simeq \mathbf{sSet}_P^{\mathfrak{c}}$  is not induced by the adjunction  $|-|_s \dashv \mathbf{Sing}_s$ , but rather as a composition of equivalences  $\mathbf{Strat}_P^{\mathfrak{c}} \xrightarrow{\mathcal{H}\mathbf{oLink}} \mathbf{Diag}_P^{\mathfrak{d}\acute{e}} \xrightarrow{\simeq} \mathbf{sSet}_P^{\mathfrak{c}}$ . At this point, it was not yet known that  $\mathbf{Sing}_s: \mathbf{Strat}_P^{\mathfrak{c}} \rightarrow \mathbf{sStrat}_P^{\mathfrak{c}}$  maps categorical equivalences to categorical equivalences. The strict homotopy equivalence of relative categories version for the cases  $\mathbf{Strat}^{\mathfrak{d},\mathfrak{p}}$  and  $\mathbf{Strat}_P^{\mathfrak{d}}$  was first shown in [DW22, Thm. 1.3]. As we will see, all other cases can be derived from this one quite readily.

Let us first show that the nomenclature for categorical equivalences in  $\mathbf{Strat}_P$  is justified:

**Lemma 3.3.3.** *Let  $f: \mathcal{X} \rightarrow \mathcal{Y} \in \mathbf{Strat}_P$  be a stratum preserving map. Then the following are equivalent:*

1.  $f$  is a categorical equivalence;
2.  $\mathbf{Sing}_s(f)$  is an equivalence in the Joyal-Kan model structure;
3. The underlying simplicial map of  $\mathbf{Sing}_s(f)$  is a categorical equivalence.

*Proof.* There is a natural isomorphism of functors of 1-categories  $\mathcal{H}\mathbf{oLink}_{\mathcal{I}} \circ \mathbf{Sing}_s \cong \mathcal{H}\mathbf{oLink}_{\mathcal{I}}$ . By definition,  $f$  is a categorical equivalence if and only if  $\mathcal{H}\mathbf{oLink}_{\mathcal{I}}(f)$  induces an equivalence in the model structure for décollages. Hence, to prove that equivalence of the first two conditions holds, it suffices to see that  $\mathcal{H}\mathbf{oLink}_{\mathcal{I}}$  creates weak equivalences, which is part of the statement of [Waa24a, Thm. 2.2.15]. The final equivalence is a consequence of [Hai23, Thm. 0.2.2].  $\square$

As an immediate corollary, we obtain:

**Corollary 3.3.4.** *Let  $f: \mathcal{X} \rightarrow \mathcal{Y} \in \mathbf{Strat}$  be a stratified map. Then the following are equivalent:*

1.  $f$  is a poset preserving categorical equivalence;
2.  $\text{Sing}_s(f)$  is an equivalence in the poset preserving Joyal-Kan model structure;
3.  $f$  is a categorical equivalence and induces an isomorphism on posets.

Furthermore, the following equivalence for diagrammatic equivalences holds:

**Lemma 3.3.5.** *A stratified map  $f: \mathcal{X} \rightarrow \mathcal{Y}$  in  $\mathbf{Strat}$  is a poset preserving diagrammatic equivalence, if and only if  $\text{Sing}_s(f)$  is a weak equivalence in  $\mathbf{sStrat}^{\text{d.p}}$ .*

*Proof.* This follows from [Waa24a, Prop. 3.4.4] together with the natural isomorphism of  $\hat{\mathcal{H}}\text{olink}_n$  with the composition of its simplicial counterpart with  $\text{Sing}_s$ .  $\square$

Furthermore, we are going to make use of the following lemma:

**Lemma 3.3.6.** *Given two categories with weak equivalences  $(C, W_C)$ ,  $(D, W_D)$  suppose that a pair of functors*

$$L: C \rightleftarrows D: R,$$

*together with natural transformations  $1_C \rightarrow RL$  and  $\eta: LR \rightarrow 1_D$  define a strict homotopy equivalence of relative categories. Let  $W'_C$  be any wide subcategory such that  $W_C \subset W'_C$ , and such that  $(C, W'_C)$  is again a category with weak equivalences. Denote  $W'_D = R^{-1}(W'_C)$ . Then the quadruple  $(L, R, \varepsilon, \eta)$  also defines a homotopy equivalence between  $(C, W'_C)$  and  $(D, W'_D)$ .*

*Proof.* Clearly,  $W_D \subset W'_D$  and  $R(W'_D) \subset W'_C$ . Really, the only thing one needs to verify is that  $L$  remains a functor of relative categories. This is an immediate consequence of applying the two-out-of-three property in a category with weak equivalences.  $\square$

*Proof of Theorem 3.3.1.* Note that all model structures on the simplicial side are obtained by localizing the diagrammatic or the poset preserving diagrammatic model categories (see [Waa24a, Prop. 2.1.3 and 3.2.25]). It follows from Lemma 3.3.6 together with Lemmas 3.3.3 and 3.3.5 and Corollary 3.3.4 that it suffices to show the cases of  $\mathbf{Strat}_P^{\text{d.p}}$  and  $\mathbf{Strat}^{\text{d.p}}$ . These were proven in [DW22, Thm. 1.3, Rem. 5.4].  $\square$


## 4 (Semi)model categories of stratified spaces

The difficulty with the results in Theorem 3.3.1 is their interpretability. Without some additional structure on the side of  $\mathbf{Strat}$ , the only real information that we have is that the homotopy theories we defined are equivalent to the combinatorial world we transferred them from. Without some additional structure, we are stuck with the difficulty of interpreting what these results actually say about classical examples of stratified spaces that is, we need to determine how the localizations of Theorem 3.3.1 relate to stratified maps, stratified homotopies or more generally the stratified simplicial mapping spaces of  $\mathbf{Strat}$ . The usual approach to relating a localization to a simplicial category is the language of simplicial model categories. In [Dou21c], such simplicial model structures were suggested for  $\mathbf{Strat}^{\text{d.p}}$  and  $\mathbf{Strat}_P^{\text{d.p}}$ . The difficulty with these model structures, however, is that classically relevant examples such as (for example) Whitney stratified spaces are not bifibrant in the latter (in fact, no stratified cone on a manifold is cofibrant). Hence, the expressive power of these model structures when it comes to investigating mapping spaces for such spaces is limited. In [Lur17, A.5] or [Nan19, Thm. 8.1.2.6.], it was shown that for essentially all

stratified spaces of classical relevance  $\mathcal{X}$  the stratified singular simplicial set  $\text{Sing}_s(\mathcal{X})$  is fibrant in  $\mathbf{sStrat}^c$ , and hence in any of the model structures for stratified simplicial sets. Furthermore, at least Whitney stratified spaces or PL-pseudo manifolds admit triangulations compatible with their stratifications ([Gor78]), and are thus stratum preserving homeomorphic to the realization of a stratified simplicial set. This suggests the approach of transferring the simplicial model structures on stratified simplicial sets to the topological setting. Constructing such a model structure in the case of  $\mathbf{Strat}^c$  was one of the stated goals and conjectures of [Nan19], which was ultimately left as a conjecture. In fact, it turns out that the model structures in the combinatorial setting do not transfer to the topological framework, which was first shown in [DW22, A.5]. We repeat the result here, in slightly different form and with a more detailed proof, as it illustrates well the difficulties which may arise in the topological but not in the combinatorial world.

**Proposition 4.0.1.** *[DW22, Prop. A.1, Rem. A.4, Rem. A.5] There does not exist a model structure on  $\mathbf{Strat}$ , for which all poset preserving diagrammatic equivalences are weak equivalences, and the inclusion  $|\Lambda_1^{[3]}| \hookrightarrow \Delta^{[3]}|_s$  is a cofibration. The analogous result for  $\mathbf{Strat}_P$ , for  $P$  non-discrete, also holds.*

*Proof.* Consider stratum preserving map over  $[p < q]$  illustrated in the following picture:



The picture shows two stratifications over  $[p < q]$  of the realization of simplicial complexes (ignoring the white path from  $a$  to  $b$  for now), embedded in  $\mathbb{R}^2$ . Denote, from right to left, the corresponding stratified spaces by  $\mathcal{X}$  and  $\mathcal{Y}$ . The  $p$ -stratum is marked in red. In  $\mathcal{X}$  the  $p$ -stratum consists of a vertical line and two sequences, given in barycentric coordinates with respect to  $a, b, b'$  and  $a, c, c'$  by  $(1 - \frac{1}{2^n}, \frac{1}{2^{n+1}}, \frac{1}{2^{n+1}})$ . In  $\mathcal{Y}$ , the vertical line is replaced by a single point. Denote these sequences, respectively, by  $b_n$  and  $c_n$ . The map  $r$  is given by convexly extending the map which keeps all vertices besides  $a'$  the same and maps the latter to  $a$ . This map is not a categorical equivalence. Indeed, consider the induced map of simplicial sets

$$\text{Sing}_s(\mathcal{X}) \rightarrow \text{Sing}_s(\mathcal{Y})$$

and the induced map of hom-sets

$$\text{hoSing}_s(\mathcal{X})(a, b) \xrightarrow{r_*} \text{hoSing}_s(\mathcal{Y})(a, b)$$

in the associated homotopy categories. First, note that  $\text{Sing}_s(\mathcal{Y})$  is, in fact, a quasi-category. This can be seen, for example, from the fact that every non-trivially stratified horn  $|\Lambda^{\mathcal{J}}|_s \rightarrow \mathcal{X}$ ,  $\mathcal{J} = [p \leq p \leq \dots \leq q]$  is constant on the  $p$ -stratum. It follows that it suffices to show that  $\mathcal{Y}$  admits extensions with respect to  $|\Lambda^{\mathcal{J}}/\Delta^{\mathcal{J}_p}|_s \hookrightarrow |\Delta^{\mathcal{J}}/\Delta^{\mathcal{J}_p}|_s$ , which always admits a stratified retraction. Consequently, elements of  $\text{Sing}_s(\mathcal{Y})(a, b)$  are given by stratified homotopy classes (rel boundary) of exit paths from  $a$  to  $b$ . It turns out that  $r_*$  is not surjective: Consider a path  $\gamma$  from  $a$  to  $b$  that (described as starting from  $b$ ) ascends monotonously in height and passes to the left of  $b_1$ , to the right of  $c_2$ , to the left of  $b_3$  etc., as illustrated in the right picture. The class of  $\gamma$  does not lie in the image of  $r_*$ . Roughly speaking, this is due to the fact that no exit path in  $\mathcal{X}$  can oscillate

around the points of both sequences  $(b_n)$  and  $(c_n)$ . For a formal proof, see Appendix C. Next, let  $\mathcal{J} = [p \leq p < q \leq q]$  and  $\mathcal{J}' = [p < q \leq q]$  consider the following diagram of pushout squares

$$\begin{array}{ccccc}
|\Lambda_1^{[3]}|_s & \hookrightarrow & |\Delta^{[3]}|_s & \xrightarrow{|s^1|_s} & |\Delta^{[2]}|_s \\
\downarrow & & \downarrow & & \downarrow \\
|\Lambda_1^{\mathcal{J}}|_s & \hookrightarrow & |\Delta^{\mathcal{J}}|_s & \longrightarrow & |\Delta^{\mathcal{I}}|_s \\
\downarrow & & \downarrow & & \downarrow \\
\mathcal{X} & \longrightarrow & \mathcal{Z} & \xrightarrow{r'} & \mathcal{Y},
\end{array} \tag{4}$$

with the upper verticals induced by the identity on spaces and the flags  $\mathcal{J}$  and  $\mathcal{J}'$ , the lower left vertical given by affinely extending the map of vertices  $e_0 \mapsto a'$ ,  $e_1 \mapsto a$  and  $e_2 \mapsto b$  and  $e_3 \mapsto c$ , and with  $s^1$  the degeneracy map collapsing the edge  $\{1, 2\}$  to  $\{1\}$ . The bottom composition is precisely  $r$ , by construction of the latter. Now, the middle left horizontal is the realization of a diagrammatic equivalence and hence a diagrammatic equivalence. It follows that if a model structure as claimed in the statement of the proposition existed, then the left lower horizontal would be an acyclic cofibration. Now, suppose that the lower right vertical  $r'$  was also a categorical equivalence. Then it would follow that  $r$  is also a categorical equivalence, in contradiction to what we have just shown. In fact, the lower left vertical is even a stratum preserving homotopy equivalence, and hence a categorical equivalence. To see this, note that  $r'$  admits a section, induced by the pushout

$$\begin{array}{ccc}
|\Delta^{\mathcal{I}}|_s & \xleftarrow{|d^0|_s} & |\Delta^{\mathcal{J}}|_s \\
\downarrow & & \downarrow \\
\mathcal{Y} & \xrightarrow{s'} & \mathcal{Z}.
\end{array} \tag{5}$$

$|d_0|_s$  is even a stratum preserving strong deformation retract (that is, admits a deformation retraction which is also stratum preserving). It follows that  $s'$  is also a stratum preserving deformation retraction, and in particular a stratum preserving homotopy equivalence. Since  $r'$  is a retraction of  $s'$ , it follows that  $r'$  is a stratified homotopy equivalence. For the case of a fixed poset  $P$ , construct  $\mathcal{X}$  and  $\mathcal{Y}$  over any trivial relation  $p < q$  in  $P$ , and simply repeat the argument omitting the first line in Diagram (4).  $\square$

#### 4.1 A transfer lemma for cofibrantly generated semimodel structures

If one takes a precise look at the counterexample in Proposition 4.0.1, one will notice that it involves a pushout of an acyclic cofibration along a stratified map whose target is somewhat pathologically stratified. In particular, we cannot expect the target to be cofibrant. Hence, what one may nevertheless hope to obtain is a left semimodel structure. Semimodel structures provide a weaker version of model structures in which one can only expect pushouts of acyclic cofibrations to remain acyclic if all objects involved are cofibrant. Despite having slightly weaker axioms, for almost all intents and purposes semimodel categories (cofibrantly generated ones to be more precise) are just as useful as regular model categories. Essentially, every theorem about model categories admits an analogue in the world of semimodel categories, as long as one takes care that sources of morphisms often need to be assumed to be cofibrant (see, for example, [WB23] for an overview of some results). We will not go through the ordeal of reproducing every necessary result from the model category world in the semimodel category world here. Instead, when necessary, we will often reference a proof in the model-categorical setting and explain what needs to be adapted in the setting of semimodel categories. Since we only ever make use of the cofibrantly generated



scenario in this paper, and the latter tends to be significantly more well-behaved, we are going to use the following definition. Recall first the notion of weak factorization system (for example, from [Rie14, Ch. 11])

**Definition 4.1.1.** Let  $\mathbf{M}$  be a bicomplete category. A cofibrantly generated left semimodel structure on  $\mathbf{M}$  is the data of three classes  $(C, W, F)$ , called respectively cofibrations, weak equivalences, and fibrations, such that:

1.  $W$  contains all isomorphisms and is closed under two-out-of-three and retracts;
2.  $(C, W \cap F)$  is a (functorial) weak factorization system;
3.  $F$  consists exactly of those morphisms which have the right lifting property with respect to all morphisms in  $W \cap C$  with cofibrant source;
4. Every morphism with cofibrant source factors (functorially) into a morphism in  $W \cap C$ , followed by a fibration.

Furthermore, we assume that there exist sets of cofibrations  $I$  and of acyclic cofibrations with cofibrant source  $J$ , such that  $F$  is the class of morphisms which have the right lifting property with respect to  $I$ , and  $F \cap W$  is precisely the class of morphisms which have the right lifting property with respect to  $J$ . A (cofibrantly generated, left) semimodel category is the data of a complete and cocomplete category  $\mathbf{C}$ , together with a (cofibrantly generated) model structure on  $\mathbf{C}$ . Since we are only concerned with left semimodel categories in this work, we will just speak of semimodel categories, when we mean left semimodel category.

It is a routine exercise to show that since we assumed cofibrant generation, our definition is equivalent to [WB23, Def. 2.1]. Note that this definition is slightly stronger than the one used, for example, in [Fre09, p. 12.1.1]. We use the following definition of simpliciality:

**Definition 4.1.2.** Let  $\mathbf{M}$  be a bi-complete simplicial category, admitting powers and tensors, with the underlying 1-category equipped with a semimodel structure. We say that  $\mathbf{M}$  is a **simplicial semimodel category**, if for every cofibration  $i: A \hookrightarrow B$  in the Kan-Quillen model structure on simplicial sets (that is, for every monomorphism), and for every fibration  $f: X \rightarrow Y$  in  $\mathbf{M}$ , the induced morphism

$$Y^B \rightarrow X^B \times_{X^A} Y^A$$

is a fibration, and furthermore a weak equivalence if  $i$  or  $f$  is a weak equivalence.

**Remark 4.1.3.** Note that this is a slightly different definition of simplicial semimodel category than the one in [WB23, Def. 2.3] or [GH05, Def. 1.1.8]. Our definition is a priori stronger than the ones in [WB23; GH05]. The latter only implies that when  $i$  or  $f$  is a cofibration, then  $Y^B \rightarrow X^B \times_{X^A} Y^A$  has the right lifting property with respect to all acyclic cofibrations with cofibrant source. However, since in our cases all relevant semimodel categories are generated by (acyclic) cofibrations with cofibrant source, for the semimodel categories we are concerned with all of these definitions agree.

**Definition 4.1.4.** Let  $\mathbf{D}$  be a bicomplete category. Consider an adjunction

$$L: \mathbf{C} \rightleftarrows \mathbf{D}: R$$

and assume that  $\mathbf{C}$  is a model category. We say that the **right transferred semimodel structure on  $\mathbf{D}$  along  $R$  exists**, if the following classes form a semi model structure on  $\mathbf{D}$ :

1. Fibrations (weak equivalences) in  $\mathbf{D}$  are precisely such  $f \in \mathbf{D}$ , for which  $R(f)$  is a fibration (weak equivalence).

2. Cofibrations in  $\mathbf{D}$  are those morphisms that have the left lifting property with respect to all acyclic fibrations (i.e. morphisms which are both a weak equivalence and a fibration).

We then call this structure the **right transferred semimodel structure** on  $\mathbf{D}$  (along  $R$ ).

Given a class of morphisms  $I$  in a bicomplete category  $\mathbf{C}$ , we denote by  $\text{cell}(I)$  the class of relative cell complexes with respect to  $I$  (see [Hir09, Def. 10.5.8]). Similarly to the scenario of regular model categories (see, for example, [nLa23] for a proof), one obtains:

**Lemma 4.1.5.** *In the situation of Definition 4.1.2, suppose that  $\mathbf{C}$  is cofibrantly generated by cofibrations  $I$  and acyclic cofibrations with cofibrant source  $J$ . Suppose furthermore that:*

1. *Every morphism in  $\mathbf{D}$  factors into an element of  $\text{cell}L(I)$ , followed by an acyclic fibration.*
2. *Every morphism in  $\mathbf{D}$  with cofibrant source factors into an element of  $\text{cell}L(J)$ , followed by a fibration.*
3. *Every element of  $\text{cell}L(J)$  with cofibrant source is a weak equivalence.*

*Then the right transferred model structure on  $\mathbf{D}$  exists and is cofibrantly generated by  $L(I)$  and  $L(J)$ .*

Let us now consider the following transfer lemma for semimodel structures.

**Lemma 4.1.6.** *Let  $\mathbf{D}$  be a complete and cocomplete category. Consider an adjunction*

$$L: \mathbf{C} \rightleftarrows \mathbf{D}: R$$

*and assume that  $\mathbf{C}$  is a cofibrantly generated model category, with cofibrations generated by a set  $I$  and acyclic cofibrations generated by a set of morphisms with cofibrant source  $J$ , arrows in which have  $\kappa$ -small source, for some cardinal  $\kappa$ . Assume furthermore the following:*

- (i)  *$R$  preserves transfinite compositions of morphisms in  $\text{cell}L(I)$  and  $\text{cell}L(J)$  respectively;*
- (ii) *Weak equivalences in  $\mathbf{C}$  are stable under transfinite composition;*
- (iii) *For any a pushout diagram in  $\mathbf{D}$*

$$\begin{array}{ccc} L(A) & \xrightarrow{L(j)} & L(B) \\ \downarrow & & \downarrow \\ X & \xrightarrow{j'} & Y \end{array} \tag{6}$$

*with  $j \in J$  and  $X \in \text{cell}(L(I))$  an absolute cell complex, the map  $j'$  is a weak equivalence.*

*Then the right transferred semi model structure on  $\mathbf{D}$  exists, and it is furthermore cofibrantly generated by  $L(I)$  and  $L(J)$ . Even more, if  $L \dashv R$  is an adjunction of simplicial categories, with  $\mathbf{D}$  admitting a simplicial tensoring and powering, and the model structure on  $\mathbf{C}$  is simplicial, then so is the induced semimodel structure on  $\mathbf{D}$ .*

*Proof.* It follows from Assumption (i) and the assumption on small sources that the classes  $L(I)$  and  $L(J)$  admit the small object argument. In fact, it follows under the adjunction  $L \dashv R$  that the sources of  $L(I)$  and  $L(J)$  are  $\kappa$  small, with respect to  $\text{cell}L(I)$  and  $\text{cell}L(J)$ . Consequently, the only remaining requirement of Lemma 4.1.5 is that every element  $j' \in \text{cell}L(J)$  with cofibrant

source is a weak equivalence. It follows by Assumptions (i) and (ii) that we only need to consider the case of a pushout diagram

$$\begin{array}{ccc} L(A) & \xrightarrow{L(j)} & L(B) \\ \downarrow & & \downarrow \\ X & \xrightarrow{j'} & Y \end{array} \quad (7)$$

with  $j \in J$  and  $X$  cofibrant. Furthermore, since  $X$  is the retract of an absolute  $L(I)$  cell complex, and weak equivalences are stable under retracts, we may reduce to the case where  $X$  is an absolute cell complex. This case is precisely the content of Assumption (iii). Simplicity follows immediately by using that

$$R(Y^A \rightarrow X^B \times_{X^A} Y^A) \cong (R(Y)^A \rightarrow R(X)^B \times_{R(X)^A} R(Y)^A),$$

and the explicit definition of (acyclic) fibrations in  $\mathbf{D}$  in terms of  $R$ .  $\square$

## 4.2 Semimodel structures on Strat

To transfer the model structures from the combinatorial world to Strat $_P$  and Strat, we now need to verify the assumptions of Lemma 4.1.6. Most of these will turn out to be fairly standard abstract arguments. The difficult part is showing Assumption (iii). This requires an in-depth study of the behavior of homotopy links of absolute cell complexes, with respect to realizations of stratified boundary inclusions, which was performed in [Waa24b].

**Definition 4.2.1.** Denote  $I := \{|\partial\Delta^{\mathcal{J}} \hookrightarrow \Delta^{\mathcal{J}}|_s \mid \mathcal{J} \in \Delta_P\} \subset \mathbf{Strat}_P$ . Absolute cell complexes with respect to  $I$  are called **cellularly stratified spaces**. If  $\mathcal{A}$  and  $\mathcal{X}$  are cellularly stratified spaces, then we call a stratum preserving map  $f: \mathcal{A} \hookrightarrow \mathcal{X}$  which is a relative  $I$ -cell complex an **inclusion of cellularly stratified spaces**.

The following result is the decisive argument to apply Lemma 4.1.6 to transfer the model structure on combinatorially stratified objects to the stratified framework. It is a corollary of the main result of [Waa24b, Thm. A], which is the main result of [Waa24b].

**Lemma 4.2.2.** *Consider a pushout diagram in Strat,*

$$\begin{array}{ccc} |\mathcal{A}|_s & \xrightarrow{|i|_s} & |\mathcal{B}|_s \\ \downarrow & & \downarrow \\ \mathcal{X} & \xrightarrow{i'} & \mathcal{Y} \end{array} \quad (8)$$

*with  $\mathcal{X}$  cellularly stratified. Suppose that  $i$  is an acyclic cofibration in  $\mathbf{sStrat}^{\mathfrak{d},\mathfrak{p}}$ . Then,  $i'$  is a weak equivalence in  $\mathbf{Strat}^{\mathfrak{d},\mathfrak{p}}$ , i.e., an isomorphism in  $\mathbf{Strat}^{\mathfrak{d},\mathfrak{p}}$ . The analogous result holds for any of the model categories of stratified simplicial sets and their corresponding topological  $\infty$ -category in Notation 4.2.5.*

Before we give a proof, note that Lemma 4.2.2 can be taken as a statement about certain pushout diagrams of stratified spaces being homotopy pushout, i.e. pushout in the associated  $\infty$ -category obtained by inverting weak equivalences. This already shows that the existence of semimodel structures is closely related with Question Q(5).

*Proof.* Let us begin with the case over a fixed poset  $P$ , and all morphisms on posets given by the identity. We claim that the diagram in the statement of the proposition is a pushout diagram in

the quasi-categories  $\mathbf{Strat}_P^{\circ}$  and  $\mathbf{Strat}_P^{\mathfrak{c}}$ . We already know that  $\mathbf{Strat}_P^{\mathfrak{c}}$  is a left-localization of  $\mathbf{Strat}_P^{\circ}$  (by Theorem 3.3.1 and the fact that  $\mathfrak{s}\mathbf{Strat}_P^{\mathfrak{c}}$  is a left Bousfield localization of  $\mathfrak{s}\mathbf{Strat}_P^{\circ}$ , together with [Lur09, Prop. 5.2.4.6]) and in particular that the natural functor  $\mathbf{Strat}_P^{\circ} \rightarrow \mathbf{Strat}_P^{\mathfrak{c}}$  is left adjoint. Therefore, by [Lur09, p. 5.2.3.5], which states that left-adjoint functors preserve colimits, it suffices to show the case of  $\mathbf{Strat}_P^{\circ}$ . Recall from Recollection 3.2.1 that taking homotopy links induces equivalences between  $\mathbf{Strat}_P^{\circ}$  and  $\mathbf{Fun}((\mathrm{sd}P)^{\mathrm{op}}, \mathbf{Spaces})$ . Thus, it suffices to show that the diagram becomes pushout after applying homotopy links. Now, by [Lur09, Cor. 5.1.2.3] a diagram in a functor quasi-category is pushout if and only if it is so after evaluating at every point. Thus, it suffices to show that the diagram in  $\mathbf{Spaces}$

$$\begin{array}{ccc} \mathcal{H}\mathrm{oLink}_{\mathcal{I}}|\mathcal{A}|_s & \longrightarrow & \mathcal{H}\mathrm{oLink}_{\mathcal{I}}|\mathcal{B}|_s \\ \downarrow & & \downarrow \\ \mathcal{H}\mathrm{oLink}_{\mathcal{I}}\mathcal{X} & \longrightarrow & \mathcal{H}\mathrm{oLink}_{\mathcal{I}}\mathcal{Y} \end{array} \quad (9)$$

is pushout. Now, the quasi-category  $\mathbf{Spaces}$  is given by the homotopy coherent nerve of  $\mathbf{Top}^{\circ}$ . [Waa24b, Thm. B] states that the diagram in  $\mathbf{Top}$  corresponding to Diagram (9) is homotopy pushout. Finally, by [Lur09, Thm. 4.2.4.1.], this is equivalent to Diagram (9) being a pushout diagram in  $\mathbf{Spaces}$ .

We have established that the diagram in the statement of the lemma is pushout in  $\mathbf{Strat}_P^{\circ}$  and  $\mathbf{Strat}_P^{\mathfrak{c}}$ . Consequently, by Theorem 3.3.1, so is its image in  $\mathfrak{s}\mathbf{Set}_P^{\circ}$  ( $\mathfrak{s}\mathbf{Set}_P^{\mathfrak{c}}$ ). By assumption,  $i$  defines an isomorphism in  $\mathfrak{s}\mathbf{Set}_P^{\circ}$  ( $\mathfrak{s}\mathbf{Set}_P^{\mathfrak{c}}$ ). It follows, again by Theorem 3.3.1, that  $\mathrm{Sing}_s(|i|_s)$  also defines an isomorphism in  $\mathfrak{s}\mathbf{Set}_P^{\circ}$  ( $\mathfrak{s}\mathbf{Set}_P^{\mathfrak{c}}$ ). Consequently, since pushouts preserve isomorphisms, it follows that  $\mathrm{Sing}_s(i')$  is an isomorphism in the quasi-category  $\mathfrak{s}\mathbf{Set}_P^{\circ}$  ( $\mathfrak{s}\mathbf{Set}_P^{\mathfrak{c}}$ ), and thus a weak equivalence in  $\mathfrak{s}\mathbf{Strat}_P^{\circ}$  ( $\mathfrak{s}\mathbf{Strat}_P^{\mathfrak{c}}$ ) (by [Hir09, Thm. 8.3.10.] and the fact that the homotopy category functor from quasi-categories to 1-categories commutes with localization). It follows that  $i$  is a weak equivalence, as was to be shown.

Next, let us cover the non-refined cases in  $\mathfrak{s}\mathbf{Strat}$ . Note that  $i$  induces an isomorphism on posets. By pushing forward with  $P_{\mathcal{A}} \rightarrow P_{\mathcal{X}}$ , and using the fact that this preserves realizations of acyclic cofibrations (by Proposition 4.2.8), we may assume that the whole diagram lies in  $\mathbf{Strat}_{P_{\mathcal{X}}}$ . Hence, this case follows from the previous cases. Finally, for the refined case, note that the refined model structures on  $\mathfrak{s}\mathbf{Strat}$  have the same acyclic cofibrations, as they are obtained via right Bousfield localization. Hence these cases were already covered in the cases of  $\mathfrak{s}\mathbf{Strat}^{\mathfrak{c};\mathfrak{p}}$  and  $\mathfrak{s}\mathbf{Strat}^{\circ;\mathfrak{p}}$ .  $\square$

We may now prove the following.

**Corollary 4.2.3.** *Let  $\underline{\mathfrak{cS}}$  be any of the simplicial model categories of stratified simplicial sets listed in Section 3.2. Let  $\underline{\mathfrak{S}}$  be correspondingly the simplicial category of topological stratified spaces  $\mathbf{Strat}_P$  or  $\mathbf{Strat}$ . Then, along the adjunction*

$$|-|_s: \underline{\mathfrak{cS}} \rightleftarrows \underline{\mathfrak{S}}: \mathrm{Sing}_s,$$

*the simplicial model structure on  $\underline{\mathfrak{cS}}$  right-transfers to a cofibrantly generated simplicial semimodel structure on  $\underline{\mathfrak{S}}$ .*

*Proof.* We verify the assumptions of Lemma 4.1.6. Note first that  $\underline{\mathfrak{cS}}$  is a cofibrantly generated simplicial model category (with acyclic generators with cofibrant source) as we have recalled in Section 3.2. Assumption (i) follows from the classical fact that any compactum in a relative cell complex only intersects finitely many cells (see, for example, [Hir09, Prop. 10.7.4]). Assumption (ii) was verified in [Waa24a, Prop. 2.1.6, 3.1.5 and 3.2.22]. Finally, Assumption (iii) follows from Lemma 4.2.2.  $\square$

**Remark 4.2.4.** We note that the analogous proof of Corollary 4.2.3 also applies to the category of surjectively stratified spaces, as investigated in [Nan19],  $\mathbf{Strat}^s$ , and one may replace  $\mathbf{sStrat}$  by  $\mathbf{sSet}$  with the Joyal-model structure (omitting the simpliciality statement). Indeed, Lemma 4.2.2 also applies to surjectively stratified spaces and realizations of Joyal acyclic cofibrations, using Proposition A.0.2 and that surjectively stratified spaces form a coreflective subcategory of  $\mathbf{Strat}$ . In particular, this also gives an affirmative answer to the semimodel structure conjectured in [Nan19, Subsec. 8.4.1]. As a consequence of Proposition A.0.2, the resulting semimodel category is Quillen equivalent to the one on all stratified spaces  $\mathbf{Strat}$  transferred from  $\mathbf{sStrat}^c$ . We also note that the alternative method suggested for the proof in [Nan19] cannot succeed, as [Nan19, Conjecture 2] is false (realizations of fibrant simplicial sets are generally not fibrant). We will provide a counterexample in future work.

**Notation 4.2.5.** We denote by  $\mathbf{Strat}_P^{\mathfrak{d}}$  ( $\mathbf{Strat}_P^c$ ) and  $\mathbf{Strat}^{\mathfrak{d},p}$  ( $\mathbf{Strat}^{\mathfrak{d}}$ ,  $\mathbf{Strat}^{c,p}$ ,  $\mathbf{Strat}^c$ ), respectively, the corresponding simplicial semimodel categories whose existence is guaranteed by Corollary 4.2.3. We, respectively, call the corresponding semi model structures the **diagrammatic (categorical, poset preserving diagrammatic, poset preserving categorical, diagrammatic, categorical)** model structures.

We use the remainder of the section to make Corollary 4.2.3 more explicit by restricting it to its special cases. From Recollections 3.2.1 and 3.2.2 and Corollary 4.2.3 we obtain:

**Theorem 4.2.6.** *Let  $P \in \mathbf{Pos}$ . The simplicial category  $\mathbf{Strat}_P$  admits the structures of cofibrantly generated simplicial semimodel categories,  $\mathbf{Strat}_P^{\mathfrak{d}}$  and  $\mathbf{Strat}_P^c$  - called the diagrammatic and categorical model structure - defined by the following classes:*

1. *Cofibrations are generated by the set of stratified boundary inclusions*

$$\{|\partial\Delta^{\mathcal{J}} \hookrightarrow \Delta^{\mathcal{J}}|_s \mid \mathcal{J} \in \Delta_P\}.$$

2. *Weak equivalences are the poset preserving diagrammatic equivalences or respectively the poset preserving categorical equivalences of  $P$ -stratified spaces.*
3. *Fibrations are the stratum preserving maps that have the right lifting property with respect to all acyclic cofibrations with cofibrant source. In  $\mathbf{Strat}^{\mathfrak{d},p}$  they are equivalently characterized by having the right lifting property with respect to all admissible stratified horn inclusions  $|\Lambda_k^{\mathcal{J}} \hookrightarrow \Delta^{\mathcal{J}}|_s$ ,  $\mathcal{J} \in \Delta_P$ .*

Next, let us consider the global version of  $\mathbf{Strat}_P$ , allowing for flexible posets. From Recollection 3.2.1 and Corollary 4.2.3 we obtain the following two results:

**Theorem 4.2.7.** *The simplicial category  $\mathbf{Strat}$  admits the structures of cofibrantly generated simplicial semimodel categories,  $\mathbf{Strat}^{\mathfrak{d},p}$  and  $\mathbf{Strat}^{c,p}$  - called the poset preserving diagrammatic and categorical model structure - with the following classes:*

1. *Cofibrations are generated by the set of stratified boundary inclusions*

$$\{|\partial\Delta^{[n]} \hookrightarrow \Delta^{[n]}|_s \mid n \in \mathbb{N}\},$$

*together with the maps of empty stratified spaces over  $[0] \sqcup [0] \hookrightarrow [0, 1]$  and  $\emptyset \hookrightarrow [0]$ .*

2. *Weak equivalences are the poset preserving diagrammatic equivalences or, respectively, the poset preserving categorical equivalences of stratified spaces.*

3. *Fibrations are the stratum preserving maps that have the right lifting property with respect to all acyclic cofibrations with cofibrant source. In  $\mathbf{Strat}^{\mathfrak{d},\mathfrak{p}}$  they are equivalently characterized by having the right lifting property with respect to all admissible stratified horn inclusions  $|\Delta_k^{\mathcal{J}} \hookrightarrow \Delta^{\mathcal{J}}|_s$ ,  $\mathcal{J} \in \Delta_{[m]}$ , for  $m \in \mathbb{N}$ .*

Similarly, to the case of stratified simplicial sets, the global diagrammatic and categorical model structures can essentially be understood entirely from the ones over a fixed poset and the functoriality of base changes along the poset. This is due to the fact that they may again be interpreted as being pieced together from the local pieces along a Cartesian bifibration:

**Proposition 4.2.8.** *The model structure on  $\mathbf{Strat}^{\mathfrak{d},\mathfrak{p}}$  ( $\mathbf{Strat}^{\mathfrak{c},\mathfrak{p}}$ ) is the unique semimodel structure on  $\mathbf{Strat}$  pieced together (in the sense of [CM20], using notation from there) from the diagrammatic (categorical) model structures on the fibers  $\mathbf{Strat}_P$  of the Grothendieck bifibration*

$$P: \mathbf{Strat} \rightarrow \mathbf{Pos}.$$

*In particular, a stratified map  $f: \mathcal{X} \rightarrow \mathcal{Y}$  is a fibration, if and only if the canonical maps  $f^{\triangleleft}: \mathcal{X} \rightarrow P(f)^* \mathcal{Y}$  is a fibration, and  $f$  is a cofibration, if and only if  $f_{\triangleright}: P(f)_! \mathcal{X} \rightarrow \mathcal{Y}$  is a cofibration.*

*Proof.* We cover the case of  $\mathbf{Strat}^{\mathfrak{d},\mathfrak{p}}$ , the case of  $\mathbf{Strat}^{\mathfrak{c},\mathfrak{p}}$  is completely analogous. Consider the forgetful functor  $P: \mathbf{Strat} \rightarrow \mathbf{Pos}$ , mapping a stratified space  $\mathcal{X}$  to its stratification poset  $P_{\mathcal{X}}$ . It is a Grothendieck bifibration ([CM20]), with the left action given by  $f \mapsto f_!$  and the right action given by  $g \mapsto g^*$ . If we equip  $\mathbf{Pos}$ , with the trivial model structure, with all morphisms bifibrations and weak equivalences given by isomorphisms, then  $P: \mathbf{Strat} \rightarrow \mathbf{Pos}$  tautologically forms a Quillen bifibration in the sense of [CM20] (replacing model categories by cofibrantly generated semimodel categories). We obtain two commutative diagrams (one for each horizontal) with diagonals Quillen bifibrations

$$\begin{array}{ccc} \mathbf{Strat}^{\mathfrak{d},\mathfrak{p}} & \xrightleftharpoons{\simeq} & \mathbf{sStrat}^{\mathfrak{d},\mathfrak{p}} \\ & \searrow & \swarrow \\ & \mathbf{Pos} & \end{array}, \quad (10)$$

where the right adjoint horizontal preserves the right action, and the left adjoint the left actions. The fibers of the right hand vertical are the model categories  $\mathbf{sStrat}_P^{\mathfrak{d}}$  (see the construction of the global model categories in [Waa24a, Subsec. 3.1]). Since the semimodel structure on  $\mathbf{Strat}^{\mathfrak{d},\mathfrak{p}}$  is transferred along the horizontal adjunction, it follows that its restriction to the fibers of the left vertical  $\mathbf{Strat}_P$ , are precisely the transfers of the model structure on  $\mathbf{sStrat}_P^{\mathfrak{d}}$  to  $\mathbf{Strat}_P$ , which is  $\mathbf{Strat}_P^{\mathfrak{d}}$ . Using the obvious cofibrantly generated semimodel categorical version of [CM20, Prop. 3.4, 3.5], which is proven identically, the characterization of cofibrations and fibrations follows.  $\square$

Finally, let us state the explicit results for the refined framework. We also are going to need the following construction.

**Construction 4.2.9.** *Consider the pushout diagram in  $\mathbf{Strat}$*

$$\begin{array}{ccc} |\partial\Delta^{[1]}|_s & \hookrightarrow & |\Delta^{[1]}|_s \\ \downarrow t & & \downarrow \\ |\Delta^{[1]}|_s & \hookrightarrow & S^1, \end{array} \quad (11)$$

*where the vertical  $t$  is given by the opposite inclusion of the boundary points of the stratified interval. The pushout is a trivially stratified  $S^1$ . In particular, the diagonal of this diagram - which is the inclusion of two points lying over separate strata into a trivially stratified  $S^1$  - is a cofibration in the refined model structures. The realization of the stratified boundary inclusion  $\partial\Delta^{[1]} \hookrightarrow \Delta^1$ , where  $\Delta^1$  is trivially stratified, is a retract of the diagonal of this diagram.*

The following are a consequence of [Waa24a, Thm. 3.2.18], which we recalled in Recollection 3.2.4, and Corollary 4.2.3. Note that, by Construction 4.2.9, the topological framework needs one less generator of cofibrations than the combinatorial framework.

**Theorem 4.2.10.** *The simplicial category  $\mathbf{Strat}$  admits the structures of cofibrantly generated simplicial semimodel categories,  $\mathbf{Strat}^{\mathfrak{d}}$  and  $\mathbf{Strat}^{\mathfrak{c}}$  - called the diagrammatic and categorical model structure - with the following classes:*

1. *Cofibrations are generated by the set of stratified boundary inclusions*

$$\{|\partial\Delta^{[n]} \hookrightarrow \Delta^{[n]}|_s \mid n \in \mathbb{N}\}.$$

2. *Weak equivalences are the diagrammatic equivalences or respectively the categorical equivalences of stratified spaces.*
3. *Fibrations are the stratum preserving maps that have the right lifting property with respect to all acyclic cofibrations with cofibrant source. In  $\mathbf{Strat}^{\mathfrak{d}}$ , they are equivalently characterized by having the right lifting property with respect to all admissible stratified horn inclusions  $|\Lambda_k^{\mathcal{J}} \hookrightarrow \Delta^{\mathcal{J}}|_s$ ,  $\mathcal{J} \in \Delta_{[m]}$ , for  $m \in \mathbb{N}$ .*

Finally, we may summarize the connections between the various semimodel categories of stratified objects following result, which follows by ([Waa24a, Prop. 2.1.3, 3.1.11, 3.2.23, 3.2.18, 3.2.25] and Proposition 5.3.7 and Theorems 3.3.1 and 4.2.10):

**Theorem 4.2.11.** *The various simplicial semimodel categories discussed in this section fit into a diagram*

$$\begin{array}{ccccc}
\mathbf{sStrat}_P^{\mathfrak{d}} & \xrightleftharpoons{\cong} & \mathbf{Strat}_P^{\mathfrak{d}} & & \\
\downarrow & \swarrow & \downarrow & \swarrow & \\
& \mathbf{sStrat}_P^{\mathfrak{c}} & \xrightleftharpoons{\cong} & \mathbf{Strat}_P^{\mathfrak{c}} & \\
\downarrow & \downarrow & \downarrow & \downarrow & \\
\mathbf{sStrat}^{\mathfrak{d},p} & \xrightleftharpoons{\cong} & \mathbf{Strat}^{\mathfrak{d},p} & & \\
\downarrow & \swarrow & \downarrow & \swarrow & \\
& \mathbf{sStrat}^{\mathfrak{c},p} & \xrightleftharpoons{\cong} & \mathbf{Strat}^{\mathfrak{c},p} & \\
\downarrow & \downarrow & \downarrow & \downarrow & \\
\mathbf{sStrat}^{\mathfrak{d}} & \xrightleftharpoons{\cong} & \mathbf{Strat}^{\mathfrak{d}} & & \\
\downarrow & \swarrow & \downarrow & \swarrow & \\
& \mathbf{sStrat}^{\mathfrak{c}} & \xrightleftharpoons{\cong} & \mathbf{Strat}^{\mathfrak{c}} & 
\end{array} \tag{12}$$

of simplicial functors, with all horizontals induced by the adjunction  $|-|_s \dashv \text{Sing}_s$ , and all remaining functors given by inclusions or the identity. This diagram has the following properties.

1. *The arrows pointing in two directions are simplicial Quillen adjunctions, where the upper (left) arrow is the left part of the adjunction.*
2. *The left adjunction part (and, respectively, the right adjunction part) together with the inclusions form a commutative diagram.*
3. *All horizontals are Quillen-equivalences, which create weak equivalences in both directions.*

4. The upper vertical maps are given by including the fibers under the Quillen bifibrations to **Pos**.
5. The diagonals pointing downward are given by left Bousfield localization at inner horn inclusions.
6. The lower verticals that point downward are given by right Bousfield localization at the refinement maps  $(-)^{\mathfrak{r}} \rightarrow 1$ .

### 4.3 Topological stratified mapping spaces

Give two layered  $\infty$ -categories  $X$  and  $Y$ , the  $\infty$ -category of functors  $Y^X$  is itself layered. Indeed, it follows from the fact that isomorphisms of functors are detected pointwise that it even suffices for  $Y$  to be layered. Together with Theorem 3.3.1, this already shows that at least the homotopy theory  $\mathbf{Strat}^{\mathfrak{f}}$  admits a notion of internal mapping space on the  $\infty$ -categorical level. Similarly, in the world of topological stratified spaces (more specifically homotopically stratified spaces) [Hug99] equipped the space of stratified maps with a natural decomposition (which generally may not be a stratification) and investigated the lifting properties of such mapping spaces.

If we are looking to bring these two notions of internal (i.e. stratified) mapping spaces - one on the  $\infty$ -categorical and one on the 1-categorical level together - the language of (semi)model categories provides the notion of a Cartesian closed (semi)model category (see, for example, [Rez10] for the case of model categories).

**Definition 4.3.1.** Let  $\mathbf{M}$  be a semimodel category the underlying 1-category of which is Cartesian closed. We say that  $\mathbf{M}$  is a **Cartesian closed semimodel category**, if for every cofibration  $i: A \rightarrow B$  and every fibration  $f: X \rightarrow Y$ , the induced morphism of exponential objects

$$X^B \rightarrow Y^B \times_{Y^A} X^A$$

is a fibration and furthermore is a weak equivalence, if additionally  $f$  is a weak equivalence, or  $i$  is a weak equivalence with cofibrant source.

In [Waa24a, Subsec. 3.5], we proved that all of the model structures for stratified homotopy theory defined on  $\mathbf{sStrat}$  (see Section 3.2) are, in fact, Cartesian closed. We may use the following lemma to transfer this result to the topological setting.

**Lemma 4.3.2.** *Let then  $\mathbf{M}$  be a Cartesian semimodel category and  $\mathbf{N}$  a Cartesian closed, bicomplete category. Suppose that there is an adjunction*

$$L: \mathbf{M} \rightleftarrows \mathbf{N}: R,$$

*such that  $L$  preserves finite products and the right transferred semimodel structure on  $\mathbf{N}$  exists. Then, together with this model structure,  $\mathbf{N}$  is a Cartesian closed semimodel category.*

*Proof.* Note first that if  $L$  preserves products, then there are canonical isomorphisms

$$R(X^{L(A)}) \cong R(X)^A$$

for  $A \in \mathbf{M}$  and  $X \in \mathbf{N}$ . Now, let us verify the hypothesis in the definition of a Cartesian closed semimodel category. Suppose that we are given a cofibration  $i: A \rightarrow B$  and a fibration  $f: X \rightarrow Y$  in the transferred semimodel structure on  $\mathbf{N}$ . In the context of Definition 4.3.1 in  $\mathbf{N}$ , let us first reduce to the case where  $A \rightarrow B$  is of the form  $L(A' \rightarrow B')$ , for  $A \rightarrow B$  a cofibration (acyclic cofibration



with cofibrant source). For the general case, we need to show that  $R(X^B \rightarrow Y^B \times_{Y^A} X^A)$  is a fibration in  $\mathbf{M}$ , or equivalently, that every lifting problem

$$\begin{array}{ccc} L(A') & \longrightarrow & X^B \\ \downarrow & \dashrightarrow & \downarrow \\ L(B') & \longrightarrow & Y^B \times_{Y^A} X^A \end{array} \quad (13)$$

with  $A' \rightarrow B'$  an (acyclic) cofibration (with cofibrant source) has a solution. Every such lifting problem is, in turn, equivalent to a lifting problem of the form

$$\begin{array}{ccc} A & \longrightarrow & X^{L(B')} \\ \downarrow & \dashrightarrow & \downarrow \\ B & \longrightarrow & Y^{L(B')} \times_{Y^{L(A')}} X^{L(A')}. \end{array} \quad (14)$$

Hence, it suffices to see that the right-hand map is an (acyclic) fibration, which implies the reduction. Finally, under the natural isomorphism we mentioned at the beginning of the proof, using the fact that right adjoint functors preserve fiber products, the map

$$R(X^{L(B')} \rightarrow R(Y^{L(B')} \times_{Y^{L(A')}} X^{L(A')}))$$

is equivalently given by the induced morphism

$$R(X)^{A'} \rightarrow R(Y)^{B'} \times_{R(Y)^{A'}} R(X)^{A'},$$

which is an (acyclic) fibration in  $\mathbf{M}$  by the assumption that the latter is a Cartesian semimodel category.  $\square$

**Corollary 4.3.3.** *Whenever  $\mathbf{Top}$  is a Cartesian closed category (i.e., in the compactly generated or  $\Delta$ -generated case), the semimodel categories  $\mathbf{Strat}^{\mathfrak{d},\mathfrak{p}}$ ,  $\mathbf{Strat}^{\mathfrak{d}}$ ,  $\mathbf{Strat}^{\mathfrak{c},\mathfrak{p}}$ ,  $\mathbf{Strat}^{\mathfrak{c}}$  are Cartesian closed.*

From this corollary, one may deduce that whenever  $\mathcal{X}$  is a fibrant stratified space and  $\mathcal{A}$  a cofibrant stratified space, in any of the semimodel structures on  $\mathbf{Strat}$ , then the stratified mapping space  $\mathcal{X}^{\mathcal{A}}$  presents the  $\infty$ -categorical internal mapping space in the corresponding  $\infty$ -category. In Section 5.4, we discuss the relationship with the stratified mapping spaces of [Hug99].

## 4.4 Applying the semimodel structures of stratified spaces

Corollary 4.2.3 opens up the study of homotopy theories of stratified spaces to all of the tools available for cofibrantly generated simplicial semimodel categories. Notably, these essentially include all of the tools available for model categories, as long as one is willing to assume cofibrancy in the appropriate places (see, for example, [Bar10; WB23] for a good overview of the literature). In this section, we are going to name a few that stand out, as they provide answers to some questions that have been open for a while (see Questions Q(1), Q(2) and Q(5) in the introduction). Let us begin, by noting that the situation is even better than what one can usually expect from a semimodel category. Namely, the simplicial structure gives us simplicial resolutions of fibrant objects (see [DK80b]), without having to cofibrantly replace first.

**Corollary 4.4.1.** *Let  $\underline{\mathbf{S}}$  be any of the simplicial semimodel categories for stratified spaces of Notation 4.2.5. Then, for any  $\mathcal{X} \in \mathbf{S}$  which is fibrant the functor*

$$\begin{aligned} \Delta^{\text{op}} &\rightarrow \mathbf{Strat} \\ [n] &\mapsto \mathcal{X}^{\Delta^n} \end{aligned}$$

*defines a simplicial resolution of  $\mathcal{X}$ .*

In particular, even non-cofibrant objects admit path objects. As a consequence of this, right-homotopy on maps with cofibrant source and fibrant target becomes an equivalence relation. We may now compute  $\infty$ -categorical mapping spaces of stratified spaces in terms of the simplicial structures.

**Corollary 4.4.2.** *Let  $\underline{\mathbf{S}}$  be any of the simplicial semimodel categories for stratified spaces defined in Section 4.2. Let  $W$  be the class of weak equivalences in  $\mathbf{S}$  and denote  $\mathbf{S}[W^{-1}]$  the quasi-categorical localization of  $\mathbf{S}$  at  $W$ . Let  $\mathcal{A} \in \mathbf{S}$  be cofibrant and  $\mathcal{X}$  be fibrant. Then there is a natural (zigzag) of weak equivalences between mapping spaces*

$$\mathbf{S}[W^{-1}](\mathcal{A}, \mathcal{X}) \simeq \underline{\mathbf{S}}(\mathcal{A}, \mathcal{X}).$$

*In particular, there is a natural bijection*

$$\pi_0(\mathbf{S}[W^{-1}](\mathcal{A}, \mathcal{X})) = [\mathcal{A}, \mathcal{X}]_s$$

*where the right hand expression denotes homotopy classes of stratified (stratum preserving when the poset is fixed) maps with respect to stratified homotopies.*

*Proof.* This follows from [GH05, Prop. 1.1.10] which is the left semimodel category version of the argument in [DK80b, §7], together with [Hin16, p. 1.2].  $\square$

Note that Corollary 4.4.2 applies to all piecewise linear pseudomanifolds or more generally conically stratified spaces, (equipped with the refined stratification, when necessary) and provides in particular a strengthened version of [DW22, Theorem 5.7].

From a global perspective, we obtain the following result, which states that the homotopy theory of stratified spaces obtained by inverting weak equivalences may equivalently be described in terms of the simplicial categories of bifibrant stratified spaces, or equivalently a localization of bifibrant stratified spaces at stratified homotopy equivalences, answering Question Q(2).

**Corollary 4.4.3.** *In the setting of Corollary 4.4.2, denote by  $\underline{\mathbf{S}}^{\circ}$  the full simplicial subcategory of bifibrant objects. Denote by  $H_s$  the class of stratified homotopy equivalences between bifibrant stratified spaces. There are canonical equivalences of  $\infty$ -categories*

$$\underline{\mathbf{S}}^{\circ} \simeq \mathbf{S}^{\circ}[H_s^{-1}] \simeq \mathbf{S}[W^{-1}]$$

*where equivalence between simplicial and quasi-categories are to be understood in terms of the Quillen equivalence between quasi-categories and simplicial categories of [Ber07a].*

*Proof.* This is the semimodel category version of [DK80b, Prop. 4.8] the proof of which is identical. We note that the first equivalence also follows from the existence of cofibrant replacements and the powering of the stratified categories over  $\mathbf{sSet}$ , using a similar flattening argument as in [DK87, p. 2.5]. Therefore, the full power of a semimodel category is not necessary to obtain this result. The second equivalence, however, needs both fibrant replacement and cofibrant replacement, as well as the Whitehead theorem in a semimodel category.  $\square$

Furthermore, we obtain the following two versions of the stratified homotopy hypothesis, providing an answer to Question Q(1).

**Theorem 4.4.4.** *Mapping a simplicial set to  $K$  to the stratified realization of its stratified simplicial set  $\mathcal{L}(K)$ , and conversely mapping a stratified space  $\mathcal{X}$  to the underlying simplicial set of  $\text{Sing}_s \mathcal{X}$ ,  $\mathcal{F}(\text{Sing}_s \mathcal{X})$ , induces a Quillen equivalence of (semi)model categories*

$$\mathbf{sSet}^{\mathcal{D}} \stackrel{\cong}{\rightleftarrows} \mathbf{Strat}^c,$$

which creates weak equivalences in both directions.

The left-hand model category  $\mathbf{sSet}^{\mathcal{D}}$  is the left Bousfield localization of  $\mathbf{sSet}^{\mathcal{J}}$  which corresponds to the full reflective subcategory  $\mathcal{Lay}_{\infty} \subset \mathbf{Cat}_{\infty}$  of small infinity categories in which every endomorphism is an isomorphism.

*Proof.* This is the combination of Corollary 4.2.3 and Theorem 4.2.10 and [Waa24a, Thm. 3.3.6]. The only thing we need to pay extra attention to is the statement that both functors create weak equivalences. In the case of the right adjoint, note that the forgetful functor

$$\mathbf{sStrat}^c \rightarrow \mathbf{sSet}^{\mathcal{D}}$$

does not create weak equivalences. However, it creates weak equivalences between such stratified simplicial sets whose strata are Kan-complexes ([Waa24a, Prop. 3.2.24]). Since  $\text{Sing}_s$  has image in this category, it follows that the composition of  $\text{Sing}_s$  with the functor forgetting the stratification creates weak equivalences. Conversely, note that  $|-|_s \circ \mathcal{L}$  is left Quillen, with source a cofibrant model category. Since the adjunction above is a Quillen equivalence (as the composition of two Quillen equivalences), the derived unit of adjunction is a weak equivalence. For  $X \in \mathbf{sSet}$ , It is given by

$$X \rightarrow \mathcal{F}(\text{Sing}_s |\mathcal{L}(X)|_s) \rightarrow \mathcal{F}\text{Sing}_s \mathcal{Y}$$

where  $|\mathcal{L}(X)|_s \rightarrow \mathcal{Y}$  is a fibrant replacement of  $|\mathcal{L}(X)|_s$ . Note, however, that as we have already shown that  $\mathcal{F} \circ \text{Sing}_s$  preserves weak equivalences, it follows by two-out-of-three that the ordinary unit of adjunction is also a weak equivalence. It follows, again by two-out-of-three, that  $f: X \rightarrow Y$  in  $\mathbf{sSet}^{\mathcal{D}}$  is a weak equivalence, if and only if  $\mathcal{F}(\text{Sing}_s |\mathcal{L}(f)|_s)$  is a weak equivalence. Finally, since  $\mathcal{F} \circ \text{Sing}_s$  creates weak equivalences, this is equivalent to  $\mathcal{L} \circ |-|_s$  creating weak equivalences.  $\square$

A quasi-categorical counterpart of this result, using Corollary 4.4.3 is:

**Corollary 4.4.5.** *Denote by  $\mathbf{Strat}^o$  the full subcategory of  $\mathbf{Strat}^c$  of bifibrant spaces. Denote by  $H_s$  the wide subcategory of stratified homotopy equivalences.  $\text{Sing}_s$  (i.e., Lurie's exit path construction of [Lur17]) induces an equivalence of quasi-categories*

$$\text{Exit}: \mathbf{Strat}^o[H_s^{-1}] \xrightarrow{\cong} \mathcal{Lay}_{\infty},$$

where  $\mathcal{Lay}_{\infty}$  denotes the quasi-category of small quasi-categories, in which every endomorphism is an isomorphism.

Finally, let us comment on Question Q(5), concerning the relationship between homotopy colimits and 1-categorical colimits of stratified spaces. The main application for the methods of regular neighborhoods in stratified cell complexes we developed in [Waa24b] was to prove that certain pushout diagrams of stratified spaces are homotopy pushout. Note that, a posteriori, this also follows from the existence of semimodel structures making inclusions of cellularly stratified spaces cofibrations. In fact, we can now use the full machinery for the computation of homotopy colimits in a simplicial (semi)model category (see, for example, [Hir09]) to compute the latter. As a particular consequence, we obtain the following result.

**Corollary 4.4.6.** *Suppose we are given a pushout diagram of stratified spaces (in  $\mathbf{Strat}_P$  or  $\mathbf{Strat}$ )*

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{c} & \mathcal{B} \\ \downarrow f & & \downarrow \\ \mathcal{X} & \hookrightarrow & \mathcal{Y}, \end{array} \tag{15}$$

*all of which are cofibrant in  $\mathbf{Strat}^{\text{d},\text{p}}$  (or equivalently  $\mathbf{Strat}^{\text{c},\text{p}}$ ) (that is, retracts of cellularly stratified spaces, by Proposition 5.2.1), with the upper vertical given by a cofibration. Then this diagram is homotopy pushout (with respect to diagrammatic and categorical weak equivalences). In particular, this holds when all spaces involved are cellularly stratified spaces and  $c$  is an inclusion of cellularly stratified spaces.*

Clearly, Lemma 4.2.2 is implied by this result. Lemma 4.2.2 was used to prove the existence of semimodel structures. Now, however, we can see that the existence of semimodel structures is essentially equivalent to diagrams of stratified cell complexes as in Diagram (15) being homotopy pushout, which illustrates the importance of Question Q(5).

## 5 (Co)fibrant stratified spaces

In the previous section, we have established the existence of semimodel structures of stratified spaces, which were transferred from the combinatorial setting of stratified simplicial sets. These were constructed with the specific goal of connecting the various homotopy theories of stratified spaces with their topology and geometry. If we are looking to deepen our understanding of this connection, the obvious question at hand is how classical examples of stratified spaces interact with these model structures. In particular, which classical examples of stratified spaces are bifibrant (Question Q(3)). For example, [Mil13; Dou21c; Nan19] all showed Whitehead theorems for certain stratified spaces, which characterize the stratified homotopy equivalences between them in terms of a priori weaker conditions. Every simplicial semimodel category comes with its own Whitehead theorem, stating that the homotopy equivalences with respect to the simplicial cylinder between the bifibrant objects are precisely the weak equivalences. The case of  $\mathbf{sStrat}^{\text{c}}$  was already proven through a direct proof in [Nan19]. Summarizing old and proving new criteria for (co)fibrancy and studying how these properties relate to more classical properties of stratified spaces is precisely the content of this section.

### 5.1 Fibrant stratified spaces and Quinn’s homotopically stratified sets

Let us begin by studying the class of fibrant objects in the model structures constructed in Section 4.2. The question of what classical examples are fibrant in  $\mathbf{Strat}^{\text{c}}$  was already investigated in [Nan19] and [Lur17], with the core results that both Quinn’s homotopically stratified spaces and Siebenmann’s conically stratified spaces (see later in this section) are such that the associated stratified singular simplicial set is a quasi-category. Hence, it turns out that most examples of stratified spaces of classical geometrical interest fall into the class of fibrant objects in the semimodel structures we defined. The central new contribution from our side in this section is that we prove that, as long as one restricts to metrizable spaces, the class of fibrant objects is in fact independent of the choice of model structure we presented here (see Proposition 5.1.4). A consequence of this is Theorem 5.1.6, which may be taken as the statement that in any geometric scenario there really is no difference between the diagrammatic homotopy theories for stratified spaces and their categorical counterparts at all.

Before we begin, note that we really only need to study fibrancy in the scenario of a fixed poset, due to the following fact:

**Lemma 5.1.1.** *Let  $\mathcal{X} \in \mathbf{Strat}$ . Then the following statements are equivalent:*

1.  $\mathcal{X}$  has the horn filling property over  $P_{\mathcal{X}}$  with respect to all admissible stratified horn inclusions. That is, it admits fillers

$$\begin{array}{ccc}
 |\Lambda_k^{\mathcal{J}}|_s & \longrightarrow & \mathcal{X} \\
 \downarrow & \nearrow \text{dashed} & \\
 |\Delta^{\mathcal{J}}|_s & & 
 \end{array}
 \tag{16}$$

in  $\mathbf{Strat}_{P_{\mathcal{X}}}$ , for all admissible pairs  $\mathcal{J} \in \Delta_P$ ,  $0 \leq k \leq n_{\mathcal{J}}$ ;

2.  $\mathcal{X}$  is fibrant in all of the semi model categories  $\mathbf{Strat}_{P_{\mathcal{X}}}^{\circ}, \mathbf{Strat}^{\circ, \text{p}}, \mathbf{Strat}^{\circ}$ ;
3.  $\mathcal{X}$  is fibrant in one of the semi model categories  $\mathbf{Strat}_{P_{\mathcal{X}}}^{\circ}, \mathbf{Strat}^{\circ, \text{p}}, \mathbf{Strat}^{\circ}$ .

Furthermore, the following analogous statements about the categorical semimodel structures holds:

1.  $\mathcal{X}$  has the horn filling property over  $P_{\mathcal{X}}$  with respect to all admissible stratified horn inclusions, and all inner stratified horn inclusions.
2.  $\mathcal{X}$  has the horn filling property over  $P_{\mathcal{X}}$  with respect to all inner stratified horn inclusions.
3.  $\mathcal{X}$  is fibrant in all of the semi model categories  $\mathbf{Strat}_{P_{\mathcal{X}}}^{\text{c}}, \mathbf{Strat}^{\text{c}, \text{p}}, \mathbf{Strat}^{\text{c}}$ ;
4.  $\mathcal{X}$  is fibrant in one of the semi model categories  $\mathbf{Strat}_{P_{\mathcal{X}}}^{\text{c}}, \mathbf{Strat}^{\text{c}, \text{p}}, \mathbf{Strat}^{\text{c}}$ .

*Proof.* The diagrammatic case is immediate by the construction of the left model structures via left-transfer, Proposition 4.2.8 together with the fact that the refined categories are obtained via a right Bousfield localization Theorem 4.2.11. For the categorical statement, the proof is almost identical; special attention only needs to be paid to the implication from the second to the first property. Note that for any stratified space  $\mathcal{X}$  the strata of the stratified-simplicial set  $\text{Sing}_s \mathcal{X}$  are Kan-complexes. Hence, the implication follows from [Hai23, Prop. 2.2.3].  $\square$

**Definition 5.1.2.** We call a stratified topological space  $\mathcal{X}$  that satisfies any of the equivalent conditions of Lemma 5.1.1 concerning the diagrammatic (categorical) semi model structures **diagrammatically (categorically) fibrant**.

Clearly, every categorically fibrant stratified space is diagrammatically fibrant. The converse must necessarily be false. If it were true, then by Corollary 4.4.3 and Theorem 3.3.1 together with [Hai23, Thm. 1.1.7] and [Dou21c, Thm. 3], it would imply that the inclusion from the  $\infty$ -category of décollages into  $\mathbf{Fun}((\text{sd}P)^{\text{op}}, \mathbf{Spaces})$  is an equivalence of categories<sup>6</sup>. However, it is surprisingly hard to provide a geometrical example for a stratified space which is diagrammatically fibrant and not categorically fibrant (depending on your definition of geometric, it is not possible at all). Before we investigate this question further, let us first recall results on which classical examples of stratified space we can expect to be fibrant.

<sup>6</sup>It is easy to write down an example of a diagram which is not a décollage. For example, take the constant diagram over  $(\text{sd}[2])^{\text{op}}$  and replace the entry at  $[0 < 1 < 2]$  by the empty set.

**Remark 5.1.3.** It was first shown in [Lur17, A.5], that all conically stratified spaces - roughly a stratified space that locally has the structure  $U \times C(\mathcal{L})$ , where  $C(\mathcal{L})$  is the stratified cone of a stratified space  $\mathcal{L}$ , are categorically fibrant. In particular, this implies that all topological pseudomanifolds (see, for example, [Ban07]), as well as all conically smooth stratified spaces (see [AFT17]) and hence also all Whitney stratified spaces ([NV23] or classically [Tho69]), are categorically and hence also diagrammatically fibrant.

It turns out, however, that fibrancy can already be obtained under significantly less geometric conditions. One of the crucial insights contributing to the foundations of stratified homotopy theory was Quinn's observation that a powerful homotopy theoretical setup for stratified spaces can already be achieved by only posing requirements on pairs of strata ([Qui88]). Quinn defined a notion of homotopically stratified set (with slightly more restrictive conditions on the stratification poset) by requiring  $\mathcal{X} \in \mathbf{Strat}$  to be metrizable, and requiring that for any two-element flag  $[p < q] \in \Delta_P$  it holds that:

1.  $\mathcal{X}_p \hookrightarrow \mathcal{X}_{p < q}$  is tame, i.e. admits a nearly strict neighborhood deformation retraction [Qui88].
2.  $\mathcal{H}\text{olink}_{p < q} \mathcal{X} \rightarrow \mathcal{X}_p$  is a Hurewicz fibration.

Together, these assumptions provided an excellent homotopy theoretic framework for classification results concerning manifold stratified spaces [Qui88; Wei94]. Note that both of these assumptions are more in line with the Hurewicz approach (see [Str72]), than with the Serre-Quillen approach to homotopy theory ([Qui67]). Since the approach we pursue here is combinatorial (i.e. following Serre and Quillen), we should expect cofibrancy in our semimodel categories to be a stronger condition than condition (1) above. Indeed, it is a consequence of [Waa24b, Prop. 4.2.4] that any locally compact cofibrant stratified space (in any of the model structures of Section 4.2) satisfies the tameness condition above. Conversely, in the two strata case it is not hard to see that  $\mathcal{H}\text{olink}_{p < q} \mathcal{X} \rightarrow \mathcal{X}_p$  being a Serre fibration is equivalent to diagrammatic and equivalently categorical fibrancy (see the proof of Proposition 5.1.4 below). Surprisingly, it follows from [Mil09, Thm. 4.9] that every homotopically stratified set is, in fact, categorically fibrant ([Nan19, Prop. 8.1.2.6]), without any restrictions on the stratification poset. Note that being fibrant is manifestly a statement about the interaction of more than two strata, which makes it so surprising that it can be achieved by only making requirements on two-strata interaction. Even more, this result does not leave much space for diagrammatically fibrant spaces that are not categorically fibrant at all (see Question Q(4)). If we assume pairwise tameness and metrizability, then it would follow that any counterexample would involve  $\mathcal{H}\text{olink}_{p < q} \mathcal{X} \rightarrow \mathcal{X}_p$  being a Serre fibration, but not a Hurewicz fibration. It turns out that no cofibrancy assumptions are necessary whatsoever, and in fact in any geometric setting diagrammatic and categorical fibrancy are equivalent conditions, which are furthermore equivalent to the Serre fibration analogue of Quinn's pairwise homotopy link condition. Furthermore, this result does not require any additional tameness assumptions.

**Proposition 5.1.4.** *Let  $\mathcal{X} \in \mathbf{Strat}$  be a metrizable stratified space. Then the following conditions are equivalent:*

- (i)  $\mathcal{X}$  is categorically fibrant;
- (ii)  $\mathcal{X}$  is diagrammatically fibrant;
- (iii) For any pair  $[p < q] \in \Delta_{P_{\mathcal{X}}}$ , the starting point evaluation map  $\mathcal{H}\text{olink}_{p < q}(\mathcal{X}) \rightarrow \mathcal{X}_p$  is a Serre fibration.
- (iv) For any pair  $[p < q] \in \Delta_{P_{\mathcal{X}}}$ , and for any flag  $\mathcal{J} = [p = \dots = p_k < q = \dots = q = p_{n_{\mathcal{J}}}]$  with  $k \geq 1$ ,  $\mathcal{X}$  has the horn filling property with respect to  $|\Lambda_k^{\mathcal{J}} \hookrightarrow \Delta^{\mathcal{J}}|_s$ ;

Before we give a proof, we note the implications of this result. By Theorem 4.2.11, the categorical setting is always obtained as a left Bousfield localization of the categorical one. In particular, the two settings have the same cofibrations, furthermore we obtain:

**Corollary 5.1.5.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathbf{Strat}$  be a metrizable stratified spaces,  $\mathcal{X}$  diagrammatically fibrant and  $f: \mathcal{Y} \rightarrow \mathcal{X}$  a stratified map. Then there are equivalences:*

1.  $f$  is a fibration in  $\mathbf{Strat}_P^{\mathfrak{d}}$   $\iff$   $f$  is a fibration in  $\mathbf{Strat}_P^{\mathfrak{c}}$ ;
2.  $f$  is a fibration in  $\mathbf{Strat}^{\mathfrak{d}, \mathfrak{p}}$   $\iff$   $f$  is a fibration in  $\mathbf{Strat}^{\mathfrak{c}, \mathfrak{p}}$ ;
3.  $f$  is a fibration in  $\mathbf{Strat}^{\mathfrak{d}}$   $\iff$   $f$  is a fibration in  $\mathbf{Strat}^{\mathfrak{c}}$ .

*Proof.* This is immediate from Proposition 5.1.4, together with Theorem 4.2.11, and the characterization of fibrations between fibrant objects in a left Bousfield localizations (the semimodel category version of [Hir09, Prop. 3.3.16.], which holds in any cofibrantly generated left-semi model category. In case the reader is worried about this claim, note that one can simply use the model categorical version, and use the fact that the semimodel structures on topological stratified spaces are transferred from model structures in the simplicial world.)  $\square$

We may hence interpret Proposition 5.1.4 as stating that in any geometrical framework the diagrammatic approach is really the same as the categorical approach, obtaining an answer to Question Q(4). More rigorously, we can phrase this insight as follows.

**Theorem 5.1.6.** *Let  $\underline{\mathbf{D}}$  be any of the diagrammatic simplicial semimodel categories of Section 4.2 and let  $\underline{\mathbf{C}}$  be its categorical pendant. Denote by  $\underline{\mathbf{D}}_m^f$  and  $\underline{\mathbf{C}}_m^f$  the respective restrictions of the semimodel structures to the full simplicial subcategory of fibrant, metrizable stratified spaces. Then equality*

$$\underline{\mathbf{D}}_m^f = \underline{\mathbf{C}}_m^f$$

*holds (on the nose). In particular, if we denote by  $\underline{\mathbf{D}}_m^{\mathfrak{o}}$  and  $\underline{\mathbf{C}}_m^{\mathfrak{o}}$  the corresponding simplicial categories of bifibrant metrizable stratified spaces, then*

$$\underline{\mathbf{D}}_m^{\mathfrak{o}} = \underline{\mathbf{C}}_m^{\mathfrak{o}}$$

*and hence the full sub- $\infty$ -categories of bifibrant metrizable stratified spaces agree.*

*Proof.* Equality on the object level follows by Proposition 5.1.4. Equality of cofibrations holds even without additional assumptions. Equality of fibrations was shown in Corollary 5.1.5. It remains to verify the equality of weak equivalences. We prove the stratum preserving case, the others are analogous: Let  $f: \mathcal{X} \rightarrow \mathcal{Y}$  be a stratum preserving map in  $\underline{\mathbf{D}}_m^f$ . We may factorize  $f$  as a cofibration and an acyclic fibration

$$\mathcal{X} \rightarrow \mathcal{Z} \rightarrow \mathcal{Y},$$

with  $\mathcal{Z}$  not necessarily metrizable. Note that since cofibrations in both structures agree, so do acyclic fibrations. By Proposition 5.1.4, it follows that  $\mathcal{Z}$  is also categorically fibrant (this did not use any metrizability assumptions on  $\mathcal{Z}$ ).  $f$  is a weak equivalence if and only if the first of these two maps is a weak equivalence. However, the latter is a map between categorically fibrant spaces. In particular, by the semimodel category version of the Whitehead theorem for Bousfield localizations (see [Hir09, Thm 3.2.13] for the model category version)<sup>7</sup>, it is a diagrammatic equivalence if and only if it is a categorical equivalence.  $\square$

<sup>7</sup>[WB23, Rem. 4.5] shows the semimodel category case for bifibrant objects in the localized model structure. Note, however, that we may always assume bifibrancy, by replacing both source and target cofibrantly first, which is done through acyclic fibrations, which are the same in both model structures. Alternatively, we can just use the full model categorical result and pass to the simplicial setting using the fact that all semimodel structures are transferred.

In other words, for geometric examples, the two homotopy theories agree. Before we provide a proof of Proposition 5.1.4, we need the following elementary set-theoretic topological lemma, the proof of which is provided in Appendix B. It is the only place where the metrizable of  $\mathcal{X}$  comes into play:

**Lemma 5.1.7.** *Let  $n \geq 0$ ,  $X$  a metrizable topological space and suppose that we are given a solid commutative diagram of (general) topological spaces of the following form:*

$$\begin{array}{ccccc}
 & & D^{n+1} \times \{0\} \cup S^n \times [0, 1] & & \\
 & \nearrow & \downarrow & \searrow f & \\
 D^{n+1} \times \{0\} \cup S^n \times [0, 1] & & D^{n+1} \times [0, 1] & \xrightarrow{\tilde{f}} & X \\
 & \searrow & \uparrow & \nearrow f' & \\
 & & D^{n+1} \times [0, 1] & & 
 \end{array} \tag{17}$$

Then there exists  $\tilde{f}: D^{n+1} \times [0, 1] \rightarrow X$  making the upper triangle commute, and furthermore such that the inclusion

$$\tilde{f}(\mathring{D}^{n+1} \times (0, 1]) \subset f'(\mathring{D}^{n+1} \times (0, 1])$$

holds.

As a consequence of Lemma 5.1.7 we obtain:

**Lemma 5.1.8.** *Let  $\mathcal{X} \in \mathbf{Strat}_P$  be a metrizable stratified space and  $\mathcal{J} = [p_0 \leq \dots \leq p_n]$  be a flag in  $P$ . Let  $0 < k < n$  and denote by  $\mathcal{I}$  the regular flag containing all  $p_i \in \mathcal{J}$  with  $i \geq k$ . If  $\mathcal{X}_{\mathcal{I}}$  is categorically fibrant, then  $\mathcal{X}$  has the horn filling property with respect to  $|\Lambda_k^{\mathcal{J}} \hookrightarrow \Delta^{\mathcal{J}}|_s$ .*

*Proof.* Let  $\mathcal{J}_k$  be the flag obtained by removing  $p_k$  from  $\mathcal{J}$ , corresponding to the  $k$ -th face of  $\Delta^{\mathcal{J}}$ . We may identify  $|\Delta^{\mathcal{J}}|_s$  with an (appropriately stratified) join  $|\Delta^{[p_k]}|_s \star |\Delta_{k}^{\mathcal{J}}|_s$ . This identification induces join coordinates  $[s, x]$ ,  $s \in [0, 1]$ ,  $x \in |\Delta_{k}^{\mathcal{J}}|_s$  on  $|\Delta^{\mathcal{J}}|_s$ , with the point  $[0, x]$  corresponding to the unique element of  $|\Delta^{[p_k]}|_s$ . Under this identification,  $|\Lambda^{\mathcal{J}}|_{s_k}$  corresponds to the join  $|\Delta^{p_k}|_s \star |\partial \Delta_k^{\mathcal{J}}|_s$ . For  $0 \leq \alpha \leq \alpha' < 1$ , we write

$$S_{\geq \alpha}^{\leq \alpha'} = \{[x, s] \in |\Delta^{[p_k]}|_s \star |\Delta_k^{\mathcal{J}}|_s \mid \alpha \leq s \leq \alpha'\},$$

and use analogous notation replacing  $\leq$  by  $<$  and  $\geq$  by  $>$ . We may then decompose  $|\Delta^{\mathcal{J}}|_s$  into

$$|\Delta^{\mathcal{J}}|_s = S_{\geq 0}^{\leq \frac{1}{2}} \cup S_{\geq \frac{1}{2}}^{\leq 1}.$$

It is immediate by the definition of the stratified simplex  $|\Delta^{\mathcal{J}}|_s$ , that the inclusion

$$|\Lambda_k^{\mathcal{J}}| \cap S_{\geq 0}^{\leq \frac{1}{2}} \hookrightarrow S_{\geq 0}^{\leq \frac{1}{2}}$$

is stratum preserving homeomorphic to the inclusion

$$|\Lambda_k^{\mathcal{J}'}|_s \hookrightarrow |\Delta^{\mathcal{J}'}|_s,$$

where  $\Delta^{\mathcal{J}'}$  is obtained by replacing every entry  $p_i$  in  $\mathcal{J}$  with  $i < k$  by  $p_k$ . In particular, by the assumption on  $\mathcal{X}_{\mathcal{I}}$ ,  $\mathcal{X}$  admits filler with respect to the latter inclusion. Thus, it suffices to show that  $\mathcal{X}$  admits fillers with respect to

$$A := S_{\geq \frac{1}{2}}^{\leq 1} \cap |\Lambda_k^{\mathcal{J}}| \cup S_{\geq \frac{1}{2}}^{\leq \frac{1}{2}} \hookrightarrow S_{\geq \frac{1}{2}}^{\leq 1} =: B.$$



Choosing any homeomorphism  $|\Delta^n| \cong D^{m+1}$ , where  $m = n - 2$ , and using the affine order preserving homeomorphism  $[\frac{1}{2}, 1] \cong [0, 1]$  we may identify the latter inclusion with a nonstratified inclusion

$$D^{m+1} \times \{0\} \cup S^m \times [0, 1] \hookrightarrow D^{m+1} \times [0, 1].$$

Since  $p_k \leq p_n$ , under this identification  $\mathring{D}^{m+1} \times (0, 1]$  is entirely contained in the  $p_n$ -stratum of  $|\Delta^{\mathcal{J}}|_s$ . Thus, given a stratum preserving map  $f: A \rightarrow \mathcal{X}$ , an extension of  $f$  to  $B$  is the same data as an extension of  $f$  to  $D^{m+1} \times [0, 1]$  mapping  $\mathring{D}^{m+1} \times (0, 1]$  to  $\mathcal{X}_{p_n}$ . We may now apply Lemma 5.1.7, from which it follows that it suffices to obtain an extension of  $f_{D^{m+1} \times \{0\} \cup S^m \times [0, 1]}$  to  $D^{m+1} \times [0, 1] \cong S_{\frac{1}{2}}^{<1}$ . Now write

$$S_{\frac{1}{2}}^{<1} = \bigcup_{l \geq 1} S_{\geq 1 - \frac{1}{2^l}}^{\leq 1 - \frac{1}{2^{l+1}}},$$

which defines a locally finite, closed cover of  $S_{\geq 1 - \frac{1}{2^l}}^{\leq 1 - \frac{1}{2^{l+1}}}$ . Inductively, we may hence reduce to solving extension problems with respect to

$$S_{\geq 1 - \frac{1}{2^l}}^{\leq 1 - \frac{1}{2^l}} \cup (|\Lambda_k^{\mathcal{J}}|_s \cap S_{\geq 1 - \frac{1}{2^l}}^{\leq 1 - \frac{1}{2^{l+1}}}) \hookrightarrow S_{\geq 1 - \frac{1}{2^l}}^{\leq 1 - \frac{1}{2^{l+1}}}.$$

Finally, if we denote by  $\mathcal{J}'_k$  the  $k$ -th face of  $\mathcal{J}'$ , then the latter inclusion is stratum preserving homeomorphic to the simplicial box product

$$|\Delta^{\mathcal{J}'_k}|_s \otimes \Delta^0 \cup_{|\partial \Delta^{\mathcal{J}'_k}|_s \otimes \Delta^0} |\partial \Delta^{\mathcal{J}'_k}|_s \otimes \Delta^1 \hookrightarrow |\Delta^{\mathcal{J}'_k}|_s \otimes \Delta^1,$$

for the inclusion of  $\Delta^0$  into  $\Delta^1$  at 0. Since  $\mathbf{Strat}_{\mathcal{J}}^c$  is a simplicial semimodel category, the latter defines an acyclic cofibration in  $\mathbf{Strat}_{\mathcal{J}}^c$ . Since  $\mathcal{X}_{\mathcal{J}}$  was assumed to be fibrant, the existence of extensions with respect to such an inclusion follows.  $\square$

We may now complete the following proof.

*Proof of Proposition 5.1.4.* Clearly, Property (i) implies Property (ii). To see that Property (ii) implies Property (iii), note that any lifting problem

$$\begin{array}{ccc} |\Lambda^n| & \longrightarrow & \mathcal{H}o\text{Link}_{p < q}(\mathcal{X}) \\ \downarrow & \nearrow \text{dashed} & \downarrow \\ |\Delta^n| & \longrightarrow & \mathcal{X}_p \end{array} \quad (18)$$

is equivalent to a lifting problem (over  $P_{\mathcal{X}}$ )

$$\begin{array}{ccc} |\Delta^{[p]}|_s \otimes \Delta^n \cup_{|\Delta^{[p]}|_s \otimes \Delta^n} |\Delta^{[p < q]}|_s \otimes \Delta^n & \longrightarrow & \mathcal{X} \\ \downarrow & \nearrow \text{dashed} & \\ |\Delta^{[p < q]}|_s \otimes \Delta^n & & \end{array} \quad (19)$$

Hence, Property (iii) follows from the simpliciality of the semi model category  $\mathbf{Strat}_{P_{\mathcal{X}}}^{\mathfrak{d}}$ . For the implication Property (iii)  $\implies$  Property (iv), note that using the identification  $|\Delta^{\mathcal{J}}|_s$  with the (appropriately stratified) join  $|\Delta^{\mathcal{J}_p}|_s \star |\Delta^{\mathcal{J}_q}|_s$ , we can interpret  $|\Delta^{\mathcal{J}}|_s$  as a stratified quotient space of

$$|\Delta^{[p < q]}|_s \otimes (\Delta^k \times \Delta^{n-k-1}),$$

obtained by collapsing  $|\Delta^{[p]}|_s \otimes (\Delta^k \times \Delta^{n-k-1})$  to  $|\Delta^{[p]}|_s \otimes \Delta^k$  and  $|\Delta^{[q]}|_s \otimes (\Delta^k \times \Delta^{n-k-1})$  to  $|\Delta^{[q]}|_s \otimes \Delta^{n-k-1}$ . Within  $|\Delta^{[p<q]}|_s \otimes (\Delta^k \times \Delta^{n-k-1})$ ,  $|\Lambda_k^{\mathcal{J}}|_s$  corresponds to

$$\begin{array}{c} |\Delta^{[p<q]}|_s \otimes (\Lambda_k^k \times \Delta^{n-k-1}) \cup |\Delta^{[p]}|_s \otimes (\Delta^k \times \Delta^{n-k-1}) \cup |\Delta^{[q]}|_s \otimes (\Delta^k \times \Delta^{n-k-1}) \\ \downarrow \\ |\Delta^{[p<q]}|_s \otimes (\Delta^k \times \Delta^{n-k-1}) \end{array} \quad (20)$$

Hence, it suffices to show that  $\mathcal{X}$  has the extension property with respect to this stratum preserving map over  $P_{\mathcal{X}}$ . Denote  $A = \Lambda_k^k \times \Delta^{n-k-1}$  and  $B = \Delta^k \times \Delta^{n-k-1}$ . Under the identification  $|\Delta^{[p<q]}|_s \cong |\Delta^{[p<q]}|_s \cup_{|\Delta^{[q]}|_s} |\Delta^{[q\leq q]}|_s$  we may decompose (20) into pushouts of inclusions

$$|\Delta^{[p]}|_s \otimes B \cup_{|\Delta^{[p]}|_s \otimes A} |\Delta^{[p<q]}|_s \otimes A \hookrightarrow |\Delta^{[p<q]}|_s \otimes B \quad (21)$$

and

$$|\partial\Delta^{[q\leq q]}|_s \otimes B \cup_{|\partial\Delta^{[q\leq q]}|_s \otimes A} |\Delta^{[q<q]}|_s \otimes A \hookrightarrow |\Delta^{[q\leq q]}|_s \otimes B. \quad (22)$$

The inclusion  $A \hookrightarrow B$  is an anodyne map of simplicial sets. It follows that the inclusion (22) which lies entirely over one stratum is given by an acyclic Quillen cofibration. Consequently,  $\mathcal{X}$  has the right lifting property with respect to the inclusion (22). Finally, lifting problems with respect to (21) are equivalent to lifting problems of the form

$$\begin{array}{ccc} |A| & \longrightarrow & \mathcal{H}o\text{Link}_{p<q}(\mathcal{X}) \\ \downarrow & \nearrow \text{dashed} & \downarrow \\ |B| & \longrightarrow & \mathcal{X}_p, \end{array} \quad (23)$$

since  $A \hookrightarrow B$  is an anodyne extension, the left-hand vertical is an acyclic cofibration in the Quillen model structure, showing the existence of a lift.

It remains to show that Property (iv) implies Property (i). It suffices to show that  $\mathcal{X}_{\mathcal{I}}$  is categorically fibrant for all regular flags  $\mathcal{I} = [q_0 < \dots < q_n]$  of  $P_{\mathcal{X}}$ . We only need to cover the case where  $n \geq 1$ , that is, where  $\mathcal{J}$  has at least two different elements (otherwise, we are in the trivially stratified case, that is, the classical case). First, note that Property (i) already implies the case where  $\mathcal{I} = [p < q]$ . Indeed, let  $\mathcal{J} = [p_0 \leq \dots \leq p_{n_{\mathcal{J}}}]$  be a flag that degenerates from  $\mathcal{I}$ . When  $p_k = p$  we may without loss of generality assume that  $k$  is maximal with the property that  $p_k = p$ , by permuting the corners of  $|\Delta^{\mathcal{J}}|_s$  by a stratum preserving homeomorphism. Then we are in the situation of Property (iv). In case when  $p_k = q$ , the inclusion  $|\Lambda_k^{\mathcal{J}} \hookrightarrow \Delta^{\mathcal{J}}|_s$  admits a stratum preserving retraction. (Consider a homeomorphism of  $|\Delta^{\mathcal{J}}|_s$  with  $D^n$ , mapping the  $k$ -th face to the northern hemisphere. Then the  $p$ -stratum is entirely contained in the equator, and projecting vertically down to the southern hemisphere defines a retraction.) We now proceed to show the horn filling property with respect to arbitrary  $\mathcal{J}$ , by induction over  $n_{\mathcal{I}}$ , the length of  $\mathcal{I}$ . We have already covered the case  $n_{\mathcal{I}} = 1$ . Now for the inductive step let  $\mathcal{J} = [p_0 \leq \dots \leq p_m]$  be a flag degenerating from a subflag of  $\mathcal{I}$ , and  $0 < k < m$ . For such  $\mathcal{J}$ , denote by  $t_{\mathcal{J}}$  the length of  $\mathcal{J}_{p_0}$  and by  $o_{\mathcal{J}}$  the length of  $\mathcal{J}_{q_{n-1}}$ . We proceed by double induction on  $t_{\mathcal{J}}$  and  $o_{\mathcal{J}}$ , keeping  $k$  flexible. If  $o_{\mathcal{J}} = -1$ , then  $\mathcal{J}$  degenerates from a proper subflag of  $\mathcal{I}$ , and we are reduced to the inductive assumption in the induction on  $n_{\mathcal{I}}$ . If  $p_0 \neq p_k$ , which is in particular the case when  $t_{\mathcal{J}} = 0$  (since  $0 < k < n$ ), then  $\{q_i \mid i \in [n_{\mathcal{I}}], q_i \geq p_k\}$  has cardinality smaller than  $\mathcal{I}$ , and by inductive assumption (in  $n_{\mathcal{I}}$ ) we can apply Lemma 5.1.8 proving the existence of a filler with respect to  $\mathcal{J}, k$ . So, suppose  $t_{\mathcal{J}} \geq 1$ ,  $p_0 = p_k$  and we have already proven the result for all cases  $\tilde{\mathcal{J}}, \tilde{k}$  where either  $t_{\tilde{\mathcal{J}}} < t_{\mathcal{J}}$  or  $o_{\tilde{\mathcal{J}}} < o_{\mathcal{J}}$ . Let  $s \in [n_{\mathcal{J}}]$  be such that  $e_s \in |\Delta^{\mathcal{J}}|_s$  is the maximal vertex lying in

the  $q_{n_{\mathcal{I}}-1}$  stratum and  $e_{n_{\mathcal{J}}} \in |\Delta^{\mathcal{J}}|_s$  the  $n_{\mathcal{J}}$ -th vertex of  $|\Delta^{\mathcal{J}}|_s$ . We may assume that  $s \geq 2$ , as  $t_{\mathcal{J}} \geq 1$  and  $n_{\mathcal{I}} \geq 2$ . Furthermore, we may assume that  $e_{n_{\mathcal{J}}}$  lies in the  $q_n$  stratum, as otherwise  $\mathcal{J}$  degenerates from a proper subflag of  $\mathcal{I}$ , and we are reduced to the inductive assumption. Now, let

$$e' = \frac{1}{2}e_s + \frac{1}{2}e_{n_{\mathcal{J}}}$$

be the halfway point between these two vertices, which by assumption also lies in the  $q_{n_{\mathcal{I}}}$  stratum. Consider the two affine stratum preserving embeddings

$$i_1: |\Delta^{\mathcal{J}}|_s \hookrightarrow |\Delta^{\mathcal{J}}|_s$$

$$e_i \mapsto \begin{cases} e_i & i < n_{\mathcal{J}} \\ e' & i = n_{\mathcal{J}} \end{cases}$$

and

$$i_2: |\Delta^{\mathcal{J}'}|_s \hookrightarrow |\Delta^{\mathcal{J}}|_s$$

$$e_i \mapsto \begin{cases} e_i & i \neq s \\ e' & i = s \end{cases}$$

where  $\mathcal{J}'$  is obtained from  $\mathcal{J}$  by replacing the  $s$ -th entry by  $q_{n_{\mathcal{I}}}$ . The images of these two embeddings cover  $|\Delta^{\mathcal{J}}|_s$  and intersect in the convex span

$$\langle \{e', e_i \mid i \in [n_{\mathcal{J}}], i \neq n_{\mathcal{J}}, s\} \rangle.$$

Under  $i_1$  this span corresponds to the  $s$ -th face of  $\Delta^{\mathcal{J}}$ , and under  $i_2$  to the  $n_{\mathcal{J}}$ -th face of  $\Delta^{\mathcal{J}'}$ , which are both given by the flag  $\mathcal{J}''$  obtained by removing  $p_s$  from  $\mathcal{J}$ . We obtain an induced stratum preserving homeomorphism (see Fig. 1 for an illustration)

$$|\Delta^{\mathcal{J}} \cup_{\Delta^{\mathcal{J}''}} \Delta^{\mathcal{J}'}|_s \cong |\Delta^{\mathcal{J}}|_s \cup_{|\Delta^{\mathcal{J}''}|_s} |\Delta^{\mathcal{J}'}|_s \xrightarrow{(i_1, i_2)} |\Delta^{\mathcal{J}}|_s.$$




Figure 1: Depiction of  $\Delta^{\mathcal{J}} \cup_{\Delta^{\mathcal{J}''}} \Delta^{\mathcal{J}'}$  with  $\Delta^{\mathcal{J}}$  spanned by  $e_0, e_1, e_2, e'$  and  $\Delta^{\mathcal{J}'}$  spanned by  $e_0, e_1, e', e_3$ .

Under this identification  $\Lambda_k^{\mathcal{J}}$  corresponds to the subcomplex  $A$  of  $\Delta^{\mathcal{J}} \cup_{\Delta^{\mathcal{J}''}} \Delta^{\mathcal{J}'} =: B$  given by removing:

- The simplices corresponding to  $\Delta^{\mathcal{J}}$  and  $\Delta^{\mathcal{J}'}$ ;
- The  $k$ -th face of  $\Delta^{\mathcal{J}}$ ,  $\Delta^{\mathcal{J}''}$ , and of  $\Delta^{\mathcal{J}'}$ ,  $\Delta^{\mathcal{J}''}$ ;

- The  $s$ -th face of  $\Delta^{\mathcal{J}}$ ,  $\Delta^{\mathcal{J}_s}$ , which is equivalently the  $n_{\mathcal{J}}$ -th face of  $\Delta^{\mathcal{J}'}$ , or the top dimensional simplex of  $\Delta^{\mathcal{J}''}$ .
- The  $k$ -th face of  $\Delta^{\mathcal{J}''}$ ,  $\Delta^{\mathcal{J}_k}$ , which is equivalently the  $(s-1)$ -th face of  $\Delta^{\mathcal{J}_k}$ .




Figure 2: Depiction of  $\Delta^{\mathcal{J}_k}$ ,  $\Delta^{\mathcal{J}}$  and  $\Delta^{\mathcal{J}'}$  with the vertices respectively opposite to  $\Delta^{\mathcal{J}_k}$ ,  $\Delta^{\mathcal{J}''}$  and  $\Delta^{\mathcal{J}_k}$  marked in green.

Next, consider the inclusions

1.  $j_1: \Lambda_{s-1}^{\mathcal{J}_k} \hookrightarrow \Delta^{\mathcal{J}_k}$ ;
2.  $j_2: \Lambda_s^{\mathcal{J}} \hookrightarrow \Delta^{\mathcal{J}}$ ;
3.  $j_3: \Lambda_k^{\mathcal{J}'} \hookrightarrow \Delta^{\mathcal{J}'}$ .

Gluing  $\Delta^{\mathcal{J}_k}$  to  $A$  along  $j_1$  adds in the missing simplices  $\Delta^{\mathcal{J}_k}$  and  $\Delta^{\mathcal{J}_k''}$ . Then, gluing in  $\Delta^{\mathcal{J}}$  along  $j_2$  adds the missing simplices  $\Delta^{\mathcal{J}}$  and  $\Delta^{\mathcal{J}''}$  (consider Fig. 2, for an illustration). Finally, gluing in  $\Delta^{\mathcal{J}'}$  along  $j_3$  adds the missing simplices  $\Delta^{\mathcal{J}'}$  and  $\Delta^{\mathcal{J}_k'}$ . We have thus exposed  $A \hookrightarrow B$  as a composition of the pushouts of horn inclusions  $j_1$ ,  $j_2$  and  $j_3$ . Since  $|-|_s$  preserves pushouts, it suffices to show that  $\mathcal{X}$  has the horn filling property with respect to (the realizations of)  $j_1$ ,  $j_2$  and  $j_3$ . Since  $s \geq 2$ ,  $j_1$  is an inner horn inclusion, and furthermore since  $p_0 = p_k$  we have  $t_{\mathcal{J}_k} < t_{\mathcal{J}}$ , which is covered by the inductive assumption.  $j_2$  is an inner horn inclusion, in which the  $s$ -th entry is not minimal, which we have already covered above through Lemma 5.1.8. Finally,  $j_3$  is an inner horn inclusion with  $o_{\mathcal{J}'} < o_{\mathcal{J}}$ , and hence is also covered by the inductive assumption. This finishes the induction.  $\square$

## 5.2 Cofibrant stratified spaces and cellularly stratified spaces

Next, let us investigate the class of cofibrant objects in the non-refined semimodel categories, both categorical and diagrammatic.

**Proposition 5.2.1.** *Let  $\mathcal{X} \in \mathbf{Strat}$ . Then the following statements are equivalent:*

1.  $\mathcal{X}$  is a retract in  $\mathbf{Strat}_{P_{\mathcal{X}}}$  of a cellularly stratified space;
2.  $\mathcal{X}$  is cofibrant in all of the semimodel categories  $\mathbf{Strat}_{P_{\mathcal{X}}}^{\mathfrak{d}}$ ,  $\mathbf{Strat}_{P_{\mathcal{X}}}^{\mathfrak{c}}$ ,  $\mathbf{Strat}^{\mathfrak{d}, \mathfrak{p}}$ ,  $\mathbf{Strat}^{\mathfrak{c}, \mathfrak{p}}$ ;
3.  $\mathcal{X}$  is cofibrant in one of the semimodel categories  $\mathbf{Strat}_{P_{\mathcal{X}}}^{\mathfrak{d}}$ ,  $\mathbf{Strat}_{P_{\mathcal{X}}}^{\mathfrak{c}}$ ,  $\mathbf{Strat}^{\mathfrak{d}, \mathfrak{p}}$ ,  $\mathbf{Strat}^{\mathfrak{c}, \mathfrak{p}}$ .

*Proof.* Note that there is no need to differentiate between the diagrammatic and categorical scenarios, as the latter are obtained from the former via left Bousfield localization (Theorem 4.2.11), and hence both settings have the same cofibrant objects. That being a retract of a cellularly stratified space is equivalent to being cofibrant over  $P_{\mathcal{X}}$  is immediate by the cofibrant generators of Theorem 4.2.7. It remains to see that  $\mathcal{X}$  being cofibrant in  $\mathbf{Strat}^{\mathfrak{d},\mathfrak{p}}$  is equivalent to  $\mathcal{X}$  being cofibrant in  $\mathbf{Strat}_{P_{\mathcal{X}}}^{\mathfrak{d}}$ . This follows immediately from Proposition 4.2.8.  $\square$

**Definition 5.2.2.** We call a stratified topological space  $\mathcal{X}$  that satisfies any of the equivalent conditions of Lemma 5.1.1 **triangularly cofibrant**.

**Remark 5.2.3.** Clearly every stratified space that admits a triangulation compatible with the stratification (i.e., in particular is the realization of a stratified simplicial set) is triangularly cofibrant. In particular, this holds for piece-linear pseudo-manifolds (see, for example, [Ban07]) or locally compact stratified spaces which are definable (with definable stratification) in some o-minimal structure on the reals ([Dri98, Thm. 1.7]). Furthermore, by [Gor78], all Whitney stratified spaces are in this class.

However, one should not restrict oneself to stratified spaces that admit a cell structure. Topological manifolds, for example, are (to the best of our knowledge) not known to admit CW structures in dimension 4 (see [KS69] for all other dimensions). However, every topological manifold is a Euclidean neighborhood retract ([Han51]) and thus a retract of a CW complex. Supposing the existence of certain stratified cylinder neighborhoods, our theory also covers examples of stratified spaces with manifold strata (see Proposition 5.2.10). Let us begin by proving that stratified neighborhood retracts of triangularly cofibrant stratified spaces are triangularly cofibrant:

**Proposition 5.2.4.** *Let  $\mathcal{X} \in \mathbf{Strat}$  be triangularly cofibrant stratified space. Let  $A \subset \mathcal{X}$  be a closed subspace such that the following holds: there is a neighborhood  $U \subset \mathcal{X}$  of  $A$ , such that  $\mathcal{A} = (A, s_{\mathcal{X}}(U), s_{\mathcal{X}}|_A: A \rightarrow s_{\mathcal{X}}(U))$  is a stratum preserving retract of  $\mathcal{U} = (U, s_{\mathcal{X}}(U), s_{\mathcal{X}}|_U: U \rightarrow s_{\mathcal{X}}(U))$ . Then  $A$  is also triangularly cofibrant.*

*Proof.* We will make frequent use of results on stratified cell complexes developed in [Waa24b]. Let  $\mathcal{X}$  be a cellularly stratified space and  $i: \mathcal{X} \hookrightarrow \mathcal{Y}$ ,  $r: \mathcal{Y} \rightarrow \mathcal{X}$  such that  $r \circ i = 1_{\mathcal{X}}$ . We claim that there is a subcomplex  $\mathcal{B}$  of a cell structure on  $\mathcal{Y}$  that contains  $i(A)$  and is entirely contained in  $r^{-1}(U)$ . Then we can compose

$$\mathcal{B} \rightarrow \mathcal{U} \rightarrow \mathcal{A},$$

where  $\mathcal{B}$  is considered a stratified space over  $P_{\mathcal{U}}$ , to obtain a retraction of  $\mathcal{A} \hookrightarrow \mathcal{B}$ . Fix a stratified cell structure  $(\sigma_i: |\Delta^{\mathcal{J}_i}|_s \rightarrow \mathcal{Y})_{i \in I}$  on  $\mathcal{Y}$  (see [Waa24b, Subsec. 2.3], for a definition and basic properties). Every cell in  $\mathcal{Y}$  defined by a map  $\sigma_i$  is contained in a finite subcomplex of  $\mathcal{Y}$  ([Waa24b, Lem. 2.3.6]). We proceed to inductively construct a subdivision of  $\mathcal{Y}$  (in the sense of [Waa24b, Def. 3.2.1]), via induction over the minimal number of cells,  $n_i$ , required in a subcomplex that contains a cell  $\sigma_i$ . Denote by  $\mathcal{Y}^{(n-1)}$  the subcomplex constructed from cells  $\sigma_i$  with  $n_i = n$ . We construct a subdivision of the cell structure of  $\mathcal{Y}$ , denoted  $\mathcal{Y}'$ , through induction on  $n$ . For  $n = 1$ , we simply take the cell structure on  $\mathcal{Y}^{(0)}$  inherited from  $\mathcal{Y}$ . Now, suppose that we have already defined a cell structure  $\mathcal{Y}'^n$  on  $\mathcal{Y}^{(n)}$  for  $n > 1$ , and a subcomplex  $\mathcal{B}^n \subset \mathcal{Y}^{(n)}$  of this cell structure, fulfilling the following properties:

1.  $r^{-1}(A) \cap \mathcal{Y}^{(n)} \subset \mathcal{B}^n$  and  $\mathcal{B}^n$  is a  $\Delta_P$ -neighborhood of  $r^{-1}(A) \cap \mathcal{Y}^{(n)}$  (in the sense of [Waa24b, Def. 2.4.1]);
2.  $\mathcal{B}^n$  is contained in the interior of  $r^{-1}(U)$ .

Now, for any cell  $\sigma_i$  with  $n_i = n + 1$ , we can barycentrically subdivide  $|\Delta^{\mathcal{J}_i}|_s$  (sufficiently many times) so that for every  $\tau \subset |\Delta^{\mathcal{J}_i}|_s$  that is contained in a simplex  $\tau'$  intersecting  $(r \circ \sigma)^{-1}(A)$  we have the following.

1.  $\tau$  is entirely contained in the interior of  $(r \circ \sigma)^{-1}(U)$ ;
2.  $\tau \cap |\partial\Delta^{\mathcal{I}_i}|_s$  is entirely contained in  $\sigma^{-1}(\mathcal{B}^n)$ .

Indeed, the first property is possible, since  $(r \circ \sigma)^{-1}(A)$  is compact and contained in the interior of  $(r \circ \sigma)^{-1}(U)$  by (inductive) assumption. It follows that  $(r \circ \sigma)^{-1}(A)$  has a positive distance (for any choice of compatible metric on  $|\Delta^{\mathcal{I}_i}|_s$ , to the complement of the interior of  $(r \circ \sigma)^{-1}(U)$ , which implies that simplices of sufficiently small radius cannot intersect  $(r \circ \sigma)^{-1}(A) \cup \sigma^{-1}(\mathcal{B}^n)$  and the complement of the interior of  $(r \circ \sigma)^{-1}(U)$ . That the second property can be assumed is argued similarly. Having chosen such a subdivision for any cell  $\sigma_i$  with  $n_i = n + 2$ , we then obtain an induced cell structure  $\mathcal{Y}'^{n+1}$  on  $\mathcal{Y}^{(n+1)}$  given by the cells of  $\mathcal{Y}'^n$  and the cells corresponding to (open) simplices in the interior of the subdivisions of  $|\Delta^{\mathcal{I}_i}|_s$ , for  $n_i = n + 2$ . Finally, we let  $\mathcal{B}^{n+1}$  be the subcomplex given by adding to  $\mathcal{B}^n$  all such cells of  $\mathcal{Y}'^{n+1} \setminus \mathcal{Y}'^n$ ,  $\sigma_i$ , which correspond to an (open) simplex in the interior of some  $|\Delta^{\mathcal{I}_i}|_s$ , contained in a closed simplex intersecting  $(r \circ \sigma)^{-1}(A)$ . Note that this does indeed define a cell complex, since the intersection of the boundary of such a simplex with the boundary of  $|\Delta^{\mathcal{I}_i}|_s$  is assumed to be mapped to  $\mathcal{B}^n$ .  $\mathcal{B}^{n+1}$  defined in this fashion fulfills the inductive assumptions by construction and [Waa24b, Lem. 2.4.4]. Then, finally, set  $\mathcal{B} = \bigcup_{n \in \mathbb{N}} \mathcal{B}^n$ , with the induced cell structure.  $\square$

**Construction 5.2.5** (Stratified mapping cylinders). *Given  $f: \mathcal{X} \rightarrow \mathcal{Y}$  in **Strat**, the stratified mapping cylinder of  $f$ ,  $M_s(f)$ , is defined as the pushout*

$$\begin{array}{ccc} \mathcal{X} \times |\Delta^0|_s & \xrightarrow{1 \times i_0} & \mathcal{X} \times |\Delta^1|_s \\ \downarrow f & & \downarrow \\ \mathcal{Y} & \hookrightarrow & M_s(f). \end{array} \quad (24)$$

Under the (nonstratified) identification  $|\Delta^1|_s = [0, 1]$ , this equips the classical mapping cylinder of  $f$  with an alternative stratification over the lower right corner of the following pushout diagram of posets.

$$\begin{array}{ccc} P_{\mathcal{X}} & \xrightarrow{1 \times i_0} & P_{\mathcal{X}} \times [1] \\ \downarrow f & & \downarrow \\ P_{\mathcal{Y}} & \hookrightarrow & P_{\mathcal{Y}} \sqcup_{\leq f} P_{\mathcal{X}}. \end{array} \quad (25)$$

Here,  $P_{\mathcal{Y}} \sqcup_{\leq f} P_{\mathcal{X}}$  is explicitly given by equipping the disjoint union  $P_{\mathcal{Y}} \sqcup P_{\mathcal{X}}$ , with an additional relation  $p \leq q$ , for  $p \in P_{\mathcal{Y}}$  and  $q \in P_{\mathcal{X}}$ , whenever  $p \leq f(q)$ . In the special case where  $P_{\mathcal{Y}}$  is a poset with one element, this amounts to adjoining a minimal element to  $P_{\mathcal{X}}$ .

**Remark 5.2.6.** Note that  $M_s(f)$  is not a mapping cylinder of  $f$  with respect to any of the model structures on **sStrat**. Indeed, the inclusion  $\mathcal{Y} \hookrightarrow M_s(f)$  can only be a weak equivalence (in any of the model structures on **Strat**) if  $\mathcal{X}$  is empty. From the perspective of  $\infty$ -categories of exit paths, no point  $x \in \mathcal{X} \times |\Delta^1|_s \subset M_s(f)$  can lie in the essential image corresponding to the map  $\mathcal{Y} \rightarrow M_s(f)$ , as this would imply a non-stratified path from a point  $y \in \mathcal{Y}$  to  $x$ .

**Corollary 5.2.7.** *Suppose that  $\mathcal{L}, \mathcal{X}, \mathcal{Y}$  are triangularly cofibrant. Let  $f: \mathcal{L} \rightarrow \mathcal{Y}, g: \mathcal{L} \rightarrow \mathcal{X}$  be stratified maps. Then the stratified double mapping cylinder  $M_s(f) \cup_g \mathcal{X}$  is triangularly cofibrant.*

*Proof.*  $M_s(f) \cup_g \mathcal{X}$  fits into a pushout diagram of stratified spaces

$$\begin{array}{ccc} \mathcal{L} \times |\partial\Delta^1|_s = \mathcal{L} \sqcup \mathcal{L} & \hookrightarrow & \mathcal{L} \times |\Delta^1|_s \\ \downarrow f \sqcup g & & \downarrow \\ \mathcal{Y} \sqcup \mathcal{X} & \hookrightarrow & M_s(f) \cup_g \mathcal{X}. \end{array} \quad (26)$$

Note that we may without loss of generality assume that we are in the  $\Delta$ -generated scenario, as every triangularly cofibrant stratified space is  $\Delta$ -generated. By Corollary 4.3.3 the upper horizontal is a cofibration, from which it follows that the lower horizontal is a cofibration.  $\mathcal{Y} \sqcup \mathcal{X}$  is cofibrant by assumption, which together with the cofibrancy  $\mathcal{Y} \sqcup \mathcal{X} \hookrightarrow M_s(f) \cup_g \mathcal{X}$  implies the claim.  $\square$

**Definition 5.2.8.** Let  $U \subset \mathcal{X}$  and  $A \subset U$  be a closed subset contained in the interior of  $U$ . Denote  $\mathcal{A} = (s_{\mathcal{X}} | A: A \rightarrow s_{\mathcal{X}}(A))$  and  $\mathcal{U} = (s_{\mathcal{X}} | U: U \rightarrow s_{\mathcal{X}}(U))$ . We say that  $U$  is a **(closed) stratified mapping cylinder neighborhood of  $A$** , if the following holds:

There exists a stratified space  $\mathcal{L}$  together with a stratified map  $f: \mathcal{L} \rightarrow \mathcal{A}$ , as well as a stratum preserving homeomorphism  $\phi: M_s(f) \hookrightarrow \mathcal{U}$  under  $\mathcal{A}$ , such that  $\phi^{-1}(\partial U) = \mathcal{L} \times \{1\}$ .

**Definition 5.2.9.** Let  $P$  be a poset and  $p \in P$ . We call

$$\sup\{n \in \mathbb{N} \mid \exists p_0, \dots, p_n: p = p_0 < p_1 < \dots < p_n\}$$

the depth of  $p$  in  $P$ . We call the supremum over the depths of all  $p \in P$  the depth of  $P$ . Given  $k \in \mathbb{N}$  and  $\mathcal{X} \in \mathbf{Strat}$ , we denote by  $\mathcal{X}_k$  the stratified subspace given by restricting  $\mathcal{X}$  along

$$(P_{\mathcal{X}})_k := \{p \in P \mid p \text{ has depth } k\} \hookrightarrow P.$$

Similarly, we denote by  $\mathcal{X}_{\leq k}$  the stratified subspace of  $\mathcal{X}$  obtained by restricting  $\mathcal{X}$  along

$$(P_{\mathcal{X}})_{\leq k} := \{p \in P \mid p \text{ has depth } \leq k\} \hookrightarrow P.$$

**Proposition 5.2.10.** *Let  $\mathcal{X}$  be a stratified space of finite depth. Suppose that the following holds:*

1. *Each stratum of  $\mathcal{X}$  is cofibrant in the Quillen model structure, i.e. a retract of a non-stratified absolute cell complex.*
2. *For each  $k \in \mathbb{N}$ , there is a family of pairwise disjoint subsets of  $\mathcal{X}$ ,  $(U^p)_{p \in (P_{\mathcal{X}})_k}$ , indexed over strata of depth  $k$  such that, for each  $p$ ,  $U^p$  is a stratified mapping cylinder neighborhood of  $\mathcal{X}_p$ .*

*Then  $\mathcal{X}$  is triangularly cofibrant.*

*Proof.* For ease of notation, we denote  $P := P_{\mathcal{X}}$ . We proceed via induction over the depth of  $\mathcal{X}$ , denoted  $d$ . In the case  $d = 0$ ,  $\mathcal{X}$  is simply a disjoint union of trivially stratified spaces, each of which is cellularly stratified by assumption. Now, for the inductive step  $d$  to  $d + 1$ : By assumption, there are stratified maps  $f: \mathcal{L}^p \rightarrow \mathcal{X}_p$ , for each  $p \in P_d$ , together with an injective (on the space level) stratified map

$$\bigsqcup_{p \in (P_{d+1})} M(f^p) \hookrightarrow \mathcal{X},$$

which restricts to an open inclusion on the open cylinders. Setting,  $\mathcal{L} = \bigsqcup_{p \in P_{d+1}} \mathcal{L}^p$  and  $f = \bigsqcup_{p \in P_d} f^p$ , we obtain a stratified open inclusion

$$i: \bigsqcup_{p \in (P_{d+1})} \mathring{M}(f^p) \cong \mathring{M}_s(f) \hookrightarrow \mathcal{X},$$

which defines a neighborhood of  $\mathcal{X}_{d+1}$ . Note that  $i$  is not necessarily an inclusion on the poset level. Now, consider  $\mathring{M}_s(f)$  as reparametrized (over  $[0, 2)$ ), denoted  $\mathring{M}'_s(f)$ , such that we may consider  $M_s(f)$  (with the usual parametrization) as a closed stratified subspace of  $\mathring{M}'_s(f)$ . With this new notation, we have inclusion  $\mathring{M}'_s(f) \hookrightarrow M_s(f) \hookrightarrow \mathring{M}'_s(f) \hookrightarrow \mathcal{X}$ . Then  $\mathring{M}'_s(f) \subset \mathring{M}'_s(f)$  is

an open neighborhood of  $\mathcal{X}_{d+1}$  with boundary  $\mathcal{L} \times \{1\}$ . We obtain a commutative diagram of stratified maps

$$\begin{array}{ccc} \mathcal{L} \times \{1\} & \xrightarrow{g} & \mathcal{X} \setminus \mathring{M}_s(f) \\ \downarrow & & \downarrow \\ M_s(f) & \hookrightarrow & \mathcal{X}, \end{array} \quad (27)$$

where  $\mathcal{X} \setminus \mathring{M}_s(f)$  is stratified over  $P_{\underline{d}}$ . Denote by  $\tilde{\mathcal{X}}$  the pushout in **Strat**. Now, on the level of topological spaces, this diagram is clearly pushout. Therefore, the induced map  $g: \tilde{\mathcal{X}} \rightarrow \mathcal{X}$  is a homeomorphism on the underlying spaces. Consider the commutative diagram

$$\begin{array}{ccc} \tilde{\mathcal{X}} & \longrightarrow & P(g)!\tilde{\mathcal{X}} \\ & \searrow & \downarrow \\ & & \mathcal{X}. \end{array} \quad (28)$$

Since  $g$  is a homeomorphism on the underlying spaces, it follows that the right vertical is an isomorphism. The upper vertical is always a cofibration (Proposition 4.2.8), hence it follows that  $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$  is a cofibration. Therefore, it suffices to show that  $\tilde{\mathcal{X}}$  is cofibrant. For this, in turn, it suffices to show that the left-hand vertical is a cofibration and  $\mathcal{X} \setminus \mathring{M}_s(f)$  is cofibrant. The latter is a retract of  $\mathcal{X}_{\underline{d}}$ . Hence, cofibrancy follows by inductive assumption. For the left vertical, by Corollary 5.2.7, it suffices to see that  $\mathcal{L}_p$  and  $\mathcal{X}_p$  are cofibrant. The latter is cofibrant by assumption. By assumption  $\mathcal{L}^p$  embeds into  $\mathcal{X}$  with an open (stratified) neighborhood of the form  $\mathcal{L}^p \times (0, 1)$ . Clearly,  $\mathcal{L}^p$  is a retract of the latter, making  $\mathcal{L}^p$  a stratified neighborhood deformation retract of  $\mathcal{X}_{\underline{d}}$ , which is cofibrant by inductive assumption. Hence, the case of  $\mathcal{L}^p$  follows by Proposition 5.2.4.  $\square$

**Remark 5.2.11.** Every topological manifold is a Euclidean neighborhood retract ([Han51]) and since every open subset of Euclidean space can be triangulated, it follows that every topological manifold is cofibrant in the Quillen model structure. Consequently, it follows by Proposition 5.2.10 that every stratified space with manifold strata, each of which admits appropriate stratified mapping cylinder neighborhoods, is cofibrant. [AFT17, Prop. 8.2.3] asserts the existence of stratified mapping cylinder neighborhoods for conically smooth stratified spaces, which would make the latter triangularly cofibrant. More generally, for homotopically stratified spaces with manifold-strata there are obstructions to the existence of (pairwise) stratified mapping cylinder neighborhoods ([Qui88, Thm. 1.7]).

### 5.3 Refined stratified spaces

Let us take a more detailed look at the cofibrant objects in the semimodel categories **Strat**<sup>o</sup> and **Strat**<sup>c</sup>. In particular, our aim is to relate them to more classical properties of stratified spaces.

**Definition 5.3.1.** A stratified space  $\mathcal{X} \in \mathbf{Strat}$  is called **refined**, if the following holds:

- $\mathcal{X}$  is **surjectively stratified**, that is,  $s_{\mathcal{X}}: X \rightarrow P_{\mathcal{X}}$  is surjective.
- For each pair  $x, y \in \mathcal{X}$ , the relation  $s_{\mathcal{X}}(x) \leq s_{\mathcal{X}}(y)$  holds if and only if there is a finite sequence of stratified maps  $\gamma_i: |\Delta^{[1]}|_s \rightarrow \mathcal{X}$ ,  $i = 1, \dots, n$ , with  $\gamma_1(0) = x$ ,  $\gamma_n(1) = y$  and  $\gamma_i(1) = \gamma_{i+1}(0)$ , for  $i < n$ .

We can think of being refined as the poset  $P_{\mathcal{X}}$  being completely reflected in the stratified paths of  $P_{\mathcal{X}}$ . Next, note the following elementary properties about refinedness:



**Proposition 5.3.2.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathbf{Strat}$  and  $\mathcal{A} \in \mathbf{sStrat}$ . Then the following holds:*

1.  $\mathcal{X}$  is refined, if and only if  $\text{Sing}_s(\mathcal{X})$  is refined;
2. If  $f: \mathcal{X} \rightarrow \mathcal{Y}$  is a categorical weak equivalence, then  $\mathcal{X}$  is refined, if and only if  $\mathcal{Y}$  is refined;
3.  $\mathcal{A}$  is refined, if and only if  $|\mathcal{A}|_s$  is refined;
4. If  $\mathcal{A} \in \mathbf{sStrat}^{c,p}$  is fibrant, i.e.  $\mathcal{A}$  a quasi category and  $A \rightarrow P$  a conservative functor, then  $\mathcal{A}$  is refined if and only if  $\mathcal{A}$  is 0-connected, in the sense of [BGH18], as an abstract stratified homotopy type.

It follows that being refined is really purely a property of the homotopy type in  $\mathbf{Strat}^{0,p}$  defined by  $\mathcal{X}$ . The model categories  $\mathbf{Strat}^c$  and  $\mathbf{Strat}^{0,p}$  may now be interpreted as the respective right Bousfield localizations presenting the full sub- $\infty$ -categories of refined stratified spaces. Even more, we may essentially construct the left adjoints of this adjunction on the 1-categorical level.

**Definition 5.3.3.** A stratified space  $\mathcal{X} \in \mathbf{Strat}$  is said to carry the  $\Delta_s$  topology if one of the following equivalent conditions holds:

1.  $X$  has the final topology with respect to the set of stratified maps  $|\Delta^{[n]}|_s \rightarrow \mathcal{X}$ , for  $n \in \mathbb{N}$ .
2.  $X$  has the final topology with respect to the set of stratified maps  $|\Delta^{[1]}|_s \rightarrow \mathcal{X}$ .

*Proof.* Let us show that these conditions are equivalent. We need to see that every stratified simplex  $|\Delta^{[n]}|_s$ , for  $n \in \mathbb{N}$ , carries the final topology with respect to stratified maps with source  $|\Delta^{[1]}|_s$ . Note that it suffices to see that for any convergent sequence in  $|\Delta^{[n]}|_s$ , there exists a subsequence  $(x_i)_{i \in \mathbb{N}}$ , as well as a stratified map  $\gamma: [0, 1] \cong |\Delta^{[1]}|_s \rightarrow |\Delta^{[n]}|_s$ , with  $\gamma(\frac{1}{2^i}) = x_i$ , for  $i \in \mathbb{N}$ . Indeed, the topology on  $|\Delta^{[n]}|_s$  is entirely determined by convergent sequences, and the latter conditions mean that these are detected by  $|\Delta^{[1]}|_s$ . So suppose that we are given a convergent sequence  $(\hat{x}_i)_{i \in \mathbb{N}}$  in  $|\Delta^{[n]}|_s$ . By passing to a subsequence, we may assume that the sequence is contained in a single stratum  $j \in [n]$ . Then, the limit point lies in some stratum  $k \leq j$ . Now define  $\gamma: [0, 1] \cong |\Delta^{[1]}|_s \rightarrow |\Delta^{[n]}|_s$  by setting  $\gamma(\frac{1}{2^i}) := x_i$ , convexly interpolating between  $x_i$  and  $x_{i+1}$  and sending 0 to the limit point of  $x_i$ . This map is continuous. Furthermore, as the strata of  $|\Delta^{[n]}|_s$  are convex, it is also stratified.  $\square$

**Definition 5.3.4.** A stratified space  $\mathcal{X} \in \mathbf{Strat}$  is called **strongly refined**, if  $\mathcal{X}$  is refined and carries the  $\Delta_s$ -topology.

**Construction 5.3.5.** For a stratified space  $\mathcal{X} \in \mathbf{Strat}$ , its **refined poset**, denoted  $P_{\mathcal{X}^\tau}$ , is the poset generated by the following elements and relations:

- An element for each  $x \in \mathcal{X}$ .
- A relation  $x \leq y$ , whenever there is a sequence of stratified paths from  $x$  to  $y$ , as in Definition 5.3.1.

Equivalently,  $P_{\mathcal{X}^\tau}$  is given by the set of path components of strata of  $\mathcal{X}$ , together with a generating relation whenever there is an exit path from one component to another. Consider the (generally noncontinuous) map  $X \rightarrow P_{\mathcal{X}^\tau}$ , given by  $x \mapsto [x]$ . It provides a factorization of  $s_{\mathcal{X}}$  through the map of posets

$$\begin{aligned} P_{\mathcal{X}^\tau} &\rightarrow P_{\mathcal{X}} \\ [x] &\mapsto s_{\mathcal{X}}(x) \end{aligned}$$

as follows:

$$\begin{array}{ccc}
 X & \dashrightarrow & P_{\mathcal{X}^\tau} \\
 \searrow & & \downarrow \\
 & s_{\mathcal{X}} & P_{\mathcal{X}}
 \end{array} \tag{29}$$

Although  $X \rightarrow P_{\mathcal{X}^\tau}$  is not necessarily continuous, its precomposition with any stratified map  $|\Delta^{[1]}|_s \rightarrow \mathcal{X}$  is continuous, by construction. We denote by  $\mathcal{X}^\tau$  the stratified space given by  $X^\tau \rightarrow P_{\mathcal{X}^\tau}$ , where  $X^\tau$  has the same underlying set as  $X$  and is equipped with the final topology with respect to stratified maps  $|\Delta^{[1]}|_s \rightarrow \mathcal{X}$ . By construction,  $X^\tau \rightarrow P_{\mathcal{X}^\tau}$  is indeed continuous and  $\mathcal{X}^\tau$  is a strongly refined stratified space. This construction induces a simplicial functor from **Strat** into the full subcategory of strongly refined stratified spaces, called **the refinement functor**, which exposes the latter as a full coreflective subcategory of **Strat**. The counit of adjunction, given by the commutative squares

$$\begin{array}{ccc}
 X^\tau & \longrightarrow & X \\
 \downarrow s_{\mathcal{X}^\tau} & & \downarrow s_{\mathcal{X}} \\
 P_{\mathcal{X}^\tau} & \longrightarrow & P_{\mathcal{X}},
 \end{array} \tag{30}$$

is called the **refinement map**.

As a consequence of the existence of the refinement functor, which makes the inclusion of strongly refined stratified spaces a left adjoint, we obtain:

**Corollary 5.3.6.** *Every colimit of strongly refined stratified spaces in **Strat** is again strongly refined.*

The refinement construction of Construction 5.3.5 is really the topological analogue of [Waa24a, Def. 3.2.11].

**Proposition 5.3.7.** *(See [Waa24a, Subsec. 3.2]), for notation.) For any  $\mathcal{X} \in \mathbf{Strat}$ , applying  $\text{Sing}_s$  to the refinement map induces a natural transformation (dashed) that makes the diagram*

$$\begin{array}{ccc}
 (\text{Sing}_s \mathcal{X})^\tau & \dashrightarrow & \text{Sing}_s(\mathcal{X}^\tau) \\
 \searrow & & \downarrow \\
 & & \text{Sing}_s(\mathcal{X})
 \end{array} \tag{31}$$

commute. The dashed transformation is an isomorphism.

*Proof.*  $\text{Sing}_s \mathcal{X}^\tau$  is refined by Proposition 5.3.2. Hence, the dashed map is induced by the fact that (simplicial) refinement is right adjoint to the inclusion of refined stratified simplicial sets. That this map is an isomorphism on posets follows immediately by the construction of  $P_{\mathcal{X}^\tau}$  in Construction 5.3.5 and [Waa24a, Prop. 3.2.8]. Furthermore, since every stratified simplex  $|\Delta^{[n]}|_s$  is refined, note that the map is an isomorphism on simplicial sets, given by

$$(\text{Sing}_s \mathcal{X})^\tau[n] = (\text{Sing}_s \mathcal{X}) = \mathbf{Strat}(|\Delta^{[n]}|_s, \mathcal{X}) = \mathbf{Strat}(|\Delta^{[n]}|_s, \mathcal{X}^\tau) = \text{Sing}_s(\mathcal{X}^\tau)[n].$$

□

We may think of Proposition 5.3.7 as stating that  $\text{Sing}_s$  sends the topological refinement map to the simplicial refinement map. As an immediate corollary, using Recollection 3.2.4, we obtain the missing part of Theorem 3.3.1:

**Corollary 5.3.8.** *The simplicial semimodel categories  $\mathbf{Strat}^\circ$  and  $\mathbf{Strat}^c$  are obtained, respectively, from  $\mathbf{Strat}^{\circ,p}$  and  $\mathbf{Strat}^{c,p}$  by right Bousfield localizing at the refinement maps  $\mathcal{X}^r \rightarrow \mathcal{X}$ .*

In particular, we have the following description of cofibrant objects in  $\mathbf{Strat}^\circ$  and  $\mathbf{Strat}^c$ :

**Proposition 5.3.9.** *Let  $\mathcal{X} \in \mathbf{Strat}$ . Then the following statements are equivalent:*

1.  $\mathcal{X}$  is cofibrant in  $\mathbf{Strat}^\circ$ ;
2.  $\mathcal{X}$  is cofibrant in  $\mathbf{Strat}^c$ ;
3.  $\mathcal{X}$  is triangularly cofibrant and refined;
4.  $\mathcal{X}$  is a retract of a refined, cellularly stratified space.

*Proof.* The equivalence between the first two statements is immediate from both semimodel categories having the same generating cofibrations. Since every cellularly stratified space carries the  $\Delta_s$ -topology (by definition), it follows that a refined cellularly stratified space is strongly refined. Since the latter form a full coreflective subcategory, by Construction 5.3.5, any retract of strongly refined spaces is strongly refined. In particular, the fourth statement implies the third. To see that the third implies the second, note that cofibrant objects  $\mathcal{X}$  in  $\mathbf{Strat}^c$  are characterized by  $\emptyset \hookrightarrow \mathcal{X}$  having the left lifting property, with respect to all stratified maps  $f: \mathcal{X} \rightarrow \mathcal{Y}$ , which are acyclic fibrations in  $\mathbf{Strat}^c$ . By Proposition 5.3.7 and [Waa24a, Thm. 3.2.18], this, in turn, is equivalent to  $f^r$  being an acyclic fibration in  $\mathbf{Strat}^{c,p}$ . Since  $\mathcal{X}$  is assumed to be refined, a lifting diagram

$$\begin{array}{ccc} & & \mathcal{X} \\ & \nearrow \text{dashed} & \downarrow f \\ \mathcal{X} & \longrightarrow & \mathcal{Y} \end{array} \quad (32)$$

admits a solution, if and only if the induced diagram

$$\begin{array}{ccc} & & \mathcal{X}^r \\ & \nearrow \text{dashed} & \downarrow f^r \\ \mathcal{X} & \longrightarrow & \mathcal{Y}^r. \end{array} \quad (33)$$

admits a solution. The latter holds, as  $\mathcal{X}$  was assumed cofibrant in  $\mathbf{Strat}^{c,p}$ . Finally, to see that the first characterization implies the fourth, note that every cofibrant object in  $\mathbf{Strat}^c$  is a retract of an absolute cell complex with respect to the stratified boundary inclusions  $\{|\partial\Delta^{[n]} \hookrightarrow \Delta^{[n]}|_s \mid n \in \mathbb{N}\}$ . It follows by Proposition 5.3.2, stratified simplices carrying the  $\Delta_s$ -topology and Corollary 5.3.6, that every such absolute cell complex is a refined cellularly stratified space.  $\square$

## 5.4 Frontier conditions and refinement

It turns out that being refined is strongly related to the way the poset structure on the strata of a stratified space is classically constructed:

**Recollection 5.4.1.** Classically, stratifications often arise from the so-called **frontier condition** (see, for example, [Mat12]). Namely, one starts with a topological space  $X$  and a locally finite decomposition into nonempty locally closed pieces  $(X_i)_{i \in I}$ . One assumes that for any  $i \in I$ , the closure  $\overline{X_i}$  is given by the disjoint union  $(X_j)_{j \in J}$ , for some subset  $J \subset I$ . Then,  $I$  naturally carries the structure of a poset, setting  $i \leq j$ , whenever  $X_i \subset \overline{X_j}$ , and  $X \rightarrow I$  defines a stratification of  $X$ .

In our framework, such poset structures induced by closure relations can be constructed as follows.

**Construction 5.4.2.** Let  $\mathcal{X} \in \mathbf{Strat}$ , and denote

$$I = \{S \subset X \mid S \neq \emptyset \exists p \in P_{\mathcal{X}}: S = \mathcal{X}_p.\}$$

We consider  $I_{\mathcal{X}}$  as equipped with the structure of a poset, by equipping it with the relation generated by

$$S \leq S' \iff S \cap \overline{S'} \neq \emptyset.$$

It follows immediately by construction that mapping  $S$  to the unique  $p \in P_{\mathcal{X}}$ , for which  $S = \mathcal{X}_p$ , induces a map of posets

$$I_{\mathcal{X}} \rightarrow P_{\mathcal{X}}.$$

**Definition 5.4.3.** We say that  $\mathcal{X}$  is **weakly frontier stratified**, if the induced map  $I_{\mathcal{X}} \rightarrow P_{\mathcal{X}}$  is an isomorphism of posets. We say that  $\mathcal{X}$  is **frontier stratified**, if in addition to this it fulfills

$$\mathcal{X}_p \cap \overline{\mathcal{X}_q} \neq \emptyset \implies \mathcal{X}_p \subset \overline{\mathcal{X}_q},$$

for all  $p, q \in P_{\mathcal{X}}$ .

**Remark 5.4.4.** If one describes a stratified space  $\mathcal{X}$  of the data of a space  $X$ , equipped with decompositions  $(X_i)_{i \in I}$  and  $(X_j)_{j \in J}$  into nonempty pieces, then classically the morphisms which are considered between such spaces are given by continuous maps  $f: X \rightarrow Y$ , such that for each  $i \in I$  there exists a  $j \in J$  with  $f(X_i) \subset Y_j$  [Hug99]. Let us call such objects decomposition spaces, and such maps decomposed maps, and denote the corresponding category by  $\mathcal{D}$ . Homotopies in this setting are defined through the cylinder given by equipping  $X \times [0, 1]$  with the decomposition  $(X_i \times [0, 1])_{i \in I}$ . There is an obvious forgetful functor from the category  $\mathcal{D}: \mathbf{Strat} \rightarrow \mathbf{D}$ , given by equipping a stratified space into its decomposition into nonempty strata. Now, if  $\mathcal{X}$  is weakly frontier stratified, then it is not hard to see that for any  $\mathcal{Y} \in \mathbf{Strat}$  the induced map

$$\mathbf{Strat}(\mathcal{X}, \mathcal{Y}) \rightarrow \mathbf{D}(\mathcal{X}, \mathcal{Y})$$

is a bijection. Furthermore, the forgetful functor commutes with cylinders. It follows (that as long as one restricts to bifibrant objects (in  $\mathbf{Strat}^{\mathbf{D}, \mathbf{P}}$ ,  $\mathbf{Strat}^{\mathbf{C}, \mathbf{P}}$ ) which are surjectively stratified and fulfill the weak frontier condition, then the resulting homotopy theory agrees with the classical homotopy theory of stratified spaces as studied in [Qui88; Hug99; Mil13]).

It turns out that the triangularly cofibrant objects that have path-connected strata and are weakly frontier stratified are precisely the cofibrant objects in the refined setting (see Proposition 5.4.5 below).

Note that for strongly frontier stratified spaces the generating relations of Construction 5.4.2 already define a partial order. This leads to the following alternative characterization of refinedness, for specific stratified spaces.

**Proposition 5.4.5.** Let  $\mathcal{X} \in \mathbf{Strat}$ , and consider the following conditions:

1.  $\mathcal{X}$  is refined.
2.  $\mathcal{X}$  has path-connected strata and is frontier stratified.
3.  $\mathcal{X}$  has path-connected strata and is weakly frontier stratified.

The first and the second property imply the third. Furthermore, if  $\mathcal{X}$  is triangularly cofibrant, then the third property implies the first. Finally, if  $\mathcal{X}$  is categorically fibrant, then the first property implies the third. In particular, for bifibrant stratified spaces in  $\mathbf{Strat}^c$  all three properties are equivalent.

*Proof.* The first two implications are obvious, using the fact that for any continuous map  $f: X \rightarrow Y$ ,  $x \in \overline{S}$  implies  $f(x) \in \overline{f(S)}$ . Suppose the third condition holds and  $\mathcal{X}$  is triangularly cofibrant. To see that  $\mathcal{X}$  is refined, we need to expose for any  $x, y \in \mathcal{X}$  with  $p := s_{\mathcal{X}}(x) < s_{\mathcal{X}}(y) := q$  a sequence of stratified paths  $\gamma_i: |\partial\Delta^{[1]}|_s \rightarrow \mathcal{X}$  from  $x$  to  $y$ , as in 5.3.1. Since  $\mathcal{X}$  is weakly frontier stratified, it suffices to show that for any pair  $p, q \in P_{\mathcal{X}}$  and  $x \in \mathcal{X}_p, y \in \mathcal{X}_q$  with

$$\mathcal{X}_p \cap \overline{\mathcal{X}_q} \neq \emptyset,$$

there is a stratified path from  $x$  to  $y$ . Furthermore, since strata are assumed to be path-connected, it suffices to construct a path from any element in  $\mathcal{X}_p$  to any element in  $\mathcal{X}_q$ . Now, let  $\mathcal{X} \xrightarrow{i} \mathcal{Y} \xrightarrow{r} \mathcal{X}$  expose  $\mathcal{X}$  as a retract of a cellularly stratified space  $\mathcal{Y}$  (Proposition 5.2.1). Any nonempty closure intersection

$$\mathcal{X}_p \cap \overline{\mathcal{X}_q} \neq \emptyset$$

in  $\mathcal{X}$  implies a non-empty closure intersection

$$\mathcal{Y}_{i(p)} \cap \overline{\mathcal{Y}_{i(q)}} \neq \emptyset.$$

Conversely, every sequence of stratified paths in  $\mathcal{Y}$ , starting and ending in  $\mathcal{X}$ , descends to a sequence of stratified paths in  $\mathcal{X}$  with the same starting and end points. Hence, it suffices to show that a closure relation

$$\mathcal{Y}_{i(p)} \cap \overline{\mathcal{Y}_{i(q)}} \neq \emptyset,$$

implies the existence of a concatenable sequence of stratified path starting in  $\mathcal{Y}_{i(p)}$  and ending in  $\mathcal{Y}_{i(q)}$ . Choose a cell structure for  $\mathcal{Y}$  (see [Waa24b, Def.2.3.1]), with open cells  $e_i$ ,  $i \in I$ . Note that each open cell is entirely contained in only one stratum. We write  $s_{\mathcal{Y}}(e_i) \in P_{\mathcal{Y}}$  to denote the latter. Furthermore, note that if  $e_i$  intersects the closure of  $e_j$ , then  $s_{\mathcal{Y}}(e_i) \leq s_{\mathcal{Y}}(e_j)$ . Suppose that  $x \in \mathcal{Y}_{i(p)} \cap \overline{\mathcal{Y}_{i(q)}}$ . In particular,  $x$  is contained in the minimal subcomplex of  $\mathcal{Y}$  that contains  $\mathcal{Y}_{i(q)}$ . This implies that there is a sequence of cells  $e_0, e_1, e_2, \dots, e_n$ , such that  $x \in e_0$  and  $e_n \subset \mathcal{Y}_q$ , and for each  $i \in [n-1]$ ,  $e_i$  intersects the closure of  $e_{i+1}$ . For  $i \in [n-1]$ , let  $x_i \in e_i \cap \overline{e_{i+1}}$ . Since  $\overline{e_{i+1}}$  is the quotient of a stratified simplex  $|\Delta^{\mathcal{J}}|_s$  over  $P_{\mathcal{X}}$ , there is a stratified path  $\gamma'_i: |\Delta^{[1]}|_s \rightarrow \mathcal{X}$ , starting in  $x_i$ , immediately entering and staying in  $e_{i+1}$ , and ending in  $x_{i+1}$ .

The converse implication is immediate. Finally, assume that  $\mathcal{X}$  is categorically fibrant and refined (and is thus weakly frontier stratified). Then whenever  $s_{\mathcal{X}}(x) \leq s_{\mathcal{X}}(y)$  any concatenable sequence of stratified paths from  $x$  to  $y$  induces a stratified path,  $\gamma: |\Delta^{[1]}|_s \rightarrow \mathcal{X}$  from  $x$  to  $y$ , which implies

$$x = \gamma(0) \in \overline{\gamma(0, 1]} \subset \overline{\mathcal{X}_{s_{\mathcal{X}}(y)}}.$$

In particular, it follows that

$$\mathcal{X}_p \cap \overline{\mathcal{X}_q} \neq \emptyset \implies \mathcal{X}_p \subset \overline{\mathcal{X}_q}.$$

□

**Remark 5.4.6.** Remark 5.4.4 also holds in a self-enriched sense, providing an answer to Question Q(6): In [Hug99] the author studied stratified notions of mapping space, given by equipping the set of decomposition maps  $\mathcal{D}(\mathcal{X}, \mathcal{Y})$  with the compact open topology, and the decomposition over  $P_{\mathcal{Y}}^{P_{\mathcal{X}}}$  induced by mapping a stratified space to its underlying map of posets. If we work in the setting where  $\mathbf{Top}$  is the category of  $\Delta$ -generated spaces or compactly generated spaces, and

thus replace the compact open topology with its respective Kelleyfication, then this construction defines the internal hom in the category of decomposition spaes. There is an obvious map of decomposition spaces

$$\mathcal{D}(\mathcal{Y}^{\mathcal{X}}) \rightarrow \mathcal{D}(\mathcal{Y})^{\mathcal{D}(\mathcal{X})},$$

given on the set level by the map  $\mathbf{Strat}(\mathcal{X}, \mathcal{Y}) \rightarrow \mathbf{D}(\mathcal{X}, \mathcal{Y})$ . Therefore, whenever  $\mathcal{X}$  is refined, the comparison map of stratified mapping spaces above is bijective. Furthermore, by construction of  $\mathcal{Y}^{\mathcal{X}}$  and [May16, Cor. 2.2.11], it is a decomposition preserving homeomorphism, whenever  $P_{\mathcal{X}}$  and  $P_{\mathcal{Y}}$  are finite. In other cases, it is the map that refines the topology in  $\mathcal{D}(\mathcal{Y})^{\mathcal{D}(\mathcal{X})}$  such that the induced map

$$\mathcal{D}(\mathcal{Y})^{\mathcal{D}(\mathcal{X})} \rightarrow P_{\mathcal{Y}}^{P_{\mathcal{X}}}$$

is continuous.

**Remark 5.4.7.** In [Hug99], Hughes studied the space of stratified paths in a stratified space  $\mathcal{X}$  with finitely many strata, which start in a closed union of strata  $\mathcal{A} \subset \mathcal{X}$ , which we denote by  $\text{Path}_{nsp}(\mathcal{X}, \mathcal{A})$ . One of the main results of [Hug99] is that when  $\mathcal{X}$  is a homotopically stratified space, then the starting point evaluation map  $\text{Path}_{nsp}(\mathcal{X}, \mathcal{A}) \rightarrow \mathcal{A}$  lifts stratified homotopies. Using the Cartesian closedness of the semimodel structures on  $\mathbf{Strat}$  we may recover a version of this result in  $\mathbf{Strat}$ : Given a categorically (respectively, diagrammatically) fibrant stratified space  $\mathcal{X} \in \mathbf{Strat}$ , the starting point evaluation map  $\mathcal{X}^{|\Delta^{[1]}|_s} \rightarrow \mathcal{X}$  is a stratified fibration. Now, for any subspace  $\mathcal{A} \subset \mathcal{X}$ , equipped with the induced stratification, we may consider the pullback diagram

$$\begin{array}{ccc} \mathcal{X}^{|\Delta^{[1]}|_s} \times_{\text{ev}_0} \mathcal{A} & \longrightarrow & \mathcal{X}^{|\Delta^{[1]}|_s} \\ \downarrow & & \downarrow \text{ev}_0 \\ \mathcal{A} & \longrightarrow & \mathcal{X}. \end{array} \quad (34)$$

It follows that the right vertical is a categorical (or, respectively, diagrammatic) fibration. In particular, it has the lifting property with respect to stratified homotopies with cofibrant source. Since  $|\Delta^{[1]}|_s$  is clearly strongly frontier stratified, the natural comparison map in (5.4.6) induces a natural continuous bijection

$$\mathcal{D}(\mathcal{X}^{|\Delta^{[1]}|_s}) \rightarrow \text{Path}_{nsp}(\mathcal{X}, \mathcal{A}),$$

which refines the topology on  $\text{Path}_{nsp}(\mathcal{X}, \mathcal{A})$  in order to make the later stratified (and compactly or  $\Delta$ -generated). Hence, one obtains a Serre (as opposed to Hurewicz-style homotopy theory) version of Hughes's result, replacing homotopically stratified spaces with the weaker condition of diagrammatic fibrancy and not requiring that  $\mathcal{A}$  is a closed union of strata, but only obtaining the homotopy lifting property with respect to triangularly cofibrant stratified spaces.

The characterization of refinedness in Proposition 5.4.5 allows us to represent all homotopy types in  $\mathbf{Strat}^f$  through the following particularly convenient stratified spaces.

**Definition 5.4.8.** A stratified space  $\mathcal{X} \in \mathbf{Strat}$  is called **CFF stratified** (C for cellular, first F for frontier, second F for fibrant), if it fulfills the following conditions:

1.  $\mathcal{X}$  is cellularly stratified.
2.  $\mathcal{X}$  has nonempty, connected strata and is frontier stratified.
3.  $\mathcal{X}$  has the horn filling property with respect to inner stratified horn inclusions  $|\Lambda_k^{[n]} \hookrightarrow \Delta^{[n]}|_s$ ,  $0 < k < n$ .

We denote the full simplicial subcategory of  $\mathbf{Strat}$ , given by CFF stratified spaces by  $\mathbf{CFF}$ .

CFF stratified spaces can be seen as an analogue to CW complexes in the classical scenario. However, note that no assumptions are made that the attaching maps of cells only maps to cells of certain dimensions. This is necessary for the small object argument to be able to produce CFF stratified spaces.

**Example 5.4.9.** Every stratified space that admits a PL structure that is compatible with triangulation is cellularly stratified. Furthermore, in increasing order of generality, every Whitney stratified space, Thom-Mather stratified space, pseudomanifold or conically stratified space (in the sense of [Lur17, A.5]) has the inner horn filling property, by [Lur17, Thm. A.6.5]. Hence, it follows that if we equip such spaces with the refined stratification (induced by the frontier condition, and by taking path components of strata) and additionally assume a piecewise linear structure, then they provide examples of CFF stratified spaces. Note also that for Thom-Mather (and hence for Whitney stratified spaces) a piecewise linear structure (compatible with the stratification) always exists ([Gor78]).

We may characterize the bifibrant objects of  $\mathbf{sStrat}^c$ . Using the small object argument on the generating classes in Theorem 4.2.10, together with Propositions 5.3.9 and 5.4.5, we obtain the following corollary.

**Corollary 5.4.10.** *A stratified space  $\mathcal{X} \in \mathbf{Strat}^c$  is bifibrant if and only if it is a retract of a CFF stratified space. Furthermore, every bifibrant stratified space is stratified homotopy equivalent to a CFF stratified space. Every stratified space is categorically equivalent to a CFF stratified space.*

In particular, it follows that the homotopy theory defined by  $\mathbf{Strat}^c$  may equivalently be interpreted in terms of CFF stratified spaces:

**Corollary 5.4.11.** *Denote by  $H_s$  the class of stratified homotopy equivalences between CFF stratified spaces. The inclusion  $\mathbf{CFF} \rightarrow \mathbf{Strat}^c$  induces an equivalence of  $\infty$ -categories*

$$\mathbf{CFF}[H_s^{-1}] \xrightarrow{\cong} \mathbf{Strat}^c.$$

Combining this result with Corollary 4.4.5, we obtain the following version of the stratified homotopy hypothesis:

**Corollary 5.4.12.** *Luriers exit-path construction induces an equivalence of  $\infty$ -categories*

$$\mathbf{CFF}[H_s^{-1}] \xrightarrow{\cong} \mathbf{Lay}_\infty$$

*between CFF stratified spaces localized at stratified homotopy equivalences and layered  $\infty$ -categories.*

## Acknowledgements

The author is supported by a PhD-stipend of the Landesgraduiertenförderung Baden-Württemberg. Furthermore, he wants to thank Sylvain Douteau and Peter Haine for the insightful discussions concerning their work.

## References

- [AFR19] David Ayala, John Francis, and Nick Rozenblyum. “A stratified homotopy hypothesis”. *J. Eur. Math. Soc. (JEMS)* 21.4 (2019), pp. 1071–1178. URL: <https://doi-org.ezp.sub.su.se/10.4171/JEMS/856>.

- [AFT17] David Ayala, John Francis, and Hiro Lee Tanaka. “Local structures on stratified spaces”. *Advances in Mathematics* 307 (2017), pp. 903–1028. URL: <https://www.sciencedirect.com/science/article/pii/S0001870816316097>.
- [Are46] Richard Arens. “Topologies for Homeomorphism Groups”. *American Journal of Mathematics* 68.4 (1946), pp. 593–610. URL: <http://www.jstor.org/stable/2371787> (visited on 07/08/2022).
- [Ban07] Markus Banagl. *Topological invariants of stratified spaces*. Springer Science & Business Media, 2007.
- [Bar10] Clark Barwick. “On left and right model categories and left and right Bousfield localizations”. *Homology Homotopy Appl.* 12.2 (2010), pp. 245–320. URL: <http://projecteuclid.org/euclid.hha/1296223884>.
- [Ber07a] Julia E. Bergner. “A model category structure on the category of simplicial categories”. *Trans. Amer. Math. Soc.* 359.5 (2007), pp. 2043–2058. URL: <https://doi.org/10.1090/S0002-9947-06-03987-0>.
- [Ber07b] Julia E. Bergner. “Three models for the homotopy theory of homotopy theories”. *Topology* 46.4 (2007), pp. 397–436. URL: <https://doi.org/10.1016/j.top.2007.03.002>.
- [BGH18] Clark Barwick, Saul Glasman, and Peter Haine. “Exodromy”. *arXiv e-prints* (July 2018). arXiv: 1807.03281 [math.AT].
- [BK12a] Clark Barwick and Daniel M. Kan. “A characterization of simplicial localization functors and a discussion of DK equivalences”. *Indag. Math. (N.S.)* 23.1-2 (2012), pp. 69–79. URL: <https://doi.org/10.1016/j.indag.2011.10.001>.
- [BK12b] Clark Barwick and Daniel M. Kan. “Relative categories: another model for the homotopy theory of homotopy theories”. *Indag. Math. (N.S.)* 23.1-2 (2012), pp. 42–68. URL: <https://doi.org/10.1016/j.indag.2011.10.002>.
- [Cis19] Denis-Charles Cisinski. *Higher categories and homotopical algebra*. Vol. 180. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2019, pp. xviii+430. URL: <https://doi.org/10.1017/9781108588737>.
- [CM20] Pierre Cagne and Paul-André Melliès. “On bifibrations of model categories”. *Adv. Math.* 370 (2020), pp. 107205, 43. URL: <https://doi-org.ezp.sub.su.se/10.1016/j.aim.2020.107205>.
- [DK70] B. J. Day and G. M. Kelly. “On topological quotient maps preserved by pullbacks or products”. *Proc. Cambridge Philos. Soc.* 67 (1970), pp. 553–558. URL: <https://doi.org/10.1017/s0305004100045850>.
- [DK80a] William G. Dwyer and Daniel M. Kan. “Calculating simplicial localizations”. *J. Pure Appl. Algebra* 18.1 (1980), pp. 17–35. URL: [https://doi.org/10.1016/0022-4049\(80\)90113-9](https://doi.org/10.1016/0022-4049(80)90113-9).
- [DK80b] William G. Dwyer and Daniel M. Kan. “Function complexes in homotopical algebra”. *Topology* 19.4 (1980), pp. 427–440. URL: [https://doi.org/10.1016/0040-9383\(80\)90025-7](https://doi.org/10.1016/0040-9383(80)90025-7).
- [DK80c] William G. Dwyer and Daniel M. Kan. “Simplicial localizations of categories”. *J. Pure Appl. Algebra* 17.3 (1980), pp. 267–284. URL: [https://doi.org/10.1016/0022-4049\(80\)90049-3](https://doi.org/10.1016/0022-4049(80)90049-3).
- [DK87] William G. Dwyer and Daniel M. Kan. “Equivalences between homotopy theories of diagrams”. *Algebraic topology and algebraic K-theory (Princeton, N.J., 1983)*. Vol. 113. Ann. of Math. Stud. Princeton Univ. Press, Princeton, NJ, 1987, pp. 180–205.



- [Dou21a] Sylvain Douteau. “A simplicial approach to stratified homotopy theory”. *Trans. Amer. Math. Soc.* 374.2 (2021), pp. 955–1006. URL: <https://doi.org/10.1090/tran/8264>.
- [Dou21b] Sylvain Douteau. “A stratified Kan-Quillen equivalence”. *arXiv e-prints* (Feb. 2021). arXiv: 2102.04876 [math.AT].
- [Dou21c] Sylvain Douteau. “Homotopy theory of stratified spaces”. *Algebr. Geom. Topol.* 21.1 (2021), pp. 507–541. URL: <https://doi.org/10.2140/agt.2021.21.507>.
- [Dri98] Lou van den Dries. *Tame topology and o-minimal structures*. Vol. 248. London Mathematical Society Lecture Note Series. Cambridge, UK: Cambridge University Press, 1998.
- [Dug03] Daniel Dugger. “Notes on delta-generated spaces” (2003).
- [DW22] Sylvain Douteau and Lukas Waas. “From homotopy links to stratified homotopy theories”. *arXiv e-prints* (2022). arXiv: 2112.02394 [math.AT].
- [Fre09] Benoit Fresse. *Modules over operads and functors*. Vol. 1967. Lecture Notes in Mathematics. Springer-Verlag, Berlin, 2009, pp. x+308. URL: <https://doi.org/10.1007/978-3-540-89056-0>.
- [Gau21] Philippe Gaucher. “Left properness of flows”. *Theory Appl. Categ.* 37 (2021), Paper No. 19, 562–612. URL: <https://doi.org/10.1002/num.22542>.
- [GH05] Paul G. Goerss and Michael J. Hopkins. “Moduli problems for structured ring spectra”. *preprint* 1086 (2005).
- [Gor78] R. Mark Goresky. “Triangulation of stratified objects”. *Proc. Amer. Math. Soc.* 72.1 (1978), pp. 193–200. URL: <https://doi.org/10.2307/2042563>.
- [Hai23] Peter J. Haine. “On the homotopy theory of stratified spaces”. *arXiv e-prints* (2023). arXiv: 1811.01119 [math.AT].
- [Han51] Olof Hanner. “Some theorems on absolute neighborhood retracts”. *Ark. Mat.* 1 (1951), pp. 389–408. URL: <https://doi.org/10.1007/BF02591376>.
- [Hen] Andre Henriques. “A model category for stratified spaces”. Note available at [http://andreghenriques.com/PDF/Model\\_Cat\\_Stratified\\_spaces.pdf](http://andreghenriques.com/PDF/Model_Cat_Stratified_spaces.pdf).
- [Hin16] Vladimir Hinich. “Dwyer-Kan localization revisited”. *Homology Homotopy Appl.* 18.1 (2016), pp. 27–48. URL: <https://doi.org/10.4310/HHA.2016.v18.n1.a3>.
- [Hir09] Philip S. Hirschhorn. *Model Categories and Their Localizations*. Mathematical surveys and monographs. American Mathematical Society, 2009. URL: <https://books.google.de/books?id=jKPN3818LwcC>.
- [Hug99] Bruce Hughes. “Stratified path spaces and fibrations”. *Proceedings of the Royal Society of Edinburgh: Section A Mathematics* 129 (Jan. 1999).
- [JT07] André Joyal and Myles Tierney. “Quasi-categories vs Segal spaces”. *Categories in algebra, geometry and mathematical physics*. Vol. 431. Contemp. Math. Amer. Math. Soc., Providence, RI, 2007, pp. 277–326. URL: <https://doi.org/10.1090/conm/431/08278>.
- [JT08] André Joyal and Myles Tierney. “Notes on simplicial homotopy theory”. *Preprint* (2008).
- [KS69] R. C. Kirby and L. C. Siebenmann. “On the triangulation of manifolds and the Hauptvermutung”. *Bull. Amer. Math. Soc.* 75 (1969), pp. 742–749. URL: <https://doi.org/10.1090/S0002-9904-1969-12271-8>.

- [Lan21] Markus Land. *Introduction to infinity-categories*. Compact Textbooks in Mathematics. Birkhäuser/Springer, Cham, 2021, pp. ix+296. URL: <https://doi.org/10.1007/978-3-030-61524-6>.
- [Lur09] Jacob Lurie. *Higher topos theory*. Vol. 170. Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 2009, pp. xviii+925. URL: <https://doi.org/10.1515/9781400830558>.
- [Lur17] Jacob Lurie. *Higher Algebra*. <http://www.math.harvard.edu/~lurie/papers/HA.pdf>. 2017.
- [Mat12] John Mather. “Notes on topological stability”. *Bull. Amer. Math. Soc. (N.S.)* 49.4 (2012), pp. 475–506.
- [Mat73] John N. Mather. “Stratifications and mappings”. *Dynamical Systems (Proc. Sympos. Univ. Bahia, Salvador, 1971)*. Ed. by M.M. PEIXOTO. New York, NY, USA: Academic Press, 1973, pp. 195–232.
- [May16] J Peter May. “Finite spaces and larger contexts”. *Unpublished book* (2016).
- [Mil09] David Miller. “Popath and Holinks”. *arXiv: Algebraic Topology* (2009). URL: <https://api.semanticscholar.org/CorpusID:2948510>.
- [Mil13] David A. Miller. “Strongly stratified homotopy theory”. *Trans. Amer. Math. Soc.* 365.9 (2013), pp. 4933–4962. URL: <https://doi.org/10.1090/S0002-9947-2013-05795-9>.
- [Mil59] John W. Milnor. “On spaces having the homotopy type of a CW-complex”. *Transactions of the American Mathematical Society* 90 (1959), pp. 272–280. URL: <https://api.semanticscholar.org/CorpusID:123048606>.
- [Nan19] Stephen J. Nand-Lal. “A Simplicial Approach to Stratified Homotopy Theory”. available at <https://livrepository.liverpool.ac.uk/3036209/>. PhD thesis. University of Liverpool, 2019.
- [nLa23] nLab authors. *transferred model structure*. <https://ncatlab.org/nlab/show/transferred+model+structure>. Revision 48. June 2023.
- [NV23] Guglielmo Nocera and Marco Volpe. “Whitney stratifications are conically smooth”. *Selecta Mathematica* 29.5 (Sept. 15, 2023), p. 68. URL: <https://doi.org/10.1007/s00029-023-00877-4>.
- [Qui67] Daniel G. Quillen. *Homotopical algebra*. Lecture Notes in Mathematics, No. 43. Springer-Verlag, Berlin-New York, 1967, iv+156 pp. (not consecutively paged).
- [Qui88] Frank Quinn. “Homotopically stratified sets”. *J. Amer. Math. Soc.* 1.2 (1988), pp. 441–499. URL: <https://doi.org/10.2307/1990924>.
- [Rez10] Charles Rezk. “A cartesian presentation of weak  $n$ -categories”. *Geometry & Topology* 14.1 (2010), pp. 521–571. URL: <https://doi.org/10.2140/gt.2010.14.521>.
- [Rez17] Charles Rezk. “COMPACTLY GENERATED SPACES”. 2017. URL: [https://ncatlab.org/nlab/files/Rezk\\_CompactlyGeneratedSpaces.pdf](https://ncatlab.org/nlab/files/Rezk_CompactlyGeneratedSpaces.pdf).
- [Rie14] Emily Riehl. *Categorical homotopy theory*. Vol. 24. New Mathematical Monographs. Cambridge University Press, Cambridge, 2014, pp. xviii+352. URL: <https://doi.org/10.1017/CB09781107261457>.
- [Str72] Arne Strøm. “The homotopy category is a homotopy category”. *Archiv der Mathematik* 23.1 (Dec. 1, 1972), pp. 435–441. URL: <https://doi.org/10.1007/BF01304912>.
- [Tho69] René Thom. “Ensembles et morphismes stratifiés”. *Bull. Amer. Math. Soc.* 75 (1969), pp. 240–284.

- [Tre09] David Treumann. “Exit paths and constructible stacks”. *Compositio Mathematica* 145.6 (2009), pp. 1504–1532.
- [Waa24a] Lukas Waas. “Combinatorial models for stratified homotopy theory” (2024). arXiv: 2403.06280 [math.AT].
- [Waa24b] Lukas Waas. “On the homotopy links of stratified cell complexes” (2024). arXiv: 2403.06272 [math.AT].
- [WB23] David White and Michael Batanin. “Left Bousfield localization without left properness”. *arXiv e-prints* (2023). arXiv: 2001.03764 [math.AT].
- [Wei94] Shmuel Weinberger. *The topological classification of stratified spaces*. Chicago, 1994.
- [Whi65] Hassler Whitney. “Tangents to an analytic variety”. *Ann. of Math. (2)* 81 (1965), pp. 496–549. URL: <https://doi.org/10.2307/1970400>.
- [Woo09] Jon Woolf. “The fundamental category of a stratified space”. *J. Homotopy Relat. Struct.* 4.1 (2009), pp. 359–387.

## A Refined stratified spaces, and Nand-Lal’s homotopy theory

In this section, we relate the homotopy theory  $\mathbf{Strat}^c$  to the work of [Nan19]:

**Recollection A.0.1.** [Nan19] To avoid dealing with empty strata, [Nan19] introduced the notion of a **surjectively stratified space**, defined as follows: Denote by  $\mathbf{Strat}_s$  the full subcategory of  $\mathbf{Strat}$  given by such stratified spaces  $\mathcal{X}$ , for which the stratification  $s_{\mathcal{X}}: \mathcal{X} \rightarrow P_{\mathcal{X}}$  is surjective. In other words, such stratified spaces whose stratification poset contains no redundant elements (but possibly redundant relations). Such a stratified space is called **surjectively stratified**. In [Nan19], Nand-Lal constructs a homotopy theory for  $\mathbf{Strat}_s$  via transfer along the composition

$$\mathbf{Strat}_s \hookrightarrow \mathbf{Strat} \xrightarrow{\text{Sing}_s} \mathbf{sStrat} \xrightarrow{\mathcal{F}} \mathbf{sSet},$$

using the Joyal model structure on  $\mathbf{sSet}$ . In other words, a weak equivalence in the resulting homotopy theory is a map of stratified spaces  $f: \mathcal{X} \rightarrow \mathcal{T}$ , for which the underlying simplicial map of the stratified simplicial map  $\text{Sing}_s(f)$  is a categorical equivalence. It follows that this class of weak equivalences is precisely the class of categorical equivalences with source and target in  $\mathbf{Strat}_s$ . We denote by  $\mathbf{Strat}_s^c$  the relative category given by  $\mathbf{Strat}_s$ , together with such weak equivalences.

Let us now relate  $\mathbf{Strat}_s^c$  to  $\mathbf{Strat}^c$ .

**Proposition A.0.2.** *The inclusion  $\mathbf{Strat}_s \hookrightarrow \mathbf{Strat}$  induces a homotopy equivalence of relative categories*

$$\mathbf{Strat}_s^c \xrightarrow{\cong} \mathbf{Strat}^c,$$

and hence an equivalence of the corresponding  $\infty$ -categories.

*Proof.* The functor  $(-)^{\text{r}}: \mathbf{Strat} \rightarrow \mathbf{Strat}$  has image in  $\mathbf{Strat}_s$ . By Proposition 5.3.7, there is a natural isomorphism  $\text{Sing}_s \circ (-)^{\text{r}} \cong (-)^{\text{r}} \circ \text{Sing}_s$ , which shows that  $(-)^{\text{r}}$  preserves refined categorical equivalences. Furthermore, since (again by Proposition 5.3.7) the refinement map is a categorical equivalence, it follows that  $(-)^{\text{r}}$  defines a homotopy inverse to the inclusion of relative categories  $\mathbf{Strat}_s^c \hookrightarrow \mathbf{Strat}^c$ . One homotopy is given by the refinement map and one by the identity.  $\square$

## B An elementary extension lemma

In this section, we give a proof of Lemma 5.1.7. Before we give a proof, let us introduce some notation and quickly illustrate where metrizable comes into play. It is not hard to see that there is a continuous map  $i: D^{n+1} \times [0, 1] \rightarrow D^{n+1} \times [0, 1] \cup S^n \times [0, 1]$ , which is the identity on  $D^{n+1} \times \{0\} \cup S^n \times [0, 1]$  and maps  $\mathring{D}^n \times (0, 1]$  into  $\mathring{D}^n \times (0, 1]$ . Hence, the obvious thing to do would be to simply glue  $f$  and  $f'$  along  $D^{n+1} \times \{0\} \cup S^n \times [0, 1]$  to a map  $\hat{f}: D^{n+1} \times [0, 1] \cup S^n \times [0, 1] \rightarrow X$  and then set  $\tilde{f} = \hat{f} \circ i$ . The issue with this approach is that the map  $\hat{f}$  obtained in this way will generally not be continuous. Indeed, it is obtained by gluing two maps defined respectively on an open and a closed subset of  $D^{n+1} \times [0, 1] \cup S^n \times [0, 1]$ . To illustrate this point a little better, consider the homeomorphism

$$D^1 \rightarrow S^0 \times [0, 1] / S^0 \times \{1\}$$

$$y \mapsto \left[ \frac{y}{\|y\|}, 1 - \|y\| \right]$$

mapping 0 to the point given by  $S^1 \times \{1\}$ . We obtain a change of coordinates  $y \hat{=} [x, s]$ . By setting

$$f'_{x,t}(s) = f'([x, s], t)$$

we can interpret the data of a map  $f': D^1 \times [0, 1] \rightarrow X$  as a continuous family of paths  $f'_{x,t}: [0, 1] \rightarrow X$ , indexed over  $(x, t) \in S^n \times [0, 1]$ , which fulfill

$$\gamma_{x,t}(1) = \gamma_{x',t}(1),$$

for all  $x, x' \in S^n, t \in [0, 1]$ . If we want  $\hat{f} = f \cup f'$  to be continuous, then we precisely need convergence

$$f'_{x_n, t_n}(s_n) \rightarrow f([x, s], 1)$$

for sequences  $([x_n, s_n], t_n) \rightarrow ([x, 0], 1)$ . Let  $d: X \times X \rightarrow [0, 1]$  denote a metric that induces the topology on  $X$  and suppose that there is a uniform bound

$$d(f'_{x,t}(s), f'_{x,t}(0)) = d(f'_{x,t}(s), f([x, s], t)) \leq \varphi(s)$$

by some continuous function  $\varphi: [0, 1] \rightarrow [0, 1]$  with  $\varphi(0) = 0$ , at least for  $(s, t) \in [0, \frac{1}{2}] \times [\frac{1}{2}, 1]$ . It follows from the triangle inequality that

$$d(f'_{x_n, t_n}(s_n), f([x, s], 1)) \leq \varphi(s_n) + d(f([x_n, 0], t_n), f([x, 0], 1)),$$

for  $(s_n, t_n)$  close to  $(0, 1)$ . In particular, it then follows from the continuity of  $f$  that this expression converges to 0. If the speed at which the paths  $f'_{x_n, t_n}$  leave  $f'_{x_n, t_n}(0)$  becomes arbitrarily large as  $(x_n, t_n) \rightarrow (x, 1)$ , then such a bound  $\varphi$  may generally not exist. We may, however, use the metrizable of  $X$  to reparametrize the paths  $f'_{x,t}$  in a way to enforce such global bounds. We proceed as follows.

*Lemma 5.1.7.* Using the notation above,  $i$  is given by

$$i: D^{n+1} \times [0, 1] \rightarrow D^{n+1} \times [0, 1] \cup S^n \times [0, 1]$$

$$([x, s], t) \mapsto ([x, s], s \frac{t}{2} + (1-s)t).$$

It remains to show that we may without loss of generality assume that  $f'$  admits a bounding function  $\varphi: [0, 1] \rightarrow [0, 1]$  as above. For two topological spaces  $T, T'$ , denote by  $T^{T'}$  the mapping

space equipped with the compact open topology. If  $T'$  is locally compact Hausdorff, then this construction defines the right adjoint to the functor  $- \times T'$  ([DK70]). Furthermore, if  $T$  is metrizable, then the topology on  $T^{T'}$  is easily seen to be the topology of uniform convergence on every compactum. Finally, denote by  $\text{Aut}(\mathbb{R}_{\geq 0})$  the subspace of self-homeomorphisms of  $\mathbb{R}_{\geq 0}$ . Consider the following three maps.

$$M^{[0,1]} \rightarrow \text{Aut}(\mathbb{R}_{\geq 0})$$

$$\gamma \mapsto \sigma_\gamma := \{s \mapsto \sup_{t \leq s, 1} d(\gamma(t), \gamma(0)) + s\};$$

$$\text{Aut}(\mathbb{R}_{\geq 0}) \rightarrow \text{Aut}(\mathbb{R}_{\geq 0})$$

$$\sigma \mapsto \sigma^{-1};$$

$$\text{Aut}(\mathbb{R}_{\geq 0}) \rightarrow [0, 1]^{[0,1]}$$

$$\sigma \mapsto \{s \mapsto \min\{\sigma(s), 1\}\}.$$

It is not hard to see, using the topology of uniform convergence, that the first and last of these maps are continuous. That inversion is continuous follows from [Are46, Thm. 4]. Now, let

$$\rho : M^{[0,1]} \rightarrow [0, 1]^{[0,1]}.$$

be the composition of these three maps. Then  $\rho$  has the following properties:

- (i) $_\rho$   $\rho(\gamma)(0) = 0$  if and only if  $s = 0$ ;
- (ii) $_\rho$   $d(\gamma(\rho(\gamma)(s)), \gamma(0)) \leq s$ ,

for all  $s \in [0, 1], \gamma \in X^{[0,1]}$ . The first property is immediate by construction of  $\rho$ . The second inequality is obtained from

$$\begin{aligned} d(\gamma(\rho(\gamma)(s)), \gamma(0)) &\leq \sup_{t \leq \rho(\gamma)(s), 1} d(\gamma(t), \gamma(0)) + \rho(\gamma)(s) = \sigma_\gamma(\rho(\gamma)(s)) \\ &\leq \sigma_\gamma(\sigma_\gamma^{-1}(s)) \\ &= s. \end{aligned}$$

Next, denote by  $\phi : [0, 1]^2 \rightarrow [0, 1]$  a function fulfilling

- (i) $_\phi$   $\phi(s, t) = 0$ ,  $(s, t) \in [0, \frac{1}{2}] \times [\frac{1}{2}, 1]$  ;
- (ii) $_\phi$   $\phi(s, t) = 1$ ,  $(s, t) \in [0, 1] \times \{0\} \cup \{1\} \times [0, 1]$ ,

and, define

$$\begin{aligned} \Phi : D^{n+1} \times [0, 1] &\rightarrow D^{n+1} \times [0, 1] \\ ([x, s], t) &\mapsto ([x, (1 - \phi(s, t))\rho(f'_{x,t}, s) + \phi(s, t)s], t). \end{aligned}$$

Note first that  $\Phi$  is well defined, that is, its value is independent of  $x \in S^n$  when  $s = 1$ , since then also  $(1 - \phi(s, t))\rho(f'_{x,t}, s) + \phi(s, t)s = 1$ . Furthermore,  $\Phi$ , has the following properties:

- (i) $_\Phi$   $\Phi([x, s], t) = ([x, s], t)$ , for  $([x, s], t) \in D^{n+1} \times \{0\} \cup S^n \times [0, 1]$ ;

$$(ii)_\Phi \quad \Phi([x, s], t)(\mathring{D}^{n+1} \times [0, 1]) \subset \mathring{D}^{n+1} \times [0, 1];$$

$$(iii)_\Phi \quad \Phi([x, s], t) = \rho(f'_{x,t}, s), \text{ for } (s, t) \in [0, \frac{1}{2}] \times [\frac{1}{2}, 1].$$

Property (i) $_\Phi$  follows from Property (i) $_\phi$  and Property (i) $_\rho$ . Property (ii) $_\Phi$  follows from Property (i) $_\rho$ , and finally Property (iii) $_\Phi$  follows from Property (ii) $_\phi$ . We may then replace  $f'$  by  $f'' = f' \circ \Phi$ , obtaining

$$(i)_{f''} \quad f''([x, s], t) = f'([x, s], t) = f([x, s], t), \text{ for } ([x, s], t) \in D^{n+1} \times \{0\} \cup S^n \times [0, 1];$$

$$(ii)_{f''} \quad f''(\mathring{D}^{n+1} \times [0, 1]) \subset f'(\mathring{D}^{n+1} \times [0, 1]);$$

$$(iii)_{f''} \quad d(f'_{x,t}(s), f'_{x,t}(0)) \leq s, \text{ for } (s, t) \in [0, \frac{1}{2}] \times [\frac{1}{2}, 1].$$

Properties (i) $_{f''}$  and (ii) $_{f''}$  are immediate by Property (i) $_\Phi$  and Property (ii) $_{f''}$  respectively, while Property (iii) $_{f''}$  follows from Property (iii) $_\Phi$  and Property (ii) $_\rho$ .  $\square$

## C Remaining part of the proof of Proposition 4.0.1

In Proposition 4.0.1 (using the notation there) we claimed that a path  $\gamma$  from  $a$  to  $b$  that (described as starting from  $b$ ) ascends monotonously in height and passes to the left of  $b_1$ , to the right of  $c_2$ , to the left of  $b_3$  etc., as illustrated, cannot lie in the image of  $r_*$  (even up to stratified homotopy). Here, we provide a rigorous proof of this statement.

*Proof.* This insight can be formalized as follows: For  $n \in \mathbb{N}$ , denote by  $\mathcal{X}_{b_n}$  the stratified spaces obtained from  $\mathcal{X}$  by taking only the point  $b_n$  as the  $p$ -stratum. The identity at the space level  $1_X$  does not induce stratified maps  $\mathcal{X}, \mathcal{Y} \rightarrow \mathcal{X}_{b_n}$ . However, it nevertheless induces (non-)stratified maps of stratified singular simplicial sets. Furthermore, by the same argument as above,  $\text{Sing}_s(\mathcal{X}_{b_n})$  are quasi-categories. In addition to this, on  $\mathcal{X}_{b_n}$ ,  $r$  is stratified homotopic to the identity relative to  $a$  and  $b$ . It follows that we obtain a commutative diagram

$$\begin{array}{ccc} \text{hoSing}_s(\mathcal{X})(a, b) & \xrightarrow{r_*} & \text{hoSing}_s(\mathcal{Y})(a, b) \\ & \searrow \tau_{c_n} & \swarrow \sigma_{b_n} \\ & \text{hoSing}_s(\mathcal{X}_{b_n})(a, b) & \end{array} \quad (35)$$

Now,  $\text{Sing}_s(\mathcal{X}_{b_n})(a, b)$  is simply the set of homotopy classes of paths in  $X \setminus \{b_n\}$ , which is homotopy equivalent to  $S^1$ . Under post-composition with the straight line path from  $b$  to  $a$ , we may thus identify

$$\text{hoSing}_s(\mathcal{X}_{b_n})(a, b) \cong \text{hoSing}_s(\mathcal{X}_{b_n})(a, a) \cong \pi_1(S^1) \cong \mathbb{Z}.$$

We may proceed mutatis mutandis replacing  $b_n$  with  $c_n$ . Then, under this identification with the integers, the sequences  $\sigma_{b_n}([\gamma]), \tau_{c_n}([\gamma]) \in \mathbb{Z}$  are (modulo signs, originating from choices of orientation) given by:

$$\begin{aligned} & (1, 0, 1, 0, 1, \dots) \\ & (0, 1, 0, 1, 0, \dots). \end{aligned}$$

However, for every morphism  $g \in \text{hoSing}_s(\mathcal{X})(a, b)$  at least one of the sequences  $\tau_{b_n}(g)$  or  $\tau_{c_n}(g)$  has only finitely many non-zero values. Indeed, assume that  $g$  is such that both  $\tau_{b_n}(g)$  and  $\tau_{c_n}(g)$  have infinitely many non-zero values. Every such  $g$  can be represented by a path  $\gamma'$  which stays in

$\mathcal{X}_p$  for  $[0, \frac{1}{2}]$  and then exits into  $\mathcal{X}_q$ . For  $\tau_{b_n}(g)$  ( $\tau_{c_n}(g)$ ) to be non-zero,  $\gamma'$  must intersect the line segment connecting  $a$  and  $b_n$  ( $c_n$ ), as otherwise  $\gamma'$  maps into a contractible subspace of  $\mathcal{X}_{b_n}$  ( $\mathcal{X}_{c_n}$ ). Since these intersection points necessarily converge to  $a$  as  $n \rightarrow \infty$ , it follows that  $a \in \gamma[\frac{1}{2}, 1]$  and therefore  $\gamma(\frac{1}{2}) = a$ . It follows that  $\gamma'(t)$  lies strictly above  $a'$  (in direction of the  $y$ -axis), for all  $t > \frac{1}{2}$  greater than some  $t_h \in \frac{1}{2}$ . Furthermore, it follows from the stratification of  $\gamma$  and the assumption on infinitely many intersection points with the line segment  $b_n$  and  $a$  above, that we find  $t_b, t_c \in (\frac{1}{2}, t_h)$ , such that  $\gamma'(t_b)$  lies on a line segment between  $a$  and  $b_n$  for some  $n > 0$ , and  $\gamma'(t_c)$  lies on a line segment between  $a$  and  $c_m$ , for some  $m > 0$ . To see this (in the case of  $(b_n)$ ), consider a sequence  $t_i$  such that each  $\gamma'(t_i)$  lies on the line segment between  $a$  and  $b_i$ . By compactness of  $[\frac{1}{2}, 1]$ , we may assume that  $t_i$  converges to some  $t' \in [\frac{1}{2}, 1]$ . Since the intersection points of the line segments from  $a$  to  $b_i$  converge to  $a$ , it follows that  $\gamma'(t') = a$ . But  $\frac{1}{2}$  is the only value in  $[\frac{1}{2}, 1]$  with  $\gamma'(t') \in \mathcal{X}_p$ , hence  $t' = \frac{1}{2}$ , and it follows that  $t_i$  converges to  $\frac{1}{2}$ . Finally, the set of points strictly above  $a'$  in  $\mathcal{X}_q$  is disconnected, with the line segments between  $a$  and  $b_m$  and  $a$  and  $c_n$  lying in different components, for all  $m, n$ , in contradiction to the connectedness of  $\gamma'(\frac{1}{2}, t_h)$ .  $\square$

## D On the proof of [Hai23, Thm. 0.1.1/3.2.4]

In this section, we take a detailed look at the proof of [Hai23, Thm. 0.1.1/3.2.4], pointing out a gap in the latter which stems from different uses of the terminology *Segal space*.

**Remark D.0.1.** On the proof of [Hai23, Thm. 0.1.1/3.2.4]: Let  $\mathbf{Strat}_P^{\text{ex}}$  denote the full subcategory of  $\mathbf{Strat}_P$  given by the fibrant stratified spaces in the categorical model structure and denote by  $W_{\text{ex}}$  the class of categorical equivalences between the latter. It was already asserted in [Hai23, Thm. 0.1.1/3.2.4] that  $\text{Sing}_s$  induces an equivalence of quasi-categories

$$\mathbf{Strat}_P^{\text{ex}}[W_{\text{ex}}^{-1}] \xrightarrow{\cong} \mathcal{A}\mathbf{Strat}_P.$$

While this result is a consequence of the existence of the categorical semimodel structure on  $\mathbf{Strat}_P$  together with Theorem 3.3.1, [Hai23] suggested a proof that did not assume the existence of such a structure. We want to point out two gaps within this proof, which seem to require a notion of fibrant replacement in  $\mathbf{Strat}_P^c$  to be closed. Let us first sketch the proof in [Hai23]. Some of the equivalences of  $\infty$ -categories in [Hai23] are not made explicit. We will choose explicit models which, to the best of our knowledge, present the intended functors of  $\infty$ -categories.

1. Denote by  $\mathbf{sStrat}_P^{\text{ex}}$  the full subcategory of  $\mathbf{sStrat}_P$  given by the (bi)fibrant objects in  $\mathbf{sStrat}_P^c$  (i.e. quasi-categories with a conservative functor into  $P$ ) and by  $H$  the class of equivalences of quasi-categories over  $P$ , between the latter. The quasi-category  $\mathbf{sStrat}_P^{\text{ex}}[H^{-1}]$  is one possible model for the  $\infty$ -category of abstract stratified homotopy types over  $P$ ,  $\mathcal{A}\mathbf{Strat}_P$ . One may then consider a diagram of 1-categories (commutative up to natural isomorphism)

$$\begin{array}{ccc} \mathbf{Strat}_P^{\text{ex}} & \longrightarrow & \mathbf{sStrat}_P^{\text{ex}} \\ \downarrow & & \downarrow \mathcal{H}\text{olink} \\ \mathbf{Strat}_P & \xrightarrow{\mathcal{H}\text{olink}} & \mathbf{Fun}((\text{sd}P)^{\text{op}}, \mathbf{sSet}) \end{array} \quad (36)$$

Now, let  $W_{\text{d}\acute{e}}$  be the class of morphisms in  $\mathbf{Fun}((\text{sd}P)^{\text{op}}, \mathbf{sSet})$  that are weak equivalences in the model structure presenting *décollages* over  $P$ . The class of categorical equivalences in  $\mathbf{Strat}_P$ , denoted  $W_c$  is precisely the inverse image of  $W_{\text{d}\acute{e}}$  under  $\mathcal{H}\text{olink}_{\mathcal{I}}$ . Furthermore, [Hai23] proves that  $H$  is precisely the inverse image of  $W_{\text{d}\acute{e}}$  under the right vertical, and that

$W_{\text{ex}}$  is precisely the inverse image of  $H$  under  $\text{Sing}_s$ . Hence, there is an induced diagram of quasi-categories

$$\begin{array}{ccc}
\mathbf{Strat}_P^{\text{ex}}[W_{\text{ex}}^{-1}] & \xrightarrow{\text{Sing}_s} & \mathbf{sStrat}_P^{\text{ex}}[H^{-1}] \simeq \mathbf{AStrat}_P \\
\downarrow & & \downarrow \mathcal{H}\text{olink} \\
\mathbf{Strat}_P^{\text{c}} = \mathbf{Strat}_P[W_c^{-1}] & \xrightarrow{\mathcal{H}\text{olink}} & \mathbf{Fun}((\text{sd}P)^{\text{op}}, \mathbf{sSet})[W_{\text{d}\acute{e}}^{-1}],
\end{array} \tag{37}$$

commutative up to natural isomorphism. It follows from [Dou21c, Thm. 3], that the left lower horizontal is an equivalence. Furthermore, it follows either from [Waa24a, Theorem 2.2.15] or from [BGH18, Thm 2.7.4] that the right vertical is an equivalence. This already proves one of the central results of [Hai23], namely the existence of an equivalence  $\mathbf{Strat}_P^{\text{c}} \simeq \mathbf{AStrat}_P$ . In addition to this result, [Hai23] aims to show that the upper vertical in Diagram (37) is also an equivalence. This follows from commutativity of Diagram (37), if one can show the following two additional claims:

2. The right vertical in Diagram (37) is fully faithful: This is not commented on further in [Hai23]. Note that there is generally no reason to assume that in a category with weak equivalences  $(\mathbf{C}, W)$  with full subcategory  $\mathbf{A}$  the natural functor  $\mathbf{A}[\mathbf{A} \cap W^{-1}] \rightarrow \mathbf{C}[W^{-1}]$  is an equivalence of quasi-categories. Consider, for example, the flattening of a quasi-category  $\mathcal{C} \in \mathbf{sSet}$ , given by the 1-category  $(\Delta_{/\mathcal{C}})^{\text{op}}$ . The flattening of  $\mathcal{C}$  has a full discrete subcategory  $\mathcal{C}_0$  given by the objects of  $\mathcal{C}$ . If we denote by  $W$  the class of morphisms in  $(\Delta_{/\mathcal{C}})^{\text{op}}$  that map 0 to 0, there is a natural equivalence of quasi-categories,  $(\Delta_{/\mathcal{C}})^{\text{op}}[W^{-1}] \simeq \mathcal{C}$ , which fixed the objects in  $\mathcal{C}_0$  (see, for example, [Lan21, Thm. 3.3.8]). However,  $\mathcal{C}_0[W \cap \mathcal{C}_0^{-1}]$  remains a discrete category. A classical condition to even ensure equivalence between  $\mathbf{C}[W^{-1}]$  and  $\mathbf{A}[\mathbf{A} \cap W^{-1}]$  is the existence of an endofunctor  $F: \mathbf{C} \rightarrow \mathbf{C}$ , with image in  $\mathbf{A}$ , together with a zig-zag of natural transformations  $1 \leftrightarrow F$ , given by morphisms in  $W$ . In the situation of Diagram (37) this amounts to constructing a fibrant replacement in  $\mathbf{Strat}_P^{\text{c}}$ , which we have done in this work, but whose existence was not yet known at the time of writing of [Hai23].
3. The right vertical in Diagram (37) is essentially surjective: Again, this claim is close to requiring fibrant replacements in  $\mathbf{Strat}_P^{\text{c}}$ . It follows from the equivalence  $\mathbf{Strat}_P^{\text{d}} \simeq \mathbf{Diag}_P$ , that up to equivalence of functors in  $\mathbf{Diag}_P$ , every d\acute{e}collage lies in the image of  $\mathbf{Strat}_P \rightarrow \mathbf{Diag}_P$ . In other words, if we denote by  $\mathbf{Strat}_P^{\text{d}\acute{e}}$  the full subcategory of stratified spaces for which  $\mathcal{H}\text{olink}_{\mathcal{I}}$  is a d\acute{e}collage, then the restricted functor  $\mathbf{Strat}_P^{\text{d}\acute{e}} \rightarrow \mathbf{Diag}_P[W_{\text{d}\acute{e}}^{-1}]$  is essentially surjective. [Hai23] then claims that  $\mathbf{Strat}_P^{\text{d}\acute{e}} = \mathbf{Strat}_P^{\text{ex}}$ , from which the result would follow. With the definition of d\acute{e}collages we used here and which is also used in [Hai23] this is incorrect. The confusion arises from two possible definitions of d\acute{e}collages (and complete Segal spaces): One is expressed purely intrinsically to the  $\infty$ -category  $\mathbf{Fun}((\text{sd}P)^{\text{op}}, \mathcal{G}\text{rpd}_{\infty})$  (and  $\mathbf{Fun}(\Delta^{\text{op}}, \mathcal{G}\text{rpd}_{\infty})$ ) and one expressed in terms of injective model structures on the 1-categories  $\mathbf{Fun}((\text{sd}P)^{\text{op}}, \mathbf{sSet})$  (and  $\mathbf{Fun}(\Delta^{\text{op}}, \mathbf{sSet})$ ). Here, we have defined d\acute{e}collages intrinsically to  $\mathbf{Fun}(\text{sd}P^{\text{op}}, \mathcal{G}\text{rpd}_{\infty})$  and complete Segal spaces in [Hai23] are also defined intrinsically to  $\mathbf{Fun}(\Delta^{\text{op}}, \mathcal{G}\text{rpd}_{\infty})$ . Alternatively, one can define d\acute{e}collages as fibrant objects in the model category  $\mathbf{Fun}(\text{sd}P^{\text{op}}, \mathbf{sSet})^{\text{d}\acute{e}}$  (defined in [Waa24a, Subsec. 2.2], as a left Bousfield localization of the injective model structure) and, as it is classically done, complete Segal spaces as (certain) fibrant objects in the injective model structure on  $\mathbf{Fun}(\Delta^{\text{op}}, \mathbf{sSet})$ . Let us call the former definition  $\infty$ -categorical and the latter 1-categorical. [Hai23] makes use of a result of Joyal and Tierney concerning the relationship of quasi-categories and 1-categorical Segal spaces [JT07, Cor. 3.6], from which it follows that  $\mathbf{Strat}_P^{\text{ex}}$  consists exactly of such stratified spaces for which  $\mathcal{H}\text{olink}_{\mathcal{I}}(\mathcal{X})$  is a 1-categorical d\acute{e}collage. Due to the overlap in terminology, [Hai23] then reasons that  $\mathbf{Strat}_P^{\text{d}\acute{e}} = \mathbf{Strat}_P^{\text{ex}}$ . This is false.



Indeed, every stratified space with two strata  $\mathcal{X}$  has the property that  $\mathcal{H}o\text{Link}_{\mathcal{I}}(\mathcal{X})$  is an  $\infty$ -categorical décollage (there are no conditions to be verified). But for  $\mathcal{H}o\text{Link}_{\mathcal{I}}(\mathcal{X})$  to be injectively fibrant the map  $\mathcal{H}o\text{Link}_{\mathcal{I}}(\mathcal{X}) \rightarrow \mathcal{X}_p$  needs to be a Serre fibration. For example,  $|\Lambda_0^{\mathcal{J}}|_s$  with  $\mathcal{J} = [p \leq p < q]$  fails to have this property.