A GEOMETRIC MODEL FOR THE MODULE CATEGORY OF A STRING ALGEBRA

KARIN BAUR AND RAQUEL COELHO SIMÕES

ABSTRACT. In this paper, we give a geometric construction of string algebras and of their module categories. Our approach uses dissections of punctured Riemann surfaces with extra data at marked points, called labels. As an application, we give a classification of support τ -tilting modules in terms of arcs in such a tiled surface. In the case when the string algebra is gentle, we recover the classification given in [31].

CONTENTS

1.	Background on string algebras	3
2.	Labelled tiling algebras	7
3.	String algebras as labelled tiling algebras	10
4.	The module category of a labelled tiling algebra	12
5.	Classification of support tau-tilting modules	24
References		32

INTRODUCTION

Gentle algebras were introduced by Assem and Skowroński in the context of iterated tilted algebras of type A and type \tilde{A} [6]. As such, gentle algebras are one level of complexity up in the family of algebras that are morally of type A. They are a particularly nice class of algebras. For example, they are tame and derived-tame [13], Gorenstein [27] and closed under derived equivalence [44]. Within the class of tame algebras, the tractability of gentle algebras means they are a useful class of algebras to serve as test cases for more general conjectures in representation theory. In recent years, there has been renewed interest in gentle algebras due to their ubiquity, having connections with symplectic geometry via partially wrapped Fukaya categories [28, 37], algebraic geometry and theoretical physics via dimer models [14], Lie theory [32] and cluster theory.

Triangulations of Riemann surfaces give a geometric framework to study cluster theory [26]. This framework provides geometric models giving detailed information regarding the representation theory of associated (Jacobian) algebras and associated (cluster) categories (see e.g. [5, 12, 16, 18, 35]). Since then, geometric models have been developed in many contexts. For example, geometric models have been used to describe not only cluster categories but also module categories of gentle algebras [10], derived categories of gentle algebras and their connection with Fukaya categories of surfaces [37, 40], module categories and derived categories of skew-gentle algebras

²⁰²⁰ Mathematics Subject Classification. Primary: 05E10, 16G20, 16G70; Secondary: 05C10.

Key words and phrases. Labelled tiling algebras; string algebras; module category; geometric model; τ -tilting.

[31, 36], a generalisation of gentle algebras which are morally of type D, and very recently module categories of semilinear locally gentle algebras [7].

String algebras are the natural next step up in complexity in the family of algebras of "type A". In particular, all gentle algebras are string algebras. Like gentle algebras, the building blocks of the representation theory over string algebras, such as indecomposable modules, irreducible morphisms and Auslander-Reiten sequences, are well understood [17, 23, 34]. However, there are notable differences. For example, string algebras need not be Gorenstein and while they are tame they need not be derived-tame. It is natural to ask therefore whether module categories of string algebras can be described using geometric models and whether these geometric models can be used to shed light on the less well studied representation theory of string algebras and to further develop connections with other areas of mathematics.

Gentle algebras can be realised in terms of so-called *tiling algebras* associated to dissections of unpunctured surfaces with boundary. This realisation can be extended to locally gentle algebras, where we drop the finite-dimensional assumption. These can be associated to dissections of surfaces which may be punctured and may not have boundary [43]. By viewing a string algebra as a quotient of a locally gentle algebra, we realise string algebras as so-called *labelled tiling algebras* associated to dissections of punctured surfaces, possibly with no boundary, together with some extra data, in terms of labels at marked points.

Theorem A (Theorem 3.1). Let A be a finite dimensional monomial algebra. Then A is a string algebra if and only if A is a labelled tiling algebra of a marked punctured surface.

We are then able to describe the building blocks of the representation theory of a string algebra combinatorially using an associated labelled tiling (or dissection) of a surface.

Theorem B (Theorems 4.6 and 4.16). Let A = A(S) be a labelled tiling algebra with associated surface S. There are explicit bijections between:

- (1) the equivalence classes of permissible arcs in S and the strings of A;
- (2) isotopy classes of certain permissible closed curves in S and powers of bands;
- (3) certain moves between permissible arcs and irreducible morphisms of string modules.

In the case where there are no labels, the associated tiling algebra is locally gentle. We recover the gentle algebras case when there are no labels and the endpoints of the dissection lie on the boundary of S.

Certain permissible arcs, called R-arcs (see Definition 4.2), permit us to give a combinatorial description of all morphisms between string modules via crossings and intersections (see Theorem 4.11, and [20] for the gentle algebras case). In upcoming [8], R-arcs are used to study maximal almost rigid objects over gentle algebras. These are new representation theoretic objects defined in [9], for algebras of type A, in connection with Cambrian lattices.

Geometric models not only play an important role in relating different areas of mathematics, they are also useful to study quite often long-standing questions in representation theory. For example, geometric models for gentle algebras have been applied to the study of:

- *d*-finite representation algebras in higher homological algebra [29];
- derived invariants for algebras [2, 3, 4, 24, 37, 39, 41];

- length hearts which are crucial to the construction of Bridgeland stability conditions [20];
- silting theory [4, 21, 22], in particular in providing counterexamples for silting completion [38, 33].

Geometric models have also been useful in τ -tilting theory, introduced in [1] as a "completion" of classical tilting theory from the point of view of mutation. The geometric models for gentle and skew gentle algebras were used to classify support τ -tilting modules over these algebras ([31], see also [15, 42]) and to study the connectedness of the mutation graph [30].

In this paper, we classify support τ -tilting modules over string algebras, using a different set of permissible arcs, called M-arcs (see Definition 4.2). In the gentle algebras case, considered in [31], support τ -tilting modules can be described in terms of non-crossing arcs. For string algebras in general we have to allow some crossings, but these are easily identifiable by a local condition with respect to labels. This is what we mean by "good crossings" below (see Definition 5.9 for the precise definition).

Theorem C (Theorem 5.13). There is a bijection between the set of support τ -tilting modules and the set of maximal collections of generalised permissible arcs with good crossings.

Acknowledgments. The authors would like to thank the Isaac Newton Institute for Mathematical Sciences, Cambridge, for support and hospitality during the programme *Cluster algebras and representation theory*, where work on this paper was undertaken. This work was supported by EPSRC grant EP/R014604/1. This project has also been supported by the EPSRC Programme Grant EP/W007509/1 and by the Royal Society Wolfson Award RSWF/R1/180004 of the first author and by European Union's Horizon 2020 research and innovation programme through the Marie Skłodowska-Curie Individual Fellowship grant 838706 of the second author.

1. Background on string algebras

Let **k** be an algebraically closed field and $Q = (Q_0, Q_1)$ be a quiver, where Q_0 is the set of vertices and Q_1 is the set of arrows. Let *I* be an admissible ideal of $\mathbf{k}Q$, and *A* be the bound quiver algebra $\mathbf{k}Q/I$.

Given $a \in Q_1$, we denote its source by s(a) and its target by t(a). We will read paths in Q from left to right, and mod(A) will denote the category of right A-modules.

The finite dimensional algebra A is said to be *monomial* if I is generated by paths of length at least two. We refer to a path in I of length k a k-relation.

Definition 1.1. [6] A finite dimensional algebra A is a *string* algebra if it admits a presentation $A = \mathbf{k}Q/I$ satisfying the following conditions:

- (S1) Each vertex of Q is the source of at most two arrows and the target of at most two arrows.
- (S2) For each arrow a in Q, there is at most one arrow b in Q such that $ab \notin I$, and there is at most one arrow c such that $ca \notin I$.
- (S3) I is generated by paths of length at least two, i.e. A is monomial.

Definition 1.2. A finite dimensional algebra $A = \mathbf{k}Q/I$ is *gentle* if is it a string algebra satisfying the following additional conditions:

- (G1) For each arrow a in Q, there is at most one arrow b' such that $ab' \in I$ and there is at most one arrow c' such that $c'a \in I$.
- (G2) I is generated by 2-relations.

A vertex v satisfying (S1) and for which the paths of length 2 going through v satisfy (S2) and (G1) is called a *gentle vertex*.

Definition 1.3. A *locally gentle algebra* is an algebra which admits a presentation $\mathbf{k}Q/I$ satisfying the conditions in Definition 1.2 but which is not necessarily finite dimensional, i.e. I is not necessarily admissible.

Example 1.4. Consider the quiver Q

$$\begin{array}{c|c} 4 & \xrightarrow{d} & 1 & \xrightarrow{e} & 5 \\ c & \downarrow a \\ 3 & \xleftarrow{b} & 2 \end{array}$$

and the following ideals:

$$I_1 = \langle de, da, abc \rangle, \ I_2 = \langle de \rangle, I_3 = \langle da \rangle$$

Then $\mathbf{k}Q/I_1$ is a string algebra which is not gentle nor locally gentle. The algebra $\mathbf{k}Q/I_2$ is locally gentle but not gentle and $\mathbf{k}Q/I_3$ is gentle. See also Example 3.3 where we consider a different ideal.

1.1. Strings and bands. Let $A = \mathbf{k}Q/I$ be a string algebra. Given $a \in Q_1$, we define a formal inverse a^{-1} such that $s(a^{-1}) = t(a)$ and $t(a^{-1}) = s(a)$. We denote by Q_1^{-1} the set of formal inverses of arrows of Q.

A walk is a sequence $w = a_1 \dots a_r$, with $a_i \in Q_1^{-1} \cup Q_1$ for each i and $t(a_i) = s(a_{i+1})$. Each of the a_i 's is called a *letter of* w. Given a walk $w = a_1 \dots a_r$, a subwalk of w is a sequence of the form $a_i a_{i+1} \dots a_j$ for some $1 \leq i \leq j \leq r$, and the *inverse of* w is the walk $a_r^{-1} \dots a_1^{-1}$.

A string is a walk which is reduced, i.e. it has no subwalks of the form aa^{-1} or $a^{-1}a$ with $a \in Q_1$, and avoids relations, i.e. it has no subwalks v with $v \in I$ or $v^{-1} \in I$. If all letters of a string lie in Q_1 (resp. Q_1^{-1}), we say the string is *direct* (resp. *inverse*).

Each vertex $i \in Q_0$ defines two *trivial strings*, denoted by 1_i^+ (or simply 1_i) and its inverse 1_i^- . For technical reasons, we also need to define the *zero string*, which will be denoted by w = 0.

Given two non-trivial (and non-zero) strings $w = a_1 \dots a_r$ and $v = b_1 \dots b_s$, the concatenation $wv = a_1 \dots a_r b_1 \dots b_s$ is well defined if $t(a_r) = s(b_1)$. Moreover, wv is a string if and only if $b_1 \neq a_r^{-1}$ and there is no relation of the form $a_i \dots a_r b_1 \dots b_j$, for some $1 \leq i \leq r$ and $1 \leq j \leq s$.

Now suppose one of the strings, say w, is trivial. Write $w = 1_x^{\gamma}$, where $\gamma \in \{+, -\}$ and $x \in Q_0$. We would like there to be at most one choice of arrow a (resp. b) with s(a) = x (resp. t(b) = x) for which $1_x^{\gamma}a$ (resp. $1_x^{\gamma}b^{-1}$) is a string. However, there could be two arrows starting (resp. ending) at x. For this reason, we need the following technical definition.

Definition 1.5. Given a string algebra $A = \mathbf{k}Q/I$, we define two sign functions $\sigma, \varepsilon: Q_1 \to \{-1, 1\}$ satisfying the following conditions:

- If $b_1, b_2 \in Q_1$ are distinct arrows with the same start, then $\sigma(b_1) \neq \sigma(b_2)$.
- If $a_1, a_2 \in Q_1$ are distinct arrows with the same target, then $\varepsilon(a_1) \neq \varepsilon(a_2)$.
- If $a, b \in Q_1$ with s(b) = t(a) and $ab \notin I$, then $\sigma(b) = -\varepsilon(a)$.

We can extend the domain of these functions to the set of all strings, as follows. If $b \in Q_1$, we define $\sigma(b^{-1}) := \varepsilon(b)$ and $\varepsilon(b^{-1}) := \sigma(b)$.

If $w = a_1 \dots a_n$ is a non-trivial string (i.e. $n \ge 1$), then $\sigma(w) := \sigma(a_1)$ and $\varepsilon(w) := \varepsilon(a_n)$.

Given $v \in Q_0$, we define $\sigma(1_v^{\pm}) := \mp 1$ and $\varepsilon(1_v^{\pm}) := \pm 1$.

We are now ready to extend the definition of concatenation of strings to involve trivial strings as well. Let $v = 1_x^{\gamma}$, where $x \in Q_0$ and $\gamma \in \{1, -1\}$, and let w be a non-trivial string. Then vw is a string if and only if s(w) = x and $\sigma(w) = -\varepsilon(v) =$ $-\varepsilon(1_x^{\gamma}) = -\gamma$. Similarly, wv is a string if and only if t(w) = x and $\varepsilon(w) = -\sigma(v) = \gamma$.

Remark 1.6. The reader can find an algorithm for choosing sign functions in [17, p. 158]. We note that if A is (locally) gentle, we always have a choice of σ, ε so that given two strings v, w, the concatenation vw is defined and is a string if and only if t(v) = s(w) and $\sigma(w) = -\varepsilon(v)$. However, this is not the case for string algebras in general. For example, consider the quiver

$$1 \xrightarrow{a} 2 \xrightarrow{b} 3 \xrightarrow{c} 4$$

bound by the relations *abd* and *bc*. For any choice of σ and ε , we must have $\sigma(d) = -\varepsilon(b)$ because *bd* is a string. However, it follows that $\sigma(d) = -\varepsilon(ab)$, but *abd* is not a string.

Every string w of A defines an indecomposable module M(w), called *string module*. If we realise the string w as a graph, the string module M(w) is given by assigning a copy of the field \mathbf{k} to each of the vertices of the graph w and a copy of the identity map to each of the arrows in graph. Given two strings w, v, we have $M(w) \simeq M(v)$ if and only if w = v or $w = v^{-1}$. The module corresponding to the zero string is the zero module.

A string $b = b_1 \dots b_r$ which is cyclic, i.e. $t(b_r) = s(b_1)$, and such that any power b^k of b is a string but b itself is not a proper power of any string, is called a *band*.

Each band $b = b_1 \dots b_r$ (up to cyclic permutation and inversion) defines a family of indecomposable modules $M(b, n, \varphi)$, called *band modules*, where $n \in \mathbb{N}$ and φ is an irreducible automorphism of \mathbf{k}^n . The band module $M(b, n, \varphi)$ is defined by assigning a copy of \mathbf{k}^n to each vertex of b, the identity map to each arrow of b apart from b_r which is assigned the automorphism φ .

Theorem 1.7. [17] Every indecomposable module over a string algebra is either a string module or a band module.

1.2. Description of morphisms between string modules. We recall here how morphisms can be described in terms of string combinatorics, using the language of [11].

Let w be a string. A factor string decomposition of w is a decomposition of the form $w = w_{\ell}w_ew_r$, such that either w_{ℓ} is a trivial string or the last letter of w_{ℓ} lies in Q_1^{-1} and either w_r is trivial or the first letter of w_r lies in Q_1 . In this case, w_e is called a factor substring of w.

Similarly, a decomposition of w of the form $w = w_{\ell}w_ew_r$, where either w_{ℓ} is a trivial string or the last letter of w_{ℓ} lies in Q_1 and either w_r is trivial or the first letter of w_r lies in Q_1^{-1} , is called a *substring decomposition*. In this case, w_e is called an *image substring* of w.

The set of equivalence classes of factor string (resp. substring) decompositions of w is denoted by Fac(w) (resp. Sub(w)).

Let v, w be two strings. We call a pair $(v_\ell v_e v_r, w_\ell w_e w_r)$ in $\mathsf{Fac}(v) \times \mathsf{Sub}(w)$ an admissible pair if $w_e = v_e$ or $w_e = v_e^{-1}$. We will also sometimes call the string w_e an overlap between w and v. Homomorphisms between string modules arise from admissible pairs as has been shown by Crawley-Boevey:

Theorem 1.8. [23] Let v, w be strings and M(v), M(w) be the corresponding string modules. Then $\dim_{\mathbf{k}} \operatorname{Hom}_{A}(M(v), M(w))$ is given by the number of admissible pairs in $\operatorname{Fac}(v) \times \operatorname{Sub}(w)$.

1.3. Irreducible morphisms and the Auslander-Reiten translate. In order to describe irreducible morphisms between string modules, we need the definition of hook and cohook.

Definition 1.9. Let $A = \mathbf{k}Q/I$ be a string algebra and let $a \in Q_1$.

- (1) Let p_a denote the direct string with $s(p_a) = s(a)$ such that it does not start with a and it is *right maximal*, i.e. there is no $b \in Q_1$ with $p_a b$ a string. We call the string $a^{-1}p_a$ the *hook of a*. If s(a) has out-valency one, then p_a is a trivial string at s(a).
- (2) Let q_a denote the direct string with $t(q_a) = t(a)$ and such that it does not end with a and it is *left maximal*, i.e. there is no $b \in Q_1$ with bq_a a string. We call the string aq_a^{-1} the *cohook of a*. If t(a) has in-valency one, then q_a is a trivial string at t(a).

Let w be a string. We define a new string $f_s(w)$ as follows:

- If there is $a \in Q_1$ such that wa^{-1} is a string, then $f_s(w)$ is obtained by adding the hook of a on the right, i.e. $f_s(w) := wa^{-1}p_a$.
- If there is no $a \in Q_1$ such that wa^{-1} is a string and w is not inverse, then $f_s(w)$ is obtained from w by removing a cohook on the right. In other words, w must be of the form $w_{\ell}aq_a^{-1}$, for some $a \in Q_1$, in which case $f_s(w) := w_{\ell}$.
- If there is no $a \in Q_1$ such that wa^{-1} is a string and w is an inverse string, then $f_s(w) := 0$.

Similarly, we define $f_t(w)$ to be of the form $p_a^{-1}aw$, if there is $a \in Q_1$ with aw a string (adding a hook on the left), or of the form w_r , where $w = q_a a^{-1} w_r$ (removing a cohook on the left), or $f_t(w) = 0$ if w is direct and left maximal.

Remark 1.10. Let A be a string algebra and w be a string over A. It is not necessarily the case that $f_t(f_s(w)) = f_s(f_t(w))$. For example, consider the quiver

 $1 \xrightarrow{a} 2 \xrightarrow{b} 3 \xrightarrow{c} 4 \xleftarrow{d} 5 \xleftarrow{e} 6 \xleftarrow{f} 7$

bound by the relations *abc* and *fed*. Consider the string $w = bcd^{-1}e^{-1}$. We have $f_s(w) = b, f_t(f_s(w)) = ab, f_t(w) = e^{-1}$ and $f_s(f_t(w)) = e^{-1}f^{-1}$.

Lemma 1.11. Let A be a string algebra and w a string over A such that M(w) is not an injective module. Then at least one of $f_s(w)$ or $f_t(w)$ is non-zero. Moreover, if both $f_s(w)$ and $f_t(w)$ are non-zero strings then $f_s(f_t(w)) = f_t(f_s(w))$.

Proof. Since M(w) is not injective, we can assume without loss of generality that there is $a \in Q_1$ such that aw is a string. In particular $f_t(w) \neq 0$. Now assume $f_s(w)$ is also non-zero.

Case 1. There is an arrow b such that awb^{-1} is a string.

In this case, we have $f_t(w) = p_a^{-1}aw$ and $f_s(f_t(w)) = p_a a^{-1}wb^{-1}p_b$. On the other hand, $f_s(w) = wb^{-1}p_b$ and so $f_t(f_s(w)) = p_a a^{-1}wb^{-1}p_b = f_s(f_t(w))$.

Case 2. There is an arrow c such that $w = w_{\ell} c q_c^{-1}$.

Then $f_t(w) = p_a^{-1}aw$ and $f_s(f_t(w)) = p_a^{-1}aw_\ell$. On the other hand, $f_s(w) = w_\ell$ and $f_t(f_s(w)) = p_a^{-1}aw_\ell$, and we are done.

The only remaining case is when $w = q_a^{-1}$. However, we would have $f_s(w) = 0$, contradicting the hypothesis.

For a string module M(w), the irreducible morphisms starting at M(w) and its Auslander-Reiten translate can be described in terms of the strings $f_s(w)$ and $f_t(w)$ as follows.

Theorem 1.12. [17] Let A be a string algebra and w a string.

- (1) There are at most two irreducible morphisms starting at M(w), with possible targets $M(f_s(w))$ and $M(f_t(w))$.
- (2) If M(w) is non-injective, suppose without loss of generality that $f_s(w) \neq 0$. Then the Auslander-Reiten sequence starting at M(w) is of the form

$$0 \longrightarrow M(w) \longrightarrow M(f_s(w)) \oplus M(f_t(w)) \longrightarrow M(f_t(f_s(w)) \longrightarrow 0.$$

In particular, $\tau^{-1}(M(w)) = M(f_t(f_s(w))).$

The remaining irreducible morphisms are between band modules of the form $M(b, n, \varphi)$ and $M(b, n+1, \varphi)$. Moreover, the Auslander-Reiten translate acts as the identity morphism on band modules, since these modules lie in homogeneous tubes.

2. LABELLED TILING ALGEBRAS

2.1. Tiled surfaces and labels. We now recall the geometric set-up appearing in the construction of locally gentle algebras and introduce labels on tiled surfaces as a means towards string algebras.

Definition 2.1. A marked surface (S, N) is given by a connected oriented Riemann surface S with a finite set N of marked points on the (possibly empty) boundary and in the interior of S. We impose that each boundary component has marked points. We write N_b for $N \cap \partial S$ and $N^\circ = N \setminus N_b$ for the marked points in the interior. The elements of N° are called punctures. We will often simply write S for the tuple (S, N).

The segments between two neighboured marked points on a boundary component are called the *boundary segments* of (S, N).

Definition 2.2. Let S = (S, N) be a marked surface.

- (1) An arc γ in S is a curve in S such that its only intersections with N are the endpoints of γ . Curves are taken up to isotopy fixing endpoints.
- (2) Two arcs γ_1 and γ_2 are *intersecting* if they have a common endpoint.
- (3) Two arcs γ_1 , γ_2 are *non-crossing* if there exist representatives in their isotopy classes which do not meet anywhere except possibly at endpoints. The arcs are called *crossing* otherwise.
- (4) A collection of arcs is *non-crossing* if none of the arcs has self-crossings in the interior of **S** and the arcs are pairwise non-crossing.

Definition 2.3. A non-crossing collection P of arcs in S is called a *dissection* (also known as *partial triangulation* or *tiling*) of S if the following holds:

- no arc of P is isotopic to a boundary segment or to a point.
- The arcs of P dissect the surface into two types of tiles described as follows:
 (i) polygonal tiles of size ≥ 3 where exactly two consecutive sides are boundary segments;
 - (ii) polygonal tiles of size ≥ 1 with no boundary segment and with a marked point in the interior.

If P is a dissection of S, the pair (S, P) is also called a *tiled surface*.

Figure 1 gives an illustration of the types of tiles. We note that tiles are allowed to be self-folded. For example, in Figure 3 the tile on the right is an hexagon with the edge 5 glued to itself.



FIGURE 1. The types of tiles in a dissection. Figure (i.b) is the special case of a tile with boundary segments and of size 3.

Note that each tile of the dissected surface contains exactly one marked point in N which is not incident to any arc in P. If the tile has a boundary segment, then this marked point lies in it, otherwise the marked point is a puncture. As illustrated in Figure 1, we will draw these "isolated" marked points as red dots, and as such we will often call them *red points*.

Notation 2.4. A dissection P of S, defines a partition $M \cup R$ of the set of marked points N where R denotes the set of red points, i.e. the marked points not incident with any of the arcs in P, and M denotes the set of the remaining points, which are drawn in the figures as black dots.

We note that in [10], unmarked boundary components were allowed in certain tiles of type (ii). Here, we replace these unmarked boundary components with punctures in R.

Let $p \in M$. If p lies in the boundary of S, let p' and p'' be two points in the same boundary component such that $p', p'' \notin N$ and p is the only marked point in N lying in the boundary segment between p' and p''. Consider a curve δ isotopic to this boundary segment but such that the only intersection with the boundary of S is at its endpoints. If p is a puncture, consider a closed simple curve δ in the interior of S around p and with no intersections with N.

The complete fan at p is defined to be the sequence of all arcs of P that δ crosses in the clockwise order. A fan at p is any subsequence of consecutive arcs of the complete fan at p.

Remark 2.5. Let $p \in M$.

- (1) If an arc in P is a loop, then it lies in exactly one complete fan. All the remaining arcs lie in two distinct complete fans.
- (2) The complete fan at $p \in \mathbb{N}^{\circ} \cap \mathbb{M}$ is an infinite periodic sequence, with period ≥ 1 .

Any two consecutive crossings of the curve δ with arcs in P define a triangle whose vertices are these two crossings and the marked point p; see Figure 2. This triangle is called an *angle at* p (also called *angle of* P), and we say the curve δ cuts this angle at p.

We will now associate an algebra to a tiled surface. We note that the subset R of the marked points N of the surface does not play a role in the definition of this algebra. The subset R will be however useful in Section 4, when describing the module category of this algebra via the surface. In the context of gentle algebras, this definition recovers



FIGURE 2. The shaded triangle is an angle at p. A puncture $p \in M$ can have a unique angle which is a self-folded triangle.

that in [10, 40] and in the context of locally gentle algebras, which includes the case where the surface has no boundary, this definition recovers that in [43].

Definition 2.6. Let (S, P) be a tiled surface. We define Q = Q(S, P) to be the quiver defined as follows:

- (1) The vertices in Q are in bijection with the arcs in P.
- (2) The arrows in Q are in bijection with the angles at the marked points in M. In other words, there is an arrow $a: x_i \to x_j$ in Q if and only if the corresponding arcs \mathbf{x}_i and \mathbf{x}_j share an endpoint $p_a \in M$ and \mathbf{x}_j is the immediate successor of \mathbf{x}_i in the complete fan at p_a .

Definition 2.7. Let (S, P) be a tiled surface.

- (1) A *label* at a point $p \in M$ is a finite fan at p of length ≥ 3 .
- (2) Given two labels ℓ and ℓ' , if ℓ is a subsequence of ℓ' , then we say the label ℓ' is *redundant*.

Remark 2.8. Each label at a point $p \in M$ defines a linearly oriented path in the quiver Q. If p is a puncture, this path may be of the form a^k , where $a \in Q_1$ is such that s(a) = t(a); see Figure 3 for an example.

Notation 2.9. We will draw a label ℓ as a red curve δ around the corresponding marked point with an orientation such that $\delta(0)$ is a point in the first arc of ℓ , $\delta(1)$ is a point in the last arc of ℓ and if ℓ contains an arc **x** multiple times, the corresponding intersections of δ with **x** are at distinct points. This is in order to make it clear what the label is from δ , especially in the case when the label starts and ends at the same arc. See Example 2.12 and Figure 3.

Definition 2.10. Let (S, P) be a tiled surface and let L be a finite set of labels. Let Q = Q(S, P) and consider the path algebra kQ. We define $I_{P,L}$ to be the ideal of kQ generated by the following paths:

- (1) ab of length 2 such that $p_a \neq p_b$ or $p_a = p_b$ and t(a) = s(b) corresponds to a loop arc.
- (2) linearly oriented paths in Q corresponding to a label in L.

The relations of type (1) arise from the tiles in (S, P), as certain compositions of two arrows within a tile. These are all the relations considered in the construction of (locally) gentle algebras via surfaces. For string algebras in general, we need also to consider labels and corresponding relations of type (2).

Definition 2.11. Let S = (S, P) be a tiled surface and let Q = Q(S, P). Let L be a finite set of labels on S such that for every $q \in N^{\circ} \cap M$, L contains a label at q. The *labelled tiling algebra* $A_{P,L}$ associated to the data (S, P, L) is the bound quiver algebra $A_{P,L} = \mathbf{k}Q/I_{P,L}$.

Note that the reason to impose labels at all punctures in M is to ensure that the algebras are finite dimensional. One could drop this and work with infinite dimensional string algebras instead.

From now on we will assume for simplicity that there are no redundant labels in L. The following example illustrates these concepts.

Example 2.12. Consider the tiling P of a disk S with four punctures in Figure 3. The quiver of this tiling is in Figure 4, where vertex *i* corresponds to the arc \mathbf{x}_i , for each $1 \leq i \leq 5$.

Consider the labels given by the sequences $\{x_2, x_1, x_2\}$, $\{x_4, x_5, x_3\}$ and $\{x_5, x_5, x_5\}$. The ideal $I_{\mathsf{P},\mathsf{L}}$ is generated by the 2-relations of type (1)

 $\{a_1a_3, a_3a_4, a_4a_5, a_5a_1, a_6a_8, a_8a_7\},\$

together with the type (2) relations: $\{a_2a_1, a_6a_7, a_8^2\}$.



FIGURE 3. Labeled tiling of a surface.



FIGURE 4. The quiver of the tiling from Figure 3. The type (1) relations are indicated by dashed (red) lines.

3. String Algebras as labelled tiling Algebras

In this section we will show that the class of labelled tiling algebras coincide with the class of string algebras. We do this by realising a string algebra as the quotient of a locally gentle algebra, by removing all relations of length ≥ 3 and a choice of 2-relations so that every vertex becomes gentle. The relations removed will correspond to the labels.

Theorem 3.1. Let $A = \mathbf{k}Q/I$ be a finite dimensional monomial algebra. The following are equivalent.

- (1) A is a string algebra.
- (2) $A = (\mathbf{k}Q/J)/(I/J)$, where $\mathbf{k}Q/J$ is a locally gentle algebra.
- (3) A is a labelled tiling algebra of a marked surface.

Proof. (1) \iff (2): Suppose $A = \mathbf{k}Q/I$ is a string algebra. Pick a choice of two sign functions σ, ε on the set of strings which satisfy the conditions in Definition 1.5. Now let U be a set of generators of I and let V be the subset of U given by all paths of length ≥ 3 and all paths ab of length 2 such that $\sigma(b) = -\varepsilon(a)$. Let $J = \langle U \setminus V \rangle$.

Clearly we have $A = (\mathbf{k}Q/J)/(I/J)$. Moreover, the algebra $B := \mathbf{k}Q/J$ is locally gentle. Indeed, if there is a vertex x in Q_0 which is not gentle in B, then we can assume without loss of generality, that there are two relations in B of the form a_1b and a_2b , with s(b) = x. But then, for the choice of sign functions above, we would have $\varepsilon(a_1) \neq \varepsilon(a_2)$, and so $\sigma(b) = -\varepsilon(a_i)$, for some $i \in \{1, 2\}$. This contradicts the fact that a_ib is a relation in B. The converse is clear.

(2) \iff (3): This correspondence follows from the bijection between the set of isomorphism classes of locally gentle algebras and the set of homeomorphism classes of tiled surface from [43, Thm. 4.10], by associating relations generating I/J with the labels on the surface.

We refer the reader to [43, Definition 4.6] for a description of the surface associated to a given locally gentle algebra. Here the authors use a pair of dual dissections of the surface. Our partial triangulation is their green dissection, and our marked points in R are the vertices of their red dissection, the dual of P.

Remark 3.2. We highlight the following features of Theorem 3.1.

- (1) Given a string algebra A, the choice of an associated locally gentle algebra corresponds to the choice of sign functions σ and ε . Fix a choice, let B be the corresponding locally gentle algebra, and assign to B the same choice of sign functions. Then we have that vw is a string in B if and only if $\sigma(w) = -\varepsilon(v)$. In particular, if a, b are arrows of A such that ab corresponds to a label in the tiled surface associated to B, then ab is not a relation in B and so we have $\sigma(b) = -\varepsilon(a)$. This observation will be used in the proofs of Theorem 4.6 and Theorem 4.16.
- (2) The number of boundary components of the tiled surface associated to a locally gentle algebra is given in terms of perfect matchings of a graph whose vertices are so called blossom vertices, see [43, Remark 4.11 (vi)] for more details. It thus follows that a string algebra A is a labelled tiling algebra of a surface with no boundary if and only if every vertex of the quiver of A has valency four.
- (3) The existence of different choices of locally gentle algebras associated to a string algebra means that the surface associated to a string algebra is not unique in general. Example 3.3 illustrates this; see Figure 5.

Example 3.3. Consider the quiver Q

$$\begin{array}{c|c} 4 & \xrightarrow{d} & 1 & \xrightarrow{e} & 5 \\ c & \downarrow a \\ 3 & \xrightarrow{b} & 2 \end{array}$$

from Example 1.4 and the admissible ideal $I = \langle da, de, abc \rangle$. The algebra $A = \mathbf{k}Q/I$ is a string algebra, but not gentle.

The two possible choices of locally gentle algebras are the finite-dimensional algebra $B_1 = \mathbf{k}Q/J_1$ with $J_1 = \langle da \rangle$ and the infinite dimensional algebra $B_2 = \mathbf{k}Q/J_2$, with $J_2 = \langle de \rangle$. The tiled surfaces corresponding to B_1 and B_2 are illustrated in Figure 5 and are not homeomorphic to each other.

The algebra A can then be associated to the surface of B_1 with labels corresponding to the relations *abc* and *de*, and to the surface of B_2 with labels corresponding to the relations *abc* and *da*.



FIGURE 5. Non-homeomorphic surfaces associated to the same string algebra.

4. The module category of a labelled tiling algebra

In this section we will describe the module category of a string algebra via surfaces. Recall that $(S, M \cup R, P, L)$ denotes a labelled tiling, where the partial triangulation P divides S into a collection of polygonal tiles with at most one boundary segment in (S, M) each. Each of these tiles has a marked point in R which lies in the boundary segment if it exists, otherwise it is a puncture (see Figure 1). Recall also that $A_{P,L}$ denotes the corresponding labelled tiling algebra and any string algebra can be realised in this way.

4.1. Permissible arcs and closed curves. We will now adapt the definition of permissible arcs and closed curves from [10] to any labelled tiling algebra. In particular, we will extend this definition to consider permissible arcs whose endpoints lie in R. These haven't been considered in [10], but they have been considered in [20] and as we will see in Subsection 4.2, sometimes it is useful to consider these arcs instead.

We first need to set up some definitions and notation.

Definition 4.1. Let γ, γ' be two curves in $(S, M \cup R)$.

- (1) We say γ consecutively crosses $\mathbf{a}_1, \mathbf{a}_2 \in \mathsf{P}$ if γ crosses \mathbf{a}_1 and \mathbf{a}_2 at the points p_1 and p_2 , and the segment of γ between the points p_1 and p_2 does not cross any other arc in P .
- (2) We denote by $I(\gamma, \gamma')$ the minimal number of transversal intersections of representatives of the isotopy classes of γ and γ' in the interior of $S \setminus (M \cup R)^{\circ}$.
- (3) The intersection vector $I_{\mathsf{P}}(\gamma)$ of γ with respect to P is the vector $(I(\gamma, \mathbf{a}_i))_{\mathbf{a}_i \in \mathsf{P}}$.
- (4) The intersection number $|I_{\mathsf{P}}(\gamma)|$ of γ with respect to P is given by $\sum_{\mathsf{a}\in\mathsf{P}} I(\gamma,\mathsf{a})$.
- (5) γ is said to be a zero arc if $I_{\mathsf{P}}(\gamma)$ is the zero vector, i.e. γ does not intersect any arc in P except possibly at an endpoint.

Note that we might have $I(\gamma, \gamma') = 0$ for arcs γ, γ' sharing an endpoint. In particular, if $\gamma \in \mathsf{P}$, then $I(\gamma, \gamma') = 0$, for every arc $\gamma' \in \mathsf{P}$.

It will be convenient for us to realise each non-zero arc γ as a concatenation of segments, defined by the consecutive crossings with P. As such, we will often denote it by $\gamma = \gamma_1 \cdots \gamma_r$, with $r \ge 2$. The segments γ_1 and γ_r are called *end segments* and the remaining ones are called *interior segments*.

Definition 4.2. Let γ be an arc in $(S, M \cup R)$.

(1) γ is an R-arc if its endpoints lie in R.

(2) γ is an M-arc if each endpoint of γ lies in M or it is a red point lying in a tile of type (i.b) (See Figure 1).

We note that all arcs in [10] are what we now call M-arcs. Indeed, in the model used in [10], the only isolated marked points, i.e. the marked points not incident with the dissection P, were the ones lying in tiles of type (i.b).

We will see in Section 4.2 that R-arcs are useful to describe morphisms between string modules. On the other hand, M-arcs will be important in the classification of support τ -tilting modules (Section 5). For this classification, we make use of irreducible morphisms and the Auslander-Reiten translate. Therefore, we will consider M-arcs in their geometric descriptions in Sections 4.3 and 4.4.

Definition 4.3. Let γ be a curve in $(S, M \cup R)$.

- (1) γ is called *permissible* if it satisfies the following conditions:
 - (a) If γ consecutively crosses two (possibly not distinct) arcs **x** and **y** of P, then **x** and **y** have a common endpoint $p \in M$, and γ cuts the corresponding angle at p; see Figure 6.
 - (b) No segment of γ cuts a label in L.
- (2) γ is a *permissible* M-*arc* (R-*arc*, resp.) if it is both permissible and an M-arc (R-arc, resp.).
- (3) If γ is a closed curve, define $\gamma^n \colon [0, n] \to \mathsf{S}$, where $n \in \mathbb{N}$, by $\gamma^n(x) \coloneqq \gamma(x \lfloor x \rfloor)$ for $x \in [0, n]$.
- (4) γ is a *permissible closed curve* if it is a closed curve which satisfies (1)(a) and such that no segment of γ^n cuts a label in L for any $n \in \mathbb{N}$.



FIGURE 6. A consecutive cross cuts an angle of P.

We note that this definition generalises that in [10]. The condition in [10] regarding the winding number is not used here. In fact, this condition was used to describe a specific permissible M-arc used in the geometric description of irreducible morphisms and the AR-translate. However, we can give a different description of these M-arcs, without referring to the winding number (see Definition 4.15).

Remark 4.4. It follows from the definition that no closed curve which wraps around a puncture in M (see Figure 7) is permissible, because every puncture in M has a label.

We will regard permissible arcs up to equivalence, which requires more than just isotopy. The following definition of equivalence of permissible arcs coincides with that in [10].

Definition 4.5. Two permissible arcs γ and γ' in $(S, M \cup R)$ are said to be *equivalent* if we can write $\gamma = \gamma_1 \cdots \gamma_r$, $\gamma' = \gamma'_1 \cdots \gamma'_r$, where γ_i and γ'_i cut the same angle, for each $2 \leq i \leq r-1$.



FIGURE 7. Curves wrapping around a single puncture are not permissible.

Note that the end-segments do not play a role in the definition of equivalence of permissible arcs. The point of this definition is that equivalent permissible arcs correspond to the same string (see Theorem 4.6). Moreover, each equivalence class of permissible arcs contains exactly one permissible R-arc up to isotopy (see Figure 8 for an illustration).



FIGURE 8. These three arcs are permissible and equivalent to each other. The arc γ_1 (in blue) is the unique R-arc in this equivalence class. The arc γ_3 is an M-arc and γ_2 is neither an M-arc nor an R-arc.

Theorem 4.6. Let A be a string algebra and $(S, M \cup R, P, L)$ a labelled tiling associated to A. There are one-to-one correspondences between:

- (1) equivalence classes of permissible arcs in $(S, M \cup R)$ and the strings of $A = A_{P,L}$.
- (2) isotopy classes of permissible closed curves c in $(S, M \cup R)$ with $|I_P(c)| \ge 2$ and powers of bands of $A = A_{P,L}$.

Proof. To prove this we use a similar strategy as in the gentle case (see [10, Theorems 3.8 and 3.9]). First, we associate the zero string to any zero arc. Note that this includes boundary segments and arcs in P. For non-zero permissible arcs with intersection number with respect to P greater or equal to 2, each consecutive crossing of a curve with P corresponds to an arrow, and since all intersections with P are considered to be transversal, we obtain a reduced walk. Now suppose that this reduced walk contains a direct path $\alpha_1 \cdots \alpha_r$ which is a relation. Again by transversality of the intersections with P and by condition 1(a) in Definition 4.3, this relation is not a relation in A_P . Therefore, the path $\alpha_1 \cdots \alpha_r$ corresponds to a label, contradicting condition 1(b) in Definition 4.3.

The correspondence between (permissible) arcs with only one intersection with P and the trivial strings of $A_{\mathsf{P},\mathsf{L}}$ is as follows. Let σ and ε be the sign functions used to define A_{P} (see proof of Theorem 3.1).

Given an arc $\gamma = \gamma_1 \gamma_2$ with only one intersection with P, we associate to it the trivial string 1_x^{π} , where x and π are as in Figure 9. Note that not all of the four arrows a_1, b_1, a_2, b_2 have to exist, but if A is not an algebra of type A_1 , at least one of them does exist.



FIGURE 9. The arc $\gamma = \gamma_1 \gamma_2$ corresponds to the trivial string 1_x^{π} , where $\pi = \varepsilon(a_1) = \sigma(b_1) = -\varepsilon(a_2) = -\sigma(b_2)$.



FIGURE 10. Permissible closed curves corresponding to b_1 (on the left) and to b_2 (on the right).

Conversely, given a string 1_x^{π} , we associate to it an arc which intersects P once at arc **x**, and we orient this arc in such a way that γ satisfies the equalities described in Figure 9. Note that these equalities hold due to Remark 3.2 (1).

Finally, given a permissible closed curve γ , pick a crossing c with P, let $c = \gamma(0)$ and pick an orientation of γ . The corresponding power of a band b^k , for some $k \ge 1$, is given by the arrows corresponding to the angles that γ cuts from $\gamma(0) = c$ to $\gamma(1) = c$. \Box

Remark 4.7. The correspondence between permissible arcs and strings in Theorem 4.6 is such that:

- the inverse γ^{-1} of a permissible arc γ corresponds to w_{γ}^{-1} ,
- the intersection vector of a permissible arc γ gives the dimension vector of the corresponding string module.

Given a permissible curve γ in $(S, M \cup R)$, we will denote by w_{γ} the corresponding string of A. If the curve γ is an arc, the corresponding string module will be denoted by M_{γ} .

Remark 4.8. It follows from Theorem 4.6 that a labelled tiling algebra is of finite representation type if and only if the intersection number of any permissible closed curve with respect to P is at most one. We note that in the gentle case, it is enough to check this property for simple closed curves, cf. [10, Corollary 3.10]. However, this is not the case for string algebras in general. For example, the string algebra given in Example 2.12 is of infinite representation type, as it has two bands: $b_1 := a_7 a_4 a_6 a_8^{-1}$ and $b_2 := a_2 a_5^{-1} a_6 a_8^{-1} a_7 a_3^{-1}$. These bands correspond to the closed curves in Figure 10. Note that in this example there are no simple closed curves which are permissible.

4.2. Morphisms via crossings and intersections. In this section, we provide a combinatorial description of morphisms between string modules. We will make use of permissible R-arcs, as they permit us to give a simple description of morphisms via crossings and intersections. Such a description has been given in [20, Section 2.5] in the case when the algebra is gentle. The proof given in [20] is via an embedding of

the module category of a gentle algebra in the bounded derived category and therefore cannot be generalised to string algebras. Our proof is different and it relies on string combinatorics, namely Theorem 1.8.

Definition 4.9. Let γ and β be two permissible R-arcs, and suppose there is a crossing c between γ and β in the interior of S. Then there must be an arc \mathbf{x} in P crossed by both arcs γ and β . The arcs \mathbf{x}, γ and β enclose a simply connected triangle δ with c as one of its vertices. We say the crossing c is a crossing from β to γ if γ follows β in the clockwise order around δ . See Figure 11.



FIGURE 11. A crossing between R-arcs, from β to γ .

Note that a given crossing is incident with at most two triangles which are opposite to each other. Thus both triangles give rise to the same orientation of the crossing. Moreover, the orientation is independent of the choice of representatives of the arcs in their isotopy classes. Therefore, the orientation of the crossing is well defined.

Definition 4.10. Let γ and β be two permissible R-arcs.

- (1) We say there is an *intersection from* β to γ if there are end-segments β_0 and γ_0 of β and γ respectively sharing an endpoint $x \in \mathbb{R}$ and such that γ_0 follows β_0 in the anticlockwise direction around x.
- (2) A trivial intersection from β to β is an intersection from β to β in which the two end-segments coincide.
- (3) An intersection from β to γ is *traversed by* P if there is an arc τ in P which crosses both β and γ and such that β , γ and τ enclose a simply connected triangle. See Figure 12 for an illustration.



FIGURE 12. Left: Intersection from β to γ traversed by P. Middle: Intersection from β to γ not traversed by P. Right: Intersection from β to γ not traversed by P and intersection from γ to β traversed by P.

We note that our definition of intersections traversed by P is motivated by [8], where these are described in terms of certain angles at red points.

Denote by $C(\beta, \gamma)$ the number of crossings from β to γ and by $I_{R,P}(\beta, \gamma)$ the number of intersections from β to γ traversed by P.

Theorem 4.11. Let β and γ be permissible R-arcs, and M_{β}, M_{γ} the corresponding string modules. Then dim Hom $(M_{\beta}, M_{\gamma}) = C(\beta, \gamma) + I_{\mathsf{R},\mathsf{P}}(\beta, \gamma) - \delta_{\beta,\gamma}$.

Proof. The dimension of $\operatorname{Hom}(M_{\beta}, M_{\gamma})$ is given by the number of overlaps in $\operatorname{Fac}(\beta) \cap \operatorname{Sub}(\gamma)$. Let w_e be such an overlap, and assume first that $M_{\beta} \not\cong M_{\gamma}$.

Case 1: w_e is not a prefix or suffix of one of the strings, say w_β . Then $w_\beta = w_\ell a^{-1} w_e b w_r$, for some $a, b \in Q_1$. On the other hand we have $w_\gamma = v_\ell c w_e d^{-1} v_r$, where we may have $v_\ell c = 0$ and/or $d^{-1} v_r = 0$. In any case, we must have a crossing from β to γ , as Figure 13 illustrates.

Case 2: w_e is a prefix or suffix of both strings w_β and w_γ .

Case 2.1: w_e is non-trivial. We can assume, without loss of generality, that either w_e is a prefix of both w_β and w_γ or w_e is a prefix of w_β and suffix of w_γ . In the first case, there is an intersection from β to γ traversed by P. In the second case, we can assume $w_\beta = w_e b w_r$ and $w_\gamma = v_\ell c w_e$, with $b, c \in Q_1$, as otherwise we would be in the first case situation. As Figure 14 illustrates, we must have a crossing from β to γ .



FIGURE 13. Case where w_e is not a suffix or prefix of one of the strings, say w_{β} .



FIGURE 14. Case where w_e is a prefix of w_β and a suffix of w_γ .

Case 2.2: w_e is trivial, i.e. $w_e = 1_x^{\pi}$, with $x \in Q_0$ and $\pi \in \{+, -\}$. Without loss of generality, we can write $w_{\beta} = bw_r$ and $w_{\gamma} = d^{-1}v_r$, with $b \in Q_1$ (resp. $d^{-1} \in Q_1^{-1}$) if w_{β} (resp. w_{γ}) is non-trivial, such that s(b) = t(d) = x. Since $M_{\beta} \not\cong M_{\gamma}$, at least one of w_{β} and w_{γ} is non-trivial. Without loss of generality, assume the latter.

If w_{β} is trivial or db = 0 is not a label, then we have an intersection from β to γ traversed by P (see Figure 15). Otherwise, either db is a label or $db \neq 0$, in which case we have a crossing from β to γ , as Figure 16 illustrates.

If $M_{\beta} \cong M_{\gamma}$, then there are two intersections from β to itself, at both end-segments. Both intersections are trivial and traversed by P. We can associate these trivial intersections to the overlap $w_e = w_{\beta}$. The corresponding morphism is the identity morphism.

In summary, we associated one crossing or one non-trivial intersection from β to γ traversed by P to a morphism from M_{β} to M_{γ} which is not the identity morphism,



FIGURE 15. Case where w_{β} is trivial or db = 0 but not a label.



FIGURE 16. Case where db is a label or $db \neq 0$.

and we associated the two trivial intersections from β to γ to the identity morphism, in the case when $M_{\beta} \cong M_{\gamma}$.

Conversely, let c be a crossing from β to γ , and let α_i and β_j be the segments of α and β , respectively, where the crossing takes place. We can then consider the maximal sequences of consecutive segments:

$$(\gamma_{i-s},\ldots,\gamma_i,\gamma_{i+1},\ldots,\gamma_{i+r})$$
 and $(\beta_{j-s},\ldots,\beta_j,\beta_{j+1},\ldots,\beta_{j+r})$

of γ and β , respectively, which read off the same string (see Figure 17 for an illustration). Denote such a string by w_e . Note that w_e can be a trivial string, but it is always non-zero, since there is at least one arc in P that is crossed both by β and γ . Given the crossing c is from β to γ , we must have $w_{\beta} = w_{\ell}a^{-1}w_ebw_r$ and $w_{\gamma} = v_{\ell}cw_ed^{-1}v_r$, where at least one of $w_{\ell}a^{-1}$, $v_{\ell}c$ is non-zero and at least one of bw_r , $d^{-1}v_r$ is non-zero. It then follows from Theorem 1.8, that there is a morphism from M_{β} to M_{γ} corresponding to the overlap w_e .



FIGURE 17. The maximal sequence of segments reading off the same string.

Now, let c be an intersection from β to γ traversed by an arc in P. Suppose this intersection is non-trivial.

Claim: There are sequences of consecutive segments

 $(\beta_1,\ldots,\beta_r,\beta_{r+1})$ and $(\gamma_1,\ldots,\gamma_r,\gamma_{r+1})$

of β and γ respectively, for some $r \ge 1$, such that $s(\beta_1) = s(\gamma_1) = c, \beta_1 \dots \beta_r$ and $\gamma_1 \dots \gamma_r$ read off the same string w_e , and $\beta_{r+1}, \gamma_{r+1}$ read off different strings (one of which may be a zero string).

Proof of the claim. If $M_{\beta} \not\cong M_{\gamma}$, this is clear. So suppose $M_{\beta} \cong M_{\gamma}$. Then a non-trivial intersection from β to $\gamma = \beta$ would imply that both end-segments of β meet at a point $x \in \mathbb{R}$. Suppose for a contradiction that $w_{\beta} = w_{\beta}^{-1}$. Then $w_{\beta} = a_1 \cdots a_n$ is non-trivial, with either $a_{\lceil \frac{n}{2} \rceil} = a_{\lceil \frac{n}{2} \rceil}^{-1}$, if n is odd, or $a_{\frac{n}{2}} = a_{\frac{n}{2}+1}^{-1}$, if n is even, both a contradiction. Therefore $w_{\beta} \neq w_{\beta}^{-1}$, and the claim follows.

Since the intersection is from β to γ it follows that w_e is a factor substring of w_β and an image substring of w_γ . Hence, by Theorem 1.8, we have a morphism from M_β to M_γ corresponding to the overlap w_e .

Finally, if the intersection c is trivial, then it corresponds to the identity morphism. But for each arc β there are two such trivial intersections, so we are counting the identity morphism twice.

Remark 4.12. Let A be a string module, A = B/I, where B is a locally gentle algebra, and let $(S, M \cup R, P, L)$ be the corresponding labelled tiling. Given two string A-modules M_{β} and M_{γ} , with R-arcs β and γ then $C(\beta, \gamma)$ is the number of overlap extensions from M_{β} to M_{γ} in mod(B) (see [19, 20] for more details).

Example 4.13. Consider the string algebra A given by the labelled tiling in Figure 3 (see Figure 4 for the quiver of this tiling).

Let $w = a_7^{-1}a_8$ and $v = a_8a_6^{-1}$. Figure 18 illustrates the five crossings and intersections, c_1, \ldots, c_5 , between the corresponding permissible R-arcs.



FIGURE 18. The crossings and intersections between the R-arcs γ_w, γ_v corresponding to strings w and v.

The intersections c_1 and c_5 are not traversed by P. The crossing c_2 is a crossing from γ_w to γ_v corresponding to the admissible pair $(a_7^{-1}1_5^{\pi}a_8, a_81_5^{\pi}a_6^{-1})$ in $\mathsf{Fac}(w) \times \mathsf{Sub}(v)$, with overlap 1_5^{π} , where $\pi = -\sigma(a_8)$. The crossing c_3 is from γ_w to γ_v and it corresponds to the admissible pair $(a_7^{-1}a_8, a_8a_6^{-1})$ in $\mathsf{Fac}(w) \times \mathsf{Sub}(v)$ with overlap a_8 . Finally the crossing c_4 is a crossing from γ_v to γ_w , and it corresponds to the admissible pair $(1_5^{\pi}a_8a_6^{-1}, a_7^{-1}a_81_5^{\pi})$ in $\mathsf{Fac}(v^{-1}) \times \mathsf{Sub}(w)$, with overlap 1_5^{π} , where $\pi = -\sigma(a_8)$.

The dimension of $\operatorname{Hom}(M(w), M(v)) = 2$, the dimension of $\operatorname{Hom}(M(v), M(w)) = 1$, $I_{\mathsf{P}}(v, w) = 0 = I_{\mathsf{P}}(w, v), C(w, v) = 2$ and C(v, w) = 1.

4.3. **Pivot elementary moves.** In order to describe the Auslander-Reiten translate of a string module combinatorially, we first need to give a combinatorial description

of the irreducible morphisms starting at string modules. We will give this description in terms of M-arcs instead of R-arcs because it allows us to give a more explicit classification of support tau-tilting modules (see Section 5).

The description of irreducible morphisms is split into two steps. The first step consists of choosing a specific representative in the equivalence class of permissible arcs.

Definition 4.14. Let $\gamma = \gamma_0 \gamma_1 \cdots \gamma_t$ be a permissible M-arc (or R-arc). Denote by Δ_s (resp. Δ_t) the tile in which γ_0 (resp. γ_t) lies in.

The anticlockwise-most M-arc representative $\gamma_{\mathcal{O}}$ in the equivalence class of γ is described as follows. The segments of $\gamma_{\mathcal{O}}$ which are interior segments are uniquely determined by the string defined by γ . The start segment (resp. terminal segment) of $\gamma_{\mathcal{O}}$ is such that the concatenation of $\gamma_{\mathcal{O}}$ with the side of Δ_s (resp. Δ_t) incident with the starting point (resp. ending point) of $\gamma_{\mathcal{O}}$ in the anticlockwise direction gives rise to an arc which is not equivalent to γ . Figure 19 illustrates the end segments of $\gamma_{\mathcal{O}}$.



FIGURE 19. The end segments of the representative γ_{0} .

Note that we can dually define the *clockwise-most* M-*arc representative*, which will be denoted by γ_{\heartsuit} .

We are now ready to describe the second step, which produces a new arc $f_t(\gamma)$ (resp. $f_s(\gamma)$) from γ_{\bigcirc} by performing a pivot elementary move which fixes the endpoint (resp. starting point) of γ_{\bigcirc} .

Definition 4.15. Let \mathbf{x}_1 be the first arc in P crossed by $\gamma_{\mathcal{O}}$ and \mathbf{x}_0 be the side of Δ_s which precedes \mathbf{x}_1 with respect to the clockwise direction around the tile. Denote by p_s the common endpoint of \mathbf{x}_0 and \mathbf{x}_1 , and by s the starting point of $\gamma_{\mathcal{O}}$. Denote by $f_t(\gamma)$ be the arc obtained from $\gamma_{\mathcal{O}}$ as follows:

Case 1. There is a label at p_s starting at the angle in Δ_s and such that it is *shorter* than $\gamma_{(5)}$, i.e. it crosses fewer arcs in $P \setminus \{x_0\}$ adjacent to p_s than $\gamma_{(5)}$ does.

In this case, we define $f_t(\gamma)$ to be the concatenation of γ_{\bigcirc} with \mathbf{x}_0 (see Figure 20).

Case 2. There is no label as in Case 1, but there are labels at s starting after the angle in Δ_s in the clockwise direction around s.

Consider the first label at s, when going around s in the clockwise direction from \mathbf{x}_0 , and let \mathbf{y} be the last arc crossed by that label. We define $f_t(\gamma)$ to be the concatenation of γ_{\bigcirc} with \mathbf{y} (see Figure 21).

Case 3. There are no labels as in Cases 1 and 2.

Note that this implies that s is a marked point in the boundary of S, as otherwise, there would be a label at s, and we would be in the situation of Case 2.

As such, we define $f_t(\gamma)$ to be the concatenation of γ_{\bigcirc} with the boundary segment connecting s with its predecessor in M in the anticlockwise direction around the boundary component (see Figure 22).



FIGURE 20. Pivot elementary move, case 1.



FIGURE 21. Pivot elementary move, case 2.



FIGURE 22. Pivot elementary move, case 3.

The arc $f_t(\gamma)$ is called the *pivot elementary move of* γ *at its ending point*. The *pivot elementary move* $f_s(\gamma)$ *of* γ *at its starting point* is defined dually.

Note that if the algebra is gentle then we are always in Case 3, and we recover the definition in [10].

Theorem 4.16. Let w be a string, γ be a permissible M-arc corresponding to w and M_{γ} be the corresponding string module. Each irreducible morphism starting at M_{γ} is obtained by a pivot elementary move at an endpoint of $\gamma_{(5)}$.

Proof. Given a string w, recall that the orientation of any permissible arc associated to w is well-defined if w is non-trivial. In the case when w is trivial, the orientation is determined using the fixed choice of sign functions; see Figure 9.

We will show that $f_t(\gamma_{\mathcal{O}})$ is a permissible M-arc whose corresponding string is $f_t(w)$. The proof that $f_s(\gamma_{\mathcal{O}})$ corresponds to $f_s(w)$ is similar. The result then follows from Theorem 1.12.

Case a. There is an arrow $a \in Q_1$ such that aw is a string.

Then $f_t(w)$ is obtained by adding the hook of a on the left of w, i.e. $p_a^{-1}aw$. Due to the orientation of γ , we have that $s(a) = x_0$, where x_0 is the vertex corresponding to the arc \mathbf{x}_0 in Definition 4.15. Indeed this is easy to see when w is non-trivial, and when w is trivial, it must be of the form $\mathbf{1}_{x_1}^{\varepsilon(a)}$ since aw is a string and so the orientation of $\gamma_{(5)}$ is as in Figure 23.



FIGURE 23. The arc γ_{\bigcirc} corresponding to $1_{x_1}^{\varepsilon(a)}$.

Moreover, there cannot be a label at p_s as in Case 1 of Definition 4.15, as otherwise aw would not be a string. We are therefore in either Case 2 or Case 3 of Definition 4.15. In either case, the string corresponding to $f_t(\gamma)$ does indeed correspond to $p_a^{-1}aw$. In Case 2, the path p_a is right maximal in the sense that there is $b \in Q_1$ with $s(b) = t(p_a)$ but $p_a b$ is a relation corresponding to a label. In Case 3, the path p_a is right maximal because there is no $b \in Q_1$ with $s(b) = t(p_a)$.

Case b. There is no arrow a such that aw is a string.

This means that either \mathbf{x}_0 is a boundary segment in (S, M), and so we are in Case 3 of Definition 4.15 or there is a label at p_s as in Case 1 of Definition 4.15. In both cases, we have that $f_t(\gamma)$ is the concatenation of $\gamma_{\mathcal{O}}$ with \mathbf{x}_0 .

This move removes the direct path p at the start of w which corresponds to a fan at p_s . Moreover, $f_t(\gamma)$ is either an arc with no intersections with P if w is direct or it is an arc whose corresponding string w_r is such that $w = q_b b^{-1} w_r$, where $b \in Q_1$ and $q_b b^{-1}$ is the cohook of b. In either case, the string corresponding to $f_t(\gamma)$ is indeed $f_t(w)$, and we are done.

4.4. Geometric interpretation of the Auslander-Reiten translate. We are now ready to describe the (inverse of the) Auslander-Reiten translate of a string module in the geometric model.

Definition 4.17. Let γ be a permissible arc, and consider the corresponding anticlockwise most M-arc representative γ_{\bigcirc} . Let \mathbf{a}_s and \mathbf{a}_t be the arcs of P or boundary segments of (\mathbf{S}, \mathbf{M}) such that $f_t(\gamma)$ (resp. $f_s(\gamma)$) is the concatenation of γ_{\bigcirc} with \mathbf{a}_s (resp. \mathbf{a}_t). Define $\tau^{-1}(\gamma)$ to be the concatenation of γ_{\bigcirc} with the arcs \mathbf{a}_s and \mathbf{a}_t . The arc $\tau(\gamma)$ is defined dually using γ_{\bigcirc} .

The action of τ^{-1} (or τ) on an arc can be seen as a generalisation of the "classical" description, in the Jacobian algebras case, of the Auslander-Reiten translate in terms of rotation of the arc. The following proposition generalises Theorem 3.9 and Corollary 3.17 of [10].

Proposition 4.18. Let M be an indecomposable module over a labelled tiling algebra A associated to (S, M, P, L) and γ a curve corresponding to M.

- (1) M is injective at vertex v if and only if $\tau^{-1}(\gamma) = v \in P$.
- (2) *M* is a non-injective string module if and only if $\tau^{-1}(\gamma)$ has non-zero intersection with P, in which case the module corresponding to $\tau^{-1}(\gamma)$ is $\tau^{-1}(M)$.
- (3) If M is a non-injective string module and $f_s(\gamma) \neq 0$ (resp. $f_t(\gamma) \neq 0$), then $\tau^{-1}(\gamma) = f_t(f_s(\gamma))$ (resp. $\tau^{-1}(\gamma) = f_s(f_t(\gamma))$).
- (4) If M is a band module then γ corresponds also to $\tau^{-1}(M)$.

Proof. Suppose M is a string module. Then M is injective if and only if the arc γ_{\bigcirc} meets only two fans which are arranged as in Figure 24. Note that \mathbf{a}_s (resp. \mathbf{a}_t) is either the last arc crossed by the first label at p_s (resp. p_t) which starts at or after \mathbf{v} in

the anticlockwise direction around p_s (resp. p_t) or \mathbf{a}_s (resp. \mathbf{a}_t) is a boundary segment in (S, M) if there is no such label. In any case, condition (1) follows.



FIGURE 24. The arc $\gamma_{(5)}$ corresponding to I_v .

In particular, if M is injective, then $\tau^{-1}(\gamma)$ does not intersect P traversely. In order to prove condition (2) it remains to show the forward implication. So suppose M is a non-injective string module. Lemma 1.11 and Theorem 4.16 imply that $f_s(\gamma) \neq 0$ or $f_t(\gamma) \neq 0$. Assume, without loss of generality, that $f_s(\gamma) \neq 0$. By Theorem 1.12, $\tau^{-1}(M)$ corresponds to the arc $f_t(f_s(\gamma))$. So it is enough to show that $\tau^{-1}(\gamma) =$ $f_t(f_s(\gamma))$, which would also prove condition (3). We do this now.

Case a. $f_s(w_{\gamma})$ is obtained by adding a hook $a^{-1}p_a$ on the right.

In this case, we can either add a hook $p_c^{-1}c$ on the left of $f_s(w_{\gamma})$ or remove a cohook $q_c c^{-1}$ on the left of $f_s(w_{\gamma})$. The latter is always possible since $f_s(w_{\gamma}) = w a^{-1} p_a$ is not a direct string. Figure 25 illustrates the different possibilities. In any of the cases, we have $\tau^{-1}(\gamma) = f_t(f_s(\gamma)).$



FIGURE 25. Possible cases when $f_s(w_{\gamma})$ is obtained by adding a hook on the right.

Case b. w_{γ} is not direct and $f_s(\gamma)$ is obtained by removing a cohook bq_b^{-1} on the right.

In this case, since M is not injective, we must be able to add a hook $p_a^{-1}a$ on the left of w_{γ} . Figure 26 illustrates this case. It is clear that $\tau^{-1}(\gamma) = f_t(f_s(\gamma))$.

This finishes the proof of (2) and (3). Condition (4) is clear since τ^{-1} acts as the identity on band modules.

Remark 4.19. Note that $\tau^{-1}(\gamma)$ is not necessarily equivalent to $f_s(f_t(\gamma))$ or to $f_t(f_s(\gamma))$, as pointed out in Remark 1.10. Figure 27 shows permissible arcs corresponding to the strings $w, f_s(f_t(w))$ and $f_t(f_s(w))$, none of which are zero arcs. We have $f_s(f_t(w)) \neq f_t(f_s(w))$. Moreover, $\tau^{-1}(w)$ corresponds to the injective module at vertex 4 and $\tau^{-1}(\gamma)$ is the arc in P corresponding to this vertex.



FIGURE 26. Possible case when $f_s(w_{\gamma})$ is obtained by removing a cohook on the right and M is not injective.



FIGURE 27. Example where $\tau^{-1}(\gamma)$, $f_s(f_t(\gamma))$ and $f_t(f_s(\gamma))$ are pairwise non-equivalent.

5. Classification of support tau-tilting modules

In this section we will use the geometric model of string algebras to give a combinatorial classification of support τ -tilting modules over this class of algebras. We begin by giving the necessary background on τ -tilting theory, and we refer the reader to [1] for more details.

Definition 5.1. Let A be a finite-dimensional algebra and $M \in \text{mod}(A)$. Denote by |M| the number of non-isomorphic indecomposable direct summands of M.

- (1) M is τ -rigid if $\text{Hom}_A(M, \tau M) = 0$.
- (2) M is τ -tilting if M is τ -rigid and |M| = |A|.
- (3) *M* is support τ -tilting if there is an idempotent $e \in A$ such that *M* is a τ -tilting $(A/\langle e \rangle)$ -module.
- (4) A support τ -rigid pair is a pair (P, M) consisting of a projective A-module P and a τ -rigid A-module M such that $\operatorname{Hom}_A(P, M) = 0$.
- (5) A support τ -tilting pair is a τ -rigid pair (P, M) for which |M| + |P| = |A|.

Proposition 5.2. [1, Proposition 2.3] Let A be a finite-dimensional algebra and $M \in \text{mod}(A)$. The module M is support τ -tilting if and only if there is a projective A-module P such that (P, M) is a support τ -tilting pair.

Proposition 5.3. [1, Corollary 2.13] A pair (P, M) is support τ -tilting if and only if it is maximal with respect to the τ -rigid pair property.

Note that if the algebra A is a string algebra, then no band module is τ -rigid since the Auslander-Reiten translate acts as the identity. Therefore, we will only need to consider string modules.

5.1. Crossings of M-arcs. As we have seen in Section 4.2, we can describe morphisms between two string modules M_{β} and M_{γ} in terms of the crossings and intersections



FIGURE 28. Each crossing of permissible clockwise-most M-arcs defines a triangle.

between the corresponding R-arcs β and γ . In this section we will see that certain representatives of M-arcs, namely the clockwise-most ones, are more useful to describe τ -rigidity, in the sense that most crossings between these arcs detect morphisms in the Hom-space Hom $(M, \tau N)$, where M, N are string modules.

In order to see this, we first need to define an orientation for each crossing.

Lemma 5.4. Let γ and β be non-trivial permissible clockwise-most M-arcs, and suppose they cross at a point c in their interior. Then there is one arc \mathbf{x} in P and a triangle of one of the forms given in Figure 28 with no puncture in R in its interior.

Proof. Let Δ be the tile the crossing c lies in. If there is $\mathbf{x} \in \mathsf{P}$ which bounds Δ which is intersected by γ and β , then we clearly have a triangle as the one on the left of Figure 28.

Suppose the segments of γ and β in Δ where the crossing *c* occurs don't have a common intersection with P. Since they are permissible arcs which cross each other, at least one of the segments must be an end-segment. The fact that there is a local triangle δ follows from the fact that both arcs are permissible and clockwise-most M-arcs. Moreover, one of the vertices *p* of δ must be in M. Let **x** be the arc in P bounding δ . If the end-segment of γ or β follows **x** around *p* in the anticlockwise order instead, then we must be in one of the situations presented in Figure 29. Therefore, there is another triangle δ' of the form given on the right of Figure 28.



FIGURE 29. The case when γ and β don't have a common intersection with P.

The fact that the red point of Δ does not lie inside the triangle follows from the definitions of permissible arc and clockwise-most M-arc.

The existence of this triangle permits us to define an orientation for each crossing.

Definition 5.5. Let γ and β be non-trivial permissible clockwise-most M-arcs, and suppose they cross at a point c in their interior. Let Δ be a triangle as in Figure 28 associated to c. We say this crossing is from β to γ if γ follows β immediately in the clockwise order around the triangle.

Remark 5.6. The orientation of the crossing is well defined. Indeed, if a crossing is incident with two triangles as in Figure 28, then they are either opposite to each

other, in which case the orientation is the same for both triangles, or they are adjacent. Since P is a dissection, it follows that both triangles share a marked point in M. But then, at least one of the triangles wouldn't be of the form given by the right picture in Figure 28, a contradiction.

5.2. Support τ -tilting via crossings. Let M, N be string modules and γ_M, γ_N be the corresponding permissible clockwise-most M-arcs. We will now give a formula for the dimension of the Hom-space $\text{Hom}(M, \tau N)$ in terms of the crossings between γ_M and γ_N . This has been done in the context of skewed-gentle algebras in [31]. We note that in [31], the authors have the opposite orientation when defining the algebra from the surface. This implies that they use the anticlockwise-most arcs instead.

Lemma 5.7. Let β and γ be two permissible R-arcs and $\beta_{\mathcal{O}}$ and $\gamma_{\mathcal{O}}$ be the corresponding permissible clockwise-most M-arcs. If there is a crossing from β to γ , then there is a crossing from $\beta_{\mathcal{O}}$ to $\gamma_{\mathcal{O}}$.

Proof. This statement follows immediately from Definition 4.9 and Definition 5.1, since at least one of the segments where the crossing between β and γ occurs is an interior segment. Note that the triangle associated to the crossing from $\beta_{\mathcal{O}}$ to $\gamma_{\mathcal{O}}$ is as the one on the left in Figure 28.

Lemma 5.8. Let M and N be string A-modules, and denote by γ_M and γ_N the corresponding permissible clockwise-most M-arcs. If $\text{Hom}_A(M, \tau N) \neq 0$ then there is a crossing from γ_M to γ_N .

Proof. Since $\text{Hom}(M, \tau N) \neq 0$, we have in particular that $N = \tau^{-1}(\tau N) \neq 0$ and τN is not injective. Moreover, w_M and $w_{\tau N}$ are of the form:

$$w_M = (w_M)_{\ell} a^{-1} e b(w_M)_r$$
, and $w_{\tau N} = (w_{\tau N})_{\ell} c e d^{-1} (w_{\tau N})_r$,

with $a, b, c, d \in Q_1$ if $(w_M)_{\ell} a^{-1}, b(w_M)_r, (w_{\tau N})_{\ell} c, d^{-1}(w_{\tau N})_r \neq 0$, respectively.

In order to check whether γ_M crosses γ_N , we need to compute $\tau^{-1}(\tau N)$.

Case 1: $f_t(w_{\tau N}) = 0.$

Then $w_{\tau N}$ must be a left maximal direct path. Since τN is not injective, there must be an arrow d' such that $w_{\tau N}d'^{-1}$ is a string and so $f_s(w_{\tau N}) = w_{\tau N}d'^{-1}p_{d'}$, where $d'^{-1}p_{d'}$ is the hook of d'. It then follows that $w_N = f_t f_s(w_{\tau N}) = p_{d'}$.

Since $w_{\tau N}$ is a left maximal direct path, we must have $d^{-1}(w_{\tau N})_r = 0$ and $(w_{\tau N})_{\ell}ce = q_{d'}$, i.e. $(w_{\tau N})_{\ell}ced'^{-1}$ is the cohook of d'. Figure 30 illustrates this situation. In particular, we have indeed a crossing from γ_M to γ_N .



FIGURE 30. The case where $f_t(w_{\tau N}) = 0$.

Case 2: $f_s(w_{\tau N}) = 0$. This case is similar to the previous case. **Case 3:** $f_t(w_{\tau N}) \neq 0$ and $f_s(w_{\tau N}) \neq 0$. In this case we have $w_N = f_s(f_t(w_{\tau_N})) = f_t(f_s(w_{\tau_N}))$. We will calculate the former. *Case 3.1:* $w_{\tau N}$ is not left maximal or $(w_{\tau N})_\ell$ has a letter in Q_1^{-1} . Then $f_t(w_{\tau N}) = w' ced^{-1}(w_{\tau N})_r$, for some non-zero string w'. Case 3.1.1: $w_{\tau N}$ is not right maximal or $(w_{\tau N})_r$ has a letter in Q_1 .

Then $w_N = f_s(f_t(w_{\tau N})) = w'ced^{-1}w''$, for some non-zero string w''. In particular, there is a non-zero morphism from M to N associated to the overlap e, and e is neither a prefix nor a suffix of w_N . Hence, there is a crossing from the R-arc of M to the R-arc of N. The result then follows from Lemma 5.7.

Case 3.1.2: $(w_{\tau N})_r$ is a right maximal inverse string.

Suppose e has a letter in Q_1 and let x denote the last such letter, i.e. $ed^{-1}(w_{\tau N})_r = e'xq_x^{-1}$, where xq_x^{-1} is the cohook of x. Then $w_N = f_s(f_t(w_{\tau N})) = w'ce'$.

Therefore, we have a non-zero morphism from M to N associated to the overlap e'. Moreover, e' is not a suffix of w_M nor a prefix of w_N . Hence, there must be a crossing from the R-arc of M to the R-arc of N, and the result follows again from Lemma 5.7.

So suppose e is an inverse string. Then $ced^{-1}(w_{\tau N})_r$ is the cohook of c and $w_N = f_s(f_t(w_{\tau N})) = w'$. We have no morphism from M to N associated to e or a subpath of e, but there is a crossing from γ_M to γ_N , as illustrated in Figure 31.



FIGURE 31. The case where the cohook of c is removed on the right.

Case 3.2: $(w_{\tau N})_{\ell}$ is a left maximal direct string.

Case 3.2.1: e has a letter in Q_1^{-1} . Denote by x^{-1} the first such letter. Then $(w_{\tau N})_{\ell}e = q_x x^{-1}e'$, where xq_x^{-1} is the cohook of x. Thus $f_t(w_{\tau N}) = e'd^{-1}(w_{\tau N})_r$.

Case 3.2.1.1: $(w_{\tau N})_r$ is not right maximal or if it has a letter in Q_1 .

Then $w_N = f_s(f_t(w_{\tau N})) = e'd^{-1}w''$, for some non-zero string w''. Hence, there is a non-zero morphism from M to N associated to the overlap e'. Moreover, e' is not a suffix of w_N nor a prefix of w_M , meaning that there is a crossing from the R-arc of Mto the R-arc of N and the result follows from Lemma 5.7.

Case 3.2.1.2: $(w_{\tau N})_r$ is a right maximal inverse string.

Since $f_s(f_t(w_{\tau N})) = w_N \neq 0$, e' must have a letter in Q_1 . Let y be the last such letter, and write $e' = e''yq^{-1}$, where q is a direct path and e'' is a non-zero string. Then $yq^{-1}d^{-1}(w_{\tau N})_r$ is the cohook of y and $w_N = e''$. Again, there is a non-zero morphism from M to N associated to the overlap e'', which is neither a prefix nor a suffix of w_M , and so the result follows from Lemma 5.7.

Case 3.2.2: e is a direct path.

Since $f_t(w_{\tau N}) \neq 0$, we must have that $d^{-1}(w_{\tau N})_r$ is non-zero and so $d((w_{\tau N})_\ell ce)^{-1}$ is the cohook of d and $f_t(w_{\tau N}) = (w_{\tau N})_r$. Given that $f_s(f_t(w_{\tau N})) \neq 0$, we can either add a hook on the right of $(w_{\tau N})_r$ or $(w_{\tau N})_r$ has a letter in Q_1 . This case is similar to the case illustrated in Figure 31, and the crossing from γ_M to γ_N takes place in the tile associated to the arrow d.

Definition 5.9. Let M, N be two string modules and γ_M, γ_N be the corresponding clockwise-most M-arcs. A crossing from γ_M to γ_N which is, up to isotopy, of the form shown in Figure 32 is called a *good crossing*. Given such a crossing, we have the following conditions:



FIGURE 32. Crossings from γ_M to γ_N not associated with morphisms from M to τN .

- $w_N = w_\ell a_1 \dots a_{r-1}$, where w_ℓ is a non-zero string, and $a_1 \dots a_{r-1}$ is a direct path of length ≥ 1 ;
- there exists $a_r \in Q_1$ such that $a_1 \dots a_{r-1} a_r$ corresponds to a label;
- γ_M intersects the arc in P corresponding to $t(a_r)$.

We denote by $X_{M,N}$ the number of good crossings from γ_M to γ_N .

The following proposition gives the dimension of the Hom-space $\text{Hom}(M, \tau N)$, where M and N are string modules, in terms of crossings of clockwise-most M-arcs. The case in common with [31] is when the algebra is gentle. In this case, there are no good crossings and the dimension of $\text{Hom}(M, \tau N)$ is precisely the number of crossings from γ_M to γ_N .

Proposition 5.10. Let M, N be string A-modules, and denote by γ_M and γ_N the corresponding permissible clockwise-most M-arcs. We have

$$\dim \operatorname{Hom}_A(M, \tau N) = \mathsf{C}(\gamma_M, \gamma_N) - \mathsf{X}_{M,N}.$$

Proof. Take a crossing c from γ_M to γ_N which is not a good crossing. We will first associate a non-zero morphism from M to τN to the crossing c.

If c defines a triangle as the one on the left in Figure 28, then γ_M and γ_N have a common intersection with P and we have $w_M = (w_M)_{\ell} a^{-1} eb(w_M)_r$ and $w_N = (w_N)_{\ell} ced^{-1}(w_N)_r$, where the overlap e is defined by the crossing c as illustrated in Figure 17 and where $a, b, c, d \in Q_1$ are such that at least one of $(w_M)_{\ell} a^{-1}$ and $(w_N)_{\ell} c$ (resp. one of $b(w_M)_r$ and $d^{-1}(w_N)_r$) is non-zero.

In order to calculate $w_{\tau N}$, we add a cohook or otherwise remove a hook, both at the start and end of w_N .

Case 1: $(w_N)_{\ell} c \neq 0$ and $d^{-1}(w_N)_r \neq 0$.

In this case, e is still a substring of $w_{\tau N}$. Therefore there is a non-zero morphism from M to τN associated to the overlap e.

Case 2: $(w_N)_{\ell} c \neq 0$ but $d^{-1}(w_N)_r = 0$.

Then we have $b(w_M)_r \neq 0$. Moreover, since c is not a good crossing, $f_s(w_N)$ is obtained by adding the cohook of b at the end of w_N , i.e. $f_s(w_N) = w_N b q_b^{-1}$. Since $c \neq 0$, it follows that there is a morphism from M to τN corresponding to the overlap ebp^{-1} , where p^{-1} is a prefix of q_b^{-1} .

Case 3: $(w_N)_{\ell}c = 0$ and $d^{-1}(w_N)_r \neq 0$.

This case is similar to Case 2.

Case 4: $(w_N)_{\ell}c = 0$ and $d^{-1}(w_N)_r = 0$.

Then we have $a, b \neq 0$ and $a^{-1}eb$ is a string. Therefore $w_{\tau N} = q_a a^{-1}ebq_b^{-1}$ and we have a non-zero morphism from M to τN with associated overlap $p_1 a^{-1}ebp_2^{-1}$, where p_1 is a suffix of q_a and p_2^{-1} is a prefix of q_b^{-1} .

Now suppose c defines a triangle as the one on the right in Figure 28. The only case which is not isotopic to the crossings analysed above is presented in Figure 33.



FIGURE 33. A crossing from γ_M to γ_N not defining a triangle as the one on the left in Figure 28.

In this case, we have an arrow $a \in Q_1$ such that $s(a) = t(w_N)$ and $w_N a$ is a string, since the *c* is not a good crossing. Therefore $f_s(w_N)$ is obtained by adding the cohook aq_a^{-1} on the right of w_N , and q_a^{-1} a substring of $w_{\tau N}$. Hence, there is a non-zero morphism from *M* to τN associated to an overlap which is a prefix of q_a^{-1} .

Now let e be an overlap giving rise to a non-zero morphism from M to τN . By Lemma 5.8, there is a crossing from M to N. If it is a good crossing then one of the segments must be an end-segment. The only cases in the proof of Lemma 5.8 where this occurs are the cases 1, 2, 3.1.2 and 3.2.2. But as we can see from Figures 30 and 31, which illustrate these cases, the corresponding crossing is not of form given in Figure 32.

It is clear that the mappings defined above between basis elements of $\text{Hom}(M, \tau N)$ and crossings from γ_M to γ_N which are not good are inverse of each other, and so the result follows.

The following corollary follows immediately from Proposition 5.10 since each crossing has a unique orientation.

Corollary 5.11. Let M, N be string A-modules, and denote by γ_M and γ_N the corresponding permissible clockwise-most M-arcs. We have

 $\dim \operatorname{Hom}_{A}(M, \tau N) + \dim \operatorname{Hom}_{A}(N, \tau M) = \mathsf{C}(\gamma_{M}, \gamma_{N}) + \mathsf{C}(\gamma_{N}, \gamma_{M}) - \mathsf{X}_{M,N} - \mathsf{X}_{N,M}.$

Definition 5.12. A generalised permissible arc is either an arc in P or a permissible clockwise-most M-arc.

We are now ready to give a combinatorial classification of support τ -tilting modules over a string algebra, which recovers the case where the algebra is gentle (cf. [31]). We note that there is a different classification of support τ -tilting modules over string algebras in [25].

Theorem 5.13. Let A be a string algebra. There is a one-to-one correspondence between the set of support τ -tilting pairs and the set of maximal collections of generalised permissible arcs for which each crossing is good.

Proof. This statement follows immediately from Proposition 5.10 and Proposition 5.3. The arcs in P correspond to the summands of the projective module in the pair and the remaining arcs correspond to the summands of the τ -rigid module in the pair. \Box

Remark 5.14. We can reformulate this classification in terms of R-arcs as follows. A collection of arcs in P together with permissible R-arcs corresponds to a support τ -tilting pair if and only if it is maximal with respect to the following properties:

- (1) all crossings are of the form given in Figure 34, and
- (2) if two R-arcs satisfy the condition shown in Figure 35, then there is a label as in Figure 36.



FIGURE 34. Crossings between R-arcs allowed in support τ -tilting modules.



FIGURE 35. Two local non-crossing configurations which are not allowed unless we are in the situation of Figure 36.



FIGURE 36. A local configuration which is allowed.

Remark 5.15. Consider the algebra G_1 given by the quiver $1 \xrightarrow{a} 2 \xrightarrow{b} 3$, bound by the relation ab. Then G_1 is gentle and it is the tiling algebra of the surface in Figure 37.

Consider the projective module P_2 and the simple module S_1 . We have that $P_2 \oplus S_1$ is not τ -rigid, since $\operatorname{Hom}_{G_1}(P_2, \tau S_1) \neq 0$. In fact, the dimension of $\operatorname{Hom}_{G_1}(P_2, \tau S_1)$ is one.

On the one hand, we have no crossings or intersections between the permissible Rarcs associated to P_2 and S_1 , and on the other hand, as expected from Proposition 5.10,



FIGURE 37. Surface with tiling algebra G_1 .

there is one crossing between the corresponding clockwise-most permissible M-arcs; see Figure 38.



FIGURE 38. The R-arcs and clockwise-most M-arcs associated to S_1 and P_2 .

Now consider the algebra G_2 given by the quiver $1 \xrightarrow{a} 2 \xrightarrow{b} 3 \xrightarrow{c} 4$ bound by the relation *abc*. Figure 39 shows the R-arcs associated to the projective modules P_1 and P_2 . We clearly have that $P_1 \oplus P_2$ is a τ -rigid G_2 -module.



FIGURE 39. The surface associated to G_2 and the R-arcs corresponding to ab and bc.

These examples illustrate the reason why we consider M-arcs to describe the Auslander-Reiten translate and classify support τ -tilting modules. On the one hand, M-arcs allow us to avoid considering local configurations which are not given by crossings or intersections. On the other hand, using R-arcs in the description of τ -rigidity does not exclude certain crossings.

Example 5.16. Consider the algebra given in Example 2.12. In Figure 40, we have the collection of clockwise-most M-arcs together with the highlighted arc in P associated to the support τ -tilting pair $(P_3, S_1 \oplus M(a_5) \oplus M(a_5^{-1}a_6) \oplus M(a_1a_2))$. Note that there is a self-crossing of the arc associated to $M(a_1a_2)$. However, this crossing is of the form shown in Figure 32, and indeed $M(a_1a_2) = P_1$ is a τ -rigid module.



FIGURE 40. An example of a support τ -tilting pair.

References

- Takahide Adachi, Osamu Iyama, and Idun Reiten. τ-tilting theory. Compos. Math., 150(3):415– 452, 2014.
- [2] Claire Amiot and Yvonne Grimeland. Derived invariants for surface algebras. J. Pure Appl. Algebra, 220(9):3133–3155, 2016.
- [3] Claire Amiot, Daniel Labardini-Fragoso, and Pierre-Guy Plamondon. Derived invariants for surface cut algebras of global dimension 2 II: the punctured case. *Comm. Algebra*, 49(1):114– 150, 2021.
- [4] Claire Amiot, Pierre-Guy Plamondon, and Sibylle Schroll. A complete derived invariant for gentle algebras via winding numbers and Arf invariants. *Selecta Math. (N.S.)*, 29(2):Paper No. 30, 36, 2023.
- [5] Ibrahim Assem, Thomas Brüstle, Gabrielle Charbonneau-Jodoin, and Pierre-Guy Plamondon. Gentle algebras arising from surface triangulations. Algebra Number Theory, 4(2):201–229, 2010.
- [6] Ibrahim Assem and Andrzej Skowroński. Iterated tilted algebras of type \mathbf{A}_n . Math. Z., 195(2):269–290, 1987.
- [7] Esther Banaian, Raphael Bennett-Tennenhaus, Karin M. Jacobsen, and Kayla Wright. A geometric model for semilinear locally gentle algebras. 2024.
- [8] Emily Barnard, Raquel Coelho Simões, Emily Gunawan, and Ralf Schiffler. Maximal almost rigid modules over gentle algebras. *in preparation*, 2024.
- [9] Emily Barnard, Emily Gunawan, Emily Meehan, and Ralf Schiffler. Cambrian combinatorics on quiver representations (type A). Adv. in Appl. Math., 143:Paper No. 102428, 37, 2023.
- [10] Karin Baur and Raquel Coelho Simões. A geometric model for the module category of a gentle algebra. Int. Math. Res. Not. IMRN, (15):11357–11392, 2021.
- [11] Karin Baur and Raquel Coelho Simões. Corrigendum: A geometric model for the module category of a gentle algebra. *International Mathematics Research Notices*, 2023(7):6291–6298, 01 2023.
- [12] Karin Baur and Bethany R. Marsh. A geometric description of m-cluster categories. Trans. Amer. Math. Soc., 360(11):5789–5803, 2008.
- [13] Viktor Bekkert and Héctor A. Merklen. Indecomposables in derived categories of gentle algebras. Algebr. Represent. Theory, 6(3):285–302, 2003.
- [14] Nathan Broomhead. Dimer models and Calabi-Yau algebras. Mem. Amer. Math. Soc., 215(1011):viii+86, 2012.
- [15] Thomas Brüstle, Guillaume Douville, Kaveh Mousavand, Hugh Thomas, and Emine Yıldı rım. On the combinatorics of gentle algebras. *Canad. J. Math.*, 72(6):1551–1580, 2020.
- [16] Thomas Brüstle and Jie Zhang. On the cluster category of a marked surface without punctures. Algebra Number Theory, 5(4):529–566, 2011.
- [17] Michael C. R. Butler and Claus Michael Ringel. Auslander-Reiten sequences with few middle terms and applications to string algebras. Comm. Algebra, 15(1-2):145–179, 1987.
- [18] Philippe Caldero, Frédéric Chapoton, and Ralf Schiffler. Quivers with relations arising from clusters (A_n case). Trans. Amer. Math. Soc., 358(3):1347–1364, 2006.
- [19] Ilke Çanakçı, David Pauksztello, and Sibylle Schroll. On extensions for gentle algebras. Canad. J. Math., 73(1):249–292, 2021.

- [20] Wen Chang. Geometric models for the hearts in the derived category of a gentle algebra, 2023.
- [21] Wen Chang, Haibo Jin, and Sibylle Schroll. Recollements of partially wrapped fukaya categories and surface cuts, 2023.
- [22] Wen Chang and Sibylle Schroll. A geometric realization of silting theory for gentle algebras. Math. Z., 303(3):Paper No. 67, 37, 2023.
- [23] William W. Crawley-Boevey. Maps between representations of zero-relation algebras. J. Algebra, 126(2):259–263, 1989.
- [24] Lucas David-Roesler. Derived equivalence of surface algebras in genus 0 via graded equivalence. Algebr. Represent. Theory, 17(1):1–30, 2014.
- [25] Florian Eisele, Geoffrey Janssens, and Theo Raedschelders. A reduction theorem for τ-rigid modules. Math. Z., 290(3-4):1377–1413, 2018.
- [26] Sergey Fomin, Michael Shapiro, and Dylan Thurston. Cluster algebras and triangulated surfaces.
 I. Cluster complexes. Acta Math., 201(1):83–146, 2008.
- [27] Christof Geiß and Idun Reiten. Gentle algebras are Gorenstein. In Representations of algebras and related topics, volume 45 of Fields Inst. Commun., pages 129–133. Amer. Math. Soc., Providence, RI, 2005.
- [28] Fabian Haiden, Ludmil Katzarkov, and Maxim Kontsevich. Flat surfaces and stability structures. Publ. Math. Inst. Hautes Études Sci., 126:247–318, 2017.
- [29] Johanne Haugland, Karin M. Jacobsen, and Sibylle Schroll. The role of gentle algebras in higher homological algebra. *Forum Math.*, 34(5):1255–1275, 2022.
- [30] Ping He, Yu Zhou, and Bin Zhu. Mutation graph of support τ -tilting modules over a skew-gentle algebra, 2022.
- [31] Ping He, Yu Zhou, and Bin Zhu. A geometric model for the module category of a skew-gentle algebra. *Math. Z.*, 304(1):Paper No. 18, 41, 2023.
- [32] Ruth Stella Huerfano and Mikhail Khovanov. A category for the adjoint representation. J. Algebra, 246(2):514–542, 2001.
- [33] Haibo Jin, Sibylle Schroll, and Zhengfang Wang. A complete derived invariant and silting theory for graded gentle algebras, 2023.
- [34] Henning Krause. Maps between tree and band modules. J. Algebra, 137(1):186–194, 1991.
- [35] Daniel Labardini-Fragoso. Quivers with potentials associated to triangulated surfaces. Proc. Lond. Math. Soc. (3), 98(3):797–839, 2009.
- [36] Daniel Labardini-Fragoso, Sibylle Schroll, and Yadira Valdivieso. Derived categories of skewgentle algebras and orbifolds. *Glasg. Math. J.*, 64(3):649–674, 2022.
- [37] YankıLekili and Alexander Polishchuk. Derived equivalences of gentle algebras via Fukaya categories. Math. Ann., 376(1-2):187–225, 2020.
- [38] Yu-Zhe Liu and Yu Zhou. A negative answer to complement question for presilting complexes, 2023.
- [39] Sebastian Opper. On auto-equivalences and complete derived invariants of gentle algebras, 2019.
- [40] Sebastian Opper, Pierre-Guy Plamondon, and Sibylle Schroll. A geometric model for the derived category of gentle algebras, 2018.
- [41] Sebastian Opper and Alexandra Zvonareva. Derived equivalence classification of Brauer graph algebras. *Adv. Math.*, 402:Paper No. 108341, 59, 2022.
- [42] Yann Palu, Vincent Pilaud, and Pierre-Guy Plamondon. Non-kissing complexes for gentle algebras. Sém. Lothar. Combin., 80B:Art. 39, 12, 2018.
- [43] Yann Palu, Vincent Pilaud, and Pierre-Guy Plamondon. Non-kissing and non-crossing complexes for locally gentle algebras. J. Comb. Algebra, 3(4):401–438, 2019.
- [44] Jan Schröer and Alexander Zimmermann. Stable endomorphism algebras of modules over special biserial algebras. Math. Z., 244(3):515–530, 2003.

Email address: ka.baur@me.com

FAKULTÄT FÜR MATHEMATIK, RUHR-UNIVERSITÄT BOCHUM

School of Mathematics, University of Leeds

Email address: r.coelhosimoes@lancaster.ac.uk

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF LANCASTER