

# On $t$ -structures adjacent and orthogonal to weight structures

Mikhail V. Bondarko \*

March 13, 2024

## Abstract

We study  $t$ -structures (on triangulated categories) that are closely related to weight structures. A  $t$ -structure couple  $t = (\underline{\mathcal{C}}_{t \leq 0}, \underline{\mathcal{C}}_{t \geq 0})$  is said to be adjacent to a weight structure  $w = (\underline{\mathcal{C}}_{w \leq 0}, \underline{\mathcal{C}}_{w \geq 0})$  if  $\underline{\mathcal{C}}_{t \geq 0} = \underline{\mathcal{C}}_{w \geq 0}$ . For a triangulated category  $\underline{\mathcal{C}}$  that satisfies the Brown representability property we prove that  $t$  that is adjacent to  $w$  exists if and only if  $w$  is *smashing* (i.e.,  $\underline{\mathcal{C}}_{w \geq 0}$  is  $\coprod_{\underline{\mathcal{C}}}$ -closed). The heart  $Ht$  of this  $t$  is the category of those functors  $Hw^{op} \rightarrow \mathbf{Ab}$  that respect products (here  $Hw$  is the heart of  $w$ ); the result has important applications. We prove several more statements on constructing  $t$ -structures starting from weight structures; we look for a *strictly orthogonal*  $t$ -structure  $t$  on some  $\underline{\mathcal{C}}'$  (where  $\underline{\mathcal{C}}, \underline{\mathcal{C}}'$  are triangulated subcategories of a common  $\underline{\mathcal{D}}$ ) such that  $\underline{\mathcal{C}}'_{t \leq 0}$  (resp.  $\underline{\mathcal{C}}'_{t \geq 0}$ ) is characterized by the vanishing of morphisms from  $\underline{\mathcal{C}}_{w \geq 1}$  (resp.  $\underline{\mathcal{C}}_{w \leq -1}$ ). Some of these results generalize properties of semi-orthogonal decompositions proved in the previous paper, and can be applied to various derived categories of (quasi)coherent sheaves on a scheme  $X$  that is projective over an affine noetherian one. We also study hearts of orthogonal  $t$ -structures and their restrictions, and prove some statements on "reconstructing" weight structures from orthogonal  $t$ -structures.

The main tool of this paper is the notion of virtual  $t$ -truncations of (cohomological) functors; these are defined in terms of weight structures and "behave as if they come from  $t$ -truncations of representing objects" whether  $t$  exists or not.

**2020 Mathematics Subject Classification.** 18G80, 14F08 (Primary) 18F20, 18G05, 18E10, 14A15, 14G40 (Secondary)

**Key words and phrases** Triangulated category, weight structure,  $t$ -structure, virtual  $t$ -truncation, adjacent structures, orthogonal structures, pure functor, coherent sheaves, perfect complexes, saturated category, coproductive extension.

## Contents

<b>1</b>	<b>A reminder on <math>t</math>-structures and weight structures</b>	<b>5</b>
1.1	Some categorical and $t$ -structure notation . . . . .	5
1.2	Some basics on weight structures . . . . .	9
1.3	On orthogonal and adjacent structures . . . . .	12

---

\*The work is supported by the Leader (Leading scientist Math) grant no. 22-7-1-13-1.

<b>2</b>	<b>On virtual <math>t</math>-truncations and their relation to orthogonal <math>t</math>-structures</b>	<b>13</b>
2.1	Virtual $t$ -truncations of (co)homological functors . . . . .	14
2.2	On weight range . . . . .	16
2.3	Some properties and restrictions of strictly orthogonal $t$ -structures	20
2.4	On coproductive extensions of weight structures . . . . .	22
<b>3</b>	<b>On <math>t</math>-structures orthogonal to (co)smashing weight structures</b>	<b>23</b>
3.1	On smashing weight structures and related notions . . . . .	24
3.2	On the existence of $t$ -structures adjacent to smashing weight structures	25
3.3	Weight structures extended from subcategories of compact objects, and orthogonal $t$ -structures	27
3.4	$t$ -structures anti-orthogonal to cosmashing weight structures . . .	30
<b>4</b>	<b>On functors and <math>t</math>-structures related to (essentially) bounded weight structures</b>	<b>32</b>
4.1	On essentially bounded weight structures and the relation to cohomological functors	33
4.2	More on the construction of $t$ -structures . . . . .	36
<b>5</b>	<b>The existence of orthogonal weight structures</b>	<b>39</b>
5.1	On hearts and $\pm$ -orthogonality of structures . . . . .	39
5.2	On the existence of orthogonal weight structures . . . . .	41
5.3	Some adjacent weight structures in the saturated case . . . . .	45

## Introduction

This paper is devoted to the study of those  $t$ -structures that are "closely related" to weight structures (on various triangulated categories).

Let us recall that  $t$ -structures on triangulated categories have become important tools in homological algebra since their introduction in [BBD82]. Respectively, their study and construction is an actual and non-trivial question. Next, in [Pau08] and [Bon10a] a rather similar notion of a weight structure  $w$  on a triangulated category  $\underline{C}$  was introduced. Moreover, in *ibid.* a  $t$ -structure  $t = (\underline{C}_{t \leq 0}, \underline{C}_{t \geq 0})$  was said to be *adjacent* to  $w$  if  $\underline{C}_{t \geq 0} = \underline{C}_{w \geq 0}$ , and certain examples of adjacent structures were constructed.<sup>1</sup> Furthermore, in [Bon10b] for a  $t$ -structure  $t$  on a triangulated category  $\underline{C}'$  that is related to  $\underline{C}$  by means of a *duality* bi-functor a more general notion of *orthogonality* of a weight structure  $w$  on  $\underline{C}$  to  $t$  was introduced. Also, the relationship between the hearts of adjacent and orthogonal structures was studied in detail.

Next, if  $w$  is adjacent to  $t$  then it determines  $t$  uniquely and vice versa. Yet the only previously existing way of constructing  $t$  if  $w$  is given was to use certain "nice generators" of  $w$  (see Definition 3.2.1(5), Theorem 3.2.2 of [Bon21], and of [Bon22a]). However, already in [Bon10a] the notion of virtual  $t$ -truncations for (co)homological functors was introduced, and it was proved that virtual  $t$ -truncations possess several nice properties. In particular, it was demonstrated that these are closely related to  $t$ -structures (whence the name) even though they are defined in terms of weight structures only.

In the current paper we propose a new construction method. We prove that adjacent and orthogonal  $t$ -structures can be constructed using virtual  $t$ -truncations whenever certain "Brown representability-type" assumptions on  $\underline{C}$

---

<sup>1</sup>In contrast to *ibid.* and [Bon10b], in the current paper we use the so-called homological convention for  $t$  and  $w$ , and say that  $t$  is left adjacent to  $w$  if  $\underline{C}_{t \geq 0} = \underline{C}_{w \geq 0}$ .

(and  $\underline{C}'$ ) are known. Respectively, our results yield the existence of some new families of  $t$ -structures.

Let us formulate one of these results. For a triangulated category  $\underline{C}$  that is *smashing*, i.e., closed with respect to (small) coproducts, and a weight structure  $w$  on it we will say that  $w$  is *smashing* whenever  $\underline{C}_{w \geq 0}$  is closed with respect to  $\underline{C}$ -coproducts (note that  $\underline{C}_{w \leq 0}$  is  $\coprod$ -closed automatically).

**Theorem 0.1** (See Theorem 3.2.3). Let  $\underline{C}$  be a smashing triangulated category that satisfies the following Brown representability property: any functor  $\underline{C}^{op} \rightarrow \mathbf{Ab}$  that respects  $(\underline{C}^{op})$ -products is representable.

Then for a weight structure  $w$  on  $\underline{C}$  there exists a  $t$ -structure  $t$  adjacent to it if and only if  $w$  is smashing. Moreover, the heart of  $t$  (if  $t$  exists) is equivalent to the category of all those additive functors  $\underline{H}w^{op} \rightarrow \mathbf{Ab}$  that respect products; here  $\underline{H}w$  is the *heart* of  $w$ .

Note here that (smashing) triangulated categories satisfying the Brown representability property are currently really popular in several areas of mathematics (thanks to the foundational results of A. Neeman and others); in particular, this property holds if either  $\underline{C}$  or  $\underline{C}^{op}$  is *compactly generated*. Moreover, it is easy to construct vast families of smashing weight structures on  $\underline{C}$  (at least) if  $\underline{C}$  is compactly generated; see Remark 3.2.4(1) below.

Certainly the dual to Theorem 0.1 is valid as well. Moreover, if  $\underline{C}^{op}$  satisfies the dual Brown representability property and  $w$  is both cosmashing and smashing then the right adjacent  $t$ -structure  $t$  (i.e.,  $\underline{C}_{t \leq 0} = \underline{C}_{w \leq 0}$ ) restricts to the subcategory of compact objects of  $\underline{C}$  as well as to all other "levels of smallness" for objects (see Theorem 3.4.1(3)). Combining this statement with an existence of weight structures Theorem 3.1 of [KeN13] we obtain a statement on  $t$ -structures extending Theorem 7.1 of *ibid*.

We also prove some alternative versions of Theorem 0.1 that can be applied to "quite small" triangulated categories. Instead of the Brown representability condition for  $\underline{C}$  one can demand it to satisfy the *R-saturatedness* one instead (see Definition 4.2.1(2) below; this is an "*R*-linear finite" version of the Brown representability). Then for any *bounded*  $w$  on  $\underline{C}$  there will exist a  $t$ -structure adjacent to it. Moreover, this statement also holds for *essentially bounded weight structures*; those generalize both bounded weight structures and semi-orthogonal decompositions. According to a saturatedness statement from [Nee18] this result can be applied to the derived category  $D^{perf}(X)$  of perfect complexes on a regular scheme that is proper over the spectrum of a Noetherian ring  $R$  (see Corollary 4.2.2(3)). We also study restrictions of orthogonal  $t$ -structures to subcategories corresponding to certain bounds and support conditions; see Propositions 2.3.5 and 3.3.4, Corollaries 4.2.2(2) and 4.2.5, and Theorem 4.2.4. We demonstrate that this result can be applied to certain derived categories of (quasi)coherent sheaves; this includes the case of a singular  $X$  (see Propositions 3.3.5 and 4.2.6).

Lastly, we try to prove some converse of these results. So we study the question when a (fixed)  $t$ -structure  $t$  on a triangulated category  $\underline{C}$  is orthogonal to a weight structure  $w$  on a "large enough" subcategory  $\underline{C}' \subset \underline{C}$ . Roughly, this is the case if and only if the heart  $\underline{H}t$  has enough projectives and those "lift" to  $\underline{H}w$  (and the latter condition follows from duals to certain Brown representability-type conditions). However, it appears to be difficult to find the conditions that ensure the existence of an adjacent weight structure (so, it is on  $\underline{C}$  itself). We prove some statements of this sort; yet these results are more

difficult to apply than the aforementioned "converse" ones, and the methods of their proofs are less original.

*Remark 0.2.* The author certainly does not claim that the methods of the current paper are the most general among the existing methods for constructing  $t$ -structures. In particular, if  $\underline{C}$  is a *well generated* triangulated category (categories of this sort are "very common") then the recent Theorem 2.3 of [Nee21] gives all those  $t$ -structures that are generated by sets of objects of  $\underline{C}$  in the sense of Definition 3.2.1(5) below.

Now, if  $\underline{C}$  is well generated then is generated by a set of its objects (as its own localizing subcategory; see Definition 3.2.1(3) below). Thus for any smashing weight structure  $w$  on it Proposition 2.3.2(10) of [Bon21] essentially says the following: there exists a set  $\mathcal{P} \subset \underline{C}_{w \geq 0}$  such that the class  $\underline{C}_{w \geq 0}$  is the smallest cocomplete pre-aisle that is "generated" by  $\mathcal{P}$  in the sense of [Nee21, §0] (cf. Discussion 1.16 of *ibid.*). Hence Theorem 2.3 of [Nee21] says that there exists a  $t$ -structure on  $\underline{C}$  such that  $\underline{C}_{t \geq 0} = \underline{C}_{w \geq 0}$  (cf. Remark 1.1.6(2) below). Thus *loc. cit.* generalizes the existence of  $t$  part of our Theorem 0.1.

On the other hand, note that neither *loc. cit.* nor the well-known Theorem A.1 of [AJS03] says anything on the hearts of  $t$ -structures. Moreover, there appears to be no chance to extend these existence results to  $R$ -saturated categories (in any way).

Let us now describe the contents of the paper. Some more information of this sort may be found in the beginnings of sections.

In §1 we give some definitions and conventions, and recall some basics on  $t$ -structures, weight structures, and semi-orthogonal decompositions.

In §2 we define virtual  $t$ -truncations of functors and prove several nice properties for them. We also relate the existence of orthogonal  $t$ -structures to virtual  $t$ -truncations; this gives general if and only if criteria for the existence of adjacent  $t$ -structures.

In §3 we study smashing triangulated categories along with the existence of  $t$ -structures adjacent to weight structures on them and certain restrictions of these  $t$ -structures. We also consider weight structures extended from subcategories of compact objects.

In §4 we discuss certain bounded versions of the results of the previous section. Our main statement concern the existence of adjacent and orthogonal  $t$ -structures in  $R$ -saturated triangulated categories and the restriction of orthogonal  $t$ -structures to subcategories corresponding to certain bounds and support conditions. We apply our general statements to various derived categories of (quasi)coherent sheaves. To formulate our statements in a more general case we recall the notions of essentially bounded (above, below, or both) objects and weight structures. In particular, we are able to generalize the central results of [Bon24].

In §5 we try to answer the question which  $t$ -structures are adjacent to weight structures. In certain situations we prove that a weight structure adjacent to a given  $t$  exists whenever  $\underline{Ht}$  has enough projectives and certain additional assumptions are fulfilled.

The author is deeply grateful to prof. A. Neeman for calling his attention to [KeN13] as well as for writing his extremely interesting texts that are crucial for the current paper.

*Remark 0.3.* 1. The current text is a certain modification of the preprint [Bon19]. The main idea is to avoid the somewhat ad hoc (and restrictive) notion of reflection of categories (see Definition 2.2.1 of *ibid.*); cf. Remark 2.4.5 below. Moreover, several new general results were added; this includes the theory of coproductive extensions, (most of) §4, §5.3, and subcategories corresponding to certain support conditions (see Propositions 2.3.5, 3.3.4, 3.3.5, and 4.2.6(2), and Corollary 4.2.2(2)).

On the other hand, the current text does not include Proposition 2.2.7, Corollary 4.1.4(2), and §4.4 of *ibid.* Probably, these statements will be included into a succeeding paper.

Note also that the terminology of the current text is substantially distinct from the one of *ibid.*

2. Since [Bon19] was already cited in some papers, it makes sense to note the following: Theorem 3.2.3(I) of *ibid.* coincides with Theorem 0.1 of the current paper, Theorem 3.3.1 of *ibid.* almost coincides with Theorem 3.3.2 below, Theorem 5.3.1(I.1) of *ibid.* is contained in Proposition 5.2.1, Theorem 5.3.1(IV) of *ibid.* was generalized to Theorem 5.3.3, and strongly  $\aleph_0$ -well generated weight structures (that were treated in §3.3 of [Bon19]) are discussed in §3.3 (yet see Remark 3.3.3(1) below).

3. The work on [Bon19] inspired the author to write [Bon24]. We note that semi-orthogonal decompositions (that are the main subject of *ibid.*) are particular case of weight structures; see Proposition 1.2.6 below for more detail. Next, both in [Bon24] and in (the justification of) Remark 4.2.7(1) below the main idea is to extend a weight structure to a bigger (smashing) category and to pass to an orthogonal  $t$ -structure (see Definitions 1.2.2(4) and 1.3.1 below). It is no wonder that applying these methods in the setting of (general) weight structures is somewhat more difficult than doing this for semi-orthogonal decompositions. On the other hand, these difficulties make the current paper somewhat more interesting and also "more original".

## 1 A reminder on $t$ -structures and weight structures

In this section we recall the notions of  $t$ -structures and weight structures, along with orthogonality and adjacency for them.

In §1.1 we introduce some categorical notation and recall some basics on  $t$ -structures.

In §1.2 we recall some of the theory of weight structures and semi-orthogonal decompositions.

In §1.3 we recall the definitions of adjacent and orthogonal weight and  $t$ -structures that are central for this paper.

### 1.1 Some categorical and $t$ -structure notation

- All products and coproducts in this paper will be small.
- Given a category  $C$  and  $X, Y \in \text{Obj } C$  we will write  $C(X, Y)$  for the set of morphisms from  $X$  to  $Y$  in  $C$ .

- All subcategories in this paper will be strictly full. Respectively, for categories  $C', C$  we write  $C' \subset C$  if  $C'$  is a strictly full subcategory of  $C$ .
- For  $S_1, S_2 \subset C$  we will write  $S_1 \cap S_2$  for the (strictly full) subcategory of  $C$  whose object class equals  $\text{Obj } S_1 \cap \text{Obj } S_2$ .
- Given a category  $C$  and  $X, Y \in \text{Obj } C$ , we say that  $X$  is a *retract* of  $Y$  if  $\text{id}_X$  can be factored through  $Y$ .
- Given an additive subcategory  $\underline{H}$  of an additive  $C$ , the (strictly full) subcategory  $\text{Kar}_C(\underline{H})$  of  $C$  whose objects are all  $C$ -retracts of objects of  $\underline{H}$  will be called *the retraction-closure* of  $\underline{H}$  in  $C$ .  
 $\underline{H}$  is said to be *retraction-closed* in  $C$  if it coincides with  $\text{Kar}_C(\underline{H})$ .
- We will say that  $C$  is *Karoubian* if any idempotent morphism yields a direct sum decomposition in it.
- The symbol  $\underline{C}$  below will always denote some triangulated category; it will often be endowed with a weight structure  $w$ . The symbols  $\underline{C}', \underline{D}$ , and  $\underline{C}^0$  are also reserved for triangulated categories only. Moreover, we will often assume  $\underline{C}, \underline{C}' \subset \underline{D}$  and  $\underline{C}^0 \subset \underline{C}$ .
- For any  $A, B, C \in \text{Obj } \underline{C}$  we will say that  $C$  is an *extension* of  $B$  by  $A$  if there exists a distinguished triangle  $A \rightarrow C \rightarrow B \rightarrow A[1]$ .
- For any  $D, E \subset \text{Obj } \underline{C}$  we will write  $D \star E$  for the class of all extensions of elements of  $E$  by elements of  $D$ .
- A class  $\mathcal{P} \subset \text{Obj } \underline{C}$  is said to be *extension-closed* if it is closed with respect to extensions and contains 0.
- The smallest additive retraction-closed extension-closed class of objects of  $\underline{C}$  containing  $\mathcal{P}$  will be called the *envelope* of  $\mathcal{P}$ .
- For  $X, Y \in \text{Obj } \underline{C}$  we will write  $X \perp Y$  if  $\underline{C}(X, Y) = \{0\}$ .  
For  $D, E \subset \text{Obj } \underline{C}$  we write  $D \perp E$  if  $X \perp Y$  for all  $X \in D, Y \in E$ .  
Given  $D \subset \text{Obj } \underline{C}$  we will write  $D^\perp$  for the class

$$\{Y \in \text{Obj } \underline{C} : X \perp Y \ \forall X \in D\}.$$

Dually,  ${}^\perp D$  is the class  $\{Y \in \text{Obj } \underline{C} : Y \perp X \ \forall X \in D\}$ .

- Given  $f \in \underline{C}(X, Y)$ , where  $X, Y \in \text{Obj } \underline{C}$ , we will call the third vertex of (any) distinguished triangle  $X \xrightarrow{f} Y \rightarrow Z$  a *cone* of  $f$ .
- Below the symbols  $\underline{A}, \underline{A}^0$ , and  $\underline{A}'$  will always denote some abelian categories;  $\underline{B}$  is an additive category.
- We will say that  $\underline{A}^0 \subset \underline{A}$  is an abelian subcategory of  $\underline{A}$  if  $\underline{A}$  is its (strictly full) subcategory that contains the  $\underline{A}$ -kernel and the  $\underline{A}$ -cokernel of any  $\underline{A}^0$ -morphism.

Moreover, we will say that  $\underline{A}^0$  is a *weak Serre* subcategory of  $\underline{A}$  if  $\underline{A}^0$  is an abelian subcategory that is closed with respect to  $\underline{A}$ -extensions.

- All complexes in this paper will be cohomological.
  - We will say that an additive covariant (resp. contravariant) functor from  $\underline{\mathcal{C}}$  into  $\underline{\mathcal{A}}$  is *homological* (resp. *cohomological*) if it converts distinguished triangles into long exact sequences.
- For a (co)homological functor  $H$  and  $i \in \mathbb{Z}$  we will write  $H_i$  (resp.  $H^i$ ) for the composition  $H \circ [-i]$ .

*Remark 1.1.* Obviously, if  $\underline{\mathcal{C}}_1, \underline{\mathcal{C}}_2$  are triangulated subcategories of  $\underline{\mathcal{C}}$  then  $\underline{\mathcal{C}}_1 \cap \underline{\mathcal{C}}_2$  is its triangulated subcategory as well; recall the strictness assumption.

Similarly, if  $\underline{\mathcal{A}}_1, \underline{\mathcal{A}}_2$  are weak Serre subcategory of  $\underline{\mathcal{A}}$  then  $\underline{\mathcal{A}}_1 \cap \underline{\mathcal{A}}_2$  is its weak Serre subcategory as well.

To generalize our formulations and relate them to the results of [Bon10b] we recall Definition 2.5.1(1) of *ibid*.

**Definition 1.1.1.** We will call a (covariant) bi-functor  $\Phi : \underline{\mathcal{C}}^{op} \times \underline{\mathcal{C}}' \rightarrow \underline{\mathcal{A}}$  a *duality* between  $\underline{\mathcal{C}}$  and  $\underline{\mathcal{C}}'$  if  $\Phi$  is homological with respect to both arguments; and is equipped with a (bi)natural bi-additive transformation  $\Phi(X, Y) \cong \Phi(X[1], Y[1])$ .

The following example of a duality will be the main one throughout this paper.

**Lemma 1.1.2.** Assume  $\underline{\mathcal{C}}, \underline{\mathcal{C}}' \subset \underline{\mathcal{D}}$ . Then the restriction of the bi-functor  $\underline{\mathcal{D}}(-, -)$  to  $\underline{\mathcal{C}}^{op} \times \underline{\mathcal{C}}'$  gives a duality  $\underline{\mathcal{C}}^{op} \times \underline{\mathcal{C}}' \rightarrow \text{Ab}$ .

*Proof.* This statement is obvious; it also easily follows from Proposition 2.5.6(1) of [Bon10b].  $\square$

Let us also recall the notion of a  $t$ -structure (mainly to fix notation).

**Definition 1.1.3.** A couple of subclasses  $\underline{\mathcal{C}}_{t \leq 0}, \underline{\mathcal{C}}_{t \geq 0} \subset \text{Obj } \underline{\mathcal{C}}$  will be said to be a  $t$ -structure  $t$  on  $\underline{\mathcal{C}}$  if they satisfy the following conditions:

- (i)  $\underline{\mathcal{C}}_{t \leq 0}$  and  $\underline{\mathcal{C}}_{t \geq 0}$  are strict, i.e., contain all objects of  $\underline{\mathcal{C}}$  isomorphic to their elements.
- (ii)  $\underline{\mathcal{C}}_{t \leq 0} \subset \underline{\mathcal{C}}_{t \leq 0}[1]$  and  $\underline{\mathcal{C}}_{t \geq 0}[1] \subset \underline{\mathcal{C}}_{t \geq 0}$ .
- (iii)  $\underline{\mathcal{C}}_{t \geq 0}[1] \perp \underline{\mathcal{C}}_{t \leq 0}$ .
- (iv) For any  $M \in \text{Obj } \underline{\mathcal{C}}$  there exists a  $t$ -decomposition distinguished triangle

$$L_t M \rightarrow M \rightarrow R_t M \rightarrow L_t M[1] \quad (1.1.1)$$

such that  $L_t M \in \underline{\mathcal{C}}_{t \geq 0}, R_t M \in \underline{\mathcal{C}}_{t \leq 0}[-1]$ .

We will also give some auxiliary definitions.

**Definition 1.1.4.** 1. For any  $i \in \mathbb{Z}$  we will use the notation  $\underline{\mathcal{C}}_{t \leq i}$  (resp.  $\underline{\mathcal{C}}_{t \geq i}$ ) for the class  $\underline{\mathcal{C}}_{t \leq 0}[i]$  (resp.  $\underline{\mathcal{C}}_{t \geq 0}[i]$ ).

2.  $\underline{Ht}$  is the (full) subcategory of  $\underline{\mathcal{C}}$  whose object class is  $\underline{\mathcal{C}}_{t=0} = \underline{\mathcal{C}}_{t \leq 0} \cap \underline{\mathcal{C}}_{t \geq 0}$ .

3. We will say that  $t$  is *bounded below* if  $\cup_{i \in \mathbb{Z}} \underline{\mathcal{C}}_{t \geq i} = \text{Obj } \underline{\mathcal{C}}$ .

Moreover, we say that  $t$  is *bounded* if the equality  $\cup_{i \in \mathbb{Z}} \underline{\mathcal{C}}_{t \leq i} = \text{Obj } \underline{\mathcal{C}}$  is valid as well.

4. Let  $\underline{\mathcal{C}}^0$  be a full triangulated subcategory of  $\underline{\mathcal{C}}$ .

We will say that  $t$  *restricts to*  $\underline{\mathcal{C}}^0$  whenever the couple  $t_0 = (\underline{\mathcal{C}}_{t \leq 0} \cap \text{Obj } \underline{\mathcal{C}}^0, \underline{\mathcal{C}}_{t \geq 0} \cap \text{Obj } \underline{\mathcal{C}}^0)$  is a  $t$ -structure on  $\underline{\mathcal{C}}^0$ .

Let us recall a few properties of  $t$ -structures.

**Proposition 1.1.5.** Let  $t$  be a  $t$ -structure on a triangulated category  $\underline{\mathcal{C}}$ . Then the following statements are valid.

1. The triangle (1.1.1) is canonically and functorially determined by  $M$ . Moreover,  $L_t$  is right adjoint to the embedding  $\underline{\mathcal{C}}_{t \geq 0} \rightarrow \underline{\mathcal{C}}$  (if we consider  $\underline{\mathcal{C}}_{t \geq 0}$  as a full subcategory of  $\underline{\mathcal{C}}$ ) and  $R_t$  is left adjoint to the embedding  $\underline{\mathcal{C}}_{t \leq -1} \rightarrow \underline{\mathcal{C}}$ .

2.  $\underline{Ht}$  is an abelian category with short exact sequences corresponding to distinguished triangles in  $\underline{\mathcal{C}}$ .

3. For any  $n \in \mathbb{Z}$  we will use the notation  $t_{\geq n}$  for the functor  $[n] \circ L_t \circ [-n]$ , and  $t_{\leq n} = [n+1] \circ R_t \circ [-n-1]$ .

Then there is a canonical isomorphism of functors  $t_{\leq 0} \circ t_{\geq 0} \cong t_{\geq 0} \circ t_{\leq 0}$  (if we consider these functors as endofunctors of  $\underline{\mathcal{C}}$ ), and the composite functor  $H^t = H_0^t$  actually takes values in the subcategory  $\underline{Ht}$  of  $\underline{\mathcal{C}}$ . Furthermore, this functor  $H^t : \underline{\mathcal{C}} \rightarrow \underline{Ht}$  is homological.

4.  $\underline{\mathcal{C}}_{t \leq 0} = \underline{\mathcal{C}}_{t \geq 1}^\perp$  and  $\underline{\mathcal{C}}_{t \geq 0} = {}^\perp(\underline{\mathcal{C}}_{t \leq -1})$ .

Consequently, these classes are retraction-closed and extension-closed in  $\underline{\mathcal{C}}$ .

Thus  $t$  is uniquely determined both by  $\underline{\mathcal{C}}_{t \geq 0}$  and by  $\underline{\mathcal{C}}_{t \leq 0}$ .

5.  $\underline{\mathcal{C}}_{t \geq 0}$  is closed with respect to all (small) coproducts that exist in  $\underline{\mathcal{C}}$ .

*Proof.* Assertions 1–4 were essentially established in §1.3 of [BBD82] (yet see Remark 1.1.6(2) below).

Assertion 5 immediately follows from assertion 4.  $\square$

*Remark 1.1.6.* 1. The notion of a  $t$ -structure is clearly self-dual, that is, the couple  $t^{op} = (\underline{\mathcal{C}}_{t \geq 0}, \underline{\mathcal{C}}_{t \leq 0})$  gives a  $t$ -structure on the category  $\underline{\mathcal{C}}^{op}$ . We will say that the latter  $t$ -structure is *opposite* to  $t$ .

2. Even though in [BBD82] where  $t$ -structures were introduced and in several preceding papers of the author the "cohomological convention" for  $t$ -structures was used, in the current text we use the homological convention; the reason for this is that it is coherent with the homological convention for weight structures (see Remark 1.2.3(3) below). Respectively, our notation  $\underline{\mathcal{C}}_{t \geq 0}$  corresponds to the class  $\underline{\mathcal{C}}^{t \leq 0}$  in the cohomological convention.

We will also need two simple statements on restrictions of  $t$ -structure.

**Lemma 1.1.7.** Assume that  $\underline{\mathcal{C}}$  is endowed with a  $t$ -structure  $t$ , and  $\underline{\mathcal{C}}^0, \underline{\mathcal{C}}^1$  are triangulated subcategories of  $\underline{\mathcal{C}}$ .

1.  $t$  restricts to  $\underline{\mathcal{C}}^0$  if and only if the functor  $L_t$  sends  $\underline{\mathcal{C}}^0$  into itself.

2. Take  $\underline{\mathcal{C}}^2 = \underline{\mathcal{C}}^0 \cap \underline{\mathcal{C}}^1$  (that is, the full subcategory of  $\underline{\mathcal{C}}$  whose object class equals  $\text{Obj } \underline{\mathcal{C}}^0 \cap \text{Obj } \underline{\mathcal{C}}^1$ ). Then  $\underline{\mathcal{C}}^2$  is triangulated.

Moreover, if  $t$  restricts both to  $\underline{\mathcal{C}}^0$  and  $\underline{\mathcal{C}}^1$  then it restricts to  $\underline{\mathcal{C}}^2$  as well, and the heart of this restriction equals the intersections of the hearts for  $\underline{\mathcal{C}}^0$  and  $\underline{\mathcal{C}}^1$ .



*Proof.* 1. If  $t$  restricts to  $\underline{\mathcal{C}}^0$  then the uniqueness of the triangles (1.1.1) (see Proposition 1.1.5(1)) implies that  $L_t$  sends  $\underline{\mathcal{C}}^0$  into itself indeed.

Conversely, to check whether  $t$  restricts to  $\underline{\mathcal{C}}^0$  it clearly suffices to verify that any object  $M$  of  $\underline{\mathcal{C}}^0$  possesses a decomposition of the sort (1.1.1) inside  $\underline{\mathcal{C}}^0$ . Now, if  $L_t(M) \in \text{Obj } \underline{\mathcal{C}}^0$  then  $R_t(M)$  is an object of  $\underline{\mathcal{C}}^0$  as well since  $\underline{\mathcal{C}}^0$  is a triangulated subcategory of  $\underline{\mathcal{C}}$ .

2.  $\underline{\mathcal{C}}^2$  is triangulated since  $\underline{\mathcal{C}}^0$  and  $\underline{\mathcal{C}}^1$  are (fully) strict subcategories of  $\underline{\mathcal{C}}$ , whereas different choices of cones of a  $\underline{\mathcal{C}}$ -morphisms are connected by (non-unique) isomorphisms.

The existence of the restricted weight structure statement is easy as well; one can immediately deduce this statement from assertion 1. Lastly, the heart assertion is an immediate consequence of our definitions.  $\square$

*Remark 1.1.8.* One may easily add some more equivalent conditions to Lemma 1.1.7(1); cf. Theorem 2.3.3(I) below.

## 1.2 Some basics on weight structures

Let us recall some basic definitions of the theory of weight structures.

**Definition 1.2.1.** I. A pair of subclasses  $\underline{\mathcal{C}}_{w \leq 0}, \underline{\mathcal{C}}_{w \geq 0} \subset \text{Obj } \underline{\mathcal{C}}$  will be said to define a weight structure  $w$  on a triangulated category  $\underline{\mathcal{C}}$  if they satisfy the following conditions.

(i)  $\underline{\mathcal{C}}_{w \leq 0}$  and  $\underline{\mathcal{C}}_{w \geq 0}$  are retraction-closed in  $\underline{\mathcal{C}}$  (i.e., contain all  $\underline{\mathcal{C}}$ -retracts of their objects).

(ii) **Semi-invariance with respect to translations.**

$$\underline{\mathcal{C}}_{w \leq 0} \subset \underline{\mathcal{C}}_{w \leq 0}[1], \underline{\mathcal{C}}_{w \geq 0}[1] \subset \underline{\mathcal{C}}_{w \geq 0}.$$

(iii) **Orthogonality.**

$$\underline{\mathcal{C}}_{w \leq 0} \perp \underline{\mathcal{C}}_{w \geq 0}[1].$$

(iv) **Weight decompositions.**

For any  $M \in \text{Obj } \underline{\mathcal{C}}$  there exists a distinguished triangle

$$L_w M \rightarrow M \rightarrow R_w M \rightarrow L_w M[1] \quad (1.2.1)$$

such that  $L_w M \in \underline{\mathcal{C}}_{w \leq 0}$  and  $R_w M \in \underline{\mathcal{C}}_{w \geq 0}[1]$ .

We will also need the following definitions.

**Definition 1.2.2.** Let  $i, j \in \mathbb{Z}$ ; assume that a triangulated category  $\underline{\mathcal{C}}$  is endowed with a weight structure  $w$ .

1. The full subcategory  $\underline{Hw}$  of  $\underline{\mathcal{C}}$  whose objects are  $\underline{\mathcal{C}}_{w=0} = \underline{\mathcal{C}}_{w \geq 0} \cap \underline{\mathcal{C}}_{w \leq 0}$  is called the *heart* of  $w$ .
2.  $\underline{\mathcal{C}}_{w \geq i}$  (resp.  $\underline{\mathcal{C}}_{w \leq i}$ ,  $\underline{\mathcal{C}}_{w=i}$ ) will denote the class  $\underline{\mathcal{C}}_{w \geq 0}[i]$  (resp.  $\underline{\mathcal{C}}_{w \leq 0}[i]$ ,  $\underline{\mathcal{C}}_{w=0}[i]$ ).
3.  $\underline{\mathcal{C}}_{[i,j]}$  denotes  $\underline{\mathcal{C}}_{w \geq i} \cap \underline{\mathcal{C}}_{w \leq j}$ .
4. Let  $\underline{\mathcal{C}}^0$  be a (strictly full) triangulated subcategory of  $\underline{\mathcal{C}}$ .

We will say that  $w$  *restricts* to  $\underline{\mathcal{C}}^0$  whenever the couple  $w_0 = (\underline{\mathcal{C}}_{w \leq 0} \cap \text{Obj } \underline{\mathcal{C}}^0, \underline{\mathcal{C}}_{w \geq 0} \cap \text{Obj } \underline{\mathcal{C}}^0)$  is a weight structure on  $\underline{\mathcal{C}}^0$ .

Moreover, in this case we will also say that  $w$  is an *extension* of  $w_0$  (to  $\underline{\mathcal{C}}$ ).

5. We will call  $\cup_{i \in \mathbb{Z}} \underline{\mathcal{C}}_{w \geq i}$  (resp.  $\cup_{i \in \mathbb{Z}} \underline{\mathcal{C}}_{w \leq i}$ ) the class of *w-bounded below* (resp., *w-bounded above*) objects of  $\underline{\mathcal{C}}$ .

Moreover, we say that  $w$  is *bounded below* (resp. *bounded above*, resp. *bounded*) if all objects of  $\underline{\mathcal{C}}$  are bounded below (resp. bounded above, resp. are bounded both below and above).

*Remark 1.2.3.* 1. A weight decomposition (of any  $M \in \text{Obj } \underline{\mathcal{C}}$ ) is almost never canonical (see Proposition 1.2.6 below for more detail).

Still for any  $m \in \mathbb{Z}$  the axiom (iv) gives the existence of a distinguished triangle

$$w_{\leq m}M \rightarrow M \rightarrow w_{\geq m+1}M \rightarrow (w_{\leq m}M)[1] \quad (1.2.2)$$

with some  $w_{\leq m}M \in \underline{\mathcal{C}}_{w \leq m}$  and  $w_{\geq m+1}M \in \underline{\mathcal{C}}_{w \geq m+1}$ ; we will call it an *m-weight decomposition* of  $M$ .

We will often use this notation below (even though  $w_{\geq m+1}M$  and  $w_{\leq m}M$  are not canonically determined by  $M$ ); we will call any possible choice either of  $w_{\geq m+1}M$  or of  $w_{\leq m}M$  (for any  $m \in \mathbb{Z}$ ) a *weight truncation* of  $M$ . Moreover, when we will write arrows of the type  $w_{\leq m}M \rightarrow M$  or  $M \rightarrow w_{\geq m+1}M$  we will always assume that they come from some *m-weight decomposition* of  $M$ .

2. In the current paper (along with several previous ones) we use the “homological convention” for weight structures, whereas in [Bon10a] and [Bon10b], the cohomological convention was used. In the latter convention the roles of  $\underline{\mathcal{C}}_{w \leq 0}$  and  $\underline{\mathcal{C}}_{w \geq 0}$  are interchanged, i.e., one takes  $\underline{\mathcal{C}}^{w \leq 0} = \underline{\mathcal{C}}_{w \geq 0}$  and  $\underline{\mathcal{C}}^{w \geq 0} = \underline{\mathcal{C}}_{w \leq 0}$ .

We also recall that weight structures were independently introduced in [Pau08]; D. Pauksztello has called them co-t-structures.

**Proposition 1.2.4.** Let  $m \leq n \in \mathbb{Z}$ ,  $M, M' \in \text{Obj } \underline{\mathcal{C}}$ ,  $g \in \underline{\mathcal{C}}(M, M')$ .

1. The axiomatics of weight structures is self-dual, i.e., on  $\underline{\mathcal{C}}' = \underline{\mathcal{C}}^{op}$  (so  $\text{Obj } \underline{\mathcal{C}}' = \text{Obj } \underline{\mathcal{C}}$ ) there exists the (opposite) weight structure  $w'$  for which  $\underline{\mathcal{C}}'_{w' \leq 0} = \underline{\mathcal{C}}_{w \geq 0}$  and  $\underline{\mathcal{C}}'_{w' \geq 0} = \underline{\mathcal{C}}_{w \leq 0}$ .
2.  $\underline{\mathcal{C}}_{w \geq 0} = (\underline{\mathcal{C}}_{w \leq -1})^\perp$  and  $\underline{\mathcal{C}}_{w \leq 0} = {}^\perp \underline{\mathcal{C}}_{w \geq 1}$ .
3.  $\underline{\mathcal{C}}_{w \leq 0}$  is closed with respect to all coproducts that exist in  $\underline{\mathcal{C}}$ .
4. For any (fixed) *m-weight decomposition* of  $M$  and an *n-weight decomposition* of  $M'$  (see Remark 1.2.3(2))  $g$  can be extended to a morphism of the corresponding distinguished triangles:

$$\begin{array}{ccccc} w_{\leq m}M & \xrightarrow{c} & M & \longrightarrow & w_{\geq m+1}M \\ \downarrow h & & \downarrow g & & \downarrow j \\ w_{\leq n}M' & \longrightarrow & M' & \longrightarrow & w_{\geq n+1}M' \end{array} \quad (1.2.3)$$

Moreover, if  $m < n$  then this extension is unique (provided that the rows are fixed).

5. If  $A \rightarrow B \rightarrow C \rightarrow A[1]$  is a  $\underline{\mathcal{C}}$ -distinguished triangle and  $A, C \in \underline{\mathcal{C}}_{w=0}$  then this distinguished triangle splits; hence  $B \cong A \oplus C \in \underline{\mathcal{C}}_{w=0}$ .
6. If  $M$  belongs to  $\underline{\mathcal{C}}_{w \leq 0}$  (resp. to  $\underline{\mathcal{C}}_{w \geq 0}$ ) then it is a retract of any choice of  $w_{\leq 0}M$  (resp. of  $w_{\geq 0}M$ ).

7. If  $M \in \underline{\mathcal{C}}_{w \geq m}$  then  $w_{\leq n} M \in \underline{\mathcal{C}}_{[m, n]}$  (for any  $n$ -weight decomposition of  $M$ ).
8. The class  $\underline{\mathcal{C}}_{[m, l]}$  is the extension-closure of  $\cup_{m \leq j \leq l} \underline{\mathcal{C}}_{w=j}$ .  
Consequently,  $\underline{\mathcal{C}}_{w=0}$  is silting if  $w$  is bounded.
9. Let  $v$  be another weight structure for  $\underline{\mathcal{C}}$ ; assume  $\underline{\mathcal{C}}_{w \leq 0} \subset \underline{\mathcal{C}}_{v \leq 0}$  and  $\underline{\mathcal{C}}_{w \geq 0} \subset \underline{\mathcal{C}}_{v \geq 0}$ .<sup>2</sup> Then  $w = v$  (i.e., these inclusions are equalities).

*Proof.* All these statements were essentially proved in [Bon10a] (yet pay attention to Remark 1.2.3(3) above!).  $\square$

Now we pass to couples that are simultaneously weight structures and  $t$ -structures.

**Definition 1.2.5.** Assume that  $\mathcal{B}, \mathcal{A}$  are strictly full triangulated subcategories of  $\underline{\mathcal{C}}$ .

Then the couple  $D = (\mathcal{A}, \mathcal{B})$  is a (length 2) *semi-orthogonal decomposition* of  $\underline{\mathcal{C}}$  (or just gives a decomposition of  $\underline{\mathcal{C}}$ ) if  $\text{Obj } \mathcal{B} \perp \text{Obj } \mathcal{A}$  and  $\text{Obj } \mathcal{B} \star \text{Obj } \mathcal{A} = \text{Obj } \underline{\mathcal{C}}$ .

Below we will need some well-known properties of semi-orthogonal decompositions.

**Proposition 1.2.6.** The following assumptions on a couple  $(C_1, C_2)$  of subclasses of  $\text{Obj } \underline{\mathcal{C}}$  are equivalent.

1. The full subcategories of  $\underline{\mathcal{C}}$  corresponding to  $C_2$  and  $C_1$  give a decomposition of  $\underline{\mathcal{C}}$ .
2.  $(C_2, C_1)$  is a  $t$ -structure and  $C_1[1] = C_1$ .
3. The full subcategory  $\underline{\mathcal{C}}_1$  of  $\underline{\mathcal{C}}$  corresponding to  $C_1$  is strict, triangulated, and *right admissible*, that is, there exists a right adjoint to the embedding  $\underline{\mathcal{C}}_1 \rightarrow \underline{\mathcal{C}}$ ;  $C_2 = C_1^\perp$ .
4.  $(C_1, C_2)$  is a weight structure and  $C_1[1] = C_1$ .
5.  $(C_1, C_2)$  is a weight structure,  $C_1[1] = C_1$ , and  $C_2[1] = C_2$ .
6.  $(C_1, C_2)$  is a weight structure  $w$  and  $\underline{Hw} = \{0\}$ .
7.  $(C_1, C_2)$  is a weight structure  $w$  and the triangle (1.2.1) is functorially determined by  $M$ .
8.  $(C_1, C_2)$  is a weight structure  $w$  and the triangle (1.2.1) is determined by  $M$  up to a non-canonical isomorphism.

*Proof.* The equivalence of conditions 3, 2, and 2 can be easily obtained by means of combining Propositions 3.4(4) and 3.2(1,2) of [BoV19] (pay attention to Remark 1.1.6(2) above!).

The equivalence of conditions 1 and 3 is very well-known; see Proposition 2.5 of [Bon24].

---

<sup>2</sup>I.e., the identity on  $\underline{\mathcal{C}}$  is weight-exact with respect to  $(w, v)$ .

Next, condition 5 clearly implies 4, and applying Proposition 1.2.4(2) (or Proposition 3.2(2) of [BoV19]) we obtain the converse implication.

Now, if condition 5 is fulfilled then  $\underline{C}_{w=0} = \underline{C}_{w=-1} \perp \underline{C}_{w=0}$ ; thus we obtain condition 6. Conversely, if condition 6 is fulfilled then Proposition 1.2.4(7) easily implies that  $C_2 \subset C_2[1]$ . Thus we can apply Proposition 1.2.4(2) once again to obtain condition 4.

Next, if  $0 \neq C \in \underline{C}_{w=0}$  then the triangles  $0 \rightarrow 0 \rightarrow 0 \rightarrow 0$  and  $C \rightarrow 0 \rightarrow C[1] \rightarrow C[1]$  clearly give two non-isomorphic weight decompositions of 0. Hence condition 8 implies condition 6.

Clearly, condition 7 implies condition 8.

Lastly, assume that condition 5 is fulfilled. Then any weight decomposition triangle is clearly a 1-weight decomposition triangle as well. Consequently, the uniqueness provided by Proposition 1.2.4(4) yields condition 7; cf. Remark 1.2.6 of [Bon22b] where compositions of the corresponding morphisms between decomposition triangles is discussed.  $\square$

*Remark 1.2.7.* 1. Roughly, semi-orthogonal decompositions are the weight structures that are also  $t$ -structures. One may also characterize them both as shift-stable weight structures and as shift-stable  $t$ -structures.

2. Note also that this notion is obviously self-dual (that is, permuting the classes in the couple yields a semi-orthogonal decomposition in the category  $\underline{C}^{op}$  cf. Proposition 1.2.4(1)).

3. Below we will also need the following simple criterion: if  $\underline{C}$  is smashing (cf. Theorem 0.1) then  $w$  is smashing if and only if the right adjoint functor provided by condition 3 of Proposition 1.2.6 respects coproducts; see Proposition 3.4(5) of [BoV19].

### 1.3 On orthogonal and adjacent structures

Now let us give a certain definition of orthogonality for weight and  $t$ -structures (cf. Remark 1.3.4(1) below).

**Definition 1.3.1.** Assume that  $\underline{C}$  and  $\underline{C}'$  are (full) triangulated subcategories of a triangulated category  $\underline{D}$ ,  $w$  is a weight structure on  $\underline{C}$  and  $t$  is a  $t$ -structure on  $\underline{C}'$ .

1. We will say that  $w$  and  $t$  are *orthogonal* if  $\underline{C}_{w \leq 0} \perp_{\underline{D}} \underline{C}'_{t \geq 1}$  and  $\underline{C}_{w \geq 0} \perp_{\underline{D}} \underline{C}'_{t \leq -1}$ .

Dually, we say that  $w$  and  $t$  are *anti-orthogonal* whenever  $\underline{C}'_{t \geq 1} \perp_{\underline{D}} \underline{C}_{w \leq 0}$  and  $\underline{C}'_{t \leq -1} \perp_{\underline{D}} \underline{C}_{w \geq 0}$ .

2. If  $\underline{C} = \underline{C}' = \underline{D}$  and  $w$  is (anti) orthogonal to  $t$  then we will also say that  $w$  and  $t$  are left (resp. right) *adjacent*.

3. We will say that  $t$  is *strictly orthogonal* to  $w$  if  $\underline{C}'_{t \geq 1} = \underline{C}_{w \leq 0}^{\perp_{\underline{D}}} \cap \text{Obj } \underline{C}'$  and  $\underline{C}'_{t \leq -1} = \underline{C}_{w \geq 0}^{\perp_{\underline{D}}} \cap \text{Obj } \underline{C}'$ .

*Remark 1.3.2.* Our main statements below yield  $t$ -structures that are strictly orthogonal to the corresponding weight structures. Note however that strictness is not automatic; in particular, it is clearly not fulfilled if  $\underline{C} \perp \underline{C}'$ ; then arbitrary  $w$  and  $t$  would be orthogonal.

Let us now relate the latter definition to the notion of adjacent structures introduced in [Bon10a].

**Proposition 1.3.3.** For  $\underline{C}$ ,  $w$ , and  $t$  as in Definition 1.3.1(2) we have the following:  $w$  and  $t$  are (left) adjacent if and only if  $\underline{C}_{w \geq 0} = \underline{C}_{t \geq 0}$ .

*Proof.* If  $\underline{C}_{w \geq 0} = \underline{C}_{t \geq 0}$  then  $w$  and  $t$  are (left) adjacent immediately from the orthogonality axioms of weight and  $t$ -structures (see Definition 1.1.3(iii) and Definition 1.2.1(iii)). Conversely, if  $w$  is adjacent to  $t$  then combining the orthogonality conditions with Proposition 1.2.4(2) and Proposition 1.1.5(4) we obtain that  $\underline{C}_{t \geq 0} \subset \underline{C}_{w \geq 0}$  and  $\underline{C}_{w \geq 0} \subset \underline{C}_{t \geq 0}$ . Hence  $\underline{C}_{w \geq 0} = \underline{C}_{t \geq 0}$  as desired.  $\square$

*Remark 1.3.4.* 1. In Definition 2.5.1(3) of [Bon10b] and §5.2 of [Bon19] orthogonality was defined with respect to an arbitrary duality  $\Phi$  between  $\underline{C}$  and  $\underline{C}'$ ; see Definition 1.1.1. Moreover, in the current paper we (essentially) restrict this definition to dualities provided by Lemma 1.1.2.

2. Proposition 1.3.3 says that our definition of adjacent "structures" is essentially equivalent to the original Definition 4.4.1 of [Bon10a].

We also make a simple observation concerning semi-orthogonal decomposition.

**Proposition 1.3.5.** Assume that  $w$  and  $t$  satisfy the assumptions of Definition 1.3.1,  $t$  is strictly orthogonal to  $w$  (note that this is condition follows from orthogonality whenever  $\underline{C}' = \underline{C}$ ; see Theorem 2.3.3(I) below), and the couple  $(\underline{C}_{w \leq 0}, \underline{C}_{w \geq 0})$  is a semi-orthogonal decomposition (see Definition 1.2.5). Then  $(\underline{C}'_{t \geq 0}, \underline{C}'_{t \leq 0})$  is a semi-orthogonal decomposition as well.

*Proof.* Applying Proposition 1.2.6 (see conditions 4 and 1 in it) along with the definition of strict orthogonality we obtain that  $\underline{C}'_{t \geq 0} = \underline{C}'_{t \geq 0}[1]$ . Hence  $(\underline{C}'_{t \geq 0}, \underline{C}'_{t \leq 0})$  is a semi-orthogonal decomposition indeed; see condition 2 in Proposition 1.2.6.  $\square$

## 2 On virtual $t$ -truncations and their relation to orthogonal $t$ -structures

This section is devoted to virtual  $t$ -truncations of functors (these come from weight structures) and their general relationship with orthogonal  $t$ -structures.

In §2.1 we recall the definition of virtual  $t$ -truncations of (co)homological functors.

In §2.2 we define weight range of functors and relate it to virtual  $t$ -truncations.

In §2.3 we prove that if  $\underline{C}' \subset \underline{C}$  then for a weight structure  $w$  on  $\underline{C}$  there exists at most one  $t$ -structure on  $\underline{C}'$  (strictly) orthogonal to it; see Theorem 2.3.3. Moreover, if  $\underline{C}^0 \subset \underline{C}$  is  $R$ -linear,  $w^0$  is a weight structure on  $\underline{C}^0$ , and  $t$  is a  $t$ -structure on  $\underline{C}$  strictly orthogonal to  $w^0$  then for any weak Serre subcategory  $\underline{A}$  of  $R - \text{Mod}$  one can specify a subcategory  $\underline{C}_{\underline{A}}$  such that  $t$  restricts to it.

In §2.4 we define *coproductive extensions* of weight structures (from subcategories). For  $(w, t, \underline{C}, \underline{C}')$  as above we prove that if  $w$  is a *coproductive extension* of a weight structure  $w^0$  on  $\underline{C}^0 \subset \underline{C}$  then  $w$  is strictly orthogonal to  $t$  if and only if  $w^0$  is (see Corollary 2.4.4); hence  $w^0$  determines  $t$  uniquely.

## 2.1 Virtual $t$ -truncations of (co)homological functors

**Definition 2.1.1.** Assume that  $\underline{\mathcal{C}}$  is endowed with a weight structure  $w$ ,  $n \in \mathbb{Z}$ , and  $\underline{\mathcal{A}}$  is an abelian category.

1. Let  $H$  be a cohomological functor from  $\underline{\mathcal{C}}$  into  $\underline{\mathcal{A}}$ .

We define the *virtual  $t$ -truncation* functor  $\tau_{\leq n}(H)$  (resp.  $\tau_{\geq n}(H)$ ) by the correspondence

$$M \mapsto \text{Im}(H(w_{\leq n+1}M) \rightarrow H(w_{\leq n}M));$$

(resp.  $M \mapsto \text{Im}(H(w_{\geq n}M) \rightarrow H(w_{\geq n-1}M))$ ); here we take arbitrary choices of the corresponding weight truncations of  $M$  and connect them using Proposition 1.2.4(4) in the case  $g = \text{id}_M$ .

2. Let  $H' : \underline{\mathcal{C}} \rightarrow \underline{\mathcal{A}}$  be a homological functor. Then we will write  $\tau_{\leq n}(H')$  for the correspondence  $M \mapsto \text{Im}(H'(w_{\leq n}M) \rightarrow H'(w_{\leq n+1}M))$  and  $\tau_{\geq n}(H') = M \mapsto \text{Im}(H'(w_{\geq n-1}M) \rightarrow H'(w_{\geq n}M))$  (here we take the same connecting arrows between weight truncations of  $M$  as above).

3. Assume that  $\underline{\mathcal{C}}$  is a triangulated subcategory of a triangulated category  $\underline{\mathcal{D}}$ . Then for any  $M \in \text{Obj } \underline{\mathcal{D}}$  we will write  $H_M = H_M^{\underline{\mathcal{C}}}$  (resp.  $H^M = H_{\underline{\mathcal{C}}}^M$ ) for the restriction of the functor (co)represented by  $M$  to  $\underline{\mathcal{C}}$  (thus  $H_M$  and  $H^M$  are functors from  $\underline{\mathcal{C}}$  into  $\text{Ab}$ ). Moreover, sometimes we will say that these functors are  $\underline{\mathcal{D}}$ -Yoneda ones, and that  $H_M$  (resp.  $H^M$ ) is  $\underline{\mathcal{D}}$ -(co)represented by  $M$ .

We recall the main properties of these constructions.

**Proposition 2.1.2.** In the notation of the previous definition the following statements are valid.

1. The objects  $\tau_{\leq n}(H)(M)$  and  $\tau_{\geq n}(H)(M)$  are  $\underline{\mathcal{C}}$ -functorial in  $M$  (and essentially do not depend on any choices).
2. The functors  $\tau_{\leq n}(H)$  and  $\tau_{\geq n}(H)$  are cohomological.
3. There exist natural transformations that yield a long exact sequence

$$\begin{aligned} \cdots \rightarrow \tau_{\leq n-1}(H) \circ [-1] \rightarrow \tau_{\geq n}(H) \xrightarrow{a_3} H \\ \xrightarrow{b_{n-1}^{-1}} \tau_{\leq n-1}(H) \rightarrow \tau_{\geq n}(H) \circ [1] \rightarrow H^{-1} \rightarrow \cdots \end{aligned} \quad (2.1.1)$$

(i.e., the result of applying this sequence to any object of  $\underline{\mathcal{C}}$  is a long exact sequence); the shift of this exact sequence by 3 positions is given by composing the functors with  $-[1]$ .

4. Assume that there exists a  $t$ -structure  $t$  that is left adjacent to  $w$  (for certain  $\underline{\mathcal{C}}'$  and  $\underline{\mathcal{D}}$  as in Definition 1.3.1). Then for any  $M \in \text{Obj } \underline{\mathcal{C}}'$  the functors  $\tau_{\geq n}(H_M)$  and  $\tau_{\leq n}(H_M)$  (where  $H_M$  is defined in Definition 2.1.1(3)) are  $\underline{\mathcal{D}}$ -represented (on  $\underline{\mathcal{C}}$ ) by  $t_{\geq n}M$  and  $t_{\leq n}M$  (see Proposition 1.1.5(3)), respectively.
5. Let  $H^{op}$  be the cohomological functor obtained from  $H$  by means of inverting arrows both in  $\underline{\mathcal{C}}$  and  $\underline{\mathcal{A}}$ . Then there exist canonical isomorphisms  $\tau_{\leq n}^{op}(H^{op}) \cong (\tau_{\geq -n}(H))^{op}$  and  $\tau_{\geq n}^{op}(H^{op}) \cong (\tau_{\leq -n}(H))^{op}$ ; here  $\tau^{op}$  denotes virtual  $t$ -truncations with respect to  $w^{op}$ .

6. The correspondence  $\tau_{\geq n}(H')$  gives a well-defined homological functor, and there exists a homological analogue of the long exact sequence (2.1.1).

Moreover, if there exists a  $t$ -structure  $t$  on a triangulated category  $\underline{\mathcal{C}}'$  that is orthogonal to  $w$  (with respect to a triangulated category  $\underline{\mathcal{D}}$  containing  $\underline{\mathcal{C}}$  and  $\underline{\mathcal{C}}'$ ),  $\underline{\mathcal{A}} = \text{Ab}$ , and the functor  $H$  is  $\underline{\mathcal{D}}$ -corepresented by an object  $N$  of  $\underline{\mathcal{C}}'$ , then  $\tau_{\geq n}(H')$  is  $\underline{\mathcal{D}}$ -corepresented by  $t_{\geq n}N$  and  $\tau_{\leq n}(H')$  is  $\underline{\mathcal{D}}$ -corepresented by  $t_{\leq n}N$ .

*Proof.* Assertions 1–3 are essentially given by Theorem 2.3.1 of [Bon10b]; yet pay attention to Remark 1.2.3(3). Assertion 4 is given by Proposition 2.5.4(1) of *ibid.*; here one should apply Lemma 1.1.2 and recall Remark 1.3.4(1). Assertion 6 is easily seen to be dual to assertions 1–3.

Assertion 5 can be obtained by means of applying Proposition 1.2.4(1) quite straightforwardly. Firstly, reversing arrows converts any  $m$ -weight decomposition of  $M \in \text{Obj } \underline{\mathcal{C}}$  into its  $-m-1$ -weight decomposition in  $\underline{\mathcal{C}}^{op}$  (with respect to  $w^{op}$ ). Moreover, if we reverse arrows in (1.2.3) then we obtain a similar diagram in  $\underline{\mathcal{C}}^{op}$ ; if  $m < n$  in the initial diagram then we obtain the unique morphism of triangles which corresponds to  $g$  and connects (the dual)  $-n-1$ -weight decomposition of  $M'$  with the corresponding  $-m-1$ -weight decomposition of  $M$ . Thus if we reverse in the definitions  $\tau_{\geq -n}(H)$  and  $\tau_{\leq -n}(H)$  (for any  $n \in \mathbb{Z}$ ) then we obtain the morphisms that compute  $\tau_{\leq n}^{op}(H^{op})$  and  $\tau_{\geq n}^{op}(H^{op})$ , respectively.  $\square$

Some more useful properties of virtual  $t$ -truncations follow from our definitions easily.

**Proposition 2.1.3.** Let  $n \in \mathbb{Z}$ ; assume that  $\underline{\mathcal{C}}$  is endowed with a weight structure  $w$  and  $H$  is a cohomological functor from  $\underline{\mathcal{C}}$  into  $\underline{\mathcal{A}}$ .

I.1. For any  $i \in \mathbb{Z}$  we have  $\tau_{\leq n+i}(H) \cong \tau_{\leq n}(H \circ [i]) \circ [-i]$  and  $\tau_{\geq n+i}(H) \cong \tau_{\geq n}(H \circ [i]) \circ [-i]$ , and also  $\tau_{\leq n+i}(H') \cong \tau_{\leq n}(H' \circ [i]) \circ [-i]$  and  $\tau_{\geq n+i}(H') \cong \tau_{\geq n}(H' \circ [i]) \circ [-i]$ .

2. Assume that  $F : \underline{\mathcal{A}} \rightarrow \underline{\mathcal{A}}'$  is an exact functor. Then  $\tau_{\leq n}(F \circ H) \cong F \circ \tau_{\leq n}(H)$  and  $\tau_{\geq n}(F \circ H) \cong F \circ \tau_{\geq n}(H)$ .

II. Assume that  $\underline{\mathcal{C}}^0$  is a triangulated category endowed with a weight structure  $w^0$ ; let  $F : \underline{\mathcal{C}}^0 \rightarrow \underline{\mathcal{C}}$  be a weight-exact functor.

1. Then  $\tau_{\leq n}^0(H \circ F) \cong (\tau_{\leq n}(H)) \circ F$  and  $\tau_{\geq n}^0(H \circ F) \cong (\tau_{\geq n}(H)) \circ F$ ; here  $\tau_{\leq n}^0(-)$  and  $\tau_{\geq n}^0(-)$  denote the corresponding virtual  $t$ -truncations with respect to  $w^0$ .

2. Assume that  $F$  is a full embedding, that is,  $w$  restricts to a weight structure  $w^0$  on  $\underline{\mathcal{C}}^0$ .

Then  $\tau_{\leq n}^0(H|_{\underline{\mathcal{C}}^0}) \cong \tau_{\leq n}(H)|_{\underline{\mathcal{C}}^0}$  and  $\tau_{\geq n}^0(H|_{\underline{\mathcal{C}}^0}) \cong \tau_{\geq n}(H)|_{\underline{\mathcal{C}}^0}$ .

3. In addition to the assumptions of the previous assertion, suppose that  $\underline{\mathcal{A}}^0$  is a (strictly full) abelian subcategory of  $\underline{\mathcal{A}}$  (see §1.1).

Then if the restricted functor  $H|_{\underline{\mathcal{C}}^0}$  takes its values in  $\underline{\mathcal{A}}^0$  then the same is true for the restrictions to  $\underline{\mathcal{C}}^0$  of all virtual  $t$ -truncations of  $H$ .

*Proof.* I.1. Obvious.

II. Very easy as well; recall that exact functors respect images of morphisms.

II.1. Very easy; note that for any  $M \in \text{Obj } \underline{\mathcal{C}}^0$  one can compute all  $(\tau_{\leq n}(H)) \circ (F(M))$  and  $(\tau_{\geq n}(H)) \circ (F(M))$  by means of applying  $H$  to the  $w$ -truncations

of  $F(M)$  that are computed as images with respect to  $F$  with respect to  $w^0$ -truncations of  $M$ .

2. This is just a particular case of the previous assertion; note that  $H|_{\underline{\mathcal{C}}^0} = H \circ F$ .

3. According to the previous assertion, it suffices to look at the values of (all the functors)  $\tau_{\leq n}^0(H|_{\underline{\mathcal{C}}^0})$  and  $\tau_{\geq n}^0(H|_{\underline{\mathcal{C}}^0})$ . For the latter purpose it suffices to apply assertion I.2 to the functors  $H|_{\underline{\mathcal{C}}^0}$  and the exact embedding functor  $\underline{A}^0 \rightarrow \underline{A}$ .  $\square$

Now let us list a few easy properties of  $R$ -linear categories, where  $R$  is a commutative ring.

**Lemma 2.1.4.** Let  $\underline{B} \subset \underline{B}'$  be additive categories, and assume that  $\underline{B}$  is  $R$ -linear.

1. Then the category  $\text{Fun}_R(\underline{B}, R - \text{Mod})$  of  $R$ -linear functors is equivalent to  $\text{AddFun}(\underline{B}, \text{Ab})$ ; here we define the multiplication by  $r \in R$  on  $F(B)$  for  $B \in \text{Obj } \underline{B}$  by means of applying  $F(r \text{id}_B)$ .

The equivalence is obtained by means of composing with the forgetful functor  $F_R : R - \text{Mod} \rightarrow \text{Ab}$ .

2. A functor  $\underline{B}^{op} \rightarrow R - \text{Mod}$  is representable by an object of  $\underline{B}'$  whenever its composition with  $F_R$  becomes representable if we consider  $\underline{B}'$  as (just) an additive category.

3. Assume that  $\underline{B}$  is also a triangulated category,  $w$  is a weight structure on it,  $n \in \mathbb{Z}$ .

Then for an  $R$ -linear cohomological functor  $H_R$  from  $\underline{B}$  into  $R - \text{Mod}$  the functors  $\tau_{\leq n}(F_R \circ H)$  and  $\tau_{\geq n}(F_R \circ H)$  are isomorphic to  $F_R \circ \tau_{\leq n}(H)$  and  $F_R \circ \tau_{\geq n}(H)$ , respectively.

Consequently, the functor  $\tau_{\leq n}(H)$  is representable in  $\underline{B}$  if and only if  $\tau_{\leq n}(F_R \circ H)$  (resp.  $\tau_{\geq n}(F_R \circ H)$ ) is (cf. assertion 2).

*Proof.* Assertion 1 is very easy; this is Lemma 3.2.8(1) of [Bon24]. Assertion 2 follows from it immediately.

Next, the first part of assertion 3 is a particular case of Proposition 2.1.3(I.2). Hence it remains to apply assertion 2 to conclude the proof.  $\square$

## 2.2 On weight range

Now we define weight range and relate it to virtual  $t$ -truncations; some of these statements will be applied elsewhere (only).

**Definition 2.2.1.** Let  $m, n \in \mathbb{Z}$ ; let  $H$  be a cohomological functor from  $\underline{\mathcal{C}}$  into  $\underline{A}$ .

Then we will say that  $H$  is of *weight range*  $\geq m$  (resp.  $\leq n$ , resp.  $[m, n]$ ) if it annihilates  $\underline{\mathcal{C}}_{w \leq m-1}$  (resp.  $\underline{\mathcal{C}}_{w \geq n+1}$ , resp. both of these classes).

Moreover, functors of weight range  $[0, 0]$  will also said to be *pure*; cf. Remark 2.2.3(2) below.

**Proposition 2.2.2.** Once again, assume that  $\underline{\mathcal{C}}$  is endowed with a weight structure  $w$ ,  $m, n \in \mathbb{Z}$ , and  $H$  is a cohomological functor from  $\underline{\mathcal{C}}$  into  $\underline{A}$ .



1. Then the functor  $\tau_{\leq n}(H)$  is of weight range  $\leq n$  and  $\tau_{\geq m}(H)$  is of weight range  $\geq m$ .
2. We have  $\tau_{\leq n}(H) \cong H$  (resp.  $\tau_{\geq m}(H) \cong H$ ) if and only if  $H$  is of weight range  $\leq n$  (resp. of weight range  $\geq m$ ).
3. We have  $\tau_{\leq n}(\tau_{\geq m})(H) \cong \tau_{\geq m}(\tau_{\leq n})(H)$ .
4. If  $H$  is of weight range  $\geq m$  then  $\tau_{\leq n}(H)$  is of weight range  $[m, n]$ .  
Dually, if  $H$  is of weight range  $\leq n$  then  $\tau_{\geq m}(H)$  is of weight range  $[m, n]$ .
5. The (not necessarily locally small) category of pure cohomological functors from  $\underline{C}$  into  $\underline{A}$  is equivalent to  $\text{AddFun}(\underline{Hw}^{op}, \underline{A})$  in the obvious natural way.
6. If  $H$  is of weight range  $\geq m$  (resp.  $\leq m$ ) then the morphism  $H(w_{\geq m}M) \rightarrow H(M)$  is epimorphic (resp. the morphism  $H(M) \rightarrow H(w_{\leq m}M)$  is monomorphic); here we take arbitrary choices of the corresponding weight decompositions of  $M$  and apply  $H$  to the connecting morphisms.
7. Assume that  $m > n$ . Then the only functors of weight range  $[m, n]$  are zero ones; thus if  $H$  is of weight range  $\leq n$  (resp.  $\geq m$ ) then  $\tau_{\geq m}(H) = 0$  (resp.  $\tau_{\leq n}(H) = 0$ ).
8. The (representable) functor  $H_M = \underline{C}(-, M) : \underline{C}^{op} \rightarrow \text{Ab}$  if of weight range  $\geq m$  if and only if  $M \in \underline{C}_{w \geq m}$ .
9. Assume that  $H$  is pure and its restriction to  $\underline{Hw}$  takes its values in an abelian subcategory  $\underline{A}^0$  of  $\underline{A}$ . Then the values of  $H$  belong to  $\underline{A}$  as well.
10. Assume that  $t$  is a  $t$ -structure on  $\underline{C}'$  (for some  $\underline{C}' \subset \underline{D}$ , where  $\underline{D}$  contains  $\underline{C}$ ) that is orthogonal to  $w$ . Then for  $N \in \underline{C}'_{t \leq 0}$  (resp.  $N \in \underline{C}'_{t \geq 0}$ , resp.  $N \in \underline{C}'_{t=0}$ ) the corresponding  $\underline{D}$ -Yoneda functor  $H_N : \underline{C}^{op} \rightarrow \text{Ab}$  (see Definition 2.1.1(3)) is of weight range  $\leq 0$  (resp.  $\geq 0$ , resp.  $[0, 0]$ ).

*Proof.* 1. Let  $M \in \underline{C}_{w \geq n+1}$ . Then we can take  $w_{\leq n}(M) = 0$ . Thus  $\tau_{\leq n}(H)(M) = 0$ , and we obtain the first part of the assertion. Its second part is easily seen to be dual to the first part.

2. This is precisely Theorem 2.3.1(III.2,3) of [Bon10b] (up to change of notation); assertion 3 is given by part II.3 of that theorem.

4. Let  $H$  be of weight range  $\geq m$ . Then  $\tau_{\leq n}(H) \cong \tau_{\leq n}(\tau_{\geq m})(H) \cong \tau_{\geq m}(\tau_{\leq n})(H)$  (according to the two previous assertions). It remains to apply assertion 1 to obtain the first statement in the assertion, whereas its second part is easily seen to be the dual of the first part (and certainly can be proved similarly).

5. Immediate from Theorem 2.1.2(2) of [Bon21].

7. For any  $l \in \mathbb{Z}$  and any cohomological  $H$  any choice of an  $l$ -weight decomposition triangle (cf. (1.2.2)) for  $M$  gives a long exact sequence

$$\begin{aligned} \cdots \rightarrow H((w_{\leq l}M)[1]) \rightarrow H(w_{\geq l+1}M) \rightarrow H(M) \\ \rightarrow H(w_{\leq l}M) \rightarrow H((w_{\geq l+1}M)[-1]) \rightarrow \cdots \end{aligned} \quad (2.2.1)$$

The exactness of this sequence in  $H(M)$  for  $l = n$  immediately gives the first part of the assertion. Next, the second part is straightforward from the first one combined with assertion 4.

Assertion 8 is immediate from Proposition 1.2.4(2).

Assertion 6 is a straightforward consequence of assertion 8; just apply (2.2.1) for  $l = m$  and for  $l = m - 1$ , respectively.

Assertion 9 immediately follows from Lemma 2.1.4 of [Bon21].

10. For  $N \in \underline{\mathcal{C}}'_{t \leq 0}$  and  $N \in \underline{\mathcal{C}}'_{t \geq 0}$  the weight range estimates for the functor  $H_N$  prescribed by the assertion are given by the definition of orthogonality, and to obtain the claim for  $N \in \underline{\mathcal{C}}'_{t=0}$  one should combine the first two weight range statements.  $\square$

*Remark 2.2.3.* 1. Roughly, the statements above say that virtual  $t$ -truncations of functors behave as if they corresponded to  $t$ -truncations of objects in a certain triangulated "category of functors" (whence the name; certainly, another justification of this idea is provided by the existence of orthogonal  $t$ -structures statements that will be proved below).

2. In particular, one can "slice" any functor of weight range  $[m, n]$  for  $m \leq n$  into "pieces" of weight  $[i, i]$  for  $m \leq i \leq n$ . Next, composing a "slice" of weight range  $[i, i]$  with  $[i]$  one obtains a pure functor.

Recall also that pure functors were studied in detail in (§2.1 of) [Bon21]; they were called pure ones due to the relation to Deligne's purity (cf. Remark 2.1.3(3–4) of *ibid.*).

Let us also prove some more complicated statements of this sort.

**Proposition 2.2.4.** Adopt the notation and conventions of Proposition 2.2.2, and assume that  $H'$  and  $H''$  are cohomological functors from  $\underline{\mathcal{C}}$  into  $\underline{\mathcal{A}}$  (as well).

I. Then the following statement are fulfilled.

1. If  $H'$  is of weight range  $\geq n$  (resp. of weight range  $\leq n - 1$ ) then any transformation  $T : H' \rightarrow H$  (resp.  $H \rightarrow H'$ ) factors through the transformation  $a_n$  (resp.  $b_{n-1}$ ) in the sequence (2.1.1).

2. If  $H'$  (resp.  $H''$ ) is of weight range  $\geq n$  (resp. of weight range  $\leq n - 1$ ) then there are no non-zero transformations from  $H'$  into  $H''$ .

II. Assume in addition that we have an exact sequence of cohomological functors from  $\underline{\mathcal{C}}$  into  $\underline{\mathcal{A}}$

$$H' \xrightarrow{a} H \xrightarrow{b} H'' \quad (2.2.2)$$

(that is, applying it to any  $M \in \text{Obj } \underline{\mathcal{C}}$  we obtain a half-exact  $\underline{\mathcal{A}}$ -sequence) and for any  $M \in \text{Obj } \underline{\mathcal{C}}$  there exists an isomorphism  $\text{Ker } a(M[-1]) \rightarrow \text{Coker } b(M)$ .

1. If  $H'$  is of weight range  $\geq n$ ,  $H''$  is of weight range  $\leq n - 1$ , then there exists a canonical isomorphism of chains of morphisms of functors  $(2.2.2) \cong \tau_{\geq n}(H) \xrightarrow{a_n} H \xrightarrow{b_{n-1}} \tau_{\leq n-1}(H)$ ; the latter one is a part of the sequence (2.1.1).

2.  $a$  factorizes as  $H' \xrightarrow{i'} \tau_{\geq n}(H) \xrightarrow{a_n} H$ , where  $i'$  is an isomorphism of functors, if and only if  $b$  factorizes as  $H \xrightarrow{b_{n-1}} \tau_{\leq n-1}(H) \xrightarrow{i''} H''$ , where  $i''$  is an isomorphism.

III. Assume that  $\Phi : \underline{\mathcal{C}}^{op} \times \underline{\mathcal{C}}' \rightarrow \underline{\mathcal{A}}$  is a duality (in the sense of Definition 1.1.1) and  $H = \Phi(-, M)$  for some object  $M$  of  $\underline{\mathcal{C}}'$ . Then the transformation  $a_n$  corresponding to  $H$  equals  $\Phi(-, f)$  for some  $\underline{\mathcal{C}}'$ -morphism  $f : L \rightarrow M$  if and only if  $b_{n-1} = \Phi(-, g)$  for a  $\underline{\mathcal{C}}'$ -morphism  $g : M \rightarrow R$ . Moreover, the morphisms  $L \xrightarrow{f} M \xrightarrow{g} R$  can be completed to a distinguished triangle in  $\underline{\mathcal{C}}'$ .

*Proof.* I.1. The two statements in the assertion are easily seen to be dual to each other; hence it suffices to consider the case where  $H'$  is of weight range  $\geq n$ . Next, the obvious functoriality of the definition of virtual  $t$ -truncations gives the following commutative square of transformations:

$$\begin{array}{ccc} \tau_{\geq n} H' & \xrightarrow{\tau_{\geq n} T} & \tau_{\geq n} H \\ \downarrow i' & & \downarrow i \\ H' & \xrightarrow{T} & H \end{array} \quad (2.2.3)$$

(cf. (2.1.1)).

Applying assertion 2 we obtain that the transformation  $i'$  is an equivalence. Hence the transformation  $\tau_{\geq n} T$  yields the factorization in question.

2. According to assertion I.1, any transformation in question factors through  $\tau_{\geq n}(H'')$ ; thus it is zero according to Proposition 2.2.2(7).

II.1. This is just a re-formulation of [Bon10b, Theorem 2.3.1(III.4)].

2. Assume that  $a = a_n \circ i'$ , where  $i'$  is an isomorphism of functors. According to the previous assertion, to prove that  $b = i'' \circ b_{n-1}$  (where  $i''$  is an isomorphism) it suffices to verify that  $H''$  is of weight range  $\leq n-1$ .

Combining our assumptions on (2.2.2) with Proposition 2.1.2(3) we obtain that for any  $M \in \text{Obj } \underline{\mathcal{C}}$  both  $H''(M)$  and  $\tau_{\leq n-1}(H)$  can be presented as certain extensions of the object  $\text{Ker } a(M[-1]) \cong \text{Ker } i'(M[-1])$  by  $\text{Coker } a(M) \cong \text{Coker } i'(M)$ . Consequently,  $H''(M) = 0$  if and only if  $\tau_{\leq n-1}(H)$ . Thus  $H''$  is of weight range  $\leq n-1$  since  $\tau_{\leq n-1}(H)$  (see Proposition 2.2.2(1)).

Similarly, if  $b = i'' \circ b_{n-1}$  then one can easily verify that  $H'(M) = 0$  if and only if  $\tau_{\geq n}(H)(M) = 0$ . Consequently,  $H'$  is of weight range  $\geq n$  since  $\tau_{\geq n}(H)$  is (see Proposition 2.2.2(1)), and applying assertion II.1 we conclude that  $a = a_n \circ i'$ .

III. Assume that  $f \in \underline{\mathcal{C}}'(L, M)$ . We complete  $f$  to a distinguished triangle

$$L \xrightarrow{f} M \xrightarrow{g} R \rightarrow L[1]. \quad (2.2.4)$$

Then the sequence  $\Phi(-, L \xrightarrow{f} M \xrightarrow{g} R)$  clearly satisfies our assumptions on (2.2.2) in assertion II. Then assertion II.2 implies that  $b_{n-1} = \Phi(-, g)$ . We also obtain the "moreover" statement in our assertion.

Conversely, if  $b_{n-1} = \Phi(-, g)$  then we also can complete  $g$  to a distinguished triangle of the form (2.2.4). We apply assertion II.2 once again to obtain  $a_n = \Phi(-, f)$ .  $\square$

We also prove a simple statement that was used in [BoS19].

**Proposition 2.2.5.** For  $M \in \text{Obj } \underline{\mathcal{C}}$  the following conditions are equivalent.

- (i)  $M \in \underline{\mathcal{C}}_{w \geq 0}$ .
- (ii)  $H(M) = 0$  for any cohomological functor  $H$  from  $\underline{\mathcal{C}}$  into (an abelian category)  $\underline{\mathcal{A}}$  that is of weight range  $\leq -1$ .
- (iii)  $(\tau_{\leq -1} H_N)(M) = \{0\}$  for any  $N \in \text{Obj } \underline{\mathcal{C}}$ .
- (iv)  $(\tau_{\leq -1} H_M)(M) = \{0\}$ .

*Proof.* Condition (i) implies condition (ii) by definition; clearly, (iii)  $\implies$  (iv). Next, condition (ii) implies condition (iii) according to Proposition 2.2.2(1).

Lastly, if  $(\tau_{\leq -1} H_M)(M) = \{0\}$  then the long exact sequence (2.1.1) yields that  $(\tau_{\geq -0} H_M)(M)$  surjects onto  $\underline{\mathcal{C}}(M, M)$ . Hence the morphism  $\text{id}_M$  factors through  $w_{\geq 0} M$ ; thus  $M$  belongs to  $\underline{\mathcal{C}}_{w \geq 0}$ .  $\square$

### 2.3 Some properties and restrictions of strictly orthogonal $t$ -structures

**Proposition 2.3.1.** Assume that  $\underline{\mathcal{C}}', \underline{\mathcal{C}} \subset \underline{\mathcal{D}}$  and  $w$  is a weight structure on  $\underline{\mathcal{C}}$ . Take

$$C_1 = \underline{\mathcal{C}}_{w \leq -1}^{\perp \underline{\mathcal{D}}} \cap \text{Obj } \underline{\mathcal{C}}'; \quad C_2 = \underline{\mathcal{C}}_{w \geq 0}^{\perp \underline{\mathcal{D}}} \cap \text{Obj } \underline{\mathcal{C}}' \quad (2.3.1)$$

1. Suppose that  $C_1 \perp C_2$  and a  $t$ -structure  $t$  on  $\underline{\mathcal{C}}'$  is orthogonal to  $w$ . Then  $t$  is also strictly orthogonal to  $w$ ; thus it equals  $(C_2[1], C_1)$ .
2. Suppose that a  $t$ -structure  $t$  on  $\underline{\mathcal{C}}'$  is strictly orthogonal to  $w$ . Then  $C_1 \perp C_2$ .

Consequently,  $t$  is the only  $t$ -structure on  $\underline{\mathcal{C}}'$  that is orthogonal to  $w$ .

*Proof.* 1. We should verify that  $t = (C_2[1], C_1)$ . The definition of orthogonality says that  $\underline{\mathcal{C}}'_{t \leq 0} \subset C_2[1]$  and  $\underline{\mathcal{C}}'_{t \geq 0} \subset C_1$ . On the other hand, recall that  $\underline{\mathcal{C}}'_{t \geq 0} = {}^{\perp \underline{\mathcal{C}}'} \underline{\mathcal{C}}'_{t \leq -1}$  and  $\underline{\mathcal{C}}'_{t \leq 0} = (\underline{\mathcal{C}}'_{t \geq 1})^{\perp \underline{\mathcal{C}}'}$  (see Proposition 1.1.5(4)). Since  $C_1 \perp C_2$ , we obtain that the converse inclusions are valid as well.

2. Since  $(C_2[1], C_1)$  is a  $t$ -structure,  $C_1 \perp C_2$ ; see the orthogonality axiom (iii) in Definition 1.1.3. Next, if a  $t$ -structure  $t'$  on  $\underline{\mathcal{C}}'$  is orthogonal to  $w$  as well, then assertion 1 implies that  $t' = t$ .  $\square$

*Remark 2.3.2.* 1. Let us formulate a simple general statement that we essentially used in the proof of Proposition 2.3.1(1); we will apply it below.

Assume that  $C_1, C_2, C'_1, C'_2 \subset \text{Obj } \underline{\mathcal{C}}'$  (for some triangulated category  $\underline{\mathcal{C}}'$ ),  $C_1 \perp C_2$ ,  $C'_1 \subset C_1$ ,  $C'_2 \subset C_2$ ,  $C'_2 = C'_1{}^\perp$ , and  $C'_1 = {}^\perp C'_2$ . Then we clearly have  $C'_1 \supset C_1$  and  $C'_2 \supset C_2$ ; thus  $C'_1 = C_1$  and  $C'_2 = C_2$ .

2. In Definition 3.2 of [SaS11] a couple  $(C'_1, C'_2)$  such that  $C'_1, C'_2 \subset \text{Obj } \underline{\mathcal{C}}'$ ,  $C'_2 = C'_1{}^\perp$ , and  $C'_1 = {}^\perp C'_2$ , was said to be a *Hom-orthogonal pair*. It is well known that Hom-orthogonal pairs are closely related both to  $t$ -structures and to weight structures; see Definition 3.6 of *ibid.* and §3 of [BoV19].

Note here that the main object of study of *ibid.* are Hom-orthogonal pairs satisfying a certain "decomposition" axiom very similar to axioms (iv) in Definitions 1.2.1 and 1.1.3. They are said to be *torsion theories*; see Definition 2.1 of *ibid.*

Now we prove our first abstract criterion on the existence of an orthogonal  $t$ -structure.

**Theorem 2.3.3.** Assume that  $\underline{\mathcal{C}}' \subset \underline{\mathcal{C}}$ ,  $\underline{\mathcal{D}} = \underline{\mathcal{C}}$ , and  $w$  is a weight structure on  $\underline{\mathcal{C}}$ .

- I. Then the following conditions are equivalent.
  - (i). There exists a  $t$ -structure  $t$  on  $\underline{\mathcal{C}}'$  orthogonal to  $w$ .
  - (ii). There exists a  $t$ -structure  $t$  on  $\underline{\mathcal{C}}'$  strictly orthogonal to  $w$ .
  - (ii'). There exists a  $t$ -structure  $t$  on  $\underline{\mathcal{C}}'$  strictly orthogonal to  $w$ , and this is the only  $t$ -structure  $t$  on  $\underline{\mathcal{C}}'$  orthogonal to  $w$ .
  - (iii). The functor  $\tau_{\geq 0} H_M^{\underline{\mathcal{C}}} = \tau_{\geq 0} H_M$  is  $\underline{\mathcal{C}}$ -representable by an object of  $\underline{\mathcal{C}}'$  for any object  $M$  of  $\underline{\mathcal{C}}'$  (that is,  $\tau_{\geq 0} H_M \cong H_L$  for some object  $L$  of  $\underline{\mathcal{C}}'$ ).

(iv). If  $M$  is an object of  $\underline{\mathcal{C}}'$  then the functor  $\tau_{\leq -1}H_M$  is isomorphic to  $H_R$  for some  $R \in \text{Obj } \underline{\mathcal{C}}'$ .

(v). For any object  $M$  of  $\underline{\mathcal{C}}'$  and  $i \in \mathbb{Z}$  the functors  $\tau_{\geq i}H_M$  and  $\tau_{\leq i}H_M$  are  $\underline{\mathcal{C}}$ -representable by objects of  $\underline{\mathcal{C}}'$ .

II. If the conditions of assertion I are fulfilled, then the corresponding Yoneda-type functor  $\underline{H}t \rightarrow \text{AddFun}(\underline{H}w^{op}, \text{Ab})$ ,  $M \mapsto H_M^{\underline{\mathcal{C}}}$  (see Definition 2.1.1(3)) is fully faithful.

Consequently, if the category  $\underline{H}w$  is  $R$ -linear then  $\underline{H}t$  embeds into  $\text{Fun}_R(\underline{H}w^{op}, R\text{-Mod})$  via the equivalence provided by Lemma 2.1.4(1).

*Proof.* Condition (ii) clearly implies condition (i). Conditions (ii) and (ii') are equivalent according to Proposition 2.3.1. Next, Proposition 1.2.4(2) implies that  $C_2 \subset \underline{\mathcal{C}}_{w \leq -1}$  (see (2.3.1) for the notation); hence  $C_1 \perp C_2$  by the definition of  $C_2$ . Hence Proposition 2.3.1 implies that (i)  $\implies$  (ii).

Let us verify that condition (iv) implies (ii). So, we should check that the couple  $(C_2[1], C_1)$  is a  $t$ -structure. Now, these classes are clearly closed with respect to  $\underline{\mathcal{C}}'$ -isomorphisms, and the "shift" axiom (ii) of Definition 1.1.3 is obviously fulfilled as well. Moreover, we have just proved that  $C_1 \perp C_2$ ; this gives the orthogonality axiom (iii). Hence it remains to check the existence of  $t$ -decompositions (this is axiom (iv) of  $t$ -structures). For an object  $M$  of  $\underline{\mathcal{C}}'$  we assume  $\tau_{\leq -1}H_M \cong H_R$  for some  $R \in \text{Obj } \underline{\mathcal{C}}'$ . Take  $\Phi$  to be the duality provided by Lemma 1.1.2. By the Yoneda lemma, the transformation  $b_{-1}$  in (2.1.1) equals  $\Phi(-, g)$  for some  $g \in \underline{\mathcal{C}}'(M, R)$ . Then Proposition 2.2.4(III) gives a  $\underline{\mathcal{C}}'$ -distinguished triangle  $L \rightarrow M \rightarrow R \rightarrow L[1]$ , where  $L \in \text{Obj } \underline{\mathcal{C}}'$  and  $H_L \cong \tau_{\geq 0}H$ . It remains to note that  $L \in C_1$  and  $R \in C_2$ ; see Proposition 2.2.2(1).

Next, (i) implies condition (v) according to Proposition 2.1.2(4). Moreover, conditions (v) clearly implies conditions (iii) and (iv).

Lastly, assume that our condition (iii) is fulfilled; we fix  $M \in \text{Obj } \underline{\mathcal{C}}'$  and assume that  $\tau_{\geq 0}H \cong H_L = \Phi(-, L)$  for some  $L \in \text{Obj } \underline{\mathcal{C}}'$ . We argue similarly to the proof of (iv)  $\implies$  (ii) above. By the Yoneda lemma, the corresponding transformation  $a_0 : \tau_{\geq 0}H \rightarrow L$  can be presented as  $\Phi(-, f)$  for some  $f \in \underline{\mathcal{C}}'(L, M)$ . Then Proposition 2.2.4(III) yields that  $\tau_{\leq -1}H_M \cong H_R$  for the corresponding  $R$ . Thus condition (iii) implies (iv), and this concludes the proof.

II. If  $M \in \underline{\mathcal{C}}'_{t=0}$  then the definition of orthogonality implies that the functor  $H_M : \underline{\mathcal{C}}^{op} \rightarrow \text{Ab}$  (see Definition 2.1.1(3)) is of weight range  $[0, 0]$ . Thus it suffices to apply Proposition 2.2.2(5) along with the Yoneda lemma and Lemma 2.1.4(1).  $\square$

Now we will describe certain restrictions of  $t$  that is strictly orthogonal to  $w$ . We start from some notation.

**Definition 2.3.4.** Throughout this paper  $R$  will be an associative commutative unital ring. Assume that  $\underline{\mathcal{C}}^0, \underline{\mathcal{C}} \subset \underline{\mathcal{D}}$  and  $\underline{\mathcal{C}}^0$  is  $R$ -linear.

1. Then for  $M \in \text{Obj } \underline{\mathcal{C}}^0$ ,  $N \in \text{Obj } \underline{\mathcal{D}}$  we define the structure of an  $R$ -module on  $\underline{\mathcal{D}}(M, N)$  as follows: to multiply morphisms in this group by  $r \in R$  we compose them with  $r \text{id}_M$ .

2. Let  $\underline{\mathcal{A}}$  be a weak Serre subcategory of  $R\text{-Mod}$ . Then we will write  $\underline{\mathcal{C}}_{\underline{\mathcal{A}}}$  for the full subcategory of  $\underline{\mathcal{C}}$  that consists of those  $N \in \text{Obj } \underline{\mathcal{C}}$  such that for any  $M \in \text{Obj } \underline{\mathcal{C}}^0$  the  $R$ -module  $\underline{\mathcal{D}}(M, N)$  belongs to  $\underline{\mathcal{A}}$ .

3. In the case where  $R$  is Noetherian,  $\beta$  is an infinite cardinal, and  $\underline{A}$  is the category of  $R$ -modules with less than  $\beta$  generators (see Proposition 2.3.5(3) below) we will modify this notation and write  $\underline{C}_\beta$  for the corresponding  $\underline{C}_\underline{A} \subset \underline{C}$ .

**Proposition 2.3.5.** Assume that  $\underline{C}^0, \underline{C} \subset \underline{D}$ ,  $\underline{C}^0$  is  $R$ -linear, a weight structure  $w^0$  on  $\underline{C}^0$  is strictly orthogonal to a  $t$ -structure  $t$  on  $\underline{C}$ , and  $\underline{A}$  and  $\underline{A}'$  are weak Serre subcategories of  $R - \text{Mod}$ .

1. Then  $\underline{C}_\underline{A}$  is triangulated and  $t$  restricts to it. The heart  $\underline{Ht}_\underline{A}$  of this restriction  $t_\underline{A}$  consists of those objects of  $\underline{Ht}$  such that the corresponding functors  $\underline{Hw}^{0op} \rightarrow R - \text{Mod}$  (see Proposition 2.2.2(5,10) and Lemma 2.1.4(1)) take values in  $\underline{A}$ .

2.  $\underline{C}_{\underline{A} \cap \underline{A}'} = \underline{C}_{\underline{A} \cap \underline{A}'} \cap \underline{C}_{\underline{A} \cap \underline{A}'}$  (cf. Remark 1.1).

3. If  $R$  is Noetherian then the category of  $R$ -modules with less than  $\beta$  generators is a weak Serre subcategory of  $R - \text{Mod}$ .

*Proof.* 1. Since for any  $M \in \text{Obj } \underline{C}$  the functor  $\underline{D}(M, -)$  is homological,  $\underline{C}_\underline{A}$  is triangulated indeed.

Next, to verify that  $t$  restricts to  $\underline{C}_\underline{A}$  it suffices to that for any  $M \in \text{Obj } \underline{C}_\underline{A}$  the object  $L_t M = t_{\geq 0} M$  belongs to  $\underline{C}_\underline{A}$  as well; see Lemma 1.1.7. The latter statement easily follows from Proposition 2.1.3(II.3) along with Lemma 2.1.4(3).

Lastly, any object of  $\underline{Ht}_\underline{A}$  yields a functor  $\underline{Hw}^{0op} \rightarrow \underline{A}$  just by the definition of  $\underline{C}_\underline{A}$ . Conversely, if an object of  $\underline{Ht}$  gives a functor  $\underline{Hw}^{0op} \rightarrow \underline{A}$  then the corresponding functor  $\underline{C}^{0op} \rightarrow R - \text{Mod}$  takes values in  $\underline{A}$  as well; see Proposition 2.2.2(9).

2. Obvious.

3. A well-known fact. □

We will describe some geometric examples for this proposition later.

## 2.4 On coproductive extensions of weight structures

In the following definition we do not assume that  $\underline{C}$  and  $\underline{A}$  are closed with respect to (small) coproducts.

**Definition 2.4.1.** 1. Let  $w^0$  be a weight structure on  $\underline{C}^0 \subset \underline{C}$ . We will say that (a weight structure)  $w = (\underline{C}_{w \leq 0}, \underline{C}_{w \geq 0})$  is the coproductive extension of  $w^0$  to  $\underline{C}$  if  $\underline{C}_{w \leq 0}$  (resp.  $\underline{C}_{w \geq 0}$ ) equals the smallest retraction-closed and extension-closed class of objects of  $\underline{C}$  that is closed with respect to  $\underline{C}$ -coproducts (that exist in  $\underline{C}$ ) and contains  $\underline{C}_{w^0 \leq 0}^0$  (resp.  $\underline{C}_{w^0 \geq 0}^0$ ).

2. It will be convenient for us to use the following somewhat clumsy terminology: a cohomological functor  $H'$  from  $\underline{C}$  into  $\underline{A}$  will be called a *cp* functor if it converts all  $\underline{C}$ -coproducts (that is, all those small coproduct diagrams that exist in  $\underline{C}$ ) into  $\underline{A}$ -products.

*Remark 2.4.2.* 1. If  $w$  is the coproductive extension of  $w^0$  to  $\underline{C}$  and  $w'$  is an arbitrary extension of  $w^0$  to  $\underline{C}$  then Proposition 1.2.4(2) is easily seen to imply that  $\underline{C}_{w \leq 0} \subset \underline{C}_{w' \leq 0}$ ; cf. Proposition 1.2.4(3). Moreover, if  $w$  is an extension of  $w^0$  to  $\underline{C}$  such that  $\underline{C}_{w \leq 0} \subset \underline{C}_{w' \leq 0}$  for any other extension  $w'$  then this assumption determines it canonically (since  $\underline{C}_{w \geq 0} \supset \underline{C}_{w' \geq 0}$ ). Yet the existence of  $w$  of this sort does not imply that  $w$  is the coproductive extension of  $w^0$  to  $\underline{C}$  since the coproductive extension of  $w^0$  to  $\underline{C}$  does not have to exist. Indeed, the

coproductive extension of  $w^0$  does not exist in the case  $\underline{C}^0 = \{0\}$  and  $\underline{C} \neq \{0\}$ ; yet  $w = (0, \underline{C})$  clearly satisfies the aforementioned "minimality" property.

2. The author is not sure that Definition 2.4.1(1) is really clever. However, one can construct plenty of examples for it (see Theorem 3.3.2 below), and it also fits well with the following simple statements.

**Proposition 2.4.3.** Assume that  $\underline{C}^0 \subset \underline{C}$ ,  $w^0$  is a weight structure on  $\underline{C}^0$ ,  $w$  is its coproductive extension to  $\underline{C}$ .

1. Let  $H$  be cp (cohomological) functor from  $\underline{C}$  into  $\underline{A}$ .

Then  $H$  is of  $w$ -weight range  $\geq 0$  (resp.  $\leq 0$ ) if and only if its restriction to  $\underline{C}^0$  is of  $w^0$ -weight range  $\geq 0$  (resp.  $\leq 0$ ).

2. Assume that  $\underline{C}, \underline{C}' \subset \underline{D}$ .

Then a  $t$ -structure  $t$  on  $\underline{C}'$  is (strictly) orthogonal to  $w$  if and only if it is (strictly) orthogonal to  $w^0$ .

*Proof.* 1. The "only if" part of the assertion is obvious; it only requires  $w$  to be an extension of  $w^0$ .

The converse implication immediately from the descriptions of  $\underline{C}_{w \leq -1}$  and  $\underline{C}_{w \geq 1}$  provided by Definition 2.4.1.

2. It clearly suffices to verify that the classes  $C_1 = \underline{C}_{w \leq -1}^{\perp \underline{D}} \cap \text{Obj } \underline{C}'$  and  $C_2 = \underline{C}_{w \geq 0}^{\perp \underline{D}} \cap \text{Obj } \underline{C}'$  (cf. (2.3.1)) coincide with

$$C_1^0 = \underline{C}_{w^0 \leq -1}^{\perp \underline{D}} \cap \text{Obj } \underline{C}' \text{ and } C_2^0 = \underline{C}_{w^0 \geq 0}^{\perp \underline{D}} \cap \text{Obj } \underline{C}' \quad (2.4.1)$$

respectively. The latter easy statement can be obtained by means of applying assertion 1 to functors from  $\underline{C}$  that are represented by objects of  $\underline{C}'$ .  $\square$

**Corollary 2.4.4.** Assume that  $\underline{C}^0, \underline{C}' \subset \underline{C}$ ,  $w^0$  is a weight structure on  $\underline{C}^0$  and  $w$  is its coproductive extension to  $\underline{C}$ .

Then the equivalent conditions of Theorem 2.3.3(I) are also equivalent to the existence of a  $t$ -structure  $t$  on  $\underline{C}'$  that is orthogonal to  $w^0$ . Moreover, this  $t$ -structure equals  $(C_2^0[1], C_1^0)$  in the notation of (2.4.1), that is,  $t$  is strictly orthogonal to  $w^0$ .

*Proof.* This is an obvious combination of Proposition 2.4.3(2) with (the equivalence of conditions (i) and (ii) in) Theorem 2.3.3(I).  $\square$

*Remark 2.4.5.* Corollary 2.4.4 is a certain substitute for Theorem 2.2.5 of [Bon19]; see Remark 0.3 above. Though the author suspects that the results of the current text are not sufficient to prove loc. cit. itself, they can be successfully applied to treat all the examples of loc. cit. known to the author; cf. Proposition 3.3.5 below.

### 3 On $t$ -structures orthogonal to (co)smashing weight structures

In this section we study the existence of adjacent weight and  $t$ -structures in triangulated categories closed with respect to (co)products (these are called smashing and cosmashing ones).

In §3.1 we consider smashing weight structures (on smashing triangulated categories); these are the ones that "respect coproducts".

In §3.2 we recall the notion of Brown representability for smashing triangulated categories, and prove Theorem 0.1, i.e., that a weight structure  $w$  on a category satisfying this condition is left adjacent to a  $t$ -structure if and only if  $w$  is smashing.

In §3.3 we study extensions of weight structures from subcategories of compact objects and the corresponding orthogonal  $t$ -structures.

In §3.4 we formulate the dual to Theorem 0.1 and give some applications for it. If  $w$  is both cosmashing and smashing then the left adjacent  $t$ -structure  $t$  restricts to the subcategory of compact objects of  $\underline{\mathcal{C}}$  as well as to all other "levels of smallness" for objects. Combining this statement with an existence of weight structures theorem from [KeN13] we obtain a statement on  $t$ -structures extending yet another result of *ibid.* We will not apply these results in the current paper.

### 3.1 On smashing weight structures and related notions

We will need a few definitions.

- Definition 3.1.1.** 1. We will say that a triangulated category  $\underline{\mathcal{C}}$  is *(co)smashing* if it is closed with respect to (small) coproducts (resp., products).
2. We will say that a weight structure  $w$  on  $\underline{\mathcal{C}}$  is *(co)smashing* if  $\underline{\mathcal{C}}$  is *(co)smashing* and the class  $\underline{\mathcal{C}}_{w \geq 0}$  is closed with respect to  $\underline{\mathcal{C}}$ -coproducts (resp.,  $\underline{\mathcal{C}}_{w \leq 0}$  is closed with respect to  $\underline{\mathcal{C}}$ -products; cf. Proposition 1.2.4(3)).
3. We will say that a  $t$ -structure  $t$  on  $\underline{\mathcal{C}}$  is *(co)smashing* if  $\underline{\mathcal{C}}$  is *(co)smashing* and the class  $\underline{\mathcal{C}}_{t \leq 0}$  is closed with respect to  $\underline{\mathcal{C}}$ -coproducts (resp.,  $\underline{\mathcal{C}}_{t \geq 0}$  is closed with respect to  $\underline{\mathcal{C}}$ -products; cf. Proposition 1.1.5(4)).
4. For an infinite cardinal  $\alpha$  and a smashing  $\underline{\mathcal{C}}$  a homological functor  $H' : \underline{\mathcal{C}} \rightarrow \underline{\mathcal{A}}$  is said to be  $\alpha$ -*small* if for any family  $N_i$ ,  $i \in I$ , we have  $H'(\coprod_{i \in I} N_i) = \varinjlim_{J \subset I, \#J < \alpha} H'(\coprod_{j \in J} N_j)$  (i.e., the obvious morphisms  $H'(\coprod_{j \in J} N_j) \rightarrow H'(\coprod_{i \in I} N_i)$  form a colimit diagram; note that this colimit is filtered).

Let us now prove some properties of these notions and relate them to virtual  $t$ -truncations.

**Proposition 3.1.2.** Assume that  $w$  is a smashing weight structure on  $\underline{\mathcal{C}}$ ,  $H' : \underline{\mathcal{C}} \rightarrow \underline{\mathcal{A}}$  is a homological functor (where  $\underline{\mathcal{A}}$  is an abelian category),  $n \in \mathbb{Z}$ , and  $\alpha$  is an infinite cardinal. Then the following statements are valid.

1. If  $\alpha' \geq \alpha$  then any  $\alpha$ -small functor is also  $\alpha'$ -small.
2.  $H'$  is  $\aleph_0$ -small if and only if it respects coproducts.
3. The class  $\underline{\mathcal{C}}_{w=0}$  is closed with respect to  $\underline{\mathcal{C}}$ -coproducts.
4. Coproducts of  $w$ -decompositions are weight decompositions as well.
5. Assume that  $\underline{\mathcal{A}}$  is an AB4\* category and a cohomological functor  $H$  from  $\underline{\mathcal{C}}$  into  $\underline{\mathcal{A}}$  is a cp one. Then  $\tau_{\geq n}(H)$  and  $\tau_{\leq n}(H)$  are cp functors as well.



6. Assume that  $\underline{A}$  is an AB5 category and  $H'$  is an  $\alpha$ -small functor. Then the functors  $\tau_{\geq n}(H')$  and  $\tau_{\leq n}(H')$  are  $\alpha$ -small as well.

*Proof.* 1. Assume that  $H'$  is an  $\alpha$ -small functor; fix an index set  $I$  and certain  $N_i \in \text{Obj } \underline{C}$ . Then for any  $J \subset I$  we have  $H'(\coprod_{j \in J} N_j) = \varinjlim_{J' \subset J, \#J' < \alpha} H'(\coprod_{j' \in J'} N_{j'})$ . Combining these facts for all  $J \subset I$  one easily obtains that  $H'(\coprod_{i \in I} N_i) = \varinjlim_{J \subset I, \#J < \alpha} H'(\coprod_{j \in J} N_j)$ .

2. Since  $H'$  is additive,  $H'(\coprod N_i) = \varinjlim_{J \subset I, \#J < \aleph_0} H'(\coprod_{j \in J} N_j)$  if and only if  $H'$  respects coproducts (since this colimit will not change if one will consider only those  $J$  that consist of a single element only).

3. This is an easy consequence of Proposition 1.2.4(2); see Proposition 2.3.2(1) of [Bon21].

4. This is an easy consequence of Proposition 1.2.4(3) along with Remark 1.2.2 of [Nee01]; it is given by Proposition 2.3.2(3) of [Bon21].

5. According to Proposition 2.1.3(I.1), it suffices to verify that the functors  $\tau_{\geq 2}(H)$  and  $\tau_{\leq 0}(H)$  are cp ones for any cp functor  $H$ . For a family  $\{M_i\}$  of objects of  $\underline{C}$  we choose certain 0 and 1-weight decompositions for all of the  $M_i$  (see Remark 1.2.3(2)) and all  $j \in \mathbb{Z}$ . According to the previous assertion, for  $M = \coprod M_i$  we can take  $w_{\leq j}M = \coprod w_{\leq j}M_i$  and  $w_{\geq j}M = \coprod w_{\geq j}M_i$  for  $j = 0, 1, 2$ . Moreover, one clearly can describe the unique morphisms  $w_{\leq 0}(\coprod M_i) \rightarrow w_{\leq 1}(\coprod M_i)$  and  $w_{\geq 2}(\coprod M_i) \rightarrow w_{\geq 1}(\coprod M_i)$  compatible with these decomposition triangles (see Proposition 1.2.4(4) and Definition 2.1.1(1)) as the coproducts of the corresponding morphisms for  $M_i$ . Applying our assumptions on  $H$  and  $\underline{A}$  we obtain that  $\tau_{\geq 2}(H)(\coprod M_i) \cong \coprod \tau_{\geq 2}(H)(M_i)$  and  $\tau_{\leq 0}(H)(\coprod M_i) \cong \coprod \tau_{\leq 0}(H)(M_i)$ .

Similarly, to prove assertion 6 it suffices to verify that the functors  $\tau_{\geq 2}(H')$  and  $\tau_{\leq 0}(H')$  are  $\alpha$ -small whenever  $H'$  is. One takes the same weight decompositions along with their coproducts corresponding to all subsets  $J$  of  $I$  of cardinality less than  $\alpha$ . Since the colimits in question are filtered ones, the AB5 assumption on  $\underline{A}$  allows to compute  $\varinjlim_{J \subset I, \#J < \alpha} \text{Im}(H'(\coprod_{j \in J} w_{\leq 0}N_j) \rightarrow H'(\coprod_{j \in J} w_{\leq 1}N_j))$  and  $\varinjlim_{J \subset I, \#J < \alpha} \text{Im}(H'(\coprod_{j \in J} w_{\geq 2}N_j) \rightarrow H'(\coprod_{j \in J} w_{\geq 1}N_j))$  as the corresponding images of colimits to obtain the statement in question.  $\square$

### 3.2 On the existence of $t$ -structures adjacent to smashing weight structures

To formulate the main results of this section and discuss examples to it we need some more definitions.

**Definition 3.2.1.** Let  $\underline{C}$  be a smashing triangulated category,  $\mathcal{P}$  is a subclass of  $\text{Obj } \underline{C}$ ,  $\underline{C}'$  is an arbitrary triangulated category.

1. We will say that  $\underline{C}$  satisfies the *Brown representability* property whenever any cp functor from  $\underline{C}$  into  $\text{Ab}$  is representable.

Dually, we will say that  $\underline{C}'$  satisfies the *dual Brown representability* property if  $\underline{C}'$  is cosmashing and any functor from  $\underline{C}'$  into  $\text{Ab}$  that respects products is corepresentable (i.e., if  $\underline{C}'^{op}$  satisfies the Brown representability assumption).

2. For an infinite cardinal  $\alpha$  an object  $M$  of  $\underline{\mathcal{C}}$  is said to be  $\alpha$ -small (in  $\underline{\mathcal{C}}$ ) if the functor  $H^M = \underline{\mathcal{C}}(M, -) : \underline{\mathcal{C}} \rightarrow \mathbf{Ab}$  is  $\alpha$ -small (see Definition 3.1.1(4)). We will write  $\underline{\mathcal{C}}^{(\alpha)}$  for the (full) subcategory  $\underline{\mathcal{C}}^{(\alpha)}$  of  $\underline{\mathcal{C}}$  that consists of  $\alpha$ -small objects.  
Moreover,  $\aleph_0$ -small objects of  $\underline{\mathcal{C}}$  (those correspond to functors that respect coproducts) will also said to be *compact*.
3. We will say that a triangulated subcategory  $\underline{\mathcal{C}}^0$  of  $\underline{\mathcal{C}}$  is *localizing* whenever it is closed with respect to  $\underline{\mathcal{C}}$ -coproducts. Respectively, we will call the smallest localizing subcategory of  $\underline{\mathcal{C}}$  that contains a given class  $\mathcal{P} \subset \mathbf{Obj} \underline{\mathcal{C}}$  the *localizing subcategory of  $\underline{\mathcal{C}}$  generated by  $\mathcal{P}$* . We will say that both  $\mathcal{P}$  and the corresponding full subcategory of  $\underline{\mathcal{C}}$  *generate  $\underline{\mathcal{C}}$* .
4. We will say that a generating class of objects  $\mathcal{P}$  as well as the subcategory  $\mathcal{C}$  of  $\underline{\mathcal{C}}$  with  $\mathbf{Obj} \mathcal{C} = \mathcal{P}$  *compactly generate  $\underline{\mathcal{C}}$*  if  $\mathcal{P}$  is also essentially small and consists of compact objects (of  $\underline{\mathcal{C}}$ ).
5. We will say that a class  $\mathcal{P}' \subset \mathbf{Obj} \underline{\mathcal{C}}'$  *generates* a weight structure  $w$  (resp. a  $t$ -structure  $t$ ) on  $\underline{\mathcal{C}}'$  whenever  $\underline{\mathcal{C}}'_{w \geq 0} = (\cup_{i > 0} \mathcal{P}'[-i])^\perp$  (resp.  $\underline{\mathcal{C}}'_{t \leq 0} = (\cup_{i > 0} \mathcal{P}'[i])^\perp$ ).

*Remark 3.2.2.* 1. Recall that  $\underline{\mathcal{C}}$  satisfies both the Brown representability property and its dual whenever it is compactly generated; see Proposition 8.4.1, Proposition 8.4.2, Theorem 8.6.1, and Remark 6.4.5 of [Nee01]. Moreover, the Brown representability property is fulfilled whenever  $\underline{\mathcal{C}}$  is just  $\aleph_1$ -perfectly generated (see Definition 8.1.4 and Theorem 8.3.3 of *ibid.*).

Recall also that any triangulated category possessing a combinatorial (Quillen) model satisfies the dual Brown representability property; see §0 of [Nee08] (the statement is given by the combination of Theorems 0.17 and 0.14 of *ibid.*).

2. A class  $\mathcal{P}'$  as above is easily seen to determine weight and  $t$ -structures it generates on  $\underline{\mathcal{C}}'$  (if any) completely; see either of Proposition 2.4(1) (along with §3) of [BoV19] or Proposition 1.1.5(4) and Proposition 1.2.4(2) above.

Now we prove our first "practical" existence of  $t$ -structures results; see Definitions 3.1.1 and 3.2.1 for the notions mentioned in our theorem.

**Theorem 3.2.3.** Let  $w$  be a weight structure on  $\underline{\mathcal{C}}$ , where  $\underline{\mathcal{C}}$  (is smashing and) satisfies the Brown representability property.

Then there exists a  $t$ -structure  $t^r$  left adjacent to  $w$  if and only if  $w$  is smashing. Moreover,  $t^r$  is cosmashing (if exists) and its heart is equivalent to the category of those additive functors  $\underline{H}w^{op} \rightarrow \mathbf{Ab}$  that respect products.

*Proof.* The "only if" assertion is essentially given by Proposition 2.4(6) of [BoV19] (cf. §3. of *ibid.*; the statement is also very easy for itself).

Conversely, assume that  $w$  is smashing. According to Theorem 2.3.3(I) the existence of  $t^r$  is equivalent to the representability of  $\tau_{\leq 0}H_M$  for any representable functor  $H_M$ . Next, Proposition 3.1.2(5) says that  $\tau_{\leq 0}H_M$  is a cp functor since  $H_M$  is. Hence  $\tau_{\leq 0}H_M$  is representable by the Brown representability assumption, and we obtain that  $t^r$  exists indeed.

Next, the category  $\underline{\mathcal{C}}$  is cosmashing according to Proposition 8.4.6 of [Nee01] (since it satisfies the Brown representability property). Moreover,  $t^r = (C_2[1], C_1)$ ,

where  $C_1 = \underline{\mathcal{C}}_{w \leq -1}^\perp$  and  $C_2 = \underline{\mathcal{C}}_{w \geq 0}^\perp$  according to Theorem 2.3.3(I) (alternatively, one can apply Proposition 1.3.3). Hence the class  $\underline{\mathcal{C}}_{t^r \geq 0}$  is closed with respect to  $\underline{\mathcal{C}}$ -products; thus  $t^r$  is cosmashing as well.

Lastly, since  $t^r = (C_2[1], C_1)$ , the class  $\underline{\mathcal{C}}_{t=0}$  equals  $(\underline{\mathcal{C}}_{w \geq 1} \cup \underline{\mathcal{C}}_{w \leq -1})^\perp$ ; hence  $Ht$  can be calculated using Proposition 2.3.2(8) of [Bon21] (see also Remark 2.1.3(2) of *ibid.* and Proposition 2.2.2(5) above; cf. Theorem 2.3.3(II)).  $\square$

*Remark 3.2.4.* 1. Now let us discuss examples to Theorem 3.2.3.

According to Theorem 5 of [Pau12], any set  $\mathcal{P}$  of compact objects of  $\underline{\mathcal{C}}$  generates a (unique) smashing weight structure (see Definition 3.2.1(5) and Remark 3.2.2(2)). Moreover, Theorem 4.5(2) of [PoS16] and Theorem 3.2.1 of [Bon22a] (we will mention these statements in the proof of Corollary 3.4.3(2) below) enable one to check whether two weight structures obtained this way are distinct. Thus one may say that there are lots of smashing weight structures on  $\underline{\mathcal{C}}$  whenever there are "plenty" of compact objects in it (see Theorem 4.15 of [PoS16] for a certain justification of this claim for derived categories of commutative rings). Thus our theorem yields a rich collection of  $t$ -structures, and the author does not know of any other methods that give all of them (cf. Remark 0.2).

Other notable statements related to the construction of smashing weight structures are Theorem 2.3.4(3) of [Bon22a] (it says that any *perfect* set of objects in a smashing triangulated category generates a smashing weight structure) and Theorem 3.1.3 and 3.2.2 of [BoS19] that treat weight-exact localization functors.

2. Now let us apply our theorem in the case of semi-orthogonal decompositions (see Definition 1.2.5 and Remark 1.2.6).

If  $w$  is a semi-orthogonal decomposition of  $\underline{\mathcal{C}}$  (where  $\underline{\mathcal{C}}$  satisfies the Brown representability property) that is smashing as a weight structure then for the corresponding  $t^r$  the couple  $(\underline{\mathcal{C}}_{t^r \geq 0}, \underline{\mathcal{C}}_{t^r \leq 0})$  is a semi-orthogonal decomposition as well; see Proposition 1.3.5. Note also that  $(\underline{\mathcal{C}}_{t^r \geq 0}, \underline{\mathcal{C}}_{t^r \leq 0})$  is a (cosmashing) weight structure.

Next, recall that the full subcategory  $\underline{\mathcal{C}}_1$  of  $\underline{\mathcal{C}}$  corresponding to  $C_1 = \underline{\mathcal{C}}_{w \leq 0}$  is triangulated and the corresponding right adjoint (to the embedding  $\underline{\mathcal{C}}_1 \rightarrow \underline{\mathcal{C}}$ ) respects coproducts; see condition 3 in Proposition 1.3.5 and Remark 1.2.7(3). Since  $w^{op}$  is a semi-orthogonal decomposition in the category  $\underline{\mathcal{C}}^{op}$  (see Proposition 1.2.4(1)), we obtain that the embedding  $\underline{\mathcal{C}}_2 \rightarrow \underline{\mathcal{C}}$  possesses a left adjoint; here  $\underline{\mathcal{C}}_2$  the subcategory of  $\underline{\mathcal{C}}$  corresponding to  $\underline{\mathcal{C}}_{w \geq 0}$ .

Thus if we apply Proposition 1.3.5 to the semi-orthogonal decomposition  $(\underline{\mathcal{C}}_{t^r \geq 0}, \underline{\mathcal{C}}_{t^r \leq 0})$  we obtain that the embedding  $\underline{\mathcal{C}}_2 \rightarrow \underline{\mathcal{C}}$  possesses a right adjoint as well; note that  $\underline{\mathcal{C}}_{t^r \geq 0} = \underline{\mathcal{C}}_{w \geq 0}$ . Thus  $\underline{\mathcal{C}}_2$  is *admissible* in  $\underline{\mathcal{C}}$  in the sense of [BoK89, Definition 2.5] and the embedding  $\underline{\mathcal{C}}' \rightarrow \underline{\mathcal{C}}$  may be completed to a *gluing datum* (cf. [BBD82, §1.4] or [Nee01, §9.2]).

So we generalize Corollary 2.4 of [NiS09] to arbitrary categories that satisfy the Brown representability property.

### 3.3 Weight structures extended from subcategories of compact objects, and orthogonal $t$ -structures

To construct certain weight structures we need the following statements that appear to be rather well-known.

**Proposition 3.3.1.** Assume that  $A$  and  $B$  are extension-closed classes of objects of  $\underline{\mathcal{C}}$ .

I. Assume that  $A \perp B[1]$ . Then the class  $A \star B$  (of all extensions of elements of  $B$  by elements of  $A$ ) is extension-closed as well.

II. Assume in addition that  $\underline{\mathcal{C}}$  is smashing, and  $A$  and  $B$  are closed with respect to  $\underline{\mathcal{C}}$ -coproducts.

1. Then  $A \star B$  is closed with respect to  $\underline{\mathcal{C}}$ -coproducts as well.

2. Assume that  $A$  is closed either with respect to  $[-1]$  or with respect to  $[1]$ .

Then  $A$  is retraction-closed in  $\underline{\mathcal{C}}$ .

*Proof.* All of these statements are rather easy.

Assertions I and II.1 immediately follow from Proposition 2.1.1 of [BoS19]. Assertion II.2 is a straightforward consequence of Corollary 2.1.3(2) of *ibid.* (see Remark 2.1.4(4) of *ibid.*)  $\square$

Now we prove that weight structures extend from subcategories of compact objects to the localizing subcategories they generate. The (proof of the) first part of the following theorem is quite similar to the corresponding arguments in §2 of [BoS19].

**Theorem 3.3.2.** Assume that  $\underline{\mathcal{C}}$  is a smashing triangulated category, a subcategory  $\underline{\mathcal{C}}^0$  of  $\underline{\mathcal{C}}^{(\aleph_0)}$  generates  $\underline{\mathcal{C}}$ , and  $w^0$  is a weight structure on  $\underline{\mathcal{C}}^0$ .

I.1. Then  $w^0$  extends uniquely to a smashing weight structure  $w$  on  $\underline{\mathcal{C}}$ .

2.  $w$  is the coproductive extension of  $w^0$  to  $\underline{\mathcal{C}}$ .

3.  $\underline{\mathcal{C}}_{w=0}$  consists of all retract of all (small)  $\underline{\mathcal{C}}$ -coproducts of elements of  $\underline{\mathcal{C}}_{w^0=0}^0$ .

4. Assume that  $\alpha$  is a *regular* cardinal, that is,  $\alpha$  cannot be presented as a sum of less than  $\alpha$  cardinals that are less than  $\alpha$ . Take  $\underline{\mathcal{C}}_\alpha$  to be the smallest triangulated subcategory of  $\underline{\mathcal{C}}$  that contains  $\underline{\mathcal{C}}^0$  and is closed with respect to coproducts of less than  $\alpha$  objects. Then there exists a coproductive extension  $w_\alpha$  of  $w^0$  to  $\underline{\mathcal{C}}_\alpha$ . Moreover,  $\underline{\mathcal{C}}_{w_\alpha=0}$  consists of all retracts of all coproducts of less than  $\alpha$  elements of  $\underline{\mathcal{C}}_{w^0=0}^0$ .

II. Assume in addition that the category  $\underline{\mathcal{C}}^0$  is essentially small.

1. Then there exists a  $t$ -structure  $t$  (left) adjacent to  $w$  (on  $\underline{\mathcal{C}}$ ).

2.  $t$  is strictly orthogonal to  $w^0$ ; hence  $t$  is both smashing and cosmashing.

3.  $\underline{Ht}$  is equivalent (in the obvious way) to the category  $\text{AddFun}(\underline{Hw}^{0op}, \text{Ab})$ .

*Proof.* I.1,2. Proposition 1.2.4(9) clearly yields the following: if a coproductive extension of  $w^0$  to  $\underline{\mathcal{C}}$  exists then it equals the only smashing extension of  $w^0$  to  $\underline{\mathcal{C}}$ .

Set  $E_1$  (resp.  $E_2$ ) for the smallest extension-closed subclass of  $\text{Obj } \underline{\mathcal{C}}$  that is closed with respect to coproducts and contains  $\underline{\mathcal{C}}_{w^0 \leq 0}^0$  (resp.  $\underline{\mathcal{C}}_{w^0 \geq 0}^0$ ). To prove our assertions, it suffices to verify that  $(E_1, E_2)$  is a weight structure on  $\underline{\mathcal{C}}$ .

Firstly, axiom (ii) of Definition 1.2.1 (for  $w^0$ ) easily implies that  $E_1 \subset E_1[1]$  and  $E_2[1] \subset E_2$ . Combining this statement with Proposition 3.3.1(II.2) we obtain that  $E_1$  and  $E_2$  are retraction-closed in  $\underline{\mathcal{C}}$ .

Next, the compactness of the elements of  $\underline{\mathcal{C}}_{w^0 \leq 0}^0$  in  $\underline{\mathcal{C}}$  implies that the class  $\underline{\mathcal{C}}_{w^0 \leq 0}^0 \perp$  is closed with respect to coproducts. Since it is also extension-closed and contains  $\underline{\mathcal{C}}_{w^0 \geq 1}^0$  by the axiom (iii) of Definition 1.2.1, this orthogonal contains  $E_2[1]$ , i.e.,  $\underline{\mathcal{C}}_{w \leq 0} \perp E_2[1]$ . Thus  $\underline{\mathcal{C}}_{w \leq 0} \subset \perp E_2[1]$ , and since the latter class

is closed with respect to coproducts and extensions, we obtain that  $E_1 \perp E_2[1]$  (cf. the proof of [BoK18, Lemma 1.1.1(2)]).

Let us now prove the existence of weight decompositions, i.e., for  $E = E_1 \star E_2$  we should prove  $E = \text{Obj } \underline{\mathcal{C}}$ . Now,  $E$  clearly contains  $\text{Obj } \underline{\mathcal{C}}^0$ , and Proposition 3.3.1(I, II.1) implies that it is also extension-closed and closed with respect to  $\underline{\mathcal{C}}$ -coproducts. Hence  $E = \text{Obj } \underline{\mathcal{C}}$ , and we obtain that  $w = (E_1, E_2)$  is a weight structure on  $\underline{\mathcal{C}}$  indeed. Clearly, this weight structure is smashing.

3. Denote our candidate for  $\underline{\mathcal{C}}_{w=0}$  by  $C$ . Firstly we note that  $C \subset \underline{\mathcal{C}}_{w=0}$  since the latter class is closed with respect to coproducts according to Proposition 3.1.2(3).

Applying Proposition 1.2.4(5) we obtain that  $C$  is extension-closed (since this class is clearly additive); obviously, it is also closed with respect to coproducts.

Next we apply Proposition 3.3.1(I, II.1) once again to obtain that the class  $C \star \underline{\mathcal{C}}_{w \geq 1}$  is extension-closed and closed with respect to coproducts; hence this class coincides with  $\underline{\mathcal{C}}_{w \geq 0}$ . Thus for any  $M \in \underline{\mathcal{C}}_{w=0}$  there exists its weight decomposition  $LM \rightarrow M \rightarrow RM \rightarrow LM[1]$  with  $LM \in C$ . Since  $M$  is a retract of  $LM$  according to Proposition 1.2.4(6), we obtain that  $M \in C$ .

4. The proof is similar to that of the preceding assertions; cf. also the proof of [BoS19, Theorem 2.2.1].

II.1. The category  $\underline{\mathcal{C}}$  is compactly generated by  $\underline{\mathcal{C}}^0$  in this case; hence  $\underline{\mathcal{C}}$  satisfies the Brown representability property (see Remark 3.2.2(1)). Next,  $w$  is smashing; thus a  $t$ -structure  $t$  adjacent to it exists according to Theorem 3.2.3.

2.  $t$  is cosmashing according to Theorem 3.2.3. It is strictly orthogonal to  $w^0$  by Corollary 2.4.4; thus  $\underline{\mathcal{C}}_{t \leq 0} = \underline{\mathcal{C}}_{w^0 \geq 1}^\perp$ . Since for any object  $N$  of  $\underline{\mathcal{C}}$  the class  $N^\perp$  is closed with respect to coproducts,  $t$  is smashing.

3. According to Theorem 3.2.3, the category  $\underline{Ht}$  is equivalent to the category of those functors from  $\underline{Hw}^{op}$  into  $\text{Ab}$  that respect products. Thus it remains to apply the description of  $\underline{Hw}$  provided by assertion I.3.  $\square$

*Remark 3.3.3.* 1. According to the well-known Lemma 4.4.5 of [Nee01], the assumptions of Theorem 3.3.2 imply that  $\underline{\mathcal{C}}^{(\aleph_0)}$  is the retraction-closure (see §1.1) of the category  $\underline{\mathcal{C}}^0$ . Note however that  $w^0$  does not extend to  $\underline{\mathcal{C}}^{(\aleph_0)}$  in general (even though it extends to  $\underline{\mathcal{C}}!$ ; see §3.1 of [BoS18].<sup>3</sup> Consequently, it does make some sense not to assume  $\underline{\mathcal{C}}^0 = \underline{\mathcal{C}}^{(\aleph_0)}$  in our theorem.

For the same reason, the weight structures provided by our theorem don't have to be *strongly  $\aleph_0$ -well generated* in the sense of Remark 3.3.4(1) of [Bon22a] (even if one assumes that  $\underline{\mathcal{C}}^0$  is essentially small; note that this is equivalent to the compact generation of  $\underline{\mathcal{C}}$ ). This observation possibly suggests that the notion of strong  $\aleph_0$ -well generation should be slightly generalized.

2. The restriction of our theorem to the case where  $w^0$  is bounded was essentially established in §4.5 of [Bon10a]; cf. Theorem 3.2.2 of [Bon21] and Remark 2.3.2(2) of [BoS19] for some more detail.

3. Note that one can easily construct plenty of examples for our theorem such that  $w^0$  is unbounded.

Indeed, there exist lots of unbounded weight structures on (essentially) small triangulated categories; in particular, one can start from a (say, bounded) weight structure on some small non-zero  $\underline{\mathcal{C}}^0 \subset \underline{\mathcal{C}}^{(\aleph_0)} \subset \underline{\mathcal{C}}$ , choose a regular  $\alpha > \aleph_0$ ,

<sup>3</sup>In any example of this sort  $w^0$  is neither bounded above nor below; see Theorem 2.2(II.2) of *ibid*.

and apply Theorem 3.3.2(I.4). Next, if  $\underline{\mathcal{C}}^0 = h\mathcal{C}_0$  is the homotopy category of a small stable  $\infty$ -category  $\mathcal{C}_0$  then one can take  $\underline{\mathcal{C}} = h\operatorname{Ind} \mathcal{C}_0$ ; see Remark 1.4.4.3 of [Lur17] (cf. also the proof of Corollary 1.4.4.2 of loc. cit.). Consequently, if  $\underline{\mathcal{C}}^0 = h\mathcal{C}_0$  is a non-zero small category endowed with a weight structure  $w^0$  then one can extend it to an unbounded weight structure on some triangulated category  $\underline{\mathcal{C}}'_0 \supset \underline{\mathcal{C}}^0$ , and  $\underline{\mathcal{C}}'_0 \subset \underline{\mathcal{C}}'^{(\aleph_0)} \subset \underline{\mathcal{C}}'$  for some compactly generated category  $\underline{\mathcal{C}}'$ .

Now let us combine Proposition 2.3.5 with Theorem 3.3.2(II).

**Proposition 3.3.4.** Adopt the notation and assumptions of Theorem 3.3.2(II); assume in addition that  $\underline{\mathcal{C}}^0$  is  $R$ -linear and  $\underline{\mathcal{A}}$  is a weak Serre subcategory of  $R - \operatorname{Mod}$ .

1. Then  $\underline{\mathcal{C}}_{\underline{\mathcal{A}}}$  is triangulated and  $t$  restricts to it.
2. The heart of this restriction is naturally equivalent to the category of  $R$ -linear functors  $\underline{H}w^{0op} \rightarrow \underline{\mathcal{A}}$ .
3. Let  $\beta$  be an infinite cardinal; assume that  $R$  is Noetherian and the  $R$ -module  $\underline{\mathcal{C}}(M, N)$  has less than  $\beta$  generators for any  $M, N \in \operatorname{Obj} \underline{\mathcal{C}}^0$ . Then  $\underline{\mathcal{C}}^0 \subset \underline{\mathcal{C}}_{\beta}$  (cf. Proposition 2.3.5(3)).

*Proof.* 1. This is particular case of Proposition 2.3.5(1).

2. Immediate from Theorem 3.3.2(II.3) combined with Proposition 2.2.2(9).

3. Obvious.  $\square$

Now we recall a "geometric" setting where Proposition 3.3.4 can be applied.

**Proposition 3.3.5.** Assume that  $R$  is Noetherian and a scheme  $X$  is projective over  $\operatorname{Spec} R$  (that is,  $X$  is a closed subscheme of the projectivization  $Y$  of a vector bundle  $\mathcal{E}$  over  $\operatorname{Spec} R$ ); take  $\underline{\mathcal{C}} = D(\operatorname{Qcoh}(X))$  (the derived categories of quasicoherent sheaves on  $X$ ) and  $\underline{\mathcal{C}}^0 = \underline{\mathcal{C}}^{(\aleph_0)}$ .

1. Then  $\underline{\mathcal{C}}$  is compactly generated,  $\underline{\mathcal{C}}^0 = D_{\operatorname{perf}}(X) \subset \underline{\mathcal{C}}$  (the subcategory of perfect complexes on  $X$ ), and  $\underline{\mathcal{C}}_{\aleph_0} = D_{\operatorname{coh}}(\operatorname{Qcoh}(X))$ ; here we use the notation of Definition 2.3.4, whereas a complex  $N \in \operatorname{Obj} D(\operatorname{Qcoh}(X))$  belongs to  $D_{\operatorname{coh}}(\operatorname{Qcoh}(X))$  whenever all of its cohomology sheaves  $H^i(N)$  are coherent.
2. Take  $T$  to be a subset of  $S = \operatorname{Spec} R$  stable under specialization, that is,  $T$  is a union of closed subsets of  $S$  (see [stacks, Tags 00L1]). Then the category  $\underline{\mathcal{A}}^T$  of  $R$ -modules supported on  $T$  is a weak Serre subcategory of  $R - \operatorname{Mod}$ , and the corresponding category  $\underline{\mathcal{C}}_{\underline{\mathcal{A}}^T}$  (see Definition 2.3.4(2)) consists of all those objects of  $\underline{\mathcal{C}}$  the sections of whose cohomology sheaves (note that those are  $R$ -modules) are supported on  $T$ .

Consequently, the category  $\underline{\mathcal{C}}_{\underline{\mathcal{A}}^T \cap R - \operatorname{mod}}$  consists of those objects of  $\underline{\mathcal{C}}$  whose cohomology sheaves are coherent and whose sections are supported on  $T$ .

*Proof.* The statement that  $\underline{\mathcal{C}}$  is compactly generated by  $D_{\operatorname{perf}}(X)$  is well-known; see Theorem 4.2.2(1) of [Bon24]. The calculation of  $\underline{\mathcal{C}}_{\aleph_0}$  is given by part 3 of loc. cit., and the calculation of  $\underline{\mathcal{C}}_{\aleph_0}$  is given by Proposition 4.2.9(2) of ibid. It remains to apply Proposition 2.3.5(3) to conclude the proof.  $\square$

### 3.4 $t$ -structures anti-orthogonal to cosmashing weight structures

Once again, we refer to Definitions 3.1.1 and 3.2.1.

**Theorem 3.4.1.** Assume that  $w$  is a cosmashing weight structure on  $\underline{\mathcal{C}}$ , and  $\underline{\mathcal{C}}$  satisfies the dual Brown representability property; let  $\alpha$  be an infinite cardinal.

1. Then there exists a smashing  $t$ -structure  $t^l$  right adjacent to  $w$  and  $\underline{H}t^l$  is equivalent to the category of those additive functors  $\underline{H}w \rightarrow \mathbf{Ab}$  that respect products.

2. The category  $\underline{\mathcal{C}}^{(\alpha)} \subset \underline{\mathcal{C}}$  is triangulated.

3. Assume that  $w$  is also smashing. Then for any infinite cardinal  $\alpha$  the  $t$ -structure  $t^l$  given by assertion 1 restricts to the subcategory  $\underline{\mathcal{C}}^{(\alpha)}$  of  $\underline{\mathcal{C}}$ . Moreover, this restricted  $t$ -structure  $t^{(\alpha)}$  is the only  $t$ -structure on  $\underline{\mathcal{C}}^{(\alpha)}$  that is anti-orthogonal to  $w$ .

*Proof.* 1. This is just the categorical dual to Theorem 3.2.3.

2. This is the easy Lemma 4.1.4 of [Nee01].<sup>4</sup>

3. The uniqueness of a  $t$ -structure on  $\underline{\mathcal{C}}^{(\alpha)}$  that is anti-orthogonal to  $w$  is given by the dual to Corollary 2.4.4. Next, for any object  $M$  of  $\underline{\mathcal{C}}^{(\alpha)}$  the functor  $H^M$  is  $\alpha$ -small by the definition of  $\underline{\mathcal{C}}^{(\alpha)}$ . Now, for  $M' = t_{\geq 0}^l M$  Proposition 2.1.2(6) says that  $H^{M'} \cong \tau_{\geq 0} H^M$ . Hence the functor  $H^{M'}$  is  $\alpha$ -small as well according to Proposition 3.1.2(6), and we obtain that  $M'$  is an object of  $\underline{\mathcal{C}}^{(\alpha)}$ . Thus  $t^l$  restricts to  $\underline{\mathcal{C}}^{(\alpha)}$  indeed; see Lemma 1.1.7(1).

*Remark 3.4.2.* Dualizing Theorem 3.4.1(3) one obtains that the  $t$ -structure  $t^r$  provided by Theorem 3.2.3 restricts to the levels of a certain cosmallness filtration on  $\underline{\mathcal{C}}$ . Yet it appears that all the levels of this filtration are zero in "reasonable cases".

Let us now verify that Theorem 3.1 of [KeN13] (that essentially generalizes Theorem 3.2 of [Pau08]) gives an example for the setting of Theorem 3.2.3(II.2), and study the corresponding structures in detail.

**Corollary 3.4.3.** Let  $\underline{\mathcal{A}}$  be an additive subcategory of the subcategory  $\underline{\mathcal{C}}^{(\aleph_0)}$  of  $\underline{\mathcal{C}}$  that compactly generates  $\underline{\mathcal{C}}$ , and assume that the category  $\mathbf{AddFun}(\underline{\mathcal{A}}, \mathbf{Ab})$  is semi-simple and  $\mathbf{Obj} \underline{\mathcal{A}} \perp_{\underline{\mathcal{C}}} \cup_{i < 0} \mathbf{Obj} \underline{\mathcal{A}}[i]$ .

1. Then there exist a smashing and cosmashing weight structure  $w$  and a  $t$ -structure  $t$  on  $\underline{\mathcal{C}}$  that are generated by  $\mathbf{Obj} \underline{\mathcal{A}}$ , and they are right adjacent.

2.  $t$  restricts to the subcategory  $\underline{\mathcal{C}}^{(\alpha)}$  (cf. Theorem 3.4.1(2)) for any infinite cardinal  $\alpha$ . Moreover (for  $\alpha = \aleph_0$ ) the corresponding class  $\underline{\mathcal{C}}_{t^{(\aleph_0)} \leq 0}^{(\aleph_0)}$  (resp.  $\underline{\mathcal{C}}_{t^{(\aleph_0)} \geq 0}^{(\aleph_0)}$ ) is the envelope of  $\cup_{i \leq 0} \mathbf{Obj} \underline{\mathcal{A}}[i]$  (resp. of  $\cup_{i \geq 0} \mathbf{Obj} \underline{\mathcal{A}}[i]$ ) in  $\underline{\mathcal{C}}$  (see §1.1).

*Proof.* 1. By Theorem 3.1 of [KeN13] there exists a weight structure  $w$  on  $\underline{\mathcal{C}}$  such that  $\underline{\mathcal{C}}_{w \geq 0} = (\cup_{i > 0} \mathcal{P}[-i])^\perp$  (i.e.  $w$  is generated by  $\mathcal{P} = \mathbf{Obj} \underline{\mathcal{A}}$ ) and  $\underline{\mathcal{C}}_{w \leq 0} = (\cup_{i > 0} \mathcal{P}[i])^\perp$ . Since the category  $\underline{\mathcal{C}}$  is compactly generated by  $\underline{\mathcal{A}}$ , it satisfies the dual Brown representability property (by the aforementioned Theorem 8.6.1 and Remark 6.4.5 of [Nee01]). Next,  $w$  is obviously smashing and cosmashing. Applying Theorem 3.2.3(II.1), we obtain the existence of a smashing  $t$ -structure  $t$  that is right adjacent to  $w$ . Since  $\underline{\mathcal{C}}_{w \leq 0} = \underline{\mathcal{C}}_{t \leq 0}$ , we obtain that  $t$  is generated by  $\mathcal{P}$  (as a  $t$ -structure) as well.

2.  $t$  restricts to the subcategory  $\underline{\mathcal{C}}^{(\alpha)}$  for any infinite cardinal  $\alpha$  according to Theorem 3.4.1(3). Thus it remains to prove that the classes  $\underline{\mathcal{C}}_{t \leq 0} \cap \mathbf{Obj} \underline{\mathcal{C}}^{(\aleph_0)} =$

<sup>4</sup>Which actually does not require any weight structure, where  $\underline{\mathcal{C}}$  is only required to be smashing.

$\underline{\mathcal{C}}_{w \leq 0} \cap \text{Obj } \underline{\mathcal{C}}^{(\mathbb{N}_0)}$  and  $\underline{\mathcal{C}}_{t \geq 0} \cap \text{Obj } \underline{\mathcal{C}}^{(\mathbb{N}_0)}$  are the envelopes in question. The latter statement easily follow from Theorem 3.2.1(2) of [Bon22a] (see Remark 3.2.2(2) of *ibid.*; note also that Theorem 4.5(1,2) of [PoS16] gives this statement under the assumption that  $\underline{\mathcal{C}}$  is a "stable derivator" triangulated category).  $\square$

$\square$

*Remark 3.4.4.* 1. Thus we obtain a serious generalization of the existence of a (certain)  $t$ -structure on  $\underline{\mathcal{C}}^{(\mathbb{N}_0)}$  statement provided by Theorem 7.1 of [KeN13]. In particular, we do not need any differential graded algebras. Note however that our arguments do not give the description of  $t^{\mathbb{N}_0}$  in terms of "generators" that would be similar to that in *loc. cit.*

2. Now we try to study the question which  $t$ -structures on  $\underline{\mathcal{C}}^{(\mathbb{N}_0)}$  extend to examples for our corollary.

So, assume that  $\underline{\mathcal{C}}$  is an arbitrary triangulated category and  $t'$  is a  $t$ -structure on  $\underline{\mathcal{C}}^{(\mathbb{N}_0)}$ , and take  $\underline{A}' = \underline{H}t'$ . Then we have  $\text{Obj } \underline{A}' \perp (\cup_{i>0} \text{Obj } \underline{A}'[-i])$  by the orthogonality axiom for  $t'$ .

Thus any essentially small abelian subcategory  $\underline{A}$  of  $\underline{H}t'$  whose objects are semi-simple satisfies all the assumptions of our corollary except the one that  $\underline{A}$  compactly generates  $\underline{\mathcal{C}}$ . Hence we can apply our corollary to the localizing subcategory  $\underline{\mathcal{C}}'$  of  $\underline{\mathcal{C}}$  generated by  $\text{Obj } \underline{A}$ .

3. Now assume in addition that  $\underline{\mathcal{C}}$  is compactly generated,  $t'$  is bounded, and  $\underline{H}t'$  is a length category (cf. Theorem 7.1 of [KeN13]). Then the category  $\underline{\mathcal{C}}^{(\mathbb{N}_0)}$  is essentially small according to Lemma 4.4.5 of [Nee01]; hence  $\underline{H}t'$  is essentially small as well, and we can take  $\underline{A}$  to be its subcategory of semi-simple objects.

Since  $t'$  is bounded and  $\underline{H}t'$  is a length category, the category  $\underline{\mathcal{C}}^{(\mathbb{N}_0)}$  is densely generated by  $\text{Obj } \underline{A}$ ; hence  $\underline{A}$  is easily seen to generate  $\underline{\mathcal{C}}$  (in the sense of Definition 3.2.1(3)). Thus one can apply Corollary 3.4.3 to this setting. Moreover, it is easily seen that our assumptions on  $t'$  (combined with part 2 of our corollary) imply that the corresponding  $t$ -structure  $t^{\mathbb{N}_0}$  coincides with  $t'$ .

4. It would be interesting to find which assumptions on a general  $t$ -structure  $t'$  on  $\underline{\mathcal{C}}^{(\mathbb{N}_0)}$  ensure that  $t'$  extends to a  $t$ -structure on  $\underline{\mathcal{C}}$  and there exists a weight structure  $w$  that is right adjacent to  $t$ .

## 4 On functors and $t$ -structures related to (essentially) bounded weight structures

The main goal of this section is to describe some more orthogonal  $t$ -structures in the case where the corresponding weight structures satisfy certain boundedness assumptions.

To formulate our statements in a more general case we recall (in §4.1) the notions of essentially bounded (above, below, or both) objects and weight structures; we motivate this "almost generalization" in Remarks 4.1.5(2,3), 4.2.3(2),



and 4.2.7 below. We also study weight structures of these types as well as virtual  $t$ -truncations with respect to them and their relation with certain "essential weight-boundedness" for functors.

In §4.2 we apply the aforementioned results to the construction of certain adjacent and orthogonal  $t$ -structures. We apply our general statements to various derived categories of (quasi)coherent sheaves. In particular, we are able to generalize some central results of [Bon24].

#### 4.1 On essentially bounded weight structures and the relation to cohomological functors

Now we will prove some bounded versions of the results of §2.2. To formulate them in the most general case we need some (somewhat clumsy) definitions and statements closely related to earlier papers of the author. Yet the reader can avoid much of this theory if she applies it to  $w$ -bounded below (resp. above) objects and weight structures only; note that those are clearly *essentially bounded* below (resp. above).

**Definition 4.1.1.** Assume that  $w$  is weight structure on  $\underline{\mathcal{C}}$ ;  $M \in \text{Obj } \underline{\mathcal{C}}$ .

1. We say that  $M$  is left (resp., right)  $w$ -degenerate (or *weight-degenerate* if the choice of  $w$  is clear) if  $M$  belongs to  $\cap_{i \in \mathbb{Z}} \underline{\mathcal{C}}_{w \geq i}$  (resp. to  $\cap_{i \in \mathbb{Z}} \underline{\mathcal{C}}_{w \leq i}$ ).
2. We say that  $M$  is *essentially  $w$ -positive* (resp. *essentially  $w$ -negative*) if for  $\tilde{M} = M \oplus M[1]$  (resp.  $\tilde{M} = M \oplus M[-1]$ ) there exists a distinguished triangle

$$RD(\tilde{M}) \rightarrow \tilde{M} \rightarrow Y \rightarrow RD(\tilde{M})[1] \quad (\text{resp. } X \rightarrow \tilde{M} \rightarrow LD(\tilde{M}) \rightarrow X[1]) \quad (4.1.1)$$

such that  $Y \in \underline{\mathcal{C}}_{w \geq 0}$  (resp.  $X \in \underline{\mathcal{C}}_{w \leq 0}$ ) and  $RD(\tilde{M})$  (resp.  $LD(\tilde{M})$ ) is right (resp. left) degenerate.

3. We say that  $M$  is *essentially  $w$ -bounded below* (resp. *above*) if  $M[-n]$  is essentially  $w$ -positive (resp. essentially  $w$ -negative) for some  $n \in \mathbb{Z}$ . We will call the shift of the triangle (4.1.1) corresponding to  $M[-n]$  by  $[n]$  (cf. (4.1.2) below) a  *$w$ -degenerate decomposition* of  $\tilde{M}$ .

4.  $w$  is said to be *essentially bounded below* (resp. *above*) if all objects of  $\underline{\mathcal{C}}$  are essentially  $w$ -bounded below (resp. above).

We say that  $w$  is *essentially bounded* if it is essentially bounded both above and below.

5. A cohomological functor  $H$  from  $\underline{\mathcal{C}}$  will be said to be *locally bounded below* (resp. *above*)<sup>5</sup> if for any  $M \in \text{Obj } \underline{\mathcal{C}}$  we have  $H^i(M) = 0$  for all  $i \ll 0$  (resp.  $i \gg 0$ ).

$H$  is said to be *locally bounded* if it is locally bounded both above and below.

Some nice properties of essential boundedness are closely related to earlier papers.

**Proposition 4.1.2.** Assume that  $w$  is weight structure on  $\underline{\mathcal{C}}$ ;  $M \in \text{Obj } \underline{\mathcal{C}}$ .

1. Assume that  $M$  is essentially  $w$ -bounded below (resp. above). Then the  $w$ -degenerate decomposition of  $\tilde{M}$  is canonical. Being more precise, the distinguished triangle

$$RD(\tilde{M}) \rightarrow \tilde{M} \rightarrow Y \rightarrow RD(\tilde{M})[1] \quad (\text{resp. } X \rightarrow \tilde{M} \rightarrow LD(\tilde{M}) \rightarrow X[1]), \quad (4.1.2)$$

---

<sup>5</sup>This terminology was inspired by [Nee18, Remark 0.2].

where  $RD(\tilde{M})$  is right (resp.  $LD(\tilde{M})$  is left)  $w$ -degenerate and  $Y$  (resp.  $X$ ) is  $w$ -bounded above (resp. below) is canonically determined by  $M$  (and does not depend on the choice of the corresponding  $n$ ).

2. Take those couples  $(m, n)$  such that  $m \leq n \in \mathbb{Z}$  and  $n < 0$  (resp.  $m > 0$ ). The following conditions are equivalent.

1.  $M$  is essentially  $w$ -positive (resp. negative).
2.  $M$  is a retract of some  $M' \in \text{Obj } \underline{\mathcal{C}}$  such that there exists a distinguished triangle

$$RD(M') \rightarrow M' \rightarrow Y \rightarrow RD(M')[1] \text{ (resp. } X \rightarrow M' \rightarrow LD(M') \rightarrow X[1]),$$

where  $Y \in \underline{\mathcal{C}}_{w \geq 0}$  (resp.  $X \in \underline{\mathcal{C}}_{w \leq 0}$ ) and  $RD(M')$  (resp.  $LD(M')$ ) is right (resp. left) degenerate.

3.  $H(M) = 0$  if  $H$  is (a cohomological functor) of weight range  $[m, n]$  and  $(m, n)$  satisfies the conditions above.
4.  $H(M) = \{0\}$  for  $H = \tau_{\leq n}(H_I)$  whenever  $I \in \underline{\mathcal{C}}_{w \geq m}$  and any  $(m, n)$  as above.
5.  $H(M) = \{0\}$ , where  $H = \tau_{\leq n}(H_{I_0})$  and  $I_0$  is a fixed choice of  $w_{\geq m}M$  and any  $(m, n)$  as above.

*Proof.* 1. Assume first that  $M$  is essentially  $w$ -bounded below and there exist distinguished triangles

$$T_i = (RD_i(\tilde{M}) \rightarrow \tilde{M} \rightarrow Y_i \rightarrow RD_i(\tilde{M})[1])$$

of the type (4.1.2) for  $i = 1, 2$ . Assume that  $Y_i \in \underline{\mathcal{C}}_{w \geq n_i}$  for  $i = 1, 2$  and some  $n_i \in \mathbb{Z}$ . We should prove that these triangles are canonically isomorphic.

We can assume that  $n_1 \geq n_2$ . Then both  $T_1$  and  $T_2$  are  $m$ -weight decompositions of  $\tilde{M}$  for any  $n < n_2$ . Applying the uniqueness statement in Proposition 1.2.4(4) one can easily obtain  $T_1 \cong T_2$ . In particular, one can note that both  $T_1$  and  $T_2$  give weight decompositions of  $\tilde{M}$  that *avoid weight*  $n_2 - 1$ ; see Theorem 2.2.1(9) of [Bon22b]. Now we applying loc. cit. to obtain  $T_1 \cong T_2$ .

The case of a  $w$ -bounded above  $M$  is just the categorical dual of assertion 1; see Proposition 1.2.4(1).

2. One may say that this is a "one-sided unbounded version of" [BoV23, Theorem 2.5(I)].

Firstly, conditions 1 and 2 are equivalent by Corollary 3.1.4 (resp. Theorem 3.1.3) of [Bon22b]. Moreover, these statements also yield that condition 2 follows from 3; recall here that pure functors (cf. condition 7 of Corollary 3.1.4 and condition 8 of Theorem 3.1.3 in ibid.) are the functors of weight range  $[0, 0]$ .<sup>6</sup> The converse implication is valid as well; see conditions 2 and Theorem 2.5(I) of [BoV23].

Lastly, loc. cit. says that conditions 3–5 become equivalent if one fixes any  $m \leq n \in \mathbb{Z}$ . Combining these equivalences for all  $m \leq n \in \mathbb{Z}$  such that  $n < 0$  (resp.  $m > 0$ ) we obtain that our versions of conditions 3–5 are equivalent as well.  $\square$

---

<sup>6</sup>Actually, in loc. cit. mentions homological pure functors  $\underline{\mathcal{C}} \rightarrow \underline{\mathcal{A}}$ ; yet one can just reverse the arrows in the target category.

Let us apply this proposition to pure functors.

**Corollary 4.1.3.** Assume that  $w^0$  is an essentially bounded below (resp. above, both) weight structure on  $\underline{C}^0$ , and  $H$  is a pure functor from  $\underline{C}^0$ .

Then  $H$  is locally bounded below (resp. above, both).

*Proof.* This statement is an immediate consequence of our definitions along with Proposition 4.1.2(II); see condition 3 in it.  $\square$

**Proposition 4.1.4.** Assume that  $w$  is weight structure on  $\underline{C}$ ,  $M$  is essentially  $w$ -bounded below (resp. above), and  $H$  is a cohomological functor from  $\underline{C}$ .

1. Assume that  $M[-n]$  is essentially  $w$ -positive (resp.  $w$ -negative, for some  $n \in \mathbb{Z}$ ), and  $m < n - 1$  (resp.  $m > n + 1$ ) and  $H$  is a cohomological functor from  $\underline{C}$ . Then  $\tau_{\leq m}(H)(\tilde{M}) \cong H(RD(\tilde{M}))$  (resp.  $\tau_{\geq m}(H)(\tilde{M}) \cong H(LD(\tilde{M}))$ ). Consequently,  $\tau_{\leq m}(H)(\tilde{M})$  (resp.  $\tau_{\geq m}(H)(\tilde{M})$ ) is a retract of  $H(RD(\tilde{M}))$  (resp. of  $H(LD(\tilde{M}))$ ).

2. Assume that  $w$  is essentially bounded below (resp. above) and for any right (resp. left) weight-degenerate  $D \in \text{Obj } \underline{C}$  we have  $H^i(D) = 0$  for  $i \ll 0$  (resp. for  $i \gg 0$ ). Then for any  $m \in \mathbb{Z}$  the functor  $\tau_{\leq m}(H)$  (resp.  $\tau_{\geq m}(H)$ ) is locally bounded below (resp. above).

3. Assume that  $w$  is essentially bounded below (resp. above) and  $H$  is locally bounded below (resp. above). Then all  $\tau_{\leq m}(H)$  and  $\tau_{\geq m}(H)$  are locally bounded below (resp. above) as well.

4. Assume that  $w$  is essentially bounded and  $H$  is locally bounded. Then all  $\tau_{\leq m}(H)$  and  $\tau_{\geq m}(H)$  are locally bounded as well.

*Proof.* 1. Since  $M$  is a retract of  $\tilde{M}$ , it suffices to prove the first part of the assertion.

Now, assume that  $M[-n]$  is essentially  $w$ -bounded below. Similarly to the argument above, note that we can take  $w_{\leq i}(\tilde{M}) = RD(\tilde{M})$  for  $i < n$ . Recalling the definition of  $\tau_{\leq m}(H)$  (see Definition 2.1.1(1)), we obtain that  $\tau_{\leq m}(H)(\tilde{M}) \cong \text{Im}(\text{id}_{H(RD(\tilde{M}))}) = H(RD(\tilde{M}))$ .

The case of an essentially  $w$ -bounded above  $M$  can be treated similarly. Moreover, it can be easily deduced from the essentially bounded below case; see Propositions 1.2.4(1) and 2.1.2(5).

2. Once again, we give the proof in the essentially bounded below case; the essentially bounded above case is very much similar and can be deduced from it.

According to the previous assertion, it suffices to verify that  $(\tau_{\leq m}(H))^i(RD(\tilde{M})) = 0$  for  $i \ll 0$  and any  $M \in \text{Obj } \underline{C}$ . Thus it remains to recall that  $RD(\tilde{M})$  is right weight-degenerate.

3. This assertion immediately follows from the previous one if we recall the exact sequence (2.1.1).

4. This assertion can be immediately obtained by means of combining the bounded above and below cases of the previous one.  $\square$

*Remark 4.1.5.* If  $w$  is *weight-Karoubian*, that is, if  $\underline{H}w$  is Karoubian then one modify the definition of essentially  $w$ -bounded above and below objects by setting  $\tilde{M} = M$  in the triangles (4.1.1); see Theorems 2.3.4 and 3.1.3 and Corollary 3.1.4 of [Bon22b]. Respectively, one can compute  $\tau_{\leq m}(H)(M)$  for

any  $M \in \text{Obj } \underline{\mathcal{C}}$  using the corresponding canonical decomposition similarly to Proposition 4.1.4(1).

2. Obviously, any semi-orthogonal decomposition couple is an essentially bounded weight structure; see condition 4 of Proposition 1.2.6.

Furthermore, one may say that general essentially bounded above and below weight structures are "mixes" of bounded above and below ones and semi-orthogonal decompositions. Indeed, if  $w$  is weight-Karoubian and essentially bounded above or below then the statements mentioned in part 1 of this remark yield the existence of a certain semi-orthogonal decomposition.

3. Respectively, the main results of this section can be used to generalize some statement on semi-orthogonal decompositions from [Bon24]. We will say more on this in Remark 4.2.7(2) below.

## 4.2 More on the construction of $t$ -structures

Now we pass to the existence of  $t$ -structures orthogonal to essentially bounded weight structures. We start with a nice "bounded  $R$ -linear analogue" of Theorem 3.2.3. Once again, we assume that  $R$  is a commutative Noetherian ring).

**Definition 4.2.1.** Assume that  $\underline{\mathcal{C}}$  is an  $R$ -linear category.

We will say that  $\underline{\mathcal{C}}$  is  $R$ -saturated if the representable functors from  $\underline{\mathcal{C}}$  into  $R - \text{Mod}$  are exactly the ones that are locally bounded and take values in the subcategory  $R - \text{mod}$  of finitely generated  $R$ -modules.

**Corollary 4.2.2.** 1. Assume that  $\underline{\mathcal{C}}$  is an  $R$ -saturated category, and  $w$  is an essentially bounded weight structure on it.

Then there exists a  $t$  structure  $t$  (left) adjacent to  $w$  (on  $\underline{\mathcal{C}}$ ). Its heart is equivalent to the category of  $R$ -linear functors from  $\underline{Hw}^{op}$  into the category of finitely generated  $R$ -modules.

2. Take  $T$  to be a union of closed subsets of  $S = \text{Spec } R$ , and  $\underline{A}^T$  to be the category of  $R$ -modules supported on  $T$  (cf. Proposition 3.3.5(2)).

Then the category  $\underline{\mathcal{C}}_{\underline{A}^T}$  corresponding to  $\underline{\mathcal{C}}^0 = \underline{\mathcal{C}}$  (see Definition 2.3.4(2)) is triangulated and  $t$  restricts to it. Moreover, the heart of this restriction is equivalent to the category of  $R$ -linear functors from  $\underline{Hw}^{op}$  into the category of finitely generated  $R$ -modules supported on  $T$ .

3. Assume that  $X$  is regular and proper over  $\text{Spec } R$ . Then the category  $\underline{\mathcal{C}} = D^{perf}(X)$  is  $R$ -saturated. Moreover,  $\underline{\mathcal{C}}$  is equivalent to  $\underline{\mathcal{C}}^{op}$ .

Consequently, if  $w$  is an essentially bounded weight structure on  $\underline{\mathcal{C}}$  then there exists a  $t$ -structure on  $\underline{\mathcal{C}}$  that is left adjacent to  $w$  and also a  $t$ -structure that is right adjacent to it.

*Proof.* 1. According to Theorem 2.3.3(I), it suffices to verify that for any representable functor  $H$  the functor  $\tau_{\leq 0}H$  is representable as well (in the  $R$ -linear sense; see Lemma 2.1.4(2,3)). Now,  $\tau_{\leq 0}H$  is locally bounded according to Proposition 4.1.4(4), and takes its values in (the abelian subcategory of) finitely generated  $R$ -modules according to Proposition 2.1.3(II.3).

Next,  $\underline{Ht}$  embeds into the category of  $R$ -linear functors  $\underline{Hw}^{op} \rightarrow R - \text{Mod}$  according to Theorem 2.3.3(II). Now, any functor  $\underline{Hw}^{op} \rightarrow R - \text{mod}$  factors through a pure cohomological functor from  $\underline{\mathcal{C}}$  into  $R - \text{mod}$ ; see Proposition 2.2.2(9). Lastly, any functor of this sort is locally bounded by Corollary 4.1.3; hence it is representable (see Definition 4.2.1(1)).

2. Recall that  $\underline{A}^T$  is a weak Serre subcategory of  $R - \text{Mod}$ ; see Proposition 3.3.5(2). Hence it remains to apply Proposition 2.3.5(1).

3. The first part of the assertion is a particular case of [Nee18, Corollary 0.5]; note that in this case  $D^{perf}(X)$  is equivalent to the bounded derived category of  $X$ ; cf. Remark 1.4(3) of [Bon24].

Next, the self-duality of  $\underline{C}$  is a well-known consequence of Grothendieck duality; see [stacks, Tags 0AU3, 0DWG, 0BFQ]. Thus it remains to apply assertion 1 to conclude the proof (see Proposition 1.2.4(1); note that  $w^{op}$  is clearly essentially bounded as well).  $\square$

*Remark 4.2.3.* 1. In all the examples of  $R$ -saturated categories known to the author one can achieve the same result by means of applying (the somewhat more clumsy) Corollary 4.2.5 instead; see Theorem 4.2.2(1) and Proposition 4.2.6 of [Bon24].

2. Similarly to Remark 3.2.4(3) one can apply Corollary 4.2.2(1,2) to semi-orthogonal decompositions. One obtains that if  $\underline{C}$  is saturated,  $\underline{C}' = \underline{C}_{\underline{A}^T}$  (note that  $\underline{C}' = \underline{C}$  if  $T = \text{Spec } R$ ), and  $(\mathcal{A}, \mathcal{B})$  is a semi-orthogonal decomposition of  $\underline{C}$ , then the couple  $((\text{Obj } \mathcal{A})^\perp \cap \text{Obj } \underline{C}', (\text{Obj } \mathcal{B})^\perp \cap \text{Obj } \underline{C}' = \mathcal{A} \cap \text{Obj } \underline{C}')$  is a semi-orthogonal decomposition of  $\underline{C}'$ . Consequently, if  $\mathcal{A}$  is a left admissible subcategory of  $\underline{C}$  then it is also admissible in it. This statement generalizes Proposition 2.6 of [BoK89], where the case  $R = \mathbb{C}$  was considered.

Now, it is well-known that the categories of the type  $D_{perf}(X)$  often admit non-trivial semi-orthogonal decompositions (in particular, in the case where  $R$  is a field). Hence there can exist unbounded weight structures on them; see Remark 4.1.5(2). On the other hand the author suspects that any weight structure on  $D_{perf}(X)$  is essentially bounded; hence one can apply Corollary 4.2.2(1,2) to it.

This observation motivated the author to consider essentially bounded weight structures in this section; cf. Remark 4.2.7 below.

3. Clearly, if  $\underline{C}$  is  $R$ -saturated and the category  $\underline{A}$  is a weak Serre subcategory of  $R - \text{Mod}$  then the corresponding category  $\underline{C}_{\underline{A}}$  depends on  $\underline{A} \cap R - \text{mod}$  only. Now,  $\underline{A} \cap R - \text{mod}$  is a weak Serre subcategory of  $R - \text{mod}$  (cf. Remark 1.1), and any weak Serre subcategory of  $R - \text{mod}$  of  $R - \text{mod}$  consists of finitely generated  $R$ -modules supported at some  $T$  as in Corollary 4.2.2(2); see Theorem A of [Tak08].<sup>7</sup> Consequently, it does not make sense to consider any  $\underline{A}$  distinct from  $\underline{A}^T$  in Corollary 4.2.2(2).

Now we pass to orthogonal structures (in distinct categories).

**Theorem 4.2.4.** Assume that  $\underline{C}^0, \underline{C}' \subset \underline{C}$ ,  $w^0$  is an essentially bounded below (resp. above, both) weight structure on  $\underline{C}^0$ ,  $w$  is its coproductive extension to  $\underline{C}$ , and there exists a  $t$ -structure  $t$  on  $\underline{C}'$  that is orthogonal to  $w^0$ . Set  $\underline{C}'^0_+$  (resp.  $\underline{C}'^0_-$ ,  $\underline{C}'^0_b$ ) to be the subcategory of  $\underline{C}'$  that consists of those  $M \in \text{Obj } \underline{C}'$  such that the functor  $H_M^{C^0}$  is locally finite below (resp. above, both).

Then the category  $\underline{C}'^0_+$  (resp.  $\underline{C}'^0_-$ ,  $\underline{C}'^0_b$ ) is triangulated and  $t$  restricts to it. Moreover, this restriction of  $t$  is strictly orthogonal to  $w^0$ , and its heart coincides with  $Ht$ .

---

<sup>7</sup>Note that extension closed abelian subcategories are called *coherent* ones in (Definition 2.3(1) of) [Tak08].

*Proof.* The subcategories  $\underline{C}'_+, \underline{C}'_-, \underline{C}'_b$  of  $\underline{C}'$  are easily seen to be triangulated regardless of any weight structures. Next, the strict orthogonality assertion immediately follows from Corollary 2.4.4.

It remains to prove the existence of the corresponding restricted  $t$ -structures and calculate the heart. Similarly to the proof of Proposition 2.3.5(1), Lemma 1.1.7(1) reduces the existence of this restriction to the following statement: for  $M$  that belongs  $\underline{C}'_+$  (resp.  $\underline{C}'_-, \underline{C}'_b$ ) the functor  $\tau_{\geq 0} H_M^{\underline{C}} = \tau_{\geq 0} H_M$  is  $\underline{C}$ -representable by an object of the corresponding subcategory. Now,  $\tau_{\geq 0} H_M \cong H_L$  for some object  $L$  of  $\underline{C}'$  (see Proposition 2.1.2(4)). To prove that this  $L$  actually belongs to  $\underline{C}'_+$  (resp.  $\underline{C}'_-, \underline{C}'_b$ ) we should check that the restricted functor  $\tau_{\geq 0} H_M^{\underline{C}^0}$  (see Proposition 2.1.3(II.2)) is locally finite below (resp. above, both). The latter statement is given by Proposition 4.1.4(3–4).

Lastly, Corollary 4.1.3 implies that any object of  $\underline{H}t$  belongs to  $\underline{C}'_+$  (resp.  $\underline{C}'_-, \underline{C}'_b$ ) indeed.  $\square$

**Corollary 4.2.5.** Assume that  $\underline{C}$  is a smashing  $R$ -linear triangulated category, an essentially small subcategory  $\underline{C}^0$  of  $\underline{C}^{(\aleph_0)}$  generates it, and  $w^0$  is an essentially bounded below (resp. above, both) weight structure on  $\underline{C}^0$ . Choose a weak Serre subcategory  $\underline{A} \subset R\text{-Mod}$ .

Then the  $t$ -structure  $t$  on  $\underline{C}_{\underline{A}}$  provided by Proposition 2.3.5(1) restricts to the intersection subcategory  $\underline{C}_{\underline{A}} \cap \underline{C}'_+$  (resp.  $\underline{C}_{\underline{A}} \cap \underline{C}'_-, \underline{C}_{\underline{A}} \cap \underline{C}'_b$ ), and the heart of this restriction is naturally equivalent to the category of  $R$ -linear functors from  $\underline{H}w^{0op}$  into  $\underline{A}$ .

*Proof.* According to Lemma 1.1.7(2), it suffices to combine Proposition 2.3.5(1) with Theorem 4.2.4.  $\square$

Now we relate the example provided by Proposition 3.3.5 to the conditions of Corollary 4.2.5.

**Proposition 4.2.6.** Assume that a scheme  $X$  is projective (see Proposition 3.3.5) over  $\text{Spec } R$  (where  $R$  is Noetherian); take  $\underline{C} = D(\text{Qcoh}(X))$  (the derived categories of quasicoherent sheaves on  $X$ ), and  $\underline{C}^0 = \underline{C}^{(\aleph_0)} = D_{\text{perf}}(X)$  (see Proposition 3.3.5 once again).

1. Then the intersection subcategories  $\underline{C}_{\aleph_0} \cap \underline{C}'_+, \underline{C}_{\aleph_0} \cap \underline{C}'_-,$  and  $\underline{C}_{\aleph_0} \cap \underline{C}'_b$  equal  $D_{\text{coh}}^+(\text{Qcoh}(X)), D_{\text{coh}}^-(\text{Qcoh}(X)),$  and  $D_{\text{coh}}^b(\text{Qcoh}(X))$ , respectively; here a complex  $N \in D_{\text{coh}}(\text{Qcoh}(X))$  belongs to  $D_{\text{coh}}^+(\text{Qcoh}(X))$  (resp.  $D_{\text{coh}}^-(\text{Qcoh}(X)), D_{\text{coh}}^b(\text{Qcoh}(X))$ ) whenever its (coherent) cohomology sheaves  $H^i(N)$  vanish for  $i \ll 0$  and (resp. for  $i \gg 0$ , in both cases).

2. Take  $T$  to be a union of closed subsets of  $S$ . Then to characterize the intersections of corresponding category  $\underline{C}_{\underline{A}^T \cap R\text{-mod}}$  (see Definition 2.3.4(2)) with  $\underline{C}'_+, \underline{C}'_-,$  and  $\underline{C}'_b$  one should add to the aforementioned conditions the assumption that the cohomology sheaves of  $N$  are supported on  $T$ .

*Proof.* 1. The calculation of these categories is given by Theorem 4.2.2(3) of [Bon24].

2. Immediate from assertion 1 combined with Propositions 3.3.5(2) and Proposition 2.3.5(2).  $\square$

*Remark 4.2.7.* 1. Thus one can apply Corollary 4.2.5 to all the intersection subcategories mentioned in Proposition 4.2.6.

2. Corollary 4.2.5 also implies that for any essentially bounded weight structure on  $D_{coh}^b(\mathrm{Qcoh}(X)) \cong D^b(\mathrm{coh}(X))$  (see [stacks, Tag 0FDA]) there exists an anti-orthogonal  $t$ -structure on  $D_{perf}(X)$ ; see Proposition 4.2.6 of [Bon24].

The author conjectures that there do not exist bounded weight structures on  $D^b(\mathrm{coh}(X))$  if  $X$  is singular. On the other hand, one may "start from" a bounded weight structure on a component of a semi-orthogonal decomposition of  $X$  (of arbitrary length; see Definition 2.3.1(II.2) of *ibid.*) to obtain an essentially bounded weight structure on the whole  $\underline{C}$ ; see Remark 4.1.5(2) and Proposition 2.3.2 of *ibid.*

3. Applying Remark 4.1.5(2) one can prove that Corollary 4.2.5 allows to generalize all the statements in Theorem 4.2.7 of *ibid.* except the bijection statement in part II.1 of *loc. cit.* from semi-orthogonal decompositions to essentially bounded (above, below, or both) weight structures. Probably, this statement and examples for it will be studied in a succeeding paper.

## 5 The existence of orthogonal weight structures

In this section we (try to) construct certain weight structures orthogonal to given  $t$ -structures.

For this purpose, in §5.1 we (recall and) prove some statements related to the hearts of weight structures.

In §5.2 we prove that the existence of an adjacent weight structure is closely related to the existence of enough projectives in the heart of  $t$ . This assumption ensures the existence of weight structure on certain ("rather large") subcategories of  $\underline{C}$  that are orthogonal to  $t$ . Under certain assumptions, we also obtain a weight structure adjacent to  $t$ .

In §5.3 we apply the results of the previous section to obtain a weight structure adjacent to a  $t$ -structure in the case where the category  $\underline{C}^{op}$  is  $R$ -saturated (whenever certain assumptions on  $t$  are fulfilled).

### 5.1 On hearts and $\pm$ -orthogonality of structures

**Definition 5.1.1.** We will say that a subcategory  $\underline{H} \subset \underline{C}$  is *connective* (in  $\underline{C}$ ) if  $\mathrm{Obj} \underline{H} \perp (\cup_{i>0} \mathrm{Obj}(\underline{H}[i]))$ .

We will say that  $\underline{H}$  is *silting* if it is connective and densely generates  $\underline{C}$  (cf. §3.1 of [KoY14]).

We recall rather well-known statements.

**Proposition 5.1.2.** 1. If  $w$  is a weight structure on  $\underline{C}$  and  $\underline{H} \subset \underline{H}w$  then  $\underline{H}$  is connective.

2. If  $\underline{H}$  is a silting subcategory of  $\underline{C}$  then the envelopes (see §1.1)  $\underline{C}_{v \leq 0}$  and  $\underline{C}_{v > 0}$  of the classes  $\cup_{i \leq 0} \mathrm{Obj} \underline{B}[i]$  and  $\cup_{i > 0} \mathrm{Obj} \underline{B}[i]$ , respectively, give a bounded weight structure  $v$  on  $\underline{C}$ , and  $\underline{H}v$  is the  $\underline{C}$ -retraction closure of the additive subcategory generated by  $\underline{H}$  (i.e., of the subcategory whose objects are  $\bigoplus H_i$  for finite sets of  $H_i \in \mathrm{Obj} \underline{H}$ ).

We will say that this  $v$  is *densely generated by  $\underline{H}$* .

*Proof.* Assertion 1 immediately follows from the orthogonality axiom in Definition 1.2.1, and assertion 2 is contained in Corollary 2.1.2 of [BoS18]; see also Theorem 5.5 of [HVVS13].  $\square$

Till the end of this section we assume that  $\underline{C}' \subset \underline{C}$  and  $t$  is a  $t$ -structure on  $\underline{C}$ .

Let us now "split" Definition 1.3.1(3).

**Definition 5.1.3.** Assume that  $w$  is a weight structure on  $\underline{C}'$ .

1. We will say that  $w$  and  $t$  are *-orthogonal* (resp. *+orthogonal*) if  $\underline{C}_{t \leq -1} = \underline{C}'_{w \geq 0}{}^{\perp_{\underline{C}}}$  (resp.  $\underline{C}_{t \geq 1} = \underline{C}'_{w \leq 0}{}^{\perp_{\underline{C}}}$ ).
2. We will write  $P_t$  for  ${}^{\perp_{\underline{C}}}(\underline{C}_{t \leq -1} \cup \underline{C}_{t \geq 1})$ ;  $P'_t = P_t \cap \text{Obj } \underline{C}'$ .
3. We will say that  $t$  is *left* (resp. *right*) *non-degenerate* if  $\cap_{i \in \mathbb{Z}} \underline{C}_{t \geq i} = \{0\}$  (resp.  $\cap_{i \in \mathbb{Z}} \underline{C}_{t \leq i} = \{0\}$ ).

Let us establish some properties of these definitions.

**Proposition 5.1.4.** Let  $t$  be a  $t$ -structure on  $\underline{C}$ ,  $P \in P_t$ .

1. The functor  $\underline{C}(P, -)$  restricts to an exact functor  $E^P : \underline{H}t \rightarrow \text{Ab}$ , and we have  $\underline{C}(P, -) \cong E^P \circ H_0^t$ .
- 2.

$$\underline{C}(P, -) \cong \underline{C}(P, H_0^t(-)) \cong \underline{H}t(H_0^t(P), H_0^t(-));$$

the first of these transformations comes from the transformations  $\text{id}_{\underline{C}} \rightarrow t_{\leq 0}$  and  $H_0^t \rightarrow t_{\leq 0}$  (see Proposition 1.1.5(1,3)). Moreover,  $H_0^t(P)$  is projective in  $\underline{H}t$ .

3. Assume that  $t$  is cosmashing and  $\underline{C}$  satisfies the dual Brown representability property. Then  $H_0^t$  gives an equivalence of (the full subcategory of  $\underline{C}$  given by)  $P_t$  with the subcategory of projective objects of  $\underline{H}t$ .

*Proof.* I. If  $0 \rightarrow A_1 \rightarrow A_2 \rightarrow A_3 \rightarrow 0$  is a short exact sequence in  $\underline{H}t$  then  $A_1 \rightarrow A_2 \rightarrow A_3 \rightarrow A_1[1]$  is a distinguished triangle according to Proposition 1.1.5(2). Applying the functor  $\underline{C}(P, -)$  to it and recalling the definition of  $P_t$  we obtain an exact sequence  $\{0\} = \underline{C}(P, A_3[-1]) \rightarrow E^P(A_1) \rightarrow E^P(A_2) \rightarrow E^P(A_3) \rightarrow E^P(A_1[1]) = \{0\}$ ; hence  $E^P$  is exact indeed. Moreover, the functors  $\Phi(P, -)$  and  $E^P \circ H_0^t$  are homological and annihilate both  $\underline{C}_{t \leq -1}$  and  $\underline{C}_{t \geq 1}$ ; hence they are isomorphic.

2. The first of these isomorphisms is provided by assertion 1. The remaining statements are rather easy; they are given by Lemma 1.3 of [AST08].

3. We should prove that any projective object  $P_0$  of  $\underline{H}t$  "lifts" to  $P_t$ .

Now, the functor  $H_0^t$  respects products according to the easy Lemma 1.4 of [Nee18] (applied in the dual form; cf. also Proposition 3.4(2) of [BoV19]); thus the composition  $G^{P_0} = \underline{H}t(P_0, -) \circ H_0^t : \underline{C} \rightarrow \text{Ab}$  respects products as well. Moreover,  $G^{P_0}$  is obviously a homological functor. Thus it is corepresentable by some  $P \in \text{Obj } \underline{C}$  that clearly belongs to  $P_t$ , and it remains to apply the previous assertion.  $\square$

**Proposition 5.1.5.** Assume that  $t$  is a  $t$ -structure on  $\underline{C}$  orthogonal to a weight structure  $w$  on  $\underline{C}' \subset \underline{C}$ .

1. Then  $\underline{C}'_{w \geq 0} = \underline{C}_{t \geq 0} \cap \text{Obj } \underline{C}$ ,  $\underline{C}'_{w \leq 0} = {}^{\perp}(\underline{C}_{t \geq 1}) \cap \text{Obj } \underline{C}$ , and  $\underline{C}'_{w=0} = P'_t$ .
2. If  $t$  is  $+$ -orthogonal to  $w$  then  $\underline{C}_{t \geq 0}$  is closed with respect to  $\underline{C}$ -products.



3. Assume that  $t$  is  $-$  or  $+$ -orthogonal to  $w$ . Then  $\{E^P : P \in \underline{C}_{w=0}\}$  is a conservative collection of functors  $\underline{H}t \rightarrow \text{Ab}$  (cf. Remark 5.1.6 below).

4. Conversely, assume that functors of the type  $E^P$  for  $P \in \underline{C}_{w=0}$  form a conservative collection and  $t$  is right (resp. left) non-degenerate. Then  $t$  is  $-$  (resp.  $+$ ) right orthogonal to  $w$ .

*Proof.* 1. The argument is rather similar to the proof of Theorem 2.3.3(I). We will use the notation  $(C_1, C_2)$  for the couple  $({}^\perp \underline{C}_{t \geq 1} \cap \text{Obj } \underline{C}', \underline{C}_{t \geq 1} \cap \text{Obj } \underline{C}')$ .

Since  $w$  is orthogonal to  $t$ , we have  $\underline{C}'_{w \leq 0} \subset C_1$  and  $\underline{C}'_{w \geq 1} \subset C_2$ . Next,  $C_1 \perp C_2$ , and applying Proposition 1.2.4(2) we obtain  $w = (C_1, C_2[-1])$ ; see Remark 2.3.2(1). Hence  $\underline{C}'_{w=0} = C_1 \cap C_2[-1] = P'_t$ .

2. Obvious.

3. Since all of these functors are exact (see Proposition 5.1.4(1)), it suffices to verify that for any non-zero  $N \in \underline{C}_{t=0}$  there exists  $M \in \underline{C}'_{w=0}$  such that  $\underline{C}(M, N) \neq \{0\}$ . Now, if  $M \perp N$  for all  $M \in \underline{C}'_{w=0}$  then combining Proposition 2.2.2(5) with the description of  $w$ -pure functors provided by Theorem 2.1.2(2) of [Bon21] we obtain  $\text{Obj } \underline{C} \perp N$ . Combining this statement with either the  $-$  or the  $+$ -orthogonality of  $t$  to  $w$  we immediately obtain  $N = 0$  (i.e., a contradiction).

4. If  $t$  is right (resp. left) non-degenerate, it suffices to verify that  $H_i^t(N) = 0$  whenever  $i < 0$  (resp.  $i > 0$ ),  $N \in \text{Obj } \underline{C}$ , and  $M \perp N$  for all  $M \in \underline{C}'_{w=i}$ . For this purpose it is clearly sufficient to check that  $H_0^t(N) = 0$  whenever  $M \perp N$  for all  $M \in \underline{C}'_{w=0}$ . The latter statement immediately follows from our assumption on  $\underline{C}'_{w=0}$  along with assertion 1.  $\square$

*Remark 5.1.6.* Since the functors of the type  $E^P$  that we consider in parts 3–4 of our proposition are exact (on  $\underline{H}t$ ), the conservativity of  $\{E^P : P \in \underline{C}'_{w=0}\}$  is fulfilled if and only if for any non-zero  $N \in \underline{C}_{t=0}$  there exists  $P \in \underline{C}'_{w=0}$  such that  $E^P(N) \neq \{0\}$ .

## 5.2 On the existence of orthogonal weight structures

Now we study the question which weight structures are adjacent to  $t$ -structures; yet in certain cases we are only able to construct an orthogonal weight structure on a subcategory of the corresponding category.

**Proposition 5.2.1.** Assume that  $\underline{C}' \subset \underline{C}$ ,  $w$  is a weight structure on  $\underline{C}'$  that is orthogonal to a  $t$ -structure  $t$  on  $\underline{C}$ , and  $\underline{C}_{t=0} \subset \text{Obj } \underline{C}'$ .

1. Then there are enough projectives in  $\underline{H}t$ , for any  $M \in \underline{C}_{t=0}$  there exists an  $\underline{H}t$ -epimorphism from  $H_0^t(P)$  into  $M$  for some  $P \in \underline{C}'_{w=0}$ , and the functor  $H_0^t$  induces an equivalence of  $\text{Kar}(\underline{H}w)$  with the category of projective objects of  $\underline{H}t$ .

2. Moreover,  $\underline{H}w$  is equivalent to the latter category whenever the category  $\underline{H}w$  is Karoubian. In particular, this is the case if the class  $\text{Obj } \underline{C}'$  is retraction-closed in  $\underline{C}$  and  $\underline{C}$  is Karoubian.

3. Assume in addition that  $t$  is left non-degenerate. Then  $t$  is  $+$ -orthogonal to  $w$ ; hence  $\underline{C}_{t \leq 0}$  is closed with respect to  $\underline{C}$ -products.

*Proof.* 1. Fix  $M \in \underline{C}_{t=0}$  and consider its  $w$ -decomposition  $P \xrightarrow{p} M \rightarrow w_{\geq 1}M \rightarrow P[1]$ . Since  $M \in \underline{C}_{t=0}$ , Proposition 5.1.5(1) implies that  $P$  belongs to  $\underline{C}'_{w \geq 0}$ ; hence  $P$  belongs to  $\underline{C}'_{w=0}$  according to Proposition 1.2.4(7)). Next,  $P \in \underline{C}'_{w \geq 0} \subset$

$\underline{\mathcal{C}}_{t \geq 0}$  (see Proposition 5.1.5(1) once again); hence the object  $P_0 = t_{\leq 0}P$  equals  $H_0^t(P)$  (see Proposition 1.1.5(3)). Therefore  $P_0$  is projective in  $\underline{Ht}$  according to Proposition 5.1.4(2).

Next, the adjunction property for the functor  $t_{\leq 0}$  (see Proposition 1.1.5(1)) implies that  $p$  factors through the  $t$ -decomposition morphism  $P \rightarrow P_0$ . Now we check that the corresponding morphism  $P_0 \rightarrow M$  is an  $\underline{Ht}$ -epimorphism. This is clearly fulfilled if and only if  $C = \text{Cone}(P_0 \rightarrow M)$  belongs to  $\underline{\mathcal{C}}_{t \geq 1}$ . The octahedral axiom of triangulated categories gives a distinguished triangle  $(t_{\geq 1}P)[1] \rightarrow w_{\geq 1}M \rightarrow C \rightarrow (t_{\geq 1}P)[2]$ ; it yields the assertion in question since  $w_{\geq 1}M \in \underline{\mathcal{C}}'_{w \geq 1} \subset \underline{\mathcal{C}}_{t \geq 1}$  and the class  $\underline{\mathcal{C}}_{t \geq 1}$  is extension-closed (see Proposition 1.1.5(4)). Thus we obtain that  $\underline{Ht}$  has enough projectives.

Now, the category of projective objects of  $\underline{Ht}$  is clearly Karoubian. As we have just verified, for any projective object  $Q$  of  $\underline{Ht}$  there exists an  $\underline{Ht}$ -epimorphism  $H_0^t(S) \rightarrow Q$  for some  $S \in \underline{\mathcal{C}}'_{w=0}$ . Since  $H_0^t(S)$  is projective in  $\underline{Ht}$  according to Proposition 5.1.5(2), this epimorphism splits, i.e.,  $Q$  equals the image of some idempotent endomorphism of  $H_0^t(S)$ . Applying Proposition 5.1.5(2) once again and lifting this endomorphism to  $\underline{Hw}$  we obtain that  $\text{Kar}(\underline{Hw})$  is equivalent to the category of projective objects of  $\underline{Ht}$  indeed.

2. If  $\underline{Hw}$  is Karoubian,  $\underline{Hw} \cong \text{Kar}(\underline{Hw})$ ; hence we obtain the equivalence in question according to assertion 1.

Next,  $\underline{\mathcal{C}}'_{w=0}$  is retraction-closed in  $\underline{\mathcal{C}}'$  since  $\underline{\mathcal{C}}'_{w \leq 0}$  and  $\underline{\mathcal{C}}'_{w \geq 0}$  are. Consequently, if  $\text{Obj } \underline{\mathcal{C}}'$  is retraction-closed in  $\underline{\mathcal{C}}$  and  $\underline{\mathcal{C}}$  is Karoubian then  $\underline{Hw}$  is Karoubian as well, and this concludes the proof.

3. If  $t$  is  $+$ -orthogonal to  $w$  then  $\underline{\mathcal{C}}_{t \leq 0}$  is closed with respect to  $\underline{\mathcal{C}}$ -products according to Proposition 5.1.5(2).

Applying Proposition 5.1.5(3) we obtain that it remains to verify that the functors of the form  $E^M$  for  $M \in \underline{\mathcal{C}}'_{w=0}$  give a conservative family of functors  $\underline{Ht} \rightarrow \text{Ab}$ . Now, for any object  $N$  of  $\underline{Ht}$  our assumptions give the existence of a projective object  $P_0$  of  $\underline{Ht}$  that surjects onto it. Moreover, applying Proposition 5.1.4(2) we obtain the existence of  $P \in \underline{\mathcal{C}}'_{w=0}$  and a morphism  $h$  from  $P$  such that  $H_0^t(h)$  is isomorphic to this surjection  $P_0 \rightarrow N$ . Hence  $E^P(N) \neq 0$  if  $N$  is non-zero, and we obtain the conservativity in question (see Remark 5.1.6).  $\square$

**Theorem 5.2.2.** Assume that  $t$  is a  $t$ -structure on  $\underline{\mathcal{C}}$  and there are enough projectives in  $\underline{Ht}$ .

I. Assume that for any projective object  $P$  of  $\underline{Ht}$  there exists  $P' \in P_t$  (see Definition 5.1.3(2)) along with an  $\underline{Ht}$ -epimorphism  $H_0^t(P') \rightarrow P$ .

1. Then the full subcategory  $\underline{\mathcal{C}}_+$  of  $\underline{\mathcal{C}}$  whose object class equals  $\cup_{i \in \mathbb{Z}} \underline{\mathcal{C}}_{t \geq i}$  is triangulated, and there exists a weight structure  $w_+$  on  $\underline{\mathcal{C}}'$  such that  $\underline{\mathcal{C}}_{+, w_+ \geq 0} = \underline{\mathcal{C}}_{t \geq 0}$  and  $t$  is  $-$ -orthogonal to  $w_+$ .

2. Furthermore, one can extend (see Definition 1.2.2(4))  $w_+$  as above to a weight structure  $w$  on  $\underline{\mathcal{C}}$  that is (left) adjacent to  $t$  whenever any of the following additional assumptions is fulfilled:

- a.  $t$  is bounded below (see Definition 1.1.4(3)).
- b. There exists an integer  $n$  such that  $\underline{\mathcal{C}}_{t \leq 0} \perp \underline{\mathcal{C}}_{t \geq n}$ .

3. Assume that  $\underline{\mathcal{C}}$  is smashing. Then there also exists a smashing weight structure  $\tilde{w}$  on the localizing subcategory  $\tilde{\mathcal{C}}$  of  $\underline{\mathcal{C}}$  that is generated by  $\underline{\mathcal{C}}_{t \leq 0}$  such that  $\underline{\mathcal{C}}_{\tilde{w} \leq 0} = \underline{\mathcal{C}}_{t \leq 0}$ , and  $\underline{H\tilde{w}}$  is equivalent to the subcategory of projective objects of  $\underline{Ht}$ .

II. Assume in addition that  $\underline{\mathcal{C}}$  satisfies the dual Brown representability property (see Definition 3.2.1(1)),  $t$  is cosmashing and  $\underline{Ht}$  has enough projectives. Then the category  $\underline{\mathcal{C}}$  is smashing and for any projective object  $P$  of  $\underline{Ht}$  there exists  $P' \in P_t$  (see Definition 5.1.3(2)) such that  $H_0^t(P') \cong P$ .

Consequently, there exists a smashing weight structure  $\tilde{w}$  on the subcategory  $\tilde{\mathcal{C}}$  of  $\underline{\mathcal{C}}$  mentioned in assertion I.3, such that  $\underline{\mathcal{C}}_{\tilde{w} \leq 0} = \underline{\mathcal{C}}_{\leq 0}$ , and  $\underline{H\tilde{w}}$  is equivalent to the subcategory of projective objects of  $\underline{Ht}$ .

*Proof.* I.1.  $\underline{\mathcal{C}}_+$  is triangulated since it is obviously shift-stable and all  $\underline{\mathcal{C}}_{t \geq i}$  are extension-closed.

Next we take  $W_1 = {}^\perp \underline{\mathcal{C}}_{t \geq 1} \cap \text{Obj } \underline{\mathcal{C}}_+$ ,  $W_2 = \underline{\mathcal{C}}_{t \geq 0}$ , and prove that  $(W_1, W_2)$  is a weight structure on  $\underline{\mathcal{C}}_+$  (cf. Theorem 2.3.3(I)).

The only non-trivial axiom check here is the existence of  $w_+$ -decompositions for all objects of  $\underline{\mathcal{C}}$ . Let us verify the existence of a  $w$ -decomposition for any  $M \in \underline{\mathcal{C}}_{t \geq i}$  by downward induction on  $i$ . The statement is obvious for  $i > 0$  since  $M \in \underline{\mathcal{C}}_{+, w_+ \geq 1} = \underline{\mathcal{C}}_{t \geq 1}$  and we can take a "trivial" weight decomposition  $0 \rightarrow M \rightarrow M \rightarrow 0$ .

Now assume that existence of  $w_+$ -decompositions is known for any  $M \in \underline{\mathcal{C}}_{t \geq j}$  for some  $j \in \mathbb{Z}$ . We should verify the existence of weight decomposition of an element  $N$  of  $\underline{\mathcal{C}}_{t \geq j-1}$ . Clearly,  $N$  is an extension of  $N'[-j-1] = H_0^t(N[j+1])[-j-1]$  by  $t_{\geq j}N$  (see Proposition 1.1.5(3) for the notation). Since the latter object possesses a weight decomposition, Proposition 3.3.1(I) (with  $A = W_1$  and  $B = W_2[1]$ ) allows us to verify the existence of a weight decomposition of  $N'[-j-1]$  (instead of  $N$ ). Our assumptions imply that there exists an epimorphism  $H_0^t(P) \rightarrow N'$  with  $P \in P_t' = W_1 \cap W_2$ . Then a cone  $C$  of the corresponding composed morphism  $P \rightarrow N'$  is easily seen to belong to  $\underline{\mathcal{C}}_{t \geq 1}'$ . Since both  $P$  and  $C$  possess weight decompositions, applying Proposition 3.3.1(I) once again we obtain the statement in question.

Lastly,  $t$  is  $--$ -orthogonal to  $w$  immediately from Proposition 1.1.5(4).

2. If assumption a is fulfilled then we can just take  $w = w_+$  since  $\underline{\mathcal{C}}$  obviously equals  $\underline{\mathcal{C}}_+$ .

Now suppose that assumption b is fulfilled. Similarly to the previous proof, it suffices to verify that for the couple  $w = ({}^\perp \underline{\mathcal{C}}_{t \geq 1}, \underline{\mathcal{C}}_{t \geq 0})$  the corresponding decompositions exist for all objects of  $\underline{\mathcal{C}}$ .

Since  $w$  is an extension of  $w_+$ , assertion I.1 gives the existence of  $w$ -decompositions for all elements of  $\underline{\mathcal{C}}_{t \geq 2-n}$ . Next, our orthogonality assumption on  $t$  yields that  $\underline{\mathcal{C}}_{t \leq 1-n} \subset \underline{\mathcal{C}}_{w \leq 0}$ ; hence one can take trivial  $w$ -decompositions for elements of  $\underline{\mathcal{C}}_{t \leq 1-n}$ . It remains to note that  $\text{Obj } \underline{\mathcal{C}} = \underline{\mathcal{C}}_{t \geq 2-n} \star \underline{\mathcal{C}}_{t \leq 1-n} = \text{Obj } \underline{\mathcal{C}}$  by axiom (iv) of  $t$ -structures, and apply Proposition 3.3.1(I) once again.

3. We argue similarly to the proof of part I.1 and verify that the couple  $(\tilde{W}_1 = {}^\perp \underline{\mathcal{C}}_{t \geq 1} \cap \text{Obj } \tilde{\mathcal{C}}, \tilde{W}_2 = \underline{\mathcal{C}}_{t \geq 0})$  gives a weight structure  $\tilde{w}$  on  $\tilde{\mathcal{C}}$ . Indeed, this weight structure would be smashing since the class  $W_2 = \underline{\mathcal{C}}_{t \geq 0}$  is closed with respect to  $\underline{\mathcal{C}}$ -coproducts; see Proposition 1.1.5(5).

Once again, for this purpose it suffices to verify that the class  $\tilde{C} = \tilde{C}_1 \star \tilde{C}_2[1]$  equals  $\text{Obj } \tilde{\mathcal{C}}$ . Immediately from assertion I.1,  $\tilde{C}$  contains  $\underline{\mathcal{C}}_{t \geq j}$  for all  $j \in \mathbb{Z}$ . Moreover,  $\tilde{C}$  is extension-closed and closed with respect to  $\underline{\mathcal{C}}$ -coproducts according to Proposition 3.3.1(I, II.1); hence  $\tilde{C}$  equals  $\text{Obj } \tilde{\mathcal{C}}$  indeed.

Lastly,  $\text{Obj } \tilde{\mathcal{C}}$  is retraction-closed in  $\underline{\mathcal{C}}$  and  $\underline{\mathcal{C}}$  is Karoubian according to Proposition 3.3.1(II.2); hence  $\underline{H\tilde{w}}$  is equivalent to the subcategory of projective objects of  $\underline{Ht}$  according to Proposition 5.1.5(2).

II.  $\underline{\mathcal{C}}$  is smashing according to Proposition 8.4.6 of [Nee01] (applied in the dual form). Applying Proposition 5.1.5(2–3) we obtain that  $\underline{\mathcal{C}}$  and  $t$  satisfy the assumptions of assertion I.3.  $\square$

*Remark 5.2.3.* 1. Clearly, parts I.3 and II of our theorem become more interesting in the case  $\tilde{\mathcal{C}} = \underline{\mathcal{C}}$ . We will discuss certain assumptions that ensure this equality in Proposition 5.2.4 below.

2. Moreover, Proposition 5.2.1 along with Theorem 5.2.2(I.3,II) can be considered as a certain complement to Theorem 3.2.3(I). So we obtain that the class of  $t$ -structures adjacent to smashing weight ones is "closely related" to the one of cosmashing  $t$ -structures such that  $\underline{Ht}$  has enough projectives.

3. The condition  $\underline{\mathcal{C}}_{t \leq 0} \perp \underline{\mathcal{C}}_{t \geq n}$  for  $n \gg 0$  (see part I.2 of our theorem) is a natural generalization of the finiteness of the cohomological dimension condition (for an abelian category).

**Proposition 5.2.4.** Let  $\underline{\mathcal{C}}$  be a smashing category generated by some set of its objects.<sup>8</sup>

1. Assume that  $w$  is a smashing weight structure on  $\underline{\mathcal{C}}$ . Then there exists a set  $S$  of objects of  $\underline{\mathcal{C}}$  such that  $\underline{\mathcal{C}}_{w \geq 0}$  equals the smallest class of objects of  $\underline{\mathcal{C}}$  that is closed with respect to extensions and coproducts and contains  $S$ .

2. Assume that  $\underline{\mathcal{C}}$  is *well generated* in the sense of [Nee01, Remark 8.1.7] (and §6.3 of [Kra10]),  $t$  is a right non-degenerate  $t$ -structure on  $\underline{\mathcal{C}}$ , and there exists a set  $S$  of objects of  $\underline{\mathcal{C}}$  such that  $\underline{\mathcal{C}}_{t \geq 0}$  equals the smallest class of objects of  $\underline{\mathcal{C}}$  that is closed with respect to extensions and coproducts and contains  $S$ .

Then the localizing subcategory  $\tilde{\mathcal{C}}$  of  $\underline{\mathcal{C}}$  that is generated by  $\underline{\mathcal{C}}_{t \leq 0}$  is  $\underline{\mathcal{C}}$  itself.

*Proof.* 1. Immediate from Proposition 2.3.2(10) of [Bon21].

2. Since  $t$  is right non-degenerate and  $\cap_{i \in \mathbb{Z}} \underline{\mathcal{C}}_{t \leq i} = \cap_{i \in \mathbb{Z}} \underline{\mathcal{C}}_{t \geq i}^\perp$  (see Proposition 1.1.5(4)), we obtain  $\tilde{\mathcal{C}}^{\perp_{\underline{\mathcal{C}}}} = \{0\}$ .

Next, assume that the embedding  $\tilde{\mathcal{C}} \rightarrow \underline{\mathcal{C}}$  possesses a right adjoint. Then this functor is the localization by the subcategory  $\tilde{\mathcal{C}}^{\perp_{\underline{\mathcal{C}}}}$ ; see Propositions 4.9.1 and 4.10.1 of [Kra10].

Consequently, it remains to verify that the embedding  $\tilde{\mathcal{C}} \rightarrow \underline{\mathcal{C}}$  possesses a right adjoint indeed. Since  $\tilde{\mathcal{C}}$  is generated by  $S$  as a localizing subcategory of  $\underline{\mathcal{C}}$ , the latter statement is a well-known combination of Theorems 7.2.1(2) and 5.1.1(2) of loc. cit.  $\square$

*Remark 5.2.5.* 1. Consequently, if  $\underline{\mathcal{C}}$  is well generated, satisfies the dual Brown representability property,  $t$  is a right non-degenerate cosmashing  $t$ -structure on  $\underline{\mathcal{C}}$  and  $\underline{Ht}$  contains enough projectives then there exists a weight structure adjacent to  $t$  if and only if there exists a set that "generates"  $S$  as in Proposition 5.2.4(2); see Theorem 5.2.2(II).

2. Recall also that Theorem 3.9 of [CGR14] ensures the existence of the right adjoint to the embedding  $\tilde{\mathcal{C}} \rightarrow \underline{\mathcal{C}}$  under certain assumptions that do not depend on the existence of a set of generators  $S$  as in Proposition 5.2.4(2).

---

<sup>8</sup>Actually, if  $\underline{\mathcal{C}}$  is generated by a set  $\{C_i\}$  then it is also generated by the single object  $\coprod C_i$ . This observation also extends to all generation assumptions of this proposition; see Corollary 2.1.3(2) of [BoS19].

### 5.3 Some adjacent weight structures in the saturated case

Now we proceed towards extending Theorem 5.2.2(2a) to a rather nice statement on saturated categories. To formulate it in a rather general form we will define some new types of  $t$ -structures.

**Definition 5.3.1.** Let  $t$  be a  $t$ -structure on  $\underline{\mathcal{C}}$ .

1. We will say that  $t$  is *essentially bounded (below)* whenever for any  $M \in \text{Obj } \underline{\mathcal{C}}$  we have  $H_i^t = 0$  for  $|i| \gg 0$  (resp. for  $i \ll 0$ ; see Proposition 1.1.5(3)).
2. We define  $\underline{\mathcal{C}}_{t=-\infty}$  as the full subcategory of  $\underline{\mathcal{C}}$  whose object class equals  $\bigcap_{i \in \mathbb{Z}} \underline{\mathcal{C}}_{t \leq i}$ .

Let us now relate these notions to semi-orthogonal decompositions.

**Proposition 5.3.2.** Let  $t$  be a  $t$ -structure  $t$  on  $\underline{\mathcal{C}}$ .

- I.1. Then the subcategory  $\underline{\mathcal{C}}_{t=-\infty}$  is triangulated and  $t$  restricts to them.
2. The restriction  $t_+$  of  $t$  to  $\underline{\mathcal{C}}_+$  is bounded. Moreover,  $t_+$  is essentially bounded if  $t$  is.
3.  $\text{Obj } \underline{\mathcal{C}}_+ \perp \text{Obj } \underline{\mathcal{C}}_{t=-\infty}$ .
- II. Assume that  $t$  is essentially bounded below.  
Then  $(\underline{\mathcal{C}}_{t=-\infty}, \underline{\mathcal{C}}_+)$  is a semi-orthogonal decomposition of  $\underline{\mathcal{C}}$ .

*Proof.* I. All of these statements are quite simple.

1. The subcategory  $\underline{\mathcal{C}}_{t=-\infty}$  is triangulated since it is obviously shift-stable and all  $\underline{\mathcal{C}}_{t \leq i}$  are extension-closed (see Proposition 1.1.5(4)).  $t$  restricts to them since the functors  $L_t$  and  $R_t$  respect all  $\underline{\mathcal{C}}_{t \leq i}$ ; here one can apply axiom (ii) of Definition 1.1.3 and Proposition 1.1.5(1,3).

2,3. Obvious.

II. We should (see Definition 1.2.5) verify that for any  $M \in \text{Obj } \underline{\mathcal{C}}$  there exists a distinguished triangle

$$B \rightarrow M \rightarrow A \rightarrow B[1] \quad (5.3.1)$$

with  $B \in \text{Obj } \underline{\mathcal{C}}_+$  and  $A \in \text{Obj } \underline{\mathcal{C}}_{t=-\infty}$ . Since  $t$  is essentially bounded below, there exists  $m \in \mathbb{Z}$  such that  $H_i^t = 0$  for  $i < m$ . Then for the corresponding distinguished triangle

$$t_{\geq m} M \rightarrow M \rightarrow t_{\leq m-1} M \rightarrow (t_{\geq m} M)[1]$$

(this is the shift of the  $t$ -decomposition triangle for  $M[m]$  by  $[-m]$ ; cf. Proposition 1.1.5(3)) we have  $t_{\geq m} M \in \underline{\mathcal{C}}_{t \geq m} \subset \text{Obj } \underline{\mathcal{C}}_+$ , whereas the object  $A = t_{\leq m-1} M$  equals  $t_{\leq m-2} M = t_{\leq m-3} M = \dots$ ; hence  $A \in \text{Obj } \underline{\mathcal{C}}_{t=-\infty}$  indeed.  $\square$

**Theorem 5.3.3.** Assume that  $\underline{\mathcal{C}}^{op}$  is an  $R$ -saturated category,  $t$  is an essentially bounded below  $t$ -structure on  $\underline{\mathcal{C}}$ , and there are enough projectives in  $\underline{Ht}$ .

1. If  $t$  is also essentially bounded then for any projective object  $P$  of  $\underline{Ht}$  there exists  $P' \in P_t$  (see Definition 5.1.3(2)) such that  $H_0^t(P') \cong P$ .
2. Assume that for any projective object  $P$  of  $\underline{Ht}$  there exists  $P' \in P_t$  (see Definition 5.1.3(2)) along with an  $\underline{Ht}$ -epimorphism  $H_0^t(P') \rightarrow P$ .

Then there exists an essentially bounded below weight structure  $w$  on  $\underline{\mathcal{C}}$  (left) adjacent to  $t$ .

*Proof.* 1. It suffices to note that the functor  $\underline{H}t(P, -) \circ H_0^t$  is locally bounded (as a functor from  $\underline{\mathcal{C}}^{op}$  into  $R - \text{Mod}$ ) and takes values in  $R - \text{mod}$ .

2. We construct this weight structure as an extension of the weight structure  $w_+$  provided by Theorem 5.2.2(I.1). As we have noted in Remark 4.2.3(2), applying Corollary 4.2.2(1) to the category  $\underline{\mathcal{C}}^{op}$  one can obtain a semi-orthogonal decomposition  $(\underline{\mathcal{C}}_+, \underline{\mathcal{C}}_0)$  of  $\underline{\mathcal{C}}$  (see Remark 1.2.7(2)) with  $\text{Obj } \underline{\mathcal{C}}_0 = {}^\perp(\text{Obj } \underline{\mathcal{C}}_+)$ . Then Proposition 3.2(5) of [BoV19] implies that  $w = (\text{Obj } \underline{\mathcal{C}}_0 \star \underline{\mathcal{C}}_{+, w_+ \leq 0}, \underline{\mathcal{C}}_{t \geq 0})$  is a weight structure. Here we apply Proposition 5.3.2(II) and duality to obtain the existence of the adjoint functor mentioned in loc. cit. and recall that  $\underline{\mathcal{C}}_{t \geq 0} = \underline{\mathcal{C}}_{+, w_+ \geq 0}$ .

Lastly, this  $w$  is clearly adjacent to  $t$  (see Proposition 1.3.3).  $\square$

*Remark 5.3.4.* 1. Clearly, all the statements and definitions of this section can be dualized.

2. It would be interesting to argue similarly to the proof of Theorem 5.3.3 in the context of Theorem 5.2.2(II). The main problem is to find conditions that would enforce  $\tilde{w}$  to be closed with respect to  $\underline{\mathcal{C}}$ -products.

## References

- [AJS03] Alonso L., Jeremías A., Souto M.J., Construction of  $t$ -structures and equivalences of derived categories// Trans. of the AMS, 355(6), 2003, 2523–2543.
- [AJPV08] Alonso Tarrió L., Jeremías López A., Pérez Rodríguez M., Vale Gonçalves M.J., The derived category of quasi-coherent sheaves and axiomatic stable homotopy// Adv. Math. 218(4), 2008, 1224–1252.
- [AST08] Assem I., Souto M.J., Trepode S., Ext-projectives in suspended subcategories// J. Pure Appl. Algebra 212 (2008), 423–434.
- [BBD82] Beilinson A., Bernstein J., Deligne P., Faisceaux pervers// Asterisque 100 (1982), 5–171.
- [BoK89] Bondal A.I., Kapranov M.M., Representable functors, Serre functors, and mutations// Izvestiya Rossiiskoi Akademii Nauk. Seriya Matematicheskaya, 53(6), 1989, 1183–1205, transl. in Izvestiya: Mathematics 35.3 (1990), 519–541.
- [BVd03] Bondal A.I., Van den Bergh M., Generators and representability of functors in commutative and noncommutative geometry// Mosc. Math. J. 3(1), 2003, 1–36.
- [Bon10a] Bondarko M.V., Weight structures vs.  $t$ -structures; weight filtrations, spectral sequences, and complexes (for motives and in general)// J. of K-theory, v. 6(3), 2010, 387–504, see also <http://arxiv.org/abs/0704.4003>
- [Bon10b] Bondarko M.V., Motivically functorial coniveau spectral sequences; direct summands of cohomology of function fields// Doc. Math., extra volume: Andrei Suslin’s Sixtieth Birthday (2010), 33–117; see also <http://arxiv.org/abs/0812.2672>

- [Bon19] Bondarko M.V., From weight structures to (orthogonal)  $t$ -structures and back, preprint, 2019, <https://arxiv.org/abs/1907.03686>
- [Bon21] Bondarko M.V., On weight complexes, pure functors, and detecting weights// J. of Algebra 574 (2021), 617–668.
- [Bon22a] Bondarko M.V., On perfectly generated weight structures and adjacent  $t$ -structures// Math. Zeit., 300, 2022, 1421–1454.
- [Bon22b] Bondarko M.V., On morphisms killing weights and stable Hurewicz-type theorems, J. Math. Jussieu, 2022, <https://doi.org/10.1017/S1474748022000470>
- [Bon24] Bondarko M.V., [https://www.researchgate.net/publication/371875245\\_Producing\\_new\\_semi](https://www.researchgate.net/publication/371875245_Producing_new_semi) to appear in Mat. Sbornik, vol. 215(4); see also <https://arxiv.org/abs/2203.07315>
- [BoK18] Bondarko M.V., Kumallagov D.Z., On Chow weight structures without projectivity and resolution of singularities (Russian)// Algebra i Analiz 30(5), 2018, 57–83, transl. in St. Petersburg Math. J. 30 (2019), no. 5, 803–819.
- [BoS18] Bondarko M.V., Sosnilo V.A., On constructing weight structures and extending them to idempotent extensions// Homology, Homotopy and Appl., vol 20(1), 2018, 37–57.
- [BoS19] Bondarko M.V., Sosnilo V.A., On purely generated  $\alpha$ -smashing weight structures and weight-exact localizations// J. of Algebra 535 (2019), 407–455.
- [BoV19] Bondarko M.V., Vostokov S.V., On torsion theories, weight and  $t$ -structures in triangulated categories (Russian)// Vestnik St.-Petersbg. Univ. Mat. Mekh. Astron., vol 6(64), iss. 1, 27–43, 2019; transl. in Vestnik St. Peters. Univers., Mathematics, 2019, vol. 52(1), 19–29.
- [BoV23] Bondarko M.V., Vostokov S.V., Killing Weights from the Perspective of  $t$ -Structures// Proceedings of the Steklov Institute of Math. 320 (2023), 51–61.
- [CGR14] Casacuberta C., Gutierrez J.J., Rosicky J., Are all localizing subcategories of stable homotopy categories coreflective// Advances in Math. 252 (2014), 158–184.
- [IyY08] Iyama O., Yoshino Y., Mutation in triangulated categories and rigid Cohen-Macaulay modules// Inv. math. 172(1), 2008, 117–168.
- [KeN13] Keller B., Nicolas P., Weight structures and simple dg modules for positive dg algebras // Int. Math. Res. Not., vol. 2013(5), 2013, 1028–1078.
- [Kra10] Krause H., Localization theory for triangulated categories, in: Triangulated Categories, London Math. Soc. Lecture Note Ser. 375 (2010), 161–235.
- [KoY14] Koenig S., Yang D., Silting Objects, Simple-Minded Collections,  $t$ -Structures and Co- $t$ -Structures for Finite-Dimensional Algebras// Doc. Math. 19 (2014), 403–438.

- [Lur17] Lurie J., Higher Algebra, September 2017, <http://www.math.harvard.edu/~lurie/papers/HA.pdf>
- [HVVS13] Hernández O.M., Valadez E.C.S., Vargas V.S., Salorio M.J.S., Auslander-Buchweitz context and co-t-structures// Applied Categorical Structures 21(5), 2013, 417–440.
- [Nee01] Neeman A., Triangulated Categories, Annals of Mathematics Studies 148 (2001), Princeton University Press, viii+449 pp.
- [Nee08] Neeman A., Brown representability follows from Rosický’s theorem// J. of Topology, 2.2 (2009), 262–276.
- [Nee18] Neeman A., Triangulated categories with a single compact generator and a Brown representability theorem, preprint, 2018, <https://arxiv.org/abs/1804.02240>
- [Nee21] Neeman A., The  $t$ -structures generated by objects// Trans. Amer. Math. Soc. 374 (2021), 8161–8175.
- [NiS09] Nicolas P., Saorin M., Parametrizing recollement data for triangulated categories// J. of Algebra 322 (2009), 1220–1250.
- [Pau08] Pauksztello D., Compact corigid objects in triangulated categories and co-t-structures// Central European Journal of Mathematics, vol. 6(1), 2008, 25–42.
- [Pau12] Pauksztello D., A note on compactly generated co-t-structures// Comm. in Algebra, vol. 40(2), 2012, 386–394.
- [PoS16] Pospisil D., Šťovíček J., On compactly generated torsion pairs and the classification of co-t-structures for commutative Noetherian rings// Trans. Amer. Math. Soc. 368 (2016), 6325–6361.
- [Ros05] Rosicky J., Generalized Brown representability in homotopy categories// Theory and applications of categories, vol. 14(19), 2005, 451–479.
- [SaS11] Saorin M., Šťovíček J., On exact categories and applications to triangulated adjoints and model structures// Adv. Math. 228(2), 968–1007, 2011.
- [Tak08] Takahashi R., Classifying subcategories of modules over a Noetherian ring// J. Lond. Math. Soc. 78 (3), 2008, 767–782.
- [stacks] The Stacks project, <https://stacks.math.columbia.edu/browse>