

# VARIETIES OF MONOIDS WITH A DISTRIBUTIVE SUBVARIETY LATTICE

SERGEY V. GUSEV

ABSTRACT. A monoid is aperiodic if all its subgroups are trivial. We completely classify all varieties of aperiodic monoids whose subvariety lattice is distributive.

## 0. BACKGROUND AND OVERVIEW

A *variety* is a class of universal algebras of a fixed type that is closed under the formation of homomorphic images, subalgebras and arbitrary direct products. A variety  $\mathbf{V}$  is *distributive* if its lattice  $\mathfrak{L}(\mathbf{V})$  of subvarieties is distributive.

There was genuine interest to investigate distributive varieties of groups. The first example of a non-distributive variety of groups was discovered by Higman [13] in the mid-1960s; see also Neumann [22, 54.24]. At the turn of the 1960s and 1970s, there were a lot of articles on this topic; see Cossey [2], Kovács and Newman [20] and Roman'kov [25], for instance. However, over time, the activity began to fade. The general problem of describing distributive varieties of groups turned out to be highly infeasible. Here it suffices to refer to the following result by Kozhevnikov [21]: there exist continuum many varieties of groups whose lattice of subvarieties is isomorphic to the 3-element chain.

In 1979, Shevrin [32, Problem 2.60a] posed the problem of classifying distributive varieties of semigroups. In the early 1990s, in a series of papers, Volkov solved this problem in a very wide partial case. In particular, Volkov completely classified all distributive varieties of *aperiodic* semigroups, i.e., semigroups all whose subgroups are trivial. In view of the above-mentioned result by Kozhevnikov [21], a classification of distributive varieties is hardly possible in general. See the survey article by Shevrin et al [31, Section 11] for more details.

The present article is concerned with the distributive varieties of *monoids*, i.e., semigroups with an identity element. Even though monoids are very similar to semigroups, the story turns out to be very different and difficult. Until recently, distributive varieties of monoids have not been studied systematically, although non-trivial examples of such varieties have long been known; see Head [12] and Wismath [33]. Over the past years, the number of accumulated examples has increased significantly; see Gusev and O.Sapir [9], Jackson [14], Jackson and Lee [15], Lee [16–18], and Zhang and Luo [34], for instance. This has paved the way for the systematic study of distributive varieties of monoids. More information and many references can be found in the recent survey [8, Section 6.3].

---

URAL FEDERAL UNIVERSITY, INSTITUTE OF NATURAL SCIENCES AND MATHEMATICS, LENINA 51, EKATERINBURG 620000, RUSSIA

*E-mail address:* sergey.gusb@gmail.com.

2010 *Mathematics Subject Classification.* 20M07 (Primary), 08B15 (Secondary).

*Key words and phrases.* Monoid, aperiodic monoid, variety, subvariety lattice, distributive lattice.

Supported by the Ministry of Science and Higher Education of the Russian Federation (project FEUZ-2023-0022).

As in the semigroup case, in view of the result by Kozhevnikov [21], one can hardly hope to describe distributive varieties of monoids in general. Therefore, first of all, it is logical to focus specifically on the class of aperiodic monoids, in which, at least hypothetically, one can hope for an exhaustive description of distributive varieties.

**Problem 1** ([8, Problem 6.10]). Describe distributive varieties of aperiodic monoids.

The article [10] describing varieties of aperiodic monoids whose subvariety lattice is a chain can be considered as the first attempt to take the first step in solving Problem 1, since being a chain is a stronger property than satisfying the distributive law. However, the systematic study of just distributive varieties of aperiodic monoids is started by the author in [6], where distributive varieties are described within the class of aperiodic monoids with central idempotents. In the follow-up paper [7], this result is extended to the class of aperiodic monoids with commuting idempotents. The present article, the third and the last work of the cycle, completes the story. We completely classify all distributive varieties of aperiodic monoids, solving Problem 1.

We provide an equational description of such varieties. Namely, we present 5 countably infinite series of varieties and 29 “sporadic” varieties such that every distributive variety of aperiodic monoids is contained in one of them. Notice that the proof this result implies that set of all distributive varieties of aperiodic monoids is countably infinite, although the set of all distributive varieties of monoids is uncountably infinite [21].

This paper is structured as follows. In Section 1, we formulate and discuss our main result. Some background results are first given in Section 2. In Section 3, exclusion identities are found for a number of certain varieties of monoids. Section 4 contains a series of examples of non-distributive varieties of monoids. Finally, Section 5 is devoted to the proof of our main result.

## 1. OUR MAIN RESULT

Let us briefly recall a few notions that we need to formulate our main result. Let  $\mathfrak{X}$  be a countably infinite set called an *alphabet*. As usual, let  $\mathfrak{X}^*$  denote the free monoid over the alphabet  $\mathfrak{X}$ . Elements of  $\mathfrak{X}$  are called *letters* and elements of  $\mathfrak{X}^*$  are called *words*. We treat the identity element of  $\mathfrak{X}^*$  as *the empty word*, which is denoted by 1. Words and letters are denoted by small Latin letters. However, words unlike letters are written in bold. An identity is written as  $\mathbf{u} \approx \mathbf{v}$ , where  $\mathbf{u}, \mathbf{v} \in \mathfrak{X}^*$ ; it is *non-trivial* if  $\mathbf{u} \neq \mathbf{v}$ .

As usual,  $\mathbb{N}$  denote the set of all natural numbers. For any  $n \in \mathbb{N}$ , we denote by  $S_n$  the full symmetric group on the set  $\{1, \dots, n\}$ . For any  $n, m, k \in \mathbb{N}$ ,  $\rho \in S_{n+m}$  and  $\tau \in S_{n+m+k}$ , we define the words:

$$\begin{aligned} \mathbf{a}_{n,m}[\rho] &:= \left( \prod_{i=1}^n z_i t_i \right) x \left( \prod_{i=1}^{n+m-1} z_i \rho y_i^2 \right) z_{(n+m)} \rho x \left( \prod_{i=n+1}^{n+m} t_i z_i \right), \\ \bar{\mathbf{a}}_{n,m}[\rho] &:= \left( \prod_{i=1}^n z_i t_i \right) x \left( \prod_{i=1}^{n+m-1} z_i \rho y_i^2 x \right) z_{(n+m)} \rho x \left( \prod_{i=n+1}^{n+m} t_i z_i \right), \\ \mathbf{c}_{n,m,k}[\tau] &:= \left( \prod_{i=1}^n z_i t_i \right) x y t \left( \prod_{i=n+1}^{n+m} z_i t_i \right) x \left( \prod_{i=1}^{n+m+k-1} z_i \tau y_i^2 \right) z_{(n+m+k)} \tau y \left( \prod_{i=n+m+1}^{n+m+k} t_i z_i \right). \end{aligned}$$

Let  $\mathbf{c}'_{n,m,k}[\tau]$  denote the word obtained from  $\mathbf{c}_{n,m,k}[\tau]$  by swapping the first occurrences of  $x$  and  $y$ . We denote also by  $\mathbf{d}_{n,m,k}[\tau]$  and  $\mathbf{d}'_{n,m,k}[\tau]$  the words obtained from the words  $\mathbf{c}_{n,m,k}[\tau]$  and  $\mathbf{c}'_{n,m,k}[\tau]$ , respectively, when reading the last words from right to left. We fix notation

for the following three identities:

$$\sigma_1 : xyzxt y \approx yxzxt y, \quad \sigma_2 : xzytx y \approx xzytyx, \quad \sigma_3 : xzxyty \approx xzyxt y.$$

Let

$$\begin{aligned} \Phi &:= \{x^2 \approx x^3, x^2 y^2 \approx y^2 x^2\}, \\ \Phi_1 &:= \{\mathbf{c}_{k,\ell,m}[\rho] \approx \mathbf{c}'_{k,\ell,m}[\rho], \mathbf{d}_{k,\ell,m}[\rho] \approx \mathbf{d}'_{k,\ell,m}[\rho] \mid k, \ell, m \in \mathbb{N}, \rho \in S_{k+\ell+m}\}, \\ \Phi_2 &:= \{\mathbf{a}_{k,\ell}[\rho] \approx \bar{\mathbf{a}}_{k,\ell}[\rho] \mid k, \ell \in \mathbb{N}, \rho \in S_{k+\ell}\}. \end{aligned}$$

Let  $\text{var } \Sigma$  denote the monoid variety given by a set  $\Sigma$  of identities. If  $\mathbf{V}$  is a monoid variety, then  $\mathbf{V}^\delta$  denotes the variety *dual* to  $\mathbf{V}$ , i.e., the variety consisting of monoids anti-isomorphic to monoids from  $\mathbf{V}$ .

Our main result is the following

**Theorem 1.1.** *A variety of aperiodic monoids is distributive if and only if it is contained in one of the varieties*

$$\begin{aligned} \mathbf{B} &:= \text{var} \{x \approx x^2\}, \\ \mathbf{D}_1 &:= \text{var} \{\Phi, xyx \approx xyx^2, x^2 y \approx x^2 yx\}, \\ \mathbf{D}_2 &:= \text{var} \{\Phi, \Phi_1, \Phi_2, xyx \approx xyx^2\}, \\ \mathbf{D}_3 &:= \text{var} \{\Phi, \sigma_2, \sigma_3, x^2 y \approx x^2 yx \approx xyx^2, xyxzx \approx xyxzx^2\}, \\ \mathbf{D}_4 &:= \text{var} \{\Phi, \sigma_2, \sigma_3, x^2 y \approx x^2 yx, x^2 yty \approx yx^2 ty, xyzx^2 ty^2 \approx yxzx^2 ty^2, xyxzx \approx xyx^2 zx\}, \\ \mathbf{D}_5 &:= \text{var} \{\Phi, \sigma_2, \sigma_3, xyx^2 \approx x^2 yx, xyzx^2 y \approx yxzx^2 y, xyxzx \approx xyxzx^2\}, \\ \mathbf{D}_6 &:= \text{var} \{\Phi, \sigma_2, \sigma_3, x^2 yx \approx x^2 yx^2, x^2 yty \approx yx^2 ty, xyzx^2 ty \approx yxzx^2 ty, xyzylx^2 \approx yxzytx^2\}, \\ \mathbf{D}_7 &:= \text{var} \left\{ \begin{array}{l} \Phi, \sigma_2, \sigma_3, x^2 yx \approx x^2 yx^2, x^2 yty \approx yx^2 ty, xyzx^2 ty^2 \approx yxzx^2 ty^2 \\ xyzx^2 y \approx yxzx^2 y, xyxzx \approx xyx^2 zx \end{array} \right\}, \\ \mathbf{D}_8 &:= \text{var} \left\{ \begin{array}{l} \Phi, \sigma_2, \sigma_3, x^2 yx \approx x^2 yx^2, x^2 yty \approx yx^2 ty, xyzx^2 ty^2 \approx yxzx^2 ty^2 \\ xyzx^2 y \approx yxzx^2 y, xyzx^2 tysx \approx yxzx^2 tysx, xyxzx \approx xyxzx^2 \end{array} \right\}, \\ \mathbf{D}_9 &:= \text{var} \{\Phi, \sigma_1, \sigma_3, x^2 yx \approx x^2 yx^2, ylx^2 y \approx ytyx^2\}, \\ \mathbf{D}_{10} &:= \text{var} \{\Phi, \sigma_1, \sigma_3, x^2 yx \approx x^2 yx^2, xyxzx \approx xyxzx^2, x^2 zytxy \approx x^2 zyt yx\}, \\ \mathbf{D}_{11} &:= \text{var} \{\Phi, \Phi_1, \Phi_2, xyx^2 \approx x^2 yx, xyxzx \approx xyxzx^2\}, \\ \mathbf{D}_{12} &:= \text{var} \left\{ \begin{array}{l} \Phi, \Phi_1, \Phi_2, x^2 yx \approx x^2 yx^2, ylx^2 y \approx ytyx^2, x^2 yty \approx yx^2 ty \\ xyzx^2 ty \approx yxzx^2 ty, xyzylx^2 \approx yxzytx^2, yzytxx^2 \approx yzxytx^2 \end{array} \right\}, \\ \mathbf{D}_{13} &:= \text{var} \left\{ \begin{array}{l} \Phi, \Phi_1, \Phi_2, x^2 yx \approx x^2 yx^2, yx^2 ty \approx xyx^2 ty, ytyx^2 \approx ytxyx^2, xyxzx \approx xyx^2 zx \\ x^2 yty^2 \approx yx^2 ty^2, x^2 yzytx \approx yx^2 zytx, x^2 yzxt y \approx yx^2 zxt y, x^2 zyt yx \approx x^2 zyt yx \\ yzx^2 ytx \approx yzyx^2 tx, xyzx^2 ty^2 \approx yxzx^2 ty^2 \end{array} \right\}, \\ \mathbf{D}_{14} &:= \text{var} \left\{ \begin{array}{l} \Phi, \Phi_1, \Phi_2, x^2 yx \approx x^2 yx^2, yx^2 ty \approx xyx^2 ty, ytyx^2 \approx ytxyx^2, xyxzx \approx xyxzx^2 \\ x^2 yty^2 \approx yx^2 ty^2, x^2 yzytx \approx yx^2 zytx, x^2 yzxt y \approx yx^2 zxt y, x^2 zyt yx \approx x^2 zyt yx \\ yzx^2 ytx \approx yzyx^2 tx, xyzx^2 ty^2 \approx yxzx^2 ty^2, xyzx^2 tysx \approx yxzx^2 tysx \end{array} \right\}, \\ \mathbf{D}_{15} &:= \text{var} \{\Phi_1, xyx \approx x^2 yx \approx xyx^2, (xy)^2 \approx (yx)^2\}, \\ \mathbf{P}_n &:= \text{var} \{\Phi_1, \Phi_2, x^n \approx x^{n+1}, x^2 y \approx yx^2\}, \\ \mathbf{Q}_n &:= \text{var} \{x^n \approx x^{n+1}, x^n y \approx yx^n, x^2 y \approx xyx\}, \\ \mathbf{R}_n &:= \text{var} \{\sigma_1, \sigma_2, x^n \approx x^{n+1}, x^2 y \approx yx^2\}, \quad n \in \mathbb{N}, \end{aligned}$$

or the dual ones.

A variety  $\mathbf{V}$  is *self-dual* if  $\mathbf{V} = \mathbf{V}^\delta$ . Notice that only the varieties  $\mathbf{B}$ ,  $\mathbf{D}_{11}$ ,  $\mathbf{D}_{15}$  and  $\mathbf{P}_n$  are self-dual among the varieties listed in Theorem 1.1. Thus, every distributive variety of aperiodic monoids is contained in either a variety in 5 countably infinite series or 29 “sporadic” varieties.

In [4], Green introduces five equivalence relations on a semigroup. These relations are collectively referred to as Green’s relations, and play a fundamental role in studying semigroups. Recall that two elements of a monoid  $M$  are in the Green’s relation  $\mathcal{J}$  if they generate the same principal ideal in  $M$ . A monoid  $M$  is  *$\mathcal{J}$ -trivial* if the Green’s relation  $\mathcal{J}$  is the equality relation in  $M$ . It is well known and can be easily verified that any variety of  $\mathcal{J}$ -trivial monoids satisfies the identities  $x^n \approx x^{n+1}$  and  $(xy)^n \approx (yx)^n$  for some  $n \in \mathbb{N}$ ; see Proposition 4.4 in [24] and its proof, for instance. It is easy to see that all the varieties listed in Theorem 1.1 except  $\mathbf{B}$  satisfy these identities. The variety  $\mathbf{B}$  is nothing but the variety of all band monoids. Thus, every distributive variety of aperiodic monoids is either a variety of bands or a variety of  $\mathcal{J}$ -trivial monoids.

A variety is *locally finite* if every finitely generated member in  $\mathbf{V}$  is finite. It follows from the fundamental result of M.Sapir [26, Theorem P] that all the varieties listed in Theorem 1.1 are locally finite. Hence every distributive variety of aperiodic monoids is so. In contrast, there is a non-locally finite distributive variety of groups [21].

In view of the result of Wismath [33], the lattice  $\mathfrak{L}(\mathbf{B})$  is countably infinite. As noted in [7], if  $\mathbf{X}$  is one of the varieties  $\mathbf{D}_1$ – $\mathbf{D}_{14}$ ,  $\mathbf{P}_n$ ,  $\mathbf{Q}_n$  or  $\mathbf{R}_n$ , then each subvariety of  $\mathbf{X}$  may be given within  $\mathbf{X}$  by a finite set of identities. Finally, it follows from the proof of Theorem 1.1 in Section 5 that the variety  $\mathbf{D}_{15}$  contains countably many subvarieties. Consequently, the set of all distributive varieties of aperiodic monoids is countably infinite.

## 2. PRELIMINARIES

**2.1. Distributive varieties of aperiodic monoids with commuting idempotents.** First, we remind the main result of [7].

**Proposition 2.1** ([7, Theorem 1.1]). *A variety of aperiodic monoids with commuting idempotents is distributive if and only if it is contained in one of the varieties  $\mathbf{D}_1$ – $\mathbf{D}_{14}$ ,  $\mathbf{P}_n$ ,  $\mathbf{Q}_n$ ,  $\mathbf{R}_n$ ,  $n \in \mathbb{N}$ , or the dual ones.*  $\square$

**2.2. Words, identities, deduction.** The *alphabet* of a word  $\mathbf{w}$ , i.e., the set of all letters occurring in  $\mathbf{w}$ , is denoted by  $\text{alph}(\mathbf{w})$ . For a word  $\mathbf{w}$  and a letter  $x$ , let  $\text{occ}_x(\mathbf{w})$  denote the number of occurrences of  $x$  in  $\mathbf{w}$ . A letter  $x$  is called *simple* [*multiple*] *in a word*  $\mathbf{w}$  if  $\text{occ}_x(\mathbf{w}) = 1$  [respectively,  $\text{occ}_x(\mathbf{w}) > 1$ ]. The set of all simple [multiple] letters in a word  $\mathbf{w}$  is denoted by  $\text{sim}(\mathbf{w})$  [respectively  $\text{mul}(\mathbf{w})$ ]. If  $\mathbf{w}$  is a word and  $x_1, \dots, x_k \in \text{alph}(\mathbf{w})$ , then we denote by  $\mathbf{w}(x_1, \dots, x_k)$  the word obtained from  $\mathbf{w}$  by removing all occurrences of letters from  $\text{alph}(\mathbf{w}) \setminus \{x_1, \dots, x_k\}$ .

A variety  $\mathbf{V}$  *satisfies* an identity  $\mathbf{u} \approx \mathbf{v}$ , if for any monoid  $M \in \mathbf{V}$  and any substitution  $\phi: \mathfrak{X} \rightarrow M$ , the equality  $\phi(\mathbf{u}) = \phi(\mathbf{v})$  holds in  $M$ . An identity  $\mathbf{u} \approx \mathbf{v}$  is *directly deducible* from an identity  $\mathbf{s} \approx \mathbf{t}$  if there exist some words  $\mathbf{a}, \mathbf{b} \in \mathfrak{X}^*$  and substitution  $\phi: \mathfrak{X} \rightarrow \mathfrak{X}^*$  such that  $\{\mathbf{u}, \mathbf{v}\} = \{\mathbf{a}\phi(\mathbf{s})\mathbf{b}, \mathbf{a}\phi(\mathbf{t})\mathbf{b}\}$ . A non-trivial identity  $\mathbf{u} \approx \mathbf{v}$  is *deducible* from a set  $\Sigma$  of identities if there exists some finite sequence  $\mathbf{u} = \mathbf{w}_0, \dots, \mathbf{w}_m = \mathbf{v}$  of words such that each identity  $\mathbf{w}_i \approx \mathbf{w}_{i+1}$  is directly deducible from some identity in  $\Sigma$ .

**Proposition 2.2** (Birkhoff’s Completeness Theorem for Equational Logic; see [1, Theorem 1.4.6]). *A variety  $\text{var } \Sigma$  satisfies an identity  $\mathbf{u} \approx \mathbf{v}$  if and only if  $\mathbf{u} \approx \mathbf{v}$  is deducible from  $\Sigma$ .*  $\square$

**2.3. Rees quotient monoids.** The following construction was introduced by Perkins [23] to build the first example of a finite semigroup generating non-finitely based variety. We use  $W^{\leq}$  to denote the closure of a set of words  $W \subset \mathfrak{X}^*$  under taking factors. For any set  $W$  of words, let  $M(W)$  denote the Rees quotient monoid of  $\mathfrak{X}^*$  over the ideal  $\mathfrak{X}^* \setminus W^{\leq}$  consisting of all words that are not factors of any word in  $W$ .

Given a variety  $\mathbf{V}$ , a word  $\mathbf{u}$  is called an *isoterm for  $\mathbf{V}$*  if the only word  $\mathbf{v}$  such that  $\mathbf{V}$  satisfies the identity  $\mathbf{u} \approx \mathbf{v}$  is the word  $\mathbf{u}$  itself.

**Lemma 2.3** ([14, Lemma 3.3]). *Let  $\mathbf{V}$  be a monoid variety and  $W$  a set of words. Then  $M(W)$  lies in  $\mathbf{V}$  if and only if each word in  $W$  is an isoterm for  $\mathbf{V}$ .*  $\square$

**2.4. Construction of O. Sapir.** In [28, 29], O. Sapir introduced a generalization of  $M(W)$  construction. This development of O. Sapir plays a critical role in the study of limit varieties of monoids [9, 29, 30]. It is also a key tool in the present article.

Let  $\alpha$  be a congruence on the free monoid  $\mathfrak{X}^*$ . The elements of the quotient monoid  $\mathfrak{X}^*/\alpha$  are called  $\alpha$ -classes. The factor relation on  $\mathfrak{X}^*$  can be naturally extended to  $\alpha$ -classes as follows: given two  $\alpha$ -classes  $\mathbf{u}, \mathbf{v} \in \mathfrak{X}^*/\alpha$  we write  $\mathbf{v} \leq_{\alpha} \mathbf{u}$  if  $\mathbf{u} = \mathbf{p}\mathbf{v}\mathbf{s}$  for some  $\mathbf{p}, \mathbf{s} \in \mathfrak{X}^*/\alpha$ . Given a set  $\mathbb{W}$  of  $\alpha$ -classes, we define  $\mathbb{W}^{\leq_{\alpha}}$  as closure of  $\mathbb{W}$  in quasi-order  $\leq_{\alpha}$ . For any set  $\mathbb{W}$  of  $\alpha$ -classes, let  $M_{\alpha}(\mathbb{W})$  denote the Rees quotient of  $\mathfrak{X}^*/\alpha$  over the ideal  $(\mathfrak{X}^*/\alpha) \setminus \mathbb{W}^{\leq_{\alpha}}$ . Evidently, if  $\alpha$  is the trivial congruence on  $\mathfrak{X}^*$ , then  $M_{\alpha}(\mathbb{W})$  is nothing but  $M(\mathbb{W})$ .

In [29], O. Sapir introduced several certain congruences on  $\mathfrak{X}^*$ . One of them, the congruence  $\gamma$ , plays a critical role in the present paper. It is defined as follows: for every  $\mathbf{u}, \mathbf{v} \in \mathfrak{X}^*$ ,  $\mathbf{u} \gamma \mathbf{v}$  if and only if  $\text{sim}(\mathbf{u}) = \text{sim}(\mathbf{v})$  and  $\mathbf{u}$  can be obtained from  $\mathbf{v}$  by changing the individual exponents of letters.

A set  $W$  of words is *stable* with respect to a monoid variety  $\mathbf{V}$  if for each  $\mathbf{u} \in W$ , we have  $\mathbf{v} \in W$  whenever  $\mathbf{V}$  satisfies  $\mathbf{u} \approx \mathbf{v}$ . The following lemma is an analogue of Lemma 2.3 for monoids of the form  $M_{\gamma}(\mathbb{W})$ .

**Lemma 2.4** ([29, Corollary 3.6]). *Let  $\mathbf{V}$  be a monoid variety and  $\mathbb{W}$  a set of  $\gamma$ -classes. Then  $M_{\gamma}(\mathbb{W})$  lies in  $\mathbf{V}$  if and only if each  $\gamma$ -class in  $\mathbb{W}$  is stable with respect to  $\mathbf{V}$ .*  $\square$

Given a set  $\mathbb{W}$  of  $\alpha$ -classes, let  $\mathbf{M}_{\alpha}(\mathbb{W})$  denote the variety generated by the monoid  $M_{\alpha}(\mathbb{W})$ . For brevity, if  $\mathbf{w}_1, \dots, \mathbf{w}_k \in \mathfrak{X}^*/\alpha$ , then we write  $M_{\alpha}(\mathbf{w}_1, \dots, \mathbf{w}_k)$  [respectively,  $\mathbf{M}_{\alpha}(\mathbf{w}_1, \dots, \mathbf{w}_k)$ ] rather than  $M_{\alpha}(\{\mathbf{w}_1, \dots, \mathbf{w}_k\})$  [respectively,  $\mathbf{M}_{\alpha}(\{\mathbf{w}_1, \dots, \mathbf{w}_k\})$ ].

Given a word  $\mathbf{u} \in \mathfrak{X}^*$ , let  $[\mathbf{u}]^{\gamma}$  denote the  $\gamma$ -class containing  $\mathbf{u}$ . We use regular expressions to describe sets of words, in particular,  $\gamma$ -classes. Given a letter  $x \in \mathfrak{X}^*$ , we let  $x^+ := \{x^n \mid n \in \mathbb{N}\}$ . Using this notation, one can represent  $\gamma$ -classes as words in the alphabet  $\{x, x^+ \mid x \in \mathfrak{X}^*\}$ . For example,  $[x^2y^2]^{\gamma} = \{x^n y^k \mid n, k \geq 2\}$ . Using regular expressions, we calculate non-zero elements of  $M_{\gamma}([x^2y^2]^{\gamma})$ :

$$\{[x^2y^2]^{\gamma}\}^{\leq \gamma} = \{xx^+yy^+\}^{\leq \gamma} = \{1, x, y, xx^+, yy^+, xy, xx^+y, xyy^+, xx^+yy^+\}.$$

**2.5. Two known results.** Let  $\mathbf{w}$  be a word with  $\text{sim}(\mathbf{w}) = \{t_1, \dots, t_m\}$ . We may assume without loss of generality that  $\mathbf{w}(t_1, \dots, t_m) = t_1 \cdots t_m$ . Then  $\mathbf{w} = \mathbf{w}_0 t_1 \mathbf{w}_1 \cdots t_m \mathbf{w}_m$  for some words  $\mathbf{w}_0, \dots, \mathbf{w}_m$ . The words  $\mathbf{w}_0, \dots, \mathbf{w}_m$  are called *blocks* of the word  $\mathbf{w}$ . The representation of the word  $\mathbf{w}$  as a product of alternating simple letters and blocks is called *decomposition* of  $\mathbf{w}$ .

The following statement is a combination of [17, Lemma 5.1] and [29, Theorem 4.3(iv)].

**Lemma 2.5.** *Let  $\mathbf{u} \approx \mathbf{v}$  be an identity of  $M_{\gamma}(x^+yzx^+)$ . If  $\mathbf{u}_0 t_1 \mathbf{u}_1 \cdots t_m \mathbf{u}_m$  is the decomposition of the word  $\mathbf{u}$ , then the decomposition of the word  $\mathbf{v}$  has the form  $\mathbf{v}_0 t_1 \mathbf{v}_1 \cdots t_m \mathbf{v}_m$  and  $\text{alph}(\mathbf{u}_i) = \text{alph}(\mathbf{v}_i)$ ,  $i = 1, \dots, m$ .*  $\square$

**Lemma 2.6** ([34, Fig. 1], [29, Fig. 1]). *The lattice  $\mathfrak{L}(\mathbf{M}_\gamma(x^+t yy^+x^+) \vee \mathbf{M}_\gamma(x^+yy^+tx^+))$  is shown in Fig. 1.*  $\square$




FIGURE 1. The lattice  $\mathfrak{L}(\mathbf{M}_\gamma(x^+t yy^+x^+) \vee \mathbf{M}_\gamma(x^+yy^+tx^+))$

### 3. EXCLUSION IDENTITIES

Here we provide exclusion identities which are satisfied by varieties not containing given monoid varieties.

**Lemma 3.1.** *Let  $\mathbf{V}$  be a monoid variety satisfying the identity  $x^n \approx x^{n+1}$  for some  $n \in \mathbb{N}$ . If  $\mathbf{M}_\gamma(xx^+yy^+) \notin \mathbf{V}$ , then  $\mathbf{V}$  satisfies the identity  $x^n y^n \approx (x^n y^n)^n$ .*

*Proof.* In view of Lemma 2.4, the variety  $\mathbf{V}$  satisfies an identity  $x^p y^q \approx \mathbf{w}$ , where  $p, q \geq 2$  and  $\mathbf{w} \notin xx^+yy^+$ . We may assume without any loss that  $\text{alph}(\mathbf{w}) = \{x, y\}$ . Then  $\mathbf{w}$  contains  $yx$  as a factor. We substitute  $x^n$  for  $x$  and  $y^n$  for  $x$  in the identity  $x^p y^q \approx \mathbf{w}$  and then multiply the identity by  $x^n$  on the left and  $y^n$  on the right. We obtain an identity which is equivalent modulo  $x^n \approx x^{n+1}$  to  $x^n y^n \approx (x^n y^n)^m$  for some  $m \geq 2$ . It is clear that the latter identity together with  $x^n \approx x^{n+1}$  imply  $x^n y^n \approx (x^n y^n)^n$ .  $\square$

**Lemma 3.2.** *Let  $\mathbf{V}$  be a monoid variety satisfying the identities*

$$(3.1) \quad xyx \approx xyx^2,$$

$$(3.2) \quad xyx \approx x^2yx,$$

$$(3.3) \quad (xy)^2 \approx (yx)^2.$$

If  $M_\gamma(x^+tyy^+x^+) \notin \mathbf{V}$ , then  $\mathbf{V}$  satisfies the identity

$$(3.4) \quad xy^2x \approx xtxy^2x.$$

*Proof.* A combination of Theorem 4.3(iii) in [29] and the dual to Lemma 4.5 in [9] implies that the variety  $\mathbf{V}$  satisfies the identity

$$(3.5) \quad xty^2x \approx xt(yx)^2.$$

Then, since

$$xy^2x \stackrel{(3.5)}{\approx} xt(yx)^2 \stackrel{(3.3)}{\approx} xt(xy)^2 \stackrel{(3.1)}{\approx} txx(xy)^2 \stackrel{(3.3)}{\approx} txx(yx)^2 \stackrel{(3.5)}{\approx} xtxy^2x,$$

the identity (3.4) holds in  $\mathbf{V}$  as well.  $\square$

For any  $x, y \in \mathfrak{X}$ , we denote by  $\overline{x^+y^+}$  the set of all words in the alphabet  $\{x, y\}$  which are not in  $x^+y^+$ . In other words,  $\overline{x^+y^+} := \{\mathbf{w} \in \mathfrak{X}^* \mid \text{alph}(\mathbf{w}) = \{x, y\}, \mathbf{w} \notin x^+y^+\}$ . For any  $n \in \mathbb{N}$ , let

$$\mathbf{c}_n := xytxy_1^2 \cdots y_n^2y \quad \text{and} \quad \mathbf{c}'_n := yxtxy_1^2 \cdots y_n^2y.$$

For brevity, put

$$\begin{aligned} \mathbf{c}_n &:= [\mathbf{c}_n]^\gamma = x^+y^+tx^+y_1y_1^+ \cdots y_ny_n^+y^+, \\ \mathbf{c}'_n &:= [\mathbf{c}'_n]^\gamma = y^+x^+tx^+y_1y_1^+ \cdots y_ny_n^+y^+. \end{aligned}$$

**Lemma 3.3.** *Let  $\mathbf{V}$  be a variety satisfying the identities (3.1), (3.2) and (3.3). Assume that  $\mathbf{V}$  does not contain the monoids  $M_\gamma(\mathbf{c}_p)$  and  $M_\gamma(\mathbf{c}'_p)$  for all  $p \in \mathbb{N}$ . Then  $\mathbf{V}$  satisfies the identity  $\mathbf{c}_{n,m,k}[\rho] \approx \mathbf{c}'_{n,m,k}[\rho]$  for any  $n, m, k \in \mathbb{N}$  and  $\rho \in S_{n+m+k}$ .*

*Proof.* Let  $r := 2(n+m+k-1) + 1$ . Since

$$\begin{aligned} \mathbf{c}_{n,m,k}[\rho] &\stackrel{\{(3.1), (3.2)\}}{\approx} \left( \prod_{i=1}^n z_i t_i \right) xyt \left( \prod_{i=n+1}^{n+m} z_i t_i \right) x \left( \prod_{i=1}^{n+m+k-1} z_{i\rho}^2 y_i^2 \right) z_{(n+m+k)\rho}^2 y \left( \prod_{i=n+m+1}^{n+m+k} t_i z_i \right) \\ &\stackrel{\mathbf{c}_r \approx \mathbf{c}'_r}{\approx} \left( \prod_{i=1}^n z_i t_i \right) yxt \left( \prod_{i=n+1}^{n+m} z_i t_i \right) x \left( \prod_{i=1}^{n+m+k-1} z_{i\rho}^2 y_i^2 \right) z_{(n+m+k)\rho}^2 y \left( \prod_{i=n+m+1}^{n+m+k} t_i z_i \right) \\ &\stackrel{\{(3.1), (3.2)\}}{\approx} \mathbf{c}'_{n,m,k}[\rho], \end{aligned}$$

it suffices to show that  $\mathbf{c}_r \approx \mathbf{c}'_r$  is satisfied by  $\mathbf{V}$ .

Suppose first that  $M_\gamma(x^+tyy^+x^+) \notin \mathbf{V}$ . Then the identity (3.4) holds in  $\mathbf{V}$  by Lemma 3.2. In this case,  $\mathbf{V}$  satisfies the identity

$$(3.6) \quad xytxy_1^2 \cdots y_r^2y \approx (xy)^2txy_1^2 \cdots y_r^2y$$

because

$$xytxy_1^2 \cdots y_r^2y \stackrel{(3.4)}{\approx} xytxyy_1^2 \cdots y_r^2y \stackrel{(3.2)}{\approx} (xy)^2txyy_1^2 \cdots y_r^2y.$$

By similar arguments, we can show that  $\mathbf{V}$  satisfies

$$(3.7) \quad yxtxy_1^2 \cdots y_r^2y \approx (yx)^2txy_1^2 \cdots y_r^2y.$$

Then the identities

$$(3.8) \quad \mathbf{c}_r \stackrel{(3.6)}{\approx} (xy)^2txy_1^2 \cdots y_r^2y \stackrel{(3.3)}{\approx} (yx)^2txy_1^2 \cdots y_r^2y \stackrel{(3.7)}{\approx} \mathbf{c}'_r$$

are satisfied by  $\mathbf{V}$  and we are done.

Suppose now that  $M_\gamma(x^+yy^+tx^+) \notin \mathbf{V}$ . Then, by the dual to Lemma 3.2, the variety  $\mathbf{V}$  satisfies the identity  $xy^2tx \approx xy^2txx$ , which is evidently implies both the identities (3.6) and (3.7), and so  $\mathbf{V}$  satisfies (3.8).

Finally, suppose that  $\mathbf{M}_\gamma(x^+yy^+tx^+) \vee \mathbf{M}_\gamma(x^+tyy^+x^+) \subseteq \mathbf{V}$ . By the condition of the lemma,  $M_\gamma(c_r) \notin \mathbf{V}$ . According to Lemma 2.4, the  $\gamma$ -class  $c_r$  is not stable with respect to  $\mathbf{V}$ . This implies that  $\mathbf{V}$  satisfies an identity  $\mathbf{u} \approx \mathbf{v}$  such that  $\mathbf{u} \in c_r$  and  $\mathbf{v} \notin c_r$ . Then, since  $M_\gamma(x^+tyy^+x^+) \in \mathbf{V}$ , Lemmas 2.4 and 2.6 imply that the  $\gamma$ -classes  $x^+tyy^+x^+$ ,  $xx^+yy^+$  and  $x^+yzx^+$  are stable with respect to  $\mathbf{V}$ . Hence

$$\mathbf{v}(x, t) \in x^+tx^+, \mathbf{v}(x, y_1) \in xx^+y_1y_1^+, \mathbf{v}(y_i, y_{i+1}) \in y_iy_i^+y_{i+1}y_{i+1}^+, \mathbf{v}(y, t, y_r) \in y^+tyry_r^+y^+,$$

$i = 1, \dots, r-1$ . It follows that  $\mathbf{v} \in \overline{x^+y^+tx^+y_1y_1^+ \cdots y_r y_r^+ y^+}$ . Then, substituting  $yt$  for  $t$  in the identity  $x\mathbf{u} \approx x\mathbf{v}$ , we result some identity which is equivalent modulo  $\{(3.1), (3.2)\}$  to (3.6). By similar arguments, we can deduce from the condition  $M_\gamma(c'_r) \notin \mathbf{V}$  that  $\mathbf{V}$  satisfies (3.7). Then the identities (3.8) are satisfied by  $\mathbf{V}$  and we are done.  $\square$

#### 4. CERTAIN VARIETIES WITH A NON-DISTRIBUTIVE SUBVARIETY LATTICE

**Proposition 4.1.** *The lattice  $\mathfrak{L}(\mathbf{M}(xyx) \vee \mathbf{M}_\gamma(xx^+yy^+))$  is not modular.*

*Proof.* In view of [27, Fact 3.1(i)], the words  $xzxyty$  and  $xzyxty$  can only form an identity of  $M(xyx)$  with each other. It is easy to see that  $\mathbf{M}_\gamma(xx^+yy^+)$  violates  $\sigma_3$ . Hence  $xzxyty$  is an isoterm for  $\mathbf{M}(xyx) \vee \mathbf{M}_\gamma(xx^+yy^+)$ . Now Lemma 2.3 applies, yielding that

$$(\mathbf{M}(xyx) \vee \mathbf{M}_\gamma(xx^+yy^+)) \wedge \mathbf{M}(xzxyty) = \mathbf{M}(xzxyty).$$

Further, in view of [14, Lemmas 4.4 and 5.10] and Lemma 2.6,

$$\mathbf{M}(xyx) \vee (\mathbf{M}_\gamma(xx^+yy^+) \wedge \mathbf{M}(xzxyty)) = \mathbf{M}(xyx) \vee \mathbf{M}(xy) = \mathbf{M}(xyx) \subset \mathbf{M}(xzxyty).$$

Therefore, the lattice  $\mathfrak{L}(\mathbf{M}(xyx) \vee \mathbf{M}_\gamma(xx^+yy^+))$  is not modular.  $\square$

**Proposition 4.2.** *The lattices  $\mathfrak{L}(\mathbf{M}_\gamma(c_n))$  and  $\mathfrak{L}(\mathbf{M}_\gamma(c'_n))$  are not modular for any  $n \in \mathbb{N}$ .*

*Proof.* It is easy to see that  $\mathbf{M}_\gamma(c_{k+1}) \subseteq \mathbf{M}_\gamma(c_k)$  for any  $k \in \mathbb{N}$ . This allows us to assume that  $n \geq 3$ . For any  $\xi, \eta \in S_2$ , we define the word:

$$\mathbf{v}_{\xi, \eta} := a_1 b_1^2 a_2 b_2^2 x_{1\xi} x_{2\xi} b^2 y_{1\eta} y_{2\eta} b_3^2 a_3 b_4^2 a_4 \mathbf{q},$$

where

$$\mathbf{q} := tx_1 c_1^2 a_1 \left( \prod_{i=2}^{n-1} c_i^2 \right) a_3 c_n^2 x_2 c^2 y_1 d_1^2 a_2 \left( \prod_{i=2}^{n-1} d_i^2 \right) a_4 d_n^2 y_2.$$

Let  $\varepsilon$  [respectively,  $v$ ] denote the identity [respectively, unique non-identity] element in  $S_2$ . Put

$$\mathbf{v}_{\varepsilon, \varepsilon} := a_1^+ b_1 b_1^+ a_2^+ b_2 b_2^+ x_1^+ x_2^+ b b^+ y_1^+ y_2^+ b_3 b_3^+ a_3^+ b_4 b_4^+ a_4^+ \mathbf{q},$$

$$\mathbf{v}_{v, \varepsilon} := a_1^+ b_1 b_1^+ a_2^+ b_2 b_2^+ \overline{x_1^+ x_2^+} b b^+ y_1^+ y_2^+ b_3 b_3^+ a_3^+ b_4 b_4^+ a_4^+ \mathbf{q},$$

$$\mathbf{v}_{\varepsilon, v} := a_1^+ b_1 b_1^+ a_2^+ b_2 b_2^+ x_1^+ x_2^+ b b^+ \overline{y_1^+ y_2^+} b_3 b_3^+ a_3^+ b_4 b_4^+ a_4^+ \mathbf{q},$$

$$\mathbf{v}_{v, v} := a_1^+ b_1 b_1^+ a_2^+ b_2 b_2^+ \overline{x_1^+ x_2^+} b b^+ \overline{y_1^+ y_2^+} b_3 b_3^+ a_3^+ b_4 b_4^+ a_4^+ \mathbf{q},$$

where

$$\mathbf{q} := tx_1^+ c_1 c_1^+ a_1^+ \left( \prod_{i=2}^{n-1} c_i c_i^+ \right) a_3^+ c_n c_n^+ x_2^+ c c^+ y_1^+ d_1 d_1^+ a_2^+ \left( \prod_{i=2}^{n-1} d_i d_i^+ \right) a_4^+ d_n d_n^+ y_2^+.$$

Notice that  $\mathbf{v}_{\xi, \eta} \in \mathbf{v}_{\xi, \eta}$  for any  $\xi, \eta \in S_2$ . We need the following two auxiliary facts.

**Lemma 4.3.** *Let  $\mathbf{u} \approx \mathbf{v}$  be an identity of  $\mathbf{M}_\gamma(c_n) \wedge \text{var}\{\mathbf{v}_{\varepsilon, \varepsilon} \approx \mathbf{v}_{\varepsilon, v} \approx \mathbf{v}_{v, \varepsilon} \approx \mathbf{v}_{v, v}\}$  with  $\mathbf{u} \in \mathbf{v}_{\varepsilon, \varepsilon} \cup \mathbf{v}_{v, \varepsilon} \cup \mathbf{v}_{\varepsilon, v} \cup \mathbf{v}_{v, v}$ . Then  $\mathbf{v} \in \mathbf{v}_{\varepsilon, \varepsilon} \cup \mathbf{v}_{v, \varepsilon} \cup \mathbf{v}_{\varepsilon, v} \cup \mathbf{v}_{v, v}$ .*



*Proof.* The evident inclusion

$$\mathbf{M}_\gamma(x^+yy^+tx^+) \vee \mathbf{M}_\gamma(x^+tyy^+x^+) \subseteq \mathbf{M}_\gamma(\mathbf{c}_n) \wedge \text{var}\{\mathbf{v}_{\varepsilon,\varepsilon} \approx \mathbf{v}_{\varepsilon,v} \approx \mathbf{v}_{v,\varepsilon} \approx \mathbf{v}_{v,v}\}$$

implies that  $\mathbf{u} \approx \mathbf{v}$  is also satisfied by the variety  $\mathbf{M}_\gamma(x^+yy^+tx^+) \vee \mathbf{M}_\gamma(x^+tyy^+x^+)$ . It follows from Lemmas 2.4 and 2.6 that the  $\gamma$ -classes  $x^+tyy^+x^+$ ,  $x^+yy^+tx^+$  and  $xx^+yy^+$  are stable with respect to  $\mathbf{M}_\gamma(x^+yy^+tx^+) \vee \mathbf{M}_\gamma(x^+tyy^+x^+)$ . It is routinely deduced from this fact that  $\mathbf{v} \in \mathbf{v}_{\varepsilon,\varepsilon} \cup \mathbf{v}_{v,\varepsilon} \cup \mathbf{v}_{\varepsilon,v} \cup \mathbf{v}_{v,v}$ . For instance,  $\mathbf{v}(a_1, b_1, t) \in a_1^+b_1^+ta_1^+$  because  $\mathbf{u}(a_1, b_1, t) \in a_1^+b_1^+ta_1^+$ , while  $\mathbf{v}(b_1, a_2) \in b_1^+a_2^+$  because  $\mathbf{u}(b_1, a_2) \in b_1^+a_2^+$ .  $\square$

**Lemma 4.4.** *Let  $\xi, \eta, \xi_1, \eta_1, \xi_2, \eta_2 \in S_2$ ,  $\mathbf{u} \in \mathbf{v}_{\xi,\eta}$  and  $\mathbf{v} \notin \mathbf{v}_{\xi,\eta}$ . If the identity  $\mathbf{u} \approx \mathbf{v}$  is directly deducible from the identity  $\mathbf{v}_{\xi_1,\eta_1} \approx \mathbf{v}_{\xi_2,\eta_2}$ , then either  $\mathbf{u} \in \mathbf{v}_{\xi_1,\eta_1}$  and  $\mathbf{v} \in \mathbf{v}_{\xi_2,\eta_2}$  or  $\mathbf{v} \in \mathbf{v}_{\xi_1,\eta_1}$  and  $\mathbf{u} \in \mathbf{v}_{\xi_2,\eta_2}$ .*

*Proof.* We may assume without loss of generality that  $(\mathbf{u}, \mathbf{v}) = (\mathbf{a}\phi(\mathbf{v}_{\xi_1,\eta_1})\mathbf{b}, \mathbf{a}\phi(\mathbf{v}_{\xi_2,\eta_2})\mathbf{b})$  for some  $\mathbf{a}, \mathbf{b} \in \mathcal{X}^*$  and  $\phi: \mathcal{X} \rightarrow \mathcal{X}^*$ . Now Lemma 4.3 implies that  $\mathbf{v} \in \mathbf{v}_{\xi',\eta'}$  for some  $\xi', \eta' \in S_2$ . Since  $\mathbf{u} \in \mathbf{v}_{\xi,\eta}$  and  $\mathbf{v} \notin \mathbf{v}_{\xi,\eta}$ , we have  $(\xi, \eta) \neq (\xi', \eta')$ . By symmetry, we may assume that  $\xi \neq \xi'$  and, moreover,  $(\xi, \xi') = (\varepsilon, v)$ . Then  $\mathbf{u}(x_1, x_2, t) \in x_1^+x_2^+tx_1^+x_2^+$  and  $\mathbf{v}(x_1, x_2, t) \in x_1^+x_2^+tx_1^+x_2^+$ . This is only possible when the image of  $t$  under  $\phi$  contains  $t$ , and one of the following holds:

- (a)  $\xi_1 = \xi = \varepsilon$ ,  $\xi_2 = \xi' = v$  and  $\phi(x_i) \in x_i^+$ ,  $i = 1, 2$ ;
- (b)  $\eta_1 = \xi = \varepsilon$ ,  $\eta_2 = \xi' = v$  and  $\phi(y_i) \in x_i^+$ ,  $i = 1, 2$ .

We notice also that if  $x$  and  $y$  are distinct letters and  $xy$  occurs in  $\mathbf{u}$  as a factor at least twice, then  $\{x, y\} = \{y_1, y_2\}$  and all these factors are between  $b$  and  $b_3$ . Since  $t \in \text{alph}(\phi(t))$ , this fact implies that

(\*)  $\phi(v)$  is either the empty word or a power of letter for any  $v \in \text{mul}(\mathbf{v}_{\xi_1,\eta_1})$ .

Suppose that (a) holds. Then

$$\phi(c_1^2a_1c_2^2 \cdots c_{n-1}^2a_{n-1}c_n^2) \in c_1c_1^+a_1^+c_2c_2^+ \cdots c_{n-1}c_{n-1}^+a_{n-1}^+c_nc_n^+.$$

It follows from (\*) that  $\phi(a_1) \in a_1^+$ ,  $\phi(a_3) \in a_3^+$  and  $\phi(c_i^2) \in c_ic_i^+$  for any  $i = 1, \dots, n$ . Then

$$\phi(a_1b_1^2a_2b_2^2x_1x_2b^2y_1\eta_1y_2\eta_1b_3^2a_3) \in a_1^+b_1b_1^+a_2^+b_2b_2^+x_1^+x_2^+bb^+y_\eta b_3b_3^+a_3^+,$$

where

$$(4.1) \quad y_\eta := \begin{cases} y_1^+y_2^+ & \text{if } \eta = \varepsilon, \\ y_1^+y_2^+ & \text{if } \eta = v. \end{cases}$$

Now we apply (\*) again, yielding that  $y_\eta = y_{1\eta}^+y_{2\eta}^+$  and

$$\begin{aligned} \phi(b_1) &\in b_1b_1^+, \phi(a_2) \in a_2^+, \phi(b_2) \in b_2b_2^+, \phi(b) \in bb^+, \\ \phi(y_{1\eta_1}) &= y_{1\eta}^+, \phi(y_{2\eta_1}) \in y_{2\eta}^+, \phi(b_3) \in b_3b_3^+. \end{aligned}$$

Finally, since  $\mathbf{v}_{\xi_1,\eta_1}(y_1, y_2, t) \in y_{1\eta_1}^+y_{2\eta_1}^+ty_{1\eta_1}^+y_{2\eta_1}^+$ , we have  $\mathbf{u}(y_1, y_2, t) \in y_{1\eta_1}^+y_{2\eta_1}^+ty_{1\eta_1}^+y_{2\eta_1}^+$ . But  $\mathbf{u}(y_1, y_2, t) \in y_{1\eta}^+y_{2\eta}^+ty_{1\eta}^+y_{2\eta}^+$ . This implies that  $\eta_1 = \eta$ . Thus,  $\mathbf{u} \in \mathbf{v}_{\xi_1,\eta_1}$  and  $\mathbf{v} \in \mathbf{v}_{\xi_2,\eta_2}$ .

Suppose that (b) holds. Then

$$\phi(d_1^2a_2d_2^2 \cdots d_{n-1}^2a_nd_n^2) \in c_1c_1^+a_1^+c_2c_2^+ \cdots c_{n-1}c_{n-1}^+a_n^+c_nc_n^+.$$

It follows from (\*) that  $\phi(a_2) \in a_1^+$ ,  $\phi(a_4) \in a_3^+$  and  $\phi(d_i^2) \in c_ic_i^+$  for any  $i = 1, \dots, n$ . Then

$$\phi(a_2b_2^2x_1\xi_1x_2\xi_1b^2y_1y_2b_3^2a_3b_4^2a_4) \in a_1^+b_1b_1^+a_2^+b_2b_2^+x_1^+x_2^+bb^+y_\eta b_3b_3^+a_4^+,$$

where  $y_\eta$  is defined by the formula (4.1). Now we apply (\*) again and obtain that  $\phi(y_2) \in bb^+$  contradicting (b). Therefore, (b) is impossible.  $\square$

One can return to the proof of Proposition 4.2. Let

$$\begin{aligned}\mathbf{X} &:= \mathbf{M}_\gamma(c_n) \wedge \text{var}\{\mathbf{v}_{\varepsilon,\varepsilon} \approx \mathbf{v}_{v,\varepsilon}, \mathbf{v}_{\varepsilon,v} \approx \mathbf{v}_{v,v}\}, \\ \mathbf{Y} &:= \mathbf{M}_\gamma(c_n) \wedge \text{var}\{\mathbf{v}_{v,\varepsilon} \approx \mathbf{v}_{v,v}\}, \\ \mathbf{Z} &:= \mathbf{M}_\gamma(c_n) \wedge \text{var}\{\mathbf{v}_{\varepsilon,\varepsilon} \approx \mathbf{v}_{\varepsilon,v}, \mathbf{v}_{v,\varepsilon} \approx \mathbf{v}_{v,v}\}.\end{aligned}$$

Consider an identity  $\mathbf{u} \approx \mathbf{u}'$  of  $\mathbf{X}$  with  $\mathbf{u} \in \mathbf{v}_{\varepsilon,\varepsilon} \cup \mathbf{v}_{v,\varepsilon}$ . We are going to show that  $\mathbf{u}' \in \mathbf{v}_{\varepsilon,\varepsilon} \cup \mathbf{v}_{v,\varepsilon}$ . In view of Proposition 2.2, we may assume without loss of generality that either  $\mathbf{u} \approx \mathbf{u}'$  holds in  $\mathbf{M}_\gamma(c_n)$  or  $\mathbf{u} \approx \mathbf{u}'$  is directly deducible from  $\mathbf{v}_{\varepsilon,\varepsilon} \approx \mathbf{v}_{v,\varepsilon}$  or  $\mathbf{v}_{\varepsilon,v} \approx \mathbf{v}_{v,v}$ . According to Lemma 4.4,  $\mathbf{u} \approx \mathbf{u}'$  cannot be directly deducible from  $\mathbf{v}_{\varepsilon,v} \approx \mathbf{v}_{v,v}$  and if  $\mathbf{u} \approx \mathbf{u}'$  is directly deducible from  $\mathbf{v}_{\varepsilon,\varepsilon} \approx \mathbf{v}_{v,\varepsilon}$ , then  $\mathbf{u}' \in \mathbf{v}_{\varepsilon,\varepsilon} \cup \mathbf{v}_{v,\varepsilon}$ . Therefore, it remains to consider the case when  $\mathbf{u} \approx \mathbf{u}'$  is satisfied by  $\mathbf{M}_\gamma(c_n)$ . In this case, it follows from Lemma 4.3 that  $\mathbf{u}' \in \mathbf{v}_{\varepsilon,\varepsilon} \cup \mathbf{v}_{v,\varepsilon} \cup \mathbf{v}_{\varepsilon,v} \cup \mathbf{v}_{v,v}$ . Further, it is routine to check that  $\mathbf{M}_\gamma(c_p)$  satisfies the identities

$$(4.2) \quad (xy)^2 \approx xy^2x \approx (yx)^2 \approx yx^2y$$

and (3.7) for any  $r \in \mathbb{N}$ . If  $\mathbf{u}' \in \mathbf{v}_{\varepsilon,v} \cup \mathbf{v}_{v,v}$ , then the identity

$$\mathbf{u}(y_1, y_2, t, d_1, \dots, d_n) \approx \mathbf{u}'(y_1, y_2, t, d_1, \dots, d_n)$$

is equivalent modulo (3.7) and (4.2) to the identity  $\mathbf{c}_n \approx \mathbf{c}'_n$ . But this is impossible because  $\mathbf{M}_\gamma(c_p)$  violates the latter identity by Lemma 2.4. We see that  $\mathbf{u}' \in \mathbf{v}_{\varepsilon,\varepsilon} \cup \mathbf{v}_{v,\varepsilon}$  in either case and, therefore, the set  $\mathbf{v}_{\varepsilon,\varepsilon} \cup \mathbf{v}_{v,\varepsilon}$  is stable with respect to  $\mathbf{X}$ . By similar arguments, we can show that the sets  $\mathbf{v}_{\varepsilon,\varepsilon}$  and  $\mathbf{v}_{\varepsilon,\varepsilon} \cup \mathbf{v}_{\varepsilon,v}$  are stable with respect to  $\mathbf{Y}$  and  $\mathbf{Z}$ , respectively. This implies that the  $\gamma$ -class  $\mathbf{v}_{\varepsilon,\varepsilon}$  is stable with respect to  $(\mathbf{X} \vee \mathbf{Z}) \wedge \mathbf{Y}$ . Clearly, both  $\mathbf{X} \wedge \mathbf{Y}$  and  $\mathbf{Z}$  satisfy the identity  $\mathbf{v}_{\varepsilon,\varepsilon} \approx \mathbf{v}_{\varepsilon,v}$ . Therefore, the  $\gamma$ -class  $\mathbf{v}_{\varepsilon,\varepsilon}$  is not stable with respect to  $(\mathbf{X} \wedge \mathbf{Y}) \vee \mathbf{Z}$ . Since  $\mathbf{Z} \subseteq \mathbf{Y}$ , we have

$$(\mathbf{X} \wedge \mathbf{Y}) \vee \mathbf{Z} \subset (\mathbf{X} \vee \mathbf{Z}) \wedge \mathbf{Y}.$$

It follows that the lattice  $\mathfrak{L}(\mathbf{M}_\gamma(c_n))$  is not modular. A similar argument works for the case of  $\mathfrak{L}(\mathbf{M}_\gamma(c'_n))$ .  $\square$

## 5. PROOF OF THEOREM 1.1

**Necessity.** Let  $\mathbf{V}$  be a distributive variety of aperiodic monoids. Then  $\mathbf{V}$  satisfies the identity  $x^n \approx x^{n+1}$  for some  $n \in \mathbb{N}$ . If  $n = 1$ , then  $\mathbf{V} \subseteq \mathbf{B}$  and we are done. Let now  $n > 1$  and  $\mathbf{V}$  is not a variety of bands. Then  $M(x) \in \mathbf{V}$  by [10, Lemma 2.5]. According to [16, Proposition 4.1], the lattice  $\mathfrak{L}(\mathbf{M}(x) \vee \mathbf{LRB})$  [respectively,  $\mathfrak{L}(\mathbf{M}(x) \vee \mathbf{RRB})$ ] is not modular, where  $\mathbf{LRB}$  [respectively,  $\mathbf{RRB}$ ] is the variety generated by the 2-element left-zero [respectively, right-zero] semigroup with a new identity element adjoined. Hence  $\mathbf{LRB}, \mathbf{RRB} \not\subseteq \mathbf{V}$ . It follows from Proposition 10.10.2c) in [1], its proof and the dual statement that  $\mathbf{V}$  satisfies the identities  $(x^n y^n)^n \approx (x^n y^n)^n x^n \approx (y^n x^n)^n$ . Further, if  $M_\gamma(xx^+yy^+) \notin \mathbf{V}$ , then, by Lemma 3.1, the identity  $x^n y^n \approx (x^n y^n)^n$  and so the identity  $x^n y^n \approx y^n x^n$  hold in  $\mathbf{V}$ . In this case,  $\mathbf{V}$  is a variety of aperiodic monoids with commuting idempotents and, therefore, we can apply Proposition 2.1. So, we may further assume that  $M_\gamma(xx^+yy^+) \in \mathbf{V}$ .

In view of [5, Lemma 2] and [29, Theorem 4.1(i)], the lattice  $\mathfrak{L}(\mathbf{M}(x^2) \vee \mathbf{M}_\gamma(yxx^+))$  is not modular. Since the monoid  $M_\gamma(yxx^+)$  lies in the variety  $\mathbf{M}_\gamma(xx^+yy^+)$  by Lemma 2.6, this implies that  $M(x^2) \notin \mathbf{V}$ . Using Lemma 2.3, one can easily deduce from this that  $\mathbf{V}$  satisfies  $x^2 \approx x^3$ . According to Proposition 4.1,  $M(xyxx) \notin \mathbf{V}$ . Hence the identity (3.1) holds in  $\mathbf{V}$  by [7, Corollary 6.8(i)]. Combining Lemma 4.4 in [7], Proposition 6.5 in [9] and the dual statements, we yield that the identities  $xyx^2 \approx x^2yx^2$ ,  $x^2yx \approx x^2yx^2$  are satisfied by  $\mathbf{V}$ .

The latter three identities together with  $(x^n y^n)^n \approx (y^n x^n)^n$  imply  $\{(3.2), (3.3)\}$ . Now we apply Lemma 3.3, Proposition 4.2 and the dual statements, yielding that  $\mathbf{V} \subseteq \mathbf{D}_{15}$ .

**Sufficiency.** By symmetry, it suffices to show that the varieties  $\mathbf{B}$ ,  $\mathbf{D}_1$ – $\mathbf{D}_{15}$ ,  $\mathbf{P}_n$ ,  $\mathbf{Q}_n$  and  $\mathbf{R}_n$  are distributive. The variety  $\mathbf{B}$  is distributive by [33, Proposition 4.7]. In view of Proposition 2.1, the varieties  $\mathbf{D}_1$ – $\mathbf{D}_{14}$ ,  $\mathbf{P}_n$ ,  $\mathbf{Q}_n$  and  $\mathbf{R}_n$  also have this property. So, it remains to prove the distributivity of  $\mathbf{D}_{15}$ .

In view of Lemma 3.1, a subvariety  $\mathbf{X}$  of  $\mathbf{D}_{15}$  satisfies  $x^2 y^2 \approx (x^2 y^2)^2$  and so  $x^2 y^2 \approx (x^2 y^2)^2 \approx (y^2 x^2)^2 \approx y^2 x^2$  whenever  $M(xx^+ yy^+) \notin \mathbf{X}$ . In this case,  $\mathbf{X} \subseteq \mathbf{M}_\gamma(x^+ y z x^+)$  by [29, Theorem 4.3(i)] and [19, Proposition 4.3]. Further, it follows from [7, Lemma 4.8] that a subvariety  $\mathbf{X}$  of  $\mathbf{D}_{15}$  satisfies the identity  $x y z x \approx x y x z x$  whenever  $\mathbf{M}_\gamma(x^+ y z x^+) \notin \mathbf{X}$  and  $M_\gamma(xx^+ yy^+) \in \mathbf{X}$ . In this case,  $\mathbf{X} = \mathbf{M}_\gamma(xx^+ yy^+)$  by [29, Theorem 4.1(iv)] and [3, Proposition 3.2a)]. We see that the lattice  $\mathfrak{L}(\mathbf{D}_{15})$  is a union of the lattice  $\mathfrak{L}(\mathbf{M}_\gamma(x^+ y z x^+) \vee \mathbf{M}_\gamma(xx^+ yy^+))$  and the interval  $[\mathbf{M}_\gamma(x^+ y z x^+) \vee \mathbf{M}_\gamma(xx^+ yy^+), \mathbf{V}]$ .

For brevity, we will denote by  $\mathbf{V}\Sigma$  the subvariety of a variety  $\mathbf{V}$  defined by a set  $\Sigma$  of identities. According to Lemma 2.6, it remains to verify that the interval  $[\mathbf{M}_\gamma(x^+ y z x^+) \vee \mathbf{M}_\gamma(xx^+ yy^+), \mathbf{D}_{15}]$  is distributive. By [11, Lemma 2.19], it suffices to find a set  $\Gamma$  of identities such that:

- (i) if  $\mathbf{X} \in [\mathbf{M}_\gamma(x^+ y z x^+) \vee \mathbf{M}_\gamma(xx^+ yy^+), \mathbf{D}_{15}]$ , then  $\mathbf{X} = \mathbf{D}_{15}\Gamma'$  for some subset  $\Gamma'$  of  $\Gamma$ ;
- (ii) if  $\mathbf{X}, \mathbf{Y} \in [\mathbf{M}_\gamma(x^+ y z x^+) \vee \mathbf{M}_\gamma(xx^+ yy^+), \mathbf{D}_{15}]$  and  $\mathbf{X} \wedge \mathbf{Y}$  satisfies an identity  $\sigma \in \Gamma$ , then  $\sigma$  holds in either  $\mathbf{X}$  or  $\mathbf{Y}$ .

We will show in Proposition 5.1 and Lemma 5.4 below that the set  $\Gamma$  consisting of all identities of the form

$$(5.1) \quad \left( \prod_{i=1}^k a_i t_i \right) x^2 y^2 \left( \prod_{i=k+1}^{k+\ell} t_i a_i \right) \approx \left( \prod_{i=1}^k a_i t_i \right) (xy)^2 \left( \prod_{i=k+1}^{k+\ell} t_i a_i \right),$$

where  $k, \ell \geq 0$  and  $a_1, \dots, a_{k+\ell} \in \{1, x, y\}$ , is relevant. Namely, the claim (i) is true by Proposition 5.1, while the claim (ii) holds by Lemma 5.4. Theorem 1.1 is thus proved.  $\square$

**Proposition 5.1.** *Each variety in the interval  $[\mathbf{M}_\gamma(x^+ y z x^+) \vee \mathbf{M}_\gamma(xx^+ yy^+), \mathbf{D}_{15}]$  can be defined within  $\mathbf{D}_{15}$  by the identities of the form (5.1), where  $k, \ell \geq 0$  and  $a_1, \dots, a_{k+\ell} \in \{1, x, y\}$ .*

To prove Proposition 5.1, we need two auxiliary results.

**Lemma 5.2.** *If  $\mathbf{w} := \mathbf{p}xy\mathbf{q}x\mathbf{r}y\mathbf{s}$  and  $\text{alph}(\mathbf{r}) \subseteq \text{mul}(\mathbf{w})$ , then  $\mathbf{D}_{15}$  satisfies the identity  $\mathbf{w} \approx \mathbf{p}y\mathbf{x}q\mathbf{x}r\mathbf{y}\mathbf{s}$ .*

*Proof.* The variety  $\mathbf{D}_{15}$  satisfies the identities

$$\mathbf{w} \stackrel{\{(3.1), (3.2)\}}{\approx} \mathbf{p}xy\mathbf{q}x\mathbf{r}'y\mathbf{s} \stackrel{\Phi_1}{\approx} \mathbf{p}y\mathbf{x}q\mathbf{x}r'y\mathbf{s} \stackrel{\{(3.1), (3.2)\}}{\approx} \mathbf{p}y\mathbf{x}q\mathbf{x}r'y\mathbf{s},$$

where  $\mathbf{r}'$  is the word obtained from  $\mathbf{r}$  by replacing each letter to its square.  $\square$

**Lemma 5.3.** *If  $\mathbf{w} := \mathbf{p}x\mathbf{q}_1\mathbf{q}_2\mathbf{x}r$  and  $\text{alph}(\mathbf{q}_1\mathbf{q}_2) \subseteq \text{mul}(\mathbf{w})$ , then  $\mathbf{D}_{15}$  satisfies the identity  $\mathbf{w} \approx \mathbf{p}x\mathbf{q}_1x\mathbf{q}_2\mathbf{x}r$ .*

*Proof.* In view of Lemma 5.2 and the dual statement, the variety  $\mathbf{D}_{15}$  satisfies the identities

$$\mathbf{w} \stackrel{\{(3.1), (3.2)\}}{\approx} \mathbf{p}x^2\mathbf{q}'_1\mathbf{q}_2\mathbf{x}r \stackrel{\Phi_1}{\approx} \mathbf{p}x\mathbf{q}'_1x\mathbf{q}_2\mathbf{x}r \stackrel{\{(3.1), (3.2)\}}{\approx} \mathbf{p}x\mathbf{q}_1x\mathbf{q}_2\mathbf{x}r,$$

where  $\mathbf{q}'_1$  is the word obtained from  $\mathbf{q}_1$  by replacing each letter to its square.  $\square$

*Proof of Proposition 5.1.* Recall that  $\Gamma$  denotes the set of all identities of the form (5.1), where  $k, \ell \geq 0$  and  $a_1, \dots, a_{k+\ell} \in \{1, x, y\}$ . Let  $\mathbf{V} \in [\mathbf{M}_\Gamma(x^+yzx^+) \vee \mathbf{M}_\Gamma(xx^+yy^+), \mathbf{D}_{15}]$ . Take an arbitrary identity  $\mathbf{u} \approx \mathbf{u}'$  of  $\mathbf{V}$ . We need to verify that  $\mathbf{u} \approx \mathbf{u}'$  is equivalent within  $\mathbf{D}_{15}$  to some subset of  $\Gamma$ .

Let  $\mathbf{u}_0 t_1 \mathbf{u}_1 \cdots t_m \mathbf{u}_m$  be the decomposition of  $\mathbf{u}$ . Lemma 2.5 implies that the decomposition of  $\mathbf{u}'$  has the form  $\mathbf{u}'_0 t_1 \mathbf{u}'_1 \cdots t_m \mathbf{u}'_m$  and  $\text{alph}(\mathbf{u}_i) = \text{alph}(\mathbf{u}'_i)$ ,  $i = 0, \dots, m$ . Taking into account Lemma 5.3, we can use the identities of  $\mathbf{D}_{15}$  to convert the words  $\mathbf{u}$  and  $\mathbf{u}'$  into some words  $\mathbf{w}$  and  $\mathbf{w}'$ , respectively, such that the following hold:

- the decompositions of the words  $\mathbf{w}$  and  $\mathbf{w}'$  are of the form  $\mathbf{w}_0 t_1 \mathbf{w}_1 \cdots t_m \mathbf{w}_m$  and  $\mathbf{w}'_0 t_1 \mathbf{w}'_1 \cdots t_m \mathbf{w}'_m$ , respectively;
- $\text{occ}_x(\mathbf{w}_i) = \text{occ}_x(\mathbf{w}'_i) = 2$  for any  $x \in \text{alph}(\mathbf{w}_i \mathbf{w}'_i)$  and  $i = 0, \dots, m$ .

Thus, it suffices to show that the identity  $\mathbf{w} \approx \mathbf{w}'$  is equivalent within  $\mathbf{D}_{15}$  to some subset of  $\Gamma$ .

We call an identity  $\mathbf{c} \approx \mathbf{d}$  *1-invertible* if  $\mathbf{c} = \mathbf{e}' x y \mathbf{e}''$  and  $\mathbf{d} = \mathbf{e}' y x \mathbf{e}''$  for some words  $\mathbf{e}', \mathbf{e}'' \in \mathfrak{X}^*$  and letters  $x, y \in \text{alph}(\mathbf{e}' \mathbf{e}'')$ . Let  $j > 1$ . An identity  $\mathbf{c} \approx \mathbf{d}$  is called *j-invertible* if there is a sequence of words  $\mathbf{c} = \mathbf{w}_0, \dots, \mathbf{w}_j = \mathbf{d}$  such that the identity  $\mathbf{w}_i \approx \mathbf{w}_{i+1}$  is 1-invertible for each  $i = 0, \dots, j-1$  and  $j$  is the least number with such a property. For convenience, we will call the trivial identity *0-invertible*.

Notice that the identity  $\mathbf{w} \approx \mathbf{w}'$  is *r-invertible* for some  $r \geq 0$  because  $\text{occ}_x(\mathbf{w}_i) = \text{occ}_x(\mathbf{w}'_i)$  for any  $x \in \mathfrak{X}$  and  $i = 0, \dots, m$ . We will use induction by  $r$ .

**Induction base.** If  $r = 0$ , then  $\mathbf{w} = \mathbf{w}'$ , whence  $\mathbf{D}_{15}\{\mathbf{w} \approx \mathbf{w}'\} = \mathbf{D}_{15}\{\emptyset\}$ .

**Induction step.** Let  $r > 0$ . Obviously,  $\mathbf{w}_s \neq \mathbf{w}'_s$  for some  $s \in \{0, \dots, m\}$ . Then there are letters  $x$  and  $y$  such that, for some  $p, q \in \{1, 2\}$ ,

- the  $p$ th occurrence of  $x$  precedes the  $q$ th occurrence of  $y$  in  $\mathbf{w}_s$ ;
- there are no other letters between these two occurrences of  $x$  and  $y$  in  $\mathbf{w}_s$ ;
- the  $q$ th occurrence of  $y$  precedes the  $p$ th occurrence of  $x$  in  $\mathbf{w}'_s$ .

In this case,  $\mathbf{w}_s = \mathbf{a}_s x y \mathbf{b}_s$  for some  $\mathbf{a}_s, \mathbf{b}_s \in \mathfrak{X}^*$ . We denote by  $\hat{\mathbf{w}}$  the word obtained from  $\mathbf{w}$  by swapping of the  $p$ th occurrence of  $x$  and the  $q$ th occurrence of  $y$  in the block  $\mathbf{w}_s$ . In other words,  $\hat{\mathbf{w}} := \mathbf{a}_s y x \mathbf{b}_s$ , where

$$\mathbf{a} := \left( \prod_{i=1}^s \mathbf{w}_{i-1} t_i \right) \mathbf{a}_s \quad \text{and} \quad \mathbf{b} := \mathbf{b}_s \left( \prod_{i=s+1}^m t_i \mathbf{w}_i \right).$$

To complete the proof, it suffices to show that  $\mathbf{w} \approx \hat{\mathbf{w}}$  holds in  $\mathbf{D}_{15}\{\mathbf{w} \approx \mathbf{w}'\}$  and  $\mathbf{D}_{15}\{\mathbf{w} \approx \hat{\mathbf{w}}\} = \mathbf{D}_{15}\Gamma_1$  for some  $\Gamma_1 \subseteq \Gamma$ . Indeed, in this case, the identity  $\hat{\mathbf{w}} \approx \mathbf{w}'$  is  $(r-1)$ -invertible. By the induction assumption,  $\mathbf{D}_{15}\{\hat{\mathbf{w}} \approx \mathbf{w}'\} = \mathbf{D}_{15}\Gamma_2$  for some  $\Gamma_2 \subseteq \Gamma$ , whence

$$\mathbf{D}_{15}\{\mathbf{w} \approx \mathbf{w}'\} = \mathbf{D}_{15}\{\mathbf{w} \approx \hat{\mathbf{w}}, \hat{\mathbf{w}} \approx \mathbf{w}'\} = \mathbf{D}_{15}\{\Gamma_1, \Gamma_2\},$$

and we are done.

Suppose that  $p = q = 1$ . Then, since  $\text{occ}_x(\mathbf{w}_s) = \text{occ}_x(\mathbf{w}'_s) = \text{occ}_y(\mathbf{w}_s) = \text{occ}_y(\mathbf{w}'_s) = 2$ , Lemma 5.2 implies that  $\mathbf{D}_{15}$  satisfies the identity  $\mathbf{w} \approx \hat{\mathbf{w}}$ . By a similar argument, we can show that  $\mathbf{D}_{15}$  satisfies the identity  $\mathbf{w} \approx \hat{\mathbf{w}}$  whenever  $p = q = 2$ . So, it remains to consider the case when  $\{p, q\} = \{1, 2\}$ .

If  $x, y \in \text{alph}(\mathbf{w}_{s'}) = \text{alph}(\mathbf{w}'_{s'})$  for some  $s' \neq s$ , then Lemma 5.2 or the dual statement implies that  $\mathbf{D}_{15}$  satisfies the identity  $\mathbf{w} \approx \hat{\mathbf{w}}$ . Thus, we may further assume that at most one of the letters  $x$  and  $y$  occurs in  $\text{alph}(\mathbf{w}_i)$  for any  $i \neq s$ . Denote this letter by  $a_i$  (if

$x, y \notin \text{alph}(\mathbf{w}_i)$ , then  $a_i$  denotes the empty word). For brevity, put

$$\mathbf{h} := \prod_{i=1}^s a_{i-1} t_i \text{ and } \mathbf{t} := \prod_{i=s+1}^m t_i a_i.$$

Suppose that  $(p, q) = (2, 1)$ . Then  $\mathbf{w}(x, y, t_1, \dots, t_m) \approx \mathbf{w}'(x, y, t_1, \dots, t_m)$  is equivalent modulo  $\{(3.1), (3.2)\}$  to  $\mathbf{h}x^2y^2\mathbf{t} \approx \mathbf{hpt}$  for some  $\mathbf{p} \in \{(xy)^2, xy^2x, (yx)^2, yx^2y, y^2x^2\}$ . The identities (4.2) hold in  $\mathbf{D}_{15}$ , whence  $\mathbf{D}_{15}\{\mathbf{w} \approx \mathbf{w}'\}$  satisfies either  $\mathbf{h}x^2y^2\mathbf{t} \approx \mathbf{h}(xy)^2\mathbf{t}$  or  $\mathbf{h}x^2y^2\mathbf{t} \approx \mathbf{h}y^2x^2\mathbf{t}$ . Further, since the identities

$$\mathbf{h}x^2y^2\mathbf{t} \stackrel{(3.1)}{\approx} \mathbf{h}x^4y^4\mathbf{t} \approx \mathbf{h}(x^2y^2)^2\mathbf{t} \stackrel{\{(3.1), (3.2)\}}{\approx} \mathbf{h}(xy)^2\mathbf{t}$$

hold in  $\mathbf{D}_{15}\{\mathbf{h}x^2y^2\mathbf{t} \approx \mathbf{h}y^2x^2\mathbf{t}\}$ , the identity  $\mathbf{h}x^2y^2\mathbf{t} \approx \mathbf{h}(xy)^2\mathbf{t}$  is satisfied by  $\mathbf{D}_{15}\{\mathbf{w} \approx \mathbf{w}'\}$  in either case. It is clear that the identity  $\mathbf{w}(x, y, t_1, \dots, t_m) \approx \hat{\mathbf{w}}(x, y, t_1, \dots, t_m)$  is nothing but the identity  $\mathbf{h}x^2y^2\mathbf{t} \approx \mathbf{h}(xy)^2\mathbf{t}$ . The variety  $\mathbf{D}_{15}\{\mathbf{h}x^2y^2\mathbf{t} \approx \mathbf{h}(xy)^2\mathbf{t}\}$  satisfies the identities

$$\begin{aligned} \mathbf{w} &= \mathbf{a}x\mathbf{y}\mathbf{b} \\ &\approx \mathbf{a}x^2y^2\mathbf{b} && \text{by } \{(3.1), (3.2)\} \\ &\approx \mathbf{a}(xy)^2\mathbf{b} && \text{by } \mathbf{h}x^2y^2\mathbf{t} \approx \mathbf{h}(xy)^2\mathbf{t} \\ &\approx \mathbf{a}y\mathbf{x}\mathbf{b} = \hat{\mathbf{w}} && \text{by Lemma 5.3.} \end{aligned}$$

Therefore, the identity  $\mathbf{h}x^2y^2\mathbf{t} \approx \mathbf{h}(xy)^2\mathbf{t}$  is equivalent within  $\mathbf{D}_{15}$  to the identity  $\mathbf{w} \approx \hat{\mathbf{w}}$ . It remains to notice that the identity  $\mathbf{h}x^2y^2\mathbf{t} \approx \mathbf{h}(xy)^2\mathbf{t}$  belongs to  $\Gamma$ .

The case when  $(p, q) = (1, 2)$  is quite similar and we omit the corresponding considerations.  $\square$

**Lemma 5.4.** *If  $\mathbf{X}, \mathbf{Y} \in [\mathbf{M}_\gamma(x^+yzx^+) \vee \mathbf{M}_\gamma(xx^+yy^+), \mathbf{V}]$  and  $\mathbf{X} \wedge \mathbf{Y}$  satisfies an identity (5.1) with  $k, \ell \geq 0$  and  $a_1, \dots, a_{k+\ell} \in \{1, x, y\}$ , then (5.1) is satisfied by either  $\mathbf{X}$  or  $\mathbf{Y}$ .*

*Proof.* In view of Proposition 2.2, there exists a finite sequence  $\mathbf{u} = \mathbf{w}_0, \dots, \mathbf{w}_m = \mathbf{v}$  of words such that each identity  $\mathbf{w}_j \approx \mathbf{w}_{j+1}$  holds in either  $\mathbf{X}$  or  $\mathbf{Y}$ . According to Lemma 2.5,

$$\mathbf{w}_j \in \left( \prod_{i=1}^k a_{i-1}^+ t_i \right) \mathbf{a}_j \left( \prod_{i=k+1}^{k+\ell} t_i a_i^+ \right),$$

where  $\text{alph}(\mathbf{a}_j) = \{x, y\}$  for any  $j = 0, \dots, m$ . Then there is  $s \in \{0, \dots, m-1\}$  such that  $\mathbf{a}_s \in x^+y^+$  but  $\mathbf{a}_{s+1} \in \overline{x^+y^+}$ . Now substitute  $t_k x$  for  $t_k$  and  $y t_{k+1}$  for  $t_{k+1}$  in  $\mathbf{w}_s \approx \mathbf{w}_{s+1}$ , resulting the identity which is equivalent modulo  $\{(3.1), (3.2)\}$  to (5.1). We see that (5.1) holds in either  $\mathbf{X}$  or  $\mathbf{Y}$ .  $\square$

**Acknowledgments.** The author is grateful to Mikhail V. Volkov for useful discussions.

#### REFERENCES

- [1] Almeida, J.: Finite Semigroups and Universal Algebra. World Scientific, Singapore (1994)
- [2] Cossey, J.: Critical groups and the lattice of varieties. Proc. Amer. Math. Soc. **20**, 217–221 (1969)
- [3] Edmunds, C.C.: On certain finitely based varieties of semigroups. Semigroup Forum **15**, 21–39 (1977)
- [4] Green, J.A.: On the structure of semigroups. Ann. Math. (2) **54**, 163–172 (1951)
- [5] Gusev, S.V.: Standard elements of the lattice of monoid varieties. Algebra i Logika **59**, 615–626 (2020) (in Russian; Engl. translation: Algebra and Logic **59**, 415–422 (2021))
- [6] Gusev, S.V.: Varieties of aperiodic monoids with central idempotents whose subvariety lattice is distributive. Monatsh. Math. **201**, 79–108 (2023)
- [7] Gusev, S.V.: Varieties of aperiodic monoids with commuting idempotents whose subvariety lattice is distributive. Semigroup Forum, submitted; see also Preprint: <http://arxiv.org/abs/2308.11544v2> (2023)

- [8] Gusev, S.V., Lee, E.W.H., Vernikov, B.M.: The lattice of varieties of monoids. *Japan. J. Math.* **17**, 117–183 (2022)
- [9] Gusev, S.V., Sapir, O.: Classification of limit varieties of  $J$ -trivial monoids. *Comm. Algebra* **50**, 3007–3027 (2022)
- [10] Gusev, S.V., Vernikov, B.M.: Chain varieties of monoids. *Dissert. Math.* **534**, 1–73 (2018)
- [11] Gusev, S.V., Vernikov, B.M.: Two weaker variants of congruence permutability for monoid varieties. *Semigroup Forum* **103**, 106–152 (2021)
- [12] Head, T.J.: The varieties of commutative monoids. *Nieuw Arch. Wiskunde. III Ser.* **16**, 203–206 (1968)
- [13] Higman, G.: Representations of general linear groups and varieties of groups. *Proc. Internat. Conf. Theory of Groups*. Austral. Nat. Univ. Canberra 1965. Gordon and Breach 1967.
- [14] Jackson, M.: Finiteness properties of varieties and the restriction to finite algebras. *Semigroup Forum* **70**, 159–187 (2005)
- [15] Jackson, M., Lee, E.W.H.: Monoid varieties with extreme properties. *Trans. Amer. Math. Soc.* **370**, 4785–4812 (2018)
- [16] Lee, E.W.H.: Varieties generated by 2-testable monoids. *Studia Sci. Math. Hungar.* **49**, 366–389 (2012)
- [17] Lee, E.W.H.: On certain Cross varieties of aperiodic monoids with commuting idempotents, *Results Math.* **66**, 491–510 (2014)
- [18] Lee, E.W.H.: A minimal pseudo-complex monoid, *Arch. Math.* **120**, 15–25 (2023)
- [19] Lee, E.W.H., Li, J.R.: Minimal non-finitely based monoids, *Dissert. Math.*, **475**, 1–65 (2011)
- [20] Kovács, L.G., Newman, M.F.: On non-cross varieties of groups. *J. Austral. Math. Soc.* **12**, 129–144 (1971)
- [21] Kozhevnikov, P.A.: On nonfinitely based varieties of groups of large prime exponent. *Comm. Algebra* **40**, 2628–2644 (2012)
- [22] Neumann, H.: Varieties of groups. Springer-Verlag, Berlin (1967)
- [23] Perkins, P.: Bases for equational theories of semigroups. *J. Algebra* **11**, 298–314 (1969)
- [24] Pin, J.-É.: Varieties of Formal Languages. North Oxford Academic, London (1986)
- [25] Roman'kov, V.A.: Nondistributivity of the lattice of varieties of nilpotent groups. *Algebra i Logika* **9**, 67–72 (1970) (in Russian)
- [26] Sapir, M.V.: Problems of Burnside type and the finite basis property in varieties of semigroups. *Izv. Akad. Nauk SSSR, Ser. Mat.* **51**, 319–340 (1987) (in Russian; English translation: *Mathematics of the USSR–Izv.* **30**, 295–314 (1988))
- [27] Sapir, O.: Non-finitely based monoids. *Semigroup Forum* **90**, 557–586 (2015)
- [28] Sapir, O.: Lee monoids are non-finitely based while the sets of their isoterms are finitely based. *Bull. Aust. Math. Soc.* **97**, 422–434 (2018)
- [29] Sapir, O.: Limit varieties of  $J$ -trivial monoids. *Semigroup Forum* **103**, 236–260 (2021)
- [30] Sapir, O.: Limit varieties generated by finite non- $J$ -trivial aperiodic monoids. *Semigroup Forum* **107**, 732–750 (2023)
- [31] Shevrin, L.N., Vernikov, B.M., Volkov, M.V.: Lattices of semigroup varieties. *Izv. Vyssh. Uchebn. Zaved. Matem. (3)*, 3–36 (2009) (in Russian; Engl. translation: *Russian Math. (Iz. VUZ)* **53**(3), 1–28 (2009))
- [32] The Sverdlovsk Notebook, 2nd ed., Ural State University, Sverdlovsk (1979) (in Russian)
- [33] Wismath, S.L.: The lattice of varieties and pseudovarieties of band monoids. *Semigroup Forum*, **33**, 187–198 (1986)
- [34] Zhang W.T., Luo, Y.F.: A new example of limit variety of aperiodic monoids. Preprint: <http://arxiv.org/abs/1901.02207> (2019)