

Chiral spin state and nematic ferromagnet in the spin-1 Kitaev- Γ model

Qiang Luo,^{1,2,*} Jize Zhao,^{3,4,†} Jinbin Li,^{1,2} and Xiaoqun Wang^{5,‡}

¹College of Physics, Nanjing University of Aeronautics and Astronautics, Nanjing, 211106, China

²Key Laboratory of Aerospace Information Materials and Physics (NUAA), MIIT, Nanjing, 211106, China

³School of Physical Science and Technology & Key Laboratory of Quantum

Theory and Applications of MoE, Lanzhou University, Lanzhou 730000, China

⁴Lanzhou Center for Theoretical Physics, Key Laboratory of Theoretical Physics of Gansu Province, Lanzhou University, Lanzhou 730000, China

⁵School of Physics, Zhejiang University, Hangzhou 310058, China

(Dated: March 14, 2024)

The higher-spin Kitaev magnets, in which the Kitaev interaction and off-diagonal exchange couplings are overwhelmingly large, have emerged as a fertile avenue to explore exotic phases and unusual excitations. In this work, we study the quantum phase diagram of the spin-1 Kitaev- Γ model on the honeycomb lattice using density-matrix renormalization group. It harbours six distinct phases and the intriguing findings are three magnetically ordered phases in which both time-reversal symmetry and lattice symmetry albeit of different sort are broken spontaneously. The chiral spin state originates from the order-by-disorder effect and exhibits an almost saturated scalar spin chirality at the quantum level. Depending on the relative strength of the two interactions, it also features columnar or plaquette valence-bond-solid-like pattern as a consequence of the translational symmetry breaking. In parallel, the nematic ferromagnets are situated at ferromagnetic Kitaev side and possess small but finite ferromagnetic ordering. The lattice-rotational symmetry breaking enforces nonequivalent bond energy along one of the three bonds. Although the intrinsic difference between the two nematic ferromagnets remains elusive, the discontinuities in the von Neumann entropy, hexagonal plaquette operator, and Wilson loop operator convincingly suggest that they are separated via a first-order phase transition.

I. INTRODUCTION

The celebrated Kitaev honeycomb model [1], which consists of bond-directional Ising couplings, plays an essential role in the understanding of exotic phases of matter, e.g., quantum spin liquid (QSL) possessing topological order and fractionalized excitations [2, 3]. The QSL is a nonmagnetic phase which escapes from any spontaneous symmetry breaking down to the lowest temperature [4] and has an intimate relation to high-temperature superconductivity in cuprates and quantum computations. In this endeavor, great efforts have been made to realize the superbly Kitaev interaction in real materials. Among them, Jackeli and Khaliulin pointed out that the strong spin-orbit coupling together with proper electronic correlations could give rise to this interaction [5]. In the quest for these “Kitaev materials” on the honeycomb lattice, which include iridates [6–8], α - RuCl_3 [9–17], and $\text{Na}_2\text{Co}_2\text{TeO}_6$ and $\text{Na}_3\text{Co}_2\text{SbO}_6$ [18–21], it turns out that they usually display magnetic orderings at low temperatures for the possible existence of competing interactions. In spite of this, signatures of continuum spectrum [22] and half-quantized thermal Hall conductivity [23] have been reported in α - RuCl_3 , while an intriguing triple- Q order that is proximity to hidden $\text{SU}(2)$ point is studied in $\text{Na}_2\text{Co}_2\text{TeO}_6$ [24].

On the other hand, devising alternative Kitaev materials with $S > 1/2$ could potentially expand the field of seeking interesting phases. So far, the highly desirable Kitaev interaction has been identified in some higher-spin magnets on the honeycomb lattice [25]. In the spin-1 Ni-based antimonates $\text{A}_3\text{Ni}_2\text{SbO}_6$ ($\text{A} = \text{Li}, \text{Na}$), the thermal entropy of $\sim \frac{1}{2} \ln 3$ per Ni element over the Neel temperatures firmly hints the existence of the Kitaev interaction [26]. In their allotropic $\text{Na}_3\text{Ni}_2\text{BiO}_6$, the Kitaev interaction is proposed to enhance exchange frustration and stabilize the one-third magnetization plateau in the partial spin-flop ferrimagnetic order [27]. In KNiAsO_4 , inclusion of the Kitaev interaction is essential to adequately explain the zigzag magnetic order and other thermodynamical quantities [28]. In the spin-3/2 Cr-based van der Waals magnet CrI_3 , CrSiTe_3 , and CrGeTe_3 [29–31], the Kitaev interaction is found to be induced by heavy ligands of I/Te despite a weak spin-orbit coupling related to Cr irons. Although the compelling evidences to advocate the Kitaev QSL remain unclear [32], there are proposals to impair the non-Kitaev terms in CrSiTe_3 and CrGeTe_3 by strain engineering [30]. In addition, identification of the Kitaev interaction in triangular-lattice compounds has also been reported, which include the spin-1 NiI_2 [33] and the spin-3/2 $1T$ - CrTe_2 [34]. In the latter, the Kitaev interaction is crucial to understand the orientation of the magnetic moment that is 70° off the normal direction of the plane [35].

According to the proposal by Baskaran, Sen, and Shankar [36], the extensive number of local conserved quantities guarantee the vanishing of the spin-spin correlation functions beyond the nearest-neighbor bonds in

*qiangluo@nuaa.edu.cn

†zhaojz@lzu.edu.cn

‡xiaoqunwang@zju.edu.cn

the pure Kitaev limit. Therefore, the ground state of the spin- S Kitaev model should always be a sort of QSL. Hitherto, the specific heat and thermal entropy [37, 38], the topological entanglement entropy [39], and the excitation spectrum [40] of the spin-1 Kitaev QSL have been widely studied. A magnetic-field-induced intermediate nonmagnetic phase sandwiched between the Kitaev QSL and the polarized phase is reported [41–43]. In the spin-3/2 analogy, the spin can be reexpressed by the $SO(6)$ Majorana representation and the Kitaev QSL is found to couple to a static \mathbb{Z}_2 gauge field [44]. Recently, there has been a surge interest in higher-spin models with Kitaev-type interaction. In the spin-1 Kitaev model, it is found that the Kitaev QSL survives against finite Heisenberg interaction [45], and intervals of the Kitaev QSLs shrink rapidly as S increases [46]. Furthermore, the interplay of Kitaev interaction and other competing terms, e.g., bi-quadratic exchange [47] and single-ion anisotropy [48], is anticipated to promote inspiring phases and unconventional quantum phase transitions (QPTs).

Motivated by the Kitaev materials with $j_{\text{eff}} = 1/2$, the Kitaev- Γ model is broadly recognized as their minimal model [49]. Despite great efforts, the full quantum phase diagram of this model remains unsettled, especially when K and Γ have opposite signs [3, 50–54]. As $S > 1/2$, the quantum fluctuations can be suppressed in some sense, leaving the possibility of figuring out a more precise quantum phase diagram. Our work thus serves as a guide to understand the novel phases in higher-spin Kitaev magnets. By virtue of the density-matrix renormalization group (DMRG) method [55–57], we elucidate the quantum phase diagram by systematically changing the ratio between the Kitaev and Γ interactions. We uncover six phases in total and three of them are non-magnetic, which are ferromagnetic (FM) and antiferromagnetic (AFM) Kitaev QSLs and AFM Γ (AF) phase that is surrounded in the Γ limit. In addition, there are three magnetically ordered states, which are the chiral spin state, and two emergent nematic ferromagnets near the FM Kitaev QSL.

The layout of the rest of the paper is as follows. In Sec. II, we explain the model and the associated operators, introduce the numerical methods, and present the quantum phase diagram. The detailed analysis of the chiral spin state and the nematic ferromagnets are shown in Sec. III and Sec. IV, respectively. Finally, conclusion and some further discussion are given in Sec. V.

II. MODEL AND PHASE DIAGRAM

We study the spin-1 Kitaev- Γ model on the honeycomb lattice, which has the Hamiltonian defined as

$$\mathcal{H} = \sum_{\langle ij \rangle \parallel \gamma} \left[K S_i^\gamma S_j^\gamma + \Gamma (S_i^\alpha S_j^\beta + S_i^\beta S_j^\alpha) \right], \quad (1)$$

where S_i^γ ($\gamma = x, y$, and z) is the γ -component of a spin operator at site i and $\langle ij \rangle \parallel \gamma$ denotes the γ -type nearest-




FIG. 1: (a) Illustration of the hexagonal plaquette operator \hat{W}_p . The nearest-neighbor links are distinguished as \mathbf{X} (red), \mathbf{Y} (green), and \mathbf{Z} (blue) bonds, respectively. (b) Illustration of the Wilson loop operator \hat{W}_l . The yellow loop denotes the Wilson loop operator along the $\mathbf{a}_2(1/2, \sqrt{3}/2)$ direction. (c) The classical phase diagram of the Kitaev- Γ model, based on previous Monte Carlo results [58, 59]. (d) The quantum phase diagram of the spin-1 Kitaev- Γ model, see the main text for explanation.

neighbor bond connecting sites i and j . α and β are the two remaining bonds under a cyclic permutation of $\{x, y, z\}$. In addition, K and Γ stand for the Kitaev and Γ interactions, respectively. We consider a trigonometric parametrization of these interactions such that $K = \cos \phi$ and $\Gamma = \sin \phi$ with $0 \leq \phi \leq \pi$.

We have studied the quantum phase diagram of this model by using the DMRG method on two different clusters. One is a C_3 -symmetric 24-site hexagonal cluster, which is primarily adopted since it retains the symmetry of the Hamiltonian in Eq. (1). The other is a $2 \times L_x \times L_y$ rhombic cluster with $N = 2L_x L_y$ sites in total, which is utilized for comparison and extrapolation. Full periodic boundary conditions are used in both clusters, with the exception of the long cylinder in the latter case. During the computation, we keep as many as $m = 4000$ block states and exert up to 24 sweeps until the typical truncation error is less than 10^{-6} .

Different from the spin-1/2 Kitaev model, the spin-1 counterpart does not seem to have an exact solution up to now. Nevertheless, it has been verified that this model can be characterized by a couple of conserved \mathbb{Z}_2 quantities. One of such quantities is the hexagonal plaquette operator [36]

$$\hat{W}_p = e^{i\pi(S_1^y + S_2^z + S_3^x + S_4^y + S_5^z + S_6^x)}, \quad (2)$$

where S_j^γ is the outgoing bond around the hexagon path at site j , see Fig. 1(a). Another representative quantity




FIG. 3: (a) The scalar spin chirality χ as a function of ϕ on the 24-site hexagonal cluster. Inset: Illustration of the spin patterns on the center spin \mathbf{S}_0 and its three nearest-neighbor spins $\{\mathbf{S}_i, \mathbf{S}_j, \mathbf{S}_k\}$. (b) and (c) sketch the patterns of the columnar and plaquette VBS-like chiral spin state at $\phi/\pi = 0.1$ and $\phi/\pi = 0.4$, respectively. The wider bond has a higher intensity of the bond energy.

III. CHIRAL SPIN STATE

A. Order-by-disorder effect

We start by considering the unfrustrated case where $K, \Gamma > 0$. Following the recipe proposed by Rouschatzakis *et al.* [62, 63], we can choose and parametrize a central spin as $\mathbf{S}_0 = (\eta_a a, \eta_b b, \eta_c c)$, where (η_a, a) [respectively, (η_b, b) and (η_c, c)] stand for the sign and intensity of S_0^x (respectively, S_0^y and S_0^z). Here, $a^2 + b^2 + c^2 = S^2$ and η s are independent Ising variables that can be ± 1 . By noticing the peculiar structures of the Kitaev and Γ interactions, the three neighbor spins subject to different bond types can be determined as $\mathbf{S}_i = -(\eta_a a, \eta_c c, \eta_b b)$, $\mathbf{S}_j = -(\eta_c c, \eta_b b, \eta_a a)$, and $\mathbf{S}_k = -(\eta_b b, \eta_a a, \eta_c c)$ [see the inset of Fig. 3(a)]. In doing so, all the spins can be determined gradually, formulating specific spin textures with finite unit cells. Given that there are only three different η s, the ground-state degeneracy is eight in addition to the free choice of $\{a, b, c\}$. Typically, the unit cell contains six spins except for the case of two-sublattice AFM phase in which $\eta_a = \eta_b = \eta_c = \pm 1$ and $a = b = c = S/\sqrt{3}$.

Irrelevant of the choice of $\{\eta_a, \eta_b, \eta_c\}$ and $\{a, b, c\}$, the ground-state energy per site is $e_{cl} = -(\Gamma + K/2)S^2$ classically. When the fluctuations are involved, the real-space perturbation theory is amenable to address the issue of choosing $\{a, b, c\}$. The second-order calculation shows that the energy correlation is [3, 64]

$$e_{cl}^{(2)} = -\frac{(\Gamma - K)^2 S}{32[\Gamma + 2K]} [(a/S)^4 + (b/S)^4 + (c/S)^4]. \quad (5)$$

Thus, when $K \neq \Gamma$, the total energy is minimized as long as one of the elements in $\{a, b, c\}$ is unitary while the rest are zero. Meanwhile, only one of the η s that adheres to the nonzero element survives. This leads to the so-called

Cartesian states which map to dimer coverings of the honeycomb lattice [36].

The order-by-disorder effect can also be verified by the linear spin-wave theory. Using the Holstein-Primakoff transformation, the spin operator at i -th site can be approximately rewritten as $\hat{\mathbf{S}}_i = (\sqrt{S/2}(b_i + b_i^\dagger), -i\sqrt{S/2}(b_i - b_i^\dagger), S - b_i^\dagger b_i)$, and the Hamiltonian in Eq. (1) is readily to be expressed into a quadratic form. The spin-wave energy is given by [52]

$$e_{sw} = \left(1 + \frac{1}{S}\right) e_{cl} + \frac{S}{2} \sum_{\mathbf{v}\mathbf{q}} \omega_{\mathbf{v}\mathbf{q}}. \quad (6)$$

When the exchange couplings are fixed, the spin-wave energy e_{sw} can vary slightly for different choice of $\{a, b, c\}$. Figure 4 shows the spherical plots of the spin-wave energy for different values of $\{a, b, c\}$ at $\phi/\pi = 0.1$ and 0.4 . In both cases, the energy takes the minimal value on the points where one of the three spin components is $\pm S$. This result is fairly consistent with that of the real-space perturbation theory shown in Eq. (5). As a byproduct, spin-wave energy gains a maximum when $|a| = |b| = |c| = S/\sqrt{3}$, showing that the AFM-like phase is not favored by quantum fluctuations.

B. Scalar spin chirality

Before disclosing patterns of dimer coverings, we wish to note that the underlying spin patterns support a magnetically ordered phase featuring finite scalar spin chirality. On the one hand, if we define the magnetic order parameter as $M(\mathbf{Q}) = \lim_{N \rightarrow \infty} \sqrt{\mathbf{S}_N(\mathbf{Q})/N}$, it is directly to find that [65]

$$M^2(\Gamma') = \frac{S^2}{4} + \frac{\eta_a \eta_b a b + \eta_b \eta_c b c + \eta_c \eta_a c a}{2} = \frac{S^2}{4} \quad (7)$$




FIG. 4: Spherical plots of the spin-wave energy ϵ_{sw} at (a) $\phi/\pi = 0.1$ and (b) $\phi/\pi = 0.4$, respectively. The central spin \mathbf{S}_0 is parameterized as $(S^x, S^y, S^z) = (\sin \vartheta \cos \varphi, \sin \vartheta \sin \varphi, \cos \vartheta)$.

and

$$M^2(\mathbf{K}) = \frac{S^2}{6} - \frac{\eta_a \eta_b ab + \eta_b \eta_c bc + \eta_c \eta_a ca}{6} = \frac{S^2}{6}, \quad (8)$$

where we have used the fact that $\eta_a \eta_b ab + \eta_b \eta_c bc + \eta_c \eta_a ca = 0$ since only one of the elements in $\{a, b, c\}$ is nonzero. The above two equations also imply that $M(\mathbf{K})/M(\Gamma') = \sqrt{6}/3 \approx 0.8165$, which is verified by our DMRG calculation shown in Fig. 2(b). On the other hand, the scalar spin chirality is usually defined as [65, 66]

$$\chi_{ijk}^\Delta = \langle \hat{\mathbf{S}}_i \cdot (\hat{\mathbf{S}}_j \times \hat{\mathbf{S}}_k) \rangle, \quad (9)$$

where sites (i, j, k) form an equilateral triangle (Δ) in the anticlockwise direction. For the spins shown in the inset of Fig. 3(a), it follows that

$$\chi_{ijk}^\Delta = (\eta_a a)^3 + (\eta_b b)^3 + (\eta_c c)^3 - 3\eta_a \eta_b \eta_c abc = S^3. \quad (10)$$

The DMRG calculation of the scalar spin chirality presented in Fig. 3(a) indeed suggests that it is close to 1 when away from the phase boundaries. Thus, our results justify this spin texture as a magnetically ordered chiral spin state.

Finally, we turn to discuss the real space patterns of dimer coverings. In the Cartesian states, there have and only have one sort of interaction for each bond, favoring either Kitaev bond with an energy of $-KS^2$ or Γ bond with an energy of $-\Gamma S^2$. Due to the unique structures of the two interactions, the Γ bonds form a set of hexagonal loops while the Kitaev bonds connect these isolated hexagonal loops. Therefore, this gives a columnar-like valence-bond solid (VBS) when Kitaev interaction is dominant and a plaquette-like VBS otherwise, see Fig. 3(b) and Fig. 3(c). In the honeycomb lattice, we note that there is a dual (triangular) lattice which is constructed by placing a vertex inside each hexagon and connecting new vertices. Thus, this forms a triangular Ising magnet with a three-sublattice $\{\eta_a, \eta_b, \eta_c\}$ unit cell. For each equally-weighted hexagonal loop (which is centered at η_b in Fig. 3(b) and η_a in Fig. 3(c)) in the two

VBSs, the nonzero component of each spin coincides with the type of outgoing bond, rendering the corresponding plaquette operator $\langle \hat{W}_p \rangle$ be unitary. By contrast, the plaquette operators at the remaining two cases are zero. Therefore, the flux-like density $\langle \bar{W}_p \rangle = \langle \hat{W}_p \rangle / N_p$ ($N_p = N/2$) is found to be $1/3$. Such a trimerization phenomenon is precisely confirmed by our DMRG calculation (see Fig. 2(a)). We note in passing that evidence of trimerization in spin-1/2 counterpart is also present, albeit $\langle \bar{W}_p \rangle$ equals to $-1/3$ [64].

It is worthwhile to note that these physical pictures should break down when $K = \Gamma (> 0)$, which is a hidden $SU(2)$ Heisenberg point [67]. The ground state turns out to be a dual FM phase via a six-sublattice transformation. In other context, the chiral spin state is also dubbed counter-rotating spiral order, which is identified in the polymorphs of the three-dimensional Li_2IrO_3 [68].

IV. NEMATIC FERROMAGNETS

The aim of this Section is to unveil the nematic ferromagnets. To this end, we start by using the von Neumann entropy, the hexagonal plaquette operator, and the Wilson loop operator to determine the QPTs. Next, we pin down the nematic ferromagnets by examining nematic and magnetic order parameters. Our findings are advocated by comparing results on 24-site hexagonal cluster and rhombic cluster. Finally, we also present the landscape of bond energy on long cylinder at representative parameter points to verify the robustness of lattice nematicity.

A. Nematic order parameter

The FM Kitaev QSL is widely believed to be fragile against perturbations. For example, the out-of-plane magnetic field of the strength $\sim 0.01K$ can drive the ground state into the polarized phase [41], while the Heisenberg interaction of the strength $\sim 0.05K$ can change the ground state into the long-range orderings [45]. Therefore, one needs to resort sensitive quantities such as the entanglement entropy to probe the QPTs. The entanglement entropy describes quantum correlations existed in the ground states and usually displays different behaviors when crossing quantum critical points. Specifically, the von Neumann entropy is defined as $\mathcal{S}_{\text{vN}}(l) = -\text{tr}(\rho_l \ln \rho_l)$ where ρ_l is the reduced density matrix of the targeted subsystem with length l . The length l is typically fixed to be $N/2$ so that the blocks of the system and environment are equal in the DMRG calculation.

Figure 5(a) shows the von Neumann entropy \mathcal{S}_{vN} in the window of $\phi/\pi \in [0.92, 1.00]$ on the 24-site hexagonal cluster. \mathcal{S}_{vN} has three discontinuities at $\phi_3^{\text{FM}}/\pi \approx 0.952$, $\phi_2^{\text{FM}}/\pi \approx 0.979$, and $\phi_1^{\text{FM}}/\pi \approx 0.996$, respectively, signifying three first-order QPTs. Noteworthily, the transi-




FIG. 5: (a) The von Neumann entropy \mathcal{S}_{vN} (left axis) and the flux-like density $\langle \bar{W}_p \rangle$ (right axis) as a function of $\phi/\pi \in [0.92, 1.00]$ on the 24-site hexagonal cluster. The shaded areas denote the regions of the two nematic ferromagnets. (b) The bond energy E_γ [$\gamma = x$ (red), y (green), and z (blue)] (left panel) and the nematic order parameter \mathcal{O}_{NF} (right panel) in the regime of interest. (c) Illustration of the patterns of the nematic ferromagnets for $\phi/\pi = 0.97$ (left panel) and $\phi/\pi = 0.99$. The thickness of the bond is proportional to cube of the relative bond energy E_{ij} , i.e., $\propto (E_{ij}/\max(E_{ij}))^3$, for the guide of eyes. (d)-(f) and the same as these of (a)-(c) but for the $2 \times 4 \times 3$ cluster.

tion points are in accordance with these obtained from the flux-like density $\langle \bar{W}_p \rangle$. While the emergent phases in the narrow region of $\phi_3^{\text{FM}} < \phi < \phi_1^{\text{FM}}$ are tempting, it is challenging to disclose them for the lack of classical analogues. Recalling that the SSFs shown in Fig. 2(c) unambiguously indicate the broken of C_3 rotational symmetry, it is thus reasonable to suspect that the lattice nematicity may arise. To proceed further, we calculate the bond energy $E_\gamma = \langle H_{ij}^\gamma \rangle$ along the three Kitaev bonds. As can be seen from the left panel in Fig. 5(b), there is an apparent difference in the bond energy in the region of $\phi_3^{\text{FM}} < \phi < \phi_1^{\text{FM}}$. Typically, the absolute values of the bond energy along two of the three bonds are close and are bigger than the remaining one. In our DMRG calculation along a specific path through the hexagonal cluster, the weak bond is of the **X**-type when $\phi/\pi = 0.97$ and is of the **Z**-type when $\phi/\pi = 0.99$, see Fig. 5(c). We thus define the nematic order parameter as the difference of the bond energy, i.e., $\mathcal{O}_{\text{NF}} = |\min(E_\gamma) - \max(E_\gamma)|$. The \mathcal{O}_{NF} is indeed finite in the region of interest. It seems to be smoothly evolve into the Kitaev QSL at ϕ_1^{FM} , while it undergoes a jump at ϕ_3^{FM} .

To check for the possible cluster dependence, we compare the results to these obtained on a $2 \times 4 \times 3$ rhom-

bic cluster, see Figs. 5(d)-(f). Comparing Fig. 5(d) to Fig. 5(a), \mathcal{S}_{vN} also acquires three jumps or kinks despite that the turning points are slightly different, i.e., $\phi_3^{\text{FM}}/\pi \approx 0.962$, $\tilde{\phi}_2^{\text{FM}}/\pi \approx 0.984$, and $\phi_1^{\text{FM}}/\pi \approx 0.997$. The Wilson loop operator $\langle W_l \rangle$ decreases gradually and goes down to zero when away from the Kitaev limit. It has a steepest descent near ϕ_1^{FM} and an evident jump at $\tilde{\phi}_2^{\text{FM}}$, followed by an inconspicuous hump with a maxima around $\tilde{\phi}_3^{\text{FM}}$ (see the inset). Pertaining the lattice nematicity, we also confirm that there is a difference in the three bond energy and one of the bond energy is obvious distinguished from the others. However, it is the **Y**-type bond that always has the weakest strength in both nematic ferromagnets, in contrast to the former case of C_3 hexagonal cluster. Irrelevant to this difference, the maximum of \mathcal{O}_{NF} in the nematic ferromagnets are rather close, with a value of 0.063 (hexagonal cluster) and 0.061 (rhombic cluster), respectively. Nevertheless, the major difference lies in that the AF phase also exhibits a finite lattice nematicity in the rhombic geometry, as opposite to C_3 -rotational hexagonal one. This outcome is somewhat akin to its spin-1/2 analogy where the so-called nematic paramagnet is reported by the infinite DMRG calculation [54].




FIG. 6: (a) The SSF $\mathbb{S}_N(\Gamma)$ as a function of ϕ/π on three $2 \times L_x \times 3$ clusters with $L_x = 2$ ($N = 12$, red circle), 3 ($N = 18$, green triangle), and 4 ($N = 24$, blue square). (b) Extrapolations of the FM order parameter $M_N(\Gamma)$ with respect to four clusters of $2 \times 2 \times 3$ ($N = 12$), $2 \times 3 \times 3$ ($N = 18$), $2 \times 4 \times 3$ ($N = 24$), and $2 \times 4 \times 4$ ($N = 32$). The parameter ϕ/π equals to 0.97 (left), 0.99 (middle), and 1.00 (right), and the error bars are less than the size of the symbols.

B. Magnetic order parameter

A parallel question to address is the magnetic ordering in the nematic region. According to the results on the 24-site hexagonal cluster, the SSF has a primary peak at the Γ point and a competing peak at \mathbf{X} (or \mathbf{X}') point. It is then curious to know how the SSF at Γ point will change as the system size varies. Given the limitation of the DMRG calculation, we turn to utilize the rhombic clusters. While these clusters are not equipped with the C_3 rotational symmetry of the Hamiltonian, the Γ point with a zero momentum is always accessible, advocating the reliability of the magnetic ordering after a proper extrapolation.

Figure 6(a) shows the ϕ -dependence of SSF $\mathbb{S}_N(\Gamma)$ on three $2 \times L_x \times 3$ clusters with $L_x = 2$ (red circle), 3 (green triangle), and 4 (blue square). $\mathbb{S}_N(\Gamma)$ has a hump in the nematic regions and its value decreases consecutively with the increase of N from 12 to 24. To proceed further, we make a quadratic extrapolation of the magnetic order parameter $M_N(\Gamma) = \sqrt{\mathbb{S}_N(\Gamma)/N}$ versus $1/\sqrt{N}$, see Fig. 6(b). We note that the magnetic order parameter on the $2 \times 4 \times 4$ cluster with $N = 32$ is taken into account, yet with the largest error bar. The extrapolations at $\phi/\pi = 0.97$ (left) and 0.99 (middle) in the nematic regions

yield small but finite values around 0.05. As a comparison, $M_N(\Gamma)$ at $\phi/\pi = 1$ in the QSL phase is estimated to be zero within the numerical precision. Therefore, we conclude that the nematic regions indeed possess FM ordering and are thus termed nematic ferromagnets.

C. Nematicity on long cylinder

Previously, we have unveiled a nematic region in the vicinity of the FM Kitaev QSL on two different 24-site clusters. It is thus imperative to check for the tendency of nematicity on larger system size. In this regard, we calculate the nearest-neighbor bond energy on a $2 \times 18 \times 3$ cylinder where open boundary condition is employed on $\mathbf{a}_1(1,0)$ direction, see Figs. 7(a) and (b). The difference in the thickness of the three bonds clearly demonstrates a lattice nematicity at $\phi/\pi = 0.97$ and 0.99, and the intensity contrast is more conspicuous in the former than the latter. When $\phi/\pi = 0.97$, the DMRG calculation suggests that the bond energy in the middle region of the cylinder along the \mathbf{X} , \mathbf{Y} , and \mathbf{Z} bonds are -0.4863 , -0.4214 , and -0.4863 , respectively, yielding a nematic order parameter of 0.0649. As a comparison, the nematic order parameter at $\phi/\pi = 0.99$ is only 0.0128, nearly one fifth of the former. Despite the difference in intensity, the values of the nematic order parameters are exceedingly close to those obtained on 24-site clusters (cf. Fig. 5). The robustness of the nematicity unambiguously emphasizes the lattice-rotational symmetry breaking in the nematic region.

To further reveal the nature of the nematic ferromagnets, we present the behaviors of von Neumann entropy $\mathcal{S}_{\text{VN}}(l)$ at $\phi/\pi = 0.97$ and 0.99 in Figs. 7(c) and (d), respectively. The curves are quite alike and the lowest branches in which l is a multiply of six are fairly flat, indicative of a vanishing central charge c in conformal field theory. In other words, the excitations in the nematic ferromagnets seem to be gapped. However, the difference of the two may lie in the distribution of entanglement spectrum. It is found that the nematic ferromagnet I at $\phi/\pi = 0.97$ is characterized by a nearly doubly degenerate entanglement spectrum at least for the lowest part, while the entanglement spectrum in the nematic ferromagnet II at $\phi/\pi = 0.99$ is odd (not shown). Nevertheless, such a difference does not seem to be a universal character as it is not observed in the 24-site clusters. Our results imply that the intrinsic divide between the two nematic ferromagnets is still obscure and calls for future study.

V. CONCLUSION

We studied the quantum phase diagram of the spin-1 Kitaev- Γ model on the honeycomb lattice. Through DMRG calculations on various clusters, we showed that the model harbours six distinct phases, which are the FM and AFM Kitaev QSLs, the nonmagnetic AF phase that




FIG. 7: (a) and (b) show the landscapes of bond energy E_{ij} on a $2 \times 18 \times 3$ long cylinder for $\phi/\pi = 0.97$ and $\phi/\pi = 0.99$, respectively. The thickness of the bond is proportional to cube of the relativity bond energy, i.e., $\propto (E_{ij}/\max(E_{ij}))^3$, for the guide of eyes. (c) and (d) show the von Neumann entropy $S_{vN}(l)$ of a consecutive segment of length l on a $2 \times 18 \times 3$ long cylinder for $\phi/\pi = 0.97$ and $\phi/\pi = 0.99$, respectively. The solid symbols of the lowest branch represent the neat edge-cutting with l being a multiply of six (i.e., the number of the sites along each column).

embraces the pure Γ limit, the chiral spin state, and two nematic ferromagnets. The Kitaev QSLs in the absence of Γ term are characterized by a unitary hexagonal plaquette operator $\langle \hat{W}_p \rangle$ and Wilson loop operator $\langle \hat{W}_l \rangle$, and the Γ -driven QPTs can be captured by the singularities in $\langle \hat{W}_p \rangle$ and $\langle \hat{W}_l \rangle$. In accordance with the spin-1 Kitaev- Γ chain [69], our result also indicates that the AFM Kitaev QSL is more robust against the Γ interaction than its FM analogy.

Beyond these nonmagnetic phases, the main findings of this work are two categories of magnetically ordered states. To begin with, when the Kitaev and Γ interactions are both positive, the chiral spin state is selected from the infinitely degenerate manifold of all possible states via order-by-disorder mechanism. As a result, the ground state exhibits a noncoplanar pattern in which all spins point along cubic axes in the spin space. It has a six-sublattice unit cell and displays peaks at \mathbf{K} and $\mathbf{\Gamma}'$ points in the Brillouin zone. Noteworthily, this phase exhibits an almost saturated scalar spin chirality, signifying the broken of time-reversal symmetry. As the Kitaev and Γ interactions are different, the real-space bond energy displays columnar (when $K > \Gamma$) or plaquette (when $K < \Gamma$) VBS-like pattern.

In addition, there are two emergent nematic ferromagnets in the vicinity of the dominating FM Kitaev regime. These phases feature multiple- Q structures where several Bragg peaks coexist in the reciprocal space. Among these peaks, the peak at $\mathbf{\Gamma}$ point seems to have the strongest intensity in a wide interval and the proper extrapolations of this magnetic order parameter yield small but finite

values, indicative of magnetically ordered states. In addition, the landscapes of the nearest-neighbor bond energy show apparent nonequivalence among the three bonds, in which two of the bond energy are nearly equal but are much larger than the remaining. The resulting nematic order parameter defined as the difference in the bond energy keeps finite in a certain region and the values are robust even on long cylinders. However, the jumps in von Neumann entropy and other quantities clearly demonstrate a first-order QPT in between. While the differences in the preferred bond and entanglement spectrum are possible evidences to reveal the distinction, decisive characters to distinguish the two remains unclear. On all accounts, these results are unambiguous to identify two nematic ferromagnets as the ground states, and our work may stimulate resurgent study on the controversial region near the FM Kitaev QSL in the spin-1/2 model [3].

Acknowledgments

We are indebted to H.-Y. Kee for the previous collaboration on related works. This work is supported by the National Program on Key Research Project (Grant No. MOST2022YFA1402700), the Natural Science Foundation of Jiangsu Province (Grant No. BK20220876), the National Natural Science Foundation of China (Grants No. 12304176, No. 12274187, No. 12247183, and No. 12247101). Q.L. also acknowledges the startup Fund of Nanjing University of Aeronautics and Astronautics

(Grant No. YAH21129). The computations are partially supported by High Performance Computing Platform of

Nanjing University of Aeronautics and Astronautics.

-
- [1] A. Kitaev, Anyons in an exactly solved model and beyond, *Ann. Phys. (NY)* **321**, 2 (2006).
- [2] J. Wen, S.-L. Yu, S. Li, W. Yu, and J.-X. Li, Experimental identification of quantum spin liquids, *npj Quantum Mater.* **4**, 12 (2019).
- [3] I. Rousochatzakis, N. B. Perkins, Q. Luo, and H.-Y. Kee, Beyond Kitaev physics in strong spin-orbit coupled magnets, *Rep. Prog. Phys.* **87**, 026502 (2024).
- [4] L. Balents, Spin liquids in frustrated magnets, *Nature (London)* **464**, 199 (2010).
- [5] G. Jackeli and G. Khaliullin, Mott insulators in the strong spin-orbit coupling limit: From Heisenberg to a quantum compass and Kitaev models, *Phys. Rev. Lett.* **102**, 017205 (2009).
- [6] F. Ye, S. Chi, H. Cao, B. C. Chakoumakos, J. A. Fernandez-Baca, R. Custelcean, T. F. Qi, O. B. Korneta, and G. Cao, Direct evidence of a zigzag spin-chain structure in the honeycomb lattice: A neutron and x-ray diffraction investigation of single-crystal Na_2IrO_3 , *Phys. Rev. B* **85**, 180403(R) (2012).
- [7] S. K. Choi, R. Coldea, A. N. Kolmogorov, T. Lancaster, I. I. Mazin, S. J. Blundell, P. G. Radaelli, Y. Singh, P. Gegenwart, K. R. Choi, S.-W. Cheong, P. J. Baker, C. Stock, and J. Taylor, Spin Waves and Revised Crystal Structure of Honeycomb Iridate Na_2IrO_3 , *Phys. Rev. Lett.* **108**, 127204 (2012).
- [8] S. H. Chun, J.-W. Kim, J. Kim, H. Zheng, C. C. Stoumpos, C. D. Malliakas, J. F. Mitchell, K. Mehlawat, Y. Singh, Y. Choi, T. Gog, A. Al-Zein, M. M. Sala, M. Krisch, J. Chaloupka, G. Jackeli, G. Khaliullin, and B. J. Kim, Direct evidence for dominant bond-directional interactions in a honeycomb lattice iridate Na_2IrO_3 , *Nat. Phys.* **11**, 462 (2015).
- [9] J. A. Sears, M. Songvilay, K. W. Plumb, J. P. Clancy, Y. Qiu, Y. Zhao, D. Parshall, and Y.-J. Kim, Magnetic order in $\alpha\text{-RuCl}_3$: A honeycomb-lattice quantum magnet with strong spin-orbit coupling, *Phys. Rev. B* **91**, 144420 (2015).
- [10] J. A. Sears, Y. Zhao, Z. Xu, J. W. Lynn, and Y.-J. Kim, Phase Diagram of $\alpha\text{-RuCl}_3$ in an in-plane Magnetic Field, *Phys. Rev. B* **95**, 180411(R) (2017).
- [11] S.-H. Baek, S.-H. Do, K.-Y. Choi, Y. S. Kwon, A. U. B. Wolter, S. Nishimoto, J. van den Brink, and B. Büchner, Evidence for a Field-Induced Quantum Spin Liquid in $\alpha\text{-RuCl}_3$, *Phys. Rev. Lett.* **119**, 037201 (2017).
- [12] A. U. B. Wolter, L. T. Corredor, L. Janssen, K. Nenkov, S. Schönecker, S.-H. Do, K.-Y. Choi, R. Albrecht, J. Hunger, T. Doert, M. Vojta, and B. Büchner, Field-induced quantum criticality in the Kitaev system $\alpha\text{-RuCl}_3$, *Phys. Rev. B* **96**, 041405(R) (2017).
- [13] Z. Wang, S. Reschke, D. Hüvonen, S.-H. Do, K.-Y. Choi, M. Gensch, U. Nagel, T. Room, and A. Loidl, Magnetic Excitations and Continuum of a Possibly Field-Induced Quantum Spin Liquid in $\alpha\text{-RuCl}_3$, *Phys. Rev. Lett.* **119**, 227202 (2017).
- [14] J. Zheng, K. Ran, T. Li, J. Wang, P. Wang, B. Liu, Z.-X. Liu, B. Normand, J. Wen, and W. Yu, Gapless Spin Excitations in the Field-Induced Quantum Spin Liquid Phase of $\alpha\text{-RuCl}_3$, *Phys. Rev. Lett.* **119**, 227208 (2017).
- [15] S.-H. Do, S.-Y. Park, J. Yoshitake, J. Nasu, Y. Motome, Y. S. Kwon, D. T. Adroja, D. J. Voneshen, K. Kim, T.-H. Jang, J.-H. Park, K.-Y. Choi, and S. Ji, Majorana fermions in the Kitaev quantum spin system $\alpha\text{-RuCl}_3$, *Nat. Phys.* **13**, 1079 (2017).
- [16] K. Ran, J. Wang, W. Wang, Z.-Y. Dong, X. Ren, S. Bao, S. Li, Z. Ma, Y. Gan, Y. Zhang, J. T. Park, G. Deng, S. Danilkin, S.-L. Yu, J.-X. Li, and J. Wen, Spin-Wave Excitations Evidencing the Kitaev Interaction in Single Crystalline $\alpha\text{-RuCl}_3$, *Phys. Rev. Lett.* **118**, 107203 (2017).
- [17] S. M. Winter, K. Riedl, P. A. Maksimov, A. L. Chernyshev, A. Honecker, and R. Valenti, Breakdown of magnons in a strongly spin-orbital coupled magnet, *Nat. Commun.* **8**, 1152 (2018).
- [18] G. Lin, *et al.*, Field-induced quantum spin disordered state in spin-1/2 honeycomb magnet $\text{Na}_2\text{Co}_2\text{TeO}_6$, *Nat. Commun.* **12**, 5559 (2021).
- [19] X. Li, Y. Gu, Y. Chen, V. O. Garlea, K. Iida, K. Kamazawa, Y. Li, G. Deng, Q. Xiao, X. Zheng, Z. Ye, Y. Peng, I. A. Zaliznyak, J. M. Tranquada, and Y. Li, Giant Magnetic In-Plane Anisotropy and Competing Instabilities in $\text{Na}_3\text{Co}_2\text{SbO}_6$, *Phys. Rev. X* **12**, 041024 (2022).
- [20] H. Takeda, J. Mai, M. Akazawa, K. Tamura, J. Yan, K. Moovendaran, K. Raju, R. Sankar, K.-Y. Choi, and M. Yamashita, Planar thermal Hall effects in the Kitaev spin liquid candidate $\text{Na}_2\text{Co}_2\text{TeO}_6$, *Phys. Rev. Research* **4**, L042035 (2022).
- [21] S. Guang, N. Li, R. L. Luo, Q. Huang, Y. Wang, X. Yue, K. Xia, Q. Li, X. Zhao, G. Chen, H. Zhou, and X. Sun, Thermal transport of fractionalized antiferromagnetic and field-induced states in the Kitaev material $\text{Na}_2\text{Co}_2\text{TeO}_6$, *Phys. Rev. B* **107**, 184423 (2023).
- [22] A. Banerjee, C. A. Bridges, J.-Q. Yan, A. A. Aczel, L. Li, M. B. Stone, G. E. Granroth, M. D. Lumsden, Y. Yiu, J. Knolle, S. Bhattacharjee, D. L. Kovrizhin, R. Moessner, D. A. Tennant, D. G. Mandrus, and S. E. Nagler, Proximate Kitaev quantum spin liquid behaviour in a honeycomb magnet, *Nat. Mater.* **15**, 733 (2016).
- [23] Y. Kasahara, T. Ohnishi, Y. Mizukami, O. Tanaka, Sixiao Ma, K. Sugii, N. Kurita, H. Tanaka, J. Nasu, Y. Motome, T. Shibauchi, and Y. Matsuda, Majorana quantization and half-integer thermal quantum Hall effect in a Kitaev spin liquid, *Nature (London)* **559**, 227 (2010).
- [24] W. G. F. Krüger, W. Chen, X. Jin, Y. Li, and L. Janssen, Triple-q Order in $\text{Na}_2\text{Co}_2\text{TeO}_6$ from Proximity to Hidden-SU(2)-Symmetric Point, *Phys. Rev. Lett.* **131**, 146702 (2023).
- [25] P. P. Stavropoulos, D. Pereira, and H.-Y. Kee, Microscopic Mechanism for a Higher-Spin Kitaev Model, *Phys. Rev. Lett.* **123**, 037203 (2019).
- [26] E. A. Zvereva, M. I. Stratan, Y. A. Ovchennikov, V. B. Nalbandyan, J.-Y. Lin, E. L. Vavilova, M. F. Iakovleva, M. Abdel-Hafiez, A. V. Silhanek, X.-J. Chen, A. Stroppa, S. Picozzi, H. O. Jeschke, R. Valentí, and A. N. Vasiliev,

- Zigzag antiferromagnetic quantum ground state in monoclinic honeycomb lattice antimonates *Phys. Rev. B* **92**, 144401 (2015).
- [27] Y. Shangguan, S. Bao, Z.-Y. Dong, N. Xi, Y.-P. Gao, Z. Ma, W. Wang, Z. Qi, S. Zhang, Z. Huang, *et al.*, A one-third magnetization plateau phase as evidence for the Kitaev interaction in a honeycomb-lattice antiferromagnet, *Nat. Phys.* **19**, 1883-1889 (2023).
- [28] K. M. Taddei, V. O. Garlea, A. M. Samarakoon, L. D. Sanjeewa, J. Xing, T. W. Heitmann, C. dela Cruz, A. S. Sefat, and D. Parker, Zigzag magnetic order and possible Kitaev interactions in the spin-1 honeycomb lattice KNiAsO_4 , *Phys. Rev. Research* **5**, 013022 (2023).
- [29] C. Xu, J. Feng, H. Xiang, and L. Bellaiche, Interplay between Kitaev interaction and single ion anisotropy in ferromagnetic CrI_3 and CrGeTe_3 monolayers, *npj Comput. Mater.* **4**, 57 (2018).
- [30] C. Xu, J. Feng, M. Kawamura, Y. Yamaji, Y. Nahas, S. Prokhorenko, Y. Qi, H. Xiang, and L. Bellaiche, Possible Kitaev quantum spin liquid state in 2D materials with $S = 3/2$, *Phys. Rev. Lett.* **124**, 087205 (2020).
- [31] P. P. Stavropoulos, X. Liu, and H.-Y. Kee, Magnetic anisotropy in spin-3/2 with heavy ligand in honeycomb mott insulators: Application to CrI_3 , *Phys. Rev. Research* **3**, 013216 (2021).
- [32] Z. Zhou, K. Chen, Q. Luo, H.-G. Luo, and J. Zhao, Strain-induced phase diagram of the $S = \frac{3}{2}$ Kitaev material CrSiTe_3 , *Phys. Rev. B* **104**, 214425 (2021).
- [33] Q. Song, C. A. Occhialini, E. Ergeçen, B. Ilyas, D. Amoroso, P. Barone, J. Kapeghian, K. Watanabe, T. Taniguchi, A. S. Botana, S. Picozzi, N. Gedik, and R. Comin, Evidence for a single-layer van der Waals multiferroic, *Nature* **602**, 601-605 (2022).
- [34] D. C. Freitas, R. Weht, A. Sulpice, G. Remenyi, P. Strobel, F. Gay, J. Marcus, and M. Núñez-Regueiro, Ferromagnetism in layered metastable $1T\text{-CrTe}_2$, *J. Phys.: Condens. Matter* **27**, 176002 (2015).
- [35] C. Huang, B. Liu, L. Jiang, Y. Pan, J. Fan, D. Shi, C. Ma, Q. Luo, and Y. Zhu, Evidence of Kitaev interaction in monolayer $1T\text{-CrTe}_2$, *Phys. Rev. B* **108**, 094433 (2023).
- [36] G. Baskaran, Diptiman Sen, and R. Shankar, Spin- S Kitaev model: Classical ground states, order from disorder, and exact correlation functions, *Phys. Rev. B* **78**, 115116 (2008).
- [37] A. Koga, H. Tomishige, and J. Nasu, Ground-state and Thermodynamic Properties of an $S = 1$ Kitaev Model, *J. Phys. Soc. Jpn.* **87**, 063703 (2018).
- [38] J. Oitmaa, A. Koga, and R. R. P. Singh, Incipient and well-developed entropy plateaus in spin- S Kitaev models, *Phys. Rev. B* **98**, 214404 (2018).
- [39] H.-Y. Lee, N. Kawashima, and Y. B. Kim, Tensor network wave function of $S = 1$ Kitaev spin liquids, *Phys. Rev. Research* **2**, 033318 (2020).
- [40] Y.-H. Chen, J. Genzor, Y. B. Kim, and Y.-J. Kao, Excitation spectrum of spin-1 Kitaev spin liquids, *Phys. Rev. B* **105**, L060403 (2022).
- [41] Z. Zhu, Z.-Y. Weng, and D. N. Sheng, Magnetic field induced spin liquids in $S = 1$ Kitaev honeycomb model, *Phys. Rev. Research* **2**, 022047(R) (2020).
- [42] C. Hickey, C. Berke, P. P. Stavropoulos, H.-Y. Kee, and S. Trebst, Field-driven gapless spin liquid in the spin-1 Kitaev honeycomb model, *Phys. Rev. Research* **2**, 023361 (2020).
- [43] I. Khait, P. P. Stavropoulos, H.-Y. Kee, and Y. B. Kim, Characterizing spin-one Kitaev quantum spin liquids, *Phys. Rev. Research* **3**, 013160 (2021).
- [44] H.-K. Jin, W. M. H. Natori, F. Pollmann, and J. Knolle, Unveiling the $S = 3/2$ Kitaev honeycomb spin liquids, *Nat. Commun.* **13**, 3813 (2022).
- [45] X.-Y. Dong and D. Sheng, Spin-1 Kitaev-Heisenberg model on a honeycomb lattice, *Phys. Rev. B* **102**, 121102(R) (2020).
- [46] K. Fukui, Y. Kato, J. Nasu, and Y. Motome, Ground-state phase diagram of spin- S Kitaev-Heisenberg models, *Phys. Rev. B* **106**, 174416 (2022).
- [47] R. Pohle, N. Shannon, and Y. Motome, Spin nematics meet spin liquids: Exotic quantum phases in the spin-1 bilinear-biquadratic model with Kitaev interactions, *Phys. Rev. B* **107**, L140403 (2023).
- [48] Ayushi Singhanian, Jeroen van den Brink, and Satoshi Nishimoto, Emergence of vortex state in the $S = 1$ Kitaev-Heisenberg model with single-ion anisotropy, *arXiv:2311.14161*.
- [49] W. Wang, Z.-Y. Dong, S.-L. Yu, and J.-X. Li, Theoretical investigation of magnetic dynamics in $\alpha\text{-RuCl}_3$, *Phys. Rev. B* **96**, 115103 (2017).
- [50] J. G. Rau, E. K.-H. Lee, and H.-Y. Kee, Generic Spin Model for the Honeycomb Iridates beyond the Kitaev Limit, *Phys. Rev. Lett.* **112**, 077204 (2014).
- [51] J. Wang, B. Normand, and Z.-X. Liu, One Proximate Kitaev Spin Liquid in the $K\text{-J}\text{-}\Gamma$ Model on the Honeycomb Lattice, *Phys. Rev. Lett.* **123**, 197201 (2019).
- [52] Q. Luo, J. Zhao, H.-Y. Kee, and X. Wang, Gapless quantum spin liquid in a honeycomb Γ magnet, *npj Quantum Mater.* **6**, 57 (2021).
- [53] H.-Y. Lee, R. Kaneko, L. E. Chern, T. Okubo, Y. Yamaji, N. Kawashima, and Y. B. Kim, Magnetic-field induced quantum phases in tensor network study of Kitaev magnets, *Nat. Commun.* **11**, 1639 (2020).
- [54] M. Gohlke, L. E. Chern, H.-Y. Kee, and Y. B. Kim, Emergence of nematic paramagnet via quantum order-by-disorder and pseudo-Goldstone modes in Kitaev magnets, *Phys. Rev. Research* **2**, 043023 (2020).
- [55] S. R. White, Density matrix formulation for quantum renormalization groups, *Phys. Rev. Lett.* **69**, 2863 (1992).
- [56] I. Peschel, X. Q. Wang, M. Kaulke, and K. Hallberg, *Density-Matrix Renormalization* (Springer, Berlin, 1999).
- [57] U. Schollwöck, The density-matrix renormalization group, *Rev. Mod. Phys.* **77**, 259 (2005).
- [58] A. Rayyan, Q. Luo, and H.-Y. Kee, Extent of frustration in the classical Kitaev- Γ model via bond anisotropy, *Phys. Rev. B* **104**, 094431 (2021).
- [59] K. Chen, Q. Luo, Z. Zhou, S. He, B. Xi, C. Jia, H.-G. Luo and J. Zhao, Triple-meron crystal in high-spin Kitaev magnets, *New J. Phys.* **25**, 023006 (2023).
- [60] P. P. Stavropoulos, Y. Yang, I. Rousochatzakis, and N. B. Perkins, Complex orders and chirality in the classical Kitaev- Γ model, *arXiv:2311.00037* (2023).
- [61] S.-S. Zhang, G. B. Halász, W. Zhu, and C. D. Batista, Variational study of the Kitaev-Heisenberg-Gamma model, *Phys. Rev. B* **104**, 014411 (2021).
- [62] I. Rousochatzakis and N. B. Perkins, Classical Spin Liquid Instability Driven By Off-Diagonal Exchange in Strong Spin-Orbit Magnets, *Phys. Rev. Lett.* **118**, 147204 (2017).
- [63] I. Rousochatzakis, Y. Sizyuk, and N. B. Perkins, Quan-

- tum spin liquid in the semiclassical regime, *Nat. Commun.* **9**, 1575 (2018).
- [64] I. Rousochatzakis, The spin-1/2 K - Γ Honeycomb Model: Semiclassical, strong coupling and exact diagonalization results, *KITP Program: Correlated Systems with Multicomponent Local Hilbert Spaces* (Sep. 28 - Dec. 18, 2020).
- [65] Q. Luo, P. P. Stavropoulos, J. S. Gordon, and H.-Y. Kee, Spontaneous chiral-spin ordering in spin-orbit coupled honeycomb magnets, *Phys. Rev. Research* **4**, 013062 (2022).
- [66] S.-Y. Yu, H. Li, Q.-R. Zhao, Y. Gao, X.-Y. Dong, Z.-X. Liu, W. Li, S.-S. Gong, Nematic chiral spin liquid in a Kitaev magnet under external magnetic field, *arXiv:2304.00555* (2023).
- [67] J. Chaloupka and G. Khaliullin, Hidden symmetries of the extended Kitaev-Heisenberg model: Implications for the honeycomb-lattice iridates $A_2\text{IrO}_3$, *Phys. Rev. B* **92**, 024413 (2015).
- [68] P. P. Stavropoulos, A. Catuneanu, and H.-Y. Kee, Counter-rotating spiral order in three-dimensional iridates: Signature of hidden symmetry in the Kitaev- Γ model, *Phys. Rev. B* **98**, 104401 (2018).
- [69] Q. Luo, S. Hu, and H.-Y. Kee, Unusual excitations and double-peak specific heat in a bond-alternating spin-1 K - Γ chain, *Phys. Rev. Research* **3**, 033048 (2021).