

# On harvesting physical predictions from asymptotically safe quantum field theories

Frank Saueressig\* and Agustín Silva†

*High Energy Physics Department, Institute for Mathematics, Astrophysics,  
and Particle Physics, Radboud University, Nijmegen, The Netherlands.*

(Dated: April 8, 2024)

Asymptotic safety is a powerful mechanism for obtaining a consistent and predictive quantum field theory beyond the realm of perturbation theory. It hinges on an interacting fixed point of the Wilsonian renormalization group flow which controls the microscopic dynamics. Connecting the fixed point to observations requires constructing the set of effective actions compatible with this microscopic dynamics. Technically, this information is stored in the UV-critical surface of the fixed point. In this work, we describe a novel approach for extracting this information based on analytical and pseudo-spectral methods. Our construction is illustrated at the level of the two-dimensional Ising model and easily generalizes to any asymptotically safe quantum field theory. It also constitutes an important step towards setting up a well-founded swampland program within the gravitational asymptotic safety program.

## I. INTRODUCTION

The Wilsonian renormalization group (RG) [1] provides a powerful tool for understanding the impact of statistical or quantum fluctuations in a given physical system [2–5]. It organizes fluctuations in momentum shells and integrates out the modes, starting from the most energetic ones and subsequently moving to lower energies. This procedure creates RG trajectories which connect effective descriptions of the same system at different values of the coarse-graining scale.

A major success of this approach is a comprehensive picture of critical phenomena and universality which are readily explained in terms of RG fixed points which control the theory’s infrared (IR) behavior. Typically, one has to tune a small number of parameters so that a RG trajectory is dragged into the fixed point in the limit where all fluctuations are integrated out. These trajectories then “forget” their microscopic origin and physical quantities like correlation functions are completely dictated by the properties of the IR fixed point.

From a physics perspective, one may also be interested in situations where a RG fixed point provides the microscopic description of the system. This situation applies to asymptotically free quantum field theories like quantum chromodynamics where the relevant fixed point is the free theory. Asymptotic safety, first proposed in [6, 7], generalizes this construction to interacting fixed points, also called non-Gaussian fixed points (NGFPs). In both cases, the fixed point equips the construction with predictive power. By definition, RG trajectories whose ultraviolet (UV) completion is provided by the fixed point span its UV-critical surface  $\mathcal{S}_{UV}$ . In the vicinity of the fixed point, this surface can be found by linearizing the RG flow and identifying the UV-attractive directions. Typically,  $\mathcal{S}_{UV}$  is finite-dimensional and embedded in a larger

space called the theory space  $\mathcal{T}$ . This allows to predict values for renormalized couplings based on the free parameters specifying a RG trajectory within  $\mathcal{S}_{UV}$ . These predictions are obtained from the end-point of the RG trajectory where all fluctuations have been accounted for. In the case of an IR fixed point, this limit is controlled by the fixed point itself, so that its properties are directly related to observables. For an UV fixed point, this link is highly non-trivial though and requires the construction of  $\mathcal{S}_{UV}$ . Depending on the dimensions of  $\mathcal{S}_{UV}$  and  $\mathcal{T}$  this becomes cumbersome rather quickly. A prototypical example is provided by asymptotically safe quantum gravity supplemented by the matter degrees of freedom of the standard model, where  $\dim(\mathcal{S}_{UV})$  is expected to be of order twenty [8]. This surface should then be embedded in an even larger space also comprising couplings associated with beyond the standard model physics for making predictions. It is clear that the canonical approach of mapping out  $\mathcal{S}_{UV}$  based on a shooting-method is intractable in such cases. Nevertheless, a solid knowledge about the effective actions within  $\mathcal{S}_{UV}$  is crucial for making predictions and potentially falsifying the asymptotic safety hypothesis.

The goal of this letter is to introduce novel strategies which allow to map out  $\mathcal{S}_{UV}$  in a computationally efficient way using pseudo-spectral methods [9, 10] which are readily boosted by machine-learning algorithms to improve convergence. We illustrate the generic algorithm based on the RG flow of a scalar field theory in less than four dimensions which has already been extensively studied, e.g., in [11–18]. Owing to the flexibility of the RG [3], our algorithm is applicable to a much wider range of settings, including the study of phase transitions in statistical physics [19], estimates of the rate of spontaneous nucleation [20], strongly interacting and high temperature theories [21], up to quantum gravity building on the gravitational asymptotic safety program [22, 23].

\* f.saueressig@science.ru.nl

† agustin.silva@ru.nl

## II. RG FLOWS, FIXED POINTS AND FREE PARAMETERS

A primary tool for computing Wilsonian RG flows is the Wetterich equation [24–27],<sup>1</sup>

$$k\partial_k\Gamma_k = \frac{1}{2}\text{Tr}\left[\left(\Gamma_k^{(2)} + \mathcal{R}_k\right)^{-1}k\partial_k\mathcal{R}_k\right], \quad (1)$$

which encodes the dependence of the effective average action  $\Gamma_k$  on the coarse-graining scale  $k$ , see [29–31] for more details. The effective average action lives on theory space  $\mathcal{T}$ . By definition,  $\mathcal{T}$  contains all action functionals which can be constructed from the field content of the theory and are compatible with its symmetry requirements. For the purpose of this work, we are interested in complete (approximate) solutions of (1),  $k \mapsto \Gamma_k$ , which interpolate between a NGFP in the limit  $k \rightarrow \infty$  (asymptotic safety) and the effective action  $\Gamma \equiv \lim_{k \rightarrow 0} \Gamma_k$  associated with observables.

Introducing a basis  $\{\mathcal{O}_n\}$  on  $\mathcal{T}$ ,  $\Gamma_k$  can be expanded as

$$\Gamma_k = \sum_n \tilde{g}_k^n \mathcal{O}_n. \quad (2)$$

Here the  $\tilde{g}_k^n$  are the dimensionful couplings of the theory which depend on  $k$ . Denoting the mass-dimension of  $\tilde{g}_k^n$  by  $d_n$ , these couplings are conveniently traded for their dimensionless counterparts

$$g_k^n \equiv \tilde{g}_k^n k^{-d_n}. \quad (3)$$

In practice, the expansion (2) retains a finite number of operators,  $N$ , only. The couplings  $g^n$ ,  $n = 1, \dots, N$  can then be read as coordinates on  $\mathbb{R}^N$ . Substituting (2) into (1), the beta functions capturing the  $k$ -dependence of these couplings  $g_k^n$  can be read off as the coefficients multiplying the basis elements  $\{\mathcal{O}_n\}$ :

$$k\partial_k g_k^n = \beta^n(g^m). \quad (4)$$

In the sequel, we will assume that the beta functions have been computed and admit a NGFP suitable for asymptotic safety.

By definition, fixed points  $\{g_*^n\}$  satisfy  $\beta^n(g_*^m) = 0, \forall n$ . The dimension of  $\mathcal{S}_{UV}$  associated with the fixed point is readily inferred by linearizing the beta functions

$$k\partial_k g_k^n \simeq \sum_m B^n_m (g_k^m - g_*^m), \quad B^n_m \equiv \left. \frac{\partial \beta^n}{\partial g^m} \right|_{g=g_*}. \quad (5)$$

These equations are readily solved in terms of the stability coefficients  $\theta_I$  and right-eigenvectors  $V_I$ , satisfying  $BV_I = -\theta_I V_I$ . Separating the stability coefficients




FIG. 1. Using the choice of couplings introduced below eq. (12), the  $k \rightarrow 0$  limit of the couplings  $u^\alpha$  is  $\{u^\alpha\} = 0$ . Every complete RG trajectory emanating from the NGFP gives rise to an effective action whose parameters  $v^\mu$  are read off from the intersect of  $\mathcal{S}_{UV}$  with the axes  $\{u^\alpha\} = 0$ .

$\{\theta_I\} \rightarrow \{\theta_\alpha, \tilde{\theta}_\mu\}$  where  $\alpha = 1, \dots, \dim(\mathcal{S}_{UV})$  and  $\mu = 1, \dots, N - \dim(\mathcal{S}_{UV})$  enumerate the stability coefficients with  $\text{Re}(\theta_\alpha) > 0$  and  $\text{Re}(\tilde{\theta}_\mu) < 0$ , respectively, one has

$$g_k^n \approx g_*^n + \sum_\alpha C_\alpha V_\alpha^n \left(\frac{k_0}{k}\right)^{\theta_\alpha} + \sum_\mu \tilde{C}_\mu \tilde{V}_\mu^n \left(\frac{k_0}{k}\right)^{\tilde{\theta}_\mu}. \quad (6)$$

Here  $k_0$  is a reference scale and  $C_\alpha, \tilde{C}_\mu$  are integration constants labelling the specific solutions. The eigendirections  $V_\alpha$  ( $\tilde{V}_\mu$ ) are attractive (repulsive) in the limit  $k \rightarrow \infty$ . Thus asymptotic safety fixes  $\tilde{C}_\mu = 0$  while the  $C_\alpha$  are free parameters. RG trajectories where  $\tilde{C}_\mu = 0$  span  $\mathcal{S}_{UV}$ .

Subsequently, we reformulate the conditions on the integration constants in terms of the dimensionless couplings. For this purpose, we split

$$g^n \mapsto \{u^\alpha, v^\mu\}, \quad \beta^n \mapsto \{\beta^\alpha, \beta^\mu\}. \quad (7)$$

The basic idea illustrated in Fig. 1 is to use the  $u^\alpha$  as coordinates on  $\mathcal{S}_{UV}$ .<sup>2</sup> Its embedding into  $\mathbb{R}^N$  is then given by a set of generating functions

$$F_\nu(u, v) = 0, \quad \nu = 1, \dots, N - \dim(\mathcal{S}_{UV}). \quad (8)$$

The generating functions are combinations of couplings whose values are conserved along the RG flow. We adopt

<sup>1</sup> Our construction is readily adapted to other RG equations like the Polchinski equation [28]. Essentially, it applies to any situation where a vector field is used to generate a constraint surface also including the case of IR-repulsive surfaces of IR-fixed points.

<sup>2</sup> Depending on the structure of  $\mathcal{S}_{UV}$ , the initial coordinate system may not cover the entire surface. In this case,  $\mathcal{S}_{UV}$  has to be covered with multiple coordinate patches. A simple illustration is provided by the surface  $x^2 + y^2 - 1 = 0$  embedded into  $\mathbb{R}^2$  where there is no single-valued function  $y(x)$  that covers the entire surface. In the context of the RG such coordinate changes have recently been discussed in [32].

the convention that  $\mathcal{S}_{\text{UV}}$  is generated by setting the conserved quantity to zero. Solving these relations for  $v^\mu$  then determines the values of the couplings  $v^\mu$  in terms of the coordinates on the surface

$$v^\mu|_{\mathcal{S}_{\text{UV}}} = v^\mu(u^\alpha). \quad (9)$$

In the vicinity of the NGFP, the functions  $F_\nu$  can be found as follows. The vectors  $\{V_\alpha, \tilde{V}_\mu\}$  span the tangent space at  $\{g_*^n\}$ . The vectors  $V_\alpha$  are tangent to  $\mathcal{S}_{\text{UV}}$  while, in general, the  $\tilde{V}_\mu$  are not normal to the surface. A set of normal vectors can then be constructed in two steps. First, we project the  $V_\alpha$  into a orthogonal basis  $t_\alpha$ , carrying out a Gram-Schmidt process. Starting from the vectors  $\tilde{V}_\mu$ , the set of normals is obtained as

$$n_\mu = \tilde{V}_\mu - \sum_\alpha t_\alpha \frac{\langle t_\alpha | \tilde{V}_\mu \rangle}{\langle t_\alpha | t_\alpha \rangle}, \quad (10)$$

where  $\langle x | y \rangle$  is the standard vector product on  $\mathbb{R}^N$ . At the linearized level, the constrained equations (8) are then given by

$$F_\mu^*(g^n) = \sum_n n_\mu^n (g^n - g_*^n). \quad (11)$$

Here \* emphasizes that this the expansion of the full  $F_\mu$  to linear order at the NGFP. One may then single out couplings  $u^\alpha$  according to (7) and solve these linear relations to express the  $v^\mu$  in terms of these parameters.

### III. CONSTRUCTING $\mathcal{S}_{\text{UV}}$

Finding the IR-endpoints of asymptotically safe RG trajectories requires the construction of  $\mathcal{S}_{\text{UV}}$  beyond the linear approximation (11). Technically, it is then convenient to work with a compact subregion of  $\mathbb{R}^N$  which contains the NGFP and the endpoints. In order to achieve this, it is instructive to discuss the  $k \rightarrow 0$ -limit of eq. (3). Keeping  $\lim_{k \rightarrow 0} \tilde{g}_k^n$  fixed and finite, which is a natural assumption for couplings appearing in the effective action, the dimensionless couplings with non-zero mass-dimension  $d_n$  either approach zero or diverge

$$\begin{aligned} \lim_{k \rightarrow 0} g_k^n &= 0, & d_n < 0, \\ \lim_{k \rightarrow 0} g_k^n &= \pm\infty, & d_n > 0. \end{aligned} \quad (12)$$

Couplings with  $d_n = 0$  take the finite value of the coupling appearing in the effective action. Based on these insights, we choose the couplings  $\tilde{u}^\alpha$  to have a negative mass dimension. This can always be achieved by making the identification  $\tilde{u}^n = 1/\tilde{g}^n$  if the initial coupling comes with a positive mass dimension. This has the advantage that the IR-endpoints are located at  $\{u^\mu\} = 0$ . Furthermore, we can take the  $v^\mu$  to be dimensionless by constructing appropriate ratios of the couplings, see (20) for an explicit example. The physical predictions for the couplings  $v^\mu$  are then read off at  $\{u^\alpha\} = 0$ .

### A. Master equation and explicit form

We now derive our master equation encoding the structure of  $\mathcal{S}_{\text{UV}}$ . Taking the  $k$ -derivative of the generating functions  $F_\mu(u, v)$  and using the definition of the beta functions (4) leads to

$$\sum_\alpha \frac{\partial F_\mu}{\partial u^\alpha} \beta^\alpha + \sum_\nu \frac{\partial F_\mu}{\partial v^\nu} \beta^\nu = 0. \quad (13)$$

At this point we assume that  $\det(\frac{\partial F_\mu}{\partial v^\nu}) \neq 0$ , so that the implicit function theorem guarantees that the system  $F_\mu = 0$  has a (local) solution in the form (9).

The master equation can be turned into an system of non-linear partial differential equations for the functions (9). Substituting  $F_\mu(u, v) = v^\mu - v^\mu(u)|_{\mathcal{S}_{\text{UV}}}$ , which obviously satisfies  $F_\mu|_{\mathcal{S}_{\text{UV}}} = 0$ , leads to the linear partial differential equation

$$\sum_\alpha \beta^\alpha \frac{\partial v^\mu}{\partial u^\alpha} = \beta^\mu. \quad (14)$$

The boundary conditions for the system are given in the linearized regime, eq. (11). Solutions can then be obtained either by solving the system (13) or (14) using numerical methods or extending the power series (11) to include higher-order terms. The latter strategy requires checking whether the endpoint of interest is within the radius of convergence of the series.

### B. Implicit solutions

The key strength of the master equation (13) is that it is a multi-linear, first order system of partial differential equations for the generating functions  $F_\mu(u, v)$ . Finding the implicit form of  $\mathcal{S}_{\text{UV}}$  by solving this system leads to a significant reduction in complexity as compared to solving the explicit system (14). In particular, the linear nature of the master equation makes the use of pseudo-spectral methods and collocation techniques [33, 34] a highly efficient tool for obtaining solutions.<sup>3</sup>

The basic idea is to select a dense set of functions  $\{\psi_i(u, v)\}$ . Typical examples used in practice include Chebyshev polynomials [9] or radial basis functions [39]. The generating functions are then approximated by expanding in this basis

$$F_\mu(u, v) \simeq \sum_{i=1}^{N_p} p_{\mu,i} \psi_i(u, v), \quad (15)$$

i.e., the functions are determined by the  $N_p \times (N - \dim(\mathcal{S}_{\text{UV}}))$  free parameters  $p_{\mu,i}$ . Since (13) contains

<sup>3</sup> For earlier applications of pseudo-spectral methods in the context of the functional renormalization group see [35–38].

derivatives of  $F_\mu(u, v)$  only, it determines solutions up to an additive constant only. Following (11), we fix this freedom by demanding that

$$F_\mu(u_*, v_*) = 0. \quad (16)$$

Secondly, the first derivatives of  $F_\mu$  evaluated at the fixed point are taken to agree with the linearized approximation

$$\left. \frac{\partial F_\mu}{\partial u^\alpha} \right|_{g_*^\alpha} = \frac{\partial F_\mu^*}{\partial u^\alpha}, \quad \left. \frac{\partial F_\mu}{\partial v^\nu} \right|_{g_*^\nu} = \frac{\partial F_\mu^*}{\partial v^\nu}. \quad (17)$$

This enforces that some of the expansion coefficients in (15) must take non-vanishing values thereby eliminating the trivial solution.

Subsequently, one chooses  $N_p$  collocation points. It is natural to take the NGFP as one of these points and construct a grid covering the parameter space of interest. Substituting the expansion (15) into (13) and evaluating the resulting expressions at the collocation points then leads to a system of algebraic equations which determines the coefficients  $p_{\mu,i}$ . The predictions  $v^\mu$  compatible with asymptotic safety are then found by evaluating the approximate solution at  $\{u^\alpha\} = 0$ .

*Systematic improvements.* In the simplest case, one may arrange the collocation points  $(u_{c,i}^\alpha, v_{c,i}^\mu)$  in a regular lattice and expand the generating functions in terms of Multivariate Cauchy Distributions (MCDs)

$$\psi_i(u, v) = \left( 1 + \sum_\alpha \frac{(u^\alpha - u_{c,i}^\alpha)^2}{\sigma^2} + \sum_\mu \frac{(v^\mu - v_{c,i}^\mu)^2}{\sigma^2} \right)^{-1}. \quad (18)$$

Here  $\sigma$  is a smoothness parameter whose typical value is chosen to be of the order of the average separation between the collocation points. Since the substitution of (15) into the system (13) results in a linear equation, taking one collocation point for each MCD leads to a unique solution for the parameters  $p_{\mu,i}$ .

Depending on the concrete application, this “basic” setup may be improved by working with less parameters than collocation points, and/or with base functions more suitable for describing sharp edges, periodicity, or optimized convergence [40]. Moreover, the positioning of the collocation points may be optimized near interesting regions. In the case of using less parameters than collocation points, the construction is overdetermined and finding solutions turns into an optimization problem, where one could use machine learning [41] techniques, such as gradient descent methods [42]. Since an exhaustive survey of all these possibilities is beyond the scope of this letter, we just limit the discussion to the simplest setting, which may then be seen as a starting point for improvements.

#### IV. EXAMPLE: SCALAR FIELD IN TWO DIMENSIONS

We illustrate the working of our general algorithm based on a scalar field theory in a two-dimensional Euclidean spacetime. The RG flow possesses a NGFP, corresponding to the Ising universality class. This fixed point is already seen in simple approximations and comes with a single UV-attractive eigendirection. For our purpose, it is then sufficient to work with the local potential approximation (LPA) [24]. In this case, the effective average action is approximated by

$$\Gamma_k[\phi] \simeq \int d^2x \left[ \frac{1}{2}(\partial\phi)^2 + U_k[\phi] \right], \quad (19)$$

where  $\phi(x)$  is a real scalar field with mass-dimension zero. Implementing our convention on the couplings, we expand the potential according to

$$U_k[\phi] = \frac{1}{4! \tilde{u}_k} \phi^4 + \sum_{0 < n, n \neq 2}^{N_{\max}} \frac{v_k^n}{(2n)! \tilde{u}_k} (\phi^2)^n. \quad (20)$$

This singles out the coupling  $\tilde{u}_k$  (taken with a negative mass dimension) associated with the  $\phi^4$ -term as the free parameter allowed by asymptotic safety. The IR-value of the dimensionless couplings  $v^n = (v^1, v^3, v^4, \dots)$  will be predicted based on the UV-critical surface of the NGFP.

In Table I we report results up to order  $N_{\max} = 10$ .<sup>4</sup> For simplicity, we limit our exposition to the simplest case where  $N_{\max} = 2$ , i.e., we are dealing with two couplings  $(u, v)$  and refer to the supplementary notebook for the detailed implementation of this case.

We start from the beta functions given in [23]. Adapting to our parameterization of the potential and setting  $N_{\max} = 2$  and  $d = 2$ , we have

$$\beta_u = 2u - \frac{3u^3}{2\pi(v+u)^3}, \quad \beta_v = -\frac{u^2(7v+u)}{4\pi(v+u)^3}. \quad (21)$$

The NGFP of the system is located at  $(u_*, v_*) = (\frac{343}{288\pi}, -\frac{49}{288\pi})$ . The stability coefficients and right-eigenvectors entering into the linearized solution (6) are

$$\theta_1 = \frac{1}{3}(1 + \sqrt{43}), \quad V_1 = \left( \frac{1}{21}(8 + \sqrt{43}), 1 \right)^T, \\ \tilde{\theta}_1 = \frac{1}{3}(1 - \sqrt{43}), \quad \tilde{V}_1 = \left( \frac{1}{21}(8 - \sqrt{43}), 1 \right)^T. \quad (22)$$

Thus we are dealing with a saddle-point with one UV-attractive and one UV-repulsive eigendirection. The UV-

<sup>4</sup> These results serve as an illustration of the method and should not be mistaken as an attempt of a precision computation. The latter may be achieved by extending the approximation along the lines of the derivative expansion used for the Wilson-Fisher fixed point [18, 43].

$N_{\max}$	$u_*$	$v_*^1$	$v_*^3$	$v_*^4$	$v_*^5$	$\theta_1$	$\tilde{\theta}_1$	$\tilde{\theta}_2$	$\tilde{\theta}_3$	$\tilde{\theta}_4$	$v^1$	$v^3$	$v^4$	$v^5$
4	0.3790	-0.0541				1.852	-2.519				0.15			
6	0.2765	-0.0746	16.3346			1.699	-2.875	-25.381			0.08	31		
8	0.2695	-0.0961	24.1984	941.6		1.573	-3.043	-22.469	-78.936		0.07	46	3209	
10	0.2809	-0.1174	28.1641	1561.8	110325	1.468	-3.134	-21.111	-66.552	-173.426	0.066	55	5538	701866

TABLE I. Properties of the NGFP associated with the Ising universality class obtained from the LPA, eq. (20), up to order  $N_{\max} = 10$ . The first and second block give the position and stability coefficients of the fixed point, respectively. The values  $v^\mu \equiv \lim_{k \rightarrow 0} v_k^\mu$  predicted by asymptotic safety are summarized in the third block.

critical surface is one-dimensional and we expect a prediction for  $v$  from asymptotic safety. Following the procedure leading to (11), the generating function for the UV-critical surface is

$$F^*(u, v) = \left(172 - 137\sqrt{43}\right) u + \left(44\sqrt{43} + 215\right) v + \frac{49(1003\sqrt{43} - 989)}{288\pi}. \quad (23)$$

Based on this input we compute the prediction for  $v$  using 1) the traditional shooting method, 2) a solution of the explicit system (14) based on series expansions of orders 10 to 15, and 3) a spectral solution evaluating the implicit equation (13) for a linear combination of 500 MCDs on a regular lattice centered on the NGFP. All methods yield

$$v \equiv \lim_{k \rightarrow 0} v_k \simeq 0.15, \quad (24)$$

with the numerical difference appearing in the third relevant digit. Fig. 2 illustrates the structures underlying this prediction.

## V. SUMMARY AND OUTLOOK

The Wilsonian renormalization group (RG) is a powerful tool to study the effect of fluctuations in statistical and quantum systems. Typically, such investigations proceed along the following steps. 1) one computes the beta functions of the theory in a suitable approximation. On this basis, one determines the RG fixed points and their stability properties. 2) In the case of asymptotic safety, where the UV-completion is provided by an interacting fixed point, one constructs the UV-critical surface of the fixed point in order to obtain the set of effective actions compatible with this UV-completion. 3) The output of step 2) is used to construct physical observables which allow to confront the predictions with observations.

Our work proposes an efficient algorithm for carrying out the second step in this procedure, by solving multi-linear partial differential equations for the generating functions encoding the embedding of the UV-critical surface. This constitutes a significant improvement compared to the usual shooting method (employed e.g., in




FIG. 2. Phase diagram generated by the beta functions (21) with arrows pointing towards lower values of  $k$ . The NGFP and the endpoint of  $\mathcal{S}_{UV}$  giving rise to the prediction (24) are marked by \*. The UV-critical surface obtained from the shooting method is marked by hollow green dots, the solution of the implicit method by brown dashes, and the one of the explicit method by a solid magenta line. For completeness we added the RG trajectories generated by  $\tilde{V}_1$  as solid orange dots.

[44–51]) which tracing individual RG trajectories. We illustrated the working of our algorithm based on the interacting fixed point corresponding to the Ising universality class. Interpreting this fixed point as the UV-completion of the RG flow, we predicted the couplings of the effective potential based on the single free parameter allowed by asymptotic safety (see Table I).

We stress that the methods described in this letter are applicable in a much broader context. Potential applications beyond the scope of the present work are the gravitational asymptotic safety program [22, 23], including its extension by matter degrees of freedom [8, 52], asymptotically safe gauge theories [53–57] and their su-

persymmetric extensions [58–62], and asymptotically safe quantum electrodynamics [63–65]. The authors have already tested its applicability for asymptotically safe scalar-tensor theories [66–83] and the resulting shapes of the effective potentials will be reported in [84].

## ACKNOWLEDGEMENTS

We thank M. Becker, C. Laporte, R. Loll, and J. Wang for useful discussions during the process of developing these ideas.

- 
- [1] K. G. Wilson and J. B. Kogut, The Renormalization group and the epsilon expansion, *Phys. Rept.* **12**, 75 (1974).
- [2] J. Berges, N. Tetradis, and C. Wetterich, Nonperturbative renormalization flow in quantum field theory and statistical physics, *Physics Reports* **363**, 223 (2002).
- [3] N. Dupuis, L. Canet, A. Eichhorn, W. Metzner, J. M. Pawłowski, M. Tissier, and N. Wschebor, The nonperturbative functional renormalization group and its applications, *Phys. Rept.* **910**, 1 (2021), arXiv:2006.04853 [cond-mat.stat-mech].
- [4] M. Reuter and F. Saueressig, Quantum Einstein Gravity, *New J. Phys.* **14**, 055022 (2012), arXiv:1202.2274 [hep-th].
- [5] J. Zinn-Justin, *Quantum field theory and critical phenomena*, Vol. 171 (Oxford university press, 2021).
- [6] S. Weinberg, Critical phenomena for field theorists, in *Understanding the fundamental constituents of matter* (Springer, 1978) pp. 1–52.
- [7] S. Weinberg, Ultraviolet divergences in quantum theories of gravitation, in *General relativity* (1979).
- [8] A. Pastor-Gutiérrez, J. M. Pawłowski, and M. Reichert, The Asymptotically Safe Standard Model: From quantum gravity to dynamical chiral symmetry breaking, *SciPost Phys.* **15**, 105 (2023), arXiv:2207.09817 [hep-th].
- [9] J. Boyd, *Chebyshev and Fourier Spectral Methods: Second Revised Edition*, Dover Books on Mathematics (Dover Publications, 2013).
- [10] J. Shen, T. Tang, and L.-L. Wang, *Spectral methods: algorithms, analysis and applications*, Vol. 41 (Springer Science & Business Media, 2011).
- [11] N. Tetradis and C. Wetterich, Critical exponents from effective average action, *Nucl. Phys. B* **422**, 541 (1994), arXiv:hep-ph/9308214.
- [12] T. R. Morris, On truncations of the exact renormalization group, *Phys. Lett. B* **334**, 355 (1994), arXiv:hep-th/9405190.
- [13] J. A. Adams, J. Berges, S. Bornholdt, F. Freire, N. Tetradis, and C. Wetterich, Solving nonperturbative flow equations, *Mod. Phys. Lett. A* **10**, 2367 (1995), arXiv:hep-th/9507093.
- [14] D. F. Litim, Critical exponents from optimized renormalization group flows, *Nucl. Phys. B* **631**, 128 (2002), arXiv:hep-th/0203006.
- [15] L. Canet, B. Delamotte, D. Mouhanna, and J. Vidal, Nonperturbative renormalization group approach to the Ising model: A Derivative expansion at order partial\*\*4, *Phys. Rev. B* **68**, 064421 (2003), arXiv:hep-th/0302227.
- [16] A. Jüttner, D. F. Litim, and E. Marchais, Global Wilson–Fisher fixed points, *Nucl. Phys. B* **921**, 769 (2017), arXiv:1701.05168 [hep-th].
- [17] G. De Polsi, I. Balog, M. Tissier, and N. Wschebor, Precision calculation of critical exponents in the  $O(N)$  universality classes with the nonperturbative renormalization group, *Phys. Rev. E* **101**, 042113 (2020), arXiv:2001.07525 [cond-mat.stat-mech].
- [18] I. Balog, H. Chaté, B. Delamotte, M. Marohnic, and N. Wschebor, Convergence of Nonperturbative Approximations to the Renormalization Group, *Phys. Rev. Lett.* **123**, 240604 (2019), arXiv:1907.01829 [cond-mat.stat-mech].
- [19] J. Berges and C. Wetterich, Equation of state and coarse grained free energy for matrix models, *Nuclear Physics B* **487**, 675 (1997).
- [20] A. Strumia, N. Tetradis, and C. Wetterich, The region of validity of homogeneous nucleation theory, *Physics Letters B* **467**, 279 (1999).
- [21] N. Tetradis and C. Wetterich, High temperature phase transitions without infrared divergences, *International Journal of Modern Physics A* **9**, 4029 (1994).
- [22] R. Percacci, *An introduction to covariant quantum gravity and asymptotic safety*, Vol. 3 (World Scientific, 2017).
- [23] M. Reuter and F. Saueressig, *Quantum gravity and the functional renormalization group: the road towards asymptotic safety* (Cambridge University Press, 2019).
- [24] C. Wetterich, Exact evolution equation for the effective potential, *Physics Letters B* **301**, 90 (1993).
- [25] T. R. Morris, The exact renormalization group and approximate solutions, *International Journal of Modern Physics A* **9**, 2411 (1994).
- [26] M. Reuter and C. Wetterich, Effective average action for gauge theories and exact evolution equations, *Nucl. Phys. B* **417**, 181 (1994).
- [27] M. Reuter, Nonperturbative evolution equation for quantum gravity, *Phys. Rev. D* **57**, 971 (1998), arXiv:hep-th/9605030.
- [28] J. Polchinski, Renormalization and effective lagrangians, *Nuclear Physics B* **231**, 269 (1984).
- [29] J. M. Pawłowski, Aspects of the functional renormalisation group, *Annals Phys.* **322**, 2831 (2007), arXiv:hep-th/0512261.
- [30] O. J. Rosten, Fundamentals of the Exact Renormalization Group, *Phys. Rept.* **511**, 177 (2012), arXiv:1003.1366 [hep-th].
- [31] F. Saueressig, The Functional Renormalization Group in Quantum Gravity (2023) arXiv:2302.14152 [hep-th].
- [32] D. Buccio and R. Percacci, Renormalization group flows between Gaussian fixed points, *JHEP* **10**, 113, arXiv:2207.10596 [hep-th].
- [33] B. Fornberg and D. M. Sloan, A review of pseudospectral methods for solving partial differential equations, *Acta numerica* **3**, 203 (1994).
- [34] M. Y. Hussaini, C. L. Streett, and T. A. Zang, *Spectral methods for partial differential equations*, Tech. Rep. (1983).
- [35] J. Borchardt and B. Knorr, Solving functional flow equations with pseudo-spectral methods, *Phys. Rev. D* **94**,

- 025027 (2016), arXiv:1603.06726 [hep-th].
- [36] J. Borchardt and B. Knorr, Global solutions of functional fixed point equations via pseudospectral methods, *Phys. Rev. D* **91**, 105011 (2015), [Erratum: *Phys.Rev.D* 93, 089904 (2016)], arXiv:1502.07511 [hep-th].
- [37] T. Draper, B. Knorr, C. Ripken, and F. Saueressig, Finite Quantum Gravity Amplitudes: No Strings Attached, *Phys. Rev. Lett.* **125**, 181301 (2020), arXiv:2007.00733 [hep-th].
- [38] J. Fehre, D. F. Litim, J. M. Pawłowski, and M. Reichert, Lorentzian Quantum Gravity and the Graviton Spectral Function, *Phys. Rev. Lett.* **130**, 081501 (2023), arXiv:2111.13232 [hep-th].
- [39] M. D. Buhmann, Radial basis functions, *Acta numerica* **9**, 1 (2000).
- [40] R. Pachón and L. N. Trefethen, Barycentric-remez algorithms for best polynomial approximation in the chebfun system, *BIT Numerical Mathematics* **49**, 721 (2009).
- [41] M. J. D. Powell, *Approximation theory and methods* (Cambridge university press, 1981).
- [42] L. Bottou, Large-scale machine learning with stochastic gradient descent, in *Proceedings of COMPSTAT'2010: 19th International Conference on Computational Statistics Paris France, August 22-27, 2010 Keynote, Invited and Contributed Papers* (Springer, 2010) pp. 177–186.
- [43] B. Delamotte, G. De Polsi, M. Tissier, and N. Wschebor, Conformal invariance and composite operators: A strategy for improving the derivative expansion of the nonperturbative renormalization group, (2024), arXiv:2401.02517 [cond-mat.stat-mech].
- [44] T. R. Morris, Three-dimensional massive scalar field theory and the derivative expansion of the renormalization group, *Nucl. Phys. B* **495**, 477 (1997), arXiv:hep-th/9612117.
- [45] C. Bervillier, B. Boisseau, and H. Giacomini, Analytical approximation schemes for solving exact renormalization group equations in the local potential approximation, *Nucl. Phys. B* **789**, 525 (2008), arXiv:0706.0990 [hep-th].
- [46] M. Reuter and H. Weyer, Renormalization group improved gravitational actions: A Brans-Dicke approach, *Phys. Rev. D* **69**, 104022 (2004), arXiv:hep-th/0311196.
- [47] M. Shaposhnikov, Asymptotic safety of gravity and the Higgs-boson mass, *Theor. Math. Phys.* **170**, 229 (2012).
- [48] A. Eichhorn and A. Held, Top mass from asymptotic safety, *Phys. Lett. B* **777**, 217 (2018), arXiv:1707.01107 [hep-th].
- [49] G. Gubitosi, R. Ooijer, C. Ripken, and F. Saueressig, Consistent early and late time cosmology from the RG flow of gravity, *JCAP* **12**, 004, arXiv:1806.10147 [hep-th].
- [50] I. Basile and A. Platania, Asymptotic Safety: Swamp-land or Wonderland?, *Universe* **7**, 389 (2021), arXiv:2107.06897 [hep-th].
- [51] A. Baldazzi, K. Falls, Y. Kluth, and B. Knorr, Robustness of the derivative expansion in Asymptotic Safety, (2023), arXiv:2312.03831 [hep-th].
- [52] A. Eichhorn and M. Schiffer, Asymptotic safety of gravity with matter, (2022), arXiv:2212.07456 [hep-th].
- [53] D. F. Litim and F. Sannino, Asymptotic safety guaranteed, *JHEP* **12**, 178, arXiv:1406.2337 [hep-th].
- [54] A. D. Bond, G. Hiller, K. Kowalska, and D. F. Litim, Directions for model building from asymptotic safety, *JHEP* **08**, 004, arXiv:1702.01727 [hep-ph].
- [55] G. Hiller, C. Hormigos-Feliu, D. F. Litim, and T. Steudtner, Model Building from Asymptotic Safety with Higgs and Flavor Portals, *Phys. Rev. D* **102**, 095023 (2020), arXiv:2008.08606 [hep-ph].
- [56] A. D. Bond and D. F. Litim, Asymptotic safety guaranteed for strongly coupled gauge theories, *Phys. Rev. D* **105**, 105005 (2022), arXiv:2202.08223 [hep-th].
- [57] D. F. Litim, N. Riyaz, E. Stamou, and T. Steudtner, Asymptotic safety guaranteed at four-loop order, *Phys. Rev. D* **108**, 076006 (2023), arXiv:2307.08747 [hep-th].
- [58] K. Intriligator and F. Sannino, Supersymmetric asymptotic safety is not guaranteed, *JHEP* **11**, 023, arXiv:1508.07411 [hep-th].
- [59] B. Bajc and F. Sannino, Asymptotically Safe Grand Unification, *JHEP* **12**, 141, arXiv:1610.09681 [hep-th].
- [60] B. Bajc, N. A. Dondi, and F. Sannino, Safe SUSY, *JHEP* **03**, 005, arXiv:1709.07436 [hep-th].
- [61] A. D. Bond and D. F. Litim, Asymptotic safety guaranteed in supersymmetry, *Phys. Rev. Lett.* **119**, 211601 (2017), arXiv:1709.06953 [hep-th].
- [62] G. Hiller, D. F. Litim, and K. Moch, Fixed points in supersymmetric extensions of the standard model, *Eur. Phys. J. C* **82**, 952 (2022), arXiv:2202.01264 [hep-ph].
- [63] H. Gies and J. Ziebell, Asymptotically Safe QED, *Eur. Phys. J. C* **80**, 607 (2020), arXiv:2005.07586 [hep-th].
- [64] H. Gies, K. K. K. Tam, and J. Ziebell, Pauli-term-induced fixed points in d-dimensional QED, *Eur. Phys. J. C* **83**, 955 (2023), arXiv:2210.11927 [hep-th].
- [65] H. Gies and K. K. K. Tam, Fermionic Fixed-Point Structure of Asymptotically Safe QED with a Pauli Term, (2024), arXiv:2403.02980 [hep-th].
- [66] G. Narain and R. Percacci, Renormalization Group Flow in Scalar-Tensor Theories. I, *Class. Quant. Grav.* **27**, 075001 (2010), arXiv:0911.0386 [hep-th].
- [67] A. Eichhorn, Quantum-gravity-induced matter self-interactions in the asymptotic-safety scenario, *Phys. Rev. D* **86**, 105021 (2012), arXiv:1204.0965 [gr-qc].
- [68] T. Henz, J. M. Pawłowski, A. Rodigast, and C. Wetterich, Dilaton Quantum Gravity, *Phys. Lett. B* **727**, 298 (2013), arXiv:1304.7743 [hep-th].
- [69] R. Percacci and G. P. Vacca, Search of scaling solutions in scalar-tensor gravity, *Eur. Phys. J. C* **75**, 188 (2015), arXiv:1501.00888 [hep-th].
- [70] P. Labus, R. Percacci, and G. P. Vacca, Asymptotic safety in  $O(N)$  scalar models coupled to gravity, *Phys. Lett. B* **753**, 274 (2016), arXiv:1505.05393 [hep-th].
- [71] T. Henz, J. M. Pawłowski, and C. Wetterich, Scaling solutions for Dilaton Quantum Gravity, *Phys. Lett. B* **769**, 105 (2017), arXiv:1605.01858 [hep-th].
- [72] D. Becker, C. Ripken, and F. Saueressig, On avoiding Ostrogradski instabilities within Asymptotic Safety, *JHEP* **12**, 121, arXiv:1709.09098 [hep-th].
- [73] A. Eichhorn, Y. Hamada, J. Lumma, and M. Yamada, Quantum gravity fluctuations flatten the Planck-scale Higgs potential, *Phys. Rev. D* **97**, 086004 (2018), arXiv:1712.00319 [hep-th].
- [74] J. M. Pawłowski, M. Reichert, C. Wetterich, and M. Yamada, Higgs scalar potential in asymptotically safe quantum gravity, *Phys. Rev. D* **99**, 086010 (2019), arXiv:1811.11706 [hep-th].
- [75] C. Wetterich, Effective scalar potential in asymptotically safe quantum gravity, *Universe* **7**, 45 (2021), arXiv:1911.06100 [hep-th].

- [76] C. Wetterich and M. Yamada, Variable Planck mass from the gauge invariant flow equation, *Phys. Rev. D* **100**, 066017 (2019), arXiv:1906.01721 [hep-th].
- [77] A. Eichhorn and M. Pauly, Constraining power of asymptotic safety for scalar fields, *Phys. Rev. D* **103**, 026006 (2021), arXiv:2009.13543 [hep-th].
- [78] N. Ohta and M. Yamada, Higgs scalar potential coupled to gravity in the exponential parametrization in arbitrary gauge, *Phys. Rev. D* **105**, 10.1103/PhysRevD.105.026013 (2022), arXiv:2110.08594 [hep-th].
- [79] C. Laporte, A. D. Pereira, F. Saueressig, and J. Wang, Scalar-tensor theories within Asymptotic Safety, *JHEP* **12**, 001, arXiv:2110.09566 [hep-th].
- [80] C. Laporte, N. Locht, A. D. Pereira, and F. Saueressig, Evidence for a novel shift-symmetric universality class from the functional renormalization group, *Phys. Lett. B* **838**, 137666 (2023), arXiv:2207.06749 [hep-th].
- [81] B. Knorr, Safe essential scalar-tensor theories, (2022), arXiv:2204.08564 [hep-th].
- [82] C. Wetterich, Quantum Gravity and Scale Symmetry in Cosmology (2023) arXiv:2211.03596 [gr-qc].
- [83] G. P. de Brito, B. Knorr, and M. Schiffer, On the weak-gravity bound for a shift-symmetric scalar field, *Phys. Rev. D* **108**, 026004 (2023), arXiv:2302.10989 [hep-th].
- [84] A. Silva, Emergence of inflaton potential from asymptotically safe gravity, In preparation.