

Generating functional of correlators of twist-2 operators in $\mathcal{N} = 1$ SUSY Yang-Mills theory, I

Marco Bochicchio,^a Mauro Papinutto,^{b,a} Francesco Scardino^{b,a}

^a*Physics Department, INFN Roma1,
Piazzale A. Moro 2, Roma, I-00185, Italy*

^b*Physics Department, Sapienza University,
Piazzale A. Moro 2, Roma, I-00185, Italy*

E-mail: marco.bochicchio@roma1.infn.it, mauro.papinutto@roma1.infn.it,
francesco.scardino@roma1.infn.it

ABSTRACT: Extending our previous work in pure Yang-Mills (YM) theory, we compute the generating functional of correlators of collinear twist-2 operators that enter the components of balanced superfields – i.e., superfields with an equal number of dotted and undotted indices in their spinor representation – in $\mathcal{N} = 1$ SUSY $SU(N)$ YM theory in Minkowskian and Euclidean space-time, in the conformal limit and renormalization-group improved form, and to the leading and next-to-leading order in the large- N expansion. The latter calculation sets strong UV asymptotic constraints on the nonperturbative solution of large- N $\mathcal{N} = 1$ SUSY YM theory that may be a pivotal guide for the search of such a solution.

Contents

1	Introduction, physics motivations and conclusions	1
2	Plan of the paper	2
3	$\mathcal{N} = 1$ SUSY YM theory in the light-cone gauge	2
4	Twist-2 balanced superfields in $\mathcal{N} = 1$ SUSY YM theory	4
4.1	Collinear twist-2 operators	4
4.2	Balanced superfields	7
5	Generating functional of Minkowskian conformal correlators	7
5.1	Connected generating functional $\mathcal{W}_{\text{conf}}$ as the log of a superdeterminant of a quadratic form	8
5.2	Connected generating functional Γ_{conf} as the log a Fredholm superdeterminant	14
5.3	Connected generating functional in the momentum representation	15
6	Generating functional of Euclidean conformal correlators	17
6.1	Analytic continuation to Euclidean space-time	17
6.2	Analytic continuation of $\mathcal{W}_{\text{conf}}$	19
6.3	Analytic continuation of Γ_{conf}	21
7	RG-improvement of the Euclidean generating functional	22
7.1	Operator mixing	22
7.2	Anomalous dimensions of twist-2 operators in $\mathcal{N} = 1$ SUSY YM theory	24
7.3	Conformal generating functional of correlators of supermultiplet operators	25
7.4	RG-improved generating functional of correlators of supermultiplet operators	27
8	Minkowskian conformal correlators	29
8.1	O^A and \tilde{O}^A correlators	30
8.2	O^λ and \tilde{O}^λ correlators	30
8.3	M and \bar{M} correlators	34
8.4	Vanishing correlators	40
9	Euclidean conformal correlators	40
9.1	O^{AE} correlators	40
9.2	$O^{\lambda E}$ correlators	41
9.3	M^E and \bar{M}^E correlators	41
A	Notation and Wick rotation	42
B	Jacobi and Gegenbauer polynomials	43

C Normalization of 2-point correlators	45
C.1 2-point correlators of gluino-gluino operators	45
C.2 2-point correlators of gluon-gluino operators	47
D An equality of determinants	49
E Conformal properties of the standard basis	51
F Nonresonant diagonal renormalization scheme	53

1 Introduction, physics motivations and conclusions

Recently, we have computed [1] the ultraviolet (UV) asymptotics of the generating functional of correlators of collinear twist-2 operators in pure $SU(N)$ Yang-Mills (YM) theory.

The present paper is the first of two installments, where we extend the above computation to $\mathcal{N} = 1$ supersymmetric (SUSY) $SU(N)$ YM theory for twist-2 operators that are the components of balanced superfields – in the present paper – and the components of unbalanced ones – in the second installment. We refer to operators – and superfields as well – as either balanced or unbalanced [1, 2] if in their spinor representation they carry either an equal or a different number of dotted and undotted indices, respectively. Actually, our computation involves twist-2 operators that have maximal-spin component along the light-cone direction p_+ in Minkowskian space-time – which we refer to as collinear twist-2 operators [1, 2] – and their analytic continuation to Euclidean space-time, and it is based on a number of innovations [1, 2] described below.

Indeed, we have computed in YM theory by Feynman diagrams the Minkowskian conformal correlators to the lowest order of perturbation theory and reconstructed from them the corresponding generating functional in the balanced and unbalanced sectors separately [2]. We have also computed in a much simpler way the complete generating functional of conformal correlators by functional-integral methods [1] and found perfect agreement with our previous computation [2], the equivalence occurring thanks to a dictionary established by means of a tedious but straightforward calculation [1]. Moreover, we have lifted the generating functional of the Euclidean conformal correlators to the generating functional of the UV asymptotic correlators [1] by means of the renormalization-group (RG) improvement and a careful choice of the renormalization scheme – dubbed nonresonant diagonal according to the Poincaré-Dulac theorem [3] – that involves a novel geometric interpretation of operator mixing [3], where the collinear twist-2 operators become multiplicatively renormalizable to all orders of perturbation theory [1]. Remarkably, the RG-improved correlators should be asymptotic to the exact correlators thanks to the asymptotic freedom (AF) [1, 4]. Finally, we have expanded the latter generating functional to the leading and next-to-leading order in the large- N expansion.

All the above techniques and results extend straightforwardly to the supersymmetric case, as we briefly recall in the present paper, the only difference being that the final

result for the generating functional of connected correlators in the SUSY YM case has the structure of the logarithm of a functional superdeterminant, instead of a determinant as in the pure YM case. Perturbatively, the logarithm of a (super)determinant occurs because in the light-cone gauge collinear twist-2 operators are quadratic in the fundamental fields and thus their generating functional to the lowest order arises from a Gaussian integral [1] – also involving anticommuting variables in the SUSY case. The above structure of the logarithm of a (super)determinant is then inherited by the RG-improved generating functional as well. Nonperturbatively, the logarithm of a (super)determinant should arise to the next-to-leading order in the large- N expansion from the glueball(gluinoball) one-loop effective action, as we have already discussed at length in the pure YM case [1, 4], which our leading-nonplanar [1, 4] RG-improved generating functional should be UV asymptotic to because of the AF. Therefore, our computation of the UV asymptotic generating functional in the large- N expansion sets strong constraints on the nonperturbative solution of large- N $\mathcal{N} = 1$ SUSY YM theory and it may be a pivotal guide for the search of such a solution.

2 Plan of the paper

In section 3 we recall the definition of $\mathcal{N} = 1$ SUSY YM theory in the light-cone gauge.

In section 4 we define the twist-2 operators that enter the components of balanced superfields.

In section 5 we evaluate to the lowest perturbative order the corresponding Minkowskian conformal generating functional as a superdeterminant of a quadratic form. We also rewrite it as a Fredholm superdeterminant employing the aforementioned dictionary, and work out the generating functional in the momentum representation.

In section 6 we analytic continue the Minkowskian conformal generating functional to Euclidean space-time.

In section 7 we work out the generating functional of Euclidean RG-improved correlators.

In section 8, as a further application, we explicitly compute the Minkowskian conformal correlators in several sectors by means of the generating functional.

In section 9 we compute the corresponding Euclidean conformal correlators.

In the appendices we report ancillary computations.

3 $\mathcal{N} = 1$ SUSY YM theory in the light-cone gauge

The action of $\mathcal{N} = 1$ SUSY $SU(N)$ YM theory in Mikowskian space-time reads [5, 6]:

$$S = \int -\frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu} + \frac{i}{2} \bar{\chi}^a \gamma^\mu (D_\mu \chi)^a d^4x \quad (3.1)$$

with $a = 1, \dots, N^2 - 1$, where χ is in the adjoint representation of the Lie algebra of $SU(N)$ and a Majorana spinor $\chi = C\bar{\chi}^T$, with C the charge conjugation, and:

$$\begin{aligned} F_{\mu\nu} &= \partial_\mu A_\nu - \partial_\nu A_\mu + i \frac{g}{\sqrt{N}} [A_\mu, A_\nu] \\ D_\mu \chi &= \partial_\mu \chi + i \frac{g}{\sqrt{N}} [A_\mu, \chi] \end{aligned} \quad (3.2)$$

in matrix notation, with $g^2 = g_{YM}^2 N$ the 't Hooft coupling [7]. The action is invariant under the SUSY transformations [5, 8]:

$$\begin{aligned} \delta A_\mu^a &= -i\bar{\zeta}\gamma_\mu\chi^a \\ \delta\chi^a &= \frac{i}{2}\sigma^{\mu\nu}F_{\mu\nu}^a\zeta \end{aligned} \quad (3.3)$$

with $\sigma^{\mu\nu} = \frac{i}{2}[\gamma^\mu, \gamma^\nu]$ and ζ a Majorana spinor. A Majorana spinor can be decomposed into two complex-conjugate Weyl spinors, λ_α and $\bar{\lambda}_{\dot{\alpha}}$:

$$\chi = \begin{pmatrix} \lambda_\alpha \\ \bar{\lambda}_{\dot{\alpha}} \end{pmatrix} \quad \bar{\chi} = \chi^\dagger \gamma^0 = \begin{pmatrix} \lambda^\alpha \bar{\lambda}_{\dot{\alpha}} \end{pmatrix} \quad (3.4)$$

Correspondingly, the action reads:

$$S = - \int \frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu} + i\bar{\lambda}^a \bar{\lambda}^{\dot{\alpha}} \sigma_{\alpha\dot{\alpha}}^\mu (D_\mu \lambda^\alpha)^a d^4x \quad (3.5)$$

The fundamental fields, A_μ and χ , respectively interpolate for the gluon and gluino. In the light-cone gauge $A_+ = 0$,

$$\chi_- = \Pi_- \chi = \begin{pmatrix} 0 \\ \lambda_2 \\ \bar{\lambda}_2 \\ 0 \end{pmatrix} \quad (3.6)$$

and A_- can be integrated out, with the projectors $\Pi_\pm = \frac{1}{2}\gamma^\mp\gamma^\pm$. The corresponding action is expressed in terms of the physical fields only – the transverse components of the gluon, A and \bar{A} :

$$\begin{aligned} A &\equiv \frac{1}{\sqrt{2}}(A_1 + iA_2) \\ \bar{A} &\equiv \frac{1}{\sqrt{2}}(A_1 - iA_2) \end{aligned} \quad (3.7)$$

and the plus component of the gluino, χ_+ :

$$\chi_+ = \Pi_+ \chi = \begin{pmatrix} \lambda_1 \\ 0 \\ 0 \\ -\bar{\lambda}_1 \end{pmatrix} \quad (3.8)$$

For simplicity we introduce two anticommuting fields, λ and $\bar{\lambda}$, so that their normalization:

$$\lambda \equiv \frac{1}{2^{\frac{1}{4}}}\lambda_1 \quad \bar{\lambda} \equiv \frac{1}{2^{\frac{1}{4}}}\bar{\lambda}_1 \quad (3.9)$$

matches the canonical normalization of the kinetic term of the action in the light-cone gauge [6]:

$$\begin{aligned} S(A, \bar{A}, \lambda, \bar{\lambda}) = & \int -\bar{A}^a \square A^a - i\bar{\lambda}^a \square \partial_+^{-1} \lambda^a \\ & - 2\frac{g}{\sqrt{N}} f^{abc} (A^a \partial_+ \bar{A}^b \bar{\partial} \partial_+^{-1} A^c + \bar{A}^a \partial_+ A^b \partial \partial_+^{-1} \bar{A}^c) - 2\frac{g^2}{N} f^{abc} f^{ade} \partial_+^{-1} (A^b \partial_+ \bar{A}^c) \partial_+^{-1} (\bar{A}^d \partial_+ A^e) \\ & - 2i\frac{g}{\sqrt{N}} f^{abc} \bar{\lambda}^a \lambda^b (\bar{\partial} A^c + \partial \bar{A}^c) - 2i\frac{g^2}{N} f^{abe} f^{cde} \partial_+^{-1} (\bar{A}^a \partial_+ A^b + A^a \partial_+ \bar{A}^b) \partial_+^{-1} (\bar{\lambda}^c \lambda^d) \\ & + 2\frac{g^2}{N} f^{abe} f^{cde} \partial_+^{-1} (\bar{\lambda}^a \lambda^b) \partial_+^{-1} (\bar{\lambda}^c \lambda^d) + 2i\frac{g}{\sqrt{N}} f^{abc} \bar{A}^a \bar{\lambda}^b \partial_+^{-1} \partial \lambda^c + 2i\frac{g}{\sqrt{N}} f^{abc} A^a \lambda^b \partial_+^{-1} \bar{\partial} \bar{\lambda}^c \\ & - 2i\frac{g^2}{N} f^{abe} f^{cde} \bar{A}^a \bar{\lambda}^b \partial_+^{-1} (A^c \lambda^d) d^4 x \end{aligned} \quad (3.10)$$

with:

$$\square = g^{\mu\nu} \partial_\mu \partial_\nu = \partial_0^2 - \sum_{i=1}^3 \partial_i^2 \quad (3.11)$$

where we employ the mostly minus metric $g_{\mu\nu}$ in Minkowskian space-time (appendix A). The v.e.v. of a product of local gauge-invariant operators \mathcal{O}_i that do not depend on A_- and χ_- reads:

$$\langle \mathcal{O}_1(x_1) \dots \mathcal{O}_n(x_n) \rangle = \frac{1}{Z} \int \mathcal{D}A \mathcal{D}\bar{A} \mathcal{D}\lambda \mathcal{D}\bar{\lambda} e^{iS(A, \bar{A}, \lambda, \bar{\lambda})} \mathcal{O}_1(x_1) \dots \mathcal{O}_n(x_n) \quad (3.12)$$

To the leading perturbative order the above v.e.v. reduces to:

$$\langle \mathcal{O}_1(x_1) \dots \mathcal{O}_n(x_n) \rangle = \frac{1}{Z} \int \mathcal{D}A \mathcal{D}\bar{A} \mathcal{D}\lambda \mathcal{D}\bar{\lambda} e^{\int d^4x (-i\bar{A}^a \square A^a + \bar{\lambda}^a \square \partial_+^{-1} \lambda^a)} \mathcal{O}_1(x_1) \dots \mathcal{O}_n(x_n) \quad (3.13)$$

We obtain for the corresponding conformal generating functional to the leading order:

$$\mathcal{Z}_{\text{conf}}[J_{\mathcal{O}}] = \frac{1}{Z} \int \mathcal{D}A \mathcal{D}\bar{A} \mathcal{D}\lambda \mathcal{D}\bar{\lambda} e^{\int d^4x (-i\bar{A}^a \square A^a + \bar{\lambda}^a \square \partial_+^{-1} \lambda^a)} \exp \left(\int d^4x \sum_i J_{\mathcal{O}_i} \mathcal{O}_i \right) \quad (3.14)$$

that is the supersymmetric generalization of the conformal generating functional in YM theory [1].

4 Twist-2 balanced superfields in $\mathcal{N} = 1$ SUSY YM theory

4.1 Collinear twist-2 operators

We list the gauge-invariant collinear twist-2 operators that enter the components of balanced superfields – to be defined below – in $\mathcal{N} = 1$ SUSY YM theory [6, 8]:

- gluon-gluon operators

$$O_s^A = \frac{1}{2} \partial_+ \bar{A}^a (i \vec{\partial}_+ + i \overleftarrow{\partial}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left(\frac{\vec{\partial}_+ - \overleftarrow{\partial}_+}{\vec{\partial}_+ + \overleftarrow{\partial}_+} \right) \partial_+ A^a \quad s = 2, 4, 6, \dots$$

$$\equiv \frac{1}{2} \bar{A}^a \mathcal{Y}_{s-2}^{\frac{5}{2}} (\vec{\partial}_+, \overleftarrow{\partial}_+) A^a \quad (4.1)$$

$$\tilde{O}_s^A = \frac{1}{2} \partial_+ \bar{A}^a (i \vec{\partial}_+ + i \overleftarrow{\partial}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left(\frac{\vec{\partial}_+ - \overleftarrow{\partial}_+}{\vec{\partial}_+ + \overleftarrow{\partial}_+} \right) \partial_+ A^a \quad s = 3, 5, 7, \dots$$

$$\equiv \frac{1}{2} \bar{A}^a \mathcal{H}_{s-2}^{\frac{5}{2}} (\vec{\partial}_+, \overleftarrow{\partial}_+) A^a \quad (4.2)$$

with $PT = (-1)^s$ and $C = +1$, where P is parity, T time-reversal and C charge conjugation.

- gluino-gluino operators

$$O_s^\lambda = \frac{1}{2} \bar{\lambda}^a (i \vec{\partial}_+ + i \overleftarrow{\partial}_+)^{s-1} C_{s-1}^{\frac{3}{2}} \left(\frac{\vec{\partial}_+ - \overleftarrow{\partial}_+}{\vec{\partial}_+ + \overleftarrow{\partial}_+} \right) \lambda^a \quad s = 2, 4, 6, \dots$$

$$\equiv \frac{1}{2} \bar{\lambda}^a \mathcal{Y}_{s-1}^{\frac{3}{2}} (\vec{\partial}_+, \overleftarrow{\partial}_+) \lambda^a \quad (4.3)$$

$$\tilde{O}_s^\lambda = \frac{1}{2} \bar{\lambda}^a (i \vec{\partial}_+ + i \overleftarrow{\partial}_+)^{s-1} C_{s-1}^{\frac{3}{2}} \left(\frac{\vec{\partial}_+ - \overleftarrow{\partial}_+}{\vec{\partial}_+ + \overleftarrow{\partial}_+} \right) \lambda^a \quad s = 1, 3, 5, \dots$$

$$\equiv \frac{1}{2} \bar{\lambda}^a \mathcal{H}_{s-1}^{\frac{3}{2}} (\vec{\partial}_+, \overleftarrow{\partial}_+) \lambda^a \quad (4.4)$$

with $PT = (-1)^s$ and $C = +1$.

- gluon-gluino operators

$$M_s = \frac{1}{2} \partial_+ A^a (i \vec{\partial}_+ + i \overleftarrow{\partial}_+)^{s-1} P_{s-1}^{(2,1)} \left(\frac{\vec{\partial}_+ - \overleftarrow{\partial}_+}{\vec{\partial}_+ + \overleftarrow{\partial}_+} \right) \lambda^a \quad s = 1, 2, 3, \dots$$

$$\equiv \frac{1}{2} A^a \mathcal{G}_{s-1}^{(2,1)} (\vec{\partial}_+, \overleftarrow{\partial}_+) \lambda^a = \frac{1}{2} (-1)^{s-1} \lambda^a \mathcal{G}_{s-1}^{(1,2)} (\vec{\partial}_+, \overleftarrow{\partial}_+) A^a \quad (4.5)$$

$$\bar{M}_s = \frac{1}{2} \bar{\lambda}^a (i \vec{\partial}_+ + i \overleftarrow{\partial}_+)^{s-1} P_{s-1}^{(1,2)} \left(\frac{\vec{\partial}_+ - \overleftarrow{\partial}_+}{\vec{\partial}_+ + \overleftarrow{\partial}_+} \right) \partial_+ \bar{A}^a \quad s = 1, 2, 3, \dots$$

$$\equiv \frac{1}{2} \bar{\lambda}^a \mathcal{G}_{s-1}^{(1,2)} (\vec{\partial}_+, \overleftarrow{\partial}_+) \bar{A}^a = \frac{1}{2} (-1)^{s-1} \bar{A}^a \mathcal{G}_{s-1}^{(2,1)} (\vec{\partial}_+, \overleftarrow{\partial}_+) \bar{\lambda}^a \quad (4.6)$$

where C_n^α are Gegenbauer polynomials, $P_n^{(\alpha, \beta)}$ Jacobi polynomials (appendix B) and we have defined:

$$\begin{aligned}
\mathcal{Y}_{s-2}^{\frac{5}{2}}(\vec{\partial}_+, \overleftarrow{\partial}_+) &= \overleftarrow{\partial}_+ (i\vec{\partial}_+ + i\overleftarrow{\partial}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left(\frac{\vec{\partial}_+ - \overleftarrow{\partial}_+}{\vec{\partial}_+ + \overleftarrow{\partial}_+} \right) \vec{\partial}_+ \\
&= \frac{\Gamma(3)\Gamma(s+3)}{\Gamma(5)\Gamma(s+1)} i^{s-2} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-1)^{s-k} \overleftarrow{\partial}_+^{s-k-1} \vec{\partial}_+^{k+1} \\
\mathcal{Y}_{s-1}^{\frac{3}{2}}(\vec{\partial}_+, \overleftarrow{\partial}_+) &= (i\vec{\partial}_+ + i\overleftarrow{\partial}_+)^{s-2} C_{s-1}^{\frac{3}{2}} \left(\frac{\vec{\partial}_+ - \overleftarrow{\partial}_+}{\vec{\partial}_+ + \overleftarrow{\partial}_+} \right) \\
&= \frac{(s+1)}{2} i^{s-1} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-1)^{s-k-1} \overleftarrow{\partial}_+^{s-k-1} \vec{\partial}_+^k
\end{aligned} \tag{4.7}$$

for even s , and:

$$\begin{aligned}
\mathcal{H}_{s-2}^{\frac{5}{2}}(\vec{\partial}_+, \overleftarrow{\partial}_+) &= \overleftarrow{\partial}_+ (i\vec{\partial}_+ + i\overleftarrow{\partial}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left(\frac{\vec{\partial}_+ - \overleftarrow{\partial}_+}{\vec{\partial}_+ + \overleftarrow{\partial}_+} \right) \vec{\partial}_+ \\
&= \frac{\Gamma(3)\Gamma(s+3)}{\Gamma(5)\Gamma(s+1)} i^{s-2} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-1)^{s-k} \overleftarrow{\partial}_+^{s-k-1} \vec{\partial}_+^{k+1} \\
\mathcal{H}_{s-1}^{\frac{3}{2}}(\vec{\partial}_+, \overleftarrow{\partial}_+) &= (i\vec{\partial}_+ + i\overleftarrow{\partial}_+)^{s-2} C_{s-1}^{\frac{3}{2}} \left(\frac{\vec{\partial}_+ - \overleftarrow{\partial}_+}{\vec{\partial}_+ + \overleftarrow{\partial}_+} \right) \\
&= \frac{(s+1)}{2} i^{s-1} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-1)^{s-k-1} \overleftarrow{\partial}_+^{s-k-1} \vec{\partial}_+^k
\end{aligned} \tag{4.8}$$

for odd s , and:

$$\begin{aligned}
\mathcal{G}_{s-1}^{(1,2)}(\vec{\partial}_+, \overleftarrow{\partial}_+) &= (i\vec{\partial}_+ + i\overleftarrow{\partial}_+)^{s-1} P_{s-1}^{(1,2)} \left(\frac{\vec{\partial}_+ - \overleftarrow{\partial}_+}{\vec{\partial}_+ + \overleftarrow{\partial}_+} \right) \vec{\partial}_+ \\
&= i^{s-1} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s+1}{k+2} (-1)^{s-k-1} \overleftarrow{\partial}_+^{s-k-1} \vec{\partial}_+^{k+1} \\
\mathcal{G}_{s-1}^{(2,1)}(\vec{\partial}_+, \overleftarrow{\partial}_+) &= \overleftarrow{\partial}_+ (i\vec{\partial}_+ + i\overleftarrow{\partial}_+)^{s-1} P_{s-1}^{(2,1)} \left(\frac{\vec{\partial}_+ - \overleftarrow{\partial}_+}{\vec{\partial}_+ + \overleftarrow{\partial}_+} \right) \\
&= i^{s-1} \sum_{k=0}^{s-1} \binom{s+1}{k} \binom{s}{k+1} (-1)^{s-k-1} \overleftarrow{\partial}_+^{s-k} \vec{\partial}_+^k
\end{aligned} \tag{4.9}$$

The spin projection on the p_+ direction is s for O_s^A, \tilde{O}_s^A and $O_s^\lambda, \tilde{O}_s^\lambda$, while it is $s + \frac{1}{2}$ for M_s and \bar{M}_s . The bosonic operators O_s^A, \tilde{O}_s^A and $O_s^\lambda, \tilde{O}_s^\lambda$ are Hermitian. The fermionic operators M_s and \bar{M}_s are Hermitian conjugate of each other.

4.2 Balanced superfields

Suitable linear combinations of the above operators form irreducible representations of the superalgebra of SUSY transformations restricted to the light-cone [8, 9]:

$$\begin{aligned}\delta A &= -2i\bar{\lambda}\zeta \\ \delta \bar{A} &= 2i\bar{\zeta}\lambda \\ \delta \lambda &= 2\zeta\partial_+\bar{A} \\ \delta \bar{\lambda} &= 2\bar{\zeta}\partial_+A\end{aligned}\tag{4.10}$$

where ζ is Majorana spinor satisfying $\Pi_+\zeta = 0$. Indeed, since the restricted SUSY transformations are linear, the set of the above operators is closed under their action. The operators that form supermultiplets – i.e., the aforementioned representations – read as follows [6, 9].

For even spin with $s \geq 2$:

$$\begin{aligned}S_s^{(1)} &= \frac{6}{s-1}O_s^A - O_s^\lambda \\ S_s^{(2)} &= \frac{6}{s+2}O_s^A + O_s^\lambda\end{aligned}\tag{4.11}$$

and for odd spin with $s \geq 3$:

$$\begin{aligned}\tilde{S}_s^{(1)} &= -\frac{6}{s-1}\tilde{O}_s^A - \tilde{O}_s^\lambda \\ \tilde{S}_s^{(2)} &= -\frac{6}{s+2}\tilde{O}_s^A + \tilde{O}_s^\lambda\end{aligned}\tag{4.12}$$

where \tilde{O}_s^A is not defined for $s = 1$, while \tilde{O}_1^λ it is, and $\tilde{S}_1^{(2)} = \tilde{O}_1^\lambda$. The above operators also diagonalize the matrix of anomalous dimension to order g^2 , where $\mathcal{N} = 1$ SUSY YM theory is actually conformal invariant in the conformal scheme (appendix E). The components of the above supermultiplets enter the balanced superfields ¹:

$$\mathbb{W}_s(x, \theta, \bar{\theta}) \sim \mathbb{S}_{s+1}^{(2)} + \theta\bar{M}_{s+1} + \bar{\theta}M_{s+1} + \theta\bar{\theta}\mathbb{S}_{s+2}^{(1)}\tag{4.13}$$

where $\mathbb{S}^{(i)} = \{S^{(i)}, \tilde{S}^{(i)}\}$ include both even- and odd-spin operators.

5 Generating functional of Minkowskian conformal correlators

The corresponding generating functional reads to the lowest perturbative order:

$$\begin{aligned}\mathcal{Z}_{\text{conf}}[J_{O^A}, J_{\tilde{O}^A}, J_{O^\lambda}, J_{\tilde{O}^\lambda}, \bar{J}_M, J_{\bar{M}}] &= \frac{1}{Z} \int \mathcal{D}A \mathcal{D}\bar{A} \mathcal{D}\lambda \mathcal{D}\bar{\lambda} e^{-\int d^4x i\bar{A}^a \square A^a - \bar{\lambda}^a \square \partial_+^{-1} \lambda^a} \\ &\exp \left(\int d^4x \sum_s O_s^A J_{O_s^A} + \tilde{O}_s^A J_{\tilde{O}_s^A} + O_s^\lambda J_{O_s^\lambda} + \tilde{O}_s^\lambda J_{\tilde{O}_s^\lambda} + \bar{J}_M M_s + \bar{M}_s J_{\bar{M}_s} \right)\end{aligned}\tag{5.1}$$

¹G. Santoni and F. Scardino, to appear on arXiv.

The sources $J_{O_s^A}$, $J_{\tilde{O}_s^A}$, $J_{O_s^\lambda}$ and $J_{\tilde{O}_s^\lambda}$ are bosonic, while $J_{\bar{M}_s}$ and \bar{J}_{M_s} are fermionic, so that we make the formal association:

$$\begin{aligned} \frac{\delta}{\delta J_{O_s^A}(x)} &\longleftrightarrow O_s^A(x) \\ \frac{\delta}{\delta J_{O_s^\lambda}(x)} &\longleftrightarrow O_s^\lambda(x) \\ \frac{\delta}{\delta \bar{J}_{M_s}(x)} &\longleftrightarrow M_s(x) \\ -\frac{\delta}{\delta J_{\bar{M}_s}(x)} &\longleftrightarrow \bar{M}_s(x) \end{aligned} \quad (5.2)$$

The generating functional of connected correlators $\mathcal{W}_{\text{conf}} = \log \mathcal{Z}_{\text{conf}}$ follows. For example:

$$\langle O_{s_1}^A(x) O_{s_2}^A(y) \rangle = \left. \frac{\delta}{\delta J_{O_{s_1}^A}(x)} \frac{\delta}{\delta J_{O_{s_2}^A}(y)} \mathcal{W}_{\text{conf}} \right|_{J=0} \quad (5.3)$$

and:

$$\langle M_{s_1}(x) \bar{M}_{s_2}(y) \rangle = \left. \frac{\delta}{\delta \bar{J}_{M_{s_1}}(x)} \left(-\frac{\delta}{\delta J_{\bar{M}_{s_2}}(y)} \right) \mathcal{W}_{\text{conf}} \right|_{J=0} \quad (5.4)$$

For brevity we will omit from now on the specification $|_{J=0}$.

5.1 Connected generating functional $\mathcal{W}_{\text{conf}}$ as the log of a superdeterminant of a quadratic form

The above functional integral is quadratic in the elementary fields and therefore it may be computed exactly [1]. Employing the symmetry properties [1] of the Gegenbauer polynomials (appendix B) we get:

$$\begin{aligned} \mathcal{Z}_{\text{conf}} &[J_{O^A}, J_{\tilde{O}^A}, J_{O^\lambda}, J_{\tilde{O}^\lambda}, \bar{J}_M, \bar{J}_{\bar{M}}] \\ &= \frac{1}{Z} \int \mathcal{D}A \mathcal{D}\bar{A} \mathcal{D}\lambda \mathcal{D}\bar{\lambda} \exp \left(-\frac{1}{2} \int d^4x \left(\bar{A}^a(x) A^a(x) \right) \mathcal{M}_{AA}^{ab} \begin{pmatrix} A^b(x) \\ \bar{A}^b(x) \end{pmatrix} \right. \\ &\quad \left. - \frac{1}{2} \int d^4x \left(\bar{\lambda}^a(x) \lambda^a(x) \right) \mathcal{M}_{\lambda\lambda}^{ab} \begin{pmatrix} \lambda^b(x) \\ \bar{\lambda}^b(x) \end{pmatrix} \right. \\ &\quad \left. + \frac{1}{4} \int d^4x \sum_s \left(\bar{\lambda}^a(x) \lambda^a(x) \right) \mathcal{M}_{\lambda A}^{ab} \begin{pmatrix} A^b(x) \\ \bar{A}^b(x) \end{pmatrix} \right. \\ &\quad \left. + \frac{1}{4} \int d^4x \sum_s \left(\bar{A}^a(x) A^a(x) \right) \mathcal{M}_{A\lambda}^{ab} \begin{pmatrix} \lambda^b(x) \\ \bar{\lambda}^b(x) \end{pmatrix} \right) \end{aligned} \quad (5.5)$$

with:

$$\begin{aligned}
\mathcal{M}_{AA}^{ab} &= \delta^{ab} \begin{pmatrix} i\square - \frac{1}{2} \sum_s J_{O_s^A} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} - \frac{1}{2} \sum_s J_{\tilde{O}_s^A} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}} & 0 \\ 0 & i\square - \frac{1}{2} \sum_s J_{O_s^A} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} + \frac{1}{2} \sum_s J_{\tilde{O}_s^A} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}} \end{pmatrix} \\
\mathcal{M}_{\lambda\lambda}^{ab} &= \delta^{ab} \begin{pmatrix} -\square\partial_+^{-1} - \frac{1}{2} \sum_s J_{O_s^\lambda} \otimes \mathcal{Y}_{s-1}^{\frac{3}{2}} - \frac{1}{2} \sum_s J_{\tilde{O}_s^\lambda} \otimes \mathcal{H}_{s-1}^{\frac{3}{2}} & 0 \\ 0 & -\square\partial_+^{-1} - \frac{1}{2} \sum_s J_{O_s^\lambda} \otimes \mathcal{Y}_{s-1}^{\frac{3}{2}} + \frac{1}{2} \sum_s J_{\tilde{O}_s^\lambda} \otimes \mathcal{H}_{s-1}^{\frac{3}{2}} \end{pmatrix} \\
\mathcal{M}_{A\lambda}^{ab} &= \frac{\delta^{ab}}{2} \begin{pmatrix} 0 & -J_{\bar{M}_s} \otimes \mathcal{G}_{s-1}^{(2,1)} (-1)^{s-1} \\ \bar{J}_{M_s} \otimes \mathcal{G}_{s-1}^{(2,1)} & 0 \end{pmatrix} \\
\mathcal{M}_{\lambda A}^{ab} &= \frac{\delta^{ab}}{2} \begin{pmatrix} 0 & J_{\bar{M}_s} \otimes \mathcal{G}_{s-1}^{(1,2)} \\ -\bar{J}_{M_s} \otimes \mathcal{G}_{s-1}^{(1,2)} (-1)^{s-1} & 0 \end{pmatrix}
\end{aligned} \tag{5.6}$$

where the symbol \otimes implies that the right and left derivatives do not act on the sources J . We group the above matrices into a single supermatrix:

$$\mathcal{X}^{ab} = \begin{pmatrix} \mathcal{M}_{\lambda\lambda}^{ab} & \mathcal{M}_{\lambda A}^{ab} \\ \mathcal{M}_{A\lambda}^{ab} & \mathcal{M}_{AA}^{ab} \end{pmatrix} \tag{5.7}$$

so that the generating functional reads:

$$\begin{aligned}
&\mathcal{Z}_{\text{conf}} [J_{O^A}, J_{\tilde{O}^A}, J_{O^\lambda}, J_{\tilde{O}^\lambda}, \bar{J}_M, J_{\bar{M}}] \\
&= \frac{1}{Z} \int \mathcal{D}A \mathcal{D}\bar{A} \mathcal{D}\lambda \mathcal{D}\bar{\lambda} \exp \left(-\frac{1}{2} \int d^4x \left(\bar{\lambda}^a(x) \lambda^a(x) \bar{A}^a(x) A^a(x) \right) \mathcal{X}^{ab} \begin{pmatrix} \lambda^b(x) \\ \bar{\lambda}^b(x) \\ A^b(x) \\ \bar{A}^b(x) \end{pmatrix} \right)
\end{aligned} \tag{5.8}$$

that reduces to the superdeterminant [10]:

$$\mathcal{Z}_{\text{conf}} [J_{O^A}, J_{\tilde{O}^A}, J_{O^\lambda}, J_{\tilde{O}^\lambda}, \bar{J}_M, J_{\bar{M}}] = \text{sDet}^{\frac{1}{2}}(\mathcal{X}) \tag{5.9}$$

More explicitly:

$$\begin{aligned}
\mathcal{Z}_{\text{conf}} [J_{O^A}, J_{\tilde{O}^A}, J_{O^\lambda}, J_{\tilde{O}^\lambda}, \bar{J}_M, J_{\bar{M}}] &= \text{Det}^{\frac{1}{2}}(\mathcal{M}_{\lambda\lambda}) \text{Det}^{-\frac{1}{2}}(\mathcal{M}_{AA} - \mathcal{M}_{A\lambda} \mathcal{M}_{\lambda\lambda}^{-1} \mathcal{M}_{\lambda A}) \\
&= \text{Det}^{-\frac{1}{2}}(\mathcal{M}_{AA}) \text{Det}^{\frac{1}{2}}(\mathcal{M}_{\lambda\lambda} - \mathcal{M}_{\lambda A} \mathcal{M}_{AA}^{-1} \mathcal{M}_{A\lambda})
\end{aligned} \tag{5.10}$$

where the two results depend on first integrating either on the fermionic or bosonic variables.

The entries of $\mathcal{M}_{A\lambda}\mathcal{M}_{\lambda\lambda}^{-1}\mathcal{M}_{\lambda A}$ read:

$$\begin{aligned}
[\mathcal{M}_{\lambda A}\mathcal{M}_{AA}^{-1}\mathcal{M}_{A\lambda}]_{11} &= \\
\frac{1}{4}\mathcal{G}_{s-1}^{(1,2)}(-1)^{s-1} \otimes \bar{J}_{M_s}(i\square - \frac{1}{2}J_{O_{s_1}^A} \otimes \mathcal{Y}_{s_1-2}^{\frac{5}{2}} - \frac{1}{2}J_{\tilde{O}_{s_1}^A} \otimes \mathcal{H}_{s_1-2}^{\frac{5}{2}})^{-1}\mathcal{G}_{s_2-1}^{(2,1)}(-1)^{s_2-1} \otimes J_{\bar{M}_{s_2}} \\
[\mathcal{M}_{\lambda A}\mathcal{M}_{AA}^{-1}\mathcal{M}_{A\lambda}]_{12} &= 0 \\
[\mathcal{M}_{\lambda A}\mathcal{M}_{AA}^{-1}\mathcal{M}_{A\lambda}]_{21} &= 0 \\
[\mathcal{M}_{\lambda A}\mathcal{M}_{AA}^{-1}\mathcal{M}_{A\lambda}]_{22} &= \\
\frac{1}{4}\mathcal{G}_{s-1}^{(1,2)} \otimes J_{\bar{M}_s}(i\square - \frac{1}{2}J_{O_{s_1}^A} \otimes \mathcal{Y}_{s_1-2}^{\frac{5}{2}} + \frac{1}{2}J_{\tilde{O}_{s_1}^A} \otimes \mathcal{H}_{s_1-2}^{\frac{5}{2}})^{-1}\mathcal{G}_{s_2-1}^{(2,1)} \otimes \bar{J}_{M_{s_2}} & \quad (5.11)
\end{aligned}$$

and similarly:

$$\begin{aligned}
[\mathcal{M}_{A\lambda}\mathcal{M}_{\lambda\lambda}^{-1}\mathcal{M}_{\lambda A}]_{11} &= \\
\frac{1}{4}\mathcal{G}_{s-1}^{(2,1)}(-1)^{s-1} \otimes J_{\bar{M}_s}(-\square\partial_+^{-1} - \frac{1}{2}J_{O_{s_1}^\lambda} \otimes \mathcal{Y}_{s_1-1}^{\frac{3}{2}} + \frac{1}{2}J_{\tilde{O}_{s_1}^\lambda} \otimes \mathcal{H}_{s_1-1}^{\frac{3}{2}})^{-1}\mathcal{G}_{s_2-1}^{(1,2)}(-1)^{s_2-1} \otimes \bar{J}_{M_{s_2}} \\
[\mathcal{M}_{A\lambda}\mathcal{M}_{\lambda\lambda}^{-1}\mathcal{M}_{\lambda A}]_{12} &= 0 \\
[\mathcal{M}_{A\lambda}\mathcal{M}_{\lambda\lambda}^{-1}\mathcal{M}_{\lambda A}]_{21} &= 0 \\
[\mathcal{M}_{A\lambda}\mathcal{M}_{\lambda\lambda}^{-1}\mathcal{M}_{\lambda A}]_{22} &= \\
\frac{1}{4}\mathcal{G}_{s-1}^{(2,1)} \otimes \bar{J}_{M_s}(-\square\partial_+^{-1} - \frac{1}{2}J_{O_{s_1}^\lambda} \otimes \mathcal{Y}_{s_1-1}^{\frac{3}{2}} - \frac{1}{2}J_{\tilde{O}_{s_1}^\lambda} \otimes \mathcal{H}_{s_1-1}^{\frac{3}{2}})^{-1}\mathcal{G}_{s_2-1}^{(1,2)} \otimes J_{\bar{M}_{s_2}} & \quad (5.12)
\end{aligned}$$

Evaluating the determinant of block matrices [1] by means of:

$$\text{Det} \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \text{Det}(A) \text{ Det}(D) \text{ Det}(1 - D^{-1}CA^{-1}B) \quad (5.13)$$

provided that A^{-1} and D^{-1} exist, we get for the connected generating functional:

$$\begin{aligned}
\mathcal{W}_{\text{conf}} &[J_{O^A}, J_{\tilde{O}^A}, J_{O^\lambda}, J_{\tilde{O}^\lambda}, \bar{J}_M, J_{\bar{M}}] \\
&= -\frac{1}{2} \log \text{Det} \left(\mathcal{I} + \frac{1}{2}i\square^{-1}J_{O_s^A} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} + \frac{1}{2}i\square^{-1}J_{\tilde{O}_s^A} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}} \right) \\
&\quad - \frac{1}{2} \log \text{Det} \left(\mathcal{I} + \frac{1}{2}i\square^{-1}J_{O_s^\lambda} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} - \frac{1}{2}i\square^{-1}J_{\tilde{O}_s^\lambda} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}} \right) \\
&\quad + \frac{1}{2} \log \text{Det} \left(\mathcal{I} + \frac{1}{2}\partial_+\square^{-1}J_{O_s^\lambda} \otimes \mathcal{Y}_{s-1}^{\frac{3}{2}} + \frac{1}{2}\partial_+\square^{-1}J_{\tilde{O}_s^\lambda} \otimes \mathcal{H}_{s-1}^{\frac{3}{2}} \right) \\
&\quad + \frac{1}{2} \log \text{Det} \left(\mathcal{I} + \frac{1}{2}\partial_+\square^{-1}J_{O_s^\lambda} \otimes \mathcal{Y}_{s-1}^{\frac{3}{2}} - \frac{1}{2}\partial_+\square^{-1}J_{\tilde{O}_s^\lambda} \otimes \mathcal{H}_{s-1}^{\frac{3}{2}} \right) \\
&\quad + \frac{1}{2} \log \text{Det} \left(\mathcal{I} - \frac{1}{4}(\mathcal{I} + \frac{1}{2}\partial_+\square^{-1}J_{O_s^\lambda} \otimes \mathcal{Y}_{s-1}^{\frac{3}{2}} - \frac{1}{2}\partial_+\square^{-1}J_{\tilde{O}_s^\lambda} \otimes \mathcal{H}_{s-1}^{\frac{3}{2}})^{-1} \right. \\
&\quad \left. \partial_+\square^{-1}\bar{J}_{M_{s_1}} \otimes \mathcal{G}_{s_1-1}^{(1,2)}(-1)^{s_1-1} \right. \\
&\quad \left. (\mathcal{I} + \frac{1}{2}i\square^{-1}J_{O_{s_2}^A} \otimes \mathcal{Y}_{s_2-2}^{\frac{5}{2}} + \frac{1}{2}i\square^{-1}J_{\tilde{O}_{s_2}^A} \otimes \mathcal{H}_{s_2-2}^{\frac{5}{2}})^{-1}i\square^{-1}J_{\bar{M}_{s_3}} \otimes \mathcal{G}_{s_3-1}^{(2,1)}(-1)^{s_3-1} \right) \\
&\quad + \frac{1}{2} \log \text{Det} \left(\mathcal{I} - \frac{1}{4}(\mathcal{I} + \frac{1}{2}\partial_+\square^{-1}J_{O_s^\lambda} \otimes \mathcal{Y}_{s-1}^{\frac{3}{2}} + \frac{1}{2}\partial_+\square^{-1}J_{\tilde{O}_s^\lambda} \otimes \mathcal{H}_{s-1}^{\frac{3}{2}})^{-1} \right. \\
&\quad \left. \partial_+\square^{-1}J_{\bar{M}_{s_1}} \otimes \mathcal{G}_{s_1-1}^{(1,2)} \right. \\
&\quad \left. (\mathcal{I} + \frac{1}{2}i\square^{-1}J_{O_{s_2}^A} \otimes \mathcal{Y}_{s_2-2}^{\frac{5}{2}} - \frac{1}{2}i\square^{-1}J_{\tilde{O}_{s_2}^A} \otimes \mathcal{H}_{s_2-2}^{\frac{5}{2}})^{-1}i\square^{-1}\bar{J}_{M_{s_3}} \otimes \mathcal{G}_{s_3-1}^{(2,1)} \right) \quad (5.14)
\end{aligned}$$

and:

$$\begin{aligned}
& \mathcal{W}_{\text{conf}} [J_{O^A}, J_{\tilde{O}^A}, J_{O^\lambda}, J_{\tilde{O}^\lambda}, \bar{J}_M, J_{\bar{M}}] \\
&= \frac{1}{2} \log \text{Det} \left(\mathcal{I} + \frac{1}{2} \partial_+ \square^{-1} J_{O_s^\lambda} \otimes \mathcal{Y}_{s-1}^{\frac{3}{2}} + \frac{1}{2} \partial_+ \square^{-1} J_{\tilde{O}_s^\lambda} \otimes \mathcal{H}_{s-1}^{\frac{3}{2}} \right) \\
&+ \frac{1}{2} \log \text{Det} \left(\mathcal{I} + \frac{1}{2} \partial_+ \square^{-1} J_{O_s^\lambda} \otimes \mathcal{Y}_{s-1}^{\frac{3}{2}} - \frac{1}{2} \partial_+ \square^{-1} J_{\tilde{O}_s^\lambda} \otimes \mathcal{H}_{s-1}^{\frac{3}{2}} \right) \\
&- \frac{1}{2} \log \text{Det} \left(\mathcal{I} + \frac{1}{2} i \square^{-1} J_{O_s^A} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} + \frac{1}{2} i \square^{-1} J_{\tilde{O}_s^A} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}} \right) \\
&- \frac{1}{2} \log \text{Det} \left(\mathcal{I} + \frac{1}{2} i \square^{-1} J_{O_s^A} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} - \frac{1}{2} i \square^{-1} J_{\tilde{O}_s^A} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}} \right) \\
&- \frac{1}{2} \log \text{Det} \left(\mathcal{I} - \frac{1}{4} (\mathcal{I} + \frac{1}{2} i \square^{-1} J_{O_s^A} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} + \frac{1}{2} i \square^{-1} J_{\tilde{O}_s^A} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}})^{-1} \right. \\
&\quad \left. i \square^{-1} J_{\bar{M}_{s_1}} \otimes \mathcal{G}_{s_1-1}^{(2,1)} (-1)^{s_1-1} \right. \\
&\quad \left. (\mathcal{I} + \frac{1}{2} \partial_+ \square^{-1} J_{O_{s_2}^\lambda} \otimes \mathcal{Y}_{s_2-1}^{\frac{3}{2}} - \frac{1}{2} \partial_+ \square^{-1} J_{\tilde{O}_{s_2}^\lambda} \otimes \mathcal{H}_{s_2-1}^{\frac{3}{2}})^{-1} \partial_+ \square^{-1} \bar{J}_{M_{s_3}} \otimes \mathcal{G}_{s_3-1}^{(1,2)} (-1)^{s_3-1} \right) \\
&- \frac{1}{2} \log \text{Det} \left(\mathcal{I} - \frac{1}{4} (\mathcal{I} + \frac{1}{2} i \square^{-1} J_{O_s^A} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} - \frac{1}{2} i \square^{-1} J_{\tilde{O}_s^A} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}})^{-1} \right. \\
&\quad \left. i \square^{-1} \bar{J}_{M_{s_1}} \otimes \mathcal{G}_{s_1-1}^{(2,1)} \right. \\
&\quad \left. (\mathcal{I} + \frac{1}{2} \partial_+ \square^{-1} J_{O_{s_2}^\lambda} \otimes \mathcal{Y}_{s_2-1}^{\frac{3}{2}} + \frac{1}{2} \partial_+ \square^{-1} J_{\tilde{O}_{s_2}^\lambda} \otimes \mathcal{H}_{s_2-1}^{\frac{3}{2}})^{-1} \partial_+ \square^{-1} J_{\bar{M}_{s_3}} \otimes \mathcal{G}_{s_3-1}^{(1,2)} \right) \quad (5.15)
\end{aligned}$$

with \mathcal{I} the identity in both color and space-time, and the sum over repeated spin indices understood. After rescaling the operators:

$$\begin{aligned}
O_s'^A(x) &= \frac{1}{N} \frac{2\Gamma(5)\Gamma(s+1)}{\Gamma(3)\Gamma(s+3)} O_s^A(x) & \tilde{O}_s'^A(x) &= \frac{1}{N} \frac{2\Gamma(5)\Gamma(s+1)}{\Gamma(3)\Gamma(s+3)} \tilde{O}_s^A(x) \\
O_s'^\lambda(x) &= \frac{1}{N} \frac{4}{s+1} O_s^\lambda(x) & \tilde{O}_s'^\lambda(x) &= \frac{1}{N} \frac{4}{s+1} \tilde{O}_s^\lambda(x) \\
M'_s &= \frac{2}{N} M_s
\end{aligned} \quad (5.16)$$

so that their 2-point correlators are of order 1 for large N , we obtain more explicitly:

$$\begin{aligned}
\mathcal{W}_{\text{conf}} [J_{O'^A}, J_{\tilde{O}'^A}, J_{O'^\lambda}, J_{\tilde{O}'^\lambda}, \bar{J}_{M'}, J_{\bar{M}'}] = & \\
& -\frac{N^2-1}{2} \log \text{Det} \left(I + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (i \vec{\partial}_+)^{s-k-1} i \square^{-1} (J_{O'_s A} + J_{\tilde{O}'_s A}) (i \vec{\partial}_+)^{k+1} \right) \\
& -\frac{N^2-1}{2} \log \text{Det} \left(I + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (i \vec{\partial}_+)^{s-k-1} i \square^{-1} (J_{O'_s A} - J_{\tilde{O}'_s A}) (i \vec{\partial}_+)^{k+1} \right) \\
& +\frac{N^2-1}{2} \log \text{Det} \left(I - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (i \vec{\partial}_+)^{s-k} i \square^{-1} (J_{O'_s \lambda} + J_{\tilde{O}'_s \lambda}) (i \vec{\partial}_+)^k \right) \\
& +\frac{N^2-1}{2} \log \text{Det} \left(I - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (i \vec{\partial}_+)^{s-k} i \square^{-1} (J_{O'_s \lambda} - J_{\tilde{O}'_s \lambda}) (i \vec{\partial}_+)^k \right) \\
& -\frac{N^2-1}{2} \log \text{Det} \left[I + \frac{1}{N^2} \left(I - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (i \vec{\partial}_+)^{s-k} i \square^{-1} (J_{O'_s \lambda} - J_{\tilde{O}'_s \lambda}) (i \vec{\partial}_+)^k \right)^{-1} \right. \\
& \left. \sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1+1}{k_1+2} (-1)^{s_1-1} (i \vec{\partial}_+)^{s_1-k_1} i \square^{-1} \bar{J}_{M'_{s_1}} (i \vec{\partial}_+)^{k_1+1} \right. \\
& \left. \left(I + \frac{1}{N} \sum_{k_2=0}^{s_2-2} \binom{s_2}{k_2} \binom{s_2}{k_2+2} (i \vec{\partial}_+)^{s_2-k_2-1} i \square^{-1} (J_{O'_{s_2} A} + J_{\tilde{O}'_{s_2} A}) (i \vec{\partial}_+)^{k_2+1} \right)^{-1} \right. \\
& \left. \sum_{k_3=0}^{s_3-1} \binom{s_3+1}{k_3} \binom{s_3}{k_3+1} (-1)^{s_3-1} (i \vec{\partial}_+)^{s_3-k_3} i \square^{-1} J_{\bar{M}'_{s_3}} (i \vec{\partial}_+)^{k_3} \right] \\
& -\frac{N^2-1}{2} \log \text{Det} \left[I + \frac{1}{N^2} \left(I - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (i \vec{\partial}_+)^{s-k} i \square^{-1} (J_{O'_s \lambda} + J_{\tilde{O}'_s \lambda}) (i \vec{\partial}_+)^k \right)^{-1} \right. \\
& \left. \sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1+1}{k_1+2} (i \vec{\partial}_+)^{s_1-k_1} i \square^{-1} J_{\bar{M}'_{s_1}} (i \vec{\partial}_+)^{k_1+1} \right. \\
& \left. \left(I + \frac{1}{N} \sum_{k_2=0}^{s_2-2} \binom{s_2}{k_2} \binom{s_2}{k_2+2} (i \vec{\partial}_+)^{s_2-k_2-1} i \square^{-1} (J_{O'_{s_2} A} - J_{\tilde{O}'_{s_2} A}) (i \vec{\partial}_+)^{k_2+1} \right)^{-1} \right. \\
& \left. \sum_{k_3=0}^{s_3-1} \binom{s_3+1}{k_3} \binom{s_3}{k_3+1} (i \vec{\partial}_+)^{s_3-k_3} i \square^{-1} \bar{J}_{M'_{s_3}} (i \vec{\partial}_+)^{k_3} \right]
\end{aligned} \tag{5.17}$$

and:

$$\begin{aligned}
& \mathcal{W}_{\text{conf}} [J_{O'^A}, J_{\tilde{O}'^A}, J_{O'^\lambda}, J_{\tilde{O}'^\lambda}, \bar{J}_{M'}, J_{\bar{M}'}] = \\
& -\frac{N^2-1}{2} \log \text{Det} \left(I + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (i \vec{\partial}_+)^{s-k-1} i \square^{-1} (J_{O'_s} + J_{\tilde{O}'_s}) (i \vec{\partial}_+)^{k+1} \right) \\
& -\frac{N^2-1}{2} \log \text{Det} \left(I + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (i \vec{\partial}_+)^{s-k-1} i \square^{-1} (J_{O'_s} - J_{\tilde{O}'_s}) (i \vec{\partial}_+)^{k+1} \right) \\
& +\frac{N^2-1}{2} \log \text{Det} \left(I - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (i \vec{\partial}_+)^{s-k} i \square^{-1} (J_{O'_s} + J_{\tilde{O}'_s}) (i \vec{\partial}_+)^k \right) \\
& +\frac{N^2-1}{2} \log \text{Det} \left(I - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (i \vec{\partial}_+)^{s-k} i \square^{-1} (J_{O'_s} - J_{\tilde{O}'_s}) (i \vec{\partial}_+)^k \right) \\
& +\frac{N^2-1}{2} \log \text{Det} \left[I + \frac{1}{N^2} \left(I + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (i \vec{\partial}_+)^{s-k-1} i \square^{-1} (J_{O'_s} - J_{\tilde{O}'_s}) (i \vec{\partial}_+)^{k+1} \right)^{-1} \right. \\
& \left. \sum_{k_1=0}^{s_1-1} \binom{s_1+1}{k_1} \binom{s_1}{k_1+1} (-1)^{s_1-1} (i \vec{\partial}_+)^{s_1-k_1} i \square^{-1} J_{\bar{M}'_{s_1}} (i \vec{\partial}_+)^{k_1} \right. \\
& \left. \left(I - \frac{1}{N} \sum_{k_2=0}^{s_2-1} \binom{s_2}{k_2} \binom{s_2}{k_2+1} (i \vec{\partial}_+)^{s_2-k_2} i \square^{-1} (J_{O'_{s_2}} + J_{\tilde{O}'_{s_2}}) (i \vec{\partial}_+)^{k_2} \right)^{-1} \right. \\
& \left. \sum_{k_3=0}^{s_3-1} \binom{s_3}{k_3} \binom{s_3+1}{k_3+2} (-1)^{s_3-1} (i \vec{\partial}_+)^{s_3-k_3} i \square^{-1} \bar{J}_{M'_{s_3}} (i \vec{\partial}_+)^{k_3+1} \right] \\
& +\frac{N^2-1}{2} \log \text{Det} \left[I + \frac{1}{N^2} \left(I + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (i \vec{\partial}_+)^{s-k-1} i \square^{-1} (J_{O'_s} + J_{\tilde{O}'_s}) (i \vec{\partial}_+)^{k+1} \right)^{-1} \right. \\
& \left. \sum_{k_1=0}^{s_1-1} \binom{s_1+1}{k_1} \binom{s_1}{k_1+1} (i \vec{\partial}_+)^{s_1-k_1} i \square^{-1} \bar{J}_{M'_{s_1}} (i \vec{\partial}_+)^{k_1} \right. \\
& \left. \left(I - \frac{1}{N} \sum_{k_2=0}^{s_2-1} \binom{s_2}{k_2} \binom{s_2}{k_2+1} (i \vec{\partial}_+)^{s_2-k_2} i \square^{-1} (J_{O'_{s_2}} - J_{\tilde{O}'_{s_2}}) (i \vec{\partial}_+)^{k_2} \right)^{-1} \right. \\
& \left. \sum_{k_3=0}^{s_3-1} \binom{s_3+1}{k_3} \binom{s_3}{k_3+1} (i \vec{\partial}_+)^{s_3-k_3} i \square^{-1} J_{\bar{M}'_{s_3}} (i \vec{\partial}_+)^{k_3} \right] \tag{5.18}
\end{aligned}$$

where I is the identity in space-time with kernel:

$$I \rightarrow \delta^{(4)}(x - y) \tag{5.19}$$

and, after performing the color trace, we have employed the definitions in eqs. (4.7), (4.8), (4.9) and [1]:

$$i \square^{-1} \overleftrightarrow{\partial}_+^{s-k-1} = (-1)^{s-k-1} \overrightarrow{\partial}_+^{s-k-1} i \square^{-1} \tag{5.20}$$

that follows from (minus) the propagator in the coordinate representation [2]:

$$i \square^{-1} \rightarrow \frac{1}{4\pi^2} \frac{1}{|x - y|^2 - i\epsilon} \tag{5.21}$$

5.2 Connected generating functional Γ_{conf} as the log a Fredholm superdeterminant

By means of the dictionary [1] we rewrite the generating functional $\mathcal{W}_{\text{conf}}$ – that is actually the logarithm of a (super)determinant of a quadratic form as remarked above – as the logarithm of a (super)determinant Γ_{conf} of integral operators formally of Fredholm type:

$$\begin{aligned} & \Gamma_{\text{conf}} [j_{O^A}, j_{\tilde{O}^A}, j_{O^\lambda}, j_{\tilde{O}^\lambda}, \bar{j}_M, j_{\bar{M}}] \\ &= -\frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} + \mathcal{D}_A^{-1} j_{O^A} + \mathcal{D}_A^{-1} j_{\tilde{O}^A}] - \frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} + \mathcal{D}_A^{-1} j_{O^A} - \mathcal{D}_A^{-1} j_{\tilde{O}^A}] \\ &+ \frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} - \mathcal{D}_\lambda^{-1} j_{O^\lambda} - \mathcal{D}_\lambda^{-1} j_{\tilde{O}^\lambda}] + \frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} - \mathcal{D}_\lambda^{-1} j_{O^\lambda} + \mathcal{D}_\lambda^{-1} j_{\tilde{O}^\lambda}] \\ &+ \frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} + (\mathbb{I} - \mathcal{D}_\lambda^{-1} j_{O^\lambda} - \mathcal{D}_\lambda^{-1} j_{\tilde{O}^\lambda})^{-1} \mathcal{D}_M^{-1} \bar{j}_M (\mathbb{I} + \mathcal{D}_A^{-1} j_{O^A} + \mathcal{D}_A^{-1} j_{\tilde{O}^A})^{-1} \mathcal{D}_{\bar{M}}^{-1} j_{\bar{M}}] \\ &+ \frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} + (\mathbb{I} - \mathcal{D}_\lambda^{-1} j_{O^\lambda} + \mathcal{D}_\lambda^{-1} j_{\tilde{O}^\lambda})^{-1} \mathcal{D}_M^{-1} j_{\bar{M}} (\mathbb{I} + \mathcal{D}_A^{-1} j_{O^A} - \mathcal{D}_A^{-1} j_{\tilde{O}^A})^{-1} \mathcal{D}_{\bar{M}}^{-1} \bar{j}_M] \end{aligned} \quad (5.22)$$

and:

$$\begin{aligned} & \Gamma_{\text{conf}} [j_{O^A}, j_{\tilde{O}^A}, j_{O^\lambda}, j_{\tilde{O}^\lambda}, \bar{j}_M, j_{\bar{M}}] \\ &= -\frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} + \mathcal{D}_A^{-1} j_{O^A} + \mathcal{D}_A^{-1} j_{\tilde{O}^A}] - \frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} + \mathcal{D}_A^{-1} j_{O^A} - \mathcal{D}_A^{-1} j_{\tilde{O}^A}] \\ &+ \frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} - \mathcal{D}_\lambda^{-1} j_{O^\lambda} - \mathcal{D}_\lambda^{-1} j_{\tilde{O}^\lambda}] + \frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} - \mathcal{D}_\lambda^{-1} j_{O^\lambda} + \mathcal{D}_\lambda^{-1} j_{\tilde{O}^\lambda}] \\ &- \frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} + (\mathbb{I} + \mathcal{D}_A^{-1} j_{O^A} + \mathcal{D}_A^{-1} j_{\tilde{O}^A})^{-1} \mathcal{D}_{\bar{M}}^{-1} j_{\bar{M}} (\mathbb{I} - \mathcal{D}_\lambda^{-1} j_{O^\lambda} - \mathcal{D}_\lambda^{-1} j_{\tilde{O}^\lambda})^{-1} \mathcal{D}_M^{-1} \bar{j}_M] \\ &- \frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} + (\mathbb{I} + \mathcal{D}_A^{-1} j_{O^A} - \mathcal{D}_A^{-1} j_{\tilde{O}^A})^{-1} \mathcal{D}_{\bar{M}}^{-1} \bar{j}_M (\mathbb{I} - \mathcal{D}_\lambda^{-1} j_{O^\lambda} + \mathcal{D}_\lambda^{-1} j_{\tilde{O}^\lambda})^{-1} \mathcal{D}_M^{-1} j_{\bar{M}}] \end{aligned} \quad (5.23)$$

where \mathbb{I} is the identity in space-time and discrete indices defined below. The corresponding kernels are defined as follows.

- gluon-gluon kernel

$$\begin{aligned} \mathcal{D}_A^{-1} &= \frac{1}{2} \frac{\Gamma(3)\Gamma(s_1 + 3)}{\Gamma(5)\Gamma(s_1 + 1)} \binom{s_1}{k_1} \binom{s_2}{k_2 + 2} (-i\partial_+)^{s_1 - k_1 + k_2} i\Box^{-1} \\ &\rightarrow \mathcal{D}_{A s_1 k_1, s_2 k_2}^{-1}(x - y) = \frac{1}{8\pi^2} \frac{\Gamma(3)\Gamma(s_1 + 3)}{\Gamma(5)\Gamma(s_1 + 1)} \binom{s_1}{k_1} \binom{s_2}{k_2 + 2} (-i\partial_+)^{s_1 - k_1 + k_2} \frac{1}{|x - y|^2 - i\epsilon} \end{aligned} \quad (5.24)$$

- gluino-gluino kernel

$$\begin{aligned} \mathcal{D}_\lambda^{-1} &= \frac{1}{2} \frac{s_1 + 1}{2} \binom{s_1}{k_1} \binom{s_2}{k_2 + 1} (-i\partial_+)^{s_1 - k_1 + k_2} i\Box^{-1} \\ &\rightarrow \mathcal{D}_{\lambda s_1 k_1, s_2 k_2}^{-1}(x - y) = \frac{1}{8\pi^2} \frac{s_1 + 1}{2} \binom{s_1}{k_1} \binom{s_2}{k_2 + 1} (-i\partial_+)^{s_1 - k_1 + k_2} \frac{1}{|x - y|^2 - i\epsilon} \end{aligned} \quad (5.25)$$

- gluon-gluino kernels

$$\begin{aligned} \mathcal{D}_M^{-1} &= -\frac{i}{2} \binom{s_1+1}{k_1+2} \binom{s_2+1}{k_2} (i\partial_+)^{s_1-k_1+k_2+1} i\square^{-1} \\ \rightarrow \mathcal{D}_{Ms_1k_1,s_2k_2}^{-1}(x-y) &= -\frac{i}{8\pi^2} \binom{s_1+1}{k_1+2} \binom{s_2+1}{k_2} (i\partial_+)^{s_1-k_1+k_2+1} \frac{1}{|x-y|^2 - i\epsilon} \end{aligned} \quad (5.26)$$

$$\begin{aligned} \mathcal{D}_{\bar{M}}^{-1} &= -\frac{i}{2} \binom{s_1}{k_1} \binom{s_2}{k_2+1} (i\partial_+)^{s_1-k_1+k_2} i\square^{-1} \\ \rightarrow \mathcal{D}_{\bar{M}s_1k_1,s_2k_2}^{-1}(x-y) &= -\frac{i}{8\pi^2} \binom{s_1}{k_1} \binom{s_2}{k_2+1} (i\partial_+)^{s_1-k_1+k_2} \frac{1}{|x-y|^2 - i\epsilon} \end{aligned} \quad (5.27)$$

The kernels are coupled to the currents $j_{\mathcal{O}_{sk}}$ that are dual to the component operators \mathcal{O}_{sk} employed to construct the conformal operators \mathcal{O}_s [2]:

$$\mathcal{O}_s = \sum_{k=0}^l \mathcal{O}_{sk} \quad (5.28)$$

Consequently, the n -point correlators are defined [1] by:

$$\langle \mathcal{O}_{s_1}(x_1) \dots \mathcal{O}_{s_n}(x_n) \rangle = \sum_{k_1=0}^{l_1} \frac{\delta}{\delta j_{\mathcal{O}_{s_1 k_1}}(x_1)} \dots \sum_{k_n=0}^{l_n} \frac{\delta}{\delta j_{\mathcal{O}_{s_n k_n}}(x_n)} \Gamma_{\text{conf}}[j_{\mathcal{O}}] \quad (5.29)$$

and the following identity holds:

$$\sum_{k_1=0}^{l_1} \frac{\delta}{\delta j_{\mathcal{O}_{s_1 k_1}}(x_1)} \dots \sum_{k_n=0}^{l_n} \frac{\delta}{\delta j_{\mathcal{O}_{s_n k_n}}(x_n)} \Gamma_{\text{conf}}[j_{\mathcal{O}}] = \frac{\delta}{\delta J_{\mathcal{O}_{s_1}}(x_1)} \dots \frac{\delta}{\delta J_{\mathcal{O}_{s_n}}(x_n)} \mathcal{W}_{\text{conf}}[J_{\mathcal{O}}] \quad (5.30)$$

according to the dictionary [1].

5.3 Connected generating functional in the momentum representation

The conformal generating functional in the momentum representation is defined by the functional integral:

$$\mathcal{Z}_{\text{conf}}[J_{\mathcal{O}}] = \frac{1}{Z} \int \mathcal{D}A \mathcal{D}\bar{A} \mathcal{D}\lambda \mathcal{D}\bar{\lambda} e^{\int d^4x (-i\bar{A}^a \square A^a + \bar{\lambda}^a \square \partial_+^{-1} \lambda^a)} \exp \left(\int \frac{d^4p}{(2\pi)^4} \sum_i J_{\mathcal{O}_i}(-p) \mathcal{O}_i(p) \right) \quad (5.31)$$

Hence, we get for the correlators in the momentum representation [1]:

$$\langle \mathcal{O}_{s_1}(p_1) \dots \mathcal{O}_{s_n}(p_n) \rangle = (2\pi)^4 \frac{\delta}{\delta J_{\mathcal{O}_{s_1}}(-p_1)} \dots (2\pi)^4 \frac{\delta}{\delta J_{\mathcal{O}_{s_n}}(-p_n)} \mathcal{W}[J_{\mathcal{O}}] \quad (5.32)$$

The generating functional is obtained by means of the kernels in the momentum representation:

$$i\square^{-1} \rightarrow \frac{-i}{|q|^2 + i\epsilon} \quad (5.33)$$

and:

$$(i\partial_+)^k i\square^{-1} \rightarrow (-q_+)^k \frac{-i}{|q|^2 + i\epsilon} \quad (5.34)$$

where we employ the measure in momentum space [1]:

$$\int \frac{d^4 q}{(2\pi)^4} \quad (5.35)$$

We also get for the identity in space-time:

$$I \rightarrow (2\pi)^4 \delta^{(4)}(q_1 - q_2) \quad (5.36)$$

for the sources:

$$J_{\mathcal{O}} \rightarrow J_{\mathcal{O}}(q_1 - q_2) \quad (5.37)$$

and the differential operators:

$$\begin{aligned} \mathcal{Y}_{s-2}^{\frac{5}{2}}(\vec{\partial}_+, \overleftarrow{\partial}_+) &\rightarrow \mathcal{Y}_{s-2}^{\frac{5}{2}}(q_{2+}, q_{1+}) = q_{1+}(q_{2+} - q_{1+})^{s-2} C_{s-2}^{\frac{5}{2}} \left(\frac{q_{2+} + q_{1+}}{q_{2+} - q_{1+}} \right) q_{2+} \\ \mathcal{Y}_{s-1}^{\frac{3}{2}}(\vec{\partial}_+, \overleftarrow{\partial}_+) &\rightarrow \mathcal{Y}_{s-1}^{\frac{3}{2}}(q_{2+}, q_{1+}) = (q_{2+} - q_{1+})^{s-2} C_{s-1}^{\frac{3}{2}} \left(\frac{q_{2+} + q_{1+}}{q_{2+} - q_{1+}} \right) \\ \mathcal{H}_{s-2}^{\frac{5}{2}}(\vec{\partial}_+, \overleftarrow{\partial}_+) &\rightarrow \mathcal{H}_{s-2}^{\frac{5}{2}}(q_{2+}, q_{1+}) = q_{1+}(q_{2+} - q_{1+})^{s-2} C_{s-2}^{\frac{5}{2}} \left(\frac{q_{2+} + q_{1+}}{q_{2+} - q_{1+}} \right) q_{2+} \\ \mathcal{H}_{s-1}^{\frac{3}{2}}(\vec{\partial}_+, \overleftarrow{\partial}_+) &\rightarrow \mathcal{H}_{s-1}^{\frac{3}{2}}(q_{2+}, q_{1+}) = (q_{2+} - q_{1+})^{s-2} C_{s-1}^{\frac{3}{2}} \left(\frac{q_{2+} + q_{1+}}{q_{2+} - q_{1+}} \right) \\ \mathcal{G}_{s-1}^{(1,2)}(\vec{\partial}_+, \overleftarrow{\partial}_+) &\rightarrow \mathcal{G}_{s-1}^{(1,2)}(q_{2+}, q_{1+}) = i(q_{2+} - q_{1+})^{s-1} P_{s-1}^{(1,2)} \left(\frac{q_{2+} + q_{1+}}{q_{2+} - q_{1+}} \right) q_{2+} \\ \mathcal{G}_{s-1}^{(2,1)}(\vec{\partial}_+, \overleftarrow{\partial}_+) &\rightarrow \mathcal{G}_{s-1}^{(2,1)}(q_{2+}, q_{1+}) = -iq_{1+}(q_{2+} - q_{1+})^{s-1} P_{s-1}^{(2,1)} \left(\frac{q_{2+} + q_{1+}}{q_{2+} - q_{1+}} \right) \end{aligned} \quad (5.38)$$

The explicit expression in the momentum representation [1] follows from eq. (B.7). We report the generating functional restricted to several sectors:

$$\begin{aligned} \mathcal{W}_{\text{conf}}[J_{O^A}, 0, 0, 0, 0, 0] \\ = -(N^2 - 1) \log \text{Det} \left((2\pi)^4 \delta^{(4)}(q_1 - q_2) - \frac{1}{2} \frac{i}{|q_1|^2 + i\epsilon} J_{O_s^A}(q_1 - q_2) \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}}(q_{2+}, q_{1+}) \right) \end{aligned} \quad (5.39)$$

where by a slight abuse of notation we have displayed as argument of the functional determinant the corresponding kernel in the momentum representation [1].

We write the generating functional of the correlators of fermionic operators, M_s and \bar{M}_s , as a single object by means of an equality of determinants (appendix D):

$$\mathcal{W}_{\text{conf}}[0, 0, 0, 0, \bar{J}_M, J_{\bar{M}}] = \log \text{Det} \left(\mathcal{I} + \frac{1}{4} i\partial_+ i\square^{-1} J_{\bar{M}_{s_1}} \otimes \mathcal{G}_{s_1-1}^{(1,2)} i\square^{-1} \bar{J}_{M_{s_2}} \otimes \mathcal{G}_{s_2-1}^{(2,1)} \right) \quad (5.40)$$

Then, we get:

$$\begin{aligned} \mathcal{W}_{\text{conf}} [0, 0, 0, 0, \bar{J}_M, J_{\bar{M}}] &= (N^2 - 1) \log \text{Det} \left((2\pi)^4 \delta^{(4)}(q_1 - q_2) \right. \\ &\quad \left. - \frac{1}{4} \int \frac{d^4 q}{(2\pi)^4} q_{1+} \frac{i}{|q_1|^2 + i\epsilon} J_{\bar{M}s_1}(q_1 - q) \otimes \mathcal{G}_{s_1-1}^{(1,2)}(q_+, q_{1+}) \frac{i}{|q|^2 + i\epsilon} \bar{J}_{M s_2}(q - q_2) \otimes \mathcal{G}_{s_2-1}^{(2,1)}(q_{2+}, q_+) \right) \end{aligned} \quad (5.41)$$

Besides:

$$\begin{aligned} \mathcal{W}_{\text{conf}} [J_{S^{(2)}}] &= \\ &\quad - (N^2 - 1) \log \text{Det} \left((2\pi)^4 \delta^{(4)}(q_1 - q_2) - \frac{1}{2} \frac{i}{|q_1|^2 + i\epsilon} \frac{6}{s+2} J_{S_s^{(2)}}(q_1 - q_2) \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}}(q_{1+}, q_{2+}) \right) \\ &\quad + (N^2 - 1) \log \text{Det} \left((2\pi)^4 \delta^{(4)}(q_1 - q_2) - \frac{1}{2} \frac{i q_{1+}}{|q_1|^2 + i\epsilon} J_{S_s^{(2)}}(q_1 - q_2) \otimes \mathcal{Y}_{s-1}^{\frac{3}{2}}(q_{1+}, q_{2+}) \right) \end{aligned} \quad (5.42)$$

Explicitly:

$$\begin{aligned} \mathcal{W}_{\text{conf}} [J_{S^{(2)}}] &= \\ &\quad - (N^2 - 1) \log \text{Det} \left((2\pi)^4 \delta^{(4)}(q_1 - q_2) - \frac{i}{|q_1|^2 + i\epsilon} \frac{s+1}{4} J_{S_s^{(2)}}(q_1 - q_2) \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} q_{1+}^{s-k-1} q_{2+}^{k+1} \right) \\ &\quad + (N^2 - 1) \log \text{Det} \left((2\pi)^4 \delta^{(4)}(q_1 - q_2) - \frac{i}{|q_1|^2 + i\epsilon} \frac{s+1}{4} J_{S_s^{(2)}}(q_1 - q_2) \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} q_{1+}^{s-k} q_{2+}^k \right) \end{aligned} \quad (5.43)$$

Correspondingly, $\Gamma_{\text{conf}} [J_{S^{(2)}}]$ reads:

$$\begin{aligned} \Gamma_{\text{conf}} [J_{S^{(2)}}] &= \\ &\quad - (N^2 - 1) \log \text{Det} \left(\delta_{s_1 k_1, s_2 k_2} (2\pi)^4 \delta^{(4)}(q_1 - q_2) + \mathcal{D}_{A s_1 k_1, s_2 k_2}^{-1}(q_1) \frac{6}{s_2 + 2} J_{S_{s_2 k_2}^{(2)}}(q_1 - q_2) \right) \\ &\quad + (N^2 - 1) \log \text{Det} \left(\delta_{s_1 k_1, s_2 k_2} (2\pi)^4 \delta^{(4)}(q_1 - q_2) + \mathcal{D}_{\lambda s_1 k_1, s_2 k_2}^{-1}(q_1) J_{S_{s_2 k_2}^{(2)}}(q_1 - q_2) \right) \end{aligned} \quad (5.44)$$

with:

$$\begin{aligned} \mathcal{D}_{A s_1 k_1, s_2 k_2}^{-1}(p) &= \frac{1}{2} \frac{\Gamma(3)\Gamma(s_1 + 3)}{\Gamma(5)\Gamma(s_1 + 1)} \binom{s_1}{k_1} \binom{s_2}{k_2 + 2} p_+^{s_1 - k_1 + k_2} \frac{-i}{|p|^2 + i\epsilon} \\ \mathcal{D}_{\lambda s_1 k_1, s_2 k_2}^{-1}(p) &= \frac{1}{2} \frac{s_1 + 1}{2} \binom{s_1}{k_1} \binom{s_2}{k_2 + 1} p_+^{s_1 - k_1 + k_2} \frac{-i}{|p|^2 + i\epsilon} \end{aligned} \quad (5.45)$$

6 Generating functional of Euclidean conformal correlators

6.1 Analytic continuation to Euclidean space-time

The Minkowskian correlators can be analytically continued [2] to Euclidean space-time by substituting (appendix A):

$$x_+ \rightarrow -ix_z \quad (6.1)$$

and:

$$\frac{1}{|x|^2 - i\epsilon} \rightarrow -\frac{1}{x^2} \quad (6.2)$$

The analytically continued operators read:

- gluon-gluon operators

$$O_s^A \rightarrow (-1)^{s+1} \text{Tr } \partial_z \bar{A}^E (\overrightarrow{\partial}_z + \overleftarrow{\partial}_z)^{s-2} C_{s-2}^{\frac{5}{2}} \left(\frac{\overrightarrow{\partial}_z - \overleftarrow{\partial}_z}{\overrightarrow{\partial}_z + \overleftarrow{\partial}_z} \right) \partial_z A^E = O_s^{AE} \quad (6.3)$$

- gluino-gluino operators

$$O_s^\lambda \rightarrow (-1)^{s-1} \text{Tr } \bar{\lambda}^E (\overrightarrow{\partial}_z + \overleftarrow{\partial}_z)^{s-1} C_{s-1}^{\frac{3}{2}} \left(\frac{\overrightarrow{\partial}_z - \overleftarrow{\partial}_z}{\overrightarrow{\partial}_z + \overleftarrow{\partial}_z} \right) \lambda^E = O_s^{\lambda E} \quad (6.4)$$

- gluon-gluino operators

$$M_s \rightarrow i(-1)^{s-1} \partial_z \text{Tr } A^E (\overrightarrow{\partial}_z + i\overleftarrow{\partial}_z)^{s-1} P_{s-1}^{(2,1)} \left(\frac{\overrightarrow{\partial}_z - \overleftarrow{\partial}_z}{\overrightarrow{\partial}_z + \overleftarrow{\partial}_z} \right) \bar{\lambda}^E = M_s^E \quad (6.5)$$

$$\bar{M}_s \rightarrow i(-1)^{s-1} \text{Tr } \lambda^E (\overrightarrow{\partial}_z + \overleftarrow{\partial}_z)^{s-1} P_{s-1}^{(1,2)} \left(\frac{\overrightarrow{\partial}_z - \overleftarrow{\partial}_z}{\overrightarrow{\partial}_z + \overleftarrow{\partial}_z} \right) \partial_z \bar{A}^E = \bar{M}_s^E \quad (6.6)$$

6.2 Analytic continuation of $\mathcal{W}_{\text{conf}}$

Therefore, the generating functional of Euclidean correlators reads:

$$\begin{aligned}
& \mathcal{W}_{\text{conf}}^E [J_{O'^A E}, J_{\tilde{O}'^A E}, J_{O'^\lambda E}, J_{\tilde{O}'^\lambda E}, \bar{J}_{M'E}, J_{\bar{M}'E}] = \\
& -\frac{N^2 - 1}{2} \log \text{Det} \left(I + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-\vec{\partial}_z)^{s-k-1} \Delta^{-1} (J_{O'_s E} + J_{\tilde{O}'_s E}) (-\vec{\partial}_z)^{k+1} \right) \\
& -\frac{N^2 - 1}{2} \log \text{Det} \left(I + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-\vec{\partial}_z)^{s-k-1} \Delta^{-1} (J_{O'_s E} - J_{\tilde{O}'_s E}) (-\vec{\partial}_z)^{k+1} \right) \\
& +\frac{N^2 - 1}{2} \log \text{Det} \left(I - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-\vec{\partial}_z)^{s-k} \Delta^{-1} (J_{O'_s^\lambda E} + J_{\tilde{O}'_s^\lambda E}) (-\vec{\partial}_z)^k \right) \\
& +\frac{N^2 - 1}{2} \log \text{Det} \left(I - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-\vec{\partial}_z)^{s-k} \Delta^{-1} (J_{O'_s^\lambda E} - J_{\tilde{O}'_s^\lambda E}) (-\vec{\partial}_z)^k \right) \\
& -\frac{N^2 - 1}{2} \log \text{Det} \left[I + \frac{1}{N^2} \left(I - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-\vec{\partial}_z)^{s-k} \Delta^{-1} (J_{O'_s^\lambda E} - J_{\tilde{O}'_s^\lambda E}) (-\vec{\partial}_z)^k \right)^{-1} \right. \\
& \left. \sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1+1}{k_1+2} (-1)^{s_1-1} (-\vec{\partial}_z)^{s_1-k_1} \Delta^{-1} \bar{J}_{M'E_{s_1}} (-\vec{\partial}_z)^{k_1+1} \right. \\
& \left. \left(I + \frac{1}{N} \sum_{k_2=0}^{s_2-2} \binom{s_2}{k_2} \binom{s_2}{k_2+2} (-\vec{\partial}_z)^{s_2-k_2-1} \Delta^{-1} (J_{O'_{s_2}^A E} + J_{\tilde{O}'_{s_2}^A E}) (-\vec{\partial}_z)^{k_2+1} \right)^{-1} \right. \\
& \left. \sum_{k_3=0}^{s_3-1} \binom{s_3+1}{k_3} \binom{s_3}{k_3+1} (-1)^{s_3-1} (-\vec{\partial}_z)^{s_3-k_3} \Delta^{-1} J_{M'E_{s_3}} (-\vec{\partial}_z)^{k_3} \right] \\
& -\frac{N^2 - 1}{2} \log \text{Det} \left[I + \frac{1}{N^2} \left(I - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-\vec{\partial}_z)^{s-k} \Delta^{-1} (J_{O'_s^\lambda E} + J_{\tilde{O}'_s^\lambda E}) (-\vec{\partial}_z)^k \right)^{-1} \right. \\
& \left. \sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1+1}{k_1+2} (-\vec{\partial}_z)^{s_1-k_1} \Delta^{-1} J_{M'E_{s_1}} (-\vec{\partial}_z)^{k_1+1} \right. \\
& \left. \left(I + \frac{1}{N} \sum_{k_2=0}^{s_2-2} \binom{s_2}{k_2} \binom{s_2}{k_2+2} (-\vec{\partial}_z)^{s_2-k_2-1} \Delta^{-1} (J_{O'_{s_2}^A E} - J_{\tilde{O}'_{s_2}^A E}) (-\vec{\partial}_z)^{k_2+1} \right)^{-1} \right. \\
& \left. \sum_{k_3=0}^{s_3-1} \binom{s_3+1}{k_3} \binom{s_3}{k_3+1} (-\vec{\partial}_z)^{s_3-k_3} \Delta^{-1} \bar{J}_{M'E_{s_3}} (-\vec{\partial}_z)^{k_3} \right]
\end{aligned} \tag{6.7}$$

and:

$$\begin{aligned}
& \mathcal{W}_{\text{conf}}^E [J_{O'AE}, J_{\tilde{O}'AE}, J_{O'\lambda E}, J_{\tilde{O}'\lambda E}, \bar{J}_{M'E}, J_{\bar{M}'E}] = \\
& -\frac{N^2-1}{2} \log \text{Det} \left(I + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-\vec{\partial}_z)^{s-k-1} \Delta^{-1} (J_{O'_s AE} + J_{\tilde{O}'_s AE}) (-\vec{\partial}_z)^{k+1} \right) \\
& -\frac{N^2-1}{2} \log \text{Det} \left(I + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-\vec{\partial}_z)^{s-k-1} \Delta^{-1} (J_{O'_s AE} - J_{\tilde{O}'_s AE}) (-\vec{\partial}_z)^{k+1} \right) \\
& + \frac{N^2-1}{2} \log \text{Det} \left(I - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-\vec{\partial}_z)^{s-k} \Delta^{-1} (J_{O'_s \lambda E} + J_{\tilde{O}'_s \lambda E}) (-\vec{\partial}_z)^k \right) \\
& + \frac{N^2-1}{2} \log \text{Det} \left(I - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-\vec{\partial}_z)^{s-k} \Delta^{-1} (J_{O'_s \lambda E} - J_{\tilde{O}'_s \lambda E}) (-\vec{\partial}_z)^k \right) \\
& + \frac{N^2-1}{2} \log \text{Det} \left[I + \frac{1}{N^2} \left(I + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-\vec{\partial}_z)^{s-k-1} \Delta^{-1} (J_{O'_s AE} - J_{\tilde{O}'_s AE}) (-\vec{\partial}_z)^{k+1} \right)^{-1} \right. \\
& \quad \sum_{k_1=0}^{s_1-1} \binom{s_1+1}{k_1} \binom{s_1}{k_1+1} (-1)^{s_1-1} (-\vec{\partial}_z)^{s_1-k_1} \Delta^{-1} J_{M'E_{s_1}} (-\vec{\partial}_z)^{k_1} \\
& \quad \left. \left(I - \frac{1}{N} \sum_{k_2=0}^{s_2-1} \binom{s_2}{k_2} \binom{s_2}{k_2+1} (-\vec{\partial}_z)^{s_2-k_2} \Delta^{-1} (J_{O'_{s_2} \lambda E} + J_{\tilde{O}'_{s_2} \lambda E}) (-\vec{\partial}_z)^{k_2} \right)^{-1} \right. \\
& \quad \left. \sum_{k_3=0}^{s_3-1} \binom{s_3}{k_3} \binom{s_3+1}{k_3+2} (-1)^{s_3-1} (-\vec{\partial}_z)^{s_3-k_3} \Delta^{-1} \bar{J}_{M'E_{s_3}} (-\vec{\partial}_z)^{k_3+1} \right] \\
& + \frac{N^2-1}{2} \log \text{Det} \left[I + \frac{1}{N^2} \left(I + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-\vec{\partial}_z)^{s-k-1} \Delta^{-1} (J_{O'_s AE} + J_{\tilde{O}'_s AE}) (-\vec{\partial}_z)^{k+1} \right)^{-1} \right. \\
& \quad \sum_{k_1=0}^{s_1-1} \binom{s_1+1}{k_1} \binom{s_1}{k_1+1} (-\vec{\partial}_z)^{s_1-k_1} \Delta^{-1} \bar{J}_{M'E_{s_1}} (-\vec{\partial}_z)^{k_1} \\
& \quad \left. \left(I - \frac{1}{N} \sum_{k_2=0}^{s_2-1} \binom{s_2}{k_2} \binom{s_2}{k_2+1} (-\vec{\partial}_z)^{s_2-k_2} \Delta^{-1} (J_{O'_{s_2} \lambda E} - J_{\tilde{O}'_{s_2} \lambda E}) (-\vec{\partial}_z)^{k_2} \right)^{-1} \right. \\
& \quad \left. \sum_{k_3=0}^{s_3-1} \binom{s_3+1}{k_3} \binom{s_3}{k_3+1} (-\vec{\partial}_z)^{s_3-k_3} \Delta^{-1} J_{M'E_{s_3}} (-\vec{\partial}_z)^{k_3} \right] \tag{6.8}
\end{aligned}$$

with:

$$\Delta = \delta_{\mu\nu} \partial_\mu \partial_\nu = \partial_4^2 + \sum_{i=1}^3 \partial_i^2 \tag{6.9}$$

and:

$$-\Delta^{-1} \rightarrow \frac{1}{4\pi^2} \frac{1}{(x-y)^2} \tag{6.10}$$

6.3 Analytic continuation of Γ_{conf}

The Euclidean conformal generating functional is obtained by performing the analytic continuation of eqs.(5.22) and (5.23):

$$\begin{aligned}
& \Gamma_{\text{conf}}^E [j_{O^A E}, j_{\tilde{O}^A E}, j_{O^\lambda E}, j_{\tilde{O}^\lambda E}, \bar{j}_{M^E}, j_{\bar{M}^E}] \\
&= -\frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} + \mathcal{D}_{EA}^{-1} j_{O^A E} + \mathcal{D}_{EA}^{-1} j_{\tilde{O}^A E}] \\
&\quad - \frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} + \mathcal{D}_{EA}^{-1} j_{O^A E} - \mathcal{D}_{EA}^{-1} j_{\tilde{O}^A E}] \\
&\quad + \frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} - \mathcal{D}_{E\lambda}^{-1} j_{O^\lambda E} - \mathcal{D}_{E\lambda}^{-1} j_{\tilde{O}^\lambda E}] \\
&\quad + \frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} - \mathcal{D}_{E\lambda}^{-1} j_{O^\lambda E} + \mathcal{D}_{E\lambda}^{-1} j_{\tilde{O}^\lambda E}] \\
&\quad + \frac{N^2 - 1}{2} \log \text{Det} \left[\mathbb{I} + (\mathbb{I} - \mathcal{D}_{E\lambda}^{-1} j_{O^\lambda E} - \mathcal{D}_{E\lambda}^{-1} j_{\tilde{O}^\lambda E})^{-1} \mathcal{D}_{EM}^{-1} \bar{j}_M \right. \\
&\quad \quad \left. (\mathbb{I} + \mathcal{D}_{EA}^{-1} j_{O^A E} + \mathcal{D}_{EA}^{-1} j_{\tilde{O}^A E})^{-1} \mathcal{D}_{E\bar{M}}^{-1} j_{\bar{M}} \right] \\
&\quad + \frac{N^2 - 1}{2} \log \text{Det} \left[\mathbb{I} + (\mathbb{I} - \mathcal{D}_{E\lambda}^{-1} j_{O^\lambda E} + \mathcal{D}_{E\lambda}^{-1} j_{\tilde{O}^\lambda E})^{-1} \mathcal{D}_{EM}^{-1} j_M \right. \\
&\quad \quad \left. (\mathbb{I} + \mathcal{D}_{EA}^{-1} j_{O^A E} - \mathcal{D}_{EA}^{-1} j_{\tilde{O}^A E})^{-1} \mathcal{D}_{E\bar{M}}^{-1} \bar{j}_M \right]
\end{aligned} \tag{6.11}$$

and:

$$\begin{aligned}
& \Gamma_{\text{conf}}^E [j_{O^A E}, j_{\tilde{O}^A E}, j_{O^\lambda E}, j_{\tilde{O}^\lambda E}, \bar{j}_{M^E}, j_{\bar{M}^E}] \\
&= -\frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} + \mathcal{D}_{EA}^{-1} j_{O^A E} + \mathcal{D}_{EA}^{-1} j_{\tilde{O}^A E}] \\
&\quad - \frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} + \mathcal{D}_{EA}^{-1} j_{O^A E} - \mathcal{D}_{EA}^{-1} j_{\tilde{O}^A E}] \\
&\quad + \frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} - \mathcal{D}_{E\lambda}^{-1} j_{O^\lambda E} - \mathcal{D}_{E\lambda}^{-1} j_{\tilde{O}^\lambda E}] \\
&\quad + \frac{N^2 - 1}{2} \log \text{Det} [\mathbb{I} - \mathcal{D}_{E\lambda}^{-1} j_{O^\lambda E} + \mathcal{D}_{E\lambda}^{-1} j_{\tilde{O}^\lambda E}] \\
&\quad - \frac{N^2 - 1}{2} \log \text{Det} \left[\mathbb{I} + (\mathbb{I} + \mathcal{D}_{EA}^{-1} j_{O^A E} + \mathcal{D}_{EA}^{-1} j_{\tilde{O}^A E})^{-1} \mathcal{D}_{E\bar{M}}^{-1} j_{\bar{M}} \right. \\
&\quad \quad \left. (\mathbb{I} - \mathcal{D}_{E\lambda}^{-1} j_{O^\lambda E} - \mathcal{D}_{E\lambda}^{-1} j_{\tilde{O}^\lambda E})^{-1} \mathcal{D}_{M\bar{E}}^{-1} \bar{j}_M \right] \\
&\quad - \frac{N^2 - 1}{2} \log \text{Det} \left[\mathbb{I} + (\mathbb{I} + \mathcal{D}_{EA}^{-1} j_{O^A E} - \mathcal{D}_{EA}^{-1} j_{\tilde{O}^A E})^{-1} \mathcal{D}_{E\bar{M}}^{-1} \bar{j}_M \right. \\
&\quad \quad \left. (\mathbb{I} - \mathcal{D}_{E\lambda}^{-1} j_{O^\lambda E} + \mathcal{D}_{E\lambda}^{-1} j_{\tilde{O}^\lambda E})^{-1} \mathcal{D}_{EM}^{-1} j_M \right]
\end{aligned} \tag{6.12}$$

with the kernels:

- gluon-gluon kernel

$$\begin{aligned}\mathcal{D}_{EA}^{-1} &= \frac{1}{2} \frac{\Gamma(3)\Gamma(s_1+3)}{\Gamma(5)\Gamma(s_1+1)} \binom{s_1}{k_1} \binom{s_2}{k_2+2} \partial_z^{s_1-k_1+k_2} \Delta^{-1} \\ \rightarrow \mathcal{D}_{EA s_1 k_1, s_2 k_2}^{-1}(x-y) &= -\frac{1}{8\pi^2} \frac{\Gamma(3)\Gamma(s_1+3)}{\Gamma(5)\Gamma(s_1+1)} \binom{s_1}{k_1} \binom{s_2}{k_2+2} \partial_z^{s_1-k_1+k_2} \frac{1}{(x-y)^2}\end{aligned}\quad (6.13)$$

- gluino-gluino kernel

$$\begin{aligned}\mathcal{D}_{E\lambda}^{-1} &= \frac{1}{2} \frac{s_1+1}{2} \binom{s_1}{k_1} \binom{s_2}{k_2+1} \partial_z^{s_1-k_1+k_2} \Delta^{-1} \\ \rightarrow \mathcal{D}_{E\lambda s_1 k_1, s_2 k_2}^{-1}(x-y) &= -\frac{1}{8\pi^2} \frac{s_1+1}{2} \binom{s_1}{k_1} \binom{s_2}{k_2+1} \partial_z^{s_1-k_1+k_2} \frac{1}{(x-y)^2}\end{aligned}\quad (6.14)$$

- gluon-gluino kernels

$$\begin{aligned}\mathcal{D}_{EM}^{-1} &= -\frac{i}{2} \binom{s_1+1}{k_1+2} \binom{s_2+1}{k_2} (-\partial_z)^{s_1-k_1+k_2+1} \Delta^{-1} \\ \rightarrow \mathcal{D}_{EM s_1 k_1, s_2 k_2}^{-1}(x-y) &= \frac{i}{8\pi^2} \binom{s_1+1}{k_1+2} \binom{s_2+1}{k_2} (-\partial_z)^{s_1-k_1+k_2+1} \frac{1}{(x-y)^2}\end{aligned}\quad (6.15)$$

$$\begin{aligned}\mathcal{D}_{E\bar{M}}^{-1} &= -\frac{i}{2} \binom{s_1}{k_1} \binom{s_2}{k_2+1} (-\partial_z)^{s_1-k_1+k_2} \Delta^{-1} \\ \rightarrow \mathcal{D}_{E\bar{M} s_1 k_1, s_2 k_2}^{-1}(x-y) &= \frac{i}{8\pi^2} \binom{s_1}{k_1} \binom{s_2}{k_2+1} (-\partial_z)^{s_1-k_1+k_2} \frac{1}{(x-y)^2}\end{aligned}\quad (6.16)$$

7 RG-improvement of the Euclidean generating functional

7.1 Operator mixing

We briefly summarize the construction of the RG-improved generating functional following [1, 3]. The renormalized Euclidean correlators:

$$\langle \mathcal{O}_{k_1}(x_1) \dots \mathcal{O}_{k_n}(x_n) \rangle = G_{k_1 \dots k_n}^{(n)}(x_1, \dots, x_n; \mu, g(\mu)) \quad (7.1)$$

satisfy the Callan-Symanzik equation:

$$\begin{aligned}&\left(\sum_{\alpha=1}^n x_\alpha \cdot \frac{\partial}{\partial x_\alpha} + \beta(g) \frac{\partial}{\partial g} + \sum_{\alpha=1}^n D_{\mathcal{O}_\alpha} \right) G_{k_1 \dots k_n}^{(n)} + \\ &\sum_a \left(\gamma_{k_1 a}(g) G_{a k_2 \dots k_n}^{(n)} + \gamma_{k_2 a}(g) G_{k_1 a k_3 \dots k_n}^{(n)} \dots + \gamma_{k_n a}(g) G_{k_1 \dots a}^{(n)} \right) = 0\end{aligned}\quad (7.2)$$

with solution:

$$\begin{aligned}G_{k_1 \dots k_n}^{(n)}(\lambda x_1, \dots, \lambda x_n; \mu, g(\mu)) &= \\ &= \sum_{j_1 \dots j_n} Z_{k_1 j_1}(\lambda) \dots Z_{k_n j_n}(\lambda) \lambda^{-\sum_{i=1}^n D_{\mathcal{O}_i}} G_{j_1 \dots j_n}^{(n)}(x_1, \dots, x_n; \mu, g\left(\frac{\mu}{\lambda}\right))\end{aligned}\quad (7.3)$$

where $D_{\mathcal{O}_i}$ is the canonical dimension of $\mathcal{O}_i(x)$ and $\gamma(g) = \gamma_0 g^2 + \dots$ the matrix of the anomalous dimensions, with:

$$\left(\frac{\partial}{\partial g} + \frac{\gamma(g)}{\beta(g)} \right) Z(\lambda) = 0 \quad (7.4)$$

in matrix notation, and:

$$Z(\lambda) = P \exp \left(\int_{g(\mu)}^{g(\frac{\mu}{\lambda})} \frac{\gamma(g')}{\beta(g')} dg' \right) \quad (7.5)$$

The corresponding UV asymptotics as $\lambda \rightarrow 0$ reads:

$$\begin{aligned} & G_{k_1 \dots k_n}^{(n)}(\lambda x_1, \dots, \lambda x_n; \mu, g(\mu)) \\ & \sim \sum_{j_1 \dots j_n} Z_{k_1 j_1}(\lambda) \dots Z_{k_n j_n}(\lambda) \lambda^{-\sum_{i=1}^n D_{\mathcal{O}_i}} G_{\text{conf } j_1 \dots j_n}^{(n)}(x_1, \dots, x_n) \end{aligned} \quad (7.6)$$

The above result greatly simplifies if a renormalization scheme exists where $Z(\lambda)$ is diagonalizable to all orders of perturbation theory, and specifically one-loop exact, with eigenvalues $Z_{\mathcal{O}_i}(\lambda)$ [3]:

$$Z_{\mathcal{O}_i}(\lambda) = \left(\frac{g(\mu)}{g(\frac{\mu}{\lambda})} \right)^{\frac{\gamma_0 \mathcal{O}_i}{\beta_0}} \quad (7.7)$$

where $\gamma_0 \mathcal{O}_i$ are the eigenvalues of γ_0 . In this case the UV asymptotics reads:

$$\langle \mathcal{O}_{j_1}(\lambda x_1) \dots \mathcal{O}_{j_n}(\lambda x_n) \rangle \sim \frac{Z_{\mathcal{O}_{j_1}}(\lambda) \dots Z_{\mathcal{O}_{j_n}}(\lambda)}{\lambda^{D_{\mathcal{O}_1} + \dots + D_{\mathcal{O}_n}}} G_{\text{conf } j_1 \dots j_n}^{(n)}(x_1, \dots, x_n) \quad (7.8)$$

with:

$$\begin{aligned} g^2\left(\frac{\mu}{\lambda}\right) & \sim \frac{1}{\beta_0 \log\left(\frac{\mu^2}{\lambda^2 \Lambda_{YM}^2}\right)} \left(1 - \frac{\beta_1}{\beta_0^2} \frac{\log \log\left(\frac{\mu^2}{\lambda^2 \Lambda_{YM}^2}\right)}{\log\left(\frac{\mu^2}{\lambda^2 \Lambda_{YM}^2}\right)} \right) \\ & \sim \frac{1}{\beta_0 \log\left(\frac{1}{\lambda^2}\right)} \left(1 - \frac{\beta_1}{\beta_0^2} \frac{\log \log\left(\frac{1}{\lambda^2}\right)}{\log\left(\frac{1}{\lambda^2}\right)} \right) \end{aligned} \quad (7.9)$$

as $\lambda \rightarrow 0$. We refer to the above scheme as nonresonant diagonal [3] according to the Poincaré-Dulac theorem that is involved in the following differential-geometric interpretation of operator mixing [3]. We interpret a finite change of basis of the renormalized operators:

$$\mathcal{O}'(x) = S(g)\mathcal{O}(x) \quad (7.10)$$

in matrix notation as a real-analytic invertible gauge transformation $S(g)$ that depends on g . Then, the matrix:

$$A(g) = -\frac{\gamma(g)}{\beta(g)} = \frac{1}{g} \left(\frac{\gamma_0}{\beta_0} + \dots \right) \quad (7.11)$$

that enters the differential equation for $Z(\lambda)$:

$$\left(\frac{\partial}{\partial g} - A(g) \right) Z(\lambda) = 0 \quad (7.12)$$

defines a connection $A(g)$:

$$A(g) = \frac{1}{g} \left(A_0 + \sum_{n=1}^{\infty} A_{2n} g^{2n} \right) \quad (7.13)$$

with a regular singularity at $g = 0$ that transforms as:

$$A'(g) = S(g)A(g)S^{-1}(g) + \frac{\partial S(g)}{\partial g}S^{-1}(g) \quad (7.14)$$

under the gauge transformation $S(g)$, with:

$$\mathcal{D} = \frac{\partial}{\partial g} - A(g) \quad (7.15)$$

the corresponding covariant derivative. Consequently, $Z(\lambda)$ is a Wilson line that transforms as:

$$Z'(\lambda) = S(g(\mu))Z(\lambda)S^{-1}(g(\frac{\mu}{\lambda})) \quad (7.16)$$

It follows from the Poincaré-Dulac theorem [3] that, if any two eigenvalues $\lambda_1, \lambda_2, \dots$ of the matrix $\frac{\gamma_0}{\beta_0}$, in nonincreasing order $\lambda_1 \geq \lambda_2 \geq \dots$, do not differ by a positive even integer:

$$\lambda_i - \lambda_j - 2k \neq 0 \quad (7.17)$$

for $i \leq j$ and k a positive integer – i.e., they are nonresonant – then a gauge transformation exists that sets $A(g)$ in the canonical nonresonant form [3]:

$$A'(g) = \frac{\gamma_0}{\beta_0} \frac{1}{g} \quad (7.18)$$

that is one-loop exact to all orders of perturbation theory (appendix F). Hence, if in addition $\frac{\gamma_0}{\beta_0}$ is diagonalizable, eq. (7.7) follows.

7.2 Anomalous dimensions of twist-2 operators in $\mathcal{N} = 1$ SUSY YM theory

The eigenvalues of γ_0 for $\mathcal{O}_s = S^{(1)}, S^{(2)}, \tilde{S}^{(1)}, \tilde{S}^{(2)}, M_s, \bar{M}_s$ read [6]:

$$\gamma_{0\mathcal{O}_s} = \frac{1}{4\pi^2} \left(\tilde{\gamma}_{\mathcal{O}_s}^0 - \frac{3}{2} \right) \quad (7.19)$$

with:

$$\begin{aligned} \tilde{\gamma}_{0S_s^{(1)}} &= \psi(s+2) + \psi(s-1) - 2\psi(1) - \frac{2(-1)^s}{(s+1)s(s-1)} \\ \tilde{\gamma}_{0S_s^{(2)}} &= \psi(s+3) + \psi(s) - 2\psi(1) + \frac{2(-1)^s}{(s+2)(s+1)s} \end{aligned} \quad (7.20)$$

and [6, 8]:

$$\begin{aligned} \tilde{\gamma}_{0\tilde{S}_s^{(i)}} &= \tilde{\gamma}_{0S_s^{(i)}} \\ \tilde{\gamma}_{0M_s} &= \tilde{\gamma}_{0S_s^{(2)}} \end{aligned} \quad (7.21)$$

Besides [6]:

$$\gamma_{0\tilde{O}_1^\lambda} = \frac{1}{4\pi^2} \frac{2}{3} = \frac{1}{6\pi^2} \quad (7.22)$$

We have verified that the first 10^4 eigenvalues of $\frac{\gamma_0}{\beta_0}$ are nonresonant, with $\beta_0 = \frac{3}{(4\pi)^2}$.

7.3 Conformal generating functional of correlators of supermultiplet operators

By noticing that:

$$S_s^{(1)'} E J_{S_s^{(1)'} E} + S_s^{(2)'} E J_{S_s^{(2)'} E} = 6 O_s'^{A E} \left(\frac{J_{S_s^{(1)'} E}}{s-1} + \frac{J_{S_s^{(2)'} E}}{s+2} \right) + O_s'^{\lambda E} \left(-J_{S_s^{(1)'} E} + J_{S_s^{(2)'} E} \right) \quad (7.23)$$

and:

$$\tilde{S}_s^{(1)'} E J_{\tilde{S}_s^{(1)'} E} + \tilde{S}_s^{(2)'} E J_{\tilde{S}_s^{(2)'} E} = -6 \tilde{O}_s'^{A E} \left(\frac{J_{\tilde{S}_s^{(1)'} E}}{s-1} + \frac{J_{\tilde{S}_s^{(2)'} E}}{s+2} \right) + \tilde{O}_s'^{\lambda E} \left(-J_{\tilde{S}_s^{(1)'} E} + J_{\tilde{S}_s^{(2)'} E} \right) \quad (7.24)$$

so that:

$$\begin{aligned} J_{O_s'^{A E}} &= 6 \left(\frac{J_{S_s^{(1)'} E}}{s-1} + \frac{J_{S_s^{(2)'} E}}{s+2} \right) \\ J_{\tilde{O}_s'^{A E}} &= -6 \left(\frac{J_{\tilde{S}_s^{(1)'} E}}{s-1} + \frac{J_{\tilde{S}_s^{(2)'} E}}{s+2} \right) \\ J_{O_s'^{\lambda E}} &= -J_{S_s^{(1)'} E} + J_{S_s^{(2)'} E} \\ J_{\tilde{O}_s'^{\lambda E}} &= -J_{\tilde{S}_s^{(1)'} E} + J_{\tilde{S}_s^{(2)'} E} \end{aligned} \quad (7.25)$$

and substituting the above formulas into eqs. (6.7) and (6.8), we obtain for the generating functional of Euclidean conformal correlators of supermultiplet operators:

$$\begin{aligned}
& \mathcal{W}_{\text{conf}}^E \left[J_{\tilde{O}_1^\lambda E}, J_{S^{(1)\prime} E}, J_{\tilde{S}^{(1)\prime} E}, J_{S^{(2)\prime} E}, J_{\tilde{S}^{(2)\prime} E}, \bar{J}_{M'E}, J_{\bar{M}'E} \right] = \\
& -\frac{N^2-1}{2} \log \text{Det} \left(I + \frac{1}{N} 6 \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-\vec{\partial}_z)^{s-k-1} \Delta^{-1} \left(\frac{J_{S_s^{(1)\prime} E}}{s-1} + \frac{J_{S_s^{(2)\prime} E}}{s+2} - \frac{J_{\tilde{S}_s^{(1)\prime} E}}{s-1} - \frac{J_{\tilde{S}_s^{(2)\prime} E}}{s+2} \right) (-\vec{\partial}_z)^{k+1} \right) \\
& -\frac{N^2-1}{2} \log \text{Det} \left(I + \frac{1}{N} 6 \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-\vec{\partial}_z)^{s-k-1} \Delta^{-1} \left(\frac{J_{S_s^{(1)\prime} E}}{s-1} + \frac{J_{S_s^{(2)\prime} E}}{s+2} + \frac{J_{\tilde{S}_s^{(1)\prime} E}}{s-1} + \frac{J_{\tilde{S}_s^{(2)\prime} E}}{s+2} \right) (-\vec{\partial}_z)^{k+1} \right) \\
& + \frac{N^2-1}{2} \log \text{Det} \left(I - \frac{1}{N} \Delta^{-1} J_{\tilde{O}_1^\lambda E} \right. \\
& \quad \left. - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-\vec{\partial}_z)^{s-k} \Delta^{-1} (-J_{S_s^{(1)\prime} E} + J_{S_s^{(2)\prime} E} - J_{\tilde{S}_s^{(1)\prime} E} + J_{\tilde{S}_s^{(2)\prime} E}) (-\vec{\partial}_z)^k \right) \\
& + \frac{N^2-1}{2} \log \text{Det} \left(I + \frac{1}{N} \Delta^{-1} J_{\tilde{O}_1^\lambda E} \right. \\
& \quad \left. - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-\vec{\partial}_z)^{s-k} \Delta^{-1} (-J_{S_s^{(1)\prime} E} + J_{S_s^{(2)\prime} E} + J_{\tilde{S}_s^{(1)\prime} E} - J_{\tilde{S}_s^{(2)\prime} E}) (-\vec{\partial}_z)^k \right) \\
& - \frac{N^2-1}{2} \log \text{Det} \left[I \right. \\
& \quad \left. + \frac{1}{N^2} \left(I + \frac{1}{N} \Delta^{-1} J_{\tilde{O}_1^\lambda E} - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-\vec{\partial}_z)^{s-k} \Delta^{-1} (-J_{S_s^{(1)\prime} E} + J_{S_s^{(2)\prime} E} + J_{\tilde{S}_s^{(1)\prime} E} - J_{\tilde{S}_s^{(2)\prime} E}) (-\vec{\partial}_z)^k \right)^{-1} \right. \\
& \quad \left. \sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1+1}{k_1+2} (-1)^{s_1-1} (-\vec{\partial}_z)^{s_1-k_1} \Delta^{-1} \bar{J}_{M'_{s_1} E} (-\vec{\partial}_z)^{k_1+1} \right. \\
& \quad \left. \left(I + \frac{1}{N} 6 \sum_{k_2=0}^{s_2-2} \binom{s_2}{k_2} \binom{s_2}{k_2+2} (-\vec{\partial}_z)^{s_2-k_2-1} \Delta^{-1} \left(\frac{J_{S_{s_2}^{(1)\prime} E}}{s_2-1} + \frac{J_{S_{s_2}^{(2)\prime} E}}{s_2+2} - \frac{J_{\tilde{S}_{s_2}^{(1)\prime} E}}{s_2-1} - \frac{J_{\tilde{S}_{s_2}^{(2)\prime} E}}{s_2+2} \right) (-\vec{\partial}_z)^{k_2+1} \right)^{-1} \right. \\
& \quad \left. \sum_{k_3=0}^{s_3-1} \binom{s_3+1}{k_3} \binom{s_3}{k_3+1} (-1)^{s_3-1} (-\vec{\partial}_z)^{s_3-k_3} \Delta^{-1} J_{\bar{M}'_{s_3} E} (-\vec{\partial}_z)^{k_3} \right] \\
& - \frac{N^2-1}{2} \log \text{Det} \left[I \right. \\
& \quad \left. + \frac{1}{N^2} \left(I - \frac{1}{N} \Delta^{-1} J_{\tilde{O}_1^\lambda E} - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-\vec{\partial}_z)^{s-k} \Delta^{-1} (-J_{S_s^{(1)\prime} E} + J_{S_s^{(2)\prime} E} - J_{\tilde{S}_s^{(1)\prime} E} + J_{\tilde{S}_s^{(2)\prime} E}) (-\vec{\partial}_z)^k \right)^{-1} \right. \\
& \quad \left. \sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1+1}{k_1+2} (-\vec{\partial}_z)^{s_1-k_1} \Delta^{-1} J_{\bar{M}'_{s_1} E} (-\vec{\partial}_z)^{k_1+1} \right. \\
& \quad \left. \left(I + \frac{1}{N} \frac{6}{s_2-1} \sum_{k=0}^{s_2-2} \binom{s_2}{k} \binom{s_2}{k+2} (-\vec{\partial}_z)^{s_2-k_2-1} \Delta^{-1} \left(\frac{J_{S_{s_2}^{(1)\prime} E}}{s_2-1} + \frac{J_{S_{s_2}^{(2)\prime} E}}{s_2+2} + \frac{J_{\tilde{S}_{s_2}^{(1)\prime} E}}{s_2-1} + \frac{J_{\tilde{S}_{s_2}^{(2)\prime} E}}{s_2+2} \right) (-\vec{\partial}_z)^{k_2+1} \right)^{-1} \right. \\
& \quad \left. \sum_{k_3=0}^{s_3-1} \binom{s_3+1}{k_3} \binom{s_3}{k_3+1} (-\vec{\partial}_z)^{s_3-k_3} \Delta^{-1} \bar{J}_{M'_{s_3} E} (-\vec{\partial}_z)^{k_3} \right]
\end{aligned} \tag{7.26}$$

and:

$$\begin{aligned}
& \mathcal{W}_{\text{conf}}^E \left[J_{\tilde{O}_1^\lambda E}, J_{S^{(1)'} E}, J_{\tilde{S}^{(1)'} E}, J_{S^{(2)'} E}, J_{\tilde{S}^{(2)'} E}, \bar{J}_{M'E}, J_{\bar{M}'E} \right] = \\
& -\frac{N^2-1}{2} \log \text{Det} \left(I + \frac{1}{N} 6 \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-\vec{\partial}_z)^{s-k-1} \Delta^{-1} \left(\frac{J_{S_s^{(1)'} E}}{s-1} + \frac{J_{S_s^{(2)'} E}}{s+2} - \frac{J_{\tilde{S}_s^{(1)'} E}}{s-1} - \frac{J_{\tilde{S}_s^{(2)'} E}}{s+2} \right) (-\vec{\partial}_z)^{k+1} \right) \\
& -\frac{N^2-1}{2} \log \text{Det} \left(I + \frac{1}{N} 6 \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-\vec{\partial}_z)^{s-k-1} \Delta^{-1} \left(\frac{J_{S_s^{(1)'} E}}{s-1} + \frac{J_{S_s^{(2)'} E}}{s+2} + \frac{J_{\tilde{S}_s^{(1)'} E}}{s-1} - \frac{J_{\tilde{S}_s^{(2)'} E}}{s+2} \right) (-\vec{\partial}_z)^{k+1} \right) \\
& + \frac{N^2-1}{2} \log \text{Det} \left(I - \frac{1}{N} \Delta^{-1} J_{\tilde{O}_1^\lambda E} \right. \\
& \left. - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-\vec{\partial}_z)^{s-k} \Delta^{-1} (-J_{S_s^{(1)'} E} + J_{S_s^{(2)'} E} - J_{\tilde{S}_s^{(1)'} E} + J_{\tilde{S}_s^{(2)'} E}) (-\vec{\partial}_z)^k \right) \\
& + \frac{N^2-1}{2} \log \text{Det} \left(I + \frac{1}{N} \Delta^{-1} J_{\tilde{O}_1^\lambda E} \right. \\
& \left. - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-\vec{\partial}_z)^{s-k} \Delta^{-1} (-J_{S_s^{(1)'} E} + J_{S_s^{(2)'} E} + J_{\tilde{S}_s^{(1)'} E} - J_{\tilde{S}_s^{(2)'} E}) (-\vec{\partial}_z)^k \right) \\
& + \frac{N^2-1}{2} \log \text{Det} \left[I + \frac{1}{N^2} \left(I + \frac{1}{N} 6 \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-\vec{\partial}_z)^{s-k-1} \Delta^{-1} \left(\frac{J_{S_s^{(1)'} E}}{s-1} + \frac{J_{S_s^{(2)'} E}}{s+2} - \frac{J_{\tilde{S}_s^{(1)'} E}}{s-1} - \frac{J_{\tilde{S}_s^{(2)'} E}}{s+2} \right) (-\vec{\partial}_z)^{k+1} \right)^{-1} \right. \\
& \left. \sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1+1}{k_1+2} (-1)^{s_1-1} (-\vec{\partial}_z)^{s_1-k_1} \Delta^{-1} J_{\tilde{M}'_{s_1} E} (-\vec{\partial}_z)^{k_1+1} \right. \\
& \left. \left(I + \frac{1}{N} \Delta^{-1} J_{\tilde{O}_1^\lambda E} - \frac{1}{N} \sum_{k_2=0}^{s_2-1} \binom{s_2}{k_2} \binom{s_2}{k_2+1} (-\vec{\partial}_z)^{s_2-k_2} \Delta^{-1} (-J_{S_{s_2}^{(1)'} E} + J_{S_{s_2}^{(2)'} E} + J_{\tilde{S}_{s_2}^{(1)'} E} - J_{\tilde{S}_{s_2}^{(2)'} E}) (-\vec{\partial}_z)^{k_2} \right)^{-1} \right. \\
& \left. \sum_{k_3=0}^{s_3-1} \binom{s_3}{k_3} \binom{s_3}{k_3+1} (-1)^{s_3-1} (-\vec{\partial}_z)^{s_3-k_3} \Delta^{-1} \bar{J}_{M'_{s_3} E} (-\vec{\partial}_z)^{k_3} \right] \\
& + \frac{N^2-1}{2} \log \text{Det} \left[I + \frac{1}{N^2} \left(I + \frac{1}{N} 6 \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-\vec{\partial}_z)^{s-k-1} \Delta^{-1} \left(\frac{J_{S_s^{(1)'} E}}{s-1} + \frac{J_{S_s^{(2)'} E}}{s+2} + \frac{J_{\tilde{S}_s^{(1)'} E}}{s-1} + \frac{J_{\tilde{S}_s^{(2)'} E}}{s+2} \right) (-\vec{\partial}_z)^{k+1} \right)^{-1} \right. \\
& \left. \sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1+1}{k_1+2} (-\vec{\partial}_z)^{s_1-k_1} \Delta^{-1} \bar{J}_{M'_{s_1} E} (-\vec{\partial}_z)^{k_1+1} \right. \\
& \left. \left(I - \frac{1}{N} \Delta^{-1} J_{\tilde{O}_1^\lambda E} - \frac{1}{N} \sum_{k_2=0}^{s_2-1} \binom{s_2}{k_2} \binom{s_2}{k_2+1} (-\vec{\partial}_z)^{s_2-k_2} \Delta^{-1} (-J_{S_{s_2}^{(1)'} E} + J_{S_{s_2}^{(2)'} E} - J_{\tilde{S}_{s_2}^{(1)'} E} + J_{\tilde{S}_{s_2}^{(2)'} E}) (-\vec{\partial}_z)^{k_2} \right)^{-1} \right. \\
& \left. \sum_{k_3=0}^{s_3-1} \binom{s_3}{k_3} \binom{s_3}{k_3+1} (-\vec{\partial}_z)^{s_3-k_3} \Delta^{-1} J_{\tilde{M}'_{s_3} E} (-\vec{\partial}_z)^{k_3} \right] \tag{7.27}
\end{aligned}$$

7.4 RG-improved generating functional of correlators of supermultiplet operators

The generating functional of the Euclidean correlators decomposes into its planar $\mathcal{W}_{\text{sphere}}^E$ and LO-nonplanar $\mathcal{W}_{\text{torus}}^E$ contributions [1, 4]. The corresponding UV asymptotics as $\lambda \rightarrow 0$ of $\mathcal{W}_{\text{sphere}}^E$ and $\mathcal{W}_{\text{torus}}^E$ follows, with:

$$\mathcal{W}_{\text{asym sphere}}^E[j_{\mathcal{O}'E}, \lambda] = -N^2 \mathcal{W}_{\text{asym torus}}^E[j_{\mathcal{O}'E}, \lambda] \tag{7.28}$$

where:

$$\begin{aligned}
& \mathcal{W}_{\text{asym torus}}^E \left[J_{\tilde{O}_1^\lambda E}, J_{S^{(1)'} E}, J_{\tilde{S}^{(1)'} E}, J_{S^{(2)'} E}, J_{\tilde{S}^{(2)'} E}, \bar{J}_{M'E}, \bar{J}_{\tilde{M}'E}, \lambda \right] = \\
& + \frac{1}{2} \log \text{Det} \left(I \right. \\
& + \frac{1}{N} 6 \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-\vec{\partial}_z)^{s-k-1} \frac{\Delta^{-1}}{\lambda^{s+2}} (Z_{S^{(1)'} E}(\lambda) \frac{J_{S_s^{(1)'} E}}{s-1} + Z_{S^{(2)'} E}(\lambda) \frac{J_{S_s^{(2)'} E}}{s+2} - Z_{\tilde{S}^{(1)'} E}(\lambda) \frac{J_{\tilde{S}_s^{(1)'} E}}{s-1} - Z_{\tilde{S}^{(2)'} E}(\lambda) \frac{J_{\tilde{S}_s^{(2)'} E}}{s+2}) (-\vec{\partial}_z)^{k+1} \\
& + \frac{1}{2} \log \text{Det} \left(I \right. \\
& + \frac{1}{N} 6 \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-\vec{\partial}_z)^{s-k-1} \frac{\Delta^{-1}}{\lambda^{s+2}} (Z_{S^{(1)'} E}(\lambda) \frac{J_{S_s^{(1)'} E}}{s-1} + Z_{S^{(2)'} E}(\lambda) \frac{J_{S_s^{(2)'} E}}{s+2} + Z_{\tilde{S}^{(1)'} E}(\lambda) \frac{J_{\tilde{S}_s^{(1)'} E}}{s-1} + Z_{\tilde{S}^{(2)'} E}(\lambda) \frac{J_{\tilde{S}_s^{(2)'} E}}{s+2}) (-\vec{\partial}_z)^{k+1} \\
& - \frac{1}{2} \log \text{Det} \left(I - \frac{1}{N} \frac{Z_{\tilde{O}_1^\lambda}(\lambda)}{\lambda^3} \Delta^{-1} J_{\tilde{O}_1^\lambda E} \right. \\
& - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-\vec{\partial}_z)^{s-k} \frac{\Delta^{-1}}{\lambda^{s+2}} (-Z_{S^{(1)'} E}(\lambda) J_{S_s^{(1)'} E} + Z_{S^{(2)'} E}(\lambda) J_{S_s^{(2)'} E} - Z_{\tilde{S}^{(1)'} E}(\lambda) J_{\tilde{S}_s^{(1)'} E} + Z_{\tilde{S}^{(2)'} E}(\lambda) J_{\tilde{S}_s^{(2)'} E}) (-\vec{\partial}_z)^k \\
& - \frac{1}{2} \log \text{Det} \left(I + \frac{1}{N} \frac{Z_{\tilde{O}_1^\lambda}(\lambda)}{\lambda^3} \Delta^{-1} J_{\tilde{O}_1^\lambda E} \right. \\
& - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-\vec{\partial}_z)^{s-k} \frac{\Delta^{-1}}{\lambda^{s+2}} (-Z_{S^{(1)'} E}(\lambda) J_{S_s^{(1)'} E} + Z_{S^{(2)'} E}(\lambda) J_{S_s^{(2)'} E} + Z_{\tilde{S}^{(1)'} E}(\lambda) J_{\tilde{S}_s^{(1)'} E} - Z_{\tilde{S}^{(2)'} E}(\lambda) J_{\tilde{S}_s^{(2)'} E}) (-\vec{\partial}_z)^k \\
& + \frac{1}{2} \log \text{Det} \left[I + \frac{1}{N^2} \left(I + \frac{1}{N} \frac{Z_{\tilde{O}_1^\lambda}(\lambda)}{\lambda^3} \Delta^{-1} J_{\tilde{O}_1^\lambda E} \right. \right. \\
& - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-\vec{\partial}_z)^{s-k} \frac{\Delta^{-1}}{\lambda^{s+2}} (-Z_{S^{(1)'} E}(\lambda) J_{S_s^{(1)'} E} + Z_{S^{(2)'} E}(\lambda) J_{S_s^{(2)'} E} + Z_{\tilde{S}^{(1)'} E}(\lambda) J_{\tilde{S}_s^{(1)'} E} - Z_{\tilde{S}^{(2)'} E}(\lambda) J_{\tilde{S}_s^{(2)'} E}) (-\vec{\partial}_z)^k \left. \right]^{-1} \\
& \sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1+1}{k_1+2} (-1)^{s_1-1} (-\vec{\partial}_z)^{s_1-k_1} \frac{\Delta^{-1}}{\lambda^{s_1+2}} Z_M(\lambda) \bar{J}_{M'_{s_1} E} (-\vec{\partial}_z)^{k_1+1} \\
& \left(I + \frac{1}{N} 6 \sum_{k_2=0}^{s_2-2} \binom{s_2}{k_2} \binom{s_2}{k_2+2} (-\vec{\partial}_z)^{s_2-k_2-1} \frac{\Delta^{-1}}{\lambda^{s_2+2}} (Z_{S^{(1)'} E}(\lambda) \frac{J_{S_{s_2}^{(1)'} E}}{s_2-1} + Z_{S^{(2)'} E}(\lambda) \frac{J_{S_{s_2}^{(2)'} E}}{s_2+2} - Z_{\tilde{S}^{(1)'} E}(\lambda) \frac{J_{\tilde{S}_{s_2}^{(1)'} E}}{s_2-1} - Z_{\tilde{S}^{(2)'} E}(\lambda) \frac{J_{\tilde{S}_{s_2}^{(2)'} E}}{s_2+2}) (-\vec{\partial}_z)^{k_2+1} \right)^{-1} \\
& \sum_{k_3=0}^{s_3-1} \binom{s_3+1}{k_3} \binom{s_3}{k_3+1} (-1)^{s_3-1} (-\vec{\partial}_z)^{s_3-k_3} \frac{\Delta^{-1}}{\lambda^{s_3+2}} Z_M(\lambda) J_{\tilde{M}'_{s_3} E} (-\vec{\partial}_z)^{k_3} \\
& + \frac{1}{2} \log \text{Det} \left[I + \frac{1}{N^2} \left(I - \frac{1}{N} \frac{Z_{\tilde{O}_1^\lambda}(\lambda)}{\lambda^3} \Delta^{-1} J_{\tilde{O}_1^\lambda E} \right. \right. \\
& - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-\vec{\partial}_z)^{s-k} \frac{\Delta^{-1}}{\lambda^{s+2}} (-Z_{S^{(1)'} E}(\lambda) J_{S_s^{(1)'} E} + Z_{S^{(2)'} E}(\lambda) J_{S_s^{(2)'} E} - Z_{\tilde{S}^{(1)'} E}(\lambda) J_{\tilde{S}_s^{(1)'} E} + Z_{\tilde{S}^{(2)'} E}(\lambda) J_{\tilde{S}_s^{(2)'} E}) (-\vec{\partial}_z)^k \left. \right]^{-1} \\
& \sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1+1}{k_1+2} (-\vec{\partial}_z)^{s_1-k_1} \frac{\Delta^{-1}}{\lambda^{s_1+2}} Z_M(\lambda) J_{\tilde{M}'_{s_1} E} (-\vec{\partial}_z)^{k_1+1} \\
& \left(I + \frac{1}{N} 6 \sum_{k_2=0}^{s_2-2} \binom{s_2}{k_2} \binom{s_2}{k_2+2} (-\vec{\partial}_z)^{s_2-k_2-1} \frac{\Delta^{-1}}{\lambda^{s_2+2}} (Z_{S^{(1)'} E}(\lambda) \frac{J_{S_{s_2}^{(1)'} E}}{s_2-1} + Z_{S^{(2)'} E}(\lambda) \frac{J_{S_{s_2}^{(2)'} E}}{s_2+2} + Z_{\tilde{S}^{(1)'} E}(\lambda) \frac{J_{\tilde{S}_{s_2}^{(1)'} E}}{s_2-1} + Z_{\tilde{S}^{(2)'} E}(\lambda) \frac{J_{\tilde{S}_{s_2}^{(2)'} E}}{s_2+2}) (-\vec{\partial}_z)^{k_2+1} \right)^{-1} \\
& \sum_{k_3=0}^{s_3-1} \binom{s_3+1}{k_3} \binom{s_3}{k_3+1} (-\vec{\partial}_z)^{s_3-k_3} \frac{\Delta^{-1}}{\lambda^{s_3+2}} Z_M(\lambda) \bar{J}_{M'_{s_3} E} (-\vec{\partial}_z)^{k_3} \quad (7.29)
\end{aligned}$$

and:

$$\begin{aligned}
& \mathcal{W}_{\text{asym torus}}^E \left[J_{\tilde{O}_1^\lambda E}, J_{S^{(1)'} E}, J_{\tilde{S}^{(1)'} E}, J_{S^{(2)'} E}, J_{\tilde{S}^{(2)'} E}, \bar{J}_{M' E}, J_{\tilde{M}' E}, \lambda \right] = \\
& + \frac{1}{2} \log \text{Det} \left(I \right. \\
& + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-\vec{\partial}_z)^{s-k-1} \frac{\Delta^{-1}}{\lambda^{s+2}} (Z_{S^{(1)'} E}(\lambda) \frac{J_{S_s^{(1)'} E}}{s-1} + Z_{S^{(2)'} E}(\lambda) \frac{J_{S_s^{(2)'} E}}{s+2} - Z_{\tilde{S}^{(1)'} E}(\lambda) \frac{J_{\tilde{S}_s^{(1)'} E}}{s-1} - Z_{\tilde{S}^{(2)'} E}(\lambda) \frac{J_{\tilde{S}_s^{(2)'} E}}{s+2}) (-\vec{\partial}_z)^{k+1} \\
& + \frac{1}{2} \log \text{Det} \left(I \right. \\
& + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-\vec{\partial}_z)^{s-k-1} \frac{\Delta^{-1}}{\lambda^{s+2}} (Z_{S^{(1)'} E}(\lambda) \frac{J_{S_s^{(1)'} E}}{s-1} + Z_{S^{(2)'} E}(\lambda) \frac{J_{S_s^{(2)'} E}}{s+2} + Z_{\tilde{S}^{(1)'} E}(\lambda) \frac{J_{\tilde{S}_s^{(1)'} E}}{s-1} + Z_{\tilde{S}^{(2)'} E}(\lambda) \frac{J_{\tilde{S}_s^{(2)'} E}}{s+2}) (-\vec{\partial}_z)^{k+1} \\
& - \frac{1}{2} \log \text{Det} \left(I - \frac{1}{N} \frac{Z_{\tilde{O}_1^\lambda}(\lambda)}{\lambda^3} \Delta^{-1} J_{\tilde{O}_1^\lambda E} \right. \\
& - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-\vec{\partial}_z)^{s-k} \frac{\Delta^{-1}}{\lambda^{s+2}} (-Z_{S^{(1)'} E}(\lambda) J_{S_s^{(1)'} E} + Z_{S^{(2)'} E}(\lambda) J_{S_s^{(2)'} E} - Z_{\tilde{S}^{(1)'} E}(\lambda) J_{\tilde{S}_s^{(1)'} E} + Z_{\tilde{S}^{(2)'} E}(\lambda) J_{\tilde{S}_s^{(2)'} E}) (-\vec{\partial}_z)^k \\
& - \frac{1}{2} \log \text{Det} \left(I + \frac{1}{N} \frac{Z_{\tilde{O}_1^\lambda}(\lambda)}{\lambda^3} \Delta^{-1} J_{\tilde{O}_1^\lambda E} \right. \\
& - \frac{1}{N} \sum_{k=0}^{s-1} \binom{s}{k} \binom{s}{k+1} (-\vec{\partial}_z)^{s-k} \frac{\Delta^{-1}}{\lambda^{s+2}} (-Z_{S^{(1)'} E}(\lambda) J_{S_s^{(1)'} E} + Z_{S^{(2)'} E}(\lambda) J_{S_s^{(2)'} E} + Z_{\tilde{S}^{(1)'} E}(\lambda) J_{\tilde{S}_s^{(1)'} E} - Z_{\tilde{S}^{(2)'} E}(\lambda) J_{\tilde{S}_s^{(2)'} E}) (-\vec{\partial}_z)^k \\
& - \frac{1}{2} \log \text{Det} \left[I \right. \\
& + \frac{1}{N^2} \left(I + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-\vec{\partial}_z)^{s-k-1} \frac{\Delta^{-1}}{\lambda^{s+2}} (Z_{S^{(1)'} E}(\lambda) \frac{J_{S_s^{(1)'} E}}{s-1} + Z_{S^{(2)'} E}(\lambda) \frac{J_{S_s^{(2)'} E}}{s+2} - Z_{\tilde{S}^{(1)'} E}(\lambda) \frac{J_{\tilde{S}_s^{(1)'} E}}{s-1} - Z_{\tilde{S}^{(2)'} E}(\lambda) \frac{J_{\tilde{S}_s^{(2)'} E}}{s+2}) (-\vec{\partial}_z)^{k+1} \right)^{-1} \\
& \sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1+1}{k_1+2} (-1)^{s_1-1} (-\vec{\partial}_z)^{s_1-k_1} \frac{\Delta^{-1}}{\lambda^{s_1+2}} Z_M(\lambda) J_{\tilde{M}'_{s_1} E} (-\vec{\partial}_z)^{k_1+1} \\
& \left(I + \frac{1}{N} \frac{Z_{\tilde{O}_1^\lambda}(\lambda)}{\lambda^3} \Delta^{-1} J_{\tilde{O}_1^\lambda E} \right. \\
& - \frac{1}{N} \sum_{k_2=0}^{s_2-1} \binom{s_2}{k_2} \binom{s_2}{k_2+1} (-\vec{\partial}_z)^{s_2-k_2} \frac{\Delta^{-1}}{\lambda^{s_2+2}} (-Z_{S^{(1)'} E}(\lambda) J_{S_{s_2}^{(1)'} E} + Z_{S^{(2)'} E}(\lambda) J_{S_{s_2}^{(2)'} E} + Z_{\tilde{S}^{(1)'} E}(\lambda) J_{\tilde{S}_{s_2}^{(1)'} E} - Z_{\tilde{S}^{(2)'} E}(\lambda) J_{\tilde{S}_{s_2}^{(2)'} E}) (-\vec{\partial}_z)^{k_2} \left. \right)^{-1} \\
& \sum_{k_3=0}^{s_3-1} \binom{s_3+1}{k_3} \binom{s_3}{k_3+1} (-1)^{s_3-1} (-\vec{\partial}_z)^{s_3-k_3} \frac{\Delta^{-1}}{\lambda^{s_3+2}} Z_M(\lambda) \bar{J}_{M'_{s_3} E} (-\vec{\partial}_z)^{k_3} \\
& - \frac{1}{2} \log \text{Det} \left[I \right. \\
& + \frac{1}{N^2} \left(I + \frac{1}{N} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-\vec{\partial}_z)^{s-k-1} \frac{\Delta^{-1}}{\lambda^{s+2}} (Z_{S^{(1)'} E}(\lambda) \frac{J_{S_s^{(1)'} E}}{s-1} + Z_{S^{(2)'} E}(\lambda) \frac{J_{S_s^{(2)'} E}}{s+2} + Z_{\tilde{S}^{(1)'} E}(\lambda) \frac{J_{\tilde{S}_s^{(1)'} E}}{s-1} + Z_{\tilde{S}^{(2)'} E}(\lambda) \frac{J_{\tilde{S}_s^{(2)'} E}}{s+2}) (-\vec{\partial}_z)^{k+1} \right)^{-1} \\
& \sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1+1}{k_1+2} (-\vec{\partial}_z)^{s_1-k_1} \frac{\Delta^{-1}}{\lambda^{s_1+2}} Z_M(\lambda) \bar{J}_{M'_{s_1} E} (-\vec{\partial}_z)^{k_1+1} \\
& \left(I - \frac{1}{N} \frac{Z_{\tilde{O}_1^\lambda}(\lambda)}{\lambda^3} \Delta^{-1} J_{\tilde{O}_1^\lambda E} \right. \\
& - \frac{1}{N} \sum_{k_2=0}^{s_2-1} \binom{s_2}{k_2} \binom{s_2}{k_2+1} (-\vec{\partial}_z)^{s_2-k_2} \frac{\Delta^{-1}}{\lambda^{s_2+2}} (-Z_{S^{(1)'} E}(\lambda) J_{S_{s_2}^{(1)'} E} + Z_{S^{(2)'} E}(\lambda) J_{S_{s_2}^{(2)'} E} - Z_{\tilde{S}^{(1)'} E}(\lambda) J_{\tilde{S}_{s_2}^{(1)'} E} + Z_{\tilde{S}^{(2)'} E}(\lambda) J_{\tilde{S}_{s_2}^{(2)'} E}) (-\vec{\partial}_z)^{k_2} \left. \right)^{-1} \\
& \sum_{k_3=0}^{s_3-1} \binom{s_3+1}{k_3} \binom{s_3}{k_3+1} (-\vec{\partial}_z)^{s_3-k_3} \frac{\Delta^{-1}}{\lambda^{s_3+2}} Z_M(\lambda) J_{\tilde{M}'_{s_3} E} (-\vec{\partial}_z)^{k_3} \\
& \left. \right] \tag{7.30}
\end{aligned}$$

8 Minkowskian conformal correlators

We derive from the generating functional in eqs. (5.14) and (5.15) the n -point conformal correlators in several sectors.

8.1 O^A and \tilde{O}^A correlators

We get:

$$\begin{aligned}
& \langle O_{s_1}^A(x_1) \dots O_{s_n}^A(x_n) \tilde{O}_{s_{n+1}}^A(x_{n+1}) \dots \tilde{O}_{s_{n+2m}}^A(x_{n+2m}) \rangle = \\
& \frac{\delta}{\delta J_{O_{s_1}^A}(x_1)} \dots \frac{\delta}{\delta J_{O_{s_n}^A}(x_n)} \frac{\delta}{\delta J_{\tilde{O}_{s_{n+1}}^A}(x_{n+1})} \dots \frac{\delta}{\delta J_{\tilde{O}_{s_{n+2m}}^A}(x_{n+2m})} \mathcal{W}_{\text{conf}}[J_{O^A}, J_{\tilde{O}^A}, 0, 0, 0, 0] \\
& = -\frac{1}{2} \frac{\delta}{\delta J_{O_{s_1}^A}(x_1)} \dots \frac{\delta}{\delta J_{O_{s_n}^A}(x_n)} \frac{\delta}{\delta J_{\tilde{O}_{s_{n+1}}^A}(x_{n+1})} \dots \frac{\delta}{\delta J_{\tilde{O}_{s_{n+2m}}^A}(x_{n+2m})} \\
& \left[\log \text{Det} \left(\mathcal{I} + \frac{1}{2} i \square^{-1} J_{O_s^A} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} + \frac{1}{2} i \square^{-1} J_{\tilde{O}_s^A} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}} \right) \right. \\
& \left. + \log \text{Det} \left(\mathcal{I} + \frac{1}{2} i \square^{-1} J_{O_s^A} \otimes \mathcal{Y}_{s-2}^{\frac{5}{2}} - \frac{1}{2} i \square^{-1} J_{\tilde{O}_s^A} \otimes \mathcal{H}_{s-2}^{\frac{5}{2}} \right) \right] \tag{8.1}
\end{aligned}$$

that reproduces the known result [1, 2]:

$$\begin{aligned}
& \langle O_{s_1}^A(x_1) \dots O_{s_n}^A(x_n) \tilde{O}_{s_{n+1}}^A(x_{n+1}) \dots \tilde{O}_{s_{n+2m}}^A(x_{n+2m}) \rangle \\
& = \frac{1}{(4\pi^2)^{n+2m}} \frac{N^2 - 1}{2^{n+2m}} 2^{\sum_{l=1}^{n+2m} s_l} i^{\sum_{l=1}^{n+2m} s_l} \frac{\Gamma(3)\Gamma(s_1 + 3)}{\Gamma(5)\Gamma(s_1 + 1)} \dots \frac{\Gamma(3)\Gamma(s_{n+2m} + 3)}{\Gamma(5)\Gamma(s_{n+2m} + 1)} \\
& \sum_{k_1=0}^{s_1-2} \dots \sum_{k_{n+2m}=0}^{s_{n+2m}-2} \binom{s_1}{k_1} \binom{s_1}{k_1 + 2} \dots \binom{s_{n+2m}}{k_{n+2m}} \binom{s_{n+2m}}{k_{n+2m} + 2} \\
& \frac{(-1)^{n+2m}}{n + 2m} \sum_{\sigma \in P_{n+2m}} (s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)})! \dots (s_{\sigma(n+2m)} - k_{\sigma(n+2m)} + k_{\sigma(1)})! \\
& \frac{(x_{\sigma(1)} - x_{\sigma(2)})_+^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)}}}{(|x_{\sigma(1)} - x_{\sigma(2)}|^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)} + 1}} \dots \frac{(x_{\sigma(n+2m)} - x_{\sigma(1)})_+^{s_{\sigma(n+2m)} - k_{\sigma(n+2m)} + k_{\sigma(1)}}}{(|x_{\sigma(n+2m)} - x_{\sigma(1)}|^2)^{s_{\sigma(n+2m)} - k_{\sigma(n+2m)} + k_{\sigma(1)} + 1}} \tag{8.2}
\end{aligned}$$

where P_n is the set of permutations of $\{1, \dots, n\}$ and the $-i\epsilon$ prescription in the denominators is understood.

8.2 O^λ and \tilde{O}^λ correlators

Performing the functional derivatives in the definition:

$$\begin{aligned}
& \langle O_{s_1}^\lambda(x_1) \dots O_{s_n}^\lambda(x_n) \rangle = \frac{\delta}{\delta J_{O_{s_1}^\lambda}(x_1)} \dots \frac{\delta}{\delta J_{O_{s_n}^\lambda}(x_n)} \mathcal{W}_{\text{conf}}[0, 0, J_{O^\lambda}, 0, 0, 0] \\
& = \frac{\delta}{\delta J_{O_{s_1}^\lambda}(x_1)} \dots \frac{\delta}{\delta J_{O_{s_n}^\lambda}(x_n)} \log \text{Det} \left(\mathcal{I} + \frac{1}{2} \partial_+ \square^{-1} J_{O_s^\lambda} \otimes \mathcal{Y}_{s-1}^{\frac{3}{2}} \right) \\
& = (N^2 - 1) \frac{\delta}{\delta J_{O_{s_1}^\lambda}(x_1)} \dots \frac{\delta}{\delta J_{O_{s_n}^\lambda}(x_n)} \sum_{l=1}^{\infty} \frac{(-1)^{l+1}}{l} \frac{1}{2^l} \sum_{s'_1} \dots \sum_{s'_l} \\
& \int d^4 y_1 \dots d^4 y_l \partial_{y_1^+} \square^{-1} (y_1 - y_2) \mathcal{Y}_{s'_2-1}^{\frac{3}{2}} \otimes J_{O_{s'_2}^\lambda}(y_2) \dots \partial_{y_l^+} \square^{-1} (y_l - y_1) \mathcal{Y}_{s'_1-1}^{\frac{3}{2}} \otimes J_{O_{s'_1}^\lambda}(y_1) \tag{8.3}
\end{aligned}$$

we obtain:

$$\begin{aligned}
& \langle O_{s_1}^\lambda(x_1) \dots O_{s_n}^\lambda(x_n) \rangle \\
&= (N^2 - 1) \sum_{\sigma \in P_n} \frac{(-1)^{n+1}}{n} \frac{1}{2^n} \partial_{x_{\sigma(1)}^+} \square^{-1}(x_{\sigma(1)} - x_{\sigma(2)}) \mathcal{Y}_{s_{\sigma(2)}-1}^{\frac{3}{2}}(\overleftarrow{\partial}_{x_{\sigma(2)}^+}, \overrightarrow{\partial}_{x_{\sigma(2)}^+}) \\
&\quad \dots \partial_{x_{\sigma(n)}^+} \square^{-1}(x_{\sigma(n)} - x_{\sigma(1)}) \mathcal{Y}_{s_{\sigma(1)}-1}^{\frac{3}{2}}(\overleftarrow{\partial}_{x_{\sigma(1)}^+}, \overrightarrow{\partial}_{x_{\sigma(1)}^+})
\end{aligned} \tag{8.4}$$

Employing the definition in eq. (4.7) we get:

$$\begin{aligned}
& \langle O_{s_1}^\lambda(x_1) \dots O_{s_n}^\lambda(x_n) \rangle = \frac{N^2 - 1}{2^n} \frac{(-1)^{n+1}}{n} i^{\sum_{l=1}^n s_l - n} \frac{(s_1 + 1)}{2} \dots \frac{(s_n + 1)}{2} \\
& \sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1}{k_1 + 1} (-1)^{s_1 - k_1 - 1} \dots \sum_{k_n=0}^{s_n-1} \binom{s_n}{k_n} \binom{s_n}{k_n + 1} (-1)^{s_n - k_n - 1} \\
& \sum_{\sigma \in P_n} \partial_{x_{\sigma(1)}^+} \square^{-1}(x_{\sigma(1)} - x_{\sigma(2)}) \overleftarrow{\partial}_{x_{\sigma(2)}^+}^{s_{\sigma(2)} - k_{\sigma(2)} - 1} \overrightarrow{\partial}_{x_{\sigma(2)}^+}^{k_{\sigma(2)}} \\
& \dots \partial_{x_{\sigma(n)}^+} \square^{-1}(x_{\sigma(n)} - x_{\sigma(1)}) \overleftarrow{\partial}_{x_{\sigma(1)}^+}^{s_{\sigma(1)} - k_{\sigma(1)} - 1} \overrightarrow{\partial}_{x_{\sigma(1)}^+}^{k_{\sigma(1)}}
\end{aligned} \tag{8.5}$$

It follows from eq. (5.21):

$$\begin{aligned}
& \langle O_{s_1}^\lambda(x_1) \dots O_{s_n}^\lambda(x_n) \rangle = \frac{N^2 - 1}{2^n} \frac{(-i)^n}{(4\pi^2)^n} \frac{(-1)^{n+1}}{n} i^{\sum_{l=1}^n s_l - n} \frac{(s_1 + 1)}{2} \dots \frac{(s_n + 1)}{2} \\
& \sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1}{k_1 + 1} (-1)^{s_1 - k_1 - 1} \dots \sum_{k_n=0}^{s_n-1} \binom{s_n}{k_n} \binom{s_n}{k_n + 1} (-1)^{s_n - k_n - 1} \\
& \sum_{\sigma \in P_n} \partial_{x_{\sigma(1)}^+} \frac{1}{|x_{\sigma(1)} - x_{\sigma(2)}|^2} \overleftarrow{\partial}_{x_{\sigma(2)}^+}^{s_{\sigma(2)} - k_{\sigma(2)} - 1} \overrightarrow{\partial}_{x_{\sigma(2)}^+}^{k_{\sigma(2)}} \\
& \dots \partial_{x_{\sigma(n)}^+} \frac{1}{|x_{\sigma(n)} - x_{\sigma(1)}|^2} \overleftarrow{\partial}_{x_{\sigma(1)}^+}^{s_{\sigma(1)} - k_{\sigma(1)} - 1} \overrightarrow{\partial}_{x_{\sigma(1)}^+}^{k_{\sigma(1)}}
\end{aligned} \tag{8.6}$$

We now employ [2]:

$$\begin{aligned}
& \partial_{x^+}^a \partial_{y^+}^b \frac{1}{|x - y|^2} = \partial_{x^+}^a \partial_{y^+}^b \frac{1}{2(x - y)_+ (x - y)_- - (x - y)_\perp^2} \\
&= (-1)^a (a + b)! 2^{a+b} \frac{(x - y)_+^{a+b}}{(|x - y|^2)^{a+b+1}}
\end{aligned} \tag{8.7}$$

so that:

$$\begin{aligned}
\langle O_{s_1}^\lambda(x_1) \dots O_{s_n}^\lambda(x_n) \rangle &= \frac{N^2 - 1}{2^n} \frac{(-i)^n}{(4\pi^2)^n} \frac{(-1)^{n+1}}{n} i^{\sum_{l=1}^n s_l - n} \frac{(s_1 + 1)}{2} \dots \frac{(s_n + 1)}{2} \\
&\sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1}{k_1 + 1} (-1)^{s_1 - k_1 - 1} \dots \sum_{k_n=0}^{s_n-1} \binom{s_n}{k_n} \binom{s_n}{k_n + 1} (-1)^{s_n - k_n - 1} \\
&\sum_{\sigma \in P_n} (-1)^{k_{\sigma(1)} + 1} (s_{\sigma(2)} - k_{\sigma(2)} + k_{\sigma(1)})! 2^{s_{\sigma(2)} - k_{\sigma(2)} + k_{\sigma(1)}} \frac{(x_{\sigma(1)} - x_{\sigma(2)})_+^{s_{\sigma(2)} - k_{\sigma(2)} + k_{\sigma(1)}}}{(|x_{\sigma(1)} - x_{\sigma(2)}|^2)^{s_{\sigma(2)} - k_{\sigma(2)} + k_{\sigma(1)} + 1}} \\
&\dots (-1)^{k_{\sigma(n)} + 1} (s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(n)})! 2^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(n)}} \frac{(x_{\sigma(n)} - x_{\sigma(1)})_+^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(n)}}}{(|x_{\sigma(n)} - x_{\sigma(1)}|^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(n)} + 1}}
\end{aligned} \tag{8.8}$$

Relabeling the permutations, $\sigma(n) \rightarrow \sigma(2)$, $\sigma(n-1) \rightarrow \sigma(3)$ and so on, while keeping $\sigma(1)$ fixed, we obtain:

$$\begin{aligned}
\langle O_{s_1}^\lambda(x_1) \dots O_{s_n}^\lambda(x_n) \rangle &= \frac{N^2 - 1}{2^n} \frac{(-i)^n}{(4\pi^2)^n} \frac{(-1)^{n+1}}{n} i^{\sum_{l=1}^n s_l - n} \frac{(s_1 + 1)}{2} \dots \frac{(s_n + 1)}{2} \\
&\sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1}{k_1 + 1} (-1)^{s_1 - k_1 - 1} \dots \sum_{k_n=0}^{s_n-1} \binom{s_n}{k_n} \binom{s_n}{k_n + 1} (-1)^{s_n - k_n - 1} \\
&\sum_{\sigma \in P_n} (-1)^{k_{\sigma(1)} + 1} (s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)})! 2^{s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)}} \frac{(x_{\sigma(1)} - x_{\sigma(n)})_+^{s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)}}}{(|x_{\sigma(1)} - x_{\sigma(n)}|^2)^{s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)} + 1}} \\
&\dots (-1)^{k_{\sigma(2)} + 1} (s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)})! 2^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)}} \frac{(x_{\sigma(2)} - x_{\sigma(1)})_+^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)}}}{(|x_{\sigma(2)} - x_{\sigma(1)}|^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)} + 1}}
\end{aligned} \tag{8.9}$$

that simplifies to:

$$\begin{aligned}
\langle O_{s_1}^\lambda(x_1) \dots O_{s_n}^\lambda(x_n) \rangle &= -\frac{N^2 - 1}{2^n} \frac{1}{(4\pi^2)^n} i^{\sum_{l=1}^n s_l} 2^{\sum_{l=1}^n s_l} \frac{(s_1 + 1)}{2} \dots \frac{(s_n + 1)}{2} \\
&\sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1}{k_1 + 1} \dots \sum_{k_n=0}^{s_n-1} \binom{s_n}{k_n} \binom{s_n}{k_n + 1} \\
&\frac{1}{n} \sum_{\sigma \in P_n} (s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)})! \dots (s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)})! \\
&\frac{(x_{\sigma(1)} - x_{\sigma(2)})_+^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)}}}{(|x_{\sigma(1)} - x_{\sigma(2)}|^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)} + 1}} \dots \frac{(x_{\sigma(n)} - x_{\sigma(1)})_+^{s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)}}}{(|x_{\sigma(n)} - x_{\sigma(1)}|^2)^{s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)} + 1}}
\end{aligned} \tag{8.10}$$

The 2-point correlators follow:

$$\begin{aligned} \langle O_{s_1}^\lambda(x_1) O_{s_2}^\lambda(x_2) \rangle &= -\frac{N^2 - 1}{4} \frac{1}{(4\pi^2)^2} i^{s_1+s_2} 2^{s_1+s_2} \frac{(s_1+1)(s_2+1)}{4} \\ &\sum_{k_1=0}^{s_1-1} \sum_{k_2=0}^{s_2-1} \binom{s_1}{k_1} \binom{s_1}{k_1+1} \binom{s_2}{k_2} \binom{s_2}{k_2+1} (-1)^{s_2-k_2+k_1} (s_1-k_1+k_2)! (s_2-k_2+k_1)! \\ &\frac{(x_1-x_2)_+^{s_1+s_2}}{(|x_1-x_2|^2)^{s_1+s_2+2}} \end{aligned} \quad (8.11)$$

that can be rewritten as:

$$\begin{aligned} \langle O_{s_1}^\lambda(x_1) O_{s_2}^\lambda(x_2) \rangle &= -\frac{N^2 - 1}{4} \frac{1}{(4\pi^2)^2} i^{s_1+s_2} 2^{s_1+s_2} \frac{(s_1+1)(s_2+1)}{4} \\ &\sum_{k_1=0}^{s_1-1} \sum_{k_2=0}^{s_2-1} \binom{s_1}{k_1} \binom{s_1}{k_1+1} \binom{s_2}{k_2} \binom{s_2}{k_2+1} (-1)^{k_1+k_2+1} (s_1+s_2-k_1-k_2-1)! (k_2+k_1+1)! \\ &\frac{(x_1-x_2)_+^{s_1+s_2}}{(|x_1-x_2|^2)^{s_1+s_2+2}} \end{aligned} \quad (8.12)$$

by substituting:

$$k'_2 = s_2 - k_2 - 1 \quad (8.13)$$

into eq. (8.11) and dropping the primed index. By employing (appendix C.1):

$$\delta_{s_1 s_2} \frac{s_1}{s_1+1} = \sum_{k_1=0}^{s_1-1} \sum_{k_2=0}^{s_2-1} \binom{s_1}{k_1} \binom{s_1}{k_1+1} \binom{s_2}{k_2} \binom{s_2}{k_2+1} (-1)^{k_1+k_2} \frac{1}{\binom{s_1+s_2}{k_1+k_2+1}} \quad (8.14)$$

the 2-point correlators become:

$$\langle O_{s_1}^\lambda(x_1) O_{s_2}^\lambda(x_2) \rangle = \delta_{s_1 s_2} \frac{N^2 - 1}{4} \frac{1}{(4\pi^2)^2} (-1)^{s_1} 2^{2s_1} (2s_1)! \frac{s_1(s_1+1)}{4} \frac{(x_1-x_2)_+^{2s_1}}{(|x_1-x_2|^2)^{2s_1+2}} \quad (8.15)$$

Similarly, we obtain the correlators of O^λ and \tilde{O}^λ from their definition:

$$\begin{aligned} \langle O_{s_1}^\lambda(x_1) \dots O_{s_n}^\lambda(x_n) \tilde{O}_{s_{n+1}}^\lambda(x_{n+1}) \dots \tilde{O}_{s_{n+2m}}^\lambda(x_{n+2m}) \rangle &= \\ &\frac{\delta}{\delta J_{O_{s_1}^\lambda}(x_1)} \dots \frac{\delta}{\delta J_{O_{s_n}^\lambda}(x_n)} \frac{\delta}{\delta J_{\tilde{O}_{s_{n+1}}^\lambda}(x_{n+1})} \dots \frac{\delta}{\delta J_{\tilde{O}_{s_{n+2m}}^\lambda}(x_{n+2m})} \mathcal{W}_{\text{conf}}[0, 0, J_{O^\lambda}, J_{\tilde{O}^\lambda}, 0, 0] \\ &= \frac{1}{2} \frac{\delta}{\delta J_{O_{s_1}^\lambda}(x_1)} \dots \frac{\delta}{\delta J_{O_{s_n}^\lambda}(x_n)} \frac{\delta}{\delta J_{\tilde{O}_{s_{n+1}}^\lambda}(x_{n+1})} \dots \frac{\delta}{\delta J_{\tilde{O}_{s_{n+2m}}^\lambda}(x_{n+2m})} \\ &\left[\log \text{Det} \left(\mathcal{I} + \frac{1}{2} \partial_+ \square^{-1} J_{O_s^\lambda} \otimes \mathcal{Y}_{s-1}^{\frac{3}{2}} + \frac{1}{2} \partial_+ \square^{-1} J_{\tilde{O}_s^\lambda} \otimes \mathcal{H}_{s-1}^{\frac{3}{2}} \right) \right. \\ &\left. + \log \text{Det} \left(\mathcal{I} + \frac{1}{2} \partial_+ \square^{-1} J_{O_s^\lambda} \otimes \mathcal{Y}_{s-1}^{\frac{3}{2}} - \frac{1}{2} \partial_+ \square^{-1} J_{\tilde{O}_s^\lambda} \otimes \mathcal{H}_{s-1}^{\frac{3}{2}} \right) \right] \end{aligned} \quad (8.16)$$

that yields:

$$\begin{aligned}
& \langle O_{s_1}^\lambda(x_1) \dots O_{s_n}^\lambda(x_n) \tilde{O}_{s_{n+1}}^\lambda(x_{n+1}) \dots \tilde{O}_{s_{n+2m}}^\lambda(x_{n+2m}) \rangle = \\
& - \frac{N^2 - 1}{2^{n+2m}} \frac{1}{(4\pi^2)^{n+2m}} i^{\sum_{l=1}^{n+2m} s_l} 2^{\sum_{l=1}^{n+2m} s_l} \frac{(s_1 + 1)}{2} \dots \frac{(s_{n+2m} + 1)}{2} \\
& \sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1}{k_1 + 1} \dots \sum_{k_{n+2m}=0}^{s_{n+2m}-1} \binom{s_{n+2m}}{k_{n+2m}} \binom{s_{n+2m}}{k_{n+2m} + 1} \\
& \frac{1}{n+2m} \sum_{\sigma \in P_{n+2m}} (s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)})! \dots (s_{\sigma(n+2m)} - k_{\sigma(n+2m)} + k_{\sigma(1)})! \\
& \frac{(x_{\sigma(1)} - x_{\sigma(2)})_+^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)}}}{(|x_{\sigma(1)} - x_{\sigma(2)}|^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)} + 1}} \dots \frac{(x_{\sigma(n+2m)} - x_{\sigma(1)})_+^{s_{\sigma(n+2m)} - k_{\sigma(n+2m)} + k_{\sigma(1)}}}{(|x_{\sigma(n+2m)} - x_{\sigma(1)}|^2)^{s_{\sigma(n+2m)} - k_{\sigma(n+2m)} + k_{\sigma(1)} + 1}}
\end{aligned} \tag{8.17}$$

8.3 M and \bar{M} correlators

The correlators of gluon-gluino operators involve the same number of M_s and \bar{M}_s , otherwise they vanish, as follows from the generating functional. Therefore, the nonvanishing correlators read:

$$\begin{aligned}
& \langle M_{s_1}(x_1) \bar{M}_{s'_1}(y_1) M_{s_2}(x_2) \bar{M}_{s'_2}(y_2) \dots M_{s_n}(x_n) \bar{M}_{s'_n}(y_n) \rangle \\
& = \frac{\delta}{\delta \bar{J}_{M_{s_1}}(x_1)} \left(-\frac{\delta}{\delta J_{\bar{M}_{s'_1}}(y_1)} \right) \dots \frac{\delta}{\delta \bar{J}_{M_{s_n}}(x_n)} \left(-\frac{\delta}{\delta J_{\bar{M}_{s'_n}}(y_n)} \right) \mathcal{W}_{\text{conf}}[0, 0, J_{\bar{M}}, \bar{J}_M] \\
& = \frac{\delta}{\delta \bar{J}_{M_{s_1}}(x_1)} \left(-\frac{\delta}{\delta J_{\bar{M}_{s'_1}}(y_1)} \right) \dots \frac{\delta}{\delta \bar{J}_{M_{s_n}}(x_n)} \left(-\frac{\delta}{\delta J_{\bar{M}_{s'_n}}(y_n)} \right) \\
& \quad \frac{1}{2} \left[\log \text{Det} \left(\mathcal{I} + \frac{1}{4} i \partial_+ i \square^{-1} \bar{J}_{M_{s_1}} \otimes \mathcal{G}_{s_1-1}^{(1,2)} (-1)^{s_1-1} i \square^{-1} J_{\bar{M}_{s_3}} \otimes \mathcal{G}_{s_2-1}^{(2,1)} (-1)^{s_2-1} \right) \right. \\
& \quad \left. + \log \text{Det} \left(\mathcal{I} + \frac{1}{4} i \partial_+ i \square^{-1} J_{\bar{M}_{s_1}} \otimes \mathcal{G}_{s_1-1}^{(1,2)} i \square^{-1} \bar{J}_{M_{s_3}} \otimes \mathcal{G}_{s_3-1}^{(2,1)} \right) \right]
\end{aligned} \tag{8.18}$$

The two determinants above are equal (appendix D):

$$\begin{aligned}
& \log \text{Det} \left(\mathcal{I} + \frac{1}{4} i \partial_+ i \square^{-1} \bar{J}_{M_{s_1}} \otimes \mathcal{G}_{s_1-1}^{(1,2)} (-1)^{s_1-1} i \square^{-1} J_{\bar{M}_{s_3}} \otimes \mathcal{G}_{s_3-1}^{(2,1)} (-1)^{s_3-1} \right) \\
& = \log \text{Det} \left(\mathcal{I} + \frac{1}{4} i \partial_+ i \square^{-1} J_{\bar{M}_{s_1}} \otimes \mathcal{G}_{s_1-1}^{(1,2)} i \square^{-1} \bar{J}_{M_{s_3}} \otimes \mathcal{G}_{s_3-1}^{(2,1)} \right)
\end{aligned} \tag{8.19}$$

so that:

$$\begin{aligned}
& \langle M_{s_1}(x_1) \bar{M}_{s'_1}(y_1) M_{s_2}(x_2) \bar{M}_{s'_2}(y_2) \dots M_{s_n}(x_n) \bar{M}_{s'_n}(y_n) \rangle \\
&= \frac{\delta}{\delta \bar{J}_{M_{s_1}}(x_1)} \left(-\frac{\delta}{\delta J_{\bar{M}_{s'_1}}(y_1)} \right) \dots \frac{\delta}{\delta \bar{J}_{M_{s_n}}(x_n)} \left(-\frac{\delta}{\delta J_{\bar{M}_{s'_n}}(y_n)} \right) \mathcal{W}_{\text{conf}} [0, 0, J_{\bar{M}}, \bar{J}_M] \\
&= \frac{\delta}{\delta \bar{J}_{M_{s_1}}(x_1)} \left(-\frac{\delta}{\delta J_{\bar{M}_{s'_1}}(y_1)} \right) \dots \frac{\delta}{\delta \bar{J}_{M_{s_n}}(x_n)} \left(-\frac{\delta}{\delta J_{\bar{M}_{s'_n}}(y_n)} \right) \\
&\quad \log \text{Det} \left(\mathcal{I} + \frac{1}{4} i \partial_+ i \square^{-1} J_{\bar{M}_{s_1}} \otimes \mathcal{G}_{s_1-1}^{(1,2)} i \square^{-1} \bar{J}_{M_{s_2}} \otimes \mathcal{G}_{s_2-1}^{(2,1)} \right) \\
&= \frac{\delta}{\delta \bar{J}_{M_{s_1}}(x_1)} \left(-\frac{\delta}{\delta J_{\bar{M}_{s'_1}}(y_1)} \right) \dots \frac{\delta}{\delta \bar{J}_{M_{s_n}}(x_n)} \left(-\frac{\delta}{\delta J_{\bar{M}_{s'_n}}(y_n)} \right) \\
&\quad (N^2 - 1) \sum_{l=1}^{\infty} \frac{(-1)^{l+1}}{l} \frac{1}{2^{2l}} \int d^4 u_1 \dots d^4 u_l d^4 v_1 \dots d^4 v_l \sum_{s_1} \dots \sum_{s_l} \sum_{s'_1} \dots \sum_{s'_l} \\
&\quad i \partial_{u_1^+} i \square^{-1} (u_1 - v_1) \mathcal{G}_{s'_1-1}^{(1,2)} (\overleftarrow{\partial}_{v_1^+}, \overrightarrow{\partial}_{v_1^+}) \otimes J_{\bar{M}_{s'_1}}(v_1) \\
&\quad i \square^{-1} (v_1 - u_2) \mathcal{G}_{s_2-1}^{(2,1)} (\overleftarrow{\partial}_{u_2^+}, \overrightarrow{\partial}_{u_2^+}) \otimes \bar{J}_{M_{s_2}}(u_2) \\
&\quad \dots i \partial_{u_l^+} i \square^{-1} (u_l - v_l) \mathcal{G}_{s'_l-1}^{(1,2)} (\overleftarrow{\partial}_{v_l^+}, \overrightarrow{\partial}_{v_l^+}) \otimes J_{\bar{M}_{s'_l}}(v_l) \\
&\quad i \square^{-1} (v_l - u_1) \mathcal{G}_{s_1-1}^{(2,1)} (\overleftarrow{\partial}_{u_1^+}, \overrightarrow{\partial}_{u_1^+}) \otimes \bar{J}_{M_{s_1}}(u_1)
\end{aligned} \tag{8.20}$$

Performing the functional derivatives, we obtain:

$$\begin{aligned}
& \langle M_{s_1}(x_1) \bar{M}_{s'_1}(y_1) M_{s_2}(x_2) \bar{M}_{s'_2}(y_2) \dots M_{s_n}(x_n) \bar{M}_{s'_n}(y_n) \rangle \\
&= (N^2 - 1) \sum_{\sigma \in P_n} \sum_{\rho \in P_n} \frac{(-1)^{2n+1}}{n} \frac{1}{2^{2n}} \text{sgn}(\sigma) \text{sgn}(\rho) \\
&\quad i \partial_{x_{\sigma(1)}^+} i \square^{-1} (x_{\sigma(1)} - y_{\rho(1)}) \mathcal{G}_{s'_{\rho(1)}-1}^{(1,2)} (\overleftarrow{\partial}_{y_{\rho(1)}^+}, \overrightarrow{\partial}_{y_{\rho(1)}^+}) \\
&\quad i \square^{-1} (y_{\rho(1)} - x_{\sigma(2)}) \mathcal{G}_{s_{\sigma(2)}-1}^{(2,1)} (\overleftarrow{\partial}_{x_{\sigma(2)}^+}, \overrightarrow{\partial}_{x_{\sigma(2)}^+}) \\
&\quad \dots i \partial_{x_{\sigma(n)}^+} i \square^{-1} (x_{\sigma(n)} - y_{\rho(n)}) \mathcal{G}_{s'_{\rho(n)}-1}^{(1,2)} (\overleftarrow{\partial}_{y_{\rho(n)}^+}, \overrightarrow{\partial}_{y_{\rho(n)}^+}) \\
&\quad i \square^{-1} (y_{\rho(n)} - x_{\sigma(1)}) \mathcal{G}_{s_{\sigma(1)}-1}^{(2,1)} (\overleftarrow{\partial}_{x_{\sigma(1)}^+}, \overrightarrow{\partial}_{x_{\sigma(1)}^+})
\end{aligned} \tag{8.21}$$

where $\text{sgn}(\sigma)$ denotes the sign of the permutation σ . Substituting eq.(5.21) in the above equation, we get:

$$\begin{aligned}
& \langle M_{s_1}(x_1) \bar{M}_{s'_1}(y_1) M_{s_2}(x_2) \bar{M}_{s'_2}(y_2) \dots M_{s_n}(x_n) \bar{M}_{s'_n}(y_n) \rangle \\
&= (N^2 - 1) \frac{i^n}{(4\pi^2)^{2n}} \sum_{\sigma \in P_n} \sum_{\rho \in P_n} \frac{(-1)^{2n+1}}{n} \frac{1}{2^{2n}} \text{sgn}(\sigma) \text{sgn}(\rho) \\
&\quad \partial_{x_{\sigma(1)}^+} \frac{1}{|x_{\sigma(1)} - y_{\rho(1)}|^2} \mathcal{G}_{s'_{\rho(1)}-1}^{(1,2)} (\overleftarrow{\partial}_{y_{\rho(1)}^+}, \overrightarrow{\partial}_{y_{\rho(1)}^+}) \frac{1}{|y_{\rho(1)} - x_{\sigma(2)}|^2} \mathcal{G}_{s_{\sigma(1)}-1}^{(2,1)} (\overleftarrow{\partial}_{x_{\sigma(2)}^+}, \overrightarrow{\partial}_{x_{\sigma(2)}^+}) \\
&\quad \dots \partial_{x_{\sigma(n)}^+} \frac{1}{|x_{\sigma(n)} - y_{\rho(n)}|^2} \mathcal{G}_{s'_{\rho(n)}-1}^{(1,2)} (\overleftarrow{\partial}_{y_{\rho(n)}^+}, \overrightarrow{\partial}_{y_{\rho(n)}^+}) \frac{1}{|y_{\rho(n)} - x_{\sigma(1)}|^2} \mathcal{G}_{s_{\sigma(1)}-1}^{(2,1)} (\overleftarrow{\partial}_{x_{\sigma(1)}^+}, \overrightarrow{\partial}_{x_{\sigma(1)}^+}) \tag{8.22}
\end{aligned}$$

Employing eq. (4.9):

$$\begin{aligned}
& \langle M_{s_1}(x_1) \bar{M}_{s'_1}(y_1) M_{s_2}(x_2) \bar{M}_{s'_2}(y_2) \dots M_{s_n}(x_n) \bar{M}_{s'_n}(y_n) \rangle = (N^2 - 1) \frac{i^n}{(4\pi^2)^{2n}} \\
& i^{s'_1-1} \sum_{k'_1=0}^{s'_1-1} \binom{s'_1}{k'_1} \binom{s'_1+1}{k'_1+2} (-1)^{s'_1-k'_1-1} \dots i^{s'_n-1} \sum_{k'_n=0}^{s'_n-1} \binom{s'_n}{k'_n} \binom{s'_n+1}{k'_n+2} (-1)^{s'_n-k'_n-1} \\
& i^{s_1-1} \sum_{k_1=0}^{s_1-1} \binom{s_1+1}{k_1} \binom{s_1}{k_1+1} (-1)^{s_1-k_1-1} \dots i^{s_n-1} \sum_{k_n=0}^{s_n-1} \binom{s_n+1}{k_n} \binom{s_n}{k_n+1} (-1)^{s_n-k_n-1} \\
& \sum_{\sigma \in P_n} \sum_{\rho \in P_n} \frac{(-1)^{2n+1}}{n} \frac{1}{2^{2n}} \text{sgn}(\sigma) \text{sgn}(\rho) \\
& \partial_{x_{\sigma(1)}^+} \frac{1}{|x_{\sigma(1)} - y_{\rho(1)}|^2} \overleftarrow{\partial}_{y_{\rho(1)}^+}^{s'_{\rho(1)}-k'_{\rho(1)}-1} \overrightarrow{\partial}_{y_{\rho(1)}^+}^{k'_{\rho(1)}+1} \frac{1}{|y_{\rho(1)} - x_{\sigma(2)}|^2} \overleftarrow{\partial}_{x_{\sigma(2)}^+}^{s_{\sigma(2)}-k_{\sigma(2)}} \overrightarrow{\partial}_{x_{\sigma(2)}^+}^{k_{\sigma(2)}} \\
& \dots \partial_{x_{\sigma(n)}^+} \frac{1}{|x_{\sigma(n)} - y_{\rho(n)}|^2} \overleftarrow{\partial}_{y_{\rho(n)}^+}^{s'_{\rho(n)}-k'_{\rho(n)}-1} \overrightarrow{\partial}_{y_{\rho(n)}^+}^{k'_{\rho(n)}+1} \frac{1}{|y_{\rho(n)} - x_{\sigma(1)}|^2} \overleftarrow{\partial}_{x_{\sigma(1)}^+}^{s_{\sigma(1)}-k_{\sigma(1)}} \overrightarrow{\partial}_{x_{\sigma(1)}^+}^{k_{\sigma(1)}} \tag{8.23}
\end{aligned}$$

and eq. (8.7), we obtain:

$$\begin{aligned}
\langle M_{s_1}(x_1) \bar{M}_{s'_1}(y_1) M_{s_2}(x_2) \bar{M}_{s'_2}(y_2) \dots M_{s_n}(x_n) \bar{M}_{s'_n}(y_n) \rangle &= (N^2 - 1) \frac{i^n}{(4\pi^2)^{2n}} \\
&\quad i^{s'_1-1} \sum_{k'_1=0}^{s'_1-1} \binom{s'_1}{k'_1} \binom{s'_1+1}{k'_1+2} (-1)^{s'_1-k'_1-1} \dots i^{s'_n-1} \sum_{k'_n=0}^{s'_n-1} \binom{s'_n}{k'_n} \binom{s'_n+1}{k'_n+2} (-1)^{s'_n-k'_n-1} \\
&\quad i^{s_1-1} \sum_{k_1=0}^{s_1-1} \binom{s_1+1}{k_1} \binom{s_1}{k_1+1} (-1)^{s_1-k_1-1} \dots i^{s_n-1} \sum_{k_n=0}^{s_n-1} \binom{s_n+1}{k_n} \binom{s_n}{k_n+1} (-1)^{s_n-k_n-1} \\
&\quad \sum_{\sigma \in P_n} \sum_{\rho \in P_n} \frac{(-1)^{2n+1}}{n} \frac{1}{2^{2n}} \text{sgn}(\sigma) \text{sgn}(\rho) \\
&\quad (-1)^{k_{\sigma(1)}+1} 2^{s'_{\rho(1)}-k'_{\rho(1)}+k_{\sigma(1)}} (s'_{\rho(1)} - k'_{\rho(1)} + k_{\sigma(1)})! \frac{(x_{\sigma(1)} - y_{\rho(1)})_+^{s'_{\rho(1)}-k'_{\rho(1)}+k_{\sigma(1)}}}{(|x_{\sigma(1)} - y_{\rho(1)}|^2)^{s'_{\rho(1)}-k'_{\rho(1)}+k_{\sigma(1)}+1}} \\
&\quad (-1)^{k'_{\rho(1)}+1} 2^{s_{\sigma(2)}-k_{\sigma(2)}+k'_{\rho(1)}+1} (s_{\sigma(2)} - k_{\sigma(2)} + k'_{\rho(1)} + 1)! \frac{(y_{\rho(1)} - x_{\sigma(2)})_+^{s_{\sigma(2)}-k_{\sigma(2)}+k'_{\rho(1)}+1}}{(|y_{\rho(1)} - x_{\sigma(2)}|^2)^{s_{\sigma(2)}-k_{\sigma(2)}+k'_{\rho(1)}+2}} \\
&\quad \dots \\
&\quad (-1)^{k_{\sigma(n)+1}} 2^{s'_{\rho(n)}-k'_{\rho(n)}+k_{\sigma(n)}} (s'_{\rho(n)} - k'_{\rho(n)} + k_{\sigma(n)})! \frac{(x_{\sigma(n)} - y_{\rho(n)})_+^{s'_{\rho(n)}-k'_{\rho(n)}+k_{\sigma(n)}}}{(|x_{\sigma(n)} - y_{\rho(n)}|^2)^{s'_{\rho(n)}-k'_{\rho(n)}+k_{\sigma(n)}+1}} \\
&\quad (-1)^{k'_{\rho(n)}+1} 2^{s_{\sigma(1)}-k_{\sigma(1)}+k'_{\rho(n)}+1} (s_{\sigma(1)} - k_{\sigma(1)} + k'_{\rho(n)} + 1)! \frac{(y_{\rho(n)} - x_{\sigma(1)})_+^{s_{\sigma(1)}-k_{\sigma(1)}+k'_{\rho(n)}+1}}{(|y_{\rho(n)} - x_{\sigma(1)}|^2)^{s_{\sigma(1)}-k_{\sigma(1)}+k'_{\rho(n)}+2}}
\end{aligned} \tag{8.24}$$

that simplifies to:

$$\begin{aligned}
& \langle M_{s_1}(x_1) \bar{M}_{s'_1}(y_1) M_{s_2}(x_2) \bar{M}_{s'_2}(y_2) \dots M_{s_n}(x_n) \bar{M}_{s'_n}(y_n) \rangle \\
&= (N^2 - 1) \frac{1}{(4\pi^2)^{2n}} 2^{\sum_{l=1}^n s_l + \sum_{l=1}^n s'_l - n} i^{\sum_{l=1}^n s_l + \sum_{l=1}^n s'_l - n} (-1)^{\sum_{l=1}^n s_l + \sum_{l=1}^n s'_l} \\
&\quad \sum_{k'_1=0}^{s'_1-1} \binom{s'_1}{k'_1} \binom{s'_1+1}{k'_1+2} \dots \sum_{k'_n=0}^{s'_n-1} \binom{s'_n}{k'_n} \binom{s'_n+1}{k'_n+2} \\
&\quad \sum_{k_1=0}^{s_1-1} \binom{s_1+1}{k_1} \binom{s_1}{k_1+1} \dots \sum_{k_n=0}^{s_n-1} \binom{s_n+1}{k_n} \binom{s_n}{k_n+1} \\
&\quad \sum_{\sigma \in P_n} \sum_{\rho \in P_n} \frac{(-1)^{2n+1}}{n} \text{sgn}(\sigma) \text{sgn}(\rho) \\
&\quad (s'_{\rho(1)} - k'_{\rho(1)} + k_{\sigma(1)})! \frac{(x_{\sigma(1)} - y_{\rho(1)})_+^{s'_{\rho(1)} - k'_{\rho(1)} + k_{\sigma(1)}}}{(|x_{\sigma(1)} - y_{\rho(1)}|^2)^{s'_{\rho(1)} - k'_{\rho(1)} + k_{\sigma(1)} + 1}} \\
&\quad (s_{\sigma(2)} - k_{\sigma(2)} + k'_{\rho(1)} + 1)! \frac{(y_{\rho(1)} - x_{\sigma(2)})_+^{s_{\sigma(2)} - k_{\sigma(2)} + k'_{\rho(1)} + 1}}{(|y_{\rho(1)} - x_{\sigma(2)}|^2)^{s_{\sigma(2)} - k_{\sigma(2)} + k'_{\rho(1)} + 2}} \\
&\quad \dots \\
&\quad (s'_{\rho(n)} - k'_{\rho(n)} + k_{\sigma(n)})! \frac{(x_{\sigma(n)} - y_{\rho(n)})_+^{s'_{\rho(n)} - k'_{\rho(n)} + k_{\sigma(n)}}}{(|x_{\sigma(n)} - y_{\rho(n)}|^2)^{s'_{\rho(n)} - k'_{\rho(n)} + k_{\sigma(n)} + 1}} \\
&\quad (s_{\sigma(1)} - k_{\sigma(1)} + k'_{\rho(n)} + 1)! \frac{(y_{\rho(n)} - x_{\sigma(1)})_+^{s_{\sigma(1)} - k_{\sigma(1)} + k'_{\rho(n)} + 1}}{(|y_{\rho(n)} - x_{\sigma(1)}|^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k'_{\rho(n)} + 2}}
\end{aligned} \tag{8.25}$$

By substituting:

$$k_{\sigma(i)} \rightarrow s_{\sigma(i)} - 1 - k_{\sigma(i)} \quad k'_{\rho(i)} \rightarrow s'_{\rho(i)} - 1 - k'_{\rho(i)} \tag{8.26}$$

we get:

$$\begin{aligned}
& \langle M_{s_1}(x_1) \bar{M}_{s'_1}(y_1) M_{s_2}(x_2) \bar{M}_{s'_2}(y_2) \dots M_{s_n}(x_n) \bar{M}_{s'_n}(y_n) \rangle \\
&= (N^2 - 1) \frac{1}{(4\pi^2)^{2n}} 2^{\sum_{l=1}^n s_l + \sum_{l=1}^n s'_l - n} i^{\sum_{l=1}^n s_l + \sum_{l=1}^n s'_l - n} (-1)^{\sum_{l=1}^n s_l + \sum_{l=1}^n s'_l} \\
&\quad \sum_{k'_1=0}^{s'_1-1} \binom{s'_1+1}{k'_1} \binom{s'_1}{k'_1+1} \dots \sum_{k'_n=0}^{s'_n-1} \binom{s'_n+1}{k'_n} \binom{s'_n}{k'_n+1} \\
&\quad \sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1+1}{k_1+2} \dots \sum_{k_n=0}^{s_n-1} \binom{s_n}{k_n} \binom{s_n+1}{k_n+2} \\
&\quad \sum_{\sigma \in P_n} \sum_{\rho \in P_n} \frac{(-1)^{2n+1}}{n} \text{sgn}(\sigma) \text{sgn}(\rho) \\
&\quad (s_{\sigma(1)} - k_{\sigma(1)} + k'_{\rho(1)})! \frac{(x_{\sigma(1)} - y_{\rho(1)})_+^{s_{\sigma(1)} - k_{\sigma(1)} + k'_{\rho(1)}}}{(|x_{\sigma(1)} - y_{\rho(1)}|^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k'_{\rho(1)} + 1}} \\
&\quad (s'_{\rho(1)} - k'_{\rho(1)} + k_{\sigma(2)} + 1)! \frac{(y_{\rho(1)} - x_{\sigma(2)})_+^{s'_{\rho(1)} - k'_{\rho(1)} + k_{\sigma(2)} + 1}}{(|y_{\rho(1)} - x_{\sigma(2)}|^2)^{s'_{\rho(1)} - k'_{\rho(1)} + k_{\sigma(2)} + 2}} \\
&\quad \dots \\
&\quad (s_{\sigma(n)} - k_{\sigma(n)} + k'_{\rho(n)})! \frac{(x_{\sigma(n)} - y_{\rho(n)})_+^{s_{\sigma(n)} - k_{\sigma(n)} + k'_{\rho(n)}}}{(|x_{\sigma(n)} - y_{\rho(n)}|^2)^{s_{\sigma(n)} - k_{\sigma(n)} + k'_{\rho(n)} + 1}} \\
&\quad (s'_{\rho(n)} - k'_{\rho(n)} + k_{\sigma(1)} + 1)! \frac{(y_{\rho(n)} - x_{\sigma(1)})_+^{s'_{\rho(n)} - k'_{\rho(n)} + k_{\sigma(1)} + 1}}{(|y_{\rho(n)} - x_{\sigma(1)}|^2)^{s'_{\rho(n)} - k'_{\rho(n)} + k_{\sigma(1)} + 2}}
\end{aligned} \tag{8.27}$$

The 2-point correlators read:

$$\begin{aligned}
& \langle M_{s_1}(x) \bar{M}_{s_2}(y) \rangle = i(N^2 - 1) \frac{1}{(4\pi^2)^2} 2^{s_1+s_2-1} i^{s_1+s_2} (-1)^{s_1+s_2} \\
& \quad \sum_{k_1=0}^{s_1-1} \sum_{k_2=0}^{s_2-1} \binom{s_1}{k_1} \binom{s_1+1}{k_1+2} \binom{s_2+1}{k_2} \binom{s_2}{k_2+1} (s_1 - k_1 + k_2)! (s_2 - k_2 + k_1 + 1)! \\
& \quad \frac{(x - y)_+^{s_1 - k_1 + k_2}}{(|x - y|^2)^{s_1 - k_1 + k_2 + 1}} \frac{(y - x)_+^{s_2 - k_2 + k_1 + 1}}{(|y - x|^2)^{s_2 - k_2 + k_1 + 2}}
\end{aligned} \tag{8.28}$$

that become:

$$\begin{aligned}
& \langle M_{s_1}(x) \bar{M}_{s_2}(y) \rangle = -i(N^2 - 1) \frac{1}{(4\pi^2)^2} 2^{s_1+s_2-1} i^{s_1+s_2} \\
& \quad \sum_{k_1=0}^{s_1-1} \sum_{k_2=0}^{s_2-1} \binom{s_1}{k_1} \binom{s_1+1}{k_1+2} \binom{s_2+1}{k_2} \binom{s_2}{k_2+1} (s_1 - k_1 + k_2)! (s_2 - k_2 + k_1 + 1)! \\
& \quad (-1)^{s_1 - k_2 + k_1} \frac{(x - y)_+^{s_1 + s_2 + 1}}{(|x - y|^2)^{s_1 + s_2 + 3}}
\end{aligned} \tag{8.29}$$

Finally, employing (appendix C.2):

$$-\delta_{s_1 s_2} \frac{s_1}{s_1 + 2} = \sum_{k_1=0}^{s_1-1} \sum_{k_2=0}^{s_2-1} \binom{s_1}{k_1} \binom{s_1+1}{k_1+2} \binom{s_2+1}{k_2} \binom{s_2}{k_2+1} (-1)^{s_1-k_2+k_1} \frac{1}{\binom{s_1+s_2+1}{s_1-k_1+k_2}} \quad (8.30)$$

we obtain:

$$\langle M_{s_1}(x) \bar{M}_{s_2}(y) \rangle = \delta_{s_1 s_2} i(N^2 - 1) \frac{1}{(4\pi^2)^2} 2^{2s_1-1} (-1)^{s_1} \frac{s_1}{s_1 + 2} (2s_1 + 1)! \frac{(x-y)^{2s_1+1}}{(|x-y|^2)^{2s_1+3}} \quad (8.31)$$

8.4 Vanishing correlators

By inspecting eqs. (5.14) and (5.15), the mixed conformal correlators of O^A , \tilde{O}^A and O^λ , \tilde{O}^λ vanish for $n, m \geq 1$ and $l, k \geq 1$:

$$\langle O_{s_1}^A(x_1) \dots O_{s_n}^A(x_n) \tilde{O}_{s'_1}^A(y_1) \dots \tilde{O}_{s'_m}^A(y_m) O_{s''_1}^\lambda(z_1) \dots O_{s''_l}^\lambda(z_l) \tilde{O}_{s'''_1}^\lambda(t_1) \dots \tilde{O}_{s'''_k}^\lambda(t_k) \rangle = 0 \quad (8.32)$$

They also vanish for $l, k = 0$ and m odd or $n, m = 0$ and k odd.

9 Euclidean conformal correlators

We work out a few examples of the Euclidean n -point correlators.

9.1 O^{AE} correlators

The O^{AE} correlators read:

$$\begin{aligned} & \langle O_{s_1}^{AE}(x_1) \dots O_{s_n}^{AE}(x_n) \rangle \\ &= \frac{1}{(4\pi^2)^n} \frac{N^2 - 1}{2^n} 2^{\sum_{l=1}^n s_l} (-1)^{\sum_{l=1}^n s_l} \frac{2(s_1+1)(s_1+2)}{4!} \dots \frac{2(s_n+1)(s_n+2)}{4!} \\ & \sum_{k_1=0}^{s_1-2} \binom{s_1}{k_1} \binom{s_1}{k_1+2} \dots \sum_{k_n=0}^{s_n-2} \binom{s_n}{k_n} \binom{s_n}{k_n+2} \\ & \frac{1}{n} \sum_{\sigma \in P_n} (s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)})! \dots (s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)})! \\ & \frac{(x_{\sigma(1)} - x_{\sigma(2)}) z^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)}}}{((x_{\sigma(1)} - x_{\sigma(2)})^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)} + 1}} \dots \frac{(x_{\sigma(n)} - x_{\sigma(1)}) z^{s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)}}}{((x_{\sigma(n)} - x_{\sigma(1)})^2)^{s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)} + 1}} \end{aligned} \quad (9.1)$$

9.2 $O^{\lambda E}$ correlators

The $O^{\lambda E}$ correlators read:

$$\begin{aligned} \langle O_{s_1}^{\lambda E}(x_1) \dots O_{s_n}^{\lambda E}(x_n) \rangle &= -\frac{N^2 - 1}{2^n} \frac{1}{(4\pi^2)^n} (-1)^{\sum_{l=1}^n s_l} 2^{\sum_{l=1}^n s_l} \frac{(s_1 + 1)}{2} \dots \frac{(s_n + 1)}{2} \\ &\quad \sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1}{k_1 + 1} \dots \sum_{k_n=0}^{s_n-1} \binom{s_n}{k_n} \binom{s_n}{k_n + 1} \\ &\quad \frac{(-1)^n}{n} \sum_{\sigma \in P_n} (s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)})! \dots (s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)})! \\ &\quad \frac{(x_{\sigma(1)} - x_{\sigma(2)}) z^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)}}}{((x_{\sigma(1)} - x_{\sigma(2)})^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k_{\sigma(2)} + 1}} \dots \frac{(x_{\sigma(n)} - x_{\sigma(1)}) z^{s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)}}}{((x_{\sigma(n)} - x_{\sigma(1)})^2)^{s_{\sigma(n)} - k_{\sigma(n)} + k_{\sigma(1)} + 1}} \end{aligned} \quad (9.2)$$

9.3 M^E and \bar{M}^E correlators

The $2n$ -point correlators of M^E and \bar{M}^E read:

$$\begin{aligned} &\langle M_{s_1}^E(x_1) \bar{M}_{s'_1}^E(y_1) M_{s_2}^E(x_2) \bar{M}_{s'_2}^E(y_2) \dots M_{s_n}^E(x_n) \bar{M}_{s'_n}^E(y_n) \rangle \\ &= (N^2 - 1) \frac{1}{(4\pi^2)^{2n}} 2^{\sum_{l=1}^n s_l + \sum_{l=1}^n s'_l - n} \\ &\quad \sum_{k'_1=0}^{s'_1-1} \binom{s'_1 + 1}{k'_1} \binom{s'_1}{k'_1 + 1} \dots \sum_{k'_n=0}^{s'_n-1} \binom{s'_n + 1}{k'_n} \binom{s'_n}{k'_n + 1} \\ &\quad \sum_{k_1=0}^{s_1-1} \binom{s_1}{k_1} \binom{s_1 + 1}{k_1 + 2} \dots \sum_{k_n=0}^{s_n-1} \binom{s_n}{k_n} \binom{s_n + 1}{k_n + 2} \\ &\quad \sum_{\sigma \in P_n} \sum_{\rho \in P_n} \frac{(-1)^{2n+1}}{n} \text{sgn}(\sigma) \text{sgn}(\rho) \\ &\quad (s_{\sigma(1)} - k_{\sigma(1)} + k'_{\rho(1)})! \frac{(x_{\sigma(1)} - y_{\rho(1)}) z^{s_{\sigma(1)} - k_{\sigma(1)} + k'_{\rho(1)}}}{((x_{\sigma(1)} - y_{\rho(1)})^2)^{s_{\sigma(1)} - k_{\sigma(1)} + k'_{\rho(1)} + 1}} \\ &\quad (s'_{\rho(1)} - k'_{\rho(1)} + k_{\sigma(2)} + 1)! \frac{(y_{\rho(1)} - x_{\sigma(2)}) z^{s'_{\rho(1)} - k'_{\rho(1)} + k_{\sigma(2)} + 1}}{((y_{\rho(1)} - x_{\sigma(2)})^2)^{s'_{\rho(1)} - k'_{\rho(1)} + k_{\sigma(2)} + 2}} \\ &\quad \dots \\ &\quad (s_{\sigma(n)} - k_{\sigma(n)} + k'_{\rho(n)})! \frac{(x_{\sigma(n)} - y_{\rho(n)}) z^{s_{\sigma(n)} - k_{\sigma(n)} + k'_{\rho(n)}}}{((x_{\sigma(n)} - y_{\rho(n)})^2)^{s_{\sigma(n)} - k_{\sigma(n)} + k'_{\rho(n)} + 1}} \\ &\quad (s'_{\rho(n)} - k'_{\rho(n)} + k_{\sigma(n)} + 1)! \frac{(y_{\rho(n)} - x_{\sigma(1)}) z^{s'_{\rho(n)} - k'_{\rho(n)} + k_{\sigma(n)} + 1}}{((y_{\rho(n)} - x_{\sigma(1)})^2)^{s'_{\rho(n)} - k'_{\rho(n)} + k_{\sigma(n)} + 2}} \end{aligned} \quad (9.3)$$

Acknowledgments

We would like to thank Loris Del Grosso for participating in the early stages of this project.

A Notation and Wick rotation

We follow the conventions in [11]. We choose the mostly minus Minkowskian metric:

$$(g_{\mu\nu}) = \text{diag}(1, -1, -1, -1) \quad (\text{A.1})$$

and define the light-cone coordinates:

$$x^\pm = \frac{x^0 \pm x^3}{\sqrt{2}} = x_\mp \quad (\text{A.2})$$

The corresponding Minkowskian (squared) distance is:

$$|x|^2 = 2x^+x^- - x_\perp^2 \quad (\text{A.3})$$

with:

$$x_\perp^2 = (x^1)^2 + (x^2)^2 \quad (\text{A.4})$$

We denote the derivative with respect to x^+ by:

$$\partial_+ = \frac{\partial}{\partial x^+} = \partial_{x^+} = \frac{\partial}{\partial x_-} = \partial_{x_-} \quad (\text{A.5})$$

and define the light-like vectors n^μ and \bar{n}^μ :

$$n_\mu n^\mu = \bar{n}_\mu \bar{n}^\mu = 0 \quad n_\mu \bar{n}^\mu = 1 \quad (\text{A.6})$$

that respectively read $(n^\mu) = \frac{1}{\sqrt{2}}(1, 0, 0, 1)$ and $(\bar{n}^\mu) = \frac{1}{\sqrt{2}}(1, 0, 0, -1)$. Correspondingly, the Minkowskian metric decomposes into transverse and longitudinal components with respect to the light-like vectors:

$$g_{\mu\nu} = g_{\mu\nu}^\perp + n_\mu \bar{n}_\nu + n_\nu \bar{n}_\mu \quad (\text{A.7})$$

The Euclidean metric reads:

$$(\delta_{\mu\nu}) = \text{diag}(1, 1, 1, 1) \quad (\text{A.8})$$

with Euclidean (squared) distance:

$$x^2 = 2x^z x^{\bar{z}} + x_\perp^2 \quad (\text{A.9})$$

where:

$$x^z = \frac{x^4 + ix^3}{\sqrt{2}} = \frac{x_4 + ix_3}{\sqrt{2}} = x_{\bar{z}} \quad (\text{A.10})$$

and:

$$x^{\bar{z}} = \frac{x^4 - ix^3}{\sqrt{2}} = \frac{x_4 - ix_3}{\sqrt{2}} = x_z \quad (\text{A.11})$$

We define the Wick rotation by:

$$x^0 = x_0 \rightarrow -ix^4 = -ix_4 \quad (\text{A.12})$$

and:

$$p_0 = p^0 \rightarrow ip_4 = ip^4 \quad (\text{A.13})$$

Eq. (A.12) ensures that $\exp(iS_M) \rightarrow \exp(-S_E)$, where S_M and S_E are respectively the Minkowskian and Euclidean actions, with S_E positive definite. By defining $p \cdot x = p_\mu x^\mu$ and $\langle px \rangle = p_\mu x^\mu$ respectively in Minkowskian and Euclidean space-time, eq. (A.13) ensures that by the Wick rotation $p \cdot x \rightarrow \langle px \rangle$, so that the pairings $p \cdot x$ and $\langle px \rangle$ are actually independent of the Minkowskian and Euclidean metric. Therefore, by a slight abuse of notation, we also write $p \cdot x$ in Euclidean space-time, instead of $\langle px \rangle$. Besides, $|x|^2 \rightarrow -x^2$ and $|p|^2 \rightarrow -p^2$. As a consequence, the Wick rotation of the scalar propagator of mass m in Minkowskian space-time:

$$\langle \phi(x)\phi(y) \rangle = \int \frac{d^4 p}{(2\pi)^4} e^{ip \cdot (x-y)} \frac{i}{|p|^2 - m^2 + i\epsilon} \quad (\text{A.14})$$

reads in Euclidean space-time:

$$\langle \phi^E(x)\phi^E(y) \rangle = \int \frac{d^4 p}{(2\pi)^4} e^{ip \cdot (x-y)} \frac{i^2}{-p^2 - m^2} = \int \frac{d^4 p}{(2\pi)^4} e^{ip \cdot (x-y)} \frac{1}{p^2 + m^2} \quad (\text{A.15})$$

as it should be. The Wick rotation of the light-cone coordinates:

$$x^+ = x_- \rightarrow -ix^z = -ix_{\bar{z}} \quad (\text{A.16})$$

and:

$$x^- = x_+ \rightarrow -ix^{\bar{z}} = -ix_z \quad (\text{A.17})$$

implies the Wick rotation of the derivative with respect to x^+ :

$$\partial_+ \rightarrow i\partial_z = i\frac{\partial}{\partial x^z} \quad (\text{A.18})$$

B Jacobi and Gegenbauer polynomials

We work out some formulas for the Jacobi and Gegenbauer polynomials employed in the present paper. For x real the Jacobi polynomials $P_l^{(\alpha,\beta)}(x)$ admit the representation [12]:

$$P_l^{(\alpha,\beta)}(x) = \sum_{k=0}^l \binom{l+\alpha}{k} \binom{l+\beta}{k+\beta} \left(\frac{x-1}{2}\right)^k \left(\frac{x+1}{2}\right)^{l-k} \quad (\text{B.1})$$

with α, β real and l a natural number. Besides, they satisfy the symmetry property:

$$P_l^{(\alpha,\beta)}(-x) = (-1)^l P_l^{(\beta,\alpha)}(x) \quad (\text{B.2})$$

The Gegenbauer polynomials $C_l^{\alpha'}(x)$ are a special case of the Jacobi polynomials:

$$C_l^{\alpha'}(x) = \frac{\Gamma(l+2\alpha')\Gamma(\alpha' + \frac{1}{2})}{\Gamma(2\alpha')\Gamma(l+\alpha'+\frac{1}{2})} P_l^{(\alpha'-\frac{1}{2}, \alpha'-\frac{1}{2})}(x) \quad (\text{B.3})$$

with the symmetry property:

$$C_l^{\alpha'}(-x) = (-1)^l C_l^{\alpha'}(x) \quad (\text{B.4})$$

We set:

$$x = \frac{b-a}{a+b} \quad (\text{B.5})$$

so that:

$$\left(\frac{x-1}{2}\right)^k \left(\frac{x+1}{2}\right)^{l-k} = (-1)^{l-k} \frac{a^{l-k} b^k}{(a+b)^l} \quad (\text{B.6})$$

Eq. (B.1) becomes:

$$P_l^{(\alpha, \beta)}(x) = \sum_{k=0}^l \binom{l+\alpha}{k} \binom{l+\beta}{k+\beta} (-1)^{l-k} \frac{a^{l-k} b^k}{(a+b)^l} \quad (\text{B.7})$$

Moreover, putting $l = J - \alpha' + \frac{1}{2}$ in eq. (E.2) and $\alpha = \beta = \alpha' - \frac{1}{2}$ in eq. (B.1), we obtain:

$$\begin{aligned} C_{J-\alpha'+\frac{1}{2}}^{\alpha'}(x) &= \frac{\Gamma(J + \frac{1}{2} + \alpha') \Gamma(\alpha' + \frac{1}{2})}{\Gamma(2\alpha') \Gamma(J+1)} \\ &\quad \sum_{k=0}^{J-\alpha'+\frac{1}{2}} \binom{J}{k} \binom{J}{k+\alpha'-\frac{1}{2}} (-1)^{J-\alpha'+\frac{1}{2}-k} \frac{a^{J-\alpha'+\frac{1}{2}-k} b^k}{(a+b)^{J-\alpha'+\frac{1}{2}}} \end{aligned} \quad (\text{B.8})$$

Specializing the above equation to $J = s$ and $\alpha' = \frac{5}{2}$, we get:

$$C_{s-2}^{\frac{5}{2}}(x) = \frac{\Gamma(s+3)\Gamma(3)}{\Gamma(5)\Gamma(s+1)} \sum_{k=0}^{s-2} \binom{s}{k} \binom{s}{k+2} (-1)^{s-k} \frac{a^{s-k-2} b^k}{(a+b)^{s-2}} \quad (\text{B.9})$$

Hence, for $J = s$ and $\alpha' = \frac{1}{2}$, we obtain:

$$C_s^{\frac{1}{2}}(x) = \sum_{k=0}^s \binom{s}{k} \binom{s}{k} (-1)^{s-k} \frac{a^{s-k} b^k}{(a+b)^s} \quad (\text{B.10})$$

We restrict α, β to the natural numbers and, correspondingly, α' to the positive half-integers and J to the natural numbers. By employing the identity:

$$\binom{l+\alpha}{k} \binom{l+\beta}{k+\beta} = \frac{(l+\beta)!(l+\alpha)!}{l!(l+\alpha+\beta)!} \binom{l}{k} \binom{l+\beta+\alpha}{k+\beta} \quad (\text{B.11})$$

it follows from eq. (B.7) that:

$$P_l^{(\alpha, \beta)}(x) = \frac{(l+\beta)!(l+\alpha)!}{l!(l+\alpha+\beta)!} \sum_{k=0}^l \binom{l}{k} \binom{l+\beta+\alpha}{k+\beta} (-1)^{l-k} \frac{a^{l-k} b^k}{(a+b)^l} \quad (\text{B.12})$$

Correspondingly, eq. (E.2) reads:

$$\begin{aligned} C_l^{\alpha'}(x) &= \frac{\Gamma(l+2\alpha')\Gamma(\alpha'+\frac{1}{2})}{\Gamma(2\alpha')\Gamma(l+\alpha'+\frac{1}{2})} \frac{(l+\alpha'-\frac{1}{2})!(l+\alpha'-\frac{1}{2})!}{l!(l+2\alpha'-1)!} \\ &\quad \sum_{k=0}^l \binom{l}{k} \binom{l+2\alpha'-1}{k+\alpha'-\frac{1}{2}} (-1)^{l-k} \frac{a^{l-k} b^k}{(a+b)^l} \end{aligned} \quad (\text{B.13})$$

that reduces to:

$$C_l^{\alpha'}(x) = \frac{\Gamma(\alpha' + \frac{1}{2})\Gamma(l + \alpha' + \frac{1}{2})}{\Gamma(2\alpha')\Gamma(l+1)} \sum_{k=0}^l \binom{l}{k} \binom{l+2\alpha'-1}{k+\alpha'-\frac{1}{2}} (-1)^{l-k} \frac{a^{l-k}b^k}{(a+b)^l} \quad (\text{B.14})$$

C Normalization of 2-point correlators

We compute the normalization of the 2-point correlators by a technique [13] involving the orthogonality of Gegenbauer polynomials that makes manifest the vanishing of correlators of operators with different spins.

C.1 2-point correlators of gluino-gluino operators

The 2-point point correlators of gluino-gluino operators read:

$$\begin{aligned} \langle O_{s_1}^\lambda(x) O_{s_2}^\lambda(y) \rangle &= -\frac{1}{4} \frac{N^2 - 1}{(4\pi^2)^2} \mathcal{Y}_{s_1-1}^{\frac{3}{2}}(\partial_{x_1^+}, \partial_{x_2^+}) \mathcal{Y}_{s_2-1}^{\frac{3}{2}}(\partial_{y_1^+}, \partial_{y_2^+}) \\ &\quad \partial_{x_1^+} \partial_{x_2^+} \frac{1}{|x_1 - y_2|^2} \frac{1}{|x_2 - y_1|^2} \Big|_{x_1=x_2=x}^{y_1=y_2=y} \end{aligned} \quad (\text{C.1})$$

Initially, we restrict to $(x-y)_\perp = 0$, so that $|x-y|^2 = 2(x-y)_+(x-y)_-$. For $x_- > y_-$ we get:

$$\frac{\Gamma(k)}{(x-y)_-^k} = \int_0^\infty d\tau \tau^{k-1} e^{-\tau(x-y)_-} \quad (\text{C.2})$$

By the above equation we convert derivatives into multiplications:

$$\begin{aligned} \partial_{x_1^+} &\rightarrow -\tau_1, & \partial_{x_2^+} &\rightarrow -\tau_2 \\ \partial_{y_1^+} &\rightarrow \tau_2, & \partial_{y_2^+} &\rightarrow \tau_1 \end{aligned} \quad (\text{C.3})$$

By the symmetry properties (appendix B):

$$\begin{aligned} \mathcal{Y}_{s_1-1}^{\frac{3}{2}}(-\tau_1, -\tau_2) &= (-1)^{s_1-1} (i\tau_1 + i\tau_2)^{s_1-1} C_{s_1-1}^{\frac{3}{2}} \left(\frac{\tau_1 - \tau_2}{\tau_1 + \tau_2} \right) \\ \mathcal{Y}_{s_2-1}^{\frac{3}{2}}(\tau_2, \tau_1) &= (-1)^{s_2-1} (i\tau_1 + i\tau_2)^{s_2-1} C_{s_2-1}^{\frac{3}{2}} \left(\frac{\tau_1 - \tau_2}{\tau_1 + \tau_2} \right) \end{aligned} \quad (\text{C.4})$$

we obtain:

$$\begin{aligned} \langle O_{s_1}^\lambda(x) O_{s_2}^\lambda(y) \rangle &= -\frac{1}{4} \frac{N^2 - 1}{(4\pi^2)^2} i^{s_1+s_2-2} \frac{1}{4(x-y)_+^2} (-1)^{s_1+s_2} \\ &\quad \int_0^\infty d\tau_1 d\tau_2 (\tau_1 + \tau_2)^{s_1+s_2-2} \tau_1 \tau_2 e^{-(\tau_1 + \tau_2)(x-y)_-} C_{s_1-1}^{\frac{3}{2}} \left(\frac{\tau_1 - \tau_2}{\tau_1 + \tau_2} \right) C_{s_2-1}^{\frac{3}{2}} \left(\frac{\tau_1 - \tau_2}{\tau_1 + \tau_2} \right) \end{aligned} \quad (\text{C.5})$$

Changing variables:

$$\tau_1 = \tau\alpha, \quad \tau_2 = \tau(1-\alpha) \quad (\text{C.6})$$

we get:

$$\begin{aligned} \langle O_{s_1}^\lambda(x) O_{s_2}^\lambda(y) \rangle &= -\frac{1}{4} \frac{N^2 - 1}{(4\pi^2)^2} i^{s_1+s_2-2} \frac{1}{4(x-y)_+^2} (-1)^{s_1+s_2} \\ &\quad \int_0^\infty d\tau \int_0^1 d\alpha \tau^{s_1+s_2+1} \alpha(1-\alpha) e^{-\tau(x-y)_-} C_{s_1-1}^{\frac{3}{2}}(1-2\alpha) C_{s_2-1}^{\frac{3}{2}}(1-2\alpha) \end{aligned} \quad (\text{C.7})$$

The τ integral yields:

$$\int_0^\infty d\tau \tau^{s_1+s_2+1} e^{-\tau(x-y)_-} = \frac{\Gamma(s_1+s_2+2)}{(x-y)_-^{s_1+s_2+2}} \quad (\text{C.8})$$

The α integral:

$$\int_0^1 d\alpha \alpha(1-\alpha) C_{s_1-1}^{\frac{3}{2}}(1-2\alpha) C_{s_2-1}^{\frac{3}{2}}(1-2\alpha) \quad (\text{C.9})$$

can be written as:

$$\int_{-1}^1 \frac{du}{2} \left(\frac{1-u^2}{4} \right) C_{s_1-1}^{\frac{3}{2}}(u) C_{s_2-1}^{\frac{3}{2}}(u) \quad (\text{C.10})$$

where $u = 1 - 2\alpha$. The orthonormality property of Gegenbauer polynomials reads:

$$\int_{-1}^1 dz (1-z^2)^{\alpha-\frac{1}{2}} C_{n_1}^\alpha(z) C_{n_2}^\alpha(z) = \delta_{n_1, n_2} \frac{\pi 2^{1-2\alpha} \Gamma(n_1 + 2\alpha)}{n_1! (n_1 + \alpha) \Gamma(\alpha)^2} \quad (\text{C.11})$$

so that for $\alpha = \frac{3}{2}$ we obtain:

$$\int_{-1}^1 \frac{du}{2} \frac{(1-u^2)}{4} C_{s_1-1}^{\frac{3}{2}}(u) C_{s_2-1}^{\frac{3}{2}}(u) = \delta_{s_1, s_2} \frac{1}{4} \frac{s_1(s_1+1)}{2s_1+1} \quad (\text{C.12})$$

Collecting all the above factors:

$$\begin{aligned} \langle O_{s_1}^\lambda(x) O_{s_2}^\lambda(y) \rangle &= -\frac{1}{4} \frac{N^2 - 1}{(4\pi^2)^2} i^{s_1+s_2-2} (-1)^{s_1+s_2} \frac{1}{4(x-y)_+^2 (x-y)_-^{s_1+s_2+2}} \\ &\quad \frac{1}{4} s_1(s_1+1) \delta_{s_1, s_2} \frac{\Gamma(s_1+s_2+2)}{2s_1+1} \end{aligned} \quad (\text{C.13})$$

and employing the identity:

$$\frac{1}{4(x-y)_+^2 (x-y)_-^{s_1+s_2+2}} = 2^{s_1+s_2} \frac{(x-y)_+^{s_1+s_2}}{(|x-y|^2)^{s_1+s_2+2}} \quad (\text{C.14})$$

yields the desired result:

$$\langle O_{s_1}^\lambda(x) O_{s_2}^\lambda(y) \rangle = -\delta_{s_1 s_2} \frac{1}{4} \frac{N^2 - 1}{(4\pi^2)^2} i^{2s_1-2} (-1)^{2s_1} 2^{2s_1} \frac{(x-y)_+^{2s_1}}{(|x-y|^2)^{2s_1+2}} \frac{s_1(s_1+1)}{4} \frac{\Gamma(2s_1+2)}{2s_1+1} \quad (\text{C.15})$$

Incidentally, equating the above equation to eq. (8.12) we deduce the identity:

$$\delta_{s_1 s_2} \frac{s_1}{s_1+1} = \sum_{k_1=0}^{s_1-1} \sum_{k_2=0}^{s_2-1} \binom{s_1}{k_1} \binom{s_1}{k_1+1} \binom{s_2}{k_2} \binom{s_2}{k_2+1} (-1)^{k_1+k_2} \frac{1}{\binom{s_1+s_2}{k_1+k_2+1}} \quad (\text{C.16})$$

C.2 2-point correlators of gluon-gluino operators

The 2-point point correlators of gluon-gluino operators read:

$$\langle M_{s_1}(x) \bar{M}_{s_2}(y) \rangle = (N^2 - 1) \frac{i}{(4\pi^2)^2} \frac{1}{2^2} \partial_{y_1^+} \frac{1}{|y_1 - x_2|^2} \mathcal{G}_{s_2-1}^{(1,2)}(\partial_{y_1^+}, \partial_{y_2^+}) \frac{1}{|y_2 - x_1|^2} \mathcal{G}_{s_1-1}^{(2,1)}(\partial_{x_1^+}, \partial_{x_2^+}) \Big|_{x_1=x_2=x}^{y_1=y_2=y} \quad (\text{C.17})$$

where:

$$\begin{aligned} \mathcal{G}_{s-1}^{(1,2)}(\partial_{y_1^+}, \partial_{y_2^+}) &= (i\partial_{y_1^+} + i\partial_{y_2^+})^{s-1} P_{s-1}^{(1,2)} \left(\frac{\partial_{y_1^+} - \partial_{y_2^+}}{\partial_{y_1^+} + \partial_{y_2^+}} \right) \partial_{y_2^+} \\ \mathcal{G}_{s-1}^{(2,1)}(\partial_{x_1^+}, \partial_{x_2^+}) &= \partial_{x_1^+} (i\partial_{x_1^+} + i\partial_{x_2^+})^{s-1} P_{s-1}^{(2,1)} \left(\frac{\partial_{x_1^+} - \partial_{x_2^+}}{\partial_{x_1^+} + \partial_{x_2^+}} \right) \end{aligned} \quad (\text{C.18})$$

Employing eqs. (C.2), (C.3) and the symmetry properties (appendix B):

$$\begin{aligned} \mathcal{G}_{s-1}^{(1,2)}(-\tau_1, -\tau_2) &= (-1)^{s_1} \tau_2 (i\tau_1 + i\tau_2)^{s_1-1} P_{s_1-1}^{(1,2)} \left(\frac{\tau_1 - \tau_2}{\tau_1 + \tau_2} \right) \\ \mathcal{G}_{s-1}^{(2,1)}(\tau_2, \tau_1) &= (-1)^{s_2-1} (i\tau_1 + i\tau_2)^{s_2-1} P_{s_2-1}^{(1,2)} \left(\frac{\tau_1 - \tau_2}{\tau_1 + \tau_2} \right) \tau_2 \end{aligned} \quad (\text{C.19})$$

we obtain:

$$\begin{aligned} \langle M_{s_1}(x) \bar{M}_{s_2}(y) \rangle &= (N^2 - 1) \frac{i}{(4\pi^2)^2} \frac{1}{2^2} i^{s_1+s_2-2} \frac{1}{4(y-x)_+^2} (-1)^{s_1+s_2} \\ &\quad \int_0^\infty d\tau_1 d\tau_2 (\tau_1 + \tau_2)^{s_1+s_2-2} \tau_1 \tau_2^2 e^{-(\tau_1+\tau_2)(y-x)_-} P_{s_1-1}^{(1,2)} \left(\frac{\tau_1 - \tau_2}{\tau_1 + \tau_2} \right) P_{s_2-1}^{(1,2)} \left(\frac{\tau_1 - \tau_2}{\tau_1 + \tau_2} \right) \end{aligned} \quad (\text{C.20})$$

Changing variables:

$$\tau_1 = \tau\alpha, \quad \tau_2 = \tau(1-\alpha) \quad (\text{C.21})$$

we get:

$$\begin{aligned} \langle M_{s_1}(x) \bar{M}_{s_2}(y) \rangle &= (N^2 - 1) \frac{i}{(4\pi^2)^2} \frac{1}{2^2} i^{s_1+s_2-2} \frac{1}{4(x-y)_+^2} (-1)^{s_1+s_2} \\ &\quad \int_0^\infty d\tau \int_0^1 d\alpha \tau^{s_1+s_2+2} \alpha(1-\alpha)^2 e^{-\tau(x-y)_-} P_{s_1-1}^{(1,2)}(1-2\alpha) P_{s_2-1}^{(1,2)}(1-2\alpha) \end{aligned} \quad (\text{C.22})$$

Integrating on τ yields:

$$\int_0^\infty d\tau \tau^{s_1+s_2+2} e^{-\tau(y-x)_-} = \frac{\Gamma(s_1 + s_2 + 3)}{(y-x)_-^{s_1+s_2+3}} \quad (\text{C.23})$$

Then, we rewrite:

$$\int_0^1 d\alpha \alpha(1-\alpha)^2 P_{s_1-1}^{(1,2)}(1-2\alpha) P_{s_2-1}^{(1,2)}(1-2\alpha) \quad (\text{C.24})$$

as:

$$\int_{-1}^1 \frac{du}{2} \left(\frac{1-u}{2} \right) \left(\frac{1+u}{2} \right)^2 P_{s_1-1}^{(1,2)}(u) P_{s_2-1}^{(1,2)}(u) \quad (\text{C.25})$$

with $u = 1 - 2\alpha$. The orthonormality property of Jacobi polynomials reads:

$$\int_{-1}^1 (1-u)^\alpha (1+u)^\beta P_m^{(\alpha,\beta)}(u) P_n^{(\alpha,\beta)}(u) du = \delta_{nm} \frac{2^{\alpha+\beta+1}}{2n + \alpha + \beta + 1} \frac{\Gamma(n+\alpha+1)\Gamma(n+\beta+1)}{\Gamma(n+\alpha+\beta+1)n!} \quad (\text{C.26})$$

so that for $\alpha = 1$ and $\beta = 2$ we obtain:

$$\int_{-1}^1 \frac{du}{2} \left(\frac{1-u}{2} \right) \left(\frac{1+u}{2} \right)^2 P_{s_1-1}^{(1,2)}(u) P_{s_2-1}^{(1,2)}(u) = \delta_{s_1 s_2} \frac{1}{2s_1 + 2} \frac{\Gamma(s_1+1)\Gamma(s_1+2)}{\Gamma(s_1+3)(s_1-1)!} \quad (\text{C.27})$$

Collecting the above results:

$$\begin{aligned} \langle M_{s_1}(x) \bar{M}_{s_2}(y) \rangle &= \delta_{s_1 s_2} (N^2 - 1) \frac{i}{(4\pi^2)^2} \frac{1}{2^2} i^{2s_1-2} (-1)^{2s_1} \frac{\Gamma(2s_1+3)}{4(y-x)_+^2 (y-x)_-^{2s_1+3}} \\ &\quad \frac{1}{2s_1 + 2} \frac{\Gamma(s_1+1)\Gamma(s_1+2)}{\Gamma(s_1+3)(s_1-1)!} \end{aligned} \quad (\text{C.28})$$

and employing the identity:

$$\frac{1}{4(y-x)_+^2 (y-x)_-^{s_1+s_2+3}} = 2^{s_1+s_2+1} (-1)^{s_1+s_2+1} \frac{(x-y)_+^{s_1+s_2+1}}{(|x-y|^2)^{s_1+s_2+3}} \quad (\text{C.29})$$

we get:

$$\langle M_{s_1}(x) \bar{M}_{s_2}(y) \rangle = \delta_{s_1 s_2} (N^2 - 1) \frac{i}{(4\pi^2)^2} i^{s_1+s_2} \Gamma(2s_1+3) 2^{2s_1-1} \frac{1}{2s_1 + 2} \frac{s_1}{s_1 + 2} \frac{(x-y)_+^{2s_1+1}}{(|x-y|^2)^{2s_1+3}} \quad (\text{C.30})$$

Incidentally, equating the above equation to eq. (8.29) we deduce the identity:

$$-\delta_{s_1 s_2} \frac{s_1}{s_1 + 2} = \sum_{k_1=0}^{s_1-1} \sum_{k_2=0}^{s_2-1} \binom{s_1}{k_1} \binom{s_1+1}{k_1+2} \binom{s_2+1}{k_2} \binom{s_2}{k_2+1} (-1)^{s_1-k_2+k_1} \frac{1}{\binom{s_1+s_2+1}{s_1-k_1+k_2}} \quad (\text{C.31})$$

D An equality of determinants

We prove the equality of the two functional determinants in eq. (8.19) arising from fermionic integration. Their expansion reads:

$$\begin{aligned}
& \frac{1}{2} \log \text{Det} \left(\mathcal{I} + \frac{1}{4} i \partial_+ i \square^{-1} \bar{J}_{M_{s_1}} \otimes \mathcal{G}_{s_1-1}^{(1,2)} (-1)^{s_1-1} i \square^{-1} J_{\bar{M}_{s_3}} \otimes \mathcal{G}_{s_3-1}^{(2,1)} (-1)^{s_3-1} \right) \\
&= \frac{N^2 - 1}{2} \sum_{l=1}^{\infty} \frac{(-1)^{l+1}}{l} \frac{1}{2^{2l}} \int d^4 u_1 \dots d^4 u_l d^4 v_1 \dots d^4 v_l \sum_{s_1} \dots \sum_{s_l} \sum_{s'_1} \dots \sum_{s'_l} \\
&\quad (-1)^{\sum_{i=1}^l s_i + s'_i} \\
&\quad i \partial_{v_1^+} i \square^{-1} (v_1 - u_1) \mathcal{G}_{s_1-1}^{(1,2)} (\overleftarrow{\partial}_{u_1^+}, \overrightarrow{\partial}_{u_1^+}) \otimes \bar{J}_{M_{s_1}} (u_1) \\
&\quad i \square^{-1} (u_1 - v_2) \mathcal{G}_{s_2-1}^{(2,1)} (\overleftarrow{\partial}_{v_2^+}, \overrightarrow{\partial}_{v_2^+}) \otimes J_{\bar{M}_{s'_2}} (v_2) \\
&\quad \dots i \partial_{v_l^+} i \square^{-1} (v_l - u_l) \mathcal{G}_{s_l-1}^{(1,2)} (\overleftarrow{\partial}_{u_l^+}, \overrightarrow{\partial}_{u_l^+}) \otimes \bar{J}_{M_{s_l}} (u_l) \\
&\quad i \square^{-1} (u_l - v_1) \mathcal{G}_{s'_1-1}^{(2,1)} (\overleftarrow{\partial}_{v_1^+}, \overrightarrow{\partial}_{v_1^+}) \otimes J_{\bar{M}_{s'_1}} (v_1)
\end{aligned} \tag{D.1}$$

and:

$$\begin{aligned}
& \frac{1}{2} \log \text{Det} \left(\mathcal{I} + \frac{1}{4} i \partial_+ i \square^{-1} J_{\bar{M}_{s_1}} \otimes \mathcal{G}_{s_1-1}^{(1,2)} i \square^{-1} \bar{J}_{M_{s_3}} \otimes \mathcal{G}_{s_3-1}^{(2,1)} \right) \\
&= \frac{N^2 - 1}{2} \sum_{l=1}^{\infty} \frac{(-1)^{l+1}}{l} \frac{1}{2^{2l}} \int d^4 u_1 \dots d^4 u_l d^4 v_1 \dots d^4 v_l \sum_{s_1} \dots \sum_{s_l} \sum_{s'_1} \dots \sum_{s'_l} \\
&\quad i \partial_{u_1^+} i \square^{-1} (u_1 - v_1) \mathcal{G}_{s'_1-1}^{(1,2)} (\overleftarrow{\partial}_{v_1^+}, \overrightarrow{\partial}_{v_1^+}) \otimes J_{\bar{M}_{s'_1}} (v_1) \\
&\quad i \square^{-1} (v_1 - u_2) \mathcal{G}_{s_2-1}^{(2,1)} (\overleftarrow{\partial}_{u_2^+}, \overrightarrow{\partial}_{u_2^+}) \otimes \bar{J}_{M_{s_2}} (u_2) \\
&\quad \dots i \partial_{u_l^+} i \square^{-1} (u_l - v_l) \mathcal{G}_{s_l-1}^{(1,2)} (\overleftarrow{\partial}_{v_l^+}, \overrightarrow{\partial}_{v_l^+}) \otimes J_{\bar{M}_{s'_l}} (v_l) \\
&\quad i \square^{-1} (v_l - u_1) \mathcal{G}_{s_1-1}^{(2,1)} (\overleftarrow{\partial}_{u_1^+}, \overrightarrow{\partial}_{u_1^+}) \otimes \bar{J}_{M_{s_1}} (u_1)
\end{aligned} \tag{D.2}$$

In eq. (D.1) employing the property of the Jacobi polynomials:

$$f(x) \mathcal{G}_{s-1}^{(\alpha, \beta)} (\overleftarrow{\partial}_+, \overrightarrow{\partial}_+) g(x) = (-1)^{s-1} g(x) \mathcal{G}_{s-1}^{(\beta, \alpha)} (\overleftarrow{\partial}_+, \overrightarrow{\partial}_+) f(x) \tag{D.3}$$

and the cyclicity of the trace, we obtain:

$$\begin{aligned}
& (-1)^{\sum_{i=1}^l s_i + s'_i} \\
& i\partial_{v_1^+} i\square^{-1}(v_1 - u_1) \mathcal{G}_{s_1-1}^{(1,2)}(\overleftarrow{\partial}_{u_1^+}, \overrightarrow{\partial}_{u_1^+}) \otimes \bar{J}_{M_{s_1}}(u_1) \\
& i\square^{-1}(u_1 - v_2) \mathcal{G}_{s_2'-1}^{(2,1)}(\overleftarrow{\partial}_{v_2^+}, \overrightarrow{\partial}_{v_2^+}) \otimes J_{\bar{M}_{s'_2}}(v_2) \\
& \dots i\partial_{v_l^+} i\square^{-1}(v_l - u_l) \mathcal{G}_{s_l-1}^{(1,2)}(\overleftarrow{\partial}_{u_l^+}, \overrightarrow{\partial}_{u_l^+}) \otimes \bar{J}_{M_{s_l}}(u_l) \\
& i\square^{-1}(u_l - v_1) \mathcal{G}_{s_1'-1}^{(2,1)}(\overleftarrow{\partial}_{v_1^+}, \overrightarrow{\partial}_{v_1^+}) \otimes J_{\bar{M}_{s'_1}}(v_1) \\
& = \\
& i\partial_{u_1^+} i\square^{-1}(u_1 - v_1) \mathcal{G}_{s_1'-1}^{(1,2)}(\overleftarrow{\partial}_{v_1^+}, \overrightarrow{\partial}_{v_1^+}) \otimes J_{\bar{M}_{s'_1}}(v_1) \\
& i\square^{-1}(v_1 - u_l) \mathcal{G}_{s_l-1}^{(2,1)}(\overleftarrow{\partial}_{u_l^+}, \overrightarrow{\partial}_{u_l^+}) \otimes \bar{J}_{M_{s_l}}(u_l) \\
& i\partial_{u_l^+} i\square^{-1}(u_l - v_l) \mathcal{G}_{s_l'-1}^{(1,2)}(\overleftarrow{\partial}_{v_l^+}, \overrightarrow{\partial}_{v_l^+}) \otimes J_{\bar{M}_{s'_l}}(v_l) \\
& i\square^{-1}(v_l - u_{l-1}) \mathcal{G}_{s_{l-1}-1}^{(2,1)}(\overleftarrow{\partial}_{u_{l-1}^+}, \overrightarrow{\partial}_{u_{l-1}^+}) \otimes \bar{J}_{M_{s_{l-1}}}(u_{l-1}) \\
& \dots i\partial_{u_2^+} i\square^{-1}(u_2 - v_2) \mathcal{G}_{s_2'-1}^{(1,2)}(\overleftarrow{\partial}_{v_2^+}, \overrightarrow{\partial}_{v_2^+}) \otimes J_{\bar{M}_{s'_2}}(v_2) \\
& i\square^{-1}(v_2 - u_1) \mathcal{G}_{s_1-1}^{(2,1)}(\overleftarrow{\partial}_{u_1^+}, \overrightarrow{\partial}_{u_1^+}) \otimes \bar{J}_{M_{s_1}}(u_1)
\end{aligned} \tag{D.4}$$

where we have employed:

$$\partial_{v^+} \square^{-1}(v - u) = -\partial_{u^+} \square^{-1}(u - v) \tag{D.5}$$

and cyclically permuted the anticommuting sources, so that the factors of $(-1)^n$ arising from eq. (D.5) and the permutation of the anticommuting sources cancel out. Relabeling the variables $x_i s_i k_i \rightarrow x_{l-i+2} s_{l-i+2} k_{l-i+2}$ for $2 \leq i \leq l$ and keeping $x_1 s_1 k_1$ fixed with

$x = u, v$ we obtain:

$$\begin{aligned}
& i\partial_{u_1^+} i\square^{-1}(u_1 - v_1) \mathcal{G}_{s'_1-1}^{(1,2)} \otimes J_{\bar{M}_{s'_1}}(v_1) (\overleftarrow{\partial}_{v_1^+}, \overrightarrow{\partial}_{v_1^+}) \\
& i\square^{-1}(v_1 - u_l) \mathcal{G}_{s_l-1}^{(2,1)} (\overleftarrow{\partial}_{u_l^+}, \overrightarrow{\partial}_{u_l^+}) \otimes \bar{J}_{M_{s_l}}(u_l) \\
& i\partial_{u_l^+} i\square^{-1}(u_l - v_l) \mathcal{G}_{s'_l-1}^{(1,2)} (\overleftarrow{\partial}_{v_l^+}, \overrightarrow{\partial}_{v_l^+}) \otimes J_{\bar{M}_{s'_l}}(v_l) \\
& i\square^{-1}(v_l - u_{l-1}) \mathcal{G}_{s_{l-1}-1}^{(2,1)} (\overleftarrow{\partial}_{u_{l-1}^+}, \overrightarrow{\partial}_{u_{l-1}^+}) \otimes \bar{J}_{M_{s_{l-1}}}(u_{l-1}) \\
& \dots i\partial_{u_2^+} i\square^{-1}(u_2 - v_2) \mathcal{G}_{s_2-1}^{(1,2)} (\overleftarrow{\partial}_{v_2^+}, \overrightarrow{\partial}_{v_2^+}) \otimes J_{\bar{M}_{s_2}}(v_2) \\
& i\square^{-1}(v_2 - u_1) \mathcal{G}_{s_1-1}^{(2,1)} (\overleftarrow{\partial}_{u_1^+}, \overrightarrow{\partial}_{u_1^+}) \otimes \bar{J}_{M_{s_1}}(u_1) \\
& = \\
& i\partial_{u_1^+} i\square^{-1}(u_1 - v_1) \mathcal{G}_{s'_1-1}^{(1,2)} (\overleftarrow{\partial}_{v_1^+}, \overrightarrow{\partial}_{v_1^+}) \otimes J_{\bar{M}_{s'_1}}(v_1) \\
& i\square^{-1}(v_1 - u_2) \mathcal{G}_{s_2-1}^{(2,1)} (\overleftarrow{\partial}_{u_2^+}, \overrightarrow{\partial}_{u_2^+}) \otimes \bar{J}_{M_{s_2}}(u_2) \\
& \dots i\partial_{u_l^+} i\square^{-1}(u_l - v_l) \mathcal{G}_{s'_l-1}^{(1,2)} (\overleftarrow{\partial}_{v_l^+}, \overrightarrow{\partial}_{v_l^+}) \otimes J_{\bar{M}_{s'_l}}(v_l) \\
& i\square^{-1}(v_l - u_1) \mathcal{G}_{s_1-1}^{(2,1)} (\overleftarrow{\partial}_{u_1^+}, \overrightarrow{\partial}_{u_1^+}) \otimes \bar{J}_{M_{s_1}}(u_1)
\end{aligned} \tag{D.6}$$

It follows:

$$\begin{aligned}
& \log \text{Det} \left(\mathcal{I} + \frac{1}{4} i\partial_+ i\square^{-1} \bar{J}_{M_{s_1}} \otimes \mathcal{G}_{s_1-1}^{(1,2)} (-1)^{s_1-1} i\square^{-1} J_{\bar{M}_{s_3}} \otimes \mathcal{G}_{s_3-1}^{(2,1)} (-1)^{s_3-1} \right) \\
& = \log \text{Det} \left(\mathcal{I} + \frac{1}{4} i\partial_+ i\square^{-1} J_{\bar{M}_{s_1}} \otimes \mathcal{G}_{s_1-1}^{(1,2)} i\square^{-1} \bar{J}_{M_{s_3}} \otimes \mathcal{G}_{s_3-1}^{(2,1)} \right)
\end{aligned} \tag{D.7}$$

E Conformal properties of the standard basis

The gauge-invariant collinear twist-2 operators in the light-cone gauge that enter the balanced superfields read in the standard basis [2, 6, 8]:

$$\begin{aligned}
O_s^A &= \frac{1}{2} \partial_+ \bar{A}^a (i \overrightarrow{\partial}_+ + i \overleftarrow{\partial}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left(\frac{\overrightarrow{\partial}_+ - \overleftarrow{\partial}_+}{\overrightarrow{\partial}_+ + \overleftarrow{\partial}_+} \right) \partial_+ A^a \\
\tilde{O}_s^A &= \frac{1}{2} \partial_+ \bar{A}^a (i \overrightarrow{\partial}_+ + i \overleftarrow{\partial}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left(\frac{\overrightarrow{\partial}_+ - \overleftarrow{\partial}_+}{\overrightarrow{\partial}_+ + \overleftarrow{\partial}_+} \right) \partial_+ A^a \\
O_s^\lambda &= \frac{1}{2} \bar{\lambda}^a (i \overrightarrow{\partial}_+ + i \overleftarrow{\partial}_+)^{s-1} C_{s-1}^{\frac{3}{2}} \left(\frac{\overrightarrow{\partial}_+ - \overleftarrow{\partial}_+}{\overrightarrow{\partial}_+ + \overleftarrow{\partial}_+} \right) \lambda^a \\
\tilde{O}_s^\lambda &= \frac{1}{2} \bar{\lambda}^a (i \overrightarrow{\partial}_+ + i \overleftarrow{\partial}_+)^{s-1} C_{s-1}^{\frac{3}{2}} \left(\frac{\overrightarrow{\partial}_+ - \overleftarrow{\partial}_+}{\overrightarrow{\partial}_+ + \overleftarrow{\partial}_+} \right) \lambda^a \\
M_s &= \frac{1}{2} \partial_+ A^a (i \overrightarrow{\partial}_+ + i \overleftarrow{\partial}_+)^{s-1} P_{s-1}^{(2,1)} \left(\frac{\overrightarrow{\partial}_+ - \overleftarrow{\partial}_+}{\overrightarrow{\partial}_+ + \overleftarrow{\partial}_+} \right) \lambda^a \\
\bar{M}_s &= \frac{1}{2} \lambda^a (i \overrightarrow{\partial}_+ + i \overleftarrow{\partial}_+)^{s-1} P_{s-1}^{(1,2)} \left(\frac{\overrightarrow{\partial}_+ - \overleftarrow{\partial}_+}{\overrightarrow{\partial}_+ + \overleftarrow{\partial}_+} \right) \partial_+ A^a
\end{aligned} \tag{E.1}$$

where $C_l^{\alpha'}(x)$ are Gegenbauer polynomials that are a special case of Jacobi polynomials (appendix B):

$$C_l^{\alpha'}(x) = \frac{\Gamma(l+2\alpha')\Gamma(\alpha'+\frac{1}{2})}{\Gamma(2\alpha')\Gamma(l+\alpha'+\frac{1}{2})} P_l^{(\alpha'-\frac{1}{2}, \alpha'-\frac{1}{2})}(x) \quad (\text{E.2})$$

with the symmetry properties:

$$C_l^{\alpha'}(-x) = (-1)^l C_l^{\alpha'}(x) \quad (\text{E.3})$$

They are the restriction (up to linear combinations [2]) to the components with maximal-spin projection s along the p_+ direction of the twist-2 operators:

$$\begin{aligned} O_s^{A\mathcal{T}=2} &= \text{Tr } F_{(\rho_1}^\mu \overleftrightarrow{D}_{\rho_2} \dots \overleftrightarrow{D}_{\rho_{s-1}} F_{\rho_s)\mu} - \text{traces} \\ \tilde{O}_s^{A\mathcal{T}=2} &= \text{Tr } \tilde{F}_{(\rho_1}^\mu \overleftrightarrow{D}_{\rho_2} \dots \overleftrightarrow{D}_{\rho_{s-1}} F_{\rho_s)\mu} - \text{traces} \\ O_s^{\lambda\mathcal{T}=2} &= \text{Tr } \bar{\chi} \gamma_{(\rho_1} \overleftrightarrow{D}_{\rho_2} \dots \overleftrightarrow{D}_{\rho_{s-1})} \chi - \text{traces} \\ \tilde{O}_s^{\lambda\mathcal{T}=2} &= \text{Tr } \bar{\chi} \gamma_{(\rho_1} \gamma_5 \overleftrightarrow{D}_{\rho_2} \dots \overleftrightarrow{D}_{\rho_{s-1})} \chi - \text{traces} \\ M_{s+\frac{1}{2}}^{\mathcal{T}=2} &= \text{Tr } F_{(\rho_1}^\nu \overleftrightarrow{D}_{\rho_2} \dots \overleftrightarrow{D}_{\rho_{s-1})} \gamma_\nu \chi - \text{traces} \end{aligned} \quad (\text{E.4})$$

where the parentheses stand for symmetrization of all the indices in between and the subtraction of the traces ensures that the contraction of any two indices vanishes. The above operators transform to the leading order of perturbation theory as primary operators with respect to the conformal group [14]. By projecting on the maximal-spin component along the p_+ direction they restrict to [2]:

$$\begin{aligned} O_s^A &= \frac{1}{2} f_{11}^a (i \overrightarrow{\partial}_+ + i \overleftarrow{\partial}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left(\frac{\overrightarrow{\partial}_+ - \overleftarrow{\partial}_+}{\overrightarrow{\partial}_+ + \overleftarrow{\partial}_+} \right) f_{11}^a \\ \tilde{O}_s^A &= \frac{1}{2} f_{11}^a (i \overrightarrow{\partial}_+ + i \overleftarrow{\partial}_+)^{s-2} C_{s-2}^{\frac{5}{2}} \left(\frac{\overrightarrow{\partial}_+ - \overleftarrow{\partial}_+}{\overrightarrow{\partial}_+ + \overleftarrow{\partial}_+} \right) f_{11}^a \\ O_s^\lambda &= \frac{1}{2} \bar{\lambda}^a (i \overrightarrow{\partial}_+ + i \overleftarrow{\partial}_+)^{s-1} C_{s-1}^{\frac{3}{2}} \left(\frac{\overrightarrow{\partial}_+ - \overleftarrow{\partial}_+}{\overrightarrow{\partial}_+ + \overleftarrow{\partial}_+} \right) \lambda^a \\ \tilde{O}_s^\lambda &= \frac{1}{2} \bar{\lambda}^a (i \overrightarrow{\partial}_+ + i \overleftarrow{\partial}_+)^{s-1} C_{s-1}^{\frac{3}{2}} \left(\frac{\overrightarrow{\partial}_+ - \overleftarrow{\partial}_+}{\overrightarrow{\partial}_+ + \overleftarrow{\partial}_+} \right) \lambda^a \\ M_s &= -\frac{1}{2} f_{11}^a (i \overrightarrow{\partial}_+ + i \overleftarrow{\partial}_+)^{s-1} P_{s-1}^{(2,1)} \left(\frac{\overrightarrow{\partial}_+ - \overleftarrow{\partial}_+}{\overrightarrow{\partial}_+ + \overleftarrow{\partial}_+} \right) \bar{\lambda}^a \\ \bar{M}_s &= -\frac{1}{2} \lambda^a (i \overrightarrow{\partial}_+ + i \overleftarrow{\partial}_+)^{s-1} P_{s-1}^{(1,2)} \left(\frac{\overrightarrow{\partial}_+ - \overleftarrow{\partial}_+}{\overrightarrow{\partial}_+ + \overleftarrow{\partial}_+} \right) f_{11}^a \end{aligned} \quad (\text{E.5})$$

that are primaries with respect to the collinear conformal subgroup $SL(2, R)$ [15] and reduce in the light-cone gauge to the operators in eq. (E.1) with $f_{11} = -\partial_+ \bar{A}$. By allowing operator mixing with derivatives of operators of lower spin they may be extended to primary conformal operators to the next-to-leading order in the conformal renormalization scheme [11] that differs from the \overline{MS} scheme by a finite renormalization.

F Nonresonant diagonal renormalization scheme

To make the present paper self-contained we provide the construction order by order in perturbation theory of the nonresonant diagonal scheme [3].

The construction proceeds by induction on $k = 1, 2, \dots$ by demonstrating that, once A_0 and the first $k - 1$ matrix coefficients $A_2, \dots, A_{2(k-1)}$ in eq. (7.13) have been set in the canonical nonresonant form in eq. (7.18) – i.e., A_0 diagonal and $A_2, \dots, A_{2(k-1)} = 0$ – a real-analytic gauge transformation exists that leaves them invariant and sets the k -th coefficient A_{2k} to 0 as well.

The 0 step of the induction consists in putting A_0 in diagonal form – with the eigenvalues in nonincreasing order – by a constant gauge transformation.

At the k -th step we choose the gauge transformation:

$$S_k(g) = 1 + g^{2k} H_{2k} \quad (\text{F.1})$$

with H_{2k} a matrix to be found below. Its inverse reads:

$$S_k^{-1}(g) = (1 + g^{2k} H_{2k})^{-1} = 1 - g^{2k} H_{2k} + \dots \quad (\text{F.2})$$

where the dots represent terms of order higher than g^{2k} . The action of $S_k(g)$ on the connection $A(g)$ furnishes:

$$\begin{aligned} A'(g) &= 2kg^{2k-1}H_{2k}(1 - g^{2k}H_{2k})^{-1} + (1 + g^{2k}H_{2k})A(g)(1 - g^{2k}H_{2k})^{-1} \\ &= 2kg^{2k-1}H_{2k}(1 - g^{2k}H_{2k})^{-1} + (1 + g^{2k}H_{2k})\frac{1}{g}\left(A_0 + \sum_{n=1}^{\infty} A_{2n}g^{2n}\right)(1 - g^{2k}H_{2k})^{-1} \\ &= 2kg^{2k-1}H_{2k}(1 - \dots) + (1 + g^{2k}H_{2k})\frac{1}{g}\left(A_0 + \sum_{n=1}^{\infty} A_{2n}g^{2n}\right)(1 - g^{2k}H_{2k} + \dots) \\ &= 2kg^{2k-1}H_{2k} + \frac{1}{g}\left(A_0 + \sum_{n=1}^k A_{2n}g^{2n}\right) + g^{2k-1}(H_{2k}A_0 - A_0H_{2k}) + \dots \\ &= g^{2k-1}(2kH_{2k} + H_{2k}A_0 - A_0H_{2k}) + A_{2(k-1)}(g) + g^{2k-1}A_{2k} + \dots \end{aligned} \quad (\text{F.3})$$

where we have skipped all the terms that contribute to an order higher than g^{2k-1} , with:

$$A_{2(k-1)}(g) = \frac{1}{g}\left(A_0 + \sum_{n=1}^{k-1} A_{2n}g^{2n}\right) \quad (\text{F.4})$$

that is the part of $A(g)$ that is not affected by the gauge transformation $S_k(g)$, and therefore verifies the hypotheses of the induction – i.e., that $A_2, \dots, A_{2(k-1)}$ vanish.

Thus, by eq. (F.3) the k -th matrix coefficient A_{2k} may be eliminated from the expansion of $A'(g)$ to order g^{2k-1} provided that an H_{2k} exists such that:

$$A_{2k} + (2kH_{2k} + H_{2k}A_0 - A_0H_{2k}) = A_{2k} + (2k - ad_{A_0})H_{2k} = 0 \quad (\text{F.5})$$

with $ad_{A_0}Y = [A_0, Y]$. If the inverse of $ad_{A_0} - 2k$ exists, the unique solution for H_{2k} is:

$$H_{2k} = (ad_{A_0} - 2k)^{-1} A_{2k} \quad (\text{F.6})$$

Hence, to complete the induction, we should demonstrate that, if the eigenvalues of A_0 are nonresonant, $ad_{A_0} - 2k$ is invertible, i.e., its kernel is trivial.

Now $ad_{\Lambda} - 2k$, as a linear operator that acts on matrices, is diagonal, with eigenvalues $\lambda_i - \lambda_j - 2k$ and the matrices E_{ij} , whose only nonvanishing entries are $(E_{ij})_{ij}$, as eigenvectors. The eigenvectors E_{ij} , normalized so that $(E_{ij})_{ij} = 1$, form an orthonormal basis for the matrices. Therefore, E_{ij} belongs to the kernel of $ad_{\Lambda} - 2k$ if and only if $\lambda_i - \lambda_j - 2k = 0$. Consequently, since $\lambda_i - \lambda_j - 2k \neq 0$ for every i, j by assumption, the kernel of $ad_{\Lambda} - 2k$ only contains 0, and the construction is complete.

References

- [1] M. Bochicchio, M. Papinutto and F. Scardino, *UV asymptotics of n-point correlators of twist-2 operators in SU(N) Yang-Mills theory*, *Phys. Rev. D* **108** (2023) 054023, [[hep-th/2208.14382](#)].
- [2] M. Bochicchio, M. Papinutto and F. Scardino, *n-point correlators of twist-2 operators in SU(N) Yang-Mills theory to the lowest perturbative order*, *JHEP* **08** (2021) 142, [[hep-th/2104.13163](#)].
- [3] M. Bochicchio, *On the geometry of operator mixing in massless QCD-like theories*, *Eur. Phys. J. C* **81** (2021) 749, [[hep-th/2103.15527](#)].
- [4] M. Bochicchio, M. Papinutto and F. Scardino, *On the structure of the large-N expansion in SU(N) Yang-Mills theory*, [hep-th/2401.09312](#).
- [5] L. Brink, J. H. Schwarz and J. Scherk, *Supersymmetric Yang-Mills Theories*, *Nucl. Phys. B* **121** (1977) 77–92.
- [6] A. V. Belitsky, S. E. Derkachov, G. P. Korchemsky and A. N. Manashov, *Dilatation operator in (super-)Yang-Mills theories on the light-cone*, *Nucl. Phys. B* **708** (2005) 115–193, [[hep-th/0409120](#)].
- [7] G. 't Hooft, *A Planar Diagram Theory for Strong Interactions*, *Nucl. Phys. B* **72** (1974) 461.
- [8] A. V. Belitsky, S. E. Derkachov, G. P. Korchemsky and A. N. Manashov, *Superconformal operators in N=4 superYang-Mills theory*, *Phys. Rev. D* **70** (2004) 045021, [[hep-th/0311104](#)].
- [9] A. V. Belitsky, D. Mueller and A. Schafer, *Implications of N=1 supersymmetry for QCD conformal operators*, *Phys. Lett. B* **450** (1999) 126–135, [[hep-ph/9811484](#)].
- [10] J. Zinn-Justin, *Quantum field theory and critical phenomena*. Oxford University Press, 2021.
- [11] V. M. Braun, G. P. Korchemsky and D. Mueller, *The uses of conformal symmetry in QCD*, *Prog. Part. Nucl. Phys.* **51** (2003) 311–398, [[hep-ph/0306057](#)].
- [12] G. Szegő, *Orthogonal Polynomials*, vol. 23 of *American Mathematical Society colloquium publications*. American Mathematical Society, 1959.
- [13] V. Kazakov and E. Sobko, *Three-point correlators of twist-2 operators in N=4 SYM at Born approximation*, *JHEP* **06** (2013) 061, [[hep-th/1212.6563](#)].

- [14] N. Beisert, G. Ferretti, R. Heise and K. Zarembo, *One-loop QCD spin chain and its spectrum*, *Nucl. Phys. B* **717** (2005) 137–189, [[hep-th/0412029](#)].
- [15] A. V. Belitsky and D. Mueller, *Broken conformal invariance and spectrum of anomalous dimensions in QCD*, *Nucl. Phys. B* **537** (1999) 397–442, [[hep-ph/9804379](#)].