Relatively functionally countable subsets of products

Anton E. Lipin

Krasovskii Institute of Mathematics and Mechanics, Ural Federal University, Yekaterinburg, Russia

Abstract

A subset A of a topological space X is called *relatively functionally countable* (RFC) in X, if for each continuous function $f: X \to \mathbb{R}$ the set $f[A]$ is countable. We prove that all RFC subsets of a product \prod \bar{n} ∈ω X_n are countable, assuming that spaces X_n are Tychonoff and all RFC subsets of every X_n are countable. In particular, in a metrizable space every RFC subset is countable.

The main tool in the proof is the following result: for every Tychonoff space X and any countable set $Q \subseteq X$ there is a continuous function f: $X^{\omega} \to \mathbb{R}^2$ such that the restriction of f to Q^{ω} is injective.

Keywords: functionally countable subsets, countable products, metrizable spaces, Hilbert cube 2020 MSC: 54A25, 54B10, 54C05, 54C30

1. Introduction

This paper comes from the following

Question 1.1 (A.V. Osipov). Is there an uncountable set $A \subseteq [0,1]^\omega$ such that for every continuous function $f : [0,1]^\omega \to \mathbb{R}$ the set $f[A]$ is countable?

The author knows this problem from private communication with A.V. Osipov. In 2016 Osipov and A. Miller proved that the answer is no under $\mathfrak{b} > \omega_1$ or cov(meager) $>\omega_1$, but they did not publish these results. Apparently, no one ever raised Question [1.1](#page-0-0) in literature, although one can find some discussion at this problem on mathoverflow [\[5\]](#page-6-0), where user fedja and T. Banakh

Email address: tony.lipin@yandex.ru (Anton E. Lipin)

proved the negative answer under \neg CH. In this paper we prove the negative answer in ZFC.

Instead of the term *projectively countable* as in [\[5\]](#page-6-0), let us use the following notion.

Definition 1.2. A subset A of a topological space X is called *relatively* functionally countable (RFC) in X, if for every continuous function $f: X \rightarrow$ R the set $f[A]$ is countable.

Note that being an RFC subset is not the same as being a functionally countable subspace. For instance, if A is an uncountable discrete and X is one-point compactification of A , then A is an RFC subset and is not a functionally countable subspace. However, every functionally countable subspace is an RFC subset. A nice criterion of functional countability for perfectly normal spaces can be found in [\[4\]](#page-6-1).

Let us describe the structure of this paper. The most complicated result here is (rather technical) Theorem [3.7.](#page-4-0) Section [3](#page-2-0) is devoted entirely to its proof. Corollaries of Theorem [3.7](#page-4-0) in terms of functional countability and, in particular, the negative answer to Question [1.1](#page-0-0) can be found in Section [4.](#page-5-0)

2. Preliminaries

We assume the following notation and conventions.

- We use the term *tree* and associated notation as in [\[3,](#page-6-2) Definition III.5.1] or [\[2,](#page-6-3) Definition 9.10] with the following amendments:
	- instead of "immediate successor" we say son;
	- the set of all leaves of a tree T is denoted by $L(T)$;
	- we write $T \preceq P$ to say that T is a subtree of P.
- if a function $f: X \to \mathbb{R}^2$ is bounded, then we denote $||f||$ the uniform norm of f in any norm of the plane \mathbb{R}^2 .
- the restriction of a function $f: X \to Y$ to a set $A \subseteq X$ is denoted by $f|_A$.
- *countable* means not greater than countable.

The following obvious proposition plays a significant role in our argumentation.

Proposition 2.1. Denote $\square = [-1, 1] \times [0, 1]$ and $\bot = \{(x, y) \in [-1, 1] \times$ $[0, 1]: x = 0$ or $y = 0$. Suppose we are given a countable set $Q \subseteq [-1, 1] \times$ (0,1). Then there is a continuous function $\varphi : \Box \to \bot$ such that (whenever $x \in [-1, 1]$ and $a \in Q$):

- (1) $\varphi(x, 0) = (x, 0);$
- $(2) \varphi(x, 1) = (0, 1);$
- (3) $\varphi(a) \neq (0, 0)$.

Also let us note a trivial fact on RFC sets.

Proposition 2.2. If a subset A of a topological space X is RFC, then for every natural n and any continuous $f: X \to \mathbb{R}^n$ the set $f[A]$ is countable.

3. Constructing maps into the plane \mathbb{R}^2

In this section we use functions, whose codomains are sets of some special form in the plane \mathbb{R}^2 . To define this special form we need the following two definitions.

Definition 3.1. We say that a triple (p, I, U) is a sprig, if all the following conditions are satisfied:

- (a) *p* is a point in \mathbb{R}^2 ;
- (b) I is a line segment in \mathbb{R}^2 and p is an end-point for I;
- (c) U is a neighborhood of $I \setminus \{p\}.$

Definition 3.2. We say that a rooted tree (T, \leq) of finite height is an *oak*, if all its elements are sprigs and every $t = (p, I, U) \in T$ satisfies the following conditions:

- (A) if (q, J, V) is a son of t, then $q \in I \setminus \{p\}$, $J \cap I = \{q\}$, $\overline{V} \subseteq U$ and $\overline{V} \cap I = \{q\};$
- (B) if (q_1, J_1, V_1) and (q_2, J_2, V_2) are different sons of t, then $\overline{V_1} \cap \overline{V_2} = \emptyset$.
	- If $(T, <)$ is an oak and $T = \{(p_t, I_t, U_t) : t \in T\}$, then we denote:
	- $Y(T) = \bigcup$ t∈T $I_t;$
- $K(T) = \{p_t : t \in T\}$ (from <u>K</u>nots);
- $C(T) = \bigcup$ $t\in L(T)$ $(I_t \setminus \{p_t\})$ (from <u>C</u>rown);
- $R(T) = Y(T) \setminus C(T)$.

Note that the order \lt of an oak (T, \lt) is determined by the family T, so we lose no information writing T instead of $(T, <)$.

Definition 3.3. Suppose $T \nvert P$ are oaks and X is a space. We say that a function $g: X \to Y(P)$ is a T-evolution of a function $f: X \to Y(T)$, if for all $a \in X$ such that $f(a) \in R(T)$ we have $g(a) = f(a)$.

Definition 3.4. Suppose T is an oak, X is a space and $Q \subseteq X$. We say that a continuous function $f: X \to Y(T)$ is a (T, X, Q) -lifting, if $f[Q] \subseteq C(T)$.

Definition 3.5. Suppose T is an oak, X is a space, $Q \subseteq X$ and W is a cover of the set Q. We say that a (T, X, Q) -lifting f is (T, X, Q, W) -splitting, if for every leaf $(p, I, U) \in L(T)$ there is $W \in \mathcal{W}$ such that $f^{-1}[I] \cap Q \subseteq W$.

Lemma 3.6. Suppose T is an oak, X is a Tychonoff space, a set $Q \subseteq X$ is countable, f is a (T, X, Q) -lifting, W is an open cover of the set Q and $\varepsilon > 0$. Then there is an oak $P \succeq T$ and a (P, X, Q, W) -splitting T-evolution g of the function f such that $||f-g|| < \varepsilon$.

Proof. Fix any indexing $Q = \{a_n : n \in \omega\}$. Denote $f_0 = f$ and $T_0 = T$. Now let us presume that for some $n \in \omega$ we have an oak $T_n \succeq T$ and a continuous function $f_n: X \to Y(T_n)$ with the following properties:

- (A1) the set $T_n \setminus T$ is finite;
- (A2) $f_n[Q] \cap (R(T) \cup K(T_n)) = \emptyset;$
- (A3) for every $(p, I, U) \in T_n \setminus T$ there is $W \in \mathcal{W}$ such that $f_n^{-1}[I] \subseteq W$.

Let us take $t_n = (p_n, I_n, U_n) \in T_n$ such that $f_n(a_n) \in I_n$. If $t_n \notin T$, define $T_{n+1} = T_n$ and $f_{n+1} = f_n$. Otherwise choose any interval $H_n \subseteq I_n$ which contains the point $f_n(a_n)$ and is small enough to satisfy the following two properties:

(H1) the length of H_n is less than $\varepsilon/2^{n+2}$;

(H2) $\overline{H_n} \cap K(T_n) = \emptyset$.

Take any sprig $s_n = (q_n, J_n, V_n)$ such that

- (S1) $q_n \in H_n \setminus f_n[Q];$
- (S2) $T_{n+1} = T_n \cup \{s_n\}$ is an oak;
- (S3) the length of J_n is less than $\varepsilon/2^{n+2}$.

Suppose $a_n \in W_n \in \mathcal{W}$. Denote $O_n = W_n \cap f_n^{-1}[H_n]$; note that it is an open set. Choose any continuous function $h_n : X \to [0,1]$ such that $h_n(a_n) = 1$ and $h_n[X \setminus O_n] = \{0\}.$

Now let us take any continuous function $\varphi_n : \overline{H_n} \times [0,1] \to \overline{H_n} \cup J_n$ with the properties as in Proposition [2.1,](#page-2-1) namely that, whenever $x \in H_n$ and $a \in Q \cap O_n$:

- $(F1) \varphi_n(x,0) = x;$
- (F2) $\varphi_n(x,1) \in J_n \setminus \{q_n\};$
- (F3) $\varphi_n(f_n(a), h_n(a)) \neq q_n$.

Define the function $f_{n+1}: X \to Y(T_{n+1})$ in the following way:

$$
f_{n+1}(a) = \begin{cases} \varphi_n(f_n(a), h_n(a)), & a \in O_n; \\ f_n(a), & a \in X \setminus O_n. \end{cases}
$$

It is easy to see that T_{n+1} and f_{n+1} satisfy the recursion assumption and $||f_{n+1} - f_n|| < \varepsilon/2^{n+1}.$

Thus, the sequence $(f_n)_{n\in\omega}$ uniformly converges to some continuous function $g: X \to \mathbb{R}^2$ such that $||f - g|| < \varepsilon$. The tree $P = \bigcup$ $T_n \succeq T$ is an oak $n \boldsymbol{\bar{\in}} \omega$ (and its height is not greater than height of T plus 1). It is also not difficult to see that g is a (P, X, Q, W) -splitting T-evolution of the function f. \Box

Theorem 3.7. Suppose X is a Tychonoff space and a set $Q \subseteq X$ is countable. Then there is a continuous function $f: X^{\omega} \to \mathbb{R}^2$ such that its restriction to Q^{ω} is injective.

Proof. For every $n \in \mathbb{N}$ choose some open cover \mathcal{W}_n of the set Q^n in the space X^n in such a way that for any different points $a, b \in Q^{\omega}$ there is $n \in \mathbb{N}$ such that no element of \mathcal{W}_n contains both $a|_n$ and $b|_n$.

Take a sprig $t = (p, I, U)$, where $p = \{0, 0\}$, $I = [0, 1] \times \{0\}$ and $U = \mathbb{R}^2$. Denote $T_1 = \{t\}$ and denote $f_1(x) = (1, 0)$ for all $x \in X$.

Now suppose that for some $n \in \mathbb{N}$ we have an oak T_n and a (T_n, X^n, Q^n) lifting $f_n: X^n \to Y(T_n)$.

By Lemma [3.6](#page-3-0) there is an oak $T_{n+1} \succeq T_n$ and a $(T_{n+1}, X^n, Q^n, \mathcal{W}_n)$ splitting T_n -evolution g_n of the function f_n such that $||f_n-g_n|| < 1/2^n$. Define the function $f_{n+1} : X^{n+1} \to Y(T_{n+1})$ in such way that $f_{n+1}(x_1, ..., x_{n+1}) =$ $g_n(x_1,\ldots,x_n)$, so the pair f_{n+1}, T_{n+1} satisfies the recursion assumption.

Now for all $n \in \mathbb{N}$ define the functions $h_n : X^{\omega} \to Y(T_n)$ as follows: $h_n(x_1, x_2, \ldots) = f_n(x_1, \ldots, x_n)$. We have $||h_{n+1} - h_n|| = ||g_n - f_n|| < 1/2^n$, so the sequence $(h_n)_{n\in\mathbb{N}}$ uniformly converges to some continuous function $f: X^{\omega} \to \mathbb{R}^2$. It remains to note that f is as required. □

Of course, Theorem [3.7](#page-4-0) may be extended to products of different spaces.

Corollary 3.8. Suppose we are given Tychonoff spaces X_n for all $n \in \omega$ and countable sets $Q_n \subseteq X_n$. Then there is a continuous function $f : \prod$ $\prod_{n\in\omega}X_n\to$ \mathbb{R}^2 such that its restriction to \prod \tilde{n} ∈ω Q_n is injective.

Proof. We can suppose that all X_n are pairwise disjoint subsets of some space X. Applying Theorem [3.7](#page-4-0) to X and $Q = \bigcup$ $\bigcup_{n\in\omega}Q_n$, we obtain this corollary.

Remark 3.9. The plane \mathbb{R}^2 in Theorem [3.7](#page-4-0) can not be replaced by \mathbb{R} .

Proof. Suppose a function $f : \mathbb{R}^{\omega} \to \mathbb{R}$ is continuous. Let us show that the restriction of f to \mathbb{Q}^{ω} is not injective.

For all $x_1, \ldots, x_n \in \mathbb{Q}$ let us denote $H(x_1, \ldots, x_n) = \{s \in \mathbb{R}^\omega : s|_n =$ (x_1, \ldots, x_n) . Clearly, every set $f[H(x_1, \ldots, x_n)]$ is connected. If some one of sets $f[H(x)]$ is singleton, then $f|_{\mathbb{Q}^{\omega}}$ is not injective. Otherwise, since f is continuous, we can choose rational numbers $a_1 \neq b_1$ such that $|f[H(a_1)] \cap f[H(b_1)]| > 1$. Reasoning similarly, we get rational numbers a_2, a_3, \ldots and b_2, b_3, \ldots such that $|f[H(a_1, \ldots, a_n)] \cap f[H(b_1, \ldots, b_n)]| > 1$ whenever $n \in \mathbb{N}$. It follows that $f(a_1, a_2, \ldots) = f(b_1, b_2, \ldots)$. \Box

4. Relatively functionally countable subsets of products

Theorem 4.1. Suppose we are given Tychonoff spaces X_n for all $n \in \omega$ and all RFC subsets of every X_n are countable. Then all RFC subsets of $\prod X_n$ $n \in \omega$ are countable.

Proof. Suppose a set $A \subseteq \Pi$ \bar{n} ∈ω $X_n = Y$ is RFC. In particular, for every projection $\pi_n : Y \to X_n$, defined as $\pi_n(x_1, x_2, ...) = x_n$, the set $Q_n = \pi_n[A]$ is countable.

By Corollary [3.8](#page-5-1) there is a continuous function $f: Y \to \mathbb{R}^2$ such that its restriction to \prod Q_n (and hence to A) is injective. By Proposition [2.2](#page-2-2) the \bar{n} ∈ω set $f[A]$ is countable, so A is countable too. \Box

In particular, the answer to Question [1.1](#page-0-0) is no.

Corollary 4.2. Every RFC subset of a metrizable space is countable.

Proof. It follows from Theorem [4.1](#page-5-2) and the fact that every metrizable space can be embedded into ω -th power of a metrizable hedgehog [\[1,](#page-6-4) Theorem 4.4.9]. \Box

Clearly, there is no possibility to replace ω by ω_1 in Theorem [4.1,](#page-5-2) since the uncountable functionally countable space ω_1 can be embedded into X^{ω_1} whenever X is Hausdorff and $|X| > 1$.

Question 4.3 ($\neg CH$). Is there a set $A \subseteq [0,1]^\omega$ such that $|A| = \mathfrak{c}$ and for every continuous function $f : [0,1]^\omega \to \mathbb{R}$ we have $|f[A]| < \mathfrak{c}$?

More generally,

Question 4.4 $(\neg CH)$. Is there a metrizable space X and its subset A such that for every continuous $f: X \to \mathbb{R}$ we have $|f[A]| < |A| \leq \mathfrak{c}$?

Acknowledgements. The author is grateful to Alexander V. Osipov for constant attention to this work.

References

- [1] R. Engelking, General Topology, Revised and completed edition, Heldermann Verlag Berlin (1989).
- [2] T. Jech, Set theory: The third millenium edition (3rd ed.). Berlin: Springer-Verlag (2006).
- [3] Kenneth Kunen, Set Theory, Studies in Logic: Mathematical Logic and Foundations, Vol. 34, College Publications, London, 2011, viii $+401$ pp.
- [4] V.V. Tkachuk, A nice subclass of functionally countable spaces, Comment.Math.Univ.Carolin, 59,3 (2018) 399-409.
- [5] https://mathoverflow.net/questions/330255/on-projectively-countablesets-in-the-hilbert-cube