

Rapid hydrofracture of icy moon shells: insights from glaciology

Robert Law^{1,2}

¹University of Bergen, Bergen, Norway

²Bjerknes Centre for Climate Research, Bergen, Norway

Key Points:

- Rapid hydrofracture is possible for realistic icy-moon stress configurations given a supply of meltwater and an initial fracture.
- This mechanism would allow surface-bed material transport with implications for habitability and may explain chaos terrain formation.

arXiv:2403.09888v2 [astro-ph.EP] 20 Mar 2024

Corresponding author: Robert Law, robert.law@uib.no

Abstract

Europa’s surface features many regions of complex topography termed ‘chaos terrains’. One set of hypotheses for chaos terrain formation requires upward migration of liquid water from perched water bodies within the icy shell formed by convection and tidal heating. However, consideration of the behaviour of terrestrial ice sheets suggests the upwards movement of water from perched water bodies is uncommon. Instead, rapid downwards hydrofracture — unbounded given a sufficient supply of water — can occur in relatively low tensile stress states given a sufficiently deep initial fracture due to the negative relative buoyancy of water. I suggest that downwards, not upwards, fracture may be more reasonable for perched water bodies but show that full hydrofracture is unlikely if the perched water body is located beneath a mechanically strong icy lid. However, full hydrofracture is possible in the event of lid collapse over a perched water body and likely in the event of a meteor impact that generates sufficient meltwater and a tensile shock. This provides a possible mechanism for the transfer of biologically important nutrients to the subsurface ocean and the formation of chaos terrains.

Plain Language Summary

Jupiter’s moon Europa has a subsurface ocean surrounded by an icy shell that features numerous ‘chaos terrains’ of jumbled topography. Most theories for the formation of chaos terrains focus on the upwards motion of water pooled within the icy shell, but this contrasts with the rapid downwards motion of water in ice sheets on Earth. This paper considers the rapid downwards motion of water on Europa and other icy moons through water-driven fracture (hydrofracture), finding that it is a possibility if a large water body is not covered by a strong lid. This has implications for delivery of biologically important material from the moon’s surface to its ocean, and may help explain the formation of chaos terrains.

1 Introduction

The presence of liquid water in icy moons beyond the asteroid belt was hypothesised by Lewis (1971) and supported by surface images and magnetic field readings from the Voyager, Galileo, Cassini satellite missions (Carr et al., 1998; Khurana et al., 1998; Porco et al., 2006; Lorenz et al., 2008; Hendrix et al., 2019). The presence of liquid water in these moons offers the tantalising possibility of extraterrestrial biology, which focussed research into whether the other basic requirements of chemical energy to drive metabolism and organic matter may also be met (e.g. Kargel et al., 2000; Sephton et al., 2018), with Jupiter’s moon Europa receiving particular attention (e.g. Chyba & Phillips, 2001; NASA, 2017).

One pathway for the possibility of life on Europa requires the exchange of oxidants and organics produced on the moon’s surface with the subsurface ocean (e.g. Delitsky & Lane, 1998; Carlson et al., 1999; Chyba, 2000; Chyba & Phillips, 2001), placing an emphasis on ‘chaos terrains’ as possible locations for major melt-through events of Europa’s icy shell (Carr et al., 1998; Soderlund et al., 2013). However, direct contact between a thin layer of brittle surface ice and an underlying ocean requires an unrealistic amount of thermal energy to locally melt Europa’s icy shell (Collins et al., 2000; Goodman et al., 2004), the thickness of which is poorly constrained but may exceed 30 km (Sotin et al., 2002; Billings & Kattenhorn, 2005; Fagents et al., 2022). Chaos terrain explanations therefore frequently invoke the presence of a near surface water body emplaced through convective upwelling (e.g. Sotin et al., 2002; Figueredo et al., 2002; Pappalardo & Barr, 2004; Schmidt et al., 2011) or sills (Michaut & Manga, 2014) that then reaches the surface through brittle fracturing of the surface (e.g. Crawford & Stevenson, 1988; Collins et al., 2000; Manga & Wang, 2007; Luzzi et al., 2021). Perched water bodies are required in these chaos terrain formation scenarios because the negative buoyancy of water with

respect to ice means fractures propagating upwards from the subsurface ocean would extend at most 90% of the way through the shell before reaching hydrostatic equilibrium (Crawford & Stevenson, 1988).

However, consideration of water bodies perched over ice on Earth strongly suggests that the negative relative buoyancy of water presents a major problem to their stability. Supraglacial lakes, which can exceed 0.125 km² in surface area, pepper the margins of the Greenland Ice Sheet and are readily observed to drain rapidly through hydrofracture (Das et al., 2008; Doyle et al., 2013; Fitzpatrick et al., 2014; Williamson, Banwell, et al., 2018; Chudley et al., 2019). Theoretical considerations show that such fractures will reach the ice-sheet bed given a sufficient supply of water (Weertman, 1973; van der Veen, 2007; Krawczynski et al., 2009). In these settings an initial fracture is required for drainage through hydrofracture to occur. This may come from tensile shock from nearby hydrofracture events (Christoffersen et al., 2018), a transient strain rate increase (e.g. Poinar & C. Andrews, 2021), or reactivation of an existing weakness (Chudley et al., 2019) though it is not clear that there is a single statistically supported explanatory mechanism (Williamson, Willis, et al., 2018). Notably, supraglacial lakes form in depressions in the ice surface which are generally dominated by compressive stresses (Doyle et al., 2013; Stevens et al., 2015; Chudley et al., 2019).

Slow density-driven downward transport of temperate ice (ice containing a percentage liquid water) through an icy moon shell has been considered as a mechanism that would prevent the formation of perched water bodies (Karlstrom et al., 2014; Carnahan et al., 2022; Hesse et al., 2022; Kalousová et al., 2024), but the possible implications of rapid hydrofracture have not thus far been explored. In this paper I cover two perched water body settings: one formed through convective and tidal heating (Fig. 1a) and a second formed by a meteorite impact (Fig. 1d). Each of these water bodies may then be separated from the ice surface by a mechanically strong ice layer (Figs. 1b, e) or a weak melange of ice, water, and slush (Figs. 1c, f). Calculating the stresses governing hydrofracture in a simplified two-dimensional plane indicates that full-thickness hydrofracture is possible when a mechanically strong lid is not present given a sufficient supply of water. I suggest hydrofracture may be more likely in the event of a meteorite strike that also provides an initial fracture site. Such a mechanism may explain chaos terrain formation if violent drainage prompts surface collapse, and provides a clear pathway for the transport of organics and oxidants into the subsurface ocean. I focus on Europa, but the results are applicable to any ice shell setting.

1.1 Materials and methods

van der Veen (1998) calculates terrestrial crevasse propagation under linear elastic fracture mechanics which treats a material as containing small defects which affect its load-bearing capacity. If high stresses concentrate near a defect the defect may propagate and ultimately fracture (Broek, 1982). In a purely elastic material, an unstable crack will propagate when the energy absorbed by a small expansion of the crack is exceeded by the energy released by a small increase in crack size (Griffiths, 1921). This condition is met when the net stress intensity factor, $K_{(\text{net})}$, exceeds the fracture toughness of the material. Ice is not a purely elastic material (instead it has a non-linear viscoelastic rheology), but linear elastic fracture mechanics has proven effective in application to terrestrial ice sheets (e.g. van der Veen, 1998; Chudley et al., 2019). The outer shells of icy moons are likely much colder and therefore much stiffer than terrestrial ice (Hussmann et al., 2002), reducing the relative importance of the viscous component of ice rheology (Goldsby & Kohlstedt, 2001). Melt-formed conduits are also ubiquitous in temperate ice alongside fractures (Fountain & Walder, 1998; Gulley et al., 2009). Below I use the essential equations from Weertman (1973), van der Veen (1998), van der Veen (2007), and Krawczynski et al. (2009) and refer the reader to the original papers for more detailed context.




Figure 1. Schematic of icy-shell hydrofracture scenarios covered in the text. (a) A perched water body as hypothesised to form through thermal convection or a sill. (b) Incomplete hydrofracture in a perched water body with a rigid lid. (c) Caldera collapse and full hydrofracture. (d) Incoming meteoroid. (e) Incomplete hydrofracture following meteorite impact. (f) Full hydrofracture of meteorite-generated meltwater when the surface is not sealed following meteorite impact.




Figure 2. Schematic of fracture calculations. See text for symbol definitions. The line at the top of the fracture is the fracture surface. The dashed line indicates the top of the perched water body.

I take a single water-filled fracture unaffected by surrounding fractures (Fig. 2) and consider only Mode I fracturing. The stress intensity factor is then determined by three processes — far-field tensile or compressive stress, hydrostatic stress, and water pressure — which can be superimposed without complication unlike if Mode II and Mode III openings were also considered (van der Veen, 1998). The far-field tensile stress in $\text{Pa}^{1/2}$ is given as

$$K^{(1)} = F(\lambda)R_{xx}\sqrt{\pi d} \quad (1)$$

where $\lambda = (d + h + a)/H$, d (m) is the fracture depth, h is the height above the fracture surface (or water body surface if present) to the actual surface, a is the depth of the water body, and H is the ice shell thickness (Fig. 2). R_{xx} (Pa) is the resistive stress defined as

$$R_{xx} = \sigma_{xx} - L \quad (2)$$

where σ_{xx} (Pa) is the full stress and $L = -\rho_i g(H + h + a - z)$ (m) is the lithostatic stress where ρ_i (kg m^{-3}) is the density of ice, g (m s^{-2}) is acceleration due to gravity and z is 0 at the base of the ice shell increasing upwards. The $F(\lambda)$ function is

$$F(\lambda) = 1.12 - 0.23\lambda + 10.55\lambda^2 - 21.72\lambda^3 + 30.39\lambda^4 \quad (3)$$

obtained by polynomial curve fitting to numerically computed stress intensity factors (Tada et al., 1973; Broek, 1982; van der Veen, 1998).

The net stress intensity factor from the overburden pressure is calculated by integrating the contributions from forces acting at different depths as

$$K^{(2)} = -\frac{2g}{\sqrt{\pi d}} \int_0^d (\rho_i b + \rho_w a + \rho_i h) G(\gamma, \lambda) db \quad (4)$$

where the $\rho_w a$ and $\rho_i h$ account for the weight of overlying water and overlying mechanically strong lid respectively and w is the density of water, $\gamma = b/d$ and b is the depth from the fracture surface to the point in the fracture under consideration. I make the assumption that the ice lid is mechanically isolated from the water body and influences the hydrostatic stress (Eq. 4) but not the water pressure (Eq. 6). This is a simplification but serves to illustrate the maximum fracture-limiting influence a lid overlying a perched water body may have. The function $G(\gamma, \lambda)$ is

$$G(\gamma, \lambda) = \frac{3.52(1-\gamma)}{(1-\lambda)^{3/2}} - \frac{4.35-5.28\gamma}{(1-\lambda)^{1/2}} + \left[\frac{1.30-0.30\gamma^{3/2}}{(1-\gamma^2)} + 0.83-1.76\gamma \right] \times [1-(1-\gamma)\lambda] \quad (5)$$

again based on numerical curve fitting of numerically derived solutions (Tada et al., 1973; van der Veen, 1998).

Last, the stress intensity factor from the water pressure within the fracture following van der Veen (1998) is

$$K^{(3)} = \frac{2g}{\sqrt{\pi d}} \int_0^d (\rho_w b + \rho_w a) G(\gamma, \lambda) db. \quad (6)$$

Evaluating the two integrals numerically gives the net stress intensity factor as

$$K^{(\text{net})} = K^1 + K^2 + K^3. \quad (7)$$

The volume of water required for a given crack propagation is calculated after Krawczynski et al. (2009) and Weertman (1973). The fracture width, w , in the x direction at depth b is

$$w(b) = \frac{4\alpha\sigma}{\mu}\omega + \frac{4\alpha\rho_i g d}{\pi\mu}\omega - \frac{4\alpha\rho_w g}{\pi\mu}\omega\psi - \frac{2\alpha\rho_i g b^2}{\pi\mu} \ln\left(\frac{d+\omega}{d-\omega}\right) + \frac{2\alpha\rho_w g(b^2-a^2)}{\pi\mu} \ln\left|\frac{\psi+\omega}{\psi-\omega}\right| - \frac{4\alpha\rho_w g b a}{\pi\mu} \ln\left|\frac{a\omega+b\psi}{a\omega-b\psi}\right| + \frac{4\alpha\rho_w g a^2}{\pi\mu} \ln\left|\frac{\omega+\psi}{\omega-\psi}\right| \quad (8)$$

where μ (Pa) is the shear modulus for ice, $\alpha = 1 - \nu$ where ν is Poisson's ratio, $\omega = \sqrt{d^2 - b^2}$ and $\psi = \sqrt{d^2 - a^2}$. Here, stress, σ , is expressed as

$$\sigma = R_{xx} - \frac{2\rho_i g d}{\pi} - \rho_w g a + \frac{\rho_w g a}{\pi} \arcsin\left(\frac{a}{d}\right) + \frac{2\rho_w g}{\pi} \psi. \quad (9)$$

This is contrary to stress calculations for Eqs. 1-7 given the difference in approach between van der Veen (1998) and Krawczynski et al. (2009), but given the effectiveness of each to its respective setting I do not reconcile the differences here. The fracture volume required for full thickness hydrofracture is then

$$V_f = l \int_0^H w(b) db \quad (10)$$

which is evaluated numerically where l is the fracture length (along the y axis, into the page, and not displayed in Fig. 2). Although the water body is approximated as plane

Table 1. Values used in calculations.

Symbol	Value
g	1.31 m s ⁻²
ν	0.3
ρ_i	917 kg m ⁻³
ρ_w	1000 kg m ⁻³
μ	0.5, 1.5 GPa

in Eqs. 1-7 to calculate the water body volume I consider a spherical perched water body of radius r , the fracture length is then taken as $l = 43r$. On Earth the fracture may at first span the diameter of the water body (Chudley et al., 2019), but a more spherical geometry would result in a shorter fracture length. On Earth once a fracture is fully established the fracture usually localises to a point feature or moulin (Chudley et al., 2019). The radius of the required sphere is

$$r = \sqrt{\frac{1}{\pi} \int_0^H w(b) db}. \quad (11)$$

Values used in calculations are provided in Table 1.

2 Results

The calculations demonstrate that under certain circumstances hydrofracture from a perched water body to the subsurface ocean of Europa is possible. Fig. 3 shows the stress intensity fracture that would be experienced at the tip of the fracture against the fracture depth. If the stress intensity at the fracture tip exceeds the critical fracture toughness of ice the fracture will continue to propagate. If a mechanically strong lid is absent the rate of change of $K^{(3)}$ with depth exceeds that of $K^{(2)}$ for all depths considered (Fig. S1) meaning that once the stress intensity factor at the fracture tip exceeds the critical fracture toughness the fracture will continue to propagate unbounded, limited only by the water supplying the fracture. An initial fracture depth of only 4.1 m under a tensile stress of 0.1 MPa is required for unstable hydrofracture if the perched water body is located on the surface. If an ice lid greater than around 10 m (dependent slightly on the value of a) is present that contributes to lithostatic but not hydrostatic pressure then the rate of change of $K^{(2)}$ with depth exceeds that of $K^{(3)}$ meaning fracture propagation is not favourable. Unreasonably deep initial fractures are required in the case of a surface water body and 0 MPa tensile stress but, once a fracture is established neutral far-field stress at depth would not lead to closure (Fig. S2).

The volume of water required to facilitate a 30 km fracture under a shear modulus of 1.5 GPa is 38.8 km³ (Fig. 4), significantly below the 20,000-60,000 km³ melt lens estimate of Schmidt et al. (2011) and an order of magnitude below melt volumes modelled to form by Kalousová et al. (2024) in the event of moderately sized meteorite impacts. The required water volume is sensitive to the shear modulus and the estimates here follow lower shear modulus values.

van der Veen (1998) shows that the key control on the downward propagation of the fracture is the filling rate and that penetration velocity equals the rate of fracture filling to within a few percent. This is because $K^{(2)}$ and $K^{(3)}$ become the dominant terms in Eq. 7 giving




Figure 3. Stress intensity factor as function of fracture depth for varying ice lid thickness, perched water body depth, and tensile stress. The blue dashed line is a reasonable upper limit for the fracture toughness of ice ($0.4 \text{ MPa}^{1/2}$) from van der Veen (1998). (a) and (b) $h = 0$ m. (c) and (d) $a = 500$ m. (a) and (c) tensile stress is 0.1 MPa. (b) and (d) tensile stress is 0.2 MPa. $H = 30$ km for all panels.




Figure 4. Water volume required for a given fracture depth. Dashed line is for a shear modulus of 0.5 GPa and solid line for 1.5 GPa. Note that the upper bound shear modulus of 3.9 GPa, as used in van der Veen (1998), is not included here. Black indicates required sphere radius, blue indicates shear volume.

$$d = \left(\frac{\rho_w}{\rho_i} \right)^{\frac{2}{3}} Qt \quad (12)$$

where Q is the vertical fill rate (m s^{-1}) and t (s) is the time period in consideration to a good approximation (van der Veen, 1998). Given a large perched water body positioned over a hydrofracture could therefore be a very rapid phenomena. Supraglacial lakes in Greenland take on the order of 2-3 hours to drain, but the water bodies have an aspect ratio of 1:100 making Q limited by lateral transport from the outer reaches of the lake (Chudley et al., 2019; Das et al., 2008). As a first approximation, even if drainage took 24 hours through ice at 100 K, only 0.34 m of side wall freezing would occur (Text S1), negligible in the context of a surface opening of ~ 200 m following Eq. 8.

3 Discussion

Figs. 3 and 4 demonstrate that full hydrofracture of Europa is possible given moderate tensile stresses, sufficient water to propagate the fracture, and the absence of a mechanically strong lid. Fractures are common across Europa's surface (e.g. Dombard et al., 2013) but abundant surface water akin to the Greenland Ice Sheet is practically impossible given the very low temperature and pressure of the ice surface (Ashkenazy, 2019). Nonetheless, in this simple if unrealistic situation, hydrofracture can occur at tensile stresses much lower than the ~ 20 MPa proposed by Crawford & Stevenson (1988) and Rudolph & Manga (2009) for full 'dry' shell fracture. Stresses are probably slightly compressive on average (of order 0.1-0.4 MPa) in Europa's shell (Greenberg et al., 1998; Pappalardo et al., 1998) meaning hydrofracture of a perched water body would only occur under spatially or temporally limited stress conditions (as is the case for the Greenland Ice Sheet). Structural analysis of surface faults indicate tensile fracturing does occur at the surface of Europa (Kattenhorn & Marshall, 2006) and analysis of the stresses induced by diurnal tides suggests that tensile stresses up to ~ 0.15 MPa are possible (Helfenstein & Parmentier, 1983; Greenberg et al., 1998; Harada & Kurita, 2006; Hurford et al., 2007).

However if, as widely hypothesised (Fagents et al., 2022), a water body is entombed then the additional lithostatic stress makes rapid hydrofracture (or any fracture at all) unlikely. Though it is worth noting that mechanisms hypothesised to direct water upwards such as the pressurisation through freezing of perched water bodies (Manga & Wang, 2007; Steinbrügge et al., 2020; Lesage et al., 2022; Quick et al., 2023), ought also to work downwards and will be aided in that case by density-difference driven fracture. But, if the lid is weakened or fractured then its weight will contribute to the hydrostatic pressure and not the lithostatic pressure and hydrofracture again becomes possible. Two such scenarios may occur.

First, Schmidt et al. (2011), Walker & Schmidt (2015), and Luzzi et al. (2021) among others propose caldera collapse of perched water bodies as a formation mechanism for chaos terrains. In several of these scenarios the blocks of ice are buoyantly supported by brine, water, and slush (Fig. 1c). This effectively sets h at zero, while increasing a mean-ing hydrofracture becomes a realistic possibility. It is unclear how long this lid state may persist before refreezing occurs (Schmidt et al., 2011; Fagents et al., 2022), but while it does only a small and temporary tensile stress and a very small initial fracture would be required to initiate rapid hydrofracture.

Second, literature on the stress impact of meteor impacts on icy moons is limited, but it appears reasonable that a meteorite impact may provide the three crucial ingredients for hydrofracture — an initial fracture site, a tensile shock, and an abundant supply of impact-generated melt water (Kalousová et al., 2024) while leaving an at least transiently mechanically weak area above the water body (Fig. 1f). A meteorite may also impact sufficiently close to an existing perched water body to trigger its hydrofracture. In fact, a meteorite that impacted the Greenland Ice Sheet, likely after its initial inception, provides a terrestrial analogue for what one may expect on Europa (Kjær et al., 2018) and given the 300 m deep crater in solid bedrock, it seems unlikely that such an impact would not have led to sizable fractures forming within the ice. A meteorite trigger for chaos terrain formation and potentially hydrofracture is also supported by Cox et al. (2010) who find no small chaos areas and no craters ≥ 50 km diameter, suggesting a possible critical impact intensity required to generate sufficient meltwater and shock. If density driven sinking of melt (Carnahan et al., 2022; Kalousová et al., 2024) actually prevents the formation of a perched water body, then a meteorite impact becomes the only remaining possible water source.

If a previously entombed water body drains, the drainage event will remove hydrostatic support and lead to extensive collapse of the overlying terrain. If the fracture remains open the water level would then be expected to stabilise at $\sim 90\%$ of the ice shell thickness (Michaut & Manga, 2014), though viscous deformation and freezing may be expected to close the fracture over a short period (years or less) after active drainage has ceased (Andrews et al., 2022). It is worth noting that in Greenland the large ice blocks generated in initial hydrofracture, with more expected if an icy lid is present (Russell, 1993), may become wedged at the fracture surface (Fig. S3) so while water may leave the site quickly, solid ice blocks should remain. This supports a more neocatastrophist view of chaos terrain formation than gradual caldera collapse (Luzzi et al., 2021).

Last, it is challenging to be definitive about the thermal state of Europa’s icy shell, but several studies suggest a warm and uniform convecting interior, within around 10 K of the pressure melting point (e.g. McKinnon, 1999; Hussmann et al., 2002; Showman & Han, 2004; Mitri & Showman, 2005). These temperatures are warm relative to an average surface temperature of around 100 K (Ashkenazy, 2019), but their correspondingly lower relative viscosity should not present a problem to elastic fracture given full-thickness hydrofracture through ice of this temperature is common in the Greenland Ice Sheet (e.g. Doyle et al., 2018; Law et al., 2021). More intriguing is the possible influence of temperate ice on water transport within Europa’s shell. Carnahan et al. (2022) and Kalousová et al. (2024) do not account for the formation of conduits within the icy shell, yet there

are no temperate terrestrial glaciers where there is not some degree of englacial drainage system formation (e.g. Hooke, 1989). A body of temperate ice within an icy shell would lack a supply of surface meltwater, but tidal or convective heating may still produce water internally (e.g. Gaidos & Nimmo, 2000). Given the melt chambers hypothesised by Carnahan et al. (2022) reach ~ 5 km in height and feature melt fractions $\sim 5\%$, it seems reasonable that transport through a drainage system may lead to pooling of water at the transition between temperate and cold ice if fracture does not occur. Further integrating understanding of englacial hydrology with icy moons can advance understanding of these mechanisms.

4 Conclusions

This paper demonstrates that in the presence of a mechanically weak lid only relatively small tensile stresses, and achievable water volumes, are required for the full hydrofracture of Europa. This has direct application to other icy moons with different gravities and stress regimes, which may also have conditions facilitating full-thickness hydrofracture. It also provides a clear mechanism for the delivery to the subsurface ocean of biologically important materials created on the surface of Europa and delivered by incoming meteorites and may provide an explanation for the formation of chaos terrains. Future research directions include numerical modelling of the spatio-temporal variation of stress configurations around varied perched water body geometries and as a result of meteorite impacts, and improving understanding of the geomorphological implications of rapid hydrofracture. Hopefully, this paper opens further interdisciplinary pathways (Garcia-Lopez & Cid, 2017; Rossi et al., 2023) between glaciology and the study of ice moons.

Open Research Section

Data supporting the conclusions is derived from the calculations outlined in the text. No further data that is not already publicly available was used. Plotting scripts can be uploaded to a University of Bergen repository before publication, but this requires a few weeks notice on their side.

Acknowledgments

Thanks to Craig Walton and Claire Guimond for introducing me to this field and Andreas Born for letting me take a bit of time away from the day job. I acknowledge funding from Norges forskningsråd. There are no conflicts of interest with respect to the results of this manuscript.

References

- Andrews, L. C., Poinar, K., & Trunz, C. (2022, 6). Controls on Greenland moulin geometry and evolution from the Moulin Shape model. *Cryosphere*, 16(6), 2421–2448. doi: 10.5194/TC-16-2421-2022
- Ashkenazy, Y. (2019, 6). The surface temperature of Europa. *Helvion*, 5(6), 1908. Retrieved from <http://www.cell.com/article/S2405844018357724/fulltext><http://www.cell.com/article/S2405844018357724/abstract>[https://www.cell.com/helivon/abstract/S2405-8440\(18\)35772-4](https://www.cell.com/helivon/abstract/S2405-8440(18)35772-4) doi: 10.1016/j.helivon.2019.e01908
- Billings, S. E., & Kattenhorn, S. A. (2005, 10). The great thickness debate: Ice shell thickness models for Europa and comparisons with estimates based on flexure at ridges. *Icarus*, 177(2), 397–412. doi: 10.1016/J.ICARUS.2005.03.013
- Broek, D. (1982). Elementary engineering fracture mechanics. *Elementary engineering fracture mechanics*. doi: 10.1007/978-94-009-4333-9
- Carlson, R. W., Anderson, M. S., Johnson, R. E., Smythe, W. D., Hendrix,

- A. R., Barth, C. A., ... Matson, D. L. (1999, 3). Hydrogen Peroxide on the Surface of Europa. *Science*, 283(5410), 2062–2064. Retrieved from <https://www.science.org/doi/10.1126/science.283.5410.2062> doi: 10.1126/SCIENCE.283.5410.2062
- Carnahan, E., Vance, S. D., Cox, R., & Hesse, M. A. (2022, 12). Surface-To-Ocean Exchange by the Sinking of Impact Generated Melt Chambers on Europa. *Geophysical Research Letters*, 49(24), e2022GL100287. Retrieved from <https://onlinelibrary.wiley.com/doi/full/10.1029/2022GL100287><https://onlinelibrary.wiley.com/doi/abs/10.1029/2022GL100287><https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2022GL100287> doi: 10.1029/2022GL100287
- Carr, M. H., Belton, M. J., Chapman, C. R., Davies, M. E., Geissler, P., Greenberg, R., ... Veverka, J. (1998, 1). Evidence for a subsurface ocean on Europa. *Nature*, 391(6665), 363–365. Retrieved from <https://www.nature.com/articles/34857> doi: 10.1038/34857
- Christoffersen, P., Bougamont, M., Hubbard, A., Doyle, S. H., Grigsby, S., & Pettersson, R. (2018, 12). Cascading lake drainage on the Greenland Ice Sheet triggered by tensile shock and fracture. *Nature Communications*, 9(1), 1064. Retrieved from <http://www.nature.com/articles/s41467-018-03420-8> doi: 10.1038/s41467-018-03420-8
- Chudley, Christoffersen, P., Doyle, S., Bougamont, M., Schoonman, C., Hubbard, B., & James, M. (2019, 12). Supraglacial lake drainage at a fast-flowing Greenlandic outlet glacier. *Proceedings of the National Academy of Sciences of the United States of America*, 116(51), 25468–25477. doi: 10.1073/pnas.1913685116
- Chyba, C. F. (2000, 1). Energy for microbial life on Europa. *Nature*, 403(6768), 381–382. Retrieved from <https://www.nature.com/articles/35000281> doi: 10.1038/35000281
- Chyba, C. F., & Phillips, C. B. (2001, 1). Possible ecosystems and the search for life on Europa. *Proceedings of the National Academy of Sciences of the United States of America*, 98(3), 801–804. Retrieved from <https://www.pnas.org/doi/abs/10.1073/pnas.98.3.801> doi: 10.1073/PNAS.98.3.801/ASSET/326781B2-7546-4ECB-9B42-5FB3403EC751/ASSETS/GRAPHIC/PQ0415098001.JPEG
- Collins, G. C., Head, J. W., Pappalardo, R. T., & Spaun, N. A. (2000, 1). Evaluation of models for the formation of chaotic terrain on Europa. *Journal of Geophysical Research: Planets*, 105(E1), 1709–1716. Retrieved from <https://onlinelibrary.wiley.com/doi/full/10.1029/1999JE001143><https://onlinelibrary.wiley.com/doi/abs/10.1029/1999JE001143><https://agupubs.onlinelibrary.wiley.com/doi/10.1029/1999JE001143> doi: 10.1029/1999JE001143
- Cox, R., Ong, L. C., Arakawa, M., & Scheider, K. C. (2010, 12). Impact penetration of Europa's ice crust as a mechanism for formation of chaotic terrain. *Meteoritics & Planetary Science*, 43(12), 2027–2048. Retrieved from <https://onlinelibrary.wiley.com/doi/full/10.1111/j.1945-5100.2008.tb00659.x><https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1945-5100.2008.tb00659.x> doi: 10.1111/J.1945-5100.2008.TB00659.X
- Crawford, G. D., & Stevenson, D. J. (1988, 1). Gas-driven water volcanism and the resurfacing of Europa. *Icarus*, 73(1), 66–79. doi: 10.1016/0019-1035(88)90085-1
- Das, S. B., Joughin, I., Behn, M. D., Howat, I. M., King, M. A., Lizarralde, D., & Bhatia, M. P. (2008, 5). Fracture propagation to the base of the Greenland Ice Sheet during supraglacial lake drainage. *Science (New York, N.Y.)*, 320(5877), 778–81. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/18420900> doi: 10.1126/science.1153360
- Delitsky, M. L., & Lane, A. L. (1998, 12). Ice chemistry on the Galilean satellites.

- Journal of Geophysical Research: Planets*, 103(E13), 31391–31403. Retrieved from <https://onlinelibrary.wiley.com/doi/full/10.1029/1998JE900020><https://onlinelibrary.wiley.com/doi/abs/10.1029/1998JE900020><https://agupubs.onlinelibrary.wiley.com/doi/10.1029/1998JE900020> doi: 10.1029/1998JE900020
- Dombard, A. J., Patterson, G. W., Lederer, A. P., & Prockter, L. M. (2013, 3). Flanking fractures and the formation of double ridges on Europa. *Icarus*, 223(1), 74–81. doi: 10.1016/J.ICARUS.2012.11.021
- Doyle, S. H., Hubbard, A. L., Dow, C. F., Jones, G. A., Fitzpatrick, A., Gusmeroli, A., ... Box, J. E. (2013, 1). Ice tectonic deformation during the rapid in situ drainage of a supraglacial lake on the Greenland Ice Sheet. *The Cryosphere*, 7(1), 129–140. Retrieved from <https://www.the-cryosphere.net/7/129/2013/> doi: 10.5194/tc-7-129-2013
- Doyle, S. H., Hubbard, B., Christoffersen, P., Young, T. J., Hofstede, C., Bougamont, M., ... Hubbard, A. (2018, 2). Physical Conditions of Fast Glacier Flow: 1. Measurements From Boreholes Drilled to the Bed of Store Glacier, West Greenland. *Journal of Geophysical Research: Earth Surface*, 123(2), 324–348. Retrieved from <http://doi.wiley.com/10.1002/2017JF004529> doi: 10.1002/2017JF004529
- Fagents, S. A., Lopes, R. M., Quick, L. C., & Gregg, T. K. (2022, 1). Cryovolcanism. *Planetary Volcanism across the Solar System*, 161–234. doi: 10.1016/B978-0-12-813987-5.00005-5
- Figueredo, P. H., Chuang, F. C., Rathbun, J., Kirk, R. L., & Greeley, R. (2002, 5). Geology and origin of Europa’s “Mitten” feature (Murias Chaos). *Journal of Geophysical Research: Planets*, 107(E5), 2–1. Retrieved from <https://onlinelibrary.wiley.com/doi/full/10.1029/2001JE001591><https://onlinelibrary.wiley.com/doi/abs/10.1029/2001JE001591><https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2001JE001591> doi: 10.1029/2001JE001591
- Fitzpatrick, A. A. W., Hubbard, A. L., Box, J. E., Quincey, D. J., van As, D., Mikkelsen, A. P. B., ... Jones, G. A. (2014, 1). A decade (2002–2012) of supraglacial lake volume estimates across Russell Glacier, West Greenland. *The Cryosphere*, 8(1), 107–121. Retrieved from <http://www.the-cryosphere.net/8/107/2014/> doi: 10.5194/tc-8-107-2014
- Fountain, A. G., & Walder, J. S. (1998, 8). Water flow through temperate glaciers. *Reviews of Geophysics*, 36(3), 299–328. doi: 10.1029/97RG03579
- Gaidos, E. J., & Nimmo, F. (2000, 6). Tectonics and water on Europa. *Nature* 2000 405:6787, 405(6787), 637–637. Retrieved from <https://www.nature.com/articles/35015170> doi: 10.1038/35015170
- Garcia-Lopez, E., & Cid, C. (2017, 7). Glaciers and ice sheets as analog environments of potentially habitable icy worlds. *Frontiers in Microbiology*, 8(JUL), 267284. Retrieved from www.frontiersin.org doi: 10.3389/FMICB.2017.01407/BIBTEX
- Goldsby, D. L., & Kohlstedt, D. L. (2001, 6). Superplastic deformation of ice: Experimental observations. *Journal of Geophysical Research: Solid Earth*, 106(B6), 11017–11030. doi: 10.1029/2000jb900336
- Goodman, J. C., Collins, G. C., Marshall, J., & Pierrehumbert, R. T. (2004, 3). Hydrothermal plume dynamics on Europa: Implications for chaos formation. *Journal of Geophysical Research: Planets*, 109(E3), 3008. Retrieved from <https://onlinelibrary.wiley.com/doi/full/10.1029/2003JE002073><https://onlinelibrary.wiley.com/doi/abs/10.1029/2003JE002073><https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2003JE002073> doi: 10.1029/2003JE002073
- Greenberg, R., Geissler, P., Hoppa, G., Tufts, B. R., Durda, D. D., Pappalardo,

- R., ... Carr, M. H. (1998, 9). Tectonic Processes on Europa: Tidal Stresses, Mechanical Response, and Visible Features. *Icarus*, 135(1), 64–78. doi: 10.1006/ICAR.1998.5986
- Griffiths, A. A. (1921, 1). VI. The phenomena of rupture and flow in solids. *Philosophical Transactions of the Royal Society of London. Series A, Containing Papers of a Mathematical or Physical Character*, 4(1), 9–14. Retrieved from <https://royalsocietypublishing.org/doi/10.1098/rsta.1921.0006> doi: 10.1098/RSTA.1921.0006
- Gulley, J. D., Benn, D. I., Müller, D., Müller, M., & Luckman, A. (2009). A cut-and-closure origin for englacial conduits in uncrevassed regions of polythermal glaciers. *Journal of Glaciology*, 55(189), 66–80. Retrieved from <https://www.igsoc.org/journal/55/189/j08j039.pdf>
- Harada, Y., & Kurita, K. (2006, 2). The dependence of surface tidal stress on the internal structure of Europa: The possibility of cracking of the icy shell. *Planetary and Space Science*, 54(2), 170–180. doi: 10.1016/J.PSS.2005.12.001
- Helfenstein, P., & Parmentier, E. M. (1983, 3). Patterns of fracture and tidal stresses on Europa. *Icarus*, 53(3), 415–430. doi: 10.1016/0019-1035(83)90206-3
- Hendrix, A. R., Hurford, T. A., Barge, L. M., Bland, M. T., Bowman, J. S., Brinckerhoff, W., ... Vance, S. D. (2019, 1). The NASA Roadmap to Ocean Worlds. *Astrobiology*, 19(1), 1–27. Retrieved from <https://www.liebertpub.com/doi/10.1089/ast.2018.1955> doi: 10.1089/AST.2018.1955/ASSET/IMAGES/LARGE/FIGURE7.JPEG
- Hesse, M. A., Jordan, J. S., Vance, S. D., & Oza, A. V. (2022, 3). Downward Oxidant Transport Through Europa’s Ice Shell by Density-Driven Brine Percolation. *Geophysical Research Letters*, 49(5), e2021GL095416. Retrieved from <https://onlinelibrary.wiley.com/doi/full/10.1029/2021GL095416><https://onlinelibrary.wiley.com/doi/abs/10.1029/2021GL095416><https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021GL095416> doi: 10.1029/2021GL095416
- Hooke, R. (1989). Englacial and Subglacial Hydrology: A Qualitative Review. *Arctic and Alpine Research*, 21(3), 221–233. Retrieved from <https://www.tandfonline.com/action/journalInformation?journalCode=uaar20> doi: 10.1080/00040851.1989.12002734
- Hurford, T. A., Sarid, A. R., & Greenberg, R. (2007, 1). Cycloidal cracks on Europa: Improved modeling and non-synchronous rotation implications. *Icarus*, 186(1), 218–233. doi: 10.1016/J.ICARUS.2006.08.026
- Hussmann, H., Spohn, T., & Wiczerkowski, K. (2002, 3). Thermal Equilibrium States of Europa’s Ice Shell: Implications for Internal Ocean Thickness and Surface Heat Flow. *Icarus*, 156(1), 143–151. doi: 10.1006/ICAR.2001.6776
- Kalousová, K., Wakita, S., Sotin, C., Neish, C. D., Soderblom, J. M., Souček, O., & Johnson, B. C. (2024, 3). Evolution of Impact Melt Pools on Titan. *Journal of Geophysical Research: Planets*, 129(3), e2023JE008107. Retrieved from <https://onlinelibrary.wiley.com/doi/full/10.1029/2023JE008107><https://onlinelibrary.wiley.com/doi/abs/10.1029/2023JE008107><https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2023JE008107> doi: 10.1029/2023JE008107
- Kargel, J. S., Kaye, J. Z., Head, J. W., Marion, G. M., Sassen, R., Crowley, J. K., ... Hogenboom, D. L. (2000, 11). Europa’s Crust and Ocean: Origin, Composition, and the Prospects for Life. *Icarus*, 148(1), 226–265. doi: 10.1006/ICAR.2000.6471
- Karlstrom, L., Zok, A., & Manga, M. (2014, 3). Near-surface permeability in a supraglacial drainage basin on the Llewellyn Glacier, Juneau Icefield, British Columbia. *The Cryosphere*, 8(2), 537–546. Retrieved from <https://tc.copernicus.org/articles/8/537/2014/> doi: 10.5194/tc-8-537-2014

- Kattenhorn, S. A., & Marshall, S. T. (2006, 12). Fault-induced perturbed stress fields and associated tensile and compressive deformation at fault tips in the ice shell of Europa: implications for fault mechanics. *Journal of Structural Geology*, 28(12), 2204–2221. doi: 10.1016/J.JSG.2005.11.010
- Khurana, K. K., Kivelson, M. G., Stevenson, D. J., Schubert, G., Russell, C. T., Walker, R. J., & Polansky, C. (1998, 10). Induced magnetic fields as evidence for subsurface oceans in Europa and Callisto. *Nature*, 395(6704), 777–780. Retrieved from <https://www.nature.com/articles/27394> doi: 10.1038/27394
- Kjær, K. H., Larsen, N. K., Binder, T., Bjørk, A. A., Eisen, O., Fahnestock, M. A., ... MacGregor, J. A. (2018, 11). A large impact crater beneath Hiawatha Glacier in northwest Greenland. *Science Advances*, 4(11). Retrieved from <https://www.science.org/doi/10.1126/sciadv.aar8173> doi: 10.1126/SCIADV.AAR8173/SUPPL{_}FILE/AAR8173{_}SM.PDF
- Krawczynski, M. J., Behn, M. D., Das, S. B., & Joughin, I. (2009, 5). Constraints on the lake volume required for hydro-fracture through ice sheets. *Geophysical Research Letters*, 36(10), L10501. Retrieved from <http://doi.wiley.com/10.1029/2008GL036765> doi: 10.1029/2008GL036765
- Law, R., Christoffersen, P., Hubbard, B., Doyle, S. H., Chudley, T. R., Schoonman, C., ... Young, T. J. (2021). Thermodynamics of a fast-moving Greenlandic outlet glacier revealed by fiber-optic distributed temperature sensing. *Science Advances*, 7(20), eabe7136. doi: <https://doi.org/10.1126/sciadv.abe7136>
- Lesage, E., Massol, H., Howell, S. M., & Schmidt, F. (2022, 7). Simulation of Freezing Cryomagma Reservoirs in Viscoelastic Ice Shells. *The Planetary Science Journal*, 3(7), 170. Retrieved from <https://iopscience.iop.org/article/10.3847/PSJ/ac75bf><https://iopscience.iop.org/article/10.3847/PSJ/ac75bf/meta> doi: 10.3847/PSJ/AC75BF
- Lewis, J. S. (1971, 10). Satellites of the outer planets: Their physical and chemical nature. *Icarus*, 15(2), 174–185. doi: 10.1016/0019-1035(71)90072-8
- Lorenz, R. D., Stiles, B. W., Kirk, R. L., Allison, M. D., Del Marmo, P. P., Iess, L., ... Hensley, S. (2008, 3). Titan’s rotation reveals an internal ocean and changing zonal winds. *Science*, 319(5870), 1649–1651. Retrieved from <https://www.science.org/doi/10.1126/science.1151639> doi: 10.1126/SCIENCE.1151639/SUPPL{_}FILE/LORENZ.SOM.PDF
- Luzzi, E., Rossi, A. P., Massironi, M., Pozzobon, R., Corti, G., & Maestrelli, D. (2021, 6). Caldera Collapse as the Trigger of Chaos and Fractured Craters on the Moon and Mars. *Geophysical Research Letters*, 48(11), e2021GL092436. Retrieved from <https://onlinelibrary.wiley.com/doi/full/10.1029/2021GL092436><https://onlinelibrary.wiley.com/doi/abs/10.1029/2021GL092436><https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021GL092436> doi: 10.1029/2021GL092436
- Manga, M., & Wang, C. Y. (2007, 4). Pressurized oceans and the eruption of liquid water on Europa and Enceladus. *Geophysical Research Letters*, 34(7), 7202. Retrieved from <https://onlinelibrary.wiley.com/doi/full/10.1029/2007GL029297><https://onlinelibrary.wiley.com/doi/abs/10.1029/2007GL029297><https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2007GL029297> doi: 10.1029/2007GL029297
- McKinnon, W. B. (1999, 4). Convective instability in Europa’s floating ice shell. *Geophysical Research Letters*, 26(7), 951–954. Retrieved from <https://onlinelibrary.wiley.com/doi/full/10.1029/1999GL900125><https://onlinelibrary.wiley.com/doi/abs/10.1029/1999GL900125><https://agupubs.onlinelibrary.wiley.com/doi/10.1029/1999GL900125> doi: 10.1029/1999GL900125
- Michaut, C., & Manga, M. (2014, 3). Domes, pits, and small chaos on Europa produced by water sills. *Journal of Geophysical Research: Planets*, 119(3),

- 550–573. Retrieved from <https://onlinelibrary.wiley.com/doi/full/10.1002/2013JE004558><https://onlinelibrary.wiley.com/doi/abs/10.1002/2013JE004558><https://agupubs.onlinelibrary.wiley.com/doi/10.1002/2013JE004558> doi: 10.1002/2013JE004558
- Mitri, G., & Showman, A. P. (2005, 10). Convective–conductive transitions and sensitivity of a convecting ice shell to perturbations in heat flux and tidal-heating rate: Implications for Europa. *Icarus*, *177*(2), 447–460. doi: 10.1016/J.ICARUS.2005.03.019
- NASA. (2017). *Europa Lander Study 2016 Report* (Tech. Rep.). NASA. Retrieved from <https://europa.nasa.gov/resources/58/europa-lander-study-2016-report/>
- Pappalardo, R. T., & Barr, A. C. (2004, 1). The origin of domes on Europa: The role of thermally induced compositional diapirism. *Geophysical Research Letters*, *31*(1). Retrieved from <https://onlinelibrary.wiley.com/doi/full/10.1029/2003GL019202><https://onlinelibrary.wiley.com/doi/abs/10.1029/2003GL019202><https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2003GL019202> doi: 10.1029/2003GL019202
- Pappalardo, R. T., Head, J. W., Greeley, R., Sullivan, R. J., Pilcher, C., Schubert, G., ... Goldsby, D. L. (1998, 1). Geological evidence for solid-state convection in Europa’s ice shell. *Nature* *198* *391*:6665, *391*(6665), 365–368. Retrieved from <https://www.nature.com/articles/34862> doi: 10.1038/34862
- Poinar, K., & C. Andrews, L. (2021, 3). Challenges in predicting Greenland supraglacial lake drainages at the regional scale. *Cryosphere*, *15*(3), 1455–1483. doi: 10.5194/TC-15-1455-2021
- Porco, C. C., Helfenstein, P., Thomas, P. C., Ingersoll, A. P., Wisdom, J., West, R., ... Squyres, S. (2006, 3). Cassini observes the active south pole of Enceladus. *Science*, *311*(5766), 1393–1401. Retrieved from <https://www.science.org/doi/10.1126/science.1123013> doi: 10.1126/SCIENCE.1123013/SUPPL{ }FILE/PORCOCC.SOM.PDF
- Quick, L. C., Roberge, A., Mendoza, G. T., Quintana, E. V., & Youngblood, A. A. (2023, 10). Prospects for Cryovolcanic Activity on Cold Ocean Planets. *The Astrophysical Journal*, *956*(1), 29. Retrieved from <https://iopscience.iop.org/article/10.3847/1538-4357/ace9b6><https://iopscience.iop.org/article/10.3847/1538-4357/ace9b6/meta> doi: 10.3847/1538-4357/ACE9B6
- Rossi, C., Cianfarra, P., Lucchetti, A., Pozzobon, R., Penasa, L., Munaretto, G., & Pajola, M. (2023, 11). Deformation patterns of icy satellite crusts: Insights for tectonic balancing and fluid migration through structural analysis of terrestrial analogues. *Icarus*, *404*, 115668. doi: 10.1016/J.ICARUS.2023.115668
- Rudolph, M. L., & Manga, M. (2009, 2). Fracture penetration in planetary ice shells. *Icarus*, *199*(2), 536–541. doi: 10.1016/J.ICARUS.2008.10.010
- Russell, A. J. (1993, 1). Supraglacial lake drainage near Sendre Strømjørd, Greenland. *Journal of Glaciology*, *39*(132), 431–433. Retrieved from https://www.cambridge.org/core/product/identifier/S0022143000016105/type/journal_article doi: 10.3189/S0022143000016105
- Schmidt, B. E., Blankenship, D. D., Patterson, G. W., & Schenk, P. M. (2011, 11). Active formation of ‘chaos terrain’ over shallow subsurface water on Europa. *Nature*, *479*(7374), 502–505. Retrieved from <https://www.nature.com/articles/nature10608> doi: 10.1038/nature10608
- Sephton, M. A., Waite, J. H., & Brockwell, T. G. (2018, 7). How to detect life on icy moons. *Astrobiology*, *18*(7), 843–855. Retrieved from <https://www.liebertpub.com/doi/10.1089/ast.2017.1656> doi: 10.1089/AST.2017.1656/ASSET/IMAGES/LARGE/FIGURE5.JPEG
- Showman, A. P., & Han, L. (2004, 1). Numerical simulations of convection in Europa’s ice shell: Implications for surface features. *Journal of Geophysical Research*:

- Planets*, 109(E1), 1010. Retrieved from <https://onlinelibrary.wiley.com/doi/full/10.1029/2003JE002103><https://onlinelibrary.wiley.com/doi/abs/10.1029/2003JE002103><https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2003JE002103> doi: 10.1029/2003JE002103
- Soderlund, K. M., Schmidt, B. E., Wicht, J., & Blankenship, D. D. (2013, 12). Ocean-driven heating of Europa's icy shell at low latitudes. *Nature Geoscience*, 7(1), 16–19. Retrieved from <https://www.nature.com/articles/ngeo2021> doi: 10.1038/ngeo2021
- Sotin, C., Head, J. W., & Tobie, G. (2002, 4). Europa: Tidal heating of upwelling thermal plumes and the origin of lenticulae and chaos melting. *Geophysical Research Letters*, 29(8), 74–1. Retrieved from <https://onlinelibrary.wiley.com/doi/full/10.1029/2001GL013844><https://onlinelibrary.wiley.com/doi/abs/10.1029/2001GL013844><https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2001GL013844> doi: 10.1029/2001GL013844
- Steinbrügge, G., Voigt, J. R., Wolfenbarger, N. S., Hamilton, C. W., Soderlund, K. M., Young, D. A., ... Schroeder, D. M. (2020, 11). Brine Migration and Impact-Induced Cryovolcanism on Europa. *Geophysical Research Letters*, 47(21), e2020GL090797. Retrieved from <https://onlinelibrary.wiley.com/doi/full/10.1029/2020GL090797><https://onlinelibrary.wiley.com/doi/abs/10.1029/2020GL090797><https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2020GL090797> doi: 10.1029/2020GL090797
- Stevens, L. A., Behn, M. D., McGuire, J. J., Das, S. B., Joughin, I., Herring, T., ... King, M. A. (2015, 6). Greenland supraglacial lake drainages triggered by hydrologically induced basal slip. *Nature* 2015 522:7554, 522(7554), 73–76. Retrieved from <https://www.nature.com/articles/nature14480> doi: 10.1038/nature14480
- Tada, H., Paris, P. C., & Irwin, G. (1973). *The Stress Analysis of Cracks Handbook* (Tech. Rep.). The Materials Information Society.
- van der Veen, C. J. (1998, 6). Fracture mechanics approach to penetration of bottom crevasses on glaciers. *Cold Regions Science and Technology*, 27(3), 213–223. doi: 10.1016/S0165-232X(98)00006-8
- van der Veen, C. J. (2007, 1). Fracture propagation as means of rapidly transferring surface meltwater to the base of glaciers. *Geophysical Research Letters*, 34(1), L01501. Retrieved from <http://doi.wiley.com/10.1029/2006GL028385> doi: 10.1029/2006GL028385
- Walker, C. C., & Schmidt, B. E. (2015, 2). Ice collapse over trapped water bodies on Enceladus and Europa. *Geophysical Research Letters*, 42(3), 712–719. Retrieved from <https://onlinelibrary.wiley.com/doi/full/10.1002/2014GL062405><https://onlinelibrary.wiley.com/doi/abs/10.1002/2014GL062405><https://agupubs.onlinelibrary.wiley.com/doi/10.1002/2014GL062405> doi: 10.1002/2014GL062405
- Weertman, J. (1973). Can a water-filled crevasse reach the bottom surface of a glacier. *International Association of Scientific Hydrology*, 95(139-145). Retrieved from http://hydrologie.org/redbooks/a095/iahs_095_0139.pdf
- Williamson, A. G., Banwell, A. F., Willis, I. C., & Arnold, N. S. (2018, 4). Dual-satellite (Sentinel-2 and Landsat 8) remote sensing of supraglacial lakes in Greenland. *The Cryosphere Discussions*, 1–27. Retrieved from <https://www.the-cryosphere-discuss.net/tc-2018-56/> doi: 10.5194/tc-2018-56
- Williamson, A. G., Willis, I. C., Arnold, N. S., & Banwell, A. F. (2018, 4). Controls on rapid supraglacial lake drainage in West Greenland: an Exploratory Data Analysis approach. *Journal of Glaciology*, 64(244), 208–226. Retrieved from https://www.cambridge.org/core/product/identifier/S0022143018000084/type/journal_article doi: 10.1017/jog.2018.8