

CONVEX CO-COMPACT HYPERBOLIC MANIFOLDS ARE DETERMINED BY THEIR PLEATING LAMINATION

BRUNO DULAR AND JEAN-MARC SCHLENKER

ABSTRACT. Convex co-compact 3-dimensional hyperbolic manifolds are uniquely determined by the pleating measured lamination on the boundary of their convex core.

CONTENTS

1. Introduction	1
1.1. Background	1
1.2. Result	2
1.3. Main ideas of the proof	2
1.4. Further questions	3
Acknowledgements	3
2. Background material	4
2.1. Convex co-compact hyperbolic metrics	4
2.2. Constant curvature surfaces in hyperbolic ends	4
2.3. Duality between hyperbolic ends and de Sitter spacetimes	4
2.4. The third fundamental form and bending lamination	5
3. Analyticity	6
4. Contractibility	7
4.1. Earthquakes	7
4.2. Minima of intersection numbers	10
4.3. Fibres of a limit of homeomorphisms	10
5. Non-contractibility	12
References	12

1. INTRODUCTION

1.1. **Background.** Let M be a 3-dimensional convex co-compact hyperbolic manifold. Then M contains a smallest non-empty geodesically convex subset, called its convex core and denoted here by $C(M)$. Thurston [Thu80, Chapter 8] noticed that the boundary of $C(M)$ is homeomorphic to $\partial_\infty M$, and is a disjoint union of locally convex pleated surfaces. As a consequence, its induced metric m is hyperbolic – of constant curvature -1 – while its pleating is described by a measured lamination l which is geodesic for m , called its *pleating* or *bending* lamination. Thurston conjectured that l (resp. m) uniquely determine M . The main goal here is to prove that, indeed, l uniquely determines M .

The measured laminations on $\partial\bar{M}$ that can be realized as the pleating lamination of the convex core of a convex co-compact hyperbolic metric were determined by Bonahon and Otal [BO04].

Theorem 1.1 (Bonahon, Otal). *Let \bar{M} be a compact 3-manifold with non-empty, incompressible boundary, and such that the interior M of \bar{M} admits a complete hyperbolic metric, and with all boundary components of genus at least 2. Let l be a measured lamination on $\partial\bar{M}$. Assume that:*

- (1) *each closed leaf of l has weight less than π ,*
- (2) *if \bar{M} is not an interval bundle over a closed surface, then $i(l, \partial A) > 0$ for all essential annulus or Möbius strip A in \bar{M} ,*

Date: v0, March 18, 2024.

JMS was partially supported by FNR project O20/14766753.

- (3) if \bar{M} is an interval bundle over a closed surface S , then $i(l, p^*l') > 0$ for all non-trivial measured geodesic lamination l' , where $p : \mathcal{ML}(S) \rightarrow \mathcal{ML}(\partial M)$ is the map induced by the restriction of the bundle projection $p : \partial \bar{M} \rightarrow S$.

Then there exists a non-Fuchsian, convex co-compact hyperbolic metric on M such that l is the measured pleating lamination of the boundary of the convex core of M .

This theorem is actually stated in [BO04] in the more general case of geometrically finite hyperbolic manifolds. Lecuire [Lec06] extended it to manifolds with compressible boundary. Bonahon and Otal [BO04] also showed that *rational* laminations — measured laminations with support a disjoint union of closed curves — can be uniquely realized. Moreover, Bonahon [Bon05] showed that any small enough pair of filling measured laminations can be realized uniquely as the measured pleating lamination of the boundary of the convex core of a quasifuchsian manifold close to the Fuchsian locus. For quasifuchsian manifolds over the once punctured torus, uniqueness of the realization was proved by Series [Ser06].

Definition 1.2. We denote by $\mathcal{ML}_{\partial M}^{\text{realizable}}$ the space of measured laminations on $\partial \bar{M}$ which satisfy the hypothesis of Theorem 1.1.

1.2. Result. The main result here is the following statement, known as Thurston's bending conjecture.

Theorem 1.3. Under the hypothesis of Theorem 1.1, the hyperbolic structure on M is uniquely determined by l .

In other terms, each $l \in \mathcal{ML}_{\partial M}^{\text{realizable}}$ is the bending lamination on the boundary of the convex core for a unique convex co-compact hyperbolic metric on M .

We expect the proof to extend to convex co-compact manifolds with compressible boundary. The proof might also extend to geometrically finite, rather than convex co-compact, hyperbolic manifolds.

1.3. Main ideas of the proof. We consider a compact 3-manifold \bar{M} with incompressible boundary, and such that the interior M of \bar{M} admits a complete hyperbolic metric, with all boundary components of genus at least 2. We denote by $\mathcal{CC}(M)$ the space of convex co-compact hyperbolic metrics on M , considered up to isotopy.

We denote by $\mathcal{T}_{\partial M}$ the Teichmüller space of $\partial \bar{M}$, and by $\mathcal{ML}_{\partial M}$ the space of measured laminations on $\partial \bar{M}$. More generally, when S is a closed surface (which typically will be a boundary component of M) we denote by \mathcal{T}_S and by \mathcal{ML}_S , respectively, the Teichmüller space of S and the space of measured laminations on S . For $K \in [-1, 0)$, we denote by \mathcal{T}_S^K the space of metrics of constant curvature K on S , which is homeomorphic to \mathcal{T}_S through rescaling.

Recall that by the celebrated Ahlfors-Bers Theorem (see e.g. [Mar16, §5.1, 5.2]), when M has incompressible boundary, the map sending a convex co-compact hyperbolic structure $g \in \mathcal{CC}(M)$ to its conformal structure at infinity $c \in \mathcal{T}_{\partial M}$ is a homeomorphism from $\mathcal{CC}(M)$ to $\mathcal{T}_{\partial M}$.

Theorem 1.3 is equivalent to the statement that the following map is injective. We already know by Theorem 1.1 that this map is surjective on $\mathcal{ML}_{\partial M}^{\text{realizable}}$.

Definition 1.4. Let $\psi : \mathcal{CC}(M) \rightarrow \mathcal{ML}_{\partial M}$ be the map sending a convex co-compact hyperbolic metric $g \in \mathcal{CC}(M)$ to the measured bending lamination on the boundary of its convex core.

Theorem 1.3 follows from the following two simple lemmas.

Lemma 1.5. Let $l \in \mathcal{ML}_{\partial M}^{\text{realizable}}$. Then $\psi^{-1}(\{l\})$ is contractible.

Lemma 1.6. Let $l \in \mathcal{ML}_{\partial M}^{\text{realizable}}$. Then $\psi^{-1}(\{l\})$ is a compact analytic subvariety of $\mathcal{CC}(M)$. Therefore it is non-contractible, unless it is a point.

1.3.1. Contractibility of the fibers. To prove Lemma 1.5, we will show that ψ is a limit of homeomorphisms.

Definition 1.7. Let $K \in (-1, 0)$. We denote by $\psi_K : \mathcal{CC}(M) \rightarrow \mathcal{T}_{\partial M}^{K^*}$ the map sending a convex co-compact hyperbolic metric g on M to the third fundamental form on the closed surface S_K of constant curvature K in M .

Here S_K is in general non-connected, and has one connected component for each boundary component of M — it follows from a result of Labourie (see Theorem 2.1 below) that this K -surface is well-defined. The third fundamental for \mathcal{III}_K on S_K then has constant curvature $K^* = K/(K + 1)$, see [Lab92b, Prop. 2.3.2]. It follows from [Sch06, Theorem 0.2] that for all $K \in (-1, 0)$, ψ_K is a homeomorphism.

Lemma 1.8. *As $K \rightarrow -1$, $\psi_K \rightarrow \psi$ pointwise in the marked length spectrum topology. Moreover, for any closed curve c on $\partial\bar{M}$, $L_{\psi_K(-)}(c) \rightarrow i(\psi(-), c)$ uniformly on compact subsets of $\mathcal{CC}(M)$.*

In other words, for any closed curve c on $\partial\bar{M}$ and any $g \in \mathcal{CC}(M)$,

$$\lim_{K \rightarrow -1} L_{\psi_K(g)}(c) \rightarrow i(\psi(g), c) ,$$

where $L_{\psi_K(g)}(c)$ denotes the length of the geodesic representative of c in the constant curvature metric $\psi_K(g)$ and $i(\psi(g), c)$ denotes the intersection between the measured lamination $\psi(g)$ and c .

Since ψ is a limit of a sequence of homeomorphisms, we can use results from the decomposition theory of manifolds [Dav86] to conclude that its fibres are contractible, see Section 4.3.

1.3.2. *Non-contractibility of the fibers.* To prove that the inverse image by ψ of a point in $\mathcal{ML}_{\partial M}$ is either a point or non-contractible, we will use the following statement.

Lemma 1.9. *Let $l \in \mathcal{ML}_{\partial M}$ satisfying the hypothesis of Theorem 1.1. Then $\psi^{-1}(\{l\})$ is a real analytic variety in $\mathcal{CC}(M)$.*

Here by “real analytic variety” we mean that, for each $l \in \mathcal{ML}_{\partial M}$ satisfying the hypothesis of Theorem 1.1, there exists a real analytic function f_l with values in a real vector space, such that $\psi^{-1}(\{l\}) = f_l^{-1}(\{0\})$. The proof of this statement uses Bonahon’s shear-bent coordinates on pleated surfaces in the ends of M , see Section 3.

The proof of Lemma 1.6 then follows using a result of Sullivan [Sul71], see Section 5.

1.4. **Further questions.** We do not consider here the “dual” question: whether convex co-compact hyperbolic manifolds are uniquely determined by the induced metric on the boundary of their convex core. A slightly subtle point is that our arguments do not, at this point, allow us to conclude that convex co-compact hyperbolic manifolds are *infinitesimally rigid* with respect to the measured bending lamination on the boundary of their convex core, that is, that any first-order deformation of a convex co-compact hyperbolic manifold induces a non-zero first-order variation of the bending lamination on the boundary of its convex core. Bonahon [Bon96] noticed that this infinitesimal rigidity property is *equivalent* to the infinitesimal rigidity with respect to the induced metric on the boundary of the convex core, a statement which would presumably lead to the proof, using existing tools, that convex co-compact manifolds are uniquely determined by the induced metric on the boundary of their convex core.

Theorem 1.3 can be considered as a special case of a more general question: whether a convex hyperbolic manifold with boundary can be described in terms of geometric data on its boundary. Two types of data can be considered: the first is the induced metric, considered for polyhedra in \mathbb{H}^3 by Alexandrov [Ale05], for convex domains with smooth boundary by Alexandrov and Pogorelov [Pog73], for convex domains in hyperbolic manifolds in [Lab92a, Sch06]. Dually, one can consider a “bending” data: for ideal polyhedra in \mathbb{H}^3 it describes the dihedral angles [And71, Riv96], for compact polyhedra to their “dual metric” [HR93], and for convex domains with smooth boundary in hyperbolic manifolds it corresponds to their third fundamental form [Sch06], while for convex cores of hyperbolic manifolds it corresponds to the measured bending lamination. Recent results deal with the induced metric [Slu18, Pro22b] and dihedral angles or dual metrics [Sch01, Pro22a] of convex domains with polyhedral boundary in hyperbolic manifolds. One can also prescribe the induced metric on some boundary components of a convex co-compact manifold and the third fundamental form [CS22] or the measured lamination [Mes23] on another. Other results describe the induced metric or third fundamental forms of smooth or polyhedral surfaces in \mathbb{H}^3 invariant under a cocompact action of a Fuchsian group, see e.g. [LS00, Fil07, Fil11].

The problem of prescribing the induced metric (or third fundamental form) of the boundary of a convex hyperbolic manifold can be considered in a “universal” setting, where a mostly conjectural picture emerges, see [BDMS21, Sch21, Sch20].

The analog of Thurston’s question on the bending laminations on the boundary of the convex core was asked by Mess [Mes07, ABB⁺07] for “quasifuchsian” AdS spacetime. A partial answer was given near the Fuchsian locus [BS12], but neither existence nor uniqueness is known in general, even for rational laminations.

Acknowledgements. We would like to thank Francesco Bonsante, Cyril Lecuire and Filippo Mazzoli for useful discussions related to this paper. The second-named author would like to thank IHES, where part of this work was completed.

2. BACKGROUND MATERIAL

2.1. Convex co-compact hyperbolic metrics. We consider in this paper convex co-compact hyperbolic manifolds, which can be defined as follows: if M is the interior of a compact manifold with boundary, a convex co-compact hyperbolic metric on M is a complete hyperbolic metric g on M such that (M, g) contains a non-empty, compact, geodesically convex subset (We say that $K \subset M$ is geodesically convex if any geodesic segment in M with endpoints in K is contained in K). We always assume M to be oriented.

A convex co-compact hyperbolic manifold (M, g) contains a unique smallest, non-empty, geodesically convex subset $C(M)$, called its convex core. Except in special situations (for Fuchsian manifolds, that is, when $C(M)$ is a totally geodesic surface), $C(M)$ is a three-dimensional domain, with boundary a disjoint union of locally convex, pleated surfaces. Each boundary component of $C(M)$ is then equipped with a hyperbolic metric – induced by the ambient hyperbolic metric – and with a measured lamination, measuring its “pleating”. See [BO04, §2], [Thu80, Chapter 8].

2.2. Constant curvature surfaces in hyperbolic ends. We consider again a convex co-compact hyperbolic manifold (M, g) . Since $C(M)$ has locally convex boundary, the exponential Gauss map defines a homeomorphism between the unit normal bundle of $C(M)$ – the set of unit vectors which are normals of oriented support planes of $C(M)$ at their intersection with $C(M)$ – and $\partial_\infty M$. Moreover, the geodesic rays defined by those unit normal vectors foliate $M \setminus C(M)$.

It follows that $M \setminus C(M)$ is a disjoint union of “hyperbolic ends”: non-complete hyperbolic manifolds homeomorphic to $\partial_i M \times [0, +\infty)$, where the $\partial_i M$, $1 \leq i \leq n$, are the connected components of $\partial \bar{M}$, complete on the side corresponding to $+\infty$, and with boundary a concave pleated surface on the side corresponding to 0. We will denote the boundary at infinity of E by $\partial_\infty E$, and the concave pleated boundary by $\partial_0 E$.

Labourie [Lab91] proved that, given such a hyperbolic end E , it admits a unique foliation by closed surfaces of constant curvature K .

Theorem 2.1 (Labourie). *Let E be a hyperbolic end. It admits a unique foliation by closed surfaces of constant curvature K , with K varying from -1 close to the concave pleated boundary, to 0 near the boundary at infinity.*

As already mentioned above, the third fundamental form on a surface of constant curvature K in E is a metric of constant curvature $K^* = K/(K+1)$. As a consequence, for each $K \in (-1, 0)$, each convex co-compact hyperbolic metric g on M determines a metric of constant curvature K on $\partial \bar{M}$ – the induced metric on the surfaces of constant curvature K in the ends of M – as well as a metric of constant curvature K^* – the third fundamental form on those surfaces. We denote by $\phi_K : \mathcal{CC}(M) \rightarrow \mathcal{T}_{\partial \bar{M}}^K$ and by $\psi_K : \mathcal{CC}(M) \rightarrow \mathcal{T}_{\partial \bar{M}}^{K^*}$ the maps obtained in this manner. It follows from the main results of [Sch06] (Theorem 1.1, together with [Lab92a], for ϕ_K , Theorem 1.2 for ψ_K) that those maps are homeomorphisms.

2.3. Duality between hyperbolic ends and de Sitter spacetimes. We also need the well-known polar duality between \mathbb{H}^3 and the de Sitter space dS^3 , as well as between hyperbolic ends and globally hyperbolic maximal compact de Sitter spacetimes. We recall the main properties here, the reader can consult e.g. [RH93, Sch98, FS19] for more details.

Recall that the hyperbolic space \mathbb{H}^3 can be defined as a connected component of a quadric in the 4-dimensional Minkowski space:

$$\mathbb{H}^3 = \{x \in \mathbb{R}^{3,1} \mid \langle x, x \rangle = -1 \wedge x_0 > 0\} .$$

In the same manner, one can define the de Sitter space dS^3 as:

$$dS^3 = \{x \in \mathbb{R}^{3,1} \mid \langle x, x \rangle = 1\} ,$$

equipped with its induced metric. It is a geodesically complete, simply connected Lorentzian space of constant curvature 1.

Given an oriented totally geodesic plane $P \subset \mathbb{H}^3$, it is the intersection with \mathbb{H}^3 of a time-like linear hyperplane $H \subset \mathbb{R}^{3,1}$. The unit vector n^* positively orthogonal to H in $\mathbb{R}^{3,1}$ is then a point in dS^3 . Conversely, given any (unoriented) space-like plane P^* in dS^3 , it is the intersection with dS^3 of a unique space-like linear hyperplane $H^* \subset \mathbb{R}^{3,1}$. The future-oriented unit normal to H^* is then a point $n \in \mathbb{H}^3$.

It follows from this definition that given two oriented totally geodesic planes P_1 and P_2 in \mathbb{H}^3 , with dual points $P_1^*, P_2^* \in dS^3$:

- P_1^* and P_2^* are connected by a space-like geodesic segment s if and only if P_1 and P_2 intersect, and the length of s is then the intersection angle between P_1 and P_2 .

- P_1^* and P_2^* are connected by a light-like segment if and only if P_1 and P_2 are disjoint but at distance 0, and oriented in a compatible way in the sense that the half-space bounded by one is contained in the half-space bounded by the other.
- P_1^* and P_2^* are connected by a time-like geodesic segment s if and only if P_1 and P_2 are at finite distance and oriented in a compatible way, and the length of s is then the hyperbolic distance between P_1 and P_2 .

If $S \subset \mathbb{H}^3$ is an oriented, locally strongly convex surface (that is, its second fundamental form is positive definite at every point), one can define the dual surface $S^* \subset dS^3$ as the set of points in dS^3 dual to the tangent planes of S . A key property is that S^* is then a space-like, strongly convex surface, and that the induced metric of S^* is equal – through the identification between S and S^* coming from the duality – to the third fundamental form of S , and conversely.

Suppose now that E is a hyperbolic end. Its universal cover \tilde{E} can be identified isometrically to a domain in \mathbb{H}^3 bounded by a concave pleated surface $\partial_0\tilde{E} \subset \mathbb{H}^3$. The set of points in dS^3 dual to the oriented planes bounding half-spaces contained in \tilde{E} constitutes a *domain of dependence*, denoted by \tilde{E}^* , in dS^3 . It is a convex domain bounded by a locally convex surface $\partial_0\tilde{E}^* \subset dS^3$ which is dual to $\partial_0\tilde{E}$. In a sense which can be made precise, $\partial_0\tilde{E}^*$ is everywhere light-like, except along a real tree which is dual to the lift to $\partial_0\tilde{E}$ of the measured bending lamination on the concave pleated surface ∂_0E in E .

By construction, \tilde{E}^* is invariant under an isometric action of the fundamental group Γ of E , which is the fundamental group of the boundary component of M corresponding to E . The quotient $E^* = \tilde{E}^*/\Gamma$ is a globally hyperbolic maximal compact (GHMC) de Sitter spacetime, with initial singularity the quotient by Γ of the real tree described above.

Given $K \in (-1, 0)$ and a surface S of constant curvature K in E , S lifts to a constant curvature K surface \tilde{S} in \tilde{E} , which is locally strictly convex. The dual surface $\tilde{S}^* \subset dS^3$ is a space-like, strongly convex surface. It is also invariant under Γ , and the quotient S^* is a Cauchy surface in E^* . Moreover, as already mentioned, S^* has induced metric equal (through the identification between S and S^* through the duality map) to the third fundamental form of S , and its curvature is constant and equal to $K^* = K/(K+1)$.

In this manner, the surfaces dual in E^* to the closed surfaces of constant curvature $K \in (-1, 0)$ which foliate E define a foliation of E^* by closed surface of constant curvature $K^* \in (-\infty, -1)$. The existence and uniqueness of this foliation was proved in [BBZ11].

2.4. The third fundamental form and bending lamination. As mentioned above, we will need a slight refinement of the following statement, similar to (slightly more elaborate) results obtained in [BMS13, Section 6].

Lemma 2.2. *Let g be a fixed convex co-compact hyperbolic metric on M . Then $\lim_{K \rightarrow -1} \psi_K(g) = \psi(g)$ in the topology of the marked length spectrum, that is, for any closed curve c on $\partial\bar{M}$, $\lim_{K \rightarrow -1} L_{\mathbb{H}_K}(c) = i(l, c)$, where l is the measured pleating lamination on the boundary of the convex core of M .*

Proof. The proof follows directly, through the duality between hyperbolic and de Sitter space, from the main result in [Bel17], see above. \square

The refinement we need is that the convergence is actually uniform on compact subsets of $\mathcal{CC}(M)$. While the proof is basically the same as for Lemma 2.2, it is necessary to keep track more carefully of the convergence.

Lemma 2.3. *Let c be a closed curve on $\partial\bar{M}$. Then $L_{\mathbb{H}_K}(c) \rightarrow i(l, c)$ uniformly on compact subsets of $\mathcal{CC}(M)$ as $K \rightarrow -1$*

Proof. Notice first that if E is an end of a quasifuchsian manifold and S is a convex surface in S , then the gradient on S of the distance $d : E \rightarrow \mathbb{R}_{\geq 0}$ to the concave pleated boundary of E is bounded by 1.

As a consequence, there exists $K_0 > -1$ and a continuous function $\Delta : [-1, K_0] \rightarrow \mathbb{R}_{\geq 0}$, with $\Delta(-1) = 0$, such that, for $K \in [-1, K_0]$, the restriction of d to the surface S_K is bounded above by $\Delta(K)$. Indeed, if $d(x) \geq d_0$ at a point $x \in S_K$, then $d \geq d_0/2$ on a ball of radius $d_0/2$ around x in S_K . Since the projection from S_K to ∂_0E is contracting by a factor at least $\cosh(d)$, and since the ball of radius r in S_K has area equal to $\pi(\cosh(r/\sqrt{|K|}) - 1)$, it would follow that

$$\text{Area}(S_K) \geq \text{Area}(\partial_0E) + (\cosh(d_0/2)^2 - 1)\pi(\cosh(d_0/2\sqrt{|K|}) - 1).$$

However it follows from the Gauss-Bonnet Theorem that $\text{Area}(S_K) = \text{Area}(\partial_0E)/|K|$, leading to an upper bound on d_0 when K is close to -1 .

It then follows that, for any $x \in S_K$, the tangent plane $T_x S_K$ to S_K at x is at distance at most $\Delta(K)$ from any support plane of $\partial_0 E$, because any support plane of $\partial_0 E$ separates x from $\partial_0 E$, so that the distance from $T_x S_K$ to the support plane is at most equal to the distance from x to $\partial_0 E$. (Here we consider that two hyperbolic planes are at distance 0 if they intersect.) Through the duality between E and E^* , this means that any point of S_K^* is at time distance at most $\Delta(K)$ from the initial singularity $\partial_0 E^*$ of E^* . (In other terms, given $x^* \in S_K^*$, the maximal time distance from x^* to $\partial_0 E^*$ is at most d_0 .)

For the last step of the argument, we introduce the foliation $(\Sigma_\tau)_{\tau \in \mathbb{R}_{>0}}$ of E^* by surfaces of constant cosmological time, see [Bel17]. Recall that the cosmological time is the time distance to the initial singularity of E^* . It follows from the previous argument that S_K^* is in the past of $\Sigma_{\Delta(K)}$ for K small enough.

A key property of convex foliations in Lorentzian spacetimes (see [Bel17, Prop. 3.1]) is that given a leaf of such a foliation, for instance Σ_τ for some $\tau > 0$, the projection along the normal lines from the past of Σ_τ is expanding. As a consequence, if we denote by g_τ the induced metric on Σ_τ , we have

$$L_{\mathbb{I}K}(c) \leq L_{g_{\Delta(K)}}(c),$$

while the same argument applied to the projection on S_K along the normal lines of the foliation by K -surfaces indicates that

$$i(l, c) \leq L_{\mathbb{I}K}(c).$$

The result therefore follows from the fact that $L_{g_\tau}(c) \rightarrow i(l, c)$ uniformly on compact subsets of $\mathcal{CC}(M)$, since the metric g_τ can be expressed in terms of grafting along the measured lamination l , see [Bel17]. \square

Proof of Lemma 1.8. The lemma is a restated version, in terms of the functions ψ_K and ψ , of Lemma 2.3. \square

3. ANALYTICITY

In this section we prove the following lemma, which will be used below in two different manners: once to show that the space of hyperbolic structures with a given measured bending lamination is contractible, and another time to prove that it cannot be contractible, unless it is a point.

For simplicity, for $l \in \mathcal{ML}_{\partial M}^{\text{realizable}}$, we will denote by $\mathcal{CC}_l(M) = \psi^{-1}(\{l\})$ the set of convex co-compact hyperbolic structures on M for which the measured bending lamination on the boundary of $C(M)$ is l . Theorem 1.3 is equivalent to the statement that, for all $l \in \mathcal{ML}_{\partial M}^{\text{realizable}}$, $\mathcal{CC}_l(M)$ is reduced to one point.

Lemma 3.1. *Let $l \in \mathcal{ML}_{\partial M}^{\text{realizable}}$. Then $\mathcal{CC}_l(M)$ is a compact, analytic subvariety in $\mathcal{CC}(M)$.*

Here by ‘‘analytic subvariety’’ we mean that there exists an analytic function $F_l : \mathcal{CC}(M) \rightarrow \mathbb{R}$ such that $F_l^{-1}(\{0\}) = \mathcal{CC}_l(M)$.

Proof. The analytic structure of $\mathcal{CC}_l(M)$ essentially follows from Bonahon’s construction of the shear-bent cocycle [Bon96].

Let $\partial_1 M, \partial_2 M, \dots, \partial_n M$ be the connected components of $\partial \bar{M}$, which are closed, oriented surfaces of genus at least 2. For each $i \in \{1, \dots, n\}$, we denote by $\mathcal{CP}_{\partial_i M}$ the space of complex projective structures on $\partial_i M$, and by $\mathcal{CP}_{\partial M} = \prod_{i=1}^n \mathcal{CP}_{\partial_i M}$ the space of complex projective structures on $\partial \bar{M}$. For each i , $\mathcal{CP}_{\partial_i M}$ is a complex manifold, see e.g. [Dum08], and so is the product $\mathcal{CP}_{\partial M}$. Moreover, $\mathcal{CC}(M)$ embeds in $\mathcal{CP}_{\partial M}$ as a complex submanifold (this follows from the definition of the complex structure on $\mathcal{CC}(M)$ and from the fact that the holonomy map of a complex projective structure is a biholomorphism, see e.g. [Dum08, §5.2].)

Bonahon [Bon96] considers, in a closed, orientable surface S , a maximal lamination λ . He defines an open subset $\mathcal{R}(\lambda)$ of the space of representations of $\pi_1 S$ in $PSL(2, \mathbb{C})$ which leave invariant a pleated surface with pleating locus corresponding to λ . For each $\rho \in \mathcal{R}(\lambda)$, he defines a *shear-bend* cocycle $\Gamma_\rho \in \mathcal{H}(\lambda; \mathbb{C}/2\pi i\mathbb{Z})$ supported on λ . He shows [Bon96, Theorem D] that this map $\rho \rightarrow \Gamma_\rho$ induces a biholomorphic homeomorphism from \mathcal{CP}_S to a certain cone in the complex vector space $\mathcal{H}(\lambda; \mathbb{C}/2\pi i\mathbb{Z})$.

The construction of this map goes through pleated surfaces in the hyperbolic manifold corresponding to a complex projective structure ρ , and the imaginary part $i\beta_\rho$ of Γ_ρ corresponds precisely to the bending cocycle along λ for this pleated surface. In particular, β_ρ is positive if and only if the pleated surface corresponding to λ is locally convex.

Going back to M , the subset $\mathcal{CP}_{\partial_i M}(l_i) \subset \mathcal{CP}_{\partial_i M}$ of complex projective structures on $\partial_i M$ containing a locally convex surface with pleating lamination l_i is therefore the inverse image by $\rho \mapsto \Gamma_\rho$ (for any λ containing the support of l_i) of a totally real vector subspace of $\mathcal{H}(\lambda; \mathbb{C}/2\pi i\mathbb{Z})$. Therefore, $\mathcal{CP}_{\partial_i M}(l_i)$ is a real analytic subvariety of $\mathcal{CP}_{\partial_i M}$.

Now it follows from the definitions that

$$\mathcal{CC}_i(M) = \mathcal{CC}(M) \cap (\mathcal{CP}_{\partial_1 M}(l_1) \times \cdots \times \mathcal{CP}_{\partial_n M}(l_n)) \subset \mathcal{CP}_{\partial M} ,$$

so that it is a real analytic subvariety of $\mathcal{CC}(M)$.

The compactness of $\mathcal{CC}_i(M)$ is an immediate consequence, in the more general case of geometrically finite hyperbolic manifolds, of the main result from [Lec04], the properness of the bending map. (We believe that it also follows, in the convex co-compact case, from the arguments used by Bonahon and Otal to prove their Closing Lemma [BO04, Prop. 8].) \square

4. CONTRACTIBILITY

In this section we will show Lemma 1.5. According to Lemma 2.2, for a fixed convex co-compact metric on M , as $K \rightarrow -1$, the third fundamental forms \mathbb{III}_K on the K -surfaces in the ends of M converge, in the topology of the marked length spectrum, to the bending lamination. This can be interpreted heuristically as a sort of convergence of the homeomorphisms ψ_K defined above to ψ , and our goal is indeed to show that ψ is a limit of homeomorphisms. So an additional construction is necessary to identify in some way the spaces $\mathcal{T}_S^{K^*}$ of metrics of constant curvature $K^* = K/(K+1)$, as $K \rightarrow -1$, with \mathcal{ML}_S .

We provide one argument using earthquakes, and we sketch another incomplete one through geodesic currents. Yet another possible approach might follow the arguments appearing in [Maz22, Prop. 4.4].

4.1. Earthquakes. In this section and the next, we fix a closed surface S , which will be a connected component of $\partial\bar{M}$.

We consider in this section a fixed metric $m_0 \in \mathcal{T}_S$. We then consider the left earthquake map based at m_0 , $E_L^{m_0} : \mathcal{ML}_S \rightarrow \mathcal{T}_S$. Recall that $E_L^{m_0}$ is a homeomorphism, by Thurston's Earthquake Theorem [Thu86].

To simplify notations, we denote by $\mathcal{T}_S^{K^*}$ the space of metrics of constant curvature $K^* = K/(K+1)$ on S considered up to isotopy. This space $\mathcal{T}_S^{K^*}$ is canonically identified with \mathcal{T}_S , because if m is a hyperbolic metric on S , then $(1/|K^*|)m$ is a metric of constant curvature K^* . We use this notation since the third fundamental form on a surface of constant curvature K in M has constant curvature K^* , see [Lab92b].

Definition 4.1. We denote by $u_K : \mathcal{T}_S^{K^*} \rightarrow \mathcal{ML}_S$ the function defined as follows. Let $h \in \mathcal{T}_S^{K^*}$ be a metric of constant curvature K^* . Let $m = |K^*|h$ be the hyperbolic metric homothetic to h . We set

$$u_K(h) = (1/\sqrt{|K^*|})(E_L^{m_0})^{-1}(m) .$$

Lemma 4.2. (1) For all $K \in (-1, 0)$, u_K is a homeomorphism.

(2) Let $(K_n)_{n \in \mathbb{N}}$ be a sequence in $(-1, 0)$ converging to -1 , and let $(h_n)_{n \in \mathbb{N}}$ be a sequence of metrics with $h_n \in \mathcal{T}_S^{K_n^*}$ such that $h_n \rightarrow l \in \mathcal{ML}$ in the topology of the marked length spectrum. Then $u_{K_n}(h_n) \rightarrow l$.

In fact, point (2) of this lemma is not entirely sufficient for the proof of the main result, and we will need the following slightly refined statement, whose proof follows from the same arguments, see below.

Lemma 4.3. Let $(K_n)_{n \in \mathbb{N}}$ be a sequence in $(-1, 0)$ converging to -1 , and let $(h_n(t))_{n \in \mathbb{N}}$ be a sequence of families of metrics, with $h_n(t) \in \mathcal{T}_S^{K_n^*}$, depending on a parameter t in a topological space T . Let γ be a closed curve on S . Assume that $Lh_n(t)(\gamma) \rightarrow l(t) \in \mathcal{ML}_S$ uniformly in t on compact subsets of T , where $l(t) \in \mathcal{ML}_S$ for each $t \in T$. Then $i(u_{K_n}(h_n(t)), \gamma) \rightarrow i(l(t), \gamma)$ uniformly in t on compact subsets of T .


The proof will rely on an elementary estimate, which might be of independent interest. See [Bon92, Lemma 16] for a complementary estimate.

Lemma 4.4. Let (S, h) be a hyperbolic surface, let $l \in \mathcal{ML}_S$, and let γ be a homotopy class of simple closed curves on S . Then

$$i(\gamma, l) - L_m(\gamma) \leq L_{E_L^{m_0}(l)}(\gamma) \leq i(\gamma, l) + L_m(\gamma) ,$$

where $i(\gamma, l)$ denotes the intersection between γ and l , and $L_m(\gamma)$ denotes the length of the geodesic representative of γ for m .

Proof. We denote by \bar{S} the cover of S with fundamental group generated by γ . We still denote by m the lift of m to \bar{S} , so that (\bar{S}, m) is a hyperbolic cylinder. The proof will be done entirely in \bar{S} . We denote by $\bar{\gamma}$ the geodesic representative of γ in (\bar{S}, m) .

FIGURE 1. Proof of Lemma 4.4, in (\bar{S}, \bar{m})

Still denote by l the measured lamination on \bar{S} corresponding to l . That is, the support of this lamination l is the union of lifts of the geodesics in the support of l on S which intersect γ , with the corresponding weight.

We now assume that l is rational in S – that is, supported on a finite set of disjoint closed curves. It is sufficient to prove the lemma in this rational case, since the general case then follows by density of rational laminations in \mathcal{ML}_S .

The set of geodesics in the support of l which intersect \bar{c}' in (\bar{S}, m) is a finite set of disjoint geodesics l_1, \dots, l_n which all intersect \bar{c}' exactly once. We assume that the l_i intersect \bar{c}' in this cyclic order.

Denote by $\bar{m} = E_m^L(l)$ the image of m – the metric on \bar{S} by the left earthquake along l . We denote by c the geodesic representative of γ in (\bar{S}, \bar{m}) , and by \bar{c} the image of the geodesic representative \bar{c}' under the left earthquake along l . That is, \bar{c} is a disjoint union of geodesic segments $\bar{c}_1, \bar{c}_2, \dots, \bar{c}_n$, with \bar{c}_i having one endpoint on l_i and one on l_{i+1} . Similarly, we denote by c_i the segment of c_i between its intersections with l_i and l_{i+1} , see Figure 1. We set

$$\gamma_i = L_{\bar{m}}(c_i), \quad \bar{\gamma}_i = L_{\bar{m}}(\bar{c}_i).$$

We consider the geodesics $l_i, 1 \leq i \leq n$ as oriented (from the left to the right side of c), and denote by x_i the oriented distance along l_i between its intersection with c and its intersection with \bar{c}_i , and by y_i the oriented distance along l_i between its intersection with c and with \bar{c}_{i+1} . By definition of a left earthquake, $y_i - x_i$ is the weight of l_i in the measured lamination l . As a consequence,

$$i(\gamma, l) = \sum_{i=1}^n y_i - x_i.$$

The proof of the upper bound in the lemma directly follows, since there is a curve in (\bar{S}, \bar{m}) , in the homotopy class γ , composed of the \bar{c}_i and of segments of the l_i of length $y_i - x_i$, so that

$$L_{\bar{m}}(\gamma) = L_{\bar{m}}(c) \leq \sum_i L_{\bar{m}}(\bar{c}_i) + \sum_i (y_i - x_i) = L_m(\gamma) + i(\gamma, l).$$

We now aim at providing a lower bound on the lengths γ_i of the geodesic segments $c_i, 1 \leq i \leq n$. For this we consider three cases, as seen on Figure 1.

- $x_i \geq 0, y_{i+1} \geq 0$ (as for $i = 1$ in the figure). It then follows from the triangle inequality (or its variant for 4-gons) that

$$\gamma_i \geq y_{i+1} - x_i - \bar{\gamma}_i .$$

- $x_i \leq 0, y_{i+1} \geq 0$ (as for $i = 2$ in the figure). The triangle inequality, applied to each of the two triangles appearing in the figure, shows that

$$\gamma_i \geq y_{i+1} - x_i - \bar{\gamma}_i .$$

- $x_i \leq 0, y_{i+1} \leq 0$ (as for $i = 3$ in the figure). Again the triangle inequality, applied in a slightly different way compared to the first case, still shows that

$$\gamma_i \geq y_{i+1} - x_i - \bar{\gamma}_i .$$

We can now sum the inequalities obtained for $i = 1, \dots, n$ and obtain that

$$\sum_i \gamma_i \geq \sum_i (y_i - x_i) - \sum_i \bar{\gamma}_i,$$

so that

$$L_{\bar{m}}(\gamma) \geq i(l, \gamma) - L_m(\gamma) ,$$

as required. \square

Proof of Lemma 4.2. The first point is a direct consequence of the definition of u_K and of the Earthquake Theorem.

To prove the second point, we will use Lemma 4.4. Let m_n be the hyperbolic metric homothetic to h_n . Let $l_n \in \mathcal{ML}_S$ be the unique measured lamination such that $E_{m_0}^L(l_n) = m_n$, so that, by definition of u_K ,

$$u_{K_n}(h_n) = \frac{1}{\sqrt{|K_n^*|}} l_n .$$

It follows from Lemma 4.4 that, for all n ,

$$L_{m_n}(c) - L_{m_0}(c) \leq i(l_n, c) \leq L_{m_n}(c) + L_{m_0}(c) .$$

This can be written, using the definition of m_n , as

$$\sqrt{|K_n^*|} L_{h_n}(c) - L_{m_0}(c) \leq i(l_n, c) \leq \sqrt{|K_n^*|} L_{h_n}(c) + L_{m_0}(c) ,$$

and, dividing by $\sqrt{|K_n^*|}$,

$$L_{h_n}(c) - \frac{L_{m_0}(c)}{\sqrt{|K_n^*|}} \leq \frac{i(l_n, c)}{\sqrt{|K_n^*|}} \leq L_{h_n}(c) + \frac{L_{m_0}(c)}{\sqrt{|K_n^*|}} .$$

Since $L_{h_n}(c) \rightarrow i(l, c)$ by the hypothesis of the lemma, we can conclude that $\frac{i(l_n, c)}{\sqrt{|K_n^*|}} \rightarrow i(l, c)$, as required. \square

Proof of Lemma 4.3. We now consider a sequence of metrics $(h_n(t))_{n \in \mathbb{N}}$ of constant curvature K_n^* , depending on a parameter $t \in T$. As above we call $m_n(t)$ the hyperbolic metric in the same conformal class as $h_n(t)$, that is,

$$m_n(t) = |K_n^*| h_n(t) ,$$

so that by the hypothesis of the lemma,

$$\frac{1}{\sqrt{|K_n^*|}} L_{m_n(t)}(\gamma) \rightarrow i(l(t), \gamma)$$

uniformly in t on compact subsets of T .

Applying Lemma 4.4, we have

$$|L_{m_n(t)}(\gamma) - i(l_n(t), \gamma)| \leq L_{m_0}(\gamma) ,$$

where $l_n(t) = (E_L^{m_0})^{-1}(m_n(t))$. It follows that

$$|\sqrt{|K_n^*|} L_{h_n(t)}(\gamma) - \sqrt{|K_n^*|} i(u_{K_n}(h_n(t)), \gamma)| \leq L_{m_0}(\gamma) ,$$

so that

$$|L_{h_n(t)}(\gamma) - i(u_{K_n}(h_n(t)), \gamma)| \leq \frac{L_{m_0}(\gamma)}{\sqrt{|K_n^*|}} .$$

It follows directly that if $L_{h_n(t)}(\gamma) \rightarrow i(l, \gamma)$ uniformly in t on compact subsets of T as $K \rightarrow -1$, then $i(u_{K_n}(h_n(t)), \gamma) \rightarrow i(l, \gamma)$ uniformly in t on compact subsets of T as $K \rightarrow -1$. \square

4.2. Minima of intersection numbers. In this section, we sketch a second incomplete approach to the problem of formalizing the convergence of ψ_K to ψ .

Let \mathcal{C}_S be the space of geodesic currents on S [Bon88]. It naturally contains the space \mathcal{ML}_S of measured laminations, as well as Teichmüller space \mathcal{T}_S under the identification of a metric m with its Liouville current L_m . Similarly, it also contains $k \cdot \mathcal{T}_S$ for each $k \in (0, 1)$. This makes of \mathcal{C}_S a natural place in which identifying all those spaces one to each other in a consistent way.

The space \mathcal{C}_S is endowed with a continuous symmetric bilinear form $i: \mathcal{C}_S \times \mathcal{C}_S \rightarrow \mathbb{R}_{\geq 0}$, the *intersection form*, which generalizes simultaneously the intersection number of measured laminations and the length of curves. A geodesic current μ is *filling* if its systole, i.e. the infimum of $i(\mu, c)$ when c runs over all simple closed curves c , is strictly positive. Let $\mathcal{C}_S^{\text{fill}}$ denote the subspace of \mathcal{C}_S consisting of filling currents.

By Lemma 2.2, the maps ψ_K converge pointwise to ψ when seen as landing inside \mathcal{C}_S . In order to identify their image, one could use a projection of \mathcal{C}_S onto \mathcal{T}_S with adequate properties.

In [HS23], Hensel and Sapir use and extend results of Kerckhoff [Ker92] and Wolpert [Wol06] to define a projection

$$(1) \quad \pi: \mathcal{C}_S^{\text{fill}} \longrightarrow \mathcal{T}_S: \mu \longmapsto \underset{m \in \mathcal{T}_S}{\operatorname{argmin}} i(\mu, m),$$

which sends a filling current to the hyperbolic metric on S minimizing their intersection. Existence and uniqueness of such a minimum is ensured by the strict convexity of the function $i(\mu, -)$ on \mathcal{T}_S , with respect to the Weil-Petersson metric [Wol06, page 5]. The projection π fixes \mathcal{T}_S . By bilinearity of the intersection form i , the projection π is scaling-invariant, hence it factors through a map $\mathbb{P}\mathcal{C}_S^{\text{fill}} \rightarrow \mathcal{T}_S$, which is continuous and proper [HS23, Theorem 1.1].

Let us fix, once and for all, a filling current $\omega \in \mathcal{C}_S^{\text{fill}}$. Let $\tau_\omega: \mathcal{C}_S \hookrightarrow \mathcal{C}_S^{\text{fill}}$ be the translation sending μ to $\mu + \omega$. Consider the following composition

$$(2) \quad \pi_\omega := \pi \circ [\cdot] \circ \tau_\omega: \mathcal{C}_S \xrightarrow{\tau_\omega} \mathcal{C}_S^{\text{fill}} \xrightarrow{[\cdot]} \mathbb{P}\mathcal{C}_S^{\text{fill}} \xrightarrow{\pi} \mathcal{T}_S.$$

The map π_ω is continuous, and its restriction $\pi_\omega|_{\mathcal{ML}_S}$ is a homeomorphism onto \mathcal{T}_S , by a theorem of Kerckhoff [Ker92, Theorem 2.1]. We would need to generalize the latter theorem to Liouville currents in order to obtain that $\pi_\omega|_{k \cdot \mathcal{T}_S}$ are homeomorphisms onto \mathcal{T}_S as well. More precisely, one way of proving that would be to show the following assertion: For $m, \beta, \beta' \in \mathcal{T}_S$, if $\operatorname{Cos}_m(\beta, \alpha) = \operatorname{Cos}_m(\beta', \alpha)$ for all $\alpha \in \mathcal{ML}_S$, then $\beta = \beta'$. This would generalize one of Kerckhoff's theorems [Ker92, Theorem 2.2] to Liouville currents, using Bonahon's extension of the total cosine [Bon92]. The latter expresses the derivative of length functions along earthquake paths.

If the above is true, then $\pi_\omega \circ \psi_K: \mathcal{CC}(M) \rightarrow \mathcal{T}_{\partial M}$, for $K \in (-1, 0)$, is a sequence of homeomorphisms converging to $\pi_\omega \circ \psi$ when K tends to -1 . The fibres of ψ would then be homeomorphic to the fibres of a limit of homeomorphisms.

4.3. Fibres of a limit of homeomorphisms. We now complete the proof that the fibres of the map $\psi: \mathcal{CC}(M) \rightarrow \mathcal{ML}_{\partial M}^{\text{realizable}}$ are contractible, using that ψ is the limit of homeomorphisms and that its fibres are compact real analytic subsets of $\mathcal{CC}(M)$. The following definition is taken from [Dav86].

Definition 4.5. *A subset F of a n -dimensional manifold is cellular if any neighborhood U of F contains a closed n -ball B such that $F \subset \operatorname{int}(B)$.*

In particular a cellular subset must be compact, as it is a nested intersection of compact balls. A large part of the decomposition theory of manifolds [Dav86] is interested in characterizing the fibres of a limit of homeomorphisms, culminating in the acclaimed *Bing's shrinking criterion*. Similar in spirit to that criterion is the following result, which we use here. We state a slightly different form, better suited to the present context, but the proof is exactly the same as in [Fin67]. Since the proof is elementary, we repeat it here.

Theorem 4.6 (Finney [Fin67, Theorem 1]). *Let N, N' be two manifolds, d a metric on N' (compatible with its topology), and $(f_n)_{n \in \mathbb{N}}$ a sequence of homeomorphisms $N \rightarrow N'$, converging uniformly on compacts to the continuous map $f: N \rightarrow N'$. The compact fibres of f are cellular subsets of the manifold N .*

Proof. Let $K := f^{-1}(x) \subseteq N$ be a compact fibre of f , with $x \in N'$. Let U be a neighborhood of K in N . We need to find a closed n -ball B contained in U such that $K \subset \operatorname{int}(B)$.

Let $V \subseteq N$ be an open set with $K \subset V \subset \bar{V} \subset U$ and such that \bar{V} is compact. We prove the two following facts:

- (1) The diameter (with respect to d) of $f_n(K) \cup \{x\}$ tends to 0 as n tends to infinity.
- (2) There exists $\epsilon_0 > 0$ such that $d(f_n(K), f_n(\partial\bar{V})) \geq \epsilon_0$ for all $n \in \mathbb{N}$.

We first prove (1). Assume it does not hold, then there exists $\epsilon > 0$, a strictly increasing sequence $(n_k)_{k \in \mathbb{N}}$ in \mathbb{N} and points $y_k, z_k \in f_{n_k}(K) \cup \{x\}$ such that $d(y_k, z_k) \geq \epsilon$. We can assume that $y_k \neq x$ for all k . In particular $y_k = f_{n_k}(b_k)$ for some $b_k \in K$ and by compactness we can assume that $b_{n_k} \rightarrow b \in K$. Up to passing to a subsequence, we can assume that either $z_k = x$ for all k or $z_k \neq x$ for all k . In the first case, one has $d(y_k, z_k) = d(f_{n_k}(b_k), x) \rightarrow d(f(b), x) = 0$ by uniform convergence of $f_n \rightarrow f$ on K . In the second case, one has $z_k = f_{n_k}(c_k)$ for some $c_k \in K$ and again we can assume that $c_k \rightarrow c \in K$, so that $d(y_k, z_k) = d(f_{n_k}(b_k), f_{n_k}(c_k)) \rightarrow d(f(b), f(c)) = 0$ for the same reason. Both cases lead to a contradiction, which concludes the proof of (1).

We now prove (2). Assume it does not hold, then there exists a strictly increasing sequence $(n_k)_{k \in \mathbb{N}}$ in \mathbb{N} and points $b_k \in K$, $v_k \in \partial\bar{V}$ such that $d(f_{n_k}(b_k), f_{n_k}(v_k)) < \frac{1}{k}$ for all $k \in \mathbb{N}$. By compactness of both K and $\partial\bar{V}$, we can assume that $b_k \rightarrow b \in K$ and $v_k \rightarrow v \in \partial\bar{V}$. Then $d(f_{n_k}(b_k), f_{n_k}(v_k)) \rightarrow d(f(b), f(v)) = 0$ by uniform convergence of $f_n \rightarrow f$ on \bar{V} and by construction. Thus $x = f(b) = f(v)$, hence $v \in f^{-1}(x) = K$, which is a contradiction since $\partial\bar{K} \cap K = \emptyset$. This concludes the proof of (2).

With (1) and (2) in hands, we can now conclude the proof. Let $B' \subset N'$ be the closed ball around x of radius $\frac{\epsilon_0}{3}$, where ϵ_0 is the one given by (2). By (1), there exists $n_0 \in \mathbb{N}$ large enough such that $f_{n_0}(K) \subseteq \text{int}(B')$. By (2), we have $f_{n_0}(\partial\bar{V}) \cap B' = \emptyset$. Therefore, $B' \subseteq f_{n_0}(V)$ and, since f_{n_0} is a homeomorphism, the preimage $B := f_{n_0}^{-1}(B')$ is the required closed n -ball around K and contained in U . This concludes the proof that K is cellular. \square

In general, a cellular subset $F \subset N$ can be quite pathological and far from being contractible [Dav86, Figure 9-1]. For example, the *topologist's sine curve* is cellular but not path-connected. Under suitable regularity conditions, however, a cellular subset is contractible. Actually, it is the case when F is an absolute neighborhood retract (ANR). This follows from Corollaries 2B and 3A in [Dav86, Section 14], but we give a direct proof for the reader's convenience.

Lemma 4.7. *Let F be a cellular subset of a manifold N . If F is an absolute neighborhood retract (ANR), then F is contractible.*

Proof. Let us suppose that F is an ANR cellular subset of N . To prove that F is contractible, we show that the identity map $\text{id}_F: F \rightarrow F$ is homotopic (inside F) to a constant map. Since F is an ANR, it is a neighborhood retract in N , i.e. there is a neighborhood $U \subset N$ of F such that the inclusion $i: F \hookrightarrow U$ has a retraction $r: U \rightarrow F$. Now, by cellularity of F , there is a closed n -ball B such that $F \subset \text{int}(B) \subset B \subset U$. Since B is contractible, there is a homotopy $H: F \times [0, 1] \rightarrow B$ from the inclusion $j: F \hookrightarrow B$ to a constant map. Then the composition

$$r|_B \circ H: F \times [0, 1] \xrightarrow{H} B \xrightarrow{r|_B} F$$

is a homotopy from $\text{id}_F = r|_B \circ j$ to a constant map, as required. \square

The last ingredient of our argument is the following lemma, which follows directly from Corollary 8A in [Dav86, Section 14] combined with Sullivan's local characterization of real analytic subsets [Sul70].

Lemma 4.8. *A compact real analytic subset of \mathbb{R}^n is an ANR.*

Proof. Let $F \subset \mathbb{R}^n$ be a compact real analytic subset. Being an ANR is a local property for metrizable spaces [Han51, Theorem 3.2], which F is (as a subset of \mathbb{R}^n). By Sullivan's result [Sul70], F is locally homeomorphic to a cone over a polyhedron, hence locally an ANR. For example, this follows from [Han51, Corollary 3.5]. \square

Combining all the above sections, we can now provide a proof of Lemma 1.5.

Lemma 4.9. *The fibres of $\psi: \mathcal{CC}(M) \rightarrow \mathcal{ML}_{\partial M}^{\text{realizable}}$ are contractible.*

Proof. For each $K \in (-1, 0)$, we introduce the function $U_K: \mathcal{T}_{\partial\bar{M}}^K \rightarrow \mathcal{ML}_{\partial\bar{M}}$, defined by applying u_K for each boundary component of $\partial\bar{M}$.

It follows directly from Lemma 1.8 and from Lemma 4.3 that for any closed curve c on $\partial\bar{M}$, $i(U_K \circ \psi_K(-), c) \rightarrow i(\psi(-), c)$ uniformly on compact subsets of $\mathcal{CC}(M)$ as $K \rightarrow -1$.

Choose a finite set of closed curves c_1, \dots, c_n in $\partial\bar{M}$ such that the function $l \mapsto (i(c_1, l), i(c_2, l), \dots, i(c_n, l))$ is a topological embedding of $\mathcal{ML}_{\partial\bar{M}}$ in \mathbb{R}^n . We can use those functions to define a distance D on $\mathcal{ML}_{\partial\bar{M}}$,

for instance by

$$D(l, l') = \sum_{i=1}^n |i(c_i, l') - i(c_i, l)| .$$

Considering this distance D on $\mathcal{ML}_{\partial\bar{M}}$ it follows from the previous paragraph that $U_K \circ \psi_K$ converges uniformly to ψ on compact subsets of $\mathcal{CC}(M)$. It then follows from Lemma 3.1, Theorem 4.6, Lemma 4.7 and Lemma 4.8 that the fibres of ψ are contractible. \square

5. NON-CONTRACTIBILITY

We provide in this section the proof of Lemma 1.6. It follows quite directly from a result of Sullivan [Sul70].¹

Sullivan [Sul70, Corollary 2] notices that a subset $X \in \mathbb{R}^n$ which is defined by real analytic equations can be triangulated, and that it is locally the cone over a polyhedron of even Euler characteristic.

The following statement follows directly.

Lemma 5.1. *A compact k -dimensional real analytic subset F of \mathbb{R}^n is a mod 2 pseudo-manifold, i.e. $H_k(F, \mathbb{Z}_2) \neq 0$. In particular, if it is contractible then it must be a point.*

Proof. As remarked by Sullivan in [Sul70] below Corollary 2, the sum of all k -dimensional simplices in a triangulation of F is a k -cycle mod 2. Since F has dimension k , this cycle is not a boundary, hence it provides a nonzero element of $H_k(F, \mathbb{Z}_2)$. \square

The proof of Lemma 1.6 follows directly, since Lemma 3.1 asserts that for all $l \in \mathcal{ML}_{\partial M}^{\text{realizable}}$, $\psi^{-1}(\{l\})$ is a compact real analytic subvariety of $\mathcal{CC}(M)$. Since it is contractible by Lemma 1.5, it is a point.

REFERENCES

- [ABB⁺07] Lars Andersson, Thierry Barbot, Riccardo Benedetti, Francesco Bonsante, William M. Goldman, François Labourie, Kevin P. Scannell, and Jean-Marc Schlenker. Notes on: “Lorentz spacetimes of constant curvature” [Geom. Dedicata **126** (2007), 3–45; mr2328921] by G. Mess. *Geom. Dedicata*, 126:47–70, 2007.
- [Ale05] Alexander D. Alexandrov. *Convex polyhedra*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2005. Translated from the 1950 Russian edition by N. S. Dairbekov, S. S. Kutateladze and A. B. Sossinsky. With comments and bibliography by V. A. Zalgaller and appendices by L. A. Shor and Yu. A. Volkov.
- [And71] E.M. Andreev. On convex polyhedra of finite volume in Lobacevskii space. *Math. USSR Sbornik*, 12 (3):225–259, 1971.
- [BBZ11] Thierry Barbot, François Béguin, and Abdelghani Zeghib. Prescribing Gauss curvature of surfaces in 3-dimensional spacetimes, application to the Minkowski problem in Minkowski space. *Ann. Inst. Fourier (Grenoble)*, 61(1):511–591, 2011.
- [BDMS21] Francesco Bonsante, Jeffrey Danciger, Sara Maloni, and Jean-Marc Schlenker. The induced metric on the boundary of the convex hull of a quasicircle in hyperbolic and anti-de Sitter geometry. *Geom. Topol.*, 25(6):2827–2911, 2021. With an appendix by Boubacar Diallo.
- [Bel17] Mehdi Belraouti. Asymptotic behavior of cauchy hypersurfaces in constant curvature space-times. *Geometriae Dedicata*, 190(1):103–133, 2017.
- [BMS13] Francesco Bonsante, Gabriele Mondello, and Jean-Marc Schlenker. A cyclic extension of the earthquake flow I. *Geom. Topol.*, 17(1):157–234, 2013.
- [BO04] Francis Bonahon and Jean-Pierre Otal. Laminations mesurées de plissage des variétés hyperboliques de dimension 3. *Ann. Math.*, 160:1013–1055, 2004.
- [Bon88] Francis Bonahon. The geometry of teichmüller space via geodesic currents. *Inventiones mathematicae*, 92(1):139–162, 1988.
- [Bon92] Francis Bonahon. Earthquakes on riemann surfaces and on measured geodesic laminations. *Transactions of the American Mathematical Society*, 330(1):69–95, 1992.
- [Bon96] Francis Bonahon. Shearing hyperbolic surfaces, bending pleated surfaces and Thurston’s symplectic form. *Ann. Fac. Sci. Toulouse Math. (6)*, 5(2):233–297, 1996.
- [Bon05] Francis Bonahon. Kleinian groups which are almost Fuchsian. *J. Reine Angew. Math.*, 587:1–15, 2005.
- [BS12] Francesco Bonsante and Jean-Marc Schlenker. Fixed points of compositions of earthquakes. *Duke Math. J.*, 161(6):1011–1054, 2012.
- [CS22] Qiyu Chen and Jean-Marc Schlenker. The geometric data on the boundary of convex subsets of hyperbolic manifolds, 2022.
- [Dav86] Robert J Daverman. *Decompositions of manifolds*. Academic Press, 1986.
- [Dum08] David Dumas. Complex projective structures. In *Handbook of Teichmüller theory. Vol. II*, volume 13 of *IRMA Lect. Math. Theor. Phys.*, pages 455–508. Eur. Math. Soc., Zürich, 2008.
- [Fil07] François Fillastre. Polyhedral realisation of hyperbolic metrics with conical singularities on compact surfaces. *Ann. Inst. Fourier (Grenoble)*, 57(1):163–195, 2007.

¹This argument was mentioned by user “Moishe Kohan” (<https://mathoverflow.net/users/39654/moishe-kohan>) in a post on mathOverflow, see <https://mathoverflow.net/q/446497> (version: 2023-05-10).

- [Fil11] François Fillastre. Fuchsian polyhedra in Lorentzian space-forms. *Math. Ann.*, 350(2):417–453, 2011.
- [Fin67] Ross L Finney. Pseudo-isotopies and cellular sets. *Michigan Mathematical Journal*, 14(4):417–421, 1967.
- [FS19] François Fillastre and Andrea Seppi. Spherical, hyperbolic, and other projective geometries: convexity, duality, transitions. In *Eighteen essays in non-Euclidean geometry*, volume 29 of *IRMA Lect. Math. Theor. Phys.*, pages 321–409. Eur. Math. Soc., Zürich, 2019.
- [Han51] Olof Hanner. Some theorems on absolute neighborhood retracts. *Arkiv för Matematik*, 1(5):389–408, 1951.
- [HR93] Craig D. Hodgson and Igor Rivin. A characterization of compact convex polyhedra in hyperbolic 3-space. *Invent. Math.*, 111:77–111, 1993.
- [HS23] Sebastian Hensel and Jenya Sapir. A projection from filling currents to teichmüller space. *Proceedings of the American Mathematical Society*, 151(08):3621–3633, 2023.
- [Ker92] Steven P Kerckhoff. Lines of minima in teichmüller space. *Duke Math. J.*, 68(1):187–213, 1992.
- [Lab91] François Labourie. Problème de Minkowski et surfaces à courbure constante dans les variétés hyperboliques. *Bull. Soc. Math. France*, 119(3):307–325, 1991.
- [Lab92a] François Labourie. Métriques prescrites sur le bord des variétés hyperboliques de dimension 3. *J. Differential Geom.*, 35:609–626, 1992.
- [Lab92b] François Labourie. Surfaces convexes dans l’espace hyperbolique et CP1-structures. *J. London Math. Soc., II. Ser.*, 45:549–565, 1992.
- [Lec04] Cyril Lecuire. Bending map and strong convergence. Available on <https://lecuire.perso.math.cnrs.fr/>, 2004.
- [Lec06] Cyril Lecuire. Plissage des variétés hyperboliques de dimension 3. *Invent. Math.*, 164(1):85–141, 2006.
- [LS00] François Labourie and Jean-Marc Schlenker. Surfaces convexes fuchsiennees dans les espaces lorentziens à courbure constante. *Math. Ann.*, 316(3):465–483, 2000.
- [Mar16] Albert Marden. *Hyperbolic manifolds*. Cambridge University Press, Cambridge, 2016. An introduction in 2 and 3 dimensions.
- [Maz22] Filippo Mazzoli. The dual volume of quasi-Fuchsian manifolds and the Weil-Petersson distance. *Trans. Amer. Math. Soc.*, 375(1):695–723, 2022.
- [Mes07] Geoffrey Mess. Lorentz spacetimes of constant curvature. *Geom. Dedicata*, 126:3–45, 2007.
- [Mes23] Abderrahim Mesbah. The induced metric and bending lamination on the boundary of convex hyperbolic 3-manifolds, 2023.
- [Pog73] Aleksei V. Pogorelov. *Extrinsic Geometry of Convex Surfaces*. American Mathematical Society, 1973. Translations of Mathematical Monographs. Vol. 35.
- [Pro22a] Roman Prosanov. Dual metrics on the boundary of strictly polyhedral hyperbolic 3-manifolds, 2022. arxiv:2203.16971.
- [Pro22b] Roman Prosanov. Hyperbolic 3-manifolds with boundary of polyhedral type, 2022. arxiv:2210.17271.
- [RH93] Igor Rivin and Craig D. Hodgson. A characterization of compact convex polyhedra in hyperbolic 3-space. *Invent. Math.*, 111:77–111, 1993.
- [Riv96] Igor Rivin. A characterization of ideal polyhedra in hyperbolic 3-space. *Annals of Math.*, 143:51–70, 1996.
- [Sch98] Jean-Marc Schlenker. Métriques sur les polyèdres hyperboliques convexes. *J. Differential Geom.*, 48(2):323–405, 1998.
- [Sch01] Jean-Marc Schlenker. Hyperbolic manifolds with polyhedral boundary. math.GT/0111136, 2001.
- [Sch06] Jean-Marc Schlenker. Hyperbolic manifolds with convex boundary. *Invent. Math.*, 163(1):109–169, 2006.
- [Sch20] Jean-Marc Schlenker. On the weyl problem for complete surfaces in the hyperbolic and anti-de sitter spaces, 2020.
- [Sch21] Jean-Marc Schlenker. The Weyl problem for unbounded convex domains in \mathbb{H}^3 . *arXiv preprint arXiv:2106.02101*, 2021.
- [Ser06] C Series. Thurston’s bending measure conjecture for once punctured torus groups. *Spaces of Kleinian groups*, 329:75, 2006.
- [Slu18] Dmitriy Slutskiy. Compact domains with prescribed convex boundary metrics in quasi-fuchsian manifolds. *Bull. Soc. Math. France*, 146(2):309–353, 2018.
- [Sul70] D Sullivan. Combinatorial invariants of analytic spaces. *Lecture Notes in Math.*, 192:165–168, 1970.
- [Sul71] D. Sullivan. Combinatorial invariants of analytic spaces. In *Proceedings of Liverpool Singularities—Symposium, I (1969/70)*, volume Vol. 192 of *Lecture Notes in Math.*, pages 165–168. Springer, Berlin-New York, 1971.
- [Thu80] William P. Thurston. Three-dimensional geometry and topology. Originally notes of lectures at Princeton University, 1979. Recent version available on <http://www.msri.org/publications/books/gt3m/>, 1980.
- [Thu86] William P. Thurston. Earthquakes in two-dimensional hyperbolic geometry. In *Low-dimensional topology and Kleinian groups (Coventry/Durham, 1984)*, volume 112 of *London Math. Soc. Lecture Note Ser.*, pages 91–112. Cambridge Univ. Press, Cambridge, 1986.
- [Wol06] S Wolpert. Convexity of geodesic-length functions: a reprise. *Spaces of Kleinian groups*, 329:233–246, 2006.

BRUNO DULAR: UNIVERSITY OF LUXEMBOURG, FSTM, DEPARTMENT OF MATHEMATICS, MAISON DU NOMBRE, 6 AVENUE DE LA FONTE, L-4364 ESCH-SUR-ALZETTE, LUXEMBOURG
Email address: `bruno.dular@uni.lu`

JEAN-MARC SCHLENKER: UNIVERSITY OF LUXEMBOURG, FSTM, DEPARTMENT OF MATHEMATICS, MAISON DU NOMBRE, 6 AVENUE DE LA FONTE, L-4364 ESCH-SUR-ALZETTE, LUXEMBOURG
Email address: `jean-marc.schlenker@uni.lu`