

Second-Order Strong Optimality and Second-Order Duality for Nonsmooth Constrained Multiobjective Fractional Programming Problems

Jiawei Chen · Luyu Liu · Yibing Lv · Debdas Ghosh · Jen-Chih Yao

Received: date / Accepted: date

Abstract This paper investigates constrained *nonsmooth multiobjective fractional programming problem* (NMFP) in real Banach spaces. It derives a quotient calculus rule for computing the first- and second-order Clarke derivatives of fractional functions involving locally Lipschitz functions. A novel second-order Abadie-type regularity condition is presented, defined with the help of the Clarke directional derivative and the Páles-Zeidan second-order directional derivative. We establish both first- and second-order strong necessary optimality conditions, which contain some new information on multipliers and imply the strong KKT necessary conditions, for a Borwein-type properly efficient solution of NMFP by utilizing generalized directional derivatives. Moreover, it derives second-order sufficient optimality conditions for NMFP under a second-order generalized convexity assumption. Additionally, we derive duality results between NMFP and its second-order dual problem under some appropriate conditions.

Keywords Multiobjective fractional programming · Second-order optimality conditions · Borwein-type properly efficient solution · Second-order Abadie-type regular condition · Mond-Weir duality

Mathematics Subject Classification (2000) 90C46 · 90C29 · 49J52

1 Introduction

Multiobjective fractional programming (MFP) represents a crucial model in operations research, finding widespread applications in fields such as computer vision, portfolio opti-

Jiawei Chen · Luyu Liu
School of Mathematics and Statistics, Southwest University, Chongqing 400715, China
Emails: J.W.Chen713@163.com (J. Chen), lyliu124@163.com (L. Liu)

Yibing Lv
School of Information and Mathematics, Yangtze University, Jingzhou 434023, China.
Email: Yibinglv@yangtzeu.edu.cn

Debdas Ghosh
Department of Mathematical Sciences, Indian Institute of Technology (BHU), Varanasi, Uttar Pradesh 221005, India
Email: debdas.mat@iitbhu.ac.in

Jen-Chih Yao
Research Center for Interneural Computing, China Medical University, Taichung 40447, Taiwan
Email: yaojc@mail.cmu.edu.tw

mization, management science, image processing, and communications. Various works have investigated and utilized MFP problems, including those in [1, 2, 3, 4, 5, 6, 7, 8, 9, 10]. Due to the nonsmooth nature of the functions involved in realistic MFP problems, practical challenges emerged in fields like economics, decision theory, optimal control, scheduling, machine learning, engineering, and game theory. As a result, MFPs have been extensively examined using subdifferentials and directional derivatives, as evident from the studies in [11, 12, 13, 14, 15, 16]. Within the optimization theory of MFP, duality theory and optimality conditions constitute fundamental areas of study. Recent research has been attempting to identify both first- and second-order characterizations for efficient points of MFP problems, leveraging a range of first- and second-order directional derivatives.

First- and second-order characterizations for a local weak efficient solution of NMFP were derived in [17] by generalized derivatives as the first- and second-order approximations of the involving functions. Khanh and Tung [18] investigated the *first-order* Karush-Kuhn-Tucker (KKT) conditions for a local Borwein-type proper efficient solution of nonsmooth semi-infinite multiobjective programming by a Mangasarian-Fromovitz regularity condition. Su and Hang [19] applied Hadamard directional derivative to establish a first-order quotient rule and derived first-order conditions for local weak efficient solutions of NMFP. They also derived, in [20], second-order conditions for strict local efficient solutions of MFP by utilizing second-order Páles-Zeidan-type upper directional derivative. The duality results and first-order sufficient optimality conditions for single-objective fractional programming problems were obtained under the (F, α, ρ, d) -convexity assumption in [21]. Further, in [22], the results proposed in [21] were extended for smooth multiobjective fractional programming problems under a convexity assumption. In [23], a notion of (C, α, ρ, d) -convexity was introduced and applied to investigate the first-order sufficient optimality conditions and duality for nonsmooth minimax fractional programming problems. The first-order optimality conditions and duality for nondifferentiable MFP problems were also explored in [24]. Very recently, in [25], duality results and optimality conditions for E -minimax fractional programming were investigated assuming the E -invexity of the involved functions, and applied to multiobjective optimization.

To the best of the authors' knowledge, only very few results exist concerning the second-order duality and second-order strong necessary optimality conditions for NMFPs. This study focuses specifically on the realm of Borwein-type proper efficient solutions for NMFP problems, which are known to be more potent than (weak) efficient solutions. By leveraging the Clarke directional derivative and second-order Páles-Zeidan generalized directional derivative, we aim to explore second-order characterizations of Borwein-type proper efficient solutions and report second-order Mond-Weir duality theory for NMFP problems involving both equality and inequality constraints. Notably, our investigation considers the locally Lipschitz continuous functions rather than those that are Fréchet differentiable.

Towards deriving second-order duality results and strong optimality conditions, we propose a *generalized second-order Abadie-type regularity condition* (GSOARC) with the help of Páles-Zeidan second-order generalized directional derivatives, which do not require the continuous Fréchet differentiability and existence of second-order directional derivatives of all involved functions that were a requirement in [26]. The first- and second-order strong KKT optimality conditions for Borwein proper efficient solutions of NMFP problems are reported under mild conditions. We show that the GSOARC cannot be relaxed to the generalized second-order Guignard-type regularity condition in obtaining the derived second-order strong optimality conditions. Furthermore, sufficient optimality conditions for NMFP are

derived under a generalized second-order convexity assumption. Additionally, the Mond-Weir-type second-order duality results for NMFP are derived. The weak, strong, and converse duality results between NMFP and its second-order dual problem are identified under some suitable conditions.

The rest of the paper is demonstrated as follows. In Section 2, we recollect some basic notions and present quotient calculus for locally Lipschitz functions. We introduce a GSOARC and investigate the first- and second-order strong KKT conditions for a Borwein-type properly efficient solution of NMFP in Section 3. Subsequently, second-order sufficient optimality conditions are presented in Section 4 under some second-order generalized convexity assumptions. In Section 5, the second-order Mond-Weir-type dual problem is introduced. We also study duality results between NMFP and the corresponding second-order dual problem. Finally, we give the conclusions in Section 6.

2 Preliminaries

Let Z be a real reflexive Banach space with the norm $\|\cdot\|$, and U be a nonempty open subset of Z . For $A \subseteq Z$, its closure, topological interior, and convex hull are denoted by $\text{cl } A$, $\text{int } A$, and $\text{co } A$, respectively. Let Z^* be the topological dual space of Z , and $\langle \cdot, \cdot \rangle$ denote the coupling between Z^* and Z . We denote the nonnegative orthant and positive orthant of \mathbb{R}^n by \mathbb{R}_+^n and \mathbb{R}_{++}^n , respectively. Let

$$\begin{aligned} F &:= (F_1, F_2, \dots, F_p)^\top : Z \rightarrow \mathbb{R}^p, \\ f &:= (f_1, f_2, \dots, f_p)^\top : Z \rightarrow \mathbb{R}^p, \\ g &:= (g_1, g_2, \dots, g_m)^\top : Z \rightarrow \mathbb{R}^m \\ \text{and } h &:= (h_1, h_2, \dots, h_l)^\top : Z \rightarrow \mathbb{R}^l \end{aligned}$$

be vector-valued functions, and let $F(x) \in \mathbb{R}_{++}^p$ and $\frac{f(x)}{F(x)} := \left(\frac{f_1(x)}{F_1(x)}, \frac{f_2(x)}{F_2(x)}, \dots, \frac{f_p(x)}{F_p(x)} \right)^\top$ for all $x \in U$, where the superscript \top denotes the transpose. For $x = (x_1, x_2, \dots, x_p)^\top$ and $y = (y_1, y_2, \dots, y_p)^\top \in \mathbb{R}^p$, we undertake the following conventional notations:

$$\begin{aligned} x = y &\iff x_i = y_i, i = 1, 2, \dots, p, \\ x \leq y &\iff x - y \in -\mathbb{R}_+^p, \\ x \leq y &\iff x - y \in -\mathbb{R}_+^p \setminus \{0\}, \\ x < y &\iff x - y \in -\mathbb{R}_{++}^p, \\ z = x * y &\iff z = (x_1 y_1, x_2 y_2, \dots, x_p y_p)^\top \\ \text{and } w = x^\top y &\iff w = x_1 y_1 + x_2 y_2 + \dots + x_p y_p. \end{aligned}$$

The relations $y \geq x$, $y \geq x$, and $y > x$ simply mean $x \leq y$, $x \leq y$ and $x < y$, respectively. For brevity, we use the following notations:

$$I := \{1, 2, \dots, p\}, J := \{1, 2, \dots, m\}, K := \{1, 2, \dots, l\}$$

and for each $x \in U$, we denote

$$\begin{aligned} \left(\frac{f}{F} \right) (x) &:= \frac{f(x)}{F(x)}, \left(\frac{f_i}{F_i} \right) (x) := \frac{f_i(x)}{F_i(x)}, i \in I \\ \text{and } \left(\frac{1}{F} \right) (x) &:= \left(\frac{1}{F_1(x)}, \frac{1}{F_2(x)}, \dots, \frac{1}{F_p(x)} \right)^\top. \end{aligned}$$

In this study, we focus on the *first- and second-order Borwein proper optimality* and *second-order Mond-Weir duality* of the following nonsmooth multiobjective fractional programming problem:

$$(NMFP) \quad \begin{cases} \min & \frac{f(x)}{F(x)} \\ \text{s.t.} & g(x) \leq 0, h(x) = 0, x \in U. \end{cases} \quad (1)$$

The set of all feasible points of NMFP is denoted by X , i.e.,

$$X := \{x \in U : g(x) \leq 0, h(x) = 0\}. \quad (2)$$

We also consider the following parametric problem associated with NMFP:

$$(s\text{-MFP}) \quad \begin{cases} \min & (f - s * F)(x) \\ \text{s.t.} & g(x) \leq 0, h(x) = 0, x \in U, \end{cases} \quad (3)$$

where $s := (s_1, s_2, \dots, s_p) \in \mathbb{R}^p$ and $(f - s * F)(x) = f(x) - s * F(x)$.

We next recall some standard definitions [19] and well-known results, which will be useful later.

Definition 2.1 An element $x_0 \in X$ is said to be

(i) a *Pareto efficient* solution of NMFP iff, there is no $x \in X$ such that

$$\left(\frac{f}{F}\right)(x) \leq \left(\frac{f}{F}\right)(x_0).$$

(ii) *weak efficient solution* of NMFP iff, there is no $x \in X$ such that

$$\left(\frac{f}{F}\right)(x) < \left(\frac{f}{F}\right)(x_0).$$

Similarly, one can define (weak) Pareto efficient solutions of s -MFP by substituting $(f - s * F)$ for $\left(\frac{f}{F}\right)$.

The following result shows an equivalence between the (weak) efficient solutions of NMFP and s -MFP.

Lemma 2.1 A feasible solution $x_0 \in X$ is a (weak) Pareto efficient solution of NMFP if and only if it is a (weak) Pareto efficient solution of s -MFP, where $s := \frac{f(x_0)}{F(x_0)}$.

Proof A proof for x_0 being a weak Pareto point is given in [19, Proposition 1]. For a proof of x_0 being a Pareto efficient point, we observe for any $x \in U$ with $x \neq x_0$ that

$$\begin{aligned} & \left(\frac{f}{F}\right)(x) - \left(\frac{f}{F}\right)(x_0) \notin -\mathbb{R}_+^p \setminus \{0\} \\ \iff & \left(\frac{f}{F}\right)(x) - s \notin -\mathbb{R}_+^p \setminus \{0\} \\ \iff & \frac{1}{F(x)}(f(x) - s * F(x)) \notin -\mathbb{R}_+^p \setminus \{0\} \\ \iff & f(x) - s * F(x) \notin -\mathbb{R}_+^p \setminus \{0\} \text{ because } \mathbb{R}_+^p \text{ is a cone and } F(x) \in \mathbb{R}_{++}^p \\ \iff & (f - s * F)(x) \not\leq (f - s * F)(x_0). \end{aligned}$$

Hence, the result follows. The proof is completed.

Definition 2.2 [27] A function $\vartheta : U \rightarrow \mathbb{R}$ is said to be *Gâteaux differentiable* at $x_0 \in U$ in a direction $v_0 \in Z$ iff, the limit

$$\lim_{t \rightarrow 0^+} \frac{\vartheta(x_0 + tv_0) - \vartheta(x_0)}{t} \quad (4)$$

exists. If the limit exists, it is denoted by $\vartheta'(x_0; v_0)$ and called the Gâteaux derivative of ϑ at x_0 along the direction v_0 .

Let $x_0 \in U$. If $\vartheta'(x_0; v)$ exists for any $v \in Z$ and there is a continuous linear function $\vartheta'_G(x_0) : Z \rightarrow \mathbb{R}$ such that

$$\vartheta'(x_0; v) = \langle \vartheta'_G(x_0), v \rangle \quad \text{for each } v \in Z,$$

then we say that $\vartheta'_G(x_0)$ is the Gâteaux derivative of ϑ at x_0 .

A vector-valued function $\vartheta := (\vartheta_1, \vartheta_2, \dots, \vartheta_p)^\top : U \rightarrow \mathbb{R}^p$ is called Gâteaux differentiable at $x_0 \in U$ if its components $\vartheta_i, i \in I$, are Gâteaux differentiable at x_0 .

Remark 2.1 (i) In some instances, as discussed in [5], the limit described in (4) can be alternatively expressed with $t \rightarrow 0$. However, for consistency with other definitions of directional derivatives, we use the one-sided limit in this paper.

- (ii) It cannot be generally concluded that ϑ is continuous at x even if ϑ is Gâteaux differentiable at x .
- (iii) If ϑ is Fréchet differentiable at x_0 , then ϑ is Gâteaux differentiable at x_0 and $\nabla\vartheta(x_0) = \vartheta'_G(x_0)$, where $\nabla\vartheta(x_0)$ is the Fréchet derivative of ϑ at x_0 .

Definition 2.3 [27] Let $\vartheta : U \rightarrow \mathbb{R}$ be a real-valued function and $x_0 \in U$.

- (i) The (Clarke) *generalized directional derivative* of ϑ at $x_0 \in U$ in the direction $v_0 \in Z$ is defined by

$$\vartheta^\circ(x_0; v_0) := \limsup_{y \rightarrow x_0, t \rightarrow 0^+} \frac{\vartheta(y_0 + tv_0) - \vartheta(y_0)}{t}.$$

- (ii) ϑ is said to be *regular* in the sense of Clarke at x_0 if $\vartheta'(x_0; v)$ exists and $\vartheta'(x_0; v) = \vartheta^\circ(x_0; v)$ holds for all $v \in Z$.

For any given $x_0 \in X$, with regard to the problem (1), we denote

$$J(x_0) := \{j \in J : g_j(x_0) = 0\}, \quad J(x_0, v) := \{j \in J(x_0) : g_j^\circ(x_0; v) = 0\},$$

and

$$Q := \{x \in Z : g(x) \leq 0, h(x) = 0, f(x) \leq_C s * F(x)\}, \quad (5)$$

where $s = \left(\frac{f}{F}\right)(x_0)$ and $g_j^\circ(x_0; v)$ is the Clarke directional derivative (Definition 2.3) of g_j at x_0 in the direction $v \in Z$.

Definition 2.4 [5, 27] Let X and Q be as in (2) and (5), respectively, and $X \neq \emptyset$.

- (i) The *contingent cone* of X at $x_0 \in X$ is defined by

$$T(X, x_0) := \{d \in Z : \exists t_k \rightarrow 0^+, \exists d_k \rightarrow d \text{ such that } x_0 + t_k d_k \in X, \forall k \in \mathbb{N}\}.$$

(ii) The *linearizing cone* of Q at $x_0 \in X$ is defined by

$$C(Q, x_0) := \left\{ d \in Z : \begin{cases} \left(\frac{f}{F}\right)_i^\circ(x_0; d) \leq 0, & i \in I, \\ g_j^\circ(x_0; d) \leq 0, & j \in J(x_0), \\ h_k^\circ(x_0; d) = 0, & k \in K \end{cases} \right\}.$$

The set $T(X, x_0)$ is a nonempty and closed cone with $0 \in T(X, x_0)$. It is well-known that if $x_0 \in \text{cl}X_1 \subseteq \text{cl}X_2 \subseteq X$, then $T(X_1, x_0) \subseteq T(X_2, x_0)$; see e.g., [5].

Definition 2.5 A vector $v \in Z$ is called a *critical point* at $x_0 \in X$ iff

$$v \in T(X, x_0) \cap C(Q, x_0).$$

We denote the *set of all critical points* at $x_0 \in X$ by $D(x_0)$, i.e.,

$$D(x_0) := \{v \in Z : v \in T(X, x_0) \cap C(Q, x_0)\}.$$

Definition 2.6 [28, 29] Let $\vartheta : U \rightarrow \mathbb{R}$ be a function and $x_0 \in U$.

(i) The *second-order directional derivative* of ϑ at x_0 in the direction $v_0 \in Z$ is defined by

$$\vartheta''(x_0; v_0) := \lim_{t \rightarrow 0^+} \frac{\vartheta(x_0 + tv_0) - \vartheta(x_0) - t\vartheta'(x_0; v_0)}{\frac{1}{2}t^2}, \text{ provided limit exists.}$$

(ii) The *Páles-Zeidan second-order generalized directional derivative* of ϑ at x_0 in the direction $v_0 \in Z$ is defined by

$$\vartheta^{\circ\circ}(x_0; v_0) := \limsup_{t \rightarrow 0^+} \frac{\vartheta(x_0 + tv_0) - \vartheta(x_0) - t\vartheta^\circ(x_0; v_0)}{\frac{1}{2}t^2}.$$

Remark 2.2 [29]

(i) If ϑ is twice Fréchet differentiable at x_0 , then

$$\langle \nabla^2 \vartheta(x_0)v, v \rangle = \vartheta''(x_0; v) = \vartheta^{\circ\circ}(x_0; v), \forall v \in Z.$$

(ii) If $\vartheta'(x_0; v_0) = \vartheta^\circ(x_0; v_0)$ and $\vartheta''(x_0; v_0)$ exists, then $\vartheta''(x_0; v_0) = \vartheta^{\circ\circ}(x_0; v_0)$. Besides, for any $\beta > 0$, $(\beta\vartheta)'(x_0; v_0) = \beta\vartheta'(x_0; v_0)$, $(\beta\vartheta)^\circ(x_0; v_0) = \beta\vartheta^\circ(x_0; v_0)$, $(\beta\vartheta)''(x_0; v_0) = \beta\vartheta''(x_0; v_0)$ and $(\beta\vartheta)^{\circ\circ}(x_0; v_0) = \beta\vartheta^{\circ\circ}(x_0; v_0)$.

Definition 2.7 [27] A function $\vartheta : U \rightarrow \mathbb{R}$ is called *locally Lipschitz continuous* at $x_0 \in U$ if there exists a neighborhood $V(x_0)$ and a positive constant $L(x_0)$ such that

$$|\vartheta(y) - \vartheta(z)| \leq L(x_0)\|y - z\|, \forall y, z \in V(x_0) \cap U.$$

The function ϑ is called *locally Lipschitz continuous on U* if it is locally Lipschitz continuous at every $x \in U$. In particular, if the positive constant $L(x)$ is independent of $x \in U$, then ϑ is called *Lipschitz continuous on U* .

Definition 2.8 [27] Let $\vartheta : U \rightarrow \mathbb{R}$ be locally Lipschitz continuous on U . The *Clarke subdifferential* of ϑ at $x_0 \in U$ is defined by

$$\partial\vartheta(x_0) := \{\xi \in Z^* : \langle \xi, v \rangle \leq \vartheta^\circ(x_0; v), \forall v \in Z\}. \quad (6)$$

Lemma 2.2 [30] *Let $\vartheta : U \rightarrow \mathbb{R}$ be locally Lipschitz continuous at $x_0 \in U$. Then there exists a constant $L_0 > 0$ such that $\|\xi\| \leq L_0$ for arbitrary $\xi \in \partial\vartheta(x_0)$.*

Remark 2.3 (i) The existence of $\vartheta^\circ(x_0; v_0)$ does not necessarily imply the existence of $\vartheta'(x_0; v_0)$, and even if $\vartheta'(x_0; v_0)$ exists, they may not be equal. However, if ϑ is a continuously Fréchet differentiable function at x_0 and the Fréchet derivative is $\nabla\vartheta(x_0)$, then it is regular in the sense of Clarke, i.e.,

$$\vartheta'(x_0; v_0) = \vartheta^\circ(x_0; v_0) = \langle \nabla\vartheta(x_0), v_0 \rangle = \langle \vartheta'_G(x_0), v_0 \rangle.$$

(ii) If ϑ is locally Lipschitz continuous on U , then $\vartheta^\circ(x; v)$ exists (finite) for any $x \in U$ and $v \in Z$, which implies that $\partial\vartheta(x)$ is compact. Besides, if $\vartheta_1, \vartheta_2 : U \rightarrow \mathbb{R}$ are locally Lipschitz continuous and regular in the sense of Clarke at $x_0 \in U$ and $\vartheta_2(x) \neq 0$ for all $x \in U$, then $\frac{\vartheta_1}{\vartheta_2} : U \rightarrow \mathbb{R}$ is regular in the sense of Clarke at $x_0 \in U$, i.e., for each $v \in Z$,

$$\left(\frac{\vartheta_1}{\vartheta_2} \right)^\circ (x_0; v) = \left(\frac{\vartheta_1}{\vartheta_2} \right)' (x_0; v) = \frac{\vartheta_1'(x_0; v)}{\vartheta_2(x_0)} - \frac{\vartheta_1(x_0)}{\vartheta_2^2(x_0)} \vartheta_2'(x_0; v).$$

Next, before we end the section, we present chain rules of quotient functions involving locally Lipschitz functions in terms of generalized directional derivatives.

Proposition 2.1 *Let ϑ_1 and ϑ_2 be two real-valued locally Lipschitz continuous functions on U with ϑ_2 being positive-valued. For any $x_0 \in U$, $v \in Z$ and $\beta \in \mathbb{R}_+$, the following are true:*

- (i) $\vartheta_1 - \beta\vartheta_2$ and $\frac{\vartheta_1}{\vartheta_2}$ are locally Lipschitz continuous on U ;
If, in addition, ϑ_2 is Gâteaux differentiable at x_0 with Gâteaux derivative $\vartheta'_{2,G}(x_0)$, then
- (ii) $\left(\frac{\vartheta_1}{\vartheta_2} \right)^\circ (x_0; v) = \frac{1}{\vartheta_2(x_0)} \left[\vartheta_1^\circ(x_0; v) - \frac{\vartheta_1(x_0)}{\vartheta_2(x_0)} \langle \vartheta'_{2,G}(x_0), v \rangle \right]$;
- (iii) $(\vartheta_1 - \beta\vartheta_2)^\circ(x_0; v) = \vartheta_1^\circ(x_0; v) - \beta \langle \vartheta'_{2,G}(x_0), v \rangle$;
Moreover, if $\vartheta_2''(x_0; v)$ exists, then
- (iv) $\left(\frac{\vartheta_1}{\vartheta_2} \right)^{\circ\circ} (x_0; v) = \frac{1}{\vartheta_2(x_0)} \left[\vartheta_1^{\circ\circ}(x_0; v) - \frac{\vartheta_1(x_0)}{\vartheta_2(x_0)} \vartheta_2''(x_0; v) \right]$;
- (v) $(\vartheta_1 - \beta\vartheta_2)^{\circ\circ}(x_0; v) = \vartheta_1^{\circ\circ}(x_0; v) - \beta \vartheta_2''(x_0; v)$.

Proof (i) By calculation, one has

$$\begin{aligned} \|(\vartheta_1 - \beta\vartheta_2)(x) - (\vartheta_1 - \beta\vartheta_2)(z)\| &= \|\vartheta_1(x) - \vartheta_1(z) - \beta\vartheta_2(x) + \beta\vartheta_2(z)\| \\ &\leq \|\vartheta_1(x) - \vartheta_1(z)\| + \beta\|\vartheta_2(x) - \vartheta_2(z)\|, \end{aligned}$$

and

$$\begin{aligned} \left\| \left(\frac{\vartheta_1}{\vartheta_2} \right)(x) - \left(\frac{\vartheta_1}{\vartheta_2} \right)(z) \right\| &= \left\| \frac{\vartheta_1(x) - \vartheta_1(z)}{\vartheta_2(x)} - \frac{\vartheta_1(z) [\vartheta_2(x) - \vartheta_2(z)]}{\vartheta_2(x) \vartheta_2(z)} \right\| \\ &\leq \left\| \frac{\vartheta_1(x) - \vartheta_1(z)}{\vartheta_2(x)} \right\| + \left\| \frac{\vartheta_1(z) [\vartheta_2(x) - \vartheta_2(z)]}{\vartheta_2(x) \vartheta_2(z)} \right\|. \end{aligned}$$

By virtue of Definition 2.7 and $\vartheta_2(x), \vartheta_2(z) > 0$, we obtain that $\vartheta_1 - \beta\vartheta_2$ and $\frac{\vartheta_1}{\vartheta_2}$ are locally Lipschitz continuous on U .

(ii)–(iv) The proofs are analogous to [20, Theorem 3.1 and Theorem 3.2].

(v) If $\vartheta_2''(x_0; v)$ exists, Definition 2.6 and Remark 2.2 imply that

$$\begin{aligned}
& (\vartheta_1 - \beta\vartheta_2)^{\circ\circ}(x_0; v) \\
&= \limsup_{t \rightarrow 0^+} \frac{(\vartheta_1 - \beta\vartheta_2)(x_0 + tv) - (\vartheta_1 - \beta\vartheta_2)(x_0) - t(\vartheta_1 - \beta\vartheta_2)^\circ(x_0; v)}{\frac{1}{2}t^2} \\
&= \limsup_{t \rightarrow 0^+} \left[\frac{(\vartheta_1 - \beta\vartheta_2)(x_0 + tv) - (\vartheta_1 - \beta\vartheta_2)(x_0)}{\frac{1}{2}t^2} - \frac{t(\vartheta_1 - \beta\vartheta_2)^\circ(x_0; v)}{\frac{1}{2}t^2} \right] \\
&= \limsup_{t \rightarrow 0^+} \left[\frac{(\vartheta_1 - \beta\vartheta_2)(x_0 + tv) - (\vartheta_1 - \beta\vartheta_2)(x_0)}{\frac{1}{2}t^2} - \frac{t\vartheta_1^\circ(x_0; v) - t\beta\langle \vartheta'_{2,G}(x_0), v \rangle}{\frac{1}{2}t^2} \right] \\
&= \limsup_{t \rightarrow 0^+} \frac{\vartheta_1(x_0 + tv) - \vartheta_1(x_0) - t\vartheta_1^\circ(x_0; v)}{\frac{1}{2}t^2} - \beta \lim_{t \rightarrow 0^+} \frac{\vartheta_2(x_0 + tv) - \vartheta_2(x_0) - \langle \vartheta'_{2,G}(x_0), v \rangle}{\frac{1}{2}t^2} \\
&= \vartheta_1^{\circ\circ}(x_0; v) - \beta\vartheta_2''(x_0; v).
\end{aligned}$$

The proof is completed.

Remark 2.4 In the chain rules for quotient functions involving locally Lipschitz functions presented in [20], the function ϑ_2 is taken to be strictly differentiable on U in the sense of Clarke. In comparison, Proposition 2.1 just assumed the existence of the Gâteaux derivative $\vartheta'_{2,G}(x_0)$. Although in finite-dimensional normed spaces, the Clarke strict derivative $D_s\vartheta_2(x)$ and the Gâteaux derivative $\vartheta'_{2,G}(x)$ coincide, Proposition 2.1 being in a general Banach space is a stronger result than Theorem 3.2 in [20]. Furthermore, it is worth noting that Proposition 2.1 (iv) and (v) may not be true if $\vartheta_2''(x; v)$ is replaced by $\vartheta_2^{\circ\circ}(x; v)$ (see Example 2.1).

Example 2.1 Let $\vartheta_1, \vartheta_2 : \mathbb{R} \rightarrow \mathbb{R}$ be two functions as follows: $\vartheta_1(x) = x^2 + 1$ and

$$\vartheta_2(x) = \begin{cases} x^2 \sin \frac{1}{x} + 1, & \text{if } x \neq 0, \\ 1, & \text{if } x = 0. \end{cases}$$

It is easy to verify that both ϑ_1 and ϑ_2 are locally Lipschitz continuous on the open interval $U = (-1, 1)$. Also, ϑ_2 is Gâteaux differentiable on U and its Gâteaux derivative is

$$\vartheta'_{2,G}(x) = \begin{cases} 2x \sin \frac{1}{x} - \cos \frac{1}{x}, & \text{if } x \neq 0; \\ 0, & \text{if } x = 0. \end{cases}$$

After calculation, we obtain that

$$\begin{aligned}
& \vartheta_1^\circ(0; v) = 0, \quad \vartheta_1^{\circ\circ}(0; v) = 2v^2, \quad \vartheta_2^\circ(0; v) = \vartheta_2'(0; v) = \vartheta'_{2,G}(0)v = 0, \\
& (\vartheta_1 - \vartheta_2)^\circ(0; v) = 0, \quad \vartheta_2^{\circ\circ}(0; v) = 2v^2, \quad \text{and } \vartheta_2''(0; v) \text{ does not exist.}
\end{aligned}$$

So,

$$\frac{1}{\vartheta_2(0)} \left[\vartheta_1^{\circ\circ}(0; v) - \frac{\vartheta_1(0)}{\vartheta_2(0)} \vartheta_2^{\circ\circ}(0; v) \right] = 2v^2 - 2v^2 = 0 \quad (7)$$

and $\vartheta_1^{\circ\circ}(0; v) - \vartheta_2^{\circ\circ}(0; v) = 0$. However, by Definition 2.6 (ii) and Definition 2.3 (i), one has

$$\left(\frac{\vartheta_1}{\vartheta_2} \right)^\circ(0; v) = 0, \quad \left(\frac{\vartheta_1}{\vartheta_2} \right)^{\circ\circ}(0; v) = 4v^2 \quad \text{and} \quad (\vartheta_1 - \vartheta_2)^{\circ\circ}(0; v) = 4v^2. \quad (8)$$

So, for any $v \neq 0$, (7) and (8) imply that

$$\left(\frac{\vartheta_1}{\vartheta_2}\right)^{\circ\circ}(0; v) > \frac{1}{\vartheta_2(0)} \left[\vartheta_1^{\circ\circ}(0; v) - \frac{\vartheta_1(0)}{\vartheta_2(0)} \vartheta_2^{\circ\circ}(0; v) \right]$$

and

$$(\vartheta_1 - \vartheta_2)^{\circ\circ}(0; v) > \vartheta_1^{\circ\circ}(0; v) - \vartheta_2^{\circ\circ}(0; v) = 0.$$

Therefore, $\vartheta_2''(x; v)$ cannot be replaced by $\vartheta_2^{\circ\circ}(x; v)$ in Proposition 2.1 (iv) and (v).

Lemma 2.3 *Let $\vartheta_1, \vartheta_2 : U \rightarrow \mathbb{R}$ be locally Lipschitz continuous and regular in the sense of Clarke at $x_0 \in U$, ϑ_2 be positive-valued, Gâteaux differentiable on U and $\left(\frac{\vartheta_1}{\vartheta_2}\right)^{\circ\circ}(x_0; v)$ exist in the direction $v \in Z$. Assume that $x_n \rightarrow x_0$, $t_n \rightarrow 0^+$, $r_n \rightarrow r$ and $r_n > 0$ for all $n \in \mathbb{N}$. If $w_n = \frac{x_n - x_0 - t_n v}{\frac{1}{2} r_n^{-1} t_n^2} \rightarrow w$, then*

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \frac{\left(\frac{\vartheta_1}{\vartheta_2}\right)(x_n) - \left(\frac{\vartheta_1}{\vartheta_2}\right)(x_0) - t_n \left(\frac{\vartheta_1}{\vartheta_2}\right)^{\circ}(x_0; v)}{\frac{1}{2} r_n^{-1} t_n^2} \\ & \leq \left(\frac{\vartheta_1}{\vartheta_2}\right)^{\circ}(x_0; w) + r \left(\frac{\vartheta_1}{\vartheta_2}\right)^{\circ\circ}(x_0; v) \\ & = \left(\frac{\vartheta_1}{\vartheta_2}\right)'(x_0; w) + r \left(\frac{\vartheta_1}{\vartheta_2}\right)^{\circ\circ}(x_0; v). \end{aligned}$$

Further, if $\left(\frac{\vartheta_1}{\vartheta_2}\right)''(x_0; v)$ exists in the direction $v \in Z$, then

$$\limsup_{n \rightarrow \infty} \frac{\left(\frac{\vartheta_1}{\vartheta_2}\right)(x_n) - \left(\frac{\vartheta_1}{\vartheta_2}\right)(x_0) - t_n \left(\frac{\vartheta_1}{\vartheta_2}\right)^{\circ}(x_0; v)}{\frac{1}{2} r_n^{-1} t_n^2} \leq \left(\frac{\vartheta_1}{\vartheta_2}\right)'(x_0; w) + r \left(\frac{\vartheta_1}{\vartheta_2}\right)''(x_0; v). \quad (9)$$

Proof Let $x_n = x_0 + t_n v + \frac{1}{2} r_n^{-1} t_n^2 w_n$. From Remark 2.3 (ii) and Proposition 2.1 (i), it yields that $\frac{\vartheta_1}{\vartheta_2}$ is locally Lipschitz continuous and regular in the sense of Clarke at $x_0 \in U$. By Lebourg's Mean-Value Theorem (see [31, Theorem 2.3.7]), we get that there exist $\theta_n \in (0, 1)$, $\xi_n \in \partial \left(\frac{\vartheta_1}{\vartheta_2}\right)(x_0 + t_n v + \frac{\theta_n}{2} r_n^{-1} t_n^2 w_n)$ such that

$$\left(\frac{\vartheta_1}{\vartheta_2}\right)(x_n) - \left(\frac{\vartheta_1}{\vartheta_2}\right)(x_0 + t_n v) = \frac{1}{2} r_n^{-1} t_n^2 \langle \xi_n, w_n \rangle.$$

Combining Lemma 2.2 with the Lipschitz continuity of $\frac{\vartheta_1}{\vartheta_2}$, there exists a constant $L_0 > 0$ such that $\|\xi_n\| \leq L_0$ for n sufficiently large. Without loss of generality, assume that $\xi_n \rightarrow \xi$. Owing to the closedness of $\partial \left(\frac{\vartheta_1}{\vartheta_2}\right)(\cdot)$ (see [32, Theorem 8.6]), there exists $\xi \in \partial \left(\frac{\vartheta_1}{\vartheta_2}\right)(x_0)$ such that

$$\lim_{n \rightarrow \infty} \frac{\left(\frac{\vartheta_1}{\vartheta_2}\right)(x_n) - \left(\frac{\vartheta_1}{\vartheta_2}\right)(x_0 + t_n v)}{\frac{1}{2} r_n^{-1} t_n^2} = \langle \xi, w \rangle.$$

In turn, due to $0 < r_n \rightarrow r$, using Definition 2.6 (ii) we get

$$\limsup_{n \rightarrow \infty} \frac{\left(\frac{\vartheta_1}{\vartheta_2}\right)(x_0 + t_n v) - \left(\frac{\vartheta_1}{\vartheta_2}\right)(x_0) - t_n \left(\frac{\vartheta_1}{\vartheta_2}\right)^{\circ}(x_0; v)}{\frac{1}{2} r_n^{-1} t_n^2} \leq r \left(\frac{\vartheta_1}{\vartheta_2}\right)^{\circ\circ}(x_0; v),$$

which together with (6) yields that

$$\begin{aligned}
& \limsup_{n \rightarrow \infty} \frac{\left(\frac{\vartheta_1}{\vartheta_2}\right)(x_n) - \left(\frac{\vartheta_1}{\vartheta_2}\right)(x_0) - t_n \left(\frac{\vartheta_1}{\vartheta_2}\right)^\circ(x_0; v)}{\frac{1}{2}r_n^{-1}t_n^2} \\
& \leq \limsup_{n \rightarrow \infty} \frac{\left(\frac{\vartheta_1}{\vartheta_2}\right)(x_n) - \left(\frac{\vartheta_1}{\vartheta_2}\right)(x_0 + t_n v)}{\frac{1}{2}r_n^{-1}t_n^2} \\
& \quad + \limsup_{n \rightarrow \infty} \frac{\left(\frac{\vartheta_1}{\vartheta_2}\right)(x_0 + t_n v) - \left(\frac{\vartheta_1}{\vartheta_2}\right)(x_0) - t_n \left(\frac{\vartheta_1}{\vartheta_2}\right)^\circ(x_0; v)}{\frac{1}{2}r_n^{-1}t_n^2} \\
& \leq \langle \xi, w \rangle + r \left(\frac{\vartheta_1}{\vartheta_2}\right)^{\circ\circ}(x_0; v) \\
& \leq \left(\frac{\vartheta_1}{\vartheta_2}\right)^\circ(x_0; w) + r \left(\frac{\vartheta_1}{\vartheta_2}\right)^{\circ\circ}(x_0; v) \\
& = \left(\frac{\vartheta_1}{\vartheta_2}\right)'(x_0; w) + r \left(\frac{\vartheta_1}{\vartheta_2}\right)^{\circ\circ}(x_0; v),
\end{aligned}$$

which comes from the Clarke regularity of $\frac{\vartheta_1}{\vartheta_2}$ at $x_0 \in U$. If $\left(\frac{\vartheta_1}{\vartheta_2}\right)''(x_0; v)$ exists in the direction $v \in Z$, then (9) results from Remark 2.2 (ii). The proof is completed.

Remark 2.5 It should be pointed out that Lemma 2.3 improves Lemma 2.1 of [26] even when ϑ_1 is continuously Fréchet differentiable at x_0 , ϑ_2 is a positive constant, and $f_1^{\circ\circ}(x_0; v)$ exists in the direction $v \in Z$. Moreover, Lemma 2.3 is reduced to Lemma 2.1 of [26] when ϑ_1 is continuously Fréchet differentiable at x_0 , ϑ_2 is a positive constant, and $\vartheta_1'(x_0; v)$ exists along $v \in Z$.

3 Second-order strong KKT necessary conditions for NMFP

If the multipliers of all objective functions in the KKT conditions are positive, it is that the strong KKT conditions hold. The strong KKT condition basically implies that all objective functions are active at the point at which the necessary optimality conditions hold. This section introduces a generalized second-order Abadie regularity condition, which extends the second-order Abadie regularity conditions proposed in [26, 33]. We further study second-order strong KKT necessary conditions that contain some new information on multipliers and imply the strong KKT necessary conditions for Borwein-type properly efficient solutions of NMFP, which is stronger than (weak) Pareto efficient solution.

To begin with, we recollect the notions of Borwein properly efficient solution, projective second-order tangent cone, and projective second-order linearizing cone.

Definition 3.1 [34, 35] A point $x_0 \in X$ is said to be *Borwein properly efficient solution* of NMFP iff

$$T\left(\left(\frac{f}{F}\right)(X) + \mathbb{R}_+^p, \left(\frac{f}{F}\right)(x_0)\right) \cap (-\mathbb{R}_+^p) = \{0\}.$$

It is pointed out in [36] that a Borwein properly efficient solution defined in Definition 3.1 must also be a Pareto efficient solution.

Definition 3.2 [26] Let $X, Q \subseteq Z$ and $v \in Z$.

- (i) The *projective second-order tangent cone* of X at $x_0 \in X$ in the direction v is defined by

$$\tilde{T}^2(X, x_0, v) = \left\{ (w, r) \in Z \times \mathbb{R}_+ : \begin{array}{l} \exists t_k \rightarrow 0^+, \exists r_k \rightarrow r, \exists w_k \rightarrow w \text{ such that} \\ \frac{t_k}{r_k} \rightarrow 0^+, x_0 + t_k v + \frac{1}{2} t_k^2 w_k \in X, \forall k \in \mathbb{N} \end{array} \right\}.$$

- (ii) The *projective second-order linearizing cone* of Q at $x_0 \in X$ in the direction v is defined by

$$\tilde{C}^2(Q, x_0, v) = \left\{ (w, r) \in Z \times \mathbb{R}_+ : \begin{array}{l} \left(\frac{f}{F}\right)_i^\circ(x_0; w) + r \left(\frac{f}{F}\right)_i^{\circ\circ}(x_0; v) \leq 0, \quad i \in I, \\ g_j^\circ(x_0; w) + r g_j^{\circ\circ}(x_0; v) \leq 0, \quad j \in J(x_0, v), \\ h_k^\circ(x_0; w) + r h_k^{\circ\circ}(x_0; v) = 0, \quad k \in K \end{array} \right\}.$$

The projective second-order tangent cone $\tilde{T}^2(X, x_0, v)$ has been widely applied to study optimality conditions; see [26, 33, 37, 38, 39]. Moreover, it follows from [39, Proposition 2.1] that $\tilde{T}^2(X, x_0, 0) = T(X, x_0) \times \mathbb{R}_+$, and $v \notin T(X, x_0)$ implies $\tilde{T}^2(X, x_0, 0) = \emptyset$.

Next, we introduce a new second-order Abadie regularity condition and a new second-order Guignard-type regular condition for NMFP.

Definition 3.3 We say that

- (i) a *generalized second-order Abadie-type regularity condition* (GSOARC) holds at $x_0 \in X$ in the direction $v \in D(x_0)$ iff

$$\tilde{C}^2(Q, x_0, v) \subseteq \tilde{T}^2(X, x_0, v). \quad (10)$$

- (ii) a *generalized second-order Guignard-type regular condition* (GSOGRC) holds at $x_0 \in X$ in the direction $v \in D(x_0)$ iff

$$\tilde{C}^2(Q, x_0, v) \subseteq \text{clco} \tilde{T}^2(X, x_0, v). \quad (11)$$

Remark 3.1 (i) It is easy to see that GSOGRC is a weaker regular condition than GSOARC.

If $v = 0$, specifically, the Páles-Zeidan second-order generalized directional derivative becomes 0, then GSOARC is reduced to a degenerate form

$$C(Q, x_0) \subseteq T(X, x_0), \quad (12)$$

as $\tilde{C}^2(Q, x_0, 0) = C(Q, x_0) \times \mathbb{R}_+$ and $\tilde{T}^2(X, x_0, 0) = T(X, x_0) \times \mathbb{R}_+$.

- (ii) If all the functions involved have continuous Fréchet derivatives and the corresponding second-order directional derivatives in the direction v exist, then the GSOARC is reduced to the *second-order Abadie regularity condition* SOARC in [26].
- (iii) In [26], SOARC was proposed by utilizing Fréchet derivatives and second-order directional derivatives. Another second-order Abadie constraint qualification in [33] was introduced in terms of Clarke generalized directional derivatives and Páles-Zeidan second-order generalized directional derivatives, which did not involve the objective functions. Compared with the second-order Abadie regularity conditions in [26, 33], GSOARC is introduced in terms of Clarke generalized directional derivatives and Páles-Zeidan second-order generalized directional derivatives, which also incorporate objective functions.

Assumption 3.1 (i) For $i \in I, j \in J, k \in K$, the functions f_i, F_i, g_j and h_k are locally Lipschitz continuous and regular in the sense of Clarke on U , and $f_i^{\circ\circ}(x; v), g_j^{\circ\circ}(x; v)$ and $h_k^{\circ\circ}(x; v)$ are finite on U for all direction $v \in Z$.

- (ii) For $i \in I, F_i$'s are Gâteaux differentiable on U with Gâteaux derivative $F_{i,G}'$, and $F_i''(x; v)$ exist on U for each direction $v \in Z$.

Lemma 3.1 *Let $x_0 \in X$ be a Borwein properly efficient solution of NMFP, $v \in D(x_0)$ and $s = (s_1, s_2, \dots, s_p)$ be defined as that in Lemma 2.1. Let Assumption 3.1 be fulfilled. Then, the following system*

$$\begin{cases} f_i^\circ(x_0; w) + rf_i^{\circ\circ}(x_0; v) - s_i(\langle F'_{i,G}(x_0), w \rangle + rF_i''(x_0; v)) \leq 0, \forall i \in I, \\ f_i^\circ(x_0; w) + rf_i^{\circ\circ}(x_0; v) - s_i(\langle F'_{i,G}(x_0), w \rangle + rF_i''(x_0; v)) < 0, \exists i \in I, \\ (w, r) \in \tilde{T}^2(X, x_0, v), \end{cases} \quad (13)$$

has no solution $(w, r) \in Z \times \mathbb{R}_+$.

Proof From Proposition 2.1 (ii) and (iv), it follows that

$$\begin{aligned} & \left(\frac{f_i}{F_i}\right)^\circ(x_0; w) + r \left(\frac{f_i}{F_i}\right)^{\circ\circ}(x_0; v) \\ &= \frac{1}{F_i(x_0)} [f_i^\circ(x_0; w) - s_i \langle F'_{i,G}(x_0), w \rangle] + \frac{r}{F_i(x_0)} [f_i^{\circ\circ}(x_0; v) - s_i F_i''(x_0; v)] \\ &= \frac{1}{F_i(x_0)} [f_i^\circ(x_0; w) + rf_i^{\circ\circ}(x_0; v) - s_i(\langle F'_{i,G}(x_0), w \rangle + rF_i''(x_0; v))]. \end{aligned}$$

Due to $F_i(x_0) > 0$ for all $i \in I$, it ensures that

$$\begin{aligned} & \left(\frac{f_i}{F_i}\right)^\circ(x_0; w) + r \left(\frac{f_i}{F_i}\right)^{\circ\circ}(x_0; v) < (\leq) 0 \\ \iff & f_i^\circ(x_0; w) + rf_i^{\circ\circ}(x_0; v) - s_i(\langle F'_{i,G}(x_0), w \rangle + rF_i''(x_0; v)) < (\leq) 0. \end{aligned}$$

Therefore, the system (13) is equivalent to

$$\begin{cases} \left(\frac{f_i}{F_i}\right)^\circ(x_0; w) + r \left(\frac{f_i}{F_i}\right)^{\circ\circ}(x_0; v) \leq 0, \forall i \in I, \\ \left(\frac{f_i}{F_i}\right)^\circ(x_0; w) + r \left(\frac{f_i}{F_i}\right)^{\circ\circ}(x_0; v) < 0, \exists i \in I, \\ (w, r) \in \tilde{T}^2(X, x_0, v). \end{cases} \quad (14)$$

Consequently, invoking Lemma 2.3 by the similar proof presented in [26, Lemma 4.1], one can obtain that the system (14) has no solution $(w, r) \in Z \times \mathbb{R}_+$. This completes the proof.

We now derive the second-order strong KKT necessary conditions for a Borwein-type proper efficient solution of NMFP with the help of GSOARC.

Theorem 3.1 [Primal condition] *Let $x_0 \in U$ be a Borwein-properly efficient solution of NMFP and $s = (s_1, s_2, \dots, s_p)$ be defined as that in Lemma 2.1. Let Assumption 3.1 be fulfilled. If GSOARC holds at x_0 in the direction $v \in D(x_0)$, then for any $r \geq 0$, the system*

$$\begin{cases} f_i^\circ(x_0; w) + rf_i^{\circ\circ}(x_0; v) - s_i(\langle F'_{i,G}(x_0), w \rangle + rF_i''(x_0; v)) \leq 0, \forall i \in I, \\ f_i^\circ(x_0; w) + rf_i^{\circ\circ}(x_0; v) - s_i(\langle F'_{i,G}(x_0), w \rangle + rF_i''(x_0; v)) < 0, \exists i \in I, \\ g_j^\circ(x_0; w) + rg_j^{\circ\circ}(x_0; v) \leq 0, \forall j \in J(x_0, v), \\ h_k^\circ(x_0; w) + rh_k^{\circ\circ}(x_0; v) = 0, \forall k \in K, \end{cases} \quad (15)$$

is incompatible in $w \in Z$.

Proof From Lemma 3.1, it follows that there is no $(w, r) \in Z \times \mathbb{R}_+$ such that the system (13) is consistent. Since GSOARC holds at x_0 in the direction v , combining (10) with Definition 3.2, for all $r \geq 0$, the system (15) does not have a solution $w \in Z$. The proof is completed.

Theorem 3.2 [Dual condition] *Let $x_0 \in U$ be a Borwein properly efficient solution of NMFP and $s = (s_1, s_2, \dots, s_p)$ be defined as stated in Lemma 2.1. Let Assumption 3.1 be fulfilled and functions f_i ($i \in I$), g_j ($j \in J$) and h_k ($k \in K$) be Gâteaux differentiable at x_0 . Assume that GSOARC holds at x_0 in the direction $v \in D(x_0)$. Then, there exist $\lambda \in \mathbb{R}_{++}^p$, $\mu \in \mathbb{R}_+^m$ and $\nu \in \mathbb{R}^l$ such that*

$$\sum_{i=1}^p \lambda_i (f'_{i,G}(x_0) - s_i F'_{i,G}(x_0)) + \sum_{j=1}^m \mu_j g'_{j,G}(x_0) + \sum_{k=1}^l \nu_k h'_{k,G}(x_0) = 0, \quad (16)$$

$$\sum_{i=1}^p \lambda_i (f_i^{\circ\circ}(x_0; v) - s_i F_i''(x_0; v)) + \sum_{j=1}^m \mu_j g_j^{\circ\circ}(x_0; v) + \sum_{k=1}^l \nu_k h_k^{\circ\circ}(x_0; v) \geq 0, \quad (17)$$

$$\mu_j g_j(x_0) = 0, \quad j = 1, 2, \dots, m \quad (18)$$

$$\text{and } \mu_j g'_{j,G}(x_0)v = 0, \quad j = 1, 2, \dots, m. \quad (19)$$

Proof From Theorem 3.1, it follows that there is no $(w, r) \in Z \times \mathbb{R}$ such that

$$\left. \begin{aligned} \langle f'_{i,G}(x_0) - s_i F'_{i,G}(x_0), w \rangle + r (f_i^{\circ\circ}(x_0; v) - s_i F_i''(x_0; v)) &\leq 0, \quad \forall i \in I, \\ \langle f'_{i,G}(x_0) - s_i F'_{i,G}(x_0), w \rangle + r (f_i^{\circ\circ}(x_0; v) - s_i F_i''(x_0; v)) &< 0, \quad \exists i \in I, \\ \langle g'_{j,G}(x_0), w \rangle + r g_j^{\circ\circ}(x_0; v) &\leq 0, \quad \forall j \in J(x_0, v), \\ \langle h'_{k,G}(x_0), w \rangle + r h_k^{\circ\circ}(x_0; v) &= 0, \quad \forall k \in K, \\ -r &\leq 0. \end{aligned} \right\} \quad (20)$$

Let $J(x_0, v) = \{j_1, j_2, \dots, j_t\}$. We consider the following matrices:

$$B := \begin{pmatrix} f'_{1,G}(x_0) - s_1 F'_{1,G}(x_0) & f_1^{\circ\circ}(x_0; v) - s_1 F_1''(x_0; v) \\ f'_{2,G}(x_0) - s_2 F'_{2,G}(x_0) & f_2^{\circ\circ}(x_0; v) - s_2 F_2''(x_0; v) \\ \vdots & \vdots \\ f'_{p,G}(x_0) - s_p F'_{p,G}(x_0) & f_p^{\circ\circ}(x_0; v) - s_p F_p''(x_0; v) \end{pmatrix},$$

$$C := \begin{pmatrix} g'_{j_1,G}(x_0) & g_{j_1}^{\circ\circ}(x_0; v) \\ g'_{j_2,G}(x_0) & g_{j_2}^{\circ\circ}(x_0; v) \\ \vdots & \vdots \\ g'_{j_t,G}(x_0) & g_{j_t}^{\circ\circ}(x_0; v) \\ 0 & -1 \end{pmatrix} \quad \text{and} \quad D := \begin{pmatrix} h'_{1,G}(x_0) & h_1^{\circ\circ}(x_0; v) \\ h'_{2,G}(x_0) & h_2^{\circ\circ}(x_0; v) \\ \vdots & \vdots \\ h'_{l,G}(x_0) & h_l^{\circ\circ}(x_0; v) \end{pmatrix}.$$

Then, note that the system (20) is inconsistent in $(w, r) \in Z \times \mathbb{R}$ if and only if

$$\langle B, (w, r) \rangle \leq 0, \quad \langle C, (w, r) \rangle \leq 0, \quad \langle D, (w, r) \rangle = 0$$

has no solution $(w, r) \in Z \times \mathbb{R}$. This together with [34, Theorem 3.24] yield that there exist $\lambda \in \mathbb{R}_{++}^p$, $\mu_j \geq 0, j \in J(x_0, v)$, $\nu \in \mathbb{R}^l$ and $\mu_0 \geq 0$ such that

$$\sum_{i=1}^p \lambda_i (f'_{i,G}(x_0) - s_i F'_{i,G}(x_0)) + \sum_{j=1}^m \mu_j g'_{j,G}(x_0) + \sum_{k=1}^l \nu_k h'_{k,G}(x_0) = 0,$$

$$\text{and } \sum_{i=1}^p \lambda_i (f_i^{\circ\circ}(x_0; v) - s_i F_i''(x_0; v)) + \sum_{j=1}^m \mu_j g_j^{\circ\circ}(x_0; v) + \sum_{k=1}^l \nu_k h_k^{\circ\circ}(x_0; v) = \mu_0 \geq 0.$$

Consequently, we get (16)–(19) by picking up $\mu_j = 0$ when $j \notin J(x_0, v)$. The proof is completed.

Remark 3.2 Theorem 3.2 presents a second-order strong KKT necessary conditions, which contains some new information on multipliers (19) at a Borwein properly efficient solution of the problem (1). From the proof of Theorem 3.2, we get a first-order strong KKT condition (16) at a Borwein properly efficient solution. As indicated in Remark 3.1, when $v = 0$, GSOARC reduces to (12). In this case, while (17) is trivial, (16) and (18) remain satisfied.

Remark 3.3 (i) If for each $i \in I, j \in J, k \in K$, f_i and g_j are continuously Fréchet differentiable at $x_0 \in U$ and $F_i(x) = 1$ and $h_k(x) \equiv 0$ for all $x \in U$, and $f_i''(x_0, v)$ and $g_j''(x_0, v)$ exist for all $i \in I, j \in J(x_0, v)$, then Theorem 4.1 and Theorem 4.2 in [26] can be deduced by Theorem 3.1 and Theorem 3.2, respectively.

(ii) If for each $i \in I, j \in J, k \in K$, f_i, g_j and h_k are continuously Fréchet differentiable at $x_0 \in U$ and $F_i(x) = 1$ for all $x \in U$, and $f_i''(x_0, v), h_k''(x_0, v)$ and $g_j''(x_0, v)$ exist for all $i \in I, j \in J(x_0, v)$, then Theorem 4.1 and Theorem 4.2 in [39] can be recovered by Theorem 3.1 and Theorem 3.2, respectively.

Corollary 3.1 *Let x_0 be a Borwein properly efficient solution of NMFP and $s = (s_1, s_2, \dots, s_p)$ be defined as that in Lemma 2.1. Let Assumption 3.1 be fulfilled and functions f_i ($i \in I$), g_j ($j \in J$) and h_k ($k \in K$) be Gâteaux differentiable at x_0 . If (12) holds at x_0 , then there exist vectors $\lambda \in \mathbb{R}_{++}^p$, $\mu \in \mathbb{R}_+^m$ and $\nu \in \mathbb{R}^l$ such that (16) and (18) hold.*

Remark 3.4 It is worth noting that $\lambda_i(f_{i,G}'(x_0) - s_i F_{i,G}'(x_0))$ is identical to $\bar{\lambda}_i \left(\frac{f_i}{F_i} \right)'_G(x_0)$ by choosing $\bar{\lambda}_i = \frac{\lambda_i}{F_i(x_0)}$ according to Proposition 2.1. Analogously, $\lambda_i(f_i^{\circ\circ}(x_0; v) - s_i F_i^{\circ\circ}(x_0; v))$ is identical to $\bar{\lambda}_i \left(\frac{f}{F} \right)^{\circ\circ}(x_0; v)$.

Since GSOGRC is weaker than GSOARC, a natural question is whether the results presented in Theorem 3.2 hold under the GSOGRC assumption. The answer is not affirmative, as evident from the following Example 3.1.

Example 3.1 Let $p = m = 2$ and $l = 1$. Consider the NMFP problem where the functions $f_i, F_i, g_j, h_k : \mathbb{R}^3 \rightarrow \mathbb{R}$ are as follows:

$$\begin{aligned} f_1(x) &= -3x_1 + x_2, \quad f_2(x) = 2x_1 - 3x_2, \quad F_1(x) = F_2(x) = 1 + x_1 + x_2, \\ g_1(x) &= -x_1, \quad g_2(x) = -x_2 \quad \text{and} \quad h(x) = x_1 x_2. \end{aligned}$$

The feasible set for this problem is

$$X = \{(x_1, x_2, x_3) \in \mathbb{R}^3 : x_1 \geq 0, x_2 \geq 0, x_1 x_2 = 0\}.$$

Clearly, $x_0 = 0$ is a Borwein properly efficient solution for NMFP since

$$T \left(\left(\frac{f}{F} \right) (X) + \mathbb{R}_+^2, \left(\frac{f}{F} \right) (x_0) \right) \cap (-\mathbb{R}_+^2) = \{0\}.$$

By directly calculation, we obtain $T(X, x_0) = X$, $s = (0, 0)$,

$$\begin{aligned} C(Q, x_0) &= \{(v_1, v_2, v_3) \in \mathbb{R}^3 : v_1 \geq 0, v_1 \leq v_2 \leq 6v_1\}, \\ D(x_0) &= T(X, x_0) \cap C(Q, x_0) = \{(v_1, v_2, v_3) \in \mathbb{R}^3 : v_1 = v_2 = 0\}, \end{aligned}$$

and for arbitrary $v \in D(x_0)$,

$$\begin{aligned} \tilde{T}^2(X, x_0, v) &= \{(w, s) \in \mathbb{R}^3 \times \mathbb{R} : w \in T(X, x_0), s \geq 0\} \\ \text{and } \tilde{C}^2(Q, x_0, v) &= \{(w, s) \in \mathbb{R}^3 \times \mathbb{R} : w \in C(Q, x_0), s \geq 0\}. \end{aligned}$$

It can be seen that GSOARC (10) is not satisfied and GSOGRC (11) is satisfied at x_0 . We point out that the conclusion derived in Theorem 3.2 is not true for this example. Indeed, because if there exist $\lambda_1 > 0$, $\lambda_2 > 0$, $\mu_1 \geq 0$, $\mu_2 \geq 0$ and $\nu \in \mathbb{R}$ such that (16) is fulfilled, i.e.,

$$\begin{aligned} & \lambda_1(\nabla f_1(x_0) - s_1 \nabla F_1(x_0)) + \lambda_2(\nabla f_2(x_0) - s_2 \nabla F_2(x_0)) \\ & + \mu_1 \nabla g_1(x_0) + \mu_2 \nabla g_2(x_0) + \nu \nabla h(x_0) = 0, \end{aligned}$$

then $\lambda_1 = -\frac{1}{7}(3\mu_1 + 2\mu_2) \leq 0$ and $\lambda_2 = -\frac{1}{7}(\mu_1 + 3\mu_2) \leq 0$, which are contradictory to $\lambda_1 > 0$ and $\lambda_2 > 0$. Therefore, Theorem 3.2 is not true under the GSOGRC assumption.

4 Second-order KKT sufficient optimality conditions for NMFP

This section studies second-order sufficient optimality conditions for NMFP under second-order generalized convexity assumptions. To start with, we introduce some notions of second-order generalized convexity.

Definition 4.1 Let $x_0 \in X \subseteq U$. A locally Lipschitz continuous function $\vartheta : U \rightarrow \mathbb{R}$ is said to be

- (i) second-order *convex* at x_0 with respect to X if for any $x \in X$ there exist $v \in Z \setminus \{0\}$ and $w \in Z \setminus \{0\}$ such that

$$\vartheta(x) - \vartheta(x_0) \geq \vartheta^\circ(x_0; w) + \frac{1}{2}\vartheta^{\circ\circ}(x_0; v),$$

- (ii) second-order *pseudoconvex* at x_0 with respect to X if for any $x \in X$ there exist $v \in Z \setminus \{0\}$ and $w \in Z \setminus \{0\}$ such that

$$\vartheta^\circ(x_0; w) + \frac{1}{2}\vartheta^{\circ\circ}(x_0; v) \geq 0 \implies \vartheta(x) \geq \vartheta(x_0),$$

- (iii) second-order *quasiconvex* at x_0 with respect to X if for any $x \in X$ there exist $v \in Z \setminus \{0\}$ and $w \in Z \setminus \{0\}$ such that

$$\vartheta(x) \leq \vartheta(x_0) \implies \vartheta^\circ(x_0; w) + \frac{1}{2}\vartheta^{\circ\circ}(x_0; v) \leq 0,$$

- (iv) second-order *infine* at x_0 with respect to X if for any $x \in X$ there exist $v \in Z \setminus \{0\}$ and $w \in Z \setminus \{0\}$ such that

$$\vartheta(x) - \vartheta(x_0) = \vartheta^\circ(x_0; w) + \frac{1}{2}\vartheta^{\circ\circ}(x_0; v).$$

Remark 4.1 It is noteworthy that if a function ϑ is second-order infine at x_0 with respect to X , then ϑ is second-order convex at x_0 with respect to X . The second-order convexity of ϑ at x_0 with respect to X implies the second-order pseudoconvexity and second-order quasiconvexity of ϑ at x_0 with respect to X . Besides, if $\vartheta : U \rightarrow \mathbb{R}$ is locally Lipschitz continuous and Gâteaux differentiable at x_0 , and $w = v = x - x_0$ in Definition 4.1(i)(ii)(iii), then the second-order convexity, second-order pseudoconvexity and second-order quasiconvexity of ϑ at x_0 with respect to X reduce to the corresponding second-order convexity of ϑ at x_0 introduced in [40, Definition 4.1].

Example 4.1 (i) Consider the function $\vartheta(x) = x^2$, $x \in \mathbb{R}$, and $x_0 = 0$. By direct calculation, we get $\vartheta^\circ(0; w) = \nabla\vartheta(0)w = 0$ for all $w \in \mathbb{R}$, and $\vartheta^{\circ\circ}(0; v) = \nabla^2\vartheta(0)v^2 = 2v^2 \geq 0$. Letting $v = x - x_0 = x$, one has

$$\vartheta(x) - \vartheta(x_0) = \vartheta^\circ(x_0; w) + \frac{1}{2}\vartheta^{\circ\circ}(x_0; v).$$

Therefore, ϑ is second-order infine at x_0 with respect to \mathbb{R} . In addition, from Remark 4.1, it is obvious that ϑ is also second-order convex (pseudoconvex and quasiconvex) at x_0 with respect to \mathbb{R} .

- (ii) Consider the function $\vartheta(x) = |x|$ and $x_0 = 0$. By direct calculation, we get $\vartheta^\circ(0; w) = |w|$, and $\vartheta^{\circ\circ}(0; v) = 0$ for all $v \in \mathbb{R}$. Hence, for any $x \neq 0$, letting $w := x - x_0 = x$, we get $\vartheta^\circ(x_0; w) + \frac{1}{2}\vartheta^{\circ\circ}(x_0; v) = |x| > 0$ for all $v \in \mathbb{R}$. So, ϑ is second-order pseudoconvex at x_0 with respect to \mathbb{R} .

Definition 4.2 A function $f := (f_1, f_2, \dots, f_p)^\top : U \rightarrow \mathbb{R}^p$ is called second-order convex (respectively, pseudoconvex, quasiconvex, and infine) at x_0 with respect to X if its components f_i , $i \in I$ are second-order convex (respectively, pseudoconvex, quasiconvex, and infine) at x_0 with respect to X and common $v \in Z$ and $w \in Z$.

By direct calculation, one easily gets the following results.

Proposition 4.1 Let f_1 and f_2 be two real-valued locally Lipschitz functions on U with f_2 being positive, and $x_0 \in X \subseteq U$. Suppose that for each $x \in U$, both $f'_{2,G}(x)$ and $f''_2(x; v)$ exist. Then,

- (i) for each $\beta > 0$, βf_1 is second-order convex (pseudoconvex, quasiconvex, or infine) at x_0 with respect to X if so is f_1 ;
(ii) $f_1 + f_2$ is second-order convex (infine) at x_0 with respect to X if so are f_1 and f_2 with a common $v \in Z$ and $w \in Z$.

For simplicity, we take the following assumption to derive second-order sufficient optimality conditions.

Assumption 4.1 All the functions involved in NMFP exhibit a form of second-order convexity in the sense of Definition 4.2 with a common $v \in D(x_0)$ and $w \in Z$.

Next, we present second-order sufficient optimality conditions for NMFP.

Theorem 4.1 Let $x_0 \in X$, $s = \frac{f(x_0)}{F(x_0)}$ and let $f - s * F$ and g be second-order convex at x_0 with respect to X , h be second-order infine at x_0 with respect to X and Assumptions 3.1 and 4.1 hold. Assume that there exist $\lambda \in \mathbb{R}^p_{++}$, $\mu \in \mathbb{R}^m_+$ and $\nu \in \mathbb{R}^l$ such that for all $v \in D(x_0)$ and $w \in Z$,

$$\lambda^\top (f^\circ(x_0; w) - \langle s * F'_G(x_0), w \rangle) + \sum_{j=1}^m \mu_j g_j^\circ(x_0; w) + \sum_{k=1}^l \nu_k h_k^\circ(x_0; w) = 0, \quad (21)$$

$$\lambda^\top (f^{\circ\circ}(x_0; v) - s * F''(x_0; v)) + \sum_{j=1}^m \mu_j g_j^{\circ\circ}(x_0; v) + \sum_{k=1}^l \nu_k h_k^{\circ\circ}(x_0; v) \geq 0 \quad (22)$$

$$\text{and } \sum_{j=1}^m \mu_j g_j(x_0) = 0. \quad (23)$$

Then, x_0 is a Pareto efficient solution of NMFP.

Proof From Lemma 2.1, it suffices to prove that x_0 is a Pareto efficient solution of the problem (3). On the contrary, if x_0 is not a Pareto efficient solution of the problem (3), then there exists $\bar{x} \in X$ for which

$$(f - s * F)(\bar{x}) \leq (f - s * F)(x_0). \quad (24)$$

Due to $\bar{x}, x_0 \in X$ and $\lambda \in \mathbb{R}_{++}^p$, (23) and (24) yield that

$$\begin{aligned} & \lambda^\top (f - s * F)(x_0) + \sum_{j=1}^m \mu_j g_j(x_0) + \sum_{k=1}^l \nu_k h_k(x_0) \\ & > \lambda^\top (f - s * F)(\bar{x}) + \sum_{j=1}^m \mu_j g_j(\bar{x}) + \sum_{k=1}^l \nu_k h_k(\bar{x}). \end{aligned} \quad (25)$$

Note that $f - s * F$ and g are second-order convex at x_0 with respect to X , and h is second-order infine at x_0 with respect to X . Using Proposition 2.1, Assumption 4.1 yields that there exist $v \in D(x_0)$ and $w \in Z$ such that

$$\begin{aligned} & \lambda^\top (f - s * F)(\bar{x}) + \sum_{j=1}^m \mu_j g_j(\bar{x}) + \sum_{k=1}^l \nu_k h_k(\bar{x}) \\ \geq & \lambda^\top (f - s * F)(x_0) + \sum_{j=1}^m \mu_j g_j(x_0) + \sum_{k=1}^l \nu_k h_k(x_0) \\ & + \lambda^\top (f - s * F)^\circ(x_0; w) + \sum_{j=1}^m \mu_j g_j^\circ(x_0; w) + \sum_{k=1}^l \nu_k h_k^\circ(x_0; w) \\ & + \frac{1}{2} \lambda^\top (f - s * F)^{\circ\circ}(x_0; v) + \sum_{j=1}^m \frac{1}{2} \mu_j g_j^{\circ\circ}(x_0; v) + \sum_{k=1}^l \frac{1}{2} \nu_k h_k^{\circ\circ}(x_0; v) \\ = & \lambda^\top (f - s * F)(x_0) + \sum_{j=1}^m \mu_j g_j(x_0) + \sum_{k=1}^l \nu_k h_k(x_0) \\ & + \lambda^\top (f^\circ(x_0; w) - \langle s * F'_G(x_0), w \rangle) + \sum_{j=1}^m \mu_j g_j^\circ(x_0; w) + \sum_{k=1}^l \nu_k h_k^\circ(x_0; w) \\ & + \frac{1}{2} \left(\lambda^\top (f^{\circ\circ}(x_0; v) - s * F''(x_0; v)) + \sum_{j=1}^m \mu_j g_j^{\circ\circ}(x_0; v) + \sum_{k=1}^l \nu_k h_k^{\circ\circ}(x_0; v) \right) \\ \stackrel{(21)\&(22)}{\geq} & \lambda^\top (f - s * F)(x_0) + \sum_{j=1}^m \mu_j g_j(x_0) + \sum_{k=1}^l \nu_k h_k(x_0), \end{aligned} \quad (26)$$

which is contradictory to (25). Therefore, x_0 is a Pareto efficient solution of the problem (3), and hence the proof is completed.

Remark 4.2 From the proof of Theorem 4.1, we see that if $\lambda \in \mathbb{R}_{++}^p$ is replaced by $\lambda \in \mathbb{R}_+^p \setminus \{0\}$ and the inequality (\geq) in (22) is replaced by the strict inequality ($>$), then x_0 is a weak efficient solution of NMFP under the assumptions of Theorem 4.1.

Theorem 4.2 *Let $x_0 \in X$ and $s = \frac{f(x_0)}{F(x_0)}$. Assume that there exist $\lambda \in \mathbb{R}_{++}^p$, $\mu \in \mathbb{R}_+^m$ and $\nu \in \mathbb{R}^l$ such that (21)–(23) hold. If $\lambda^\top (f - s * F)$ is second-order pseudoconvex at x_0 with respect to X , $\mu^\top g$ is second-order quasiconvex at x_0 with respect to X , h is second-order infine at x_0 with respect to X and Assumptions 3.1 and 4.1 hold, then x_0 is a Pareto efficient solution of NMFP.*

Proof Consider the functions $\phi(x) := \lambda^\top (f - s * F)(x)$ and $\psi(x) := \mu^\top g(x)$ for $x \in X$. Then, we have

$$\psi(x) \leq \psi(x_0) \leq 0 \quad \text{and} \quad h(x) = h(x_0) = 0, \quad \forall x \in X.$$

Since $\mu^\top g$ is second-order quasiconvex at x_0 with respect to X and h is second-order infine at x_0 with respect to X , Assumptions 4.1 yields that for any $x \in X$, there exist $v \in D(x_0)$ and $w \in Z$ such that

$$\psi^\circ(x_0; w) + \frac{1}{2}\psi^{\circ\circ}(x_0; v) \leq 0, \quad h_k^\circ(x_0; w) + \frac{1}{2}h_k^{\circ\circ}(x_0; v) = 0, \quad k \in K.$$

Moreover, we have

$$\sum_{j=1}^m \mu_j g_j^\circ(x_0; w) + \frac{1}{2} \sum_{j=1}^m \mu_j g_j^{\circ\circ}(x_0; v) \leq 0 \quad \text{and} \quad \sum_{k=1}^l \nu_k h_k^\circ(x_0; w) + \frac{1}{2} \sum_{k=1}^l \nu_k h_k^{\circ\circ}(x_0; v) = 0.$$

From (21) and (22), we deduce that

$$\lambda^\top (f^\circ(x_0; w) - \langle s * F'_G(x_0), w \rangle) + \frac{1}{2} \lambda^\top (f^{\circ\circ}(x_0; v) - s * F''(x_0; v)) \geq 0,$$

i.e., $\phi^\circ(x_0; w) + \frac{1}{2}\phi^{\circ\circ}(x_0; v) \geq 0$. Since $\phi(x)$ is second-order pseudoconvex at x_0 with respect to X , we have $\phi(x) \geq \phi(x_0)$ for all $x \in X$, and so

$$\lambda^\top (f - s * F)(x) \geq \lambda^\top (f - s * F)(x_0), \quad \forall x \in X. \quad (27)$$

Due to $\lambda \in \mathbb{R}_{++}^p$, (27) implies that

$$(f - s * F)(x_0) - (f - s * F)(x) \notin \mathbb{R}_+^p \setminus \{0\}, \quad \forall x \in X.$$

It, therefore, follows from Definition 2.1 that x_0 is a Pareto efficient solution of the problem (3), and hence it is a Pareto efficient solution of NMFP by Lemma 2.1. The proof is completed.

In the following, we exemplify Theorem 4.2.

Example 4.2 Let $p = m = 2$ and $l = 1$. Consider the NMFP problem, where the functions $f_i, F_i, g_j, h_k: \mathbb{R}^2 \rightarrow \mathbb{R}$ are defined by

$$\begin{aligned} f_1(x) &:= 3x_1^4 + 5x_1^2 + 6x_2^2, \quad f_2(x) := -2x_2^2, \quad F_1(x) = F_2(x) := x_1^2 + x_2^2 + 1, \\ g_1(x) &:= -x_1^2, \quad g_2(x) := -x_2 \quad \text{and} \quad h(x) := x_1^2. \end{aligned}$$

The feasible set for the problem is $X = \{(x_1, x_2) \in \mathbb{R}^2 : x_1 = 0, x_2 \geq 0\}$, and for $x_0 = (0, 0)$, we get $s = (0, 0)$ and $D(x_0) = X$. Take $\lambda_1 = 1, \lambda_2 = 2, \mu_1 = 1, \mu_2 = 0$ and $\nu = -2$. For each $x = (x_1, x_2) \in X$, there exist $v = x \in D(x_0)$ and $w = (1, 1)$ such that $\lambda^\top (f - s * F)$ is second-order convex (and pseudoconvex) at x_0 with respect to X , $\mu^\top g$ is second-order quasiconvex at x_0 with respect to X , and h is second-order infine at x_0 with respect to X . In addition, for any $v \in D(x_0)$, we have (21)–(23). Observe that for any $x \in X$,

$$\frac{f(x_0)}{F(x_0)} - \frac{f(x)}{F(x)} = \left(-\frac{3x_1^4 + 5x_1^2 + 6x_2^2}{x_1^2 + x_2^2 + 1}, \frac{2x_2^2}{x_1^2 + x_2^2 + 1} \right)^\top = \left(-\frac{6x_2^2}{x_2^2 + 1}, \frac{2x_2^2}{x_2^2 + 1} \right)^\top \notin \mathbb{R}_+^2 \setminus \{0\}.$$

It thus implies that x_0 is a Pareto efficient solution of NMFP.

Remark 4.3 From the proof of Theorem 4.2, specifically beginning with (27), we can conclude that if $\lambda \in \mathbb{R}_{++}^p$ is replaced by $\lambda \in \mathbb{R}_+^p \setminus \{0\}$ in Theorem 4.2, then x_0 is a weak efficient solution of NMFP under the assumptions of Theorem 4.2.

5 Second-order duality

In this section, we study Mond-Weir-type second-order duality of NMFP. Duality results between NMFP and its second-order dual problem are established under the generalized second-order convexity assumptions. A *Mond-Weir-type second-order dual problem* (MWSD) of NMFP is formulated as follows:

$$\begin{aligned}
& \max \quad \left(\frac{f}{F} \right) (u) \\
& \text{s.t.} \quad \lambda^\top (f'_G(u) - s * F'_G(u)) + \sum_{j=1}^m \mu_j g'_{j,G}(u) + \sum_{k=1}^l \nu_k h'_{k,G}(u) = 0, \\
& \quad \lambda^\top (f^{\circ\circ}(u; v) - s * F''(u; v)) + \sum_{j=1}^m \mu_j g_j^{\circ\circ}(u; v) + \sum_{k=1}^l \nu_k h_k^{\circ\circ}(u; v) \geq 0, \quad \forall v \in D(u), \\
& \quad \sum_{j=1}^m \mu_j g_j(u) + \sum_{k=1}^l \nu_k h_k(u) \geq 0, \quad \sum_{i=1}^p \lambda_i = 1, \\
& \quad (u, \lambda, \mu, \nu) \in U \times \mathbb{R}_{++}^p \times \mathbb{R}_+^m \times \mathbb{R}^l,
\end{aligned} \tag{28}$$

where $s = \frac{f(u)}{F(u)}$. A vector (u, λ, μ, ν) satisfying all the constraints of MWSD is said to be a feasible solution of MWSD. The set of feasible solutions of MWSD is denoted by \mathcal{F}_M .

Throughout this section, we always assume Assumption 5.1 hold.

Assumption 5.1 (i) $f_i, F_i (i \in I), g_j (j \in J)$ and $h_k (k \in K)$ are all locally Lipschitz continuous, Gâteaux differentiable and regular in the sense of Clarke on U with Gâteaux derivative $f'_{i,G}, F'_{i,G}, g'_{j,G}$ and $h'_{k,G}$, respectively;
(ii) For $i \in I, j \in J, k \in K, f_i^{\circ\circ}(u; v), g_j^{\circ\circ}(u; v), h_k^{\circ\circ}(u; v)$ and $F_i''(u, v)$ are all finite at each $u \in U$ for all directions $v \in Z$.

We first give the weak duality between NMFP and MWSD.

Theorem 5.1 [Weak duality] *Let $x \in X$ and $(u, \lambda, \mu, \nu) \in \mathcal{F}_M$. Assume that $f - s * F$ and g are second-order convex at u with respect to U , h is second-order infine at u with respect to U and Assumption 4.1 holds. Then,*

$$\left(\frac{f}{F} \right) (x) \not\leq \left(\frac{f}{F} \right) (u).$$

Proof Suppose to the contrary that

$$\left(\frac{f}{F} \right) (x) \leq \left(\frac{f}{F} \right) (u). \tag{29}$$

Taking into account that $F(x) \in \mathbb{R}_{++}^p$ for all $x \in X$, (29) implies that

$$\begin{aligned}
\left(\frac{f}{F} \right) (x) - \left(\frac{f}{F} \right) (u) \in -\mathbb{R}_+^p \setminus \{0\} & \iff f(x) - s * F(x) \in -\mathbb{R}_+^p \setminus \{0\} \\
& \iff f(x) - s * F(x) \leq f(u) - s * F(u).
\end{aligned}$$

This together with $\lambda \in \mathbb{R}_{++}^p$ yields that

$$\lambda^\top (f - s * F)(x) < \lambda^\top (f - s * F)(u). \quad (30)$$

Since $x \in X$ and $(u, \lambda, \mu, \nu) \in \mathcal{F}_M$, we have $g(x) \leq 0$, $h(x) = 0$ and

$$\sum_{j=1}^m \mu_j g_j(u) + \sum_{k=1}^l \nu_k h_k(u) \geq 0.$$

Combining with (30), we have

$$\begin{aligned} & \lambda^\top (f - s * F)(x) + \sum_{j=1}^m \mu_j g_j(x) + \sum_{k=1}^l \nu_k h_k(x) \\ & < \lambda^\top (f - s * F)(u) + \sum_{j=1}^m \mu_j g_j(u) + \sum_{k=1}^l \nu_k h_k(u). \end{aligned} \quad (31)$$

By the similar arguments as that in deriving (26) in Theorem 4.1, we get

$$\begin{aligned} & \sum_{i=1}^p \lambda_i (f_i - s_i F_i)(x) + \sum_{j=1}^m \mu_j g_j(x) + \sum_{k=1}^l \nu_k h_k(x) \\ & \geq \sum_{i=1}^p \lambda_i (f_i - s_i F_i)(u) + \sum_{j=1}^m \mu_j g_j(u) + \sum_{k=1}^l \nu_k h_k(u), \end{aligned}$$

which is a contradictory to (31). Hence, the result follows. The proof is completed.

Theorem 5.2 [Strong duality] *Let $x_0 \in X$ be a Borwein-properly efficient solution of NMFP. Assume that all conditions of Theorem 3.2 are satisfied. Then, there exist $\bar{\lambda} \in \mathbb{R}_{++}^p$, $\bar{\mu} \in \mathbb{R}_+^m$ and $\bar{\nu} \in \mathbb{R}^l$ such that $(x_0, \bar{\lambda}, \bar{\mu}, \bar{\nu}) \in \mathcal{F}_M$. Furthermore, if all conditions of Theorem 5.1 hold, then $(x_0, \bar{\lambda}, \bar{\mu}, \bar{\nu})$ is a Pareto efficient solution of MWSD.*

Proof From Theorem 3.2 it follows that there exists $(\bar{\lambda}, \bar{\mu}, \bar{\nu}) \in \mathbb{R}_{++}^p \times \mathbb{R}_+^m \times \mathbb{R}^l$ satisfies (16)-(18). Due to $h(x_0) = 0$, (18) yields that

$$\sum_{j=1}^m \bar{\mu}_j g_j(x_0) + \sum_{k=1}^l \bar{\nu}_k h_k(x_0) = 0.$$

With no loss of generality, we can assume that $\sum_{i=1}^p \bar{\lambda}_i = 1$ since one can pick up $\tilde{\lambda}_i = \frac{\bar{\lambda}_i}{\sum_{i=1}^p \bar{\lambda}_i}$ due to $\bar{\lambda} \in \mathbb{R}_{++}^p$. Consequently, one has $(x_0, \tilde{\lambda}, \bar{\mu}, \bar{\nu}) \in \mathcal{F}_M$. By Theorem 5.1, we obtain

$$\left(\frac{f}{F} \right) (x_0) \not\leq \left(\frac{f}{F} \right) (u), \quad \forall (u, \lambda, \mu, \nu) \in \mathcal{F}_M.$$

It, therefore, implies that (x_0, λ, μ, ν) is a Pareto efficient solution of MWSD. The proof is completed.

Theorem 5.3 [Converse duality] *Let $(u, \lambda, \mu, \nu) \in \mathcal{F}_M$ be a Pareto efficient solution of MWSD with $u \in X$. Assume that $\lambda^\top (f - s * F)$ is second-order pseudoconvex at u with respect to U , $\mu^\top g$ is second-order quasiconvex at u with respect to U , h is second-order infine at u with respect to U and Assumption 4.1 holds. Then u is a Pareto efficient solution of NMFP.*

Proof If possible, suppose that u is not a Pareto efficient solution of NMFP. Then, there exists $\tilde{x} \in X$ for which

$$\left(\frac{f}{F}\right)(\tilde{x}) \leq \left(\frac{f}{F}\right)(u).$$

Therefore, $f(\tilde{x}) - s * F(\tilde{x}) \leq 0 = f(u) - s * F(u)$, and

$$\lambda^\top (f(\tilde{x}) - s * F(\tilde{x})) < \lambda^\top (f(u) - s * F(u)). \quad (32)$$

Due to $(u, \lambda, \mu, \nu) \in \mathcal{F}_M$, (28) results in

$$\begin{aligned} & \sum_{i=1}^p \lambda_i \langle f'_{i,G}(u), w \rangle - s_i \langle F'_{i,G}(u), w \rangle + \sum_{j=1}^m \mu_j \langle g'_{j,G}(u), w \rangle \\ & + \sum_{k=1}^l \nu_k \langle h'_{k,G}(u), w \rangle + \frac{1}{2} \sum_{i=1}^p \lambda_i (f_i^{\circ\circ}(u; v) - s_i F_i''(u; v)) \\ & + \frac{1}{2} \sum_{j=1}^m \mu_j g_j^{\circ\circ}(u; v) + \frac{1}{2} \sum_{k=1}^l \nu_k h_k^{\circ\circ}(u; v) \geq 0, \forall v \in D(u), w \in Z, \end{aligned} \quad (33)$$

and $\mu^\top g(u) + \nu^\top h(u) \geq 0$. This together with $u \in X$ yields that

$$\mu^\top g(u) = 0 \geq \mu^\top g(x), \quad h(u) = h(x) = 0, \forall x \in X. \quad (34)$$

Since $\mu^\top g$ is second-order quasiconvex at u with respect to U , h is second-order infine at u with respect to U , it follows from Proposition 4.1 and Assumption 4.1 that for any $x \in U$, there exist $v \in D(u)$ and $w \in Z$ such that

$$\sum_{j=1}^m \mu_j \langle g'_{j,G}(u), w \rangle + \frac{1}{2} \sum_{j=1}^m \mu_j g_j^{\circ\circ}(u; v) \leq 0 \quad (35)$$

$$\text{and } \sum_{k=1}^l \nu_k \langle h'_{k,G}(u), w \rangle + \frac{1}{2} \sum_{k=1}^l \nu_k h_k^{\circ\circ}(u; v) = \nu^\top h(x) - \nu^\top h(u) = 0. \quad (36)$$

Summing (33), (35) and (36), it yields that

$$\sum_{i=1}^p \lambda_i \langle f'_{i,G}(u), w \rangle - s_i \langle F'_{i,G}(u), w \rangle + \frac{1}{2} \sum_{i=1}^p \lambda_i (f_i^{\circ\circ}(u; v) - s_i F_i''(u; v)) \geq 0.$$

By the second-order pseudoconvexity of $\lambda^\top (f - s * F)$ at u with respect to U , one has

$$\lambda^\top (f - s * F)(x) \geq \lambda^\top (f - s * F)(u), \quad \forall x \in U \setminus \{u\},$$

which contradicts (32). Hence, the result follows. The proof is completed.

Remark 5.1 If $\lambda \in \mathbb{R}_{++}^p$ is replaced by $\lambda \in \mathbb{R}_+^p \setminus \{0\}$ in MWSD, then we can conclude the strong duality and converse duality between the weak efficient solutions of NMFP and that of MWSD.

6 Conclusions

We have presented chain rules of quotient functions involving locally Lipschitz functions in terms of first and second-order directional derivatives, which improves that of [20, 26, 39]. A new second-order Abadie-type regular condition has been introduced in terms of Clarke directional derivative and Páles-Zeidan second-order directional derivative, which is different from the second-order Abadie-type regular conditions in [26, 33]. Second-order strong KKT conditions for a Borwein-properly efficient solution of NMFP have been established. Based on s -MFP, second-order sufficient optimality conditions for a Pareto efficient solution of NMFP have been obtained under some generalized second-order convexity assumptions. Finally, we have proposed a Mond-Weir-type second-order dual problem of NMFP and obtained the weak, strong and converse duality results between NMFP and its second-order dual problem.

In the lines of the derived results, one can study Schaible-type second-order dual problem [22] of NMFP, and also can attempt to design algorithms of NMFP via s -MFP. The obtained results in this paper can further be extended by using Dini Hadamard-type second-order generalized directional derivatives. It is also interesting to extend the results presented in this paper to the vector case when \mathbb{R}_+^p is replaced by a general closed convex cone.

Acknowledgements. This paper was supported by the Natural Science Foundation of China (Nos. 12071379, 12271061), the Natural Science Foundation of Chongqing (cstc2021jcyj-msxmX0925, cstc2022ycjh-bgzxm0097), Youth Project of Science and Technology Research Program of Chongqing Education Commission of China (No. KJQN202201802) and the Southwest University Graduate Research Innovation Program (No. SWUS23058). D. Ghosh is thankful for the research funding CRG (CRG/2022/001347) and MATRICS (MTR/2021/000696), SERB, India. J.-C. Yao acknowledges MOST (No. 111-2115-M-039-001-MY2), Taiwan. The first version of the paper was finished on 10th May, 2023.

Author contributions. All authors contributed equally to this article.

Compliance with Ethical Standards It is not applicable.

Competing interests

No potential conflict of interest was reported by the authors.

References

1. Bector, C. R.: Non-linear fractional functional programming with non-linear constraints. *Z. Angew. Math. Mech.* 48, 284–286 (1968)
2. Borwein, J. M.: Fractional programming without differentiability. *Math. Program.* 11, 283–290 (1976)
3. Schaible, S.: Fractional programming. *Z. Oper. Res. Ser. A-B.* 1, 39–54 (1983)
4. Luc, D. T.: *Theory of vector optimization. Lecture Notes in Economics and Mathematical Systems.* Springer, Berlin (1989)
5. Jahn, J.: *Vector Optimization Theory, Applications and Extensions.* Springer, Berlin (2004)
6. David, R. M., Edward, C. S. and Sheldon, H. J.: An application of the branch, bound, and remember algorithm to a new simple assembly line balancing dataset. *European J. Oper. Res.* 236(2), 403–409 (2014)
7. Sun, X. K., Long, X. J., Chai, Y.: Sequential optimality conditions for fractional optimization with applications to vector optimization. *J. Optim. Theory Appl.* 164, 479–499 (2015)
8. Shen, K. M. and Yu, W.: Fractional programming for communication systems—Part I: Power control and beamforming. *IEEE Trans. Signal Process.* 66(10), 2616–2630 (2018)
9. Shen, K. M. and Yu, W.: Fractional programming for communication systems—Part II: Uplink scheduling via matching. *IEEE Trans. Signal Process.* 66(10), 2631–2644 (2018)
10. Ansari, Q. H., Köbis, E., Yao, J. C.: *Vector Variational Inequalities and Vector Optimization: Theory and Applications.* Springer, Cham (2018)

11. Singh, C. and Hanson, M. A.: Saddle point theory for nondifferentiable multiobjective fractional programming. *J. Inform. Optim. Sci.* 7, 41–48 (1986)
12. Bector, C. R., Chandra, S. and Husain, I.: Optimality conditions and duality in subdifferentiable multiobjective fractional programming. *J. Optim. Theory Appl.* 79, 105–125 (1993)
13. Liu, J. C.: Optimality and duality for multiobjective fractional programming involving nonsmooth (F, ρ) -convex functions. *Optim.* 36, 333–346 (1996)
14. Kim, D. S., Kim, S. J. and Kim, M. H.: Optimality and duality for a class of nondifferentiable multiobjective fractional programming problems. *J. Optim. Theory Appl.* 129, 131–146 (2006)
15. Jayswal, A., Izhar, A. and Kummari, K.: Optimality conditions and duality in multiobjective fractional programming involving right upper-Dini-derivative functions. *Miskolc Math. Notes.* 16, 887–906 (2015)
16. Das, K.: Sufficiency and duality of set-valued fractional programming problems via second-order contingent epiderivative. *Yugosl. J. Oper. Res.* 32, 167–188 (2022)
17. Khanh, P. Q. and Tung, L. T.: First- and second-order optimality conditions for multiobjective fractional programming. *TOP.* 23, 419–440 (2015)
18. Khanh, P. Q. and Tung, N. M.: On the Mangasarian-Fromovitz constraint qualification and Karush-Kuhn-Tucker conditions in nonsmooth semi-infinite multiobjective programming. *Optim. Lett.* 14, 2055–2072 (2020)
19. Su, T. V. and Hang, D. D.: Optimality and duality in nonsmooth multiobjective fractional programming problem with constraints. *4OR.* 20, 105–137 (2022)
20. Su, T. V. and Hang, D. D.: Second-order optimality conditions in locally Lipschitz multiobjective fractional programming problem with inequality constraints. *Optim.* 72, 1171–1198 (2023)
21. Liang, Z. A., Huang, H. X. and Pardalos, P. M.: Optimality conditions and duality for a class of nonlinear fractional programming problems. *J. Optim. Theory Appl.* 110, 611–619 (2001)
22. Liang, Z. A., Huang, H. X. and Pardalos, P. M.: Efficiency conditions and duality for a class of multiobjective fractional programming problems. *J. Global Optim.* 27, 447–471 (2003)
23. Yuan, D. H., Liu, X. L., Chinchuluun, A. and Pardalos, P. M.: Nondifferentiable minimax fractional programming problems with (C, α, ρ, d) -convexity. *J. Optim. Theory Appl.* 129, 185–199 (2006)
24. Chinchuluun, A., Yuan, D. H. and Pardalos, P. M.: Optimality conditions and duality for nondifferentiable multiobjective fractional programming with generalized convexity. *Ann. Oper. Res.* 154, 133–147 (2007)
25. Pokharna, N. and Tripathi, I. P.: Optimality and duality for E-minimax fractional programming: application to multiobjective optimization. *J. Appl. Math. Comput.* 69, 2361–2388 (2023)
26. Feng, M. and Li, S. J.: Second-order strong Karush/Kuhn-Tucker conditions for proper efficiencies in multiobjective optimization. *J. Optim. Theory Appl.* 181, 766–786 (2019)
27. Clarke, F. H.: *Optimization and Nonsmooth Analysis*. John Wiley & Sons, Inc., New York (1983)
28. Ivanov, V. I.: Second-order optimality conditions for vector problems with continuously Fréchet differentiable data and second-order constraint qualifications. *J. Optim. Theory Appl.* 166, 777–790 (2015)
29. Páles, Z. and Zeidan, V. M.: Nonsmooth optimum problems with constraints. *SIAM J. Control Optim.* 32, 1476–1502 (1994)
30. Ivanov, V. I.: Second-order optimality conditions for inequality constrained problems with locally Lipschitz data. *Optim. Lett.* 4, 597–608 (2010)
31. Lebourg, G.: Valeur moyenne pour gradient généralisé. *C. R. Acad. Sci. Paris Sér. A-B.* 281, A1, A795–A797 (1975)
32. Rockafellar, R. T. and Wets, R. J.-B.: *Variational Analysis*. Springer-Verlag, Berlin (1998)
33. Aanchal and Lalitha, C. S.: Second-order optimality conditions for locally Lipschitz vector optimization problems. *Optim.* 0, 1–20 (2023)
34. Ehrgott, M.: *Multicriteria Optimization* (second edition). Springer-Verlag, Berlin (2005)
35. Borwein, J.: Proper efficient points for maximizations with respect to cones. *SIAM J. Control Optim.* 15, 57–63 (1977)
36. Sawaragi, Y., Nakayama, H. and Tanino, T.: *Theory of Multiobjective Optimization*. Academic Press, Inc., Orlando, FL (1985)
37. Penot, J. P.: Second-order conditions for optimization problems with constraints. *SIAM J. Control Optim.* 37, 303–318 (1999)
38. Jiménez, B. and Novo, V.: Optimality conditions in differentiable vector optimization via second-order tangent sets. *Appl. Math. Optim.* 49, 123–144 (2004)
39. Feng, M. and Li, S. J.: On second-order optimality conditions for continuously Fréchet differentiable vector optimization problems. *Optim.* 67, 2117–2137 (2018)
40. Luu, D.V.: Second-order optimality conditions and duality for nonsmooth multiobjective optimization problems. *Appl. Set-Valued Anal. Optim.* 4, 41–54 (2022)