# Goodness-of-Fit for Conditional Distributions: An Approach Using Principal Component Analysis and Component Selection

Rui Cui Faculty of Economics and Management East China Normal University Yuhao Li Department of Financial and Acturial Mathematics Xi'an Jiaotong-Liverpool University

#### Abstract

This paper introduces a novel goodness-of-fit test technique for parametric conditional distributions. The proposed tests are based on a residual marked empirical process, for which we develop a conditional Principal Component Analysis. The obtained components provide a basis for various types of new tests in addition to the omnibus one. Component tests that based on each component serve as experts in detecting certain directions. Smooth tests that assemble a few components are also of great use in practice. To further improve testing efficiency, we introduce a component selection approach, aiming to identify the most contributory components. The finite sample performance of the proposed tests is illustrated through Monte Carlo experiments. (JEL: C12, C52)

# 1. Introduction

In regression analysis, conditional distributions are fundamental for the construction of numerous economic models. They not only illuminate the relationship between the response variable and the covariates but also aid in comprehending other significant indicators like expectation, variation, quantile, and hazard rate. Notably, conditional distributions are utilized in a variety of economic models, encompassing categorical or discrete choice models (such

Acknowledgments: We are deeply indebted to Miguel A. Delgado for his advice and support at all stages of this paper. The first author is particularly grateful to Winfried Stute for his guidance and care during the stay in Giessen. We also thank: Carlos Velasco, Juan Carlos Escanciano, Jesús Gonzalo, Nazarii Salish, Junyi Peng Zhou, Julius Vainora, Minghai Mao, Xiaojun Song. All errors belong to us.

E-mail: rcui@fem.ecnu.edu.cn (Cui); yuhao.li@xjtlu.edu.cn (Li)

as (ordered) logit/probit models), count data regression models (for instance, Poisson, negative-binomial, hurdle, and zero-inflation models), and duration models (for example, Cox's model and the accelerated failure time model). In these models, the conditional distribution functions are usually assumed to possess a known functional form with some unknown parameters. Accurate estimation and inferences hinge upon the correct specification of the parametric functional form.

Let Y denote the response variable, and X is a vector of covariates. This paper presents an investigation of specification tests that are designed to ascertain whether the conditional distribution of Y given X belongs to a predetermined parametric family, indexed by some unknown parameters. Specifically, the tests are for

$$H_0: \Pr[Y \le y|X] = F(y|X;\theta_0), \text{ a.s for } y \in \mathbb{R}, \text{ and some } \theta \in \Theta$$

against

$$H_1: \Pr[Y \leq y | X] \neq F(y | X; \theta_0)$$
, a.s for  $y \in \mathbb{R}$ , and some  $\theta \in \Theta$ 

where  $\theta_0$  is the limit of a consistent estimator  $\hat{\theta}$ . We assume that the parameter space  $\Theta$  lies in  $\mathbb{R}^{d_{\theta}}$ , where  $d_{\theta}$  is a fixed positive integer.

While there is a substantial amount of literature on consistent testing of conditional expectation models, including prominent works by Bierens (1982, 1990); Bierens and Ploberger (1997); Stute (1997); Stinchcombe and White (1998); Delgado and Manteiga (2001); Fan and Li (2000); Escanciano (2006); Hardle and Mammen (1993); Horowitz and Härdle (1994); Hong and White (1995); Zheng (1996); Lavergne and Vuong (2000), the volume of literature on the consistent specification testing of conditional distribution models is relatively sparse. Particularly, only a handful of papers addressing this topic have been published, including works by Andrews (1997); Zheng (2000, 2012); Bierens and Wang (2012). It is worth noting that the frameworks presented in Andrews (1997) and Bierens and Wang (2012) share the same methodology.

The basic idea consists of transforming the conditional test to a continuum of unconditional moments. The approach is commonly referred to as the Integrated Conditional Moment (ICM) framework. Meanwhile, an alternative approach, known as kernel tests, has been proposed by Zheng (2000, 2012). The kernel tests involve comparing parametric functional forms with corresponding nonparametric or semi-parametric estimates.

Both of the two type tests are consistent against all possible alternatives, in which sense they are referred to as omnibus tests. However, despite consistency, the omnibus tests often lack of performance in practice. Their limitation is that they only suggest an overall rejection, but give no sign on the specific aspects of the data that brought this rejection. This raises the question: can we gain more information from the omnibus test by partitioning it into some pieces, each measuring some distinctive aspect of the data. For instance, if the omnibus test could be decomposed into orthogonal directions, each representing a unique direction of departure, we could investigate the significance of each direction in addition to the overall significance. This would be much more informative, offering a comprehensive understanding of the nature of the deviation.

Such a decomposition was initially introduced by Durbin and Knott (1972) through Principal Component Analysis (PCA). They applied functional PCA on the classical empirical process to test the goodness-of-fit of unconditional distributions, providing an orthogonal decomposition of the omnibus test into its Principal Components (PCs). Each PC represents deviations in a unique direction, providing a basis for more effective tests, including Neyman's type smooth tests and optimal directional tests. In later work, Durbin et al. (1975) expanded the application of the PCA method for testing composite hypothesis problems. Stute (1997) further demonstrated the significant application of PCA in goodness-of-fit tests by discussing the issue of checking the mean regression model.

While PCA is an effective tool for assessing goodness-of-fit, its full potential has not been fully explored in existing literature. Previous papers often selected components to form tests based on the experiential understanding that the first few components account for most of the deviation. This approach, however, may not be reliable in more complicated situations, such as when dealing with large-dimensional covariates, as the contributions of PCs can be dispersed in these cases. An alternative approach could be to develop a data-driven criterion to help determine the contribution of each component to the test statistic. Essentially, we need to determine which components are crucial and how to assemble them. This is challenging as it requires prior information about potential deviations in the data. Therefore, even though PCA provides much more information than the omnibus test itself, it raises new questions and challenges about how to use it most effectively.

This paper intends to investigate the application of PCA for power improvement, within the context of goodness-of-fit testing for conditional distributions. We propose a testing procedure that includes two key contributions: a PCA framework for checking conditional distributions and a method for selecting significant components. In addition, we propose smooth tests that are based on the aggregation of the selected components.

Specifically, the first contribution involves applying a functional PCA for a residual marked empirical process within the ICM framework. Our Principal Component Decomposition (PCD) method, which we term as "conditional PCD", differs from existing ones, as the inclusion of covariate effect in distribution functions changes the structure of the empirical process dramatically, making previous methods invalid.

The primary reason to chose the ICM approach is due to its insensitiveness to dimension when certain weighting functions are appropriately chosen. Notably, Escanciano (2006) suggests substituting the indicator weighting function with a projection weight, while Bierens (1982) proposes the use of the characteristic function. These dimension reduction methods can be incorporated into our approach to handle high-dimensional covariates.

The underperformance of omnibus tests can be readily explained through PCA. In the original process forming the omnibus test, the weight of the components, which is determined by their eigenvalues, diminishes quickly.

This suggests that the omnibus test significantly underweights the latter components, which exactly reveal certain high-frequency departures. As a result, high-frequency departures are scarcely detectable through the omnibus test, leading to its low power. However, when we have the components, we can assign their weights freely. They can be used individually to construct tests, as experts in detecting certain directions. Reweighted sums of them may also prove beneficial against a wide range of alternatives in smooth tests. In that regard, these components serve as building blocks for more powerful tests.

However, as mentioned earlier, determining the exact way to utilize these blocks for constructing tests is challenging. Our second contribution involves a novel "learning then testing" procedure that incorporates a sample splitting technique. In the learning sample, we ascertain the contribution of each component based on the p-values of component tests and select the most significant ones. Small p-values indicate significant deviations from the null. Meanwhile, in the testing sample, we formulate smooth tests that equally aggregate the selected components. Our simulation studies demonstrate that even this simplistic assembly can substantially elevate testing efficiency.

The structure of this paper is as follows. Section 2 provides an overview of existing omnibus ICM tests for conditional distributions. Our conditional PCD method and test statistics based on the components are introduced in Section 3. In section 4 we introduce the component selection technique. In section 5, we examine the finite-sample performance of the proposed test by means of a series of Monte-Carlo exercises. Section 6 presents the asymptotic results for both the components and the test statistics. Finally, section 7 concludes the paper.

# 2. Omnibus ICM Tests for Conditional Distributions

Let us consider a sample  $\{Y_i, X_i\}, i = 1, \dots, n$  which consists of i.i.d. realizations of  $\{Y, X\}$ . Here, Y is the real-valued response variable and X is the covariate vector. Our interest lies in understanding the conditional distribution

of Y given X. The ICM framework consists of transforming the conditional test of

$$\mathbb{E}\left[\mathbb{I}\{Y \le y\} - F(y|x;\theta_0)|x\right] = 0$$

to a continuum of unconditional moments of

$$\mathbb{E}\left[\left(\mathbb{I}\{Y \le y\} - F(y|x;\theta_0)\right)h(x)\right] = 0$$

where h(x) is a weighting function.

Following this approach, we introduce a residual marked empirical process:

$$R_n(y,v) = n^{-1/2} \sum_{i=1}^n h(v, X_i) M_i(y).$$
(2.1)

Here  $v \in \mathbb{R}^{d^{\theta}}$  is the auxiliary vector inside the weighting function, and the term  $M_i(y)$  is given by

$$M_i(y) = \mathbb{1}_{\{Y_i \le y\}} - F\left(y \mid X_i, \theta\right), \qquad (2.2)$$

representing a single event process after centering. Each  $M_i(y)$  shares a common covariance kernel with the Brownian Bridge.

This process include all the empirical processes that form the existing ICM tests. For example, if the h function is chosen as the indicator function  $\mathbb{1}_{\{X_i \leq x\}}$ , it yields Andrews' process, i.e.,

$$\alpha_n(y,x) := n^{-1/2} \sum_{i=1}^n \mathbb{1}_{\{X_i \le x\}} M_i(y).$$
(2.3)

His test is a conditional Kolmogorov statistic based on the estimated process:

$$CK = \sup_{y,x} \left| \hat{\alpha}_n(y,x) \right|, \tag{2.4}$$

and he establishes the weak convergence result of this test statistic under a pseudometric defined as (3.6) in his work.

However, Andrews' test encounters the issue of the curse of dimensionality due to the selection of the indicator function. Escanciano (2006) suggested a solution to overcome high-dimensionality problem by replacing the weighting function  $\mathbb{1}_{\{X_i \leq x\}}$  with projection weights. Hence, the process he considered is

$$\zeta_n(\beta, z, y) = n^{-1/2} \sum_{i=1}^n \mathbb{1}_{\{\beta^\top X_i \le z\}} M_i(y)$$

His test is a Cramer von-Mises type statistic with the empirical distribution as the integration measure, which is defined as:

$$CvM_{es} = \int (\hat{\zeta}_n(\beta, z, y))^2 F_{n,Y}(dy) F_{n,\beta}(dz) d\beta.$$
(2.5)

Bierens (1982) presented another commonly employed empirical process for testing conditional moments. This method uses the characteristic function as its weighting function, and the corresponding Cramer von-Mises statistic with standard multivariate normal distribution as the integration measure is:

$$CvM_{ch} = \int |\hat{Z}(\xi)|^2 d\mu(\xi),.$$
 (2.6)

where

$$\hat{Z}(\xi) = \int n^{-1/2} \sum_{i=1}^{n} \exp(i\xi^{\top} X_i) M_i(y) F_{n,Y}(dy)$$

and  $\mu(\xi)$  is the standard multivariate normal distribution.

It is also necessary to acknowledge the assumptions related to the conditional distribution model and the estimation process.

(A1).  $F(y \mid X_i, \theta)$  is differentiable in  $\theta$  on a neighborhood of  $\theta_0, \forall i \ge 1$ .

(A2). 
$$\sup_{(y,v)\in\mathbb{R}^{(d'+1)}}\sup_{\theta:\|\theta-\theta_0\|\leq r_n}\left\|\frac{1}{n}\sum_{i=1}^n\frac{\partial}{\partial\theta}F(y\mid X_i,\theta)h(v,X_i)-\Delta_0(y,v)\right\|\to 0, \ a.s.$$

for all sequences of positive constants  $\{r_n : n \ge 1\}$  such that  $r_n \to 0$ , where  $\Delta_0(y,v) = \int (\partial/\partial\theta) F(y \mid s, \theta_0) h(v, s) dF_X(s)$  and  $F_X$  denotes the marginal distribution of X.

(A3). 
$$\sup_{(y,v)\in\mathbb{R}^{(d'+1)}} \|\Delta_0(y,v)\| < \infty$$

and  $\Delta_0(\cdot)$  is uniformly continuous on  $\mathbb{R}^{(d'+1)}$ .

(A4). The parametric estimator has an expression as

$$n^{1/2} \left(\hat{\theta} - \theta_0\right) = n^{-1/2} \sum_{i=1}^n l(X_i, Y_i, \theta_0) + o_p(1), \text{ conditional on } X \text{ a.s.}$$

for some function l such that it is a measurable function and satisfies  $\int l(y, x, \theta_0) F(dy \mid x, \theta_0) = 0$  for all x in the support of X and  $\int l_0(x) F_X(dx) < \infty$ , where  $l_0(x) = \int ||l(y, x, \theta_0)||^{2+\varepsilon} F(dy \mid x, \theta_0)$  for some  $\varepsilon > 0$ .

Assumptions A1-A3 are standard in the literature, see, e.g., Andrews (1997), while assumption A4 stipulates that the maximum likelihood estimator meets the necessary conditions as an example, and is also widely adopted in the literature, see.e.g., Bierens (1982); Bierens and Ploberger (1997); Bierens and Wang (2012); Delgado and Manteiga (2001).

#### 3. Principal Component Analysis

Given that the process  $R_n$  is multivariate and includes dependent components y and v, it doesn't possess an explicit Karhunen-Loève representation. In this section, we aim to develop a conditional PCD argument for  $R_n$ . This can be realized in two steps: firstly, the single event process M(y)'s PCD could be obtained while conditioning on X, and subsequently, the obtained PCs need to be added up with respect to the observations of X.

#### 3.1. Conditional Principal Component Decomposition

Let's begin with the PCD of M(y) conditional on X. Under the null model, and with X held constant, the covariance kernel of M(y) is identical to the covariance kernel of a transformed Brownian Bridge. That is,

$$\mathbb{E}(M(y_1)M(y_2) \mid X) = F(y_1 \land y_2 \mid X, \theta_0) - F(y_1 \mid X, \theta_0)F(y_2 \mid X, \theta_0). \quad (3.1)$$

For each x, we define the transformation function as:

$$T(y,x) := F(y \mid X = x, \theta_0),$$
 (3.2)

Cui and Li

where  $\theta_0$  represents the true value of the parameters. The function T is nondecreasing with regards to y, and  $T(-\infty, x) = 0$ ,  $T(\infty, x) = 1$ .

Hence, we have

$$\mathbb{E}(M(y_1)M(y_2) \mid X = x) = K(T(y_1, x), T(y_2, x)),$$
(3.3)

where  $K(s,t) = s \wedge t - st$  denotes the covariance kernel of the standard Brownian Bridge.

The eigenfunctions of M(t) can be determined by transformation. Let:

$$\mu_j = \frac{1}{(\pi j)^2}, \quad \varphi_j(y) = \sqrt{2}sin(j\pi y), \quad j = 1, 2, \cdots$$
 (3.4)

represent the eigenvalues and eigenfunctions of the standard Brownian Bridge with the covariance kernel K(s,t). For each x, let  $f_j$  be the transformation:

$$f_j(y,x) := \varphi_j(T(y,x)). \tag{3.5}$$

Therefore, for every fixed x,  $\{f_j(\cdot, x)\}_{j=1}^{\infty}$  forms an orthonormal basis for a subspace of  $L^2(\mathbb{R}, T(\cdot, x))$ , which is the Hilbert space of all square integrable functions on  $\mathbb{R}$  with an inner product defined as:

$$\langle \rho, g \rangle_x = \int_{\mathbb{R}} \rho(y) g(y) T(dy, x),$$

since

$$\langle f_j, f_h \rangle_x = \int_{\mathbb{R}} \varphi_j \left( T(y, x) \right) \varphi_h \left( T(y, x) \right) T(dy, x)$$
$$= \int_0^1 \varphi_j(u) \varphi_h(u) du = \begin{cases} 1 & j = h \\ 0 & j \neq h \end{cases}.$$

Moreover,  $\{f_j(\cdot, x)\}_{j=1}^{\infty}$  are the eigenfunctions of the covariance kernel  $K(T(y_1, x), T(y_2, x))$  with associated eigenvalues  $\{\mu_j\}_{j=1}^{\infty}$ . That is,

$$\int_{\mathbb{R}} K(T(y_1, x), T(y_2, x)) f_j(y_1, x) T(dy_1, x) = \mu_j f_j(y_2, x) f_j(y_2, x) f_j(y_2, x) f_j(y_2, x) f_j(y_2, x) f_j(y_2, x) f_j(y_1, x) f_j(y_2, x) f_j(y_2, x) f_j(y_1, x) f_j(y_2, x) f_j(y_2, x) f_j(y_2, x) f_j(y_1, x) f_j(y_2, x) f_$$

By Mercer's theorem, the covariance kernel can be decomposed as

$$K(T(y_1, x), T(y_2, x)) = \sum_{j=1}^{\infty} \mu_j f_j(y_1, x) f_j(y_2, x).$$
(3.6)

Cui and Li

The Karhunen-Loève representation of the centered single event process is

$$M_i(y) = \sum_{j=1}^{\infty} \mu_j^{1/2} z_{ij} f_j(y, X_i) \quad a.s., \ i = 1, \cdots, n$$
(3.7)

where

$$z_{ij} := \mu_j^{-1/2} \langle M_i, f_j(\cdot, X_i) \rangle_{X_i}$$
  
=  $\mu_j^{-1/2} \int_{\mathbb{R}} M_i(y) f_j(y, X_i) T(dy, X_i).$  (3.8)

The  $z_{ij}$  is the j'th PC of  $M_i(y)$  conditional on  $X_i$ . For each j and  $j \neq h$ , it has the following properties

$$\mathbb{E}(z_{ij} \mid X_i) = 0,$$
$$\mathbb{E}(z_{ij}^2 \mid X_i) = 1,$$
(3.9)

$$\mathbb{E}(z_{ij}z_{ih} \mid X_i) = 0.$$

That is, for each *i*, the PCs have a conditional zero mean and unit variance, and are uncorrelated with each other given X. The next step involves summing up the obtained conditional PCs similarly to how  $R_n(y,x)$  has summed up  $M_i(y)$ . By substituting the conditional PCD (3.7) into equation (2.1), we can decompose  $R_n$ :

$$R_n(y,v) = n^{-1/2} \sum_{i=1}^n h(v,X_i) \left( \sum_{j=1}^\infty \mu_j^{1/2} z_{ij} f_j(y,X_i) \right)$$
$$= \sum_{j=1}^\infty \mu_j^{1/2} \left[ n^{-1/2} \sum_{i=1}^n z_{ij} h(v,X_i) f_j(y,X_i) \right].$$

We call the term in the square bracket the j'th component process of  $R_n$  and denote it as

$$c_{n,j}(y,v) := n^{-1/2} \sum_{i=1}^{n} z_{ij} h(v, X_i) f_j(y, X_i).$$
(3.10)

This result is summarized in the following proposition.

**Proposition 1** Under the null hypothesis, the processes (2.1) can be decomposed into a weighted sum of component processes, i.e.,

$$R_n(y,v) = \sum_{j=1}^{\infty} \mu_j^{1/2} c_{n,j}(y,v).$$
(3.11)

The weights are the square root of the standard Brownian Bridge eigenvalues.

Actually, in this distribution scenario, each PC  $z_{ij}$  takes an explicit form as described by:

$$z_{ij} = \sqrt{2} \cos(j\pi T(Y_i, X_i)).$$
 (3.12)

To elaborate, let us first define the cosine function  $g_j$  that corresponds to  $f_j$  as:

$$g_j(y,x) := \varphi_j(T(y,x)) = \sqrt{2}cos\left(j\pi T(y,x)\right).$$

By applying integration by parts to the integral (3.8), we have

$$z_{ij} = \int_{-\infty}^{\infty} g_j(s, X_i) dM_i(s) = g_j(Y_i, X_i).$$

Repeating (3.9) we have

$$\mathbb{E}(g_j(Y_i, X_i) \mid X_i) = 0,$$
$$\mathbb{E}(g_j^2(Y_i, X_i) \mid X_i) = 1,$$
$$\mathbb{E}(g_j(Y_i, X_i)g_h(Y_i, X_i) \mid X_i) = 0.$$

This result shows that the PCs can be obtained through a simple transformation of the data, namely  $g_j(Y_i, X_i)$ , and they have nice orthogonal structures. Furthermore, for each  $j \ge 1$ ,  $g_j(Y_i, X_i)$  have the same conditional distribution on  $X_i$ , the Fourier transform of which equal to the Besselfunction of order zero. In another word,  $g_j$  is a conditional distribution-free transformation. We can rewrite the component processes in a more concise way

$$c_{n,j}(y,v) = n^{-1/2} \sum_{i=1}^{n} h(v, X_i) f_j(y, X_i) g_j(Y_i, X_i).$$
(3.13)

## 3.2. Tests based on component processes

The component processes we derived provide a basis of new specification tests for conditional distributions. Our test will be based on the component process after estimation, i.e., the process

$$\hat{c}_{n,j}(y,v) := n^{-1/2} \sum_{i=1}^{n} h(v,X_i) \hat{f}_j(y,X_i) \hat{g}_j(Y_i,X_i).$$
(3.14)

Here  $\hat{f}_j(y,x) = \varphi_j\left(\hat{T}(y,x)\right)$ , and  $\hat{g}_j(y,x) = \varphi_j\left(\hat{T}(y,x)\right)$ , with  $\hat{T}(y,x)$  being an estimator of the function T(y,x). A natural consistent one is

$$\hat{T}(y,x) = F\left(y \mid X = x, \hat{\theta}\right).$$

The CvM type statistic can be constructed based on each component process. Specifically, for each  $j = 1, 2, \cdots$ , we have what we refer to as component tests

$$CvM_{n,j} = \int \hat{c}_{n,j}^2(y,v)F_{n,Y}(dy)\mu(dv).$$

Additionally, smooth test statistics can be constructed based on the reweighted sum of component processes. By assigning weights to the components and considering the sum of the first m components, the corresponding CvM type statistic<sup>1</sup> for some fixed m are

$$KS_{n,j} = \sup_{y,v} \left| \hat{c}_{n,j}(y,v) \right|$$

<sup>1.</sup> Similarly, KS type statistics can be written as

$$CvM_{nm} = \int \left(\sum_{j=1}^{m} w_j \hat{c}_{n,j}(y,v)\right)^2 F_{n,Y}(dy)\mu(dv).$$
(3.15)

A smooth test, which is similar to Neyman's smooth test, is the one that takes  $w = (1, \dots, 1)$ . While the *j*'th component test is the one that takes *w* as the *j*'th unit vector, i.e.,  $w = (0, \dots, 1, \dots, 0)$ . The smooth tests provide a compromise between the comprehensive omnibus tests and the individual component tests. However, a suitable weight vector *w* must be selected prior to model checking.

The asymptotic theories of the component processes and our tests will be discussed in section 6. Given their convergence, in oder to estimate the critical values of our tests in practice, we follow Andrews (1997) to run a parametric bootstrap.

Recall equation (3.11), where the weight for the *j*'th component process takes the value of  $\mu_j^{1/2}$ . This weight decreases rapidly with an increase in *j*, resulting in the later components being down-weighted in the original process. However, each component represents a specific aspect of deviation from the null hypothesis. For example, high-frequency deviations are more apparent in the latter components. Therefore, the omnibus test, which assigns lower weights to latter components, has reduced power to detect these high-frequency deviations. Whereas tests based on the latter components are specifically designed to detect such alternatives, effectively serving as a counterpoint to the limitations of the omnibus test. To better understand the role of components, we end this section with a simulation example. The simulation will demonstrate how the components act as specialized experts in detecting specific deviations.

and

$$KS_{nm} = \sup_{y,v} \Big| \sum_{j=1}^{m} w_j \hat{c}_{n,j}(y,x) \Big|$$

#### 3.3. Role of components: a simple simulation example

When testing unconditional normal distributions, Durbin and Knott (1972) proposed examining each PC based on the observation that different PCs respond to departures in different moments. Specifically, the first PC of the normal empirical process is sensitive to shifts in the mean, while the second PC is sensitive to shifts in variance. Similarly, the third and fourth PCs show sensitivity to shifts in skewness and kurtosis, respectively.

In this example, we're considering conditional normal models with a conditional mean and fixed variance, denoted as,

$$H_0: Y \mid X \sim \mathcal{N}(\theta_0 + \theta_1 X, \theta_2),$$

with parameters being  $\theta = (\theta_0, \theta_1, \theta_2)$ . We'll use real-valued X for simplicity, and generate Data Generating Processes (DGPs) that exhibit shifts in mean, variance, skewness, and kurtosis, respectively. This will enable us to observe how the components serve when testing conditional normal against deviations in moments.

The first two DGPs are specified as:

- DGP of Mean shift:  $Y \sim \mathcal{N}(1 + x + 5\cos(2x), 1);$
- DGP of Heteroscedasticity:  $Y \sim \mathcal{N}(1+x, 6(x-0.5)^2+0.5)$ .

The DGPs for skewness and kurtosis shifts are derived from the following conditional distribution function

$$F(y \mid x, \gamma_3, \gamma_4) = \Phi(y - 1 - x) + \gamma_3 \sin(3\pi\Phi(y - 1 - x)) + \gamma_4 \sin(4\pi\Phi(y - 1 - x)),$$

where  $\Phi(\cdot)$  is the distribution function of standard normal variable. This distribution is a conditional version of (8.4) in Durbin et al. (1975) and  $\gamma_3$  and  $\gamma_4$  represent deviations in skewness and kurtosis, respectively:

- DGP of Skewness shift:  $\gamma_3 = 0.1, \gamma_4 = 0;$
- DGP of Kurtosis shift:  $\gamma_3 = 0, \gamma_4 = 0.1$ .

 $CvM_{n,3}$ 

 $CvM_{n,4}$ 

0.042

0.074

0.064

0.066

Here, we only scrutinize the component test statistics. Critical values are procured by implementing standard parametric bootstrap procedures, and these details can be found in Section 5. The performance of the first four component tests is shown in Table 1 and 2, with the bold type indicating the test that has the highest power.

Variance Mean n = 100200 300 n = 100200 300  $CvM_{n,1}$ 0.444 0.904 0.1940.7800.2220.178 $CvM_{n,2}$ 0.0820.1180.1140.2640.4260.476

0.006

0.048

0.006

0.050

0.002

0.056

TABLE 1. Component Test Results, Mean and Variance Deviations

 $CvM_{n,k}, k = 1, 2, \ldots, 4$  are the k-th component test with n being the sample size.

0.066

0.084

_		Skewness			Kurtosis	
	n = 100	200	300	n = 100	200	300
$CvM_{n,1}$	0.618	0.906	0.968	0.024	0.030	0.028
$CvM_{n,2}$	0.104	0.162	0.198	0.078	0.110	0.098
$CvM_{n,3}$	0.838	0.992	1.000	0.054	0.070	0.068
$CvM_{n,4}$	0.118	0.178	0.192	0.226	0.292	0.360

TABLE 2. Component Test Results, Skewness and Kurtosis Deviations

 $CvM_{n,k},\,k=1,2,\ldots,4$  are the k-th component test with n being the sample size.

Similar patterns, as in the unconditional normal case, are observed. That is, the first four component tests each specialize in testing deviations in the first four moments respectively. Therefore, if we want to test conditional normal variables against deviations in moments, the component tests would be the first choice.

From this example, it's evident that understanding which components of a given data set have the most contribution to departures can greatly enhance the efficiency of testing. In typical cases, we often expect the data to be infrequent, which allows us to concentrate mainly on the first few components. However, this situation would be drastically altered when dealing with largedimensional covariates. In such a large-dimensional problem, not only the first few components are no longer reliable, but also the pattern of normal distribution where each PC is responsible for each moment, disappears. To address this, we propose a novel 'learning then testing' procedure in the subsequent section. This procedure aids us in acquiring preliminary insights about the data and pinpointing the key components. This knowledge can then be used to guide our testing procedure, potentially improving its efficiency.

#### 4. Component Selection

In earlier sections, we show that the residual process can be theoretically decomposed into a countable number of components or directions, but this does not inherently lead to an improvement in power performance without the expression of some information about the direction of the deviation. In general, it is often challenging for researchers to ascertain, a priori, the directions in which deviations are predominantly concentrated. In this section, we outline a straightforward strategy for acquiring such information. Our approach involves splitting the full sample into learning and testing sub-samples. In the learning sample, we can estimate the p-values corresponding to each component test by parametric bootstrapping the empirical null distribution. Subsequently, we can sort these p-values and select the first m components to facilitate the test in the testing sub-sample. We use the following pseudo code to illustrate this idea.

$\mathbf{A}$	lgorith	m 1	: ]	Learning	the	P-1	Va	lues
--------------	---------	-----	-----	----------	-----	-----	----	------

- 1 Input: Z = (X, Y), Bootstrap sample size B, component number M;
- **2** Split the sample into learning and testing parts:  $\{X_{learn}, Y_{learn}\}, \{X_{test}, Y_{test}\};$
- **3** Using the learning sample, obtain a consistent estimator  $\hat{\theta}_{learn}$ , as well as M component test statistics  $\{\hat{c}_j(\hat{\theta}_{learn})\}_{j \in [M]}$ ;
- 4 Perform parametric bootstrap in the learning sample for each component process and obtain the associated empirical distribution  $\{\hat{c}_{j,b}\}_{b\in[B]}$ ;
- 5 Calculate p-values:  $p_j = mean(\hat{c}_j(\hat{\theta}_{learn}) \leq \{\hat{c}_{j,b}\}_{b \in [B]})$ ;
- **6** Sort these p-values in ascend order, select the first m components, record the corresponding indexes and form a vector  $\mathcal{W}$ ;

# Algorithm 2: Testing Procedures

- 1 **Input:**  $Z_{test} = (X_{test}, Y_{test})$ , Bootstrap sample size B, a vector of component index W learned from Algorithm 1, significance level  $\alpha$ ;
- 2 Using the testing sample, obtain a consistent estimator  $\hat{\theta}_{test}$ , as well as component test statistics  $\{\hat{c}_j(\hat{\theta}_{test})\}_{j \in \mathcal{W}}$ ;
- **3** Construct an equal weight smooth test statistic  $\bar{c} = mean(\{\hat{c}_j(\theta_{test})\}_{j \in \mathcal{W}});$
- 4 Perform parametric bootstrap in the testing sample for each component process in  $\mathcal{W}$  and obtain the associated bootstrap statistic  $\{\hat{c}_{j,b}\}_{j \in \mathcal{W}, b \in [B]}$ ;
- 5 Obtain the empirical distribution for the smooth statistic:  $\{\bar{c}_b\}_{b\in[B]}$ , where  $\bar{c}_b = mean(\{\hat{c}_{j,b}\}_{j\in\mathcal{W}})$ ;
- 6 Calculate the p-value  $p = mean(\bar{c} \leq \{\bar{c}_b\}_{b \in [B]})$ , reject the null if  $p < \alpha$ .

During the learning procedures, it is crucial to scrutinize the quality of outcomes, specifically the variability in p-values. There are several methodologies available that can be adopted to improve the overall stability of the learning process. An intuitive strategy might be to increase the size of the learning sub-sample. However, such expansion would invariably lead to a decrease in the size of the testing sub-sample, potentially causing a reduction in power. The problem of how to optimally partition these two sub-samples is a complex issue and falls outside the scope of this paper.

Another option is the implementation of bagging. This technique involves repeatedly resampling the learning sub-sample and reapplying Algorithm 1 to yield a collection of p-values for each component process. These p-values are subsequently averaged, effectively reducing the variance of the p-values for each component. However, this method does come with a downside, which is the increased computational demand.

During the testing procedures, a significant yet unresolved issue relates to the assignment of weights to each component. In this study, we adopt a simple, though somewhat naive approach of conferring equal weights. Ideally, components with smaller p-values, suggesting substantial deviations from the null hypothesis, should receive greater weights. Doing so would ensure these deviations are magnified within the smooth test statistics. However, it must be underlined that the weight assignment also influences the variations seen in the test statistics. It is plausible that an inappropriate assignment of weights could exacerbate variance, ultimately leading to reduced power. Presently, our understanding of how these weights impact variance or the optimal methodology for their determination is limited. These topics mandate further investigation.

## 5. Simulation

In this section, we will examine our proposed test statistics by using Monte Carlo simulation. We list the detailed DGPs below.

- $DGP_0: Y \sim \mathcal{N}(1 + \beta^\top x, 1);$
- $DGP_1: Y \sim \mathcal{N}(1 + \beta^\top x, (2\beta^\top (x 0.5))^2);$
- $DGP_2: Y \sim Exp(1 + \beta^\top x);$
- $DGP_3 DGP_5 : Y \sim \mathcal{N}(1 + \beta^\top x + 5\cos(k\beta^\top x), 3.5);$
- $DGP_6 DGP_8 : Y \sim \mathcal{N}(1 + 5\cos(k\beta^\top x), (2\beta^\top (x 0.5))^2);$

Each DGP generates the covariates  $x \in \mathbb{R}^d$  from a uniform distribution U(0,1). We set the dimension of x at d = 15. All coefficients are set equal to one:  $\beta = 1$ .

 $DGP_1$  and  $DGP_2$  investigate the deviations in mean and variance, as well as a distributional deviation, respectively. On the other hand,  $DGP_3 - DGP_5$ examine only the mean deviation, but with varying frequencies, specifically k = 3, 4, 5. Lastly,  $DGP_6 - DGP_8$  with k = 5, 6, 7 are employed to illustrate that while learning can generally enhance power properties, it is not universally the most potent procedure.

We analyze the proposed smooth test both with and without the application of the learning then testing procedure. The characteristic function is selected as the weighting function h(v, X), and the standard multivariate normal distribution is leveraged as the integration measure  $\mu(v)$ .

In addition to examining the proposed smooth test statistics, we also scrutinizes three omnibus test statistics, namely: (i) Andrew's CK test, (ii) CvM type test with Escanciano's projection weighting function, and (iii) CvM type test with characteristic weighting function. These tests are respectively defined by equations (2.4)-(2.6). Table 3 presents the estimated size at the 95% confidence level, with the sample size ranging from n = 100 to n = 300. These results imply that the proposed component tests, including the learning procedure, have minor size distortions. Furthermore, these distortions tend to decrease as the sample size increases.

The rest of the tables (Tables 4 to 7) report the power performance of various test statistics. We draw the following conclusions from them.

Firstly, component-based tests generally outperform omnibus tests, but they aren't consistently the most powerful. For instance, in DGPs 3-5 (Tables 5 and 6), three omnibus tests (CK,  $CvM_{ch}$  and  $CvM_{es}$ ) with identity, projected identity and characteristic functions as weighting functions, respectively, have negligible power. In contrast, component-based tests, whether using 5 or 10 components or employing the learning procedure, exhibit substantial power. On the other hand, DGPs 6-8, as seen in Tables 6-7, two omnibus tests ( $CvM_{ch}$ and  $CvM_{es}$ ) exhibit superior performance. This can be explained by the data in the appendix Tables B.3 and B.4. In these DGPs, the first two components show strong deviations. The omnibus test statistic assigns descending weights to components, with the first component receiving the highest weight. In this scenario, the most significant deviations receive the most critical weights. In contrast, component-based tests treat all components equally, resulting in a loss of power.

Secondly, in component-based tests, the learning procedure typically boosts power performance. This improvement is most observable in DGPs 6-8 (Tables 6-7). For other DGPs, it appears that the learning procedure results in virtually identical power performance to the component test with the first 5 components. However, we encountered a challenge with the learning procedure in terms of the stability of the results. The p-values for each iteration of the Monte Carlo exercises fluctuated considerably for some DGPs. This instability could potentially explain the under-performance of the learning procedure. A possible solution is to employ the application of bagging, which serves to reduce the variance of the p-values for each component. However, this approach does present a cost in the form of computational burden.

Lastly, the quantity and weighting of each component play significant roles. This importance becomes clear in  $DGP_6 - DGP_8$ , as summarized in Table 8. One could determine the number of components based on their respective p-values. For instance, in these DGPs, the p-values for the first two components are considerably lower than the others, suggesting that using only two components might enhance power performance. Our simulation results support this hypothesis, as we noted a substantial increase in power. The weighting of each component is also crucial. In these DGPs, only the first two components have significant deviation, implying that the omnibus test statistic could approximate a two-component test statistic with eigenvalue weightings. Comparing the power performance of  $CvM_{ch}$  and Smoooth\_Learn\_2 reveals that weightings significantly impact the results. However, determining the optimal weightings is a complex task and is a topic for future studies.

	C	l = 15	
	n = 100	200	300
СК	0.046	0.058	0.046
$CvM_{ch}$	0.088	0.068	0.052
$CvM_{es}$	0.074	0.044	0.056
Smooth_5	0.088	0.052	0.056
Smooth_10	0.08	0.062	0.062
Smooth_Learn_5	N/A	0.068	0.056

TABLE 3. Estimated size for DGP0 at 95%

The Smooth\_k measure is constructed by considering the first k components with equal weights. Furthermore, for learning the characteristics of the underlying testing problems, the first 50 observations are utilized. The Smooth\_Learn\_k measure takes into account the largest k components with a sample size n - 50.

		$DGP_1$			$DGP_{2}$		
-		200	300	100	- 100 200		
	n = 100	200	300	n = 100	200	300	
СК	0.598	0.446	0.258	0.018	0.038	0.048	
$CvM_{ch}$	0.978	0.994	0.988	1.000	1.000	1.000	
$CvM_{es}$	0.754	0.818	0.846	1.000	1.000	1.000	
Smooth_5	0.966	0.998	1.000	0.774	0.870	0.916	
Smooth_10	0.984	1.000	1.000	0.574	0.860	0.962	
Smooth_Learn_5	N/A	0.992	1.000	N/A	0.962	0.976	

TABLE 4. Estimated power for  $DGP_1$  and  $DGP_2$  at 95%

The Smooth\_k measure is constructed by considering the first k components with equal weights. Furthermore, for learning the characteristics of the underlying testing problems, the first 50 observations are utilized. The Smooth\_Learn\_k measure takes into account the largest k components with a sample size n - 50.

		$DGP_3$				
	n = 100	200	300	n = 100	200	300
СК	0.008	0.034	0.058	0.016	0.048	0.050
$CvM_{ch}$	0.052	0.080	0.054	0.066	0.036	0.034
$CvM_{es}$	0.046	0.054	0.048	0.042	0.028	0.026
Smooth_5	0.092	0.216	0.216	0.094	0.170	0.238
Smooth_10	0.098	0.174	0.188	0.088	0.142	0.194
Smooth_Learn_5	N/A	0.192	0.312	N/A	0.196	0.300

TABLE 5. Estimated power for  $DGP_3$  and  $DGP_4$  at 95%

The Smooth\_k measure is constructed by considering the first k components with equal weights. Furthermore, for learning the characteristics of the underlying testing problems, the first 50 observations are utilized. The Smooth\_Learn\_k measure takes into account the largest k components with a sample size n - 50.

#### 6. Asymptotic Theory

In this section, we develop asymptotic results of the component processes and the test statistics. From (3.13), each component process is a sum of i.i.d. centered random functions with variance

$$H_j(y,v) := \mathbb{E}\left[h^2(v,X)f_j^2(y,X)\right] = \int h^2(v,s)f_j^2(y,s)F_X(ds),$$

where  $F_X(\cdot)$  denotes the distribution function of X. Following Andrews (1997), we have the below theorem.

		DCP-			DGPc		
-		DGF5			DG16		
	n = 100	200	300	n = 100	200	300	
СК	0.022	0.036	0.056	0.572	0.394	0.260	
$CvM_{ch}$	0.064	0.046	0.046	0.888	0.914	0.948	
$CvM_{es}$	0.040	0.030	0.030	0.570	0.636	0.656	
Smooth_5	0.108	0.192	0.244	0.412	0.454	0.536	
Smooth_10	0.092	0.144	0.164	0.284	0.376	0.518	
Smooth_Learn_5	N/A	0.180	0.280	N/A	0.606	0.630	

TABLE 6. Estimated power for  $DGP_5$  and  $DGP_6$  at 95%

The Smooth\_k measure is constructed by considering the first k components with equal weights. Furthermore, for learning the characteristics of the underlying testing problems, the first 50 observations are utilized. The Smooth\_Learn\_k measure takes into account the largest k components with a sample size n - 50.

		$DGP_7$		$DGP_8$		
	n = 100	200	300	n = 100	200	300
СК	0.510	0.382	0.264	0.536	0.412	0.240
$CvM_{ch}$	0.856	0.910	0.920	0.906	0.918	0.932
$CvM_{es}$	0.532	0.628	0.688	0.512	0.630	0.644
Smooth_5	0.362	0.448	0.488	0.430	0.422	0.494
Smooth_10	0.250	0.370	0.466	0.278	0.348	0.488
Smooth_Learn_5	N/A	0.640	0.620	N/A	0.640	0.598

TABLE 7. Estimated power for  $DGP_7$  and  $DGP_8$  at 95%

The Smooth\_k measure is constructed by considering the first k components with equal weights. Furthermore, for learning the characteristics of the underlying testing problems, the first 50 observations are utilized. The Smooth\_Learn\_k measure takes into account the largest k components with a sample size n - 50.

**Theorem 1** Under the null hypothesis and (A1)-(A3), for each j, the process  $c_{n,j}(y,v)$  converges weakly to a centered Gaussian process

$$c_{n,j} \stackrel{d}{\to} c_{\infty,j}.$$

The limit Gaussian process  $c_{\infty,j}$  has covariance structure

$$K(y_1, y_2, v_1, v_2) = \int h(v_1, s) h(v_2, s) f_j(y_1, s) f_j(y_2, s) F_X(ds).$$

Moreover,  $c_{\infty,j}$  and  $c_{\infty,h}$  are independent for  $j \neq h$ .

		n = 300	
	$DGP_6$	$DGP_7$	$DGP_8$
$CvM_{ch}$	0.948	0.920	0.932
$CvM_{es}$	0.656	0.688	0.644
Smooth_5	0.536	0.488	0.494
Smooth_10	0.518	0.466	0.488
Smooth_Learn_5	0.630	0.620	0.598
Smooth_Learn_2	0.714	0.726	0.738

TABLE 8. Component Number and Weights

The Smooth\_k measure is constructed by considering the first k components with equal weights. Furthermore, for learning the characteristics of the underlying testing problems, the first 50 observations are utilized. The Smooth\_Learn\_k measure takes into account the largest k components with a sample size n - 50.

Now, let's consider the component process after estimation. As shown in the Appendix,  $\hat{c}_{n,j}(y,x)$  has the same asymptotic distribution as

$$\tilde{c}_{n,j}(y,v) := c_{n,j}(y,v) - A_j(y,v)n^{-1/2}\sum_{i=1}^n l(X_i, Y_i, \theta_0),$$

where

$$A_j(y,v) = \mu_j^{-1/2} \mathbb{E}\left[h(v,X)f_j(y,X)\int_{\mathbb{R}} F_{\theta}(y \mid X, \theta_0)f_j(y,X)T(dy,X)\right]$$

with  $F_{\theta}$  denoting the partial derivative w.r.t.  $\theta$ .

**Theorem 2** Under the null hypothesis and (A1)-(A4), for each  $j = 1, 2, \cdots$ , the process  $\hat{c}_{n,j}(y, v)$  converges weakly to a centered Gaussian process

$$\hat{c}_{n,j} \stackrel{d}{\to} \tilde{c}_{\infty,j}.$$

The limit Gaussian process  $\tilde{c}_{\infty,j}(y,v)$  has covariance structure

$$K(y_1, y_2, v_1, v_2) = \mathbb{E}\Big[h(v_1, X)f_j(y_1, X)g_j(Y, X) - A_j(y_1, v_1)l(X, Y, \theta_0),$$
$$h(v_2, X)f_j(y_2, X)g_j(Y, X) - A_j(y_2, v_2)l(X, Y, \theta_0)\Big].$$

As for the reweighted process  $\sum_{j=1}^{m} w_j \hat{c}_{n,j}(y,v)$ , it has the same asymptotic distribution with the process  $\sum_{j=1}^{m} w_j \tilde{c}_{n,j}(y,v)$ . Its asymptotic distribution is given in the following theorem.

**Theorem 3** Under the null hypothesis and (A1)-(A4), for any given weight  $w = \{w_j\}_{j=1}^m$ , the process  $\sum_{j=1}^m w_j \hat{c}_{n,j}(y,v)$  converges weakly to a centered Gaussian process

$$\sum_{j=1}^m w_j \hat{c}_{n,j} \stackrel{d}{\to} \tilde{c}_{\infty}^w.$$

The limit Gaussian process  $\tilde{c}^w_{\infty}(y, x)$  has covariance structure

$$K(y_1, y_2, x_1, x_2) = \mathbb{E}\left[\sum_{j=1}^m w_j \left(h(v_1, X)f_j(y_1, X)g_j(Y, X) - A_j(y_1, v_1)l(X, Y, \theta_0)\right), \\ \sum_{j=1}^m w_j \left(h(v_2, X)f_j(y_2, X)g_j(Y, X) - A_j(y_2, v_2)l(X, Y, \theta_0)\right)\right].$$

The convergence of our test statistics are then straightforward from the continuous mapping theorem.

# 7. Conclusion

The primary interest of this paper is to answer two out of three questions in the framework of checking conditional distributions: (i) whether the omnibus statistics can be partitioned into a suitable set of individual components, each measuring some distinctive aspect of the data; (ii) given the partition of the statistics, how to identify which components encapsulate more discrepancy information and (iii) how to optimally design a new test statistic informed by this identification. We address the first two of these issues in this paper by conducting a conditional PCA of the omnibus test. The components serve as a foundation for more robust specification tests. We further propose a straightforward learning procedure to address the challenge of determining the importance of each component. In rough terms, this strategy involves constructing a series of directional test statistics and bootstrapping their corresponding p-values. Small p-values indicate significant deviations in the corresponding directions, which we interpret as signals. Conversely, large p-values indicate noise in the respective directions.

In an ideal scenario, these directions should be weighted based on the quantity of signals they contain. However, determining optimal weights is a complex task and falls outside the scope of this paper. Hence, we adopt a simplistic strategy of assigning equal weights, which, despite its naivety, provides an efficient solution within the constraints of our study.

## References

- Andrews, Donald WK (1997). "A conditional Kolmogorov test." Econometrica: Journal of the Econometric Society, pp. 1097–1128.
- Bai, Jushan (2003). "Testing parametric conditional distributions of dynamic models." *Review of Economics and Statistics*, 85(3), 531–549.
- Bickel, Peter J, Chris AJ Klaassen, Peter J Bickel, Ya'acov Ritov, J Klaassen, Jon A Wellner, and YA'Acov Ritov (1993). Efficient and adaptive estimation for semiparametric models, vol. 4. Springer.
- Bickel, Peter J and Michael J Wichura (1971). "Convergence criteria for multiparameter stochastic processes and some applications." The Annals of Mathematical Statistics, 42(5), 1656–1670.
- Bierens, Herman J (1982). "Consistent model specification tests." Journal of Econometrics, 20(1), 105–134.
- Bierens, Herman J (1990). "A consistent conditional moment test of functional form." Econometrica: Journal of the Econometric Society, pp. 1443–1458.
- Bierens, Herman J and Werner Ploberger (1997). "Asymptotic theory of integrated conditional moment tests." *Econometrica: Journal of the Econometric Society*, pp. 1129– 1151.
- Bierens, Herman J and Li Wang (2012). "Integrated conditional moment tests for parametric conditional distributions." *Econometric Theory*, 28(2), 328–362.
- Billingsley, Patrick (2013). Convergence of probability measures. John Wiley & Sons.
- Delgado, Miguel A and Wenceslao González Manteiga (2001). "Significance testing in nonparametric regression based on the bootstrap." The Annals of Statistics, 29(5), 1469– 1507.
- Delgado, Miguel A and Winfried Stute (2008). "Distribution-free specification tests of conditional models." Journal of Econometrics, 143(1), 37–55.
- Durbin, James (1973). "Weak convergence of the sample distribution function when parameters are estimated." The Annals of Statistics, pp. 279–290.
- Durbin, James and Martin Knott (1972). "Components of Cramér-von Mises statistics. I." Journal of the Royal Statistical Society: Series B (Methodological), 34(2), 290–307.
- Durbin, James, Martin Knott, and CC Taylor (1975). "Components of Cramér-Von Mises Statistics. Ii." Journal of the Royal Statistical Society: Series B (Methodological), 37(2), 216–237.
- Escanciano, J Carlos (2006). "A consistent diagnostic test for regression models using projections." *Econometric Theory*, 22(6), 1030–1051.
- Escanciano, J Carlos (2009). "On the lack of power of omnibus specification tests." *Econometric Theory*, 25(1), 162–194.
- Eubank, Randall L and Jeffrey D Hart (1992). "Testing goodness-of-fit in regression via order selection criteria." The annals of Statistics, pp. 1412–1425.

- Eubank, RL and VN LaRiccia (1992). "Asymptotic comparison of Cramer-von Mises and nonparametric function estimation techniques for testing goodness-of-fit." *The Annals* of Statistics, 20(4), 2071–2086.
- Fan, Yanqin and Qi Li (2000). "Consistent model specification tests: Kernel-based tests versus Bierens' ICM tests." *Econometric Theory*, 16(6), 1016–1041.
- Grenander, Ulf (1950). "Stochastic processes and statistical inference." Arkiv för matematik, 1(3), 195–277.
- Hardle, Wolfgang and Enno Mammen (1993). "Comparing nonparametric versus parametric regression fits." The Annals of Statistics, pp. 1926–1947.
- Hong, Yongmiao and Halbert White (1995). "Consistent specification testing via nonparametric series regression." *Econometrica: Journal of the Econometric Society*, pp. 1133–1159.
- Horowitz, Joel L and Wolfgang Härdle (1994). "Testing a parametric model against a semiparametric alternative." *Econometric theory*, 10(5), 821–848.
- Huber, Peter J (1985). "Projection pursuit." The annals of Statistics, pp. 435–475.
- Inglot, Tadeusz and Teresa Ledwina (1996). "Asymptotic optimality of data-driven Neyman's tests for uniformity." The Annals of Statistics, 24(5), 1982–2019.
- Janssen, Arnold (2000). "Global power functions of goodness of fit tests." The Annals of Statistics, 28(1), 239–253.
- Jennrich, Robert I (1969). "Asymptotic properties of non-linear least squares estimators." The Annals of Mathematical Statistics, 40(2), 633–643.
- Kac, Marc and AJF Siegert (1947). "An explicit representation of a stationary Gaussian process." The Annals of Mathematical Statistics, 18(3), 438–442.
- Kallenberg, Wilbert CM and Teresa Ledwina (1997). "Data-driven smooth tests when the hypothesis is composite." *Journal of the American Statistical Association*, 92(439), 1094–1104.
- Khmaladze, Estate V (1982). "Martingale approach in the theory of goodness-of-fit tests." Theory of Probability & Its Applications, 26(2), 240–257.
- Khmaladze, Estate V (1993). "Goodness of fit problem and scanning innovation martingales." The Annals of Statistics, pp. 798–829.
- Lavergne, Pascal and Quang Vuong (2000). "Nonparametric significance testing." Econometric Theory, 16(4), 576–601.
- Ledwina, Teresa (1994). "Data-driven version of Neyman's smooth test of fit." Journal of the American Statistical Association, 89(427), 1000–1005.
- Neyman, Jerzy (1937). "Smooth test for goodness of fit." Scandinavian Actuarial Journal, 1937(3-4), 149–199.
- Pollard, David (2012). Convergence of stochastic processes. Springer Science & Business Media.

- Rosenblatt, Murray (1952). "Remarks on a multivariate transformation." The annals of mathematical statistics, 23(3), 470–472.
- Schoenfeld, David (1980). "Tests based on linear combinations of the orthogonal components of the Cramér-von Mises statistic when parameters are estimated." The Annals of Statistics, pp. 1017–1022.
- Schoenfeld, David A (1977). "Asymptotic properties of tests based on linear combinations of the orthogonal components of the Cramer-von Mises statistic." The Annals of Statistics, pp. 1017–1026.
- Stinchcombe, Maxwell B and Halbert White (1998). "Consistent specification testing with nuisance parameters present only under the alternative." *Econometric theory*, 14(3), 295–325.
- Stute, W, WL Xu, and LX Zhu (2008). "Model diagnosis for parametric regression in highdimensional spaces." *Biometrika*, 95(2), 451–467.
- Stute, Winfried (1993). "Consistent estimation under random censorship when covariables are present." Journal of Multivariate Analysis, 45(1), 89–103.
- Stute, Winfried (1997). "Nonparametric model checks for regression." The Annals of Statistics, pp. 613–641.
- Stute, Winfried and TL Anh (2012). "Principal component analysis of martingale residuals." J. Indian Statist. Assoc, 50(1-2), 263–276.
- Stute, Winfried, Silke Thies, and Li-Xing Zhu (1998). "Model checks for regression: an innovation process approach." The Annals of Statistics, 26(5), 1916–1934.
- Stute, Winfried and Li-Xing Zhu (2002). "Model checks for generalized linear models." Scandinavian Journal of Statistics, 29(3), 535–545.
- Zheng, John Xu (1996). "A consistent test of functional form via nonparametric estimation techniques." Journal of Econometrics, 75(2), 263–289.
- Zheng, John Xu (2000). "A consistent test of conditional parametric distributions." Econometric Theory, 16(5), 667–691.
- Zheng, Xu (2012). "Testing parametric conditional distributions using the nonparametric smoothing method." Metrika, 75, 455–469.

# **Appendix A: Proofs**

# **Proof of Theorem 1:**

Note that each  $f_j$  and  $g_j$  are bounded and differentiable. From (3.13) the weak convergence of  $c_{n,j}$  follows from functional CLT. The independence between  $c_{\infty,j}$  and  $c_{\infty,h}$  comes from the Gaussian property and conditional uncorrelation between  $z_{ij}$  and  $z_{ih}$ .

# **Proof of the asymptotic equivalence of** $\hat{c}_{n,j}(y,x)$ and $\tilde{c}_{n,j}(y,x)$ :

First note that the sine and cosin functions are bounded and differentiable. Since  $\hat{\theta}$  is root n-consistent,  $\hat{f}_j$  and  $\hat{g}_j$  are uniformly consistent.

We can write

$$\hat{c}_{n,j}(y,x) - n^{-1/2} \sum_{i=1}^{n} \mathbb{1}_{\{X_i \le x\}} f_j(t,X_i) g_j(Y_i,X_i)$$

$$= \left( \hat{c}_{n,j}(y,x) - n^{-1/2} \sum_{i=1}^{n} \int_0^\infty \mathbb{1}_{\{X_i \le x\}} \hat{f}_j(t,X_i) \hat{g}_j(s,X_i) dM_i(s) \right)$$

$$+ \left( n^{-1/2} \sum_{i=1}^{n} \int_0^\infty \mathbb{1}_{\{X_i \le x\}} \hat{f}_j(t,X_i) \hat{g}_j(s,X_i) dM_i(s) \right)$$

$$- n^{-1/2} \sum_{i=1}^{n} \int_0^\infty \mathbb{1}_{\{X_i \le x\}} f_j(t,X_i) g_j(s,X_i) dM_i(s) \right).$$

Cui and Li

For the first differnce, take the Taylor's expansion of  $\hat{c}_{n,j}(y,x)$  and then we have

$$\begin{aligned} \hat{c}_{n,j}(y,x) &= n^{-1/2} \sum_{i=1}^{n} \int_{-\infty}^{\infty} \mathbb{1}_{\{X_i \le x\}} \hat{f}_j(t,X_i) \hat{g}_j(s,X_i) dM_i(s) \\ &- \mu_j^{-1/2} n^{-1} \sum_{i=1}^{n} \mathbb{1}_{\{X \le x\}} \hat{f}_j(y,X) \int_{-\infty}^{\infty} F_{\theta}(y \mid X,\theta_0) \hat{f}_j(y,X) \hat{T}(dy,X) \\ &\times n^{-1/2} \sum_{i=1}^{n} l(X_i,Y_i,\theta_0) + o_p(1) \\ &= n^{-1/2} \sum_{i=1}^{n} \int_{-\infty}^{\infty} \mathbb{1}_{\{X_i \le x\}} \hat{f}_j(t,X_i) \hat{g}_j(s,X_i) dM_i(s) \\ &- A_j(y,x) n^{-1/2} \sum_{i=1}^{n} l(X_i,Y_i,\theta_0) + o_p(1) \end{aligned}$$

For the second difference, let us denote

$$K(s) = s - s^2,$$

then

 $\rightarrow$ 

$$E\left[n^{-1/2}\sum_{i=1}^{n}\int_{-\infty}^{\infty} \left(\mathbbm{1}_{\{X_{i}\leq x\}}\hat{f}_{j}(y,X_{i})\hat{g}_{j}(s,X_{i}) - \mathbbm{1}_{\{X_{i}\leq x\}}f_{j}(y,X_{i})g_{j}(s,X_{i})\right)dM_{i}(s)\right]^{2}$$

$$= E\left[n^{-1/2}\sum_{i=1}^{n}\int_{-\infty}^{\infty} \left(\mathbbm{1}_{\{X_{i}\leq x\}}\hat{f}_{j}(y,X_{i})\hat{g}_{j}(s,X_{i}) - \mathbbm{1}_{\{X_{i}\leq x\}}f_{j}(y,X_{i})g_{j}(s,X_{i})\right)^{2} \\ \times K(T(ds,X_{i}))\right]$$

$$= \int_{-\infty}^{\infty} E\left[\left(\mathbbm{1}_{\{X_{i}\leq x\}}\hat{f}_{j}(y,X_{i})\hat{g}_{j}(s,X_{i}) - \mathbbm{1}_{\{X_{i}\leq x\}}f_{j}(y,X_{i})g_{j}(s,X_{i})\right)^{2}K(T(ds,X_{i}))\right]$$

0.

$$n^{-1/2} \sum_{i=1}^{n} \int_{-\infty}^{\infty} \left( \mathbbm{1}_{\{X_i \le x\}} \hat{f}_j(y, X_i) \hat{g}_j(s, X_i) - \mathbbm{1}_{\{X_i \le x\}} f_j(y, X_i) g_j(s, X_i) \right) dM_i(s) = o_p(1).$$

Thus,

$$\hat{c}_{n,j}(y,x) = n^{-1/2} \sum_{i=1}^{n} \mathbb{1}_{\{X_i \le x\}} f_j(t,X_i) g_j(Y_i,X_i) - A_j(y,x) n^{-1/2} \sum_{i=1}^{n} l(X_i,Y_i,\theta_0) + o_p(1) d_{i,j}(y,x) - A_j(y,x) n^{-1/2} d_{i,j}(y,x) - A_j(y,x) - A_j(y,x) - A_j(y,x) - A_j(y,x) n^{-1/2} d_{i,j}(y,x) - A_j(y,x) - A_j($$

The weak convergence of  $c_{n,j}(y,x)$  and  $\hat{c}_{n,j}(y,x)$  is then easy to obtain.

# Appendix B: Component Test Results

This appendix documents size and power results for different components. These results indicate features of the underlying testing problems.

	-	$DGP_0$		1		
	n = 100	200	300	n = 100	200	300
$CvM_{n,1}$	0.098	0.034	0.042	0.972	0.982	0.986
$CvM_{n,2}$	0.082	0.054	0.054	0.998	1.000	1.000
$CvM_{n,3}$	0.064	0.076	0.056	0.304	0.296	0.536
$CvM_{n,4}$	0.082	0.060	0.066	0.946	1.000	1.000
$CvM_{n,5}$	0.052	0.046	0.056	0.404	0.626	0.822
$CvM_{n,6}$	0.076	0.052	0.060	0.882	0.994	1.000
$CvM_{n,7}$	0.056	0.072	0.054	0.472	0.738	0.864
$CvM_{n,8}$	0.084	0.056	0.076	0.828	0.976	1.000
$CvM_{n,9}$	0.060	0.062	0.036	0.414	0.736	0.904
$CvM_{n,10}$	0.094	0.028	0.062	0.750	0.944	1.000

TABLE B.1. Component Test Results,  $DGP_0$  and  $DGP_1$ 

 $PC_k, k = 1, 2, \dots, 10$  are tests using the  $k^{th}$  component.

		$DGP_2$		$DGP_3$			
	n = 100	200	300	n = 100	200	300	
$CvM_{n,1}$	0.996	1.000	1.000	0.064	0.064	0.070	
$CvM_{n,2}$	0.890	0.968	0.998	0.008	0.014	0.016	
$CvM_{n,3}$	0.676	0.978	1.000	0.176	0.306	0.392	
$CvM_{n,4}$	0.632	0.792	0.924	0.042	0.058	0.070	
$CvM_{n,5}$	0.228	0.326	0.324	0.128	0.200	0.288	
$CvM_{n,6}$	0.470	0.714	0.894	0.084	0.172	0.250	
$CvM_{n,7}$	0.102	0.120	0.062	0.068	0.100	0.172	
$CvM_{n,8}$	0.310	0.578	0.828	0.120	0.224	0.316	
$CvM_{n,9}$	0.082	0.072	0.082	0.082	0.082	0.096	
$CvM_{n,10}$	0.228	0.442	0.706	0.118	0.156	0.300	

TABLE B.2. Component Test Results,  $DGP_2$  and  $DGP_3$ 

 $PC_k, k = 1, 2, \dots, 10$  are tests using the  $k^{th}$  component.

		$DGP_4$			$DGP_5$	
	n = 100	200	300	n = 100	200	300
$CvM_{n,1}$	0.064	0.064	0.070	0.100	0.030	0.074
$CvM_{n,2}$	0.008	0.014	0.016	0.004	0.008	0.022
$CvM_{n,3}$	0.176	0.306	0.392	0.180	0.316	0.368
$CvM_{n,4}$	0.042	0.058	0.070	0.038	0.066	0.082
$CvM_{n,5}$	0.128	0.200	0.288	0.134	0.194	0.274
$CvM_{n,6}$	0.084	0.172	0.250	0.096	0.138	0.262
$CvM_{n,7}$	0.068	0.100	0.172	0.068	0.120	0.124
$CvM_{n,8}$	0.120	0.224	0.316	0.116	0.202	0.310
$CvM_{n,9}$	0.082	0.082	0.096	0.066	0.062	0.084
$CvM_{n,10}$	0.118	0.156	0.300	0.114	0.176	0.248

TABLE B.3. Component Test Results,  $DGP_6$  and  $DGP_7$ 

 $PC_k, k = 1, 2, \dots, 10$  are tests using the  $k^{th}$  component.

		$DGP_6$		DGP <sub>7</sub>		
	n = 100	200	300	n = 100	200	300
$CvM_{n,1}$	0.870	0.896	0.920	0.866	0.908	0.910
$CvM_{n,2}$	0.742	0.848	0.924	0.732	0.856	0.928
$CvM_{n,3}$	0.022	0.004	0.000	0.008	0.004	0.000
$CvM_{n,4}$	0.172	0.060	0.092	0.152	0.076	0.054
$CvM_{n,5}$	0.020	0.000	0.002	0.014	0.006	0.004
$CvM_{n,6}$	0.118	0.218	0.440	0.132	0.184	0.368
$CvM_{n,7}$	0.010	0.016	0.044	0.018	0.004	0.020
$CvM_{n,8}$	0.060	0.100	0.128	0.064	0.076	0.114
$CvM_{n,9}$	0.006	0.010	0.014	0.008	0.018	0.010
$CvM_{n,10}$	0.050	0.052	0.118	0.072	0.074	0.134

TABLE B.4. Component Test Results,  $DGP_6$  and  $DGP_7$ 

 $PC_k, k = 1, 2, \dots, 10$  are tests using the  $k^{th}$  component.

	$DGP_8$		
	n = 100	200	300
$CvM_{n,1}$	0.884	0.920	0.912
$CvM_{n,2}$	0.784	0.876	0.938
$CvM_{n,3}$	0.024	0.006	0.002
$CvM_{n,4}$	0.150	0.068	0.050
$CvM_{n,5}$	0.010	0.010	0.002
$CvM_{n,6}$	0.112	0.240	0.402
$CvM_{n,7}$	0.010	0.008	0.024
$CvM_{n,8}$	0.072	0.062	0.106
$CvM_{n,9}$	0.010	0.004	0.014
$CvM_{n,10}$	0.052	0.082	0.172

TABLE B.5. Component Test Results,  $DGP_8$ 

 $PC_k, k = 1, 2, ..., 10$  are tests using the  $k^{th}$  component.