# A Graded Schur Lemma and a graded-monoidal structure for induced modules over graded-commutative algebras

Jürgen Fuchs <sup>a</sup> and Tobias Grøsfjeld <sup>b</sup>

<sup>a</sup> Teoretisk fysik, Karlstads Universitet Universitetsgatan 21, S-65188 Karlstad

<sup>b</sup> Matematiska institutionen, Stockholms Universitet S-106 91 Stockholm

#### Abstract

We consider algebras and Frobenius algebras, internal to a monoidal category, that are graded over a finite abelian group. For the case that A is a twisted group algebra in a linear abelian monoidal category we obtain a graded generalization of the Schur Lemma for the category of induced A-modules. We further show that if the monoidal category is braided and A is commutative up to a bicharacter of the grading group, then the category of induced A-modules can be endowed with a graded-monoidal structure that is twisted by the bicharacter. In the particular case that the grading group is  $\mathbb{Z}/2\mathbb{Z}$ , these findings reproduce known results about superalgebras and super-monoidal structures.

## 1 Introduction

Algebras in monoidal categories and their representation theory have become customary tools both in mathematics and in theoretical physics. While for important aspects a Morita invariant approach via module categories over monoidal categories is convenient, there are also issues which are naturally discussed directly in terms of concrete representative algebras rather than their Morita class. Specifically, a significant property an algebra may possess that is not Morita invariant is commutativity.

The category of right modules over a commutative algebra A in a (suitably nice)  $\Bbbk$ -linear braided monoidal category C inherits a natural monoidal structure from the tensor product over A of A-bimodules. In this note we study how this feature generalizes to the case of algebras A that are graded-commutative in the sense that  $A = \bigoplus_{i \in \Gamma} A_i$  is graded over a finite abelian group  $\Gamma$  such that the multiplication on A is grade-preserving and (braided-)commutative up to a normalized two-cocycle on  $\Gamma$  (see Definitions 2.2 and 2.24). We first consider the non-full subcategory  $mod_{\Gamma}^{\circ}$ -A of the category  $mod_{\Gamma}$ -A of  $\Gamma$ -graded right A-modules that is obtained by admitting as morphisms only grade-preserving module morphisms. We show in Proposition 3.4 that if the tensor product  $\otimes_A$  of A-bimodules exists and A is graded-commutative with respect to a cocycle  $\kappa$  that is a bicharacter, then  $\otimes_A$  furnishes a monoidal structure on  $mod_{\Gamma}^{\circ}$ -A.

Our basic goal in this note is then to obtain similar results for categories of A-modules on which no restriction to grade-preserving morphisms is imposed. To achieve this we consider instead of the category  $mod_{\Gamma}$ -A of all right A-modules its full subcategory ind-A of induced Amodules. Our first main result (Proposition 4.5 and Theorem 4.6) then concerns the structure of the morphism spaces in such categories of induced modules, without reference to a braiding on C: For any twisted group algebra algebra A (as defined in Example 2.6) in a k-linear abelian monoidal category C with simple monoidal unit the space of module morphisms between any two simple induced A-modules Ind(x) and Ind(y) is either zero or isomorphic as a  $\Gamma$ -graded  $\Bbbk$ -vector space to the endomorphims of Ind(x) (as well as to those of Ind(y)) which, in turn, is isomorphic as a  $\Sigma_x$ -graded k-algebra to the group algebra k  $\Sigma_x$  twisted by a two-cocycle, where  $\Sigma_x$  is the subgroup  $\{i \in \Gamma \mid x \otimes A_i \cong x\}$  of  $\Gamma$ . Moreover, any non-zero homogeneous module morphism between simple induced A-modules is an isomorphism. This result constitutes a generalization of the Schur Lemma for abelian categories, and accordingly we refer to it as the Graded Schur Lemma. As we point out in Remark 4.9, in the case  $\Gamma = \mathbb{Z}/2\mathbb{Z}$  this comprises several variants of what is known as the Super Schur Lemma, which is e.g. relevant in the study of fermionic topological phases in condensed matter physics.

For a generic graded-commutative algebra A, it is natural to ask whether the tensor product  $\widetilde{\otimes}$  that the category  $ind^{\circ}-A$  of induced A-modules with grade-preserving module morphisms inherits as a full subcategory of  $mod_{\Gamma}^{\circ}-A$  can be extended to all of ind-A. This turns out to be possible only if A is actually commutative (see Lemma 5.4). However, this observation can be upgraded to the graded-commutative situation, thus leading to our second main result (Theorem 5.7) which states: For an algebra A in a braided monoidal category that is graded-commutative with respect to a bicharacter  $\kappa$  of a finite abelian group  $\Gamma$ , the category ind-A can be endowed with the structure of a  $(\Gamma,\kappa)$ -graded-monoidal category. Furthermore, if the tensor product of A-bimodules exists and A is coproduct-separable Frobenius, then restriction to the subcategory ind-A reproduces the tensor product  $\widetilde{\otimes}$ .

Here the relevant notion of a linear  $(\Gamma, \kappa)$ -graded-monoidal category is the following. Such

a category is, by definition, a category enriched over the category  $\operatorname{vect}_{\Gamma}^{(0)}$  of  $\Gamma$ -graded vector spaces with grade-preserving linear maps, in which the morphism spaces are graded and the tensor product of morphisms is twisted by the bicharacter  $\kappa$  [Pu, BrD]. For the details see Definitions and 2.36 and 2.37. The notion of  $(\Gamma, \kappa)$ -graded-monoidal category is thus in particular substantially different from other structures of graded category that appear in the literature, both from the one in which the objects are graded (see e.g. [EGNO, Ch. 9.13] for the case of fusion categories) and which e.g. underlies the notion of a  $\Gamma$ -equivariant category, and from the " $\Gamma$ -graded monoidal structure" as defined in [FrW] and studied in e.g. [CGO]. Instead, it is a generalization from  $\mathbb{Z}/2\mathbb{Z}$  to an arbitrary finite abelian group of the notion of a supermonoidal structure, which arises in the context of monoidal supercategories of the form considered in [BrE].

The additional assumption of A being Frobenius as well as coproduct-separable, in the sense of Definitions (2.13) and (2.14), that is made in the second part of Theorem 5.7 is in particular satisfied for twisted group algebras. Technically, its relevance derives from the fact that the tensor product of A-bimodules, if it exists, can be realized as the image of an idempotent, see Lemma 2.21. Note that coproduct-separable Frobenius algebras in monoidal categories play a crucial role in various different contexts, such as in the study of weak monoidal Morita equivalence of tensor categories [Müg] and of autonomous pseudomonoids in monoidal bicategories [St], and in two-dimensional conformal quantum field theory [FuRS1, He].

The rest of this paper is organized as follows. In Sections 2.1–2.3 we present pertinent background information on graded algebras and graded Frobenius algebras in monoidal categories, on their categories of graded modules and on the tensor product of their bimodules. This includes a few illustrating examples, in particular the twisted group algebras in Examples 2.6 and 2.16, as well as a discussion of analogues for modules (Lemma 2.12) and for Frobenius algebras (Lemma 2.18) of the cocycle-twisting of algebras. We also collect, in Convention 2.1, some conventions and notations that are used throughout the text. Section 2.4 presents the notion of a graded-commutative algebra and a few results about them; e.g. a twisted group algebra is graded-commutative iff the twisting cochain is a bicharacter (Lemma 2.26). Sections 2.5 and 2.6 provide supplementary background on monoidal structures on monads and on the appropriate notion of graded-monoidal category, respectively.

In Section 3 we exhibit the tensor product on the category  $mod_{\Gamma}^{\circ}$ -A for a graded-commutative algebra A, including the description as the image of an idempotent (Proposition 3.4(ii)) in case A is coproduct-separable Frobenius. Section 4 discusses aspects of the category ind-A of induced A-modules for the case that A is a twisted group algebra, culminating in the Graded Schur Lemma, Theorem 4.6.

Section 5 is devoted to establishing Theorem 5.7, i.e. providing the structure of a  $(\Gamma, \kappa)$ -graded-monoidal category, in the sense of Definition 2.37, on the category of induced modules over an algebra A that is graded-commutative with respect to a bicharacter  $\kappa$ . This is achieved by realizing the category ind-A as the Kleisli category over the monad  $\mathbb A$  with underlying endofunctor  $-\otimes A$ , and identifying, in Proposition 5.5, a corresponding graded variant of a monoidal structure on  $\mathbb A$  that implies a  $\kappa$ -twisted interchange law.

### 2 Preliminaries

### 2.1 Graded algebras

Let  $C = (C, \otimes, \mathbf{1}, a, u_l, u_r)$  be an additive monoidal category. Recall that a (unital, associative) algebra in C is a triple  $A = (\dot{A}, \mu, \eta)$  consisting of an object  $\dot{A} \in C$ , a multiplication morphism  $\mu \in \operatorname{Hom}(\dot{A} \otimes \dot{A}, \dot{A})$  and a unit morphism  $\eta \in \operatorname{Hom}(\mathbf{1}, \dot{A})$  satisfying  $\mu \circ (\mu \otimes id_{\dot{A}}) = \mu \circ (id_{\dot{A}} \otimes \mu) \circ a_{\dot{A}, \dot{A}, \dot{A}}$  (associativity) and  $\mu \circ (\eta \otimes id_{\dot{A}}) \circ u_{l, \dot{A}}^{-1} = id_{\dot{A}} = \mu \circ (id_{\dot{A}} \otimes \eta) \circ u_{r, \dot{A}}^{-1}$  (unit properties).

Below we will adhere to the following conventions, unless stated otherwise:

#### Convention 2.1.

- 1. Throughout this paper, k is an algebraically closed field. A generic category will be assumed to be additive and k-linear.
- 2. Whenever no confusion can arise, we suppress the associator a and unitors u<sub>l</sub> and u<sub>r</sub> of a monoidal category.
- 3. When x is an object with additional structure, such as an algebra or module, we denote the underlying plain object by  $\dot{x}$ . However, we often abuse notation by denoting the object underlying an algebra  $A = (\dot{A}, \mu, \eta)$  just by the symbol A, and similarly for modules.
- 4. For any k-linear monoidal category we assume that  $\text{Hom}(\mathbf{1},\mathbf{1}) \cong k$ , and for any algebra we assume that its underlying object is non-zero.
- 5. Throughout this paper,  $\Gamma$  is a finite abelian group and  $\kappa$  is a normalized two-cocycle on  $\Gamma$ . We write the group product of  $\Gamma$  as addition; in particular the unit element is denoted by  $0 \in \Gamma$ . If  $\Gamma$  occurs in a k-linear setting, we assume that the characteristic of k and the order of  $\Gamma$  are coprime.
- 6. In Section 4 A is a twisted group algebra in the sense of Example 2.6, while in Sections 3 and 5 A is graded-commutative in the sense of Definition 2.24 and it is assumed that the tensor product  $\otimes_A$  of A-bimodules exists.

For an object  $c \in C$  with a direct sum decomposition  $c = \bigoplus_i c_i$ , we denote by  $e_i^c$  and  $r_c^i$  the embedding and restriction morphisms for  $c_i$  as a retract of c and set  $e_i^c \circ r_c^i =: p_i^c$ . By definition we have  $r_c^i \circ e_j^c = \delta_{i,j} i d_{c_i}$ , so that the morphisms  $p_i^c$  are mutually orthogonal idempotents,  $p_i^c \circ p_i^c = \delta_{i,j} p_i^c$ . If the direct summands are labeled by the group  $\Gamma$ , i.e.

$$c = \bigoplus_{i \in \Gamma} c_i \,, \tag{2.1}$$

we call c a  $\Gamma$ -graded object. The tensor product of two  $\Gamma$ -graded objects c and d is naturally  $\Gamma$ -graded, with  $(c \otimes d)_i = \bigoplus_{j \in \Gamma} c_j \otimes d_{i-j}$ . A morphism  $f: c \to d$  between  $\Gamma$ -graded objects is grade-preserving iff it satisfies

$$r_d^j \circ f \circ e_i^c = 0 \quad \text{for all } i, j \in \Gamma \text{ with } i \neq j.$$
 (2.2)

Borrowing terminology from the case  $\Gamma = \mathbb{Z}/2\mathbb{Z}$ , we call a grade-preserving morphism synonymously also a *homogeneously even* morphism.

**Definition 2.2.** A  $\Gamma$ -graded algebra in a monoidal category C is an algebra  $A = (\dot{A}, \mu, \eta)$  in C for which the underlying object  $\dot{A} \in C$  is  $\Gamma$ -graded and the multiplication morphism  $\mu$  is homogeneously even.

Thus a  $\Gamma$ -graded algebra has a direct sum decomposition  $A = \bigoplus_{i \in \Gamma} A_i$ . When no confusion can arise, we abbreviate  $e_i^A =: e_i, \ r_A^i =: r^i$  and  $p_i^A =: p_i$ . That  $\mu$  is homogeneously even then means that the morphisms

$$\mu_{i,j}^k := r^k \circ \mu \circ (e_i \otimes e_j) \in \text{Hom}(A_i \otimes A_j, A_k) \tag{2.3}$$

satisfy  $\mu_{i,j}^k = 0$  if  $k \neq i+j$ , for  $i, j, k \in \Gamma$ . Accordingly we will use the abbreviation

$$\mu_{i,j}^{i+j} =: \mu_{i,j}$$
. (2.4)

It follows from the unit axioms that  $e_0 \circ r^0 \circ \eta = \eta$  and thus, with the natural convention that the monoidal unit is  $\Gamma$ -graded with  $\mathbf{1}_j = \delta_{j,0} \mathbf{1}$ , that  $\eta$  is automatically homogeneously even.

**Remark 2.3.** Let  $A = (\dot{A}, \mu, \eta)$  be a  $\Gamma$ -graded algebra in a  $\mathbb{k}$ -linear monoidal category C, and let  $\kappa \colon \Gamma \times \Gamma \to \mathbb{k}^{\times}$  be a normalized two-cocycle on  $\Gamma$ . Then as a direct generalization of a standard result for  $\mathbb{k}$ -algebras (compare e.g. Chapter 2 of [Ka] for the case of group algebras), the triple  $A^{[\kappa]} := (\dot{A}, \mu^{[\kappa]}, \eta)$  with cocycle-twisted multiplication

$$\mu_{i,j}^{[\kappa]} := \kappa(i,j)\,\mu_{i,j} \tag{2.5}$$

defines a structure of an associative algebra in C. To see that  $\mu^{[\kappa]}$  is an associative multiplication on  $\dot{A}$ , use  $\mu = \sum_{i,j\in\Gamma} e_{i+j} \circ \mu_{i,j} \circ (r^i \otimes r^j)$  and the analogous decomposition of  $\mu^{[\kappa]}$  to write

$$\mu^{[\kappa]} \circ (\mu^{[\kappa]} \otimes id_{A}) = \sum_{i,j,k \in \Gamma} \kappa(i,j) \, \kappa(i+j,k) \, e_{i+j+k} \circ [\mu_{i+j,k} \circ (\mu_{i,j} \otimes id_{A_{k}})] \circ (r^{i} \otimes r^{j} \otimes r^{k}) \,,$$

$$\mu^{[\kappa]} \circ (id_{A} \otimes \mu^{[\kappa]}) = \sum_{i,j,k \in \Gamma} \kappa(j,k) \, \kappa(i,j+k) \, e_{i+j+k} \circ [\mu_{i,j+k} \circ (id_{A_{i}} \otimes \mu_{j,k})] \circ (r^{i} \otimes r^{j} \otimes r^{k}) \,.$$

$$(2.6)$$

By associativity of  $\mu$  we have  $\mu_{i+j,k} \circ (\mu_{i,j} \otimes id_{A_k}) = \mu_{i,j+k} \circ (id_{A_i} \otimes \mu_{j,k})$ , while by the closedness  $d\kappa = 1$  of  $\kappa$  the cocycle factors in the two expressions are equal. That  $\eta$  is a unit for  $\mu^{[\kappa]}$  follows from  $\eta = p_0 \circ \eta$  together with the fact that  $\kappa$  is normalized, i.e.  $\kappa(i,0) = 1 = \kappa(0,i)$  for  $i \in \Gamma$ .

**Lemma 2.4.** Let  $A = (\dot{A}, \mu, \eta)$  be a  $\Gamma$ -graded algebra in a  $\mathbb{k}$ -linear monoidal category C and let  $\kappa$  and  $\kappa'$  be normalized cocycles on  $\Gamma$ . If  $\kappa$  and  $\kappa'$  are cohomologous, then the algebras  $A^{[\kappa]}$  and  $A^{[\kappa']}$  obtained as in Remark 2.3 are isomorphic by a grade-preserving morphism. Conversely, provided that the morphism  $\mu_{i,j}$  is non-zero for all  $i, j \in \Gamma$ , if  $A^{[\kappa]}$  and  $A^{[\kappa']}$  are isomorphic as algebras by a grade-preserving morphism, then  $\kappa$  and  $\kappa'$  are cohomologous.

Proof. That  $\kappa$  and  $\kappa'$  are cohomologous means that  $\kappa'(i,j) = \tau(i) \tau(j) \tau^{-1}(i+j) \kappa(i,j)$  for some function  $\tau \colon \Gamma \to \mathbb{k}^\times$  with  $\tau(0) = 1$ . Thus the multiplications on  $A^{[\kappa]}$  and on  $A^{[\kappa']}$  are related by  $\mu_{i,j}^{[\kappa']} = \tau(i) \tau(j) \tau^{-1}(i+j) \mu_{i,j}^{[\kappa]}$ . It follows that the automorphism  $\varphi := \bigoplus_{i \in \Gamma} \tau^{-1}(i) id_{A_i}$  of the object A satisfies  $(\mu^{[\kappa']} \circ (\varphi \otimes \varphi))_{i,j} = (\varphi \circ \mu^{[\kappa]})_{i,j}$  for all  $i,j \in \Gamma$  and is thus a morphism of associative algebras from  $A^{[\kappa]}$  to  $A^{[\kappa']}$ . Because of  $\tau(0) = 1$ ,  $\varphi$  is also compatible with the unit. To see the converse, assume that  $A^{[\kappa]}$  and  $A^{[\kappa']}$  are isomorphic as objects by a grade-preserving morphism  $\varphi$ . Owing to the equality  $\mu_{i,j}^{[\kappa']} = \kappa'(i,j) \kappa(i,j)^{-1} \mu_{i,j}^{[\kappa]}$ , in order that  $\varphi$  is an algebra morphism it must act as a scalar on the image of  $\mu_{i,j}$ . If  $\mu_{i,j} \neq 0$  for all  $i, j \in \Gamma$ , we obtain this way a one-cochain by which  $\kappa'$  and  $\kappa$  are cohomologous.

**Example 2.5.** For any normal subgroup H of a finite group G, the group algebra k[G] is a G/H-graded algebra in the category of k-vector spaces:  $k[G] = \bigoplus_{gH \in G/H} k[gH]$ .

**Example 2.6.** A  $\Gamma$ -graded algebra in the category vect of finite-dimensional  $\mathbb{k}$ -vector spaces can be seen as an algebra in the category  $\operatorname{vect}_{\Gamma}^{(0)}$  of finite-dimensional  $\Gamma$ -graded  $\mathbb{k}$ -vector spaces and grade-preserving  $\mathbb{k}$ -linear maps. The following generalization (compare [EGNO, Example 7.8.3]) is our main, and motivating, class of examples. Consider an object A that is pointed, i.e. a direct sum

$$A_{(\Gamma)} := \bigoplus_{i \in \Gamma} L_i \tag{2.7}$$

of invertible objects  $L_i$ , and with  $\Gamma$  a finite subgroup of the Picard group Pic(C) of C (i.e. of the group of isomorphism classes of invertible objects), and require that  $L_{i+j} \cong L_i \otimes L_j$  and  $L_i \ncong L_j$  for  $i \neq j$ . (Note that, since according to Convention 2.1.4 the monoidal unit **1** is linear-simple, i.e. with endomorphism space being isomorphic to k, every invertible object is linear-simple as well.)

The Picard subcategory  $\mathscr{P}ic(\mathcal{C})$ , i.e. the full monoidal subcategory of  $\mathcal{C}$  consisting of direct sums of invertible objects, is equivalent as an abelian category to the category  $\text{vect}_{\text{Pic}(\mathcal{C})}^{\Psi}$  of  $\text{Pic}(\mathcal{C})$ -graded  $\mathbb{k}$ -vector spaces with an associator that defines a three-cocycle  $\Psi \in Z^3(\text{Pic}(\mathcal{C}), \mathbb{k}^{\times})$ ; monoidally equivalent categories give rise to cohomologous three-cocycles. Assume further that there is a two-cochain  $\omega \in C^2(\Gamma, \mathbb{k}^{\times})$  on  $\Gamma$  that satisfies  $d\omega = \Psi|_{\Gamma}$ , i.e. in components,

$$\omega(i,j)\,\omega(i+j,k) = \omega(j,k)\,\omega(i,j+k)\,\Psi(i,j,k) \qquad \text{for } i,j,k \in \Gamma.$$
 (2.8)

Now select for each pair of elements  $i, j \in \Gamma$  a basis  $\lambda_{i,j}$  of the one-dimensional morphism space  $\text{Hom}(L_i \otimes L_j, L_{i+j})$ . Then setting

$$\mu_{i,j} := \omega(i,j) \,\lambda_{i,j} \tag{2.9}$$

for  $i, j \in \Gamma$  defines a structure of a  $\Gamma$ -graded algebra on the  $\Gamma$ -graded object  $A_{(\Gamma)} \in \mathcal{P}ic(\mathcal{C})$  (associativity precisely amounts to (2.8)). When defining  $\eta$  as the embedding of the subobject  $L_0 \cong \mathbf{1}$  into  $A_{(\Gamma)}$ , this multiplication is also unital. We refer to such a  $\Gamma$ -graded algebra as a twisted group algebra.

For  $\kappa$  any normalized two-cocycle on  $\Gamma$ , owing to  $d(\omega \kappa) = d\omega d\kappa = d\omega$ ,  $A_{(\Gamma)}$  endowed with the  $\kappa$ -twisted multiplication  $\mu^{[\kappa]}$  (2.5) is again a twisted group algebra. In fact, every semisimple indecomposable algebra in  $\text{vect}_{\text{Pic}(\mathcal{C})}^{\Psi}$  is such an algebra [Os, Ex. 2.1]. Replacing  $\mu$  by  $\mu^{[\kappa]}$  amounts to changing the chosen bases  $\lambda_{i,j}$  to  $\kappa(i,j) \lambda_{i,j}$ . Also, if  $\kappa = d\chi$  is a coboundary, then by Lemma 2.4 the twisting by  $\kappa$  can be absorbed into an automorphism  $\sum_{i \in \Gamma} \chi(i) i d_{L_i}$  of  $A_{(\Gamma)}$ , so that  $(A_{(\Gamma)}, \mu)$  and  $(A_{(\Gamma)}, \mu^{[d\chi]})$  are isomorphic as algebras.

Note that this works in fact also for non-abelian groups  $A_{(\Gamma)}$ . Later on (see Example 2.25) we will consider braided categories C, for which Pic(C), and thus  $\Gamma \leq \text{Pic}(C)$ , is abelian.

**Example 2.7.** Consider the special case  $\Gamma = \operatorname{Pic}(C) = \mathbb{Z}/2\mathbb{Z}$  of Example 2.6, and let  $L \in C$  be the (up to isomorphism unique) order-2 invertible object. Then the class of  $\Psi$  in  $H^3(\mathbb{Z}/2\mathbb{Z}, \mathbb{k}^{\times})$  is determined by the value  $F_{1,1}^{(LLL)L} \in \{\pm 1\}$  of the gauge-independent 6*j*-symbol  $F_{1,1}^{(LLL)L}$  of the semisimple category  $\operatorname{\mathscr{P}ic}(C)$ . The object  $A = \mathbf{1} \oplus L$  carries an algebra structure iff  $F_{1,1}^{(LLL)L} = 1$ ,

 $<sup>^{1}</sup>$  We use standard conventions for the 6j-symbols of a semisimple monoidal category, see e.g. Appendix A of [FuG].

and in this case the algebra is of the form described in Example 2.6 with  $\Psi$  and  $\omega$  trivial. As a concrete realization, take the semisimple modular tensor category  $\mathcal{C}(A_1^{(1)},k)$  whose objects correspond to the integrable representations of the untwisted affine Lie algebra  $A_1^{(1)}$  at nonnegative integral level k (compare e.g. [EGNO, Ex. 8.18.5 & 8.18.6]). This category has k+1 non-isomorphic simple objects  $U_{\nu}$ , labeled by  $\nu \in \{0,1,\ldots,k\}$ , has Picard group  $\mathbb{Z}/2\mathbb{Z}$  with  $L = U_k$ , and  $F_{1,1}^{(LLL)L} = (-1)^k$ . Thus the object  $A = \mathbf{1} \oplus L$  is an algebra in  $\mathcal{C}(A_1^{(1)},k)$  iff k is even.

**Example 2.8.** A category C may admit a  $\mathbb{Z}/2\mathbb{Z}$ -graded algebra with underlying object of the form  $A = A_0 \oplus A_1$  such that  $A_1$  is simple while  $A_0 \cong \mathbf{1} \oplus L$  with L an order-2 invertible object and  $L \otimes A_1 \cong A_1 \cong A_1 \otimes L$ . A category for which this happens is, in the notation of Example 2.7,  $C(A_1^{(1)}, 4)$ , with  $L = U_4$  and  $A_1 = U_2$  [FuRS1, Sect. 3.6.2]. In this case one actually has  $A \cong A_1 \otimes A_1$ , and in particular A is Morita equivalent to the trivial algebra  $\mathbf{1} \in C(A_1^{(1)}, 4)$ . We do not know of a category with a  $\mathbb{Z}/2\mathbb{Z}$ -graded algebra of this type that is not Morita equivalent to  $\mathbf{1}$ .

Convention 2.9. If for a  $\Gamma$ -graded algebra  $A = \bigoplus_i A_i$  the morphism spaces  $\operatorname{Hom}(A_i \otimes A_j, A_{i+j})$  are zero- or one-dimensional for all  $i, j \in \Gamma$ , then, given any choice  $\{\lambda_{i,j}\}$  of bases of these spaces, we set  $\mu_{i,j} \equiv \mu_{i,j}^{i+j} =: \overline{\mu}_{i,j} \lambda_{i,j}$  with numbers  $\overline{\mu}_{i,j} \in \mathbb{k}$  and call the matrix  $\boldsymbol{\mu} := (\overline{\mu}_{i,j})_{i,j \in \Gamma}$  the matrix representation of  $\mu$ . By abuse of notation we use the same symbols and terminology whenever  $\mu_{i,j} = 0$ , independently of the dimensionality of  $\operatorname{Hom}(A_i \otimes A_j, A_{i+j})$ .

**Example 2.10.** Let  $x \in C$  be arbitrary. Then the object  $X := \mathbf{1} \oplus x$  endowed with the multiplication  $\mu$  with matrix representation

$$\boldsymbol{\mu} = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \tag{2.10}$$

is a unital associative algebra, and it is  $\mathbb{Z}/2\mathbb{Z}$ -graded with  $X_0 = \mathbf{1}$  and  $X_1 = x$ . This algebra X can be seen as an analogue of the quotient ring  $\mathbb{k}[t]/\langle t^2 \rangle$ ; likewise there are analogues of  $\mathbb{k}[t]/\langle t^\ell \rangle$  for any integer  $\ell > 2$ .

Recall that a right module m over an algebra A in C consists of an object  $\dot{m} \in C$  and a representation morphism  $q: \dot{m} \otimes A \to \dot{m}$  satisfying  $q \circ (q \otimes id_A) = q \circ (id_{\dot{m}} \otimes \mu)$  and  $q \circ (id_{\dot{m}} \otimes \eta) = id_{\dot{m}}$ . Analogously, for a left A-module  $n = (\dot{n}, \rho)$  the representation morphism  $\rho: A \otimes \dot{n} \to \dot{n}$  obeys  $\rho \circ (id_A \otimes \rho) = \rho \circ (\mu \otimes id_{\dot{n}})$  and  $\rho \circ (\eta \otimes id_{\dot{n}}) = id_{\dot{n}}$ . A bimodule is both a left and a right module, with commuting left and right actions. A morphism  $f: m \to m'$  of right A-modules is a morphism of the underlying objects that commutes with the A-action, i.e.  $f: \dot{m} \to \dot{m}'$  such that  $q_n \circ (f \otimes id_A) = f \circ q_m$ ; we denote the set of such module morphisms by  $Hom_A(m, n)$ . We will often refer to a right A-module just as an A-module.

**Definition 2.11.** Let A be a  $\Gamma$ -graded algebra in a monoidal category C. A  $\Gamma$ -graded right module over A is a right A-module  $(\dot{m}, q)$  for which the object  $\dot{m} \in C$  is  $\Gamma$ -graded (i.e. has a direct sum decomposition as in (2.1)) and the action morphism q is homogeneously even. The category  $mod_{\Gamma}^{\circ}$ -A of  $\Gamma$ -graded right A-modules is the category that has  $\Gamma$ -graded right A-modules as objects and homogeneously even A-module morphisms as morphisms.

More explicitly, a  $\Gamma$ -graded right module  $(\dot{m}, \mathbf{q})$  has a direct sum decomposition  $\dot{m} = \bigoplus_{i \in \Gamma} m_i$  such that the morphisms  $\mathbf{q}_{i,j}^k := r_m^k \circ \mathbf{q} \circ (e_i^m \otimes e_j) \in \mathrm{Hom}(m_i \otimes A_j, m_k)$  satisfy  $\mathbf{q}_{i,j}^k = 0$  if  $k \neq i+j$ , for  $i, j, k \in \Gamma$ . And a homogeneously even A-module morphism is a module morphism f between  $\Gamma$ -graded right A-modules  $(\dot{m}, \mathbf{q})$  and  $(\dot{m}', \mathbf{q}')$  such that  $r_{m'}^j \circ f \circ e_i^m = 0$  if  $j \neq i$ , for  $i, j \in \Gamma$ . Similarly as done for graded algebras in (2.4), for a  $\Gamma$ -graded right module  $(c, \mathbf{q})$  we abbreviate  $\mathbf{q}_{i,j}^{i+j} =: \mathbf{q}_{i,j}$ .

The notion of cocycle-twisting of algebras in Remark 2.3 has a counterpart for modules. The two constructions are related as follows:

**Lemma 2.12.** Let  $m = (\dot{m}, \mathbf{q})$  be a  $\Gamma$ -graded right module over a  $\Gamma$ -graded algebra A in a  $\mathbb{k}$ -linear monoidal category C, and let  $\kappa$  be a normalized two-cocycle on  $\Gamma$ . The  $\kappa$ -twisted action

$$q^{[\kappa]} := \sum_{i,j \in \Gamma} e^{\dot{m}}_{i+j} \circ q^{[\kappa]}_{i,j} \circ (r^{\dot{i}}_{\dot{m}} \otimes r^{\dot{j}}) \qquad \text{with} \qquad q^{[\kappa]}_{i,j} := \kappa(i,j) \, q_{i,j} \tag{2.11}$$

endows the object  $\dot{m} \in C$  with the structure of a right  $A^{[\kappa]}$ -module

$$m^{[\kappa]} = (\dot{m}, \mathbf{q}^{[\kappa]}). \tag{2.12}$$

Analogously, for  $n = (\dot{n}, \rho)$  a  $\Gamma$ -graded left A-module, the object  $\dot{n}$  comes with the structure of a left  $A^{[\kappa]}$ -module  $n^{[\kappa]} = (\dot{n}, \rho^{[\kappa]})$  with  $\kappa$ -twisted left action  $\rho^{[\kappa]} := \sum_{i,j \in \Gamma} e^{\dot{n}}_{i+j} \circ \rho^{[\kappa]}_{i,j} \circ (r^i \otimes r^j_{\dot{n}})$  with  $\rho^{[\kappa]}_{i,j} := \kappa(i,j) \, \rho_{i,j}$ .

*Proof.* The sequence

$$q^{[\kappa]} \circ (q^{[\kappa]} \otimes id_{A}) = \sum_{i,j,k \in \Gamma} \kappa(i+j,k) \, \kappa(i,j) \, e^{\dot{m}}_{i+j+k} \circ q_{i+j,k} \circ (q_{i,j} \otimes id_{A_{k}}) \circ (r^{\dot{i}}_{\dot{m}} \otimes r^{\dot{j}} \otimes r^{\dot{k}})$$

$$= \sum_{i,j,k \in \Gamma} \kappa(i,j+k) \, \kappa(j,k) \, e^{\dot{m}}_{i+j+k} \circ q_{i,j+k} \circ (id_{\dot{m}_{i}} \otimes \mu_{j,k}) \circ (r^{\dot{i}}_{\dot{m}} \otimes r^{\dot{j}} \otimes r^{\dot{k}})$$

$$= \sum_{i,j,k \in \Gamma} e^{\dot{m}}_{i+j+k} \circ q^{[\kappa]}_{i,j+k} \circ (id_{m_{i}} \otimes \mu^{[\kappa]}_{j,k}) \circ (r^{\dot{i}}_{\dot{m}} \otimes r^{\dot{j}} \otimes r^{\dot{k}})$$

$$= q^{[\kappa]} \circ (id_{\dot{m}} \otimes \mu^{[\kappa]})$$
(2.13)

of equalities shows that  $q^{[\kappa]}$  is compatible with the twisted multiplication  $\mu^{[\kappa]}$ . Here the third equality uses the cocycle condition for  $\kappa$  as well as the compatibility of q with  $\mu$ . The compatibility of  $q^{[\kappa]}$  with the unit is seen similarly by using that the cocycle  $\kappa$  is normalized. The calculation for the left module case is analogous.

### 2.2 Frobenius algebras and separability

A (counital, coassociative) coalgebra in C is a triple  $B = (B, \Delta, \varepsilon)$  consisting of an object  $B \in C$  and morphisms  $\Delta \in \text{Hom}(B, B \otimes B)$  and  $\varepsilon \in \text{Hom}(B, \mathbf{1})$  satisfying  $(\Delta \otimes id_A) \circ \Delta = (id_A \otimes \Delta) \circ \Delta$  and  $(\varepsilon \otimes id_A) \circ \Delta = id_A = (id_A \otimes \varepsilon) \circ \Delta$ .

**Definition 2.13.** A Frobenius algebra in C is a quintuple  $F = (F, \mu, \eta, \Delta, \varepsilon)$  such that  $(F, \mu, \eta)$  is an algebra,  $(F, \Delta, \varepsilon)$  is a coalgebra, and

$$(\mu \otimes id_A) \circ (id_A \otimes \Delta) = \Delta \circ \mu = (id_A \otimes \mu) \circ (\Delta \otimes id_A), \qquad (2.14)$$

i.e. the coproduct is a morphism of F-bimodules.

**Definition 2.14.** A separable algebra in C is an algebra  $(A, \mu, \eta)$  such that the multiplication  $\mu$  splits as a morphism of A-bimodules, i.e. there is a bimodule morphism  $n \in Hom(A, A \otimes A)$  such that  $\mu \circ n = id_A$ . A coproduct-separable algebra, also called  $\Delta$ -separable algebra, in C is an algebra-coalgebra  $(A, \mu, \eta, \Delta, \varepsilon)$  which as an algebra is separable with  $n = \Delta$ .

A Frobenius algebra  $(A, \mu, \eta, \Delta, \varepsilon)$  is separable iff

$$\mu \circ (id_A \otimes \zeta) \circ \Delta = id_A \tag{2.15}$$

for some morphism  $\zeta \in \text{Hom}(\mathbf{1}, A)$  [Mul, Prop. 3.1]. In particular, for a connected Frobenius algebra, i.e. one satisfying  $\text{Hom}(\mathbf{1}, A) = \mathbb{k} \eta$ , separability and coproduct-separability are equivalent.

Dually to Definition 2.2, a  $\Gamma$ -graded coalgebra in C is a coalgebra  $A=(\dot{A},\Delta,\varepsilon)$  whose underlying object is  $\Gamma$ -graded and for which the comultiplication  $\Delta$  is homogeneously even, i.e. the morphisms  $\Delta_k^{i,j}:=(r^i\otimes r^j)\circ\Delta\circ e_k$  in  $\operatorname{Hom}(A_k,A_i\otimes A_j)$  obey  $\Delta_k^{i,j}=0$  for  $k\neq i+j$ . As in the case of the unit morphism of an algebra, with the natural convention that  $\mathbf{1}$  is  $\Gamma$ -graded with  $\mathbf{1}_j=\delta_{j,0}\,\mathbf{1}$ , the counit morphism  $\varepsilon$  is automatically homogeneously even. For a  $\Gamma$ -graded coalgebra we abbreviate  $\Delta_{i+j}^{i,j}:=\Delta^{i,j}$ . A  $\Gamma$ -graded Frobenius algebra in C is a Frobenius algebra  $A=(\dot{A},\mu,\eta,\Delta,\varepsilon)$  which is  $\Gamma$ -graded both as an algebra and as a coalgebra.

Remark 2.15. Graded Frobenius algebras in the monoidal category comod-H of comodules over a Hopf k-algebra H have been considered in [DNN]. If H is the group algebra kG of a group G, an algebra in comod-H is just a G-graded k-algebra. As shown in Corollary 4.2 of [DNN], a (strongly) graded finite-dimensional k-algebra A is graded Frobenius if and only if its neutral component  $A_0$  is Frobenius. For algebras A in a generic k-linear monoidal category, being graded Frobenius is still sufficient for  $A_0$  being Frobenius, but the opposite implication no longer holds.

**Example 2.16.** For  $\Gamma \leq \text{Pic}(C)$ , the twisted group algebra  $A_{(\Gamma)}$  described in Example 2.6 can be endowed with the structure of a  $\Gamma$ -graded coalgebra by setting [FuRS2, Lemma 3.10(iii)]

$$\Delta^{i,j} := |\Gamma|^{-1} \omega(i,j)^{-1} \, \mathsf{Y}^{i,j},\tag{2.16}$$

where we write  $Y^{i,j}$  for the basis of the one-dimensional vector space  $\operatorname{Hom}(L_{i+j}, L_i \otimes L_j)$  that is covector dual to the basis  $\lambda_{i,j}$  of  $\operatorname{Hom}(L_i \otimes L_j, L_{i+j})$  used in the definition of the multiplication of  $A_{(\Gamma)}$ . This makes  $A_{(\Gamma)}$  into a coproduct-separable  $\Gamma$ -graded Frobenius algebra.

**Example 2.17.** We define the matrix representation  $\Delta = (\overline{\Delta}_{i,j})_{i,j\in\Gamma}$  of the comultiplication of a  $\Gamma$ -graded coalgebra with zero- or one-dimensional morphism spaces  $\operatorname{Hom}(A_{i+j}, A_i \otimes A_j)$  analogously as the matrix representation of  $\mu$  in Convention 2.9. Similarly as in Example 2.10, if  $X = \mathbf{1} \oplus x$  is endowed with the comultiplication  $\Delta$  with matrix representation

$$\Delta = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}, \tag{2.17}$$

then it is a  $\mathbb{Z}/2\mathbb{Z}$ -graded counital coalgebra. However, this structure does not fit together with the multiplication (2.10) into a Frobenius structure. On the other hand, provided that x is simple one does obtain a Frobenius algebra with multiplication (2.10) by endowing X with a Frobenius form that has the same matrix representation as  $\mu$ . However, this does not yield a  $\mathbb{Z}/2\mathbb{Z}$ -graded coalgebra structure on X.

**Lemma 2.18.** Let C be a monoidal category and  $A = (\dot{A}, \mu, \eta, \Delta, \varepsilon)$  be a  $\Gamma$ -graded Frobenius algebra in C. Then  $A^{[\kappa]} := (\dot{A}, \mu^{[\kappa]}, \eta, \Delta^{[\kappa]}, \varepsilon)$  with cocycle-twisted multiplication (2.5) and cocycle-twisted comultiplication  $\Delta^{(\kappa)}{}^{i,j} := \kappa(i,j)^{-1}\Delta^{i,j}$ , and with the same unit and counit as A, is again a Frobenius algebra in C. Further, if A is separable, then so is  $A^{[\kappa]}$ .

*Proof.* That  $A^{[\kappa]}$  is an algebra is explained in Remark 2.3; that it is a coalgebra follows in the same way. Concerning the Frobenius property, by direct calculation we have (taking again for simplicity C to be strict)

$$(id_{A} \otimes \mu^{[\kappa]}) \circ (\Delta^{[\kappa]} \otimes id_{A}) = \sum_{i,j,k \in \Gamma} \kappa(i+k,-k)^{-1} \kappa(-k,j) \left( e_{i+k} \otimes e_{j-k} \right)$$

$$\circ \left( id_{A_{i+k}} \otimes \mu_{-k,j} \right) \circ (\Delta^{i+k,-k} \otimes id_{A_{j}}) \circ (r^{i} \otimes r^{j})$$

$$(2.18)$$

and

$$\Delta^{[\kappa]} \circ \mu^{[\kappa]} = \sum_{i,j,k \in \Gamma} \kappa(i+k,j-k)^{-1} \kappa(i,j) \left( e_{i+k} \otimes e_{j-k} \right) \circ \Delta^{i+k,j-k} \circ \mu_{i,j} \circ (r^i \otimes r^j). \tag{2.19}$$

Since  $\kappa$  is a two-cocycle, the products of  $\kappa$ -factors in these two expressions are equal, and hence equality of (2.18) and (2.19) is implied by the corresponding equality for the untwisted operations, i.e. by the Frobenius property of A. The second Frobenius identity follows analogously. The statement about separability follows directly from the fact that  $\mu_{i,j}^{[\kappa]} \circ \Delta^{(\kappa)}{}^{i,j} = \mu_{i,j} \circ \Delta^{i,j}$  together with the characterization (2.15) of separability.

Remark 2.19. Like for k-algebras, the property of admitting a Frobenius structure is not Morita invariant. Rather, the Morita class of a Frobenius algebra in a monoidal category may contain quasi-Frobenius algebras that are not Frobenius [Sh]. In contrast, admitting a *symmetric* Frobenius structure is a Morita invariant property. The notion of symmetric Frobenius algebra, which is defined in any pivotal monoidal category, is important in applications like those in [Müg, FuRS1, He]. This notion does not play a role for the considerations in the present paper, but it is worth noting that the full subcategory generated by the homogeneous components of a twisted group algebra, being equivalent to  $\text{vect}_{\Gamma}^{(0)}$ , admits a pivotal structure.

# 2.3 Tensor product of bimodules

The standard tensor product of bimodules over an algebra in the category of k-vector spaces readily generalizes to other k-linear categories:

**Definition 2.20.** Let A be an algebra in a strict monoidal category C, and let  $m = (\dot{m}, q)$  and  $n = (\dot{n}, \rho)$  be a right and a left A-module in C, respectively. If the coequalizer

$$m \otimes_A n := \operatorname{coequ}(\dot{m} \otimes A \otimes \dot{n} \xrightarrow[id_{\dot{m}} \otimes \rho]{\operatorname{q} \otimes id_{\dot{n}}} \dot{m} \otimes \dot{n}),$$
 (2.20)

exists, then it is called the tensor product over A of m and n.

Note that  $id_{\dot{m}} \otimes \eta \otimes id_{\dot{n}}$  is a common section of  $q \otimes id_{\dot{n}}$  and  $id_{\dot{m}} \otimes \rho$ , i.e. the coequalizer (2.20) is reflexive. Also, here and analogously for the statements below, strictness of C is assumed

only for notational convenience; the only modification in the non-strict case is that one of the two morphisms that are coequalized gets composed with an associator.

Assume now that C admits all coequalizers of the form (2.20) and that the tensor product functor  $\otimes$  of C preserves such coequalizers. Then for any two A-bimodules m and n the object  $m \otimes_A n$  admits a natural structure of A-bimodule as well. Hence the tensor product (2.20) endows the category A- $\beta imod$  of A-bimodules with the structure of a monoidal category, with A as monoidal unit and with associator and unitors inherited from those of C; compare [GJ, Sect. 4.1] for a detailed exposition in a bicategorical context, and e.g. [EGNO, Exc. 7.8.22,7.8.28] for the case that C is a multitensor category.

If the algebra A is separable Frobenius, then the tensor product  $\otimes_A$  can be realized as an image. Denote by

$$q_A: \quad \dot{m} \otimes \dot{n} \longrightarrow m \otimes_A n$$
 (2.21)

the epimorphism given by the definition of the coequalizer (2.20). Then, as a moderate generalization of Lemma 1.21 and Corollary 1.22 of [KO], we have

**Lemma 2.21.** Let A be an algebra in a strict monoidal category C, and let  $m = (\dot{m}, \gamma)$  be a right A-module and  $n = (\dot{n}, \rho)$  a left A-module. If  $\pi \equiv \pi_{m,n} \in \operatorname{End}(\dot{m} \otimes \dot{n})$  is an idempotent and obeys  $q_A \circ \pi = q_A$  as well as

$$\pi \circ (Q \otimes id_{\dot{n}}) = \pi \circ (id_{\dot{m}} \otimes \rho), \qquad (2.22)$$

then we have

$$m \otimes_A n \cong \operatorname{im}(\pi)$$
, (2.23)

and there is a direct sum decomposition  $\dot{m} \otimes \dot{n} \cong m \otimes_A n \oplus \ker(\pi)$ . Moreover, if in addition A is separable Frobenius with separability morphism n, then the endomorphism

$$p \equiv p_{m,n} := (\mathfrak{d} \otimes \rho) \circ (id_{\dot{m}} \otimes (\mathfrak{d} \circ \eta) \otimes id_{\dot{n}}) \tag{2.24}$$

satisfies these three properties.

Proof. By the universal property of  $m \otimes_A n$  as a coequalizer, validity of (2.22) implies that the kernel  $\ker(q_A)$  is a subobject of  $\ker(\pi)$ , while  $q_A \circ p = q_A$  implies that  $\ker(\pi)$  is a subobject of  $\ker(q_A)$ . Thus  $\ker(\pi) = \ker(q_A)$  and hence, using that  $q_A$  is epi,  $\operatorname{im}(\pi) \cong \operatorname{im}(q_A) = m \otimes_A n$ . Further, using that A is Frobenius it follows that the morphism p given by (2.24) satisfies (2.22), and using that A is separable Frobenius it follows that p is an idempotent. Finally, by the definition of  $m \otimes_A n$  as a coequalizer we have

$$q_A \circ p = q_A \circ \left[ id_{\dot{m}} \otimes \left( \rho \circ (id_A \otimes \rho) \circ ((n \circ \eta) \otimes id_{\dot{n}}) \right) \right]$$
 (2.25)

which when combined with the separability and the representation property of  $\rho$  simplifies to  $q_A \circ p = q_A$ .

Thus in short, if A is separable Frobenius, then we have  $m \otimes_A n \cong \operatorname{im}(p)$  with  $p \equiv p_{m,n}$  as given in (2.24). Moreover, the tensor product of morphisms is then given by

$$f \otimes_A g = p_{m',n'} \circ (f \otimes g) = (f \otimes g) \circ p_{m,n}$$
 (2.26)


for  $f: m \to m'$  and  $g: n \to n'$  morphisms of right and left A-modules, respectively.

### 2.4 Graded-commutative algebras

Let now C in addition be braided, with braiding  $\beta$ . Given an algebra  $A = (\dot{A}, \mu, \eta)$  in C, the opposite algebra  $A^{\text{op}}$ , defined as  $A^{\text{op}} := (\dot{A}, \mu \circ \beta_{\dot{A}, \dot{A}}, \eta)$ , is again a unital associative algebra in C

**Lemma 2.22.** Let A be an algebra in a braided monoidal category. The categories A-mod of left A-modules and mod-A<sup>op</sup> of right A<sup>op</sup>-modules are isomorphic. Specifically, if  $n = (\dot{n}, \rho)$  is a left A-module, then  $(\dot{n}, \rho \circ \beta_{\dot{n}, A})$  is a right A<sup>op</sup>-module, and if  $(\dot{m}, q)$  is a right A<sup>op</sup>-module, then  $(\dot{m}, q \circ \beta_{A,\dot{m}})$  is a left A-module.

*Proof.* That, given a left A-module  $(\dot{n}, \rho)$ , the morphism  $\rho \circ \beta_{\dot{n},A}$  defines a right  $A^{\text{op}}$ -action on  $\dot{m}$  follows directly with the help of the naturality of the braiding. In terms of the standard graphical string calculus for monoidal categories, the compatibility with the multiplication on  $A^{\text{op}}$  is seen as follows:



(Our conventions for the string calculus can be easily read off from these pictures.) Moreover, module morphisms with respect to  $\rho \circ \beta_{\dot{m},A}$ , as follows again from naturality of the braiding. One therefore deals with a functor from A-mod to mod- $A^{op}$ . By invertibility of the braiding, this functor is an isomorphism.

**Remark 2.23.** By iteration, for any  $\ell \in \mathbb{Z}$ ,  $A^{(\ell)} := (\dot{A}, \mu \circ (\beta_{\dot{A}, \dot{A}})^{\ell}, \eta)$  is an algebra structure on the object  $\dot{A}$ . Via appropriate even and odd powers, respectively, of the braiding, all categories  $A^{(2r)}$ -mod and mod- $A^{(2r+1)}$  for  $r \in \mathbb{Z}$  are isomorphic. Also, if  $A = (\dot{A}, \mu, \eta, \Delta, \varepsilon)$  is a Frobenius algebra, then

$$A^{(\ell)} := (\dot{A}, \mu \circ (\beta_{\dot{A}, \dot{A}})^{\ell}, \eta, (\beta_{\dot{A}, \dot{A}})^{-\ell} \circ \Delta, \varepsilon)$$
(2.28)

is again a Frobenius algebra structure on the object A. Moreover, if A is separable, then so is  $A^{(\ell)}$ , and if if A is coproduct-separable, then so is  $A^{(\ell)}$ .

A commutative algebra in a braided monoidal category C is an algebra A that is isomorphic to its opposite algebra  $A^{\text{op}}$  (and thus to all algebras  $A^{(n)}$ ). In terms of the components (2.3) of the multiplication, this means  $\mu_{j,i} \circ \beta_{A_i,A_j} = \kappa(i,j) \mu_{i,j}$ . For graded algebras we give analogously

**Definition 2.24.** Let C be a braided monoidal category. A  $\Gamma$ -graded algebra  $A = (A, \mu, \eta)$  in C is called graded-commutative iff

$$\mu_{j,i} \circ \beta_{A_i,A_j} = \kappa(i,j) \,\mu_{i,j} \tag{2.29}$$

for  $i, j \in \Gamma$  for a normalized two-cocycle  $\kappa$  on  $\Gamma$ .

A commutative algebra is thus the same as a graded-commutative algebra for  $\kappa$  the trivial two-cocycle. Note that in our terminology 'graded-commutative' we neither specify the relevant two-cocycle nor the relevant braiding explicitly; this is legitimate because in the situations we are interested in, we deal with a given fixed cocycle  $\kappa$  and a given fixed braiding  $\beta$ .

**Example 2.25.** Consider again the twisted group algebras from Example 2.6, with underlying objects  $A_{(\Gamma)} = \bigoplus_{i \in \Gamma} L_i$ , and let now  $\mathcal{C}$  be braided (so that  $\operatorname{Pic}(\mathcal{C})$ , and hence  $\Gamma$ , is necessarily abelian). The full subcategory  $C_{(\Gamma)}$  of C whose objects are direct sums of the  $L_i$  with  $i \in \Gamma$ is equivalent to the semisimple abelian category  $\text{vect}_{\Gamma}^{(0)}$  of finite-dimensional  $\Gamma$ -graded  $\mathbb{k}$ -vector spaces with homogeneously even linear maps. The inequivalent associators and braidings of a categorical group, and thus also those for the Picard category  $\mathcal{P}ic(\mathcal{C})$  and its full monoidal subcategory  $C_{(\Gamma)}$ , are classified by the third abelian cohomology [JoS, Sect. 3]. (An abelian three-cocycle  $(\Psi,\Omega) \in Z^3_{ab}(G,\mathbb{k}^{\times})$  on a group G consists of an ordinary group three-cocycle  $\Psi \in Z^3(G, \mathbb{R}^{\times})$  and a two-cochain  $\Omega \in C^2(G, \mathbb{R}^{\times})$  that satisfies compatibility relations with  $\Psi$ , see [JoS].) An abelian three-cocycle  $(\Psi, \Omega) \in Z^3_{ab}(\operatorname{Pic}(\mathcal{C}), \mathbb{k}^{\times})$  encodes precisely the restriction to  $\mathcal{P}ic(\mathcal{C})$  of the associator and braiding of  $\mathcal{C}$ . In terms of bases  $\lambda_{i,j}$  of  $\operatorname{Hom}(L_i \otimes L_j, L_{i+j})$  (cf. Example 2.6), the braiding on  $\mathcal{P}ic(\mathcal{C})$  is given by  $\lambda_{j,i} \circ \beta_{L_i,L_j} = \Omega(j,i)^{-1}\lambda_{i,j}$ , while the associator sends  $\lambda_{i+j,k} \circ (\lambda_{i,j} \otimes id_{L_k})$  to  $\Psi(i,j,k)^{-1}(\lambda_{i,j+k} \circ (id_{L_i} \otimes \lambda_{j,k}))$  [FuRS2, App. C.2]. Two abelian three-cocycles yield equivalent braided structures iff they are cohomologous. Further [FuRS2, Prop. 3.14(i), every simple special Frobenius algebra in  $(C_{(\Gamma)}; \psi, \omega)$ , with  $(\psi, \omega) = (\Psi, \Omega)|_{\Gamma}$ , is isomorphic to one of the twisted group algebras  $A_{(\Gamma)}$  with multiplication (2.9).

**Lemma 2.26.** The twisted group algebra  $A_{(\Gamma)}$  in  $(C_{(\Gamma)}; \psi, \omega)$  is graded-commutative iff the cochain  $\omega$  is a bicharacter on  $\Gamma$ .

Proof. Combining the formula (2.9) for the multiplication with the defining relation (2.29) for a graded-commutative algebra and with the expression of the braiding in terms of  $\omega$ , one finds that  $\omega = \kappa^{-1}$ . In particular, the two-cochain  $\omega$  must in fact be a two-cocycle, and as a consequence  $\psi = d\kappa = 1$ . Further, by definition of  $Z_{ab}^3$  one has  $(1, \omega) \in Z_{ab}^3(\Gamma, \mathbb{k}^{\times})$  iff  $\omega$  is a bicharacter, i.e. satisfies  $\omega(i+j,k) = \omega(i,k) \, \omega(j,k)$  and  $\omega(i,j+k) = \omega(i,j) \, \omega(i,k)$  for  $i,j,k \in \Gamma$ .

Example 2.27. Consider the  $\mathbb{Z}/2\mathbb{Z}$ -graded Frobenius algebra  $A=\mathbf{1}\oplus L$  with L an order-2 invertible object satisfying  $F_L:=F_{\mathbf{1},\mathbf{1}}^{(LL\,L)\,L}=1$ , as described in Example 2.7, in a monoidal category C with Picard group  $\mathbb{Z}/2\mathbb{Z}$ . If C is braided, then the category  $\mathscr{P}ic(C)$  is braided as well. The elements of the abelian cohomology group  $H^3_{\mathrm{ab}}(\mathbb{Z}/2\mathbb{Z}, \mathbb{k}^\times)$  can be labeled by the possible values of  $F_L$  and of the multiple of  $id_{L\otimes L}$  that constitutes the self-braiding  $\beta_{L,L}$ . Compatibility of associator and braiding requires that the latter number squares to  $F_L$ , so that in the situation at hand we have  $\beta_{L,L}=\pm id_{L\otimes L}$ . For  $\beta_{L,L}=id_{L\otimes L}$ ,  $\mathscr{P}ic(C)$  is braided equivalent to the category of finite-dimensional  $\mathbb{Z}/2\mathbb{Z}$ -graded vector spaces and  $A=\mathbf{1}\oplus L$  is a commutative algebra in C, while for  $\beta_{L,L}=-id_{L\otimes L}$ ,  $\mathscr{P}ic(C)$  is braided equivalent to the category of finite-dimensional supervector spaces and A is a supercommutative algebra in C. In the particular case of  $C=C(A_1^{(1)},k)$ , as defined in Example 2.7, one finds  $\beta_{L,L}=\mathrm{i}^k\,id_{L\otimes L}$  [EGNO, Exc. 8.18.6], hence for  $k\in 4\mathbb{Z}$  the object  $\mathbf{1}\oplus U_k$  admits a natural structure of a commutative Frobenius algebra, while for  $k\in 4\mathbb{Z}+2$  it admits a natural structure of a supercommutative Frobenius algebra.

**Example 2.28.** The  $\mathbb{Z}/2\mathbb{Z}$ -graded algebra  $X = \mathbf{1} \oplus x$  from Example 2.10 is graded-commutative with respect to any braiding on C.

### 2.5 Monoidal monads

Pertinent aspects of the representation theory of an algebra A in a monoidal category C can conveniently be studied in terms of the corresponding monad on C, whose underlying endofunctor is  $A \otimes -$ . We will use this approach in Section 5.

Recall that a monad on a category C is an algebra  $\mathbb{T} = (T, \mu^{\mathbb{T}}, \eta^{\mathbb{T}})$  in the monoidal category of endofunctors of C. Thus  $\mu^{\mathbb{T}}$  and  $\eta^{\mathbb{T}}$  are natural transformations  $\mu^{\mathbb{T}} \colon T^2 \Longrightarrow T$  and  $\eta^{\mathbb{T}} \colon Id_C \Longrightarrow T$  satisfying

$$\mu^{\mathbb{T}} \circ \mu^{\mathbb{T}} T = \mu^{\mathbb{T}} \circ T \mu^{\mathbb{T}} \quad \text{and} \quad \mu^{\mathbb{T}} \circ \eta^{\mathbb{T}} T = Id_{\mathcal{C}} = \mu^{\mathbb{T}} \circ T \eta^{\mathbb{T}},$$
 (2.30)

or, in components,  $\mu_x^{\mathbb{T}} \circ \mu_{Tx}^{\mathbb{T}} = \mu_x^{\mathbb{T}} \circ T \mu_x^{\mathbb{T}}$  and  $\mu_x^{\mathbb{T}} \circ \eta_{Tx}^{\mathbb{T}} = id_{Tx} = \mu_x^{\mathbb{T}} \circ T \eta_x^{\mathbb{T}}$  for  $x \in \mathcal{C}$ . Naturality of  $\mu^{\mathbb{T}}$  and  $\eta^{\mathbb{T}}$  means that  $\mu_{Ty}^{\mathbb{T}} \circ TTf = Tf \circ \mu_x^{\mathbb{T}}$  and that  $Tf \circ \eta_x^{\mathbb{T}} = \eta_y^{\mathbb{T}} \circ f$  for  $f : x \to y$ , respectively.

**Definition 2.29.** A module over a monad  $\mathbb{T}$  in  $\mathcal{C}$  is a pair  $(x, \rho_x)$  consisting of an object  $x \in \mathcal{C}$  and a morphism  $\rho_x \colon Tx \to x$  satisfying  $\rho_x \circ \mu_x^{\mathbb{T}} = \rho_x \circ T\rho_x$  and  $\rho_x \circ \eta_x^{\mathbb{T}} = id_x$ . A morphism of  $\mathbb{T}$ -modules from  $(x, \rho_x)$  to  $(y, \rho_y)$  is a morphism  $f \colon x \to y$  in  $\mathcal{C}$  satisfying  $f \circ \rho_x = \rho_y \circ Tf$ .

In part of the literature a  $\mathbb{T}$ -module is instead called a  $\mathbb{T}$ -algebra. In the present context this terminology would be inconvenient.

**Definition 2.30.** A monoidal structure on a monad  $\mathbb{T} = (T, \mu^{\mathbb{T}}, \eta^{\mathbb{T}})$  on a monoidal category C is a family  $t_{(2)} \equiv t_{(2)}^{\mathbb{T}}$  of morphisms

$$t_{(2)}^{x,y}: Tx \otimes Ty \longrightarrow T(x \otimes y)$$
 (2.31)

that are natural in  $x, y \in \mathcal{C}$  and make  $(T, t_{(2)}, \eta_1^{\mathbb{T}})$  into a lax monoidal functor and  $\mu^{\mathbb{T}}$  and  $\eta^{\mathbb{T}}$  into monoidal natural transformations.

Put differently, a monoidal monad is a monad in the 2-category of monoidal categories, lax monoidal functors and monoidal natural transformations. Explicitly, the conditions that the natural transformations  $\mu^{\mathbb{T}}$  and  $\eta^{\mathbb{T}}$  are monoidal are given by the commutativity of the diagrams

$$T^{2}x \otimes T^{2}y \xrightarrow{t_{(2)}^{Tx,Ty}} T(Tx \otimes Ty) \xrightarrow{T(t_{(2)}^{x,y})} T^{2}(x \otimes y)$$

$$\downarrow^{\mu_{x}^{\mathbb{T}} \otimes \mu_{y}^{\mathbb{T}}} \qquad \downarrow^{\mu_{x \otimes y}^{\mathbb{T}}}$$

$$Tx \otimes Ty \xrightarrow{t_{(2)}^{x,y}} T(x \otimes y)$$

$$(2.32)$$

and

$$x \otimes y \xrightarrow{\eta_x^{\mathbb{T}} \otimes \eta_y^{\mathbb{T}}} Tx \otimes Ty$$

$$\downarrow t_{(2)}^{x,y}$$

$$T(x \otimes y)$$

$$(2.33)$$

respectively.

Dually to the notion of a monad, a *comonad* on a category  $\mathcal{C}$  is a coalgebra  $\mathbb{T} = (T, \Delta^{\mathbb{T}}, \varepsilon^{\mathbb{T}})$  in the category of endofunctors of  $\mathcal{C}$ , i.e. the natural transformations  $\Delta^{\mathbb{T}} : T \Longrightarrow T^2$  and  $\varepsilon^{\mathbb{T}} : T \Longrightarrow Id_{\mathcal{C}}$  satisfy  $\Delta^{\mathbb{T}} T \circ \Delta^{\mathbb{T}} = T\Delta^{\mathbb{T}} \circ \Delta^{\mathbb{T}}$  and  $\varepsilon^{\mathbb{T}} T \circ \Delta^{\mathbb{T}} = Id_{\mathcal{C}} = T\varepsilon^{\mathbb{T}} \circ \Delta^{\mathbb{T}}$ .

**Definition 2.31.** [St, DP] (i) A Frobenius monad is a monad  $(T, \mu^{\mathbb{T}}, \eta^{\mathbb{T}})$  together with a comonad structure  $(T, \Delta^{\mathbb{T}}, \varepsilon^{\mathbb{T}})$  such that the equalities

$$(\mu^{\mathbb{T}} \otimes id_T) \circ (id_T \otimes \Delta^{\mathbb{T}}) = \Delta^{\mathbb{T}} \circ \mu^{\mathbb{T}} = (id_T \otimes \mu^{\mathbb{T}}) \circ (\Delta^{\mathbb{T}} \otimes id_T)$$
(2.34)

hold.

(ii) A commutative Frobenius monad is a Frobenius monad  $\mathbb{T}$  together with a natural isomorphism  $\tau \colon T^2 \Rightarrow T^2$  satisfying the hexagon identity  $T \tau \circ \tau T \circ T \tau = \tau T \circ T \tau \circ \tau T$  and being compatible with the monad and comonad structures according to  $\mu^{\mathbb{T}} \circ \tau = \mu^{\mathbb{T}}$ ,  $\tau \circ \eta^{\mathbb{T}} T = T \eta^{\mathbb{T}}$ ,  $\tau \circ \Delta^{\mathbb{T}} = \Delta^{\mathbb{T}}$  and  $\varepsilon^{\mathbb{T}} T \circ \tau = T \varepsilon^{\mathbb{T}}$ , as well as  $\tau \circ \mu^{\mathbb{T}} T = T \mu^{\mathbb{T}} \circ \tau T \circ T \tau$  and  $\Delta^{\mathbb{T}} T \circ \tau = T \tau \circ \tau T \circ T \Delta^{\mathbb{T}}$ . (iii) A separable monad is a monad whose multiplication has a one-sided inverse  $\eta^{\mathbb{T}}$ , so that  $\mu^{\mathbb{T}} \circ \eta^{\mathbb{T}} = id_T$ 

(iv) A coproduct-separable monad is a monad-comonad  $(T, \mu^{\mathbb{T}}, \eta^{\mathbb{T}}, \Delta^{\mathbb{T}}, \varepsilon^{\mathbb{T}})$  which is separable with  $\eta^{\mathbb{T}} = \Delta^{\mathbb{T}}$ , so that  $\mu^{\mathbb{T}} \circ \Delta^{\mathbb{T}} = id_T$ .

Remark 2.32. Just like for Frobenius algebras, there are several equivalent definitions of Frobenius monad [St, Thm. 1.6]. They basically boil down to the statement that the Frobenius pairing  $\varepsilon^{\mathbb{T}} \circ \mu^{\mathbb{T}}$  and Frobenius copairing  $\Delta^{\mathbb{T}} \circ \eta^{\mathbb{T}}$  furnish the counit and unit of a two-sided self-adjunction  $T \vdash T \dashv T$ . To appreciate this, recall that every adjunction  $\alpha$  of functors  $F: C \hookrightarrow \mathcal{D}: G$  (G right adjoint to F) gives rise to a monad  $(T, \mu^{\mathbb{T}}, \eta^{\mathbb{T}})$  on C, with T = GF, unit  $\eta^{\mathbb{T}}$  being the unit  $\eta^{\alpha}$  of the adjunction, and multiplication given by whiskering the counit  $\varepsilon^{\alpha}$  of the adjunction,  $\mu^{\mathbb{T}} = id_G \varepsilon^{\alpha} id_F$  [Ri, Lemma 5.1.3]. Likewise, it gives rise to a comonad  $(S, \Delta^{\mathbb{S}}, \varepsilon^{\mathbb{S}})$  on  $\mathcal{D}$ , with S = FG,  $\varepsilon^{\mathbb{S}} = \varepsilon^{\alpha}$  and  $\Delta^{\mathbb{S}} = id_F \eta^{\alpha} id_G$ . If the adjunction  $\alpha$  is two-sided, then there is accordingly both a monad and a comonad structure with underlying endofunctor T (and likewise for S); these fit together to form a Frobenius monad (this is easily seen when using bicategorical string diagrams, see e.g. [Müg, Sect. 1.2]).

To any algebra  $A = (\dot{A}, \mu, \eta)$  in a monoidal category C there is naturally associated a monad  $\mathbb{A} = (T_A, \mu^{\mathbb{A}}, \eta^{\mathbb{A}})$  on C, with  $T_A = -\otimes \dot{A}$ , and with multiplication and unit given by natural transformations  $\mu^{\mathbb{A}}$  and  $\eta^{\mathbb{A}}$  with components

$$\mu_c^{\mathbb{A}} = (id_c \otimes \mu) \circ a_{c,A,A} \quad \text{and} \quad \eta_c^{\mathbb{A}} = (id_c \otimes \eta) \circ \mathbf{u}_{r,c}^{-1}$$
 (2.35)

for  $c \in C$ . Indeed, the defining properties of  $\mu^{\mathbb{A}}$  and  $\eta^{\mathbb{A}}$  are equivalent to the associativity and unit properties of  $\mu$  and  $\eta$ . Likewise, to any coalgebra  $A = (\dot{A}, \Delta, \varepsilon)$  in C there is associated a comonad  $(T_A, \Delta^{\mathbb{A}}, \varepsilon^{\mathbb{A}})$  on C, with underlying endofunctor  $T_A$  and with natural transformations having components

$$\Delta_c^{\mathbb{A}} = a_{c,A,A}^{-1} \circ (id_c \otimes \Delta) \quad \text{and} \quad \varepsilon_c^{\mathbb{A}} = u_{r,c} \circ (id_c \otimes \varepsilon).$$
 (2.36)

By direct verification of the required equalities one checks:

**Lemma 2.33.** Let A be a Frobenius algebra in a monoidal category  $\mathcal{C}$ . The induced monad and comonad structures on the endofunctor  $T_A = -\otimes \dot{A}$  fit together to the structure of a Frobenius monad  $\mathbb{A} = (T_A, \mu^{\mathbb{A}}, \eta^{\mathbb{A}}, \Delta^{\mathbb{A}}, \varepsilon^{\mathbb{A}})$  on  $T_A$ . If A is a separable (coproduct-separable) algebra, then  $\mathbb{A}$  is a separable (coproduct-separable) monad. If  $\mathcal{C}$  is braided and A is commutative, then  $\mathbb{A}$  is a commutative Frobenius monad, with  $\tau_c = id_c \otimes \beta_{\dot{A},\dot{A}}$ .

### 2.6 Graded-enriched categories

For  $\Gamma$  a finite abelian group, the category  $\operatorname{vect}_{\Gamma}$  has as objects  $\Gamma$ -graded  $\mathbb{k}$ -vector spaces  $V = \bigoplus_{i \in \Gamma} V_i$  and as morphisms all  $\mathbb{k}$ -linear maps. The category  $\operatorname{vect}_{\Gamma}^{(0)}$  is its wide subcategory whose morphisms are the grade-preserving  $\mathbb{k}$ -linear maps. A symmetric monoidal structure on  $\operatorname{vect}_{\Gamma}$  that inherits the associativity and unit constraints from the monoidal category  $(\operatorname{vect}, \otimes)$  of  $\mathbb{k}$ -vector spaces (and which we thus still denote by  $\otimes$ ) is obtained by setting  $(V \otimes V')_i := \bigoplus_{j \in \Gamma} V_j \otimes V'_{i-j}$  for objects and  $(f \otimes f')(v \otimes v') := f(v) \otimes f'(v')$  for morphisms of  $\operatorname{vect}_{\Gamma}$ . The subcategory  $\operatorname{vect}_{\Gamma}^{(0)}$  inherits this symmetric monoidal structure.

As is well known, for  $\Gamma = \mathbb{Z}/2\mathbb{Z}$  the monoidal structure can be twisted by the non-trivial  $\mathbb{Z}/2\mathbb{Z}$ -bicharacter (see e.g. [BrE]), thereby yielding instead a *supermonoidal* structure. A natural generalization of this construction to arbitrary finite abelian groups has been presented in [Pu, BrD]. In the sequel we follow the exposition in [ScV].

We denote the grade of a homogeneous vector  $v \in V_i$  by |v| = i and the grade of a homogeneous linear map f by |f|.

**Definition 2.34.** Let  $\kappa$  be a  $\Gamma$ -bicharacter. The  $\kappa$ -twisted tensor product  $\otimes^{\kappa}$  on  $\text{vect}_{\Gamma}$  is defined on objects as  $V \otimes^{\kappa} V' := V \otimes V'$  and on morphisms as the quadrilinear extension of the prescription

$$(f \otimes^{\kappa} f')(v \otimes^{\kappa} v') := \kappa(|f'|, |v|) f(v) \otimes^{\kappa} f'(v')$$
(2.37)

for homogeneous morphisms  $f: V \to W$ ,  $f': V' \to W'$  and homogeneous vectors  $v \in V$ ,  $v' \in V'$ .

Note that, as we always work with some fixed group  $\Gamma$ , the dependence on  $\Gamma$  is suppressed in the notation and terminology. The  $\kappa$ -twisted tensor product is *not* a monoidal structure on vect $\Gamma$ . However, it enjoys properties analogous to those of a monoidal structure, except for the following:

**Lemma 2.35.** ( $\kappa$ -Twisted Interchange Law.) We have

$$(f' \otimes^{\kappa} g') \circ (f \otimes^{\kappa} g) = \kappa(|g'|, |f|) (f' \circ f) \otimes^{\kappa} (g' \circ g)$$
(2.38)

for any quadruple of homogeneous morphisms f, f', g, g' in  $\text{vect}_{\Gamma}$ .

*Proof.* This follows immediately by combining the character property of  $\kappa$  in both of its arguments with the identities |f(v)| = |f| + |v| and  $|g' \circ g| = |g'| + |g|$ .

#### Definition 2.36.

- 1. A  $\Gamma$ -graded  $\mathbb{k}$ -linear category is a category enriched over the symmetric monoidal category (vect<sub> $\Gamma$ </sub><sup>(0)</sup>,  $\otimes$ ).
- 2. A  $\Gamma$ -graded  $\mathbb{k}$ -linear functor is a functor enriched over  $(\text{vect}_{\Gamma}^{(0)}, \otimes)$ .
- 3. We denote by  $\mathbb{k}$ - $Cat_{\Gamma}$  the category having small  $\Gamma$ -graded  $\mathbb{k}$ -linear categories as objects and  $\Gamma$ -graded  $\mathbb{k}$ -linear functors as morphisms.
- 4. The  $\kappa$ -twisted Cartesian product  $C \boxtimes^{\kappa} \mathcal{D}$  of two  $\Gamma$ -graded  $\mathbb{k}$ -linear categories is defined on objects as the Cartesian product,  $Obj(C \boxtimes^{\kappa} \mathcal{D}) := Obj(C) \times Obj(\mathcal{D})$  for  $C, \mathcal{D} \in \mathbb{k}$ - $Cat_{\Gamma}$ , and on morphism spaces as the tensor product of vector spaces with composition of morphisms defined as in (2.38).

We indicate the prescription for the composition in  $\mathbb{C} \boxtimes^{\kappa} \mathcal{D}$  by writing

$$\operatorname{Hom}_{\mathbb{C}\boxtimes^{\kappa}\mathbb{D}}((c,d),(c',d')) = \operatorname{Hom}_{\mathbb{C}}(c,c') \otimes^{\kappa} \operatorname{Hom}_{\mathbb{D}}(d,d'). \tag{2.39}$$

By construction,  $C \boxtimes^{\kappa} \mathcal{D}$  is endowed with the structure of a  $\Gamma$ -graded  $\mathbb{k}$ -linear category and  $\boxtimes^{\kappa}$  becomes a monoidal structure on the category  $\mathbb{k}$ - $Cat_{\Gamma}$ .

**Definition 2.37.** [ScV, Def. 2.4] A  $(\Gamma, \kappa)$ -graded-monoidal category is a  $\Gamma$ -graded  $\kappa$ -linear category C together with a unit object and a  $\Gamma$ -graded  $\kappa$ -linear functor  $\otimes^{\kappa} : C \boxtimes^{\kappa} C \to C$  that satisfy the same unitality and associativity axioms as in the case of a strict monoidal category.

In particular, the  $\kappa$ -twisted tensor product  $\otimes^{\kappa}$  introduced in Definition 2.34 furnishes a  $\Gamma$ -graded  $\mathbb{k}$ -linear functor from  $\text{vect}_{\Gamma} \boxtimes^{\kappa} \text{vect}_{\Gamma}$  to  $\text{vect}_{\Gamma}$  and thus endows  $\text{vect}_{\Gamma}$  with the structure of a  $(\Gamma, \kappa)$ -graded-monoidal category.

# 3 A tensor product of modules over a graded-commutative algebra

Let C be a braided category,  $\Gamma$  a finite abelian group, and A a  $\Gamma$ -graded algebra in C that is graded-commutative with respect to a two-cocycle  $\kappa$  on  $\Gamma$ . Then by the defining equality (2.29), the opposite algebra  $A^{\text{op}}$  coincides with the cocycle-twisted algebra  $A^{[\kappa]}$  described in Remark 2.3. Now by Lemma 2.22, via the braiding one obtains a right  $A^{\text{op}}$ -module from a left A-module, while by Lemma 2.12, via cocycle-twisting one obtains a right  $A^{[\kappa]}$ -module from a  $\Gamma$ -graded right A-module. We combine the two procedures by setting, for a  $\Gamma$ -graded right A-module m = (m, q) with  $m = \bigoplus_i n_i$ ,

$$\rho_{i,i}^{(\beta;\kappa)} := \kappa(i,j)^{-1} \, \mathcal{Q}_{i,i} \circ \beta_{A_i,m_i} \tag{3.1}$$

for  $i, j \in \Gamma$ .

**Proposition 3.1.** Let  $m = (\dot{m}, \mathbf{q})$  be a  $\Gamma$ -graded right module over a graded-commutative algebra A.

(i) We have

$$m^{(\beta;\kappa)} := (\dot{m}, \rho^{(\beta;\kappa)}) \in A\text{-mod}, \qquad (3.2)$$

i.e.  $\kappa^{-1}$ -twisting the left  $A^{\text{op}}$ -module obtained from m by the braiding yields an ordinary left A-module.

(ii) If and only if the two-cocycle  $\kappa$  with respect to which A is graded-commutative is a bicharacter, the triple  $(\dot{m}, \rho^{(\beta;\kappa)}, \mathbf{q})$  is an A-bimodule.

*Proof.* (i) Since, as noticed above, the opposite algebra  $A^{\text{op}}$  of a graded-commutative algebra A coincides with  $A^{[\kappa]}$ , we can regard the left  $A^{\text{op}}$ -module that is obtained from m via the braiding as a left  $A^{[\kappa]}$ -module. The claim then follows by applying Lemma 2.12 'backwards' to the so obtained left  $A^{[\kappa]}$ -module.

(ii) By direct calculation one finds

$$\rho_{i,j+k}^{(\beta;\kappa)} \circ (id_{A_i} \otimes Q_{j,k}) = \kappa(i,j) \, \kappa(i,k) \, \kappa^{-1}(i,j+k) \, Q_{i+j,k} \circ (\rho_{i,j}^{(\beta;\kappa)} \otimes id_{A_k})$$
(3.3)

for  $i, j, k \in \Gamma$ . So the left and right A-actions on m commute,  $\rho^{(\beta;\kappa)} \circ (id_A \otimes \gamma) = \gamma \circ (\rho^{(\beta;\kappa)} \otimes id_A)$ , if and only if the cocycle  $\kappa$  is a character in its second argument. The statement then follows because a cocycle is a character in one argument iff it is a bicharacter.

**Remark 3.2.** Analogously, for a  $\Gamma$ -graded left A-module  $m = (\dot{m}, \rho)$  with underlying object  $\dot{m} = \bigoplus_i m_i$  we set  $\alpha_{i,j}^{(\beta;\kappa)} := \kappa(i,j)^{-1} \rho_{j,i} \circ \beta_{m_i,A_j}$  for  $i,j \in \Gamma$ . Then if A is graded-commutative,  $\kappa^{-1}$ -twisting the right  $A^{\text{op}}$ -module obtained from m by the braiding yields an ordinary right A-module,  $(\dot{m}, \alpha^{(\beta;\kappa)}) \in mod - A$ , and iff moreover  $\kappa$  is a bicharacter,  $(\dot{m}, \rho, \alpha^{(\beta;\kappa)})$  is an A-bimodule. The first statement is shown in the same way as part (i) of Proposition 3.1, while for the second statement one calculates

$$Q_{i+j,k}^{(\beta;\kappa)} \circ (\rho_{i,j} \otimes id_{A_k}) = \kappa(i,k) \, \kappa(j,k) \, \kappa^{-1}(i+j,k) \, \rho_{i,j+k} \circ (id_{A_i} \otimes Q_{i,k}^{(\beta;\kappa)})$$
(3.4)

for  $i, j, k \in \Gamma$ , meaning that for the left and right A-actions to commute,  $\kappa$  now has to be a character in its first argument, which is again equivalent to  $\kappa$  being a bicharacter.

**Remark 3.3.** By (the proof of) Lemma 2.26, the requirement that  $\kappa$  is a bicharacter is automatically satisfied (and in fact equivalent to A being graded-commutative) if the graded-commutative algebra A is a twisted group algebra.

With the help of Proposition 3.1 we obtain

**Proposition 3.4.** Let A be an algebra in a braided monoidal category C for which the tensor product  $\otimes_A$  of A-bimodules exists. Let  $\Gamma$  be a finite abelian group and let A be graded-commutative with respect to a cocycle  $\kappa$  on  $\Gamma$  that is a bicharacter on  $\Gamma$ .

- (i) The category  $mod_{\Gamma}^{\circ}$ -A of  $\Gamma$ -graded right A-modules and homogeneously even module morphisms admits a structure  $(mod_{\Gamma}^{\circ}$ - $A, \widetilde{\otimes})$  of a monoidal category.
- (ii) If A is in addition coproduct-separable Frobenius, then the tensor product  $m \otimes n$  of two  $\Gamma$ -graded right A-modules  $(\dot{m}, \mathbf{q}_m)$  and  $(\dot{n}, \mathbf{q}_n)$  is the image of the morphism

$$p_{m,n}^{[\kappa]} := (q_m \otimes \rho_n^{(\beta;\kappa)}) \circ (id_m \otimes (\Delta \circ \eta) \otimes id_n)$$
(3.5)

(which is an idempotent).

*Proof.* (i) Consider two right A-modules  $m=(\dot{m}, \mathbf{q}_m)$  and  $n=(\dot{n}, \mathbf{q}_n)$ . Invoking Proposition 3.1, we can promote m and n to A-bimodules  $(\dot{m}, \rho_m^{(\beta;\kappa)}, \mathbf{q}_m)$  and  $(\dot{n}, \rho_n^{(\beta;\kappa)}, \mathbf{q}_n)$ , which we still denote by m and n. The bimodule tensor product over A, as given in Definition 2.20, then yields an A-bimodule

$$m \widetilde{\otimes} n := (\dot{m}, \rho_m^{(\beta;\kappa)}, \mathfrak{q}_m) \otimes_A (\dot{n}, \rho_n^{(\beta;\kappa)}, \mathfrak{q}_n).$$
 (3.6)

By forgetting the left A-action, this becomes a right A-module, which we still denote by  $m \otimes n$ . Since  $\otimes_A$  is a monoidal structure on  $A\text{-}\mathit{bimod}$ , with associator and unitors inherited from C, it follows immediately that  $\widetilde{\otimes}$  is a monoidal structure on  $\mathit{mod}_{\Gamma}^{\circ}$ -A, again with associator and unitors inherited from those of C, and with A as monoidal unit.

(ii) If A is coproduct-separable Frobenius, then the tensor product  $\otimes_A$  can be described, as in Lemma 2.21, as the image of an idempotent. This idempotent is given by formula (2.24), with appropriate right and left A-actions. In the case at hand the latter are  $\mathfrak{q}_m$  and  $\rho_n^{(\beta;\kappa)}$ , respectively, thus yielding (3.5).

Remark 3.5. The following analogous results hold for left modules. For A an algebra in a braided monoidal category  $\mathcal{C}$  that is graded-commutative with respect to a cocycle  $\kappa$  on  $\Gamma$ , the category  $A\text{-}mod_{\Gamma}$  of  $\Gamma$ -graded left A-modules admits a structure  $(A\text{-}mod_{\Gamma}, \widetilde{\otimes})$  of a monoidal category. If A is in addition coproduct-separable Frobenius, then the tensor product  $m \widetilde{\otimes} n$  of two left A-modules m and n is the image of the idempotent  $(\gamma_m^{(\beta;\kappa)} \otimes \rho_n) \circ (id_m \otimes (\Delta \circ \eta) \otimes id_n)$ . In the particular case that A is a commutative algebra, this reproduces results that have been obtained, under some insignificant further conditions on  $\mathcal{C}$ , in [KO, Cor. 1.22] and (part of) [FFRS, Prop. 3.21(i)].

# 4 Induced modules over twisted group algebras

In Section 3 we have dealt with the category  $mod_{\Gamma}^{\circ}$ -A of  $\Gamma$ -graded right modules over a  $\Gamma$ -graded algebra A; by Definition 2.11,  $mod_{\Gamma}^{\circ}$ -A has homogeneously even module morphisms as morphisms. It is, however, equally natural to refrain from restricting to homogeneously even morphisms. We denote by  $mod_{\Gamma}$ -A the category having  $\Gamma$ -graded right A-modules as objects and all module morphisms between them as morphisms.

In full generality, there need not be any interesting grading on the morphisms of  $mod_{\Gamma}$ -A. However, upon induction, morphisms in C give rise to homogeneously even morphisms in  $mod_{\Gamma}$ -A, so to attain a non-trivial grading there need to be module morphisms that are not induced. Henceforth we therefore restrict our attention to a situation in which it is guaranteed that all modules under consideration are  $\Gamma$ -graded and that their morphism spaces are naturally graded as well and can be described explicitly: we require  $A = A_{(\Gamma)}$  to be a twisted group algebra in the sense of Example 2.6 and consider induced A-modules.

**Definition 4.1.** An induced (or free) right module over an algebra  $(A, \mu, \eta)$  in a monoidal category C is a right A-module of the form  $(\dot{m}, \varsigma) = Ind(x) := (x \otimes A, id_x \otimes \mu)$  for some  $x \in C$ . The category  $\dot{m}d$ -A is the full subcategory of mod-A whose objects are induced right A-modules.

Induced left modules and induced bimodules are defined analogously. The functor

$$Ind: \quad C \longrightarrow mo\ell - A \tag{4.1}$$

that maps  $x \in C$  to Ind(x) and  $f \in Hom(x, y)$  to  $f \otimes id_A \in Hom_A(Ind(x), Ind(y))$  for  $x, y \in C$  is called the *induction* (or *free*) functor. The functor Ind is faithful and is left adjoint to the forgetful functor  $U : mod - A \to C$  which acts as  $(\dot{m}, q) \mapsto \dot{m}$ . If A is Frobenius (such as in the case at hand, where A is a twisted group algebra), then Ind is also right adjoint to U, so that we have a two-sided adjunction  $Ind \vdash U \dashv Ind$ , i.e.

$$\operatorname{Hom}_{A}(\operatorname{Ind}(x), m) \cong \operatorname{Hom}(x, \dot{m}) \quad \text{and} \quad \operatorname{Hom}_{A}(m, \operatorname{Ind}(x)) \cong \operatorname{Hom}(\dot{m}, x).$$
 (4.2)

Next we make use of the fact that any  $\Gamma$ -graded algebra A is automatically  $\Gamma$ -graded as a bimodule over itself. By simply assigning grade 0 to an object  $x \in C$ , whereby  $x \otimes A$  is naturally graded as  $x \otimes A = \bigoplus_{i \in \Gamma} x_i$  with  $x_i = x \otimes A_i$ , this directly implies

**Lemma 4.2.** Every induced module over a  $\Gamma$ -graded algebra is naturally  $\Gamma$ -graded in the sense of Definition 2.11.

As a consequence, for a twisted group algebra A the category ind-A is  $\Gamma$ -graded in the sense of Definition 2.36.1. The  $\Gamma$ -grading on ind-A is in fact strong, in the sense that for every object Ind(x) and every  $i \in \Gamma$  there exists an isomorphism of grade i with domain Ind(x), namely  $Ind(x) \xrightarrow{\cong} Ind(x \otimes A_i)$ .

By construction, all morphisms in ind-A that are in the image of the induction functor (4.1) are homogeneously even. Non-even and non-homogeneous endomorphisms will be of particular interest for objects  $x \in \mathcal{C}$  for which the subset

$$\Sigma_x := \{ i \in \Gamma \mid x \otimes A_i \cong x \} \tag{4.3}$$

of  $\Gamma$  is non-trivial, i.e. which obey if  $x \otimes A_i \cong x$  for some homogeneous component of A with  $i \neq 0$ . We call  $\Sigma_x$  the *stabilizer* of x. Note that if  $x \otimes A_i \cong x \cong x \otimes A_j$  for  $i, j \in \Gamma$ , then also  $x \otimes A_{i+j} \cong x$ , i.e. the stabilizer is in fact a subgroup of  $\Gamma$ .

**Example 4.3.** Consider the semisimple modular tensor category  $C(A_1^{(1)}, k)$  introduced in Example 2.7, having non-isomorphic simple objects  $U_{\nu}$  with  $\nu \in \{0, 1, ..., k\}$  and Picard group  $\mathbb{Z}/2\mathbb{Z}$ . The stabilizers of the simple objects with respect to the group algebra  $A = U_0 \oplus U_k$  are trivial,  $\Sigma_{U_{\nu}} = \{0\}$ , unless  $\nu = k/2$ , in which case  $\Sigma_{U_k/2} = \mathbb{Z}/2\mathbb{Z}$ .

To study the structure of morphisms in ind-A in detail, we now further assume that C is k-linear abelian and that the monoidal unit of C is simple. If C has objects with non-trivial stabilizer, then ind-A is not idempotent-complete and hence not abelian, but we can still define an object  $m \in ind-A$  to be simple iff it has precisely two quotient objects, namely itself and the zero object. We refer to an induced A-module that is simple as an object of ind-A as a simple induced A-module, even if it would not be simple as an object in the category of all A-modules.

**Lemma 4.4.** Let C be k-linear abelian monoidal with simple monoidal unit, let  $\Gamma$  be an abelian group, and let  $A = A_{(\Gamma)}$  be a  $\Gamma$ -graded twisted group algebra. An object  $Ind(x) \in ind$ -A is a simple induced A-module if and only if  $x \in C$  is a simple object of C.

*Proof.* (i) If  $x \in C$  is not simple, then it has a non-trivial quotient object, so there exists a non-zero  $y \in C$ , not isomorphic to x, with an epimorphism  $f \in \text{Hom}(x,y)$ . We then also have  $x \otimes A \not\cong y \otimes A$  and hence  $Ind(x) \not\cong Ind(y)$ . Moreover, since the functor Ind is a left adjoint, it preserves epimorphisms, and hence  $Ind(f) \in \text{Hom}_A(Ind(x), Ind(y))$  is epi as well. It follows that  $Ind(x) \in ind$ -A is not simple.

(ii) Assume that the induced module Ind(x) is not simple, i.e. has a non-trivial quotient object in ind-A. Then there exists a non-zero  $m \in ind-A$ , not isomorphic to Ind(x), with an epimorphism in  $Hom_A(Ind(x), m)$ . By definition of ind-A there is thus an object  $y \in C$  with m = Ind(y). Using that Ind is left adjoint to the forgetful functor U it follows that

$$\operatorname{Hom}(x, y \otimes A) = \operatorname{Hom}(x, U\operatorname{Ind}(y)) \cong \operatorname{Hom}_A(\operatorname{Ind}(x), m) \neq 0. \tag{4.4}$$

Assume now that x is simple. If y is simple as well, then it follows from (4.4) that  $y \cong x \otimes A_i$  for some  $i \in \Gamma$ , and thus that  $Ind(x) \cong Ind(y) = m$ , which is a contradiction. In case y is not simple, then it has a non-trivial simple quotient  $y' \in C$  with an epimorphism in  $Hom_C(y, y')$ , and thus in particular  $Hom_A(Ind(y), Ind(y')) \cong Hom(y, y' \otimes A) \neq 0$ . This implies, in turn, situation is reduced to the previous one with y' in place of y.

In a k-linear abelian monoidal category C with simple monoidal unit Schur's Lemma holds (see e.g. [EGNO, Lemma 1.5.2]): any non-zero morphism between two simple objects in C is an isomorphism, and Hom(x,x) is a k-division algebra for any simple object  $x \in C$ . Thus, since we assume k to be algebraically closed, for any pair of simple objects  $x,y \in C$  we have  $Hom(x,y) \cong k$  if x and y are isomorphic, and Hom(x,y) = 0 else. We will show that in this situation there is a graded analogue of Schur's Lemma for induced modules over twisted group algebras in C. We first observe

**Proposition 4.5.** Let A be a  $\Gamma$ -graded twisted group algebra in a k-linear abelian monoidal category C with simple monoidal unit, with  $\Gamma$  a finite abelian group. Let  $x \in C$  be simple, with stabilizer  $\Sigma_x$ . Then there is an isomorphism

$$\operatorname{End}_{A}(\operatorname{Ind}(x)) \cong \mathbb{k}_{\varpi} \Sigma_{x} \tag{4.5}$$

of  $\Sigma_x$ -graded k-algebras between the endomorphims of the induced module  $Ind(x) \in ind - A$  and the group algebra  $k \Sigma_x$  twisted by a two-cocycle  $\varpi$ .

*Proof.* We first establish (4.5) as an isomorphism of vector spaces. Abbreviate  $\Sigma_x \equiv \Sigma$ . Select for each  $\Sigma$ -coset c of  $\Gamma$  a representative  $i_c \in \Gamma$ , so that  $c = i_c + \Sigma = \{i_c + s \mid s \in \Sigma\}$ . Then as a set we have  $\Gamma = \{i_c + s \mid c \in \Gamma/\Sigma, s \in \Sigma\}$  and thus

$$x \otimes A = \bigoplus_{s \in \Sigma} \bigoplus_{c \in \Gamma/\Sigma} x \otimes A_{i_c+s} \cong \bigoplus_{s \in \Sigma} \left( \bigoplus_{c \in \Gamma/\Sigma} x \otimes A_s \otimes A_{i_c} \right) \cong \bigoplus_{s \in \Sigma} \left( \bigoplus_{c \in \Gamma/\Sigma} x \otimes A_{i_c} \right). \tag{4.6}$$

In short,  $x \otimes A$  decomposes into a direct sum of  $|\Sigma|$  copies of the ' $\Gamma$ -orbit' of x which is the object  $\bigoplus_{c \in \Gamma/\Sigma} x \otimes A_{i_c}$ , where up to isomorphism the choice of representatives  $i_c$  of the cosets  $c \in \Gamma/\Sigma$  is irrelevant. It follows that as vector spaces we have

$$\operatorname{End}_{A}(\operatorname{Ind}(x)) \stackrel{(4.2)}{\cong} \operatorname{Hom}(U(\operatorname{Ind}(x)), x) = \operatorname{Hom}_{\mathcal{C}}(x \otimes A, x)$$

$$\stackrel{(4.6)}{\cong} \bigoplus_{s \in \Sigma} \bigoplus_{c \in \Gamma/\Sigma} \operatorname{Hom}(x \otimes A_{i_{c}}, x) \cong \bigoplus_{s \in \Sigma} \operatorname{Hom}(x, x) \cong \mathbb{k}^{|\Sigma|}.$$

$$(4.7)$$

To see that  $\operatorname{End}_A(\operatorname{Ind}(x))$  is isomorphic to  $\Bbbk \Sigma$  as a  $\Sigma$ -graded vector space, we provide a graded basis. For  $s \in \Sigma$  select an isomorphism  $\varphi_s \in \operatorname{Hom}(x, x \otimes A_s)$  and define an endomorphism  $\phi_s := \bigoplus_{i \in \Gamma} \phi_{s;i}$  of  $x \otimes A$  by

$$\phi_{s;i}: \quad x \otimes A_i \xrightarrow{\varphi_s \otimes id} (x \otimes A_s) \otimes A_i \xrightarrow{a_{x,A_s,A_i}} x \otimes (A_s \otimes A_i) \xrightarrow{id \otimes \mu_{s,i}} x \otimes A_{i+s}. \tag{4.8}$$

It follows directly from associativity of A that the morphisms  $\phi_s$  are in fact module endomorphisms,  $\phi_s \in \operatorname{End}_A(\operatorname{Ind}(x))$ . Also,  $\phi_s$  is homogeneous of grade s, so that in particular the set  $\{\phi_s \mid s \in \Sigma\}$  is linearly independent and is thus a graded basis of  $\operatorname{End}_A(\operatorname{Ind}(x)) \cong \mathbb{k}^{|\Sigma|}$ . Finally we have  $\phi_{s'} \circ \phi_s = \varpi(s, s') \phi_{s+s'}$  with numbers  $\varpi(s, s') \in \mathbb{k}^{\times}$  for  $s, s' \in \Sigma$ . Associativity of composition amounts to the statement that the numbers  $\varpi(s, s')$  assemble to a two-cocycle on  $\Sigma$ . Modifying the choice of isomorphisms  $\varphi_s$  changes  $\varpi$  by a two-coboundary.

#### Theorem 4.6. (Graded Schur Lemma)

Let A be a  $\Gamma$ -graded twisted group algebra in a k-linear abelian monoidal category C with

simple monoidal unit, with  $\Gamma$  a finite abelian group.

- (i) The space of module morphisms between any two simple induced A-modules Ind(x) and Ind(y) is either zero or isomorphic as a  $\Gamma$ -graded k-vector space to the spaces of endomorphisms of both Ind(x) and Ind(y).
- (ii) Any non-zero homogeneous module morphism between simple induced A-modules is an isomorphism.

*Proof.* (i) Analogously as in (4.7) we have  $\operatorname{Hom}_A(\operatorname{Ind}(x),\operatorname{Ind}(y)) \cong \operatorname{Hom}_{\mathcal{C}}(x \otimes A,y)$ . If  $x \not\cong y \otimes A_i$  for every  $i \in \Gamma$ , then this space is zero. Otherwise we have  $x \cong y \otimes A_j$  for some  $j \in \Gamma$  (which implies that  $\Sigma_y = \Sigma_x$  and thus  $\operatorname{End}_A(\operatorname{Ind}(y)) \cong \operatorname{End}_A(\operatorname{Ind}(x))$ ), and again analogously as in (4.7) we obtain

$$\operatorname{Hom}_{A}(\operatorname{Ind}(x),\operatorname{Ind}(y)) \cong \bigoplus_{s \in \Sigma_{x}} \bigoplus_{c \in \Gamma/\Sigma_{x}} \operatorname{Hom}(x \otimes A_{i_{c}+j},x). \tag{4.9}$$

Now for any  $j \in \Gamma$  there is a permutation  $\pi$  of  $\Gamma/\Sigma_x$  such that for every  $c \in \Gamma/\Sigma_x$  one has  $i_c + j = i_{\pi(c)} + s'$  with some  $s' \in \Sigma_x$ . This implies that the right hand side of (4.9) is isomorphic to  $\bigoplus_{s \in \Sigma_x} \bigoplus_{c \in \Gamma/\Sigma_x} \operatorname{Hom}(x \otimes A_{i_c}, x)$ , and hence  $\operatorname{Hom}_A(\operatorname{Ind}(x), \operatorname{Ind}(y)) \cong \operatorname{End}_A(\operatorname{Ind}(x))$  as vector spaces. It then also follows directly that  $\operatorname{Hom}_A(\operatorname{Ind}(x), \operatorname{Ind}(y))$  inherits a  $\Gamma$ -grading from  $\operatorname{End}_A(\operatorname{Ind}(x))$ .

(ii) Via composition, the space  $\operatorname{Hom}_A(\operatorname{Ind}(x),\operatorname{Ind}(y))$  has a natural structure of an  $\operatorname{End}_A(\operatorname{Ind}(x))$ - $\operatorname{End}_A(\operatorname{Ind}(y))$ -bimodule. As a consequence the homogeneous subspaces of the  $\Gamma$ -graded vector space  $\operatorname{Hom}_A(\operatorname{Ind}(x),\operatorname{Ind}(y))$  are one-dimensional.

Remark 4.7. As mentioned, for abelian monoidal categories the ordinary Schur Lemma is valid. Accordingly, if the category ind-A is abelian, then Theorem 4.6 implies the ordinary Schur Lemma. But generically, ind-A is not idempotent complete and thus not abelian. On the other hand, at least if A is a special Frobenius algebra, every A-module is a module retract of an induced module [FFRS, Lemma 4.8] and the category mod-A of all A-modules is the idempotent completion of ind-A. As a consequence, if C is abelian and A is special Frobenius, then so is mod-A, so that the ordinary Schur Lemma holds in mod-A. Also, as follows directly from the proof of Theorem 4.6, the ordinary Schur Lemma holds in the wide subcategory  $ind^{\circ}-A$  of ind-A whose morphisms are the homogeneously even morphisms of ind-A.

**Example 4.8.** For  $\Gamma = \mathbb{Z}/2\mathbb{Z}$ , i.e.  $A = \mathbf{1} \oplus L$  with L an order-2 invertible object, we have

$$\operatorname{Hom}_{A}(\operatorname{Ind}(x),\operatorname{Ind}(y)) \cong \begin{cases} \mathbb{k} & \text{if } x \otimes L \not\cong x \text{ and } (x \cong y \text{ or } x \cong y \otimes L), \\ \mathbb{k}(\mathbb{Z}/2\mathbb{Z}) & \text{if } x \otimes L \cong x \cong y, \\ 0 & \text{otherwise}. \end{cases}$$

$$\tag{4.10}$$

Remark 4.9. Particular realizations of Proposition 4.5 and Theorem 4.6, or of parts thereof, are well known, like e.g. Theorem 14 of [MaZ] which describes homogeneous automorphisms of graded modules over graded Lie algebras. In the representation theory of superalgebras, the trichotomy (4.10), then sometimes called the *Super Schur Lemma*, lists the possible dimensions of morphism spaces between simple modules over a finite-dimensional associative superalgebra, see e.g. Corollary 2.15 of [Joz] and Lemma 2.3 of [WW]. In that case, the possibility of having stabilizer  $\mathbb{Z}/2\mathbb{Z}$  is coupled to the existence of the simple superalgebras of type Q(n); accordingly, modules with stabilizer  $\mathbb{Z}/2\mathbb{Z}$  are then called modules of type Q. In the context of

spin topological quantum field theories and their applications to fermionic topological phases in condensed matter physics, objects with stabilizer  $\mathbb{Z}/2\mathbb{Z}$  are also known as *Majorana objects*, see e.g. [ALW]. It is, however, worth keeping in mind that what is relevant for the possible existence of such objects is only that the relevant algebra A is  $\mathbb{Z}/2\mathbb{Z}$ -graded, not that it admits any kind of super structure (compare also Example 2.27).

# 5 A graded-monoidal structure for induced modules

In this section we take C to be braided and A to be an algebra in C that is graded-commutative for some finite abelian group  $\Gamma$  and normalized two-cocycle  $\kappa$ .

As seen above, the category ind-A is naturally  $\Gamma$ -graded. Also, as a full subcategory of  $mod_{\Gamma}^{\circ}$ -A, the category ind°-A inherits the tensor product  $\widetilde{\otimes}$  that we considered in Section 3 (see Proposition 3.4). Indeed, the braiding of C gives a (non-canonical) isomorphism  $UInd(x) \otimes UInd(y) = x \otimes A \otimes y \otimes A \xrightarrow{\cong} x \otimes y \otimes A \otimes A = UInd(x \otimes y) \otimes A$ . This isomorphism fits together with the A-action on Ind(x) and Ind(y) in such a way that the coequalizer in the definition of  $Ind(x) \otimes_A Ind(y)$  becomes isomorphic as an object to  $x \otimes y \otimes B$  with B the coequalizer of  $id_A \otimes \mu$  and  $\mu \otimes id_A$  on  $A \otimes A \otimes A$ , i.e.  $B \cong A \otimes_A A \cong A$ ; moreover, the so obtained isomorphism from  $U(Ind(x) \otimes_A Ind(y))$  to  $x \otimes y \otimes A = UInd(x \otimes y)$  is compatible with the A-action on  $Ind(x \otimes y)$ .

An obvious question is then whether the monoidal structure  $\widetilde{\otimes}$  on  $ind^{\circ}$ -A can be extended to all of ind-A. As we will see, this is the case only if A is in fact commutative. In the general case it is natural to try to construct instead a suitably twisted monoidal structure on ind-A that makes it a  $(\Gamma, \kappa)$ -graded-monoidal category in the sense of Definition 2.37, and that reduces to the tensor product  $\widetilde{\otimes}$  for homogeneous morphisms of grade zero.

To find a candidate for such a twisted monoidal structure on ind-A, it will be convenient to consider the monad  $\mathbb{A}$  with underlying endofunctor  $T_A = -\otimes A$ , as described in (2.35), and to identify a corresponding candidate for a twisted monoidal structure on  $\mathbb{A}$ . Working with  $\mathbb{A}$ , we realize ind-A as the Kleisli category [Bo, Ch. 4.1] over  $\mathbb{A}$ .

**Definition 5.1.** The Kleisli category over a monad  $\mathbb{T} = (T, \mu^{\mathbb{T}}, \eta^{\mathbb{T}})$  on a category C is the following category  $C_{\mathbb{T}}$ : The objects of  $C_{\mathbb{T}}$  are the same as those of C – for distinction, we write  $\overline{x}$  for the object  $x \in C$  when regarded as an object of  $C_{\mathbb{T}}$ . The morphisms of  $C_{\mathbb{T}}$  are given by  $\operatorname{Hom}_{C_{\mathbb{T}}}(\overline{x}, \overline{y}) = \operatorname{Hom}_{C}(x, Ty)$ . The composition  $\operatorname{Hom}_{C_{\mathbb{T}}}(\overline{y}, \overline{z}) \times \operatorname{Hom}_{C_{\mathbb{T}}}(\overline{x}, \overline{y}) \to \operatorname{Hom}_{C_{\mathbb{T}}}(\overline{x}, \overline{z})$  of morphisms is the Kleisli composition

$$(g,f) \longmapsto \mu_z^{\mathbb{T}} \circ Tg \circ f =: g \,\overline{\circ}\, f$$
 (5.1)

for  $f \in \operatorname{Hom}_{C_{\mathbb{T}}}(\overline{x}, \overline{y})$  and  $g \in \operatorname{Hom}_{C_{\mathbb{T}}}(\overline{y}, \overline{z})$ . The identity morphism of  $\overline{x}$  is the relevant component of the unit of the monad,  $id_{\overline{x}} = \eta_x^{\overline{x}} \colon x \to Tx$ .

Analogously as for algebras, an induced (or free) module over a monad  $\mathbb T$  on C is a  $\mathbb T$ -module of the form  $Ind_{\mathbb T}(x):=(Tx,\mu_x^{\mathbb T})$  for some  $x\in C$ . Of significance to us is that the category  $\mathbb T$ -ind of induced modules over a monad  $\mathbb T$  is equivalent to the Kleisli category  $C_{\mathbb T}$  of  $\mathbb T$ . An equivalence  $C_{\mathbb T} \xrightarrow{\cong} \mathbb T$ -ind is given by mapping objects as  $\overline x \mapsto Ind_{\mathbb T}(x)$  and morphisms as  $(\overline x \xrightarrow{f} \overline y) \mapsto \mu_y^{\mathbb T} \circ Tf$  [Bo, Prop. 4.1.6]; as a quasi-inverse this has the functor given by  $Ind_{\mathbb T}(x) \mapsto \overline x$  and  $(Ind(x) \xrightarrow{g} Ind(y)) \mapsto g \circ \eta_x^{\mathbb T}$ .

Furthermore, in the case of the monad  $\mathbb{A}$ , the category ind-A of induced A-modules is isomorphic to the category of induced  $\mathbb{A}$ -modules and thus equivalent to the Kleisli category  $C_{\mathbb{A}}$ . The Kleisli composition (5.1) on  $C_{\mathbb{A}}$  reads

$$g \,\overline{\circ} \, f = (\mathrm{i} d_z \otimes \mu) \circ (g \otimes \mathrm{i} d_A) \circ f \,. \tag{5.2}$$

A monoidal structure  $t_{(2)}$  on a monoid  $\mathbb{T}$  on a monoidal category C equips its Kleisli category  $C_{\mathbb{T}}$  with a tensor product  $\overline{\otimes}$  via setting  $\overline{x} \overline{\otimes} \overline{y} := \overline{x \otimes y}$  for  $x, y \in C$  and

$$f \overline{\otimes} g := t_{(2)}^{x',y'} \circ (f \otimes g) : \quad x \otimes y \longrightarrow T(x' \otimes y')$$
 (5.3)

for morphisms  $f: x \to Tx'$  and  $g: y \to Ty'$  in C. Indeed, there is a bijection between monoidal structures on  $\mathbb{T}$  and monoidal structures on  $C_{\mathbb{T}}$  for which the induction functor from C to  $C_{\mathbb{T}}$  is strict monoidal [Se, Prop. 1.2.2]. Thus if  $\mathbb{T}$  is endowed with a monoidal structure, then there is a monoidal structure on the category of induced  $\mathbb{T}$ -modules such that the equivalence  $C_{\mathbb{T}} \xrightarrow{\simeq} \mathbb{T}$ -in $\mathcal{J}$  is strict monoidal.

Now recall from Definition 2.30 that a monoidal structure on a monad requires the multiplication of the monad to become a monoidal natural transformation. We observe

**Lemma 5.2.** Let  $t_{(2)}$  be a monoidal structure on a monad  $\mathbb{T} = (T, \mu^{\mathbb{T}}, \eta^{\mathbb{T}})$ . The monoidality of the multiplication  $\mu^{\mathbb{T}}$  is equivalent to the validity of the interchange law

$$(f' \overline{\otimes} g') \overline{\circ} (f \overline{\otimes} g) = (f' \overline{\circ} f) \overline{\otimes} (g' \overline{\circ} g)$$

$$(5.4)$$

for any quadruple of morphisms  $f: x \to Tx'$ ,  $g: y \to Ty'$ ,  $f': x' \to Tx''$  and  $g': y' \to Ty''$  in C.

*Proof.* Monoidality of  $\mu^{\mathbb{T}}$  is expressed by commutativity of the diagram (2.32), i.e. by the equality  $\mu_{x\otimes y}^{\mathbb{T}} \circ T(t_{(2)}^{x,y}) \circ t_{(2)}^{Tx,Ty} = t_{(2)}^{x,y} \circ (\mu_x^{\mathbb{T}} \otimes \mu_y^{\mathbb{T}})$ . Precomposing with  $(Tf' \otimes Tg') \circ (f \otimes g)$ , the left hand side of this equality becomes, using naturality of  $t_{(2)}$ ,

$$\mu_{x''\otimes y''}^{\mathbb{T}}\circ T(t_{(2)}^{x'',y''})\circ t_{(2)}^{Tx'',Ty''}\circ (Tf'\otimes Tg')\circ (f\otimes g)$$

$$=\mu_{x''\otimes y''}^{\mathbb{T}}\circ T(t_{(2)}^{x'',y''}\circ (f'\otimes g'))\circ (t_{(2)}^{x',y'}\circ (f\otimes g))\equiv (f'\overline{\otimes}g')\overline{\circ}(f\overline{\otimes}g),$$
(5.5)

while the right hand side becomes

$$t_{(2)}^{x'',y''} \circ (\mu_{x''}^{\mathbb{T}} \otimes \mu_{y''}^{\mathbb{T}}) \circ (Tf' \otimes Tg') \circ (f \otimes g)$$

$$= t_{(2)}^{x'',y''} \circ ((\mu_{x''}^{\mathbb{T}} \circ Tf' \circ f) \otimes (\mu_{y''}^{\mathbb{T}} \circ Tg' \circ g)) \equiv (f' \overline{\circ} f) \overline{\otimes} (g' \overline{\circ} g).$$
(5.6)

Thus commutativity of (2.32) implies (5.4). Conversely, when taking  $x = Tx' = T^2x''$  and  $y = Ty' = T^2y''$  as well as  $f = id_{Tx'}$  etc., (5.4) reduces to commutativity of (2.32).

**Remark 5.3.** In case A is a coproduct-separable Frobenius algebra (e.g. a twisted group algebra), the tensor product of A-bimodules, and hence also  $\widetilde{\otimes}$ , can be expressed as the image of the idempotent (3.5). That  $Ind(x) \widetilde{\otimes} Ind(y)$  can be identified with  $Ind(x \otimes y)$  is then understood directly by noticing that the object underlying the module  $Ind(x) \widetilde{\otimes} Ind(y)$  is the image of

$$id_x \otimes \left[ (\beta_{A,y}^{-1} \otimes id_A) \circ (id_y \otimes p_{A,A}) \circ (\beta_{A,y} \otimes id_A) \right] \in \operatorname{End}(x \otimes A \otimes y \otimes A),$$
 (5.7)

with  $p_{A,A}$  the idempotent

$$p_{A|A} := (\mu \otimes \mu) \circ (id_A \otimes (\Delta \circ \eta) \otimes id_A) = \Delta \circ \mu \tag{5.8}$$

in  $\operatorname{End}(A \otimes A)$ , by which A is canonically exhibited as a retract of  $A \otimes A$ . Also, if A is separable Frobenius, then by Lemma 2.33 the monad A is separable Frobenius in the sense of Definition 2.31, with comonad structure given by (2.36).

We take the explicit form of the idempotent (5.8) a motivation to make the ansatz

$$t_{(2)}^{x,y} := (id_x \otimes id_y \otimes \mu) \circ (id_x \otimes \beta_{A,y} \otimes id_A) \tag{5.9}$$

for  $x, y \in C$  as a candidate for a twisted analogue of a monoidal structure on the monad  $\mathbb{A}$ , even for algebras A that are not Frobenius. This ansatz is vindicated by

#### Lemma 5.4.

- (i) For  $A = (A, \mu, \eta)$  an algebra in a braided monoidal category  $(C, \otimes, \mathbf{1}; \beta)$ , the family (5.9) of morphisms furnishes a monoidal structure on the monad  $\mathbb{A}$  if and only if A is commutative.
- (ii) If  $A = (A, \mu, \eta, \Delta, \varepsilon)$  is a commutative coproduct-separable Frobenius algebra, then the monoidal structure (5.9) on  $\mathbb{A}$  amounts to a tensor product on the Kleisli category  $C_{\mathbb{A}}$  that is isomorphic to the tensor product  $\widetilde{\otimes}$  of induced A-modules.

Proof. (i) That the family (5.9) of morphisms gives a lax monoidal structure on the functor  $-\otimes A$  follows immediately from the definitions, while naturality of the family (5.9) and monoidality of the natural transformation  $\eta^{\mathbb{A}}$  are direct consequences of the naturality of the braiding  $\beta$ . Finally, that the natural transformation  $\mu^{\mathbb{A}} : \mathbb{A}^2 \to \mathbb{A}$  satisfies (2.32) boils down to the equality  $\mu \circ (\mu \otimes \mu) \circ (id_A \otimes \beta_{A,A} \otimes id_A) = \mu \circ (\mu \otimes \mu)$ . This is satisfied iff A is commutative. (ii) By coproduct-separability of A, the family  $s_{(2)}^{x,y} := (id_x \otimes \beta_{A,y}^{-1} \otimes id_A) \circ (id_x \otimes id_y \otimes \Delta)$  of morphisms is a one-sided inverse to the family (5.9),  $t_{(2)} \circ s_{(2)} = id$ . Accordingly we obtain a tensor product on the category  $C_{\mathbb{A}}$  corresponding to (5.9) that can be described as the image of the idempotent  $s_{(2)}^{x,y} \circ t_{(2)}^{x,y}$ . Moreover, the Frobenius property and commutativity of A imply that  $s_{(2)}^{x,y} \circ t_{(2)}^{x,y} = p_{Ind(x),Ind(y)}$  for all  $x, y \in C$ , with  $p_{m,n}$  as defined in (2.24), i.e.  $s_{(2)}^{x,y} \circ t_{(2)}^{x,y}$  equals the idempotent whose image is the tensor product  $Ind(x) \otimes_A Ind(y)$  of induced A-modules.  $\square$ 


According to Lemma 5.4 the family (5.9) is no longer a monoidal structure on  $\mathbb{A}$  when the algebra A is graded-commutative for a non-trivial two-cocycle  $\kappa$ . Instead we have

**Proposition 5.5.** Let A be a graded-commutative algebra in a  $\mathbb{k}$ -linear braided monoidal category C. Define  $\overline{\otimes}$  as in (5.3), with  $t_{(2)}$  given by (5.9). Then in the Kleisli category  $C_{\mathbb{A}}$  the  $\kappa$ -twisted interchange law

$$(f' \,\overline{\circ}\, f) \,\overline{\otimes}\, (g' \,\overline{\circ}\, g) = \sum_{i,j \in \Gamma} \kappa(i,j) \,(f' \,\overline{\otimes}\, g'_j) \,\overline{\circ}\, (f_i \,\overline{\otimes}\, g) \tag{5.10}$$

holds for all quadruples  $f: x \to x' \otimes A$ ,  $g: y \to y' \otimes A$ ,  $f': x' \to x'' \otimes A$  and  $g': y' \to y'' \otimes A$  of morphisms, where  $f = \sum_{i \in \Gamma} f_i$  and  $g' = \sum_{j \in \Gamma} g'_j$  are the decompositions of f and g' into sums of homogeneous morphisms  $f_i: x \to x' \otimes A_i$  and  $g'_j: y' \to y'' \otimes A_j$ , respectively.

*Proof.* Instead of writing out the relevant formulas, which are a bit lengthy, we work with string diagrams. Using associativity of A we have



If A is graded-commutative, i.e. satisfies (2.29), then the right hand sides of 5.11 and 5.12, when restricted to the components  $f_i$  and  $g'_j$ , differ by a factor  $\kappa(i, j)$ , thus proving the equality (5.10).

**Remark 5.6.** If  $\kappa$  is the trivial cocycle – and thus in particular if  $\Gamma$  is trivial, i.e. A is commutative – we obtain the ordinary interchange law and hence an ordinary monoidal structure on the category  $in \delta - A$ .

Now recall the notion of a  $(\Gamma, \kappa)$ -graded-monoidal category from Definition 2.37. The previous observations can be combined to

**Theorem 5.7.** Let  $\Gamma$  be a finite abelian group and  $\kappa$  a  $\Gamma$ -bicharacter. Let  $\mathcal{C}$  be a  $\mathbb{k}$ -linear braided monoidal category and A a  $\Gamma$ -graded algebra in  $\mathcal{C}$  that is graded-commutative with respect to  $\kappa$ .

- (i) The category ind-A can be endowed with the structure of a  $(\Gamma, \kappa)$ -graded-monoidal category, with  $\kappa$ -twisted monoidal structure obtained by transporting  $\overline{\otimes}$  from  $C_A$  to ind-A.
- (ii) If in addition the tensor product  $\otimes_A$  of A-bimodules exists and A is coproduct-separable Frobenius, then the restriction of  $\overline{\otimes}$  to  $ind^{\circ}$ -A is isomorphic to the tensor product  $\widetilde{\otimes}$  on  $ind^{\circ}$ -A that is induced from the bimodule tensor product  $\otimes_A$ .

*Proof.* The  $\kappa$ -twisted interchange law (5.10) amounts to a structure of a  $(\Gamma, \kappa)$ -graded-monoidal category  $(C^A, \overline{\otimes})$  on the Kleisli category of  $\mathbb{A}$ . This can be transported to  $\mathbb{A}$ -ind and thus to ind-A in the same way as a monoidal structure. Claim (ii) follows directly from Lemma 5.4(ii).

Remark 5.8. If the neutral component  $A_0$  of A is isomorphic to the tensor unit 1 (which is e.g. the case for twisted group algebras), then every homogeneously even morphism in ind-A is in the image of the induction functor  $Ind: C \to ind-A$ . As a consequence, Ind restricts to an equivalence from C to the wide subcategory  $ind^{\circ}-A$  of ind-A. Moreover, the restriction of the  $\kappa$ -twisted tensor product on ind-A to  $ind^{\circ}-A$  defines a tensor product on  $ind^{\circ}-A$  and, provided that the tensor product of A-bimodules exists, this coincides with the tensor product  $\widetilde{\otimes}$  that  $ind^{\circ}-A$  inherits as a full subcategory of  $mod_{\Gamma}^{\circ}-A$ . Accordingly we could equivalently have defined the tensor product on  $ind^{\circ}-A$  by setting  $Ind(x) \otimes_{ind^{\circ}-A} Ind(y) := Ind(x \otimes_{C} y)$ .

Remark 5.9. One notable inspiration for this text is the study of supercategories in [BrE]. The 1-categorical structure of a monoidal supercategory in [BrE, Def. 1.4(i)] coincides with the tensor product with which our construction would endow the  $\kappa$ -graded right modules of a superalgebra, with  $\kappa(i,j) = (-1)^{ij}$  the non-trivial  $\mathbb{Z}/2\mathbb{Z}$ -bicharacter. That we work with right rather than left modules has its origin in the fact that we then produce the same super interchange law  $(f' \otimes^{\kappa} g') \circ (f \otimes^{\kappa} g) = (-1)^{|g'||f|} (f' \circ f) \otimes^{\kappa} (g' \circ g)$  as in [BrE]. Had we used left modules instead, we would have arrived at  $(f' \otimes^{\kappa} g') \circ (f \otimes^{\kappa} g) = (-1)^{|f'||g|} (f' \circ f) \otimes^{\kappa} (g' \circ g)$ , as the braiding involved would affect the complementary components.

In the event that  $A = A_0 \oplus A_1$  is a twisted group algebra over  $\mathbb{Z}/2\mathbb{Z}$ , we also get what in [BrE] is called a  $\Pi$ -structure: it is given by tensoring with the invertible object  $Ind(A_1)$ . The latter is an *odd unit*, i.e. odd-isomorphic to the monoidal unit  $Ind(A_0)$ . Thus in this case  $in\mathcal{A}$ -A is a monoidal  $\Pi$ -supercategory in the sense of Definition 1.12 on [BrE].

#### ACKNOWLEDGMENTS:

We thank Lukas Woike for helpful comments. J.F. is supported by VR under project no. 2022-02931.

# References

- [ALW] D. Aasen, E. Lake, and K. Walker, Fermion condensation and super pivotal categories, J. Math. Phys. 60 (2019) 121901\_1-112 [cond-mat/1709.01941]
- [Bo] F. Borceux, Handbook of Categorical Algebra. Volume 2: Categories and Structures (Cambridge University Press, Cambridge 1994)
- [BrD] J. Brundan and N. Davidson, Categorical actions and crystals, Contemp. Math. 683 (2017) 105–147 [math.QA/1603.08938]
- [BrE] J. Brundan and A.P. Ellis, Monoidal supercategories, Commun. Math. Phys. 351 (2017) 1045–1089 [math.RT/1603.05928]
- [CGO] A.M. Cegarra, A.R. Garzón, and J.A. Ortega, Graded extensions of monoidal categories, J. Algebra 241 (2001) 620–657
- [DNN] S. Dăscălescu, C. Năstăsescu, and L. Năstăsescu, Frobenius algebras of corepresentations and group-graded vector spaces, J. Algebra 406 (2014) 226–250 [math.QA/1307.7304]
- [DP] K. Došen and Z. Petrić, Syntax for split preorders, Ann. Pure Appl. Logic 164 (2013) 443–481
- [EGNO] P.I. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik, *Tensor Categories* (American Mathematical Society, Providence 2015)
- [FFRS] J. Fröhlich, J. Fuchs, I. Runkel, and C. Schweigert, Correspondences of ribbon categories, Adv. Math. 199 (2006) 192–329 [math.CT/0309465]
- [FrW] A. Fröhlich and C.T.C. Wall, Graded monoidal categories, Compos. Math. 28 (1974) 229–285
- [FuG] J. Fuchs and T. Grøsfjeld, Tetrahedral symmetry of 6j-symbols in fusion categories, J. Pure Appl. Alg. 227 (2023) 107112\_1-45 [math.QA/2106.16186]
- [FuRS1] J. Fuchs, I. Runkel, and C. Schweigert, TFT construction of RCFT correlators I: Partition functions, Nucl. Phys. B 646 (2002) 353–497 [hep-th/0204148]
- [FuRS2] J. Fuchs, I. Runkel, and C. Schweigert, TFT construction of RCFT correlators III: Simple currents, Nucl. Phys. B 694 (2004) 277–353 [hep-th/0403157]
- [GJ] N. Gambino and A. Joyal, On operads, bimodules and analytic functors, Memoirs Amer. Math. Soc. 289 (2017) 1–110 [math.CT/1405.7270]
- [He] A. Henriques, Three-tier CFTs from Frobenius algebras, Contemp. Math. 613 (2014) 1–31 [math-ph/1304.7328]
- [JoS] A. Joyal and R. Street, Braided tensor categories, Adv. Math. 102 (1993) 20–78
- [Joz] T. Józefiak, Semisimple superalgebras, Springer Lecture Notes in Mathematics 1352 (1988) 96–113
- [Ka] G. Karpilovsky, Group Representations, Volume 2 (North Holland, Amsterdam 1996)
- [KO] A.A. Kirillov and V. Ostrik, On a q-analog of McKay correspondence and the ADE classification of  $\widehat{\mathfrak{sl}}(2)$  conformal field theories, Adv. Math. 171 (2002) 183–227 [math.QA/0101219]
- [MaZ] V. Mazorchuk and K. Zhao, Graded simple Lie algebras and graded simple representations, Manuscripta math. 156 (2018) 215–240 [math.RT/1504.05114]
- [Müg] M. Müger, From subfactors to categories and topology I. Frobenius algebras in and Morita equivalence of tensor categories, J. Pure Appl. Alg. 180 (2003) 81–157 [math.CT/0111204]
- [Mul] V. Mulevičius, Condensation inversion and Witt equivalence via generalised orbifolds, preprint math.QA/2206.02611
- [Os] V. Ostrik, Module categories over the Drinfeld double of a finite group, Int. Math. Res. Notices 2003 (2003) 1507–1520 [math.QA/0202130]

- [Pu] K.K. Putyra. A 2-category of chronological cobordisms and odd Khovanov homology, Banach Center Publ. 103 (2014) 291–355 [math.GT/1310.1895]
- [Ri] E. Riehl, Category Theory in Context (Dover Publications, New York 2016)
- [ScV] L. Schelstraete and P. Vaz, Odd Khovanov homology and higher representation theory, preprint math.GT/2311.14394
- [Se] G.J. Seal, *Tensors*, *monads and actions*, Theory and Appl. Cat. 28 (2013) 403–433 [math. CT/1205.0101]
- [Sh] K. Shimizu, Quasi-Frobenius algebras in finite tensor categories, preprint math.QA/2402. 02929
- [St] R. Street, Frobenius monads and pseudomonoids, J. Math. Phys. 45 (2004) 3930–3948
- [WW] J. Wan and W. Wang, Lectures on spin representation theory of symmetric groups, Bulletin of the Institute of Mathematics 7 (2012) 91–144 [math.RT/1110.0263]