HIGHER STRUCTURES FOR LIE H-PSEUDOALGEBRAS

APURBA DAS

ABSTRACT. Let H be a cocommutative Hopf algebra. The notion of Lie H-pseudoalgebra is a multivariable generalization of Lie conformal algebras. In this paper, we study some higher structures related to Lie H-pseudoalgebras where we increase the flexibility of the Jacobi identity. Namely, we first introduce L_{∞} H-pseudoalgebras (also called strongly homotopy Lie H-pseudoalgebras) as the homotopy analogue of Lie H-pseudoalgebras. We give several equivalent descriptions of such homotopy algebras and show that some particular classes of these homotopy algebras are closely related to the cohomology of Lie H-pseudoalgebras and crossed modules of Lie H-pseudoalgebras. Next, we introduce another higher structure, called Lie-2 H-pseudoalgebras which are the categorification of Lie H-pseudoalgebras. Finally, we show that the category of Lie-2 H-pseudoalgebras is equivalent to the category of certain $L_{\infty} H$ -pseudoalgebras.

Mathematics Subject Classification (2020). 17B56, 17B69, 18N40, 18N25. Keywords. Lie H-pseudoalgebras, L_{∞} H-pseudoalgebras, Lie-2 H-pseudoalgebras.

CONTENTS

1.	Introduction	1
2.	Lie H -pseudoalgebras and their cohomology	3
3.	Strongly homotopy Lie <i>H</i> -pseudoalgebras	5
4.	Examples of strongly homotopy Lie <i>H</i> -pseudoalgebras	11
5.	Skeletal and strict homotopy Lie <i>H</i> -pseudoalgebras	15
6.	Categorification of Lie <i>H</i> -pseudoalgebras	18
References		21

1. INTRODUCTION

1.1. Lie *H*-pseudoalgebras. The concept of Lie conformal algebras was introduced by Kac [13,14] as a tool to study vertex operator algebras. Lie conformal algebras encode the axiomatic description of the operator product expansion of chiral fields in conformal field theory. Subsequently, the notion of Lie Hpseudoalgebra was introduced by Bakalov, D' Andrea and Kac^[2] as a multivariable generalization of Lie conformal algebras. While Lie conformal algebras are defined on $\mathbf{k}[\partial]$ -modules, Lie H-pseudoalgebras are defined on left H-modules. Roughly, given a cocommutative Hopf algebra H, one first considers a pseudotensor category $\mathcal{M}^*(H)$. A Lie *H*-pseudoalgebra is simply a Lie algebra object in $\mathcal{M}^*(H)$. They have close connections to differential Lie algebras of Ritt [19] and Hamiltonian formalism in the theory of nonlinear evolution equations [11]. Given an ordinary Lie algebra \mathfrak{g} , one can construct a Lie H-pseudoalgebra structure on the tensor product $H \otimes \mathfrak{g}$. This is called the current Lie *H*-pseudoalgebra of \mathfrak{g} and it is denoted by $Cur(\mathfrak{g})$. On the other hand, each element in a Lie *H*-pseudoalgebra may be recast in terms of its Fourier coefficients, sometimes called creation and annihilation operators in the physical literature. The space of all such annihilation operators forms a Lie algebra (in general infinite-dimensional). Therefore, Lie Hpseudoalgebras are also useful to study infinite-dimensional Lie algebras. Structure theory, representations and cohomology of Lie *H*-pseudoalgebras are widely studied in [2-4,8,9,24] and in the references therein. See [6,12,18,25,26] for some other types of pseudoalgebras and relevant results.

APURBA DAS

1.2. Strongly homotopy algebras and categorifications. In mathematics, when one allows to increase the flexibility of a structure, one obtains higher structures. This can be done often in two ways, namely, by homotopification and categorification of given algebraic identity/identities. Higher structures, such as strongly homotopy algebras and categorifications of algebras are key objects in higher differential geometry, higher gauge theory and infinity-category theory [1,7,17,20]. The notion of an L_{∞} algebra (also called a strongly homotopy Lie algebra) which is the homotopification of a Lie algebra, was introduced by Lada and Stasheff [17] and further studied by Lada and Markl [16]. This homotopy algebra plays a significant role in many areas of mathematics and mathematical physics including deformation theory, quantization of Poisson manifolds and classical field theory [10,15,20]. On the other hand, the notion of a Lie-2 algebra was introduced by Baez and Crans [1] as the categorification of Lie algebras. Like Lie algebras are closely related to the set-theoretical solutions of the Yang-Baxter equation, Lie-2 algebras are related to the Zamolodchikov tetrahedron equation. Although strongly homotopy algebras and categorifications of algebras are two different generalizations of algebras, it has been observed in [1] that these higher structures are closely connected. More precisely, they showed that the category of 2-term L_{∞} algebras is equivalent to the category of Lie-2 algebras. Among other results, they considered and characterized skeletal and strict L_{∞} algebras.

1.3. Graded Lie *H*-pseudoalgebras and strongly homotopy Lie *H*-pseudoalgebras. The concept of graded Lie algebras are the Lie objects in the category of graded vector spaces. In [25] the author has considered graded Lie *H*-pseudoalgebras in his study of generalizations of some results of Lie *H*-pseudoalgebras in the graded context. Subsequently, representations and cohomology of graded Lie *H*-pseudoalgebras are studied in [23]. The author also generalized some results of Lie *H*-pseudobialgebras and Manin triples of Lie *H*-pseudoalgebras in the graded context.

In this paper, we aim to study two higher structures in the context of Lie *H*-pseudoalgebras. We first introduce the notion of an L_{∞} *H*-pseudoalgebra (also called a strongly homotopy Lie *H*-pseudoalgebra or a sh Lie *H*-pseudoalgebra) which is the homotopy invariant extension of graded Lie *H*-pseudoalgebras. To define such a notion, we first consider a graded pseudotensor category $\mathcal{M}_{gr}^*(H)$ whose objects are graded left *H*-modules and (compositions of) polylinear maps are defined through the iterative use of the comultiplication map of *H*. With this, an L_{∞} *H*-pseudoalgebra is simply an L_{∞} algebra in the graded pseudotensor category $\mathcal{M}_{gr}^*(H)$. We give some equivalent descriptions of L_{∞} *H*-pseudoalgebras. Among others, given a graded left *H*-module \mathcal{L} , we construct a graded Lie algebra whose Maurer-Cartan elements correspond to L_{∞} *H*-pseudoalgebra structures on \mathcal{L} (cf. Theorem 3.8). This construction generalizes the Maurer-Cartan characterization of Lie *H*-pseudoalgebras given in [24]. We also consider representations and cohomology of L_{∞} *H*-pseudoalgebras. Given an L_{∞} *H*pseudoalgebra \mathcal{L} , one can construct an L_{∞} algebra on the space $H^* \otimes_H \mathcal{L}$. This is called the annihilation algebra of \mathcal{L} . Next, let Γ be a finite group acting on *H* by Hopf algebra automorphisms. Let $\tilde{H} = H \# \mathbf{k}[\Gamma]$ be the corresponding smash product. We observe that any L_{∞} \tilde{H} -pseudoalgebra is equivalent to an L_{∞} *H*-pseudoalgebra equipped with an action of Γ preserving the structure maps.

We provide several examples of L_{∞} *H*-pseudoalgebras. First of all, we show that any L_{∞} conformal algebra introduced in [21] is an example of an L_{∞} *H*-pseudoalgebra. We also construct the current L_{∞} *H*-pseudoalgebra associated to an L_{∞} algebra. We define the notion of an A_{∞} *H*-pseudoalgebra and show that a suitable skew-symmetrization of an A_{∞} *H*-pseudoalgebra gives rise to an L_{∞} *H*-pseudoalgebra.

Next, we focus on those L_{∞} *H*-pseudoalgebras whose underlying graded left *H*-module is concentrated in arity 0 and 1. We call them 2-term L_{∞} *H*-pseudoalgebras. The collection of all 2-term L_{∞} *H*-pseudoalgebras and morphisms between them forms a category, denoted by **2shLie***H*. We also consider skeletal and strict L_{∞} *H*-pseudoalgebras and generalize some results of Baez and Crans [1] in the present context. More precisely, we show that skeletal L_{∞} *H*-pseudoalgebras correspond to third cocycles of Lie *H*-pseudoalgebras and strict L_{∞} *H*-pseudoalgebras correspond to crossed modules of Lie *H*-pseudoalgebras.

3

1.4. Categorification of Lie *H*-pseudoalgebras. In the final part of the paper, we consider the categorification of Lie *H*-pseudoalgebras. Namely, we introduce and study Lie-2 *H*-pseudoalgebras. When $H = \mathbf{k}$ is the base field, our notion coincides with the Lie-2 algebras considered by Baez and Crans. We show that the collection of all Lie-2 *H*-pseudoalgebras and morphisms between them forms a category, denoted by Lie2*H*. Finally, we prove that the category Lie2*H* is equivalent to the category 2shLie*H* (cf. Theorem 6.4).

The paper is organized as follows. In Section 2, we recall Lie *H*-pseudoalgebras and their cohomology theory. In Section 3, we introduce L_{∞} *H*-pseudoalgebras, their representations and cohomology. Various examples of L_{∞} *H*-pseudoalgebras are given in Section 4. We consider skeletal and strict L_{∞} *H*-pseudoalgebras in Section 5 and provide their characterizations. Finally, the notion of Lie-2 *H*-pseudoalgebras is introduced in Section 6 and proves Theorem 6.4.

2. Lie *H*-pseudoalgebras and their cohomology

In this section, we recall some necessary background on Lie H-pseudoalgebras that are required for the smooth reading of the present paper. More precisely, we recall Lie H-pseudoalgebras, their representations and cohomology theory. We will mainly follow the references [2], [8], [24].

First, recall that the notion of a pseudotensor category was introduced by Beilinson and Drinfeld [5] to study Lie algebras, representations, cohomology etc. from categorical points of view. Lie H-pseudoalgebras are Lie algebras in a suitable pseudotensor category associated with a cocommutative Hopf algebra H.

2.1. Definition. A pseudotensor category C is a class of objects together with the following data:

(i) For any finite non-empty set I, a family of objects $\{L_i\}_{i \in I}$ labelled by I and an object M, there is a vector space $\operatorname{Lin}(\{L_i\}_{i \in I}, M)$ of polylinear maps equipped with an action of the symmetric group S_I on $\operatorname{Lin}(\{L_i\}_{i \in I}, M)$ by permuting the factors,

(ii) for any finite non-empty sets I, J and a surjective map $\pi : J \to I$, families of objects $\{L_i\}_{i \in I}$, $\{N_i\}_{i \in J}$ and an object M, there exists a composition map

$$\operatorname{Lin}(\{L_i\}_{i\in I}, M) \otimes \bigotimes_{i\in I} \operatorname{Lin}(\{N_j\}_{j\in\pi^{-1}(i)}, L_i) \to \operatorname{Lin}(\{N_j\}_{j\in J}, M), \ \varphi \otimes (\bigotimes_{i\in I} \psi_i) \mapsto \varphi(\{\psi_i\}_{i\in I})$$
(1)

subject to satisfy the following axioms:

- (associativity) for another finite non-empty set K and a surjective map $\theta: K \to J$, families of objects $\{P_k\}_{k \in K}$ and given $\zeta_j \in \text{Lin}(\{P_k\}_{k \in \theta^{-1}(j)}, N_j)$, we have

$$\varphi\bigl(\{\psi_i(\{\zeta_j\}_{j\in J})\}_{i\in I}\bigr) = \bigl(\varphi(\{\psi_i\}_{i\in I})\bigr)(\{\zeta_j\}_{j\in J}),$$

- (unit) for any object M, there exists an element $1_M \in Lin(\{M\}, M)$ such that

$$1_M(\varphi) = \varphi(\{1_{L_i}\}_{i \in I}) = \varphi, \text{ for any } \varphi \in \operatorname{Lin}(\{L_i\}_{i \in I}, M),$$
(2)

- (equivariance) the composition maps (1) are equivariant with respect to the actions of the symmetric groups.

2.2. **Definition.** (i) Let C be a pseudotensor category. A *Lie algebra* in the pseudotensor category C is an object L equipped with a polylinear map $\beta \in \text{Lin}(\{L, L\}, L)$ that is skew-symmetric in the sense that $\beta = -\sigma_{12}\beta$ (where $\sigma_{12} = (1 \ 2) \in S_2$) and satisfies the Jacobi identity

$$\beta(\beta(\cdot, \cdot), \cdot) = \beta(\cdot, \beta(\cdot, \cdot)) - \sigma_{12}\beta(\cdot, \beta(\cdot, \cdot)).$$
(3)

We denote a Lie algebra as above simply by the pair (L, β) .

(ii) Let (L,β) be a Lie algebra in a pseudotensor category \mathcal{C} . A representation of the Lie algebra (L,β) is a pair (M,γ) , where M is an object of \mathcal{C} and $\gamma \in \text{Lin}(\{L,M\},M)$ is a polylinear map satisfying

$$\gamma(\beta(\cdot, \cdot), \cdot) = \gamma(\cdot, \gamma(\cdot, \cdot)) - \sigma_{12}\gamma(\cdot, \gamma(\cdot, \cdot)).$$

It follows that any Lie algebra is a representation of itself, called the adjoint representation.

Cohomology of a Lie algebra in a pseudotensor category C with coefficients in a representation is defined in [2]. However, we will not require this general cohomology theory. At the end of this section, we revise the cohomology of a Lie *H*-pseudoalgebra which is sufficient for the present paper.

2.3. **Remark.** It is important to remark that any pseudotensor category C is naturally an ordinary category with the hom sets $\operatorname{Hom}_{\mathcal{C}}(L, M) := \operatorname{Lin}(\{L\}, M)$, for any two objects L and M.

2.4. **Example.** Let C = Vect be the category of all vector spaces over the field **k**. Given any collection $\{L_i\}_{i\in I}$ of vector spaces labelled by a non-empty finite set I and another vector space M, we set $\operatorname{Lin}(\{L_i\}_{i\in I}, M) := \operatorname{Hom}(\otimes_{i\in I} L_i, M)$. The symmetric group S_I acts on $\operatorname{Lin}(\{L_i\}_{i\in I}, M)$ by permuting the factors in $\otimes_{i\in I} L_i$. This is an example of a pseudotensor category in which the composition map (1) is defined by $\varphi(\{\psi_i\}_{i\in I}) = \varphi \circ (\otimes_{i\in I} \psi_i)$.

2.5. **Example.** Let H be a cocommutative bialgebra and $\mathcal{M}^{l}(H)$ be the category of all left H-modules. Given any collection $\{L_i\}_{i \in I}$ of left H-modules and a left H-module M, we set $\operatorname{Lin}(\{L_i\}_{i \in I}, M) := \operatorname{Hom}_H(\otimes_{i \in I} L_i, M)$ on which the symmetric group S_I acts by permuting the factors in $\otimes_{i \in I} L_i$. This is also a pseudotensor category where the composition map is defined similarly to the previous example.

The most important example of a pseudotensor category for Lie *H*-pseudoalgebras is given by the following. Let *H* be a cocommutative Hopf algebra with the comultiplication map $\Delta : H \to H^{\otimes 2}$. Given any finite non-empty set *I*, we define the tensor product functor $\mathcal{M}^{l}(H)^{\otimes I} \to \mathcal{M}^{l}(H^{\otimes I})$ by the notation $\boxtimes_{i \in I}$. Thus, if $\{L_i\}_{i \in I}$ is a collection of left *H*-modules then $\boxtimes_{i \in I} L_i$ is nothing but $\bigotimes_{i \in I} L_i$ with the left $H^{\otimes I}$ -action given by $(\bigotimes_{i \in I} h_i)(\bigotimes_{i \in I} x_i) = \bigotimes_{i \in I} h_i x_i$, for $\bigotimes_{i \in I} h_i \in H^{\otimes I}$ and $\bigotimes_{i \in I} x_i \in \bigotimes_{i \in I} L_i$. We consider the pseudotensor category $\mathcal{M}^*(H)$ with the same objects as $\mathcal{M}^{l}(H)$ (i.e. left *H*-modules) but with different polylinear maps (hence different composition maps),

$$\operatorname{Lin}(\{L_i\}_{i\in I}, M) := \operatorname{Hom}_{H^{\otimes I}}(\boxtimes_{i\in I}L_i, H^{\otimes I} \otimes_H M).$$

Here the symmetric group S_I acts on the space $\operatorname{Lin}(\{L_i\}_{i\in I}, M)$ by simultaneously permuting the factors in $\boxtimes_{i\in I}L_i$ and $H^{\otimes I}$. This is well-defined as H is cocommutative. For any surjective map $\pi : J \twoheadrightarrow I$ between finite non-empty sets, consider the functor $\Delta^{(\pi)} : \mathcal{M}^l(H^{\otimes I}) \to \mathcal{M}^l(H^{\otimes J}), M \mapsto H^{\otimes J} \otimes_{H^{\otimes I}} M$, where $H^{\otimes I}$ acts on $H^{\otimes J}$ via the iterative comultiplication determined by π . With these, the composition of polylinear maps in the pseudotensor category $\mathcal{M}^*(H)$ is given by $\varphi(\{\psi_i\}_{i\in I}) := \Delta^{(\pi)}(\phi) \circ (\boxtimes_{i\in I}\psi_i)$.

2.6. **Definition.** A Lie *H*-pseudoalgebra (or a Lie pseudoalgebra if *H* is clear from the context) is a Lie algebra (L, β) in the pseudotensor category $\mathcal{M}^*(H)$.

It follows from the above definition that a Lie *H*-pseudoalgebra is a left *H*-module *L* equipped with a map $\beta \in \operatorname{Hom}_{H^{\otimes 2}}(L \boxtimes L, H^{\otimes 2} \otimes_H L)$, also called a *pseudobracket*, that is skew-symmetric and satisfies the Jacobi identity. We often write $\beta(x \otimes y)$ as [x * y], for $x, y \in L$. Note that the $H^{\otimes 2}$ -linearity of the pseudobracket β simply means that $[fx * gy] = ((f \otimes g) \otimes_H 1)[x * y]$, for all $x, y \in L$ and $f, g \in H$. Explicitly,

if
$$[x * y] = \sum_{i} (f_i \otimes g_i) \otimes_H e_i$$
 then $[fx * gy] = \sum_{i} (ff_i \otimes gg_i) \otimes_H e_i.$

Similarly, the pseudobracket is skew-symmetric implies that $[y * x] = -(\sigma_{12} \otimes_H 1)[x * y]$, for all $x, y \in L$. Explicitly,

if
$$[x * y] = \sum_{i} (f_i \otimes g_i) \otimes_H e_i$$
 then $[y * x] = -\sum_{i} (g_i \otimes f_i) \otimes_H e_i.$

Finally, to describe explicitly the Jacobi identity (3) for the pseudobracket, we need to compute all the compositions (in $\mathcal{M}^*(H)$) that appeared in the Jacobi identity. Let $x, y, z \in L$. If

$$[x * y] = \sum_{i} (f_i \otimes g_i) \otimes_H e_i \text{ and say } [e_i * z] = \sum_{j} (f_{ij} \otimes g_{ij}) \otimes_H e_{ij},$$

then [[x * y] * z] is the element of $H^{\otimes 3} \otimes_H L$ given by

$$[[x * y] * z] = \sum_{i,j} \left(f_i f_{ij(1)} \otimes g_i f_{ij(2)} \otimes g_{ij} \right) \otimes_H e_{ij}.$$

$$\tag{4}$$

Similarly, if $[y * z] = \sum_{i} (h_i \otimes l_i) \otimes_H d_i$ and say $[x * d_i] = \sum_{j} (h_{ij} \otimes l_{ij}) \otimes_H d_{ij}$, then

$$[x * [y * z]] = \sum_{i,j} \left(h_{ij} \otimes h_i l_{ij(1)} \otimes l_i l_{ij(2)} \right) \otimes_H d_{ij} \text{ as an element of } H^{\otimes 3} \otimes_H L.$$
(5)

Similarly, we can define the composition [y * [x * z]]. Note the Jacobi identity (3) for the pseudobracket is simply [[x * y] * z] = [x * [y * z]] - [y * [x * z]], where all the terms are explicitly given above. We often denote a Lie *H*-pseudoalgebra simply by *L* if the pseudobracket $\beta = [\cdot * \cdot]$ is clear from the context.

Let $(L, [\cdot * \cdot])$ and $(L', [\cdot * \cdot]')$ be two Lie *H*-pseudoalgebras. A morphism of Lie *H*-pseudoalgebras from L to L' is a *H*-linear map $\theta : L \to L'$ that satisfies $(\mathrm{id}_{H^{\otimes 2} \otimes_H} \theta)[x * y] = [\theta(x) * \theta(y)]'$, for all $x, y \in L$.

Let L be a Lie H-pseudoalgebra. A representation of L is a left H-module M equipped with a map $\gamma \in \operatorname{Hom}_{H^{\otimes 2}}(L \boxtimes M, H^{\otimes 2} \otimes_H M)$, called the *action map* written as $\gamma(x \otimes u) = x * u$, that satisfies

$$[x * y] * u = x * (y * u) - y * (x * u), \text{ for all } x, y \in L \text{ and } u \in M.$$
(6)

Here each composition in (6) can be explicitly defined similarly to the above discussions.

Let L be a Lie H-pseudoalgebra and M be a representation. For each $n \ge 0$, we define an abelian group $C^n_{\text{Lie}-H}(L,M)$ by

$$C^{n}_{\text{Lie-}H}(L,M) = \begin{cases} \mathbf{k} \otimes_{H} M & \text{if } n = 0, \\ \{\theta \in \text{Hom}_{H^{\otimes n}}(L^{\boxtimes n}, H^{\otimes n} \otimes_{H} M) \mid \theta \text{ is skew-symmetric, i.e.} \\ \sigma \theta = (-1)^{\text{sgn}(\sigma)} \theta, \forall \sigma \in S_n \} & \text{if } n \ge 1. \end{cases}$$

Then there is a map $\delta: C^n_{\text{Lie-}H}(L, M) \to C^{n+1}_{\text{Lie-}H}(L, M)$ given by

$$(\delta(1\otimes_H u))(x) = \sum_i \varepsilon(g_i) f_i \otimes_H u_i, \text{ if } x * u = \sum_i (f_i \otimes g_i) \otimes_H u_i$$

and

$$(\delta\theta)(x_1, \dots, x_{n+1}) = \sum_{i=1}^{n+1} (-1)^{i+1} (\sigma_{1 \to i} \otimes_H 1) \ x_i * \theta(x_1, \dots, \hat{x_i}, \dots, x_{n+1}) \\ + \sum_{1 \le i < j \le n+1} (-1)^{i+j} (\sigma_{1 \to i \atop 2 \to j} \otimes_H 1) \ \theta([x_i * x_j], x_1, \dots, \hat{x_i}, \dots, \hat{x_j}, \dots, x_{n+1}),$$

for $1 \otimes_H u \in C^0_{\text{Lie}-H}(L, M)$, $\theta \in C^{n \ge 1}_{\text{Lie}-H}(L, M)$ and $x, x_1, \ldots, x_{n+1} \in L$. Here $\sigma_{1 \to i}$ and $\sigma_{1 \to i}_{2 \to j}$ are respectively the permutations on $H^{\otimes n+1}$ given by

$$\sigma_{1 \to i}(h_i \otimes h_1 \otimes \cdots \otimes h_{i-1} \otimes h_{i+1} \otimes \cdots \otimes h_{n+1}) = h_1 \otimes \cdots \otimes h_{n+1},$$

$$\sigma_{1 \to i}(h_i \otimes h_j \otimes h_1 \otimes \widehat{h_i} \otimes \cdots \otimes \widehat{h_j} \otimes \cdots \otimes h_{n+1}) = h_1 \otimes \cdots \otimes h_{n+1}.$$

It has been observed in [2] that $\delta^2 = 0$. The cohomology groups of the cochain complex $\{C^{\bullet}_{\text{Lie-}H}(L, M), \delta\}$ are called the cohomology of L with coefficients in the representation M. We denote the corresponding cohomology groups by $H^{\bullet}_{\text{Lie-}H}(L, M)$.

3. Strongly homotopy Lie *H*-pseudoalgebras

In this section, we first consider a (graded) pseudotensor category $\mathcal{M}_{gr}^*(H)$ generalizing $\mathcal{M}^*(H)$ in the graded context. A graded Lie algebra in the pseudotensor category $\mathcal{M}_{gr}^*(H)$ is called a graded Lie *H*-pseudoalgebra. Motivated by this, we define the notion of an L_{∞} *H*-pseudoalgebra as an L_{∞} algebra in the pseudotensor category $\mathcal{M}_{ar}^*(H)$. Finally, we give some equivalent descriptions of L_{∞} *H*-pseudoalgebras.

Let grVect be the category of all \mathbb{Z} -graded vector spaces. For any collection $\{\mathcal{L}_i\}_{i\in I}$ of graded vector spaces and another graded vector space \mathcal{M} , we set $\operatorname{Lin}(\{\mathcal{L}_i\}_{i\in I}, \mathcal{M}) = \operatorname{Hom}(\otimes_{i\in I}\mathcal{L}_i, \mathcal{M})$ the set of all graded

linear maps from the graded vector space $\otimes_{i \in I} \mathcal{L}_i$ to the graded vector space \mathcal{M} . Note that $\operatorname{Lin}(\{\mathcal{L}_i\}_{i \in I}, \mathcal{M})$ is a graded vector space with the summands

$$\operatorname{Lin}(\{\mathcal{L}_i\}_{i\in I},\mathcal{M}) = \oplus_{n\in\mathbb{Z}}\operatorname{Lin}^n(\{\mathcal{L}_i\}_{i\in I},\mathcal{M}) = \oplus_{n\in\mathbb{Z}}\operatorname{Hom}^n(\otimes_{i\in I}\mathcal{L}_i,\mathcal{M}),$$

where $\operatorname{Hom}^{n}(\otimes_{i\in I}\mathcal{L}_{i},\mathcal{M})$ is the space of all degree *n* graded linear maps. The symmetric group S_{I} also acts on the graded space $\operatorname{Lin}({\mathcal{L}_{i}}_{i\in I},\mathcal{M})$ by $(\sigma f)(\otimes_{i\in I}x_{i}) = \epsilon(\sigma)f(\otimes_{i\in I}x_{\sigma^{-1}(i)})$, where $\epsilon(\sigma) = \epsilon(\sigma; \otimes_{i\in I}x_{i})$ is the standard *Koszul sign* that appears in the graded context. The composition map $\varphi({\psi_{i}}_{i\in I})$ is given by

$$\left(\varphi(\{\psi_i\}_{i\in I})\right)(\otimes_{j\in J}x_j) = \left(\varphi\circ(\otimes_{i\in I}\psi_i)\right)(\otimes_{j\in J}x_j) = \epsilon(\{\psi_i\}_{i\in I}; \otimes_{j\in J}x_j)\varphi\left(\otimes_{i\in I}\psi_i(\otimes_{j\in \pi^{-1}(i)}x_j)\right).$$
(7)

Here $\epsilon(\{\psi_i\}_{i \in I}; \otimes_{j \in J} x_j)$ is also the Koszul sign. The above composition maps turn out to be associative and equivariant with respect to the actions of the symmetric groups. Further, for any graded vector space \mathcal{L} , we have the identity map $\mathrm{id}_{\mathcal{L}} \in \mathrm{Lin}^0(\{\mathcal{L}\}, \mathcal{L})$ that satisfies (2).

Motivated by the above discussions, we introduce the following.

3.1. **Definition.** A graded pseudotensor category C_{gr} (over \mathbb{Z}) is a class of \mathbb{Z} -graded objects together with graded vector spaces

$$\operatorname{Lin}(\{\mathcal{L}_i\}_{i\in I},\mathcal{M})=\oplus_{n\in\mathbb{Z}}\operatorname{Lin}^n(\{\mathcal{L}_i\}_{i\in I},\mathcal{M}),$$

equipped with actions of the symmetric groups S_I among them by permuting the factors and there are composition maps $\varphi(\{\psi_i\}_{i \in I})$ subject to satisfy the associativity, unit condition and the equivariance condition.

Of course the category grVect is a graded pseudotensor category. More generally, let H be a cocommutative Hopf algebra and $\mathcal{M}_{gr}^{l}(H)$ be the category of all graded left H-modules. For any collection $\{\mathcal{L}_i\}$ of graded left H-modules and another graded left H-module \mathcal{M} , we set

$$\operatorname{Lin}(\{\mathcal{L}_i\}_{i\in I},\mathcal{M}) = \bigoplus_{n\in\mathbb{Z}}\operatorname{Lin}^n(\{\mathcal{L}_i\}_{i\in I},\mathcal{M}) = \bigoplus_{n\in\mathbb{Z}}\operatorname{Hom}_H^n(\otimes_{i\in I}\mathcal{L}_i,\mathcal{M}),$$

where $\operatorname{Hom}_{H}^{n}(\bigotimes_{i\in I}\mathcal{L}_{i},\mathcal{M})$ is the space of all graded *H*-linear maps of degree *n* from the graded space $\bigotimes_{i\in I}\mathcal{L}_{i}$ to \mathcal{M} . The symmetric group action and the compositions can be defined as the motivated example grVect. With these, $\mathcal{M}_{ar}^{l}(H)$ turns out to be a graded pseudotensor category.

Next, we consider a graded pseudotensor category $\mathcal{M}_{gr}^*(H)$ as follows. Its objects are the same as the objects of $\mathcal{M}_{gr}^l(H)$ (i.e. graded left *H*-modules) and polylinear maps are defined by

$$\operatorname{Lin}(\{\mathcal{L}_i\}_{i\in I},\mathcal{M})=\oplus_{n\in\mathbb{Z}}\operatorname{Lin}^n(\{\mathcal{L}_i\}_{i\in I},\mathcal{M}):=\oplus_{n\in\mathbb{Z}}\operatorname{Hom}^n_{H^{\otimes I}}(\boxtimes_{i\in I}\mathcal{L}_i,H^{\otimes I}\otimes_H\mathcal{M}).$$

The symmetric group actions and composition maps are graded analogues of the same defined for $\mathcal{M}^*(H)$.

3.2. **Definition.** Let \mathcal{C}_{gr} be a graded pseudotensor category (over \mathbb{Z}).

(i) A graded Lie algebra in C_{gr} is a pair (\mathcal{L}, β) consisting of an object \mathcal{L} with a degree 0 polylinear map $\beta \in \operatorname{Lin}^{0}({\mathcal{L}, \mathcal{L}}, \mathcal{L})$ that is graded skew-symmetric (i.e. $\beta(\cdot, \cdot) = -\epsilon(\sigma_{12})(\sigma_{12} \otimes_{H} 1)\sigma_{12}\beta(\cdot, \cdot))$ and satisfies the graded Jacobi identity

$$\beta(\beta(\cdot, \cdot), \cdot) = \beta(\cdot, \beta(\cdot, \cdot)) - \epsilon(\sigma_{12})\sigma_{12}\beta(\cdot, \beta(\cdot, \cdot)).$$

(ii) A differential graded Lie algebra in C_{gr} is a graded Lie algebra (\mathcal{L}, β) equipped with a degree -1 polylinear map $d \in \operatorname{Lin}^{-1}(\{\mathcal{L}\}, \mathcal{L})$ satisfying

$$d^2 = 0$$
 and $d\beta(\cdot, \cdot) = \beta(d \cdot, \cdot) + \epsilon(\beta; \cdot, \cdot)\beta(\cdot, d \cdot).$

3.3. **Definition.** A graded Lie *H*-pseudoalgebra is a graded Lie algebra in the graded pseudotensor category $\mathcal{M}_{gr}^{*}(\mathcal{H})$. Similarly, a differential graded Lie *H*-pseudoalgebra is a differential graded Lie algebra in the graded pseudotensor category $\mathcal{M}_{gr}^{*}(\mathcal{H})$.

Thus, a graded Lie *H*-pseudoalgebra is a pair (\mathcal{L}, β) consisting of a graded left *H*-module $\mathcal{L} = \bigoplus_{n \in \mathbb{Z}} \mathcal{L}^n$ with a degree 0 graded pseudobracket $\beta \in \operatorname{Hom}_{H^{\otimes 2}}^0(\mathcal{L} \boxtimes \mathcal{L}, H^{\otimes 2} \otimes_H \mathcal{L})$, written as $\beta(x \otimes y) = [x * y]$, satisfying

- (graded skew-symmetry) $[y * x] = -(-1)^{|x||y|} (\sigma_{12} \otimes_H 1) [x * y],$
- (graded Jacobi identity) for any homogeneous elements $x,y,z\in\mathcal{L},$

$$[[x * y] * z] = [x * [y * z]] - (-1)^{|x||y|} [y * [x * z]].$$

Further, it is a differential graded Lie *H*-pseudoalgebra if there exists a *H*-linear degree $-1 \mod d : \mathcal{L} \to H \otimes_H \mathcal{L}$ satisfies $d^2 = 0$ and $d[x * y] = [dx * y] + (-1)^{|x|}[x * dy]$, for all $x, y \in \mathcal{L}$.

In the following, we will introduce L_{∞} *H*-pseudoalgebras. Before we give the explicit description, we motivate the reader by giving the following.

3.4. **Definition.** Let C_{gr} be a graded pseudotensor category (over \mathbb{Z}). An L_{∞} algebra in C_{gr} is a pair $(\mathcal{L}, \{\beta_k\}_{k\geq 1})$ consisting of an object $\mathcal{L} = \bigoplus_{n \in \mathbb{Z}} \mathcal{L}^n$ together with a collection

$$\left\{\beta_k \in \operatorname{Lin}^{k-2}(\{\underbrace{\mathcal{L},\ldots,\mathcal{L}}_{k \text{ many}}\},\mathcal{L})\right\}_{k \ge 1}$$

of polylinear maps with specific degrees (namely, $\deg(\beta_k) = k - 2$, for $k \ge 1$) that satisfy the following axioms:

- ((graded) skew-symmetry) each β_k is skew-symmetric in the sense that $\beta_k = (-1)^{\sigma} (\sigma \beta_k)$, for all $\sigma \in S_k$,

- (higher Jacobi identities) for each $N \in \mathbb{N}$ and homogeneous elements $x_1, \ldots, x_N \in \mathcal{L}$, the following identity holds:

$$\sum_{k+l=N+1} \sum_{\sigma \in Sh(l,k-1)} (-1)^{\sigma} \epsilon(\sigma) (-1)^{l(k-1)} \beta_k \left(\beta_l(x_{\sigma(1)}, \dots, x_{\sigma(l)}), x_{\sigma(l+1)}, \dots, x_{\sigma(N)} \right) = 0.$$
(8)

Let $C_{gr} = \text{grVect}$ be the graded pseudotensor category of all graded vector spaces. It follows from the above definition that an L_{∞} algebra in grVect is a pair $(\mathcal{L}, \{\beta_k\}_{k\geq 1})$ consisting of a graded vector space $\mathcal{L} = \bigoplus_{n \in \mathbb{Z}} \mathcal{L}^n$ with a collection $\{\beta_k \in \text{Hom}^{k-2}(\mathcal{L}^{\otimes k}, \mathcal{L})\}_{k\geq 1}$ of graded skew-symmetric maps satisfying the higher Jacobi identities (8). This is a classical L_{∞} algebra considered in [16,17]. Thus, an L_{∞} algebra in a graded pseudotensor category generalizes classical L_{∞} algebras. Motivated by this observation, we now define the following.

3.5. **Definition.** An L_{∞} *H*-pseudoalgebra (or simply a strongly homotopy Lie *H*-pseudoalgebra) is an L_{∞} algebra in the graded pseudotensor category $\mathcal{M}_{gr}^{*}(H)$. More precisely, an L_{∞} *H*-pseudoalgebra is a pair $(\mathcal{L}, \{\beta_k\}_{k\geq 1})$ consisting of a graded left *H*-module $\mathcal{L} = \bigoplus_{n\in\mathbb{Z}}\mathcal{L}^n$ equipped with a collection $\{\beta_k \in \text{Hom}_{H^{\otimes k}}(\mathcal{L}^{\boxtimes k}, H^{\otimes k} \otimes_H \mathcal{L})\}_{k\geq 1}$ of polylinear maps with $\deg(\beta_k) = k-2$, for $k \geq 1$, subject to the following conditions:

- (graded skew-symmetry) each β_k is graded skew-symmetric in the sense that

1

 $\beta_k(x_1,\ldots,x_k) = (-1)^{\sigma} \epsilon(\sigma)(\sigma \otimes_H 1) \beta_k(x_{\sigma^{-1}(1)},\ldots,x_{\sigma^{-1}(k)}), \text{ for all } \sigma \in S_k,$

- (higher Jacobi identities) For each $N \in \mathbb{N}$ and homogeneous elements $x_1, \ldots, x_N \in \mathcal{L}$, the identity (8) holds.

3.6. **Remark.** (i) Any graded Lie *H*-pseudoalgebra (\mathcal{L}, β) can be realized as an L_{∞} *H*-pseudoalgebra $(\mathcal{L}, \{\beta_k\}_{k\geq 1})$ with $\beta_2 = \beta$ and $\beta_k = 0$ for all $k \neq 2$. Similarly, a differential graded Lie *H*-pseudoalgebra (\mathcal{L}, β, d) is an L_{∞} *H*-pseudoalgebra with $\beta_1 = d$, $\beta_2 = \beta$ and $\beta_k = 0$ for all $k \neq 1, 2$.

(ii) As expected, any classical L_{∞} algebra $(\mathcal{L}, \{\beta_k\}_{k\geq 1})$ is an L_{∞} *H*-pseudoalgebra for $H = \mathbf{k}$.

There is another equivalent description of an L_{∞} *H*-pseudoalgebra that renders some other simple descriptions. Let $(\mathcal{L}, \{\beta_k\}_{k\geq 1})$ be an L_{∞} *H*-pseudoalgebra. Consider the graded left *H*-module $\mathcal{L}[-1]$ given by $\mathcal{L}[-1] = \bigoplus_{n \in \mathbb{Z}} (\mathcal{L}[-1])^n = \bigoplus_{n \in \mathbb{Z}} \mathcal{L}^{n-1}$. For any $k \geq 1$, we define

$$\eta_k \in \operatorname{Hom}_{H^{\otimes k}}\left(\left(\mathcal{L}[-1]\right)^{\boxtimes k}, H^{\otimes k} \otimes_H \mathcal{L}[-1]\right) \text{ by } \eta_k = (-1)^{\frac{k(k-1)}{2}} s \circ \beta_k \circ (s^{-1})^{\otimes k}, \tag{9}$$

where $s : \mathcal{L} \to \mathcal{L}[-1]$ is the degree +1 map that identifies \mathcal{L} and $\mathcal{L}[-1]$, and $s^{-1} : \mathcal{L}[-1] \to \mathcal{L}$ is the inverse of s. Since deg $(\beta_k) = k - 2$, it follows that deg $(\eta_k) = -1$. The graded skew-symmetry of β_k is equivalent to the fact that the map η_k is graded symmetric. Furthermore, the higher Jacobi identities (8) are equivalent to the identities

$$\sum_{k+l=N+1} \sum_{\sigma \in \operatorname{Sh}(l,k-1)} \epsilon(\sigma) \eta_k \left(\eta_l(u_{\sigma(1)}, \dots, u_{\sigma(l)}), u_{\sigma(l+1)}, \dots, u_{\sigma(N)} \right) = 0,$$
(10)

for all $N \in \mathbb{N}$ and homogeneous elements $u_1, \ldots, u_N \in \mathcal{L}[-1]$. As a summary, we obtain the following.

3.7. **Proposition.** Let $\mathcal{L} = \bigoplus_{n \in \mathbb{Z}} \mathcal{L}^n$ be a graded left *H*-module. An L_{∞} *H*-pseudoalgebra structure on \mathcal{L} is equivalent to a collection of degree -1 polylinear maps $\{\eta_k \in \operatorname{Hom}_{H^{\otimes k}}^{-1}((\mathcal{L}[-1])^{\otimes k}, H^{\otimes k} \otimes_H \mathcal{L}[-1])\}_{k \geq 1}$ that are graded symmetric and satisfying the shifted higher Jacobi identities (10).

Next, we will consider another characterization of L_{∞} *H*-pseudoalgebras in terms of Maurer-Cartan elements in a suitable graded Lie algebra. Let $\mathcal{W} = \bigoplus_{n \in \mathbb{Z}} \mathcal{W}^n$ be a graded left *H*-module. For any $p \in \mathbb{Z}$ and $k \geq 1$, let $\operatorname{Hom}_{H^{\otimes k}}^p(\mathcal{W}^{\boxtimes k}, H^{\otimes k} \otimes_H \mathcal{W})$ be the set of all graded polylinear maps in $\operatorname{Hom}_{H^{\otimes k}}^p(\mathcal{W}^{\boxtimes k}, H^{\otimes k} \otimes_H \mathcal{W})$ of degree *p*. We also let sym $\operatorname{Hom}_{H^{\otimes k}}^p(\mathcal{W}^{\boxtimes k}, H^{\otimes k} \otimes_H \mathcal{W})$ be the set of all maps in $\operatorname{Hom}_{H^{\otimes k}}^p(\mathcal{W}^{\boxtimes k}, H^{\otimes k} \otimes_H \mathcal{W})$ that are graded symmetric. We define

$$\operatorname{Hom}_{\operatorname{poly}}^{p}(\mathcal{W},\mathcal{W}) := \oplus_{k\geq 1} \operatorname{Hom}_{H^{\otimes k}}^{p}(\mathcal{W}^{\boxtimes k}, H^{\otimes k} \otimes_{H} \mathcal{W}),$$

symHom_{\operatorname{poly}}^{p}(\mathcal{W},\mathcal{W}) := \oplus_{k\geq 1} \operatorname{symHom}_{H^{\otimes k}}^{p}(\mathcal{W}^{\boxtimes k}, H^{\otimes k} \otimes_{H} \mathcal{W}).

Thus, an element $\eta \in \operatorname{Hom}_{\operatorname{poly}}^{p}(\mathcal{W}, \mathcal{W})$ (resp. $\operatorname{Hom}_{\operatorname{poly}}^{p}(\mathcal{W}, \mathcal{W})$) is a formal sum $\eta = \sum_{k\geq 1} \eta_{k}$, where $\eta_{k} \in \operatorname{Hom}_{H^{\otimes k}}^{p}(\mathcal{W}^{\boxtimes k}, H^{\otimes k} \otimes_{H} \mathcal{W})$ (resp. sym $\operatorname{Hom}_{H^{\otimes k}}^{p}(\mathcal{W}^{\boxtimes k}, H^{\otimes k} \otimes_{H} \mathcal{W})$) for $k \geq 1$. It has been implicitly observed by Wu [24] that the graded space $\bigoplus_{p\in\mathbb{Z}}$ sym $\operatorname{Hom}_{\operatorname{poly}}^{p}(\mathcal{W}, \mathcal{W})$ carries a graded Lie bracket given by

$$\left[\left[\sum_{k\geq 1}\eta_k,\sum_{l\geq 1}\zeta_l\right]\right] = \sum_{N\geq 1}\sum_{k+l=N+1}\left[\left[\eta_k,\zeta_l\right]\right] := \sum_{N\geq 1}\sum_{k+l=N+1}\left(\eta_k\diamond\zeta_l - (-1)^{pq}\zeta_l\diamond\eta_k\right),\tag{11}$$

where

$$(\eta_k \diamond \zeta_l)(w_1, \dots, w_N) := \sum_{\sigma \in \mathrm{Sh}(l, k-1)} \epsilon(\sigma) \eta_k \big(\zeta_l(w_{\sigma(1)}, \dots, w_{\sigma(l)}), w_{\sigma(l+1)}, \dots, w_{\sigma(N)} \big), \tag{12}$$

for $\eta = \sum_{k \ge 1} \eta_k \in \text{symHom}_{\text{poly}}^p(\mathcal{W}, \mathcal{W}), \zeta = \sum_{l \ge 1} \zeta_l \in \text{symHom}_{\text{poly}}^q(\mathcal{W}, \mathcal{W}) \text{ and } w_1, \dots, w_N \in \mathcal{W}.$

3.8. **Theorem.** Let $\mathcal{L} = \bigoplus_{n \in \mathbb{Z}} \mathcal{L}^n$ be a graded left *H*-module. Then there is a one-to-one correspondence between L_{∞} *H*-pseudoalgebra structures on \mathcal{L} and Maurer-Cartan elements in the graded Lie algebra $(\bigoplus_{p \ge 1} \text{symHom}_{poly}^p(\mathcal{W}, \mathcal{W}), [[,]])$, where $\mathcal{W} = \mathcal{L}[-1]$.

Proof. Let $\{\beta_k \in \operatorname{Hom}_{H^{\otimes k}}^{k-2}(\mathcal{L}^{\boxtimes k}, H^{\otimes k} \otimes_H \mathcal{L})\}_{k\geq 1}$ be a collection of graded skew-symmetric polylinear maps of specific degrees. Then we have seen in (9) that the above collection is equivalent to having a collection $\{\eta_k \in \operatorname{Hom}_{H^{\otimes k}}^{-1}(\mathcal{W}^{\boxtimes k}, H^{\otimes k} \otimes_H \mathcal{W})\}_{k\geq 1}$ of graded symmetric polylinear maps. If we consider the element $\eta = \sum_{k>1} \eta_k \in \operatorname{symHom}_{\operatorname{poly}}^{-1}(\mathcal{W}, \mathcal{W})$, then we have from (11) that

$$\llbracket \eta, \eta \rrbracket = \sum_{N \ge 1} \sum_{k+l=N+1} (\eta_k \diamond \eta_l + \eta_l \diamond \eta_k) = 2 \sum_{N \ge 1} \sum_{k+l=N+1} (\eta_k \diamond \eta_l),$$

where $(\eta_k \diamond \eta_l)$ is given in (12). This show that $\eta = \sum_{k\geq 1} \eta_k$ is a Maurer-Cartan element in the graded Lie algebra $(\bigoplus_{p\geq 1} \text{symHom}_{\text{poly}}^p(\mathcal{W}, \mathcal{W}), [\![,]\!])$ if and only if the collection $\{\eta_k\}_{k\geq 1}$ satisfy the identities (10). Hence the result follows after using Proposition 3.7.

3.9. **Remark.** The above theorem generalizes the Maurer-Cartan characterization of Lie *H*-pseudoalgebras considered by Wu [24].

As a consequence of all the above discussions, we get the following equivalent descriptions of an L_{∞} *H*-pseudoalgebra:

(i) $(\mathcal{L}, \{\beta_k\}_{k\geq 1})$ is an L_{∞} *H*-pseudoalgebra,

(ii) there exists a collection of degree -1 polylinear maps $\{\eta_k \in \operatorname{Hom}_{H^{\otimes k}}((\mathcal{L}[-1])^{\otimes k}, H^{\otimes k} \otimes_H \mathcal{L}[-1])\}_{k \ge 1}$ that are graded symmetric and satisfy the shifted higher Jacobi identities (10),

- (iii) there exists a degree -1 square-zero coderivation η on the coassociative coalgebra $(T_H^c(\mathcal{L}[-1]), \Delta),$
- (iv) there exists a Maurer-Cartan element in the graded Lie algebra $(\bigoplus_{p\geq 1} \operatorname{Hom}_{polv}^{p}(\mathcal{L}[-1], \mathcal{L}[-1]), [\![,]\!]).$

3.1. Representations and cohomology of strongly homotopy Lie *H*-pseudoalgebras. In this subsection, we first consider representations of an L_{∞} *H*-pseudoalgebra. Then using the Maurer-Cartan characterization of L_{∞} *H*-pseudoalgebras, we define their cohomology (with coefficients in representations).

Let $(\mathcal{L} = \bigoplus_{n \in \mathbb{Z}} \mathcal{L}^n, \{\beta_k\}_{k \ge 1})$ be an L_{∞} *H*-pseudoalgebra. A representation of the L_{∞} *H*-pseudoalgebra is given by a pair $(\mathcal{M} = \bigoplus_{n \in \mathbb{Z}} \mathcal{M}^n, \{\gamma_k\}_{k \ge 1})$ consisting of a graded left *H*-module $\mathcal{M} = \bigoplus_{n \in \mathbb{Z}} \mathcal{M}^n$ with a collection of polylinear maps $\{\gamma_k \in \operatorname{Hom}_{H^{\otimes k}}(\mathcal{L}^{\boxtimes k-1} \boxtimes \mathcal{M}, H^{\otimes k} \otimes_H \mathcal{M})\}_{k \ge 1}$ with $\operatorname{deg}(\gamma_k) = k - 2$ for $k \ge 1$, that subject to satisfy the following conditions:

(i) each γ_k is skew-symmetric on simultaneous permutations of the factors of $\mathcal{L}^{\boxtimes k-1}$ and first k-1 factors of $H^{\otimes k}$,

(ii) for each $N \in \mathbb{N}$, the identity (8) holds when $x_1, \ldots, x_{N-1} \in \mathcal{L}$ and $x_N \in \mathcal{M}$ (with the convension that $\beta_k = \gamma_k$ when $\sigma(N) = N$ and $\beta_l = \gamma_l$ when $\sigma(l) = N$).

It follows from the above definition that an L_{∞} *H*-pseudoalgebra $(\mathcal{L}, \{\beta_k\}_{k\geq 1})$ is a representation of itself. This is called the *adjoint representation*.

3.10. **Remark.** Any representation of a Lie *H*-pseudoalgebra (see Equation (6)) can be seen as a representation of the corresponding L_{∞} *H*-pseudoalgebra concentrated in degree 0.

Given a Lie H-pseudoalgebra and a representation of it, one can construct the semidirect product Lie H-pseudoalgebra [2]. This can be generalized in the strongly homotopy context.

3.11. **Proposition.** Let $(\mathcal{L}, \{\beta_k\}_{k\geq 1})$ be an L_{∞} H-pseudoalgebra and $(\mathcal{M}, \{\gamma_k\}_{k\geq 1})$ be a representation of it. Then $(\mathcal{L} \oplus \mathcal{M}, \{\overline{\beta}_k\}_{k\geq 1})$ is an L_{∞} H-pseudoalgebra, where the structure maps $\{\overline{\beta}_k\}_{k\geq 1}$ are given by

$$(\overline{\beta}_k)((x_1, u_1), \dots, (x_k, u_k)) := (\beta_k(x_1, \dots, x_k), \sum_{i=1}^k (-1)^{k-i} \gamma_k(x_1, \dots, \widehat{x_i}, \dots, x_k, u_i)).$$
(13)

Let $(\mathcal{L}, \{\beta_k\}_{k\geq 1})$ be an L_{∞} *H*-pseudoalgebra and \mathcal{M} be a graded left *H*-module equipped with a collection of polylinear maps $\{\gamma_k \in \operatorname{Hom}_{H^{\otimes k}}^{k-2}(\mathcal{L}^{\boxtimes k-1} \boxtimes \mathcal{M}, H^{\otimes k} \otimes_H \mathcal{M})\}_{k\geq 1}$ such that each γ_k is skew-symmetric on simultaneous permutations of the factors of $\mathcal{L}^{\boxtimes k-1}$ and first k-1 factors of $H^{\otimes k}$. Then on the direct sum graded left *H*-module $\mathcal{L} \oplus \mathcal{M}$, we define a collection $\{\overline{\beta}_k\}_{k\geq 1}$ of skew-symmetric polylinear maps by (13). Then it can be easily checked that $(\mathcal{L} \oplus \mathcal{M}, \{\overline{\beta}_k\}_{k\geq 1})$ is an L_{∞} *H*-pseudoalgebra if and only if $(\mathcal{M}, \{\gamma_k\}_{k\geq 1})$ is a representation of the L_{∞} *H*-pseudoalgebra $(\mathcal{L}, \{\beta_k\}_{k\geq 1})$.

Let $(\mathcal{L}, \{\beta_k\}_{k\geq 1})$ be an L_{∞} *H*-pseudoalgebra and $(\mathcal{M}, \{\gamma_k\}_{k\geq 1})$ be a representation of it. Consider the semidirect product L_{∞} *H*-pseudoalgebra $(\mathcal{L} \oplus \mathcal{M}, \{\overline{\beta}_k\}_{k\geq 1})$ given in Proposition 3.11. This L_{∞} *H*-pseudoalgebra gives rise to the Maurer-Cartan element $\overline{\eta} = \sum_{k\geq 1} \overline{\eta}_k$ in the graded Lie algebra

$$(\bigoplus_{p\in\mathbb{Z}}\operatorname{Hom}_{\operatorname{poly}}^{p}((\mathcal{L}\oplus\mathcal{M})[-1],(\mathcal{L}\oplus\mathcal{M})[-1]),[[,]]))$$

where $\overline{\eta}_k = (-1)^{\frac{k(k-1)}{2}} s \circ \overline{\beta}_k \circ (s^{-1})^{\otimes k}$ for $k \geq 1$. This yields a cochain complex $\{C^{\bullet}_{\mathrm{shLie-}H}(\mathcal{L} \oplus \mathcal{M}, \mathcal{L} \oplus \mathcal{M}), \delta_{\overline{\eta}}\}$ induced by the Maurer-Cartan element $\overline{\eta}$. More precisely,

$$C^{n}_{\mathrm{shLie-}H}(\mathcal{L} \oplus \mathcal{M}, \mathcal{L} \oplus \mathcal{M}) := \mathrm{Hom}_{\mathrm{poly}}^{-(n-1)} \big((\mathcal{L} \oplus \mathcal{M})[-1], (\mathcal{L} \oplus \mathcal{M})[-1] \big) \text{ and} \\ \delta_{\overline{\eta}}(\eta) := (-1)^{n-1} [\![\overline{\eta}, \eta]\!], \text{ for } \eta \in C^{n}_{\mathrm{shLie-}H}.$$

For each $n \in \mathbb{Z}$, we define a subspace $C^n_{\mathrm{shLie}-H}(\mathcal{L}, \mathcal{M}) \subset C^n_{\mathrm{shLie}-H}(\mathcal{L} \oplus \mathcal{M}, \mathcal{L} \oplus \mathcal{M})$ by

$$C^{n}_{\mathrm{shLie}\text{-}H}(\mathcal{L},\mathcal{M}) := \mathrm{Hom}_{\mathrm{poly}}^{-(n-1)} \big(\mathcal{L}[-1], \mathcal{M}[-1] \big) = \bigoplus_{k \ge 1} \mathrm{Hom}_{H^{\otimes k}}^{-(n-1)} \big((\mathcal{L}[-1])^{\otimes k}, H^{\otimes k} \otimes_{H} \mathcal{M}[-1] \big).$$

Then it can be easily seen that $\delta_{\overline{\eta}}(C^n_{\mathrm{shLie}-H}(\mathcal{L},\mathcal{M})) \subset C^{n+1}_{\mathrm{shLie}-H}(\mathcal{L},\mathcal{M})$. In other words, we get a subcomplex $\{C^{\bullet}_{\mathrm{shLie}-H}(\mathcal{L},\mathcal{M}), \delta_{\overline{\eta}}\}$ of the cochain complex $\{C^{\bullet}_{\mathrm{shLie}-H}(\mathcal{L}\oplus\mathcal{M},\mathcal{L}\oplus\mathcal{M}), \delta_{\overline{\eta}}\}$. The cohomology groups of this subcomplex are called the cohomology groups of the L_{∞} *H*-pseudoalgebra \mathcal{L} with coefficients in the representation \mathcal{M} .

3.2. Γ -actions. Let H be a cocommutative Hopf algebra and Γ be a group (not necessarily finite) acting on H by Hopf algebra automorphisms. We denote the action of Γ on H by $g \cdot f$, for $g \in \Gamma$ and $f \in H$. That is, $g \cdot f = gfg^{-1}$. Consider the smash product $\tilde{H} = H \# \mathbf{k}[\Gamma]$. As an associative algebra, it is the semi-direct product of H and $\mathbf{k}[\Gamma]$, and as a coassociative coalgebra it is the tensor product of coalgebras. Note that, for any finite nonempty set I, there is a map $\delta_I : \tilde{H}^{\otimes I} \to H^{\otimes I} \otimes_H \tilde{H}$ given by

 $\delta_I(\otimes_{i\in I} f_i g_i) = (\otimes_{i\in I} f_i) \otimes g$ if all g_i 's are equal to some g

and zero otherwise. Let $\mathcal{L} = \bigoplus_{n \in \mathbb{Z}} \mathcal{L}_n$ be a graded left \widetilde{H} -module (thus, an object in $\mathcal{M}_{gr}^*(\widetilde{H})$). This is equivalent to the fact that \mathcal{L} is a graded left H-module equipped with an action of Γ on it that is compatible with the action of H. More precisely, the correspondence is given by

$$(g \cdot f)(x) = g(f(g^{-1}x)), \text{ for } f \in H, g \in \Gamma \text{ and } x \in \mathcal{L}$$

Let $\mathcal{M}^*_{gr,\Gamma}(H)$ be the graded subcategory of $\mathcal{M}^*_{gr}(H)$ whose objects are graded left \tilde{H} -modules and with those graded polylinear maps of $\mathcal{M}^*_{gr}(H)$ that commute with the action of Γ in the sense that

$$\varphi(\{g\psi_i\}_{i\in I}) = g \cdot \varphi(\{\psi_i\}_{i\in I}), \text{ for all } g \in \Gamma.$$

Then $\mathcal{M}^*_{gr,\Gamma}(H)$ is also a graded pseudotensor category. Moreover, there is a graded pseudotensor functor $\delta : \mathcal{M}^*_{gr}(\widetilde{H}) \to \mathcal{M}^*_{gr,\Gamma}(H)$ defined as follows. For an object \mathcal{L} in $\mathcal{M}^*_{gr}(\widetilde{H})$ (i.e. a graded left \widetilde{H} -module), we set $\delta(\mathcal{L}) = \mathcal{L}$. For a polylinear map $\varphi \in \operatorname{Lin}(\{\mathcal{L}_i\}_{i \in I}, \mathcal{M})$ in $\mathcal{M}^*_{gr}(\widetilde{H})$, we set $\delta(\varphi)$ to be the composition

$$\boxtimes_{i\in I}\mathcal{L}_i\xrightarrow{\varphi}\widetilde{H}^{\otimes I}\otimes_{\widetilde{H}}\mathcal{M}\xrightarrow{\delta_I\otimes_{\widetilde{H}}\mathrm{id}}(H^{\otimes I}\otimes_H\widetilde{H})\otimes_{\widetilde{H}}\mathcal{M}\xrightarrow{\cong}H^{\otimes I}\otimes_H\mathcal{M}$$

Since the maps δ_I are compatible with the actions of the symmetric groups and with the comultiplication of \tilde{H} , it follows that δ is compatible with the actions of the symmetric groups and with compositions of polylinear maps.

The proof of the following result is similar to [2, Theorem 5.1].

3.12. **Proposition.** For a finite group G, the functor $\delta : \mathcal{M}_{gr}^*(\widetilde{H}) \to \mathcal{M}_{gr,\Gamma}^*(H)$ is an equivalence of graded pseudotensor categories.

As a corollary of the above result, we get the following.

3.13. Theorem. Let H be a cocommutative Hopf algebra and Γ be a finite group. Let $\tilde{H} = H \# \mathbf{k}[\Gamma]$. Then an L_{∞} \tilde{H} -pseudoalgebra is same as an L_{∞} H-pseudoalgebra $(\mathcal{L}, \{\beta_k\}_{k\geq 1})$ equipped with an action of Γ on the graded left H-module \mathcal{L} that satisfies

$$\beta_k(gx_1,\ldots,gx_k) = g \cdot \beta_k(x_1,\ldots,x_k),\tag{14}$$

for $k \geq 1$; $x_1, \ldots, x_k \in \mathcal{L}$ and $g \in \Gamma$.

Let $\{\beta_k\}_{k\geq 1}$ be an L_{∞} *H*-pseudoalgebra structure on \mathcal{L} that satisfies (14). Then the L_{∞} \widetilde{H} -pseudoalgebra structure $\{\widetilde{\beta}_k\}_{k\geq 1}$ on \mathcal{L} is given by

$$\widetilde{\beta}_k(x_1,\ldots,x_k) = \sum_{g \in \Gamma} ((g^{-1} \otimes 1 \otimes \cdots \otimes 1) \otimes_{\widetilde{H}} 1) \beta_k(gx_1,\ldots,x_k), \text{ for } k \ge 1.$$

3.3. Annihilation algebras. Let H be a cocommutative Hopf algebra with the multiplication Δ and count ε . Let Y be an H-bimodule which is also a commutative H-differential algebra for both the left and right actions of H.

For any graded left *H*-module $\mathcal{L} = \bigoplus_{n \in \mathbb{Z}} \mathcal{L}^n$, we define $\mathcal{A}_Y \mathcal{L} := Y \otimes_H \mathcal{L}$. Then there is a left *H*-action on $\mathcal{A}_Y \mathcal{L}$ given by $h(y \otimes_H x) = hy \otimes_H x$, for $h \in H$, $y \in Y$ and $x \in \mathcal{L}$. Additionally, if $(\mathcal{L}, \{\beta_k\}_{k > 1})$ is an L_{∞} *H*-pseudoalgebra, then we can define a collection of products $\{\widetilde{\beta}_k : (\mathcal{A}_Y \mathcal{L})^{\otimes k} \to \mathcal{A}_Y \mathcal{L}\}_{k \geq 1}$ by

$$\widetilde{\beta}_k (y_1 \otimes_H x_1, \dots, y_k \otimes_H x_k) = \sum_i (y_1 f_{1i}) \cdots (y_k f_{ki}) \otimes_H e_i$$

if $\beta_k (x_1, \dots, x_k) = \sum_i (f_{1i} \otimes \dots \otimes f_{ki}) \otimes_H e_i.$

It is easy to see that these maps are well-defined. Moreover, we have the following result.

3.14. **Proposition.** Let $(\mathcal{L}, \{\beta_k\}_{k\geq 1})$ be an L_{∞} *H*-pseudoalgebra. Then the pair $(\mathcal{A}_Y \mathcal{L}, \{\widetilde{\beta}_k\}_{k\geq 1})$ is an L_{∞} algebra whose structure maps additionally satisfy

$$h\big(\widetilde{\beta}_k(y_1\otimes_H x_1,\ldots,y_k\otimes_H x_k)\big)=\widetilde{\beta}_k\big(h_{(1)}(y_1\otimes_H x_1),\ldots,h_{(k)}(y_k\otimes_H x_k)\big), \text{ for any } h\in H.$$

Proof. Let $y_1 \otimes_H x_1, \ldots, y_N \otimes_H x_N \in \mathcal{A}_Y \mathcal{L}$ and $\sigma \in Sh(l, k-1)$ be any shuffle with k+l = N+1. If

$$\beta_k \big(\beta_l (x_{\sigma(1)}, \dots, x_{\sigma(l)}), x_{\sigma(l+1)}, \dots, x_{\sigma(N)} \big) = \sum_i (f_{1i}^{\sigma} \otimes \dots \otimes f_{Ni}^{\sigma}) \otimes_H e_i$$

then we observe that

$$\begin{split} \beta_k \Big(\beta_l \big(y_{\sigma(1)} \otimes_H x_{\sigma(1)}, \dots, y_{\sigma(l)} \otimes_H x_{\sigma(l)} \big), y_{\sigma(l+1)} \otimes_H x_{\sigma(l+1)}, \dots, y_{\sigma(N)} \otimes_H x_{\sigma(N)} \Big) \\ &= \sum_i (y_{\sigma(1)} f_{1i}^{\sigma}) \cdots (y_{\sigma(N)} f_{Ni}^{\sigma}) \otimes_H e_i \\ &= (y_{\sigma(1)} \cdots y_{\sigma(N)}) (\mu_{H^{\otimes N}} \otimes 1) \beta_k \big(\beta_l (x_{\sigma(1)}, \dots, x_{\sigma(l)}), x_{\sigma(l+1)}, \dots, x_{\sigma(N)} \big). \end{split}$$

Here $\mu_{H^{\otimes N}} : H^{\otimes N} \to H$ is the multiplication of H. Hence the result follows after writing the expression of the higher Jacobi identities. The last part follows from the definition.

Note that the base field \mathbf{k} is an H-bimodule and a commutative associative H-differential algebra with the action $h1 = \varepsilon(h)$, for $h \in H$ and $1 \in \mathbf{k}$. Then it follows from the above Proposition that if $(\mathcal{L}, \{\beta_k\}_{k\geq 1})$ is an L_{∞} H-pseudoalgebra, then $(\mathcal{A}_{\mathbf{k}}\mathcal{L} = \mathbf{k} \otimes_H \mathcal{L}, \{\widetilde{\beta}_k\}_{k\geq 1})$ is an L_{∞} algebra. Explicitly, we have $\mathcal{A}_{\mathbf{k}}\mathcal{L} = \mathcal{L}/H_+\mathcal{L}$, where $H_+\mathcal{L} = \{h \cdot x \mid h \in H \text{ with } \varepsilon(h) = 0 \text{ and } x \in \mathcal{L}\}$ and the operations $\{\widetilde{\beta}_k\}_{k\geq 1}$ are given by

$$\widetilde{\beta}_k([x_1],\ldots,[x_k]) = \sum_i [\varepsilon(f_{1i})\cdots\varepsilon(f_{ki})e_i] \quad \text{if } \beta_k(x_1,\ldots,x_k) = \sum_i (f_{1i}\otimes\cdots\otimes f_{ki})\otimes_H e_i.$$

Here [x] denotes the class of the element $x \in \mathcal{L}$ in the quotient $\mathcal{L}/H_+\mathcal{L}$.

3.15. **Definition.** Let $Y = H^*$ be the dual of H. Then $Y = H^*$ is an H-bimodule and cocommutative associative H-differential algebra. For any L_{∞} H-pseudoalgebra $\mathcal{L} = (\mathcal{L}, \{\beta_k\}_{k \geq 1})$, the L_{∞} algebra

$$\mathcal{A}(\mathcal{L}) := (\mathcal{A}_{H^*}(\mathcal{L}) = H^* \otimes_H \mathcal{L}, \{\beta_k\}_{k \ge 1})$$

is called the *annihilation algebra* of \mathcal{L} .

In a forthcoming article, we will discuss annihilation algebra in more detail. More precisely, we aim to study differential L_{∞} algebras (following the work of Ritt [19] about differential Lie algebras) and find their relations with annihilation algebras.

4. Examples of strongly homotopy Lie *H*-pseudoalgebras

4.1. L_{∞} conformal algebras. The concept of Lie conformal algebras was introduced by Kac [13,14] to study the operator product expansions of chiral fields in conformal field theory. More precisely, a *Lie conformal algebra* is a $\mathbf{k}[\partial]$ -module *L* equipped with a \mathbf{k} -linear map $[\cdot_{\lambda} \cdot] : L \otimes L \to L[\lambda]$, called the λ -bracket that satisfy

- (conformal sesquilinearity) $[\partial x_{\lambda} y] = -\lambda [x_{\lambda} y]$ and $[x_{\lambda} \partial y] = (\partial + \lambda) [x_{\lambda} y]$,
- (skew-symmetry) $[y_{\lambda}x] = -[x_{-\lambda-\partial}y],$
- (conformal Jacobi identity) $[x_{\lambda}[y_{\nu}z]] = [[x_{\lambda}y]_{\lambda+\nu}z] + [y_{\nu}[x_{\lambda}z]].$

It has been observed in [2] that Lie conformal algebras are exactly Lie *H*-pseudoalgebras, where $H = \mathbf{k}[\partial]$ is the Hopf algebra of polynomials in one variable ∂ . The correspondence between the λ -bracket and the pseudobracket is given by

$$[x_{\lambda}y] = \sum_{i} p_{i}(\lambda)e_{i} \text{ if and only if } [x*y] = \sum_{i} \left(p_{i}(-\partial)\otimes 1\right)\otimes_{\mathbf{k}[\partial]} e_{i}.$$

On the other hand, L_{∞} conformal algebras were introduced in [21] as the strongly homotopy analogue of Lie conformal algebras. Recall that an L_{∞} conformal algebra is a graded $\mathbf{k}[\partial]$ -module $\mathcal{L} = \bigoplus_{n \in \mathbb{Z}} \mathcal{L}^n$ equipped with a collection

$$\{l_k: \mathcal{L}^{\otimes k} \to \mathcal{L}[\lambda_1, \dots, \lambda_{k-1}], x_1 \otimes \dots \otimes x_k \mapsto (l_k)_{\lambda_1, \dots, \lambda_{k-1}}(x_1, \dots, x_k) \text{ with } \deg(l_k) = k-2\}_{k \ge 1}$$

of k-linear maps that satisfy the following conditions:

- each l_k is conformal sesquilinear in the sense that

$$(l_k)_{\lambda_1,\dots,\lambda_{k-1}}(x_1,\dots,\partial x_i,\dots,x_{k-1},x_k) = -\lambda_i(l_k)_{\lambda_1,\dots,\lambda_{k-1}}(x_1,\dots,x_k),$$
$$(l_k)_{\lambda_1,\dots,\lambda_{k-1}}(x_1,\dots,x_{k-1},\partial x_k) = (\partial + \lambda_1 + \dots + \lambda_k)(l_k)_{\lambda_1,\dots,\lambda_{k-1}}(x_1,\dots,x_k),$$

- each l_k is graded skew-symmetric in the sense that

$$(l_k)_{\lambda_1,\dots,\lambda_{k-1}}(x_1,\dots,x_k) = \operatorname{sgn}(\sigma)\epsilon(\sigma) \ (l_k)_{\lambda_{\sigma(1)},\dots,\lambda_{\sigma(k-1)}}(x_{\sigma(1)},\dots,x_{\sigma(k-1)},x_{\sigma(k)})\big|_{\lambda_k\mapsto\lambda_k^{\dagger}}, \text{ for } \sigma\in S_k,$$

- for each $N \ge 1$ and homogeneous $x_1, \ldots, x_N \in \mathcal{L}$, the following higher conformal Jacobi identity holds:

$$\sum_{p+q=N+1} \sum_{\sigma \in \operatorname{Sh}(q,N-q)} \operatorname{sgn}(\sigma) \epsilon(\sigma) (-1)^{q(p-1)}$$

$$(15)$$

$$(l_p)_{\lambda_{\sigma}^{\dagger},\lambda_{\sigma(q+1)},\dots,\lambda_{\sigma(N-1)}} ((l_q)_{\lambda_{\sigma(1)},\dots,\lambda_{\sigma(q-1)}} (x_{\sigma(1)},\dots,x_{\sigma(q)}), x_{\sigma(q+1)},\dots,x_{\sigma(N)}) = 0.$$

Note that $\sigma \in \text{Sh}(q, N-q)$ implies that either $\sigma(q) = N$ or $\sigma(N) = N$. When $\sigma(q) = N$, we use the notation $\lambda_{\sigma}^{\dagger} = -\partial - (\lambda_{\sigma(q+1)} + \cdots + \lambda_{\sigma(N)})$ and when $\sigma(N) = N$, we use the notation $\lambda_{\sigma}^{\dagger} = \lambda_{\sigma(1)} + \cdots + \lambda_{\sigma(q)}$ in the above identities.

Then it can be checked that an L_{∞} conformal algebra is the same as an L_{∞} $\mathbf{k}[\partial]$ -pseudoalgebra. More precisely, $(\mathcal{L}, \{l_k\}_{k\geq 1})$ is an L_{∞} conformal algebra if and only if $(\mathcal{L}, \{\beta_k\}_{k\geq 1})$ is an L_{∞} $\mathbf{k}[\partial]$ -pseudoalgebra, where the correspondence between the structure maps are given by

$$(l_k)_{\lambda_1,\dots,\lambda_{k-1}}(x_1,\dots,x_k) = \sum_i p_i^{k,1}(\lambda_1)\cdots p_i^{k,k-1}(\lambda_{k-1})e_i \text{ if and only if}$$
$$\beta_k(x_1,\dots,x_k) = \left(p_i^{k,1}(-\partial)\otimes\cdots\otimes p_i^{k,k-1}(-\partial)\otimes 1\right)\otimes_{\mathbf{k}[\partial]} e_i.$$

4.2. Current L_{∞} *H*-pseudoalgebras. Given a Lie algebra \mathfrak{g} , one can define a Lie *H*-pseudoalgebra structure on the left *H*-module $H \otimes \mathfrak{g}$ with the pseudobracket

$$[(f \otimes x) * (g \otimes y)] := (f \otimes g) \otimes_H (1 \otimes [x, y]_{\mathfrak{g}}), \text{ for } f \otimes x, g \otimes y \in H \otimes \mathfrak{g}.$$

This is called the *current Lie H*-*pseudoalgebra* associated to \mathfrak{g} , and it is denoted by Cur(\mathfrak{g}). See [2] for more details. This construction can be generalized to the homotopy context. More precisely, let $(\mathcal{L}, \{l_k\}_{k\geq 1})$ be an L_{∞} algebra. Then the graded left *H*-module $H \otimes \mathcal{L}$ equipped with the graded polylinear maps

 $\left\{\beta_k \in \operatorname{Hom}_{H^{\otimes k}}^{k-2}\left((H \otimes \mathcal{L})^{\boxtimes k}, H^{\otimes k} \otimes_H (H \otimes \mathcal{L})\right)\right\}_{k \ge 1}$

forms a L_∞ $H\mbox{-}p\mbox{seudoalgebra}, where$

$$\beta_k(f_1 \otimes x_1, \dots, f_k \otimes x_k) = (f_1 \otimes \dots \otimes f_k) \otimes_H (1 \otimes l_k(x_1, \dots, x_k)), \text{ for any } k \ge 1.$$

The verification is straightforward. This is called the current L_{∞} *H*-pseudoalgebra associated to the L_{∞} algebra $(\mathcal{L}, \{l_k\}_{k\geq 1})$. We denote it by Cur (\mathcal{L}) .

More generally, let $H' \subset H$ be a Hopf subalgebra and $(\mathcal{L}, \{\beta'_k\}_{k\geq 1})$ be an L_{∞} H'-pseudoalgebra. Then it can be verified that the pair $\operatorname{Cur}_{H'}^H \mathcal{L} = (H \otimes_{H'} \mathcal{L}, \{\beta_k\}_{k\geq 1})$ is an L_{∞} H-pseudoalgebra, where the structure

maps are given by

$$\beta_k (f_1 \otimes_{H'} x_1, \dots, f_k \otimes_{H'} x_k) = \sum_i (f_1 f_{1,i} \otimes \dots \otimes f_k f_{k,i}) \otimes_H (1 \otimes_{H'} e_i)$$

whenever $\beta'_k (x_1, \dots, x_k) = \sum_i (f_{1,i} \otimes \dots \otimes f_{k,i}) \otimes_{H'} e_i.$

4.3. Graded Lie *H*-pseudoalgebras whose each arity is a free *H*-module of rank 1. Let $\mathcal{L} = \bigoplus_{n \in \mathbb{Z}} He_n$ be a graded left *H*-module whose each arity is a free *H*-module of rank 1. Note that any polylinear map $\beta_k \in \operatorname{Hom}_{H^{\otimes k}}^{k-2}(\mathcal{L}^{\boxtimes k}, H^{\otimes k} \otimes_H \mathcal{L})$ is determined by the values $\beta_k(e_{i_1}, e_{i_2}, \ldots, e_{i_k})$, or equivalently, by elements $\alpha_{i_1,\ldots,i_k} \in H^{\otimes k}$ such that $\beta_k(e_{i_1},\ldots,e_{i_k}) = \alpha_{i_1,\ldots,i_k} \otimes_H e_{i_1+\cdots+i_k+k-2}$.

Then we have the following.

4.1. **Proposition.** Let $\mathcal{L} = \bigoplus_{n \in \mathbb{Z}} He_n$ be a graded left H-module whose each arity is a free H-module of rank 1. Suppose there are maps $\{\beta_k \in \operatorname{Hom}_{H^{\otimes k}}^{k-2}(\mathcal{L}^{\boxtimes k}, H^{\otimes k} \otimes_H \mathcal{L})\}_{k \geq 1}$ which are determined by the elements α_{i_1,\ldots,i_k} $(k \geq 1 \text{ and } i_1,\ldots,i_k \in \mathbb{Z})$. Then $(\mathcal{L}, \{\beta_k\}_{k \geq 1})$ is a L_{∞} H-pseudoalgebra if and only if

$$\begin{aligned} \alpha_{i_{\sigma(1)},\dots,i_{\sigma(k)}} &= (-1)^{\sigma} \epsilon(\sigma; i_1,\dots,i_k) \alpha_{i_1,\dots,i_k}, \text{ for all } k \ge 1 \text{ and } \sigma \in S(k), \\ \sum_{k+l=N+1} \sum_{\sigma \in \operatorname{Sh}(l,k-1)} (-1)^{\sigma} \epsilon(\sigma) (-1)^{l(k-1)} \\ & \left(\alpha_{i_{\sigma(1)},\dots,i_{\sigma(l)}} \otimes 1^{\otimes k-1}\right) (\Delta^l \otimes \operatorname{id}^{\otimes k-1}) \alpha_{i_{\sigma(1)}+\dots+i_{\sigma(l)}+l-2,i_{\sigma(l+1)},\dots,i_{\sigma(N)}} = 0, \end{aligned}$$

for all $N \in \mathbb{N}$. Here $\Delta^l : H \to H^{\otimes l}$ is the iterative comultiplication map.

Proof. For any $\sigma \in Sh(l, k-1)$, note that

$$\beta_l(x_{\sigma(1)},\ldots,x_{\sigma(l)}) = \alpha_{i_{\sigma(1)},\ldots,i_{\sigma(l)}} \otimes_H e_{i_{\sigma(1)}+\cdots+i_{\sigma(l)}+l-2}$$

Therefore,

$$\beta_k(\beta_l(x_{\sigma(1)},\ldots,x_{\sigma(l)}),x_{\sigma(l+1)},\ldots,x_{\sigma(N)})$$

= $(\alpha_{i_{\sigma(1)},\ldots,i_{\sigma(l)}} \otimes 1^{\otimes k-1})(\Delta^l \otimes \mathrm{id}^{\otimes k-1})\alpha_{i_{\sigma(1)}+\cdots+i_{\sigma(l)}+l-2,i_{\sigma(l+1)},\ldots,i_{\sigma(N)}}.$

Hence the result follows by using the definition of an L_{∞} *H*-pseudoalgebra.

4.2. **Remark.** Let $\mathcal{L} = \bigoplus_{n \in \mathbb{Z}} He_n$ be a graded left *H*-module whose each arity is a free *H*-module of rank 1. Suppose there is a map $\beta \in \operatorname{Hom}_{H^{\otimes 2}}^0(\mathcal{L}^{\boxtimes 2}, H^{\otimes 2} \otimes_H \mathcal{L})$ which is determined by the elements $\alpha_{i,j} \in H^{\otimes 2}$ (for $i, j \in \mathbb{Z}$). Then (\mathcal{L}, β) is a graded Lie *H*-pseudoalgebra if and only if

$$\alpha_{i,j} = -(-1)^{ij} \alpha_{j,i},$$

$$(\alpha_{i,j} \otimes 1)(\Delta \otimes \mathrm{id}) \alpha_{i+j,k} = (1 \otimes \alpha_{j,k})(\mathrm{id} \otimes \Delta) \alpha_{i,j+k} - (-1)^{ij} (\sigma_{12} \otimes \mathrm{id}) ((1 \otimes \alpha_{i,k})(\mathrm{id} \otimes \Delta) \alpha_{j,i+k}) = 0.$$

4.4. Strongly homotopy associative *H*-pseudoalgebras. In this subsection, we first introduce the notion of A_{∞} *H*-pseudoalgebras as the strongly homotopy version of associative *H*-pseudoalgebras. In the end, we show that a suitable skew-symmetrization of an A_{∞} *H*-pseudoalgebra gives rise to an L_{∞} *H*-pseudoalgebra.

4.3. **Definition.** ([2]) Let C be a pseudotensor category. An associative algebra in C is a pair (A, μ) of an object A with a polylinear map $\mu \in \text{Lin}(\{A, A\}, A)$ satisfying the associativity $\mu(\mu(\cdot, \cdot), \cdot) = \mu(\cdot, \mu(\cdot, \cdot))$.

Let H be a cocommutative Hopf algebra (although the cocommutativity is not required to define associative H-pseudoalgebras). An associative H-pseudoalgebra is an associative algebra in the pseudotensor category $\mathcal{M}^*(H)$. It follows that an associative H-pseudoalgebra is a left H-module A with a map $\mu \in \operatorname{Hom}_{H^{\otimes 2}}(A \boxtimes A, H^{\otimes 2} \otimes_H A)$, often called the *pseudoproduct* denoted by $\mu(a \otimes b) = a * b$, satisfying (a * b) * c = a * (b * c), for all $a, b, c \in A$. Here the compositions are similar to the one given in (4) and (5).

One can also define graded associative algebras, differential graded associative algebras and A_{∞} algebras in a graded pseudotensor category. If we consider the graded pseudotensor category $\mathcal{M}_{qr}^*(H)$ associative

to a cocommutative Hopf algebra H, an A_{∞} algebra in $\mathcal{M}_{gr}^*(H)$ is simply called an A_{∞} H-pseudoalgebra. More precisely, we have the following.

4.4. **Definition.** An A_{∞} *H*-pseudoalgebra is a pair $(\mathcal{A}, \{\mu_k\}_{k\geq 1})$ consisting of a graded left *H*-module $\mathcal{A} = \bigoplus_{n \in \mathbb{Z}} \mathcal{A}^n$ equipped with a collection of polylinear maps

$$\{\mu_k \in \operatorname{Hom}_{H^{\otimes k}}^{k-2}(\mathcal{A}^{\boxtimes k}, H^{\otimes k} \otimes_H \mathcal{A})\}_{k \ge 1}$$

subject to satisfy the following higher associativity identities: for each $N \ge 1$ and homogeneous elements $a_1, \ldots, a_N \in \mathcal{A}$,

$$\sum_{k+l=N+1} \sum_{\lambda=1}^{k} (-1)^{\lambda(l+1)+l(|a_1|+\dots+|a_{\lambda-1}|)} \mu_k(a_1,\dots,a_{\lambda-1},\mu_l(a_\lambda,\dots,a_{\lambda+l-1}),a_{\lambda+l},\dots,a_N) = 0.$$
(16)

As before, μ_k is $H^{\otimes k}$ -linear simply means that $\mu_k(h_1a_1, \ldots, h_ka_k) = ((h_1 \otimes \cdots \otimes h_k) \otimes_H 1)\mu_k(a_1, \ldots, a_k)$, for all $a_1, \ldots, a_k \in \mathcal{A}$ and $h_1, \ldots, h_k \in H$. Finally, to explicitly describe the higher associativity identities, we will write down explicitly each term under the summation of (16). First, for any fixed $k, l \geq 1$ with k+l = N+1, suppose

$$\mu_l(a_{\lambda},\ldots,a_{\lambda+l-1}) = \sum_i (h_i^1 \otimes \cdots \otimes h_i^l) \otimes_H e_i \in H^{\otimes l} \otimes_H \mathcal{A}$$

and

$$\mu_k(a_1,\ldots,a_{\lambda-1},e_i,a_{\lambda+1},\ldots,a_N) = \sum_j (h_{ij}^{\lambda,1} \otimes \cdots \otimes h_{ij}^{\lambda,k}) \otimes_H e_{ij} \in H^{\otimes k} \otimes_H \mathcal{A}$$

Then $\mu_k(a_1,\ldots,a_{\lambda-1},\mu_l(a_{\lambda},\ldots,a_{\lambda+l-1}),a_{\lambda+l},\ldots,a_N)$ is the element of $H^{\otimes N}\otimes_H \mathcal{A}$ given by

$$\sum_{i,j} \left(h_{ij}^{\lambda,1} \otimes \cdots \otimes h_{ij}^{\lambda,\lambda-1} \otimes h_i^1 h_{ij(1)}^{\lambda,\lambda} \otimes \cdots \otimes h_i^l h_{ij(l)}^{\lambda,\lambda} \otimes h_{ij}^{\lambda,\lambda+1} \otimes \cdots \otimes h_{ij}^{\lambda,k} \right) \otimes_H e_{ij}$$

4.5. **Remark.** (i) When $H = \mathbf{k}$, an A_{∞} *H*-pseudoalgebra is nothing but a classical A_{∞} algebra [22].

(ii) Any associative *H*-pseudoalgebra can be regarded as an A_{∞} *H*-pseudoalgebra concentrated in arity 0.

Let $\mathcal{W} = \bigoplus_{n \in \mathbb{Z}} \mathcal{W}^n$ be a graded left *H*-module. Then the graded space $\bigoplus_{p \in \mathbb{Z}} \operatorname{Hom}_{\operatorname{poly}}^p(\mathcal{W}, \mathcal{W})$ carries a graded Lie bracket given by

$$\llbracket\sum_{k\geq 1}\nu_k, \sum_{l\geq 1}\theta_l\rrbracket^{\sim} = \sum_{N\geq 1}\sum_{k+l=N+1}\llbracket\nu_k, \theta_l\rrbracket^{\sim} := \sum_{N\geq 1}\sum_{k+l=N+1} \left(\nu_k\odot\theta_l - (-1)^{pq}\theta_l\odot\nu_k\right),$$

where

$$(\eta_k \odot \zeta_l)(w_1, \dots, w_N) := \sum_{i=1}^k (-1)^{|w_1| + \dots + |w_{i-1}|} \nu_k (w_1, \dots, w_{i-1}, \theta_l(w_i, \dots, w_{i+l-1}), w_{i+l}, \dots, w_N),$$

for $\sum_{k\geq 1} \nu_k \in \operatorname{Hom}_{\operatorname{poly}}^p(\mathcal{W}, \mathcal{W}), \sum_{l\geq 1} \theta_l \in \operatorname{Hom}_{\operatorname{poly}}^q(\mathcal{W}, \mathcal{W})$ and $w_1, \ldots, w_N \in \mathcal{W}$. Then similar to Theorem 3.8, here one can prove the following result.

4.6. **Theorem.** Let $\mathcal{A} = \bigoplus_{n \in \mathbb{Z}} \mathcal{A}^n$ be a graded left *H*-module. Then there is a one-to-one correspondence between A_{∞} *H*-pseudoalgebra structures on \mathcal{A} and Maurer-Cartan elements in the graded Lie algebra $(\bigoplus_{p \in \mathbb{Z}} \operatorname{Hom}_{polv}^p(\mathcal{W}, \mathcal{W}), [[,]]^{\sim})$, where $\mathcal{W} = \mathcal{A}[-1]$.

4.7. **Remark.** Note that the graded Lie algebra $(\bigoplus_{p \in \mathbb{Z}} \operatorname{Hom}_{\operatorname{poly}}^{p}(\mathcal{W}, \mathcal{W}), [\![,]\!]^{\sim})$ constructed above has a close connection with the graded Lie algebra $(\bigoplus_{p \in \mathbb{Z}} \operatorname{symHom}_{\operatorname{poly}}^{p}(\mathcal{W}, \mathcal{W}), [\![,]\!])$ defined in Section 3. More precisely, for any $p \in \mathbb{Z}$, the map $(-)^{\operatorname{sym}} : \operatorname{Hom}_{\operatorname{poly}}^{p}(\mathcal{W}, \mathcal{W}) \to \operatorname{symHom}_{\operatorname{poly}}^{p}(\mathcal{W}, \mathcal{W})$ defined by

$$(\sum_{k\geq 1}\nu_k)^{\text{sym}} = \sum_{k\geq 1}(\nu_k)^{\text{sym}}, \text{ where}$$
$$(\nu_k)^{\text{sym}}(w_1,\ldots,w_k) := \sum_{\sigma\in S(k)}\epsilon(\sigma)(\sigma\otimes_H 1)\nu_k(w_{\sigma(1)},\ldots,w_{\sigma(k)})$$

defines a morphism from $(\bigoplus_{p \in \mathbb{Z}} \operatorname{Hom}_{\operatorname{poly}}^{p}(\mathcal{W}, \mathcal{W}), [\![,]\!]^{\sim})$ to $(\bigoplus_{p \in \mathbb{Z}} \operatorname{symHom}_{\operatorname{poly}}^{p}(\mathcal{W}, \mathcal{W}), [\![,]\!])$.

In [2] the authors showed that an associative *H*-pseudoalgebra gives rise to a Lie *H*-pseudoalgebra if we take the skew-symmetrization of the associative pseudoproduct. More precisely, let (A, μ) be an associative *H*-pseudoalgebra, where we use the notation $\mu(a \otimes b) = a * b$, for $a, b \in A$. Then (A, β) is a Lie *H*-pseudoalgebra, where the skew-symmetric pseudobracket β is given by

$$\beta(a \otimes b) = [a * b] := a * b - (\sigma_{12} \otimes_H 1)(b * a), \text{ for } a, b \in A.$$

In the following, we will generalize this result in the strongly homotopy context.

4.8. **Theorem.** Let $(\mathcal{A}, \{\mu_k\}_{k\geq 1})$ be an A_{∞} H-pseudoalgebra. Then $(\mathcal{A}, \{\beta_k\}_{k\geq 1})$ is an L_{∞} H-pseudoalgebra, where

$$\beta_k(a_1,\ldots,a_k) = \sum_{\sigma \in S_k} (-1)^{\sigma} \epsilon(\sigma) (\sigma \otimes_H 1)(\mu_k)(a_{\sigma^{-1}(1)},\ldots,a_{\sigma^{-1}(k)}).$$
(17)

Proof. Let $\mathcal{W} = \mathcal{A}[-1]$. Since $(\mathcal{A}, \{\mu_k\}_{k\geq 1})$ is ab A_{∞} *H*-pseudoalgebra, there is a Maurer-Cartan element in the graded Lie algebra $(\bigoplus_{p\in\mathbb{Z}} \operatorname{Hom}_{\operatorname{poly}}^{p}(\mathcal{W}, \mathcal{W}), [\![,]\!]^{\sim})$. The Maurer-Cartan element is precisely given by $\nu = \sum_{k\geq 1} \nu_k \in \operatorname{Hom}_{\operatorname{poly}}^{-1}(\mathcal{W}, \mathcal{W})$, where

$$\nu_k = (-1)^{\frac{k(k-1)}{2}} \ s \circ \mu_k \circ (s^{-1})^{\otimes k}.$$

Next, consider the collection of skew-symmetric polylinear maps $\{\beta_k\}_{k\geq 1}$ as defined in (17). If $\{\eta_k\}_{k\geq 1}$ are the symmetric polylinear maps in symHom⁻¹_{poly}(\mathcal{W}, \mathcal{W}) defined by $\eta_k = (-1)^{\frac{k(k-1)}{2}} s \circ \beta_k \circ (s^{-1})^{\otimes k}$, then by a simple observation (see also [24]) shows that

$$(\nu_k)^{\text{sym}} = \eta_k$$
, for all $k \ge 1$.

Hence $(\sum_{k\geq 1} \nu_k)^{\text{sym}} = \sum_{k\geq 1} (\nu_k)^{\text{sym}} = \sum_{k\geq 1} eta_k$. Therefore,

$$\llbracket\sum_{k\geq 1}\eta_k, \sum_{k\geq 1}\eta_k\rrbracket = \llbracket(\sum_{k\geq 1}\nu_k)^{\operatorname{sym}}, (\sum_{k\geq 1}\nu_k)^{\operatorname{sym}}\rrbracket = \left(\llbracket\sum_{k\geq 1}\nu_k, \sum_{k\geq 1}\nu_k\rrbracket^{\sim}\right) = 0.$$

The result now follows as a consequence of Theorem 3.8.

5. Skeletal and strict homotopy Lie H-pseudoalgebras

Our aim in this section is to discuss and study some particular classes of L_{∞} *H*-pseudoalgebras. We first start with those L_{∞} *H*-pseudoalgebras whose underlying graded left *H*-module is concentrated in arities 0 and 1. We call them 2-term L_{∞} *H*-pseudoalgebras. Next, we consider skeletal and strict L_{∞} *H*-pseudoalgebras that are special classes of 2-term L_{∞} *H*-pseudoalgebras. We show that skeletal L_{∞} *H*-pseudoalgebras correspond to third cocycles of (ordinary) Lie *H*-pseudoalgebras. Finally, we introduce crossed modules of Lie *H*-pseudoalgebras and show that they correspond to strict L_{∞} *H*-pseudoalgebras. Our results in this section are motivated by the results developed by Baez and Crans [1] for classical L_{∞} algebras.

We have already mentioned that 2-term L_{∞} *H*-pseudoalgebras are those L_{∞} *H*-pseudoalgebras whose underlying graded left *H*-module is concentrated in arities 0 and 1. Hence by assuming $\mathcal{L} = \mathcal{L}_0 \oplus \mathcal{L}_1$ in Definition 3.5, we get the following.

5.1. **Definition.** A 2-term L_{∞} *H*-pseudoalgebra is a triple $(\mathcal{L}_1 \xrightarrow{\beta_1} \mathcal{L}_0, \beta_2, \beta_3)$ consisting of a 2-term chain complex $\mathcal{L}_1 \xrightarrow{\beta_1} \mathcal{L}_0$ of left *H*-modules, a skew-symmetric polylinear map $\beta_2 \in \operatorname{Hom}_{H^{\otimes 2}}(\mathcal{L}_i \boxtimes \mathcal{L}_j, H^{\otimes 2} \otimes_H \mathcal{L}_{i+j})$ and another skew-symmetric polylinear map $\beta_3 \in \operatorname{Hom}_{H^{\otimes 3}}(\mathcal{L}_0 \boxtimes \mathcal{L}_0, H^{\otimes 3} \otimes_H \mathcal{L}_1)$ subject to satisfy the following conditions:

(i)
$$\beta_1(\beta_2(x, u)) = \beta_2(x, \beta_1(u)),$$

(ii)
$$\beta_2(\beta_1(u), v) = \beta_2(u, \beta_1(v))$$

(iii) $\beta_2(\beta_2(x,y),z) - \beta_2(x,\beta_2(y,z)) + \beta_2(y,\beta_2(x,z)) = -\beta_1(\beta_3(x,y,z)),$

- (iv) $\beta_2(\beta_2(x,y),v) \beta_2(x,\beta_2(y,v)) + \beta_2(y,\beta_2(x,v)) = -\beta_3(x,y,\beta_1(v)),$
- (v) (the Jacobiator identity)

$$\begin{split} \beta_2(x, \beta_3(y, z, w)) &- \beta_2(y, \beta_3(x, z, w)) + \beta_2(z, \beta_3(x, y, w)) - \beta_2(w, \beta_3(x, y, z)) \\ &- \beta_3(\beta_2(x, y), z, w) + \beta_3(\beta_2(x, z), y, w) - \beta_3(\beta_2(x, w), y, z) \\ &- \beta_3(\beta_2(y, z), x, w) + \beta_3(\beta_2(y, w), x, z) - \beta_3(\beta_2(z, w), x, y) = 0, \end{split}$$

for all $x, y, z, w \in \mathcal{L}_0$ and $u, v \in \mathcal{L}_1$.

Let $\mathcal{L} = (\mathcal{L}_1 \xrightarrow{\beta_1} \mathcal{L}_0, \beta_2, \beta_3)$ and $\mathcal{L}' = (\mathcal{L}'_1 \xrightarrow{\beta'_1} \mathcal{L}'_0, \beta'_2, \beta'_3)$ be 2-term L_{∞} *H*-pseudoalgebras. A morphism between them is given by a triple $f = (f_0, f_1, f_2)$ consisting of *H*-linear maps $f_0 : \mathcal{L}_0 \to \mathcal{L}'_0$ and $f_1 : \mathcal{L}_1 \to \mathcal{L}'_1$, and a polylinear map $f_2 \in \operatorname{Hom}_{H^{\otimes 2}}(\mathcal{L}_0 \boxtimes \mathcal{L}_0, H^{\otimes 2} \otimes_H \mathcal{L}'_1)$ satisfying

$$\beta_{1} \circ f_{0} = f_{0} \circ \beta_{1}',$$

$$(\mathrm{id}_{H^{\otimes 2} \otimes_{H}} f_{0})(\beta_{2}(x,y)) - \beta_{2}'(f_{0}(x), f_{0}(y)) = (\mathrm{id}_{H^{\otimes 2} \otimes_{H}} \beta_{1}')(f_{2}(x,y)),$$

$$(\mathrm{id}_{H^{\otimes 2} \otimes_{H}} f_{1})(\beta_{2}(x,u)) - \beta_{2}'(f_{0}(x), f_{1}(u)) = f_{2}(x, \beta_{1}(u)),$$

$$\beta_{2}'(f_{2}(x,y), f_{0}(z)) - \beta_{2}'(f_{0}(x), f_{2}(y,z)) + \beta_{2}'(f_{0}(y), f_{2}(x,z)) + f_{2}(\beta_{2}(x,y),z)$$

$$-f_{2}(x, \beta_{2}(y,z)) + f_{2}(y, \beta_{2}(x,z)) = \beta_{3}'(f_{0}(x), f_{0}(y), f_{0}(z)) - (\mathrm{id}_{H^{\otimes 3} \otimes_{H}} f_{1})(\beta_{3}(x,y,z)),$$

for all $x, y, z \in \mathcal{L}_0$ and $u \in \mathcal{L}_1$. We often denote a morphism as above by $(f_0, f_1, f_2) : \mathcal{L} \rightsquigarrow \mathcal{L}'$.

Let $\mathcal{L} = (\mathcal{L}_1 \xrightarrow{\beta_1} \mathcal{L}_0, \beta_2, \beta_3), \ \mathcal{L}' = (\mathcal{L}'_1 \xrightarrow{\beta'_1} \mathcal{L}'_0, \beta'_2, \beta'_3) \text{ and } \mathcal{L}'' = (\mathcal{L}''_1 \xrightarrow{\beta''_1} \mathcal{L}''_0, \beta''_2, \beta''_3) \text{ be 2-term } L_{\infty} H$ pseudoalgebras. Suppose $f = (f_0, f_1, f_2) : \mathcal{L} \rightsquigarrow \mathcal{L}' \text{ and } g = (g_0, g_1, g_2) : \mathcal{L}' \rightsquigarrow \mathcal{L}''$ are morphisms between 2-term $L_{\infty} H$ -pseudoalgebras. Then their composition $g \circ f : \mathcal{L} \rightsquigarrow \mathcal{L}''$ is given by

$$g \circ f = (g_0 \circ f_0, g_1 \circ f_1, g_2 \circ (f_0 \otimes f_0) + (\mathrm{id}_{H^{\otimes 2}} \otimes g_1) \circ f_2).$$

With all the above definitions and notations, we have the following result.

5.2. **Proposition.** The collection of all 2-term L_{∞} H-pseudoalgebras and morphisms between them forms a category. (We denote this category by **2shLie**H).

In the following, we consider skeletal and strict L_{∞} *H*-pseudoalgebras and provide their characterizations.

5.3. Definition. Let $(\mathcal{L}_1 \xrightarrow{\beta_1} \mathcal{L}_0, \beta_2, \beta_3)$ be a 2-term L_{∞} *H*-pseudoalgebra. It is said to be

- (i) skeletal if $\beta_1 = 0$,
- (ii) strict if $\beta_3 = 0$.

Let $(\mathcal{L}_1 \xrightarrow{0} \mathcal{L}_1, \beta_2, \beta_3)$ be a skeletal L_{∞} *H*-pseudoalgebra. It follows from condition (iii) of the Definition 5.1 that the left *H*-module \mathcal{L}_0 with the skew-symmetric polylinear map $\beta_2 \in \operatorname{Hom}_{H^{\otimes 2}}(\mathcal{L}_0 \boxtimes \mathcal{L}_0, H^{\otimes 2} \otimes_H \mathcal{L}_0)$ is a Lie *H*-pseudoalgebra. Further, it follows from the condition (iv) of the same definition that the left *H*-module \mathcal{L}_1 equipped with the polylinear map $\beta_2 \in \operatorname{Hom}_{H^{\otimes 2}}(\mathcal{L}_0 \boxtimes \mathcal{L}_1, H^{\otimes 2} \otimes_H \mathcal{L}_1)$ is a representation of the Lie *H*-pseudoalgebra (\mathcal{L}_0, β_2) . Finally, the condition (v) is equivalent to $(\delta\beta_3)(x, y, z, w) = 0$, where δ is the coboundary map of the Lie *H*-pseudoalgebra (\mathcal{L}_0, β_0) with coefficients in the representation (\mathcal{L}_1, β_2) . In other words, $\beta_3 \in C^3_{\operatorname{Lie}-H}(\mathcal{L}_0, \mathcal{L}_1)$ is a 3-cocycle.

5.4. **Proposition.** There is a one-to-one correspondence between skeletal L_{∞} H-pseudoalgebras and triples of the form (L, M, θ) in which L is a Lie H-pseudoalgebra, M is a representation and θ is a 3-cocycle.

Proof. Let $(\mathcal{L}_1 \xrightarrow{0} \mathcal{L}_1, \beta_2, \beta_3)$ be a skeletal L_{∞} *H*-pseudoalgebra. Then $((\mathcal{L}_0, \beta_2), (\mathcal{L}_1, \beta_2), \beta_3)$ is the required triple.

Conversely, let (L, M, θ) be a triple in which $L = (L, \beta)$ is a Lie *H*-pseudoalgebra, $M = (M, \gamma)$ is a representation and θ is a 3-cocycle. Then $(M \xrightarrow{0} L, \beta_2, \beta_3)$ is a skeletal L_{∞} *H*-pseudoalgebra, where

 $\beta_2(x,y) = \beta(x \otimes y), \ \beta_2(x,u) = -\beta_2(u,x) = \gamma(x,u) \text{ and } \beta_3 = \theta, \text{ for } x, y \in L, u \in M.$

17

The above two correspondences are inverses to each other.

The above result has generalizations for higher cocycles. Let n > 2 be a fixed natural number and $(\mathcal{L}, \{\beta_k\}_{k\geq 1})$ be an L_{∞} *H*-pseudoalgebra whose underlying graded left *H*-module \mathcal{L} is concentrated in arities 0 and n-1. That is, \mathcal{L} and the map β_1 (which is automatically trivial in this case) can be realized as the chain complex $\mathcal{L}_{n-1} \to 0 \to \cdots \to 0 \to \mathcal{L}_0$. Further, for the degree reason, it can be easily observed that the only non-trivial operations in $\{\beta_k\}_{k\geq 1}$ are given by

$$\beta_{2} \in \operatorname{Hom}_{H^{\otimes 2}}(\mathcal{L}_{0} \boxtimes L_{0}, H^{\otimes 2} \otimes_{H} \mathcal{L}_{0}), \quad \beta_{2} \in \operatorname{Hom}_{H^{\otimes 2}}(\mathcal{L}_{0} \boxtimes L_{n-1}, H^{\otimes 2} \otimes_{H} \mathcal{L}_{n-1}) \text{ and} \\ \beta_{n+1} \in \operatorname{Hom}_{H^{\otimes n+1}}(\mathcal{L}_{0}^{\boxtimes n+1}, H^{\otimes n+1} \otimes_{H} \mathcal{L}_{n-1}).$$

Then from the higher Jacobi identities (for N = 3), it follows that the skew-symmetric polylinear map $\beta_2 \in \operatorname{Hom}_{H^{\otimes 2}}(\mathcal{L}_0 \boxtimes L_0, H^{\otimes 2} \otimes_H \mathcal{L}_0)$ makes the left *H*-module \mathcal{L}_0 into a Lie *H*-pseudoalgebra and the polylinear map $\beta_2 \in \operatorname{Hom}_{H^{\otimes 2}}(\mathcal{L}_0 \boxtimes L_{n-1}, H^{\otimes 2} \otimes_H \mathcal{L}_{n-1})$ makes the pair $(\mathcal{L}_{n-1}, \beta_2)$ into a representation of the Lie *H*-pseudoalgebra (\mathcal{L}_0, β_2) . Further, the higher Jacobi identity for N = n + 2 is equivalent to $(\delta \beta_{n+1})(x_1, \ldots, x_{n+2}) = 0$, where δ is the coboundary operator of the Lie *H*-pseudoalgebra (\mathcal{L}_0, β_2) with coefficients in the representation $(\mathcal{L}_{n-1}, \beta_2)$. We have the following result whose proof is similar to the proof of Proposition 5.4.

5.5. **Proposition.** There is a one-to-one correspondence between L_{∞} H-pseudoalgebras whose underlying graded left H-module is concentrated in arities 0 and n-1 (for n > 2), and triples of the form (L, M, θ) in which L is a Lie H-pseudoalgebra, M is a representation and θ is a (n + 1)-cocycle.

We now introduce crossed modules of Lie *H*-pseudoalgebras and find their relations with strict L_{∞} *H*-pseudoalgebras.

5.6. **Definition.** A crossed modules of Lie *H*-pseudoalgebras is a quadruple (L, L', φ, γ) in which L, L'are both Lie *H*-pseudoalgebras, $\varphi : L' \to L$ is a morphism of Lie *H*-pseudoalgebras and $\gamma \in \text{Hom}_{H^{\otimes 2}}(L \boxtimes L', H^{\otimes 2} \otimes_H L')$ is a polylinear map that makes (L', γ) into a representation of the Lie *H*-pseudoalgebra *L* satisfying additionally

$$(\mathrm{id}_{H^{\otimes 2}} \otimes_H \varphi)(\gamma(x, y')) = [x * \varphi(y')], \ \gamma(\varphi(x'), y') = [x' * y']' \text{ and} [\gamma(x, y') * z']' = \gamma(x, [y' * z']') - [y' * \gamma(x, z')]', \text{ for } x \in L, x', y', z' \in L'.$$

Here $[\cdot * \cdot]$ and $[\cdot * \cdot]'$ denote the pseudobrackets of the Lie *H*-pseudoalgebras *L* and *L'*, respectively.

Let L be a Lie H-pseudoalgebra with the pseudobracket $\beta = [\cdot * \cdot]$. Then the quadruple $(L, L, \mathrm{id}, \beta)$ is a crossed modules of Lie H-pseudoalgebras.

Let L be a Lie H-pseudoalgebra and $N \subset L$ be an ideal (i.e. $[x * n] \in H^{\otimes 2} \otimes_H N$, for all $x \in L$ and $n \in N$). Then the quadruple (L, N, i, ad) is a crossed modules of Lie H-pseudoalgebras, where $i : N \hookrightarrow L$ is the inclusion map and ad denotes the adjoint representation.

Let L, L' be two Lie *H*-pseudoalgebras and $\theta : L \to L'$ be a morphism of Lie *H*-pseudoalgebras. Then $(L, \ker \theta, i, ad)$ is a crossed modules of Lie *H*-pseudoalgebras.

The following result is straightforward, hence we omit the proof.

5.7. **Proposition.** Let (L, L', φ, γ) be a crossed modules of Lie H-pseudoalgebras. Then the direct sum left H-module $L \oplus L'$ carries a Lie H-pseudoalgebra structure with the pseudobracket

$$[(x,x')*(y,y')]_{L\oplus L'} := ([x*y], \gamma(x,y') - (\sigma_{12} \otimes_H 1)\gamma(y,x') + [x'*y']'), \text{ for } (x,x'), (y,y') \in L \oplus L'.$$

5.8. Theorem. There is a one-to-one correspondence between strict L_{∞} H-pseudoalgebras and crossed modules of Lie H-pseudoalgebras.

Proof. Let $(\mathcal{L}_1 \xrightarrow{\beta_1} \mathcal{L}_0, \beta_2, \beta_3 = 0)$ be a strict L_{∞} *H*-pseudoalgebra. Then it follows from condition (iii) of Definition 5.1 that the left *H*-module \mathcal{L}_0 equipped with the skew-symmetric polylinear map $\beta_2 \in$ $\operatorname{Hom}_{H^{\otimes 2}}(\mathcal{L}_0 \boxtimes \mathcal{L}_0, H^{\otimes} \otimes_H \mathcal{L}_0)$ is a Lie *H*-pseudoalgebra. Next, consider the left *H*-module \mathcal{L}_1 with the

APURBA DAS

skew-symmetric polylinear map $[u * v]' := \beta_2(\beta_1(u), v) = \beta_2(u, \beta_1(v))$, for $u, v \in \mathcal{L}_1$. Then it follows from condition (iv) of Definition 5.1 that \mathcal{L}_1 is a Lie *H*-pseudoalgebra. Further, the *H*-linear map $\beta_1 : \mathcal{L}_1 \to \mathcal{L}_0$ is a morphism of Lie *H*-pseudoalgebras (follows from condition (i)). Finally, the polylinear map $\beta_2 \in$ $\operatorname{Hom}_{H^{\otimes 2}}(\mathcal{L}_0 \boxtimes \mathcal{L}_1, H^{\otimes 2} \otimes_H \mathcal{L}_1)$ defines a representation of the Lie *H*-pseudoalgebra (\mathcal{L}_0, β_2) on the left *H*-module \mathcal{L}_1 . With these notations, it is easy to verify that ($\mathcal{L}_0, \mathcal{L}_1, \beta_1, \beta_2$) is a crossed modules of Lie *H*-pseudoalgebras.

On the other hand, if $((L, [\cdot * \cdot]), (L', [\cdot * \cdot]'), \varphi, \gamma)$ is a crossed modules of Lie *H*-pseudoalgebras, then it can be verified that $(L' \xrightarrow{\varphi} L, \beta_2, \beta_3 = 0)$ is a strict L_{∞} *H*-pseudoalgebra, where

 $\beta_2(x,y) = [x*y], \ \beta_2(x,x') = -\beta_2(x',x) := \gamma(x,x') \ \text{ and } \ \beta_2(x',y') = 0, \ \text{for } x,y \in L, x',y' \in L'.$

Finally, the above two correspondences are inverses to each other.

Combining Proposition 5.7 with Theorem 5.8, we get the following result.

5.9. **Proposition.** Let $(\mathcal{L}_1 \xrightarrow{\beta_1} \mathcal{L}_0, \beta_2, \beta_3 = 0)$ be a strict L_{∞} *H*-pseudoalgebra. Then $(\mathcal{L}_0 \oplus \mathcal{L}_1, [\cdot * \cdot]_{\mathcal{L}_0 \oplus \mathcal{L}_1})$ is a Lie *H*-pseudoalgebra, where

$$[(x,u)*(y,v)]_{\mathcal{L}_0\oplus\mathcal{L}_1} := (\beta_2(x,y), \beta_2(x,v) - (\sigma_{12}\otimes_H 1)\beta(y,u) + \beta_2(\beta_1(u),v)), \text{ for } (x,u), (y,v) \in \mathcal{L}_0 \oplus \mathcal{L}_1.$$

6. CATEGORIFICATION OF LIE H-PSEUDOALGEBRAS

Categorification of Lie algebras, also known as Lie-2 algebras was introduced by Baez and Crans [1] in the study of higher structures. They observed that Lie-2 algebras are closely related to L_{∞} algebras. Among others, they showed that the category of Lie-2 algebras is equivalent to the category of 2-term L_{∞} algebras. In this section, we generalize the above result in the context of Lie *H*-pseudoalgebras. For this, we first introduce Lie-2 *H*-pseudoalgebras as the categorification of Lie *H*-pseudoalgebras and show that the collection of all Lie-2 *H*-pseudoalgebras form a category, which we denote by **Lie2***H*. Finally, we prove that the category **Lie2***H* is equivalent to the category **2shLie***H* of 2-term L_{∞} *H*-pseudoalgebras, hence finds a bridge between categorification and homotopification of Lie *H*-pseudoalgebras.

Let H be a cocommutative Hopf algebra and $\mathcal{M}^{l}(H)$ be the category of left H-modules.

6.1. **Definition.** A left-2 module over H is a category internal to the category $\mathcal{M}^{l}(H)$.

It follows that a left-2 module over H is a category $C = (C_1 \rightrightarrows C_0)$, whose collection of objects C_0 and collection of morphisms C_1 are both left H-modules such that the source and target maps $s, t : C_1 \rightarrow C_0$, the identity-assigning map $i : C_0 \rightarrow C_1$ and the composition map $C_1 \times_{C_0} C_1 \rightarrow C_1$ are all H-linear maps. For any $x \in C_0$, the image i(x) of the identity-assigning map is often denoted by 1_x .

Let $C = (C_1 \Rightarrow C_0)$ and $C' = (C'_1 \Rightarrow C'_0)$ be two left-2 modules over H. A morphism between them is a functor internal to the category $\mathcal{M}^l(H)$. Thus, a morphism from C to C' consists of a pair $F = (F_0, F_1)$ of H-linear maps $F_0: C_0 \to C'_0$ and $F_1: C_1 \to C'_1$ that commute with all the structure maps of the categories C and C'. The collection of all left-2 modules over H and morphisms between them form a category, denoted by $\mathcal{M}^{l_2}(H)$.

On the other hand, a 2-term chain complex over H is a complex of the form $\mathcal{L}_1 \xrightarrow{\partial} \mathcal{L}_0$, where $\mathcal{L}_0, \mathcal{L}_1$ are both left H-modules and ∂ is a H-linear map. A morphism of 2-term chain complexes from $\mathcal{L}_1 \xrightarrow{\partial} \mathcal{L}_0$ to $\mathcal{L}'_1 \xrightarrow{\partial'} \mathcal{L}'_0$ is given by a pair (f_0, f_1) of H-linear maps $f_0 : \mathcal{L}_0 \to \mathcal{L}'_0$ and $f_1 : \mathcal{L}_1 \to \mathcal{L}'_1$ satisfying $\partial' \circ f_1 = f_0 \circ \partial$. The collection of all 2-term chain complexes over H and morphisms between them form a category, denoted by **2Term**H.

Let $C = (C_1 \Rightarrow C_0)$ be a left-2 module over H, hence an object in $\mathcal{M}^{l_2}(H)$. Then ker $(s) \xrightarrow{t_{|ker}(s)} C_0$ is a 2-term chain complex over H. Moreover, a morphism between left-2 modules over H induces a morphism between the corresponding 2-term chain complexes over H. Hence we obtain a functor $\mathcal{S} : \mathcal{M}^{l_2}(H) \to \mathbf{2Term}H$. Conversely, if $\mathcal{L}_1 \xrightarrow{\partial} \mathcal{L}_0$ is a 2-term chain complex over H (hence an object in $\mathbf{2Term}H$) then

$$C := (\mathcal{L}_0 \oplus \mathcal{L}_1 \rightrightarrows \mathcal{L}_0)$$

is a left-2 module over H, where the structure maps are given by

s(x, u) = x, $t(x, u) = x + \partial u$ and i(x) = (x, 0), for $(x, u) \in \mathcal{L}_0 \oplus \mathcal{L}_1, x \in \mathcal{L}_0$.

Similarly, a morphism of 2-term chain complexes gives rise to a morphism between the corresponding left-2 modules over H. As a consequence, we get a functor $\mathcal{T} : \mathbf{2Term} H \to \mathcal{M}^{l2}(H)$. Finally, it is easy to show that the functors \mathcal{S} and \mathcal{T} make the categories $\mathcal{M}^{l2}(H)$ and $\mathbf{2Term} H$ equivalent. See [1] for the similar observation when $H = \mathbf{k}$.

6.2. **Definition.** A Lie-2 H-pseudoalgebra is a triple $(C, [\cdot * \cdot], \mathcal{J})$ that consists of a left-2 module $C = (C_1 \rightrightarrows C_0)$ over H, a $H^{\otimes 2}$ -linear skew-symmetric functor $[\cdot * \cdot] : C \otimes C \to H^{\otimes 2} \otimes_H C$ and a $H^{\otimes 3}$ -linear skew-symmetric natural isomorphism (called the Jacobiator)

$$\mathcal{J}_{x,y,z} : [[x * y] * z] \to [x * [y * z]] - [y * [x * z]]$$

that satisfy the following Jacobiator identity:



Let $(C, [\cdot * \cdot], \mathcal{J})$ and $(C', [\cdot * \cdot]', \mathcal{J}')$ be two Lie-2 *H*-pseudoalgebras. A morphism between them is a functor $F = (F_0, F_1) : C \to C'$ between the underlying left-2 modules over *H* and a natural isomorphism

 $(F_2)^{x,y}: [F_0(x) * F_0(y)]' \to (\mathrm{id}_{H^{\otimes 2}} \otimes_H F_0)[x * y]$

such that for any $x, y, z \in C_0$, the following diagram is commutative

$$\begin{split} & [[F_{0}(x) * F_{0}(y)]' * F_{0}(z)]' \xrightarrow{\mathcal{J}'_{F_{0}(x),F_{0}(y),F_{0}(z)}} [F_{0}(x) * [F_{0}(y) * F_{0}(z)]']' - [F_{0}(y) * [F_{0}(x) * F_{0}(z)]']' \\ & \downarrow [F_{0}(x) * (F_{0})]' \downarrow \qquad \qquad \downarrow [F_{0}(x) * (F_{0})]' - [F_{0}(y) * (F_{0})]' - [F_{0}(y) * (F_{0})]' \\ & \downarrow [F_{0}(x) * (F_{0})]' = [F_{0}(x) * (F_{0})]' - [F_{0}(y) * (F_{0})]' - [F_{0}(y) * (F_{0})]' \\ & \downarrow [F_{0}(x) * (F_{0})]' = [F_{0}(x) * (F_{0})]' - [F_{0}(y) * (F_{0})]' - [F_{0}(y) * (F_{0})]' \\ & \downarrow [F_{0}(x) * (F_{0})]' = [F_{0}(x) * (F_{0})]' - [F_{0}(y) * (F_{0})]' - [F_{0}(y) * (F_{0})]' \\ & \downarrow [F_{0}(x) * (F_{0})]' = [F_{0}(x) * (F_{0})]' \\ & \downarrow [F_{0}(x) * (F_{0})]' = [F_{0}(x) * (F_{0})]' - [F_{0}(y) * (F_{0})]' \\ & \downarrow [F_{0}(x) * (F_{0})]' = [F_{0}(x) * (F_{0})]' \\ & \downarrow [F_{0}(x) * (F_{0})]' = [F_{0}(x) * (F_{0})]' \\ & \downarrow [F_{0}(x) * (F_{0})]' = [F_{0}(x) * (F_{0})]' \\ & \downarrow [F_{0}(x) * (F_{0})]' = [F_{0}(x) * (F_{0})]' \\ & \downarrow [F_{0}(x) * (F_{0})]' = [F_{0}(x) * (F_{0})]' \\ & \downarrow [F_{0}(x) * (F_{0})]' = [F_{0}(x) * (F_{0})]' \\ & \downarrow [F_{0}(x) * (F_{0})]' = [F_{0}(x) * (F_{0})]' \\ & \downarrow [F_{0}(x) * (F_{0})]' \\ & \downarrow [F_{0}(x) * (F_{0})]' = [F_{0}(x) * (F_{0})]' \\ & \downarrow [F_{0}(x) * (F_{0})]' = [F_{0}(x) * (F_{0})]' \\ & \downarrow [F_{0}(x) *$$

We often denote a morphism as above by the triple (F_0, F_1, F_2) .

6.3. **Theorem.** The collection of Lie-2 H-pseudoalgebras and morphisms between them form a category. We denote this category by Lie2H.

Proof. The verification of this result is straightforward once we define the composition of morphisms and the identity morphisms. Let $(C, [\cdot * \cdot], \mathcal{J})$ and $(C', [\cdot * \cdot]', \mathcal{J}')$ be two Lie-2 *H*-pseudoalgebras and (F_0, F_1, F_2) be a morphism from *C* to *C'*. Suppose $(C'', [\cdot * \cdot]'', \mathcal{J}'')$ is another Lie-2 *H*-pseudoalgebra and (G_0, G_1, G_2) is a morphism from *C'* to *C''*. Then their composition is defined by the triple

$$(G_0 \circ F_0, G_1 \circ F_1, (\mathrm{id}_{H^{\otimes 2}} \otimes_H G_0)(F_2^{-,-}) \circ (G_2)^{F_0(-), F_0(-)}),$$

where the last component can be understood by the following composition

$$[(G_0 \circ F_0)(x) * (G_0 \circ F_0)(y)]'' \xrightarrow{(G_2)^{F_0(x), F_0(y)}} (\operatorname{id}_{H^{\otimes 2} \otimes_H} G_0)[F_0(x) * F_0(y)]' \xrightarrow{(\operatorname{id}_{H^{\otimes 2} \otimes_H} G_0)(F_2)^{x,y}} (\operatorname{id}_{H^{\otimes 2} \otimes_H} G_0 \circ F_0)[x * y].$$

Moreover, if $(C, [\cdot * \cdot], \mathcal{J})$ is any Lie-2 *H*-pseudoalgebra, then the identity morphism on *C* is given by the triple $(\mathrm{id}_{C_0}, \mathrm{id}_{C_1}, \mathrm{id}^{-,-})$, where $\mathrm{id}^{x,y} : [x * y] \to [x * y]$ is the identity natural isomorphism. \Box

We are now in a position to prove the main result of this section.

6.4. Theorem. The categories Lie2H and 2shLieH are equivalent.

Proof. Let $(C = (C_1 \rightrightarrows C_0), [\cdot * \cdot], \mathcal{J})$ be a Lie-2 *H*-pseudoalgebra. First consider the 2-term chain complex $\ker(s) \xrightarrow{t|_{\ker(s)}} C_0$ over *H*. We define $H^{\otimes 2}$ -linear maps (both denoted by the same notation)

$$\beta_2: C_0 \otimes C_0 \to H^{\otimes 2} \otimes_H C_0 \text{ and } \beta_2: C_0 \otimes \ker(s) \to H^{\otimes 2} \otimes_H \ker(s)$$

by $\beta_2(x,y) = [x * y]$ and $\beta_2(x,u) = [1_x * u]$, for $x, y \in C_0$ and $u \in \ker(s)$. Another skew-symmetric $H^{\otimes 3}$ -linear map $\beta_3 : C_0 \otimes C_0 \otimes C_0 \to H^{\otimes 3} \otimes_H \ker(s)$ we define by

$$\beta_3(x, y, z) = pr_2 \left(\mathcal{J}_{x, y, z} : [[x * y] * z] \to [x * [y * z]] - [y * [x * z]] \right), \text{ for } x, y, z \in C_0.$$

Here $pr_2: H^{\otimes 3} \otimes_H (C_0 \oplus C_1) \to H^{\otimes 3} \otimes_H C_1$ is the projection onto the second factor. Then similar to the work of Baez and Crans [1], it can be checked that the triple $\mathcal{L} = (\ker(s) \xrightarrow{t|_{\ker(s)}} C_0, \beta_2, \beta_3)$ is a 2-term L_{∞} *H*-pseudoalgebra.

Next, let $(C = (C_1 \Rightarrow C_0), [\cdot * \cdot], \mathcal{J})$ and $(C' = (C'_1 \Rightarrow C'_0), [\cdot * \cdot]', \mathcal{J}')$ be two Lie-2 *H*-pseudoalgebras and (F_0, F_1, F_2) be a morphism between them. Consider the corresponding 2-term L_{∞} *H*-pseudoalgebras $\mathcal{L} = (\ker(s) \xrightarrow{t|_{\ker(s)}} C_0, \beta_2, \beta_3)$ and $\mathcal{L}' = (\ker(s') \xrightarrow{t'|_{\ker(s')}} C'_0, \beta'_2, \beta'_3)$. We define two *H*-linear maps $f_0: C_0 \to C'_0$ and $f_1: \ker(s) \to \ker(s')$, and a $H^{\otimes 2}$ -linear map $f_2: C_0 \otimes C_0 \to H^{\otimes 2} \otimes_H \ker(s')$ by

$$f_0(x) = F_0(x), \ f_1(u) = F_1(u) \text{ and } f_2(x,y) = (F_2)^{x,y} - 1_{s((F_2)^{x,y})} = pr_2((F_2)^{x,y}),$$

for $x, y \in C_0$ and $u \in \ker(s)$. Then a straightforward observation shows that $(f_0, f_1, f_2) : \mathcal{L} \rightsquigarrow \mathcal{L}'$ is a morphism of 2-term L_{∞} *H*-pseudoalgebras. As a consequence, we obtain a functor $S : \text{Lie}2H \rightarrow 2\text{shLie}H$.

In the following, we construct a functor \mathcal{T} : **2shLie** $H \to$ **Lie**2H. We start with a 2-term L_{∞} Hpseudoalgebra ($\mathcal{L}_1 \xrightarrow{\beta_1} \mathcal{L}_0, \beta_2, \beta_3$). First, we consider the left-2 module $C = (\mathcal{L}_0 \oplus \mathcal{L}_1 \rightrightarrows \mathcal{L}_0)$ over H. We define a $H^{\otimes 2}$ -linear skew-symmetric functor $[\cdot * \cdot] : C \otimes C \to H^{\otimes 2} \otimes_H C$ by

$$[(x, u) * (y, v)] := (\beta_2(x, y), \beta_2(x, v) - (\sigma_{12} \otimes_H 1)\beta_2(y, v) + \beta_2(\beta_1(u), v)),$$

for $(x, u), (y, v) \in \mathcal{L}_0 \oplus \mathcal{L}_1$. We define the Jacobiator by

$$\mathcal{J}_{x,y,z} = (\beta_2(\beta_2(x,y),z), \ \beta_3(x,y,z)), \text{ for } x, y, z \in \mathcal{L}_0.$$

It follows from condition (iii) of Definition 5.1 that the Jacobiator $\mathcal{J}_{x,y,z}$ has the source [[x * y] * z] and the target [x * [y * z]] - [y * [x * z]], for $x, y, z \in \mathcal{L}_0$. Note that the Jacobiator is also a natural isomorphism. Finally, a direct calculation (using the conditions of Definition 5.1) yields that the Jacobiator makes the diagram (18) commutative. See [1] for similar calculation. Hence we obtain a Lie-2 *H*-pseudoalgebra $(\mathcal{L}_0 \oplus \mathcal{L}_1 \rightrightarrows \mathcal{L}_0, [\cdot * \cdot], \mathcal{J})$.

Next, let $\mathcal{L} = (\mathcal{L}_1 \xrightarrow{\beta_1} \mathcal{L}_0, \beta_2, \beta_3)$ and $\mathcal{L}' = (\mathcal{L}'_1 \xrightarrow{\beta'_1} \mathcal{L}'_0, \beta'_2, \beta'_3)$ be 2-term L_{∞} *H*-pseudoalgebras and $(f_0, f_1, f_2) : \mathcal{L} \rightsquigarrow \mathcal{L}'$ be a morphism between them. We consider the corresponding Lie-2 *H*-pseudoalgebras

$$C = (\mathcal{L}_0 \oplus \mathcal{L}_1 \rightrightarrows \mathcal{L}_0, [\cdot * \cdot], \mathcal{J}) \text{ and } C' = (\mathcal{L}'_0 \oplus \mathcal{L}'_1 \rightrightarrows \mathcal{L}'_0, [\cdot * \cdot]', \mathcal{J}'), \text{respectively.}$$

We define *H*-linear maps $F_0 : \mathcal{L}_0 \to \mathcal{L}'_0, F_1 : \mathcal{L}_0 \oplus \mathcal{L}_1 \to \mathcal{L}'_0 \oplus \mathcal{L}'_1$ and a natural isomorphism $(F_2)^{x,y} : [F_0(x) * F_0(y)]' \to (\mathrm{id}_{H^{\otimes 2}} \otimes_H F_0)[x * y]$ by

$$F_0 = f_0, \ F_1 = f_0 \oplus f_1 \ \text{and} \ (F_2)^{x,y} = ([F_0(x) * F_0(y)]', f_2(x,y)).$$

Then (F_0, F_1, F_2) turns out to be a morphism of Lie-2 *H*-pseudoalgebras from *C* to *C'*. It follows that we get a functor $\mathcal{T} : \mathbf{2shLie}H \to \mathbf{Lie}2H$.

Finally, we will show that the functors S and \mathcal{T} make the categories **Lie**₂*H* and **2shLie***H* equivalent. In other words, we construct natural isomorphisms $\Lambda : \mathcal{T} \circ S \Rightarrow 1_{\mathbf{Lie}_{2H}}$ and $\Upsilon : S \circ \mathcal{T} \Rightarrow 1_{\mathbf{2shLie}_{H}}$. Let $(C = (C_1 \Rightarrow C_0), [\cdot * \cdot], \mathcal{J})$ be a Lie-2 *H*-pseudoalgebra. If we apply the functor S, we get the 2term L_{∞} *H*-pseudoalgebra (ker(s) $\xrightarrow{t|_{\mathrm{ker}(s)}} C_0, \beta_2, \beta_3$). Again, if we apply the functor \mathcal{T} , we get the Lie-2 *H*-pseudoalgebra

$$(C' = (C_0 \oplus \ker(s) \rightrightarrows C_0), [\cdot * \cdot]', \mathcal{J}').$$

Then it is easy to see that the triple $(\Lambda_0 = \mathrm{id}_{C_0}, \Lambda_1, \Lambda_2)$, where $\Lambda_1 : C_0 \oplus \ker(s) \to C_1, (\Lambda_1)(x, u) = 1_x + u$ and $(\Lambda_2)^{x,y} = \mathrm{id} : [\Lambda_0(x) * \Lambda_0(y)] = [x * y] \to (\mathrm{id}_{H^{\otimes 2}} \otimes_H \Lambda_0)[x * y]' = [x * y]$ is an isomorphism of Lie-2 H-pseudoalgebras from C' to C. This construction yields the natural isomorphism $\Lambda : \mathcal{T} \circ \mathcal{S} \Rightarrow \mathbf{1}_{\mathbf{Lie}2H}$. On the other hand, a direct observation shows that the composition functor $\mathcal{S} \circ \mathcal{T}$ is the identity functor $\mathbf{1}_{\mathbf{2shLie}H}$. Hence we can take $\Upsilon : \mathcal{S} \circ \mathcal{T} \Rightarrow \mathbf{1}_{\mathbf{2shLie}H}$ to be the identity natural isomorphism. This completes the proof.

Acknowledgements. The author would like to thank the Department of Mathematics, IIT Kharagpur for providing the beautiful academic atmosphere where the research has been carried out.

Data Availability Statement. Data sharing does not apply to this article as no new data were created or analyzed in this study.

References

- [1] J. C. Baez and A. S. Crans, Higher-dimensional algebra VI: Lie 2-algebras, Theor. Appl. Categ. 12 (2004), 492-528.
- [2] B. Bakalov, A. D'Andrea and V. G. Kac, Theory of finite pseudoalgebras, Adv. Math. 162 (2001), 1-140.
- [3] B. Bakalov, A. D'Andrea and V. G. Kac, Irreducible modules over finite simple Lie pseudoalgebras I. Primitive pseudoalgebras of type W and S, Adv. Math. 204 (2006), no. 1, 278-346.
- [4] B. Bakalov, A. D'Andrea and V. G. Kac, Irreducible modules over finite simple Lie pseudoalgebras II. Primitive pseudoalgebras of type K, Adv. Math. 232 (2013), no. 1, 188-237.
- [5] A. Beilinson and V. Drinfeld, Chiral algebras, Amer. Math. Soc. Colloq. Publ., 51, American Mathematical Society, Providence, RI (2004) vi+375 pp.
- [6] C. Boyallian and J. I. Liberati, On pseudo-bialgebras, J. Algebra 372 (2012), 1-34.
- [7] E. Cheng and A. Lauda, Higher-Dimensional Categories: an illustrated guide book, University of Sheffield (2004).
- [8] A. D'Andrea and G. Marchei, Representations of Lie pseudoalgebras with coefficients, J. Algebra 329 (2011) 92-106.
- [9] A. Das, Non-abelian cohomology of Lie H-pseudoalgebras and inducibility of automorphisms, arXiv preprint, http://arxiv.org/abs/2312.10341.
- [10] Y. Frégier and M. Zambon, Simultaneous deformations of algebras and morphisms via derived brackets, J. Pure Appl. Algebra 219 (2015), 5344-5362.
- [11] I. M. Gel'fand and I. Ya. Dorfman, Hamiltonian operators and the classical Yang–Baxter equation, Funct. Anal. Appl. 16 (1982), 241–248.
- [12] M. Golenishcheva-Kutuzova and V. G. Kac, Γ-conformal algebras, J. Math. Phys. 39 (1998), 2290-2305.
- [13] V. G. Kac, Vertex algebras for beginners, Univ. Lecture Ser., 10 American Mathematical Society, Providence, RI, 1998. vi+201 pp.
- [14] V. G. Kac, Formal distribution algebras and conformal algebras, https://arxiv.org/abs/q-alg/9709027.
- [15] M. Kontsevich, Deformation quantization of Poisson manifolds, Lett. Math. Phys. 66 (2003), 157-216.

APURBA DAS

- [16] T. Lada and M. Markl, Strongly homotopy Lie algebras, Comm. Algebra 23 (1995), 2147-2161.
- [17] T. Lada and J. Stasheff, Introduction to SH Lie algebras for physicists, Int. J. Theor. Phys. 32 (1993), 1087-1103.
- [18] A. Retakh, Unital associative pseudoalgebras and their representations, J. Algebra 277 (2004), 769-805.
- [19] J. F. Ritt, Associative differential operations, Ann. of Math. (2) 51 (1950), 756-765.
- [20] C. L. Rogers, 2-plectic geometry, Courant algebroids, and categorified prequantization, J. Symplectic Geom. 11 (2013), 3869-3897.
- [21] A. Sahoo and A. Das, Homotopy conformal algebras, https://doi.org/10.48550/arXiv.2308.05951
- [22] J. D. Stasheff, Homotopy associativity of H-spaces. I, II, Trans. Amer. Math. Soc. 108 (1963), 275-312.
- [23] Q. Sun, A note on graded Lie H-pseudobialgebras, Indian J. Pure Appl. Math. 49 (1) (2018), 1-21.
- [24] Z. Wu, Lie algebra structures on cohomology complexes of some H-pseudoalgebras, J. Algebra 396 (2013), 117-142.
- [25] Z. Wu, Graded left symmetric pseudo-algebras, Comm. Algebra 43 (2015), 3869-3897.
- [26] Z. Wu, Leibniz H-pseudoalgebras, J. Algebra 437 (2015), 1-33.

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY, KHARAGPUR 721302, WEST BENGAL, INDIA. Email address: apurbadas3480gmail.com, apurbadas3480gmaths.iitkgp.ac.in