THE SCHUR POLYNOMIALS IN ALL PRIMITIVE *n*TH ROOTS OF UNITY

MASAKI HIDAKA AND MINORU ITOH

ABSTRACT. We show that the Schur polynomials in all primitive nth roots of unity are 1, 0, or -1, if n has at most two distinct odd prime factors. This result can be regarded as a generalization of properties of the coefficients of the cyclotomic polynomial and its multiplicative inverse. The key to the proof is the notion of a unimodular system of vectors. Namely, this result is reduced in turn to four propositions on the unimodularity of vector systems, and the last proposition is proved by using graph theory.

1. INTRODUCTION

The following assertion on the Schur polynomials in all primitive nth roots of unity is the main theorem of this article¹:

Theorem 1.1. Let $\omega_1, \ldots, \omega_d$ be all primitive *n*th roots of unity (thus *d* is equal to $\varphi(n)$, where φ is Euler's totient function), and λ be a partition whose length is at most *d*. Moreover, we assume the following condition on *n*:

(*) n has at most two distinct odd prime factors.

Then, we have

$$s_{\lambda}(\omega_1, \dots, \omega_d) = 1, 0, \text{ or } -1.$$

Here, s_{λ} is the Schur polynomial associated to λ .

The condition (*) holds for many natural numbers. For example, all natural numbers less than $105 = 3 \cdot 5 \cdot 7$ satisfy (*).

Theorem 1.1 has been known for $\lambda = (1^k)$ and (k) as properties of the coefficients of the cyclotomic polynomial and its multiplicative inverse.

First, when $\lambda = (1^k)$, the Schur polynomial associated with λ equals the kth elementary symmetric polynomial e_k . Thus, we have

$$s_{\lambda}(\omega_1,\ldots,\omega_d)=e_k(\omega_1,\ldots,\omega_d),$$

and this equals the coefficient of x^{d-k} in the cyclotomic polynomial $\Phi_n(x)$ (up to sign), because

$$\Phi_n(x) = (x - \omega_1) \cdots (x - \omega_d).$$

As is well known, A. Migotti [Mi] showed that the coefficients of $\Phi_n(x)$ are all in the set $\{1, 0, -1\}$, if n satisfies (*).

²⁰¹⁰ Mathematics Subject Classification. 05E05, 05C50, 11B83, 11C08, 11C20, 11R18.

Key words and phrases. primitive root of unity, cyclotomic polynomial, Schur polynomial, unimodular system, graph theory, matroid.

This research was partially supported by JSPS Grant-in-Aid for Scientific Research (C) 16K05067 and 21K03209.

¹This theorem was first found in the Master's thesis of the first author [H].

Secondly, when $\lambda = (k)$, the Schur polynomial associated with λ equals the kth complete homogeneous symmetric polynomial h_k . Thus, we have

$$s_{\lambda}(\omega_1,\ldots,\omega_d) = h_k(\omega_1,\ldots,\omega_d)$$

and this equals the coefficient of x^k in $\Phi_n(x)^{-1}$. Indeed, we have

$$\Phi_n(x)^{-1} = (x - \omega_1)^{-1} \cdots (x - \omega_d)^{-1}$$

= $(-)^d \omega_1 \cdots \omega_d (1 - x \omega_1^{-1})^{-1} \cdots (1 - x \omega_d^{-1})^{-1}$
= $\sum_{k \ge 0} x^k h_k(\omega_1^{-1}, \dots, \omega_d^{-1})$
= $\sum_{k \ge 0} x^k h_k(\omega_1, \dots, \omega_d),$

because $(-)^d \omega_1 \cdots \omega_d = 1$ and $h_k(\omega_1^{-1}, \ldots, \omega_d^{-1}) = h_k(\omega_1, \ldots, \omega_d)$. P. Moree [Mo] showed that the coefficients of $\Phi_n(x)^{-1}$ are all in the set $\{1, 0, -1\}$, if n satisfies $(*)^2$.

Theorem 1.1 is a generalization of these two results.

The key to the proof is the notion of a unimodular system of vectors. Theorem 1.1 is reduced in turn to four propositions on the unimodularity of vector systems, and the last proposition is proved by using graph theory.

2. Unimodular systems

The key to the proof of Theorem 1.1 is the notion of a unimodular system of vectors ([DG]). We recall this notion in this section. Let K be a field of characteristic 0, and V an *n*-dimensional K-vector space.

Definition 2.1. Let X be a finite subset of V satisfying $0 \notin X$ and $\langle X \rangle = V$. We say that X is a unimodular system of V, if for any basis $B \subset X$ the determinant of B does not depend on B.

Here, the meaning of the determinant of B is as follows. We can express B as an n by n matrix through a linear isomorphism $f: V \to K^n$, so that we can define det B excluding the sign through this correspondence (an ambiguity of sign caused by the order of the elements of B). Let us put $[a] = \{a, -a\}$ (this is the equivalent class determined by identifying two scalars equal up to sign). From the above, we can determine $[\det B]$ relative to f. Definition 2.1 does not depend on the choice of f.

Remark. The definition of unimodular system in [DG] does not require the conditions $0 \notin X$ or $\langle X \rangle = V$.

Proposition 2.2. The following conditions are equivalent:

(1) For any basis $B \subset X$, the determinant det B does not depend on B.

(2) For any basis $B \subset X$, the set $\mathbb{Z}B$ does not depend on B.

(3) X can be identified with the columns of a totally unimodular matrix through a linear isomorphism $V \to K^n$.

Here, a totally unimodular matrix means a matrix for which the determinant of every square submatrix is 1, 0, or -1.

The root systems of type A are a typical example of a unimodular system (in fact, this is a maximal unimodular system).

²Interestingly, the proof in the case $\lambda = (k)$ is easier than that in the case $\lambda = (1^k)$.

3. Four propositions on the unimodularity of vector systems

In this section, we state four propositions on the unimodularity of vector systems (Propositions 3.1–3.4). The main theorem is reduced to these four propositions sequentially:

Theorem $1.1 \Leftarrow$ Proposition $3.1 \Leftarrow$ Proposition $3.2 \Leftarrow$ Proposition $3.3 \Leftarrow$ Proposition 3.4.

After all, to prove the main theorem, it suffices to show Proposition 3.4.

The four propositions are as follows:

Proposition 3.1. When n satisfies (*), the following set is a unimodular system of \mathbb{C}^d :

$$\Omega_n = \left\{ \left. \begin{pmatrix} \omega_1^k \\ \vdots \\ \omega_d^k \end{pmatrix} \right| \, k = 0, 1, \dots, n-1 \right\}.$$

Proposition 3.2. When n satisfies (*), the following set is a unimodular system of $\mathbb{Q}(\zeta_n)$:

$$Z_n = \left\{ z \in \mathbb{C} \mid z^n = 1 \right\}.$$

Here, ζ_n is a primitive *n*th root of unity, and we regard the cyclotomic field $\mathbb{Q}(\zeta_n)$ as a *d*-dimensional \mathbb{Q} -vector space.

Proposition 3.3. (1) If p is an odd prime, Z_p is a unimodular system of $\mathbb{Q}(\zeta_p)$. (2) If p and q are odd primes,

$$Z_p \otimes Z_q = \left\{ x \otimes y \, \big| \, x \in Z_p, \ y \in Z_q \right\}$$

is a unimodular system of $\mathbb{Q}(\zeta_p) \otimes \mathbb{Q}(\zeta_q)$, where " \otimes " means the tensor product of two \mathbb{Q} -vector spaces.

Proposition 3.4. (1) If X is a fundamental zero-sum system of V, then X is a unimodular system of V.

(2) If X and Y are fundamental zero-sum systems of V and W, respectively, then

$$X \otimes Y = \left\{ x \otimes y \, \big| \, x \in X, \ y \in Y \right\}$$

is a unimodular system of $V \otimes W$.

Here, we define the notion of a fundamental zero-sum system as follows:

Definition 3.5. For a finite subset X of a finite dimensional \mathbb{Q} -vector space V, we say that X is a fundamental zero-sum system of V, when the following conditions hold:

$$|X| = \dim(V) + 1, \qquad \langle X \rangle = V, \qquad \sum X = 0.$$

Here, we put $\sum X = \sum_{x \in X} x$.

For example, if $\{e_1, \ldots, e_n\}$ is a basis of V,

$$\left\{e_1,\ldots,e_n,-\sum_{i=1}^n e_i\right\}$$

is a fundamental zero-sum system of V. Conversely, any fundamental zero-sum system of V can be expressed in this form. Hence, any two fundamental zero-sum systems of V are interchanged by a linear automorphism.

Moreover, when p is an odd prime, Z_p is a fundamental zero-sum system of the Q-vector space $\mathbb{Q}(\zeta_p)$. Hence, Proposition 3.4 is a generalization of Proposition 3.3.

In the next section, we will explain the reduction of the main theorem Proposition 3.4 through Propositions 3.1, 3.2, and 3.3. Moreover, we prove Proposition 3.4 in Section 5.

4. Reduction of the main theorem to Proposition 3.4

In this section, we explain the reduction of the main theorem to Proposition 3.4 through Propositions 3.1, 3.2, and 3.3.

4.1. Theorem 1.1 \leftarrow **Proposition 3.1.** First, Theorem 1.1 is reduced to Proposition 3.1. Indeed, using Proposition 3.1, we can prove Theorem 1.1 as follows.

The Schur polynomial s_{λ} is expressed as

$$s_{\lambda}(x_1,\ldots,x_d) = a_{\delta+\lambda}(x_1,\ldots,x_d)/a_{\delta}(x_1,\ldots,x_d).$$

Here, a_{μ} is a Vandermonde type determinant defined by

$$a_{\mu}(x_1,\ldots,x_d) = \det(x_i^{\mu_j})_{1 \le i,j \le d}$$

for $\mu = (\mu_1, \ldots, \mu_d) \in \mathbb{Z}_{\geq 0}^d$. Moreover, we put

$$\delta = (d - 1, d - 2, \dots, 1, 0).$$

For any $\mu \in \mathbb{Z}_{>0}^d$, there exist $v_1, \ldots, v_d \in \Omega_n$ such that

$$a_{\mu}(\omega_1,\ldots,\omega_d) = \det(v_1,\ldots,v_d).$$

By Proposition 3.1, when n satisfies (*), there exists a nonzero complex number a satisfying

$$\left\{a_{\mu}(\omega_{1},\ldots,\omega_{d}) \mid \mu \in \mathbb{Z}_{\geq 0}^{d}\right\} = \left\{\det(v_{1},\ldots,v_{d}) \mid v_{1},\ldots,v_{d} \in \Omega_{n}\right\} = \{a,0,-a\}.$$

Moreover, we see $a_{\delta}(\omega_1, \ldots, \omega_d) \neq 0$ easily. Theorem 1.1 is immediate from this.

4.2. Proposition $3.1 \leftarrow$ Proposition 3.2. Proposition 3.1 can be reduced to Proposition 3.2 through a natural linear isomorphism as follows.

To prove Proposition 3.1, it suffices to show that Ω_n is a unimodular system of $\langle \Omega_n \rangle$, the \mathbb{Q} -vector space generated by Ω_n . We note that $\langle \Omega_n \rangle$ is isomorphic to the cyclotomic field $\mathbb{Q}(\zeta_n)$ (as \mathbb{Q} -vector spaces) through the correspondence

$$\begin{pmatrix} z_1 \\ \vdots \\ z_d \end{pmatrix} \mapsto z_1.$$

Moreover, Ω_n is identified with Z_n through this isomorphism. Thus, Proposition 3.1 is equivalent to Proposition 3.2.

4.3. Proposition 3.2 \leftarrow **Proposition 3.3.** Proposition 3.2 can also be reduced to Proposition 3.3 through a natural linear isomorphism.

First we note the following lemma:

Lemma 4.1. When X and Y are unimodular systems of V and W, respectively, $X \sqcup Y$ is a unimodular system of $V \oplus W$.

Next, when we have $n = p_1^{l_1} \cdots p_k^{l_k}$ where p_1, \ldots, p_k are distinct primes, we can identify $\mathbb{Q}(\zeta_n)$ with the $p_1^{l_1-1} \cdots p_k^{l_k-1}$ -fold direct sum of

$$\mathbb{Q}(\zeta_{p_1})\otimes\cdots\otimes\mathbb{Q}(\zeta_{p_k})$$

as \mathbb{Q} -vector spaces. Moreover, through this isomorphism, we can identify Z_n with the $p_1^{l_1-1}\cdots p_k^{l_k-1}$ -fold disjoint sum of

$$Z_{p_1} \otimes \cdots \otimes Z_{p_k}.$$

This follows from (1) and (2) of the following lemma:

Lemma 4.2. (1) When natural numbers a and b are coprime, there exists a linear isomorphism $\mathbb{Q}(\zeta_{ab}) \to \mathbb{Q}(\zeta_{a}) \otimes \mathbb{Q}(\zeta_{b})$ such that the image of Z_{ab} is equal to $Z_a \otimes Z_b$. (2) For any prime p, there exists a linear isomorphism $\mathbb{Q}(\zeta_{p^l}) \to \mathbb{Q}(\zeta_p)^{\oplus p^{l-1}}$ such that the image of Z_{p^l} is equal to the p^{l-1} -fold disjoint sum of Z_p .

Proof. (1) We consider the following correspondence:

$$\mathbb{Q}(\zeta_a) \otimes \mathbb{Q}(\zeta_b) \to \mathbb{Q}(\zeta_{ab}), \qquad z \otimes w \mapsto zw.$$

This gives a linear isomorphism, and the image of $Z_a \otimes Z_b$ is equal to Z_{ab} . (2) Let $\mathbb{Q}(\zeta_p)^{(j)}$ denote a copy of $\mathbb{Q}(\zeta_p)$ for $j \in 0, 1, \ldots, p^{l-1} - 1$. Moreover, we denote by $z^{(j)}$ the counterpart of $z \in \mathbb{Q}(\zeta_p)$ in $\mathbb{Q}(\zeta_p)^{(j)}$. Let us consider the following correspondence:

$$\bigoplus_{j=0}^{p^{l-1}-1} \mathbb{Q}(\zeta_p)^{(j)} \to \mathbb{Q}(\zeta_{p^l}), \qquad z^{(j)} \mapsto \zeta_{p^l}^j z.$$

This gives a linear isomorphism (it suffices to show the surjectiveness because the dimensions are equal). It is obvious that the image of $\bigsqcup_{j=0}^{p^{l-1}-1} Z_p^{(j)}$ is equal to Z_{p^l} .

Thus, we have the following isomorphism, because $\mathbb{Q}(\zeta_2) = \mathbb{Q}$:

$$\mathbb{Q}(\zeta_n) \simeq \begin{cases} \bigoplus_{i \in \Lambda} \mathbb{Q}^{(i)} \text{ (where } |\Lambda| = 2^{k-1}), & n = 2^k, \\ \bigoplus_{i \in \Lambda} \mathbb{Q}(\zeta_p)^{(i)} \text{ (where } |\Lambda| = p^{l-1}), & n = p^l, \\ \bigoplus_{i \in \Lambda} \mathbb{Q}(\zeta_p)^{(i)} \text{ (where } |\Lambda| = 2^{k-1}p^{l-1}), & n = 2^kp^l, \\ \bigoplus_{i \in \Lambda} (\mathbb{Q}(\zeta_p) \otimes \mathbb{Q}(\zeta_q))^{(i)} \text{ (where } |\Lambda| = p^{l-1}q^{m-1}), & n = p^lq^m, \\ \bigoplus_{i \in \Lambda} (\mathbb{Q}(\zeta_p) \otimes \mathbb{Q}(\zeta_q))^{(i)} \text{ (where } |\Lambda| = 2^{k-1}p^{l-1}q^{m-1}), & n = 2^kp^lq^m. \end{cases}$$

Here, p and q are odd primes. Through this isomorphism f, we can write the image $f(Z_n)$ as

$$f(Z_n) = \begin{cases} \bigsqcup_{i \in \Lambda} Z_2^{(i)}, & n = 2^k, \\ \bigsqcup_{i \in \Lambda} Z_p^{(i)}, & n = p^l, \\ \bigsqcup_{i \in \Lambda} (Z_2 \otimes Z_p)^{(i)}, & n = 2^k p^l, \\ \bigsqcup_{i \in \Lambda} (Z_p \otimes Z_q)^{(i)}, & n = p^l q^m, \\ \bigsqcup_{i \in \Lambda} (Z_2 \otimes Z_p \otimes Z_q)^{(i)}, & n = 2^k p^l q^m. \end{cases}$$

Thus, we see that Proposition 3.2 follows from Proposition 3.3. Indeed, we have $Z_2 = \{1, -1\}$ and the following lemma:

Lemma 4.3. If X is a unimodular system of V, then $\{1, -1\} \otimes X$ is also a unimodular system of V.

4.4. Proposition 3.3 \leftarrow Proposition 3.4. When *p* is an odd prime, Z_p is a fundamental zero-sum system of the Q-vector space $\mathbb{Q}(\zeta_p)$. Thus, Proposition 3.4 is a generalization of Proposition 3.3.

5. Proof of Proposition 3.4

The main theorem has been reduced to Proposition 3.4. In this section, we prove this Proposition 3.4. First, we prove Proposition 3.4 (1) in Section 5.1 and Proposition 3.4 (2) using graph theory in Sections 5.2 and 5.3.

5.1. Any fundamental zero-sum system is unimodular. First, we prove Proposition 3.4 (1). Namely, we show that any fundamental zero-sum system is unimodular.

Proof of Proposition 3.4 (1). Let X be a fundamental zero-sum system of V. We take two bases $X_1, X_2 \subset X$, and assume that $X_1 \neq X_2$. Then, there exist two different elements $x_1, x_2 \in X$ satisfying

$$X_1 = X \setminus \{x_1\} = X_0 \sqcup \{x_2\}, \qquad X_2 = X \setminus \{x_2\} = X_0 \sqcup \{x_1\}.$$

Here, we put $X_0 = X \setminus \{x_1, x_2\}$. Thus, we have $[\det X_1] = [\det X_2]$, because

$$[\det X_1] = [\det(X_0 \sqcup \{x_2\})]$$

= $[\det(X_0 \sqcup \{x_2 + \sum X_0\})]$
= $[\det(X_0 \sqcup \{-x_1\})]$
= $[\det(X_0 \sqcup \{x_1\})]$
= $[\det X_2].$

Here, the third equality is obtained by an elementary operation. Moreover, the fourth equality follows from

$$x_1 + x_2 + \sum X_0 = \sum X = 0.$$

5.2. Reduction to a claim on an equivalence relation. In the remainder of this section, we prove Proposition 3.4 (2). Namely, we show that if X and Y are fundamental zero-sum systems of V and W, respectively, $X \otimes Y$ is a unimodular system of $V \otimes W$.

We put $M = \dim V$ and $N = \dim W$, and consider the following two families of subsets of $X \otimes Y$:

$$\mathcal{A} = \{ A \subset X \otimes Y \mid |A| = MN \},\$$
$$\mathcal{B} = \{ A \subset X \otimes Y \mid A \text{ is a basis of } V \otimes W \}.$$

Note that $\mathcal{B} \subset \mathcal{A}$.

Let us consider an equivalence relation on \mathcal{A} satisfying

$$A_1 \sim A_2 \Rightarrow [\det A_1] = [\det A_2].$$

To introduce this, we first put

$$\mathcal{H} = \{ X \otimes y \, \big| \, y \in Y \} \cup \{ x \otimes Y \, \big| \, x \in X \},\$$

where

$$X \otimes y = \{ x \otimes y \mid x \in X \}, \qquad x \otimes Y = \{ x \otimes y \mid y \in Y \}.$$

Then, we have $\sum H = 0$ for any $H \in \mathcal{H}$. Indeed when $H = X \otimes y$, we have

$$\sum H = \sum (X \otimes y) = (\sum X) \otimes y = 0 \otimes y = 0$$

The proof for $H = x \otimes Y$ is almost identical. For $A_1, A_2 \in \mathcal{A}$, we define $A_1 \sim A_2$ to mean that there exist $H \in \mathcal{H}$ and $a_1, a_2 \in H$ satisfying

$$H \subset A_1 \sqcup \{a_2\} = A_2 \sqcup \{a_1\}$$

Example 5.1. When $X = \{x_1, x_2, x_3\}$ and $Y = \{y_1, y_2, y_3\}$, we have

$$\mathcal{H} = \left\{ X \otimes y_1, X \otimes y_2, X \otimes y_3, x_1 \otimes Y, x_2 \otimes Y, x_3 \otimes Y \right\}.$$

Let us put

$$A_1 = \{(x_2, y_2), (x_2, y_3), (x_3, y_2), (x_3, y_3)\}, \qquad A_2 = \{(x_2, y_2), (x_2, y_2), (x_3, y_2), (x_3, y_1)\}.$$

Then, we have $A_1 \sim A_2$, because putting $H = x_3 \otimes Y$ we have

$$H \subset A_1 \sqcup \{(x_3, y_1)\} = A_2 \sqcup \{(x_3, y_3)\}$$

Lemma 5.2. If $A_1 \sim A_2$, we have $[\det A_1] = [\det A_2]$.

Proof. Since $A_1 \sim A_2$, there exist $H \in \mathcal{H}$ and $a_1, a_2 \in H$ satisfying

$$H \subset A_1 \sqcup \{a_2\} = A_2 \sqcup \{a_1\}.$$

We put

$$A_0 = A_1 \smallsetminus \{a_1\} = A_2 \smallsetminus \{a_1\},$$

so that

$$[\det A_1] = [\det(A_0 \sqcup \{a_1\})]$$

= $[\det(A_0 \sqcup \{a_1 + \sum (H \smallsetminus \{a_1, a_2\})\})]$
= $[\det(A_0 \sqcup \{-a_2\})]$
= $[\det(A_0 \sqcup \{a_2\})]$
= $[\det A_2].$

Here, the second equality is obtained by an elementary operation by noting $H \setminus \{a_1, a_2\} \subset A_0$. Moreover, the third equality follows from

$$a_1 + a_2 + \sum (H \setminus \{a_1, a_2\}) = \sum H = 0.$$

Let \sim denote the equivalence relation on \mathcal{A} generated by the relation \sim . Then, the following is apparent from the discussion so far:

Proposition 5.3. For $A_1, A_2 \in \mathcal{A}$ satisfying $A_1 \sim A_2$, we have $[\det A_1] = [\det A_2]$. In particular, the linear independence of A_1 is equivalent to that of A_2 .

Thus, to prove that $X \otimes Y$ is unimodular, it suffices to show the following proposition:

Proposition 5.4. \mathcal{B} is an equivalence class under the relation \sim .

5.3. Graph theory. We can rephrase Proposition 5.4 neatly using the terminology of graph theory.

Henceforth, we discuss the following families of sets instead of \mathcal{A} and \mathcal{B} :

$$\mathcal{A}' = \{ A^c \, \big| \, A \in \mathcal{A} \}, \qquad \mathcal{B}' = \{ A^c \, \big| \, A \in \mathcal{B} \}.$$

Here, we put $A^c = X \otimes Y \smallsetminus A$. Since

$$|X \otimes Y| = (M+1)(N+1) = MN + M + N + 1,$$

 \mathcal{A}' is the family of all (M + N + 1)-point subsets of $X \otimes Y$:

$$\mathcal{A}' = \left\{ A \subset X \otimes Y \, \big| \, |A| = M + N + 1 \right\}.$$

We can define a natural equivalence relation \approx on \mathcal{A}' . Namely, for $A_1, A_2 \in \mathcal{A}$, we define $A_1^c \approx A_2^c$ to mean that $A_1 \sim A_2$. In other words, this relation on \mathcal{A}' is the equivalence relation generated by the relation \approx on \mathcal{A}' . Here, we define $A_1^c \approx A_2^c$ to mean that $A_1 \sim A_2$, namely, there exist $H \in \mathcal{H}$ and $a_1, a_2 \in H$ satisfying

(5.1)
$$H \cap A_1^c = \{a_1\}, \quad H \cap A_2^c = \{a_2\}, \quad A_1^c \smallsetminus \{a_1\} = A_2^c \smallsetminus \{a_2\}.$$

The final equality is equivalent to $A_2^c = (A_1^c \setminus \{a_1\}) \sqcup \{a_2\}.$

Thus, to prove Proposition 5.4, it suffices to show the following:

Proposition 5.5. \mathcal{B}' is an equivalence class under the relation \approx on \mathcal{A}' .

The terminology of graph theory is useful to discuss this. We identify $X \otimes Y$ with the complete bipartite graph with bipartition (X, Y) (from now on, we fix the vertex set, and we regard a graph as a set of edges). Thus, \mathcal{A}' is equal to the set of all bipartite graphs with M + N + 1 edges.

We call a vertex of degree 1 a leaf (and the edge meeting a leaf is called a leaf edge). Namely, $x \in X$ is a leaf of $E \in \mathcal{A}'$, if and only if there exists a unique $y \in Y$ satisfying $(x, y) \in E$. Similarly $y \in Y$ is a leaf of B, if and only if there exists a unique $x \in Y$ satisfying $(x, y) \in E$.

Using the term "leaf," we can state the definition of the relation \approx on \mathcal{A}' elegantly. Indeed, for $E_1, E_2 \in \mathcal{A}'$, we have $E_1 \approx E_2$, if and only if there exists $v \in X \sqcup Y$ satisfying

- v is both a leaf of E_1 and a leaf of E_2 .
- we have $E_1 \setminus \{a_1\} = E_2 \setminus \{a_2\}$, where a_1 is the unique leaf edge of E_1 meeting v, and a_2 is the unique leaf edge of E_2 meeting v.

Namely, we can say that $E_1 \approx E_2$, if E_2 is obtained from E_1 by an exchange of leaf edges.

Example 5.6. Let us revisit Example 5.1. We put $E_1 = A_1^c$ and $E_2 = A_2^c$. Then, we have

$$E_1 = \{(x_1, y_1), (x_1, y_2), (x_1, y_3), (x_2, y_1), (x_3, y_1)\},\$$

$$E_2 = \{(x_1, y_1), (x_1, y_2), (x_1, y_3), (x_2, y_1), (x_3, y_3)\}.$$

Since we have previously shown $A_1 \sim A_2$, we have $E_1 \approx E_2$. Of course, the condition (5.1) holds by putting

$$H = x_3 \otimes Y,$$
 $a_1 = (x_3, y_1),$ $a_2 = (x_3, y_3).$

As graphs, we can express E_1 and E_2 as follows (we denote by \bullet and \circ the elements of X and Y, respectively):



These are certainly trees. Moreover, E_2 is obtained from E_1 by an exchange of leaf edges as follows:

$$E_2 = (E_1 \smallsetminus \{(x_3, y_1)\}) \sqcup \{(x_3, y_3)\}.$$

The following relation is a key of the proof of Proposition 5.5:

Proposition 5.7. We have $\mathcal{B}' = \mathcal{T}$. Here, \mathcal{T} is the set of all bipartite graphs with bipartition (X, Y) that are trees.

Before proving Propositions 5.5 and 5.7, we recall the following fundamental fact on trees ([D]):

Lemma 5.8. For a graph E with an r-point vertex set, the following three conditions are equivalent:

- (1) E is a tree.
- (2) E is connected, and |E| = r 1.
- (3) E has no cycle, and |E| = r 1.

Therefore, \mathcal{T} is a subset of \mathcal{A}' . The following lemma is also immediate from this:

Lemma 5.9. Let *E* be a nonconnected graph with an *r*-point vertex set. If |E| = r - 1, a connected component with no cycles exists.

From these lemmas, we see the following proposition on the equivalence relation \approx on \mathcal{A}' :

Proposition 5.10. Any graph in \mathcal{A}' having a cycle is equivalent to a graph with isolated vertex with respect to \approx .

Proof. Let us take any graph $E_1 \in \mathcal{A}'$, and assume that this has a cycle but no isolated vertex. By Lemmas 5.8 and 5.9, there exists a connected component $E'_1 \subsetneq E_1$, and we can assume that E'_1 has no cycle. Then, E'_1 has a leaf v_1 . Since $E_1 \setminus E'_1 \neq \emptyset$, we can take an edge $(x_1, y_1) \in E_1 \setminus E'_1$. Noting this, we put

$$b_1 = \begin{cases} (v_1, y_1), & v_1 \in X, \\ (x_1, v_1), & v_1 \in Y. \end{cases}$$

Moreover, we denote by a_1 the unique edge of E'_1 meeting v_1 . Using these, we put

$$E_2 = (E_1 \setminus \{a_1\}) \sqcup \{b_1\}, \qquad E'_2 = E'_1 \setminus \{a_1\}.$$

Then, we have $E_2 \approx E_1$. If $E'_2 = \emptyset$, E_2 has an isolated vertex. Thus, let us assume that $E'_2 \neq \emptyset$. Then, E'_2 is a connected component of E_2 having no cycle. Hence, we can construct E_3 from E_2 similarly. Then, the sequence $|E'_1|, |E'_2|, \ldots$ decreases by one, so that there exists some $k \in \mathbb{N}$ satisfying $E'_k = \emptyset$.

Using this proposition, we see one of the inclusions in Proposition 5.7 as follows:

Proof of (\subset) of Proposition 5.7. Take an $E \in \mathcal{B}'$, and assume that $E \notin \mathcal{T}$. Then, E has a cycle by Lemma 5.8. By Proposition 5.10, E is equivalent to a graph $R \in \mathcal{A}'$ with an isolated vertex. Thus, there exists $H \in \mathcal{H}$ satisfying $H^c \supset R$, namely, $H \subset R^c$. Since His linearly dependent, $R^c \in \mathcal{A}$ is also linearly dependent. Thus, by Proposition 5.3, E^c is also linearly dependent. This contradicts with $E^c \in \mathcal{B}$.

Now that we know $\mathcal{B}' \subset \mathcal{T}$, to prove Proposition 5.5, it suffices to show that any two elements of \mathcal{T} (namely any two trees in \mathcal{A}') are equivalent with respect to \approx . This is shown as follows (this also proves (\supset) of Proposition 5.7).

We consider the following subset of \mathcal{T} :

$$\mathcal{S} = \{ (x \otimes Y) \cup (X \otimes y) \mid x \in X, \ y \in Y \}.$$

To prove Proposition 5.5, it suffices to show the following two lemmas:

Lemma 5.11. Any element of \mathcal{T} is equivalent to an element in \mathcal{S} . Indeed, for $E \in \mathcal{T}$, we have $E \approx (x \otimes Y) \cup (X \otimes y)$, if $(x, y) \in E$.

Lemma 5.12. Any two elements in \mathcal{S} are equivalent to each other.

Proof of Lemma 5.11. We take an $E_1 \in \mathcal{T}$. We fix an edge $(x_{\infty}, y_{\infty}) \in E_1$ and put $E_{\infty} = (x_{\infty} \otimes Y) \cup (X \otimes y_{\infty})$. Then, we have $E_1 \approx E_{\infty}$. To prove this, we consider the following map:

$$f: \mathcal{A}' \to \{0, 1, \dots, M + N + 1\}, \\ E \mapsto \left| \left\{ v \in X \mid (v, y_{\infty}) \in E \right\} \right| + \left| \left\{ v \in Y \mid (x_{\infty}, v) \in E \right\} \right|.$$

Then, E_{∞} is the unique element of \mathcal{A}' for which the value of f is M + N + 1. If $E_1 \neq E_{\infty}$, we construct $E_2 \in \mathcal{T}$ from E_1 as follows. Since $E_1 \neq E_{\infty}$, we can take a leaf $v_1 \in E_1$ that is not connected to either x_{∞} or y_{∞} . Noting this, we put

$$E_2 = (E_1 \smallsetminus \{a_1\}) \sqcup \{b_1\},\$$

where a_1 is the unique leaf edge of E_1 meeting v_1 , and b_1 is the edge defined by

$$b_1 = \begin{cases} (v_1, y_{\infty}), & v_1 \in X, \\ (x_{\infty}, v_1), & v_1 \in Y. \end{cases}$$

Then, we have $E_2 \approx E_1$. Moreover, we see $f(E_2) = f(E_1) + 1$. If $E_2 \neq E_{\infty}$, we can construct E_3 from E_2 similarly. Then, the sequence $f(E_1), f(E_2), \ldots$ increases by one, so that there exists some $k \in \mathbb{N}$ satisfying $E_k = E_{\infty}$.

Proof of Lemma 5.12. Take any $x_1, x_2 \in X$ and $y_1, y_2 \in Y$. Then, we have $(x_1, y_1), (x_2, y_2) \in (x_2 \otimes Y) \cup (X \otimes y_1)$. Hence, by Lemma 5.11, we have

$$(x_1 \otimes Y) \cup (X \otimes y_1) \approx (x_2 \otimes Y) \cup (X \otimes y_1) \approx (x_2 \otimes Y) \cup (X \otimes y_2).$$

6. The root system of type A is unimodular

As explained in Section 2, the root system of type A_n is unimodular. This fact can be proved in a way similar to the proof of Proposition 3.4 (2), but more easily. In this section, we will state this proof roughly. This will help us to better understand the mechanics of the discussion in the previous section.

We denote by Φ_n the root system of type A_n . Let us identify any vector $v \in \Phi_n$ with -v (this identification has no adverse effect on discussing whether Φ_n is unimodular or not). Then, we can regard Φ_n as a set of size n(n+1)/2. We identify this with the complete graph K_{n+1} with n+1 vertices (as in the previous section, we regard a graph as a set of edges). We put

$$\hat{\mathcal{A}} = \left\{ A \subset K_{n+1} \, \middle| \, |A| = n \right\}.$$

Moreover, we put

$$\tilde{\mathcal{B}} = \{ A \in \tilde{\mathcal{A}} \mid A \text{ is a basis of } \langle \Phi_n \rangle \}, \\
\tilde{\mathcal{T}} = \{ A \in \tilde{\mathcal{A}} \mid A \text{ is a tree} \}, \\
\tilde{\mathcal{H}} = \{ A \subset K_{n+1} \mid A \text{ is a cycle} \}, \\
\tilde{\mathcal{S}} = \{ A \in \tilde{\mathcal{A}} \mid A \text{ is a star} \}.$$

We consider an equivalence relation \approx on \mathcal{A} generated by exchanges of leaf edges. In other words, this is the equivalence relation generated by the relation \approx on $\tilde{\mathcal{A}}$. Here, we define $A_1 \approx A_2$ to mean that there exist $H \in \mathcal{H}$ and $a_1, a_2 \in H$ satisfying

$$H \cap A_1^c = \{a_1\}, \qquad H \cap A_2^c = \{a_2\}, \qquad A_1^c \smallsetminus \{a_1\} = A_2^c \smallsetminus \{a_2\}.$$

Then, we see that $\tilde{\mathcal{T}}$ is an equivalence class of this equivalence relation (in the proof of this fact, $\tilde{\mathcal{S}}$ plays the role of \mathcal{S}). As a consequence of this fact, we see that $\tilde{\mathcal{B}} = \tilde{\mathcal{T}}$. Thus, we see that Φ_n is unimodular.

Our proof of Proposition 3.4 (2) can be regarded as the bipartite graph version of this proof.

7. The relation to matroid theory

The discussion in the previous section has similarities to the matroid theory.

For a connected graph, the edge set forms a matroid by defining the bases by trees. This is called a graphic matroid. Moreover, we call the dual matroid of a graphic matroid a cographic matroid. In other wards, we define the bases by the condition that the complement is a tree.

We see that $X \otimes Y$ is isomorphic to the cographic matroid determined by the complete bipartite graph. Moreover, the root system of type A_n is isomorphic to the graphic matroid determined by the complete graph K_{n+1} (by identifying v with -v). Moreover, it is known that graphic matroids and cographic matroids are regular matroids. Hence, they can be realized by a unimoudlar system ([O]).

However, the discussion in the previous section is not contained in this discussion on matroids. Indeed, we see that $X \otimes Y$ is isomorphic to a unimodular system as matroids, but this fact is weaker than Proposition 3.4 (2) (this asserts that $X \otimes Y$ itself is unimodular). Thus, in this article, we directly proved Proposition 3.4 (2).

References

- [DG] V. Danilov and V. Grishukhin, Maximal unimodular systems of vectors, European J. Combin. 20 (1999), no. 6, 507–526.
- [D] R. Diestel, Graph theory. Fifth edition, Graduate Texts in Mathematics, 173. Springer, Berlin, 2017.
- [H] M. Hidaka, The Schur polynomials in primitive nth roots of unity (in Japanese), Master's thesis, Kagoshima University, 2019.
- [Mi] A. Migotti, Zur Theorie der Kreisteilungsgleichung, S.-B. der Math.-Naturwiss. Classe der Kaiser. Akad. der Wiss., Wien 87 (1883), 7–14.
- [Mo] P. Moree, Inverse cyclotomic polynomials, J. Number Theory, 129 (2009), 667–680.
- [O] J. Oxley, Matroid theory. Second edition, Oxford Graduate Texts in Mathematics, 21. Oxford University Press, Oxford, 2011.

FUTURE NET CO., LTD., FUKUOKA 810-0022, JAPAN *Email address:* k35754880gmail.com

Graduate School of Science and Engineering, Kagoshima University, Kagoshima 890-0065, Japan

Email address: itoh@sci.kagoshima-u.ac.jp