

On extended perfect codes

Konstantin Vorob'ev

*Institute of Mathematics and Informatics, Bulgarian Academy of Sciences,
Sofia 1113, Bulgaria
konstantin.vorobev@gmail.com*

March 19, 2024

Abstract

We consider extended 1-perfect codes in Hamming graphs $H(n, q)$. Such nontrivial codes are known only when $n = 2^k$, $k \geq 1$, $q = 2$, or $n = q + 2$, $q = 2^m$, $m \geq 1$. Recently, Bespalov proved nonexistence of extended 1-perfect codes for $q = 3, 4$, $n > q + 2$. In this work, we characterize all positive integers n , r and prime p , for which there exist such a code in $H(n, p^r)$.

Keywords: perfect code, extended code, Krawtchouk polynomial, Hamming graph.
2000 MSC: 05B30, 05A05, 94B25.

1 Introduction

The *Hamming graph* $H(n, q)$ is a graph whose vertices are the words of length n over the alphabet $\{0, 1, \dots, q - 1\}$. Two vertices are adjacent if and only if they differ in exactly one coordinate position, the Hamming distance $d(\bar{x}, \bar{y})$ between vertices \bar{x} , \bar{y} is the number of coordinates in which they differ. A subset C of the set of vertices of $H(n, q)$ is called a *code* in this graph. A code distance d of C is the minimum distance between distinct elements of C .

A code C in a simple graph G is called a 1-perfect code (perfect 1-error-correction code), if the vertex set of G is partitioned into balls of radius 1 centered in the code vertices. It is known [8] that if $q = p^m$ is a prime power, then there is a 1-perfect code in $H(n, q)$ if and only if $n = \frac{q^t - 1}{q - 1}$ for some integer $t \geq 1$. In the case when q is not a prime power, the problem of existence of 1-perfect codes is far from being solved. Golomb and Posner proved [4] nonexistence of 1-perfect codes in $H(7, 6)$ based on the nonexistence of a pair of orthogonal latin squares of order 6. Later Heden and Roos found [5] a necessary condition on existence of 1-perfect codes, which in particular implies the non-existence of 1-perfect codes in $H(19, 6)$.

Another interesting class of codes in $H(n, q)$ is a class of so called extended 1-perfect codes. A code C in $H(n, q)$ is an extended 1-perfect code if a puncturing (also known as a projection) of any coordinate position gives a 1-perfect code in $H(n - 1, q)$. It is known that such codes exist in $H(2^m, 2)$ and in $H(2^m + 2, 2^m)$ for any positive integer m (see [8, 3]). Formally, a one vertex in $H(2, q)$ is also a 1-perfect code (we will call it trivial). Ball proved [1] that there are no linear extended 1-perfect codes in $H(q + 2, q)$, when q is odd prime. It is also known [6] that there are no extended 1-perfect codes in $H(7, 5)$ and $H(9, 7)$. Recently, Bespalov [2] proved nonexistence of extended 1-perfect codes for $q = 3, 4$, $n > q + 2$.

In this paper we are focused on extended 1-perfect codes in Hamming graphs $H(n, q)$. Using a technique of weight distribution and some number-theoretical analysis we prove that when q is

a prime power, nontrivial extended 1-perfect codes in Hamming graphs $H(n, q)$ exist only for n, q listed above, i.e. in $H(2^m, 2)$ and $H(2^m + 2, 2^m)$, for positive integer m .

2 Preliminaries

An r -partition (C_1, C_2, \dots, C_r) of the vertex set of a graph is called *equitable* with a *quotient matrix* $S = (s_{ij})_{i,j \in \{1,2,\dots,r\}}$ if every vertex from C_i has exactly s_{ij} neighbours in C_j . The sets C_1, C_2, \dots, C_r are called *cells* of the partition.

Clearly, a 1-perfect code in $H(n, q)$ is an equitable partition with the quotient matrix $\begin{pmatrix} 0 & n(q-1) \\ 1 & n(q-1)-1 \end{pmatrix}$.

Given a code C in a simple graph $G = (V, E)$, the distance partition with respect to C is the partition $\Pi = (C = C(0), C(1), \dots, C(\rho))$ of V with respect to the distance from code C , i.e. $C(i) = \{x \in V : d(x, C) = i\}$, and all $C(i)$ are nonempty. The number $\rho = \rho(C)$ is defined by the equality $\bigcup_{i=0}^{\rho} C(i) = V$ and is called the covering radius of C . A nonempty code C of a graph is called completely regular if the partition of the vertex set with respect to the distance from C is equitable.

For our arguments we need the following auxiliary statement.

Proposition 1. [2, Theorem 2]. *Given a code C in $H(n, q)$, C is an extended 1-perfect code if and only if $(C = C(0), C(1), C(2), C(3))$ is an equitable partition in $H(n, q)$ with quotient matrix*

$$S = \begin{pmatrix} 0 & n(q-1) & 0 \\ 1 & q-2 & (n-1)(q-1) \\ 0 & n & n(q-2) \end{pmatrix}.$$

Let us define a matrix $S' = \frac{n(q-1)E-S}{q} = \frac{1}{q} \begin{pmatrix} n(q-1) & -n(q-1) & 0 \\ -1 & (n-1)(q-1)+1 & -(n-1)(q-1) \\ 0 & -n & n \end{pmatrix}$.

Our next step is to find the Jordan canonical form of S .

Lemma 1. *For $q \geq 2$ and $n \geq 2$, the equation $S' = QJQ^{-1}$ holds, where*

$$Q = \begin{pmatrix} 1 & 1 & 1 \\ \frac{q-2}{n(q-1)} & -\frac{1}{q-1} & 1 \\ -\frac{1}{(n-1)(q-1)} & \frac{1}{(q-1)^2} & 1 \end{pmatrix}, J = \begin{pmatrix} \frac{nq-n-q+2}{q} & 0 & 0 \\ 0 & n & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

$$Q^{-1} = \begin{pmatrix} \frac{n(n-1)(q-1)}{(nq-n-q+2)(n+q-2)} & \frac{n(n-1)(q-1)(q-2)}{(nq-n-q+2)(n+q-2)} & -\frac{n(n-1)(q-1)^2}{(nq-n-q+2)(n+q-2)} \\ \frac{(q-1)^2}{q(n+q-2)} & -\frac{n(q-1)^2}{q(n+q-2)} & \frac{(n-1)(q-1)^2}{q(n+q-2)} \\ \frac{1}{q(nq-n-q+2)} & \frac{n(q-1)}{q(nq-n-q+2)} & \frac{(n-1)(q-1)^2}{q(nq-n-q+2)} \end{pmatrix}.$$

Proof. The proof is obtained by direct calculations. Let us also notice that

$$\det(Q) = -\frac{q(nq-n-q+2)(n+q-2)}{n(n-1)(q-1)^3} \neq 0,$$

for $n \geq 2$ and $q \geq 2$. □

Evidently, the equation from Lemma 1 corresponds to a reduction to the Jordan canonical form of the matrix S' . Hence, diagonal elements of J are the eigenvalues of S' . For our following arguments we also need to know values of Krawtchouk polynomials on these eigenvalues.

Well-known family of orthogonal Krawtchouk polynomials is defined by the following equations

$$K_r(x, q, n) = \sum_{j=0}^r (-1)^j q^{r-j} \binom{n-r+j}{j} \binom{n-x}{r-j},$$

where n, q and degree r are nonnegative integers.

Lemma 2. *For $q, n, x \in \mathbb{Z}$, $q \geq 2$, $n \geq 2$, $0 \leq x \leq n$, we have that*

$$K_n(x, q, n) = (-1)^x (q-1)^{n-x}.$$

Proof. By definition of Krawtchouk polynomial for $r = n$, it is easy to compute

$$K_n(x, q, n) = \sum_{j=0}^n (-1)^j q^{n-j} \binom{n-x}{n-j} = (-1)^x \sum_{j'=0}^{n-x} (-1)^{j'} q^{n-x-j'} \binom{n-x}{n-x-j'} = (-1)^x (q-1)^{n-x}.$$

□

Now let us turn back to extended 1-perfect codes. We know that its distance partition Π is an equitable partition. Takes some integer i , $2 \leq i \leq n$ and consider a graph $H_i(n, q)$ with the same vertex set and two vertices being adjacent if the distance (here, Hamming distance) between them equals i . The natural question here, is Π equitable also in this new graph? The answer is positive, and not only for the Hamming graph. Actually, any equitable partition of a given distance regular graph G will be equitable for distance- i graph G , see [9, Sect. 2.2.2, 2.1.5] and [7]. Moreover, there is a way to find the corresponding quotient matrix of the partition in a new graph directly. In the following Lemma we make it for $H_n(n, q)$.

Lemma 3. *Given an extended 1-perfect code C , then the distance partition Π of C in $H(n, q)$ is equitable also in $H_n(n, q)$. Corresponding quotient matrix in the last graph equals QMQ^{-1} , where*

$$M = \begin{pmatrix} (-1)^{\frac{nq-n-q+2}{q}} (q-1)^{\frac{n+q-2}{q}} & 0 & 0 \\ 0 & (-1)^n & 0 \\ 0 & 0 & (q-1)^n \end{pmatrix},$$

and matrices Q and Q^{-1} are defined in Lemma 1.

Proof. As we discussed before, Π is equitable in $H_n(n, q)$. By [7, Theorem 3 and Example 8] the corresponding quotient matrix equals $K_n(S', q, n) = QK_n(J, q, n)Q^{-1}$. Since J is a diagonal matrix we have that

$$K_n(J, q, n) = \begin{pmatrix} K_n(nq - n - q + 2, q, n) & 0 & 0 \\ 0 & K_n(n, q, n) & 0 \\ 0 & 0 & K_n(0, q, n) \end{pmatrix}.$$

The last step it to apply Lemma 2 and we finish. □

3 Extended perfect codes

From Section 1 we remember, that extended 1-perfect codes are known only when n and q belong to the following three infinite families.

Proposition 2. *Extended 1-perfect codes in $H(n, q)$ exist for the following values of n and q :*

1. $n = 2$, $q \geq 2$ (one vertex in $H(2, q)$),

2. $n = q + 2$, $q = 2^m$, $m \geq 1$ (extended perfect code in $H(q + 2, q)$).

3. $n = 2^k$, $k \geq 1$, $q = 2$ (extended perfect code in $H(2^k, 2)$).

The main idea of this Section is to find the quotient matrix of the initial distance partition in $H_n(n, q)$ in closed form.

Theorem 1. *Let C be an extended 1-perfect code in $H(n, q)$, $q \geq 2$, $n \geq 2$. Then*

$$(q - 2) \frac{(q - 1)^x - (-1)^x}{x} \in \mathbb{N},$$

where $x = \frac{n+q-2}{q}$.

Proof. Consider a distance partition $\Pi = (C = C(0), C(1), C(2))$ defined by C , with the quotient matrix

$$S = \begin{pmatrix} 0 & n(q-1) & 0 \\ 1 & q-2 & (n-1)(q-1) \\ 0 & n & n(q-2) \end{pmatrix}.$$

By Lemma 3 we have that the quotient matrix of Π in $H_n(n, q)$ equals QMQ^{-1} , where

$$M = \begin{pmatrix} (-1)^{\frac{nq-n-q+2}{q}} (q-1)^{\frac{n+q-2}{q}} & 0 & 0 \\ 0 & (-1)^n & 0 \\ 0 & 0 & (q-1)^n \end{pmatrix},$$

and matrices Q and Q^{-1} were defined in Lemma 1. Let us count the entry of M in the third row and the first column. After all necessary calculations we have that

$$(M)_{3,1} = \frac{1}{q(nq - n - q + 2)} \left((q-1)^n - (-1)^n + q \left((-1)^n - (q-1)^{\frac{q+n-2}{q}} (-1)^{\frac{nq-n-q}{q}} \right) + \right. \\ \left. + (q-2) \frac{(q-1)^{\frac{n+q-2}{q}} (-1)^{\frac{nq-q-n+2}{q}} - (-1)^n}{\frac{n+q-2}{q}} \right).$$

By definition of weight distribution this value is nonnegative integer. Clearly, the last summand in big brackets must be integer. Therefore,

$$(q-2) \frac{(q-1)^{\frac{n+q-2}{q}} (-1)^{\frac{nq-q-n+2}{q}} - (-1)^n}{\frac{n+q-2}{q}} \in \mathbb{Z}.$$

After multiplying it by $(-1)^{\frac{nq-q-n+2}{q}}$ we have the required. \square

This new necessary condition proved in Theorem 1 works for arbitrary q . However, we are mainly interested in the case $q = p^m$, where p is a prime number and $m \geq 1$. The next step is to classify all such pairs of n and q that there exist extended 1-perfect code in $H(n, q)$ and q is a prime power.

Theorem 2. *Let C be an extended 1-perfect code in $H(n, q)$ for $q = p^m$, where p is a prime number $m \geq 1$, $n \geq 2$. Then one of the following holds:*

1. $n = 2$,
2. $p = 2$, $m = 1$ and $n = 2^t$ for some integer $t \geq 2$,

3. $p = 2$, $m \geq 1$, $n = q + 2$.

Proof. We consider the case when q is a prime power. Since 1-perfect codes exist in $H(n', q)$ only for $n' = \frac{q^k - 1}{q - 1}$ for some integer $k \geq 1$ we conclude that

$$n = \frac{q^k - 1}{q - 1} + 1, \quad (1)$$

for some integer $k \geq 1$.

The case $k = 1$ immediately lead us a to the one-vertex code in $H(n, 2)$. In the rest of the proof we suppose that $k \geq 2$. Our next step is to use Theorem 1. Indeed, for $n = \frac{q^k - 1}{q - 1} + 1$ we have that

$$r = (q - 2) \frac{(q - 1)^x - (-1)^x}{x} \in \mathbb{N}, \quad (2)$$

where

$$x = \frac{q^{k-1} + q - 2}{q - 1} \in \mathbb{N}, \quad (3)$$

Let us rewrite this equation in the following form

$$x(q - 1) = q^{k-1} + q - 2 \quad (4)$$

The following analysis depends on the parity of p .

Case $p > 2$. Evidently, it follows from (2) that x is odd too. Since $k \geq 2$, from (3) we have $x > 1$. We know that q is odd, so $(q, q - 2) = 1$ (here and further by (a, b) we mean the greatest common divisor of two positive integers a and b). Simple analysis of (2) and (4) shows that

$$(x, q) = (x, q - 1) = (x, q - 2) = 1. \quad (5)$$

Using this fact and (2) we have that $\frac{(q-1)^x - (-1)^x}{x} \in \mathbb{N}$. Let t be the smallest odd prime divisor of x . Since $x > 1$ and x is odd, t is not less than 3. Since $(q - 1)^x + 1$ is divisible by t , we conclude that $(q - 1)^{2x} - 1$ is divisible by t . Equality (5) implies $(t, q - 1) = 1$. Consequently, $(q - 1)^{t-1} - 1$ is also divisible by t . These two divisions guarantee us that there exists the smallest natural number d such that $(q - 1)^d - 1$ is divisible by t , and d divides $(t - 1, 2x)$. It follows from the choice of t that d does not have odd prime divisors.

Combining this fact with the one that $2x$ is not divisible by 4, we have that $d = 1$ or $d = 2$. In other words, q or $q(q - 2)$ is divisible by t correspondingly. These cases both contradict to (5).

Case $p = 2$. First of all, if $m = 1$ (i.e. $q = 2$) then extended 1-perfect codes exist in $H(2^k, 2)$ for all $k \geq 2$. Hence, further we consider the case $m \geq 2$. If $k = 2$ then we have $n = q + 2$ where q is a power of 2. These are exactly parameters of the codes from another known family of codes. To finish the prove we need to consider the case $k \geq 3$. The idea is to repeat the arguments from the previous case for $\frac{x}{2}$. Clearly, it follows from (3) that x is even but not divisible by 4. Hence, $\frac{x}{2}$ is odd. Since $k \geq 3$, we have $\frac{x}{2} > 1$. Again, analysis of (2) and (4) shows that

$$\left(\frac{x}{2}, q\right) = \left(\frac{x}{2}, q - 1\right) = \left(\frac{x}{2}, q - 2\right) = 1. \quad (6)$$

Define t as the smallest odd prime divisor of $\frac{x}{2}$. As before, $t \geq 3$. Now we have that $(q - 1)^x - 1$ and $(q - 1)^t - 1$ are both divisible by t . Hence, there exist the smallest natural d such that d divides $(t - 1, x)$. By choice of t and the fact that x is not divisible by 4, we have that $d = 1$ or $d = 2$. Consequently, q or $q(q - 2)$ is divisible by t correspondingly, which contradicts to (6). \square

The proof is relatively short and is based on the fact that q ia prime power. Unfortunately, the idea of the proof can not be applied to the case of arbitrary q directly. Indeed, in general case one has more complicated condition instead of equation (1). Moreover, as we discussed in Section 1, one does not know any nontrivial 1-perfect codes when q is not a prime number. So the generalization of ideas from the proof of Theorem 2 might make sense only when one find some series of 1-perfect codes in this case.

4 Conclusion

In this work we completely characterize all numbers n , p and m , for which there exists an extended 1-perfect code in $H(n, q)$, $q = p^m$. The situation for q not a prime power seems more complicated, especially since nontrivial 1-perfect codes are still not known in this case.

The crucial idea of the proof of Theorem 2 is to find analytically some elements of the quotient matrix of the distance partition of initial code in i -distance Hamming graph for some i and then analyze it. We believe that this idea will be useful for other interesting families of completely regular codes in different distance-regular graphs.

5 Acknowledgments

The author was supported by the NSP P. Beron project CP-MACT. Author is grateful to Fedor Petrov for useful discussions on the proof of main Theorem.

References

- [1] S. Ball. On sets of vectors of a finite vector space in which every subset of basis size is a basis. *J. Eur. Math. Soc.*, 14(3):733–748, 2012. DOI: 10.4171/JEMS/316.
- [2] Evgeny Beshpalov. On the non-existence of extended 1-perfect codes and mds codes. *Journal of Combinatorial Theory, Series A*, 189:105607, 2022.
- [3] J. Borges, J. Rifà, and V. A. Zinoviev. On completely regular codes. *Probl. Inf. Transm.*, 55(1):1–45, Jan. 2019. DOI: 10.1134/S0032946019010010.
- [4] S. W. Golomb and E. C. Posner. Rook domains, latin squares, and error-distributing codes. *IEEE Trans. Inf. Theory*, 10(3):196–208, 1964. DOI: 10.1109/TIT.1964.1053680.
- [5] O. Heden and C. Roos. The non-existence of some perfect codes over non-prime power alphabets. *Discrete Math.*, 311(14):1344–1348, 2011. DOI: 10.1016/j.disc.2011.03.024.
- [6] J. I. Kokkala, D. S. Krotov, and P. R. J. Östergård. On the classification of MDS codes. *IEEE Trans. Inf. Theory*, 61(12):6485–6492, December 2015. DOI: 10.1109/TIT.2015.2488659.
- [7] D. S. Krotov. On weight distributions of perfect colorings and completely regular codes. *Des. Codes Cryptography*, 61(3):315–329, 2011. DOI: 10.1007/s10623-010-9479-4.
- [8] F. J. MacWilliams and N. J. A. Sloane. *The Theory of Error-Correcting Codes*. Amsterdam, Netherlands: North Holland, 1977.
- [9] W. J. Martin. *Completely Regular Subsets*. PhD thesis, University of Waterloo, Waterloo, Ontario, Canada, 1992. Online: <http://users.wpi.edu/~martin/RESEARCH/THESIS/>.