

**FLOQUET–BLOCH FUNCTIONS ON NON-SIMPLY
CONNECTED MANIFOLDS, THE AHARONOV–BOHM
FLUXES, AND CONFORMAL INVARIANTS OF
IMMERSED SURFACES**

I.A. TAIMANOV

ABSTRACT. Spectral (Bloch) varieties of multidimensional differential operators on non-simply connected manifolds are defined. In their terms it is given a description of the analytical dependence of the spectra of magnetic Laplacians on non-simply connected manifolds on the values of the Aharonov–Bohm fluxes and a construction of analogues of spectral curves for two-dimensional Dirac operators on Riemann surfaces and, thereby, new conformal invariants of immersions of surfaces into 3- and 4-dimensional Euclidean spaces.

To S.P. Novikov on his 85-th birthday

1. INTRODUCTION

The modern theory of finite-zone (finite gap) operators goes back to Novikov’s work on the integration of the periodic Korteweg–de Vries equation [14]. The spectral curves of finite-zone one-dimensional operators introduced in it Schrödinger later formed the basis of algebraic-geometric methods for integrating soliton equations. The generalization of spectral curves to the case of two-dimensional Schrödinger operators, defined in [7], has expanded the applications of this method.

In this paper we will discuss applications of spectral (Bloch) varieties of multidimensional differential operators on non-simply connected manifolds.

The main objectives of this work are

- description of the analytical dependence of the spectra of magnetic Laplacians on non-simply connected manifolds on the values of the Aharonov–Bohm fluxes (§3, Theorem 2);
- construction of analogues of spectral curves for two-dimensional Dirac operators on Riemann surfaces and, thereby, new conformal invariants of immersions of surfaces into 3- and 4-dimensional Euclidean spaces (SS4 and 5, Theorems 4 and 5).

Let us point out as an example another nontrivial situation, when the spectral curve (one-dimensional Bloch variety) is naturally defined. In [3] there were considered Bloch eigenfunctions of the Cauchy–Riemann operator on a punctured torus

$$T^2 = \mathbb{C}/\{\mathbb{Z}e_1 + \mathbb{Z}e_2\},$$

i.e. a two-dimensional torus with punctured points p_1, \dots, p_N . They are defined as solutions to the equation

$$\bar{\partial}\psi = 0,$$

satisfying the Floquet–Bloch conditions

$$\psi(z + e_j) = e^{2\pi i\langle k, e_j \rangle} \psi(z),$$

and having asymptotics

$$\psi(z) = \frac{a_j}{z - p_j} + O(|z - p_j|).$$

It was shown in [3] that such functions exist exactly for tuples

$$(k_1, k_2, a_1, \dots, a_N) \in \Gamma,$$

belonging to some spectral curve Γ .

After our talk at a conference in Lumumba Russian People’s Friendship University in October 2023, Yu. Kordyukov drew our attention to the works of [9, 10, 18] devoted to the Floquet–Bloch theory for periodic elliptic operators on non-simply connected compact manifolds. They contain part of the results from §2, but we save their conclusion, since it was not presented separately and will be needed later. These works study a generalization of the Bloch property and its application to questions of the existence of eigenvalues of such operators. We will be more interested in Bloch varieties.

2. FLOQUET–BLOCH EIGENFUNCTIONS OF LAPLACE–BELTRAMI AND SCHRÖDINGER OPERATORS

Let M be a closed manifold with a nontrivial fundamental group $\pi_1(M)$. Consider the composition epimorphism

$$\rho : \pi_1(M) \rightarrow H_1(M) \rightarrow \mathbb{Z}^k = H_1(M)/\text{Torsion}$$

and denote by

$$p : M_0 \rightarrow M$$

the covering corresponding to the subgroup $\text{Ker } \rho$: $\pi_1(M_0) = \text{Ker } \rho$. The group $\Lambda = H_1(M)/\text{Torsion} = \mathbb{Z}^N$, where $N = \dim H_1(M; \mathbb{R})$ acts freely on M_0 , so that $M = M_0/\Lambda$.

Let us choose in Λ the generators Z_1, \dots, Z_N and in $H^1(M; \mathbb{R})$ a basis realized by closed 1-forms $\omega_1, \dots, \omega_N$, such that

$$\langle \omega_j, Z_k \rangle = \int_{Z_j} \omega_k = \delta_{jk}, \quad j, k = 1, \dots, N.$$

The preimages under covering p of 1-forms $\omega_1, \dots, \omega_N$ are the differentials of functions

$$p^*\omega_k = dh_k, \quad h_k : M_0 \rightarrow \mathbb{R}, \quad k = 1, \dots, N.$$

Let us denote by T_1, \dots, T_N generators of the action of Λ on M_0 such that

$$h_j(T_k x) = h_j(x) + \delta_{jk}.$$

Let us consider functions on M_0 satisfying the conditions

$$(1) \quad \begin{aligned} \psi &: M_0 \rightarrow \mathbb{C}, \\ \psi(T_k x) &= e^{i\kappa_m} \psi(x), \quad m = 1, \dots, N, \end{aligned}$$

in this case the quantities $\kappa_1, \dots, \kappa_N \in \mathbb{C}$ are called *quasi-momenta* and are determined up to $2\pi m, m \in \mathbb{Z}$.¹

The multipliers which correspond to such a function define a homomorphism

$$\mu : \Lambda \rightarrow \mathbb{C}^* = \mathbb{C} \setminus \{0\}, \quad \mu(T_m) = e^{i\kappa_m}, \quad m = 1, \dots, N.$$

We have

Proposition 1. *If a function ψ satisfies (1), then it can be represented as*

$$(2) \quad \psi(x) = e^{i(\kappa_1 h_1(x) + \dots + \kappa_N h_N(x))} \varphi(x),$$

where $\varphi : M_0 \rightarrow \mathbb{C}$ is a Λ -periodic function.

The Floquet-Bloch functions were defined as solutions of differential equations with periodic coefficients on \mathbb{R}^N , satisfying (1), Floquet for $N = 1$ and Bloch for arbitrary dimension N . It was assumed that they have the form (2), where $h_1(x) = x^1, \dots, h_N(x) = x^N$ — coordinate functions.

Let us consider the Laplace operator on the Riemannian manifold M . Let $g_{jk} dx^j dx^k$ denote the Riemannian metric. The Laplace–Beltrami

¹If the periods of the functions h_m are not normalized by the unit, then the quasimomenta are determined taking into account these periods. For example, in the theory of the one-dimensional periodic Schrödinger operator after a shift by period T , the Bloch function is multiplied by $e^{i\kappa T}$, where κ is the quasi-momentum [14].

operator (or, more simply, the Laplacian), corresponding to this metric, acts on smooth functions on the manifold and is given by the expression

$$(3) \quad \Delta = \frac{1}{\sqrt{g}} \frac{\partial}{\partial x^j} \sqrt{g} g^{jk} \frac{\partial}{\partial x^k},$$

where $g = \det(g_{jk})$, $g^{jk} g_{km} = \delta_m^j$ and everywhere we mean summation over repeated upper and lower indices. Its action on the function f takes the form

$$\Delta f = \frac{1}{\sqrt{g}} \frac{\partial}{\partial x^j} \left(\sqrt{g} g^{jk} \frac{\partial f}{\partial x^k} \right).$$

The Schrödinger operator on a manifold is defined as the sum of the operator $-\Delta$ and multiplication by the potential $U(x)$:

$$L = -\Delta + U.$$

We call by *Bloch eigenfunction* of the Schrödinger operator with eigenvalue E a solution to the equation

$$(4) \quad L\psi = E\psi,$$

satisfying (1). As a function it is defined on the maximal Abelian covering M_0 of $M = M_0/\mathbb{Z}^N$.

The following fact is proved by straightforward calculations.

Proposition 2. *The function $\psi : M_0 \rightarrow \mathbb{C}$ of the form (2) satisfies the equation*

$$(5) \quad L\psi = (-\Delta + U)\psi = E\psi$$

if and only if the function $\varphi : M \rightarrow \mathbb{C}$ satisfies the equation

$$(6) \quad - \left[\Delta\varphi - g^{jk} \frac{\partial}{\partial x^j} \left(\sum_{m=1}^N \kappa_m h_m \right) \frac{\partial}{\partial x^k} \left(\sum_{q=1}^N \kappa_q h_q \right) \varphi + \right. \\ \left. + i \left(\sum_{m=1}^N \kappa_m \Delta h_m \right) \varphi + 2i g^{jk} \left(\sum_{m=1}^N \kappa_m \frac{\partial h_m}{\partial x^j} \right) \frac{\partial \varphi}{\partial x^k} \right] + U\varphi = E\varphi.$$

It should be noted that the terms in (6) that include the functions h_m are correctly defined on M , although the functions h_m are defined on the cover M_0 . By definition, the function h_m is chosen such that

$$dh_m = p^* \omega_m,$$

where $\omega_m = A_{m,k} dx^k$ is a smooth 1-form on the compact manifold M . That's why

$$\frac{\partial h_m}{\partial x^k} = A_{m,k}, \quad m = 1, \dots, N, k = 1, \dots, \dim M,$$

and this quantity is correctly defined on M . Let's write the equation (6) in terms of 1-forms A :

$$(7) \quad \begin{aligned} & - \left[\Delta - g^{jk} \left(\sum_{m=1}^N \kappa_m A_{m,j} \right) \left(\sum_{q=1}^N \kappa_q A_{q,k} \right) + \right. \\ & + i \sum_{m=1}^N \kappa_m \left(g^{jk} \frac{\partial A_{m,k}}{\partial x^j} + \frac{1}{\sqrt{g}} \frac{\partial \sqrt{g} g^{jk}}{\partial x^j} A_{m,k} \right) + \\ & \left. + 2i g^{jk} \left(\sum_{m=1}^N \kappa_m A_{m,j} \right) \frac{\partial}{\partial x^k} \right] \varphi + U\varphi = E\varphi. \end{aligned}$$

Recall that $L_p(M), p \in \mathbb{Z}, p \geq 0$, denotes the space of functions M such that $\int_M |f|^p d\mu < \infty$, where $d\mu$ is the measure generated by the metric. More precisely, this is the space of equivalence classes of functions with respect to the relation $f_1 \sim f_2$ if f_1 and f_2 coincide outside a set of measure zero. In $L_p(M)$ the norm $\|f\|_p = \left(\int_M |f|^p d\mu \right)^{\frac{1}{p}}$ is defined. For C^∞ -functions on M we define norms

$$\|\nabla^k f\|^2 = \nabla^{j_1} \dots \nabla^{j_k} f \nabla_{j_1} \dots \nabla_{j_k} f,$$

where ∇ is the covariant derivative. We denote by $H_k^p(M)$ the closure of the space of C^∞ -functions with $\|\nabla^l f\| \in L_p(M)$ for $0 \leq l \leq k$ relative to the norm $\|g\|_{H_k^p} = \sum_{l=0}^k \|\nabla^l f\|_p$. If the Riemannian manifold M is compact and without boundary, then the spaces defined in this way do not depend on the choice of metric and Kondrashov's theorem holds: the embedding $H_k^q \subset L_p$ is compact for $1 \geq \frac{1}{p} > \frac{1}{q} - \frac{k}{\dim M}$. Detailed definitions and proofs of these facts are presented in [2].

Theorem 1. *In \mathbb{C}^{N+1} there is an analytic subset Γ , a spectral variety distinguished as the set zeros*

$$\Gamma = \{\mathcal{P}(E, \kappa_1, \dots, \kappa_N) = 0\}$$

of a nontrivial entire function such that a point $(E, \kappa_1, \dots, \kappa_N)$ belongs to Γ if and only if there exists a Bloch eigenfunction of the form (2) with quasimomenta $\kappa_1, \dots, \kappa_N$, satisfying (5).

The spectral variety does not depend on the choice of functions h_1, \dots, h_N such that $p^ \omega_k = dh_k, k = 1, \dots, N$, and therefore is determined only by the operator L .*

PROOF. Let us choose a constant ε such that the operator $\Delta + \varepsilon : H_2^2(M) \rightarrow L_2(M)$ is invertible and represent φ in the form

$$\varphi = (\Delta + \varepsilon)^{-1} \xi.$$

We rewrite the equation (7) in the form

$$(8) \quad \left\{ 1 + \left[(E - U - \varepsilon) + \sum_m \kappa_m B_m + \sum_{m,q} \kappa_m \kappa_q B_{mq} + \sum_m \kappa_m C_m^k \frac{\partial}{\partial x^k} \right] (\Delta + \varepsilon)^{-1} \right\} \xi = 0.$$

Composition of the multiplication by a bounded function $G_\kappa = [(E - U - \varepsilon) + \sum_m \kappa_m B_m + \sum_{m,q} \kappa_m \kappa_q B_{mq}]$ and $(\Delta + \varepsilon)^{-1}$ splits into a composition

$$L_2 \xrightarrow{(\Delta + \varepsilon)^{-1}} H_2^2 \xrightarrow{\text{embedding}} L_2 \xrightarrow{\times G_\kappa} L_2.$$

The composition of the first two operators is an operator that increases smoothness, the first being bounded, and the embedding is compact. The multiplication operator by G_κ is bounded. The composition of bounded and compact operators is compact, hence the operator

$$[(E - U - \varepsilon) + \sum_m \kappa_m B_m + \sum_{m,q} \kappa_m \kappa_q B_{mq}] (\Delta + \varepsilon)^{-1}$$

is compact for any κ and E . There is a similar decomposition

$$L_2 \xrightarrow{(\Delta + \varepsilon)^{-1}} H_2^2 \xrightarrow{\sum \kappa_m C_m^k \frac{\partial}{\partial x^k}} H_2^1 \xrightarrow{\text{embedding}} L_2,$$

in which the embedding is compact and other operators are bounded.

Therefore, the equation (8) has the form

$$(9) \quad (1 + A(E, \kappa_1, \dots, \kappa_N)) \xi = 0,$$

where $A(E, \kappa_1, \dots, \kappa_N)$ is a polynomial in $\kappa_1, \dots, \kappa_N$ and E is a pencil of compact operators from $L_2(M)$ to $L_2(M)$. By the Keldysh theorem, in this case the equation (9) is solvable then and only if $(E, \kappa_1, \dots, \kappa_N)$ belongs to some analytic subset Γ in \mathbb{C}^{N+1} , which is the zero set of an entire function \mathcal{P} .

Since for $\kappa_1 = \dots = \kappa_N = 0$ this subset intersects \mathbb{C} along the spectrum of the operator $-\Delta + E$, which is discrete, then Γ is a proper subset and the entire function \mathcal{P} is non-trivial.

Suppose we took other functions $\tilde{h}_1, \dots, \tilde{h}_N$, satisfying these conditions. Then

$$h_j(T_k x) - \tilde{h}_j(T_k x) = 0 \quad \text{for all } j, k \text{ and } x.$$

Therefore, the function $h_j - \tilde{h}_j$ is Λ -periodic and descends to a function on M .

If

$$\psi(x) = e^{i \sum_m \kappa_m h_m(x)} \varphi(x)$$

is a Bloch eigenfunction, $L\psi = E\psi$, then

$$\psi(x) = e^{i\sum_m \kappa_m \tilde{h}_m(x)} [e^{i\sum_m \kappa_m (h_m(x) - \tilde{h}_m(x))} \varphi(x)] = e^{i\sum_m \kappa_m \tilde{h}_m(x)} \tilde{\varphi}(x).$$

The converse is also true, which implies the coincidence of spectral manifolds constructed from different sets of functions $\{h_j\}$ and $\{\tilde{h}_j\}$. The theorem has been proven.

EXAMPLE. If $M = S^1$, then $\Gamma/\{\kappa \sim \kappa + 2\pi\}$ is the spectral curve of the one-dimensional Schrödinger operator with a periodic potential, and the classical Bloch eigenfunctions of this operator [14] correspond to the case $h(x) = x$.

In proof of Theorem 1, we followed our works [20, 21]. This proof for the periodic Schrödinger operator on \mathbb{R}^N and the heat operator $\partial_y - \partial_x^2 + u$, included in the Lax representation for the Kadomtsev–Petviashvili II equation, was obtained by us in 1984 and was even used and cited in [12], but we published it in [20] when we found an application of this method for defining the spectral curves of tori immersed in the three-dimensional space. Note that for the operator $i\partial_y - \partial_x^2 + u$, which enters the Lax representation for the Kadomtsev–Petviashvili equation I, the spectral curve does not exist.

3. SPECTRA OF MAGNETIC LAPLACIANS AND THE AHARONOV–BOHM FLUXES

This work arose as a continuation of [24] in the study of magnetic Laplacians on non-simply connected manifolds.

The equation (7) is rewritten as

$$(-\tilde{\Delta} + U)\varphi = E\varphi,$$

where

$$\tilde{\Delta} = \frac{1}{\sqrt{g}} \left(\frac{\partial}{\partial x^j} + i\mathcal{A}_j \right) \sqrt{g} g^{jk} \left(\frac{\partial}{\partial x^k} + i\mathcal{A}_k \right).$$

The operator $\tilde{\Delta}$ is the magnetic Laplacian (the Laplace–Bochner operator) corresponding to the magnetic field $F = \sum_{j < k} F_{jk} dx^j \wedge dx^k$ with potential

$$\mathcal{A} = \mathcal{A}_k dx^k, \quad d\mathcal{A} = F.$$

In our case, for real values of $\kappa_1, \dots, \kappa_N$ we have

$$\mathcal{A} = \sum_m \kappa_m \omega_m$$

and

$$d\mathcal{A} = 0,$$

i.e. this operator corresponds to the magnetic Laplacian with a zero magnetic field. However, with such a formal procedure for constructing the magnetic Laplacian, the spectrum of L depends on the choice of potential.

The theory of the Aharonov–Bohm effect [1] also considers the situation with a zero magnetic field on a non-simply connected manifold, which is the plane \mathbb{R}^2 with a certain number N of punctured points $\{Q_1, \dots, Q_N\}$ (in reality, of small disks). We have

$$H^1(\mathbb{R}^2 \setminus \{Q_1, \dots, Q_N\}; \mathbb{R}) = \mathbb{R}^N$$

and we can take the following closed 1-forms as generators $\omega_1, \dots, \omega_N$, of this group, such that the integral of ω_j along a small closed contour γ_k , inside which lies only one punctured point — Q_k , is equal to

$$\oint_{\gamma_k} \omega_j = \oint_{\gamma_k} A_{j,l} dx^l = \delta_{jk}, \quad j, k = 1, \dots, N.$$

When quantizing a zero magnetic field on a plane with punctured points, a choice arises of the “vector potential” $\mathcal{A} = \mathcal{A}_l dx^l$, which can be cohomologically nontrivial:

$$[\mathcal{A}] \neq 0 \in H^1(\mathbb{R}^2 \setminus \{Q_1, \dots, Q_N\}; \mathbb{R}).$$

Expanding it into generators $[\omega_1], \dots, [\omega_N]$,

$$[\Omega] = \kappa_1[\omega_1] + \dots + \kappa_N[\omega_N],$$

we obtain as coefficients the so-called Aharonov–Bohm fluxes $\kappa_1, \dots, \kappa_N$. Aharonov and Bohm indicated that these fluxes are physically observable, which was later confirmed by physical experiments.

When quantizing a system with the Hamilton function

$$H(x, p) = g^{jk}(x)p_j p_k + U(x)$$

in a cohomologically trivial magnetic field F , where $[F] = 0 \in H^2(M; \mathbb{R})$, on a non-simply connected manifold M the vector potential of the magnetic field A is expanded into the following sum

$$\mathcal{A} = \mathcal{A}_0 + \sum_m \kappa_m \omega_m, \quad d\mathcal{A}_0 = F, \quad d\mathcal{A}_j = 0, \quad j = 1, \dots, N.$$

It is natural to call the expansion coefficients $\kappa_1, \dots, \kappa_N$ the Aharonov–Bohm fluxes.

Analogously to Theorem 1, the following fact is proved.

Theorem 2. *When quantizing the motion of a charged particle in an (exact) magnetic field F , $dF = 0$, on a non-simply connected manifold*

spectrum of the corresponding magnetic Laplacian

$$\widehat{H} = -\frac{1}{\sqrt{g}} \left(\frac{\partial}{\partial x^j} + i\mathcal{A}_j \right) \sqrt{g} g^{jk} \left(\frac{\partial}{\partial x^k} + i\mathcal{A}_k \right) + U(x)$$

is related to Aharonov–Bohm flows by the analytical relation

$$(10) \quad \mathcal{P}(E, \kappa_1, \dots, \kappa_N) = 0.$$

Note that for $F \neq 0$ the choice of vector potential \mathcal{A}_0 is ambiguous. The equation

$$\mathcal{P}(E, 0, \dots, 0) = 0$$

specifies the spectrum of the magnetic Laplacian with $\mathcal{A} = \mathcal{A}_0$ and the addition of non-trivial values of the Aharonov–Bohm fluxes shows how the spectrum changes, i.e., in fact, the spectrum at the selected value of the vector potential \mathcal{A}_0 is compared with its deformations when adding to \mathcal{A} closed 1-forms. The difference between two vector potentials is a closed 1-form and the flux values κ_m correspond to it.

This relation (10) is non-trivial, which is already shown by the example of a zero magnetic field $F = 0$.

Theorem 3. *On an N -dimensional torus with $A_0 = 0$, $A_{j,l} = \delta_{jl}\kappa_l$, $j = 1, \dots, N$, the equation (10) defines the Bloch variety of the Schrödinger operator $L = -\Delta + U$ with periodic potential $U(x)$. In this case, the flows $\kappa_1, \dots, \kappa_N$ are quasimoments of Bloch eigenfunctions.*

According to the Bethe–Sommerfeld conjecture, proven in [5, 6, 15, 16, 17], for $N \geq 2$ there are a finite number of gaps on the real line such that for any E lying outside these gaps, there is a set of fluxes $\kappa_1, \dots, \kappa_N$, that E is the eigenvalue of the operator

$$-\sum_{m=1}^N \left(\frac{\partial}{\partial x^m} - i\kappa_m \right)^2 + U(x),$$

those. Different quantizations of even a zero magnetic field lead to different eigenvalues of the magnetic Laplacian.

4. SPECTRAL VARIETIES OF DIRAC OPERATORS ON RIEMANN SURFACES

Let us consider the Dirac operator on a two-dimensional spinor Riemannian manifold M —a closed Riemann surface of genus g .

Let z be a conformal parameter on M and

$$ds^2 = e^{2\alpha} dzd\bar{z}$$

be a Riemannian metric. The Dirac operator has the form

$$D_0 = e^{-3\alpha/2} \begin{pmatrix} 0 & \partial \\ -\bar{\partial} & 0 \end{pmatrix} e^{\alpha/2},$$

where $\partial = \frac{\partial}{\partial z}$ and $\bar{\partial} = \frac{\partial}{\partial \bar{z}}$. It acts on sections of spinor bundles.

We will consider a more general operator by including the potential $V(x)$ — a matrix function with scalar components that acts on spinors by multiplying:

$$(11) \quad D = D_0 + V = 2e^{-3\alpha/2} \begin{pmatrix} 0 & \partial \\ -\bar{\partial} & 0 \end{pmatrix} e^{\alpha/2} + V.$$

Since D_0 has a discrete spectrum, there is a value of E_0 such that the operator

$$D_0 - E_0,$$

acting on L_2 -sections of the spinor bundle is invertible. We can apply the same reasoning as in the proof of Theorem 1 to derive the following result. The only serious difference is the analogue of Kondrashov's theorem on compactness of embedding for sections of spinor bundles, but in the case of compact manifolds it immediately follows from a similar theorem for function spaces.

Theorem 4. *In \mathbb{C}^{2g+1} there is an analytic subset, the spectral variety, determined as the zero set*

$$S = \{\mathcal{P}(E, \kappa_1, \dots, \kappa_N) = 0\}$$

of a nontrivial entire function such that $(E, \kappa_1, \dots, \kappa_{2g})$ lies to S if and only if there is a Bloch eigenspinor of the form

$$(12) \quad \psi(x) = e^{i(\kappa_1 h_1(x) + \dots + \kappa_{2g} h_{2g}(x))} \varphi(x),$$

on M_0 , where φ is a section of the spinor bundle over M , satisfying the equation

$$D\varphi = E\varphi.$$

The spectral variety does not depend on the choice of functions h_1, \dots, h_{2g} such that $p^\omega_k = dh_k, k = 1, \dots, 2g$, and therefore is determined only by the operator D .*

5. SPECTRAL VARIETIES AS INVARIANTS OF SURFACES IN EUCLIDEAN THREE- AND FOUR-SPACES

5.1. The Weierstrass (spinor) representation of surfaces in three- and four-spaces. Any immersion of a closed oriented surface M immersed into \mathbb{R}^3 or \mathbb{R}^4 is described in terms of a solution to the Dirac equation

$$\mathcal{D}\psi = 0,$$

where

$$\mathcal{D} = \begin{pmatrix} 0 & \partial \\ -\bar{\partial} & 0 \end{pmatrix} + \begin{pmatrix} U & 0 \\ 0 & \bar{U} \end{pmatrix}.$$

For brevity, we will limit ourselves to surfaces in \mathbb{R}^3 . In this case, the potential U is real. Following [23], we identify \mathbb{R}^3 with the space of Hermitian 2×2 -matrices A : $\bar{A}^\top = -A$, and represent the immersion in the form

$$(13) \quad \begin{aligned} X(P) &= \begin{pmatrix} ix^3 & -x^1 - ix^2 \\ x^1 - ix^2 & -ix^3 \end{pmatrix} (P) = \\ &= i \int_{P_0}^P \left[\begin{pmatrix} \psi_1 \bar{\psi}_2 & -\bar{\psi}_2^2 \\ \psi_1^2 & -\psi_1 \psi_2 \end{pmatrix} dz + \begin{pmatrix} \bar{\psi}_1 \psi_2 & \bar{\psi}_1^2 \\ -\psi_2^2 & -\bar{\psi}_1 \psi_2 \end{pmatrix} d\bar{z} \right] + X(P_0). \end{aligned}$$

Here x^1, x^2, x^3 are the Euclidean coordinates in \mathbb{R}^3 , $\psi = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}$ is a solution of the Dirac equation, $X : \mathcal{U} \rightarrow \mathbb{R}^3$ is an immersion of a simply connected domain $\mathcal{U} \subset \mathbb{C}$ in \mathbb{R}^3 , z is a complex parameter (coordinate) in \mathcal{U} , P_0 is a fixed point in \mathcal{U} and the integral is taken along any path from P_0 to P lying in \mathcal{U} (its value does not depend on the path).

The vector function ψ defines the surface with up to a shift $X(P_0)$, i.e. essentially defines its Gaussian map.

These formulas for the local representation of surfaces allowing soliton deformations were derived in [11], where it was shown that the parameter z is conformal, the induced metric takes the form

$$ds^2 = (|\psi_1|^2 + |\psi_2|^2)^2 dz d\bar{z} = e^{2\alpha} dz d\bar{z}$$

and the potential U of the Dirac operator takes the form

$$U = \frac{e^\alpha}{2} H,$$

where H is the mean curvature of the surface (in the case of surfaces in \mathbb{R}^3 , the potential U is real).

In [19] it was proven that any closed oriented surface in \mathbb{R}^3 has such a representation, and that ψ_1 and ψ_2 are sections of a spinor bundle over the immersed surface.

5.2. Spectral curves of immersed tori. Let M be a torus immersed in \mathbb{R}^3 by (13). Let

$$\mathbb{C}/\{\mathbb{Z}e_1 + \mathbb{Z}e_2\} \approx M$$

be a torus conformally equivalent to M . We denote by e_1 and e_2 the generators of the period lattice Λ . We will identify them with the generators of $H_1(M; \mathbb{Z}_2)$.

The immersion is given by the spinors $\psi_1\sqrt{dz}$ and $\bar{\psi}_2\sqrt{dz}$. The spinor bundle over any closed oriented Riemann surface is given by the quadratic form

$$q : H_1(M; \mathbb{Z}_2) \rightarrow \mathbb{Z}_2,$$

where, briefly speaking, $p(w) = (-1)^{q(w)}$ is the monodromy of a spinor under a shift by w . Recall that a form on $H_1(M; \mathbb{Z}_2)$ is called quadratic if

$$q(w_1 + w_2) = q(w_1) + q(w_2) + w_1 \cdot w_2,$$

where $w_1 \cdot w_2$ is the intersection (modulo two) of the cycles w_1 and w_2 .

For example, it is easy to show that \sqrt{dz} is a section of the bundle for which

$$q_0(0) = 0, \quad q_0(e_1) = q_0(e_2) = q_0(e_1 + e_2) = 1.$$

Since the quantities $\psi_1^2 dz$, $\bar{\psi}_2^2 dz$ and $\psi_1 \bar{\psi}_2 dz$ are doubly periodic, then ψ is the Bloch eigenfunction of \mathcal{D} , which, when shifted by periods, is multiplied by $\mu(w) = \pm 1$. Note that the multiplier mapping is given by the homomorphism

$$\nu : H_1(M; \mathbb{Z}_2) \rightarrow \mathbb{Z}_2, \quad \mu(w) = (-1)^{\nu(w)}.$$

The spinors $\psi_1\sqrt{dz}$ and $\bar{\psi}_2\sqrt{dz}$ are sections of the spinor bundle, which corresponds to the quadratic form

$$q = q_0 + \nu.$$

There are four different homomorphisms $H_1(M; \mathbb{Z}_2) \rightarrow \mathbb{Z}_2$, which, using the previous formula, define four different quadratic forms on $H_1(M; \mathbb{Z}_2)$ (and thereby spinor structures on M).

From the above it follows that the spectral variety is determined independently of the spinor structure.

In [20] we introduced the spectral curve of an immersed torus as the Bloch variety of \mathcal{D} with doubly periodic potential $U(x)$ and at zero energy level $E = 0$. Bloch eigenfunctions are solutions to the equations

$$\mathcal{D}\psi = 0, \quad \psi(x + e_j) = e^{2\pi i \langle k, e_j \rangle} \psi(x) = \mu(e_j) \psi(x), \quad j = 1, 2.$$

After replacement

$$\psi \rightarrow \varphi = e^{-\alpha/2} \psi$$

this Dirac equation reduces to

$$D\varphi = 0,$$

where the operator D has the form (11) and

$$V = \begin{pmatrix} 2Ue^{-\alpha/2} & 0 \\ 0 & 2Ue^{-\alpha/2} \end{pmatrix} = \begin{pmatrix} H & 0 \\ 0 & H \end{pmatrix}.$$

It follows from Theorem 4 (after the obvious linear transformation $\kappa \rightarrow k$) that such solutions exist, exactly, for $\mathcal{P}(k_1, k_2) = 0$, where \mathcal{P} is some entire function. Since the multipliers μ are invariant under the shift k by vectors from the dual lattice:

$$\Lambda^* = \{v : \langle v, e_k \rangle \in \mathbb{Z}, k = 1, 2\},$$

then after factorization

$$S = \{\mathcal{P} = 0\} \rightarrow \{\mathcal{P} = 0\}/\Lambda = \Gamma$$

we obtain a Riemann surface Γ , which parametrizes all Bloch eigenfunctions of the operator \mathcal{D} .

This spectral curve is “almost” preserved under conformal transformations of the ambient space. Namely, the multipliers are preserved. The simplest proof is the following: it is quite obvious that the multipliers are preserved when \mathbb{R}^3 rotates (they reduce to the actions of $SO(3)$ on ψ). The inversion in terms of (13) takes the form

$$X \rightarrow X^{-1}$$

and there is a Moutard transformation connecting the Weierstrass representations of tori and their Bloch eigenfunctions while preserving the multipliers [23]. At the same time, [13] contains examples where inversion leads to the appearance of double points on the spectral curve.

For very large quasimomenta, the spectral curve “approaches” the spectral curve in the case of zero potential:

$$k_2 \approx \pm ik_1.$$

Near the asymptotic end, where $k_2 \approx ik_1$ we can introduce such a local parameter λ^{-1} that

$$(14) \quad \mu(v) = \lambda v + \frac{C_0 \bar{v}}{\lambda} + O(\lambda^{-2}),$$

where

$$(15) \quad C_0 = -\frac{1}{\text{Area}(\mathbb{C}/\Lambda)} \int_{\mathbb{C}/\Lambda} U^2 = -\frac{1}{4 \text{Area}(\mathbb{C}/\Lambda)} \int_M H^2 d\mu$$

is $\int_M H^2 d\mu$ is the Willmore functional of M .

This observation formed the basis of our proposed “spectral” approach to Willmore’s conjecture, which remained unrealized (see discussion in [19, 21, 24]).

Note that the spectral curves of immersed tori can be reducible [4].

All these statements are also true for surfaces in \mathbb{R}^4 . Only in this case can the potential $U(x)$ be complex-valued and the Weierstrass representation is given by a pair of spinors ψ and ϕ such that $\mathcal{D}\psi = 0$ and $\mathcal{D}^\vee\phi = 0$, where the operator \mathcal{D}^\vee is obtained from \mathcal{D} by permuting

U and \bar{U} on the diagonal in the matrix coefficient (see, for example, [21, 22]). The Moutard transformation corresponding to inversions in \mathbb{R}^4 was derived in [13].

5.3. Spectral varieties of higher genera surfaces. The arguments from §5.2 extend to surfaces of large genus immersed in \mathbb{R}^3 and \mathbb{R}^4 . We have

Theorem 5. *If D is a Dirac operator of the form (11), included in the Weierstrass representation closed oriented surface of genus g immersed into \mathbb{R}^3 or \mathbb{R}^4 , then analytic set*

$$\{\mathcal{P}(0, \kappa_1, \dots, \kappa_{2g}) = 0\},$$

defined by Theorem 4 and factorized by the action $\kappa \rightarrow \kappa + 2\pi n, n \in \mathbb{Z}$, is the spectral variety of Γ . The multiplier function $\mu = (e^{i\kappa_1}, \dots, e^{i\kappa_{2g}})$ invariant under conformal transformations of the ambient space.

Probably these complex manifolds should contain information about the Willmore functional of the surface similar to that given in (14) and (15). However, already for $g = 2$ the spectral variety has a complex dimension equal to three. If a surface of large genus has symmetries, then this should lead to special classes of spectral varieties.

The work was carried out with the support of the Mathematical Center in Akademgorodok, agreement with the Ministry of Science and Higher Education of the Russian Federation No. 075-15-2022-281.

REFERENCES

- [1] Aharonov, Y., Bohm, D.: Significance of electromagnetic potentials in quantum theory. *Physical Review*, **115** (3) (1959), 485–491.
- [2] Aubin, T.: *Some Nonlinear Problems in Riemannian Geometry*. Springer, Berlin, 1998.
- [3] Bohle, C., Taimanov, I.A.: Spectral curves for Cauchy-Riemann operators on punctured elliptic curves. *Functional Anal. Appl.* **47**:4 (2013), 319–322.
- [4] Bohle, Ch., Taimanov, I.A.: Euclidean minimal tori with planar ends and elliptic solitons. *Int. Math. Res. Notices*, 2015, no. 14, 5907–5932.
- [5] Veliev, O.A.: Asymptotic formulas for the eigenvalues of a periodic Schrödinger operator and the Bethe–Sommerfeld conjecture. *Funct. Anal. Appl.* **21**:2 (1987), 87–100.
- [6] Veliev, O.A.: Perturbation theory for the periodic multidimensional Schrödinger operator and the Bethe–Sommerfeld conjecture. *Int. J. Contemp. Math. Sci.* **2** (2007), no.1-4, 19–87.
- [7] Dubrovin, B.A., Krichever, I.M., Novikov, S.P.: The Schrödinger equation in a periodic field and Riemann surfaces. *Soviet Math. Dokl.* **17** (1976), 947–952.

- [8] Dubrovin, B.A., Matveev, V.B., Novikov, S.P.: Non-linear equations of Korteweg–de Vries type, finite-zone linear operators, and Abelian varieties. *Russian Math. Surveys* **31**:1 (1976), 59–146.
- [9] Kha, Minh: Green’s function asymptotics of periodic elliptic operators on abelian coverings of compact manifolds. *J. Funct. Anal.* **274**:2 (2018), 341–387.
- [10] Kobayashi, T., Ono, K., Sunada, T.: Periodic Schrödinger operators on a manifold. *Forum Math.* **1** (1989), 69–79.
- [11] Konopelchenko, B.G.: Induced surfaces and their integrable dynamics. *Stud. Appl. Math.* **96** (1996), 9–52.
- [12] Krichever, I.M.: Spectral theory of two-dimensional periodic operators and its applications. *Russian Math. Surveys* **44**:2 (1989), 145–225.
- [13] Matuev, R.M., Taimanov, I.A.: The Moutard transformation of two-dimensional Dirac operators and the conformal geometry of surfaces in four-space. *Math. Notes* **100**:6 (2016), 835–846.
- [14] Novikov, S.P.: Periodic problem for the Korteweg–de Vries equation. *Functional Anal. Appl.* **8**:3 (1998), 236–246.
- [15] Parnowski, L.: Bethe–Sommerfeld conjecture. *Ann. Henri Poincaré* **9** (2008), no.3, 457–508.
- [16] Skriganov, M.M.: Proof of the Bethe–Sommerfeld conjecture in dimension two. *Dokl. Akad. Nauk SSSR* **248**:1 (1979), 39–42 (in Russian).
- [17] Skriganov, M.M.: The spectrum band structure of the three-dimensional Schrödinger operator with periodic potential. *Invent. Math.* **80**:1 (1985), 107–121.
- [18] Sunada, T.: A periodic Schrödinger operator on an abelian cover. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **37** (3) (1990), 575–583.
- [19] Taimanov, I.A.: Modified Novikov–Veselov equation and differential geometry of surfaces. *Amer. Math. Soc. Transl. Ser. 2, V. 179, 1997, P. 133–151* [arXiv:dg-ga/9511005].
- [20] Taimanov, I.A.: The Weierstrass representation of closed surfaces in \mathbb{R}^3 . *Functional Anal. Appl.* **32**:4 (1998), 258–267.
- [21] Taimanov, I.A.: Two-dimensional Dirac operator and surface theory. *Russian Math. Surveys* **61**:1 (2006), 79–159.
- [22] Taimanov, I.A.: Surfaces in the four-space and the Davey-Stewartson equations. *J. of Geometry and Physics*, **56** (2006), 1235–1256.
- [23] Taimanov, I.A.: The Moutard transformation of two-dimensional Dirac operators and the Möbius geometry. *Math. Notes* **97**:1 (2015), 124–135.
- [24] Taimanov, I.A.: Surfaces via spinors and soliton equations. *Proc. Int. Cong. Math. 2022, Vol.4, 2638-2654, EMS Press, Berlin, 2023.*
- [25] Taimanov, I.A.: Geometry and quasiclassical quantization of magnetic monopoles. *Theoret. and Math. Phys* **218**:1 (2024), 129–144.

SOBOLEV INSTITUTE OF MATHEMATICS, NOVOSIBIRSK, 630090 RUSSIA
Email address: taimanov@math.nsc.ru