

# Common substring with shifts in $b$ -ary expansions

Xin Liao<sup>a</sup>, Dingding Yu<sup>a,\*</sup>

<sup>a</sup>*School of Mathematics and Statistics, Wuhan University, Wuhan 430072, China*

---

## Abstract

Denote by  $S_n(x, y)$  the length of the longest common substring of  $x$  and  $y$  with shifts in their first  $n$  digits of  $b$ -ary expansions. We show that the sets of pairs  $(x, y)$ , for which the growth rate of  $S_n(x, y)$  is  $\alpha \log n$  with  $0 \leq \alpha \leq \infty$ , have full Hausdorff dimension.

*Keywords:*  $b$ -ary expansions; common substring with shifts; Hausdorff dimension

---

## 1. Introduction

Fix a positive integer  $b \geq 2$ . Every  $x \in (0, 1]$  admits a unique non-terminating  $b$ -ary expansion:

$$x = \sum_{i=1}^{\infty} \frac{x_i}{b^i} := (x_1, x_2, \dots),$$

where  $x_i \in \mathcal{A} := \{0, 1, \dots, b-1\}$  for each  $i \geq 1$ . The infinite sequence  $(x_1, x_2, \dots) \in \mathcal{A}^{\mathbb{N}}$  is called the digit sequence of  $x$ .

For  $(x, y) \in (0, 1] \times (0, 1]$  and  $n \in \mathbb{N}$ , the length of the longest common substring  $L_n(x, y)$ , and the length of the longest common substring with shifts  $S_n(x, y)$  of  $x$  and  $y$  in the first  $n$  digits of  $b$ -ary expansions are defined as

$$\begin{aligned} L_n(x, y) &= \max\{l \geq 1 : x_{i+1}^{i+l} = y_{i+1}^{i+l} \text{ for some } 0 \leq i \leq n-l\}, \\ S_n(x, y) &= \max\{l \geq 1 : x_{i+1}^{i+l} = y_{j+1}^{j+l} \text{ for some } 0 \leq i, j \leq n-l\}, \end{aligned}$$

where  $x_{i+1}^{i+l}$  denotes the substring  $(x_{i+1}, \dots, x_{i+l})$ .

It is worth mentioning that the longest common substring problem concerns the shortest distance between two orbits of a dynamical system  $T : X \rightarrow X$  over a metric space  $(X, d)$ . The shortest distance between two orbits is defined as

$$d_n(x, y) = \min_{i, j=0, \dots, n-1} (d(T^i x, T^j y)).$$

---

\*corresponding author

*Email addresses:* xin\_liao@whu.edu.cn (Xin Liao), yudding\_sgr@whu.edu.cn (Dingding Yu)

Barros, Liao, and Rousseau [1, Section 3] pointed out that  $S_n(x, y) \leq -\log d_n(x, y) \leq S_{2n}(x, y)$  for almost all pairs  $(x, y)$ . Therefore,  $S_n(x, y)$  and  $-\log d_n(x, y)$  have the same asymptotic behavior. Moreover, It is showed [1, Theorem 7] that if the Rényi entropy  $H_2$  exists, then for Lebesgue almost all  $(x, y) \in (0, 1] \times (0, 1]$ ,

$$\lim_{n \rightarrow \infty} \frac{S_n(x, y)}{\log n} = \frac{2}{H_2}.$$

Li and Yang [2] proved that for Lebesgue almost all  $(x, y) \in (0, 1] \times (0, 1]$ ,

$$\lim_{n \rightarrow \infty} \frac{L_n(x, y)}{\log_b n} = 1.$$

Then, it is natural to study the points for which  $L_n(x, y)$  and  $S_n(x, y)$  increase with various speeds. For  $0 \leq \alpha \leq \infty$ , we define the level sets

$$L(\alpha) = \left\{ (x, y) \in (0, 1] \times (0, 1] : \lim_{n \rightarrow \infty} \frac{L_n(x, y)}{\log_b n} = \alpha \right\}.$$

In [2], Li and Yang proved that all these level sets have full Hausdorff dimension.

However, despite a number of contributions dealing with the sets linked with  $L_n(x, y)$ , there is few conclusion taking  $S_n(x, y)$  into account. To fill this gap, we aim to extend Li and Yang's results to the level sets associated with  $S_n(x, y)$ . Different from Li and Yang's method, we mainly exploit the estimation of the spectral radius of the matrix, see Lemma 2.2.

For  $0 \leq \alpha \leq \infty$ , we define

$$S(\alpha) = \left\{ (x, y) \in (0, 1] \times (0, 1] : \lim_{n \rightarrow \infty} \frac{S_n(x, y)}{\log_b n} = \alpha \right\}. \quad (1.1)$$

The following result shows that all the level sets associated with  $b$ -ary expansions have full Hausdorff dimension.

**Theorem 1.1.** *Let  $S(\alpha)$  be defined as in (1.1). Then*

$$\dim_{\mathcal{H}} S(\alpha) = 2, \quad \forall \alpha \in [0, \infty].$$

Denote

$$S(0, \infty) = \left\{ (x, y) \in (0, 1] \times (0, 1] : \liminf_{n \rightarrow \infty} \frac{S_n(x, y)}{\log_b n} = 0, \limsup_{n \rightarrow \infty} \frac{S_n(x, y)}{\log_b n} = \infty \right\}.$$

We also investigate the “sizes” of  $S(\alpha)$  and  $S(0, \infty)$  from a topological point of view.

**Theorem 1.2.** *For  $0 < \alpha < \infty$ , the set  $S(\alpha)$  is of the first category. The set  $S(0, \infty)$  is residual.*

The paper is organized as follows. In the next section, we compile several lemmas crucial for subsequent proofs of the main theorems. In Sections 3 and 4, we give the proofs of Theorem 1.1 and Theorem 1.2 respectively.

## 2. Preliminaries

To let our exposition be self-contained, before giving the proof of Theorem 1.1, we present some useful lemmas.

Let  $\mathbb{M}$  be a subset of  $\mathbb{N}$  and let  $\#$  denote the cardinality of a set. We say that the set  $\mathbb{M}$  is of density 0 if

$$\lim_{n \rightarrow \infty} \frac{\#\{i \in \mathbb{M} : i \leq n\}}{n} = 0.$$

Write  $\mathbb{N} \setminus \mathbb{M} = \{n_1 < n_2 < \dots\}$  and define a self-mapping  $\varphi_{\mathbb{M}}$  on  $(0, 1]$  by

$$0.x_1x_2 \dots \mapsto 0.x_{n_1}x_{n_2} \dots.$$

Let  $J \subset (0, 1]$ . The following lemma describes the relation between the Hausdorff dimensions of  $J$  and  $\varphi_{\mathbb{M}}(J) = \{\varphi_{\mathbb{M}}(x) : x \in J\}$ .

**Lemma 2.1.** (*[3, Lemma 2.3]*) *Suppose that the set  $\mathbb{M}$  is of density zero in  $\mathbb{N}$ . Then, for any set  $J \subset (0, 1]$ , we have  $\dim_{\mathcal{H}} J = \dim_{\mathcal{H}} \varphi_{\mathbb{M}}(J)$ .*

For an  $n \times n$  matrix  $A$  of 0's and 1's having a specified number  $\tau$  of 0's, the next lemma gives a lower bound of the spectral radius of  $A$  when  $\tau \leq \lfloor n/2 \rfloor \lceil n/2 \rceil$ , where  $\lfloor n \rfloor$  denotes the largest integer no larger than  $n$ , and  $\lceil n \rceil$  denotes the smallest integer no smaller than  $n$ . Denote by  $\rho(A) = \lim_{n \rightarrow \infty} \|A^n\|^{1/n}$  the spectral radius of the matrix  $A$ .

**Lemma 2.2.** (*[4, Theorem 2.1]*) *Let  $n$  be a positive integer and let  $\tau$  be an integer with  $0 \leq \tau \leq \lfloor n/2 \rfloor \lceil n/2 \rceil$ . Denote by  $C(n, \tau)$  the class of all  $n \times n$  matrices of 0's and 1's with exactly  $\tau$  0's. Then*

$$\rho(A) \geq \frac{1}{2}(n + \sqrt{n^2 - 4\tau}), \quad \forall A \in C(n, \tau).$$

The following lemma giving a refinement of the Mass Distribution Principle, is usually called Billingsley's lemma. For  $x \in (0, 1]$ , let  $I_m(x)$  denote the  $m$ -th generation, half-open  $b$ -adic interval of the form  $[\frac{j-1}{b^m}, \frac{j}{b^m})$  containing  $x$  and  $|I_m(x)|$  denote the length of the interval  $I_m(x)$ .

**Lemma 2.3.** (*[5, Lemma 1.4.1]*) *Let  $A \subset (0, 1]$  be a Borel set and  $\mu$  be a finite Borel measure on  $(0, 1]$ . Suppose  $\mu(A) > 0$ . If*

$$\beta_1 \leq \liminf_{m \rightarrow \infty} \frac{\log \mu(I_m(x))}{\log |I_m(x)|} \leq \beta_2,$$

for all  $x \in A$ , then  $\beta_1 \leq \dim_{\mathcal{H}}(A) \leq \beta_2$ .

## 3. Proof of Theorem 1.1

Theorem 1.1 follows from a series of propositions.

Let  $p \geq 2$  be a positive integer. Define

$$E_p := \{x \in (0, 1] : x_i = 0, \text{ if } i = kp^2 + j \text{ for some } k \in \mathbb{N} \text{ and } j \in [1, p + 1]\}. \quad (3.1)$$

We obtain the Hausdorff dimension of  $E_p$  by the above Billingsley's lemma immediately.

**Proposition 3.1.** *Let  $E_p$  be defined as in (3.1). Then for any  $p \geq 2$ ,*

$$\dim_{\mathcal{H}}(E_p) = \frac{p^2 - (p + 1)}{p^2}.$$

*Proof.* Denote  $J = \{kp^2 + j : k \in \mathbb{N} \text{ and } j \in [p + 2, p^2]\}$ . Then  $E_p$  can be expressed as

$$E_p = \left\{ x \in (0, 1] : x = \sum_{i \in J} \frac{x_i}{b^i}, x_i \in \mathcal{A} \right\}. \quad (3.2)$$

The set  $E_p$  is covered by exactly  $b^{\#(J \cap \{1, \dots, m\})}$  many closed  $b$ -adic intervals of generation  $m$ . Let  $\mu$  be the probability measure on  $E_p$  that assigns equal measure to the  $m$ -th generation covering intervals. That is, for any  $m \geq 1$ , we have

$$\mu(I_m(x)) = b^{-\#(J \cap \{1, \dots, m\})}.$$

This measure makes the digits  $\{x_i\}_{i \in J}$  in (3.2) to be independent and identically distributed uniform random bits. For any  $x \in E_p$ ,

$$\frac{\log \mu(I_m(x))}{\log |I_m(x)|} = \frac{\log b^{-\#(J \cap \{1, \dots, m\})}}{\log b^{-m}} = \frac{\#(J \cap \{1, \dots, m\})}{m}.$$

Obviously,  $\lim_{m \rightarrow \infty} \frac{\#(J \cap \{1, \dots, m\})}{m} = \frac{p^2 - (p + 1)}{p^2}$ . Then by Billingsley's lemma 2.3, we obtain

$$\dim_{\mathcal{H}} E_p = \lim_{m \rightarrow \infty} \frac{\log \mu(I_m(x))}{\log |I_m(x)|} = \frac{p^2 - (p + 1)}{p^2}.$$

□

Define

$$F_p := \{x \in (0, 1] : x_i^{i+p} \neq 0^{p+1}, \forall i \geq 1\}, \quad (3.3)$$

where  $0^{p+1}$  denotes the substring  $(0, 0, \dots, 0)$  with  $p + 1$  zeros. The following lemma gives the Hausdorff dimension of  $F_p$ .

**Proposition 3.2.** *Let  $F_p$  be defined as in (3.3). Then for any  $p \geq 2$ ,*

$$\dim_{\mathcal{H}} F_p = \frac{\log \rho(A)}{\log b^p},$$

where  $A$  is a matrix of size  $b^p \times b^p$ , with  $pb^{p-1} - (p - 1)b^{p-2}$  zeros and all other entries equal to one.

*Proof.* For any  $i \geq 1$  and  $x_i^{2^{p+i-1}} \in \mathcal{A}^{2^p}$ , there exist non-negative integers  $0 \leq x'_i, x'_{i+1} \leq b^p - 1$

such that  $\sum_{j=i}^{2p+i-1} \frac{x_j}{b^j} = \frac{x'_i}{b^{ip}} + \frac{x'_{i+1}}{b^{2ip}}$ , which means every  $2p$  digits in  $b$ -ary expansion can be expressed as 2 digits in  $b^p$ -ary expansion. Without loss of generality, we can assume  $i = 1$ .

- (1) If  $x_1^{p+1} = 0^{p+1}$ , then  $x'_1 = 0$  and  $x'_2 \in \{0, \dots, b^{p-1} - 1\}$ . Thus  $(x'_1, x'_2)$  has  $b^{p-1}$  values.
- (2) If  $x_2^{p+2} = 0^{p+1}$ , then  $x'_1 \in \{0, \dots, b - 1\}$  and  $x'_2 \in \{0, \dots, b^{p-2} - 1\}$ , when  $x'_1 = 0$ , the values of  $(x'_1, x'_2)$  have already been counted in case (1), so there are  $(b-1)b^{p-2}$  values for  $(x'_1, x'_2)$ , which are different from case (1).
- ...
- (3) For any  $2 \leq k < p$ , if  $x_{k+1}^{p+k+1} = 0^{p+1}$ , then  $x'_1 \in \{0, \dots, b^k - 1\}$  and  $x'_2 \in \{0, \dots, b^{p-k-1} - 1\}$ . When  $x'_1 \in \{0, \dots, b^{k-1} - 1\}$ , the values of  $(x'_1, x'_2)$  have been appeared in the previous cases, so there are  $(b^k - b^{k-1})b^{p-k-1} = (b-1)b^{p-2}$  distinct values for  $(x'_1, x'_2)$ .

If there exists  $0^{p+1}$  in  $x_1 \dots x_{2p}$ , then from the preceding discussion, we deduce that  $(x'_1, x'_2)$  can take  $pb^{p-1} - (p-1)b^{p-2}$  distinct values. Denote the set of these  $(x'_1, x'_2)$  by  $\Lambda((x'_1, x'_2))$ . Then

$$\#\Lambda((x'_1, x'_2)) = pb^{p-1} - (p-1)b^{p-2}.$$

Define a  $b^p \times b^p$  matrix  $A = (A_{ij})$ ,  $0 \leq i, j \leq b^p - 1$  as follows:

$$A_{ij} = \begin{cases} 0, & \text{if } (i, j) \in \Lambda((x'_1, x'_2)); \\ 1, & \text{otherwise.} \end{cases}$$

Then, the number of 0's in matrix  $A$  is  $pb^{p-1} - (p-1)b^{p-2}$ . Hence,  $F_p$  can be expressed as

$$F_p = \left\{ x \in (0, 1] : x = \sum_{i=1}^{\infty} \frac{x'_i(x)}{b^{ip}}, A_{x'_i x'_{i+1}} = 1, \text{ for all } i \right\}.$$

Using the theory of shifts of finite types (see [5, Example 1.3.3] for more details), we can calculate the Hausdorff dimension of  $F_p$ :

$$\dim_{\mathcal{H}} F_p = \frac{\log \rho(A)}{\log b^p}.$$

□

*Proof of Theorem 1.1.* First, let us focus on the case  $0 \leq \alpha < \infty$ .

For  $k \geq 1$ , let  $m_k = p^2 \cdot 2^k$ ,  $\ell_k = \lfloor \alpha(k+1) \log_b 2 \rfloor$  and  $t_k = \sum_{i=1}^k (m_i + \ell_i)$ . Define self-mapping

$f_k$  on  $(0, 1]$  by

$$\begin{aligned}
f_1(x) &= \sum_{i=1}^{m_1} \frac{x_i}{b^i} + \sum_{i=m_1+1}^{t_1} \frac{1}{b^i} + \sum_{i=t_1+1}^{\infty} \frac{x_{i-\ell_1}}{b^i} \\
&\dots \\
f_k(x) &= \sum_{i=1}^{t_{k-1}} \frac{x_i}{b^i} + \sum_{i=t_{k-1}+1}^{t_{k-1}+\ell_k} \frac{1}{b^i} + \sum_{i=t_{k-1}+\ell_k+1}^{\infty} \frac{x_{i-\ell_k}}{b^i}.
\end{aligned}$$

Then, for any  $x \in (0, 1]$ , the limit

$$f(x) := \lim_{k \rightarrow \infty} f_k(f_{k-1}(\dots f_1(x))), \quad (3.4)$$

exists.

For each  $x \in f(E_p)$  and  $y \in f(F_p)$ , if  $t_k \leq n < t_{k+1}$ , then

$$\ell_k \leq S_n(x, y) < \ell_{k+1} + 2p^2.$$

It follows that

$$\lim_{n \rightarrow \infty} \frac{S_n(x, y)}{\log_b n} = \alpha,$$

and hence

$$f(E_p) \times f(F_p) \subset S(\alpha).$$

Therefore, using [6, Product formula 7.2], we have

$$\dim_{\mathcal{H}} S(\alpha) \geq \dim_{\mathcal{H}} f(E_p) + \dim_{\mathcal{H}} f(F_p).$$

On the other hand, since  $\lim_{k \rightarrow \infty} \frac{\sum_{i=1}^k \ell_i}{t_k} = 0$ , the assumption of Lemma 2.1 is satisfied for the sets  $E_p$  and  $F_p$ . Hence, we obtain

$$\begin{aligned}
\dim_{\mathcal{H}} f(E_p) &= \dim_{\mathcal{H}} E_p, \\
\dim_{\mathcal{H}} f(F_p) &= \dim_{\mathcal{H}} F_p.
\end{aligned}$$

Therefore, utilizing Proposition 3.1 and 3.2, we have

$$\dim_{\mathcal{H}} S(\alpha) \geq \frac{p^2 - p - 1}{p^2} + \frac{\log \rho(A)}{\log b^p},$$

where  $A$  is defined as in Proposition 3.2. By Lemma 2.2, we have

$$\rho(A) \geq \frac{1}{2} \left( b^p + \sqrt{b^{2p} - 4(pb^{p-1} - (p-1)b^{p-2})} \right).$$

Taking  $p \rightarrow \infty$  yields

$$\dim_{\mathcal{H}} S(\alpha) \geq \lim_{p \rightarrow \infty} \frac{p^2 - p - 1}{p^2} + \frac{\log \rho(A)}{\log b^p} = 2.$$

Thus, we have shown the desired result for  $0 \leq \alpha < \infty$ .

As for  $\alpha = \infty$ , the proof needs to be modified accordingly. For this case, we define  $m_k = p^2 \cdot 2^k$  and  $\ell_k = k^2$  for  $k \geq 1$ . Then, the rest of the proof proceeds as before.  $\square$

#### 4. Proof of Theorem 1.2

For  $\alpha \in [0, \infty]$ , set

$$\underline{S}^*(\alpha) = \left\{ (x, y) \in (0, 1]^2 : \liminf_{n \rightarrow \infty} \frac{\log S_n(x, y)}{\log_b n} \leq \alpha \right\},$$

$$\overline{S}_*(\alpha) = \left\{ (x, y) \in (0, 1]^2 : \limsup_{n \rightarrow \infty} \frac{\log S_n(x, y)}{\log_b n} \geq \alpha \right\}.$$

The proof of Theorem 1.2 consists of several claims.

**Claim 1:** For any  $0 \leq \alpha \leq \infty$ ,  $S(\alpha)$  is dense in  $(0, 1]^2$ .

*Proof.* For any  $\alpha \in [0, \infty]$ , there exists  $(x', y') \in (0, 1]^2$  such that

$$\lim_{n \rightarrow \infty} \frac{\log S_n(x', y')}{\log_b n} = \alpha.$$

For any  $(x, y) \in (0, 1]^2$ , we can find a sequence of points in  $S(\alpha)$ ,

$$\tilde{x}_k := \sum_{i=1}^k \frac{x_i}{b^i} + \sum_{i=k+1}^{\infty} \frac{x'_i}{b^i},$$

$$\tilde{y}_k := \sum_{i=1}^k \frac{y_i}{b^i} + \sum_{i=k+1}^{\infty} \frac{y'_i}{b^i},$$

such that  $(\tilde{x}_k, \tilde{y}_k) \rightarrow (x, y)$  as  $n \rightarrow \infty$ . Indeed, for any  $k \geq 1$ ,  $\tilde{x}_k$  ( $\tilde{y}_k$ , respectively) and  $x'$  ( $y'$ , respectively) differ only in finitely many digits. Then

$$\lim_{n \rightarrow \infty} \frac{\log S_n(\tilde{x}_k, \tilde{y}_k)}{\log_b n} = \lim_{n \rightarrow \infty} \frac{\log S_n(x', y')}{\log_b n} = \alpha.$$

Thus  $S(\alpha)$  is dense in  $(0, 1]^2$ .  $\square$

**Claim 2:** For any  $0 < \alpha < \infty$ ,  $\underline{S}^*(\alpha)$  and  $\overline{S}^*(\alpha)$  are residual.

*Proof.* Let  $\alpha \in (0, \infty)$  be fixed. From Claim 1, we can deduce that  $\underline{S}^*(\alpha)$  and  $\overline{S}^*(\alpha)$  are dense. Since  $(0, 1]^2$  is a Baire space, it suffices to show that  $\underline{S}^*(\alpha)$  and  $\overline{S}^*(\alpha)$  are  $G_\delta$  sets.

We observe that, for any  $k > 0$ ,

$$\underline{S}^*(\alpha) = \bigcap_{k=1}^{\infty} \bigcap_{N=1}^{\infty} \bigcup_{n=N}^{\infty} B_n(\alpha, k),$$

and

$$\overline{S}^*(\alpha) = \bigcap_{k=\lfloor 1/\alpha \rfloor + 1}^{\infty} \bigcap_{N=1}^{\infty} \bigcup_{n=N}^{\infty} \hat{B}_n(\alpha, k),$$

where  $B_n(\alpha, k)$  and  $\hat{B}_n(\alpha, k)$  are defined by

$$B_n(\alpha, k) := \{(x, y) \in (0, 1]^2 : S_n(x, y) < n^{\alpha+1/k}\},$$

and

$$\hat{B}_n(\alpha, k) := \{(x, y) \in (0, 1]^2 : S_n(x, y) > n^{\alpha-1/k}\}.$$

All non-empty sets  $B_n(\alpha, k)$  and  $\hat{B}_n(\alpha, k)$  are open sets, implying that  $\underline{S}^*(\alpha)$  and  $\overline{S}^*(\alpha)$  are  $G_\delta$  sets. □

*Proof of Theorem 1.2.* For any  $K \in \mathbb{N}$ , we have

$$\underline{S}^*(0) = \bigcap_{K=1}^{\infty} \underline{S}^*(1/K) \quad \text{and} \quad \overline{S}^*(\infty) = \bigcap_{K=1}^{\infty} \overline{S}^*(K).$$

From Claim 2, we conclude that  $\underline{S}^*(1/K)$  and  $\overline{S}^*(K)$  are residual. Then  $\underline{S}^*(0)$  and  $\overline{S}^*(\infty)$  are residual. Hence the set

$$S(0, \infty) = \underline{S}^*(0) \cap \overline{S}^*(\infty)$$

is residual.

For any  $0 < \alpha < \infty$ , we have

$$S(\alpha) \subset \left( \underline{S}^*(0) \cap \overline{S}^*(\infty) \right)^c.$$

By the definition of the set of the first category, we have  $S(\alpha)$  is of first category. □

## Acknowledgments

The authors wish to express their sincere appreciation to Professor Lingmin Liao who critically read the paper and made numerous helpful suggestions.



## References

- [1] Barros, Vanessa and Liao, Lingmin and Rousseau, Jérôme. On the shortest distance between orbits and the longest common substring problem. *Advances in Mathematics*, 344: 311–339, 2019.
- [2] Li, Jinjun and Yang, Xiangfeng. On longest matching consecutive subsequence. *International Journal of Number Theory*, 15(08): 1745–1758, 2019.
- [3] Chen, Haibo and Tang, Junmin. The waiting spectra of the sets described by the quantitative waiting time indicators. *Science China Mathematics*, 57(11): 2335–2346, 2014.
- [4] Brualdi, Richard A and Solheid, Ernie S. On the minimum spectral radius of matrices of zeros and ones. *Linear Algebra and its Applications*, 85: 81–100, 1987.
- [5] Bishop, Christopher J and Peres, Yuval. *Fractals in probability and analysis*. Cambridge University Press, 162, 2017.
- [6] Falconer, Kenneth. *Fractal geometry: Mathematical Foundations and Applications*. John Wiley & Sons, 2004.