

A PRIORI ESTIMATES FOR PARABOLIC MONGE-AMPÈRE TYPE EQUATIONS

YANG ZHOU AND RUIXUAN ZHU

ABSTRACT. We prove the existence and regularity of convex solutions to the first initial-boundary value problem of the parabolic Monge-Ampère equation

$$\begin{cases} u_t = \det D^2 u & \text{in } Q_T, \\ u = \phi & \text{on } \partial_p Q_T, \end{cases}$$

where ϕ is a smooth function, $Q_T = \Omega \times (0, T]$, $\partial_p Q_T$ is the parabolic boundary of Q_T , and Ω is a uniformly convex domain in \mathbb{R}^n with smooth boundary. Our approach can also be used to prove similar results for γ -Gauss curvature flow with any $0 < \gamma \leq 1$.

1. INTRODUCTION

In this paper, we study the first initial-boundary value problem of the parabolic Monge-Ampère equation

$$\begin{cases} u_t = \det D^2 u & \text{in } Q_T, \\ u = \phi & \text{on } \partial_p Q_T, \end{cases} \quad (1.1)$$

where $Q_T = \Omega \times (0, T]$, $\partial_p Q_T$ is the parabolic boundary of Q_T , and Ω is a uniformly convex domain in \mathbb{R}^n with smooth boundary. Assume that ϕ is a smooth function in \overline{Q}_T with $\phi_t \geq c_0 > 0$ on $\partial\Omega \times [0, T]$ and $\phi(\cdot, 0)$ uniformly convex. We say a function $f : (x, t) \in \mathbb{R}^n \times \mathbb{R}^+ \rightarrow f(x, t) \in \mathbb{R}$, is parabolically convex if it is continuous and convex in x , and nondecreasing in t . If f is further uniformly convex in x , we say f uniformly parabolically convex.

We consider solutions in the following Banach space.

2000 *Mathematics Subject Classification.* 35K96, 35B65.

Key words and phrases. Parabolic Monge-Ampère equation, Gauss Curvature Flow, regularity.

The second author was supported by NSFC 12141103 and the China Scholarship Council during a visit to the Australian National University.

Definition 1.1. For $Q = \Omega \times I$, where $\Omega \subset \mathbb{R}^n$ is an open convex domain, and $I = (a, b]$, we denote by $\tilde{C}^{k,\alpha}(\overline{Q})$ the Banach space with the norm

$$\|w\|_{\tilde{C}^{k,\alpha}(\overline{Q})} = \sup_{\substack{|\gamma|+2s \leq k \\ (x,t) \in Q}} |D^\gamma D_t^s w(x,t)| + \sup_{\substack{|\gamma|+2s=k \\ (x,t),(y,\tau) \in Q}} \frac{|D^\gamma D_t^s w(x,t) - D^\gamma D_t^s w(y,\tau)|}{(|x-y|^2 + |t-\tau|)^{\alpha/2}}.$$

We shall prove

Theorem 1.2. *Assume that equation (1.1) satisfies the compatibility condition of any order (see Remark 3.2) and the above assumptions. Then there exists a unique smooth, parabolically convex solution u to the first initial-boundary value problem (1.1).*

The proof of Theorem 1.2 depends on the a priori global $\tilde{C}^{2,\alpha}$ estimates. Specifically, we have the following theorem:

Theorem 1.3. *Let $u \in \tilde{C}^4(\overline{Q}_T)$ be a parabolically convex solution of equation (1.1). Assume that Ω is a uniformly convex domain in \mathbb{R}^n with C^4 boundary, $\phi \in \tilde{C}^4(\overline{Q}_T)$, $\phi_t \geq c_0 > 0$ on $\partial\Omega \times [0, T]$, and $\phi(\cdot, 0)$ is uniformly convex. Then it holds*

$$\|u\|_{\tilde{C}^{2,\alpha}(\overline{Q}_T)} \leq C,$$

for some positive constants C and $\alpha \in (0, 1)$ depending only on $n, \Omega, c_0, \|\phi\|_{\tilde{C}^4(\overline{Q}_T)}$ and the uniform convexity of $\phi(\cdot, 0)$.

We can also prove Theorem 1.2 and 1.3 for Gauss curvature flow with boundary

$$\begin{cases} u_t = \frac{\det D^2 u}{(1 + |Du|^2)^{(n+1)/2}} & \text{in } Q_T, \\ u = \phi & \text{on } \partial_p Q_T. \end{cases} \quad (1.2)$$

Monge-Ampère equations figure in various geometric problems such as the Minkowski problem and geometric flows. The elliptic Monge-Ampère equations have received considerable study since the 1970s. One can refer to [10, 29] and references therein for more related works. In a celebrated work [2], Caffarelli, Nirenberg, and Spruck resolved the existence and global $C^{2,\alpha}$ regularity of smooth solutions to the Dirichlet problem of Monge-Ampère equations.

Parabolic Monge-Ampère equations occur in Gauss curvature flows, which were first introduced by Firey in [11] to model the shapes of worn stones. In 1976, Krylov proposed three parabolic analogs of Monge-Ampère equations [20], and the regularity problem of those equations has been studied by many researchers.

See [4, 6, 14, 16, 18, 24, 27, 30, 31, 32, 33] for regularity results of smooth and viscosity solutions, [15, 34, 35] for the Liouville problem, and [1, 7, 8] for the Gauss curvature flow with flat sides. In a remarkable work [30], Tso obtained the interior $\tilde{C}^{2,\alpha}$ estimate for the equation

$$\begin{cases} -u_t \det D^2 u = f, \\ u(x, 0) = h, \quad x \in \mathbb{R}^n. \end{cases}$$

More regularity results of the operator $\mathcal{M}u = -u_t \det D^2 u$ can be found in [16, 24]. There are also many regularity results regarding the parabolic Monge-Ampère operators $\mathcal{M}u = -u_t + (\det D^2 u)^{1/n}$ and $\mathcal{M}u = -u_t + \log \det D^2 u$. The global $\tilde{C}^{2,\alpha}$ regularity of solutions to the first initial-boundary value problems of these two operators were proved respectively in [18] and [31]. One can also find some other results in [5, 17, 26, 34].

However, the global $\tilde{C}^{2,\alpha}$ regularity for equations (1.1) and (1.2) remains open for these two classical parabolic Monge-Ampère equations. Equation (1.2) was studied in [25], where the author where the author proved the existence of a kind of self-similar solutions and obtained its gradient estimate when $\phi \equiv 0$. While the higher global regularity is open. The main obstacle is the lack of concavity of the equations.

It is worth to point out that our approach to obtaining the global C^2 and $\tilde{C}^{2,\alpha}$ estimates requires a combination of corresponding boundary estimates for equation (1.1) and interior estimates for equation (1.5), the equation of the Legendre transform of u , rather than studying only one of the equations to obtain it. Here the interior estimates of equation (1.5) depends on the concavity of this equation.

The proof of Theorem 1.3 can be divided into the following three steps:

Step 1: Establish the C^2 estimate on boundary

We first obtain the C^0 and C^1 estimates, and

$$C^{-1} \leq u_t \leq C, \tag{1.3}$$

for some positive constant C . Then, we obtain the bound of $D^2 u$ on the parabolic boundary by using the method in [2]. Thus, by (1.3) we have the estimates for the eigenvalues

$$\lambda(D^2 u) \geq C \quad \text{on} \quad \partial_p Q_T. \tag{1.4}$$

Step 2: Establish the global C^2 estimate by the duality method

Due to the lack of concavity, we cannot use the maximum principle to control the global maximum of $|D^2u|$ by $\sup_{\partial_p Q_T} |D^2u|$. Instead, we apply the estimates in **Step 1** to the equation after Legendre transform.

Let U be the Legendre transform of u , given by

$$U(y, t) = \sup\{y \cdot x - u(x, t) : x \in \Omega\}.$$

Denote $Q_T^* = \{(y, t) : y \in Du(\Omega, t), t \in (0, T]\}$. Then U satisfies

$$-U_t \det D^2U = 1 \quad \text{in } Q_T^*. \quad (1.5)$$

Since the above equation is concave, one can infer

$$\sup_{Q_T^*} |D^2U| \leq \sup_{\partial_p Q_T^*} |D^2U|,$$

which combined with (1.3) and (1.4) yields an upper bound of eigenvalues $\lambda(D^2U)$ in Q_T^* . Then, by duality we have $\lambda(D^2u) \geq C$ in Q_T . By (1.3), it follows that

$$C^{-1} \leq \lambda(D^2u) \leq C \quad \text{in } Q_T. \quad (1.6)$$

Step 3: Establish the global $\tilde{C}^{2,\alpha}$ estimate

According to **Step 2**, equation (1.5) is uniformly parabolic. Since the equation is also concave, we can obtain the space Hölder estimate for $D^2U(\cdot, t)$ for each t by the Evans-Krylov Theorem. The interior $\tilde{C}^{2,\alpha}$ regularity then follows by [9, 21] and the interior higher-order regularity of U also follows by the bootstrap technique.

As for the boundary estimate, we first use Krylov-Safonov's estimate (see [22]) to get the global space-time Hölder estimate for u_t . Then by differentiating the equations along tangential directions τ_k , we shall prove the $\tilde{C}^{1,\alpha}$ regularity of $D_{\tau_k}u$ on the lateral surface of the parabolic boundary, followed by the space-time Hölder estimate for second normal derivatives.

This paper is arranged as follows. In Section 2, we establish the *a priori* estimates in $\tilde{C}^{2,\alpha}(\Omega \times [\epsilon, T])$. We then obtain the existence and uniqueness of solutions in Section 3, completing the proof of Theorems 1.3 and 1.2. Finally, we prove the existence of a unique smooth solution to equation (1.2) in Section 4.

2. THE A PRIORI ESTIMATE

We first consider the parabolic Monge-Ampère equation

$$\begin{cases} u_t = \det D^2 u & \text{in } Q_T, \\ u = \phi & \text{on } \partial_p Q_T, \end{cases} \quad (2.1)$$

where $Q_T = \Omega \times (0, T]$, $\partial_p Q_T$ is the parabolic boundary of Q_T and Ω is a uniformly convex domain in \mathbb{R}^n with smooth boundary. Assume that ϕ is a smooth function in \overline{Q}_T with $\phi_t \geq c_0 > 0$ on $\partial\Omega \times [0, T]$ and $\phi(\cdot, 0)$ uniformly convex.

In this section, we suppose that $u \in \tilde{C}^4(\overline{Q}_T)$ is a parabolically convex solution to equation (2.1) and aim to prove the $\tilde{C}^{2,\alpha}$ a priori estimate to u . For this purpose, we shall obtain the C^2 estimate for u on the parabolic boundary first.

2.1. Step 1: C^2 estimates on the boundary.

At first, we obtain the C^0 and C^1 estimates, stated as the lemma below.

Lemma 2.1. *$u(\cdot, t)$ is strictly convex for any $t \in [0, T]$, and the following inequalities hold,*

$$\|u\|_{C^1(\overline{Q}_T)} \leq C, \quad (2.2)$$

and

$$1/C \leq u_t \leq C. \quad (2.3)$$

Here C is a universal positive constant which only depends on n , Ω , c_0 , $\|\phi\|_{C^2(\overline{Q}_T)}$ and the uniform convexity of $\phi(\cdot, 0)$.

Proof. By the convexity assumption of u , $\max_{\overline{Q}_T} u = \max_{\partial_p Q_T} u = \max_{\partial_p Q_T} \phi$. Suppose that u reaches its minimum in \overline{Q}_T at point $p \in Q_T$. Then for sufficiently small $\epsilon > 0$, $u_\epsilon = u + t\epsilon$ also attains its minimum at $p_\epsilon \in Q_T$. Then we have $\partial_t u_\epsilon(p_\epsilon) \leq 0$ and $D^2 u_\epsilon(p_\epsilon) \geq 0$. It gives that

$$u_t(p_\epsilon) < 0 \text{ and } D^2 u(p_\epsilon) \geq 0,$$

which leads to a contradiction. Thus we have $\min_{\overline{Q}_T} u = \min_{\partial_p Q_T} u = \min_{\partial_p Q_T} \phi$. Hence $\|u\|_{C^0(\overline{Q}_T)} \leq \|\phi\|_{C^0(\overline{Q}_T)} \leq C$.

As for the first derivative estimate of u w.r.t. t , note that $1/C \leq u_t \leq C$ on $\partial\Omega \times [0, T]$ for some constant C due to the boundary condition. Here C depends on the maximum and minimum value of ϕ_t on $\partial\Omega \times [0, T]$. On $\Omega \times \{0\}$, $u_t = \det D^2 u = \det D^2 \phi$ is also bounded away from zero and infinity thanks to the

uniform convexity and C^2 boundedness of ϕ . Letting $h = u_t$ and differentiating the equation (2.1) w.r.t. t on both sides, we have

$$h_t = hu^{ij}h_{ij} \quad \text{in } Q_T.$$

We claim that h reaches its positive maximum and minimum values on $\partial_p Q_T$. And this results in estimate (2.3). The strict convexity follows directly from (2.3) and smoothness of ϕ , see [3] for a reference. To prove the claim, suppose that h attains its positive maximum in Q_T , so is $h_\epsilon = h - \epsilon t$ for sufficiently small ϵ . Then at that point, $h_\epsilon > 0$, $D^2 h_\epsilon \leq 0$ and $\partial_t h_\epsilon \geq 0$. Consequently we have $h_t > 0$, $h > 0$ and $D^2 h \leq 0$ at that point. This contradicts the equation of h . The proof is the same for the minimum of h . Thus we have proved (2.3).

To estimate $|Du|$, it suffices to prove the bound for $|D_\nu u|$ on $\partial\Omega \times [0, T]$ for the outward unit normal vector ν , since $|Du|$ is bounded by $|D\phi|$ on $\Omega \times \{0\}$ and tangentially on $\partial\Omega \times [0, T]$. For any fixed $t_0 \in [0, T]$, take a smooth function w such that $w = \phi(t_0)$ on $\partial\Omega \times \{t_0\}$, and $\det D^2 w \geq C_2$. Note that such w does exist, and one example is to assume w with $(-d_{\partial\Omega} + d_{\partial\Omega}^2)\delta + \phi$ near the boundary for some positive constant δ large enough. Then by the classical comparison principle for Monge-Ampère equations, we have $u \geq w$ in $\Omega \times \{t_0\}$. Thus $D_\nu u \leq D_\nu w \leq C$, with C depending on the maximum and minimum values of ϕ_t , $D\phi$ on \overline{Q}_T , $D^2\phi(\cdot, 0)$ on $\overline{\Omega}$ and Ω . Till now we have proved formula (2.2). \square

The proof for the second-order derivative estimates on the parabolic boundary follows a similar structure as in Section 2 of [2]. We first establish the boundedness of the second tangential derivatives of u on the boundary, followed by the mixed second derivatives. The boundedness of the second normal derivatives is then derived from these estimates and the equation of u .

Theorem 2.2. *There is a positive constant C such that*

$$1/C \leq \lambda(D^2 u) \leq C \quad \text{on } \partial_p Q_T, \quad (2.4)$$

where $\lambda(D^2 u)$ denotes the eigenvalues of $D^2 u$ and the constant C depends only on $n, \Omega, c_0, \|\phi\|_{\tilde{C}^4(\overline{Q}_T)}$ and the uniform convexity of $\phi(\cdot, 0)$.

Proof. By the boundary condition, the matrix norm of $D^2 u$ is bounded by $D^2\phi(\cdot, 0)$ on $\Omega \times \{0\}$. We also note that by the uniform convexity of ϕ , there exists a positive constant C such that

$$C^{-1}I_n \leq D^2 u(x, 0) \leq CI_n$$

for $x \in \Omega$. Here C depends on $\|D^2\phi(\cdot, 0)\|_{L^\infty(\Omega)}$ and the uniform convexity of $\phi(\cdot, 0)$. Let (x_0, t_0) be an arbitrary point on $\partial\Omega \times [0, T]$, w.l.o.g. $x_0 = 0$ and $\partial\Omega$ be parameterized by $x_n = \rho(x')$ on a small neighborhood of 0 with $\rho(0) = 0$, $D\rho(0) = 0$ and $x_n \geq 0$. We claim that

$$C^{-1}I_{n-1} \leq D_{x'x'}u(0, t_0) \leq CI_{n-1} \quad \text{for any } t_0 \in [0, T]. \quad (2.5)$$

Differentiate the identity $u(x', \rho(x'), t) = \phi(x', \rho(x'), t)$ to obtain

$$(\nabla_{x'} + \nabla\rho\partial_n)(u - \phi) = 0. \quad (2.6)$$

Twice differentiation at point $(0, t_0)$ gives that

$$D_{x'x'}u(0, t_0) = -u_n(0, t_0)D^2\rho(0) + D_{x'x'}\phi(0, t_0) + D^2\rho(0)\phi_n(0, t_0).$$

We can obtain the latter inequality in (2.5) from this formula. The former inequality of (2.5) follows the same proof as in [2] if we regard u_t as a data function at the time t_0 , and the estimation depends on the maximum and minimum values of u_t .

Then we shall prove a uniform bound for the mixed second derivatives of u . Suppose that

$$\rho(x') = \frac{1}{2} \sum_{\alpha, \beta < n} B_{\alpha\beta} x_\alpha x_\beta + O(|x'|^3) \quad (2.7)$$

in a neighborhood of 0. Note that the matrix $B_{\alpha\beta}$ is positive definite because of the uniform convexity of $\partial\Omega$. Denote the operator

$$\mathcal{T} = \partial_\alpha + \sum_{\beta < n} B_{\alpha\beta} (x_\beta \partial_n - x_n \partial_\beta). \quad (2.8)$$

Differentiating equation (2.1), we obtain

$$\mathcal{L}\mathcal{T}u = 0, \quad \text{for the linear operator } \mathcal{L} =: u_t u^{ij} \partial_{ij} - \partial_t.$$

Thus, by (2.3), we have

$$|\mathcal{L}\mathcal{T}(u - \phi)| = |\mathcal{L}\mathcal{T}\phi| = |u_t u^{ij} \partial_{ij} \mathcal{T}\phi - \partial_t \mathcal{T}\phi| \leq \underline{c} \left(1 + \sum_{i=1}^n u^{ii}\right) \quad (2.9)$$

for some positive constant \underline{c} depending on $\|\phi\|_{\tilde{C}^4(\overline{Q}_T)}$ and the positive lower and upper bounds for u_t . Moreover, on $\Omega \times \{0\}$, $\mathcal{T}(u - \phi) = 0$ and

$$|\mathcal{T}(u - \phi)| = \left| - \sum_{\beta < n} B_{\alpha\beta} x_n (u_\beta - \phi_\beta) \right| \leq \bar{c} x_n, \quad \text{on } \partial\Omega \times [0, T]. \quad (2.10)$$

Consider the following barrier function

$$w(x, t) = -a|x|^2 + bx_n.$$

Then $\mathcal{L}w = -2au_t \sum_{i=1}^n u^{ii}$. Note that

$$\sum_{i=1}^n u^{ii} \geq n \left(\prod_{i=1}^n u^{ii} \right)^{1/n} \geq n (\det D^2 u)^{-1/n} \geq nC^{-1/n},$$

where C depends on the maximum of u_t . If we take a big enough, then

$$\mathcal{L}(w \pm \mathcal{T}(u - \phi)) \leq 0.$$

Since Ω is uniformly convex and bounded, there exists some universal constant c such that $x_n \geq c|x|^2$ on $\bar{\Omega}$. Thus we could take $b \geq a/c + \bar{c}$ to derive

$$w \pm \mathcal{T}(u - \phi) \geq 0 \text{ on } \partial_p Q_T.$$

By using the comparison principle for u and w , we obtain that

$$-w \leq \mathcal{T}(u - \phi) \leq w \quad \text{in } \bar{Q}_T.$$

Since $w(0, t) = \mathcal{T}(u - \phi)(0, t) = 0$, this implies that

$$|\partial_n \mathcal{T}(u - \phi)(0, t)| = |u_{\alpha n}(0, t) - \phi_{\alpha n}(0, t)| \leq b,$$

which yields that

$$|u_{\alpha n}(0, t)| \leq b + \|\phi\|_{C^2(\bar{Q}_T)} \leq C. \quad (2.11)$$

Finally, we turn to the second normal derivatives of u . From the equation (2.1), we know that

$$u_t(0, t) = \det D_{x'x'} u(0, t) u_{nn}(0, t) + P(D_{x'x'} u(0, t), D_{x'x_n} u(0, t)),$$

where P is a n -degree polynomial on $D_{x'x'} u(0, t)$ and $D_{x'x_n} u(0, t)$. Combining formula (2.3), (2.5) and (2.11), we obtain a uniform bound for u_{nn} on $\partial\Omega \times [0, T]$.

To conclude, we have proved a uniform bound for the second derivatives of u on $\partial_p Q_T$, i.e.

$$\|D^2 u\| \leq C \quad \text{on } \partial_p Q_T. \quad (2.12)$$

Since u_t is strictly positive and bounded, we get a lower bound for eigenvalues of $D^2 u$ on the parabolic boundary. \square

2.2. Step 2: global C^2 estimates.

Let U be the Legendre transform of u defined on $Q_T^* = \{(y, t) : y \in Du(\Omega, t), t \in (0, T]\}$, i.e.,

$$U(y, t) = \sup_{x \in \Omega} (y \cdot x - u(x, t)).$$

Then U satisfies

$$-U_t \det D^2 U = 1 \quad \text{in } Q_T^*. \quad (2.13)$$

Note that $Du(\Omega, t)$ is not necessarily convex, neither is Q_T^* . Here Q_T^* is bounded and simply connected due to the smoothness of u and formula (2.2). By estimates (2.3) and (2.4) above, we know that

$$C^{-1} \leq -U_t \leq C \quad \text{and} \quad \|D^2 U\|_{L^\infty(\partial_p Q_T^*)} \leq C. \quad (2.14)$$

Lemma 2.3. *If $U \in \tilde{C}^4(\overline{Q_T^*})$ satisfies equation (2.13), it holds that*

$$\sup_{Q_T^*} \|D^2 U\| \leq \sup_{\partial_p Q_T^*} \|D^2 U\|.$$

Proof. Differentiating the equation (2.13), we have

$$\frac{U_{tk}}{U_t} + U^{ij} U_{ijk} = 0 \quad \text{for any } k = 1, \dots, n,$$

and

$$\frac{U_{tkl}}{U_t} - \frac{U_{tk} U_{tl}}{U_t^2} + U^{ij} U_{ijkl} - U_{ijk} U_{lpq} U^{ip} U^{qj} = 0 \quad \text{for any } k, l = 1, \dots, n. \quad (2.15)$$

If we define $\tilde{L} = \frac{1}{U_t} \partial_t + U^{ij} \partial_{ij}$, then $\tilde{L}DU = 0$ and

$$\tilde{L}U_{kk} = \frac{U_{tkk}}{U_t} + U^{ij} U_{ijk} = \frac{U_{tk}^2}{U_t^2} + U_{ijk} U_{pqk} U^{ip} U^{qj}$$

by the identity (2.15). The last term of the above equality is nonnegative since U is convex. Hence we have $\tilde{L}U_{kk} \geq 0$. Note that $U_t < 0$ and $(U^{ij}) \geq 0$, then by the maximum principle for linear parabolic equations, we obtain that

$$\sup_{Q_T^*} U_{kk} = \sup_{\partial_p Q_T^*} U_{kk},$$

or,

$$\sup_{Q_T^*} \|D^2 U\| \leq \sup_{\partial_p Q_T^*} \|D^2 U\|.$$

□

By formula (2.14) and Lemma 2.3, the eigenvalues of D^2u have a positive lower bound in Q_T . Combined with (2.3), they are also bounded from above. To be more explicit, we have proved the following theorem:

Theorem 2.4. *It holds that*

$$1/C \leq \lambda(D^2u) \leq C \quad \text{on } \overline{Q}_T, \quad (2.16)$$

where C is a positive constant depending on $n, \Omega, c_0, \|\phi\|_{\tilde{C}^4(\overline{Q}_T)}$ and the uniform convexity of $\phi(\cdot, 0)$.

2.3. Step 3: global $\tilde{C}^{2,\alpha}$ estimates.

By regularity theory [30] for the equation (2.13), U is smooth in the interior. Then by (2.14), we have the higher regularity for u in Q_T .

Theorem 2.5. *Recall that $u \in \tilde{C}^4(\overline{Q}_T)$ is a parabolically convex solution to equation (2.1). Then, for any $Q \subset\subset Q_T$, we have*

$$\|u\|_{\tilde{C}^4(\overline{Q})} \leq C,$$

where C is a positive constant depending on n, Ω, c_0, ϕ and $\text{dist}(Q, Q_T)$.

Let $h = u_t$, then h satisfies

$$h_t = a^{ij}h_{ij}.$$

Here $a^{ij} = (\det D^2u)u^{ij}$ and h are bounded from below and above by Theorem 2.4. By theories of linear parabolic equations [24], we can obtain the \tilde{C}^α estimate for u_t on \overline{Q}_T .

Theorem 2.6. *Under the same conditions as in Theorem 2.5, we have*

$$\|u_t\|_{\tilde{C}^\alpha(\overline{Q}_T)} \leq C$$

for some positive constants C and α depending on $n, \Omega, c_0, \|\phi\|_{\tilde{C}^4(\overline{Q}_T)}$ and the uniform convexity of $\phi(\cdot, 0)$.

Remark 2.7. The interior higher regularity of u can also be obtained by using the same method in Proposition 5.3 of [4] once we get the global \tilde{C}^2 estimates and the \tilde{C}^α estimate for u_t .

We split the proof of the global $\tilde{C}^{2,\alpha}$ estimate into two parts. A short-time existence theorem gives the estimate for D^2u on $\Omega \times (0, 2\epsilon]$. As for the estimate on

the rest part of Q_T , considering the linear parabolic equation satisfied by tangential derivatives of u , we use the approach of Theorem 9.31 in [13] to deduce the Hölder regularity of the gradient, which is independent of the moduli of continuity of the principal coefficients.

Let $Y_0 = (y_0, t_0)$ be any point on $\partial\Omega \times [\epsilon, T]$ with some constant $\epsilon > 0$. Without loss of generality, assume that $y_0 = 0$, and on a small neighborhood of 0, $\partial\Omega$ is the image of $y_n = \rho(y')$. Perform the transformation of coordinates $x = \Psi(y)$, and $\Psi : B_{\rho_0}(0, t_0) \cap \overline{Q}_T \rightarrow \overline{\mathbb{R}}_n^+ \times (0, +\infty)$, with

$$x' = y', \quad x_n = y_n - \rho(y').$$

Let

$$v(x, t) = T_k(u - \phi)(x', x_n + \rho(x'), t)$$

with any $1 \leq k < n$, where

$$T_k =: \partial_k + \rho_k \partial_n \quad \text{for } k < n, \quad T_n = \partial_n.$$

Take $G = B_{R_0}^+ \times (t_0 - R_0^2, t_0]$, such that $G \subset \text{Im } \Psi$. Then by direct computations,

$$\begin{aligned} -\partial_t(T_k u) + u_t u^{ij}(T_k u)_{ij} &= u_t u^{ij} \rho_{kij} u_n + u_t u^{ij} \rho_{ki} u_{nj} + u_t u^{ij} \rho_{kj} u_{ni} \\ &=: \overline{f}. \end{aligned}$$

Thus under (x, t) coordinates,

$$-\partial_t v + u_t u^{ij} P_i P_j v = Lv - u_t u^{ij} \rho_{ij} v_n + LT_k \phi.$$

Here $L = -\partial_t + a^{ij} \partial_{ij}$, $P_k = \partial_k - \rho_k \partial_n$ for $1 \leq k < n$, $P_n = \partial_n$ and

$$a^{ij} = u_t u^{ij} + u_t u^{kl} \rho_k \rho_l \delta_n^i \delta_n^j - u_t u^{kj} \rho_k \delta_n^i - u_t u^{ik} \rho_k \delta_n^j,$$

where δ_n^i equals 1 when $n = i$, otherwise 0, and the summation over k, l is from 1 to $n - 1$. Let

$$f = u_t u^{ij} \rho_{ij} T_k u_n - LT_k \phi + \overline{f}.$$

Then, v satisfies the equation

$$\begin{cases} -v_t + a^{ij} v_{ij} = f & \text{in } G, \\ v = 0 & \text{on } F, \end{cases} \quad (2.17)$$

where $F = \partial G \cap \{x_n = 0\}$ is the flat boundary.

By Lemma 2.1 and Theorem 2.4, we have

$$\lambda I_n \leq (a^{ij}) \leq \Lambda I_n, \quad (2.18)$$

and

$$|f| \leq C_0, \quad |v| + |Dv| \leq C_1 \quad \text{in } \bar{G} \quad (2.19)$$

for some positive constants λ, Λ, C_0 and C_1 depending on $n, \Omega, c_0, \|\phi\|_{\tilde{C}^4(\bar{Q}_T)}$ and the uniform convexity of $\phi(\cdot, 0)$.

Before proving the boundary $\tilde{C}^{1,\alpha}$ regularity of v , we first give three lemmas including the weak Harnack inequality, the interior $\tilde{C}^{1,\alpha}$ regularity of v , and a technical lemma.

Lemma 2.8. *Let $v \in \tilde{C}^2(\bar{G})$ be a solution of equation (2.17) with conditions (2.18) and (2.19). Then there are constants $0 < \alpha < 1$ and $C > 0$ depending only on $n, \lambda, \Lambda, \text{diam } G, C_0$ and C_1 , such that for any $Q(X_1, R) \subset G, r \leq R$,*

$$\text{osc}_{Q(X_1, r)} Dv \leq C \left(\frac{r}{R} \right)^\alpha (\text{osc}_{Q(X_1, R)} Dv + R),$$

where $Q(X_1, R) = \{(x, t) : |x - x_1| < R, t_1 - R^2 < t \leq t_1\}$ with $X_1 = (x_1, t_1)$.

This lemma is derived from Theorem 12.1 in [24]. Note that this lemma reveals the interior $\tilde{C}^{1,\alpha}$ estimate of v . The next lemma is the weak Harnack inequality for linear parabolic equations, borrowed from Lemma 7.37 in [24].

Lemma 2.9. *Under the same conditions as in Lemma 2.8. Suppose also that v is non-negative. Then for any $Q(X_1, 4R) \subset G$, we have*

$$\left(R^{-n-2} \int_{\Theta(X_1, R)} v^p dX \right)^{1/p} \leq C \left(\inf_{Q(X_1, R)} v + R^{\frac{n}{n+1}} \|f\|_{L^{n+1}(G)} \right).$$

Here $\Theta(X_1, R) = Q((x_1, t_1 - 4R^2), R)$ and p, C are positive constants depending only on n, λ, Λ .

Additionally, the next lemma is Lemma 12.4 in [24].

Lemma 2.10. *Let $w \in C(\bar{Q})$ and $Dw \in C(Q)$ for some cylinder $Q = Q(X_0, R)$. Suppose that there are nonnegative constants \bar{C}_1, \bar{C}_2 and $\alpha < 1$ such that*

$$\text{osc}_{Q(X_1, r)} Dw \leq \bar{C}_1 \left(\frac{r}{\mu} \right)^\alpha (\text{osc}_{Q(X_1, \mu)} Dw + \bar{C}_2 \mu^\alpha),$$

whenever $Q(X_1, \mu) \subset Q$ and $r \leq \mu$. Then for any $\ell \in \mathbb{R}^n$ and $a \in \mathbb{R}$, we have

$$\sup_{Q(X_0, R/2)} |Dw - \ell| \leq C(\bar{C}_1, n, \alpha) \left(R^{-1} \sup_Q |w(X) - \ell \cdot x - a| + \bar{C}_2 R^\alpha \right).$$

We will use Lemma 2.9 to prove an oscillation estimate for $\frac{v}{x_n}$, which gives the Hölder estimate for v_n on F , the flat boundary. Combining this with the interior estimates, Lemma 2.8, and with the help of Lemma 2.10, we will derive a global $\tilde{C}^{1,\alpha}$ estimate for v in $\frac{1}{2}G := B_{R_0/2}^+ \times (t_0 - R_0^2/4, t_0]$. Recall the definition of v , then the global Hölder estimate of D^2u on $\Omega \times (\epsilon, T]$ follows from the equation.

Theorem 2.11. *Let $v \in \tilde{C}^2(\bar{G})$ be a solution of equation (2.17) with conditions (2.18) and (2.19). Then there are small positive constants r_0 and δ depending on n, λ, Λ, R_0 and C_0 such that for all $r \in (0, r_0)$ and $X_1 \in Q_{r,\delta} = \{(x', x_n, t) : |x'| < r, 0 < x_n < \delta r, t_0 - r^2 < t \leq t_0\}$, we have*

$$|Dv(X_1) - Dv(0, t_0)| \leq Cr^\alpha,$$

for some positive constants C and $\alpha < 1$ depending on $n, C_0, C_1, \lambda, \Lambda, r_0, \delta$ and R_0 .

Proof. Take $r < r_0$ small such that $r_0 + r_0/\delta \leq 1/2R_0$. $\delta < 1$ is a small constant to be chosen later. For any $X_1 = (x_1, t_1) \in Q_{r,\delta}$, let $\ell = Dv(X'_1)$, where $X'_1 = (x'_1, 0, t_1) = (x_{1,1}, \dots, x_{1,n-1}, 0, t_1) \in F$. Since $v = 0$ on F , $\ell = (0, \dots, 0, \ell_n)$. Take any point

$$X = (x, t) \in Q(X_1, \mu) = \{|x - x_1| < \mu, t_1 - \mu^2 < t \leq t_1\} \quad \text{with } \mu = \frac{1}{2}x_{1,n}.$$

If we take $\sigma = r + \mu/\delta$, then $X, X'_1 \in Q_{\sigma,\delta} \subset Q_{R_0/2,\delta}$. Consider the function $v(X) - \ell \cdot x$, then by Lemma 2.8 and Lemma 2.10,

$$\begin{aligned} |Dv(X_1) - Dv(X'_1)| &\leq \sup_{Q(X_1, \mu/2)} |Dv - \ell| \\ &\leq C \left(\mu^{-1} \sup_{Q(X_1, \mu)} |v(X) - \ell \cdot x| + \mu^\alpha \right). \end{aligned} \quad (2.20)$$

We claim that

$$\text{osc}_{Q_{\sigma,\delta}} \frac{v(X)}{x_n} \leq C\sigma^\alpha. \quad (2.21)$$

Then since $v|_F = 0$ and $X, X'_1 \in Q_{\sigma,\delta}$, we get

$$\left| \frac{v(X)}{x_n} - \ell_n \right| \leq \text{osc}_{Q_{\sigma,\delta}} \frac{v(X)}{x_n} \leq C\sigma^\alpha.$$

Thus (2.20) gives

$$\begin{aligned}
|Dv(X_1) - Dv(X'_1)| &\leq C \left(\mu^{-1} \sup_{Q(X_1, \mu)} Cx_n \sigma^\alpha + \mu^\alpha \right) \\
&\leq C(\mu^{-1} \sigma^\alpha (x_{1,n} + \mu) + \mu^\alpha) \\
&\leq C(\sigma^\alpha + \mu^\alpha) \\
&\leq Cr^\alpha.
\end{aligned}$$

Again by the claim, $|Dv(X'_1) - Dv(0, t_0)| \leq C\sigma^\alpha \leq Cr^\alpha$, then we have proved the theorem by the triangle inequality. For the rest part of this proof, we will show the claim (2.21).

First suppose that $v \geq 0$ and denote

$$F_{R, \delta} =: \{(x, t) : |x'| < R, x_n = \delta R, t_0 - R^2 < t \leq t_0\}.$$

We will prove the following inequality

$$\inf_{F_{R, \delta}} \frac{v}{x_n} \leq 10 \left(\inf_{Q_{R/2, \delta}} \frac{v}{x_n} + R \sup |f| \right) \quad \text{for } R < 1/4R_0. \quad (2.22)$$

Without loss of generality, assume that

$$R = 1, \inf_{F_{1, \delta}} \frac{v}{x_n} = 1. \quad (2.23)$$

Let

$$w(x, t) = \left[1 - |x'|^2 + (1 + \sup |f|) \frac{x_n - \delta}{\sqrt{\delta}} + t - t_0 \right] x_n,$$

then for $L = -\partial_t + a^{ij} \partial_{ij}$, we have

$$Lw \geq -\delta + \frac{2(1 + \sup |f|)}{\sqrt{\delta}} \lambda - c(n) \wedge (1 + \delta) \quad \text{in } Q_{1, \delta}.$$

Hence, $Lw \geq \sup |f| \geq Lv$ in $Q_{1, \delta}$, when δ is small enough. On $\{x_n = 0\} \cap \partial_p Q_{1, \delta}$, $w = 0 = v$. On $\{x_n = \delta\} \cap \partial_p Q_{1, \delta}$, $w \leq \delta(1 - |x'|^2) \leq \delta \leq v$ by (2.23). Additionally, on $\{|x'| = 1\} \cap \partial_p Q_{1, \delta}$ and $\{t = t_0 - 1\} \cap \partial_p Q_{1, \delta}$, $w < 0 \leq v$. Thus by maximal principle, we have $w \leq v$ in $Q_{1, \delta}$. Therefore, in $Q_{1/2, \delta}$,

$$\begin{aligned}
\frac{v}{x_n} &\geq 1 - |x'|^2 + (1 + \sup |f|) \frac{x_n \delta}{\sqrt{\delta}} + t - t_0 \\
&\geq 1 - \frac{1}{4} - \sqrt{\delta}(1 + \sup |f|) - \frac{1}{4} \\
&\geq \frac{1}{10} - \sup |f|.
\end{aligned}$$

when δ is sufficiently small. Now without the assumption (2.23), let

$$m_1 = \inf_{F_{R,\delta}} \frac{v}{x_n}, \quad \text{and} \quad \tilde{v}(x, t) = \frac{v(Rx, R^2(t - t_0) + t_0)}{m_1 R}.$$

Then $-\tilde{v}_t + a^{ij}\tilde{v}_{ij} = \tilde{f}$ in $Q_{1,\delta}$, with $\tilde{f} = \frac{R}{m_1}f$. Note that $\inf_{F_{1,\delta}} \frac{\tilde{v}}{x_n} = 1$. Then from the above arguments,

$$\frac{\tilde{v}(x, t)}{x_n} \geq 1/10 - \sup |\tilde{f}|.$$

Hence in the original coordinates, we have

$$\frac{v}{x_n} \geq \frac{m_1}{10} - R \sup |f| \quad \text{in } Q_{R/2,\delta},$$

which yields (2.22).

Denote

$$Q_{R,\delta}^* = \{|x'| < R, \frac{1}{2}\delta R < x_n < \delta R, t_0 - R^2 < t \leq t_0\},$$

$$Q'_{R,\delta} = \{|x'| < R, \frac{1}{2}\delta R < x_n < \delta R, t_0 - 3R^2 < t \leq t_0 - 2R^2\},$$

and

$$m = \inf_{Q_{2R,\delta}} \frac{v}{x_n}, \quad M = \sup_{Q_{2R,\delta}} \frac{v}{x_n}.$$

Then $Mx_n - v \geq 0$, $v - mx_n \geq 0$ in $Q_{2R,\delta}$. Applying Lemma 2.9, the weak Harnack inequality, to $Mx_n - v$ and $v - mx_n$ in $Q_{R,\delta}^*$, we have

$$\begin{aligned} \left(R^{-n-2} \int_{Q'_{R,\delta}} (Mx_n - v)^p \right)^{1/p} &\leq C \left(\inf_{Q_{R,\delta}^*} (Mx_n - v) + R^2 \sup |f| \right) \\ &\leq C \left(R \inf_{F_{R,\delta}} \left(M - \frac{v}{x_n} \right) + R^2 \sup |f| \right) \\ &\leq C \left(R \inf_{Q_{R/2,\delta}} \left(M - \frac{v}{x_n} \right) + R^2 \sup |f| \right) \\ &\leq CR \left(M - \sup_{Q_{R/2,\delta}} \frac{v}{x_n} + R \sup |f| \right). \end{aligned} \tag{2.24}$$

The third inequality above is deduced by (2.22). Similarly we have

$$\left(R^{-n-2} \int_{Q'_{R,\delta}} (v - mx_n)^p \right)^{1/p} \leq CR \left(\inf_{Q_{R/2,\delta}} \frac{v}{x_n} - m + R \sup |f| \right). \tag{2.25}$$

Add equations (2.24) and (2.25) up, and by Minkowski inequality,

$$(M - m) \left(R^{-n-2} \int_{Q'_{R,\delta}} x_n^p \right)^{1/p} \leq CR \left(M - m - \text{osc}_{Q_{R/2,\delta}} \frac{v}{x_n} + R \sup |f| \right).$$

Direct computations give that

$$\text{osc}_{Q_{R/2,\delta}} \frac{v}{x_n} \leq C \left(\text{osc}_{Q_{2R,\delta}} \frac{v}{x_n} + R \sup |f| \right).$$

Then it follows from the classical iterative arguments (see Lemma 8.23 in [13] for a reference) that

$$\text{osc}_{Q_{R,\delta}} \frac{v}{x_n} \leq C \left(\frac{R}{R_0} \right)^\beta \left(\text{osc}_{Q_{R_0,\delta}} \frac{v}{x_n} + R_0 \sup |f| \right) \quad (2.26)$$

for any $R < R_0$ and some $\beta < 1$ depending on $n, C_0, C_1, \lambda, \Lambda, \delta$. We still use α to denote $\min\{\alpha, \beta\}$. Since $|v| + |Dv|$ and $|f|$ are bounded in \bar{G} , the claim (2.21) is now proved. \square

Since $v|_F = 0$, the inequality (2.26) implies the Hölder continuity of Dv on F . With the above theorem, we infer that,

$$|Dv(X) - Dv(Y)| \leq C(|x - y| + |t - s|^{1/2})^\alpha$$

for any $X \in G$ and $Y \in F$. The \tilde{C}^α estimate for Dv on \bar{G} then follows from the triangle inequality and Lemma 2.8.

Reverting to u , we have

Corollary 2.12. *For any $\epsilon > 0$, there exist positive constants C and α such that*

$$\|D^2u\|_{\tilde{C}^\alpha(\bar{\Omega} \times (\epsilon, T])} \leq C,$$

where the constants C and α depend only on $n, \epsilon, \Omega, c_0, \|\phi\|_{\tilde{C}^4(\bar{Q}_T)}$ and the uniform convexity of $\phi(\cdot, 0)$.

Proof. Recall that $v = T_k(u - \phi)$ in $B_{\rho_0}(Y_0)$ with $T_k = \partial_k + \rho_k \partial_n$, $1 \leq k < n$, where we take ρ_0 to be small. If we define $\tilde{u}(x, t) = u(x', x_n + \rho(x'), t)$, then $\tilde{u}_{ij} = T_i T_j u$, for any $1 \leq i, j \leq n$. What we have proved above is equivalent to

$$\|\tilde{u}_{kl}\|_{\tilde{C}^\alpha(\bar{G})} \leq C \text{ for any } 1 \leq k < n \text{ and } 1 \leq l \leq n. \quad (2.27)$$

To prove the \tilde{C}^α estimate for \tilde{u}_{nn} , note that the equation (2.1) can be written as $\det(P_i P_j \tilde{u}) = \tilde{u}_t$, with $P_k = \partial_k - \rho_k \partial_n$ for $1 \leq k < n$ and $P_n = \partial_n$, which is equivalently

$$Q_1 \tilde{u}_{nn} + Q_2 = \tilde{u}_t,$$

where Q_1 and Q_2 are polynomials of \tilde{u}_{kl} with $k + l < 2n$ and ρ up to the second derivatives. By Theorem 2.4, the eigenvalues of the Hessian D^2u have positive lower and upper bounds; therefore Q_1 has a positive lower bound. This fact

together with Theorem 2.6 and (2.27) leads to the \tilde{C}^α estimate of \tilde{u}_{nn} . From this the \tilde{C}^α estimate of D^2u in $\Omega \times (\epsilon, T]$ follows by the arbitrariness of Y_0 and the interior higher regularity of u in Theorem 2.5.

□

3. EXISTENCE AND UNIQUENESS

Equation (2.1) is uniformly parabolic by virtue of Theorem 2.4. Hence with some compatibility conditions on ϕ , we have the following short-time existence theorem by the inverse function theorem. The proof is a modification of the argument in [12], Theorem 2.5.7.

Theorem 3.1. *Suppose that Ω is a uniformly convex domain with smooth boundary. ϕ is a smooth and uniformly convex function on $\bar{\Omega}$. h is a smooth function on $\partial\Omega \times [0, T]$ with $h_t > c_0 > 0$. Suppose also that the equation*

$$\begin{cases} u_t = \det D^2u, \\ u|_{t=0} = \phi, \quad u|_{\partial\Omega} = h, \end{cases} \quad (3.1)$$

satisfies the compatibility condition of order 1, i.e.,

$$h(0) = \phi|_{\partial\Omega} \quad \text{and} \quad h_t(0) = \det D^2\phi|_{\partial\Omega}.$$

Then for any $0 < \alpha \leq 1$, there is a small positive constant ϵ such that the initial boundary value problem (3.1) has a unique solution $u \in \tilde{C}^{2,\alpha}(\bar{Q}_{2\epsilon})$, which is uniformly parabolically convex. Here $Q_{2\epsilon} = \Omega \times (0, 2\epsilon]$ and $\epsilon < T$ depends on n, Ω, ϕ, h and α .

Proof. Consider the equation

$$\begin{cases} -w_t + \Delta w = \Delta\phi - \det D^2\phi, \\ w|_{t=0} = \phi, \quad w|_{\partial\Omega} = h. \end{cases} \quad (3.2)$$

According to theorem 5.2 in Chapter 4 of [23], problem (3.2) has a unique solution $w \in \tilde{C}^{2,\alpha}(\bar{Q}_{T_0})$ for some positive constant T_0 . Moreover, $w(x, t)$ is still uniformly convex in x for any $0 \leq t \leq T_0$ when T_0 is small enough. Indeed, by the $\tilde{C}^{2,\alpha}$ regularity of w , $\|w(\cdot, t) - \phi\|_{C^2(\bar{\Omega})} \leq Ct^{\alpha/2}$, then the uniform convexity of $w(\cdot, t)$ follows from that of ϕ for sufficiently small t .

Let

$$g = -w_t + \det D^2w \in \tilde{C}^\alpha(\bar{Q}_{T_0}).$$

Then by equation (3.2) and the compatibility condition, $g|_{t=0} = 0$. Let V be an open neighborhood of w in $\tilde{C}^{2,\alpha}(\overline{Q}_{T_0})$. Define the map $\Phi : V \rightarrow W$ by

$$\Phi(u) = (-u_t + \det D^2 u, u|_{t=0}, u|_{\partial\Omega \times [0, T_0]}), \quad (3.3)$$

where $W = \tilde{C}^\alpha(\overline{Q}_{T_0}) \times S$ and

$$S = \{(a, b) \in C^{2,\alpha}(\overline{\Omega}) \times \tilde{C}^{2,\alpha}(\partial\Omega \times [0, T_0]) : a|_{\partial\Omega} = b(0), \det D^2 a|_{\partial\Omega} = b_t(0)\}.$$

One can prove that V and W are Banach manifolds (see theorem 4.4 in [19]). Note that $\Phi(w) = (g, \varphi, h)$. The differential of Φ at w is a linear operator denoted by L .

$$L\eta = (-\eta_t + (\det D^2 w)w^{ij}\eta_{ij}, \eta(0), \eta|_{\partial\Omega \times [0, T_0]}), \quad \text{for any } \eta \in V.$$

The construction of w and S ensures the compatibility condition for L . Hence by Theorem 5.2 in Chapter 4 of [23], L is an isomorphism from the tangent space of V at w to that of W at (g, φ, h) . Then by the inverse function theorem, Φ is an C^1 diffeomorphism from $B_{\tilde{\rho}}(w)$, a small neighborhood of w in $\tilde{C}^{2,\alpha}(\overline{Q}_{T_0})$ to an open neighborhood U of (g, φ, h) in W . Take $\tilde{\rho}$ small enough, then the functions in $B_{\tilde{\rho}}(w)$ is still uniformly convex in x .

Let $\epsilon \in (0, T_0)$ be small, $\eta_\epsilon \in C^\infty([0, 1])$ such that $0 \leq \eta_\epsilon \leq 1$, $0 \leq \eta'_\epsilon \leq C\epsilon^{-1}$, and $\eta_\epsilon = 0$, if $0 \leq t \leq 2\epsilon$; $\eta_\epsilon = 1$, if $4\epsilon \leq t \leq 1$. Denote $g_\epsilon = g\eta_\epsilon$. Then $g_\epsilon \in \tilde{C}^\alpha(\overline{Q}_{T_0})$. One can prove that $g_\epsilon \rightarrow g$ in $\tilde{C}^\alpha(\overline{Q}_{T_0})$ when $\epsilon \rightarrow 0$. Thus by taking ϵ small enough, $(g_\epsilon, \varphi, h) \in U \subset W$, and there is a function $u \in B_{\tilde{\rho}}(w) \subset \tilde{C}^{2,\alpha}(\overline{Q}_{T_0})$ such that

$$\Phi(u) = (g_\epsilon, \varphi, h).$$

Since $g_\epsilon = 0$ for $0 \leq t \leq 2\epsilon$, then $u \in \tilde{C}^{2,\alpha}(\overline{Q}_{2\epsilon})$ is a uniformly parabolically convex solution to equation (3.1) in $\Omega \times (0, 2\epsilon]$. \square

Remark 3.2. When the initial-boundary condition of equation (3.1) satisfies the compatibility condition of order $m = [\frac{l+\alpha}{2}] + 1$, there exists a unique solution $u \in \tilde{C}^{l+2,\alpha}(\overline{Q}_\epsilon)$. The proof is almost the same as above, and here we omit it. One can refer to Chapter 4, Section 5 in [23] for the general definition of m -th order compatibility condition. To be more explicit, the compatibility condition of m -th order for our equation means that

$$u^{(k)}(x)|_{x \in \partial\Omega} = \frac{\partial^k h}{\partial t^k} \Big|_{t=0'}$$

for $k = 0, \dots, m$, where $u^{(k)}$ is recursively defined by

$$u^{(0)} = \varphi, \quad u^{(k+1)}(x) = \frac{\partial^k}{\partial t^k} \Big|_{t=0} \det D^2 u(t, x).$$

Here the right term of the second equality can be recursively expressed by derivatives of φ .

Lemma 3.3. *Suppose that $u, v \in \tilde{C}^2(Q_T) \cap C(\overline{Q}_T)$ are uniformly convex in x . If*

$$\begin{cases} -u_t + \det D^2 u \geq -v_t + \det D^2 v & \text{in } Q_T, \\ u \leq v & \text{on } \partial_p Q_T. \end{cases}$$

Then $u \leq v$ in Q_T .

Proof. Note that

$$\begin{aligned} 0 &\geq (u - v)_t + \det D^2 v - \det D^2 u \\ &= (u - v)_t + \int_0^1 \frac{d}{ds} \det D^2(sv + (1 - s)u) ds \\ &= w_t - \bar{a}^{ij} w_{ij}, \end{aligned}$$

where $w = u - v$, and the matrix (\bar{a}^{ij}) is

$$\int_0^1 \det D^2(sv + (1 - s)u)(sD^2u + (1 - s)D^2v)^{-1} ds.$$

Then $(\bar{a}^{ij}) > 0$. Since $w \leq 0$ on $\partial_p Q_T$, we have $w \leq 0$ in Q_T by comparison theorem of linear parabolic equations. \square

The uniqueness of solutions to the equation (2.1) then follows immediately from the above lemma.

Corollary 3.4. *Let Ω be a uniformly convex bounded domain with smooth boundary and ϕ be a continuous function in \overline{Q}_T . Then equation (2.1) admits at most one \tilde{C}^2 smooth and parabolically convex solution.*

We are in a position to prove our main theorems.

Proof of Theorem 1.3: By uniqueness, the a priori solution u in Corollary 2.12 coincides with the solution derived from Theorem 3.1 on $[0, 2\epsilon]$ if $\varphi = \phi|_{t=0}$ and $h = \phi|_{\partial\Omega}$. Thus, for some small positive constant ϵ , $\|u\|_{\tilde{C}^{2,\alpha}(\overline{Q}_{2\epsilon})}$ is bounded. Combined with Corollary 2.12, this gives the global $\tilde{C}^{2,\alpha}$ estimate to u on \overline{Q}_T . \square

Proof of Theorem 1.2: The existence of $\tilde{C}^{2,\alpha}$ solutions of equation (2.1) can be derived step by step from the short-time existence under the uniform global $\tilde{C}^{2,\alpha}$ estimate. It is also unique by Corollary 3.4.

Additionally, the higher order regularity of u can also be obtained by classic regularity theory of linear parabolic equations iteratively if $\partial\Omega$ and ϕ are smooth enough, and equation (2.1) admits such solution uniquely if the equation satisfies higher order compatibility conditions in Remark 3.2. Specifically, let $k \geq 2$, $0 < \beta \leq 1$. Assume that $\partial\Omega$ is of class $C^{k+2,\beta}$, $\phi \in \tilde{C}^{k+2,\beta}(\overline{Q}_T)$, and equation (2.1) satisfies the compatibility condition of order $[\frac{k+\beta}{2}] + 1$. Then the equation admits a unique parabolically convex solution $u \in \tilde{C}^{k+2,\beta}(\overline{Q}_T)$. \square

4. GAUSS CURVATURE FLOW

In this section, we consider the Gauss curvature flow with boundary

$$\begin{cases} u_t = \frac{\det D^2 u}{(1 + |Du|^2)^{\frac{n+1}{2}}} & \text{in } Q_T, \\ u = \phi & \text{on } \partial_p Q_T. \end{cases} \quad (4.1)$$

where ϕ and Q_T are defined as before. We will obtain the *a priori* estimate for equation (4.1) using the same method as before.

Theorem 4.1. *Let $u \in \tilde{C}^4(\overline{Q}_T)$ be a parabolically convex solution of equation (4.1). Suppose that Ω is a uniformly convex domain in \mathbb{R}^n , $\partial\Omega \in C^4$, $\phi \in \tilde{C}^4(\overline{Q}_T)$, $\phi_t \geq c_0 > 0$ on $\partial\Omega \times [0, T]$ and $\phi(\cdot, 0)$ is uniformly convex. Then we have*

$$\|u\|_{\tilde{C}^{2,\alpha}(\overline{Q}_T)} \leq C$$

for some positive constants C and α depending only on $n, T, \Omega, c_0, \|\phi\|_{\tilde{C}^4(\overline{Q}_T)}$ and the uniform convexity of $\phi(\cdot, 0)$.

Proof. The proof of this theorem is similar to that of Theorem 1.3, except that the linearized equation of (4.1) has lower order terms concerning Du .

(a) C^0 and C^1 estimates. Note that u reaches its maximum and minimum value on $\partial_p Q_T$, then $\|u\|_{C^0(\overline{Q}_T)} \leq \|\phi\|_{C^0(\overline{Q}_T)} \leq C$.

For convenience we denote $\beta = \frac{n+1}{2}$. Differentiating the equation (4.1), we have

$$\mathcal{L}'u_k = 0, \quad \text{for } k = 1, \dots, n, \quad \text{and} \quad \mathcal{L}'u_t = 0$$

with

$$\begin{aligned}\mathcal{L}' &= -\partial_t + \frac{\det D^2 u}{(1 + |Du|^2)^\beta} u^{ij} \partial_{ij} - 2\beta \frac{\det D^2 u}{(1 + |Du|^2)^{\beta+1}} u_i \partial_i \\ &= -\partial_t + u_t u^{ij} \partial_{ij} - \frac{2\beta}{1 + |Du|^2} u_t u_i \partial_i.\end{aligned}\tag{4.2}$$

Then \mathcal{L}' is a linear parabolic operator. It follows that $|u_t|$ is bounded by $\sup_{\partial_p Q_T} |u_t|$ and

$$C^{-1} \leq u_t \leq C$$

for some positive constant C depending on $n, c_0, \Omega, \|\phi\|_{\tilde{C}^2}$ and the uniform convexity of $\phi(\cdot, 0)$.

To prove the bound of $|Du|$, we apply the similar auxiliary function in [28]. Fix any $t_0 \in [0, T]$. Let $\tilde{\phi}$ be the solution of the equation

$$\begin{cases} \det D^2 \tilde{\phi} = 1 & \text{in } \Omega, \\ \tilde{\phi} = \phi(\cdot, t_0) & \text{on } \partial\Omega. \end{cases}$$

The existence and C^2 bound of solutions to this equation are guaranteed by [2]. We consider the barrier function

$$\bar{w} = \tilde{\phi} - \frac{1}{q} \log(1 + kd_{\partial\Omega})\tag{4.3}$$

defined in $\Omega_\delta = \{x \in \Omega : d_{\partial\Omega}(x) < \delta\}$ with small positive constant δ such that $\bar{w} \in C^4(\Omega_\delta)$.

The positive constants δ, q , and k shall be determined later. Direct computations give that

$$\begin{aligned}\frac{\det D^2 \bar{w}}{(1 + |D\bar{w}|^2)^\beta} &\geq \frac{1 + Ck^{n+1}q^{-n}(1 + kd_{\partial\Omega})^{-n-1}}{(1 + 2|D\tilde{\phi}|^2 + 2k^2q^{-2}(1 + kd_{\partial\Omega})^{-2})^\beta} \\ &\geq Cq,\end{aligned}$$

when $k^2q^{-2}(1 + k\delta)^{-2} \geq \max_{\bar{\Omega}}(1 + 2|D\tilde{\phi}|^2)$ and δ is small enough.

We first fix the sufficiently big constant q such that $Cq \geq \max u_t$. Thus

$$\frac{\det D^2 \bar{w}}{(1 + |D\bar{w}|^2)^\beta} \geq \frac{\det D^2 u}{(1 + |Du|^2)^\beta} \quad \text{in } \Omega_\delta \times \{t_0\}.$$

Then we take δ to be small enough and $k\delta \geq 1$ to make

$$k^2q^{-2}(1 + k\delta)^{-2} \geq \frac{1}{4}q^{-2}\delta^{-2} \geq \max_{\bar{\Omega}}(1 + 2|D\tilde{\phi}|^2).$$

Finally, we take k sufficiently large to satisfy $\log(k\delta) \geq q(\max_{\overline{\Omega}} |\tilde{\phi}| + \max_{\overline{Q_T}} |u|)$, then $\bar{w} \leq u$ on $\partial\Omega_\delta \times \{t_0\}$. Therefore $u \geq \bar{w}$ in $\Omega_\delta \times \{t_0\}$ by the comparison principle. Then for the exterior normal vector ν at any point x_0 on $\partial\Omega$, we have $u_\nu \leq w_\nu \leq C$. By convexity of u in x and arbitrariness of t_0 , this proves the bound for $|Du|$ on $\overline{Q_T}$.

(b) C^2 estimates on the boundary. We can prove this by the same method as before but with different barrier functions. As for the second tangential derivatives, we can also regard $u_t(1 + |Du|^2)^{\frac{n+1}{2}}$ which has positive lower and upper bounds as a data function. Then it follows that

$$C^{-1}I_{n-1} \leq D_{x'x'}u \leq CI_{n-1} \quad \text{on} \quad \partial_p Q_T. \quad (4.4)$$

To prove the uniform bound for the mixed second derivatives, we use the comparison principle for \mathcal{L}' . Denote \mathcal{T} as in (2.8). Then

$$|\mathcal{L}'\mathcal{T}(u - \phi)| \leq \underline{a} \left(1 + \sum_{i=1}^n u^{ii}\right) \quad (4.5)$$

for some positive constant \underline{a} depending on $\|\phi\|_{\tilde{C}^4(\overline{Q_T})}$, the positive lower and upper bounds for u_t and the bound for $|Du|$.

Recall that $0 \in \partial\Omega$ and the boundary $\partial\Omega$ can be represented as in (2.7) near 0. On $\Omega \times \{0\}$, $\mathcal{T}(u - \phi) = 0$ and

$$|\mathcal{T}(u - \phi)| \leq \bar{a}|x'|^2, \quad \text{on} \quad (B_\varepsilon \cap \partial\Omega) \times [0, T]. \quad (4.6)$$

In $(B_\varepsilon \cap \Omega) \times [0, T]$ with $B_\varepsilon = \{|x| < \varepsilon\}$, we shall consider the following barrier function

$$\zeta(x, t) = \frac{1}{2} \sum_{\alpha, \beta < n} (B_{\alpha\beta} - \mu\delta_{\alpha\beta})x_\alpha x_\beta + \frac{1}{2}Mx_n^2 - Nx_n,$$

where μ is a fixed small constant such that $B_{\alpha\beta} - \mu\delta_{\alpha\beta} > 0$, and M, N will be determined later.

First we take ε small, $M\varepsilon \leq 2$ and N larger than $1 + \lambda_{max}/c$, where λ_{max} is the maximum eigenvalue of the $(n-1) \times (n-1)$ matrix $(B_{\alpha\beta} - \mu\delta_{\alpha\beta})$, and c is the constant such that $x_n \geq c|x'|^2$ on $\partial\Omega$. Then we have $\zeta < 0$ in $(B_\varepsilon \cap \Omega) \times \{0\}$.

Thus

$$A\zeta \pm \mathcal{T}(u - \phi) \leq 0, \quad \text{in} \quad (B_\varepsilon \cap \Omega) \times \{0\} \quad \text{for all positive constant } A. \quad (4.7)$$

Note that we can choose a large $M \geq 2$ and a small $c_1 > 0$ such that

$$\begin{aligned} \mathcal{L}'\zeta &\geq c_1 \left(\sum_{i=1}^n u^{ii} + Mu^{nn} \right) - c_2 \\ &\geq \frac{1}{2}c_1 \sum_{i=1}^n u^{ii} - c_2 + c_3 M^{1/n}. \end{aligned} \quad (4.8)$$

Here we use the bounds for u_t and the fact that

$$\frac{1}{n} \left(\sum_{i=1}^n u^{ii} + Mu^{nn} \right) \geq \frac{1}{n} \left(\sum_{i=1}^{n-1} \lambda_i^{-1} + M\lambda_n^{-1} \right) \geq \left(M\lambda_1^{-1} \cdots \lambda_n^{-1} \right)^{1/n},$$

where $\lambda_1 \leq \cdots \leq \lambda_n$ are the eigenvalues of $\{u_{ij}\}$. c_3 is a positive constant with $c_3 \leq Cc_1$, where C depends on n and the maximum of u_t . The constant c_2 depends on the maximum of u_t , $|Du|$ and the diameter of Ω . Hence, for $A \geq \max\{1, 2\underline{a}/c_1\}$ and $M \geq ((c_1 + c_2)/c_3)^n$, we have

$$\mathcal{L}'(A\zeta \pm \mathcal{T}(u - \phi)) \geq 0, \quad \text{in } (B_\varepsilon \cap \Omega) \times (0, T]. \quad (4.9)$$

From now on M is fixed.

Next, we estimate ζ on $\partial(B_\varepsilon \cap \Omega) \times [0, T]$. For ε small enough, we have

$$\zeta \leq -\frac{1}{2}\mu \sum_{\beta < n} x_\beta^2 + O(|x'|^3) \leq -\frac{1}{4}\mu|x'|^2, \quad \text{on } (\overline{B_\varepsilon} \cap \partial\Omega) \times [0, T]. \quad (4.10)$$

Then, on $(\partial B_\varepsilon \cap \Omega) \times [0, T]$ and if $\frac{1}{4}\mu|x'|^2 > Mx_n^2$, we have

$$\begin{aligned} \zeta &\leq \frac{1}{2}(B_{\alpha\beta} - \mu\delta_{\alpha\beta})x_\alpha x_\beta + \frac{1}{2}Mx_n^2 - x_n \\ &\leq -\frac{1}{2}\mu|x'|^2 + C|x'|^3 + Mx_n^2 \\ &\leq -c_4\varepsilon^2, \quad \text{for } \varepsilon \text{ small enough.} \end{aligned} \quad (4.11)$$

Additionally, on $(\partial B_\varepsilon \cap \Omega) \times [0, T]$ and if $\frac{1}{4}\mu|x'|^2 \leq Mx_n^2$, we have $x_n \geq c_5\varepsilon$ and

$$\begin{aligned} \zeta &\leq C|x'|^2 + Mx_n^2 - x_n \\ &\leq (M + C)\varepsilon^2 - c_5\varepsilon \\ &\leq -\frac{1}{2}c_5\varepsilon, \quad \text{for } \varepsilon \text{ small enough.} \end{aligned} \quad (4.12)$$

Since $|\mathcal{T}(u - \phi)|$ has an upper bound in \overline{Q}_T , which is supposed to be c_6 , then by (4.6), (4.10), (4.11) and (4.12), we get

$$A\zeta \pm \mathcal{T}(u - \phi) \leq 0, \quad \text{on } \partial(B_\varepsilon \cap \Omega) \times [0, T], \quad (4.13)$$

for $A = \max\{1, 2\underline{a}/c_1, 4\bar{a}/\mu, c_6/(c_4\varepsilon^2), 2c_6/(c_5\varepsilon)\}$. Thus by maximum principle and equations (4.9), (4.7) and (4.13), we get

$$A\zeta \pm \mathcal{T}(u - \phi) \leq 0, \quad \text{in } (B_\varepsilon \cap \Omega) \times [0, T]. \quad (4.14)$$

Since $\zeta(0, t_0) = \mathcal{T}(u - \phi)(0, t_0) = 0$ for each $t_0 \in [0, T]$, we have

$$|\partial_n \mathcal{T}(u - \phi)(0, t_0)| = |u_{\alpha n}(0, t_0) - \phi_{\alpha n}(0, t_0)| \leq AN,$$

which yields that

$$|u_{\alpha n}(0, t_0)| \leq AN + \|\phi\|_{C^2(\overline{Q}_T)} \leq C. \quad (4.15)$$

By the equation (4.1) and estimate (4.4), we then get $u_{nn} \leq C$ on $\partial\Omega \times [0, T]$ for some universal constant C . Combining with the positive lower and upper bounds of u_t , we have

$$1/C \leq \lambda(D^2u) \leq C \quad \text{on } \partial_p Q_T \quad (4.16)$$

for some positive constant C depending on $n, \Omega, c_0, \|\phi\|_{\tilde{C}^4(\overline{Q}_T)}$ and the uniform convexity of $\phi(\cdot, 0)$.

(c) *global C^2 estimates.* The Legendre transform U of u satisfies

$$-U_t \det D^2U = (1 + |y|^2)^{-\beta} \quad \text{in } Q_T^*$$

with $C^{-1} \leq -U_t \leq C$ and $\|D^2U\|_{L^\infty(\partial_p Q_T^*)} \leq C$. Conducting the same calculation as in Lemma 2.3, we have

$$\tilde{L}U_{kk} \geq -2\beta,$$

where $\tilde{L} = \frac{1}{U_t} \partial_t + U^{ij} \partial_{ij}$. Thus $\tilde{L}(U_{kk} - Ct) \geq 0$ for some universal constant C . Hence by maximum principle and (4.16),

$$\sup_{Q_T^*} U_{kk} \leq C \left(\sup_{\partial_p Q_T^*} U_{kk} + T \right) \leq C,$$

which yields $\lambda(D^2u) \geq C$ in Q_T for some positive constant C depending on $n, \Omega, c_0, T, \|\phi\|_{\tilde{C}^4(\overline{Q}_T)}$ and uniform convexity of $\phi(\cdot, 0)$. Therefore we have proved

$$1/C \leq \lambda(D^2u) \leq C \quad \text{on } \overline{Q}_T.$$

(d) *$\tilde{C}^{2,\alpha}$ estimates.* The interior higher regularity of u follows from that of U in Q_T^* by [30]. Under the same condition of Theorem 2.6, it holds $\|u_t\|_{\tilde{C}^\alpha(\overline{Q}_T)} \leq C$ by Krylov-Safonov's estimate.

Note that if we let $v = T_k(u - \phi)$, $1 \leq k \leq n - 1$, as in subsection 2.3, we have

$$-v_t + a^{ij}v_{ij} + b^i v_i = f \quad \text{in } G = B_{R_0}^+ \times (t_0 - R_0^2, t_0].$$

The term $b^i v_i$ comes from the differential of $(1 + |Du|^2)^{-\beta}$ and the related coordinate transformation. Here a^{ij} is positive definite, and $|b^i|$, $|v_i|$ and $|f|$ are bounded due to the boundedness of u_t , Du and D^2u . This is the case we have dealt with in equation (2.17). Thus the $\tilde{C}^{2,\alpha}$ estimate of u on $\bar{\Omega} \times [\epsilon, T]$ follows from the same argument in the proof of Corollary 2.12.

The short-time existence theorem ensures the $\tilde{C}^{2,\alpha}$ estimate of u in a small time interval. The proof in this case is similar to the proof of Theorem 3.1. The Fréchet differential of the map Φ , defined similarly as (3.3), is a linear parabolic operator with extra lower order terms. It is still an isomorphism under some suitable compatibility conditions. Thus the short-time existence of the solution follows from the inverse map theorem.

In conclusion, we have proved the global $\tilde{C}^{2,\alpha}$ estimate for u , the solution of equation (4.1). \square

We also present an existence theorem for equation (4.1).

Theorem 4.2. *Assume Ω is a smooth and uniformly convex domain in \mathbb{R}^n , ϕ is a smooth function in \bar{Q}_T with $\phi_t \geq c_0 > 0$ on $\partial\Omega \times [0, T]$, and $\phi(\cdot, 0)$ is uniformly convex. If equation (4.1) satisfies the compatibility condition of any order, then there exists a unique smooth parabolically solution u to the initial boundary value problem (4.1).*

Remark 4.3. Using the above method, one can also prove the global $\tilde{C}^{2,\alpha}$ estimate for the solutions to the first initial-boundary value problem of γ -Gauss curvature flow as well as the existence of solutions when $0 < \gamma \leq 1$.

REFERENCES

- [1] B. Andrews, *Gauss curvature flow: the fate of the rolling stones*, *Invent. Math.* **138** (1999), 151–161.
- [2] L. A. Caffarelli, L. Nirenberg, J. Spruck, *The Dirichlet problem for nonlinear second-order elliptic equations, I. Monge-Ampère equation*, *Comm. Pure Appl. Math.* **37** (1984), no. 3, 369–402.
- [3] L. A. Caffarelli, *A localization property of viscosity solutions to the Monge-Ampère equation and their strict convexity*, *Ann. of Math. (2)* **131** (1990), no. 1, 129–134.
- [4] K. Choi, P. Daskalopoulos, L. Kim, K.-A. Lee, *The evolution of complete non-compact graphs by powers of Gauss curvature*, *J. Reine Angew. Math.* **757** (2019), 131–158.
- [5] B. Chow, D. H. Tsai, *Nonhomogeneous Gauss curvature flows*, *Indiana Univ. Math. J.* **47** (1998), no. 3, 965–994.

- [6] P. Daskalopoulos, O. Savin, $C^{1,\alpha}$ regularity of solutions to parabolic Monge-Ampère equations, *Amer. J. Math.* **134** (2012), 1051–1087.
- [7] P. Daskalopoulos, R. Hamilton, *The free boundary in the Gauss curvature flow with flat sides*, *J. Reine Angew. Math.* **510** (1999), 187–227.
- [8] P. Daskalopoulos, K. A. Lee, *Worn stones with flat sides all time regularity of the interface*, *Invent. Math.* **156** (2004), no. 3, 445–493.
- [9] L. Evans, *Classical solutions of fully nonlinear, convex, second order elliptic equations*, *Comm. Pure Appl. Math.* **25** (1982), 333–362.
- [10] A. Figalli, *The Monge-Ampère equation and its applications*, *Zurich Lect. Adv. Math.*, European Mathematical Society (EMS), Zürich, 2017.
- [11] W. J. Firey, *Shapes of worn stones*, *Mathematika* **21** (1974), 1–11.
- [12] C. Gerhardt, *Curvature problems*, *Series in geometry and topology*, International Press, 2006.
- [13] D. Gilbarg, N. S. Trudinger, *Elliptic partial differential equations of second order*, vol. 224, Springer-Verlag, Berlin-New York, 1977.
- [14] P. Guan, L. Ni, *Entropy and a convergence theorem for Gauss curvature flow in high dimension*, *J. Eur. Math. Soc.* **19** (2017), no. 12, 3735–3761.
- [15] C. E. Gutiérrez, Q. Huang, *A generalization of a theorem by Calabi to the parabolic Monge-Ampère equation*, *Indiana Univ. Math. J.* (1998), 1459–1480.
- [16] C. E. Gutiérrez, Q. Huang, $W^{2,p}$ estimates for the parabolic Monge-Ampère equation, *Arch. Ration. Mech. Anal.* **159** (2001), 137–177.
- [17] Q. Huang, G. Lu, *On a priori $C^{1,\alpha}$ and $W^{2,p}$ estimates for a parabolic Monge-Ampère equation in the Gauss curvature flows*, *Amer. J. Math.* **128** (2006), 453–480.
- [18] N. M. Ivochkina, O.A. Ladyzhenskaia, *Parabolic equations generated by symmetric functions of the eigenvalues of the Hessian or by the principal curvatures of a surface, I. Parabolic Monge-Ampère equations*, *St. Petersburg Math. J.* **6** (1995), 575–594.
- [19] J. Kitagawa, *A parabolic flow toward solutions of the optimal transportation problem on domains with boundary*, *J. Reine Angew. Math.* **672** (2012), 127–160.
- [20] N. V. Krylov, *Sequences of convex functions, and estimates of the maximum of the solution of a parabolic equation*, *Sibirskii Matematicheskii Zhurnal* **17** (1976), 290–303.
- [21] N. V. Krylov, *Boundedly nonhomogeneous elliptic and parabolic equations in a domain* (Russian). *Izv. Akad. Nauk SSSR Ser. Mat.* **47** (1983), 75–108. English translation in *Math. USSR. Izv.* **20** (1983), 459–492.
- [22] N. V. Krylov, M. V. Safonov, *A property of the solutions of parabolic equations with measurable coefficients*, (Russian) *Izv. Akad. Nauk SSSR Ser. Mat.* **44** (1980), no. 1, 161–175.
- [23] O.A. Ladyzhenskaia, V.A. Solonnikov, and N.N. Ural'tseva, *Linear and quasi-linear equations of parabolic type*, Nauka, Moscow, 1968.
- [24] G. Lieberman. *Second order parabolic differential equations*, World Scientific, Singapore, 1996.
- [25] V. Oliker, *Evolution of nonparametric surfaces with speed depending on curvature, I. The Gauss curvature case*, *Indiana Univ. Math. J.* **40** (1991), no.1, 237–258.
- [26] O. C. Schnürer, K. Smoczyk, *Neumann and second boundary value problems for hessian and Gauss curvature flows*, *Ann. Inst. Henri Poincaré (C) Anal. Non Linéaire.* **20** (2003), no. 6, 1043–1073.

- [27] L. Tang, *Regularity results on the parabolic Monge-Ampère equation with VMO type data*, J. Differ. Equ. **255** (2013), 1646–1656.
- [28] N. S. Trudinger, J. I. Urbas, *The Dirichlet problem for the equation of prescribed Gauss curvature*, Bull. Aust. Math. Soc. **28** (1983), 217–231.
- [29] N. S. Trudinger, X. J. Wang, *The Monge-Ampère equation and its geometric applications*, Handbook of Geometric Analysis, Adv. Lect. Math., 7, pp. 467–524. Int. Press, Somerville, MA, 2008.
- [30] K. Tso, *Deforming a hypersurface by its Gauss-Kronecker curvature*, Comm. Pure Appl. Math. **38** (1985), no. 6, 867–882.
- [31] K. Tso, *On a real Monge-Ampère functional*, Invent. Math. **101** (1990), no. 2, 425–448.
- [32] R. Wang, G. Wang, *On existence, uniqueness and regularity of viscosity solutions for the first initial boundary value problems to parabolic Monge-Ampère equation*, Northeast. Math. J. **8** (1992), 417–446.
- [33] R. Wang, G. Wang, *The geometric measure theoretical characterization of viscosity solutions to parabolic Monge-Ampère type equation*. J. Partial Diff. Equ. **6** (1993), 237–254.
- [34] J. Xiong, J. Bao, *On Jörgens, Calabi, and Pogorelov type theorem and isolated singularities of parabolic Monge-Ampère equations*, J. Differ. Equ. **250** (2011), 367–385.
- [35] W. Zhang, J. Bao, B. Wang, *An extension of Jörgens–Calabi–Pogorelov theorem to parabolic Monge-Ampère equation*, Calc. Var. Partial Differ. Equ. **57** (2018), 1–36.

(Yang Zhou) SCHOOL OF MATHEMATICS AND APPLIED STATISTICS, UNIVERSITY OF WOLLONGONG, WOLLONGONG, NSW 2522, AUSTRALIA

Email address: yang.zhou18798@outlook.com

(Ruixuan Zhu) DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING 100084, CHINA

Email address: zhurx19@mails.tsinghua.edu.cn