

Norm-induced Cuts: Optimization with Lipschitzian Black-box Functions

ADRIAN GÖß, ALEXANDER MARTIN, SEBASTIAN POKUTTA, KARTIKEY SHARMA

ABSTRACT. In this paper, we consider a finite dimensional optimization problem minimizing a continuous objective on a compact domain subject to a multi-dimensional constraint function. For the latter, we only assume the availability of a Lipschitz property. In recent literature methods based on non-convex outer approximation are proposed for tackling one dimensional equality constraints on bounded polyhedral domains, which are Lipschitz with respect to the maximum norm. To the best of our knowledge, however, there exists no non-convex outer approximation method for a general problem class. We introduce a meta-level solution framework to solve such problems and tackle the underlying theoretical foundations. Considering the feasible domain without the constraint function as manageable, our method relaxes the multidimensional constraint and iteratively refines the feasible region by means of norm-induced cuts, relying on an oracle for the resulting sub-problems. We show the method's correctness and investigate the problem complexity. In order to account for discussions about functionality, limits, and extensions, we present computational examples including illustrations.

1. INTRODUCTION

In today's applications, we are faced with a steady increase in global connectivity and a massive availability of data; see [1]. Thus, when modeling these conditions in terms of a mathematical optimization problem, its formulation grows in complexity and data-based approaches can be leveraged, respectively. Hereby, the resulting constraint functions may lack differentiability, which motivates a retreat to their treatment as black-box functions with a Lipschitz property. Hence, we address the following general form of an optimization problem.

$$\begin{aligned} \min_x \quad & f(x) \\ \text{s.t.} \quad & r(x) \leq \mathbf{0}, \\ & x \in \Omega. \end{aligned} \tag{P}$$

We assume Ω to be a non-empty, compact subset of the finite dimensional vector space X over a field F , i.e., $\Omega \subset X \cong F^n$, $n \in \mathbb{N}$. The space X is associated with a norm $\|\cdot\|_X$. That is, in particular, $(X, \|\cdot\|_X)$ is a normed space. Set Ω serves as a domain for the function $r : \Omega \rightarrow \mathbb{R}^m$ mapping it into the m -dimensional real space for some $m \in \mathbb{N}$. We mention that $\mathbf{0} \in \mathbb{R}^m$ denotes the vector of m zeros and the inequality in (P) is to be interpreted component-wise. In addition, function r is assumed to be Lipschitz continuous on Ω with a known Lipschitz constant $L > 0$, i.e., for any $x, y \in \Omega$, it holds that

$$\|r(x) - r(y)\|_{\mathbb{R}^m} \leq L\|x - y\|_X,$$

Date: March 19, 2024.

2010 Mathematics Subject Classification. 90C26, 90C30, 90C56, 90C60 .

Key words and phrases. Global optimization, Lipschitz optimization, Black-box function evaluation, Derivative-free optimization.

with $\|\cdot\|_{\mathbb{R}^m}$ denoting a monotonous norm in \mathbb{R}^m . Lastly, the objective function $f : \Omega \rightarrow \mathbb{R}$ is assumed to be continuous on Ω .

Note that the value of the constant L depends on the specific choice of norms. We emphasize that only the evaluation of r at any point $x \in \Omega$ needs to be possible. Furthermore, we investigate the problem (P) for an arbitrary field F , as the general formulation of our method allows for it. As mentioned below, the field can even be substituted (partially) by a ring such as the integers \mathbb{Z} , extending the applicability to mixed-integer problems.

In our problem formulation, it is assumed that we can evaluate the constraint function r without having an explicit representation. Such problems are tackled in the field of *black-box optimization* or, more commonly, *derivative-free optimization*. Here, the applicability of any particular approach also depends on the structure of Ω . For example, if Ω is defined by linear constraints only, a directional direct-search approach can be leveraged which applies a Lagrangian reformulation; see [12]. For the case when non-smooth constraints define Ω , filtering techniques as proposed in [2] may be more suitable. Further, in terms of direct search, a well-known method called *mesh adaptive direct search (MADS)* (see [3]) is capable of handling general constraints in Ω , too. Alternatively, Berghen [4] proposes a trust-region interpolation-based approach to tackle such problems. For a general overview on constrained derivative-free optimization, we refer to [7, Chapter 13].

However, assuming no further information about the involved functions can be quite limiting. For example, even when minimizing a continuous function over a simple box-constraint domain, in a worst-case situation, a finite bound on the number of iterations to find an optimal solution cannot be expected, see, e.g., [20, Chapter 6]. Hence, we assume the availability of more information, in particular, that the function r is Lipschitz continuous and that we know its constant. Thus, we touch the field of *Lipschitz optimization*. Traditionally, problems with univariate, Lipschitzian objectives are tackled by saw-tooth cover methods, see [15] for the original source. Here, the constraint functions are iteratively bounded by means of the Lipschitz constant. More general problems, similar to (P), are solved by branch-and-bound schemes or outer approximation, see [11, Chapter XI] for an overview.

Some works in recent literature have leveraged the Lipschitz property for the constraints by relaxing them and iteratively re-incorporating this information. For example, the method proposed in [17] approximates the original constraint function by a parallelepipedal shape and refines this in every iteration. In contrast, the approach of [8, 18] entirely omits the constraint function at first and improves a non-convex outer approximation of the remaining domain step by step. This is achieved by bounding the distance to found incumbents in terms of the infinity norm.

The algorithmic scheme presented in [18] tackles a simplified version of problem (P) limited to polyhedral constraints on a mixed-integer domain in the sub-problem along with one-dimensional implicit Lipschitz constraints. However, this particular scheme cannot tackle the general problem (P) that we propose, as the resulting sub-problem may not be polyhedral or additional variables and constraints may be too costly or undesired.

In terms of applications, our approach in this paper is particularly well suited for non-linear optimization problems which use surrogate models as constraints along with simpler, e.g., convex, domains. Such a domain can represent an additional source of uncertainty which increases the complexity of the problem. For instance, linear constraints can be associated with ellipsoidal uncertainty sets which

are equivalent to second-order cone programming (SOCP) problems. This is particularly relevant for engineering design applications where you need to manage implementation uncertainty along with complex non-linear response behaviors [6, 14].

To the best of our knowledge, there exists no extended approach to handle the posed general problem classes, so we aim for one and investigate its properties in the abstract setting. We are also not restricted to a particular norm, as long as the respective Lipschitz property is available and the norm is monotonous. Furthermore, the resulting sub-problems are not explicitly constructed, but assumed to be solved by an oracle due to the generality of the problem class.

Our contribution includes (i) the introduction of a meta-level method tackling a problem with Lipschitzian black-box constraints in the generality of (P). (ii) We also investigate the approach theoretically in terms of correctness and problem complexity. (iii) Additionally, we visualize the functionality of the method, its limits considering local solutions of the sub-problems, and the benefit of having information about Lipschitz constants for components of the constraint function.

The remainder of the paper is structured as follows. In Section 2, we give basic definitions in order to formulate the introduced method clearly and to keep the proof of correctness in Section 3 concise. Section 4 is dedicated to an analysis regarding a termination bound in the case of an infeasible problem w.r.t. the properties of r and Ω , as well as to a discussion about the problem complexity. In Section 5, we illustrate the procedure, present an example which shows its weakness in terms of local solutions, and show the value of more information in terms of Lipschitz constants of components. We close the paper with Section 6 by summarizing the content and pointing to further research directions.

2. NORM-INDUCED CUT METHOD

In this section, we formulate a method which is able to tackle Problem (P). From now on, we will omit the subscripts for each norm, where it is unambiguous. We start by giving precise definitions which clarify the occurring mathematical objects. Afterwards, the solution method is introduced, iteratively solving a relaxed problem and adding cuts in order to enforce feasibility. The cuts are determined by a point in Ω , the norm, and the function r with its Lipschitz constant L . We define them formally below.

Definition 2.1. For $z \in \mathbb{R}^m$, let $z_+ = (\max(z_1, 0), \dots, \max(z_m, 0))^T$ define the component-wise maximum of the elements of z and 0. Now, let $y \in \Omega$ such that $r(y) \not\leq \mathbf{0}$ and let $L > 0$ define the Lipschitz constant of function r . Then, for $x \in \Omega$, we call the inequality

$$\frac{\|r(y)_+\|}{L} \leq \|x - y\|, \quad (\text{NC})$$

a norm-induced cut in y .

Whenever referring to the *radius of a norm-induced cut*, we consider the constant term $\|r(y)_+\|/L$ of (NC) for a given $y \in \Omega$. In the case of $X \cong \mathbb{R}^n$ we note that a norm-induced cut is non-convex in variable x independent of the choice of norm.

Note that the norm-induced cuts force the variable x to have a minimum distance from the current incumbent. This distance is the radius of the norm-induced cut and includes the violation of $r(x) \leq \mathbf{0}$ at the incumbent and the Lipschitz constant L . Informally, having a greater violation or a smaller L increases the radius of the norm-induced cut and, thus, leading to a larger “excluded area” in the next iteration.

Now, let the feasible set of (P) be denoted by

$$Q := \{x \in \Omega \mid r(x) \leq \mathbf{0}\} \subseteq \Omega.$$

In the following, we define a family of sets which involve norm-induced cuts and which are relaxations to Q .

Definition 2.2. Let $k \in \mathbb{N}$ be a positive integer and $(x^i)_{i=0, \dots, k-1} \subseteq \Omega$ be a sequence such that $r(x^i) \not\leq \mathbf{0}$ for all $i = 0, \dots, k-1$. Then, we define

$$Q_k := \left\{ x \in \Omega \mid \forall i \in \{0, \dots, k-1\} : \frac{\|r(x^i)_+\|}{L} \leq \|x - x^i\| \right\},$$

the k th relaxed set of Q . It is obtained from Ω by relaxing the inequality constraint $r(x) \leq \mathbf{0}$ and adding norm-induced cuts for each element x^i of the sequence for a total of k cuts. For $k = 0$, we naturally define $Q_0 := \Omega$.

In Lemma 3.3, we show that these Q_k are indeed relaxations of Q . The resulting family of relaxed problems is denoted as

$$\min_{x \in Q_k} f(x), \tag{P}_k$$

for $k \in \mathbb{N}_0$. We interpret k as the iteration index of our method. The following remark shows a connection between two consecutive relaxed problems (P_k) and (P_{k+1}) , in particular, between their feasible sets Q_k and Q_{k+1} , respectively.

Remark 2.3. Let $k \in \mathbb{N}_0$ and let $(x^i)_{i=0, \dots, k} \subseteq \Omega$ be the sequence to define the k th and the $(k+1)$ th relaxed set Q_k and Q_{k+1} , respectively. Then, Q_{k+1} is given by

$$Q_{k+1} = Q_k \cap \left\{ x \in \Omega \mid \frac{\|r(x^k)_+\|}{L} \leq \|x - x^k\| \right\}.$$

In other words, the $(k+1)$ th relaxed set results from the k th relaxed set by adding the *norm-induced cut* in x^k , i.e.,

$$\frac{\|r(x^k)_+\|}{L} \leq \|x - x^k\|.$$

Now, we are able to state our method for solving problem (P). It is called the *Norm-induced Cutting Method (NIC)*, following its main functionality of iteratively adding norm-induced cuts.

In particular, it starts by relaxing the inequality constraint in (P), $r(x) \leq \mathbf{0}$, and solves the resulting relaxed problem (P_k) for $k = 0$. It is assumed that we have a suitable method to solve (P_k) for all $k \in \mathbb{N}_0$. If the relaxed problem is infeasible, so is the original one and a message is returned accordingly. Otherwise, the method checks if the solution of the relaxed problem x^0 is also feasible for the original problem by evaluating function r at x^0 and checking for the inequality. If it is satisfied, the current incumbent x^0 is returned as a solution to the original problem. Otherwise, a norm-induced cut in incumbent x^0 is added. This defines the set Q_1 . The resulting problem (P_1) is then solved in the next iteration.

In conclusion, the method adds norm-induced cuts as long as the incumbents are infeasible for the original problem (P). Below we give a formal description of the NIC method in pseudo code. We point out that any specific stopping criterion is omitted, as we examine the method's behavior for an infinite number of iterations in Section 3.

As mentioned in the introduction, a reader familiar with the field of Lipschitz optimization might consider our approach as a generalization of the work [18]. Although originated independently, this is a reasonable consideration which we want to discuss in the following.

Norm-induced Cutting Method (NIC)

Input: Set Ω , function r , Lipschitz constant L

Output: A message of (\mathbf{P}) 's infeasibility, a solution to (\mathbf{P}) , or a sequence $(x^i)_{i \in \mathbb{N}_0}$

```

1: set  $k \leftarrow 0$ 
2: solve  $(\mathbf{P}_k)$ 
3: if  $(\mathbf{P}_k)$  is infeasible then
4:   return ‘ $(\mathbf{P})$  is infeasible’
5: else
6:   denote  $x^k$  as the solution of  $(\mathbf{P}_k)$ 
7: end if
8: if  $r(x^k) \leq 0$  then
9:   return solution  $x^k$  to  $(\mathbf{P})$ 
10: end if
11: set  $k \leftarrow k + 1$ 
12: goto (2)

```

Remark 2.4. In [18] the authors consider a problem originated from mixed-integer non-linear optimization. In particular, the problem consists of a linear objective function with linear constraints, as well as implicitly given one dimensional non-linear equality constraints. Hereby, the optimizing variable x has finite bounds and is part of a mixed-integer space, where the non-linear functions depend on its continuous part only.

In their algorithm, they iteratively add a cut for each function which does not satisfy its equality condition (approximately). The cut is analogous to the norm-induced cuts considered here, but uses the infinity norm in the continuous part of the x -space and a right-hand side of the absolute violation, as well as the global Lipschitz constant. Due to the infinity norm, they re-model this cut by introduction of additional binary variables and linear constraints which upholds the character of a mixed-integer linear problem for the resulting sub-problems. As a consequence, the dimension of the sub-problems grows by a factor of iteration times number of non-linear constraints. For this algorithm, the authors prove correctness and give a worst-case termination bound for obtaining an approximate solution.

We investigate a more general problem class, by considering a compact domain in a vector space over some field (and/or \mathbb{Z} , compare Remark 3.9) instead of the linear constraints in a mixed-integer domain. Regarding the non-linear functions, we go one step further and, for the individual components, assume a lack of information regarding the Lipschitz constant.

Our method is also based on adding cuts, but with respect to a general monotonous norm. Hence, the resulting sub-problems inherit an equally general structure involving the set Ω . This, however, prevents an explicit re-modeling the cuts in order to keep the sub-problems in their original problem class, but only allows for a discussion in this regard.

In terms of theoretical contribution, our proofs of correctness, at some points, follow a different approach and open a new perspective on the method. In addition, our termination bound is analogous to the results in [18] yet more general to account for the more general setting. We also comment on the availability of a kind of local Lipschitz constant in Remark 3.7, answering a question posed in the conclusion of their paper, and showcase the advantage of leveraging component-wise Lipschitz constants on an example (see Example 5.3).

Now, in order to discuss properties of the **NIC** method in the following sections, we introduce the notation of a ball and the terms ε -packing/-covering. We start with the former.

Definition 2.5. Let $\varepsilon > 0$, $\bar{x} \in X$, and $\|\cdot\|$ a norm on X . Then, we define the ε -ball around \bar{x} as

$$B_\varepsilon(\bar{x}) := \{x \in X \mid \|x - \bar{x}\| < \varepsilon\}.$$

For the case $\bar{x} = \mathbf{0}$, we write the short form

$$B_\varepsilon := B_\varepsilon(\mathbf{0}) = \{x \in X \mid \|x\| < \varepsilon\}.$$

For the fundamental mathematics regarding ε -packing/-covering, we refer to [19]. We use the notation from the lecture notes [10] and adapt the definition to fit our question.

Definition 2.6. Let $k \in \mathbb{N}_0$, $\varepsilon > 0$, and $\|\cdot\|$ a norm on X . Further, consider $A \subseteq X$ and $P := \{x^0, \dots, x^k\} \subseteq A$ a set of points in A .

- a) If $A \subseteq \bigcup_{i=0}^k B_\varepsilon(x^i)$, we call P an ε -covering of A . Furthermore, we define the minimal cardinality of such a P as

$$N(A, \|\cdot\|, \varepsilon) := \min \{k \mid \exists \varepsilon\text{-covering of } A \text{ with size } k\},$$

the covering number of A w.r.t. ε .

- b) If $\{B_{\varepsilon/2}(x^i)\}_{i=0, \dots, k}$ are pairwise disjoint, i.e., $\|x^i - x^j\| \geq \varepsilon$ for all $i, j = 0, \dots, k$, $i \neq j$, we call P an ε -packing of A . We also define the maximal cardinality of such a P , i.e.,

$$M(A, \|\cdot\|, \varepsilon) := \max \{k \mid \exists \varepsilon\text{-packing of } A \text{ with size } k\},$$

the packing number of A w.r.t. ε .

3. CONVERGENCE AND OPTIMALITY RESULTS

This section is dedicated to show the correctness of **NIC**. Depending on the feasibility of the original problem **(P)**, we show feasibility and even optimality of a returned solution or of the accumulation points of the sequence of incumbents produced by the method. In the end, we give a statement regarding the availability of local solutions in step 2. We start by remarking on the boundedness of **(P)** in case of feasibility.

Remark 3.1. As r is Lipschitz continuous, it is also continuous. By re-writing the feasible set of **(P)** as

$$Q = \Omega \cap r^{-1}((-\infty, 0]^m),$$

where $r^{-1}(\cdot)$ notes the pre-image of r , we can use the closure of $(-\infty, 0]^m$ to derive the closure of $r^{-1}((-\infty, 0]^m)$. Together with the compactness of Ω , this implies that Q is compact. Therefore, Problem **(P)** minimizes a continuous function over a compact set and, thus, if $Q \neq \emptyset$, the problem has a finite solution value.

With this in mind, we state two lemmata. The first one ensures that **NIC** does not find any incumbent twice, the second one implies that **(P_k)** is a relaxation of **(P)** for each $k \in \mathbb{N}$, as mentioned in Definition 2.2. In other terms, we construct an outer approximation of the feasible region.

Lemma 3.2. Consider some $k \in \mathbb{N}_0$ and a solution x^k to **(P_k)** such that $r(x^k) \not\leq \mathbf{0}$. Then, $x^k \notin Q_{k+1}$, i.e., the incumbent x^k is infeasible for problem **(P_{k+1})**.

Proof. As x^k is a solution to (\mathbf{P}_k) , we have $x^k \in Q_k$. As $L > 0$ and there exists a strictly positive component of $r(x^k)$, it follows

$$\|x^k - x^k\| = 0 < \frac{\|r(x^k)_+\|}{L}.$$

Then, Remark 2.3 directly implies that $x^k \notin Q_{k+1}$ and the claim follows. \square

Lemma 3.3. *For any $k \in \mathbb{N}_0$, it holds that $Q \subseteq Q_k$.*

Proof. In the following, we use the short notation $[m] := \{1, \dots, m\}$. For a vector $z \in \mathbb{R}^m$ and a subset of indices $I \subseteq [m]$, let $z_I \in \mathbb{R}^m$ such that

$$(z_I)_p = \begin{cases} z_p, & p \in I, \\ 0, & \text{else,} \end{cases} \quad \text{for } p \in [m].$$

Now, if $k = 0$, $Q_k = \Omega$ by construction and the claim follows from Definition 2.2.

Otherwise, let $k \in \mathbb{N}$ arbitrary but fixed and let $(x^i)_{i=0, \dots, k-1}$ be a sequence to determine Q_k . Then, it holds true that

$$x^i \in \Omega \quad \wedge \quad r(x^i) \not\leq \mathbf{0},$$

for $i = 0, \dots, k-1$, where the second condition is equivalent to $\|r(x^i)_+\| > 0$. We define the set of non-negative indices of $r(x^i)$ as $P^i := \{p \in [m] \mid r(x^i)_p \geq 0\}$. Further, consider some $\bar{x} \in Q$, thus, $\bar{x} \in \Omega$ and $r(\bar{x}) \leq \mathbf{0}$. We can conclude that

$$0 < \frac{\|r(x^i)_+\|}{L} \leq \frac{1}{L} \|(r(\bar{x}))_{P^i} - r(x^i)_+\| \leq \frac{1}{L} \|r(\bar{x}) - r(x^i)\| \leq \|\bar{x} - x^i\|,$$

for all $i = 1, \dots, k-1$, where the second and third inequalities follow from the assumed monotonicity of the norm. In particular, the components of the argument in the norm are non-decreasing in absolute terms from left to right, therefore, the norm is too. Hence, for all $i = 0, \dots, k-1$, \bar{x} satisfies the norm-induced cut in x^i . With Definition 2.2 it follows $\bar{x} \in Q_k$ and the claim is proven. \square

Lemma 3.3 shows that infeasibility of Problem (\mathbf{P}_k) implies the infeasibility of Problem (\mathbf{P}) , in particular, $Q_k = \emptyset \Rightarrow Q = \emptyset$. Therefore, step 4 associated with the corresponding if-clause is reasonable and NIC gives a correct answer if it stops in step 4. Furthermore, it also follows from Lemma 3.3 that (\mathbf{P}_k) is a relaxation of (\mathbf{P}) for $k \in \mathbb{N}_0$. Therefore, the solution value of (\mathbf{P}_k) , which is obtained during NIC, gives a lower bound to the value of (\mathbf{P}) .

In the following, we prove that NIC either converges to a feasible solution in finite time or produces a sequence with feasible accumulation points for (\mathbf{P}) .

Theorem 3.4. *Let (\mathbf{P}) be feasible, i.e., $Q \neq \emptyset$. Then, either*

- a) *NIC stops with a feasible solution x^* for (\mathbf{P}) in step 9, or*
- b) *NIC creates a sequence of incumbents $(x^i)_{i \in \mathbb{N}_0}$, which has at least one convergent subsequence, each of which has a limit feasible for (\mathbf{P}) . That is, $(x^i)_{i \in \mathbb{N}_0}$ has at least one accumulation point and all accumulation points of $(x^i)_{i \in \mathbb{N}_0}$ are feasible for (\mathbf{P}) .*

Proof. With the assumption and Lemma 3.3, it is clear that

$$\emptyset \neq Q \subseteq \bigcap_k Q_k. \quad (1)$$

Here, k indicates the iteration index and is specified in the case distinction below. From (1) we have that $Q_k \neq \emptyset$ and, thus, NIC does not stop in step 4. Hence, it either stops in step 9 or runs for infinite time. We start with the former case.

a) Let $k \in \mathbb{N}_0$ be the index of the stopping iteration. According to Definition 2.2, $Q_k \subseteq \Omega$ and, thus, the incumbent satisfies $x^k \in \Omega$. Since stopping in step 9 requires $r(x^k) \leq \mathbf{0}$, it follows that **NIC** returns $x^* := x^k \in Q$, i.e., a feasible point for (P).

b) **NIC** runs for infinite time, i.e., stopping in step 4 is not possible and we assume that we never satisfy the criterion in step 9. Again, as $Q_k \subseteq \Omega$ for all $k \in \mathbb{N}_0$, we receive a sequence $(x^i)_{i \in \mathbb{N}_0} \subseteq \Omega$. With Ω being compact, $(x^i)_{i \in \mathbb{N}_0}$ has at least one convergent subsequence $(x^{i_j})_{j \in \mathbb{N}_0}$ with

$$\lim_{j \rightarrow \infty} x^{i_j} = x^* \in \Omega.$$

We prove the feasibility of x^* for (P) by contradiction and assume $x^* \notin Q$, i.e., $r(x^*) \not\leq \mathbf{0}$. This directly implies $\|r(x^*)_+\| > 0$. As **NIC** is assumed to not terminate, we have

$$\forall j \in \mathbb{N}_0 : r(x^{i_j}) \not\leq \mathbf{0},$$

which analogously implies $\|r(x^{i_j})_+\| > 0$. Furthermore, having compactness of Ω and the (Lipschitz) continuity of r on Ω , we can deduce the existence of an $\varepsilon > 0$, such that

$$\mathbf{0} < 2\varepsilon < \|r(x^*)_+\|.$$

For a vector $x \in \Omega$, let $P^+(x) := \{p \in [m] \mid r(x)_p > 0\}$ denote the positive indices of r at x . Due to the convergence of $(x^{i_j})_{j \in \mathbb{N}_0}$ to x^* and the continuity of r , there exists a $j(\varepsilon) \in \mathbb{N}_0$ such that

$$P^+(x^*) \subseteq P^+(x^{i_j}) \quad \wedge \quad \|x^* - x^{i_j}\| < \frac{\varepsilon}{L}, \quad \text{for all } j \geq j(\varepsilon). \quad (2)$$

That is, for all $j \geq j(\varepsilon)$, the element x^{i_j} of the subsequence violates the constraint $r(x) \leq \mathbf{0}$ at least on the same indices as x^* and it is inside an ε/L -ball around x^* . Using the triangle inequality, the monotonicity of the norm, and the results from above, we conclude

$$\begin{aligned} 2\varepsilon &< \|r(x^*)_+\| = \|r(x^*)_+ - r(x^{i_j})_+ + r(x^{i_j})_+\| \\ &\leq \|r(x^*)_+ - r(x^{i_j})_+\| + \|r(x^{i_j})_+\| \\ &\leq \|r(x^*) - r(x^{i_j})\| + \|r(x^{i_j})_+\| \\ &\leq L\|x^* - x^{i_j}\| + \|r(x^{i_j})_+\| \leq \varepsilon + \|r(x^{i_j})_+\|, \end{aligned} \quad (3)$$

for all $j \geq j(\varepsilon)$. In conclusion, we see that

$$\|r(x^{i_j})_+\| > \varepsilon, \quad \text{for all } j \geq j(\varepsilon).$$

That is, for all $j \geq j(\varepsilon)$, a norm-induced cut in $x^{i_j} \in Q_{i_j}$ of the form

$$\|x - x^{i_j}\| \geq \frac{\|r(x^{i_j})_+\|}{L} > \frac{\varepsilon}{L} \quad (4)$$

is added, resulting in Q_{i_j+1} . In order to show termination, Lemma 3.2 and (4) imply that it suffices to prove that only a finite number t of x^{i_j} can be contained in the original feasible set $Q_0 = \Omega$, having a pairwise distance (w.r.t. the norm) of at least ε/L . After adding all the respective cuts, **NIC** would end up with an empty Q_k for some k . This is a contradiction and, thus, we derive termination.

In other terms, we aim to find a maximal ε/L -packing of Ω . This motivates us to set $M(\delta) = M(\Omega, \|\cdot\|, \delta)$ and $N(\delta) = N(\Omega, \|\cdot\|, \delta)$ the packing and covering number of Ω w.r.t. some $\delta > 0$, respectively, compare Definition 2.6. For normed spaces, it is well known that $M(2\delta) \leq N(\delta)$ (see, e.g., [19]) and, thus, $M(\varepsilon/L) \leq N(\varepsilon/(2L))$. That is, our claim of a finite termination is proven if there exists a finite $\varepsilon/(2L)$ -covering.

Now, as X is a finite dimensional vector space, it is isomorphic to \mathbb{R}^n with an isomorphism $\phi : X \rightarrow \mathbb{R}^n$. As Ω is compact, $\phi(\Omega)$ is also compact and analogously

$\phi(B_{\varepsilon/(2L)})$ is open. By the theorem of Heine-Borel (see, e.g., [16, Theorem 2.41]) there exists a finite cover of $\phi(\Omega)$ with translations of $\phi(B_{\varepsilon/(2L)})$. Therefore, the equivalent statement holds true for Ω and $B_{\varepsilon/(2L)}$, and it follows $N(\varepsilon/(2L)) < \infty$. This shows $M(\varepsilon/L) < \infty$. Hence, **NIC** terminates in a finite amount of iterations, a contradiction. Therefore, $x^* \in \Omega$ and claim b) is proven. \square

In other terms, this shows that **NIC** solves the feasibility problem of **(P)** in case it really is feasible. The following remark takes the availability of possibly smaller Lipschitz constants into account and comments on their effects on Theorem 3.4.

Remark 3.5. *Let's assume we have access to the local Lipschitz constants L_x of r for each $x \in \Omega$. That is,*

$$\forall x \in \Omega \exists L_x > 0 : \|r(x) - r(y)\| \leq L_x \|x - y\|, \quad \text{for all } y \in \Omega. \quad (5)$$

First, note that the statements of Lemma 3.2 and Lemma 3.3 still hold when using the local Lipschitz constants L_{x^k} and L_{x^i} in the proofs, respectively. Second, as the global Lipschitz constant L is an upper bound to every local one, i.e., $L \geq L_x$ for all $x \in \Omega$, the inequality in (2) can be re-written to $\|x^ - x^{ij}\| < \varepsilon/L_{x^{ij}}$, leading to the same result as in inequality (3) and, thus, a possible greater right-hand side in inequality (4). Therefore, the availability of local Lipschitz constants can lead to stronger cuts and may result in faster convergence.*

*Further, during the execution of **NIC** in iteration $k \in \mathbb{N}_0$, we have to consider the Lipschitz continuity of r only on Q_k . We note that, while this has no influence on the behavior of the algorithm per se, it might be possible to leverage a bounded domain such as Q_k to compute the Lipschitz constant efficiently, as well as obtain a lower constant.*

*It is also possible to combine locality with the current feasible region. In particular, if we are in iteration $k \in \mathbb{N}_0$ and have solved **(P_k)** to receive x^k , we could compute L_{x^k} for $y \in Q_k$ instead of doing so for all $y \in \Omega$ as denoted in (5). Then, we could proceed and add the cut with respect to L_{x^k} .*

Besides the question of feasibility of the output, its quality/optimality also needs to be investigated. Note that Theorem 3.4 does not require the solution in step 2 to be optimal per se. Therefore, **NIC** produces a feasible output even if **(P_k)** is only solved for a feasible solution. However, when investigating the optimality of the returned solution or the accumulation points, we assume the global optimality of x^k for **(P_k)** in step 2 and apply a case distinction as in Theorem 3.4.

Theorem 3.6. *Let **(P)** be feasible, i.e., $Q \neq \emptyset$, and let step 2 produce a globally optimal point x^k for **(P_k)**, $k \in \mathbb{N}_0$. Then, either*

- a) ***NIC** stops with a globally optimal solution x^* of **(P)** in step 9, or*
- b) ***NIC** creates a sequence of incumbents $(x^i)_{i \in \mathbb{N}_0}$, which has at least one convergent subsequence, each of which has a limit that is globally optimal for **(P)**. That is, $(x^i)_{i \in \mathbb{N}_0}$ has at least one accumulation point and all accumulation points of $(x^i)_{i \in \mathbb{N}_0}$ are globally optimal for **(P)**.*

Proof. With **(P)** being feasible, Theorem 3.4 shows that **NIC** either stops in step 9 or iterates infinitely often. We start with the former case.

a) Let $k \in \mathbb{N}_0$ be the index of the stopping iteration, in which x^k is returned. According to Theorem 3.4, x^k is feasible for **(P)**, i.e., $x^k \in Q$, and, thus, it is

$$\min_{x \in Q} f(x) \leq f(x^k).$$

Further, by assumption x^k is optimal for (P_k) and Lemma 3.3 states $Q \subseteq Q_k$ which implies

$$f(x^k) = \min_{x \in Q_k} f(x) \leq \min_{x \in Q} f(x).$$

Together, we conclude

$$\min_{x \in Q} f(x) = f(x^k),$$

which proves claim a).

b) Assuming **NIC** iterates infinitely, we denote $(x^i)_{i \in \mathbb{N}_0}$ as the resulting sequence of incumbents. Theorem 3.4 states that it has at least one accumulation point and that all its accumulation points are feasible for (P) . Hence, let $(x^{i_j})_{j \in \mathbb{N}_0}$ denote a convergent subsequence of $(x^i)_{i \in \mathbb{N}_0}$ with limit $x^* \in Q$. The feasibility of x^* for (P) immediately gives

$$\min_{x \in Q} f(x) \leq f(x^*).$$

Applying Lemma 3.3 on the index of the subsequence, leads to $Q \subseteq Q_{i_j}$ and, thus,

$$f(x^{i_j}) = \min_{x \in Q_{i_j}} f(x) \leq \min_{x \in Q} f(x),$$

for $j \in \mathbb{N}_0$. Leveraging the continuity of f when taking the limit, we can derive

$$\lim_{j \rightarrow \infty} f(x^{i_j}) = f(\lim_{j \rightarrow \infty} x^{i_j}) = f(x^*) \leq \min_{x \in Q} f(x).$$

Therefore, $f(x^*) = \min_{x \in Q} f(x)$ which concludes the proof. \square

So far, we have proven that **NIC** is suited to tackle feasible problems which have a structure like (P) . We can further extend the method when either Lipschitz information is available for components of the constraint function r or the function in question is defined on a lower dimensional subspace of the feasible set. The combination of these two extension is also treatable. A formalization of this is given below.

Remark 3.7. If $r = (r_p)_{p=1, \dots, R}$ for $R \in \mathbb{N}$ and Lipschitz constants $L_p > 0$ are available for the functions $r_p : \Omega \rightarrow \mathbb{R}^{m_p}$, $m_p \in \mathbb{N}$, we can re-write (P) as

$$\begin{aligned} \min_x f(x) \\ \text{s.t. } r_p(x) \leq \mathbf{0}, \quad \text{for } p = 1, \dots, R, \\ x \in \Omega. \end{aligned} \tag{P_R}$$

Then, we can perform **NIC** analogously by checking for each $r_p(x^k) \leq \mathbf{0}$ before step 9. Moreover, if the condition is not satisfied for all $p = 1, \dots, R$, we can add one norm-induced cut for the p which has $r_p(x^k) \not\leq \mathbf{0}$ and maximizes $\|r_p(x^k)_+\|/L_p$. In particular, such a cut is given by

$$\|x - x^k\| \geq \|r_p(x^k)_+\|/L_p.$$

One can see that the proofs above hold equally for such a procedure. We visualize this case on a specific instance in Example 5.3.

Apart from that, if r in (P) maps from a lower dimensional subset of Ω into the \mathbb{R}^m , we can re-write $x = (x', x'') \in \Omega = \Omega' \times \Omega''$ with $\dim(\Omega'), \dim(\Omega'') \geq 1$, $\dim(\Omega') + \dim(\Omega'') = n$, and $r : \Omega' \rightarrow \mathbb{R}^m$. Denoting the first $\dim(\Omega')$ elements of a vector with the prime, we can investigate the case of an incumbent x^k with $r((x^k)') \not\leq \mathbf{0}$. As a vector $((x^k)', x'')$, for all $x'' \in \Omega''$, can never be a valid solution for (P) , the norm-induced cut in x^k can be added in $(x^k)'$ instead. That is, we can add

$$\|x' - (x^k)'\| \geq \|r((x^k)')_+\|/L,$$

with an appropriate norm for Ω' . This can lead to faster convergence, as cuts are added in a possible smaller subspace Ω' .

In conclusion, one can even combine several inequalities depending on different subvectors of x . The only necessary condition is the availability of respective Lipschitz constants and norms.

We showed that **NIC** works correctly when we have a global solution method for (\mathbf{P}_k) . But the question arises whether the algorithm still fulfills the desired behavior when only a local solution method is available. In the following, we verify that if only local solutions can be found in step 2, a point returned in step 9 is indeed a locally optimal point.

Theorem 3.8. *Let (\mathbf{P}) be feasible, i.e., $Q \neq \emptyset$, and let step 2 produce a locally optimal point x^k for (\mathbf{P}_k) , $k \in \mathbb{N}_0$. Then, if **NIC** stops in step 9, it returns a locally optimal solution x^* .*

Proof. Assume that **NIC** stops in step 9 in iteration $k \in \mathbb{N}_0$. As x^k is a locally optimal solution for (\mathbf{P}_k) , there exists an $\varepsilon > 0$ such that

$$f(x^k) = \min_{x \in B_\varepsilon(x^k) \cap Q_k} f(x).$$

By Theorem 3.4 $x^k \in Q$ and, thus, $x^k \in Q \cap B_\varepsilon(x^k)$. We infer

$$\min_{x \in Q \cap B_\varepsilon(x^k)} f(x) \leq f(x^k).$$

Further, the result from Lemma 3.3 implies $Q \cap B_\varepsilon(x^k) \subseteq Q_k \cap B_\varepsilon(x^k)$. We can derive

$$f(x^k) = \min_{x \in Q_k \cap B_\varepsilon(x^k)} f(x) \leq \min_{x \in Q \cap B_\varepsilon(x^k)} f(x),$$

and, thus, conclude that

$$\min_{x \in Q \cap B_\varepsilon(x^k)} f(x) = f(x^k).$$

Therefore, x^k is a local optimum for (\mathbf{P}) and the proof is complete. \square

If **NIC** does not terminate in step 9 with returning a point, we can in general not guarantee the local optimality of the accumulation points. On the contrary, the sequence might even converge to the worst possible point, as we show in Example 5.2.

We conclude the section by a remark regarding alternatives for the field F .

Remark 3.9. *Note that replacing F with the ring \mathbb{Z} does not change the validity of the computations above. That is, one can consider Ω as a subset of a mixed-integer domain as well.*

4. TERMINATION AND PROBLEM COMPLEXITY

This section discusses a termination criterion for **NIC** based on the properties of r in case of infeasibility in (\mathbf{P}) and the problem complexity. We restrict ourselves to $F = \mathbb{R}$.

4.1. Termination for Infeasible Problems. Let (\mathbf{P}) be infeasible, i.e., $Q = \emptyset$. Hence, $r(x) \not\leq \mathbf{0}$ and $\|r(x)_+\| > 0$ for all $x \in \Omega$. Due to the compactness of Ω and the (Lipschitz) continuity of r , there exists $\delta > 0$ such that $\delta < \min_{x \in \Omega} \|r(x)_+\|$. In analogy to the proof of Theorem 3.4, we can deduce inequality (4) regarding the norm-induced cuts for a sequence of incumbents $(x^i)_{i \in \mathbb{N}_0}$, i.e.,

$$\|x - x^i\| \geq \frac{\|r(x^i)_+\|}{L} > \frac{\delta}{L}, \quad \text{for all } i \in \mathbb{N}_0.$$

Ω	$\ \cdot\ $	T
$\Omega_1 := \prod_{j=1}^n [a_j, b_j]$	$\ \cdot\ _\infty$	$\prod_{j=1}^n [(L/\delta)(b_j - a_j) + 1]$
$\Omega_2 := \prod_{j=1}^n [a_j, b_j] \cap \mathbb{Z}^n$	$\ \cdot\ _\infty$	$ \Omega_2 $
$\Omega_3 := \{x \mid \ x\ _2 < D\}$	$\ \cdot\ _2$	$(2LD/\delta + 1)^n$

TABLE 1. Examples for T in (6) for $\delta/L < 1$ depending on the structure of Ω and the choice of norm.

Now, any new solution x^i must satisfy the upper inequality with respect to all previous incumbents x^0, \dots, x^{i-1} ; compare Remark 2.3. In particular, for all $l \in \mathbb{N}$, it holds

$$\|x^l - x^i\| > \frac{\delta}{L}, \quad \text{for all } i = 0, \dots, l-1.$$

Thus, the number of iterations until termination can be bounded by the maximal number of points in Ω which have a pairwise distance, with respect to the norm, of more than δ/L . This refers to a maximal δ/L -packing of Ω and was formalized in Definition 2.6.

In particular, our aim is to compute $M(\delta/L)$ or an upper bound for it. To the best of our knowledge there is no such bound for a general normed finite-dimensional vector space. Though, when restricting X to \mathbb{R}^n , the situation differs.

Remark 4.1 ([10, 19]). *Let $X = \mathbb{R}^n$ and $\text{vol}(\cdot)$ denote the Lebesgue measure (“volume”). Further, we define $B := B_1$ as the unit ball under the considered norm $\|\cdot\|$ and set $B/2 := B_{1/2}$. Then, it holds*

$$M(\delta/L) \leq T, \quad (6)$$

where

$$T := \frac{\text{vol}(\Omega + (\delta/L)B/2)}{\text{vol}((\delta/L)B/2)},$$

and the sum of two sets is considered the Minkowski sum.

Depending on the structure of Ω and the specific choice of norm, this leads to upper bounds dependent on δ and L . We have computed some examples in Appendix A and give an overview of the results in Table 1.

4.2. Problem Complexity. It is inadequate to expect finding exact solutions for abstract problem classes, as we consider them, in finite time; see [13, 20]. Hence, for the investigation of problem complexity, we only consider approximate solutions.

Definition 4.2. *Let $\varepsilon > 0$ and let r_q denote the q -th component of r , $q = 1, \dots, m$. Then, we call $x^* \in \Omega$ an ε -approximate solution for (P) if and only if*

$$\forall q = 1, \dots, m : r_q(x^*) \leq \varepsilon. \quad (7)$$

As shown in [20], continuity of f is not sufficient to find an approximate solution in a finite amount of steps. Thus, we assume the objective f to be also Lipschitz. Additionally, we assume the interior of Ω to be non-empty, and that there exists an $\varepsilon > 0$ such that an optimal solution x^* to (P) is contained in an ε -ball inside Ω . This ensures that if NIC adds norm-induced cuts with a radius of ε , it is not possible to cut off every neighboring point in Ω .

For the lower bound on the problem complexity, we use a result from [13]. We define the following constants regarding Ω .

Definition 4.3 ([13]). *Let $\varepsilon > 0$ and let Ω have a non-empty interior. Then, we set the radius of Ω as*

$$\rho(\Omega) := \min\{s \mid \exists x \in \Omega : \|x - y\| \leq s, \text{ for all } y \in \Omega\}.$$

Further, the asphericity of Ω is defined as

$$\alpha(\Omega) = \inf\{\beta \mid \exists x, y \in \Omega, s \geq 0 : B_s(x) \subseteq \Omega \subseteq B_{\beta s}(y)\}.$$

In other terms, the asphericity defines the ratio of radii of the minimal ball containing Ω and the maximal ball contained in Ω .

Since the magnitude of each r_q has a great influence on how to choose a proper approximation guarantee ε , we use the radius $\rho(\Omega)$ and assume r to be divided by the constant $\rho(\Omega)L$, “normalizing” the inequality, i.e., $r \leftarrow r/(\rho(\Omega)L)$.

Now, when dealing with problem complexity, we consider the number of oracle calls (step 2) until achieving an approximate solution. We denote this number by $T(\varepsilon)$ for $\varepsilon > 0$. So far, the NIC method has been developed to find exact solutions. As such, we modify it as follow:

- a) We only check for the component-wise inequalities (7) in the if-clause before step 9,
- b) and we add the norm-induced cut from Definition 2.1 with respect to a slightly changed radius of $\max\{\|r(x^k)_+\|/L, \varepsilon\}$.

Given these modifications, the number of oracle calls required to find the ε -approximate solution is given by the following theorem.

Theorem 4.4 ([13]). *Let $\varepsilon > 0$ and let Ω have a non-empty interior. Then,*

$$T(\varepsilon) \geq \left(\frac{c}{\alpha(\Omega)}\right)^n \frac{1}{\varepsilon^n},$$

with some constant $c > 0$.

For the upper bound, we use Remark 4.1 with ε instead of δ/L . That is, instead of giving a termination bound dependent on the values of r , we do it in terms of the approximation guarantee ε . We formalize this in the following.

Theorem 4.5. *Let $\varepsilon > 0$ and let Ω have a non-empty interior. Then,*

$$T(\varepsilon) \leq (2\rho(\Omega) + \varepsilon)^n \frac{1}{\varepsilon^n}.$$

Proof. With the definition of the radius in Definition 4.3, there exists $\bar{x} \in \Omega$ such that $\Omega \subseteq B_{\rho(\Omega)}(\bar{x})$. Together with Remark 4.1, we conclude

$$\begin{aligned} T(\varepsilon) &\leq \frac{\text{vol}(\Omega + \varepsilon B/2)}{\text{vol}(\varepsilon B/2)} \leq \frac{\text{vol}(B_{\rho(\Omega)} + \varepsilon B/2)}{\text{vol}(\varepsilon B/2)} \\ &= \frac{\text{vol}(B_{\rho(\Omega)+\varepsilon/2})}{\text{vol}(\varepsilon B/2)} = \left(\frac{\rho(\Omega) + \varepsilon/2}{\varepsilon/2}\right)^n \\ &= \left(\frac{2\rho(\Omega)}{\varepsilon} + 1\right)^n = (2\rho(\Omega) + \varepsilon)^n \frac{1}{\varepsilon^n}. \end{aligned}$$

□

In conclusion, considering the dimension n fixed, we observe that the lower and upper complexity bound show identical order of magnitude in ε .

5. COMPUTATIONAL EXAMPLES AND ILLUSTRATIONS

Having investigated the problem complexity, we now aim to illustrate the behavior of NIC. We give an example of a two dimensional problem, involving non-linear constraints in the form of a sine function. We note that such a problem can not be treated by standard solvers such as Gurobi [9] without reformulation.

k	(x_1^k, x_2^k)	$r(x^k)$	$ r(x^k) /L$
0	$(-1, -1)$	≈ 1.84	≈ 1.30
1	$\approx (-0.08, -0.08)$	≈ 0.16	≈ 0.11
2	$\approx (4 \cdot 10^{-5}, 4 \cdot 10^{-5})$	$\approx 8 \cdot 10^{-5}$	-

TABLE 2. Incumbents and violations for Example 5.1.




FIGURE 1. Illustration of Example 5.1

Example 5.1. Consider the non-linear problem

$$\begin{aligned} \min_{x_1, x_2} & |x_1 - x_2| + x_1 \\ \text{s.t.} & -\sin(x_1) - x_2 \leq 0, \\ & x_1, x_2 \in [-1, 1]. \end{aligned} \quad (P_{\sin})$$

Following our notation, we have $f(x) = f(x_1, x_2) = |x_1 - x_2| + x_1$, $r(x) = -\sin(x_1) - x_2$, and $\Omega = [-1, 1]^2$. The optimal solution of Problem (P_{\sin}) is $x^* = (0, 0)$. See Figure 1a for a graphical representation of x^* and the feasible region $Q = \Omega \cap \{x \in \mathbb{R}^2 \mid r(x) \leq 0\}$. In the following, we omit the subscript ‘+’, as r is 1-dimensional.

We specify an accuracy of $\varepsilon = 10^{-4}$. Further, we consider the \mathbb{R}^2 with the Euclidean norm $\|\cdot\|_2 = \sqrt{x_1^2 + x_2^2}$ and the \mathbb{R} with the absolute value $|\cdot|$.

As r is differentiable, we can apply the mean value theorem and calculate its global Lipschitz constant L on Ω as the maximal norm of the gradient of r over Ω . In a formal manner, this reads

$$L = \max_{x \in \Omega} \|\nabla r(x)\|_2 = \max_{x \in \Omega} \left\| \begin{pmatrix} -\cos(x_1) \\ -1 \end{pmatrix} \right\|_2 = \sqrt{2}.$$

In the first iteration, the algorithm initializes $k \leftarrow 0$ and solves $\min \{f(x) \mid x \in Q_0 = \Omega\}$. We summarize the resulting incumbents, values of $r(x^k)$ and the radius of the norm-induced cuts in Table 2.

We illustrate the cutting procedure in Figure 1b. The first cut in iteration 0 is represented with a dashed circle and its center x^0 with a black dot in the lower left corner. A dotted circle visualizes the second cut in iteration 1 around its center x^1 . The next found point x^2 is visualized in a different color and is the ε -optimal solution.

Note that we globally solved the subproblems tackled in the above example. In Theorem 3.8 we have shown that a point returned in step 9 under a local method for step 2 is a locally optimal point for (P). If no such point is returned, i.e., NIC creates an infinite series converging to the worst possible point in worst case. In the following, we give an example which shows this behavior.

Example 5.2. *Here, we consider a one dimensional problem in \mathbb{R} with the absolute value $|\cdot|$ which reads*

$$\begin{aligned} \min \quad & -|x| \\ \text{s.t.} \quad & \frac{1}{3}x^3 \geq 0, \\ & x \in [-1, 1]. \end{aligned} \tag{P_{\text{bad}}}$$

Following our notation, we have $f(x) = -|x|$, $r(x) = -\frac{1}{3}x^3$, and $\Omega = [-1, 1]$. When we calculate the Lipschitz constant of r on Ω , we can use the differentiability of r and the fraction to get $L = \max_{x \in \Omega} |r'(x)| = \max_{x \in [-1, 1]} |x^2| = 1$. Again, we omit the subscript ‘+’ due to the 1-dimensional r .

In order to understand Problem (P_{bad}), we can re-write the problem as $\max \{x \mid x \in [0, 1]\}$ and see that its optimal point is at $x^* = 1$. In contrast, the “worst” possible point in terms of the optimization problem would be $\hat{x} = 0$, as this is the maximum of Problem (P_{bad}). We will show that NIC converges to \hat{x} if the solution method used in step 2 is only a local one.

We assume step 2 to always return a locally optimal solution in $[-1, 0)$ if it exists. Therefore, relaxing the inequality $r(x) \leq 0$ in (P_{bad}), we receive the incumbent $x^0 = -1$ with a violation of $r(x^0) = 1/3$. Adding the cut $|x - x^0| \geq r(x^0)/L = 1/3$, we end up with $Q_1 = [-2/3, 1]$ as the feasible set of the next problem to investigate. We can show by induction that the incumbents produced by the solution method satisfy the following relation

$$x^{k+1} = x^k + r(x^k) = x^k - \frac{1}{3}(x^k)^3, \tag{8}$$

for $k \in \mathbb{N}_0$. Further, using induction once more, we can prove that indeed $x^k \in [-1, 0)$ for all $k \in \mathbb{N}_0$.

Now, we define the right-hand side of (8) as function $\nu(x) := x - \frac{1}{3}x^3$. Note that this is a contraction on $[-1, 0]$. Therefore, we can use Banach’s fixed-point theorem to see that ν has a unique fixed-point \hat{x} on $[-1, 0]$. This fixed-point is indeed $\hat{x} = 0$, which can be seen by $\nu(0) = 0$. Further, Banach’s theorem allows to show that by starting with any $x^0 \in [-1, 0]$, the sequence defined by $x^{k+1} = \nu(x^k)$, which is equivalent to (8), converges to \hat{x} . Therefore, $\lim_{k \rightarrow \infty} x^k = \hat{x} = 0$.

In summary, this shows that the sequence produced by NIC converges to the “worst” point. If we have a solution method returning a global optimal point, then the algorithm would return $x^* = 1$ in its second iteration at the latest.

Finally, we want to compare the performance when considering r itself versus the components of r separately. This was discussed in Remark 3.7 and is made explicit here.

Example 5.3. *We consider an example from the literature, in particular the continuous relaxation of Example 2.2 in [5]. Following previous notation, we set*

$$r(x) = r(x_1, x_2) = \begin{pmatrix} r_1(x_1, x_2) \\ r_2(x_1, x_2) \end{pmatrix} := \begin{pmatrix} \cos(6x_1)/2 - x_2 + 1.8 \\ -2 \sin(4x_1)/\sqrt{x_1} + x_2 - 2 \end{pmatrix},$$

which allows to give the problem in [5] as

$$\begin{aligned} \min_{x_1, x_2} \quad & x_1 + 4x_2 \\ \text{s.t.} \quad & r(x_1, x_2) \leq \mathbf{0}, \\ & x_1 \in [1, 10], x_2 \in [0, 4]. \end{aligned} \tag{P}_{\text{comp}}$$

In order to use Gurobi [9] as a solver for the sub-problems, as mentioned in the introducing comment of this section, we tackle Problem (P_{comp}) with NIC with respect to the Euclidean norm. That is, we added the norm-induced cuts in a squared form, resulting in non-convex quadratic cuts, for which Gurobi is known to perform well.

Regarding the Lipschitz constants, we compute their (squared) value for r_1 and r_2 by maximizing the (squared) Euclidean norm of the respective gradient over the box $[1, 10] \times [0, 4]$. As these are box-constrained non-linear optimization problems, their solution can be obtained by Newton-type methods. This results in (squared) Lipschitz constants $L_1^2 = 10$ and $L_2^2 \approx 42.83$ for r_1 and r_2 , respectively. The Lipschitz constant L for r itself can be obtained by maximizing the squared Euclidean operator norm of its Jacobian. In this case, by investigating the derivatives of r_1 and r_2 with respect to x_2 (as part of the Jacobian) we can show that the maximal value of the squared derivatives with respect to x_1 is obtained by $L_i^2 - 1$, $i = 1, 2$, in particular, $L^2 = L_1^2 - 1 + L_2^2 - 1 \approx 50.83$.

Now, we leverage these values to relax constraint $r(x_1, x_2) \leq \mathbf{0}$ and apply norm-induced cuts of the form

$$\|x - x^k\|_2^2 \geq \frac{\|r(x_1^k, x_2^k)_+\|_2^2}{L^2},$$

when considering r as one function, or

$$\|x - x^k\|_2^2 \geq \frac{|r_1(x_1^k, x_2^k)|^2}{L_1^2}, \text{ and } \|x - x^k\|_2^2 \geq \frac{|r_2(x_1^k, x_2^k)|^2}{L_2^2},$$

if $r_1(x_1^k, x_2^k) > 0$ and $r_2(x_1^k, x_2^k) > 0$, respectively, when considering the components of r separately. Running NIC for 100 iterations with Gurobi, we can compare the convergence of both alternatives to the optimal value in Figure 2. We note that the optimal objective value of Problem (P_{comp}) is approximately 6.76.

One can observe that using the components of r separately gives the opportunity of faster convergence to the optimal objective value. In iteration 100, the relative difference of the optimal value of (P_{comp}) to the current value of the relaxation (P_k) is approximately 0.58% when using L_1 , L_2 , and 18.55% when using just L . This results from the additional amount of information, i.e., the Lipschitz constants of the components. In particular, when only one constraint $r_1 \leq 0$ or $r_2 \leq 0$ is violated, considering them separately adds cuts with smaller Lipschitz constants, thus, larger right-hand side than when considering only r . Hence, the former leads to a tighter approximation of the feasible region and, hence, leads to faster convergence.

However, one must recognize that in practice the Lipschitz constant might not be available for the components of a multi-dimensional function, but only for the function itself.

6. CONCLUSION

We have introduced the NIC method in order to tackle the abstract problem class (P). Besides a continuous objective on a compact domain, the method only requires a Lipschitz property of the constraint functions. Leveraging an oracle for the global solution of (P_k), we have proven the correctness of NIC in terms of returning an optimal solution when terminating or, if not, creating a sequence whose




FIGURE 2. Objective value per iteration of NIC when considering entire r or its components with respective Lipschitz constants

accumulation points are all optimal. This has been complemented by discussions about the termination bound in the infeasible case and the problem complexity, showing asymptotic convergence of the bounds in the latter. The theoretical content has been supplemented by examples, illustrating the method, visualizing the possible effects of a local oracle, and demonstrating the benefit of knowing Lipschitz constants for the components of the constraint function.

While there exist optimization software like Gurobi [9] which successfully tackle quadratic constraint problems, they struggle when facing general non-linear constraints. Our approach enables such software to solve even more general problems by executing NIC using them as an oracle.

As observed especially in the examples, the magnitude of the radius of the norm-induced cuts depends on the Lipschitz constant L in the denominator and is crucial. If an evaluation of the constraint function is cheap, one can compute an approximation of the local Lipschitz constant and, thus, reduce its value. Apart from that, applications with small L seem reasonable. For example, the field of optimal control of power networks include sine or cosine constraints with small L and, thus, could be tackled by NIC. This gives rise to another interesting topic of research.

ACKNOWLEDGEMENTS

We thank the Deutsche Forschungsgemeinschaft for their support within project A05 in the *Sonderforschungsbereich/Transregio 154 Mathematical Modelling, Simulation and Optimization using the Example of Gas Networks*.

Further, we thank Martin Schmidt for a fruitful discussion about the distinction to the present article.

REFERENCES

- [1] S. A. Altman and C. R. Bastian. “DHL Global Connectedness Index 2022.” In: (2023). DOI: <https://doi.org/10.58153/wjkw-vjs44>.
- [2] C. Audet and J. E. Dennis. “A Pattern Search Filter Method for Nonlinear Programming without Derivatives.” In: *SIAM Journal on Optimization* 14.4 (2004), pp. 980–1010. DOI: [10.1137/S105262340138983X](https://doi.org/10.1137/S105262340138983X).

- [3] C. Audet and J. E. Dennis. “Mesh Adaptive Direct Search Algorithms for Constrained Optimization.” In: *SIAM Journal on Optimization* 17.1 (2006), pp. 188–217. DOI: [10.1137/040603371](https://doi.org/10.1137/040603371).
- [4] F. V. Berghen. “CONDOR, a parallel, direct, constrained optimizer for high-computing-load, black box objective functions.” In: *Proceedings of the third MIT conference on Computational Fluid and Solid Mechanics, Elsevier*. 2005.
- [5] T. Berthold. “Heuristic algorithms in global MINLP solvers.” PhD thesis. 2014, p. 366. URL: <http://www.zib.de/berthold/Berthold2014.pdf>.
- [6] D. Bertsimas, O. Nohadani, and K. M. Teo. “Robust optimization for unconstrained simulation-based problems.” In: *Operations research* 58.1 (2010), pp. 161–178.
- [7] A. R. Conn, K. Scheinberg, and L. N. Vicente. *Introduction to Derivative-Free Optimization*. Philadelphia: SIAM, 2009. DOI: [10.1137/1.9780898718768](https://doi.org/10.1137/1.9780898718768). eprint: <https://epubs.siam.org/doi/pdf/10.1137/1.9780898718768>.
- [8] J. Grübel, R. Krug, M. Schmidt, and W. Wollner. “A Successive Linear Relaxation Method for MINLPs with Multivariate Lipschitz Continuous Nonlinearities.” In: (2023). arXiv: [2208.06444](https://arxiv.org/abs/2208.06444) [math.OC].
- [9] Gurobi Optimization, LLC. *Gurobi Optimizer Reference Manual*. 2023. URL: <https://www.gurobi.com>.
- [10] Y. Han. *Lecture notes in Informatic-theoretic Lower Bounds in Data Science (Lecture 12)*. May 2021. URL: https://web.stanford.edu/class/ee378c/scribe_lec12.pdf.
- [11] R. Horst and H. Tuy. *Global optimization: Deterministic approaches*. Third. Heidelberg: Springer Berlin, 1996. DOI: [10.1007/978-3-662-03199-5](https://doi.org/10.1007/978-3-662-03199-5).
- [12] R. M. Lewis and V. Torczon. “Pattern Search Algorithms for Bound Constrained Minimization.” In: *SIAM Journal on Optimization* 9.4 (1999), pp. 1082–1099. DOI: [10.1137/S1052623496300507](https://doi.org/10.1137/S1052623496300507).
- [13] A. S. Nemirovskij and D. B. Yudin. *Problem complexity and method efficiency in optimization*. Norwich: Wiley-Interscience, 1983.
- [14] B. Ozturk and A. Saab. “Optimal aircraft design decisions under uncertainty via robust signomial programming.” In: *AIAA Aviation 2019 Forum*. 2019, p. 3351.
- [15] S. Piyavskii. “An algorithm for finding the absolute extremum of a function.” In: *USSR Computational Mathematics and Mathematical Physics* 12.4 (1972), pp. 57–67. DOI: [https://doi.org/10.1016/0041-5553\(72\)90115-2](https://doi.org/10.1016/0041-5553(72)90115-2).
- [16] W. Rudin. *Principles of Mathematical Analysis*. International series in pure and applied mathematics. McGraw-Hill, 1976. URL: <https://books.google.de/books?id=kwqzPAAACAAJ>.
- [17] M. Schmidt, M. Sirvent, and W. Wollner. “A decomposition method for MINLPs with Lipschitz continuous nonlinearities.” In: *Mathematical Programming* 178.1 (2019), pp. 449–483. DOI: [10.1007/s10107-018-1309-x](https://doi.org/10.1007/s10107-018-1309-x).
- [18] M. Schmidt, M. Sirvent, and W. Wollner. “The cost of not knowing enough: mixed-integer optimization with implicit Lipschitz nonlinearities.” In: *Optimization Letters* 16.5 (2022), pp. 1355–1372.
- [19] V. M. Tikhomirov. “ ε -Entropy and ε -Capacity of Sets in Functional Spaces.” In: *Selected Works of A. N. Kolmogorov: Volume III: Information Theory and the Theory of Algorithms*. Ed. by A. N. Shirayev. Dordrecht: Springer, 1993, pp. 86–170. DOI: [10.1007/978-94-017-2973-4_7](https://doi.org/10.1007/978-94-017-2973-4_7).
- [20] S. A. Vavasis. *Nonlinear Optimization: Complexity Issues*. New York: Oxford University Press, 1991.

APPENDIX A. EXAMPLES FOR FORMULA (6)

Let $L > 0$ be the (upper bound to the) Lipschitz constant and $\delta > 0$, such that $\delta/L < 1$. We calculate the examples as appearing in Table 1. For the sake of clarity, we define the unit ball $B = B_1$ with respect to the considered norm.

- a) Let $\Omega_1 = \prod_{j=1}^n [a_j, b_j]$ be an n -dimensional box with $a_j, b_j \in \mathbb{R}$ and $a_j < b_j$ for $j = 1, \dots, n$. We consider the maximum-norm, i.e., $\|x\|_\infty = \max_{i \in [n]} |x_i|$ for $x \in \mathbb{R}^n$ and $[n] := \{1, \dots, n\}$.

In this case, we have $B = [-1, 1]^n$ and, thus, $\text{vol}(B) = 2^n$. It follows, $\text{vol}((\delta/L)B/2) = (\delta/L)^n$ and

$$\Omega_1 + (\delta/L)B/2 = \prod_{j=1}^n [a_j - \frac{\delta}{2L}, b_j + \frac{\delta}{2L}].$$

Therefore, $\text{vol}(\Omega_1 + (\delta/L)B/2) = \prod_{j=1}^n (b_j - a_j + \delta/L)$ and we conclude

$$T = \frac{\text{vol}(\Omega_1 + (\delta/L)B/2)}{\text{vol}((\delta/L)B/2)} = \prod_{j=1}^n \left[\frac{L}{\delta} (b_j - a_j) + 1 \right].$$

- b) Let us consider the maximum-norm again and $\Omega_2 = \Omega_1 \cap \mathbb{Z}^n$ to be the integral (lattice) points in Ω_1 . Then, we can calculate the number of lattice points as

$$|\Omega_2| = \prod_{j=1}^n ([b_j] - [a_j] + 1).$$

Since we assumed $\delta/L < 1$ in the beginning, it holds true that

$$(\{z_1\} + (\delta/L)B/2) \cap (\{z_2\} + (\delta/L)B/2) = \emptyset,$$

for $z_1, z_2 \in \Omega_2$, $z_1 \neq z_2$. Therefore, we can derive the volume $\text{vol}(\Omega_2 + (\delta/L)B/2) = |\Omega_2| \text{vol}((\delta/L)B/2)$ and, thus,

$$T = \frac{\text{vol}(\Omega_2 + (\delta/L)B/2)}{\text{vol}((\delta/L)B/2)} = \frac{|\Omega_2| \text{vol}((\delta/L)B/2)}{\text{vol}((\delta/L)B/2)} = |\Omega_2|.$$

- c) Here, we consider the Euclidean or 2-norm, i.e., $\|x\|_2 = \sqrt{\left(\sum_{j=1}^n x_j^2\right)}$. For some radius $r > 0$, we have B_r and can derive from standard literature that

$$\text{vol}(B_r) = \frac{\pi^{n/2}}{\Gamma(\frac{n}{2} + 1)} r^n = \text{vol}(B) r^n,$$

where $\Gamma(\cdot)$ denotes the gamma function. Now, for $D > 0$, let $\Omega_3 = B_D$ and it follows that

$$\text{vol}(\Omega_3 + (\delta/L)B/2) = \text{vol}(B_{D+(\delta/(2L))}) = \text{vol}(B)(D + (\delta/(2L)))^n.$$

Therefore, we can conclude

$$T = \frac{\text{vol}(B_D + (\delta/L)B/2)}{\text{vol}((\delta/L)B/2)} = \frac{(D + (\delta/(2L)))^n}{(\delta/(2L))^n} = \left(\frac{2L}{\delta} D + 1\right)^n.$$

(A. Göß) UNIVERSITY OF TECHNOLOGY NUREMBERG, ANALYTICS & OPTIMIZATION LAB, UL-MENSTR. 52I, 90433 NÜRNBERG, GERMANY

Email address: adrian.goess@utn.de

(A. Martin) UNIVERSITY OF TECHNOLOGY NUREMBERG, ANALYTICS & OPTIMIZATION LAB, UL-MENSTR. 52I, 90443 NÜRNBERG, GERMANY

Email address: alexander.martin@utn.de

(S. Pokutta) ZUSE INSTITUTE BERLIN, MATHEMATICAL ALGORITHMIC INTELLIGENCE,
TAKUSTR. 7, 14195 BERLIN, GERMANY
Email address: `pokutta@zib.de`

(K. Sharma) ZUSE INSTITUTE BERLIN, MATHEMATICAL ALGORITHMIC INTELLIGENCE,
TAKUSTR. 7, 14195 BERLIN, GERMANY
Email address: `kartikey.sharma@zib.de`