

Shadow Hamiltonians of structure-preserving integrators for Nambu mechanics

Atsushi Horikoshi*

Department of Natural Sciences, Tokyo City University, Tokyo 158-8557, Japan

Symplectic integrators are widely implemented numerical integrators for Hamiltonian mechanics, which preserve the Hamiltonian structure (symplecticity) of the system. Although the symplectic integrator does not conserve the energy of the system, it is well known that there exists a conserving modified Hamiltonian, called the shadow Hamiltonian. For the Nambu mechanics, which is one of the generalized Hamiltonian mechanics, we can also construct structure-preserving integrators by the same procedure used to construct the symplectic integrators. In the structure-preserving integrator, however, the existence of shadow Hamiltonians is non-trivial. This is because the Nambu mechanics is driven by multiple Hamiltonians and it is non-trivial whether the time evolution by the integrator can be cast into the Nambu mechanical time evolution driven by multiple shadow Hamiltonians. In the present paper we construct structure-preserving integrators for a simple Nambu mechanical system, and derive the shadow Hamiltonians in two ways. This is the first attempt to derive shadow Hamiltonians of structure-preserving integrators for Nambu mechanics. We show that the fundamental identity, which corresponds to the Jacobi identity in Hamiltonian mechanics, plays an important role to calculate the shadow Hamiltonians using the Baker-Campbell-Hausdorff formula. It turns out that the resulting shadow Hamiltonians have indefinite forms depending on how the fundamental identities are used. This is not a technical artifact, because the exact shadow Hamiltonians obtained independently have the same indefiniteness.

I. INTRODUCTION

In 1973, Nambu proposed a generalized Hamiltonian mechanics, now called the Nambu mechanics [1]. He extended the phase space spanned by the canonical doublet (q, p) to one spanned by a multiplet (x_1, x_2, \dots, x_N) , where $N \geq 3$, and using the Liouville theorem as a guiding principle, he generalized the Hamiltonian H to multiple Hamiltonians $(H_1, H_2, \dots, H_{N-1})$, the Poisson bracket to the Nambu bracket, and the Hamilton's equations to the Nambu equations. It was later found that the Jacobi identity for the Poisson bracket should be generalized to the fundamental identity for the Nambu bracket [2, 3].

As an example of the Nambu mechanics, Nambu showed that the Euler equations for the free rigid body can be derived from the Nambu equations [1]. However, applications of the Nambu mechanics have been limited to some particular systems so far, although there have been some interesting applications in the field of fluid mechanics (see, for example, Refs.[4, 5]). This is because the systems described by the Nambu mechanics should have multiple conserved quantities as the Hamiltonians. Recently we have proposed a novel approach to the Nambu mechanics, the hidden Nambu formalism, and revealed that the Nambu mechanical structures are hidden in any Hamiltonian systems [6]. For example, a Hamiltonian system of (q, p) with a Hamiltonian H can be described by the $N = 3$ Nambu system of $(x_1, x_2, x_3) = (q, p, q^2)$ with the Hamiltonians $H_1 = H$ and $H_2 = x_3 - x_1^2$. Here the second Hamiltonian H_2 works as a constraint between variables. This formalism allows us to investigate the properties of the Nambu mechanics in various systems including semiclassical systems [7–9].

As for numerical integration algorithms for solving ordinary differential equations, a number of structure-preserving integrators have been proposed over the decades [10]. For Hamiltonian systems, the symplectic integrators are well known, which are numerical integration algorithms that preserve the Hamiltonian structure of the systems. The symplectic integrator can be constructed by splitting the Hamiltonian H into some pieces and composing the time evolution operators associated with those pieces. Therefore the symplectic integrator does not conserve the energy, i.e., the original Hamiltonian H . However it is also well known that the symplectic integrator conserve a modified Hamiltonian, called the shadow Hamiltonian H_S , which is slightly different from the original Hamiltonian [10, 11]. The shadow Hamiltonians are in general calculated order by order using the Baker-Campbell-Hausdorff (BCH) formula and the Jacobi identity. For the harmonic oscillator, some exact expressions of the shadow Hamiltonians have been found by calculations without the BCH formula [12, 13]. Since the existence of the shadow Hamiltonian guarantees that the error of the original Hamiltonian does not grow secularly, the symplectic integrators have been used for long-time simulations of various Hamiltonian systems [10, 11]. Symplectic integrators with various degrees of precision

* horikosi@tcu.ac.jp

have been proposed, and one can choose the appropriate integrators for the purpose (for the Python package, see [14], for example).

On the other hand, for the Nambu systems, we can also construct structure-preserving integrators by the splitting and composing procedure used to construct symplectic integrators for Hamiltonian systems [15]. Here we referred to those integrators not as the symplectic integrators, but as the more general name, because the Nambu systems do not have symplecticity. Similar to the symplectic integrators, the structure-preserving integrator preserve the Nambu mechanical structure of the system but does not conserve the original Hamiltonians. However, in contrast to the symplectic integrators, it is non-trivial whether the shadow Hamiltonians exist in the structure-preserving integrator. This is because it is non-trivial that the time evolution by the structure-preserving integrator can be described by a time evolution operator associated with the shadow Hamiltonians. Indeed, it has been suggested in Ref [15] that in the Nambu system corresponding to the Euler equations for the free rigid body, there are no shadow Hamiltonians of the structure-preserving integrator constructed by splitting all the Hamiltonians.

In the present paper, we study the shadow Hamiltonians of structure-preserving integrators for a simple $N = 3$ Nambu system, a harmonic oscillator described by three variables $(x_1, x_2, x_3) = (q, p, q^2)$, which is given by the hidden Nambu formalism [6, 7]. We attempt to calculate the shadow Hamiltonians using the BCH formula and the fundamental identity, and also we seek to find the exact expressions of the shadow Hamiltonians referring to the exact shadow Hamiltonians of the symplectic integrators [12, 13].

The outline of the paper is as follows. In Sect. 2 we give a slightly lengthy review of the symplectic integrators and related issues to emphasize the similarities and differences between the symplectic integrators and the structure-preserving integrators presented later. In Sect. 3 we review the Nambu mechanics and present a harmonic oscillator system described by three variables $(x_1, x_2, x_3) = (q, p, q^2)$ as an example of $N = 3$ Nambu system. In Sect. 4 we present the structure-preserving integrators for the $N = 3$ harmonic oscillator and show the details of the derivations of the shadow Hamiltonians with some numerical results. Conclusions are given in the last section.

II. HAMILTONIAN MECHANICS

A. Hamilton's equation

Consider a one degree of freedom Hamiltonian system consisting of a set of canonical variables: $\mathbf{x} = {}^t(x_1, x_2) = {}^t(q, p)$. Let $H(x_1, x_2)$ be the Hamiltonian of the system, then the Hamilton's equation of motion can be written as

$$\frac{df}{dt} = \{f, H\} \quad (1)$$

for any function $f(x_1, x_2)$. Here

$$\{A, B\} = \frac{\partial(A, B)}{\partial(x_1, x_2)} = \epsilon_{ij} \frac{\partial A}{\partial x_i} \frac{\partial B}{\partial x_j} \quad (2)$$

is the Poisson bracket and ϵ_{ij} is the 2D Levi-Civita symbol.

For a variable transformation $\mathbf{x} \rightarrow \mathbf{y}$, the equation of motion written in the new variable \mathbf{y} has the same form as in Eq. (1) if the transformation satisfies the condition

$$\frac{\partial y_i}{\partial x_k} \epsilon_{kl} \frac{\partial y_j}{\partial x_l} = \epsilon_{ij}. \quad (3)$$

This is called the symplectic condition. The time evolution along the exact solution of the Hamilton's equation (1) satisfies this condition. Under the exact short time evolution $\mathbf{x}(t) \rightarrow \mathbf{x}(t+h)$, where h is a small time step, the infinitesimal phase space volume is conserved: $dx_1(t+h)dx_2(t+h) = dx_1(t)dx_2(t)$. This is the Liouville theorem, which follows from the property that the velocity field $\dot{\mathbf{x}} = d\mathbf{x}/dt$ is divergence-free, $\text{div } \dot{\mathbf{x}} = 0$. The Liouville theorem can also be derived from the symplectic condition (3).

B. Symplectic integrators

Among various numerical integration schemes for Hamiltonian systems, the symplectic integrators are the ones that preserve symplecticity of the system [10, 11]. We define the Liouville operator X_H as

$$X_H = \epsilon_{ij} \frac{\partial H}{\partial x_j} \frac{\partial}{\partial x_i}, \quad (4)$$

and write the Hamilton's equation (1) as $df/dt = X_H f$, then the short time evolution of $f(t)$ can be formally written as

$$f(t+h) = e^{hX_H} f(t). \quad (5)$$

In case that the Hamiltonian is separable, $H(x_1, x_2) = T(x_2) + V(x_1)$, the Liouville operator X_H can be split into $X_H = X_T + X_V$, and each short time evolution can be evaluated as

$$e^{hX_T} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_1 + h \frac{\partial T}{\partial x_2} \\ x_2 \end{pmatrix}, \quad (6)$$

$$e^{hX_V} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_1 \\ x_2 - h \frac{\partial V}{\partial x_1} \end{pmatrix}. \quad (7)$$

Since these are exact expressions, each dynamics is symplectic.

Composing exact solutions (6)-(7), we can define various symplectic integrators for X_H . For example, the well-known position Verlet integrator can be constructed by composing three exact solutions,

$$\Phi_h^{TVT} = e^{\frac{h}{2}X_T} \circ e^{hX_V} \circ e^{\frac{h}{2}X_T}. \quad (8)$$

Since symplecticity is preserved in each time evolution, this is a symplectic integrator. Using the Trotter formula [11], the precision of this integrator can be evaluated as

$$e^{hX_H} = e^{\frac{h}{2}X_T} e^{hX_V} e^{\frac{h}{2}X_T} + O(h^3). \quad (9)$$

That is, the position Verlet integrator is a second-order symplectic integrator. This integrator also has the important property, the time-reversal symmetry: $\Phi_{-h}^{TVT} \circ \Phi_h^{TVT} = 1$. In the same way, we can construct another well-known second-order symplectic integrator, the velocity Verlet integrator $\Phi_h^{VTV} = e^{\frac{h}{2}X_V} \circ e^{hX_T} \circ e^{\frac{h}{2}X_V}$.

C. Shadow Hamiltonian

Although the symplectic integrator dose not conserve the energy (the original Hamiltonian H), it is known to conserve a quantity close to the original Hamiltonian. This means that the product of exponentials can be represented by an effective Liouville operator X_{eff} (taking the position Verlet integrator (8) as an example),

$$e^{hX_{\text{eff}}} = e^{\frac{h}{2}X_T} e^{hX_V} e^{\frac{h}{2}X_T}, \quad (10)$$

and there exists a modified Hamiltonian, called the *shadow Hamiltonian* H_S , such that X_{eff} can be written as

$$X_{\text{eff}} = X_{H_S} = \epsilon_{ij} \frac{\partial H_S}{\partial x_j} \frac{\partial}{\partial x_i}. \quad (11)$$

The functional form of X_{eff} can be obtained using the Baker-Campbell-Hausdorff (BCH) formula [10]. For example, applying the second order BCH formula to Eq. (10) yields

$$X_{\text{eff}} = X_T + X_V - \frac{h^2}{24} \left([X_T, [X_T, X_V]] - 2[X_V, [X_V, X_T]] \right) + O(h^3), \quad (12)$$

where $[X, Y] = XY - YX$ is the commutator. The commutator of two Liouville operators can be expressed in terms of a single Liouville operator,

$$\begin{aligned} [X_A, X_B]f &= X_A\{f, B\} - X_B\{f, A\} = \{\{f, B\}, A\} - \{\{f, A\}, B\} \\ &= \{f, \{B, A\}\} = X_{\{B, A\}}f. \end{aligned} \quad (13)$$

Here we used the Jacobi identity: $\{\{f, B\}, A\} = \{\{f, A\}, B\} + \{f, \{B, A\}\}$ or $\{\{f, A\}, B\} = \{\{f, B\}, A\} + \{f, \{A, B\}\}$. With either identity, we obtain the same result (13). Using Eq. (13) for the BCH formula (12) and identifying $X_{\text{eff}} = X_{H_S}$, we obtain the shadow Hamiltonian H_S for the position Verlet integrator (8),

$$H_S = H - \frac{h^2}{24} \left(\{\{V, T\}, T\} - 2\{\{T, V\}, V\} \right) + O(h^3). \quad (14)$$

Although the original Hamiltonian H is not conserved in the time evolution by the position Verlet integrator Φ_h^{TVT} , the shadow Hamiltonian H_S is conserved.

D. Example: Harmonic oscillator

We show an example of a harmonic oscillator,

$$H(x_1, x_2) = \frac{1}{2m}x_2^2 + \frac{m\omega^2}{2}x_1^2. \quad (15)$$

This Hamiltonian is separable, $H(x_1, x_2) = T(x_2) + V(x_1)$, where $T(x_2) = (1/2m)x_2^2$ and $V(x_1) = (m\omega^2/2)x_1^2$, and the Liouville operator X_H can be split into $X_H = X_T + X_V$, where

$$X_T = \frac{1}{m}x_2 \frac{\partial}{\partial x_1}, \quad X_V = -m\omega^2 x_1 \frac{\partial}{\partial x_2}. \quad (16)$$

Here we consider the position Verlet integrator (8). Using the BCH formula (12), the shadow Hamiltonian (14) can be evaluated as

$$H_S = H - \frac{\omega^2}{24m}h^2x_2^2 + \frac{m\omega^4}{12}h^2x_1^2 + O(h^3). \quad (17)$$

On the other hand, it can be shown by explicit calculation that the position Verlet integrator conserves the following quantity [13],

$$H_c = H - \frac{\omega^2}{8m}h^2x_2^2. \quad (18)$$

Although H_c is conserved, it is NOT the shadow Hamiltonian. That is, the Liouville operator associated with H_c ,

$$X_{H_c} = \frac{1}{m} \left(1 - \frac{\omega^2 h^2}{4}\right) x_2 \frac{\partial}{\partial x_1} - m\omega^2 x_1 \frac{\partial}{\partial x_2}, \quad (19)$$

is not the effective operator X_{eff} . The exact shadow Hamiltonian H_S^e in the case of the harmonic oscillator has been derived by some authors [12, 13],

$$X_{\text{eff}} = X_{H_S^e} = F(\omega h) X_{H_c}, \quad (20)$$

$$H_S^e = F(\omega h) H_c, \quad (21)$$

where the factor $F(x)$ is given by

$$F(x) = \sum_{n=0}^{\infty} \frac{(n!)^2}{(2n+1)!} x^{2n} = \frac{2 \arcsin(\frac{x}{2})}{x \sqrt{1 - \frac{x^2}{4}}}, \quad (22)$$

$$= 1 + \frac{1}{6}x^2 + \frac{1}{30}x^4 + O(x^6), \quad (23)$$

for $0 < |x| < 2$.

III. NAMBU MECHANICS

A. Nambu equation

Nambu generalized the Hamiltonian mechanics by generalizing the canonical doublet q and p to an N -plet, where $N \geq 3$ [1]. Here we consider $N = 3$ system: $\mathbf{x} = {}^t(x_1, x_2, x_3)$. Nambu then generalized the Hamilton's equation of motion (1) to the Nambu equation,

$$\frac{df}{dt} = \{f, H, G\}. \quad (24)$$

Here H and G are *Hamiltonians*, and $\{*, *, *\}$ is the Nambu bracket,

$$\{A, B, C\} = \frac{\partial(A, B, C)}{\partial(x_1, x_2, x_3)} = \epsilon_{ijk} \frac{\partial A}{\partial x_i} \frac{\partial B}{\partial x_j} \frac{\partial C}{\partial x_k}, \quad (25)$$

where ϵ_{ijk} is the 3D Levi-Civita symbol. Nambu mechanics is N -variable dynamics driven by $N - 1$ Hamiltonians, and those Hamiltonians are conserved.

For a variable transformation $\mathbf{x} \rightarrow \mathbf{y}$, the equation of motion written in the new variable \mathbf{y} has the same form as in Eq. (24) if the transformation satisfies the condition

$$\epsilon_{lmn} \frac{\partial y_i}{\partial x_l} \frac{\partial y_j}{\partial x_m} \frac{\partial y_k}{\partial x_n} = \epsilon_{ijk}. \quad (26)$$

This is similar to the symplectic condition (3), but not the same as it. The time evolution along the exact solution of the Nambu equation (24) satisfies this condition (26). As with the Hamiltonian mechanics, the Nambu mechanics is divergence-free, $\text{div } \dot{\mathbf{x}} = 0$, and therefore the Liouville theorem also holds in the Nambu mechanics.

B. Example: $N = 3$ harmonic oscillator

The systems described by the Nambu mechanics have been known only in limited examples, such as the Euler equations for the free rigid body [1]. In 2013 we proposed the hidden Nambu formalism and showed that systems extended to include composite variables can be described by the Nambu mechanics [6, 7]. As an example, here we consider a harmonic oscillator (15) with $N = 3$ variables, one of them being a composite variable,

$$x_1 = q, \quad x_2 = p, \quad x_3 = q^2. \quad (27)$$

Defining two Hamiltonians as

$$H(x_1, x_2, x_3) = \frac{1}{2m} x_2^2 + \frac{m\omega^2}{2} x_3, \quad (28)$$

$$G(x_1, x_2, x_3) = x_3 - x_1^2, \quad (29)$$

the Nambu equations (24) read,

$$\frac{d}{dt} x_1 = \frac{1}{m} x_2, \quad \frac{d}{dt} x_2 = -m\omega^2 x_1, \quad \frac{d}{dt} x_3 = \frac{2}{m} x_1 x_2. \quad (30)$$

These are consistent with the Hamilton equations for q , p , and q^2 .

IV. NUMERICAL INTEGRATORS FOR NAMBU MECHANICS

A. Structure-preserving integrators

In the Nambu mechanics, the condition (26) holds instead of the symplectic condition (3). In this paper we refer to the integrators that satisfy the condition (26) as structure-preserving integrators. Let us take the $N = 3$ harmonic oscillator presented in the previous section as an example to see how to construct the structure-preserving integrators. We define the generalized Liouville operator $X_{H,G}$ as

$$X_{H,G} = \epsilon_{ijk} \frac{\partial H}{\partial x_j} \frac{\partial G}{\partial x_k} \frac{\partial}{\partial x_i}. \quad (31)$$

Then the Nambu equation (24) can be written as $df/dt = X_{H,G}f$ and the short time evolution of $f(t)$ can be formally expressed as

$$f(t+h) = e^{hX_{H,G}} f(t). \quad (32)$$

For the $N = 3$ harmonic oscillator system, two Hamiltonians (28)-(29) are separable,

$$H(x_1, x_2, x_3) = A(x_2) + B(x_3), \quad (33)$$

$$G(x_1, x_2, x_3) = C(x_3) + D(x_1), \quad (34)$$

where

$$A(x_2) = \frac{1}{2m} x_2^2, \quad B(x_3) = \frac{m\omega^2}{2} x_3, \quad (35)$$

$$C(x_3) = x_3, \quad D(x_1) = -x_1^2. \quad (36)$$

The Liouville operator $X_{H,G}$ can be split into $X_{H,G} = X_{A,C} + X_{A,D} + X_{B,C} + X_{B,D}$. Noting that $X_{B,C} = 0$, we define the following three Liouville operators such that $X_{H,G} = X_1 + X_2 + X_3$,

$$X_1 = X_{A,C} = \frac{1}{m}x_2 \frac{\partial}{\partial x_1}, \quad (37)$$

$$X_2 = X_{B,D} = -m\omega^2 x_1 \frac{\partial}{\partial x_2}, \quad (38)$$

$$X_3 = X_{A,D} = \frac{2}{m}x_1 x_2 \frac{\partial}{\partial x_3}. \quad (39)$$

Here X_1 and X_2 correspond to X_T and X_V in Eqs. (16), respectively. Each Liouville operator gives exact short time evolution as follows,

$$e^{hX_1} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} x_1 + \frac{1}{m}hx_2 \\ x_2 \\ x_3 \end{pmatrix}, \quad (40)$$

$$e^{hX_2} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} x_1 \\ x_2 - m\omega^2 hx_1 \\ x_3 \end{pmatrix}, \quad (41)$$

$$e^{hX_3} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 + \frac{2}{m}hx_1x_2 \end{pmatrix}. \quad (42)$$

In each time evolution, the condition (26) is satisfied and the Liouville theorem holds.

Composing exact solutions (40)-(42), we can define various integrators for $X_{H,G}$. For example, by composing five exact solutions symmetrically, we can construct an integrator,

$$\Phi_h^{12321} = e^{\frac{h}{2}X_1} \circ e^{\frac{h}{2}X_2} \circ e^{hX_3} \circ e^{\frac{h}{2}X_2} \circ e^{\frac{h}{2}X_1}. \quad (43)$$

Since the condition (26) is satisfied by each step, the integrator Φ_h^{12321} keeps the condition (26). Therefore Φ_h^{12321} is a structure-preserving integrator. This is a second-order integrator because its precision can be evaluated using the Trotter formula [11],

$$e^{hX_{H,G}} = e^{\frac{h}{2}X_1} e^{\frac{h}{2}X_2} e^{hX_3} e^{\frac{h}{2}X_2} e^{\frac{h}{2}X_1} + O(h^3). \quad (44)$$

This integrator also has the time-reversal symmetry: $\Phi_h^{12321} \circ \Phi_h^{12321} = 1$. We can construct a total of $3! = 6$ types of second-order structure-preserving integrators by composing five exact solutions symmetrically. It should be noted here that $X_1 = X_T$ and $X_2 = X_V$, and they do not depend on x_3 . That is, the time evolution of x_3 is decoupled from the time evolutions of x_1 and x_2 , yielding

$$\Phi_h^{12321} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \Phi_h^{13231} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \Phi_h^{31213} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \Phi_h^{TVTV} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \quad (45)$$

$$\Phi_h^{23132} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \Phi_h^{21312} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \Phi_h^{32123} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \Phi_h^{VTVV} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}. \quad (46)$$

B. Shadow Hamiltonians

The product of exponentials can be represented by an effective Liouville operator X_{eff} . For the integrator Φ_h^{12321} ,

$$e^{hX_{\text{eff}}} = e^{\frac{h}{2}X_1} e^{\frac{h}{2}X_2} e^{hX_3} e^{\frac{h}{2}X_2} e^{\frac{h}{2}X_1}. \quad (47)$$

However, it is non-trivial whether there exist *shadow Hamiltonians* H_S and G_S such that X_{eff} can be written as

$$X_{\text{eff}} = X_{H_S, G_S} = \epsilon_{ijk} \frac{\partial H_S}{\partial x_j} \frac{\partial G_S}{\partial x_k} \frac{\partial}{\partial x_i}. \quad (48)$$

Using the BCH formula [10], X_{eff} can be written as

$$\begin{aligned}
X_{\text{eff}} = X_1 + X_2 + X_3 - \frac{h^2}{24} & \left([X_1, [X_1, X_2]] + [X_1, [X_1, X_3]] + [X_1, [X_2, X_3]] \right. \\
& - 2[X_2, [X_2, X_1]] + [X_2, [X_2, X_3]] + [X_2, [X_1, X_3]] \\
& \left. - 2[X_3, [X_3, X_1]] - 2[X_3, [X_3, X_2]] + 3[X_3, [X_1, X_2]] \right) \\
& + O(h^3).
\end{aligned} \tag{49}$$

Here $X_1 + X_2 + X_3 = X_{H,G}$, and in the present case we can see that some commutators are zero, $[X_1, [X_1, X_3]] = [X_2, [X_2, X_3]] = [X_3, [X_3, X_1]] = [X_3, [X_3, X_2]] = [X_3, [X_1, X_2]] = 0$. Then Eq. (49) can be simplified as

$$\begin{aligned}
X_{\text{eff}} = X_{H,G} - \frac{h^2}{24} & \left([X_{A,C}, [X_{A,C}, X_{B,D}]] + [X_{A,C}, [X_{B,D}, X_{A,D}]] \right. \\
& \left. - 2[X_{B,D}, [X_{B,D}, X_{A,C}]] + [X_{B,D}, [X_{A,C}, X_{A,D}]] \right) + O(h^3),
\end{aligned} \tag{50}$$

where we used the original notations (37)-(39). In order for X_{eff} to be written in the form of Eq. (48), first of all, the second-order correction terms in Eq. (50) must be represented by a sum of Liouville operators. In the case of symplectic integrators for the Hamiltonian mechanics, we used the Jacobi identity to derive Eq. (13), whereas in the case of the structure preserving integrators for Nambu mechanics, we use the fundamental identity [2, 3]. Let us consider to represent $[X_{A,B}, [X_{C,D}, X_{E,F}]]$ by a sum of Liouville operators, where $A \sim F$ are any Hamiltonians. First, the commutator of $X_{C,D}$ and $X_{E,F}$ can be written as

$$\begin{aligned}
[X_{C,D}, X_{E,F}]f & = X_{C,D}\{f, E, F\} - X_{E,F}\{f, C, D\} \\
& = \{\{f, E, F\}, C, D\} - \{\{f, C, D\}, E, F\}.
\end{aligned} \tag{51}$$

If we use the fundamental identity for the first term in Eq. (51), $\{\{f, E, F\}, C, D\} = \{\{f, C, D\}, E, F\} + \{f, \{E, C, D\}, F\} + \{f, E, \{F, C, D\}\}$, then we obtain

$$[X_{C,D}, X_{E,F}]f = \left(X_{\{E,C,D\},F} + X_{E,\{F,C,D\}} \right) f. \tag{52}$$

On the other hand, if we use the fundamental identity for the second term in Eq. (51), $\{\{f, C, D\}, E, F\} = \{\{f, E, F\}, C, D\} + \{f, \{C, E, F\}, D\} + \{f, C, \{D, E, F\}\}$, then we obtain

$$[X_{C,D}, X_{E,F}]f = \left(-X_{\{C,E,F\},D} - X_{C,\{D,E,F\}} \right) f. \tag{53}$$

That is, the expression of $[X_{C,D}, X_{E,F}]$ is not unique and depends on which term in Eq. (51) the fundamental identity is used for. Here we choose the first term in Eq. (51) and proceed with the expression (52). Then we have

$$[X_{A,B}, [X_{C,D}, X_{E,F}]]f = [X_{A,B}, X_{\{E,C,D\},F}]f + [X_{A,B}, X_{E,\{F,C,D\}}]f, \tag{54}$$

where

$$[X_{A,B}, X_{\{E,C,D\},F}]f = \{\{f, \{E, C, D\}, F\}, A, B\} - \{\{f, A, B\}, \{E, C, D\}, F\}, \tag{55}$$

$$[X_{A,B}, X_{E,\{F,C,D\}}]f = \{\{f, E, \{F, C, D\}\}, A, B\} - \{\{f, A, B\}, E, \{F, C, D\}\}. \tag{56}$$

Here we should make two more choices on the use of the fundamental identity. If we use the fundamental identities for the first terms in Eq. (55) and Eq. (56), we get

$$\begin{aligned}
[X_{A,B}, [X_{C,D}, X_{E,F}]]f & = \left(X_{\{\{E,C,D\}A,B\},F} + X_{\{E,C,D\},\{F,A,B\}} \right. \\
& \left. + X_{\{E,A,B\},\{F,C,D\}} + X_{E,\{\{F,C,D\},A,B\}} \right) f.
\end{aligned} \tag{57}$$

There are a total of three two-choices for the use of the fundamental identity, and therefore the term $[X_{A,B}, [X_{C,D}, X_{E,F}]]f$ has $2^3 = 8$ types of expressions, including Eq. (57). Now let us evaluate the second order terms in Eq. (50) using eight types of expressions and the specific forms of the Hamiltonians (35)-(36). On the term $[X_{A,C}, [X_{A,C}, X_{B,D}]]$,

two of the eight expressions give $X_{B,-\frac{2}{m^2}x_2^2}$, and six give $X_{\frac{\omega^2}{m}x_2^2,C}$. The other three terms also have two different representations. The results can be summarized as follows,

$$[X_{A,C}, [X_{A,C}, X_{B,D}]] = X_{B,-\frac{2}{m^2}x_2^2} \quad \text{or} \quad X_{\frac{\omega^2}{m}x_2^2,C}, \quad (58)$$

$$[X_{A,C}, [X_{B,D}, X_{A,D}]] = X_{A, 2\omega^2x_1^2} \quad \text{or} \quad X_{-\frac{\omega^2}{m}x_2^2,D}, \quad (59)$$

$$[X_{B,D}, [X_{B,D}, X_{A,C}]] = X_{B,-2\omega^2x_1^2} \quad \text{or} \quad X_{m\omega^4x_1^2,C}, \quad (60)$$

$$[X_{B,D}, [X_{A,C}, X_{A,D}]] = X_{A, 2\omega^2x_1^2} \quad \text{or} \quad X_{-\frac{\omega^2}{m}x_2^2,D}. \quad (61)$$

There are various possible representations of X_{eff} . If we choose

$$[X_{A,C}, [X_{A,C}, X_{B,D}]] = 3X_{B,-\frac{2}{m^2}x_2^2} - 2X_{\frac{\omega^2}{m}x_2^2,C}, \quad (62)$$

$$[X_{A,C}, [X_{B,D}, X_{A,D}]] = X_{-\frac{\omega^2}{m}x_2^2,D}, \quad (63)$$

$$[X_{B,D}, [X_{B,D}, X_{A,C}]] = X_{m\omega^4x_1^2,C}, \quad (64)$$

$$[X_{B,D}, [X_{A,C}, X_{A,D}]] = X_{-\frac{\omega^2}{m}x_2^2,D}, \quad (65)$$

then Eq. (50) can be rewritten as follows,

$$X_{\text{eff}} = X_{H,G} + X_{H, \frac{1}{4m^2}h^2x_2^2} + X_{\frac{\omega^2}{12m}h^2x_2^2 + \frac{m\omega^4}{12}h^2x_1^2, G} + O(h^3), \quad (66)$$

where we used $X_{A,-\frac{2}{m^2}x_2^2} = 0$ and $X_{m\omega^4x_1^2,D} = 0$. Therefore, if we choose (62)-(65) the effective Liouville operator for the integrator $\Phi_{12321}(h)$ can be written as $X_{\text{eff}} = X_{H_S, G_S}$ with the shadow Hamiltonians:

$$H_S = H + \frac{\omega^2}{12m}h^2x_2^2 + \frac{m\omega^4}{12}h^2x_1^2 + O(h^3), \quad (67)$$

$$G_S = G + \frac{1}{4m^2}h^2x_2^2 + O(h^3). \quad (68)$$

Note that these shadow Hamiltonians are not unique. For example, if we choose

$$[X_{A,C}, [X_{A,C}, X_{B,D}]] = X_{B,-\frac{2}{m^2}x_2^2}, \quad (69)$$

$$[X_{A,C}, [X_{B,D}, X_{A,D}]] = X_{A, 2\omega^2x_1^2}, \quad (70)$$

$$[X_{B,D}, [X_{B,D}, X_{A,C}]] = X_{B,-2\omega^2x_1^2}, \quad (71)$$

$$[X_{B,D}, [X_{A,C}, X_{A,D}]] = X_{A, 2\omega^2x_1^2}, \quad (72)$$

and use $X_{A,-\frac{2}{m^2}x_2^2} = 0$, then we obtain other types of shadow Hamiltonians:

$$H_S = H + O(h^3), \quad (73)$$

$$G_S = G + \frac{1}{12m^2}h^2x_2^2 - \frac{\omega^2}{6}h^2x_1^2 + O(h^3). \quad (74)$$

We refer to the shadow Hamiltonians calculated using the BCH formula as the BCH shadow Hamiltonians. Although we could calculate the BCH Hamiltonians here, they have indefinite expressions.

C. Conserved quantities and exact shadow Hamiltonians

It can be shown by explicit calculation that the structure-preserving integrator Φ_h^{12321} conserves the following quantities,

$$H_c = H, \quad (75)$$

$$G_c = G + \frac{1}{4m^2}h^2x_2^2. \quad (76)$$

Although H_c and G_c are conserved, they are not the shadow Hamiltonians. That is, the Liouville operator associated with H_c and G_c ,

$$X_{H_c, G_c} = \frac{1}{m} \left(1 - \frac{\omega^2 h^2}{4} \right) x_2 \frac{\partial}{\partial x_1} - m\omega^2 x_1 \frac{\partial}{\partial x_2} + \frac{2}{m} x_1 x_2 \frac{\partial}{\partial x_3}, \quad (77)$$

is not the effective Liouville operator X_{eff} . Since $\Phi_h^{12321} = \Phi_h^{TVT}$ for the time evolutions of x_1 and x_2 as shown in Eq. (45), the x - and y -derivative terms in the effective Liouville operator X_{eff} for Φ_h^{12321} must agree with the effective Liouville operator (20) for Φ_h^{TVT} . From this consistency condition, we have

$$X_{\text{eff}} = F(\omega h)X_{H_c, G_c}, \quad (78)$$

where the factor $F(x)$ is same as Eq. (22). The exact shadow Hamiltonians H_S^e and G_S^e are given by identifying $X_{\text{eff}} = X_{H_S^e, G_S^e}$. However, depending on the way to distribute the factor $F(\omega h)$ to H_c and G_c , the exact shadow Hamiltonians take different forms. In general, we can define them with a real parameter α ,

$$H_S^e = F(\omega h)^\alpha H_c = H + \frac{\alpha\omega^2}{12m}h^2x_2^2 + \frac{\alpha m\omega^4}{12}h^2x_3 + O(h^3), \quad (79)$$

$$G_S^e = F(\omega h)^{1-\alpha}G_c = G + \frac{1}{4m^2}h^2x_2^2 + \frac{(1-\alpha)\omega^2}{6}h^2x_3 - \frac{(1-\alpha)\omega^2}{6}h^2x_1^2 + O(h^3). \quad (80)$$

On the other hand, for the BCH shadow Hamiltonians (50), we noted that there is arbitrariness due to the choices for the use of the fundamental identity. Here we represent the effective Liouville operator (50) as $X_{\text{eff}} = \alpha X_{\text{eff}} + (1-\alpha)X_{\text{eff}}$ with a real parameter α . Using Eqs. (62)-(65) for the first term αX_{eff} and using Eqs. (69)-(72) for the second term $(1-\alpha)X_{\text{eff}}$, the effective Liouville operator can be written as $X_{\text{eff}} = X_{H_S, G_S}$ with the shadow Hamiltonians,

$$H_S = H + \frac{\alpha\omega^2}{12m}h^2x_2^2 + \frac{\alpha m\omega^4}{12}h^2x_1^2 + O(h^3), \quad (81)$$

$$G_S = G + \frac{1}{12m^2}h^2x_2^2 + \frac{\alpha}{6m^2}h^2x_2^2 - \frac{(1-\alpha)\omega^2}{6}h^2x_1^2 + O(h^3). \quad (82)$$

If we replace x_3 in the third term of Eq. (79) using $G_S^e = \text{Const}$, replace x_3 in the third term of Eq. (80) using $H_S^e = \text{Const}$, and neglect the constant terms, then the BCH shadow Hamiltonians (81)-(82) are consistent with the exact shadow Hamiltonians (79)-(80) up to the second order of h .

D. Summary of the second order integrators

So far we have considered one of the second order structure-preserving integrators Φ_h^{12321} for the $N = 3$ harmonic oscillator described by the three variables (27). We have calculated the BCH shadow Hamiltonians, and found conserved quantities and the exact shadow Hamiltonians. Similar analyses can be performed for the other five integrators. In Table I we summarize the results of conserved quantities H_c and G_c for six integrators. Note that the conserved quantities are the same in Φ_h^{12321} and Φ_h^{13231} . This is because they give the same time evolution, $\Phi_h^{12321}\mathbf{x}(t) = \Phi_h^{13231}\mathbf{x}(t)$. For the same reason, the conserved quantities in the integrator Φ_h^{21312} are same as those in Φ_h^{23132} . The exact shadow Hamiltonians have indefinite expressions, and they can be given by $H_S^e = F(\omega h)^\alpha H_c$ and $G_S^e = F(\omega h)^{1-\alpha}G_c$ with a real parameter α .

TABLE I. Conserved quantities H_c and G_c of the second order integrators for the $N = 3$ harmonic oscillator described by variables $(x_1, x_2, x_3) = (q, p, q^2)$. The exact shadow Hamiltonians can be given by $H_S^e = F(\omega h)^\alpha H_c$ and $G_S^e = F(\omega h)^{1-\alpha}G_c$ with a real parameter α .

	H_c	G_c
Φ_h^{12321}	H	$G + \frac{1}{4m^2}h^2x_2^2$
Φ_h^{13231}	H	$G + \frac{1}{4m^2}h^2x_2^2$
Φ_h^{31213}	$H - \frac{\omega^2}{4m}h^2x_2^2$	$G - \frac{1}{4m^2}h^2x_2^2$
Φ_h^{21312}	$H - \frac{m\omega^4}{8}h^2x_1^2$	G
Φ_h^{23132}	$H - \frac{m\omega^4}{8}h^2x_1^2$	G
Φ_h^{32123}	$H + \frac{m\omega^4}{8}h^2x_1^2$	$G + \frac{\omega^2}{2}h^2x_1^2$

E. Numerical results

Finally we give a numerical demonstration of a second order integrator constructed in this work. We present here the results of one of the six integrators, Φ_h^{32123} . The original Hamiltonians H and G are given in Eqs. (28)-(29), the conserved quantities are $H_c = H + \frac{m\omega^4}{8}h^2x_1^2$ and $G_c = G + \frac{\omega^2}{2}h^2x_1^2$, and the exact shadow Hamiltonians can be written as $H_S^e = F(\omega h)^\alpha H_c$ and $G_S^e = F(\omega h)^{1-\alpha} G_c$ with a real parameter α . Here we set the parameters $m = \omega = 1$ and the small time step $h = 0.1$. We evolve the harmonic oscillator in time by n steps using the integrator Φ_h^{32123} with initial conditions $(x_1(0), x_2(0), x_3(0)) = (1, 1, 1)$, and obtain numerical solutions $\mathbf{x}^{32123}(t) = (\Phi_h^{32123})^n \mathbf{x}(0)$, where $t = nh$. Then using the solutions $\mathbf{x}^{32123}(t)$ we evaluate the original Hamiltonians $H(t)$ and $G(t)$ and the conserved quantities $H_c(t)$ and $G_c(t)$. Figure 1 shows the error in the original Hamiltonian, $H(t) - H(0)$, and the error in the conserved quantity, $H_c(t) - H_c(0)$. Figure 2 also shows the errors in the original Hamiltonian, $G(t) - G(0)$, and in the conserved quantity, $G_c(t) - G_c(0)$. It can be seen that neither $H(t)$ nor $G(t)$ is conserved; however, either of them oscillates with no increase in error. That is, the integrator Φ_h^{32123} gives stable time evolution and can be used for long-time simulations.




FIG. 1. Time evolutions by Φ_h^{32123} : the error in the original Hamiltonian $H(t) - H(0)$ (solid line) and the error in the conserved quantity $H_c(t) - H_c(0)$ (dashed line).




FIG. 2. Time evolutions by Φ_h^{32123} : the error in the original Hamiltonian $G(t) - G(0)$ (solid line) and the error in the conserved quantity $G_c(t) - G_c(0)$ (dashed line).

It can also be seen from Figs. 1-2 that H_c and G_c are both conserved. However, note that they are not the shadow Hamiltonians, but just conserved quantities. To see this point clearly, we calculate the exact solutions of the Nambu equations with original Hamiltonians, the conserved quantities, and the exact shadow Hamiltonians, and compare those three solutions with the numerical solutions computed by the integrator Φ_h^{32123} . The Nambu equations with the original Hamiltonians H and G are given by Eqs. (30). On the other hand, the Nambu equations with the conserved quantities H_c and G_c as the Hamiltonians are written as

$$\frac{d}{dt}x_1 = \frac{1}{m}x_2, \quad \frac{d}{dt}x_2 = -m\omega^2 a^2 x_1, \quad \frac{d}{dt}x_3 = \frac{2}{m}b^2 x_1 x_2, \quad (83)$$

where $a = \sqrt{1 - \frac{\omega^2 h^2}{4}}$ and $b = \sqrt{1 - \frac{\omega^2 h^2}{2}}$. Then, the Nambu equations with the exact shadow Hamiltonians H_S^e and G_S^e are given by

$$\frac{d}{dt}x_1 = \frac{1}{m}F(\omega h)x_2, \quad \frac{d}{dt}x_2 = -m\omega^2 a^2 F(\omega h)x_1, \quad \frac{d}{dt}x_3 = \frac{2}{m}b^2 F(\omega h)x_1 x_2. \quad (84)$$

We denote the solutions of Eqs. (30) by $\mathbf{x}^o(t)$, the solutions of Eqs. (83) by $\mathbf{x}^c(t)$, and the solutions of Eqs. (84)




FIG. 3. The difference between the two dynamics: $x_3^o(t) - x_3^{32123}(t)$ (solid line). Here data are plotted every 100 steps.




FIG. 4. The differences between the two dynamics: $x_3^c(t) - x_3^{32123}(t)$ (solid line) and $x_3^s(t) - x_3^{32123}(t)$ (dashed line).

by $\mathbf{x}^s(t)$. Here we give only the third components of the solutions,

$$x_3^o(t) = \left(\frac{1}{2}x_1(0)^2 - \frac{1}{2m^2\omega^2}x_2(0)^2 \right) \cos 2\omega t + \frac{1}{m\omega}x_1(0)x_2(0) \sin 2\omega t + x_3(0) - \frac{1}{2}x_1(0)^2 + \frac{1}{2m^2\omega^2}x_2(0)^2, \quad (85)$$

$$x_3^c(t) = b^2 \left[\left(\frac{1}{2}x_1(0)^2 - \frac{1}{2m^2\tilde{\omega}^2}x_2(0)^2 \right) \cos 2\tilde{\omega}t + \frac{1}{m\tilde{\omega}}x_1(0)x_2(0) \sin 2\tilde{\omega}t + \frac{1}{b^2}x_3(0) - \frac{1}{2}x_1(0)^2 + \frac{1}{2m^2\tilde{\omega}^2}x_2(0)^2 \right], \quad (86)$$

$$x_3^s(t) = b^2 \left[\left(\frac{1}{2}x_1(0)^2 - \frac{1}{2m^2\tilde{\omega}^2}x_2(0)^2 \right) \cos 2\tilde{\omega}t + \frac{1}{m\tilde{\omega}}x_1(0)x_2(0) \sin 2\tilde{\omega}t + \frac{1}{b^2}x_3(0) - \frac{1}{2}x_1(0)^2 + \frac{1}{2m^2\tilde{\omega}^2}x_2(0)^2 \right], \quad (87)$$

where $\tilde{\omega} = a\omega$ and $\tilde{t} = F(\omega h)t$. Figure 3 shows the difference between the exact solution of the Nambu mechanics with the original Hamiltonians (85) and the numerical solution given by the integrator Φ_h^{32123} , $x_3^o(t) - x_3^{32123}(t)$. Although both of $x_3^o(t)$ and $x_3^{32123}(t)$ are periodic, the frequencies are slightly different and we can observe a beat over a very long time domain. Figure 4 show two more differences, $x_3^c(t) - x_3^{32123}(t)$ and $x_3^s(t) - x_3^{32123}(t)$. Since a beat is also observed in the difference $x_3^o(t) - x_3^{32123}(t)$, the conserved quantities H_c and G_c are not the shadow Hamiltonians to reproduce the time evolution given by the integrator Φ_h^{32123} . On the other hand, it can be seen that $x_3^s(t) - x_3^{32123}(t)$ remains zero all the time, indicating that H_s^e and G_s^e work as the shadow Hamiltonians of this integrator.

V. CONCLUSIONS AND FUTURE WORKS

We have constructed the structure-preserving integrators for an $N = 3$ Nambu system, the harmonic oscillator described by three variables $(x_1, x_2, x_3) = (q, p, q^2)$. We have shown that the shadow Hamiltonians of the constructed integrators can be calculated by using the BCH formula and the fundamental identity, and also the exact expressions of the shadow Hamiltonians can be found in terms of the consistency with the exact shadow Hamiltonian of the symplectic integrator. This is the first work to derive the explicit forms of BCH shadow Hamiltonians and exact shadow Hamiltonians in a Nambu system. However, we have also found that they both have indefinite expressions. The BCH shadow Hamiltonians are indefinite due to the freedom in the choices for the use of the fundamental identity, while the exact shadow Hamiltonians are indefinite due to the way of distribution of the factor $F(\omega h)$. Although they are both indefinite, we have confirmed that they are consistent with each other up to the order \hbar^2 .

In the present paper, we focused on second order structure-preserving integrators for a simple Nambu system given by the hidden Nambu formalism [6, 7]. It would be interesting to study other Nambu systems and other types of integrators. For example, in the Hamiltonian mechanics, it has been an important problem to construct symplectic integrators in non-separable Hamiltonian systems (see, e.g., [17]). In the Nambu mechanics, we often see non-separable Hamiltonian(s) in multiple Hamiltonians: for example, the $N = 3$ harmonic oscillator with three composite variables, $(x_1, x_2, x_3) = (q^2, p^2, qp)$, is driven by the Hamiltonians $H = \frac{1}{2m}x_2 + \frac{m\omega^2}{2}x_1$ and $G = 2x_3^2 - 2x_1x_2$ [7]; the latter

Hamiltonian is non-separable. It would be a challenging work to construct structure-preserving integrators for such Nambu systems with non-separable Hamiltonian(s).

We have seen that the fundamental identity plays an important role in calculating the shadow Hamiltonians using the BCH formula. However, it is well known that the fundamental identity does not hold in many degrees of freedom systems [3, 8, 16]. Therefore it would be interesting to see if shadow Hamiltonians exist in many degrees of freedom Nambu systems.

The integrators we have constructed in the present paper are those with time-reversal symmetry. It would be important to consider the role of time-reversal symmetry in structure-preserving integrators for the Nambu mechanics [15].

Finally, it is known that the Nambu mechanics can always be represented in the form of the noncanonical Hamiltonian mechanics with the noncanonical Poisson bracket defined by the Nambu bracket [3, 8, 18]. Therefore it would also be interesting to study the relationship between the structure-preserving integrators for Nambu mechanics and the Lie-Poisson integrators for noncanonical Hamiltonian mechanics [10, 19].

ACKNOWLEDGMENTS

This work was partly supported by a scientific grant from the Ministry of Education, Culture, Sports, Science and Technology under Grant No. 26610134.

-
- [1] Y. Nambu, Phys. Rev. D **7**, 2405 (1973).
 - [2] D. Sahoo and M. C. Valsakumar, Phys. Rev. A **46**, 4410 (1992).
 - [3] L. Takhtajan, Commun. Math. Phys. **160**, 295 (1994).
 - [4] P. N evier and R. Blender, J. Phys. A: Math. Gen. **26**, L1189 (1993).
 - [5] Y. Suzuki, Prog. Theor. Exp. Phys. **2022**, ptac094 (2022).
 - [6] A. Horikoshi and Y. Kawamura, Prog. Theor. Exp. Phys. **2013**, 073A01 (2013) [arXiv:1304.2086 [math-ph]].
 - [7] A. Horikoshi, Prog. Theor. Exp. Phys. **2019**, 123A02 (2019) [arXiv:1908.03567 [quant-ph]].
 - [8] A. Horikoshi, Prog. Theor. Exp. Phys. **2021**, 12C106 (2021) [arXiv:2103.00688 [math-ph]].
 - [9] C. Chandre and A. Horikoshi, J. Math. Phys. **64**, 052702 (2023) [arXiv:2109.13663 [math.DS]].
 - [10] E. Hairer, C. Lubich, G. Wanner, Geometric Numerical Integration: Structure-Preserving Algorithms for Ordinary Differential Equations (Springer, 2010).
 - [11] D. Frenkel and B. Smit, Understanding Molecular Simulation: From Algorithms to Applications (Academic Press, 2023).
 - [12] D. J. Hardy and D. I. Okunbor, J. Chem. Phys. **102**, 8978 (1995).
 - [13] H. Kobayashi, Phys. Lett. A **371**, 360 (2007) [arXiv:0705.4523 [math-ph]].
 - [14] pyHamSys, <https://pypi.org/project/pyhamsys/>
 - [15] K. Modin, J. Gen. Lie Theory Appl. **3**, 39 (2009).
 - [16] P.-M. Ho and Y. Matsuo, Prog. Theor. Exp. Phys. **2016**, 06A104 (2016) [arXiv:1603.09534 [hep-th]].
 - [17] M. Tao, Phys. Rev. E **94**, 043303 (2016) [arXiv:1609.02212 [math.NA]].
 - [18] I. Bialynicki-Birula and P. J. Morrison, Phys. Lett. A **158**, 453 (1991).
 - [19] R. I. McLachlan, Phys. Rev. Lett. **71**, 3043 (1993).