Semidefinite programming on population clustering: a local analysis

Shuheng Zhou

University of California, Riverside, CA 92521

Abstract

In this paper, we consider the problem of partitioning a small data sample of size n drawn from a mixture of 2 sub-gaussian distributions. In particular, we design and analyze two computational efficient algorithms to partition data into two groups approximately according to their population of origin given a small sample in a recent paper (Zhou 2023a). Our work is motivated by the application of clustering individuals according to their population of origin using markers, when the divergence between any two of the populations is small. Moreover, we are interested in the case that individual features are of low average quality γ , and we want to use as few of them as possible to correctly partition the sample. Here we use $p\gamma$ to denote the ℓ_2^2 distance between two population centers (mean vectors), namely, $\mu^{(1)}, \mu^{(2)} \in \mathbb{R}^p$. We allow a full range of tradeoffs between n, p, γ in the sense that partial recovery (success rate < 100%) is feasible once the signal to noise ratio $s^2 := \min\{np\gamma^2, p\gamma\}$ is lower bounded by a constant. Our work builds upon the semidefinite relaxation of an integer quadratic program that is formulated essentially as finding the maximum cut on a graph, where edge weights in the cut represent dissimilarity scores between two nodes based on their p features in Zhou (2023a). More importantly, we prove that the misclassification error decays exponentially with respect to the SNR s^2 in the present paper. The significance of such an exponentially decaying error bound is: when $s^2 = \Omega(\log n)$, perfect recovery of the cluster structure is accomplished. This result was introduced in Zhou (2023a) without a proof. We therefore present the full proof in the present work.

1 Introduction

We explore a type of classification problem that arises in the context of computational biology. The biological context for this problem is we are given DNA information from n individuals from k populations of origin and we wish to classify each individual into the correct category. DNA contains a series of markers called SNPs (Single Nucleotide Polymorphisms), each of which has two variants (alleles). We use bit 1 and bit 0 to denote them. The problem is that we are given a small sample of size n, e.g., DNA of n individuals (think of n in the hundreds or thousands), each described by the values of p features or markers, e.g., SNPs (think of p as an order of magnitude larger than n). Our goal is to use these features to classify the individuals according to their population of origin. Given the population of origin of an individual, the genotypes can be reasonably assumed to be generated by drawing alleles independently from the appropriate distribution.

Suppose we are given a data matrix $X \in \mathbb{R}^{n \times p}$ with samples from two populations $\mathcal{C}_1, \mathcal{C}_2$, such that

$$\forall i \in \mathcal{C}_g, \ \mathbb{E}(X_{ij}) = \mu_j^{(g)} \quad g = 1, 2, \forall j = 1, \dots, p.$$

$$\tag{1}$$

Our goal is to estimate the group membership vector $u_2 \in \{-1, 1\}^n$ such that

$$u_{2,j} = 1$$
 for $j \in \mathcal{C}_1$ and $u_{2,j} = -1$ for $j \in \mathcal{C}_2$, (2)

where the sizes of clusters $|\mathcal{C}_{\ell}| =: n_{\ell}, \forall \ell$ may not be the same. Our ultimate goal is to estimate the solution to the discrete optimization problem:

maximize
$$x^T R x$$
 subject to $x \in \{-1, 1\}^n$, (3)

where R is a static reference matrix to be specified. It was previously shown that, in expectation, among all balanced cuts in the complete graph formed among n vertices (sample points), the cut of maximum weight corresponds to the correct partition of the n points according to their distributions in the balanced case ($n_1 = n_2 = n/2$). Here the weight of a cut is the sum of weights across all edges in the cut, and the edge weight equals the Hamming distance between the bit vectors of the two endpoints; See [10, 38]. Under suitable conditions, the statement above also holds with high probability (w.h.p.); The analyses in [10] and [38] focused on the high dimensional setting, where $p \gg n$.

Features have slightly different probabilities depending on which population the individual belongs to. Denote by Δ^2 the ℓ_2^2 distance between two population centers (mean vectors), namely, $\mu^{(1)}, \mu^{(2)} \in \mathbb{R}^p$. We focus on the case where p > n, although it is not needed. Note that Δ measures the Euclidean distance between $\mu^{(1)}$ and $\mu^{(2)}$ and thus represents their separation. The objective we consider is to minimize the total data size D = np needed to correctly classify the individuals in the sample as a function of the "average quality" γ of the features:

$$\gamma := \Delta^2/p, \text{ where } \Delta^2 := \sum_{k=1}^p (\mu_k^{(1)} - \mu_k^{(2)})^2 \text{ and } \mu^{(i)} = (\mu_1^{(i)}, \dots, \mu_p^{(i)}) \in \mathbb{R}^p, i = 1, 2.$$
 (4)

In other words, in the context of population clustering, it has been previously shown one can use a random instance of the integer quadratic program:

(Q) maximize
$$x^T A x$$
 subject to $x \in \{-1, 1\}^n$ (5)

to identify the correct partition of nodes according to their population of origin w.h.p. so long as the data size D is sufficiently large and the separation metric is at the order of $\Delta^2 = \Omega(\log n)$. Here $A = (a_{ij})$ is an $n \times n$ symmetric matrix, where for $1 \leq i, j \leq n, a_{ij} = a_{ji}$ denotes the edge weight between nodes i and j, computed from the individuals' bit vectors. This result is structural, rather than algorithmic.

The integer quadratic program (3) (or (5)) is NP-hard [24]. In a groundbreaking paper [19], Goemans and Williamson show that one can use semidefinite program (SDP) as relaxation to solve these approximately. See references therein for earlier works. In the present work, we use semidefinite relaxation of the graph cut problem (5), which was originally formulated in [10, 38] in the context of population clustering. More generally, one may consider semidefinite relaxations for the following sub-gaussian mixture model with k centers (implicitly, with rank-k mean matrix embedded), where we have n observations $X_1, X_2 \dots, X_n \in \mathbb{R}^p$:

$$X_i = \mu^{(\psi_i)} + \mathbb{Z}_i,\tag{6}$$

where $\mathbb{Z}_1, \ldots, \mathbb{Z}_n \in \mathbb{R}^p$ are independent, sub-gaussian, mean-zero, random vectors and $\psi_i : i \to \{1, \ldots, k\}$ assigns node *i* to a group \mathcal{C}_j with the mean $\mu^{(j)} \in \mathbb{R}^p$ for some $j \in [k]$. Here we denote by [k] the set of integers $\{1, \ldots, k\}$. Hence, each row vector X_i of data matrix X is a *p*-dimensional sub-gaussian random vector and we assume rows are independent.

The proposed semidefinite relaxation framework in [39] was inspired by [20]. The important distinction of the present work from that as considered in [20] is: to estimate the group membership vector u_2 , we replace the random adjacency matrix stochastic block models with an instance of symmetric matrix computed from the centered data matrix, which we now introduce. First, we define the global center of the data. Let X be as in (1) with row vectors $X_1, \ldots, X_n \in \mathbb{R}^p$.

Definition 1.1. (The global center) Denote by $\mathbb{Z}_i = X_i - \mathbb{E}(X_i)$. Then the global center is the average over n row vectors $X_1, \ldots, X_n \in \mathbb{R}^p$, for X as in (1),

$$\widehat{\mu}_n = \frac{1}{n} \sum_{i=1}^n X_i, \quad and \quad \widehat{\mu}_n - \mathbb{E}\widehat{\mu}_n := \frac{1}{n} \sum_{i=1}^n \mathbb{Z}_i = \frac{1}{n} \sum_{i=1}^n X_i - \mathbb{E}(X_i).$$
(7)

Estimators and convex relaxation. Let $\mathbf{1}_n = [1, \ldots, 1] \in \mathbb{R}^n$ denote a vector of all 1s and $E_n := \mathbf{1}_n \mathbf{1}_n^T$. As in many statistical problems, one simple but crucial step is to first obtain the centered data Y. In the present work, to find a convex relaxation, we will first construct a matrix Y by subtracting the sample mean $\hat{\mu}_n \in \mathbb{R}^p$ as computed from (7) from each row vector X_i of the data matrix X. Denote by

$$Y = X - P_1 X, \quad \text{where} \quad P_1 := \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^T = E_n / n \tag{8}$$

is a projection matrix. Loosely speaking, this procedure is called "global centering" in the statistical literature, for example, see [21]. Given matrix Y, we construct:

$$A := YY^T - \lambda(E_n - I_n), \text{ where } \lambda = \frac{2}{n(n-1)} \sum_{i < j} \langle Y_i, Y_j \rangle$$
(9)

and I_n denotes the identity matrix. To estimate the group membership vector $u_2 \in \{-1, 1\}^n$, we consider the following semidefinite optimization problem from [39]:

(SDP) maximize
$$\langle A, Z \rangle$$
 subject to $Z \succeq 0, I_n \succeq \operatorname{diag}(Z)$. (10)

Here $Z \succeq 0$ indicates that the matrix Z is constrained to be positive semidefinite, $A \succeq B$ means that $A - B \succeq 0$, and the inner product of matrices $\langle A, B \rangle = tr(A^T B)$. Here and in the sequel, denote by

$$\mathcal{M}_G^+ := \{ Z : Z \succeq 0, I_n \succeq \operatorname{diag}(Z) \} \subset [-1, 1]^{n \times n}$$

the set of positive semidefinite matrices whose entries are bounded by 1 in absolute value. Since all possible solutions of (Q) are feasible for SDP, the optimal value of SDP is an upper bound on the optimal value of (Q). Moreover, SDP (10) is shown to be equivalent to the optimization problem

(SDP1) maximize
$$\langle YY^T - \lambda E_n, Z \rangle$$
 subject to $Z \succeq 0, \operatorname{diag}(Z) = I_n.$ (11)

See the supplementary Proposition D.1. The (global) analysis framework for the semidefinite relaxation in [20] was set in the context of community detection in sparse networks, where A

represents the symmetric adjacency matrix of a random graph. In other words, they study the semidefinite relaxation of the integer program (5), where an $n \times n$ random matrix A (observed) is used to replace the hidden static R in the original problem (3) such that $\mathbb{E}(A) = R$. The innovative proof strategy of [20] is to apply the Grothendieck's inequality for the random error $A - \mathbb{E}(A)$ rather than the original matrix A as considered in the earlier literature. We call this approach the global analysis in [39], following [12]. Throughout this paper, denote by $w_{\min} := \min_{j=1,2} w_j$, where $w_j = n_j/n$. We use $n_{\min} := nw_{\min}$ and $n_{\max} := nw_{\max}$ to represent the size of the smallest and the largest cluster respectively. For a symmetric matrix M, let $\lambda_{\max}(M)$ and $\lambda_{\min}(M)$ be the largest and the smallest eigenvalue of M respectively. The operator norm $||M||_2$ is defined to be $\sqrt{\lambda_{\max}(M^T M)}$.

1.1 Our approach and contributions

Our approach of using the centered data matrix Y in SDP1 is novel to the best of our knowledge. Following [39], we use YY^T and the corresponding A as the input to our optimization algorithms, ensuring both computational efficiency and statistical convergence, especially in the low signal-tonoise ratio (SNR) case when $s^2 \approx (p\gamma) \wedge (np\gamma^2) = o(\log n)$. The concentration bounds on the operator norm $||YY^T - \mathbb{E}YY^T||_2$ imply that, up to a constant factor, the same bounds also hold for $||A - \mathbb{E}A||_2$, in view of Lemma 5.4 and (35). This is a desirable property of the SDP (10) for its global analysis. Denote by C_0 the ψ_2 -constant of $\mathbb{Z}_i \in \mathbb{R}^p$, cf. (19). When the independent sub-gaussian random vectors \mathbb{Z}_i in (6) are isotropic, we use the following notion of signal-to-noise ratio:

SNR isotropic
$$s^2 = (p\gamma/C_0^2) \wedge (np\gamma^2/C_0^4).$$
 (12)

This quantity appears in [18] explicitly and implicitly in [10, 38], and is compatible with the total sample size (np) lower bound in (23), to ensure partial recovery of the clusters when $s^2 = o(\log n)$.

In the present work: (a) we will present a transparent and unified global and local analysis framework for the semidefinite programming optimization problem (10); and (b) we prove that the error decays exponentially with respect to the signal-to-noise ratio (SNR) in Theorem 2.3, even for the sub-gaussian design with dependent features so long as the lower bounds in (30) hold. In particular, the implication of such an exponentially decaying error bound is: when $s^2 = \Omega(\log n)$, perfect recovery of the cluster structure is accomplished. See [35, 15, 18] and references therein. Although the result in Theorem 2.3 is in the same spirit as that in [18], we prove these error bounds for the SDP (10), which is motivated by the graph partition problem (5), while they establish such error bounds for the semidefinite relaxation based on the k-means criterion (40) directly following [31]; See Section 3, cf. (43) and (47).

In other words, although our general result in Theorem 2.3 coincides with that of [18] for k = 2, we prove it for the much simpler SDP1 (and SDP) with the only set of constraints being $\{Z_{ii} = 1, \forall i \in [n]\}$ besides Z being positive semidefinite. This improvement may be of theoretical interest. In particular, unlike [18], we do not enforce entries of Z in SDP and SDP1 to be nonnegative, which enables faster computation. Moreover, we do not need an explicit de-biasing step so long as assumption (A2) holds, showing that SDP has an inherent tolerance on the variance discrepancy between the two component distributions, at least up to a certain threshold as specified in (33). Finally, through the refined local analysis in the present work, we gained new insights that for the balanced case $(n_1 = n_2 = n/2)$, the bias may have been substantially reduced due to our centering and adjustment steps in the construction of A as in (9) and an OracleSDP, cf. (16), and hence we expect that assumption (A2) can be substantially relaxed for balanced partitions.

Construction of a reference matrix. Let Y be a centered matrix with row vectors Y_1, \ldots, Y_n as in (8). The SDP1 estimator as in (11) crucially exploits the geometric properties of the two mean vectors in matrix Y, resulting in a natural choice of R. By linearity of expectation,

$$\mathbb{E}\widehat{\mu}_{n} := w_{1}\mu^{(1)} + w_{2}\mu^{(2)}, \quad \text{where} \quad w_{i} = |\mathcal{C}_{i}| / n, i = 1, 2, \\
\mathbb{E}(Y_{i}) := \mathbb{E}X_{i} - \mathbb{E}\widehat{\mu}_{n} = \begin{cases} w_{2}(\mu^{(1)} - \mu^{(2)}) & \text{if } i \in \mathcal{C}_{1}; \\ w_{1}(\mu^{(2)} - \mu^{(1)}) & \text{if } i \in \mathcal{C}_{2}. \end{cases}$$
(13)

Definition 1.2. (The reference matrix) W.l.o.g., we assume that the first n_1 rows of X belong to C_1 and the rest belong to C_2 . Denote by $n_1 = w_1 n$ and $n_2 = w_2 n$. For Y as defined in (8), we construct

$$R := \mathbb{E}(Y)\mathbb{E}(Y)^T = p\gamma \begin{bmatrix} w_2^2 E_{n_1} & -w_1 w_2 E_{n_1 \times n_2} \\ -w_1 w_2 E_{n_2 \times n_1} & w_1^2 E_{n_2} \end{bmatrix}.$$
 (14)

Recall the overall goal of convex relaxation is to: (a) estimate the solution of the integer quadratic problem (3) with an appropriately chosen reference matrix R such that solving the integer quadratic problem (3) will recover the cluster *exactly*; and (b) choose the convex set \mathcal{M}_G^+ (resp. \mathcal{M}_{opt} , cf. (15)) so that the semidefinite relaxation of the static problem (3) is tight. This means that when we replace A (resp. $\tilde{A} := YY^T - \lambda E_n$) with R in SDP (10) (resp. SDP1 (11)) as done in Lemma 1.3, we obtain a solution Z^* which can then be used to recover the clusters exactly. Here, $Z^* = u_2 u_2^T$ will maximize the reference objective function $\langle R, Z \rangle$ among all $Z \in [-1, 1]^{n \times n}$, and naturally among all $Z \in \mathcal{M}_{opt} \subset [-1, 1]^{n \times n}$, given that $Z^* \in \mathcal{M}_{opt}$.

Lemma 1.3. [39] Let $\mathcal{M}_{opt} \subseteq \mathcal{M}_G^+ \subset [-1,1]^{n \times n}$ be as defined in (15):

$$\mathcal{M}_{opt} = \{ Z : Z \succeq 0, \operatorname{diag}(Z) = I_n \} \subset \mathcal{M}_G^+.$$
(15)

Then for R as in Definition 1.2 and $u_2 \in \{-1, 1\}^n$ as in (2),

$$Z^* = \underset{Z \in \mathcal{M}_{opt}}{\operatorname{arg\,max}} \langle R, Z \rangle = u_2 u_2^T.$$

Construction of an OracleSDP. A straightforward calculation leads to the expression of the reference matrix $R := \mathbb{E}(Y)\mathbb{E}(Y)^T$ as in Definition 1.2. However, unlike the settings of [20], $\mathbb{E}A \neq R$, resulting in a large bias. It is crucial to bridge the gap between YY^T and the reference matrix $R = \mathbb{E}(Y)\mathbb{E}(Y)^T$. A remedy was proposed in [39] to transform (10) into an equivalent **OracleSDP** formulation:

(**OracleSDP**) maximize
$$\langle B, Z \rangle$$
 subject to $Z \in \mathcal{M}_{opt}$ where (16)

$$B := A - \mathbb{E}\tau I_n \quad \text{for} \quad \tau = \frac{1}{n} \sum_{i=1}^n \langle Y_i, Y_i \rangle , \qquad (17)$$

where the adjustment term $\mathbb{E}\tau I_n$ plays no role in optimization, since the extra trace term $\propto \langle I_n, Z \rangle = \operatorname{tr}(Z)$ is a constant function of Z on the feasible set \mathcal{M}_{opt} . In other words, optimizing the original SDP (10) over the larger constraint set \mathcal{M}_G^+ is equivalent to maximizing $\langle B, Z \rangle$ over $Z \in \mathcal{M}_{opt}$ as in (16). We emphasize that our algorithm solves SDP1 (11) rather than the oracle SDP (16). However, formulating the OracleSDP helps us with the global and local analyses, in controlling the operator norm of the bias term, namely, $\|\mathbb{E}B - R\|_2$ and a related quantity given in Lemma 4.2.

A unified framework for the local and global analyses. Given this OracleSDP formulation, the bias analysis on $\mathbb{E}B - R$ as in Lemma 2.6, and the concentration of measure bounds on $YY^T - \mathbb{E}YY^T$ have already enabled simultaneous analyses of the SDP and the closely related spectral clustering method in [39]; cf. Section 3. In the local analysis in the present paper, cf. Lemma 4.3, we will obtain a high probability uniform control over $|\langle YY^T - \mathbb{E}(YY)^T, \hat{Z} - Z^* \rangle|$ in a local neighborhood of Z^* ,

$$\sup_{\widehat{Z}\in\mathcal{M}_{opt}\cap(Z^*+r_1B_1^{n\times n})} \left| \langle YY^T - \mathbb{E}(YY^T), \widehat{Z} - Z^* \rangle \right| \le f(r_1) \quad (W_1), \tag{18}$$

where

$$\left\|\widehat{Z} - Z^*\right\|_1 := \sum_{i,j} \left|\widehat{Z}_{ij} - Z^*_{ij}\right| \le r_1, \quad \text{ for } \widehat{Z}, Z^* \in \mathcal{M}_{\text{opt}},$$

and $f(r_1)$ is a function that depends on the ℓ_1 radius $r_1, r_1 \leq 2n(n-1)$. Lemma 4.3 is one of the main technical contributions in this paper, which in turn depends on the geometry of the constraint set \mathcal{M}_{opt} and the sharp concentration of measure bounds on $YY^T - \mathbb{E}YY^T$ in Theorem 2.7. This approach also gives rise to the notion of a *local analysis*, following [12]. Similar to the global analysis, the performance of SDP also crucially depends on controlling the bias term $\mathbb{E}B - R$; however, we now control $\langle \mathbb{E}B - R, \widehat{Z} - Z^* \rangle$ uniformly over all $\widehat{Z} \in \mathcal{M}_{opt}$ in Lemma 4.2, rather than the operator norm $\|\mathbb{E}B - R\|_2$. The bias and variance tradeoffs are described in the sequel. Combining these results leads to an exponentially decaying error bound with respect to the SNR s^2 to be presented in Theorem 2.3.

We need to introduce some notation. Recall for a random variable X, the ψ_2 -norm of X, denoted by $||X||_{\psi_2}$, is $||X||_{\psi_2} = \inf\{t > 0 : \mathbb{E}\exp(X^2/t^2) \leq 2\}$. We use $\mathbb{Z} = (z_{ij})$ to denote the meanzero random matrix with independent, mean-zero, sub-gaussian row vectors $\mathbb{Z}_1, \ldots, \mathbb{Z}_n \in \mathbb{R}^p$ as considered in (6), where for a constant C_0 and $\operatorname{Cov}(\mathbb{Z}_j) := \mathbb{E}(\mathbb{Z}_j \mathbb{Z}_j^T)$,

$$\forall j = 1, \dots, n, \| \langle \mathbb{Z}_j, x \rangle \|_{\psi_2} \leq C_0 \| \langle \mathbb{Z}_j, x \rangle \|_{L_2} \text{ for any } x \in \mathbb{R}^p,$$
(19)

where
$$\| \langle \mathbb{Z}_j, x \rangle \|_{L_2}^2 := x^T \mathbb{E}(\mathbb{Z}_j \mathbb{Z}_j^T) x = x^T \mathsf{Cov}(\mathbb{Z}_j) x.$$
 (20)

Throughout this paper, we use V_i to denote the trace of covariance $Cov(\mathbb{Z}_i)$, for all $j \in C_i$:

$$V_1 = \mathbb{E} \langle \mathbb{Z}_j, \mathbb{Z}_j \rangle \quad \forall j \in \mathcal{C}_1 \text{ and } V_2 = \mathbb{E} \langle \mathbb{Z}_j, \mathbb{Z}_j \rangle \quad \forall j \in \mathcal{C}_2.$$

$$(21)$$

(A1) Let $\mathbb{Z} = X - \mathbb{E}X$. Let $\mathbb{Z}_i = X_i - \mathbb{E}X_i, i \in [n]$ be independent, mean-zero, sub-gaussian random vectors with independent coordinates such that for all $i, j, ||X_{ij} - \mathbb{E}X_{ij}||_{\psi_2} \leq C_0$. (A2) The two distributions have bounded discrepancy in their variance profiles:

$$|V_1 - V_2| \leq \xi n p \gamma/3$$
 for some $w_{\min}^2/8 > 2\xi = \Omega(1/n_{\min}).$ (22)

Although (A1) assumes that the random matrix \mathbb{Z} has independent sub-gaussian entries, matching the separation (and SNR) condition (23), we emphasize that the conclusions as stated in Theorems 1.4 and 2.3 hold for the general two-group model as considered in Lemma 2.5 so long as (A2) holds, upon adjusting the lower bounds in (23). In particular, we allow each population to have distinct covariance structures, with diagonal matrices as special cases.

1.2 Prior results

For completeness, we first state in Theorem 1.4 the main result under assumptions (A1) and (A2), using the global analysis [39, 20], in order to set the context for local analysis in this paper. For a matrix $B = (b_{ij})$ of size $n \times n$, let vec {B} be formed by concatenating columns of matrix B into a vector of size n^2 ; we use $||B||_1 = \sum_{i,j=1}^n |b_{ij}|$ to denote the ℓ_1 norm of vec {B} and $||B||_F = (\sum_{i,j} b_{ij}^2)^{1/2}$ to denote the ℓ_2 norm of vec {B}.

Theorem 1.4. [39] Let $C_j \subset [n]$ denote the group membership, with $|C_j| = n_j$ and $\sum_j n_j = n$. Suppose that for $j \in C_i$, $\mathbb{E}X_j = \mu^{(i)}$, where i = 1, 2. Let \widehat{Z} be a solution of SDP1 (and SDP). Suppose that (A1) and (A2) hold, and for some absolute constants C, C',

$$p\gamma = \Delta^2 \ge \frac{C'C_0^2}{\xi^2}$$
 and $pn \ge \frac{CC_0^4}{\xi^2\gamma^2}$, where ξ is the same as in (22). (23)

Then with probability at least $1 - 2\exp(-cn)$, we have for u_2 as in (2),

$$\left\|\widehat{Z} - u_2 u_2^T\right\|_1 / n^2 = \delta \le 2K_G \xi / w_{\min}^2 \text{ and } \left\|\widehat{Z} - u_2 u_2^T\right\|_F / n^2 \le 4K_G \xi / w_{\min}^2.$$
(24)

The parameter ξ in (A2) is understood to be set as a fraction of the parameter δ appearing in Theorem 1.4, which in turn is assumed to be lower bounded: $1 > \delta = \Omega(1/n)$, since we focus on the partial recovery of the group membership vector u_2 in the present work. In particular, ξ^2 can be chosen conservatively to be inversely proportional to s^2 ; cf. Section 2. This choice enables the bias and variance tradeoffs as elaborated in Section 2.

1.3 Related work

In the theoretical computer science literature, earlier work focused on learning from mixture of wellseparated Gaussians (component distributions), where one aims to classify each sample according to which component distribution it comes from; See for example [13, 5, 36, 3, 22, 25]. In earlier works [13, 5], the separation requirement depends on the number of dimensions of each distribution; this has recently been reduced to be independent of p, the dimensionality of the distribution for certain classes of distributions [3, 23]. While our aim is different from those results, where n > pis almost universal and we focus on cases p > n (a.k.a. high dimensional setting), we do have one common axis for comparison, the ℓ_2 -distance between any two centers of the distributions as stated in (30) (or (23)), which is essentially optimal. Results in [38, 10] were among the first such results towards understanding rigorously and intuitively why their proposed algorithms and previous methods in [30, 33] work with low sample settings when $p \gg n$ and np satisfies (23). However, such results were only known to exist for balanced max-cut algorithms [38, 10], and hence these were structural as no polynomial time algorithms were given for finding the max-cut. The main contribution of the present work and [39] is: we use the proposed SDP (10) and the related spectral algorithms to find the partition, and prove quantitatively tighter bounds than those in [38] and [10] by removing these logarithmic factors entirely. Recently, these barriers have also been broken down by a sequence of work [35, 15, 18]. For example, [15, 16] have also established such bounds, but they focus on balanced clusters and require an extra $\sqrt{\log n}$ factor in (25) in the second component:

In [15], cf. eq.(8):
$$\Delta^2 = p\gamma = \Omega\left(1 + \sqrt{p\log n/n}\right)$$
 or (25)

In [16], cf. eq.(13):
$$\Delta^2 = p\gamma = \Omega\left((1 \lor p/n) + \sqrt{p \log n/n}\right).$$
(26)

As a result, in (26), a lower bound on the sample size is imposed: $n \ge 1/\gamma$ in case p > n, and moreover, the size of the matrix $np = \Omega(\log n/\gamma^2)$, similar to the bounds in [8]; cf. Theorem 1.2 therein. Hence, these earlier results still need the SNR to be at the order of $s^2 = \Omega(\log n)$. We compare with [18] in Section 3. Importantly, because of these new concentration of measure bounds, our theory works for the small sample and low SNR (n cases forboth SDP (in its local and global analyses) and the spectral algorithm under (A2), which is perhaps the more surprising case. For example, previously, the spectral algorithms in [8] partition sample points based on the top few eigenvectors of the gram matrix XX^T , following an idea that goes back at least to [17]. In [8], the two parameters n, p are assumed to be roughly at the same order. hence not allowing a full range of tradeoffs between the two dimensions as considered in the present work. Such a lower bound on n was deemed to be unnecessary given the empirical evidence [8]. In contrast, the spectral analysis in [39] uses the leading eigenvector of the gram matrix YY^T , based on centered data, which will directly improve the results in [8] as we can now remove the lower bound on n. We also refer to [22, 34, 20, 28, 1, 6, 11, 7, 18, 26, 16, 27, 2, 29] and references therein for related work on the Stochastic Block Models (SBM), mixture of (sub)Gaussians and clustering in more general metric spaces.

Organization. The rest of the paper is organized as follows. In Section 2, we present the subgaussian mixture models and our main theoretical results. In Section 3, we highlight the connections and key differences between our approach and convex relaxation algorithms based on the k-means criterion, and other related work. We present preliminary results for the local analysis (for proving Theorem 2.3) in Section 4. We prove Theorem 2.3 in Section 4.1. We provide a proof sketch for the key lemmas for Theorem 2.3 in Section 5. We conclude in Section 6. We place all additional technical proofs in the supplementary material.

Notation. Let \mathbf{B}_{2}^{n} and \mathbb{S}^{n-1} be the unit Euclidean ball and the unit sphere in \mathbb{R}^{n} respectively. Let e_{1}, \ldots, e_{n} be the canonical basis of \mathbb{R}^{n} . For a set $J \subset \{1, \ldots, n\}$, denote $E_{J} = \operatorname{span}\{e_{j} : j \in J\}$. For a vector $v \in \mathbb{R}^{n}$, we use v_{J} to denote the subvector $(v_{j})_{j \in J}$. For a vector x, $||x||_{\infty} := \max_{j} |x_{j}|$, $||x||_{1} := \sum_{j} |x_{j}|$, and $||x||_{2} := \sqrt{\sum_{j} x_{j}^{2}}$; diag(x) denotes the diagonal matrix whose main diagonal entries are the entries of x. For a matrix $B \in \mathbb{R}^{n \times n}$, $\operatorname{tr}(B) = \sum_{i=1}^{n} B_{ii}$. Denote by $B_{1}^{n \times n} = \{(b_{ij}) \in \mathbb{R}^{n \times n} : \sum_{i} \sum_{j} |b_{ij}| \leq 1\}$ the ℓ_{1} unit ball over vec $\{B\}$. Let diag(A) and offd(A) be the diagonal and the off-diagonal part of matrix A respectively. For a matrix A, let $||A||_{\infty} = \max_{i} \sum_{j=1}^{n} |a_{ij}|$ denote the maximum absolute row sum; Let $||A||_{\max} = \max_{i,j} |a_{ij}|$ denote the component-wise max norm. For two numbers $a, b, a \wedge b := \min(a, b)$, and $a \lor b := \max(a, b)$. We write $a \asymp b$ if $ca \leq b \leq Ca$ for some positive absolute constants c, C which are independent of n, p, and γ . We write f = O(h) or $f \ll h$ if $|f| \leq Ch$ for some absolute constant $C < \infty$ and $f = \Omega(h)$ or $f \gg h$ if h = O(f). We

write f = o(h) if $f/h \to 0$ as $n \to \infty$, where the parameter n will be the size of the matrix under consideration. In this paper, $C, C_1, C_2, C_4, c, c', c_1$, etc, denote various absolute positive constants which may change line by line.

2 Theory

In this section, we first state the main result derived from the local analysis, namely, Theorem 2.3, where we present an error bound (31) that decays exponentially in the SNR parameter s^2 as defined in (29). We then present Corollary 2.4, followed by discussions. We prove Theorem 2.3 in Section 4.1. First, we have two definitions.

Definition 2.1. A random vector $W \in \mathbb{R}^m$ is called sub-gaussian if the one-dimensional marginals $\langle W, h \rangle$ are sub-gaussian random variables for all $h \in \mathbb{R}^m$: (1) W is called isotropic if for every $h \in \mathbb{S}^{m-1}$, $\mathbb{E} |\langle W, h \rangle|^2 = 1$; (2) W is ψ_2 with a constant C_0 if for every $h \in \mathbb{S}^{m-1}$, $||\langle W, h \rangle||_{\psi_2} \leq C_0$. The sub-gaussian norm of $W \in \mathbb{R}^m$ is denoted by

$$\|W\|_{\psi_2} := \sup_{h \in \mathbb{S}^{m-1}} \|\langle W, h \rangle \|_{\psi_2}.$$
 (27)

Definition 2.2. (Data generative process.) Let H_1, \ldots, H_n be deterministic $p \times m$ matrices, where we assume that $m \ge p$. Suppose that random matrix $\mathbb{W} = (w_{jk}) \in \mathbb{R}^{n \times m}$ has $W_1, \ldots, W_n \in \mathbb{R}^m$ as independent row vectors, where W_j , for each j, is an isotropic sub-gaussian random vector with independent entries satisfying

$$\forall j \in [n], \quad \operatorname{Cov}(W_j) := \mathbb{E}(W_j W_j^T) = I_m, \quad \mathbb{E}[w_{jk}] = 0, \quad and \quad \max_{jk} \|w_{jk}\|_{\psi_2} \le C_0, \quad \forall k.$$
(28)

Suppose that we have for row vectors $\mathbb{Z}_1, \ldots, \mathbb{Z}_n \in \mathbb{R}^p$ of the noise matrix $\mathbb{Z} \in \mathbb{R}^{n \times p}$,

$$\forall j = 1, \dots, n, \quad \mathbb{Z}_j^T = W_j^T H_j^T, \quad where \quad H_j \in \mathbb{R}^{p \times m}, \quad 0 < \|H_j\|_2 < \infty,$$

and H_j 's are allowed to repeat, for example, across rows from the same cluster C_i for some i = 1, 2. Throughout this paper, we assume that $m \ge p$ to simplify our exposition, although this is not necessary.

Signal-to-noise ratios. This notion of SNR (12) can be properly adjusted when coordinates in \mathbb{Z}_i are dependent in view of (19) and (20):

SNR anisotropic:
$$s^2 = \frac{\Delta^2}{C_0^2 \max_j \|\mathsf{Cov}(\mathbb{Z}_j)\|_2} \wedge \frac{np\gamma^2}{C_0^4 \max_j \|\mathsf{Cov}(\mathbb{Z}_j)\|_2^2}.$$
 (29)

Theorem 2.3. Suppose that for $j \in C_i \subset [n]$, $\mathbb{E}X_j = \mu^{(i)}$, where i = 1, 2. Let \widehat{Z} be a solution of SDP1. Let s^2 be as defined in (29). Let C_0 be as defined in (28). Suppose the noise matrix $\mathbb{Z} = X - \mathbb{E}(X)$ is generated according to Definition 2.2:

 $\forall j \in \mathcal{C}_i, \quad \mathbb{Z}_j = H_i W_j, \quad where \quad H_i \in \mathbb{R}^{p \times m} \quad is \ deterministic \ and \quad 0 < \|H_i\|_2 < \infty,$ for $i \in \{1, 2\}$. Suppose that for some absolute constants C, C_1 and some $0 < \xi \le w_{\min}^2/16$,

$$p\gamma \ge \frac{CC_0^2 \max_j \|Cov(\mathbb{Z}_j)\|_2}{\xi^2} \quad and \quad np \ge \frac{C_1 C_0^4 \max_j \|Cov(\mathbb{Z}_j)\|_2^2}{\gamma^2 \xi^2}.$$
 (30)

Suppose (A2) holds, where the parameter ξ in (22) is understood to be the same as the parameter ξ in (30). Then with probability at least $1-2\exp(-c_1n)-c_2/n^2$, for some absolute constants c_0, c_1, c_2 ,

$$\left\| \widehat{Z} - u_2 u_2^T \right\|_1 / n^2 \le \exp\left(-c_0 s^2 w_{\min}^4 \right).$$
(31)

Corollary 2.4. (Exponential decay in s^2) Let \hat{x} denote the eigenvector of \hat{Z} corresponding to the largest eigenvalue, with $\|\hat{x}\|_2 = \sqrt{n}$. Denote by $\theta_{SDP} = \angle(\hat{x}, u_2)$, the angle between \hat{x} and u_2 , where recall $u_{2j} = 1$ if $j \in C_1$ and $u_{2j} = -1$ if $j \in C_2$. In the settings of Theorem 2.3, with probability at least $1 - 2\exp(-cn) - 2/n^2$, for some absolute constants c, c_0, c_1 ,

$$\sin(\theta_{SDP}) \leq 2 \left\| \widehat{Z} - u_2 u_2^T \right\|_2 / n \leq \exp(-c_1 s^2 w_{\min}^4) \quad and$$
$$\min_{\alpha = \pm 1} \left\| (\alpha \widehat{x} - u_2) / \sqrt{n} \right\|_2 \leq 2^{3/2} \left\| \widehat{Z} - u_2 u_2^T \right\|_2 / n \leq 4 \exp(-c_0 s^2 w_{\min}^4 / 2). \tag{32}$$

Corollary 2.4 follows from the Davis-Kahan Theorem and Theorem 2.3, and is given in [39]; cf. Proof of Corollary 2.8 therein. Our local and global analyses also show the surprising result that Theorems 1.4 and 2.3 do not depend on the clusters being balanced, nor do they require identical variance or covariance profiles, so long as (A2) holds.

The bias and variance tradeoffs. Roughly speaking, to ensure that $||\mathbb{E}B - R||_2$ is bounded, we require

$$(A2') \quad |V_1 - V_2| \le w = O\left(n\Delta \lor \sqrt{np}\right), \text{ where } \Delta = \sqrt{p\gamma}.$$
(33)

Notice that $|V_1 - V_2| = O(p)$ by definition, and hence (33) holds trivially in the large sample setting where n > p. Given a fixed average quality parameter γ , the tolerance on $|V_1 - V_2|$ depends on the sample size n, the total data size D := np, and the separation parameter Δ . In particular, the tolerance parameter w is chosen to be at the same order as the upper bound we obtain on $||YY^T - \mathbb{E}YY^T||_2$ in Theorem 2.7. In the supplementary Theorem D.2, we show that the operator norm for B - R is controlled essentially at the same order as $||YY^T - \mathbb{E}(YY^T)||_2$. To see this, first, we have for A as in (10),

$$\|B - \mathbb{E}B\|_{2} \leq \|YY^{T} - \mathbb{E}YY^{T}\|_{2} + |\lambda - \mathbb{E}\lambda| \|E_{n} - I_{n}\|_{2}, \quad \text{since}$$
(34)

$$B - \mathbb{E}B := A - \mathbb{E}A = YY^T - \mathbb{E}YY^T - (\lambda - \mathbb{E}\lambda)(E_n - I_n).$$
(35)

Combining Lemmas 2.6, 5.4, Theorem 2.7, with (34), we have with probability at least $1 - 2\exp(-cn)$,

$$\|B - R\|_{2} \leq \|B - \mathbb{E}B\|_{2} + \|\mathbb{E}B - R\|_{2} \leq 2 \|YY^{T} - \mathbb{E}(YY^{T})\|_{2} + \|\mathbb{E}B - R\|_{2} \leq \xi np\gamma.$$
(36)

First, we state Lemma 2.5, which characterizes the two-group design matrix covariance structures as considered in Theorems 1.4 and 2.3, with isotropic design as the special case.

Lemma 2.5. (Two-group sub-gaussian mixture model) Denote by X the two-group design matrix as considered in (1). Let $W_1, \ldots, W_n \in \mathbb{R}^m$ be independent, mean-zero, isotropic, subgaussian random vectors satisfying (28). Let $\mathbb{Z}_j = X_j - \mathbb{E}X_j = H_i W_j$, for all $j \in C_i$, where $H_i \in \mathbb{R}^{p \times m}$ is deterministic and $0 < ||H_i||_2 < \infty$, for each $i \in \{1, 2\}$. Then $\mathbb{Z}_1, \ldots, \mathbb{Z}_n$ are independent sub-gaussian random vectors with $\mathsf{Cov}(\mathbb{Z}_i)$ satisfying (19) and (20), where

$$\forall j \in \mathcal{C}_i, \quad \operatorname{Cov}(\mathbb{Z}_j) := \mathbb{E}(\mathbb{Z}_j \mathbb{Z}_j^T) = \mathbb{E}(H_i W_j W_j^T H_i^T) = H_i H_i^T \text{ and } V_i := \|H_i\|_F^2 = \operatorname{tr}(\Sigma_i).$$
(37)

Lemma 2.6. [39] Let X be as in Lemma 2.5. Suppose (A2) holds. Suppose that $\xi \geq \frac{1}{2n} (4 \vee \frac{1}{w_{\min}})$ and $n \geq 4$. Then we have $\|\mathbb{E}B - R\|_2 \leq \frac{2}{3} \xi n p \gamma$. Finally, when $V_1 = V_2$, we have $\|\mathbb{E}B - R\|_2 \leq p \gamma/3$.

Theorem 2.7. (Anisotropic design matrix) [39] Let X be as in Lemma 2.5. Let Y be as in (8). Let $\mu = \frac{\mu^{(1)} - \mu^{(2)}}{\sqrt{p\gamma}} \in \mathbb{S}^{p-1}$. Then with probability at least $1 - 2\exp(-c_8n)$, for C_0 as defined in (28),

$$\|YY^T - \mathbb{E}(YY^T)\|_2 \le C_3(C_0 \max_i \|H_i^T \mu\|_2) n\sqrt{p\gamma} + C_4(C_0 \max_i \|H_i\|_2)^2(\sqrt{pn} \lor n) \le \frac{1}{6}\xi np\gamma.$$

See the supplementary Section D for a proof sketch of Theorem 2.7 and the proof of Lemma 2.6, which we include for self-containment. The error bound in (36) was crucial in the global analysis leading to Theorem 1.4, as well as the simultaneous analysis of the spectral clustering method; cf. Section 3, Variation 2. Indeed, our previous and current results show that even when n is small, by increasing p such that the total sample size satisfies the second condition in (23) and (30), we ensure partial recovery of cluster structures using the SDP (10) or the spectral algorithm as described in [39]. On the other hand, once there are enough features such that the distance between the two centers are bounded below by a constant, adding more samples (individuals) to the clusters will reduce the number of features we need to do partial recovery. Now (30) implies that

$$256/w_{\min}^{4} \leq \frac{1}{\xi^{2}} \leq \frac{p\gamma}{CC_{0}^{2}\max_{j}\|\mathsf{Cov}(\mathbb{Z}_{j})\|_{2}} \wedge \frac{np\gamma^{2}}{C_{1}C_{0}^{4}\max_{j}\|\mathsf{Cov}(\mathbb{Z}_{j})\|_{2}^{2}} \asymp s^{2}, \quad \text{and hence}$$
(38)
$$\xi np\gamma/2 \geq C_{0}^{2}\max_{i}\|\mathsf{Cov}(\mathbb{Z}_{i})\|_{2}\left(\frac{n}{\xi} \vee \sqrt{np}\right) =: w, \quad \text{given that} \quad \frac{1}{\xi} \leq \frac{\sqrt{p\gamma}}{CC_{0}\max_{j}\|\mathsf{Cov}(\mathbb{Z}_{j})\|_{2}^{1/2}},$$

resulting in (33) being a sufficient condition for (A2) to hold. Given fixed p and γ , increasing the sample size n also leads to a *higher tolerance* on the variance discrepancy $|V_1 - V_2|$ between the two component distributions as shown in (33). To control the misclassification error using the global analysis, the parameters (δ, ξ) in Theorem 1.4 can be chosen to satisfy the following relations:

$$\xi^2 \asymp \frac{C_0^2}{p\gamma} \vee \frac{C_0^4}{np\gamma^2} = \frac{1}{s^2} \quad \text{and} \quad \delta = \frac{2K_G\xi}{w_{\min}^2},\tag{39}$$

in view of (23) and (24). Thus we obtain in Theorem 1.4 that the misclassification error is inversely proportional to the square root of the SNR parameter s^2 as in (12) in view of (39), while in Theorem 2.3 and Corollary 2.4, the error decays exponentially in the SNR parameter s^2 as defined in (29).

3 The k-means criterion of a partition

We now discuss the k-means criterion and its semidefinite relaxations. Denote by $X \in \mathbb{R}^{n \times p}$ the data matrix with row vectors X_i as in (40) (see also (6)). The k-means criterion of a partition $\mathcal{C} = \{\mathcal{C}_1, \ldots, \mathcal{C}_k\}$ of sample points $\{1, \ldots, n\}$ is based on the total sum-of-squared Euclidean distances from each point $X_i \in \mathbb{R}^p$ to its assigned cluster centroid \mathbf{c}_j , namely,

$$g(X, \mathcal{C}, k) := \sum_{j=1}^{k} \sum_{i \in \mathcal{C}_j} \|X_i - \mathbf{c}_j\|_2^2, \quad \text{where} \quad \mathbf{c}_j := \frac{1}{|\mathcal{C}_j|} \sum_{\ell \in \mathcal{C}_j} X_\ell \in \mathbb{R}^p.$$
(40)

Getting a global solution to (40) through an integer programming formulation as in [32] and [31] is NP-hard and it is NP-hard for k = 2 [14, 4]. The partition C can be represented by a block diagonal matrix of size $n \times n$, defined as: $\forall i, j \in [k]^2, \forall a, b \in C_i \times C_j$,

$$B_{ab}^* = 1/|\mathcal{C}_j|$$
 if $i = j$ and $B_{ab}^* = 0$ otherwise.

The collection of such matrices can be described by

$$\mathcal{P}_k = \{ B \in \mathbb{R}^{n \times n} : B = B^T, B \ge 0, B^2 = B, \operatorname{tr}(B) = k, B\mathbf{1}_n = \mathbf{1}_n \},$$
(41)

where $B \ge 0$ means that all elements of B are nonnegative. Minimizing the k-means objective $g(X, \mathcal{C}, k)$ is equivalent to [37, 32, 31],

maximize
$$\langle XX^T, Z \rangle$$
 subject to $Z \in \mathcal{P}_k$. (42)

Peng-Wei relaxations. Let Ψ_n denote the linear space of real *n* by *n* symmetric matrices. Now consider the semidefinite relaxation of the *k*-means objective (42),

maximize
$$\langle XX^T, Z \rangle$$
 subject to $Z \in \mathcal{M}_k$, (43)
where $\mathcal{M}_k = \{ Z \in \Psi_n : Z \ge 0, Z \succeq 0, \operatorname{tr}(Z) = k, Z \mathbf{1}_n = \mathbf{1}_n \}$

as considered in [31]. The key differences between this and the SDP (10) are: (a) In the convex set \mathcal{M}_{opt} (15), we do not enforce that all entries are nonnegative, namely, $Z_{ij} \geq 0, \forall i, j$. This allows faster computation; (b) In order to derive concentration of measure bounds that are sufficiently tight, we make a natural, yet important data processing step, where we center the data according to their column mean following Definition 1.1 before computing A as in (9); (c) We do not enforce $Z\mathbf{1}_n = \mathbf{1}_n$. Note that for \mathcal{M}_k as in (43), when $Z_{ij} \geq 0, \forall i, j$, row sum $||Z_{j,\cdot}||_1 = 1, \forall j$ and hence $||Z||_2 \leq ||Z||_{\infty} = 1$.

Variation 2. To speed up computation, one can drop the nonnegative constraint on elements of Z leading to the following semidefinite relaxation [31]:

maximize
$$\langle XX^T, Z \rangle$$
 subject to $I_n \succeq Z \succeq 0$, $\operatorname{tr}(Z) = k$, and $Z\mathbf{1}_n = \mathbf{1}_n$. (44)

Now when Z is a feasible solution to (44), define

$$Z_1 := Z - \mathbf{1}_n \mathbf{1}_n^T / n \text{ and hence } Z_1 := (I - P_1) Z = (I - P_1) Z (I - P_1),$$
(45)

where recall $\mathbf{1}_n/\sqrt{n}$ is the unit-norm leading eigenvector of Z. Now [31] shows that the set of feasible solutions to (44) have immediate connections to the SVD of YY^T through the following reduction:

maximize
$$\langle YY^T, Z_1 \rangle$$
 subject to $I_n \succeq Z_1 \succeq 0$, $\operatorname{tr}(Z_1) = k - 1$. (46)

Then the algorithm for solving (46) and (44) is given as follows [31]:

Spectral Clustering: (a) Using singular value decomposition (SVD) method to compute the first k-1 largest eigenvalues of YY^T and their corresponding eigenvectors v_1, \ldots, v_{k-1} ;

and **(b)** Set
$$Z_1 = \sum_{j=1}^{k-1} v_j v_j^T$$
; and return $Z = \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^T + Z_1$ as a solution to (44).

SVD-based algorithms allow even faster computation. In particular, for k = 2, we have the optimal solution of (44) being $Z_1 = v_1 v_1^T$, where v_1 is the leading eigenvector of YY^T . The signs of the coefficients of v_1 correctly estimate the partition of the vertices, up to $O(\xi^2 n)$ misclassified vertices, where we set $\xi^2 \simeq 1/s^2$ in [39]; cf. (39). Hence the misclassification error is bounded to be inversely proportional to the SNR parameter s^2 ; cf. (39). This should be compared with Theorem 2.3 and Corollary 2.4, where the misclassification error is improved to $O(n \exp(-c_0 s^2 w_{\min}^4))$. Moreover, one can sort the values of v_1 and find the *nearly optimal* partition according to the k-means criterion; See Algorithm 2 and numerical examples in [39].

Discussions. The main issue with the k-means relaxation is that the solutions tend to put sample points into groups of the same sizes, and moreover, the diagonal matrix Γ can cause a bias, where

$$\Gamma = (\mathbb{E}[\langle \mathbb{Z}_i, \mathbb{Z}_j \rangle])_{i,j} = \operatorname{diag}([\operatorname{tr}(\operatorname{Cov}(\mathbb{Z}_1)), \dots, \operatorname{tr}(\operatorname{Cov}(\mathbb{Z}_n))])_{i,j})$$

especially when V_1, V_2 differ from each other. The authors of [35], [18], and [9] propose a preliminary estimator of Γ , denoted by $\widehat{\Gamma}$, and consider

$$\widehat{Z} \in \arg \max_{Z \in \mathcal{M}_k} \langle X X^T - \widehat{\Gamma}, Z \rangle , \quad \text{where } \mathcal{M}_k \text{ is as in } (43) , \qquad (47)$$

instead of the original Peng-Wei SDP relaxation (43). Consequently, besides computation, another main advantage of our SDP and spectral formulation in [39] is that we do not need to have a separate estimator for $tr(\Sigma_j)$, where $\Sigma_j, j = 1, 2$ denotes the covariance matrices of sub-gaussian random vectors $\mathbb{Z}_j, j \in [n]$, so long as (A2) holds. In some sense, SDP1 has a certain inherent tolerance on the variance discrepancy between the two populations in the sense of (33). When (A2) is violated, we may adopt similar ideas in [35, 18] to make adjustments to SDP1 to correct the bias. It is an arguably simpler task to deal with $|V_1 - V_2|$ than actual values of V_1, V_2 . See Remark 5.5.

4 Proof sketch for Theorem 2.3

In this section, we need to state some intermediate results followed by the actual proof of Theorem 2.3 in Section 4.1. Let the reference matrix R be constructed as in Definition 1.2. Lemma 4.1 follows from definitions of B and R, and the lower bound in Lemma 4.4.

Lemma 4.1. (Elementary Inequality) By optimality of $\widehat{Z} \in \mathcal{M}_{opt}$, we have

$$p\gamma w_{\min}^{2} \left\| \widehat{Z} - Z^{*} \right\|_{1} \leq \langle R, Z^{*} - \widehat{Z} \rangle \leq \langle YY^{T} - \mathbb{E}(YY^{T}), \widehat{Z} - Z^{*} \rangle + \langle \mathbb{E}B - R, \widehat{Z} - Z^{*} \rangle - \langle (\lambda - \mathbb{E}\lambda)(E_{n} - I_{n}), \widehat{Z} - Z^{*} \rangle =: W_{1} + W_{2} + W_{3}.$$

$$(48)$$

We prove Lemma 4.1 in Section 5.1. To bound the excess risk $\langle R, Z^* - \hat{Z} \rangle$, we present an upper bound on the bias term $W_2 := \left| \langle \mathbb{E}B - R, \hat{Z} - Z^* \rangle \right|$ as well as the large deviation bound on $W_3 := \left| \langle (\lambda - \mathbb{E}\lambda)(E_n - I_n), \hat{Z} - Z^* \rangle \right|$ over all $\hat{Z} \in \mathcal{M}_{opt}$. Lemma 4.2 shows that each term takes out at most a fraction of the total signal strength on the LHS of (48) respectively. We then obtain an upper bound on the variance term W_1 in Lemma 4.3 in the local neighborhood of Z^* .

Lemma 4.2. Suppose all conditions in Theorem 2.3 hold. Let $\frac{1}{4(n-1)} < \xi \le w_{\min}^2/16$. Then

$$\forall \widehat{Z} \in \mathcal{M}_{opt}, \quad \left| \left\langle \mathbb{E}B - R, \widehat{Z} - Z^* \right\rangle \right| =: |W_2| \leq 2p\gamma \left\| \widehat{Z} - Z^* \right\|_1 \left(\xi + \frac{1}{4(n-1)} \right); \quad (49)$$

Under the settings of Theorem 2.7, we have with probability at least $1 - \exp(cn)$,

$$\forall \quad \widehat{Z} \in \mathcal{M}_{opt}, \quad \left| \left\langle (\lambda - \mathbb{E}\lambda)(E_n - I_n), \widehat{Z} - Z^* \right\rangle \right| =: |W_3| \leq \frac{1}{3} \xi p \gamma \left\| \widehat{Z} - Z^* \right\|_1$$

Lemma 4.3. Suppose all conditions in Theorem 2.7 hold. Let $r_1 \leq 2qn$ for a positive integer $1 \leq q < n$. Then on event \mathcal{G}_1 , where $\mathbb{P}(\mathcal{G}_1) \geq 1 - c/n^2$, we have for $\xi \leq w_{\min}^2/16$,

$$\sup_{\widehat{Z}\in\mathcal{M}_{opt}\cap(Z^*+r_1B_1^{n\times n})} \left| \langle YY^T - \mathbb{E}(YY^T), \widehat{Z} - Z^* \rangle \right| \leq \frac{5}{6} \xi p\gamma \left\| \widehat{Z} - Z^* \right\|_1 + C'(C_0 \max_i \|H_i\|_2) \left(n\sqrt{p\gamma} + (C_0 \max_i \|H_i\|_2)\sqrt{np} \right) \left[\frac{r_1}{2n} \right] \sqrt{\log(2en/\lceil \frac{r_1}{2n} \rceil)}.$$

The final result we need for both the local and global analyses is to verify a non-trivial global curvature of the excess risk $\langle R, Z^* - \hat{Z} \rangle$ for the feasible set \mathcal{M}_{opt} at the maximizer Z^* .

Lemma 4.4. (Excess risk lower bound) [39] Let R be as in Definition 1.2 and $Z^* = u_2 u_2^T$. Then

for every
$$Z \in \mathcal{M}_{opt}$$
, $\langle R, Z^* - Z \rangle \ge p\gamma w_{\min}^2 \|Z - Z^*\|_1$. (50)

This is already stated as the lower bound in (48). Hence we can control $\xi \leq w_{\min}^2/16$ and the sample size lower bound in (30) so that each component W_2, W_3 is only a fraction of the signal at the level of $p\gamma w_{\min}^2 ||(Z^* - Z)||_1$. We prove Lemma 4.2 in Section 5.2. We prove Lemma 4.3 in Section 5.3. In particular, the proof of Lemma 4.3 requires the large deviation bound in Theorem 2.7.

4.1 Proof of Theorem 2.3

Let $\lceil r_1/(2n) \rceil =: q < n$ and $r_1 := \left\| \widehat{Z} - Z^* \right\|_1$. Then, we have by Lemma 4.3, on event \mathcal{G}_1 ,

$$\mathcal{G}_{1}: \sup_{\widehat{Z}\in\mathcal{M}_{\text{opt}}\cap(Z^{*}+r_{1}B_{1}^{n\times n})} \left| \langle YY^{T} - \mathbb{E}(YY^{T}), \widehat{Z} - Z^{*} \rangle \right| \leq \frac{5}{6} \xi p\gamma \left\| \widehat{Z} - Z^{*} \right\|_{1} + C'C_{0} \max_{i} \|H_{i}\|_{2} \left(n\sqrt{p\gamma} + (C_{0} \max_{i} \|H_{i}\|_{2})\sqrt{np} \right) \left\lceil \frac{r_{1}}{2n} \right\rceil \sqrt{\log(2en/\lceil \frac{r_{1}}{2n}\rceil)},$$
(51)

and by Lemma 4.2, $\forall \widehat{Z} \in \mathcal{M}_{opt}$,

on event
$$\mathcal{G}_2$$
: $|W_3| = \left| \langle (\lambda - \mathbb{E}\lambda)(E_n - I_n), \widehat{Z} - Z^* \rangle \right| \leq \frac{1}{3} \xi p \gamma \left\| \widehat{Z} - Z^* \right\|_1.$ (52)

Let $C, C', C_1, c, c_1, c_2, \ldots$ be absolute positive constants. For the rest of the proof, we assume $\mathcal{G}_1 \cap \mathcal{G}_2$ holds, where $\mathbb{P}(\mathcal{G}_2 \cap \mathcal{G}_1) \geq 1 - \exp(cn) - c/n^2$. By Lemmas 4.1 and 4.4, (49), (51), and (52), we have for all $\widehat{Z} \in \mathcal{M}_{opt} \cap (Z^* + r_1 B_1^{n \times n})$,

$$0 \leq p\gamma w_{\min}^{2} \left\| Z^{*} - \widehat{Z} \right\|_{1} \leq \langle R, Z^{*} - \widehat{Z} \rangle \leq \left| \langle YY^{T} - \mathbb{E}(YY^{T}), \widehat{Z} - Z^{*} \rangle \right| + |W_{2}| + |W_{3}|$$

$$\leq (10/3)\xi p\gamma \left\| \widehat{Z} - Z^{*} \right\|_{1} + C'(C_{0} \max_{i} \|H_{i}\|_{2})q\sqrt{\log(2en/q)} \left(n\sqrt{p\gamma} + C_{0}(\max_{i} \|H_{i}\|_{2})\sqrt{np} \right)$$

By moving $(10/3)\xi p\gamma \left\| \widehat{Z} - Z^* \right\|_1 < \frac{5}{24}p\gamma w_{\min}^2 r_1$, where $\xi \le w_{\min}^2/16$, to the LHS of (53), we have

$$\frac{19}{24} p \gamma w_{\min}^2 r_1 \leq p \gamma w_{\min}^2 \left\| Z^* - \widehat{Z} \right\|_1 - (10/3) \xi p \gamma \left\| \widehat{Z} - Z^* \right\|_1 \\
\leq C C_0(\max_i \|H_i\|_2) \sqrt{\log(2en/q)} \left(2nq \sqrt{p\gamma} + C_0(\max_i \|H_i\|_2) q \sqrt{np} \right).$$
(54)

Then q must satisfy one of the following two conditions in order to guarantee (54): for $r_1 \leq 2nq$,

1. Under the assumption that $\Delta^2 w_{\min}^4 = p\gamma w_{\min}^4 = \Omega((C_0 \max_i ||H_i||_2)^2)$, suppose

$$p\gamma w_{\min}^2 \leq C_1 C_0(\max_i \|H_i\|_2) \sqrt{\log(2en/q)} \sqrt{p\gamma}, \quad \text{which implies that}$$
$$\frac{cp\gamma w_{\min}^4}{(C_0 \max_i \|H_i\|_2)^2} \leq \log(2en/q) \text{ and hence } q \leq n \exp\Big(-\frac{c_1 \Delta^2 w_{\min}^4}{(C_0 \max_i \|H_i\|_2)^2}\Big).$$

2. Alternatively, under the assumption that $np\gamma^2 w_{\min}^4 = \Omega(C_0^4)$, we require

$$p\gamma w_{\min}^2 r_1 \le 2qnp\gamma w_{\min}^2 \le C_2(C_0 \max_i ||H_i||_2)^2 \sqrt{\log(2en/q)} q\sqrt{np}$$

and hence $np\gamma^2 w_{\min}^4 \leq C_3 (C_0 \max_i ||H_i||_2)^4 \log(2en/q)$, which implies that

$$\log(q/2en) \leq -\frac{C'np\gamma^2 w_{\min}^4}{(C_0 \max_i \|H_i\|_2)^4}, \quad \text{resulting in} \quad q \leq n \exp\big(-\frac{c_2 np\gamma^2 w_{\min}^4}{(C_0 \max_i \|H_i\|_2)^4}\big).$$

Putting things together, we have

$$\left\| Z^* - \widehat{Z} \right\|_{1} := r_1 \le 2qn \le n^2 \exp(-Cs^2 w_{\min}^4), \quad \text{where } s^2 \text{ is as defined in (29)},$$
(55)
and $q \le n \exp(-Cs^2 w_{\min}^4)$ for $s^2 = \frac{\Delta^2}{(C_0 \max_i \|H_i\|_2)^2} \wedge \frac{np\gamma^2}{(C_0 \max_i \|H_i\|_2)^4}.$

5 Proof of key lemmas for Theorem 2.3

First we state two facts. We prove Fact 5.2 in Section B.

Fact 5.1. By the supplementary Proposition D.1, we have for the optimal solution \hat{Z} as in SDP (10),

 $\operatorname{diag}(\widehat{Z}) = \operatorname{diag}(Z^*) = I_n \quad and \quad \langle I_n, \widehat{Z} - Z^* \rangle = \operatorname{tr}(\widehat{Z}) - \operatorname{tr}(Z^*) = 0.$

Fact 5.2. Recall $|C_1| = n_1$ and $|C_2| = n_2$. Let Z^* be as defined in Lemma 1.3:

$$Z^* = u_2 u_2^T = \begin{bmatrix} E_{n_1} & -E_{n_1 \times n_2} \\ -E_{n_2 \times n_1} & E_{n_2} \end{bmatrix}.$$
 (56)

Then, for $P_2 = Z^*/n$, we have $\operatorname{tr}(P_2(Z^* - \widehat{Z})) = \frac{1}{n} \operatorname{tr}(Z^*(Z^* - \widehat{Z})) = \frac{1}{n} \left\| Z^* - \widehat{Z} \right\|_1 \le 2(n-1).$

5.1 Proof of Lemma 4.1

Denote by $\widetilde{A} = YY^T - \lambda E_n$. Now for B as in (17) and $R = \mathbb{E}(Y)\mathbb{E}(Y)^T$, we have

$$\mathbb{E}B - R := \mathbb{E}(YY^T) - \mathbb{E}(Y)\mathbb{E}(Y)^T - \mathbb{E}\lambda(E_n - I_n) - \mathbb{E}\tau I_n.$$
(57)

Clearly $\widehat{Z}, Z^* \in \mathcal{M}_{opt}$ by definition, we have $\langle \widehat{Z}, I_n \rangle = \langle I_n, Z^* \rangle = n$. By optimality of $\widehat{Z} \in \mathcal{M}_{opt}$, we have by (17) and (10),

$$\langle B, \widehat{Z} - Z^* \rangle := \langle YY^T - \lambda(E_n - I_n) - \mathbb{E}\tau I_n, \widehat{Z} - Z^* \rangle$$

$$:= \langle YY^T - \lambda E_n, \widehat{Z} - Z^* \rangle \ge 0,$$
 (58)

where (58) holds since $\widehat{Z} := \arg \max_{Z \in \mathcal{M}_{opt}} \langle \widetilde{A}, Z \rangle$ and $\langle I_n, \widehat{Z} - Z^* \rangle = 0$. Now we have by (57),

$$M := \mathbb{E}(YY^{T}) - \mathbb{E}(Y)\mathbb{E}(Y)^{T} - \lambda(E_{n} - I_{n}) - \mathbb{E}\tau I_{n}$$

= $\mathbb{E}(YY^{T}) - \mathbb{E}(Y)\mathbb{E}(Y)^{T} - \mathbb{E}\lambda(E_{n} - I_{n}) - \mathbb{E}\tau I_{n} - (\lambda - \mathbb{E}\lambda)(E_{n} - I_{n})$
= $(\mathbb{E}B - R) - (\lambda - \mathbb{E}\lambda)(E_{n} - I_{n}).$ (59)

Then we have by (58) and (59),

$$\begin{array}{ll} \langle R, Z^* - \widehat{Z} \rangle &:= \langle -\mathbb{E}(Y)\mathbb{E}(Y)^T, \widehat{Z} - Z^* \rangle \\ &\leq & \langle YY^T - \lambda(E_n - I_n) - \mathbb{E}\tau I_n, \widehat{Z} - Z^* \rangle - \langle \mathbb{E}(Y)\mathbb{E}(Y)^T, \widehat{Z} - Z^* \rangle \\ &= & \langle YY^T - \mathbb{E}(YY^T), \widehat{Z} - Z^* \rangle + \langle \mathbb{E}B - R, \widehat{Z} - Z^* \rangle - \langle (\lambda - \mathbb{E}\lambda)(E_n - I_n), \widehat{Z} - Z^* \rangle . \end{array}$$

Thus we have the RHS of (48) holds. Now the LHS of (48) holds by Lemma 4.4. \Box

5.2 Proof of Lemma 4.2

To be fully transparent, we now decompose the bias term into three components. Intuitively, W_0, W_2 and W arise due to the imbalance in variance profiles. Hence (A2) is needed to control this bias.

Proposition 5.3. (Bias decomposition) [39] Denote by

$$W_0 = (V_1 - V_2) \begin{bmatrix} w_2 I_{n_1} & 0\\ 0 & -w_1 I_{n_2} \end{bmatrix} \quad and \quad W_2 := \frac{V_1 - V_2}{n} \begin{bmatrix} E_{n_1} & 0\\ 0 & -E_{n_2} \end{bmatrix}.$$
(60)

Then for W_0, W_2 as defined in (60),

$$\mathbb{E}B - R = W_0 - \mathbb{W} - \frac{\operatorname{tr}(R)}{(n-1)} (I_n - \frac{E_n}{n}), \quad where$$
(61)

$$\mathbb{W} := W_2 + \frac{(V_1 - V_2)(w_2 - w_1)}{n} E_n, \quad and when \ V_1 = V_2, W_0 = \mathbb{W} = 0.$$
(62)

Lemma 5.4. (Deterministic bounds) Let Y be as specified in Definition 1.1. By definition of τ and λ ,

$$(n-1)\left|\lambda - \mathbb{E}\lambda\right| = \left|\tau - \mathbb{E}\tau\right| \le \left\|YY^T - \mathbb{E}(YY^T)\right\|_2.$$
(63)

By Lemma 5.4 and Theorem 2.7, we have with probability at least $1 - 2\exp(-c_6 n)$,

$$\left| (\lambda - \mathbb{E}\lambda) \left\langle (E_n - I_n), \widehat{Z} - Z^* \right\rangle \right| \leq \frac{\left\| YY^T - \mathbb{E}(YY^T) \right\|_2}{n - 1} \left| \sum_{i \neq j} (\widehat{Z} - Z^*)_{ij} \right| \leq \frac{1}{3} \xi p \gamma \left\| \widehat{Z} - Z^* \right\|_1.$$

Now we have by Proposition 5.3,

$$\langle \mathbb{E}B - R, \widehat{Z} - Z^* \rangle = \langle W_0 - \mathbb{W}, \widehat{Z} - Z^* \rangle - \frac{\operatorname{tr}(R)}{n-1} \langle I_n - E_n/n, \widehat{Z} - Z^* \rangle,$$

where W_0 is a diagonal matrix as in (60). Hence by Fact 5.1, (A2), we have for \mathbb{W} as in (62),

$$\langle W_0, \widehat{Z} - Z^* \rangle = \langle W_0, \operatorname{diag}(\widehat{Z} - Z^*) \rangle = 0;$$
(64)

$$\left\| \left\langle \mathbb{W}, \widehat{Z} - Z^* \right\rangle \right\| \leq \|\mathbb{W}\|_{\max} \left\| \widehat{Z} - Z^* \right\|_1 \leq \frac{2|v_1 - v_2|}{n} \| (Z^* - Z) \|_1 \leq \xi p \gamma \| (Z^* - Z) \|_1.$$
(65)

Hence, by the triangle inequality and $tr(R)/n = p\gamma w_1 w_2$,

$$\langle \mathbb{E}B - R, \widehat{Z} - Z^* \rangle \leq \left| \langle \mathbb{W}, \widehat{Z} - Z^* \rangle \right| + \left| \frac{\operatorname{tr}(R)}{n(n-1)} \langle E_n, \widehat{Z} - Z^* \rangle \right|$$

$$\leq \frac{2 |V_1 - V_2|}{n} \left\| \widehat{Z} - Z^* \right\|_1 + \frac{p \gamma w_1 w_2}{n-1} \left\| \widehat{Z} - Z^* \right\|_1$$

$$\leq p \gamma \left\| \widehat{Z} - Z^* \right\|_1 \left(\xi + \frac{1}{4(n-1)} \right) \leq 2\xi p \gamma \left\| \widehat{Z} - Z^* \right\|_1.$$

The lemma thus holds. \Box

Remark 5.5. Balanced Partitions Suppose that $w_1 = w_2$, then we may hope to obtain a tighter bound for $\langle \mathbb{E}B - R, \hat{Z} - Z^* \rangle$. Indeed, it is our conjecture that (65) can be substantially tightened for the balanced case, where $\mathbb{W} = W_2$. For $w_1 = w_2$, we have (a) $\frac{1}{n} \mathbf{1}_n^T Z^* u_2 = \mathbf{1}_n^T u_2 = 0$, and (b) for $\mathbb{W} = W_2$,

$$\frac{1}{|V_1 - V_2|} \left| \left\langle \mathbb{W}, \widehat{Z} - Z^* \right\rangle \right| = \frac{1}{n} \left| \sum_{i,j \in \mathcal{C}_1, j \neq i} (Z_{ij}^* - \widehat{Z}_{ij}) - \sum_{i,j \in \mathcal{C}_2, j \neq i} (Z_{ij}^* - \widehat{Z}_{ij}) \right|$$

$$(66)$$

$$= \frac{1}{n} \left| \left\langle \begin{bmatrix} E_{n_1} & 0\\ 0 & -E_{n_2} \end{bmatrix}, Z^* - \widehat{Z} \right\rangle \right| = \frac{1}{n} \left| \mathbf{1}_n^T (\widehat{Z} - Z^*) u_2 \right| = \frac{1}{n} \left| u_2^T \widehat{Z}(\mathbf{B}_n) \right|$$

$$= \frac{1}{n} \left| \sum_{i \in \mathcal{C}_1} \sum_{j \in [n], j \neq i} \widehat{Z}_{ij} - \sum_{i \in \mathcal{C}_2} \sum_{j \in [n], j \neq i} \widehat{Z}_{ij} \right|, \tag{68}$$

where recall diag(\widehat{Z}) = diag(Z^*) = I_n and (67) holds by symmetry and (a). Suppose that we add an additional constraint such that the row sums of \widehat{Z} are equal, namely, for all $i \in [n]$, $\sum_{j \in \{[n] \setminus i\}} \widehat{Z}_{ij} = r$ for some chosen r, then (68) becomes 0, since the average row sum for each cluster will be the same. Without this constraint, a bound tighter than $\|Z^* - \widehat{Z}\|_1/n$, cf. (65) and the supplementary Fact B.2, is expected for (67) due to cancellations, since $\forall(i,j) \in C_k, k = 1, 2$, we have $(Z_{ij}^* - \widehat{Z}_{ij}) \ge 0$ for the two block-wise sums in (66). Although we do not have a proof, it is our conjecture that for the balanced case where $n_1 = n_2$, the bias term essentially becomes a small order term compared with the variance; cf. the supplementary Facts D.3, B.2 and (64). See also the related Fact 5.2, where the bound is tight.

5.3 Proof of Lemma 4.3

Let $\Lambda = YY^T - \mathbb{E}YY^T$ and $\Psi := \mathbb{Z}\mathbb{Z}^T - \mathbb{E}\mathbb{Z}\mathbb{Z}^T$. It remains to obtain an upper bound for the quantity in (18). The local analysis on (18) relies on the operator norm bound we obtained in Theorem 2.7 as already shown in Lemma 4.2. The key distinction from the global analysis in [39] is, we need to analyze the projection operators in the following sense. Following the proof idea of [15], we define the following projection operator:

$$\mathcal{P}: M \in \mathbb{R}^{n \times n} \to P_2 M + M P_2 - P_2 M P_2 \quad \text{for} \quad P_2 = Z^*/n,$$

$$\mathcal{P}^{\perp}: M \in \mathbb{R}^{n \times n} \to M - \mathcal{P}(M) = (I_n - P_2) M (I_n - P_2).$$

Now suppose that we use the following decomposition to bound for all $\widehat{Z} \in \mathcal{M}_{opt} \cap (Z^* + r_1 B_1^{n \times n})$, where $\mathcal{M}_{opt} := \{Z : Z \succeq 0, \operatorname{diag}(Z) = I_n\} \subset [-1, 1]^{n \times n}$:

$$\langle \Lambda, \widehat{Z} - Z^* \rangle = \langle \Lambda, \mathcal{P}(\widehat{Z} - Z^*) \rangle + \langle \Lambda, \mathcal{P}^{\perp}(\widehat{Z} - Z^*) \rangle =: S_1(\widehat{Z}) + S_2(\widehat{Z}),$$

where $S_1(\widehat{Z}) = \langle \Lambda, \mathcal{P}(\widehat{Z} - Z^*) \rangle = \langle \mathcal{P}(\Lambda), \widehat{Z} - Z^* \rangle,$ and $S_2(\widehat{Z}) := \langle \Lambda, \mathcal{P}^{\perp}(\widehat{Z} - Z^*) \rangle = \langle \Lambda, \mathcal{P}^{\perp}(\widehat{Z}) \rangle \leq \|\Lambda\|_2 \|Z^* - \widehat{Z}\|_1 / n.$

First, we control $S_1(\widehat{Z})$ uniformly for all $\widehat{Z} \in \mathcal{M}_{opt} \cap (Z^* + r_1 B_1^{n \times n})$ in Lemma 5.6. We then obtain an upper bound for $S_2(\widehat{Z})$ in Lemma 5.7, uniformly for all $\widehat{Z} \in \mathcal{M}_{opt}$. Lemmas 5.6 and 5.7 show that the total *noise* contributed by the second term, namely, $S_2(\widehat{Z})$ and a portion of that by $S_1(\widehat{Z})$ is only a fraction of the total signal strength on the LHS of (48). This allows us to prove (55), in combination with the deterministic and high probability bounds in Lemma 4.2.

Lemma 5.6. Suppose all conditions in Theorem 2.7 hold. Then, with probability at least $1 - \frac{c'}{n^2}$, for some absolute constants c', C,

$$\sup_{\widehat{Z} \in \mathcal{M}_{opt} \cap (Z^* + r_1 B_1^{n \times n})} S_1(\widehat{Z}) \leq \frac{2}{3} \xi p \gamma \left\| \widehat{Z} - Z^* \right\|_1 + CC_0(\max_i \left\| H_i^T \right\|_2) \left\lceil \frac{r_1}{2n} \right\rceil \sqrt{\log(2en/\lceil \frac{r_1}{2n} \rceil)} \left(n\Delta + C_0(\max_i \left\| H_i^T \right\|_2) \sqrt{np} \right)$$

Lemma 5.7. Suppose all conditions in Theorem 2.7 hold. Let $\xi \leq w_{\min}^2/16$. Then, with probability at least $1 - \exp(cn)$, for all $\widehat{Z} \in \mathcal{M}_{opt}$,

$$S_2(\widehat{Z}) \leq \|YY^T - \mathbb{E}(YY^T)\|_2 \|Z^* - \widehat{Z}\|_1 / n \leq \frac{1}{6}\xi p\gamma \|\widehat{Z} - Z^*\|_1.$$

We give an outline for Lemma 5.6 in Section 5.4. We prove Lemma 5.7 in the current section. Lemma 4.3 follows immediately from Lemmas 5.6 and 5.7. \Box

Proof of Lemma 5.7. The proof of Lemma 5.7 follows from arguments in [15]. Denote the noise matrix by $\Lambda := YY^T - \mathbb{E}(YY^T)$. Since $\widehat{Z} \succeq 0$, we have by properties of projection,

$$\mathcal{P}^{\perp}(\widehat{Z}) = (I - P_2)(\widehat{Z})(I - P_2) \succeq 0 \text{ and } \mathcal{P}^{\perp}(Z^*) = (I_n - P_2)Z^*(I_n - P_2) = 0,$$

where $P_2 = Z^*/n$. Thus we have for $\operatorname{tr}(\widehat{Z}) = \operatorname{tr}(Z^*) = n$,

$$\begin{aligned} \left\| \mathcal{P}^{\perp}(\widehat{Z}) \right\|_{*} &:= \left\| (I - P_{2})(\widehat{Z})(I - P_{2}) \right\|_{*} = \operatorname{tr}((I - P_{2})(\widehat{Z})(I - P_{2})) \\ &= \operatorname{tr}(Z^{*}(Z^{*} - \widehat{Z}))/n = \left\| Z^{*} - \widehat{Z} \right\|_{1}/n, \end{aligned}$$

where $\|\cdot\|_*$ denotes the nuclear norm, the last inequality holds by Fact 5.2. Thus we have

$$S_2(Z) = \langle \Lambda, \mathcal{P}^{\perp}(\widehat{Z}) \rangle \leq \|\Lambda\|_2 \left\| \mathcal{P}^{\perp}(\widehat{Z}) \right\|_* = \left\| YY^T - \mathbb{E}YY^T \right\|_2 \left\| Z^* - \widehat{Z} \right\|_1 / n.$$

The lemma thus holds in view of Theorem 2.7. $\hfill\square$

5.4 Proof outline for Lemma 5.6

Throughout this section, we consider $\widehat{Z} \in \mathcal{M}_{opt} \cap (Z^* + r_1 B_1^{n \times n})$, where $r_1 \leq 2qn$ for some positive integer $q \in [n]$. As predicted, the decomposition and reduction ideas for the global analysis [39], cf. the supplementary Section D.2, are useful for the local analysis as well. We now briefly introduce them. First, we decompose

$$YY^{T} - \mathbb{E}(YY^{T}) = YY^{T} - \mathbb{E}(Y)\mathbb{E}(Y)^{T} + \mathbb{E}(Y)\mathbb{E}(Y)^{T} - \mathbb{E}(YY^{T})$$

$$= \mathbb{E}(Y)(Y - \mathbb{E}(Y))^{T} + (Y - \mathbb{E}(Y))(\mathbb{E}(Y))^{T} + \widehat{\Sigma}_{Y} - \Sigma_{Y},$$
(69)

where $\widehat{\Sigma}_Y = (Y - \mathbb{E}(Y))(Y - \mathbb{E}(Y))^T$ and

$$\widehat{\Sigma}_Y - \Sigma_Y = (Y - \mathbb{E}(Y))(Y - \mathbb{E}(Y))^T + \mathbb{E}(Y)\mathbb{E}(Y)^T - \mathbb{E}(YY^T) = (I - P_1)(\mathbb{Z}\mathbb{Z}^T - \mathbb{E}(\mathbb{Z}\mathbb{Z}^T))(I - P_1).$$
(70)

Then we have by the triangle inequality,

$$S_{1} := \left\langle \mathcal{P}(\Lambda), \widehat{Z} - Z^{*} \right\rangle \leq \left| \left\langle \mathcal{P}((Y - \mathbb{E}(Y))\mathbb{E}(Y)^{T}), \widehat{Z} - Z^{*} \right\rangle \right| \\ \left| \left\langle \mathcal{P}(\mathbb{E}Y(Y - \mathbb{E}(Y))^{T}), \widehat{Z} - Z^{*} \right\rangle \right| + \left| \left\langle \mathcal{P}(\widehat{\Sigma}_{Y} - \Sigma_{Y}), \widehat{Z} - Z^{*} \right\rangle \right|.$$

We need to first present the following results.

Lemma 5.8. Suppose all conditions in Theorem 2.7 hold. Then, with probability at least $1 - \frac{c}{n^2}$,

$$\sup_{\widehat{Z}\in\mathcal{M}_{opt}\cap(Z^*+r_1B_1^{n\times n})} \left| \left\langle \mathcal{P}(\mathbb{E}(Y)(Y-\mathbb{E}(Y))^T), \widehat{Z}-Z^* \right\rangle \right|$$

$$= \sup_{\widehat{Z}\in\mathcal{M}_{opt}\cap(Z^*+r_1B_1^{n\times n})} \left| \left\langle \mathcal{P}((Y-\mathbb{E}(Y))\mathbb{E}(Y)^T), \widehat{Z}-Z^* \right\rangle \right|$$

$$\leq C_5C_0(\max_i \left\| H_i^T \mu \right\|_2) \sqrt{p\gamma} r_1 \sqrt{\log(2en/\lceil \frac{r_1}{2n} \rceil)}.$$

Lemma 5.9. Under the conditions in Theorem 2.7, we have with probability at least $1 - \frac{c'}{n^2}$,

$$\sup_{\widehat{Z}\in\mathcal{M}_{opt}\cap(Z^*+r_1B_1^{n\times n})} \left| \left\langle \mathcal{P}(\widehat{\Sigma}_Y-\Sigma_Y),\widehat{Z}-Z^*\right\rangle \right| \\ \leq \frac{2}{3}\xi p\gamma \left\| \widehat{Z}-Z^* \right\|_1 + C_{10}(C_0\max_i \|H_i\|_2)^2 \lceil \frac{r_1}{2n} \rceil \left(\sqrt{np\log(2en/\lceil \frac{r_1}{2n}\rceil)} + \sqrt{r_1}\log(2en/\lceil \frac{r_1}{2n}\rceil) \right) \right|.$$

Combining Lemmas 5.8 and 5.9, We have with probability at least $1 - \frac{c}{n^2}$, for $\Delta^2 = p\gamma$,

$$\begin{split} \sup_{\widehat{Z}\in\mathcal{M}_{\text{opt}}\cap(Z^*+r_1B_1^{n\times n})} S_1(\widehat{Z}) &\leq \sup_{\widehat{Z}\in\mathcal{M}_{\text{opt}}\cap(Z^*+r_1B_1^{n\times n})} \left| \left\langle \mathcal{P}(\widehat{\Sigma}_Y - \Sigma_Y), \widehat{Z} - Z^* \right\rangle \right| \\ &+ \sup_{\widehat{Z}\in\mathcal{M}_{\text{opt}}\cap(Z^*+r_1B_1^{n\times n})} 2 \left| \left\langle \mathcal{P}((Y - \mathbb{E}(Y))\mathbb{E}(Y)^T), \widehat{Z} - Z^* \right\rangle \right| \\ &\leq \frac{2}{3}\xi p\gamma \left\| \widehat{Z} - Z^* \right\|_1 + 2C_5C_0(\max_i \|H_i^T\mu\|_2) \lceil \frac{r_1}{2n} \rceil \sqrt{\log(2en/\lceil \frac{r_1}{2n}\rceil)n\Delta} + \\ C_{10}(C_0\max_i \|H_i\|_2)^2 \lceil \frac{r_1}{2n} \rceil \sqrt{\log(2en/\lceil \frac{r_1}{2n}\rceil)} \left(\sqrt{np} + \sqrt{r_1}\sqrt{\log(2en/\lceil \frac{r_1}{2n}\rceil)} \right) \\ &\leq \frac{2}{3}\xi p\gamma \left\| \widehat{Z} - Z^* \right\|_1 + C'C_0(\max_i \|H_i^T\|_2) \lceil \frac{r_1}{2n} \rceil \sqrt{\log(2en/\lceil \frac{r_1}{2n}\rceil)} \left(n\Delta + C_0(\max_i \|H_i^T\|_2) \sqrt{np} \right), \end{split}$$

where in the last inequality, we have by (30), for $r_1 \leq 2qn$ and $q = \lceil \frac{r_1}{2n} \rceil$,

$$C_{0}(\max_{i} \|H_{i}^{T}\|_{2})\sqrt{r_{1}}\sqrt{\log(2en/\lceil\frac{r_{1}}{2n}\rceil)} \leq C_{0}(\max_{i} \|H_{i}^{T}\|_{2})\sqrt{2nq\log(2en/q)} \\ \leq C_{0}(\max_{i} \|H_{i}^{T}\|_{2})n\sqrt{2\log(2e)} = O(n\sqrt{p\gamma}),$$

where the first inequality holds since $\max_{q \in [n]} q \log(2en/q) = n \log(2e)$, given that $q \log(2en/q)$ is a monotonically increasing function of q, for $1 \leq q < n$, and the second inequality holds since $p\gamma = \Omega((C_0 \max_i ||H_i^T||_2)^2)$. Lemma 5.6 thus holds. We prove Lemmas 5.8 and 5.9 in Sections 5.5 and 5.6 respectively. \Box

5.5 Proof of Lemma 5.8

First, we have

$$\langle (Y - \mathbb{E}(Y))\mathbb{E}(Y)^T P_2, \widehat{Z} - Z^* \rangle = \langle P_2 \mathbb{E}(Y)(Y - \mathbb{E}(Y))^T, \widehat{Z} - Z^* \rangle,$$
(71)

$$\langle P_2(Y - \mathbb{E}(Y))\mathbb{E}(Y)^T, \hat{Z} - Z^* \rangle = \langle \mathbb{E}(Y)(Y - \mathbb{E}(Y))^T P_2, \hat{Z} - Z^* \rangle.$$
 (72)

Then we have by the triangle inequality, (72) and (71),

$$\left\langle \mathcal{P}((Y - \mathbb{E}(Y))\mathbb{E}(Y)^T), \widehat{Z} - Z^* \right\rangle = \left| \left\langle \mathcal{P}(\mathbb{E}(Y)(Y - \mathbb{E}(Y))^T), \widehat{Z} - Z^* \right\rangle \right|$$
(73)

$$\leq \left| \left\langle \left(\mathbb{E}(Y)(Y - \mathbb{E}(Y))^T \right) P_2, \widehat{Z} - Z^* \right\rangle - \left\langle P_2 \mathbb{E}(Y)(Y - \mathbb{E}(Y))^T P_2, \widehat{Z} - Z^* \right\rangle \right| \\ + \left| \left\langle P_2 \mathbb{E}(Y)(Y - \mathbb{E}(Y))^T, \widehat{Z} - Z^* \right\rangle \right| =: |T_1| + |T_2|.$$

We will bound the two terms T_1, T_2 in Lemmas 5.10 and 5.11 respectively. We prove Lemma 5.11 in Section 5.7. We prove Lemma 5.10 in the supplementary Section C.1, where we also show that (71) and (72) hold.

Lemma 5.10. Under the conditions in Theorem 2.7, we have for all $\widehat{Z} \in \mathcal{M}_{opt}$, with probability at least $1 - 2 \exp(-cn)$,

$$|T_1| := \left| \left\langle (I - P_2) \mathbb{E}(Y) (Y - \mathbb{E}(Y))^T P_2, \widehat{Z} - Z^* \right\rangle \right| \le C_4 C_0(\max_i \left\| H_i^T \mu \right\|_2) \sqrt{p\gamma} \left\| \widehat{Z} - Z^* \right\|_1.$$

Moreover, when $w_1 = w_2$, $T_1 = 0$.

Lemma 5.11. Suppose all conditions in Theorem 2.7 hold. Suppose $r_1 \leq 2qn$ for some positive integer $1 \leq q < n$. Let $\Delta = \sqrt{p\gamma}$. Then, with probability at least $1 - \frac{c}{n^2}$,

$$\sup_{\widehat{Z}\in\mathcal{M}_{opt}\cap(Z^*+r_1B_1^{n\times n})}\langle P_2\mathbb{E}(Y)(Y-\mathbb{E}(Y))^T,\widehat{Z}-Z^*\rangle \leq C_3C_0(\max_i \left\|H_i^T\mu\right\|_2)\Delta r_1\sqrt{\log(2en/\left\lceil\frac{r_1}{2n}\right\rceil)}.$$

Putting things together, we have an uniform upper bound on $\langle \mathcal{P}(\mathbb{E}(Y)(Y - \mathbb{E}(Y))^T), \hat{Z} - Z^* \rangle$.

5.6 Proof of Lemma 5.9

Upon obtaining the bounds in Lemma 5.8, there are only two unique terms left, which we bound in Lemmas 5.13 and 5.12 respectively. Notice that by the triangle inequality, we have for symmetric matrices $\widehat{\Sigma}_Y - \Sigma_Y = (I - P_1)\Psi(I - P_1)$ and $\widehat{Z} - Z^*$,

$$\langle \mathcal{P}(\widehat{\Sigma}_Y - \Sigma_Y), \widehat{Z} - Z^* \rangle \mid \leq 2 | \langle P_2(\widehat{\Sigma}_Y - \Sigma_Y), \widehat{Z} - Z^* \rangle | + | \langle P_2(\widehat{\Sigma}_Y - \Sigma_Y)P_2, \widehat{Z} - Z^* \rangle |.$$

As the first step, we first obtain in Lemma 5.12 a deterministic bound on $\langle P_2(\widehat{\Sigma}_Y - \Sigma_Y)P_2, \widehat{Z} - Z^* \rangle$. Then, in combination with Theorem 2.7, we obtain the high probability bound as well. We prove Lemma 5.12 in the supplementary Section C.2. Next we state Lemma 5.13, which we prove in Section 5.8.

Lemma 5.12. Denote by $\Psi := (\mathbb{Z}\mathbb{Z}^T - \mathbb{E}(\mathbb{Z}\mathbb{Z}^T))$. Then

$$\left| \langle P_2(\widehat{\Sigma}_Y - \Sigma_Y) P_2, \widehat{Z} - Z^* \rangle \right| \leq (1 + |w_1 - w_2|)^2 \|\Psi\|_2 \|\widehat{Z} - Z^*\|_1 / n.$$

Moreover, under the conditions in Theorem 2.7, we have with probability at least $1 - \exp(cn)$,

$$\sup_{\widehat{Z}\in\mathcal{M}_{opt}} \left| \left\langle P_2(\widehat{\Sigma}_Y - \Sigma_Y) P_2, \widehat{Z} - Z^* \right\rangle \right| \leq \frac{1}{3} \xi p \gamma \left\| \widehat{Z} - Z^* \right\|_1$$

Lemma 5.13. Suppose $r_1 \leq 2qn$ for some positive integer $1 \leq q < n$. Suppose all conditions in Theorem 2.3 hold. Then, with probability at least $1 - \frac{c}{n^2}$,

$$\sup_{\widehat{Z} \in \mathcal{M}_{opt} \cap (Z^* + r_1 B_1^{n \times n})} 2 \left| \left\langle P_2(\widehat{\Sigma}_Y - \Sigma_Y), \widehat{Z} - Z^* \right\rangle \right| \le \frac{1}{3} \xi p \gamma \left\| \widehat{Z} - Z^* \right\|_1 + C_{10}(C_0 \max_i \|H_i\|_2)^2 \lceil \frac{r_1}{2n} \rceil \left(\sqrt{np \log(2en/\lceil \frac{r_1}{2n} \rceil)} + \sqrt{r_1} \log(2en/\lceil \frac{r_1}{2n} \rceil) \right)$$

Lemma 5.9 follows from Lemmas 5.12 and 5.13, since with probability at least $1 - \frac{c}{n^2} - \exp(cn)$,

$$\sup_{\widehat{Z}\in\mathcal{M}_{opt}\cap(Z^{*}+r_{1}B_{1}^{n\times n})} \left| \left\langle \mathcal{P}(\widehat{\Sigma}_{Y}-\Sigma_{Y}),\widehat{Z}-Z^{*}\right\rangle \right|$$

$$\leq \sup_{\widehat{Z}\in\mathcal{M}_{opt}\cap(Z^{*}+r_{1}B_{1}^{n\times n})} \left(2\left| \left\langle P_{2}(\widehat{\Sigma}_{Y}-\Sigma_{Y}),\widehat{Z}-Z^{*}\right\rangle \right| + \left| \left\langle P_{2}(\widehat{\Sigma}_{Y}-\Sigma_{Y})P_{2},\widehat{Z}-Z^{*}\right\rangle \right| \right)$$

$$\leq \frac{2}{3}\xi p\gamma \left\| \widehat{Z}-Z^{*} \right\|_{1} + C_{10}(C_{0}\max_{i}\|H_{i}\|_{2})^{2} \lceil \frac{r_{1}}{2n} \rceil \left(\sqrt{np\log(2en/\lceil \frac{r_{1}}{2n}\rceil)} + \sqrt{r_{1}\log(2en/\lceil \frac{r_{1}}{2n}\rceil)} \right).$$

5.7 Proof of Lemma 5.11

We state in Lemmas 5.14 and 5.15 two reduction steps. Denote by $L_i := \langle \mu^{(1)} - \mu^{(2)}, \mathbb{Z}_i \rangle$ throughout this section. For all $\widehat{Z} \in M_{\text{opt}} \cap (Z^* + r_1 B_1^{n \times n})$, we have $\widehat{w} \in B_{\infty}^n \cap \lceil r_1/(2n) \rceil B_1^n$, where

$$\widehat{w}_j := \frac{1}{2n} \left\| (\widehat{Z} - Z^*)_{\cdot j} \right\|_1 \le 1 \text{ and } \sum_j \widehat{w}_j \le \lceil r_1 / (2n) \rceil,$$
(74)

where B_{∞}^n and B_1^n denote the unit ℓ_{∞} ball and ℓ_1 ball respectively; cf. Fact B.1. Denote by $L_1^* \geq L_2^* \geq \ldots \geq L_n^*$ (resp. $\widehat{w}_1^* \geq \widehat{w}_2^* \geq \ldots \geq \widehat{w}_n^*$) the non-decreasing arrangement of $|L_j|$ (resp. \widehat{w}_j). Lemma 5.14 is analogous to the supplementary Lemma A.3 for the global analysis. We then have the reduction as in Lemma 5.15. We prove Lemmas 5.14 and 5.15, in the supplementary sections C.4 and C.5 respectively.

Lemma 5.14. (Deterministic bounds) Let \hat{w}_j be as in (74). Then

$$\left| \left\langle P_2(\mathbb{E}(Y)(Y - \mathbb{E}(Y))^T), \widehat{Z} - Z^* \right\rangle \right| \le 4w_1 w_2 n \sum_{j=1}^n |L_i| \,\widehat{w}_j + 4w_1 w_2 \left| \sum_{i=1}^n L_i \right| \left\| \widehat{Z} - Z^* \right\|_1 / (2n).$$
(75)

Lemma 5.15. (Reduction to order statistics) Under the settings of Lemma 5.14, we have

$$\sup_{\widehat{Z}\in\mathcal{M}_{opt}\cap(Z^*+r_1B_1^{n\times n})} \left| \left\langle P_2(\mathbb{E}(Y)(Y-\mathbb{E}(Y))^T), \widehat{Z}-Z^* \right\rangle \right| \le 3n \sum_{j=1}^{|r_1/2n|} L_j^*.$$
(76)

It remains to bound the sum on the RHS of (76), namely, $\sum_{j=1}^{\lceil r_1/(2n)\rceil} L_j^*$. To do so, we state in Proposition 5.16 a high probability bound on $\sum_{j=1}^{q} L_j^*$, which holds simultaneously for all $q \in [n]$.

Proposition 5.16. Suppose all conditions in Lemma 2.5 hold. Let random matrix \mathbb{Z} satisfy the conditions as stated therein. Let $1 \leq q \leq n$ denote a positive integer. Then for some absolute constants C_5 , c, we have with probability at least $1 - \frac{c}{n^2}$, simultaneously for all $q \in [n]$,

$$\sum_{j=1}^{q} L_{j}^{*} \leq C_{5} C_{0}(\max_{i} \left\| H_{i}^{T} \mu \right\|_{2}) \sqrt{p \gamma} q \sqrt{\log(2en/q)},$$

where H_i and $\mu = (\mu^{(1)} - \mu^{(2)})/\sqrt{p\gamma}$ are as defined in Theorem 2.7.

We prove Proposition 5.16 in Section C.6. It remains to prove Lemma 5.11. Indeed, we have with probability at least $1 - \frac{c}{n^2}$, by (75) and (76), and Proposition 5.16, for $w_1w_2 \leq 1/4$,

$$\sup_{\widehat{Z}\in\mathcal{M}_{opt}\cap(Z^*+r_1B_1^{n\times n})} \left| \left\langle P_2(\mathbb{E}(Y)(Y-\mathbb{E}(Y))^T), \widehat{Z}-Z^* \right\rangle \right| \\ \leq 3n \sum_{j=1}^{\lceil r_1/(2n)\rceil} L_j^* \leq C_3 C_0(\max_i \left\| H_i^T \mu \right\|_2) \Delta r_1 \sqrt{\log(2en/\lceil \frac{r_1}{2n}\rceil)},$$

where it is understood that we set $q = \lceil \frac{r_1}{2n} \rceil \in [n]$ for some $r_1 > 0$. The lemma is thus proved. \Box

5.8 Proof of Lemma 5.13

First we note that by (70),

$$\langle (\widehat{\Sigma}_Y - \Sigma_Y) P_2, \widehat{Z} - Z^* \rangle = \langle P_2(\widehat{\Sigma}_Y - \Sigma_Y), \widehat{Z} - Z^* \rangle$$

$$= \langle P_2 \Psi, \widehat{Z} - Z^* \rangle - \langle P_2 P_1 \Psi, \widehat{Z} - Z^* \rangle$$

$$+ \langle P_2 P_1 \Psi P_1, \widehat{Z} - Z^* \rangle - \langle P_2 \Psi P_1, \widehat{Z} - Z^* \rangle .$$

$$(77)$$

We state the following deterministic bounds, which we need in the proof of Lemmas 5.12 and 5.18. Moreover, we need Lemma 5.18. We prove Lemmas 5.17 and 5.18 in the supplementary Sections C.7 and C.8 respectively.

Lemma 5.17. (Deterministic bounds) Let $\Psi = \mathbb{Z}\mathbb{Z}^T - \mathbb{E}(\mathbb{Z}\mathbb{Z}^T)$. Then

$$\left| \left\langle P_2 \Psi P_1, \widehat{Z} - Z^* \right\rangle \right| \leq \|\Psi\|_2 \left\| \widehat{Z} - Z^* \right\|_1 / n,$$
(78)

$$\left| \langle P_2 P_1 \Psi P_1, \widehat{Z} - Z^* \rangle \right| \leq |w_1 - w_2| \|\Psi\|_2 \left\| \widehat{Z} - Z^* \right\|_1 / n,$$
(79)

and
$$\left| \left\langle P_2 P_1 \Psi P_1 P_2, \widehat{Z} - Z^* \right\rangle \right| \leq |w_1 - w_2|^2 \|\Psi\|_2 \left\| \widehat{Z} - Z^* \right\|_1 / n.$$
 (80)

Finally, we will show the following bounds:

$$\langle P_2 \Psi P_2, \widehat{Z} - Z^* \rangle \leq \|\Psi\|_2 \left\| \widehat{Z} - Z^* \right\|_1 / n, \tag{81}$$

$$\left| \langle P_2 \Psi P_1 P_2, \widehat{Z} - Z^* \rangle \right| = \left| \langle P_2 P_1 \Psi P_2, \widehat{Z} - Z^* \rangle \right| \le \|\Psi\|_2 |w_1 - w_2| \left\| \widehat{Z} - Z^* \right\|_1 / n.$$
 (82)

When $w_1 = w_2$, the RHS of (79), (80) and (82) all become 0, since $P_2P_1 = 0$ in that case.

Lemma 5.18. (Deterministic bounds) Denote by $\Psi = \mathbb{Z}\mathbb{Z}^T - \mathbb{E}(\mathbb{Z}\mathbb{Z}^T)$ and off $d(\Psi)$ its off-diagonal component, where the diagonal elements are set to be 0. Denote by

$$\begin{aligned} Q_j &:= & \sum_{t \in \mathcal{C}_1, t \neq j} \Psi_{jt} - \sum_{t \in \mathcal{C}_2, t \neq j} \Psi_{jt} = \sum_{t \in \mathcal{C}_1} \operatorname{offd}(\Psi)_{jt} - \sum_{t \in \mathcal{C}_2} \operatorname{offd}(\Psi)_{jt}, \\ S_j &:= & \sum_{t \in [n], t \neq j} \Psi_{jt} = \sum_{t \in [n]} \operatorname{offd}(\Psi)_{jt}. \end{aligned}$$

Let \widehat{w}_i be the same as in (74). Then for $\|\text{diag}(\Psi)\|_{\max} := \max_j |\Psi_{jj}|$,

$$\left| \left\langle P_2 \operatorname{diag}(\Psi), \widehat{Z} - Z^* \right\rangle \right| \leq \left\| \operatorname{diag}(\Psi) \right\|_{\max} \left\| \widehat{Z} - Z^* \right\|_1 / n$$
(83)

$$\left| \left\langle P_2 P_1 \operatorname{diag}(\Psi), \widehat{Z} - Z^* \right\rangle \right| \leq 2 \left| w_1 - w_2 \right| \left\| \operatorname{diag}(\Psi) \right\|_{\max} \left\| \widehat{Z} - Z^* \right\|_1 / n \tag{84}$$

$$\left| \langle P_2 \text{offd}(\Psi), \widehat{Z} - Z^* \rangle \right| \leq 2 \sum_{j \in [n]} |Q_j| \, \widehat{w}_j, \quad and$$

$$(85)$$

$$\langle P_2 P_1 \text{offd}(\Psi), \widehat{Z} - Z^* \rangle \mid \leq 2 |w_1 - w_2| \sum_k |S_k| \cdot \widehat{w}_k.$$
 (86)

Denote by $Q_1^* \ge Q_2^* \ge \ldots \ge Q_n^*$ (resp. $\widehat{w}_1^* \ge \widehat{w}_2^* \ge \ldots \ge \widehat{w}_n^*$) the non-decreasing arrangement of $|Q_i|$ (resp. \widehat{w}_j). Denote by $S_1^* \ge S_2^* \ge \ldots \ge S_n^*$ the non-decreasing arrangement of $|S_i|$ (resp. \widehat{w}_j).

The RHS of (86) and (85) are bounded to be at the same order; cf. Proposition 5.19; Moreover, the RHS of (84) and (86) are both 0 in case we have balanced clusters. We finish the proof of Lemma 5.13 in the supplementary Section C.3. Finally, we state Proposition 5.19, which we prove in the supplementary Section C.9.

Proposition 5.19. Let q denote a positive integer. Let C_4, C_5, c be absolute constants. Under the conditions in Theorem 2.7, we have with probability at least $1 - \frac{c}{n^2}$, simultaneously for all positive $q \in [n]$,

$$\sum_{j=1}^{q} Q_{j}^{*} \leq C_{4}(C_{0} \max_{i} \|H_{i}\|_{2})^{2} q \left(\sqrt{np \log(2en/q)} + \sqrt{nq} \log(2en/q)\right),$$
(87)

$$\sum_{j=1}^{q} S_{j}^{*} \leq C_{5}(C_{0} \max_{i} \|H_{i}\|_{2})^{2} q \left(\sqrt{np \log(2en/q)} + \sqrt{nq} \log(2en/q)\right)$$
(88)

6 Conclusion

Our work is motivated by the two threads of work in combinatorial optimization and in community detection, and particularly by [20] to revisit the max-cut problem (5) and its convex relaxations. Centering the data matrix X plays a key role in the statistical analysis and in understanding the roles of sample size lower bounds for partial recovery of the clusters [39]. As in [39], we focus on the sample size lower bound and show that a full range of tradeoffs between the sample size and the number of features are feasible so long as s^2 is lower bounded in the sense of (23) and (30). In the present work, we further elaborate upon the roles of Theorem 2.7 in the local analysis of the SDP in Section 5.3, and the bias and variance tradeoffs. More importantly, we prove that the misclassification error decays exponentially with respect to the SNR s^2 in the present paper. We elaborated upon the connections and differences between our work and those in [15, 18].

A Preliminary results

We present preliminary results for the global and local analyses in this section. These results are either proved in [39], or follow from results therein.

Remarks on covariance structures. Lemma 2.5 characterizes the two-group design matrix variance and covariance structures to be considered in Theorems 1.4 and 2.3. It is understood that when H_i is a symmetric square matrix, it can be taken as the unique square root of the corresponding covariance matrix. Expressions in (27), (19), and (20) are compatible with each other [39]. More explicitly, we use the construction in Definition 2.2 to generate $\mathbb{Z}_j \in \mathbb{R}^p$ with certain covariance structures. When we allow each population to have distinct covariance structures following Theorem 2.3, we have for some universal constant C, and for all $j \in C_i$,

$$\|\mathbb{Z}_{j}\|_{\psi_{2}} := \sup_{h \in \mathbb{S}^{p-1}} \| \langle \mathbb{Z}_{j}, h \rangle \|_{\psi_{2}} \leq \|W_{j}\|_{\psi_{2}} \|H_{i}\|_{2} \leq CC_{0} \max_{i} \|H_{i}\|_{2} \text{ since}$$
(89)

$$\forall h \in \mathbb{S}^{p-1}, \quad \| \langle \mathbb{Z}_j, h \rangle \|_{\psi_2} = \| \langle H_i W_j, h \rangle \|_{\psi_2} \le \| W_j \|_{\psi_2} \| H_i^T h \|_2, \tag{90}$$

where $||W_j||_{\psi_2} \leq CC_0$ by definition of (28). Without loss of generality (w.l.o.g.), one may assume that $C_0 = 1$, as one can adjust H_i to control the upper bound in (89) through $||H_i||_2$.

Theorem A.1. (Hanson-Wright inequality for anisotropic random vectors) [39] Let H_1, \ldots, H_n be deterministic $p \times m$ matrices, where we assume that $m \ge p$. Let $\mathbb{Z}_1^T, \ldots, \mathbb{Z}_n^T \in \mathbb{R}^p$ be row vectors of \mathbb{Z} . We generate \mathbb{Z} according to Definition 2.2. Then we have for t > 0, for any $A = (a_{ij}) \in \mathbb{R}^{n \times n}$,

$$\mathbb{P}\left(\left|\sum_{i=1}^{n}\sum_{j\neq i}^{n} \langle \mathbb{Z}_{i}, \mathbb{Z}_{j} \rangle a_{ij}\right| > t\right) \\
\leq 2\exp\left(-c\min\left(\frac{t^{2}}{(C_{0}\max_{i}\|H_{i}\|_{2})^{4}p\|A\|_{F}^{2}}, \frac{t}{(C_{0}\max_{i}\|H_{i}\|_{2})^{2}\|A\|_{2}}\right)\right), \quad (91)$$

where $\max_{i} \|\mathbb{Z}_{i}\|_{\psi_{2}} \leq CC_{0} \max_{i} \|H_{i}\|_{2}$ in the sense of (89).

Denote by

$$\mu := \frac{\mu^{(1)} - \mu^{(2)}}{\left\| \mu^{(1)} - \mu^{(2)} \right\|_2} = \frac{\mu^{(1)} - \mu^{(2)}}{\sqrt{p\gamma}} \in \mathbb{S}^{p-1}.$$
(92)

Lemma A.2. (Projection for anisotropic sub-gaussian random vectors) [39] Suppose all conditions in Theorem 2.3 hold. Let $R_i = H_i^T$. Let μ be as defined in (92). Then

$$\| \langle \mathbb{Z}_{j}, \mu \rangle \|_{\psi_{2}}^{2} \leq C_{0}^{2} \| \langle \mathbb{Z}_{j}, \mu \rangle \|_{L_{2}}^{2} := C_{0}^{2} \mu^{T} \operatorname{Cov}(\mathbb{Z}_{j}) \mu$$

$$where \ \mu^{T} \operatorname{Cov}(\mathbb{Z}_{j}) \mu = \mu^{T} H_{i} H_{i}^{T} \mu = \| R_{i} \mu \|_{2}^{2} \quad for \ each \ j \in \mathcal{C}_{i}, i = 1, 2.$$

$$(93)$$

Thus for any t > 0, for some absolute constants c, c', we have for each $r = (r_1, \ldots, r_n) \in \mathbb{S}^{n-1}$ and $u = (u_1, \ldots, u_n) \in \{-1, 1\}^n$ the following tail bounds:

$$\mathbb{P}\left(\left|\sum_{i=1}^{n} r_i \left\langle \mathbb{Z}_i, \mu \right\rangle \right| \ge t\right) \le 2 \exp\left(-\frac{ct^2}{(C_0 \max_i \|R_i\mu\|_2)^2}\right), \quad and \tag{94}$$

$$\mathbb{P}\left(\left|\sum_{i=1}^{n} u_i \left\langle \mathbb{Z}_i, \mu \right\rangle \right| \ge t\right) \le 2 \exp\left(-\frac{c't^2}{n(C_0 \max_i \|R_i\mu\|_2)^2}\right).$$
(95)

First, we state Lemma A.3. Upon obtaining (13), Lemma A.3 is deterministic and does not depend on the covariance structure of \mathbb{Z} . See Lemma 5.4 in [39].

Lemma A.3. (Reduction: a deterministic comparison lemma) [39] Let X be as in Lemma 2.5. Let Y be as in Definition 1.1. Let $\mathbb{Z}_j, j \in [n]$ be row vectors of $X - \mathbb{E}X$ and $\hat{\mu}_n$ be as defined in (7). For $x_i \in \{-1, 1\}$,

$$\sum_{i=1}^{n} x_i \left\langle Y_i - \mathbb{E}Y_i, \mu^{(1)} - \mu^{(2)} \right\rangle \leq \frac{2(n-1)}{n} \sum_{i=1}^{n} \left| \left\langle \mathbb{Z}_i, \mu^{(1)} - \mu^{(2)} \right\rangle \right|.$$
(96)

Corollary A.4. Let X be as in Lemma 2.5. Let Y be as in Definition 1.1. Under the conditions in Theorem 2.7, we have with probability at least $1 - \exp(-c'n)$,

$$\left| \left\langle \mu^{(1)} - \mu^{(2)} / \sqrt{p\gamma}, \sum_{j \in C_1} (Y_j - \mathbb{E}(Y_j)) - \sum_{j \in C_2} (Y_j - \mathbb{E}(Y_j)) \right\rangle \\ \leq 2 \sum_{i=1}^n \left| \left\langle \frac{\mu^{(1)} - \mu^{(2)}}{\sqrt{p\gamma}}, \mathbb{Z}_i \right\rangle \right| \leq C C_0 n \max_i \|R_i \mu\|_2.$$

Proof. Now for $t = nCC_0 \max_i ||R_i\mu||_2/2$

ī.

$$\mathbb{P}\left(\max_{u\in\{-1,1\}^n}\sum_{j=1}^n u_i\langle\mu,\mathbb{Z}_j\rangle \geq t\right) \leq 2^n 2\exp\left(-\frac{c't^2}{n(C_0\max_i\|R_i\mu\|_2)^2}\right) \leq 2\exp(-cn),$$

where the last step follows from (95). And hence with probability at least $1 - \exp(-c'n)$, we have by Lemma A.3,

$$\left| \left\langle \mu^{(1)} - \mu^{(2)} / \sqrt{p\gamma}, \sum_{j \in C_1} (Y_j - \mathbb{E}(Y_j)) - \sum_{j \in C_2} (Y_j - \mathbb{E}(Y_j)) \right\rangle \right.$$

$$\leq \max_{x \in \{-1,1\}^n} \sum_{i=1}^n x_i \left\langle \mu, Y_j - \mathbb{E}(Y_j) \right\rangle$$

$$\leq 2 \max_{u \in \{-1,1\}^n} \sum_{j=1}^n u_j \left\langle \mu, \mathbb{Z}_j \right\rangle \leq nCC_0 \max_i ||R_i\mu||_2.$$

B Additional proofs for Theorem 2.3

First we prove Fact 5.2.

Proof of Fact 5.2. One can check that $Z_{ij}^* \cdot ((Z^* - \widehat{Z})_{ij}) \ge 0$ for all i, j. On the diagonal blocks in (56), we have $(Z^* - \widehat{Z})_{ij} \ge 0$, since $Z_{ij}^* = 1$; on the off-diagonal blocks, we have $(Z^* - \widehat{Z})_{ij} \le 0$, since $Z_{ij}^* = -1$. Now $\left\|\widehat{Z} - Z^*\right\|_1 \le 2n(n-1)$ by Fact 5.1. \Box

Next we state Facts B.1 and B.2.

Fact B.1. Let Z_{j} represent the j^{th} column of Z. Let $\widehat{Z} \in \mathcal{M}_{opt}$ and $\left\|\widehat{Z} - Z^*\right\|_1 \leq r_1 \leq 2qn$, where $q = \left\lceil \frac{r_1}{2n} \right\rceil < n$ is a positive integer and Z^* is as defined in (56). Denote by

$$\widehat{w}_{j} = \frac{1}{2n} \sum_{i \in [n]} \left| (\widehat{Z} - Z^{*})_{ij} \right| := \left\| (\widehat{Z} - Z^{*})_{\cdot j} \right\|_{1} / (2n) < 1$$
(97)

thus we have
$$\sum_{j=1}^{n} \widehat{w}_j := \left\| \widehat{Z} - Z^* \right\|_1 / (2n) \le q < n.$$
 (98)

Hence the weight vector $\widehat{w} = (\widehat{w}_1, \dots, \widehat{w}_n) \in B_{\infty}^n \cap qB_1^n$, where $B_{\infty}^n = \{y \in \mathbb{R}^n : |y_j| \leq 1\}$ and $B_1^n = \{y \in \mathbb{R}^n : \sum_j |y_j| \leq 1\}$ denote the unit ℓ_{∞} ball and ℓ_1 ball respectively. Moreover, we have for \widehat{w}_j as in (97),

$$\forall j \in \mathcal{C}_1, \ \widehat{w}_j = -\frac{1}{2n} \big(\sum_{i \in \mathcal{C}_1} (\widehat{Z} - Z^*)_{ij} - \sum_{i \in \mathcal{C}_2} (\widehat{Z} - Z^*)_{ij} \big) \quad and \\ \forall j \in \mathcal{C}_2, \ \widehat{w}_j = \frac{1}{2n} \big(\sum_{i \in \mathcal{C}_1} (\widehat{Z} - Z^*)_{ij} - \sum_{i \in \mathcal{C}_2} (\widehat{Z} - Z^*)_{ij} \big).$$

Proof. By definition of Z^* , we have $\forall j \in \mathcal{C}_1$,

$$\forall i \in \mathcal{C}_2, \quad (\widehat{Z} - Z^*)_{ij} \ge 0 \quad \text{and} \quad \forall i \in \mathcal{C}_1, \ (\widehat{Z} - Z^*)_{ij} \le 0,$$

and hence

$$\widehat{w}_j = \frac{1}{2n} \Big(\sum_{i \in \mathcal{C}_2} (\widehat{Z} - Z^*)_{ij} - \sum_{i \in \mathcal{C}_1} (\widehat{Z} - Z^*)_{ij} \Big) = \left\| (\widehat{Z} - Z^*)_{\cdot j} \right\|_1 / (2n) < 1.$$

Similarly, we have $\forall j \in \mathcal{C}_2$,

$$\forall i \in \mathcal{C}_1, \quad (\widehat{Z} - Z^*)_{ij} \ge 0 \text{ and } \quad \forall i \in \mathcal{C}_2, (\widehat{Z} - Z^*)_{ij} \le 0,$$

and hence

$$\widehat{w}_j = \frac{1}{2n} \Big(\sum_{i \in \mathcal{C}_1} (\widehat{Z} - Z^*)_{ij} - \sum_{i \in \mathcal{C}_2} (\widehat{Z} - Z^*)_{ij} \Big) = \left\| (\widehat{Z} - Z^*)_{\cdot j} \right\|_1 / (2n) < 1.$$

Fact B.2. We have by symmetry of $\hat{Z} - Z^*$, under the settings in Fact B.1,

$$\frac{1}{2n} \left| \mathbf{1}_n^T (\widehat{Z} - Z^*) u_2 \right| = \left| \frac{1}{2n} \sum_{i=1}^n \left(\sum_{j \in \mathcal{C}_1} (\widehat{Z} - Z^*)_{ij} - \sum_{j \in \mathcal{C}_2} (\widehat{Z} - Z^*)_{ij} \right) \right| \\ \leq \sum_{i=1}^n |\widehat{w}_i| = \frac{1}{2n} \left\| (\widehat{Z} - Z^*) \right\|_1 = \frac{1}{2n} \sum_j \left\| (\widehat{Z} - Z^*)_{j\cdot} \right\|_1.$$

C Technical proofs for Section 5

C.1 Proof outline of Lemma 5.10

Recall by (13),

$$\mathbb{E}(Y_i) = \begin{cases} w_2(\mu^{(1)} - \mu^{(2)}) & \text{if } i \in \mathcal{C}_1, \\ w_1(\mu^{(2)} - \mu^{(1)}) & \text{if } i \in \mathcal{C}_2. \end{cases}$$

First, we show that (71) holds. To see this, we have

$$\langle (Y - \mathbb{E}(Y))\mathbb{E}(Y)^T P_2, \widehat{Z} - Z^* \rangle = \operatorname{tr}((\widehat{Z} - Z^*)(Y - \mathbb{E}(Y))\mathbb{E}(Y)^T P_2)$$

= $\langle P_2 \mathbb{E}(Y)(Y - \mathbb{E}(Y))^T, \widehat{Z} - Z^* \rangle.$

Similarly, (72) holds since

$$\langle P_2(Y - \mathbb{E}(Y))\mathbb{E}(Y)^T, \widehat{Z} - Z^* \rangle = \operatorname{tr}((\widehat{Z} - Z^*)P_2(Y - \mathbb{E}(Y))\mathbb{E}(Y)^T)$$

= $\langle \mathbb{E}(Y)(Y - \mathbb{E}(Y))^T P_2, \widehat{Z} - Z^* \rangle.$ (99)

Denote by $V = (\mathbb{E}(Y)(Y - \mathbb{E}(Y))^T)u_2$, which is a vector with coordinates V_1, \ldots, V_n such that

$$\begin{aligned} \forall i \in \mathcal{C}_1, \quad V_i &= -w_2 \langle \mu^{(2)} - \mu^{(1)}, \sum_{j \in C_1} (Y_j - \mathbb{E}(Y_j)) - \sum_{j \in C_2} (Y_j - \mathbb{E}(Y_j)) \rangle \\ \forall i \in \mathcal{C}_2, \quad V_i &= w_1 \langle \mu^{(2)} - \mu^{(1)}, \sum_{j \in C_1} (Y_j - \mathbb{E}(Y_j)) - \sum_{j \in C_2} (Y_j - \mathbb{E}(Y_j)) \rangle , \end{aligned}$$

where $Y_j \in \mathbb{R}^p$ represents the j^{th} row vector of matrix Y. Similarly, we have by (99),

$$\langle P_{2}(Y - \mathbb{E}(Y))\mathbb{E}(Y)^{T}, \widehat{Z} - Z^{*} \rangle = \operatorname{tr}((\widehat{Z} - Z^{*})\mathbb{E}(Y)(Y - \mathbb{E}(Y))^{T}P_{2})$$

$$= \operatorname{tr}(P_{2}(\widehat{Z} - Z^{*})\mathbb{E}(Y)(Y - \mathbb{E}(Y))^{T}) = \operatorname{tr}(\frac{u_{2}u_{2}^{T}}{n}(\widehat{Z} - Z^{*})(\mathbb{E}(Y)(Y - \mathbb{E}(Y))^{T}))$$

$$= \operatorname{tr}(\frac{u_{2}^{T}}{n}(\widehat{Z} - Z^{*})(\mathbb{E}(Y)(Y - \mathbb{E}(Y))^{T})u_{2}) = \langle \frac{1}{n} \big(\sum_{i \in \mathcal{C}_{1}} (\widehat{Z} - Z^{*})_{i \cdot} - \sum_{i \in \mathcal{C}_{2}} (\widehat{Z} - Z^{*})_{i \cdot} \big), V \rangle$$

$$= 2\sum_{k} V_{k} \cdot \frac{1}{2n} \big(\sum_{i \in \mathcal{C}_{1}} (\widehat{Z} - Z^{*})_{ik} - \sum_{i \in \mathcal{C}_{2}} (\widehat{Z} - Z^{*})_{ik} \big) =: S.$$

Now

$$\begin{aligned} \forall k \in \mathcal{C}_{1}, \quad \frac{1}{2n} \Big(\sum_{i \in \mathcal{C}_{1}} (\widehat{Z} - Z^{*})_{ik} - \sum_{i \in \mathcal{C}_{2}} (\widehat{Z} - Z^{*})_{ik} \Big) V_{k} \\ &= w_{2} \widehat{w}_{k} \left\langle \mu^{(2)} - \mu^{(1)}, \sum_{j \in C_{1}} (Y_{j} - \mathbb{E}(Y_{j})) - \sum_{j \in C_{2}} (Y_{j} - \mathbb{E}(Y_{j})) \right\rangle, \\ &\text{and} \quad \forall k \in \mathcal{C}_{2}, \quad \frac{1}{2n} \Big(\sum_{i \in \mathcal{C}_{1}} (\widehat{Z} - Z^{*})_{ik} - \sum_{i \in \mathcal{C}_{2}} (\widehat{Z} - Z^{*})_{ik} \Big) V_{k} \\ &= w_{1} \widehat{w}_{k} \left\langle \mu^{(2)} - \mu^{(1)}, \sum_{j \in C_{1}} (Y_{j} - \mathbb{E}(Y_{j})) - \sum_{j \in C_{2}} (Y_{j} - \mathbb{E}(Y_{j})) \right\rangle. \end{aligned}$$

Hence by (97) and Fact B.1,

$$S = 2C_{S} \cdot \langle \mu^{(2)} - \mu^{(1)}, \sum_{j \in C_{1}} (Y_{j} - \mathbb{E}(Y_{j})) - \sum_{j \in C_{2}} (Y_{j} - \mathbb{E}(Y_{j})) \rangle, \qquad (100)$$

where
$$C_{S} := w_{2} \sum_{k \in C_{1}} \widehat{w}_{k} + w_{1} \sum_{k \in C_{2}} \widehat{w}_{k} \leq \sum_{k \in [n]} \widehat{w}_{k} = \left\| \widehat{Z} - Z^{*} \right\|_{1} / (2n).$$

Next we define

$$W := 2u_2^T \mathbb{E}(Y)(Y - \mathbb{E}(Y))^T u_2/n = 2u_2^T V/n = \frac{2}{n} (\sum_{i \in \mathcal{C}_1} V_i - \sum_{i \in \mathcal{C}_2} V_i)$$
(101)
$$= \langle \mu^{(1)} - \mu^{(2)}, \sum_{j \in \mathcal{C}_1} (Y_j - \mathbb{E}Y_j) - \sum_{j \in \mathcal{C}_2} (Y_j - \mathbb{E}Y_j) \rangle 4w_1 w_2.$$

Then

$$\langle P_2 \mathbb{E}(Y)(Y - \mathbb{E}(Y))^T P_2, \widehat{Z} - Z^* \rangle = \operatorname{tr}(P_2(\widehat{Z} - Z^*)P_2 \mathbb{E}(Y)(Y - \mathbb{E}(Y))^T)$$

$$= \frac{1}{2n} u_2^T (\widehat{Z} - Z^*) u_2 (2u_2^T \mathbb{E}(Y)(Y - \mathbb{E}(Y))^T u_2/n) =: -W \sum_k \widehat{w}_k, \qquad (102)$$

where (102) holds by (101) and Fact 5.2, since for $P_2 = u_2 u_2^T / n = Z^* / n$,

$$\frac{u_2^T}{2n}(\widehat{Z} - Z^*)u_2 = \operatorname{tr}(P_2(\widehat{Z} - Z^*))/2 = -\left\|\widehat{Z} - Z^*\right\|_1/(2n) = -\sum_k \widehat{w}_k.$$

Putting things together, we have by (98), (100) and (102),

$$T_{1} := \langle \mathbb{E}(Y)(Y - \mathbb{E}(Y))^{T}P_{2}, \widehat{Z} - Z^{*} \rangle - \langle P_{2}\mathbb{E}(Y)(Y - \mathbb{E}(Y))^{T}P_{2}, \widehat{Z} - Z^{*} \rangle$$

$$= S + W \sum_{k} \widehat{w}_{k} = 2 \langle \mu^{(2)} - \mu^{(1)}, \sum_{j \in C_{1}} (Y_{j} - \mathbb{E}(Y_{j})) - \sum_{j \in C_{2}} (Y_{j} - \mathbb{E}(Y_{j})) \rangle \cdot$$

$$(w_{2} \sum_{k \in C_{1}} \widehat{w}_{k} + w_{1} \sum_{k \in C_{2}} \widehat{w}_{k} - 2w_{1}w_{2} \sum_{k \in [n]} \widehat{w}_{k}),$$

where

$$\begin{aligned} w_2 \sum_{k \in \mathcal{C}_1} \widehat{w}_k + w_1 \sum_{k \in \mathcal{C}_2} \widehat{w}_k - 2w_1 w_2 \sum_{k \in [n]} \widehat{w}_k \\ &\leq \left\| u_2 \sum_{k \in \mathcal{C}_1} \widehat{w}_k + w_1 \sum_{k \in \mathcal{C}_2} \widehat{w}_k \right\| \lor \left| 2w_1 w_2 \sum_{k \in [n]} \widehat{w}_k \right| \\ &\leq \left\| \widehat{Z} - Z^* \right\|_1 / (2n) \le q < n. \end{aligned}$$

Now, for $w_1 = w_2 = 1/2$, the above sum is 0 and hence $T_1 = 0$. Let $r_1 \leq 2qn$. Then, by Corollary A.4, we have with probability at least $1 - 2\exp(-cn)$, for all $\widehat{Z} \in \mathcal{M}_{opt} \cap (Z^* + r_1B_1^{n \times n})$,

$$\begin{aligned} |T_{1}| &= \left| \left\langle \left(\mathbb{E}(Y)(Y - \mathbb{E}(Y))^{T} \right) P_{2}, \widehat{Z} - Z^{*} \right\rangle - \left\langle P_{2}\mathbb{E}(Y)(Y - \mathbb{E}(Y))^{T} P_{2}, \widehat{Z} - Z^{*} \right\rangle \right| \\ &\leq \left| \left\langle \mu^{(2)} - \mu^{(1)}, \sum_{j \in C_{1}} (Y_{j} - \mathbb{E}(Y_{j})) - \sum_{j \in C_{2}} (Y_{j} - \mathbb{E}(Y_{j})) \right\rangle \right| \left\| \widehat{Z} - Z^{*} \right\|_{1} / n \\ &\leq C_{4}C_{0}(\max_{i} \|R_{i}\mu\|_{2}) \sqrt{p\gamma} \left\| \widehat{Z} - Z^{*} \right\|_{1} \leq C_{4}C_{0}(\max_{i} \|R_{i}\mu\|_{2}) r_{1}\sqrt{p\gamma}. \end{aligned}$$

C.2 Proof of Lemma 5.12

Theorem C.1. [39] In the settings of Theorem 2.7, we have with probability at least $1-2\exp(-c_6n)$,

$$\left\|\widehat{\Sigma}_{Y} - \mathbb{E}\widehat{\Sigma}_{Y}\right\|_{2} \leq \left\|\mathbb{Z}\mathbb{Z}^{T} - \mathbb{E}\mathbb{Z}\mathbb{Z}^{T}\right\|_{2} \leq C_{2}C_{0}^{2}\max_{j}\left\|\operatorname{Cov}(Z_{j})\right\|_{2}(\sqrt{np} \lor n).$$
(103)

Proof of Lemma 5.12. First,

$$\langle P_2(\widehat{\Sigma}_Y - \Sigma_Y) P_2, \widehat{Z} - Z^* \rangle = \langle P_2(I - P_1) \Psi(I - P_1) P_2, \widehat{Z} - Z^* \rangle$$

$$= \langle P_2 \Psi P_2, \widehat{Z} - Z^* \rangle + \langle P_2 P_1 \Psi P_1 P_2, \widehat{Z} - Z^* \rangle$$

$$- \langle P_2 P_1 \Psi P_2, \widehat{Z} - Z^* \rangle - \langle P_2 \Psi P_1 P_2, \widehat{Z} - Z^* \rangle .$$

$$(104)$$

Hence the deterministic statement in Lemma 5.12 holds in view of (104), (80), (81), and (82),

$$\langle P_{2}(\widehat{\Sigma}_{Y} - \Sigma_{Y})P_{2}, \widehat{Z} - Z^{*} \rangle | \leq | \langle P_{2}\Psi P_{2}, \widehat{Z} - Z^{*} \rangle | + | \langle P_{2}P_{1}\Psi P_{1}P_{2}, \widehat{Z} - Z^{*} \rangle | + \\ 2 | \langle P_{2}P_{1}\Psi P_{2}, \widehat{Z} - Z^{*} \rangle | \\ \leq \frac{1}{n} ||\widehat{Z} - Z^{*}||_{1} ||\Psi||_{2} (1 + (w_{1} - w_{2})^{2} + 2 |w_{2} - w_{1}|) \\ \leq \frac{1}{n} ||\widehat{Z} - Z^{*}||_{1} ||\Psi||_{2} (1 + |w_{1} - w_{2}|)^{2}.$$

Finally, we have by Theorem C.1, with probability at least $1 - \exp(c'n)$,

$$\|\Psi\|_{2} = \left\|\mathbb{Z}\mathbb{Z}^{T} - \mathbb{E}\mathbb{Z}\mathbb{Z}^{T}\right\|_{2} \leq C_{2}C_{0}^{2} \max_{j} \|\mathsf{Cov}(\mathbb{Z}_{j})\|_{2} \left(\sqrt{np} \lor n\right) \leq \frac{1}{12}\xi np\gamma,$$

where the last step holds so long as (30) holds for sufficiently large constants C, C_1 and $\xi \leq \frac{w_{\min}^2}{16}$. Thus, we have with probability at least $1 - \exp(c'n)$, for all $\widehat{Z} \in \mathcal{M}_{opt}$,

$$|\langle P_2(\widehat{\Sigma}_Y - \Sigma_Y)P_2, \widehat{Z} - Z^*\rangle| \le 8 \|\Psi\|_2 \|\widehat{Z} - Z^*\|_1/(2n) \le \frac{1}{3}\xi p\gamma \|\widehat{Z} - Z^*\|_1.$$

The lemma is thus proved. \Box

C.3 Proof of Lemma 5.13

Corollary C.2. Under the settings in Proposition 5.19, we have with probability at least $1 - \frac{c}{n^2}$,

$$\sup_{\widehat{Z}\in\mathcal{M}_{opt}\cap(Z^*+r_1B_1^{n\times n})} \left| \left\langle P_2 \text{offd}(\Psi), \widehat{Z}-Z^* \right\rangle \right| \le 4 \sum_{j=1}^{\lceil r_1/(2n)\rceil} Q_j^*$$

$$\le 4C_4 (C_0 \max_i \|H_i\|_2)^2 \lceil \frac{r_1}{2n} \rceil \left(\sqrt{np \log(2en/\lceil \frac{r_1}{2n}\rceil)} + \sqrt{r_1} \log(2en/\lceil \frac{r_1}{2n}\rceil) \right),$$
(105)

and

$$\sup_{\widehat{Z} \in \mathcal{M}_{opt} \cap (Z^* + r_1 B_1^{n \times n})} \left| \langle P_2 P_1 \text{offd}(\Psi), \widehat{Z} - Z^* \rangle \right| \le 4 |w_1 - w_2| \sum_{j=1}^{\lceil r_1/(2n) \rceil} S_j^*$$
(106)
$$\le 4 |w_1 - w_2| C_5 (C_0 \max_i ||H_i||_2)^2 \lceil \frac{r_1}{2n} \rceil \left(\sqrt{np \log(2en/\lceil \frac{r_1}{2n} \rceil)} + \sqrt{r_1} \log(2en/\lceil \frac{r_1}{2n} \rceil) \right).$$

Proof of Lemma 5.13. Let $\widehat{Z} \in \mathcal{M}_{opt} \cap (Z^* + r_1 B_1^{n \times n})$, where $r_1 \leq 2qn$ for some positive integer $1 \leq q \leq n$. With probability at least $1 - \exp(c_4 n)$, by Lemma 5.17 (cf. (78) and (79)), we have for all $\widehat{Z} \in \mathcal{M}_{opt}$,

$$W_{1} := \left| \left\langle P_{2}\Psi P_{1}, \widehat{Z} - Z^{*} \right\rangle \right| + \left| \left\langle P_{2}P_{1}\Psi P_{1}, \widehat{Z} - Z^{*} \right\rangle \right| \\ \leq \left(1 + |w_{1} - w_{2}|\right) \|\Psi\|_{2} \frac{1}{n} \left\| \widehat{Z} - Z^{*} \right\|_{1}.$$

Thus we have by the triangle inequality, by (77),

$$\left| \left\langle P_2(\widehat{\Sigma}_Y - \Sigma_Y), \widehat{Z} - Z^* \right\rangle \right| \leq \left| \left\langle P_2 \Psi P_1, \widehat{Z} - Z^* \right\rangle \right| + \left| \left\langle P_2 P_1 \Psi P_1, \widehat{Z} - Z^* \right\rangle \right| + \left| \left\langle P_2 \Psi, \widehat{Z} - Z^* \right\rangle \right| + \left| \left\langle P_2 P_1 \Psi, \widehat{Z} - Z^* \right\rangle \right| =: W_1 + W_2,$$

where we decompose

$$W_{2} := \left| \left\langle P_{2}\Psi, \widehat{Z} - Z^{*} \right\rangle \right| + \left| \left\langle P_{2}P_{1}\Psi, \widehat{Z} - Z^{*} \right\rangle \right| \leq \left| \left\langle P_{2}\operatorname{diag}(\Psi), \widehat{Z} - Z^{*} \right\rangle \right| + \left| \left\langle P_{2}P_{1}\operatorname{diag}(\Psi), \widehat{Z} - Z^{*} \right\rangle \right| \quad (W_{2}(\operatorname{diag})) + \left| \left\langle P_{2}\operatorname{offd}(\Psi), \widehat{Z} - Z^{*} \right\rangle \right| + \left| \left\langle P_{2}P_{1}\operatorname{offd}(\Psi), \widehat{Z} - Z^{*} \right\rangle \right| \quad (W_{2}(\operatorname{offd})).$$

Then, we have for all $\widehat{Z} \in Z^* + r_1 B_1^{n \times n}$, by (79), (78), (83), and (84),

$$W_{2}(\text{offd}) + W_{1} = \left| \langle P_{2} \text{diag}(\Psi), \widehat{Z} - Z^{*} \rangle \right| + \left| \langle P_{2} P_{1} \text{diag}(\Psi), \widehat{Z} - Z^{*} \rangle \right| + \left| \langle P_{2} \Psi P_{1}, \widehat{Z} - Z^{*} \rangle \right| \\ \leq (1 + |w_{1} - w_{2}|) \max_{k} (|\Psi_{kk}| + ||\Psi||_{2}) \left\| \widehat{Z} - Z^{*} \right\|_{1} / n \\ \leq 4 ||\Psi||_{2} \left\| \widehat{Z} - Z^{*} \right\|_{1} / n,$$
(107)

and moreover, we have by (105), and (106),

$$\sup_{\widehat{Z}\in\mathcal{M}_{opt}\cap(Z^{*}+r_{1}B_{1}^{n\times n})} W_{2}(offd)$$

$$:= \sup_{\widehat{Z}\in\mathcal{M}_{opt}\cap(Z^{*}+r_{1}B_{1}^{n\times n})} \left(\left| \langle P_{2}offd(\Psi),\widehat{Z}-Z^{*} \rangle \right| + \left| \langle P_{2}P_{1}offd(\Psi),\widehat{Z}-Z^{*} \rangle \right| \right)$$

$$\leq 4 \sup_{\widehat{w}\in B_{\infty}^{n}\cap\lceil r_{1}/(2n)\rceil} \left(\sum_{j=1}^{\lceil r_{1}/(2n)\rceil} Q_{j}^{*} + |w_{1}-w_{2}| \sum_{j=1}^{\lceil r_{1}/(2n)\rceil} S_{j}^{*} \right).$$
(108)

Thus we have for $\Psi = \mathbb{Z}\mathbb{Z}^T - \mathbb{E}(\mathbb{Z}\mathbb{Z}^T)$, by (107) and (108), with probability at least $1 - c_2/n^2$, and $q = \lceil r_1/(2n) \rceil$,

$$\sup_{\widehat{Z}\in\mathcal{M}_{opt}\cap(Z^{*}+r_{1}B_{1}^{n\times n})} \left| \left\langle P_{2}(\widehat{\Sigma}_{Y}-\Sigma_{Y}),\widehat{Z}-Z^{*}\right\rangle \right| \leq \sup_{\widehat{Z}\in\mathcal{M}_{opt}\cap(Z^{*}+r_{1}B_{1}^{n\times n})} 4 \|\Psi\|_{2} \|\widehat{Z}-Z^{*}\|_{1} / n$$

$$+ 4 \sup_{\widehat{w}\in B_{\infty}^{n}\cap\lceil r_{1}/(2n)\rceil B_{1}^{n}} \left(\sum_{j=1}^{\lceil r_{1}/(2n)\rceil} Q_{j}^{*} + |w_{1}-w_{2}| \sum_{j=1}^{\lceil r_{1}/(2n)\rceil} S_{j}^{*}\right)$$

$$\leq \frac{1}{3}\xi p\gamma \|\widehat{Z}-Z^{*}\|_{1} + C_{10}(C_{0}\max_{i}\|H_{i}\|_{2})^{2} \lceil \frac{r_{1}}{2n} \rceil \left(\sqrt{np\log(2en/\lceil \frac{r_{1}}{2n}\rceil)} + \sqrt{r_{1}}\log(2en/\lceil \frac{r_{1}}{2n}\rceil)\right)$$

•

The lemma thus holds. $\hfill\square$

Proof of Corollary C.2. For any positive integer $q \in [n]$ and \hat{w} as in (74),

$$\sup_{\widehat{w}\in B_{\infty}^{n}\cap qB_{1}^{n}}\sum_{j\in[n]}|Q_{j}|\,\widehat{w}_{j} \leq \sup_{\widehat{w}\in B_{\infty}^{n}\cap qB_{1}^{n}}\sum_{j\in[n]}Q_{j}^{*}\widehat{w}_{j}^{*} = \sup_{\widehat{w}\in B_{\infty}^{n}\cap qB_{1}^{n}}\left(\sum_{j=1}^{q}Q_{j}^{*}\widehat{w}_{j}^{*} + \sum_{j=q+1}^{n}Q_{j}^{*}\widehat{w}_{j}^{*}\right) \\
\leq \sum_{j=1}^{q}Q_{j}^{*} + qQ_{q+1}^{*} \leq 2\sum_{j=1}^{q}Q_{j}^{*}.$$
(109)

Following the same arguments in Lemma 5.11, we have by (85) and (74),

$$\sup_{\widehat{Z} \in M_{\text{opt}} \cap Z^* + r_1 B_1^{n \times n}} \left| \left\langle P_2 \text{offd}(\Psi), \widehat{Z} - Z^* \right\rangle \right| \leq 2 \sup_{\widehat{w} \in B_\infty^n \cap \lceil r_1/(2n) \rceil} \sum_{B_1^n} |Q_j| \, \widehat{w}_j$$

Now (105) follows from (85), (109), and (87), where we set $q = \lceil r_1/(2n) \rceil$. Notice that for any positive integer $q \in [n]$, following the same argument, we have

$$\sup_{\widehat{w}\in B_{\infty}^{n}\cap qB_{1}^{n}}\sum_{j\in[n]}|S_{j}|\,\widehat{w}_{j} \leq \sup_{\widehat{w}\in B_{\infty}^{n}\cap qB_{1}^{n}}\sum_{j\in[n]}S_{j}^{*}\widehat{w}_{j}^{*}\leq 2\sum_{j=1}^{q}S_{j}^{*}.$$
(110)

Hence (106) follows from (86), (110), (88), and Proposition 5.19. \Box

C.4 Proof of Lemma 5.14

The proof of Lemma 5.14 is entirely deterministic. Recall by (13), we have

$$\frac{1}{n}\mathbb{E}(Y)^T u_2 = \frac{1}{n} (\sum_{i \in \mathcal{C}_1} \mathbb{E}(Y_i) - \sum_{i \in \mathcal{C}_2} \mathbb{E}(Y_i)) = 2w_1 w_2 (\mu^{(1)} - \mu^{(2)}).$$

Throughout this proof, we denote by $\mu := (\mu^{(1)} - \mu^{(2)})/\sqrt{p\gamma}$. Let Z^* be as defined in (56). Recall $P_2 = u_2 u_2^T/n = Z^*/n$. Thus we have by symmetry of $\widehat{Z} - Z^*$ and definition of \widehat{w}_j in (97),

$$\langle P_2(\mathbb{E}(Y)(Y - \mathbb{E}(Y))^T), \widehat{Z} - Z^* \rangle = \frac{1}{n} u_2^T (\mathbb{E}(Y)(Y - \mathbb{E}(Y))^T) (\widehat{Z} - Z^*) u_2$$

$$= 2w_1 w_2 \sqrt{p\gamma} \sum_{j \in [n]} \langle \mu, Y_j - \mathbb{E}(Y_j) \rangle \cdot \left(\sum_{i \in \mathcal{C}_1} (\widehat{Z} - Z^*)_{ji} - \sum_{i \in \mathcal{C}_2} (\widehat{Z} - Z^*)_{ji} \right)$$

$$= 4n w_1 w_2 \sqrt{p\gamma} \left(\sum_{j \in \mathcal{C}_2} \widehat{w}_j \langle \mu, Y_j - \mathbb{E}(Y_j) \rangle - \sum_{j \in \mathcal{C}_1} \widehat{w}_j \langle \mu, Y_j - \mathbb{E}(Y_j) \rangle \right),$$

where $\widehat{w}_j \geq 0$ is as defined in Fact B.1,

$$\forall j \in \mathcal{C}_2, \quad \widehat{w}_j := \frac{1}{2n} \big(\sum_{i \in \mathcal{C}_1} (\widehat{Z} - Z^*)_{ij} - \sum_{i \in \mathcal{C}_2} (\widehat{Z} - Z^*)_{ij} \big), \\ \forall j \in \mathcal{C}_1, \quad \widehat{w}_j := -\frac{1}{2n} \big(\sum_{i \in \mathcal{C}_1} (\widehat{Z} - Z^*)_{ij} - \sum_{i \in \mathcal{C}_2} (\widehat{Z} - Z^*)_{ij} \big).$$

Moreover, we have for each $j \in [n]$,

$$n \langle \mu, Y_j - \mathbb{E}(Y_j) \rangle = \sum_{i=1, i \neq j}^n \langle \mu, \mathbb{Z}_j - \mathbb{Z}_i \rangle = \sum_{i=1, i \neq j}^n \langle \mu, -\mathbb{Z}_i \rangle + \sum_{i=1, i \neq j}^n \langle \mu, \mathbb{Z}_j \rangle$$
$$= \sum_{i=1}^n \langle \mu, -\mathbb{Z}_i \rangle + \langle \mu, \mathbb{Z}_j \rangle + (n-1) \langle \mu, \mathbb{Z}_j \rangle$$
$$= n \langle \mu, \mathbb{Z}_j \rangle - \sum_{i=1}^n \langle \mu, \mathbb{Z}_i \rangle .$$

Hence we have by (98), and $L_{j}=\sqrt{p\gamma}\left\langle \left. \mu,Z_{j}\right. \right\rangle$,

$$\langle P_{2}(\mathbb{E}(Y)(Y - \mathbb{E}(Y))^{T}), \widehat{Z} - Z^{*} \rangle = 4w_{1}w_{2} \Big(\sum_{j \in \mathcal{C}_{2}} \widehat{w}_{j} nL_{j} - \sum_{j \in \mathcal{C}_{1}} \widehat{w}_{j} nL_{j} \Big) - 4w_{1}w_{2} \Big(\sum_{i=1}^{n} L_{i} \Big) \Big(\sum_{j \in \mathcal{C}_{2}} \widehat{w}_{j} - \sum_{j \in \mathcal{C}_{1}} \widehat{w}_{j} \Big) \leq 4w_{1}w_{2}n \sum_{j \in [n]} |L_{i}| \, \widehat{w}_{j} + 4w_{1}w_{2} \left| \sum_{i=1}^{n} L_{i} \right| \left\| (\widehat{Z} - Z^{*}) \right\|_{1} / (2n).$$

C.5 Proof of Lemma 5.15

Let $1 \leq q < n$ denote a positive integer. Then for the first component on the RHS of (75) in Lemma 5.14, we have

$$4w_{1}w_{2}\sum_{j\in[n]}|L_{j}|\,\widehat{w}_{j} = 4w_{1}w_{2}\sum_{j\in[n]}\left|\langle\mu^{(1)}-\mu^{(2)},Z_{i}\rangle\right|\widehat{w}_{j}$$

$$\leq \sum_{j\in[n]}L_{j}^{*}\widehat{w}_{j}^{*} = \sum_{j=1}^{q}L_{j}^{*}\widehat{w}_{j}^{*} + \sum_{j=q+1}^{n}L_{j}^{*}\widehat{w}_{j}^{*}, \qquad (111)$$

$$\sup \sum_{j\in[n]}|L_{j}|\,\widehat{w}_{j} \leq \sup \sum_{j\in[n]}L_{j}^{*}\widehat{w}_{j}^{*} + \sup \sum_{j=1}^{n}L_{j}^{*}\widehat{w}_{j}^{*}$$

and
$$\sup_{\widehat{w}\in B_{\infty}^{n}\cap qB_{1}^{n}}\sum_{j\in[n]}|L_{j}|\,\widehat{w}_{j} \leq \sup_{\widehat{w}\in B_{\infty}^{n}}\sum_{j=1}L_{j}^{*}\widehat{w}_{j}^{*} + \sup_{\widehat{w}\in qB_{1}^{n}}\sum_{j=q+1}L_{j}^{*}\widehat{w}_{j}^{*}$$
$$\leq \sum_{j=1}^{q}L_{j}^{*} + qL_{q+1}^{*} \leq 2\sum_{j=1}^{q}L_{j}^{*}.$$
(112)

By Fact B.1, we have $\widehat{w} = (\widehat{w}_1, \dots, \widehat{w}_n) \in B_{\infty}^n \cap \lceil \frac{r_1}{2n} \rceil B_1^n$, for all $\widehat{Z} \in \mathcal{M}_{opt} \cap (Z^* + r_1 B_1^{n \times n})$. By setting $q = \lceil \frac{r_1}{2n} \rceil \in [n]$ in(112), we obtain

$$\sup_{\widehat{w}\in B_{\infty}^{n}\cap\lceil\frac{r_{1}}{2n}\rceil B_{1}^{n}}\sum_{j\in[n]}|L_{j}|\,\widehat{w}_{j}\leq 2\sum_{j=1}^{\lceil r_{1}/(2n)\rceil}L_{j}^{*}.$$
(113)

Moreover, for the second component on the RHS of (75), we have by (98),

$$\sup_{\widehat{Z} \in \mathcal{M}_{opt} \cap Z^* + r_1 B_1^{n \times n}} \left| \frac{1}{n} \sum_{j=1}^n L_j \right| \left\| (\widehat{Z} - Z^*) \right\|_1 / (2n) \\
\leq \sup_{\widehat{w} \in B_\infty^n \cap \lceil \frac{r_1}{2n} \rceil B_1^n} \frac{1}{n} \sum_{j=1}^n |L_j| \left| \sum_j \widehat{w}_j \right| \leq \frac{\lceil r_1 / 2n \rceil}{n} \sum_{j=1}^n L_j^* \leq \sum_{j=1}^{\lceil r_1 / (2n) \rceil} L_j^*, \quad (114)$$

where (114) holds since in the second sum $\sum_{j=1}^{q} L_{j}^{*}$, where $q = \lceil \frac{r_{1}}{2n} \rceil$, we give a unit weight to each of the q largest components $L_{j}^{*}, j \in [q]$, while in $\frac{q}{n} \sum_{i=1}^{n} L_{j}^{*}$, we give the same weight $q/n \leq 1$ to each of n components $L_{j}^{*}, j \in [n]$. Putting things together, we have by (75), (113), and (114),

$$\sup_{\widehat{Z}\in\mathcal{M}_{\text{opt}}\cap(Z^*+r_1B_1^{n\times n})} \left| \left\langle P_2(\mathbb{E}(Y)(Y-\mathbb{E}(Y))^T), \widehat{Z}-Z^* \right\rangle \right|$$

$$\leq \sup_{\widehat{w}\in B_\infty^n\cap \lceil \frac{r_1}{2n}\rceil B_1^n} n \sum_{j\in[n]} |L_j| \,\widehat{w}_j + \sup_{\widehat{w}\in B_\infty^n\cap \lceil \frac{r_1}{2n}\rceil B_1^n} \left| \sum_{j=1}^n L_j \right| \sum_j \widehat{w}_j \le 3 \sum_{j=1}^{\lceil \frac{r_1}{2n}\rceil} L_j^*.$$

The lemma thus holds. \Box

C.6 Proof of Proposition 5.16

Recall by (92),

$$L_j := \langle \mu^{(1)} - \mu^{(2)}, \mathbb{Z}_j \rangle = \sqrt{p\gamma} \langle \mu, \mathbb{Z}_j \rangle.$$

First, we bound for any integer $q \in [n]$, and $\tau > 0$,

$$\mathbb{P}\left(\sum_{j=1}^{q} L_{j}^{*} > \tau\right) \leq \mathbb{P}\left(\exists J \in [n], |J| = q, \sum_{j \in J} |L_{j}| > \tau\right) \\
= \mathbb{P}\left(\max_{J \in [n], |J| = q} \max_{u = (u_{1}, \dots, u_{q}) \in \{-1, 1\}^{q}} \sum_{j \in J} u_{j}L_{j} > \tau\right) \\
\leq \sum_{J: |J| = q} \sum_{u \in \{-1, 1\}^{q}} \mathbb{P}\left(\sum_{j \in J} u_{j}L_{j}/\Delta > \tau/\Delta\right) \\
\leq \binom{n}{q} 2^{q} \exp\left(-\frac{(\tau/\Delta)^{2}}{C_{1}C_{0}^{2}(\max_{j} ||R_{j}\mu||_{2}^{2})q}\right),$$
(115)

where (115) holds by the union bound and the sub-gaussian tail bound; To see this, notice that for each fixed index set $J \in [n]$ and vector $u \in \{-1, 1\}^q$, following the proof of Lemma A.2, we obtain for $\Delta^2 = p\gamma$ and $\mu \in \mathbb{S}^{p-1}$,

$$\left\| \sum_{j \in J, |J|=q} u_j L_j / \sqrt{p\gamma} \right\|_{\psi_2}^2 \leq \sum_{j \in J, |J|=q} \|L_j / \sqrt{p\gamma}\|_{\psi_2}^2 = \sum_{j \in J, |J|=q} \left\| \langle \mu^{(1)} - \mu^{(2)} / \sqrt{p\gamma}, \mathbb{Z}_j \rangle \right\|_{\psi_2}^2$$
$$= \sum_{j \in J, |J|=q} \|\langle \mu, \mathbb{Z}_j \rangle \|_{\psi_2}^2 \leq C_1 q (C_0 \max_j \|R_j \mu\|_2)^2.$$
(116)

Hence by the sub-gaussian tail bound, cf. Lemma A.2, upon replacing n with $1 \le q < n$, we have for each fixed index set $J \subset [n]$,

$$\mathbb{P}\left(\sum_{j\in J:|J|=q} u_j L_j/\Delta > \tau/\Delta\right) \le \exp\left(-\frac{(\tau/\Delta)^2}{C_1(C_0 \max_j \|R_j\mu\|_2)^2 q}\right).$$
(117)

Set $\tau_q = C_3 C_0(\max_j ||R_j\mu||_2) \Delta q \sqrt{\log(2en/q)}$ for some absolute constant $C_3 > c \sqrt{C_1}$ for $c \ge 2$. Then by (115) and (117), we have

$$\mathbb{P}\left(\sum_{j=1}^{q} L_{j}^{*} > \tau_{q}\right) \leq \binom{n}{q} 2^{q} \exp\left(-\frac{C_{3}^{2}(C_{0} \max_{j} ||R_{j}\mu||_{2})^{2} q^{2} \log(2en/q)}{C_{1}(C_{0} \max_{j} ||R_{j}\mu||_{2})^{2} q}\right) \leq (2en/q)^{q} \exp\left(-cq \log(2en/q)\right) \leq e^{q \log(2en/q)} e^{-cq \log(2en/q)} \leq e^{-c'q \log(2en/q)},$$

where $\binom{n}{q}2^q \leq (en/q)^q 2^q = (2en/q)^q = \exp(q\log(2en/q))$. Following the calculation as done in [18], and the inequality immediately above, we have

$$\sum_{q=1}^{n} \binom{n}{q} 2^{q} e^{-\tau_{q}^{2}/(C_{1}C_{0}^{2}q\Delta^{2})} \leq \sum_{q=1}^{n} e^{-c'q\log(2en/q)} \leq \frac{c}{n^{2}}.$$

The proposition thus holds. \Box

C.7 Proof of Lemma 5.17

Throughout this proof, denote by $\Psi = \mathbb{Z}\mathbb{Z}^T - \mathbb{E}(\mathbb{Z}\mathbb{Z}^T)$. Let u_2 be the group membership vector as in (2). Let $\mathbf{1}_n = (1, \ldots, 1)$ and $\mathbf{1}_n / \sqrt{n} \in \mathbb{S}^{n-1}$. Hence we have $u_2^T \mathbf{1}_n = |C_1| - |C_2|$. First, we have by Fact B.2, (78) holds since

$$\left| \left\langle P_2 \Psi P_1, \widehat{Z} - Z^* \right\rangle \right| = \left| \operatorname{tr}((\widehat{Z} - Z^*) P_2 \Psi P_1) \right| = \left| \operatorname{tr}(P_2 \Psi P_1(\widehat{Z} - Z^*)) \right|$$
$$= \left| \operatorname{tr}(\frac{u_2 u_2^T}{n} \Psi \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^T (\widehat{Z} - Z^*)) \right|$$
$$\leq \left| \frac{2u_2^T \Psi \mathbf{1}_n}{n} \right| \left| \frac{\mathbf{1}_n^T (\widehat{Z} - Z^*) u_2}{2n} \right| \leq \left\| \Psi \right\|_2 \left\| \widehat{Z} - Z^* \right\|_1 / n.$$

Second, (79) holds since $\mathbf{1}_n^T \Psi \mathbf{1}_n / n \leq \|\Psi\|_2$ and

$$\langle P_2 P_1 \Psi P_1, \widehat{Z} - Z^* \rangle = \langle \frac{u_2 u_2^T}{n} \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^T \Psi \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^T, \widehat{Z} - Z^* \rangle$$

$$= \langle \frac{u_2}{n} (u_2^T \frac{1}{n} \mathbf{1}_n) \frac{1}{n} (\mathbf{1}_n^T \Psi \mathbf{1}_n) \mathbf{1}_n^T, \widehat{Z} - Z^* \rangle$$

$$= 2 \frac{|C_1| - |C_2|}{n} \frac{\mathbf{1}_n^T \Psi \mathbf{1}_n}{n} \frac{1}{2n} \mathbf{1}_n^T (\widehat{Z} - Z^*) u_2.$$

Hence by Fact B.2,

$$\left| \langle P_2 P_1 \Psi P_1, \widehat{Z} - Z^* \rangle \right| \leq |w_1 - w_2| \|\Psi\|_2 \|\widehat{Z} - Z^*\|_1 / n$$

To bound (80), we have by Fact 5.2,

$$\langle P_2 P_1 \Psi P_1 P_2, \widehat{Z} - Z^* \rangle = \langle \frac{u_2 u_2^T}{n} \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^T \Psi \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^T \frac{u_2 u_2^T}{n}, \widehat{Z} - Z^* \rangle$$

$$= \langle \frac{u_2}{n} (u_2^T \mathbf{1}_n/n) \frac{(\mathbf{1}_n^T \Psi \mathbf{1}_n)}{n} (\mathbf{1}_n^T u_2/n) u_2^T, \widehat{Z} - Z^* \rangle$$

$$= (w_1 - w_2)^2 \frac{\mathbf{1}_n^T \Psi \mathbf{1}_n}{n} \langle \frac{u_2 u_2^T}{n}, \widehat{Z} - Z^* \rangle ,$$

and hence

$$\left| \langle P_2 P_1 \Psi P_1 P_2, \widehat{Z} - Z^* \rangle \right| \leq (w_1 - w_2)^2 \|\Psi\|_2 \|\widehat{Z} - Z^*\|_1 / n.$$

Finally, it remains to show (81) and (82). First, we rewrite the inner product as follows:

$$\langle \widehat{Z} - Z^*, P_2 \Psi P_2 \rangle = \operatorname{tr}((\widehat{Z} - Z^*)^T \frac{u_2 u_2^T}{n} \Psi \frac{u_2 u_2^T}{n}) = \frac{u_2^T (\widehat{Z} - Z^*)^T u_2}{n} \frac{u_2^T \Psi u_2}{n} = -\frac{1}{n} \left\| \widehat{Z} - Z^* \right\|_1 \frac{u_2^T \Psi u_2}{n};$$

Then (81) holds by Fact 5.2. Finally, we compute

$$\langle P_2 P_1 \Psi P_2, \widehat{Z} - Z^* \rangle = \operatorname{tr}((\widehat{Z} - Z^*) \frac{u_2 u_2^T}{n} \frac{\mathbf{1}_n \mathbf{1}_n^T}{n} \Psi \frac{u_2 u_2^T}{n})$$

$$= \operatorname{tr}(\frac{u_2^T}{n} (\widehat{Z} - Z^*) u_2) \frac{u_2^T \mathbf{1}_n}{n} \frac{\mathbf{1}_n^T \Psi u_2}{n},$$

and $\langle P_2 \Psi P_1 P_2, \widehat{Z} - Z^* \rangle = \operatorname{tr}((\widehat{Z} - Z^*) \frac{u_2 u_2^T}{n} \Psi \frac{\mathbf{1}_n \mathbf{1}_n^T}{n} \frac{u_2 u_2^T}{n})$

$$= \operatorname{tr}(P_2(\widehat{Z} - Z^*)) \frac{u_2^T \Psi \mathbf{1}_n}{n} \frac{\mathbf{1}_n^T u_2}{n}.$$

Then

$$\left| \left\langle P_2 P_1 \Psi P_2, \widehat{Z} - Z^* \right\rangle \right| = \left| \left\langle P_2 \Psi P_1 P_2, \widehat{Z} - Z^* \right\rangle \right|$$

$$\leq |w_1 - w_2| \frac{1}{n} \left\| \widehat{Z} - Z^* \right\|_1 \|\Psi\|_2.$$

Hence (82) holds and the lemma is proved. \Box

C.8 Proof of Lemma 5.18

First, we have

$$\langle P_2 P_1 \Psi, \widehat{Z} - Z^* \rangle = \operatorname{tr}((\widehat{Z} - Z^*) P_2 P_1 \Psi)$$

$$= \operatorname{tr}((\widehat{Z} - Z^*) \frac{u_2 u_2^T}{n} \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^T \Psi) = \operatorname{tr}((\widehat{Z} - Z^*) \frac{u_2}{n} (u_2^T \frac{1}{n} \mathbf{1}_n) \mathbf{1}_n^T \Psi)$$

$$= \frac{|C_1| - |C_2|}{n} \operatorname{tr}((\widehat{Z} - Z^*) \frac{u_2}{n} \mathbf{1}_n^T \Psi) = (w_1 - w_2) \mathbf{1}_n^T \Psi (\widehat{Z} - Z^*) \frac{u_2}{n}.$$

Thus

$$\begin{aligned} \left| \langle P_2 P_1 \text{offd}(\Psi), \widehat{Z} - Z^* \rangle \right| &= |w_1 - w_2| \left| \mathbf{1}_n^T \text{offd}(\Psi) (\widehat{Z} - Z^*) \frac{w_2}{n} \right| \\ &\leq 2 |w_1 - w_2| \sum_{k=1}^n \left| \left(\sum_{j \neq k}^n \Psi_{kj} \right) \right| \left| \frac{1}{2n} \left(\sum_{j \in \mathcal{C}_1} (\widehat{Z} - Z^*)_{kj} - \sum_{j \in \mathcal{C}_2} (\widehat{Z} - Z^*)_{kj} \right) \right| \\ &= 2 |w_1 - w_2| \sum_k |S_k| \, \widehat{w}_k. \end{aligned}$$

Therefore, we have shown (86). Now for the diagonal component, denote by

$$|S_{\text{diag}}| = \left| \langle P_2 P_1 \text{diag}(\Psi), \widehat{Z} - Z^* \rangle \right| = |w_1 - w_2| \left| \mathbf{1}_n^T \text{diag}(\Psi) (\widehat{Z} - Z^*) \frac{u_2}{n} \right|$$

$$:= 2 |w_1 - w_2| \left| \sum_{k \in C_1} \Psi_{kk}(-\widehat{w}_k) + \sum_{k \in C_2} \Psi_{kk} \widehat{w}_k \right|$$

$$\leq 2 |w_1 - w_2| \sum_{k \in [n]} |\Psi_{kk}| \, \widehat{w}_k \leq |w_1 - w_2| \max_k |\Psi_{kk}| \left\| \widehat{Z} - Z^* \right\|_1 / n,$$

where recall for u_2 as in (2),

$$\frac{(\widehat{Z}-Z^*)u_2}{n} = (-\widehat{w}_1,\ldots,-\widehat{w}_{n_1},\widehat{w}_{n_1+1},\ldots,\widehat{w}_n).$$

Now

$$\langle P_{2}\Psi, \widehat{Z} - Z^{*} \rangle = \operatorname{tr}((\widehat{Z} - Z^{*})P_{2}\Psi) = \operatorname{tr}(P_{2}\Psi(\widehat{Z} - Z^{*})) = \langle \Psi, (\widehat{Z} - Z^{*})P_{2} \rangle$$

$$= \frac{1}{n}(u_{2}^{T}\Psi(\widehat{Z} - Z^{*})u_{2}) = \langle \Psi u_{2}, (\widehat{Z} - Z^{*})u_{2}/n \rangle$$

$$= 2\sum_{k=1}^{n} \left(\sum_{t \in \mathcal{C}_{1}} \Psi_{kt} - \sum_{t \in \mathcal{C}_{2}} \Psi_{kt}\right) \cdot \frac{1}{2n} \left(\sum_{j \in \mathcal{C}_{1}} (\widehat{Z} - Z^{*})_{kj} - \sum_{j \in \mathcal{C}_{2}} (\widehat{Z} - Z^{*})_{kj}\right)$$

$$= 2\left(\sum_{k \in \mathcal{C}_{1}} \Psi_{kk}(-\widehat{w}_{k}) - \sum_{t \in \mathcal{C}_{2}} \Psi_{tt}\widehat{w}_{t}\right) + 2\left(\sum_{j \in \mathcal{C}_{1}} Q_{j}(-\widehat{w}_{j}) + \sum_{j \in \mathcal{C}_{2}} Q_{j}\widehat{w}_{j}\right) = Q_{\text{diag}} + Q_{\text{offd}}.$$

For the diagonal component, we have

$$\begin{aligned} |Q_{\text{diag}}| &:= \left| \left\langle P_2 \text{diag}(\Psi), \widehat{Z} - Z^* \right\rangle \right| &= 2 \left| \sum_{k \in C_1} \Psi_{kk}(-\widehat{w}_k) + \sum_{k \in C_2} (-\Psi_{kk}) \widehat{w}_k \right| \\ &\leq 2 \sum_{k \in [n]} |\Psi_{kk}| \, \widehat{w}_k \leq \max_k |\Psi_{kk}| \left\| \widehat{Z} - Z^* \right\|_1 / n. \end{aligned}$$

Moreover, for the off-diagonal component, we have

$$|Q_{\text{offd}}| = 2 \left| \sum_{j \in \mathcal{C}_1} Q_j(-\widehat{w}_j) + \sum_{j \in \mathcal{C}_2} Q_j \widehat{w}_j \right| \le 2 \sum_{j \in [n]} |Q_j| \, \widehat{w}_j.$$

Hence (85) holds, given that

$$\left| \langle P_2 \text{offd}(\Psi), \widehat{Z} - Z^* \rangle \right| \leq 2 \sum_{k \in [n]} \widehat{w}_k |Q_k|.$$

C.9 Proof of Proposition 5.19

Denote by $\Psi = \mathbb{Z}\mathbb{Z}^T - \mathbb{E}(\mathbb{Z}\mathbb{Z}^T)$. Recall

$$\forall j \in [n] \quad Q_j := \sum_{t \in \mathcal{C}_1, t \neq j} \Psi_{tj} - \sum_{t \in \mathcal{C}_2, t \neq j} \Psi_{tj} \quad \text{and} \quad S_j := \sum_{i \neq j}^n \Psi_{ji}.$$

It remains to show (87) and (88). The two bounds follow identical steps, and hence we will only prove (87). First, we bound for any positive integer $1 \le q < n$ and any $\tau > 0$,

$$\mathbb{P}\left(\sum_{j=1}^{q} Q_{j}^{*} > \tau\right) = \mathbb{P}\left(\exists J \in [n], |J| = q, \sum_{j \in J} |Q_{j}| > \tau\right) \\
= \mathbb{P}\left(\max_{J \in [n], |J| = q} \max_{y = (y_{1}, \dots, y_{q}) \in \{-1, 1\}^{q}} \sum_{j \in J} y_{j} Q_{j} > \tau\right) \\
\leq \sum_{J: |J| = q} \sum_{y \in \{-1, 1\}^{q}} \mathbb{P}\left(\sum_{j \in J} y_{j} Q_{j} > \tau\right) = q_{0}(\tau).$$
(118)

We now bound $q_0(\tau)$. Denote by $w \in \{0, -1, 1\}^n$ the 0-extended vector of $y \in \{-1, 1\}^{|J|}$, that is, for the fixed index set $J \subset [n]$, we have $w_j = y_j, \forall j \in J$ and $w_j = 0$ otherwise. Now we can write for $A = (a_{tj}) = w \otimes u_2$, where u_2 is the group membership vector:

$$\forall t \in \mathcal{C}_1, j \in J, \quad a_{jt} = u_{2,t}y_j = y_j, \\ \forall t \in \mathcal{C}_2, j \in J, \quad a_{jt} = u_{2,t}y_j = -y_j,$$

and 0 otherwise. In other words, we have nq number of non-zero entries in A and each entry has absolute value 1. Thus $||A||_2 \leq ||A||_F = \sqrt{qn}$ and

$$\sum_{j\in J} y_j \left(\sum_{t\in\mathcal{C}_1, t\neq j} \left\langle \mathbb{Z}_t, \mathbb{Z}_j \right\rangle - \sum_{t\in\mathcal{C}_2, t\neq j} \left\langle \mathbb{Z}_t, \mathbb{Z}_j \right\rangle \right) = \sum_{j\in[n]} \sum_{t\neq j} a_{jt} \left\langle \mathbb{Z}_t, \mathbb{Z}_j \right\rangle,$$

since each row of offd(A) in J has n-1 nonzero entries and |J| = q. Set

$$\tau_q = C_4 (C_0 \max_i \|H_i\|_2)^2 q(\sqrt{np \log(2en/q)} + \sqrt{nq} \log(2en/q)).$$

Now for each fixed index set $J \in [n]$ with |J| = q, and a fixed vector $y \in \{-1, 1\}^q$,

$$\mathbb{P}\left(\sum_{j\in J} y_j Q_j > \tau_q\right) = \mathbb{P}\left(\sum_{j\in J} y_j \left(\sum_{t\in\mathcal{C}_1, t\neq j} \Psi_{tj} - \sum_{t\in\mathcal{C}_2, t\neq j} \Psi_{tj}\right) > \tau_q\right) \\
= \mathbb{P}\left(\sum_j \sum_{t\neq j} a_{jt} \langle \mathbb{Z}_t, \mathbb{Z}_j \rangle > \tau_q\right) \\
\leq 2\exp\left(-c(C_4^2 \wedge C_4)q\log(2en/q)\right).$$
(119)

It remains to show the inequality in (119). By Theorem A.1, we have for $q \in [n]$, $A = (a_{ij}) \in \mathbb{R}^{n \times n}$, and $\tau_q = C_4(C_0 \max_i ||H_i||_2)^2 q(\sqrt{np \log(2en/q)} + \sqrt{nq} \log(2en/q))$,

$$\begin{aligned} & \mathbb{P}\left(\left|\sum_{i=1}^{n}\sum_{j\neq i}^{n}\left\langle\mathbb{Z}_{i},\mathbb{Z}_{j}\right\rangle a_{ij}\right| > \tau_{q}\right) \\ & \leq 2\exp\left(-c\min\left(\frac{\tau_{q}^{2}}{(C_{0}\max_{i}\|H_{i}\|_{2})^{4}p\|A\|_{F}^{2}}, \frac{\tau_{q}}{(C_{0}\max_{i}\|H_{i}\|_{2})^{2}\|A\|_{2}}\right)\right) \\ & \leq 2\exp\left(-c\min\left(\frac{(C_{4}q\sqrt{np\log(2en/q)})^{2}}{pnq}, \frac{C_{4}q\sqrt{nq}\log(2en/q)}{\sqrt{qn}}\right)\right) \\ & \leq 2\exp\left(-c(C_{4}^{2}\wedge C_{4})q\log(2en/q)\right), \end{aligned}$$

where $\max_i ||Z_i||_{\psi_2} \leq CC_0 \max_i ||H_i||_2$ in the sense of (89). Thus (119) holds. Similarly, we can prove the same bound holds for (105), following an identical sequence of arguments. Now we have for by (118) and (119),

$$q_0(\tau_q) \leq \binom{n}{q} 2^q \exp\left(-c(C_4^2 \wedge C_4)q\log(2en/q)\right) \\ = e^{q\log(2en/q)} \exp\left(-c(C_4^2 \wedge C_4)q\log(2en/q)\right) \\ \leq 2\exp\left(-c'(C_4^2 \wedge C_4)q\log(2en/q)\right),$$

where $\binom{n}{q}2^q \leq (en/q)^q 2^q = (2en/q)^q = \exp(q\log(2en/q))$. Hence following the calculation as done in [18], we have

$$\sum_{q=1}^{n} q_0(\tau_q) \le \sum_{q=1}^{n} \binom{n}{q} 2^q \exp\left(-c(C_4^2 \wedge C_4)q \log(2en/q)\right) \le \sum_{q=1}^{n} e^{-c'q \log(2en/q)} \le \frac{c'}{n^2}.$$

The proposition thus holds. \Box

D The oracle estimator and its properties

The presentation of this section follows from [39], which we include for self-containment. Both A and B are defined to bridge the gap between YY^T and $R = \mathbb{E}(Y)\mathbb{E}(Y)^T$, and the bias term

$$\mathbb{E}B - R = \mathbb{E}YY^T - \mathbb{E}(Y)\mathbb{E}(Y)^T - \mathbb{E}\lambda(E_n - I_n) - \mathbb{E}\tau I_n$$

is substantially smaller than $\mathbb{E}A - R$ in the operator norm, under assumption (A2). Proposition D.1 shows that the optimization goals of SDP1 and SDP (10) are equivalent to that of **Oracle SDP**.

Proposition D.1. [39] The optimal solutions \hat{Z} as in SDP (10) must have their diagonals set to I_n . Thus, the set of optimal solutions \hat{Z} in (10) coincide with those on the convex subset \mathcal{M}_{opt} as in (15),

$$\underset{Z \in \mathcal{M}_{G}^{+}}{\operatorname{arg\,max}} \langle A, Z \rangle = \underset{Z \in \mathcal{M}_{opt}}{\operatorname{arg\,max}} \langle A, Z \rangle = \underset{Z \in \mathcal{M}_{opt}}{\operatorname{arg\,max}} \langle \widetilde{A}, Z \rangle$$
(120)

$$= \underset{Z \in \mathcal{M}_{opt}}{\operatorname{arg\,max}} (\langle A, Z \rangle - \mathbb{E}\tau \langle I_n, Z \rangle).$$
(121)

Theorem D.2 follows immediately from Lemma 2.6 and Theorem 2.7.

Theorem D.2. (*R* is the leading term) [39] Suppose the conditions in Theorem 2.3 hold. Then with probability at least $1 - 2\exp(-cn)$, for the oracle *B* as in (17),

 $||B - R||_2 \leq \xi np\gamma$, where the reference R is as defined in (14).

D.1 Proof of Lemma 2.6

The proof follows from [39], Section H. We include it here for self-containment. We have the following facts about R as defined in (14).

Fact D.3. First, $\mathbf{1}_n^T R \mathbf{1}_n = 0$. Hence

$$\mathbf{1}_{n}^{T} \text{offd}(R) \mathbf{1}_{n} = \mathbf{1}_{n}^{T} R \mathbf{1}_{n} - \text{tr}(R) = -\text{tr}(R) = -np\gamma w_{2}w_{1}, \text{ where}$$
(122)
$$\text{tr}(R)/(p\gamma) = nw_{2}^{2}w_{1} + nw_{1}^{2}w_{2} = w_{1}w_{2}n \text{ and} ||R||_{2} = \text{tr}(R) = w_{1}w_{2}np\gamma.$$

Thus $\operatorname{tr}(R)/n = p\gamma w_1 w_2$ and for $n \ge 4$,

$$\left\|\frac{\operatorname{tr}(R)}{n-1}\left(I_n - E_n/n\right)\right\|_2 \leq \frac{n}{n-1}p\gamma w_1 w_2 \leq p\gamma/3,\tag{123}$$

where $||I_n - E_n/n||_2 = 1$ since $I_n - E_n/n$ is a projection matrix.

Proof of Lemma 2.6. We have by Proposition 5.3,

$$\|\mathbb{E}B - R\| = \left\| W_0 - \mathbb{W} - \frac{\operatorname{tr}(R)}{(n-1)} (I_n - \frac{E_n}{n}) \right\| \\ \leq \|W_0\| + \|\mathbb{W}\| + \left\| \frac{\operatorname{tr}(R)}{(n-1)} (I_n - \frac{E_n}{n}) \right\|,$$
(124)

where the $\|\cdot\|$ is understood to be either the operator or the cut norm. Then

$$\|W_0\|_2 \leq |V_1 - V_2| (w_2 \vee w_1).$$
(125)

Moreover,

$$\|\mathbb{W}\|_{2} \leq |V_{1} - V_{2}| (w_{1} \vee w_{2}).$$
(126)

Combining (123), (124), (125), and (126), we have for $w_{\min} := w_1 \wedge w_2$ and $n\xi \ge \frac{1}{2w_{\min}}$,

$$\begin{split} \|\mathbb{E}B - R\|_{2} &\leq \|W_{0}\|_{2} + \|\mathbb{W}\|_{2} + \left\|\frac{\operatorname{tr}(R)}{(n-1)}(I_{n} - \frac{E_{n}}{n})\right\|_{2} \\ &\leq 2|V_{1} - V_{2}|(w_{1} \lor w_{2}) + p\gamma/3 \\ &= \frac{2}{3}\xi np\gamma(1 - w_{\min}) + p\gamma/3 \leq \frac{2}{3}\xi np\gamma, \end{split}$$

where we use the fact that

$$\frac{2}{3}\xi p\gamma nw_{\min} \ge p\gamma/3$$
 since $2\xi nw_{\min} \ge 1$.

The lemma thus holds for the general setting; when $V_1 = V_2$, we show the improved bounds in Lemma H.7. [39]. \Box

D.2 Proof sketch of Theorem 2.7

Let Y be as in Definition 1.1. We provide a proof outline for Theorem 2.7 in this section. The idea of decomposition and reduction appears in the local analysis as well; See Proofs for Lemma 5.6 in Section 5.4, as well as Corollary A.4. We have by (69) and (70), and the triangle inequality,

$$\|\Lambda\|_{2} = \|YY^{T} - \mathbb{E}(YY^{T})\|_{2} \leq \|\widehat{\Sigma}_{Y} - \Sigma_{Y}\|_{2} + \|\mathbb{E}(Y)(Y - \mathbb{E}(Y))^{T} + (Y - \mathbb{E}(Y))\mathbb{E}(Y)^{T}\|_{2},$$

where $\widehat{\Sigma}_{Y} = (Y - \mathbb{E}(Y))(Y - \mathbb{E}(Y))^{T}$ and
 $\|\widehat{\Sigma}_{Y} - \Sigma_{Y}\|_{2} \leq \|I - P_{1}\|_{2} \|\mathbb{Z}\mathbb{Z}^{T} - \mathbb{E}(\mathbb{Z}\mathbb{Z}^{T})\|_{2} \|I - P_{1}\|_{2}.$ (127)

To control $||M_Y|| = ||\mathbb{E}(Y)(Y - \mathbb{E}(Y))^T + (Y - \mathbb{E}(Y))(\mathbb{E}(Y))^T||$, we need to bound the projection of each mean-zero random vector $\mathbb{Z}_j, \forall j \in [n]$, along the direction of $v := \mu^{(1)} - \mu^{(2)}$. In other words, a particular direction for which we compute the one-dimensional marginals is the direction between the two centers, namely, $\mu^{(1)}$ and $\mu^{(2)}$. Let $c, c', c_1, c_5, C_3, C_4, \ldots$ be absolute constants.

Lemma D.4. (Projection: probabilistic view) [39] Suppose conditions in Theorem 2.7 hold. Then we have with probability at least $1 - 2 \exp(-cn)$,

$$\begin{aligned} \left\| \mathbb{E}(Y)(Y - \mathbb{E}(Y))^T + (Y - \mathbb{E}(Y))(\mathbb{E}(Y))^T \right\|_2 &\leq 4\sqrt{n}\sqrt{w_1w_2} \sup_{q \in S^{n-1}} \left| \sum_i q_i \left\langle \mathbb{Z}_i, \mu^{(1)} - \mu^{(2)} \right\rangle \right| \\ &\leq 2C_3C_0(\max_i \left\| H_i^T \mu \right\|_2)n\sqrt{p\gamma}. \end{aligned}$$

Hence Theorem 2.7 follows immediately from Lemma D.4 and Theorem C.1. Lemma D.4 follows from Lemma A.3 and the sub-gaussian concentration of measure bound in (94). On the other hand, controlling the second component in (69) amounts to the problem of covariance estimation given the mean matrix $\mathbb{E}(Y)$, in view of (127) and Theorem C.1. See [39] for proofs; cf. Lemma 7.1. and Theorem 7.2 therein.

References

- [1] ABBE, E. (2017). Community detection and the stochastic block model: recent developments. Journal of Machine Learning Research 4.
- [2] ABBE, E., FAN, J. and WANG, K. (2022). An ℓ_p theory of PCA and spectral clustering. Ann. Statist. 50 2359–2385.
- [3] ACHLIOPTAS, D. and MCSHERRY, F. (2005). On spectral learning of mixtures of distributions. In *Proceedings of the 18th Annual COLT*. (Version in http://www.cs.ucsc.edu/ optas/papers/).
- [4] ALOISE, D., DESHPANDE, A., HANSEN, P. and POPAT, P. (2009). NP-hardness of Euclidean sum-of-squares clustering. *Machine Learning* 75 245–248.
- [5] ARORA, S. and KANNAN, R. (2001). Learning mixtures of arbitrary gaussians. In *Proceedings* of 33rd ACM Symposium on Theory of Computing.
- [6] BALAKRISHNAN, S., WAINWRIGHT, M. J. and YU, B. (2017). Statistical guarantees for the EM algorithm: From population to sample-based analysis. *The Annals of Statistics* **45** 77–120.
- [7] BANKS, J., MOORE, C., VERSHYNIN, R., VERZELEN, N. and XU, J. (2018). Informationtheoretic bounds and phase transitions in clustering, sparse PCA, and submatrix localization. *IEEE Trans. Inform. Theory* 64 4872–4994.
- [8] BLUM, A., COJA-OGHLAN, A., FRIEZE, A. and ZHOU, S. (2009). Separating populations with wide data: a spectral analysis. *Electronic Journal of Statistics* **3** 76–113.
- [9] BUNEA, F., GIRAUD, C., LUO, X., ROYER, M. and VERZELEN, N. (2020). Model assisted variable clustering: minimax-optimal recovery and algorithms. *The Annals of Statistics* 48 111–137.
- [10] CHAUDHURI, K., HALPERIN, E., RAO, S. and ZHOU, S. (2007). A rigorous analysis of population stratification with limited data. In *Proceedings of the 18th ACM-SIAM SODA*.
- [11] CHEN, X. and YANG, Y. (2021). Hanson-Wright inequality in Hilbert spaces with application to K-means clustering for non-Euclidean data. *Bernoulli* 27 586–614.
- [12] CHRÉTIEN, S., CUCURINGU, M., LECUÉ, G. and NEIRAC, L. (2021). Learning with semidefinite programming: statistical bounds based on fixed point analysis and excess risk curvature. *Journal of Machine Learning Research* 22.
- [13] DASGUPTA, S. and SCHULMAN, L. J. (2000). A two-round variant of em for gaussian mixtures. In Proceedings of the 16th Conference on Uncertainty in Artificial Intelligence (UAI).
- [14] DRINEAS, P., FRIEZE, A., KANNAN, R., VEMPALA, S. and VINAY, V. (2004). Clustering large graphs via the singular value decomposition. *Machine Learning* 9–33.
- [15] FEI, Y. and CHEN, Y. (2018). Hidden integrality of SDP relaxations for sub-Gaussian mixture models. In *Conference On Learning Theory*.
- [16] FEI, Y. and CHEN, Y. (2021). Hidden integrality and semi-random robustness of SDP relaxation for Sub-Gaussian mixture model.

- [17] FIEDLER, M. (1973). Algebraic connectivity of graphs. Czechoslovak Mathematical Journal 23 298-305. URL http://eudml.org/doc/12723
- [18] GIRAUD, C. and VERZELEN, N. (2019). Partial recovery bounds for clustering with the relaxed k-means. Mathematical Statistics and Learning 1 317–374.
- [19] GOEMANS, M. and WILLIAMSON, D. (1995). Improved approximation algorithms for maximum cut and satisfiability problems using semidefinite programming. JACM 42 1115–1145.
- [20] GUEDON, O. and VERSHYNIN, R. (2016). Community detection in sparse networks via grothendieck's inequality. *Probability Theory and Related Fields* 165 1025–1049.
- [21] HORNSTEIN, M., FAN, R., SHEDDEN, K. and ZHOU, S. (2019). Joint mean and covariance estimation for unreplicated matrix-variate data. *Journal of the American Statistical Association* (*Theory and Methods*) **114** 682–696.
- [22] KALAI, A. T., MOITRA, A. and VALIANT, G. (2010). Efficiently learning mixtures of two gaussians. In Proceedings of the Forty-second ACM Symposium on Theory of Computing. ACM.
- [23] KANNAN, R., SALMASIAN, H. and VEMPALA, S. (2005). The spectral method for general mixture models. In *Proc. of the 18th Annual COLT*.
- [24] KARP, R. M. (1972). Reducibility among combinatorial problems. In Proceedings of a symposium on the Complexity of Computer Computations, held March 20-22, 1972, at the IBM Thomas J. Watson Research Center, Yorktown Heights, New York, USA (R. E. Miller and J. W. Thatcher, eds.). The IBM Research Symposia Series, Plenum Press, New York.
- [25] KUMAR, A. and KANNAN, R. (2010). Clustering with spectral norm and the k-means algorithm. In Proceedings of the 2010 IEEE 51st Annual Symposium on Foundations of Computer Science. IEEE Computer Society.
- [26] LI, X., LI, Y., LING, S., STROHMER, T. and WEI, K. (2020). When do birds of a feather flock together? k-means, proximity, and conic programming. *Mathematical Programming* 179 295–341.
- [27] LÖFFLER, M., ZHANG, A. Y. and ZHOU, H. H. (2021). Optimality of spectral clustering in the Gaussian mixture model. *The Annals of Statistics* 49 2506 – 2530.
- [28] LU, Y. and ZHOU, H. H. (2016). Statistical and computational guarantees of Lloyd's algorithm and its variants.
- [29] NDAOUD, M. (2022). Sharp optimal recovery in the two component gaussian mixture model. Ann. Statist. 50 2096–2126.
- [30] PATTERSON, N., PRICE, A. and REICH, D. (2006). Population structure and eigenanalysis. PLoS Genet 2. Doi:10.1371/journal.pgen.0020190.
- [31] PENG, J. and WEI, Y. (2007). Approximating K-means-type clustering via semidefinite programming. SIAM Journal on Optimization 18 186–205.

- [32] PENG, J. and XIA, Y. (2005). A new theoretical framework for K-means-type clustering. In Foundations and Advances in Data Mining. Springer.
- [33] PRICE, A., PATTERSON, N., PLENGE, R., WEINBLATT, M., SHADICK, N. and REICH, D. (2006). Principal components analysis corrects for stratification in genome-wide association studies. *nature genetics* **38** 904–909.
- [34] ROHE, K., CHATTERJEE, S. and YU, B. (2011). Spectral clustering and the high-dimensional stochastic blockmodel. Ann. Statist. 39 1878–1915.
- [35] ROYER, M. (2017). Adaptive clustering through semidefinite programming. Advances in Neural Information Processing Systems 1795–1803.
- [36] VEMPALA, V. and WANG, G. (2002). A spectral algorithm of learning mixtures of distributions. In *Proceedings of the 43rd IEEE FOCS*.
- [37] ZHA, H., HE, X., DING, C., SIMON, H. and GU, M. (2002). Spectral relaxation for k-means clustering. In Advances in Neural Information Processing Systems 14. MIT Press.
- [38] ZHOU, S. (2006). *Routing, disjoint Paths, and classification*. Ph.D. thesis, Carnegie Mellon University, Pittsburgh, PA.
- [39] ZHOU, S. (2023). Semidefinite programming on population clustering: a global analysis.