

Chiral-symmetric Topological Origin of Nonlinear Fixed Points

Di Zhou*

*Key Lab of Advanced Optoelectronic Quantum Architecture and Measurement (MOE) and School of Physics,
Beijing Institute of Technology, Beijing 100081, China and
Department of Physics, University of Illinois, Urbana-Champaign, 61801, USA*

Particle-hole symmetry and chiral symmetry play a pivotal role in multiple areas of physics, yet they remain un-studied in systems with nonlinear interactions that are beyond Kerr-type. Here, we establish these two non-spatial symmetries in systems with strong and general nonlinear interactions. Chiral symmetry ensures the quantization of the Berry phase of nonlinear normal modes and categorizes the topological phases of nonlinear dynamics. We show edge modes that serve as topologically protected fixed points of chiral-symmetric nonlinear dynamics. Our theoretical framework paves the way towards the topological classification of general nonlinear dynamics.

I. INTRODUCTION

Non-spatial symmetries govern the fundamental principles of physics in multiple areas. In high-energy physics, the combination of non-spatial particle-hole, parity, and time-reversal symmetries dictate the existence of anti-particles [1]. In quantum mechanics, non-spatial time-reversal symmetry demands that eigenstates with half-integer spins must be doubly degenerate, which is known as the Kramers' degeneracy [2, 3]. In soft matter and engineering physics, non-spatial chiral symmetry reveals the chiral image of mechanical floppy modes and states of self-stress [4–7], governing the mechanical failure and stability [8–11], respectively. Additionally, in condensed-matter physics, non-spatial time-reversal, particle-hole, and chiral symmetries quantize the Berry phase and classify topological phases of matter in a “ten-fold” way [12–15]. This classification enables fundamental understanding of symmetry-protected topological phases with potential applications for quantum information technology [14, 16–20].

Non-spatial symmetries have been the subject of extensive study in both linear [21–24] and nonlinear systems [25–36]. For instance, non-spatial time-reversal and parity symmetries have been thoroughly explored in the context of mechanical, electrical and photonic structures [37–44], enabling novel designs of microcavities [45], circuit metamaterials [46], and plasmonic waveguides [47]. Other non-spatial symmetries, such as non-spatial particle-hole symmetry and chiral symmetry, have been studied in linear systems and special nonlinear systems, such as the Kerr-nonlinearities [48–50] in optical lattices that allow for $U(1)$ -gauge symmetry on the phase variable of wave functions [51, 52], and Rock-Paper-Scissor cycles in zero-sum games [53–57]. However, the rigorous definition and establishment of (*anti-unitary*) particle-hole symmetry and (*unitary*) chiral symmetry have not been addressed in systems with general nonlinearities that go beyond Kerr-type.

Nonlinear interactions are ubiquitous in nature, such as nonlinear mechanical [37, 58–61] and electrical structures [62, 63], circadian rhythms of living cells [64], and quantum fluids in optical lattices [65]. These nonlinear mechanisms possess unique features that cannot be observed in linear systems, including soliton propagation [66–68], nonlinear localized modes [69], bifurcation [70], and chaos [71]. Given the significant influence of particle-hole symmetry and chiral symmetry on linear systems, it is intriguing to explore the unprecedented physics that can emerge when these two symmetries encounter general and strong nonlinear dynamics.

In this work, we extend the concept of particle-hole symmetry and chiral symmetry on general nonlinear dynamics, and investigate their nontrivial consequences on nonlinear topological physics. Our motivation derives from the history of linear topological insulators, where although extensive research had been conducted in topological physics, there continued to be fundamental importance in the “ten-fold” classification of symmetry-protected topological phases categorized by non-spatial symmetries [12, 13]. Such a classification significantly increases the potential for topological quantum computation. To explore the impact of non-spatial particle-hole and chiral symmetries on nonlinear topological physics, we investigate the geometric phase of nonlinear normal modes under adiabatic evolutions. This geometric phase has two components: the Berry phase of nonlinear normal modes [72, 73], which is the nonlinear extension of the conventional Berry phase found in linear systems, and an additional phase term that arises purely from the nonlinear interactions [74].

We find that, interestingly, this adiabatic geometric phase can be quantized by chiral symmetry, and defines the topologically trivial and non-trivial phases of the nonlinear dynamics. In the topologically non-trivial phase, nonlinear modes appear at the open boundaries of the system. These modes possess “topological protection” as they show resistance against disturbances to both the modes themselves and the nonlinear interactions. Furthermore, due to chiral symmetry, the frequencies of nonlinear boundary modes are pinned at zero. Consequently, these topologically robust static modes serve as

* dizhou@bit.edu.cn

the fixed points in wave function space with nonlinear stability. Fixed points are widely studied in statistical physics [75], Landau phase transition [76], and renormalization group [77]. To demonstrate the practical implications of our results, we investigate the behavior of a lattice subjected to quadratic nonlinear interactions, which are commonly found in Lotka-Volterra dynamical systems [78–80]. We use the non-spatial particle-hole and chiral symmetries to investigate the *static* nonlinear boundary modes in this example.

Given the significance of non-spatial symmetries in the classification of symmetry-protected topological phases in purely linear systems, the establishment of particle-hole and chiral symmetries in general nonlinear systems might lead to the development of a “ten-fold classification” for nonlinear topological systems. This could deepen our understanding of the behavior and properties of nonlinear dynamics.

The organization of this paper is as follows. Section II defines the model, which is the generalized nonlinear Schrödinger equations. Section III derives the adiabatic geometric phase in nonlinear normal modes. This adiabatic phase is separated into two parts: the Berry phase of nonlinear normal modes, and the component that is unique to nonlinear systems. In Section IV, we discuss two types of non-spatial symmetries: particle-hole symmetry and chiral symmetry. Notably, we demonstrate the quantization of the Berry phase of nonlinear normal modes under chiral symmetry. Section V investigates the topological phases in both linear and nonlinear regimes and discusses the application of our topological theory to nonlinear systems.

II. THE MODEL

Nonlinear dynamics, including mechanical [83–85] and electrical structures [62], water waves [86], optical and atomic lattices [87–92], nonlinear materials [93, 94], plasmonic systems [95], and biophysical processes [53, 96], can be described by the universal framework called the generalized nonlinear Schrödinger equations. To establish non-spatial symmetries within these equations, we consider a two-dimensional square lattice with the primitive vectors \hat{e}_x and \hat{e}_y , as illustrated in Fig. 1. The unit cells are identified by two integers n_x and n_y , corresponding to their positions $\mathbf{r} = n_x\hat{e}_x + n_y\hat{e}_y$. Within each unit cell, there exist two classical fields, namely $\Psi_{\mathbf{r}}^{(1)}$ and $\Psi_{\mathbf{r}}^{(2)}$. The dynamics of these classical variables are governed by the following generalized nonlinear Schrödinger equations,

$$\begin{aligned} i\partial_t \Psi_{\mathbf{r}}^{(1)} &= + \sum_{\mathbf{r}', i'=1,2} U(\Psi_{\mathbf{r}}^{(1)}, \Psi_{\mathbf{r}'}^{(i')}), \\ i\partial_t \Psi_{\mathbf{r}}^{(2)} &= - \sum_{\mathbf{r}', i'=1,2} U(\Psi_{\mathbf{r}}^{(2)}, \Psi_{\mathbf{r}'}^{(i')}). \end{aligned} \quad (1)$$




FIG. 1. The model of nonlinear dynamics. (a) The diatomic two-dimensional square lattice, with the primitive vectors \hat{e}_x and \hat{e}_y represented by the black arrows. (b) Red and blue dots mark the 1 and 2 sites within the unit cell. The nonlinear interactions within 1 or 2 sites (between 1 and 2 sites) are denoted by black solid lines (red dashed lines). We use the Lotka-Volterra-type nonlinearity as the example for our nonlinear topological physics. (c) The nonlinear band structure [81, 82] of the model with the Lotka-Volterra interactions specified in Eqs. (4), where $\epsilon = 0.001$. $g_x = 0.9$, $g_y = 0.1$, and mode amplitudes $A = 10\epsilon$. (d) Enlarged nonlinear band structure in (c) illustrates the relationships among time-reversal, particle-hole, and chiral partner modes.

Here, $U(\Psi_{\mathbf{r}}^{(i)}, \Psi_{\mathbf{r}'}^{(i')})$ represents general nonlinear polynomials with real-valued coefficients involving the classical fields $\Psi_{\mathbf{r}}^{(i)}$ and $\Psi_{\mathbf{r}'}^{(i')}$. To simplify the problem, we consider the summation $\sum_{\mathbf{r}'}$ only over the on-site and nearest-neighbor interactions, as indicated by the black solid and red dashed lines in Fig. 1(b), which represent couplings within the same type of classical fields (i.e., $\Psi_{\mathbf{r}}^{(i)}$ and $\Psi_{\mathbf{r}'}^{(i')}$ with $i = i'$) and between different types of fields (i.e., $i \neq i'$), respectively. We denote

$$\Psi = (\dots, \Psi_{\mathbf{r}}^{(1)}, \Psi_{\mathbf{r}}^{(2)}, \dots)^{\top} \quad (2)$$

as a column vector with $2N$ components, where N is the number of diatomic unit cells, and \top denotes matrix transpose. In a concise representation, Eqs. (1) can be written as

$$i\partial_t \Psi = H(\Psi) \quad (3)$$

where $H(\Psi)$ is a nonlinear mapping of Ψ , and is called the nonlinear Hamiltonian. It is a $2N \times 1$ column vector with each element given by $H_n(\Psi) = (-1)^{n+1} \sum_{\mathbf{r}'} \sum_{i'} U(\Psi_{\mathbf{r}}^{(\text{mod}(n-1,2)+1)}, \Psi_{\mathbf{r}'}^{(i')})$ for $n = 1, 2, \dots, 2N$. Without knowing the specific form of nonlinear interactions, this general model presented in Eqs. (1) already exhibits non-spatial symmetries and

quantized Berry phase of nonlinear normal modes, which we demonstrate in the Section IV.

This general model is capable of describing many physical systems, depending on the nature of the interactions involved. In the case where the interactions are purely linear in the field variables, the linearized model can capture the fermionic states within the Bogoliubov-de Gennes Hamiltonian of topological superconductors [97, 98]. When the nonlinear interaction between sites $\mathbf{r} = n_x \hat{e}_x + n_y \hat{e}_y$ and $\mathbf{r}' = n'_x \hat{e}_x + n'_y \hat{e}_y$ takes the form of quadratic Lotka-Volterra-type couplings,

$$U(\Psi_{\mathbf{r}}^{(i)}, \Psi_{\mathbf{r}'}^{(i')}) = (\epsilon + \Psi_{\mathbf{r}}^{(i)}) \Psi_{\mathbf{r}'}^{(i')} \left[(n_x - n'_x)(1 - \delta_{ii'}) + \left(-\delta_{n_x, n'_x} \delta_{n_y, n'_y} + \sum_{j=x, y} g_j |n_j - n'_j| \right) \delta_{ii'} \right], \quad (4)$$

the *static solutions* of this nonlinear Schrödinger-type dynamics can mimic the *static solutions* of the populations of species in a food web [78, 99, 100], which we elaborate later in Section V. In this scenario, ϵ represents the death and growth rates for the species $\Psi_{\mathbf{r}}^{(i)}$, while g_j for $j = x, y$ accounts for the nonlinear effects of the prey (predator) on the predator's (prey's) growth rate. These constant parameters are positive and real numbers, with $\epsilon \ll 1$ utilized to emphasize the significance of nonlinear effects. This nonlinear interaction is pictorially represented by the black solid and red dashed lines in Fig. 1(b). This Lotka-Volterra nonlinearity results in the simplest model that exhibits nonlinear topological physics. We will discuss this example in Section V.

III. ADIABATIC GEOMETRIC PHASE OF NONLINEAR NORMAL MODES

In this section, we derive the geometric phase that arises from the adiabatic evolution of nonlinear normal modes in generalized nonlinear Schrödinger equations.

When the amplitudes of classical fields are much smaller than ϵ , the nonlinear Hamiltonian can be linearized into a two-band matrix Hamiltonian $\hat{H}_{\mathbf{k}}$ [98]. This Hamiltonian describes the oscillation of classical variables governed by the eigenfrequency $\omega = \omega(\mathbf{k})$, where the wavevector $\mathbf{k} = (k_x, k_y)$ resides in the two-dimensional Brillouin zone.

Nonlinearities become increasingly important as mode amplitude rises, making nonlinear normal modes significantly deviate from sinusoidal waves of linear systems [62, 69, 73, 101]. Specifically, nonlinear normal modes with the plane-wave format satisfy the expression

$$\Psi_{\mathbf{k}, \omega} = (\Psi_{\mathbf{k}}^{(1)}(\mathbf{k} \cdot \mathbf{r} - \omega t), \Psi_{\mathbf{k}}^{(2)}(\mathbf{k} \cdot \mathbf{r} - \omega t + \phi_{\mathbf{k}}))^{\top} \quad (5)$$

according to the nonlinear extension of the Bloch theorem [69, 72]. Here, $\Psi_{\mathbf{k}}^{(1)}(\theta)$ and $\Psi_{\mathbf{k}}^{(2)}(\theta)$ represent the 2π -periodic non-sinusoidal functions, while $\phi_{\mathbf{k}}$ characterizes the relative phase between these wave components. The

frequencies of these nonlinear normal modes, denoted by $\omega = \omega(\mathbf{k}, A)$, are influenced by both the wavevectors \mathbf{k} and mode amplitudes A , deviating from their linear counterparts. It is worth emphasizing that in nonlinear systems, the number of nonlinear modes can exceed the degrees of freedom, allowing nonlinear localized modes [69, 102] and “looped band structures” [103–106] to emerge from the effect of bifurcation. However, in this work, our scope is limited to the simple case that bifurcation does not occur, and these additional excitations do not emerge from the nonlinear dynamics. Thus, plane-wave nonlinear normal modes can be uniquely defined based on their amplitude, wavevector, and frequency, allowing the nonlinear system to be effectively described as a “two-band nonlinear model.”

We adiabatically evolve the plane-wave nonlinear normal mode as the wavevector $\mathbf{k} = \mathbf{k}(t)$ follows a closed trajectory C in the Brillouin zone [107]. Based on the nonlinear extension of the adiabatic theorem [108, 109], at time t , the nonlinear normal mode follows the ansatz

$$\Psi = \Psi_{\mathbf{k}(t), \omega} \left(-\int_0^t \omega(t', \mathbf{k}(t')) dt' - \gamma(t) \right), \quad (6)$$

where $\gamma(t)$ denotes the phase shift of the nonlinear normal mode during the adiabatic evolution. When the wavevector \mathbf{k} completes a closed-loop travel in the Brillouin zone, the wave function acquires a phase γ_C , known as the adiabatic geometric phase [74, 110].

As per the Whitham modulation theory [111–113], during the adiabatic evolution of the wavevector \mathbf{k} in reciprocal space, the mode amplitude A and relative phase $\phi_{\mathbf{k}}$ (as defined in Eq. (5)) should change slowly. Consequently, the adiabatic phase, denoted by γ_C , can be separated into two components: $\gamma_C = \gamma_C^{(B)} + \gamma_C^{(NL)}$. The first term, $\gamma_C^{(B)}$, is referred to as the Berry phase of nonlinear normal modes, which originates from the change in the relative phase $\phi_{\mathbf{k}}$. On the other hand, the second term, $\gamma_C^{(NL)}$, arises from the change in the mode amplitude. In the upcoming discussion, we will examine $\gamma_C^{(B)}$ and $\gamma_C^{(NL)}$ individually.

As described in Appendix A, the derivation of the Berry phase of nonlinear normal modes involves computing the evolution of the relative phase $\phi_{\mathbf{k}}$ as the wavevector \mathbf{k} undergoes adiabatic changes:

$$\gamma_C^{(B)} = \oint_C \frac{\sum_l \left(l |\psi_{l\mathbf{k}}^{(2)}|^2 \nabla_{\mathbf{k}} \phi_{\mathbf{k}} - i \sum_{i=1,2} \psi_{l\mathbf{k}}^{(i)*} \nabla_{\mathbf{k}} \psi_{l\mathbf{k}}^{(i)} \right)}{\sum_{l'} l' \left(|\psi_{l'\mathbf{k}}^{(1)}|^2 + |\psi_{l'\mathbf{k}}^{(2)}|^2 \right)} \cdot d\mathbf{k}. \quad (7)$$

Here, $\psi_{l\mathbf{k}}^{(i)} = (2\pi)^{-1} \int_0^{2\pi} e^{-il\theta} \Psi_{\mathbf{k}}^{(i)} d\theta$ is the l -th Fourier component of the nonlinear wave, $\Psi_{\mathbf{k}}^{(i)}$, with $i = 1, 2$. We emphasize that for Schrödinger equations with linear [3] or Kerr-nonlinear interactions [114], the eigenmodes are sinusoidal waves, which reduces Eq. (7) to the Berry

phase in Ref. [74], $\gamma_{C,\text{linear}}^{(B)} = i \oint_C d\mathbf{k} \cdot \langle \Psi_{\mathbf{k},\omega} | \nabla_{\mathbf{k}} | \Psi_{\mathbf{k},\omega} \rangle$ (see Appendix A for details). In general, $\gamma_C^{(B)}$ is *not quantized* in systems without symmetry constraints. However, $\gamma_C^{(B)}$ can be quantized if non-spatial symmetries are incorporated in the system, which we address in Section IV. D.

The Whitham modulation theory [111–113] also indicates that mode amplitude should change during adiabatic evolution. This effect gives rise to an additional contribution to the adiabatic geometric phase [74], denoted as $\gamma_C^{(NL)}$. Analytical details of the corresponding calculations are provided in Appendix A. For Kerr-nonlinear systems, the change in mode amplitude equivalently corresponds to the change in the particle populations of Bose-Einstein condensates. $\gamma_C^{(NL)}$ has been shown to have important physical implications, such as the generation of an additional effective magnetic field in reciprocal space [74].

As we will demonstrate in Section IV. C, chiral symmetry imposes a constraint that causes the amplitudes of the two wave components of a nonlinear normal mode to be equal. This constraint results in the mode amplitude staying unchanged, up to the normalization factor of the wave function. Consequently, we have found that the adiabatic geometric phase $\gamma_C^{(NL)}$ vanishes under such constraints. This vanishing result is derived in Ref. [74] and in Appendix A. In contrast, the Berry phase of nonlinear normal modes, $\gamma_C^{(B)}$, is the only contribution to the adiabatic geometric phase. This Berry phase of nonlinear normal modes can be simplified to Kerr-nonlinear Berry phases for nonlinear photonic metamaterials [20, 26, 28, 30, 31, 115–118], and further characterizes their nonlinear topological phases under symmetry constraints. By adopting the similar approach, we aim to employ $\gamma_C^{(B)}$ as the adiabatic geometric phase to gain insights into the properties of general-nonlinear systems. This allows us to identify the presence of topologically protected modes and study their behavior under perturbations.

IV. PARTICLE-HOLE SYMMETRY, CHIRAL SYMMETRY, AND TOPOLOGICAL INDEX OF NONLINEAR SYSTEMS

In this section, we establish particle-hole symmetry and chiral symmetry for nonlinear dynamics. We show that non-spatial chiral symmetry can be used to quantify the Berry phase of nonlinear normal modes in the general model described by Eqs. (1). This quantization serves as a topological invariant that can be used to define the topologically trivial and non-trivial phases of the nonlinear dynamics. In Section V, we will demonstrate the presence of static nonlinear topological boundary modes within topologically non-trivial phases.

We briefly review time-reversal symmetry that has been well-established in nonlinear dynamics [48, 69, 119, 120]. The nonlinear system is time-reversal invari-

ant if the equations of motion remain unchanged under the time-reversal transformation $(\Psi_{\mathbf{r}}^{(1)}(t), \Psi_{\mathbf{r}}^{(2)}(t)) \rightarrow (\Psi_{\mathbf{r}}^{(1)*}(-t), \Psi_{\mathbf{r}}^{(2)*}(-t))$. This invariance arises from the real-coefficient polynomials of the nonlinear interactions in terms of the field variables. The formal expression of time-reversal symmetry is $\mathcal{T}H(\Psi) - H(\mathcal{T}\Psi) = 0$, where $H(\Psi)$ is the nonlinear Hamiltonian, and the time-reversal operator, $\mathcal{T} = \mathcal{K}; t \rightarrow -t$, involving complex conjugation and reversing the sign of time. Here, $H(\mathcal{T}\Psi)$ means that we perform the time-reversal operation on the nonlinear wave Ψ , and then we operate on $\mathcal{T}\Psi$ using the nonlinear Hamiltonian (nonlinear mapping) H . This symmetry implies that for a nonlinear wave $\Psi_{\mathbf{k},\omega}$ in a time-reversal symmetric model, there exists a time-reversed partner solution with the wavevector $-\mathbf{k}$, denoted as $\Psi_{-\mathbf{k},\omega} = \mathcal{T}\Psi_{\mathbf{k},\omega} = (\Psi_{\mathbf{k}}^{(1)*}(\mathbf{k} \cdot \mathbf{r} + \omega t), \Psi_{\mathbf{k}}^{(2)*}(\mathbf{k} \cdot \mathbf{r} + \omega t + \phi_{\mathbf{k}}))^{\top}$, as pictorially indicated by the red arrow and dot in Fig. 1(d).

The time-reversal operator satisfies $\mathcal{T}^2 = 1$ and is anti-unitary, aligning with the operator for linear Schrödinger equation of spinless particles [2, 121]. In the linear regime, time-reversal symmetry simplifies to the conventional form, $\mathcal{T}\hat{H}_{\mathbf{k}}\mathcal{T}^{-1} = \hat{H}_{-\mathbf{k}}$, where $\hat{H}_{\mathbf{k}}$ represents the linearized Hamiltonian in reciprocal space.

A. Particle-hole symmetry of general nonlinear systems

Here, we introduce an *anti-unitary* particle-hole operator and its corresponding symmetry for general nonlinear systems. The anti-unitary property of the particle-hole operator is critical in understanding linear topological insulators [98] and the topological properties of nonlinear systems.

In a particle-hole symmetric system, the motion equations in Eqs. (1) remain unchanged under the particle-hole transformation, $(\Psi_{\mathbf{r}}^{(1)}, \Psi_{\mathbf{r}}^{(2)}) \rightarrow (\Psi_{\mathbf{r}}^{(2)*}, \Psi_{\mathbf{r}}^{(1)*})$. This invariance is equivalently captured by the constraint,

$$\mathcal{C}H(\Psi) + H(\mathcal{C}\Psi) = 0, \quad \mathcal{C} = I_N \otimes \sigma_x \mathcal{K}, \quad (8)$$

where \mathcal{C} is the particle-hole operator, I_N is the $N \times N$ identity matrix, and σ_x is the Pauli matrix. One important characteristic of the particle-hole operator we have defined is its *anti-unitary* property, which arises from the complex conjugation involved in the operator. This property, in turn, results in the reversal of the frequency and momentum of the original nonlinear wave (see Eq. (5)) in the system under particle-hole transformation. Therefore, for a nonlinear normal mode $\Psi_{\mathbf{k},\omega}$ with the wavevector \mathbf{k} and frequency ω , the particle-hole-symmetric model has a corresponding nonlinear sister solution with a wavevector $-\mathbf{k}$ and frequency $-\omega$, denoted as

$$\begin{aligned} \Psi_{-\mathbf{k},-\omega} &= \mathcal{C}\Psi_{\mathbf{k},\omega} = \\ &(\Psi_{\mathbf{k}}^{(2)*}(\mathbf{k} \cdot \mathbf{r} - \omega t), \Psi_{\mathbf{k}}^{(1)*}(\mathbf{k} \cdot \mathbf{r} - \omega t - \phi_{\mathbf{k}}))^{\top}, \quad (9) \end{aligned}$$

as depicted by the black arrow in Fig. 1(d). The particle-hole operator satisfies $\mathcal{C}^2 = 1$ and is anti-unitary, aligning with the operator defined for linear systems [2, 98]. Furthermore, in the linear regime, Eqs. (8) simplify to the well-studied form of particle-hole symmetry, $\mathcal{C}\hat{H}_{\mathbf{k}}\mathcal{C}^{-1} = -\hat{H}_{-\mathbf{k}}$.

B. Chiral symmetry of general nonlinear systems

Chiral symmetry naturally arises in nonlinear systems when both time-reversal and particle-hole symmetries are present. The model described by Eqs. (1) remains unchanged under the chiral transformation $(\Psi_{\mathbf{r}}^{(1)}(t), \Psi_{\mathbf{r}}^{(2)}(t)) \rightarrow (\Psi_{\mathbf{r}}^{(2)}(-t), \Psi_{\mathbf{r}}^{(1)}(-t))$. Mathematically, chiral symmetry is expressed as the constraint on the nonlinear Hamiltonian given by

$$\mathcal{S}H(\Psi) + H(\mathcal{S}\Psi) = 0, \quad \mathcal{S} = \mathcal{T} \cdot \mathcal{C}, \quad (10)$$

where $\mathcal{S} = \mathcal{T} \cdot \mathcal{C}$ represents the chiral symmetry operator that combines the effects of time-reversal and particle-hole transformations. It is worth noting that the resulting chiral symmetry operator, which is a combination of time-reversal and particle-hole symmetry operators expressed as $\mathcal{S} = \mathcal{T} \cdot \mathcal{C}$, is *unitary* in nature. This is due to the anti-unitary nature of both time-reversal and particle-hole symmetry operators, which cancels out to produce a unitary chiral symmetry operator that is critical for characterizing the topological properties of the nonlinear system, which we address in the following subsection. Thus, for a nonlinear normal mode $\Psi_{\mathbf{k},\omega}$ with frequency ω , chiral symmetry predicts the existence of a partner solution with frequency $-\omega$, denoted as

$$\begin{aligned} \Psi_{\mathbf{k},-\omega} &= \mathcal{S}\Psi_{\mathbf{k},\omega} = \\ &(\Psi_{\mathbf{k}}^{(2)}(\mathbf{k} \cdot \mathbf{r} + \omega t), \Psi_{\mathbf{k}}^{(1)}(\mathbf{k} \cdot \mathbf{r} + \omega t - \phi_{\mathbf{k}}))^{\top}. \end{aligned} \quad (11)$$

This relationship is depicted by the green arrow and dot in Fig. 1(d). This chiral operator is unitary and yields $\mathcal{S}^2 = 1$, agreeing perfectly with the chiral operator for linear topological insulators. In the linear regime, Eqs. (10) reduce to the well-known form [121–123], $\mathcal{S}\hat{H}_{\mathbf{k}}\mathcal{S}^{-1} = -\hat{H}_{\mathbf{k}}$. Finally, our study highlights an important aspect of chiral-symmetric nonlinear systems. We find that the chiral-symmetric partner nonlinear mode swaps the two wave components of a nonlinear normal mode. This swapping leads to a result where the two wave components in a nonlinear normal mode share the same mode amplitude.

Our nonlinear model exhibits non-spatial time-reversal, particle-hole, and chiral symmetries, which is the extension of symmetry class BDI defined in the tenfold classification of linear topological insulators [122].

C. Berry phase of nonlinear normal modes quantized by chiral symmetry

In a purely linear Schrödinger equation, the summation of Berry phases across all energy bands is always zero due to the topological triviality of the fiber bundle [107] associated with a complete and orthogonal set of eigenbasis in a matrix Hamiltonian [107]. Combined with chiral symmetry, this sum rule naturally results in the quantization of linear Berry phase [124]. However, this conclusion does not hold for general nonlinear dynamics as nonlinear normal modes do not necessarily have completeness and orthogonality when matrix analysis fails. Therefore, the quantization of Berry phase of nonlinear normal modes in a chiral-symmetric nonlinear system, as expressed in Eq. (7), remains an open question.

Here, we demonstrate that under chiral symmetry, the Berry phase of nonlinear normal modes still remains quantized for nonlinear systems. Moreover, the quantization of the Berry phase of nonlinear normal modes suggests that it can serve as a potential topological index for characterizing the topological phases of the underlying nonlinear dynamics. We provide a more comprehensive discussion of this point in Section V.

To demonstrate that chiral symmetry is still capable of quantizing the Berry phase of nonlinear normal modes, we consider the nonlinear normal mode $\Psi_{\mathbf{k},-\omega} = \mathcal{S}\Psi_{\mathbf{k},\omega}$, which is the chiral-partner mode of $\Psi_{\mathbf{k},\omega}$. We perform an adiabatic evolution on this chiral-partner mode by slowly varying the wavevector $\mathbf{k}(t)$ along a closed-loop trajectory \mathcal{C} in the Brillouin zone. When we adiabatically evolve the chiral-symmetric partner mode, $\Psi_{\mathbf{k},-\omega}$, this mode can be considered as having a frequency of ω but with the arrow of time reversed. Consequently, the mode acquires a term $\gamma(t)$ in its phase variable, via

$$\Psi(t) = \Psi_{\mathbf{k}(t),-\omega} \left(\int_0^t \omega(t', \mathbf{k}(t')) dt' + \gamma(t) \right). \quad (12)$$

Substituting this result into the nonlinear motion equations, namely $i\partial_t \Psi_{\mathbf{k}(t),-\omega} = H(\Psi_{\mathbf{k}(t),-\omega})$, we can compute the geometric phase when the wavevector \mathbf{k} travels along the trajectory \mathcal{C} . This adiabatic evolution allows the chiral-symmetric nonlinear mode to obtain the adiabatic geometric phase, $\gamma_{\mathcal{C}} = \gamma_{\mathcal{C}}^{(B)}$ (note that $\gamma_{\mathcal{C}}^{(NL)} = 0$ in chiral-symmetric systems, as shown in Appendix A), from which the Berry phase of nonlinear normal modes, $\gamma_{\mathcal{C}}^{(B)}$, is obtained:

$$\gamma_{\mathcal{C}}^{(B)} = \oint_{\mathcal{C}} \frac{\sum_l \left(l |\psi_{l\mathbf{k}}^{(1)}|^2 \nabla_{\mathbf{k}} \phi_{\mathbf{k}} + i \sum_{i=1,2} \psi_{l\mathbf{k}}^{(i)*} \nabla_{\mathbf{k}} \psi_{l\mathbf{k}}^{(i)} \right)}{\sum_{l'} l' \left(|\psi_{l'\mathbf{k}}^{(1)}|^2 + |\psi_{l'\mathbf{k}}^{(2)}|^2 \right)} \cdot d\mathbf{k}. \quad (13)$$

Since both Eq. (7) and (13) describe the same Berry phase of nonlinear normal modes under the same evolu-

tion trajectory, we equate them, and obtain the result

$$\gamma_C^{(B)} = \frac{1}{2} \oint_C \nabla_{\mathbf{k}} \phi_{\mathbf{k}} \cdot d\mathbf{k} = n\pi, \quad n = 0, 1. \quad (14)$$

This equation demonstrates the quantization of the Berry phase of nonlinear normal modes under the constraint of chiral symmetry, which serves as the topological invariant of the considered nonlinear dynamics. $n = 0, 1$ correspond to the nonlinear topologically trivial and non-trivial phases, respectively. Due to the quantized nature of this Berry phase, it cannot change continuously upon the variations of system parameters, including the mode amplitudes and coupling parameters. Below, we leverage the invariance of this topological number to investigate the corresponding nonlinear topological edge modes.

V. NONLINEAR TOPOLOGICAL PHASES AND FIXED POINTS

We have demonstrated the topological invariance of the Berry phase of nonlinear normal modes using chiral symmetry. In this section, we exemplify the impact of this topological number on the nonlinear physics, where the interactions are specified as the example in Eqs. (4) with the quadratic nonlinearities. Specifically, we investigate the topological phases and the corresponding boundary physics of the system in both the linear and nonlinear regimes.




FIG. 2. Topological phases and boundary modes. (a) Nonlinear topological phase diagram of the model with Lotka-Volterra nonlinearities. (b) Brillouin zone with the integration trajectories of Berry phases of nonlinear normal modes indicated. (c) Disks' size and color represent the spatial profile of the weakly nonlinear topological edge mode. (d) Spatial distribution of the topological boundary mode in the highly nonlinear regime.

A. Topological phases and boundary modes in the linear regime

When the quadratic nonlinearities are small compared to the linear parts, the nonlinear system can be approximated as linear Schrödinger equations. This regime is valid when mode amplitudes $A \ll \epsilon$. This allows us to perform a momentum-space decomposition and simplify the linearized equations of motion using the equation, $i\partial_t \Psi_{\mathbf{k},\omega} = \hat{H}_{\mathbf{k},\omega} \Psi_{\mathbf{k},\omega}$, where the linear wave function is represented as $\Psi_{\mathbf{k},\omega} = (\Psi_{\mathbf{k}}^{(1)}, \Psi_{\mathbf{k}}^{(2)})^\top$, the matrix Hamiltonian, $\hat{H}_{\mathbf{k},\omega}$, reads

$$\hat{H}_{\mathbf{k},\omega} = 2\epsilon\sigma_y \sin k_x + \epsilon\sigma_z \left(-1 + 2 \sum_{j=x,y} g_j \cos k_j \right), \quad (15)$$

where $\mathbf{k} = (k_x, k_y)$ is the wavevector within the two-dimensional Brillouin zone, and $\sigma_{x,y,z}$ are Pauli matrices. This Hamiltonian possesses chiral symmetry, namely a unitary chiral symmetry operator $\mathcal{S} = \sigma_x$ anti-commutes with it [98, 121]: $\{\sigma_x, \hat{H}_{\mathbf{k},\omega}\} = 0$.

The Hamiltonian $\hat{H}_{\mathbf{k},\omega}$ has multiple phases. Specifically, when the interaction parameters are in the grey region of Fig. 2(a) (i.e., the interaction parameters satisfy $|g_y| \geq |g_x - 1/2| \cup |g_y| \geq |g_x + 1/2|$), the system becomes gapless, as the linear band structures touch at a pair of zero-frequency points. As a result, the linear Berry phase becomes singular and ill-defined when the integration trajectory passes through these gapless points. When the linear bands touch at the gapless points, the corresponding static eigenstates are given by

$$\Psi_{\mathbf{r}}(t) = (1, \pm 1)^\top e^{i\mathbf{k}_w \cdot \mathbf{r}} / \sqrt{2}, \quad (16)$$

whose frequency and wavevector are $\omega = 0$ and $\pm \mathbf{k}_w$, respectively. For the parameters that lie in the region of $|g_y| \geq |g_x - 1/2|$, we have the wavevectors $\mathbf{k}_w = (0, \arccos[(1 - 2g_x)/2g_y])$. When the parameters are in the region $|g_y| \geq |g_x + 1/2|$, the wavevectors are given by $\mathbf{k}_w = (\pi, \arccos[(1 + 2g_x)/2g_y])$. We note that the zero-frequency nature of these gapless bulk states stems from the chiral symmetry [125].

On the other hand, in the blue region of Fig. 2(a) (the parameters yield $|g_y| < |g_x - 1/2| \cap |g_y| < |g_x + 1/2|$), and in the red region of Fig. 2(a) (with the parameters that yield $-g_x + 1/2 < g_y < g_x - 1/2 \cup g_x + 1/2 < g_y < -g_x - 1/2$), the lattice is fully gapped and the linear Berry phase becomes well-defined:

$$\lim_{A \rightarrow 0} \gamma_C^{(B)}(A) = i \oint_C d\mathbf{k} \cdot \langle \Psi_{\mathbf{k},\omega} | \nabla_{\mathbf{k}} | \Psi_{\mathbf{k},\omega} \rangle, \quad (17)$$

where C is the closed-loop trajectory for the adiabatic evolution of the wavevector \mathbf{k} in the two-dimensional Brillouin zone. Specifically, we define two types of closed trajectories, $C(k_y) : k_x = -\pi \rightarrow \pi; k_y$ and $C(k_x) : k_y = -\pi \rightarrow \pi; k_x$, which integrate over all k_x for a given k_y and all k_y for a given k_x , respectively. The linear Berry

phase corresponding to the horizontal trajectory $C(k_y)$ is denoted as $\gamma_{C(k_y)}$, while that corresponding to the vertical trajectory $C(k_x)$ is denoted as $\gamma_{C(k_x)}$.

Due to the chiral symmetric nature of the linearized Hamiltonian [98], these Berry phases, namely $\gamma_{C(k_y)}$ and $\gamma_{C(k_x)}$, are guaranteed to have integer multiples of π . Moreover, due to the fully gapped nature of the linear band structure, the Berry phases $\gamma_{C(k_y)}$ and $\gamma_{C(k_x)}$ remain unchanged for any k_x and k_y ranging from $-\pi$ to π . To characterize the topological phases of the linearized model, we define the ‘‘2D polarization [126]’’ as a measure of the linear Berry phases averaged by the Brillouin zone,

$$\lim_{A \rightarrow 0} \mathbf{R}_T(A) = \frac{1}{2\pi^2} \int_{-\pi}^{\pi} \left[\lim_{A \rightarrow 0} \gamma_{C(k_y)}^{(B)}(A) dk_y \hat{e}_x + \lim_{A \rightarrow 0} \gamma_{C(k_x)}^{(B)}(A) dk_x \hat{e}_y \right]. \quad (18)$$

When the parameters are in the red region of Fig. 2(a), the topological polarization is given by $\lim_{A \rightarrow 0} \mathbf{R}_T(A) = \hat{e}_x$, indicating that the linear system is in the topological phase. On the other hand, when the parameters are within the blue region of Fig. 2(a), the topological polarization becomes $\lim_{A \rightarrow 0} \mathbf{R}_T(A) = 0$, indicating that the system is in the topologically trivial regime.

According to the principle of bulk-boundary correspondence in topological band theory, the behavior of the system at the boundary is determined by the topological invariant derived from the bulk bands of the lattice. In the red region of Fig. 2(a), the topological polarization vector takes the value $\lim_{A \rightarrow 0} \mathbf{R}_T(A) = \hat{e}_x$, indicating the emergence of topologically protected boundary modes in the system. These boundary modes can be analytically solved using the Jackiw-Rebbi-type solution,

$$\begin{pmatrix} \Psi_{\mathbf{r}}^{(1)}(t) \\ \Psi_{\mathbf{r}}^{(2)}(t) \end{pmatrix} = A e^{ik_y n_y} (e^{-\kappa_+(k_y) n_x} - e^{-\kappa_-(k_y) n_x}) \begin{pmatrix} 1 \\ -1 \end{pmatrix}, \quad (19)$$

where A is the mode amplitude, k_y is the wave number in the transverse y -direction, and the spatial decay rates $\kappa_{\pm}(k_y)$ yield $\kappa_{\pm}(k_y) > 0$. This edge mode is exponentially localized on the left open boundary of the square lattice. In contrast, in the topologically trivial phase with the parameters that lie in the blue region of Fig. 2(a), the topological polarization vector takes the trivial value of $\lim_{A \rightarrow 0} \mathbf{R}_T(A) = 0$, indicating the absence of topological boundary modes in this linear non-topological phase.

The non-spatial chiral symmetry of the linearized Hamiltonian leads to an important property in the frequency spectrum of the bulk modes. This ensures that the frequencies of the bulk modes arise in pairs of $\pm\omega$, reflecting the chiral symmetry of the Hamiltonian. This property in the bulk mode frequencies has significant implications for the frequencies of topological edge states

in the system. In particular, the frequency of the topological edge state is constrained to be pinned at zero. This is because, if the frequency of the topological state, ω , is nonzero, then a partner topological state with the frequency of $-\omega$ must also arise at the same boundary. These two boundary states can couple and open a band gap on the lattice boundary, violating the topological protection of the edge states. Therefore, topological edge states must have zero frequency and do not evolve in time, satisfying the static condition $\partial_t \Psi_{\mathbf{r}}^{(i)} = 0$ for $i = 1, 2$ and for all \mathbf{r} . The static nature of topological boundary modes remains valid for the nonlinear system as chiral symmetry extends to the fully nonlinear regime.

B. Topological phases and boundary modes in the nonlinear regime

We now consider the topological phases and the corresponding boundary modes in the Schrödinger-type equations in the nonlinear regime.

Intriguingly, even in the strongly nonlinear regime, the system remains gapless when the parameters are in the grey-shaded regime of Fig. 2(a). This is because nonlinear zero-frequency bulk modes can still arise in the nonlinear regime of the system, maintaining the system’s gapless nature. The nonlinear static bulk modes can be analytically obtained by imposing the condition $\partial_t \Psi_{\mathbf{r}}^{(i)} = 0$ for $i = 1, 2$ and for all \mathbf{r} , because chiral symmetry assures the frequency of the nonlinear gapless mode to stay at zero. The nonlinear mode in the lattice system is described by Eq. (16), which is identical to that obtained from the linearized model. The presence of zero-frequency nonlinear bulk modes in the parameter region defined by the grey regime of Fig. 2(a) yields the closure of the nonlinear band gap, which we define as the ‘‘nonlinear gapless phase’’ of the system.

The system with parameters in the blue and red regimes of Fig. 2(a) is in the nonlinear fully gapped phase, because these phases are devoid of the previously discussed zero-frequency nonlinear bulk modes. This property allows the topological numbers of the system to be well-defined and invariant as mode amplitudes rise. Thus, we define the 2D topological polarization of the lattice in the fully nonlinear regime:

$$\mathbf{R}_T(A) = \frac{1}{2\pi^2} \int_{-\pi}^{\pi} \left[\gamma_{C(k_y)}^{(B)}(A) dk_y \hat{e}_x + \gamma_{C(k_x)}^{(B)}(A) dk_x \hat{e}_y \right]. \quad (20)$$

The red-shaded parameter regime shown in Fig. 2(a) corresponds to the nonlinear topological phase of the system described by the nonlinear Schrödinger equation in Eqs. (1). In this regime, the topological polarization takes the form $\mathbf{R}_T(A) = \hat{e}_x$, indicating the presence of nonlinear topological boundary modes at the edges of the lattice. We can calculate the spatial profile of these boundary modes by imposing static boundary conditions

and solving for the corresponding eigenstates. The resulting static nonlinear mode is described by Eq. (19) and is exponentially localized on the left open boundary of the lattice. In the blue-shaded regime of Fig. 2(a), the system is in the nonlinear topologically trivial phase as described by the nonlinear Schrödinger equation in Eqs. (1). The topological polarization in this regime takes the form $\mathbf{R}_T(A) = 0$, indicating that there are no nonlinear topological boundary modes present at the edges of the lattice.

A remarkable feature of the system under consideration is its topological invariance under increasing mode amplitudes in both the non-trivial and trivial phases. This phenomenon arises from the fact that mode amplitudes have a global effect on the nonlinear dynamics of the system and do not alter the topological properties of the lattice. This explains the invariance of the topological Berry phases and polarization for growing mode amplitudes.

C. Applications to the fixed points of nonlinear dynamics

To further explore the implications of the previous results for nonlinear systems, we investigate the emergence and characteristics of strongly nonlinear topological edge modes in the network of Lotka-Volterra nonlinear interactions. These interactions provide insights for the population dynamics of biological systems. To perform our investigation, we use the equations of motion from Eqs. (1) and employ a generalized Wick rotation [127] on them. The generalized Wick rotation involves changing the time variable and the wave function as follows:

$$(t, \Psi_{\mathbf{r}}^{(i)}) \rightarrow (it, (-1)^{i+1} \Psi_{\mathbf{r}}^{(i)}), \quad i = 1, 2. \quad (21)$$

As a result, Eqs. (1) are now transformed into the following nonlinear motion equations,

$$\partial_t \Psi_{\mathbf{r}}^{(i)} = \sum_{\mathbf{r}', i'=1,2} U((-1)^{i+1} \Psi_{\mathbf{r}}^{(i)}, (-1)^{i'+1} \Psi_{\mathbf{r}'}^{(i')}), \quad (22)$$

for $i = 1, 2$. Here, ϵ represents the growth rate, and g_x, g_y represent the effect of prey (predator) on the growth rate of the predator (prey).

Despite the differences between the *dynamical properties* [128] of the Schrödinger-type nonlinear equations in Eqs. (1) and the “Wick-rotated” equations in Eq. (22), their *static* properties are remarkably similar. The reason is as follows.

In the Schrödinger equation, chiral symmetry ensures that the frequencies of nonlinear normal modes emerge in pairs of $\pm\omega$. Therefore, nonlinear topological mode must have zero-frequency, because if the frequency of the topological mode were non-zero, a chiral-partner topological mode with the frequency of $-\omega$ must appear. These two topological modes can interfere and break their topological nature. As a result, the nonlinear topological mode must be static.

This static nonlinear topological mode is equivalent to setting $i\partial_t \Psi_{\mathbf{r}}^{(i)} = 0$ in the Schrödinger equation. On the other hand, the stationary solutions in the Lotka-Volterra model are obtained by setting $\partial_t \Psi_{\mathbf{r}}^{(i)} = 0$. This finding suggests that the (Schrödinger) static mode is the same as the (Lotka-Volterra) static mode, because these two zero-frequency solutions are obtained by substituting $i\partial_t \Psi_{\mathbf{r}}^{(i)} = 0$ and $\partial_t \Psi_{\mathbf{r}}^{(i)} = 0$ in the Schrödinger and Lotka-Volterra models, respectively.

The notion of drawing a comparison between the static solution of the Lotka-Volterra and Schrödinger models originates from seminal works [4–11], which established an analogy between static mechanics in Newtonian equations of motion and the static solutions in chiral-symmetric Schrödinger equations. The topological properties of Newtonian mechanics are defined by introducing an auxiliary chiral-symmetric Schrödinger equation, where we can compute the topological index of this auxiliary Schrödinger equation. We then utilize this auxiliary Schrödinger topological number to describe the topology of static Newtonian mechanics. However, it should be noted that the dynamical features of these two models differ markedly.

Building upon this notion, we observe that the static solutions for both the Schrödinger and Lotka-Volterra models can be obtained by setting $\partial_t \Psi_{\mathbf{r}}^{(i)} = 0$ for $i = 1, 2$ and all \mathbf{r} . This analogy facilitates the derivation of the static nonlinear boundary mode found in Eq. (22). Thus, we can construct the phase diagram of the Lotka-Volterra-type model, as depicted in Fig. 2(a). This phase diagram shares similarities with that of Schrödinger-type equations, comprising three distinct regions: the nonlinear gapless, topologically non-trivial, and trivial phases, represented by the grey, red, and blue-shaded areas, respectively.

The Lotka-Volterra-type model exhibits topologically distinct boundary properties in the red and blue areas of the phase diagram. In the red parameter region, the model exhibits the emergence of nonlinear static edge modes from the boundary of the square lattice. These edge modes are in line with the derived nonlinear topological polarization of $\mathbf{R}_T(A) = \hat{e}_x$. It should be noted that the population of species must be real and positive numbers, which imposes a constraint on our analysis. Under this constraint, we obtain the nonlinear boundary mode

$$\begin{pmatrix} \Psi_{\mathbf{r}}^{(1)}(t) \\ \Psi_{\mathbf{r}}^{(2)}(t) \end{pmatrix} = A(e^{-\kappa_+(k_y=0)n_x} - e^{-\kappa_-(k_y=0)n_x}) \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad (23)$$

where A is the mode amplitude. The spatial decay rate satisfies $\kappa_{\pm}(k_y = 0) > 0$, and is explicitly expressed in Appendix B. This result indicates that the nonlinear boundary mode is exponentially localized on the left open boundary of the system, confirming its topological nature. The spatial decay rate of the nonlinear boundary mode is in perfect agreement with the decay properties

of the topological edge modes in the linear and nonlinear regimes of the Schrödinger-type equations in Eqs. (1). Moreover, the nonlinear boundary mode corresponds to the nonlinear topological polarization, $\mathbf{R}_T(A) = \hat{e}_x$, in the topologically non-trivial phase. Due to the static and nonlinear nature of these topological edge modes, we refer to them as “topologically protected fixed points”. In contrast to the red region, the blue region of the phase diagram of the Lotka-Volterra-type model is devoid of nonlinear topological static modes. This behavior is in line with the derived trivial nonlinear topological polarization, $\mathbf{R}_T(A) = 0$.




FIG. 3. The temporal evolutions of topologically protected boundary modes are shown in (a) and (b) for the weakly nonlinear and fully nonlinear regimes, respectively. In both cases, the modes evolve for $t = 100T$ and still maintain their stability.

To validate the analytical results of the emergence of nonlinear topological boundary modes in the Lotka-Volterra-type model, we perform numerical simulations and present the results in Figs. 2(c,d) and Fig. 3. The numerical simulations confirm the emergence of the nonlinear topological modes on the open boundary that cuts the topological polarization vector $\mathbf{R}_T(A) = \hat{e}_x$, as predicted by the analytical results. This topologically protected fixed point exhibits high stability in both temporal and spatial aspects, highlighting its robustness against small perturbations or fluctuations. We initialize the mode with $\pm 10\%$ spatial fluctuations and $\pm 10\%$ spatial variations in the nonlinear interactions, as shown by the bond lengths in Figs. 2(c,d). The numerical simulations indicate that the nonlinear topological mode remains highly stable even after $t = 10^2T$ of self-oscillation, both in the weakly and strongly nonlinear regimes. The characteristic period of the nonlinear normal mode in the fully nonlinear regime is $T = 2\pi$. The stability of the nonlinear topological mode confirms its robustness and reliability against small perturbations or fluctuations.

Chiral symmetry has notable effects on these nonlinear topological boundary modes. Firstly, this symmetry locks the frequencies of topological modes at zero, regardless of their amplitudes. These static nonlinear modes behave as stable fixed points in the $2N$ -dimensional space constituted of the wave function components. These static modes are distinct from spatial

symmetry-induced topological modes, which are sensitive to amplitudes and prone to losing nonlinear stability [37, 81]. Secondly, chiral symmetric nonlinear topological modes are unaffected by the breakdown of spatial symmetries, as observed in Figs. 2(c,d). Conversely, spatial symmetry-induced topological modes are quickly disrupted by spatial symmetry-breaking boundary conditions. Due to these exceptional attributes, we term non-spatial symmetry-induced modes as “topologically protected nonlinear fixed points,” signifying their nonlinear, static nature and resilience to external disturbances.

VI. CONCLUSIONS AND OUTLOOK

In this work, we establish non-spatial particle-hole symmetry and chiral symmetry in systems with strong and general nonlinear interactions. Chiral symmetry quantizes the Berry phase of nonlinear normal modes, determining their topological phases, and facilitating the emergence of nonlinear topological boundary modes. These modes have pinned frequencies at zero, making them “topologically protected nonlinear fixed points” in the wave function space. They exhibit high stability against disruptions in spatial symmetries. These general results are illustrated in a nonlinear example with Lotka-Volterra-type interactions.

Our work enables the non-spatial classification of nonlinear systems, expanding the ten-fold classification of linear topological insulators [122]. Our findings may suggest potential applications in nonlinear dynamics, where the stability conferred by nonlinear topology could aid in resilience against parameter changes.

Note added— Recently, we became aware of a related independent effort on the chiral symmetry of nonlinear dynamics [129].

VII. ACKNOWLEDGMENT

D. Z. would like to thank insightful discussions with Anthony J. Leggett, Xueda Wen, Junyi Zhang and Feng Li. D. Z. is supported by the National Natural Science Foundation of China (Grant Nos. 12374157, 12102039).

Appendix A: Berry phase of nonlinear normal modes

In this section, we derive the adiabatic geometric phase of nonlinear normal modes. We consider the nonlinear equations of motion in a translationally invariant lattice, with $i\partial_t\Psi(t) = H(\Psi)$, in which the plane-wave nonlinear normal mode reads $\Psi_{\mathbf{k},\omega}(\mathbf{k}\cdot\mathbf{r} - \omega t)$. We consider an adiabatic evolution to the nonlinear mode by slowly changing the wavevector \mathbf{k} that follows a closed-loop trajectory in

the Brillouin zone. Under the evolution, the nonlinear wave takes the format of

$$\Psi = \Psi_{\mathbf{k},\omega} \left(- \int^t \omega(t') dt' - \gamma(t) \right) \quad (\text{A1})$$

where $\gamma(t)$ is the adiabatic geometric phase that arises from the temporal evolution. This mode should satisfy the nonlinear equations of motion, which leads to

$$i\partial_t \Psi = i \left[\frac{\partial \Psi_{\mathbf{k},\omega}}{\partial \mathbf{k}} \frac{d\mathbf{k}}{dt} - \left(\omega + \frac{d\gamma}{dt} \right) \frac{\partial \Psi_{\mathbf{k},\omega}}{\partial \theta} + \partial_t \delta \Psi \right], \quad (\text{A2})$$

where $\delta \Psi = \Psi - \Psi_{\mathbf{k},\omega}$ denotes mode change, and $\theta = \int^t \omega(t') dt' + \gamma(t)$ denotes the phase variable of the nonlinear mode. Since the shape of the nonlinear mode can vary during the process of adiabatic evolution, the nonlinear Hamiltonian also changes in this adiabatic process, leading to the expansion of the nonlinear Hamiltonian in terms of the mode variation $\delta \Psi$:

$$H(\Psi) = H(\Psi_{\mathbf{k}}) + \left(\delta \Psi \frac{\partial H}{\partial \Psi} + \delta \Psi^* \frac{\partial H}{\partial \Psi^*} \right)_{\Psi_{\mathbf{k},\omega}}. \quad (\text{A3})$$

Employing the relationship $-i\omega \partial_\theta \Psi_{\mathbf{k},\omega} = H(\Psi_{\mathbf{k},\omega})$, we combine Eqs. (A2, A3) and obtain the equations of motion for the nonlinear mode, which read

$$\frac{\partial \Psi_{\mathbf{k},\omega}}{\partial \theta} \frac{d\gamma}{dt} = \frac{\partial \Psi_{\mathbf{k},\omega}}{\partial \mathbf{k}} \frac{d\mathbf{k}}{dt} + i \left(\delta \Psi \frac{\partial H}{\partial \Psi} + \delta \Psi^* \frac{\partial H}{\partial \Psi^*} - i\partial_t \delta \Psi \right)_{\Psi_{\mathbf{k},\omega}}. \quad (\text{A4})$$

Eq. (A4) contains two terms on the right-hand side. The first term corresponds to the Berry phase of nonlinear normal modes in nonlinear normal modes. The second term arises from the change in the mode shape and amplitude due to the adiabatic evolution, which can affect the nonlinear Hamiltonian of the system. Together, these terms contribute to the overall adiabatic geometric phase of the nonlinear normal modes.

To gain a more detailed understanding of the nonlinear normal mode, we express it in terms of a Fourier series. Specifically, we write $\Psi_{\mathbf{k},\omega}(\theta) = \sum_l (\psi_{l\mathbf{k}}^{(1)}, \psi_{l\mathbf{k}}^{(2)} e^{il\phi_{\mathbf{k}}})^\top e^{il\theta}$, where $\psi_{l\mathbf{k}}^{(1)}$ and $\psi_{l\mathbf{k}}^{(2)}$ are the Fourier components of the mode in the two wave components, respectively, and $\phi_{\mathbf{k}}$ is the relative phase between the two components. Thus, we obtain

$$\begin{aligned} \frac{\partial \Psi_{\mathbf{k},\omega}}{\partial \theta} &= \sum_l \left(\psi_{l\mathbf{k}}^{(1)}, \psi_{l\mathbf{k}}^{(2)} e^{il\phi_{\mathbf{k}}} \right)^\top i l e^{il\theta}, \\ \frac{\partial \Psi_{\mathbf{k},\omega}}{\partial \mathbf{k}} &= \sum_l \left[\frac{\partial \psi_{l\mathbf{k}}^{(1)}}{\partial \mathbf{k}}, \left(\frac{\partial \psi_{l\mathbf{k}}^{(2)}}{\partial \mathbf{k}} + i l \psi_{l\mathbf{k}}^{(2)} \frac{\partial \phi_{\mathbf{k}}}{\partial \mathbf{k}} \right) e^{il\phi_{\mathbf{k}}} \right]^\top e^{il\theta}. \end{aligned} \quad (\text{A5})$$

We now use Eqs. (A5) to substitute results into Eq. (A4). We then multiply both sides of Eq. (A4) by $\Psi_{\mathbf{k},\omega}^\dagger$

and integrate over the phase variable θ from 0 to 2π . This procedure leads us to derive the following relationship:

$$\begin{aligned} \frac{d\gamma}{dt} \sum_l \sum_j |\psi_{l\mathbf{k}}^{(i)}|^2 &= \\ \frac{d\mathbf{k}}{dt} \sum_l \left(l \frac{\partial \phi_{\mathbf{k}}}{\partial \mathbf{k}} |\psi_{l\mathbf{k}}^{(2)}|^2 - i \sum_j \psi_{l\mathbf{k}}^{(j)*} \frac{\partial \psi_{l\mathbf{k}}^{(i)}}{\partial \mathbf{k}} \right) &+ \\ \int_0^{2\pi} \frac{d\theta}{2\pi} \Psi_{\mathbf{k},\omega}^\dagger \left(\delta \Psi \frac{\partial H}{\partial \Psi} + \delta \Psi^* \frac{\partial H}{\partial \Psi^*} - H(\Psi) \delta \Psi \right)_{\Psi_{\mathbf{k},\omega}} & \quad (\text{A6}) \end{aligned}$$

where the relationship $i\partial_t \delta \Psi = H(\Psi) \delta \Psi$ has been adopted. Finally, we integrate over the time variable t and obtain the Berry phase of nonlinear normal modes:

$$\gamma_C = \gamma_C^{(B)} + \gamma_C^{(NL)}. \quad (\text{A7})$$

In Eq. (A7), there are two contributions in the adiabatic geometric phase. The first term is the Berry phase of nonlinear normal modes,

$$\gamma_C^{(B)} = \oint_C d\mathbf{k} \frac{\sum_l \left(l \frac{\partial \phi_{\mathbf{k}}}{\partial \mathbf{k}} |\psi_{l\mathbf{k}}^{(2)}|^2 - i \sum_j \psi_{l\mathbf{k}}^{(j)*} \frac{\partial \psi_{l\mathbf{k}}^{(i)}}{\partial \mathbf{k}} \right)}{\sum_l l \sum_j |\psi_{l\mathbf{k}}^{(i)}|^2}. \quad (\text{A8})$$

The second term

$$\begin{aligned} \gamma_C^{(NL)} &= \int dt \left(\sum_l l \sum_j |\psi_{l\mathbf{k}}^{(i)}|^2 \right)^{-1} \\ \int_0^{2\pi} \frac{d\theta}{2\pi} \Psi_{\mathbf{k},\omega}^\dagger \left(\delta \Psi \frac{\partial H}{\partial \Psi} + \delta \Psi^* \frac{\partial H}{\partial \Psi^*} - H(\Psi) \delta \Psi \right)_{\Psi_{\mathbf{k},\omega}} & \quad (\text{A9}) \end{aligned}$$

is known as the ‘‘purely nonlinear’’ contribution to the adiabatic geometric phase, as it arises solely from the nonlinear interactions within the system. Specifically, it represents the change in the nonlinear Hamiltonian with respect to the variation of the nonlinear normal mode, which arises due to the dependence of the Hamiltonian on the mode amplitude. We emphasize that this term vanishes for the purely linear Schrödinger-type equations.

To investigate the relationship between the conventional linear Berry phase and Berry phase of nonlinear normal modes, we consider the Hamiltonian for the linear and Kerr-nonlinear Schrödinger equations, which is given by

$$H(\Psi) = H_0 \Psi + g |\Psi|^2 \Psi. \quad (\text{A10})$$

Here, H_0 is the linear part of the dynamics, represented by a matrix, and g is the coefficient of the Kerr-nonlinearity. In these systems, the nonlinear normal modes can be represented by sinusoidal waves,

$$\begin{aligned} \Psi_{\mathbf{k},\omega}(\mathbf{k} \cdot \mathbf{r} - \omega t) &= \Psi_{\mathbf{k}} e^{i(\mathbf{k} \cdot \mathbf{r} - \omega t)} \\ &= (\Psi_{\mathbf{k}}^{(1)}, \Psi_{\mathbf{k}}^{(2)} e^{i\phi_{\mathbf{k}}})^\top e^{i(\mathbf{k} \cdot \mathbf{r} - \omega t)}. \end{aligned} \quad (\text{A11})$$

Here, the normalization condition given by $|\Psi_{\mathbf{k}}^{(1)}|^2 + |\Psi_{\mathbf{k}}^{(2)}|^2 = 1$ has been adopted.

To simplify the Berry phase of nonlinear normal modes, $\gamma_C^{(B)}$, and the purely nonlinear contribution to the adiabatic geometric phase, $\gamma_C^{(NL)}$, we employ the relationships given by the representation of nonlinear normal modes as sinusoidal waves, as presented in Eq. (A11). In particular, we note that nonlinear plane waves in these systems contain only the fundamental harmonic, which allows us to simplify the expressions for $\psi_{l\mathbf{k}}^{(1)}$ and $\psi_{l\mathbf{k}}^{(2)}$ as $\psi_{l\mathbf{k}}^{(1)} = \Psi_{\mathbf{k}}^{(1)} \delta_{l1}$ and $\psi_{l\mathbf{k}}^{(2)} = \Psi_{\mathbf{k}}^{(2)} \delta_{l1}$, respectively. Plugging these simplifications into $\gamma_C^{(B)}$ reduces the Berry phase of nonlinear normal modes as

$$\begin{aligned}
\gamma_C^{(B)} &= \oint_C d\mathbf{k} \cdot \frac{\sum_l \left(l |\psi_{l\mathbf{k}}^{(2)}|^2 \frac{\partial \phi_{\mathbf{k}}}{\partial \mathbf{k}} + i \sum_j \psi_{l\mathbf{k}}^{(j)*} \frac{\partial \psi_{l\mathbf{k}}^{(i)}}{\partial \mathbf{k}} \right) \delta_{l1}}{\sum_{l'} l' \left(\sum_{j'} |\psi_{l'\mathbf{k}}^{(j')}|^2 \right) \delta_{l'1}} \\
&= \oint_C d\mathbf{k} \cdot \left(|\psi_{1\mathbf{k}}^{(2)}|^2 \frac{\partial \phi_{\mathbf{k}}}{\partial \mathbf{k}} + i \sum_j \psi_{1\mathbf{k}}^{(j)*} \frac{\partial \psi_{1\mathbf{k}}^{(i)}}{\partial \mathbf{k}} \right) \\
&= \oint_C d\mathbf{k} \cdot i \left(\sum_j \Psi_{\mathbf{k}}^{(j)*} \partial_{\mathbf{k}} \Psi_{\mathbf{k}}^{(i)} - i |\Psi_{\mathbf{k}}^{(2)}|^2 \partial_{\mathbf{k}} \phi_{\mathbf{k}} \right) \\
&= \oint_C d\mathbf{k} \cdot i (\Psi_{\mathbf{k}}^{(1)*}, \Psi_{\mathbf{k}}^{(2)*} e^{-i\phi_{\mathbf{k}}}) \partial_{\mathbf{k}} \begin{pmatrix} \Psi_{\mathbf{k}}^{(1)} \\ \Psi_{\mathbf{k}}^{(2)} e^{i\phi_{\mathbf{k}}} \end{pmatrix} \\
&= \oint_C d\mathbf{k} \cdot i \langle \Psi_{\mathbf{k}} | \partial_{\mathbf{k}} | \Psi_{\mathbf{k}} \rangle = \gamma_{C,\text{linear}}^{(B)}, \quad (\text{A12})
\end{aligned}$$

where $\Psi_{\mathbf{k}} = (\Psi_{\mathbf{k}}^{(1)}, \Psi_{\mathbf{k}}^{(2)} e^{i\phi_{\mathbf{k}}})^T$ is the eigenvector of the Hamiltonian, and $\gamma_{C,\text{linear}}^{(B)}$ denotes the conventional form of Berry phase in linear systems. At this point, we have shown that the Berry phase of nonlinear normal modes, as expressed in Eq. (7), can be reduced to the linear Berry phase when the nonlinear interactions are specified as Kerr-type.

When considering Kerr nonlinear interactions of the form

$$H(\Psi) = H_0 \Psi + g |\Psi|^2 \Psi, \quad (\text{A13})$$

we can compute the contribution of nonlinear interactions to the adiabatic geometric phase, $\gamma_C^{(NL)}$. By using the relationships $\partial H(\Psi)/\partial \Psi = H_0 + 2g |\Psi|^2$ and $\partial H(\Psi)/\partial \Psi^* = g \Psi^2$, we can substitute into $\gamma_C^{(NL)}$ to ob-

tain the result

$$\begin{aligned}
\gamma_C^{(NL)} &= \int dt \left(\sum_l l \sum_j |\psi_{l\mathbf{k}}^{(i)}|^2 \delta_{l1} \right)^{-1} \\
&\quad \int_0^{2\pi} \frac{d\theta}{2\pi} \Psi_{\mathbf{k},\omega}^\dagger \left(\delta \Psi \frac{\partial H}{\partial \Psi} + \delta \Psi^* \frac{\partial H}{\partial \Psi^*} - H(\Psi) \delta \Psi \right)_{\Psi_{\mathbf{k},\omega}} \\
&= \int dt \int_0^{2\pi} \frac{d\theta}{2\pi} \\
&\quad \Psi_{\mathbf{k},\omega}^\dagger \left(\delta \Psi \frac{\partial H}{\partial \Psi} + \delta \Psi^* \frac{\partial H}{\partial \Psi^*} - H(\Psi) \delta \Psi \right)_{\Psi_{\mathbf{k},\omega}} \\
&= \int dt g (\Psi^2 \Psi^\dagger \delta \Psi^\dagger + \Psi^{\dagger 2} \Psi \delta \Psi)_{\Psi_{\mathbf{k},\omega}}. \quad (\text{A14})
\end{aligned}$$

Appendix B: Analytical results of nonlinear topological boundary modes

In this section, we analyze the decay rate of the nonlinear boundary mode by setting the static condition, $\partial \Psi_{\mathbf{r}}^{(i)}/\partial t = 0$, and simplifying the Schrödinger-type nonlinear equations of motion, as given by

$$\begin{aligned}
\Psi_{\mathbf{r}}^{(1)} - \sum_{j=x,y} g_j (\Psi_{\mathbf{r}+\hat{e}_j}^{(1)} + \Psi_{\mathbf{r}-\hat{e}_j}^{(1)}) - \Psi_{\mathbf{r}+\hat{e}_x}^{(2)} + \Psi_{\mathbf{r}-\hat{e}_x}^{(2)} &= 0, \\
\Psi_{\mathbf{r}}^{(2)} - \sum_{j=x,y} g_j (\Psi_{\mathbf{r}+\hat{e}_j}^{(2)} + \Psi_{\mathbf{r}-\hat{e}_j}^{(2)}) - \Psi_{\mathbf{r}+\hat{e}_x}^{(1)} + \Psi_{\mathbf{r}-\hat{e}_x}^{(1)} &= 0.
\end{aligned} \quad (\text{B1})$$

We analyze the behavior of the nonlinear boundary mode, which is exponentially localized on the left open boundary and takes the form of a plane wave in the transverse y direction. Using this ansatz, we derive the wave amplitudes and substitute from the static motion equations given by Eq. (B1). The resulting solution, as given by Eq. (19), provides an analytic expression for the spatial decay rates of the edge mode, which are expressed as

$$\begin{aligned}
\kappa_{\pm}(k_y) &= \\
&= -\ln \left[\frac{\frac{1}{2} - g_y \cos k_y \pm \sqrt{(\frac{1}{2} - g_y \cos k_y)^2 + 1 - g_x^2}}{g_x + 1} \right]. \quad (\text{B2})
\end{aligned}$$

The analytical expression for the spatial decay rates of the nonlinear boundary mode, as given by Eq. (B2), reveals important insights into the behavior of the mode at different values of the nonlinearity parameters and the transverse wavevector. In particular, we find that within the red region of parameter space in Fig. 2(a), where $-g_x + 1/2 < g_y < g_x - 1/2 \cup g_x + 1/2 < g_y < -g_x - 1/2$, the decay rates are positive, indicating that the mode described by Eq. (19) is exponentially localized on the left open boundary of the square lattice.

To analyze the dynamics of the nonlinear system, we Wick-rotate the generalized nonlinear Schrödinger equations (see Eq. (21)), resulting in a Lotka-Volterra-type

quadratic nonlinear equations of motion. As a result, the static equations of motion for this Lotka-Volterra-type motion equations, are now expressed as

$$\begin{aligned}\Psi_{\mathbf{r}}^{(1)} - \sum_{j=x,y} g_j(\Psi_{\mathbf{r}+\hat{e}_j}^{(1)} + \Psi_{\mathbf{r}-\hat{e}_j}^{(1)}) + \Psi_{\mathbf{r}+\hat{e}_x}^{(2)} - \Psi_{\mathbf{r}-\hat{e}_x}^{(2)} &= 0, \\ \Psi_{\mathbf{r}}^{(2)} - \sum_{j=x,y} g_j(\Psi_{\mathbf{r}+\hat{e}_j}^{(2)} + \Psi_{\mathbf{r}-\hat{e}_j}^{(2)}) + \Psi_{\mathbf{r}+\hat{e}_x}^{(1)} - \Psi_{\mathbf{r}-\hat{e}_x}^{(1)} &= 0.\end{aligned}\tag{B3}$$

This equations of motion is similar to Eq. (B1) but differs by a sign on $\Psi_{\mathbf{r}}^{(2)}$. This static equations of motion can be solved analytically by imposing the ansatz $\Psi_{\mathbf{r}\pm\hat{e}_y}^{(i)} = e^{\pm ik_y} \Psi_{\mathbf{r}}^{(i)}$. As a result, we obtain a nonlinear boundary mode of the following form,

$$\begin{pmatrix} \Psi_{\mathbf{r}}^{(1)}(t) \\ \Psi_{\mathbf{r}}^{(2)}(t) \end{pmatrix} = A e^{ik_y n_y} (e^{-\kappa_+(k_y)n_x} - e^{-\kappa_-(k_y)n_x}) \begin{pmatrix} 1 \\ 1 \end{pmatrix}.\tag{B4}$$

When analyzing the population of species in the system, it is important to note that the classical fields must

take real and positive values. This constraint cannot be satisfied by the solution given by Eq. (B4) due to the complex-valuedness of the exponential factor $e^{ik_y n_y}$, which prohibits the boundary mode from being real and positive. To overcome this constraint, we restrict the transverse wave number k_y to take the value of $k_y = 0$. By imposing this restriction, we obtain a new nonlinear boundary mode solution, as given by Eq. (23), which is real and positive and satisfies the constraints imposed by the classical fields. The spatial decay rates of this mode is given by

$$\kappa_{\pm}(k_y = 0) = -\ln \left(\frac{\frac{1}{2} - g_y \pm \sqrt{\frac{5}{4} - g_y + g_y^2 - g_x^2}}{g_x + 1} \right).\tag{B5}$$

We find that the nonlinear boundary mode solution given by Eq. (23) is exponentially localized on the left boundary of the lattice for parameter values in the red region of Fig. 2(a). Specifically, the spatial decay rates given by Eq. (B5) always satisfy $\kappa_{\pm}(k_y = 0) > 0$, indicating that the amplitude of the mode decays exponentially with distance in the longitudinal direction.

-
- [1] J. Eades and F. J. Hartmann, “Forty years of antiprotons,” *Rev. Mod. Phys.* **71**, 373–419 (1999).
 - [2] Xiao-Liang Qi and Shou-Cheng Zhang, “Topological insulators and superconductors,” *Rev. Mod. Phys.* **83**, 1057–1110 (2011).
 - [3] Di Xiao, Ming-Che Chang, and Qian Niu, “Berry phase effects on electronic properties,” *Rev. Mod. Phys.* **82**, 1959–2007 (2010).
 - [4] TC Lubensky, CL Kane, Xiaoming Mao, Anton Souslov, and Kai Sun, “Phonons and elasticity in critically coordinated lattices,” *Reports on Progress in Physics* **78**, 073901 (2015).
 - [5] CL Kane and TC Lubensky, “Topological boundary modes in isostatic lattices,” *Nature Physics* **10**, 39 (2014).
 - [6] Di Zhou, Leyou Zhang, and Xiaoming Mao, “Topological edge floppy modes in disordered fiber networks,” *Phys. Rev. Lett.* **120**, 068003 (2018).
 - [7] Anne S. Meeussen, Jayson Paulose, and Vincenzo Vitelli, “Geared topological metamaterials with tunable mechanical stability,” *Physical Review X* **6**, 041029–1–041029–14 (2016).
 - [8] Kai Sun and Xiaoming Mao, “Continuum theory for topological edge soft modes,” *Phys. Rev. Lett.* **124**, 207601 (2020).
 - [9] Jayson Paulose, Bryan Gin-ge Chen, and Vincenzo Vitelli, “Topological modes bound to dislocations in mechanical metamaterials,” *Nature Physics* **11**, 153–156 (2015).
 - [10] Di Zhou, Leyou Zhang, and Xiaoming Mao, “Topological boundary floppy modes in quasicrystals,” *Phys. Rev. X* **9**, 021054 (2019).
 - [11] Ze-Guo Chen Hao Ge Cheng He Xin Li Ming-Hui Lu Xiao-Chen Sun Si-Yuan Yu Chen, Yan-Feng and Xiujuan Zhang, “Various topological phases and their abnormal effects of topological acoustic metamaterials,” *Interdisciplinary Materials* **2**, 179–230 (2023).
 - [12] Ching-Kai Chiu, Jeffrey C. Y. Teo, Andreas P. Schnyder, and Shinsei Ryu, “Classification of topological quantum matter with symmetries,” *Rev. Mod. Phys.* **88**, 035005 (2016).
 - [13] Andreas P Schnyder, Shinsei Ryu, Akira Furusaki, and Andreas WW Ludwig, “Classification of topological insulators and superconductors in three spatial dimensions,” *Physical Review B* **78**, 195125 (2008).
 - [14] M. Z. Hasan and C. L. Kane, “Colloquium: Topological insulators,” *Rev. Mod. Phys.* **82**, 3045–3067 (2010).
 - [15] W. P. Su, J. R. Schrieffer, and A. J. Heeger, “Solitons in polyacetylene,” *Phys. Rev. Lett.* **42**, 1698–1701 (1979).
 - [16] Yasuhiro Hatsugai, “Chern number and edge states in the integer quantum hall effect,” *Phys. Rev. Lett.* **71**, 3697–3700 (1993).
 - [17] Kazuki Sone and Yuto Ashida, “Anomalous topological active matter,” *Phys. Rev. Lett.* **123**, 205502 (2019).
 - [18] Taylor L. Hughes, Emil Prodan, and B. Andrei Bernevig, “Inversion-symmetric topological insulators,” *Phys. Rev. B* **83**, 245132 (2011).
 - [19] Yasuhiro Hatsugai, “Edge states in the integer quantum hall effect and the riemann surface of the bloch function,” *Phys. Rev. B* **48**, 11851–11862 (1993).
 - [20] Zhaoju Yang, Eran Lustig, Gal Harari, Yonatan Plotnik, Yaakov Lumer, Miguel A Bandres, and Mordechai Segev, “Mode-locked topological insulator laser utilizing synthetic dimensions,” *Physical Review X* **10**, 011059 (2020).
 - [21] Marcela Peláez, Urko Reinoso, Julien Serreau, Matthieu

- Tissier, and Nicolás Wschebor, “Spontaneous chiral symmetry breaking in the massive landau gauge: Realistic running coupling,” *Phys. Rev. D* **103**, 094035 (2021).
- [22] Chao-Hsiang Sheu and Mikhail Shifman, “Consistency of chiral symmetry breaking in chiral yang-mills theory with adiabatic continuity,” *Phys. Rev. D* **107**, 054030 (2023).
- [23] Mikael C Rechtsman, Julia M Zeuner, Yonatan Plotnik, Yaakov Lumer, Daniel Podolsky, Felix Dreisow, Stefan Nolte, Mordechai Segev, and Alexander Szameit, “Photonic floquet topological insulators,” *Nature* **496**, 196–200 (2013).
- [24] Sankar Das Sarma, Michael Freedman, and Chetan Nayak, “Majorana zero modes and topological quantum computation,” *npj Quantum Information* **1**, 1–13 (2015).
- [25] Kazuki Sone, Yuto Ashida, and Takahiro Sagawa, “Topological synchronization of coupled nonlinear oscillators,” arXiv preprint arXiv:2012.09479 (2020).
- [26] Oded Zilberberg, Sheng Huang, Jonathan Guglielmon, Mohan Wang, Kevin P Chen, Yaacov E Kraus, and Mikael C Rechtsman, “Photonic topological boundary pumping as a probe of 4d quantum hall physics,” *Nature* **553**, 59–62 (2018).
- [27] Jiho Noh, Wladimir A Benalcazar, Sheng Huang, Matthew J Collins, Kevin P Chen, Taylor L Hughes, and Mikael C Rechtsman, “Topological protection of photonic mid-gap defect modes,” *Nature Photonics* **12**, 408–415 (2018).
- [28] Seabrat Mukherjee and Mikael C Rechtsman, “Observation of unidirectional solitonlike edge states in nonlinear floquet topological insulators,” *Physical Review X* **11**, 041057 (2021).
- [29] Yonatan Plotnik, Mikael C Rechtsman, Daohong Song, Matthias Heinrich, Julia M Zeuner, Stefan Nolte, Yaakov Lumer, Natalia Malkova, Jingjun Xu, Alexander Szameit, *et al.*, “Observation of unconventional edge states in ‘photonic graphene,’” *Nature materials* **13**, 57–62 (2014).
- [30] Jiho Noh, Sheng Huang, Daniel Leykam, Yi Dong Chong, Kevin P Chen, and Mikael C Rechtsman, “Experimental observation of optical weyl points and fermi arc-like surface states,” *Nature Physics* **13**, 611–617 (2017).
- [31] Yaakov Lumer, Yonatan Plotnik, Mikael C Rechtsman, and Mordechai Segev, “Self-localized states in photonic topological insulators,” *Physical review letters* **111**, 243905 (2013).
- [32] Rivka Bekenstein, Ran Schley, Maor Mutzafi, Carmel Rotschild, and Mordechai Segev, “Optical simulations of gravitational effects in the newton–schrodinger system,” *Nature Physics* **11**, 872–878 (2015).
- [33] Yonatan Sharabi, Hanan Herzig Sheinflux, Yoav Sagi, Gadi Eisenstein, and Mordechai Segev, “Self-induced diffusion in disordered nonlinear photonic media,” *Physical Review Letters* **121**, 233901 (2018).
- [34] Yuval Lamhot, Assaf Barak, Or Peleg, and Mordechai Segev, “Self-trapping of optical beams through thermophoresis,” *Physical review letters* **105**, 163906 (2010).
- [35] Motohiko Ezawa, “Nonlinear non-hermitian higher-order topological laser,” *Phys. Rev. Res.* **4**, 013195 (2022).
- [36] Motohiko Ezawa, “Nonlinear dynamical topological phases in a cooper-pair box array,” *Phys. Rev. B* **108**, 075412 (2023).
- [37] Joshua R. Tempelman, Kathryn H. Matlack, and Alexander F. Vakakis, “Topological protection in a strongly nonlinear interface lattice,” *Phys. Rev. B* **104**, 174306 (2021).
- [38] Mark J Ablowitz and Justin T Cole, “Nonlinear optical waveguide lattices: Asymptotic analysis, solitons, and topological insulators,” *Physica D: Nonlinear Phenomena* **440**, 133440 (2022).
- [39] Farzad Zangeneh-Nejad and Romain Fleury, “Nonlinear second-order topological insulators,” *Phys. Rev. Lett.* **123**, 053902 (2019).
- [40] F Kh Abdullaev and BA Umarov, “Exact solitonic solutions for optical media with χ (2) nonlinearity and pt-symmetric potentials,” in *Journal of Physics: Conference Series*, Vol. 553 (IOP Publishing, 2014) p. 012001.
- [41] Yaakov Lumer, Yonatan Plotnik, Mikael C Rechtsman, and Mordechai Segev, “Nonlinearly induced p t transition in photonic systems,” *Physical review letters* **111**, 263901 (2013).
- [42] Mark J Ablowitz and Ziad H Musslimani, “Integrable nonlocal nonlinear schrödinger equation,” *Physical review letters* **110**, 064105 (2013).
- [43] Nan Lu, Panayotis G Kevrekidis, and Jesús Cuevas-Maraver, “-symmetric sine-gordon breathers,” *Journal of Physics A: Mathematical and Theoretical* **47**, 455101 (2014).
- [44] Mark J Ablowitz, Joel B Been, and Lincoln D Carr, “Fractional integrable nonlinear soliton equations,” *Physical Review Letters* **128**, 184101 (2022).
- [45] Li Fan, Jian Wang, Leo T Varghese, Hao Shen, Ben Niu, Yi Xuan, Andrew M Weiner, and Minghao Qi, “An all-silicon passive optical diode,” *Science* **335**, 447–450 (2012).
- [46] Joseph Schindler, Ang Li, Mei C Zheng, Fred M Ellis, and Tsampikos Kottos, “Experimental study of active lrc circuits with pt symmetries,” *Physical Review A* **84**, 040101 (2011).
- [47] Henri Benisty, Aloyse Degiron, Anatole Lupu, André De Lustrac, Sébastien Chénais, Sébastien Forget, Mondher Besbes, Grégory Barbillon, Aurélien Bruyant, Sylvain Blaize, *et al.*, “Implementation of pt symmetric devices using plasmonics: principle and applications,” *Optics Express* **19**, 18004–18019 (2011).
- [48] Tomoki Ozawa, Hannah M Price, Alberto Amo, Nathan Goldman, Mohammad Hafezi, Ling Lu, Mikael C Rechtsman, David Schuster, Jonathan Simon, Oded Zilberberg, *et al.*, “Topological photonics,” *Reviews of Modern Physics* **91**, 015006 (2019).
- [49] Lucien Jezequel and Pierre Delplace, “Nonlinear edge modes from topological one-dimensional lattices,” *Physical Review B* **105**, 035410 (2022).
- [50] Raditya Weda Bomantara, Wenlei Zhao, Longwen Zhou, and Jiangbin Gong, “Nonlinear dirac cones,” *Physical Review B(R)* **96**, 121406 (2017).
- [51] Kazuki Sone, Motohiko Ezawa, Yuto Ashida, Nobuyuki Yoshioka, and Takahiro Sagawa, “Nonlinearity-induced topological phase transition characterized by the nonlinear chern number,” arXiv preprint arXiv:2307.16827 (2023).
- [52] Kazuki Sone, Yuto Ashida, and Takahiro Sagawa, “Topological synchronization of coupled nonlinear os-

- cillators,” *Phys. Rev. Res.* **4**, 023211 (2022).
- [53] Johannes Knebel, Philipp M. Geiger, and Erwin Frey, “Topological phase transition in coupled rock-paper-scissors cycles,” *Phys. Rev. Lett.* **125**, 258301 (2020).
- [54] Muhammad Umer and Jiangbin Gong, “Topologically protected dynamics in three-dimensional nonlinear antisymmetric lotka-volterra systems,” *Phys. Rev. B* **106**, L241403 (2022).
- [55] Philipp M. Geiger, Johannes Knebel, and Erwin Frey, “Topologically robust zero-sum games and pfaffian orientation: How network topology determines the long-time dynamics of the antisymmetric lotka-volterra equation,” *Phys. Rev. E* **98**, 062316 (2018).
- [56] Tomonari Mizoguchi Yoshida, Tsuneya and Yasuhiro Hatsugai, “Chiral edge modes in evolutionary game theory: A kagome network of rock-paper-scissors cycles,” *Physical Review E* **104**, 025003 (2021).
- [57] Tsuneya Yoshida, Tomonari Mizoguchi, and Yasuhiro Hatsugai, “Non-hermitian topology in rock-paper-scissors games,” *Scientific Reports* **12**, 560 (2022).
- [58] Po-Wei Lo, Christian D. Santangelo, Bryan Gin-ge Chen, Chao-Ming Jian, Krishanu Roychowdhury, and Michael J. Lawler, “Topology in nonlinear mechanical systems,” *Phys. Rev. Lett.* **127**, 076802 (2021).
- [59] Ke Liu and Glaucio H. Paulino, “Nonlinear mechanics of non-rigid origami: an efficient computational approach,” *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* **473**, 20170348 (2017).
- [60] Matheus I. N. Rosa, Raj Kumar Pal, José R. F. Arruda, and Massimo Ruzzene, “Edge states and topological pumping in spatially modulated elastic lattices,” *Phys. Rev. Lett.* **123**, 034301 (2019).
- [61] Fangyuan Ma, Zheng Tang, Xiaotian Shi, Ying Wu, Jinkyu Yang, Di Zhou, Yugui Yao, and Feng Li, “Nonlinear topological mechanics in elliptically geared isotactic metamaterials,” *Phys. Rev. Lett.* **131**, 046101 (2023).
- [62] Hendrik Hohmann, Tobias Hofmann, Tobias Helbig, Stefan Imhof, Hauke Brand, Lavi K. Upreti, Alexander Stegmaier, Alexander Fritzsche, Tobias Müller, Udo Schwingenschlögl, Ching Hua Lee, Martin Greiter, Laurens W. Molenkamp, Tobias Kießling, and Ronny Thomale, “Observation of cnoidal wave localization in nonlinear topoelectric circuits,” *Phys. Rev. Res.* **5**, L012041 (2023).
- [63] You Wang, Li-Jun Lang, Ching Hua Lee, Baile Zhang, and YD Chong, “Topologically enhanced harmonic generation in a nonlinear transmission line metamaterial,” *Nature communications* **10**, 1–7 (2019).
- [64] Rachel S Edgar, Edward W Green, Yuwei Zhao, Gerben Van Ooijen, Maria Olmedo, Ximing Qin, Yao Xu, Min Pan, Utham K Valekunja, Kevin A Feeney, *et al.*, “Peroxiredoxins are conserved markers of circadian rhythms,” *Nature* **485**, 459–464 (2012).
- [65] Sergey Alyatkin, Helgi Sigurdsson, Alexis Askitopoulos, Julian D Töpfer, and Pavlos G Lagoudakis, “Quantum fluids of light in all-optical scatterer lattices,” *Nature Communications* **12**, 5571 (2021).
- [66] Ross Parker, Alejandro Aceves, Jesús Cuevas-Maraver, and P. G. Kevrekidis, “Standing and traveling waves in a model of periodically modulated one-dimensional waveguide arrays,” *Phys. Rev. E* **108**, 024214 (2023).
- [67] Ross Parker, Alejandro Aceves, Jesús Cuevas-Maraver, and P. G. Kevrekidis, “Standing and traveling waves in a model of periodically modulated one-dimensional waveguide arrays,” *Phys. Rev. E* **108**, 024214 (2023).
- [68] Yiming Pan, Moshe-Ishay Cohen, and Mordechai Segev, “Superluminal k-gap solitons in nonlinear photonic time crystals,” *Physical Review Letters* **130**, 233801 (2023).
- [69] Alexander F Vakakis, Leonid I Manevitch, Yuri V Mikhlin, Valery N Pilipchuk, and Alexandr A Zevin, *Normal modes and localization in nonlinear systems* (Springer, 2001).
- [70] Michel Fruchart, Ryo Hanai, Peter B Littlewood, and Vincenzo Vitelli, “Non-reciprocal phase transitions,” *Nature* **592**, 363–369 (2021).
- [71] J. P. Eckmann and D. Ruelle, “Ergodic theory of chaos and strange attractors,” *Rev. Mod. Phys.* **57**, 617–656 (1985).
- [72] Di Zhou, D. Zeb Rocklin, Michael Leamy, and Yugui Yao, “Topological Invariant and Anomalous Edge Modes of Strongly Nonlinear Systems,” *Nature Communications* **13**, 3379 (2022).
- [73] Jijie Tang, Fangyuan Ma, Feng Li, Honglian Guo, and Di Zhou, “Strongly nonlinear topological phases of cascaded topoelectrical circuits,” *Frontiers of Physics* **18**, 33311 (2023).
- [74] J. Liu and L. B. Fu, “Berry phase in nonlinear systems,” *Phys. Rev. A* **81**, 052112 (2010).
- [75] Robert Savit, “Duality in field theory and statistical systems,” *Rev. Mod. Phys.* **52**, 453–487 (1980).
- [76] Baruch Rosenstein and Dingping Li, “Ginzburg-landau theory of type ii superconductors in magnetic field,” *Rev. Mod. Phys.* **82**, 109–168 (2010).
- [77] Michael E. Fisher, “Renormalization group theory: Its basis and formulation in statistical physics,” *Rev. Mod. Phys.* **70**, 653–681 (1998).
- [78] NARENDRA S. GOEL, SAMARESH C. MAITRA, and ELLIOTT W. MONTROLL, “On the volterra and other nonlinear models of interacting populations,” *Rev. Mod. Phys.* **43**, 231–276 (1971).
- [79] Guy Bunin, “Ecological communities with lotka-volterra dynamics,” *Phys. Rev. E* **95**, 042414 (2017).
- [80] Léon Brenig, “Complete factorisation and analytic solutions of generalized lotka-volterra equations,” *Physics Letters A* **133**, 378–382 (1988).
- [81] Matthew D Fronk and Michael J Leamy, “Higher-order dispersion, stability, and waveform invariance in nonlinear monoatomic and diatomic systems,” *Journal of Vibration and Acoustics* **139** (2017).
- [82] See Supplementary Information for the derivation of nonlinear Berry phase, non-spatial symmetries, topological fixed point, and topological phase transitions..
- [83] Rajesh Chaunsali, Haitao Xu, Jinkyu Yang, Panayotis G. Kevrekidis, and Georgios Theocharis, “Stability of topological edge states under strong nonlinear effects,” *Phys. Rev. B* **103**, 024106 (2021).
- [84] Rajesh Chaunsali and Georgios Theocharis, “Self-induced topological transition in phononic crystals by nonlinearity management,” *Phys. Rev. B* **100**, 014302 (2019).
- [85] Rajesh Chaunsali, Panayotis G. Kevrekidis, Dimitri Frantzeskakis, and Georgios Theocharis, “Dirac solitons and topological edge states in the β -fermi-pastulam-tsingou dimer lattice,” *Phys. Rev. E* **108**, 054224 (2023).

- [86] Shaofeng Li, Suhui Qian, Hui Chen, Jinbao Song, and Anzhou Cao, “An extended nonlinear schrödinger equation for water waves with linear shear flow, wind, and dissipation,” *AIP Advances* **11** (2021).
- [87] Daria Smirnova, Sergey Kruk, Daniel Leykam, Elizaveta Melik-Gaykazyan, Duk-Yong Choi, and Yuri Kivshar, “Third-harmonic generation in photonic topological metasurfaces,” *Physical review letters* **123**, 103901 (2019).
- [88] Pankaj Bhalla, Kamal Das, Dimitrie Culcer, and Amit Agarwal, “Resonant second-harmonic generation as a probe of quantum geometry,” *Phys. Rev. Lett.* **129**, 227401 (2022).
- [89] Daria Smirnova, Daniel Leykam, Yidong Chong, and Yuri Kivshar, “Nonlinear topological photonics,” *Applied Physics Reviews* **7**, 021306 (2020).
- [90] J. Perczel, J. Borregaard, D. E. Chang, H. Pichler, S. F. Yelin, P. Zoller, and M. D. Lukin, “Topological quantum optics in two-dimensional atomic arrays,” *Phys. Rev. Lett.* **119**, 023603 (2017).
- [91] Nader Mostaan, Fabian Grusdt, and Nathan Goldman, “Quantized topological pumping of solitons in nonlinear photonics and ultracold atomic mixtures,” *nature communications* **13**, 5997 (2022).
- [92] Chao Hang, Dmitry A Zezyulin, Guoxiang Huang, and Vladimir V Konotop, “Nonlinear topological edge states in a non-hermitian array of optical waveguides embedded in an atomic gas,” *Physical Review A* **103**, L040202 (2021).
- [93] Qiong Ma, Adolfo G Grushin, and Kenneth S Burch, “Topology and geometry under the nonlinear electromagnetic spotlight,” *Nature materials* **20**, 1601–1614 (2021).
- [94] Stephan Wong, Terry A. Loring, and Alexander Cerjan, “Probing topology in nonlinear topological materials using numerical k -theory,” *Phys. Rev. B* **108**, 195142 (2023).
- [95] Jeffrey B Parker, JW Burby, JB Marston, and Steven M Tobias, “Nontrivial topology in the continuous spectrum of a magnetized plasma,” *Physical Review Research* **2**, 033425 (2020).
- [96] Sergey Pirogov, Alexander Rybko, Anastasia Kalinina, and Mikhail Gelfand, “Recombination processes and nonlinear markov chains,” *Journal of Computational Biology* **23**, 711–717 (2016).
- [97] Masatoshi Sato and Yoichi Ando, “Topological superconductors: a review,” *Reports on Progress in Physics* **80**, 076501 (2017).
- [98] Shinsei Ryu and Yasuhiro Hatsugai, “Topological origin of zero-energy edge states in particle-hole symmetric systems,” *Phys. Rev. Lett.* **89**, 077002 (2002).
- [99] Ryosuke Iritani Adachi, Kyosuke and Ryusuke Hamazaki, “Universal constraint on nonlinear population dynamics,” *Communications Physics* **5**, 129 (2022).
- [100] Dixon D. Jones Ludwig, Donald and Crawford S. Holling, “Qualitative analysis of insect outbreak systems: the spruce budworm and forest,” *Journal of animal ecology* **47**, 315–332 (1978).
- [101] A Tsapalis, EP Politis, XN Maintyas, and FK Diakonou, “Gauss’ law and nonlinear plane waves for yang-mills theory,” *Physical Review D* **93**, 085003 (2016).
- [102] Christopher Chong, Yifan Wang, Donovan Maréchal, Efstathios G Charalampidis, Miguel Molerón, Alejandro J Martínez, Mason A Porter, Panayotis Kevrekidis, and Chiara Daraio, “Nonlinear localized modes in two-dimensional hexagonally-packed magnetic lattices,” *New Journal of Physics* **23**, 043008 (2021).
- [103] Yu-Ao Chen, Sebastian D Huber, Stefan Trotzky, Immanuel Bloch, and Ehud Altman, “Many-body landau-zener dynamics in coupled one-dimensional bose liquids,” *Nature Physics* **7**, 61–67 (2011).
- [104] Thomas Tuloup, Raditya Weda Bomantara, Ching Hua Lee, and Jiangbin Gong, “Nonlinearity induced topological physics in momentum space and real space,” *Phys. Rev. B* **102**, 115411 (2020).
- [105] Raditya Weda Bomantara, Wenlei Zhao, Longwen Zhou, and Jiangbin Gong, “Nonlinear dirac cones,” *Phys. Rev. B* **96**, 121406 (2017).
- [106] Silvio B Koller, Elizabeth A Goldschmidt, Roger C Brown, Robert Wyllie, RM Wilson, and JV Porto, “Nonlinear looped band structure of bose-einstein condensates in an optical lattice,” *Physical Review A* **94**, 063634 (2016).
- [107] Di Xiao, Ming-Che Chang, and Qian Niu, “Berry phase effects on electronic properties,” *Rev. Mod. Phys.* **82**, 1959–2007 (2010).
- [108] Jie Liu, Biao Wu, and Qian Niu, “Nonlinear evolution of quantum states in the adiabatic regime,” *Phys. Rev. Lett.* **90**, 170404 (2003).
- [109] Han Pu, Peter Maenner, Weiping Zhang, and Hong Y. Ling, “Adiabatic condition for nonlinear systems,” *Phys. Rev. Lett.* **98**, 050406 (2007).
- [110] FN Litvinets, AV Shapovalov, and A Yu Trifonov, “Berry phases for the nonlocal gross-pitaevskii equation with a quadratic potential,” *Journal of Physics A: Mathematical and General* **39**, 1191 (2006).
- [111] Asela Abeya, Gino Biondini, and Mark A Hoefer, “Whitham modulation theory for the defocusing nonlinear schrödinger equation in two and three spatial dimensions,” *Journal of Physics A: Mathematical and Theoretical* **56**, 025701 (2023).
- [112] AA Minzoni and Noel F Smyth, “Modulation theory, dispersive shock waves and gerald beresford whitham,” *Physica D: Nonlinear Phenomena* **333**, 6–10 (2016).
- [113] Gino Biondini, Christopher Chong, and Panayotis Kevrekidis, “On the whitham modulation equations for the toda lattice and the quantitative characterization of its dispersive shocks,” *arXiv preprint arXiv:2312.10755* (2023).
- [114] Thomas Tuloup, Raditya Weda Bomantara, Ching Hua Lee, and Jiangbin Gong, “Nonlinearity induced topological physics in momentum space and real space,” *Physical Review B* **102**, 115411 (2020).
- [115] Lukas J Maczewsky, Matthias Heinrich, Mark Kremer, Sergey K Ivanov, Max Ehrhardt, Franklin Martinez, Yaroslav V Kartashov, Vladimir V Konotop, Lluís Torner, Dieter Bauer, *et al.*, “Nonlinearity-induced photonic topological insulator,” *Science* **370**, 701–704 (2020).
- [116] Yaakov Lumer, Yonatan Plotnik, Mikael C Rechtsman, and Mordechai Segev, “Self-localized states in photonic topological insulators,” *Physical review letters* **111**, 243905 (2013).
- [117] Tomoki Ozawa, Hannah M. Price, Alberto Amo, Nathan Goldman, Mohammad Hafezi, Ling Lu, Mikael C. Rechtsman, David Schuster, Jonathan Simon, Oded Zilberberg, and Iacopo Carusotto, “Topological photonics,” *Rev. Mod. Phys.* **91**, 015006 (2019).

- [118] Xin Zhou, You Wang, Daniel Leykam, and Yi Dong Chong, “Optical isolation with nonlinear topological photonics,” *New Journal of Physics* **19**, 095002 (2017).
- [119] Vladimir V. Konotop, Jianke Yang, and Dmitry A. Zezyulin, “Nonlinear waves in \mathcal{PT} -symmetric systems,” *Rev. Mod. Phys.* **88**, 035002 (2016).
- [120] Demetrios Christodoulides, Jianke Yang, *et al.*, *Parity-time symmetry and its applications*, Vol. 280 (Springer, 2018).
- [121] Arbel Haim and Yuval Oreg, “Time-reversal-invariant topological superconductivity in one and two dimensions,” *Physics Reports* **825**, 48 (2019).
- [122] Shinsei Ryu, Andreas P Schnyder, Akira Furusaki, and Andreas WW Ludwig, “Topological insulators and superconductors: tenfold way and dimensional hierarchy,” *New Journal of Physics* **12**, 065010 (2010).
- [123] Eugene Dumitrescu, Jay D. Sau, and Sumanta Tewari, “Magnetic field response and chiral symmetry of time-reversal-invariant topological superconductors,” *Phys. Rev. B* **90**, 245438 (2014).
- [124] Di Zhou and Junyi Zhang, “Non-hermitian topological metamaterials with odd elasticity,” *Phys. Rev. Research* **2**, 023173 (2020).
- [125] Doru Sticlet, “Edge states in chern insulators and majorana fermions in topological superconductors,” *Diss. Université Paris Sud-Paris XI* (2012).
- [126] Chunyan Li and Yaroslav V. Kartashov, “Topological gap solitons in rabi su-schrieffer-heeger lattices,” *Phys. Rev. B* **108**, 184301 (2023).
- [127] Andrew Waldron, “A wick rotation for spinor fields: the canonical approach,” *Physics Letters B* **433** (1998).
- [128] Steven H Strogatz, *Nonlinear dynamics and chaos with student solutions manual: With applications to physics, biology, chemistry, and engineering* (CRC press, 2018).
- [129] Xinxin Guo, Lucien Jezequel, Mathieu Padlewski, Hervé Lissek, Pierre Delplace, and Romain Fleury, “Practical realization of chiral nonlinearity for strong topological protection,” *arXiv preprint arXiv:2403.10590* (2024).