A consistent test of spherical symmetry for multivariate and high-dimensional data via data augmentation

Bilol Banerjee and Anil K. Ghosh

Theoretical Statistics and Mathematics Unit, Indian Statistical Institute, Kolkata

Abstract

We develop a test for spherical symmetry of a multivariate distribution P that works even when the dimension of the data d is larger than the sample size n. We propose a non-negative measure $\zeta(P)$ such that $\zeta(P) = 0$ if and only if P is spherically symmetric. We construct a consistent estimator of $\zeta(P)$ using the data augmentation method and investigate its large sample properties. The proposed test based on this estimator is calibrated using a novel resampling algorithm. Our test controls the Type-I error, and it is consistent against general alternatives. We also study its behaviour for a sequence of alternatives $(1 - \delta_n)F + \delta_n G$, where $\zeta(G) = 0$ but $\zeta(F) > 0$, and $\delta_n \in [0, 1]$. When $\limsup \delta_n < 1$, for any G, the power of our test converges to unity as n increases. However, if $\limsup \delta_n = 1$, the asymptotic power of our test depends on $\lim n(1 - \delta_n)^2$. We establish this by proving the minimax rate optimality of our test over a suitable class of alternatives and showing that it is Pitman efficient when $\dim n(1 - \delta_n)^2 > 0$. Moreover, our test is provably consistent for high-dimensional data even when d is larger than n. Our numerical results amply demonstrate the superiority of the proposed test over some state-of-the-art methods. **Keywords:** Consistency; Contiguous alternatives; Data augmentation;

Minimax rate optimality; Pitman efficiency; Spherical symmetry.

1 Introduction

An inherent property of nature is that it tends to exhibit some form of symmetry within itself. In the nineteenth century, such symmetric patterns were approximated by the normal distribution (see, e.g., Lehmann, 2012, for a history of statistical methods). But with time, more general notions of symmetry were introduced. One of the most popular notions is spherical symmetry or elliptic symmetry (i.e., spherical symmetry after standardization) (see, e.g. Chmielewski, 1981; Fang et al., 1990; Fourdrinier et al., 2018). A *d*-dimensional (d > 1) random vector **X** is said to follow a spherically symmetric distribution if **X** has the same distribution as **HX** (i.e., $\mathbf{X} \stackrel{D}{=} \mathbf{HX}$) for any $d \times d$ orthogonal matrix **H**. This is an important class of distributions, and several statistical methods have been developed motivated by the sphericity or the ellipticity of the underlying distribution (see, e.g., Randles, 1989; Chaudhuri and Sengupta, 1993; Jörnsten, 2004; Ghosh and Chaudhuri, 2005). Therefore, testing the sphericity of a distribution is an important statistical problem, and we investigate it here based on a sample $\mathcal{D} = {\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_n}$ of n independent realizations of the random vector **X**.

Fang et al. (1993) considered this problem and proposed an asymptotically distribution-free test for spherical symmetry using the projection pursuit technique. They implemented the Wilcoxon-Mann-Whitney test for several pairs of projection directions and used the minimum over all such projection-pairs as the test statistic. However, this is only a necessary test for sphericity and does not have consistency under general alternatives. Koltchinskii and Li (1998) proposed a test based on the difference between the empirical spatial rank function and the theoretical spatial rank function under spherical symmetry, where the unknown components of these theoretical ranks were estimated from the data. The authors studied the large sample

behaviour of the test statistic and proposed a bootstrap method for calibration. Smith (1977) proposed a test statistic for bivariate data that uses the fact that if X is spherically symmetric then $\|X\|$ and $X/\|X\|$ are independent and $\mathbf{X}/\|\mathbf{X}\|$ follows a uniform distribution over the perimeter of the unit circle in \mathbb{R}^2 . Later, Baringhaus (1991) modified the test statistic and generalized the test to any arbitrary dimension. However, this test involves a complex function of dimension d, which makes it difficult to study its high dimensional behaviour. Diks and Tong (1999) proposed a Monte Carlo test for multivariate spherical symmetry conditionally on minimal sufficient statistics, but the consistency of this test against general alternatives is missing from the literature. Liang et al. (2008) proposed some necessary tests for spherical symmetry by using the fact that under H_0 , the null hypothesis of spherical symmetry, $\mathbf{X}/\|\mathbf{X}\|$ is uniformly distributed on the surface of the unit sphere in \mathbb{R}^d , but they did not consider the independence between and $\|\mathbf{X}\|$ and $\mathbf{X}/\|\mathbf{X}\|$. Hence et al. (2014) proposed a test based on characteristic functions and calibrated the test using a bootstrap algorithm. However, their test requires generation of uniform grid over the unit sphere, which becomes computationally prohibitive even in moderately large dimensions. Albisetti et al. (2020) proposed another test utilizing the fact that X is spherically symmetric if and only if $\mathbb{E}\{\mathbf{u}^{\top} \mathbf{X} \mid \mathbf{v}^{\top} \mathbf{X}\} = 0$ for all **u** and **v** with $\mathbf{u}^{\top}\mathbf{v} = 0$. They constructed a Kolmogorov-Smirnov-type test statistic over suitable choices of test functions and studied its theoretical properties. Recently, Huang and Sen (2023) proposed tests for different notions of symmetry using optimal transport for multivariate data. They showed that the Pittman efficiency of their test is never worse than the efficiency of Hotelling's T^2 test for Gaussian alternatives if the reference distribution in the optimal transport step is chosen to be Gaussian.

Most of these above tests are large-sample consistent for any fixed dimension d and applicable when the dimension of the data d is small compared to the sample size n (i.e., d << n). However, the applicability of these tests for high-dimensional data (i.e., when d is comparable to or larger than n) is not clear. Zou et al. (2014) and Feng and Liu (2017) proposed tests of sphericity for high-dimensional data where the test statistics were constructed using the multivariate sign function assuming ellipticity of the underlying distribution. Ding (2020) proposed a test based on the ratio of traces of different powers of the sample variance-covariance matrix and established its high dimensional consistency. However, these tests may fail when the underlying distribution is not spherically symmetric but $\mathbf{X}/||\mathbf{X}||$ is uniformly distributed over the surface of the unit sphere in \mathbb{R}^d (e.g. angular symmetric) or the variance-covariance matrix is a constant multiple of the identity matrix. This motivates us to construct a test of spherical symmetry which does not have such limitations. The test we propose in this article, is applicable to high-dimensional data even when d is significantly larger than n, and it has consistency for a certain class of nonparametric alternatives including some shrinking alternatives.

In Section 2, we propose a new measure of spherical asymmetry $\zeta(P)$ by using the fact that $\mathbf{X} \sim P$ is spherically symmetric if and only if **X** and $\|\mathbf{X}\|\mathbf{U}$ are identically distributed, where $\|\mathbf{X}\|$ and **U** are independent and $\mathbf{U} \sim \text{Unif}(\mathcal{S}^{d-1})$, the uniform distribution on the surface of the *d*-dimensional unit sphere. The measure $\zeta(P)$ is non-negative, and it takes the value zero if and only if P is spherically symmetric. However, the measure involves some terms that are not estimable from the observed data \mathcal{D} only. To overcome this limitation, we propose a data augmentation approach to estimate $\zeta(P)$. We study the large sample properties of this estimator and build a test based on it. In Section 3, we study the asymptotic behaviour of the resulting test under mixture alternatives of the form $(1 - \delta_n)F + \delta_n G$, where G is spherically symmetric, but F is not spherical. For any fixed d, F and a sequence $\{\delta_n\}$ that remains bounded away from 1, we prove that our test is consistent in the sense of the minimum power over the class of spherical distributions $\mathcal{G} = \{G : \zeta(G) = 0\}$, i.e., for any $G \in \mathcal{G}$, the power of our test converges to 1 as n increases. We also establish that our test is minimax rate optimal over a certain class of alternatives and Pitman efficient against contamination alternatives for which $n(1-\delta_n)^2$ converges to a positive number. This in turn proves the consistency of the test over a fairly general class of shrinking alternatives when d and n both diverge to infinity. Extensive simulation studies are carried out in Section 4 to compare the performance of our test with some state-of-the-art methods. All proofs and some auxiliary theoretical results are deferred to the Appendix.

2 The proposed methodology

We know that \mathbf{X} is spherically symmetric if and only if \mathbf{X} and $\mathbf{H}\mathbf{X}$ have the same distribution for any orthogonal matrix \mathbf{H} . However, we can also characterize spherical symmetry using the following lemma.

Lemma 2.1. A d-dimensional random vector \mathbf{X} is spherically symmetric if and only if $\mathbf{X} \stackrel{D}{=} \|\mathbf{X}\| \mathbf{U}$ where \mathbf{U} is uniformly distributed over the surface of the unit sphere \mathcal{S}^{d-1} and is independent of $\|\mathbf{X}\|$.

For proof of the above lemma, see p.31 in Fang et al. (1990). Using this characterization, we propose a new measure of spherical asymmetry, which is applicable to general nonparametric distributions. Let φ_1 be the characteristic function of $\mathbf{X} \sim \mathbf{P}$ and φ_2 be that of its sprherically symmetric variant $\|\mathbf{X}\| \mathbf{U}$, where $\mathbf{U} \sim \text{Unif}(\mathcal{S}^{d-1})$ and $\|\mathbf{X}\|$ are independent. We define

$$\zeta(\mathbf{P}) = \int \left|\varphi_1(\mathbf{t}) - \varphi_2(\mathbf{t})\right|^2 \frac{d^{d/2}}{(2\pi)^{d/2}} e^{-d\|\mathbf{t}\|^2/2} \mathrm{d}\mathbf{t},\tag{1}$$

where the integral is taken in the principal value sense. The following proposition proves that $\zeta(P)$ characterizes spherical symmetry by showing $\zeta(P) \ge 0$, where the equality holds if and only if P is spherically symmetric.

Proposition 2.1. $\zeta(P)$ is non-negative and $\zeta(P) = 0$ if and only if P is a spherically symmetric distribution.

Since, $\zeta(\mathbf{P})$ is the integration of $|\varphi_1(\mathbf{t}) - \varphi_2(\mathbf{t})|^2$ with respect to a Gaussian distribution in \mathbb{R}^d , one can also derive an alternative form for $\zeta(\mathbf{P})$ using the following lemma.

Lemma 2.2. For any two non-zero vectors $\mathbf{X}_1, \mathbf{X}_2 \in \mathbb{R}^d$,

$$\int \exp\left\{i\langle \mathbf{t}, \mathbf{X}_1 - \mathbf{X}_2\rangle\right\} \frac{d^{d/2}}{(2\pi)^{d/2}} e^{-d\|\mathbf{t}\|^2/2} \mathrm{d}\mathbf{t} = \exp\left\{-\frac{1}{2d}\|\mathbf{X}_1 - \mathbf{X}_2\|^2\right\}$$

Using Lemma 2.2 we get the following alternative representation of $\zeta(P)$.

Theorem 2.1. If $\mathbf{X}_1, \mathbf{X}_2$ are independent copies of $\mathbf{X} \sim \mathbf{P}$, then $\zeta(\mathbf{P})$ can be expressed as

$$\zeta(\mathbf{P}) = \mathbb{E}\left\{\exp\{-\frac{1}{2d}\|\mathbf{X}_1 - \mathbf{X}_2\|^2\}\right\} + \mathbb{E}\left\{\exp\{-\frac{1}{2d}\|\mathbf{X}_1' - \mathbf{X}_2'\|^2\}\right\} - 2\mathbb{E}\left\{\exp\{-\frac{1}{2d}\|\mathbf{X}_1 - \mathbf{X}_2'\|^2\}\right\},$$

where $\mathbf{X}'_i = \|\mathbf{X}_i\|\mathbf{U}_i$ for $i = 1, 2, \mathbf{U}_1, \mathbf{U}_2$ independent and identically distributed (i.i.d.) as Unif (\mathcal{S}^{d-1}) , which are independent of \mathbf{X}_1 and \mathbf{X}_2 .

Remark 1. Theorem 2.1 shows that $\zeta(\mathbf{P})$ can be expressed as a function of the pairwise distance between $\mathbf{X}_1, \mathbf{X}_2$ and their symmetric variants $\mathbf{X}'_1, \mathbf{X}'_2$. Since, the pairwise distances are invariant under orthogonal transformation, $\zeta(\mathbf{P})$ is also invariant under rotation. In particular, we have $\zeta(N(\mathbf{0}, \boldsymbol{\Sigma})) = \zeta(N(\mathbf{0}, \mathbf{H}\boldsymbol{\Sigma}\mathbf{H}^T))$ for any orthogonal matrix \mathbf{H} . The exact expression of $\zeta(N(\mathbf{0}, \boldsymbol{\Sigma}))$ for any positive-semidefinite variance-covariance matrix $\boldsymbol{\Sigma}$ is interesting. Interested readers are referred to Section A.1 in the Appendix for a detailed derivation.

Remark 2. $\zeta(\mathbf{P})$ can also be viewed as the energy distance (see,e.g. Székely and Rizzo, 2004) or the maximum mean discrepancy (MMD) (see, e.g., Gretton et al., 2007) between the distributions of \mathbf{X} and $\|\mathbf{X}\|\mathbf{U}$. MMD is used to quantify the distributional difference using embedding into a Reproducing Kernel Hilbert Space (RKHS). Here $K(\mathbf{x}, \mathbf{y}) = \exp\{-\|\mathbf{x} - \mathbf{y}\|^2/(2d)\}$ is the kernel associated with that RKHS.

2.1 Estimation of $\zeta(P)$

Let $\mathcal{D} = \{\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_n\}$ be a random sample of size *n* from a *d*-dimensional distribution P. Note that the term $\mathbb{E}\{\exp\{-\frac{1}{2d}\|\mathbf{X}_1 - \mathbf{X}_2\|^2\}\}$ in $\zeta(\mathbf{P})$ can be easily estimated by its empirical analog, but $\zeta(\mathbf{P})$ involves two other terms $\mathbb{E}\{\exp\{-\frac{1}{2d}\|\mathbf{X}_1' - \mathbf{X}_2'\|^2\}\}$ and $\mathbb{E}\{\exp\{-\frac{1}{2d}\|\mathbf{X}_1 - \mathbf{X}_2'\|^2\}\}$, which are not estimable from \mathcal{D} only. Therefore, we adopt a data augmentation approach. We generate $\mathbf{U}_1, \mathbf{U}_2, \dots, \mathbf{U}_n$, a random

sample of size *n* from Unif(\mathcal{S}^{d-1}) and define $\mathbf{X}'_i = \|\mathbf{X}_i\| \mathbf{U}_i$ for i = 1, 2, ..., n. Using the augmented data $\mathcal{D}' = \{(\mathbf{X}_1, \mathbf{X}'_1), (\mathbf{X}_2, \mathbf{X}'_2), ..., (\mathbf{X}_n, \mathbf{X}'_n)\}$ we construct an unbiased estimator of $\zeta(\mathbf{P})$ given by

$$\hat{\zeta}_n = \binom{n}{2}^{-1} \sum_{i=1}^n \sum_{j=i+1}^n \Big\{ \exp\{-\frac{1}{2d} \|\mathbf{X}_i - \mathbf{X}_j\|^2\} + \exp\{-\frac{1}{2d} \|\mathbf{X}_i' - \mathbf{X}_j'\|^2\} - 2\exp\{-\frac{1}{2d} \|\mathbf{X}_i - \mathbf{X}_j'\|^2\} \Big\}.$$

Clearly $\hat{\zeta}_n$ can be viewed as a U-statistic with the symmetrized kernel function

$$g((\mathbf{x}_1, \mathbf{x}_1'), (\mathbf{x}_2, \mathbf{x}_2')) = K(\mathbf{x}_1, \mathbf{x}_2) + K(\mathbf{x}_1', \mathbf{x}_2') - K(\mathbf{x}_1, \mathbf{x}_2') - K(\mathbf{x}_2, \mathbf{x}_1'),$$
(2)

where $K(\mathbf{x}, \mathbf{y}) = \exp\{-\|\mathbf{x} - \mathbf{y}\|^2/(2d)\}$. Here g is a bounded kernel, and using the bounded difference inequality, we can establish the following bound for the deviation of $\hat{\zeta}_n$ from its population analog $\zeta(\mathbf{P})$.

Theorem 2.2. If $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_n$ are independent copies of a d-dimensional random vector $\mathbf{X} \sim \mathbf{P}$, then

$$\mathbb{P}\left\{\left|\hat{\zeta}_n - \zeta(\mathbf{P})\right| > \epsilon\right\} \le 2\exp\{-\frac{n\epsilon^2}{32}\},\$$

and this inequality holds irrespective of the dimension d.

Theorem 2.2 shows that $\hat{\zeta}_n$ is a strongly consistent estimator of $\zeta(\mathbf{P})$ and the exponential bound is free from d. The large sample distribution of $\hat{\zeta}_n$ can be derived using the theory of U-statistics (see Lee, 1990). The asymptotic null distribution of $\hat{\zeta}_n$ is given by the following theorem.

Theorem 2.3. (Asymptotic null distribution) Let $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_n$ be independent copies of the random vector \mathbf{X} , which follows a spherically symmetric distribution \mathbf{P} . Define $\mathbf{X}'_i = \|\mathbf{X}_i\|\mathbf{U}_i$ for $i = 1, 2, \ldots, n$, where $\mathbf{U}_1, \mathbf{U}_2, \ldots, \mathbf{U}_n$ are i.i.d. Unif (S^{d-1}) . Let λ_k $(k = 1, 2, \ldots)$ be the eigenvalue corresponding to the eigenfunction ψ_k of the integral equation

$$\mathbb{E}\{g((\mathbf{x}, \mathbf{x}'), (\mathbf{X}, \mathbf{X}'))\psi_k(\mathbf{X}, \mathbf{X}')\} = \lambda_k \psi_k(\mathbf{x}, \mathbf{x}'),\$$

where g is as in equation (2). Then as n goes to infinity, $n\hat{\zeta}_n \xrightarrow{D} \sum_{k=1}^{\infty} \lambda_k(Z_i^2 - 1)$, where $\{Z_k : k \ge 1\}$ is a sequence of *i.i.d.* standard normal variables.

In Theorem 2.3, the limiting null distribution of $\hat{\zeta}_n$ (after appropriate adjustment for location and scale) turns out to be a weighted sum of independent chi-squares due to the first order degeneracy of g under H_0 . However, under H_1 , g is a non-degenerate kernel. Therefore, the limiting distribution of $\hat{\zeta}_n$ (after appropriate adjustment for location and scale) turns out to be Gaussian, which is asserted by the following theorem.

Theorem 2.4. (Asymptotic alternative distribution) Let $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_n$ be independent copies of a random vector \mathbf{X} , which follows a distribution P that is not spherically symmetric. Define $\mathbf{X}'_i = \|\mathbf{X}_i\|\mathbf{U}_i$ where $\mathbf{U}_1, \mathbf{U}_2, \ldots, \mathbf{U}_n$ is an i.i.d. $Unif(S^{d-1})$. Then as n goes to infinity, $\sqrt{n}(\hat{\zeta}_n - \zeta(P)) \xrightarrow{D} N(0, 4\sigma^2)$, where $\sigma^2 = Var \Big\{ \mathbb{E} \big\{ g\big((\mathbf{X}_1, \mathbf{X}'_1), (\mathbf{X}_2, \mathbf{X}'_2)\big) \mid (\mathbf{X}_1, \mathbf{X}'_1) \big\} \Big\}$, and g is as defined in equation (2).

2.2 Test of Spherical Symmetry

Theorem 2.2 shows that $\hat{\zeta}_n$ is a strongly consistent estimator of $\zeta(\mathbf{P})$. Now, from Proposition 2.1, it is clear that under $H_0 \hat{\zeta}_n$ converges almost surely to zero, but under H_1 , it converges to a positive constant. Therefore, the power of a test that rejects H_0 for higher values of $\hat{\zeta}_n$ converges to unity as the sample size increases. However, it is difficult to find the critical value based on the asymptotic null distribution of $n\hat{\zeta}_n$ (see Theorem 2.3) since it involves an ℓ_2 sequence $\{\lambda_k\}$, which depends on the underlying distribution P. So, we propose a novel resampling algorithm to compute the cut-off. The algorithm is given below. Resampling algorithm

- (I) Given the augmented data \mathcal{D}' compute the test statistic $\hat{\zeta}_n$.
- (II) Let $\pi = (\pi(1), \pi(2), \dots, \pi(n))$ be an element in $\{0, 1\}^n$. Define $\mathbf{Y}_i = \pi(i)\mathbf{X}_i + (1 \pi(i))\mathbf{X}'_i$ and $\mathbf{Y}'_i = (1 \pi(i))\mathbf{X}_i + \pi(i)\mathbf{X}'_i$ for $i = 1, 2, \dots, n$. Use $(\mathbf{Y}_1, \mathbf{Y}'_1), \dots, (\mathbf{Y}_n, \mathbf{Y}'_n)$ to compute $\hat{\zeta}_n(\pi)$, the resampling analogue of $\hat{\zeta}_n$.

(III) Repeat step (II) for all possible π to get the critical value for a level α ($0 < \alpha < 1$) test as

$$c_{1-\alpha} = \inf\{t \in \mathbb{R} : \frac{1}{2^n} \sum_{\pi \in \{0, 1\}^n} \mathbf{I}[\hat{\zeta}_n(\pi) \le t] \ge 1-\alpha\}.$$

This resampling algorithm is motivated by the fact that under H_0 , the \mathbf{X}_i and its spherically symmetric variant \mathbf{X}'_i (i = 1, 2, ..., n) are exchangeable. Therefore, under H_0 , the random swap between \mathbf{X}_i and \mathbf{X}'_i does not change the distribution of $\hat{\zeta}_n$. We can reject H_0 if $\hat{\zeta}_n$ is larger than $c_{1-\alpha}$. Note that the p-value of this conditional test is given by

$$p_n = \frac{1}{2^n} \sum_{\pi \in \{0, 1\}^n} \mathbf{I}[\hat{\zeta}_n(\pi) \ge \hat{\zeta}_n].$$
(3)

So, alternatively, we can reject H_0 if $p_n < \alpha$. The following lemma shows that this resampling algorithm gives a valid level α test.

Lemma 2.3. Let $\hat{\zeta}_n$ be the estimator of $\zeta(\mathbf{P})$ as defined in Section 2.2. If p_n denotes the conditional p-value as defined in (3), then under $H_0: \zeta(\mathbf{P}) = 0$ we have $\mathbb{P}\{p_n < \alpha\} \leq \alpha$ irrespective of n and d.

Interestingly, we can also control the threshold $c_{1-\alpha}$ by a deterministic sequence that does not depend on d and converges to zero with increasing sample size. This is asserted by the following theorem.

Lemma 2.4. If $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_n$ are independent copies of a d-dimensional random variable $\mathbf{X} \sim \mathbf{P}$, then for any α (0 < α < 1), the inequality $c_{1-\alpha} \leq 2(\alpha(n-1))^{-1}$ holds with probability one.

So, irrespective of the value of d, $c_{1-\alpha}$ is of order $O_P(n^{-1})$ and it converges to zero almost surely as n diverges to infinity. For any fixed d, we can also show that for any distribution P, $n\hat{\zeta}_n(\pi)$ converges in distribution to a weighted sum of independent chi-squares as n diverges to infinity. Since, under H_1 , $\hat{\zeta}_n$ converges to a positive number, the conditional p-value p_n converges to zero as n increases. Hence, the power of the resulting conditional test converges to one. This is formally stated in the following theorem.

Theorem 2.5. The power of the conditional test based on p_n converges to one as n diverges to infinity.

Though the above resampling algorithm leads to a consistent level α test, it has a computational complexity of the order $O(n^2 2^n)$. So, it is not computationally feasible to implement even if the sample size is moderately large. Therefore, in practice, we propose to generate $\pi_1, \pi_2, \ldots, \pi_B$ uniformly from $\{0, 1\}^n$ and compute the randomized p-value

$$p_{n,B} = \frac{1}{(B+1)} \Big(\sum_{i=1}^{B} I\{\hat{\zeta}_n(\pi_i) \ge \hat{\zeta}_n\} + 1 \Big).$$

We reject H_0 if $p_{n,B} < \alpha$. The following theorem shows that $p_{n,B}$ closely approximates p_n for large B and thereby justifies the use $p_{n,B}$ for the practical implementation of the test.

Theorem 2.6. Given the augmented data \mathcal{D}' , $p_{n,B}$ converges to p_n almost surely as B diverges to infinity.

3 Asymptotic properties of the test

In this section, we study some large sample properties of our test. First, we investigate the robustness of our test against contamination alternatives. Next, we establish its minimax rate optimality against a suitable class of nonparametric alternatives and prove its consistency even when the dimension of data increases with the sample size. Finally, we show that our test is efficient in the Pittman sense under contiguous contamination alternatives.

3.1 Robustness

Consider a distribution F, which is not spherically symmetric. Let $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_n$ be independent and identically distributed random vectors from a contaminated distribution $F_{\delta} = (1 - \delta)F + \delta G$ for some $0 < \delta < 1$. The following lemma shows the effect of this contamination on $\zeta(\cdot)$ by providing a relation among $\zeta(F), \zeta(G)$ and $\zeta(F_{\delta})$.

Lemma 3.1. For any δ ($0 < \delta < 1$), we have $\zeta(F_{\delta}) = (1 - \delta)^2 \zeta(F) + \delta^2 \zeta(G) + 2\delta(1 - \delta)\zeta'(G, F)$, where

$$\zeta'(G,F) = \mathbb{E}\{K(\mathbf{X}_1,\mathbf{Y}_1)\} + \mathbb{E}\{K(\mathbf{X}_1',\mathbf{Y}_1')\} - \mathbb{E}\{K(\mathbf{X}_1,\mathbf{Y}_1')\} - \mathbb{E}\{K(\mathbf{X}_1',\mathbf{Y}_1)\}.$$

Here $\mathbf{X}_1 \sim G$ and $\mathbf{Y}_1 \sim F$ are independent, $\mathbf{X}'_1 = \|\mathbf{X}_1\|\mathbf{U}_1, \mathbf{Y}'_1 = \|\mathbf{Y}_1\|\mathbf{U}_2$ for $\mathbf{U}_1, \mathbf{U}_2$ are *i.i.d.* Unif (\mathcal{S}^{d-1}) and $K(\mathbf{x}, \mathbf{y}) = \exp\{-\frac{1}{2d}\|\mathbf{x} - \mathbf{y}\|^2\}$.

Note that if G is spherically symmetric, $\zeta(G) = 0$ and $\zeta'(G, F) = 0$. Therefore, for any fixed δ , we have $\zeta(F_{\delta}) = (1 - \delta)^2 \zeta(F)$. This shows that $\zeta(.)$ has bounded Gateaux derivative. The following theorem also shows that for any $\delta \in (0, 1)$ and G spherically symmetric, the power of our test for the contaminated alternative F_{δ} converges to one as the sample size increases.

Theorem 3.1. Let $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_n$ be independent copies of $\mathbf{X} \sim F_{\delta} = (1 - \delta)F + \delta G$, where $0 < \delta < 1$, and $\zeta(F) > 0$. Then, the minimum power of the proposed test over the class of the spherical distributions G, i.e., $\inf_{G:\zeta(G)=0} \mathbb{P}\{\hat{\zeta}_n > c_{1-\alpha}\}$, converges to one as the sample size diverges to infinity.

This theorem shows that our test is asymptotically robust against outliers for any fixed proportion of contamination. Since $\zeta(F_{1-\delta}) = \delta^2 \zeta(F)$, it shows the convergence of the power of our test for $F_{1-\delta}$ as well. So, if a sample from a spherically symmetric distribution has a small proportion of contamination by observations from a non-spherical distribution, our test can successfully detect the presence of those contaminations when the sample size is large. The result in Theorem 3.1 holds even for a sequence $\{\delta_n\}$ that remains bounded away from one. However, if that is not the case, the asymptotic power of the test will depend on the rate of convergence of $1 - \delta_n$ and may yield non-trivial limits for certain choices of $\{\delta_n\}$. This is explored in the following subsection.

3.2 Minimax rate optimality and high-dimensional behaviour

Let us consider a testing problem involving a pair of hypotheses $H_0 : \zeta(\mathbf{P}) = 0$ and $H'_1 : \zeta(\mathbf{P}) > \epsilon(n)$ where $\epsilon(n)$ is a positive number that depends on the sample size n. Let $\mathcal{F}(\epsilon(n)) := \{\mathbf{P} \mid \zeta(\mathbf{P}) > \epsilon(n)\}$ be the class of alternatives under H'_1 and $\mathbb{T}_n(\alpha)$ be the class of all level α test. The minimax type II error rate for this problem is defined as

$$R_n(\epsilon(n)) = \inf_{\phi \in \mathbb{T}_n(\alpha)} \sup_{F \in \mathcal{F}(\epsilon(n))} \mathbb{P}_{F^n} \{\phi = 0\},$$

where \mathbb{P}_{F^n} denotes the probability corresponding to the joint distribution of $(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_n)$, where the \mathbf{X}_i s are independent copies of $\mathbf{X} \sim F$. Here we want to find an optimum choice of $\epsilon(n)$ (call it $\epsilon_0(n)$) such that the following two conditions hold.

(a) For any $0 < \beta < 1 - \alpha$, there exists a constant $c(\alpha, \beta) > 0$ such that for all $0 < c < c(\alpha, \beta)$, we have $\liminf_{n \in \mathcal{N}} R_n(c \ \epsilon_0(n)) \ge \beta$.

(b) There exists a level α test ϕ_0 such that for any $0 < \beta < 1 - \alpha$, there exists $C(\alpha, \beta) > 0$ for which $\lim_{n \to \infty} \sup_{F \in \mathcal{F}(c \ \epsilon_0(n))} \mathbb{P}_{F^n} \{ \phi_0 = 0 \} \leq \beta \text{ for all } c > C(\alpha, \beta), \text{ or in other words, } \limsup_{n \to \infty} R_n(c \ \epsilon_0(n)) \leq \beta \text{ for all } c > C(\alpha, \beta).$

This optimal rate $\epsilon_0(n)$ is called the minimax rate of separation for the above problem, and the test ϕ_0 is called the minimax rate optimal test. Theorem 3.2 below shows that $\epsilon_0(n)$ cannot be of smaller order than $O(n^{-1})$. So, for any $0 < \beta < 1 - \alpha$ and any $\phi \in \mathbb{T}_n(\alpha)$, we can always find a distribution F with $\zeta(F)$ of the order $O(n^{-1})$ or smaller such that the type II error rate of the test ϕ , i.e., $\mathbb{P}_{F^n}\{\phi=0\}$ is more than β .

Theorem 3.2. For $0 < \beta < 1 - \alpha$, there exists a constant $c_0(\alpha, \beta)$ such that the minimax type II error rate $R_n(c/n)$ is lower bounded by β for all n and all $0 < c < c_0(\alpha, \beta)$.

Remark 3. In particular, consider the distribution $F_{\delta_n} = (1 - \delta_n)F + \delta_n G$ where $\zeta(G) = 0$ and $\zeta(F) > 0$ (note that $\zeta(F_{\delta_n}) = (1 - \delta_n)^2 \zeta(F)$). If δ_n is such that $n(1 - \delta_n)^2 \to 0$ as $n \to \infty$ (i.e. $\zeta(F_{\delta_n})$ is of smaller asymptotic order than $O(n^{-1})$), then the power of any level α test will fall below the nominal level α .

In the next theorem, we establish that in the case of $\epsilon_0(n) = 1/n$, our test based on $\hat{\zeta}_n$ satisfies the condition (b) stated above. Therefore, these two theorems (Theorem 3.2 and 3.3) together show that the minimax rate of separation is $\epsilon_0(n) = n^{-1}$, and our proposed test has the minimax rate optimality for the class of alternatives $\mathcal{F}(\epsilon(n))$.

Theorem 3.3. For $0 < \beta < 1 - \alpha$, there exists a constant $C_0(\alpha, \beta)$ (that does not depend on d) such that asymptotically the maximum type II error of the test based on $\hat{\zeta}_n$ over $\mathcal{F}(c/n)$ is uniformly bounded above by β for all $c > C_0(\alpha, \beta)$, i.e., $\limsup_{n \to \infty} \sup_{F \in \mathcal{F}(c\lambda(n))} \mathbb{P}_{F^n}(\hat{\zeta}_n \leq c_{1-\alpha}) \leq \beta$ for all $c > C_0(\alpha, \beta)$.

Remark 4. Consider the same example as in Remark 3. Since $\zeta(F_{\delta_n}) = (1 - \delta_n)^2 \zeta(F)$, from Theorem 3.3 it is clear that if $n(1 - \delta_n)^2 \to \infty$ as $n \to \infty$ (i.e. $\zeta(F_{\delta_n})$ is of higher asymptotic order than $O(n^{-1})$), the power of our proposed test converges to one.

Note that the constant $C(\alpha, \beta)$ in Theorem 3.3 does not depend on the dimension d. However, $\zeta(F)$ may vary with the dimension. We know that under certain regularity conditions (see, e.g., Hall et al., 2005; Ahn et al., 2007; Jung and Marron, 2009), as the dimension increases, pairwise distances among the observations (after appropriate scaling) converge to a constant, and all observations tend to lie on the surface of a sphere of increasing radius. So, in such situations, $\zeta(F)$ converges to 0 as d increases. Therefore, one may be curious to know how this test will perform if the dimension and the sample size increase simultaneously. Theorem 3.3 answers this question. It shows that as long as $n\zeta(F)$ diverges to infinity, irrespective of whether d is fixed or it increases with the sample size, the power of our test converges to one. This high-dimensional consistency of our test is asserted by the following theorem.

Theorem 3.4. Suppose that $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_n$ are independent copies of $\mathbf{X} \sim \mathbf{P}^{(d)}$, a d-dimensional distribution. If d = d(n) grows with the sample size n in such a way that $n\zeta(\mathbf{P}^{(d)})$ diverges to infinity as n increases. Then, the power of our test converges to one as n and d both diverge to infinity.

So, even if $\zeta(\mathbf{P}^{(d)})$ converges to 0 as d increases with n, the power of our test converges to unity as long as $\zeta(\mathbf{P}^{(d)})$ converges at a slower rate than $O(n^{-1})$. However, if the distance convergence does not hold and we have $\liminf_{d\to\infty} \zeta(\mathbf{P}^{(d)}) > 0$, the power of our test converges to one even if the sample size increases at a very slow rate. An example of such a distribution is given in Section 4 (see Example 4(a)). For such examples, one can expect the test to have good performance even in the High Dimension Low Sample Size (HDLSS) setup, where n is fixed (but suitably large) and d diverges to infinity. However, in the case of distance concentration in the HDLSS set-up, where we have $\liminf_{d\to\infty} \zeta(\mathbf{P}^{(d)}) = 0$, we need to increase the sample size suitably to get good performance. This is further explored in our simulation studies.

3.3 Pitmann Efficiency

Now, consider the alternative $F_{1-\beta_n n^{-1/2}} = (1-\beta_n n^{-1/2})G + (\beta_n n^{-1/2})F$, but assume that β_n is a sequence of positive numbers converging to some $\beta \in (0, \infty)$. Let f and g be the densities corresponding to G and

F, respectively. To study the asymptotic behaviour of our test for such an alternative, we first study the asymptotic behaviour of $n\hat{\zeta}_n$ and its resample analog $n\hat{\zeta}_n(\pi)$. The following result shows that under suitable assumption on F and G, the sequence of alternative asymmetric distributions $F_{1-\beta_n n^{-1/2}}$ is contiguous and locally asymptotically normal.

Proposition 3.1. Let $\mathbf{X}_1, \ldots, \mathbf{X}_n$ be independent copies of $\mathbf{X} \sim G$. Then, under the assumption that $\int (f(\mathbf{u})/g(\mathbf{u}) - 1)^2 g(\mathbf{u}) d\mathbf{u} < \infty$ as n grows to infinity, we have

$$\left|\log\left\{\prod_{i=1}^{n}\left(1+\frac{\beta_{n}}{\sqrt{n}}\left\{\frac{f(\mathbf{X}_{i})}{g(\mathbf{X}_{i})}-1\right\}\right)\right\}-\frac{\beta_{n}}{\sqrt{n}}\sum_{i=1}^{n}\left\{\frac{f(\mathbf{X}_{i})}{g(\mathbf{X}_{i})}-1\right\}+\frac{\beta_{n}^{2}}{2}\mathbb{E}\left\{\frac{f(\mathbf{X}_{1})}{g(\mathbf{X}_{1})}-1\right\}^{2}\right|\xrightarrow{P}0.$$

Now using Proposition 3.1 and Le Cam's third lemma, we establish the local asymptotic behaviour of $n\hat{\zeta}_n$ in the following theorem.

Theorem 3.5. (Local limit distribution) Let $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_n$ be independent copies of $\mathbf{X} \sim F_{1-\beta_n n^{-1/2}}$ and $\mathbf{X}'_i = \|\mathbf{X}_i\|\mathbf{U}_i$ where $\mathbf{U}_1, \mathbf{U}_2, \ldots, \mathbf{U}_n$ are i.i.d. Unif (\mathcal{S}^{d-1}) . Also let λ_k be the eigenvalues with corresponding eigenfunction ψ_k ($k = 1, 2, \ldots$) of the integral equation $\mathbb{E}\{g((\mathbf{x}_1, \mathbf{x}'_1), (\mathbf{X}_1, \mathbf{X}'_1))\psi_k((\mathbf{X}_1, \mathbf{X}'_1))\} = \lambda_k \psi_k((\mathbf{x}_1, \mathbf{x}'_1))$, where g is as in equation (2). Then, as n tends to infinity,

$$n\hat{\zeta}_n \xrightarrow{D} \sum_{k=1}^{\infty} \lambda_k \left(\left(Z_k + \beta \mathbb{E}_F \{ \psi_i(\mathbf{X}_1, \mathbf{X}_1') \} \right)^2 - 1 \right),$$

where Z_i is a sequence of *i.i.d.* standard normal random variables.

Theorem 3.5 shows that for $\beta > 0$, the local limit distribution of $n\hat{\zeta}_n$ is stochastically larger than its limiting null distribution as in Theorem 2.3. The following theorem establishes that the local limiting distribution of the permuted statistic $n\hat{\zeta}_n(\pi)$ under the sequence of alternatives $F_{1-\beta_n n^{-1/2}}$ is identical to the asymptotic null distribution of $n\hat{\zeta}_n$.

Theorem 3.6. Let $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_n$ be independent copies of $\mathbf{X} \sim \mathbf{P}_n$ and $\hat{\zeta}_n(\pi)$ be the resampling analog of our test statistic obtained using Algorithm (I)-(III) in Section 2.2. Then, under any fixed alternative (i.e., $\mathbf{P}_n = F$ for some asymmetric distribution F) or a contiguous alternative (i.e., $\mathbf{P}_n = F_{1-\beta_n n^{-1/2}}$), as n grows to infinity, $n\hat{\zeta}_n(\pi) \xrightarrow{D} \sum_{k=1}^{\infty} \lambda_k(Z_k^2 - 1)$, where $\{Z_k\}$ is a sequence of i.i.d standard normal random variables and $\{\lambda_k\}$ is a square sumable sequence of real numbers.

Theorems 3.5 and 3.6 together show that under $F_{1-\beta_n n^{-1/2}}$, the power of our test converges to a nontrivial limit, and as β starts increasing from zero, the power of our test gradually increases from α to one. This establishes that our test is efficient in the Pitman sense. However, the exact expression of the limit is not analytically tractable.

We now present a small simulation study where we generate n observations from $F_{1-\beta_n n^{-1/2}}$ in \mathbb{R}^{10} where G is the standard normal distribution and F is a normal distribution with mean zero and variance-covariance matrix $\Sigma = (0.5) \mathbf{I} + 0.5 \mathbf{11}^{\top}$, where \mathbf{I} is the 10×10 identity matrix, $\mathbf{1}$ is the 10-dimensional vector with all elements equal to one. We consider three different sequences (a) $\beta_n = 5n^{-0.1}$, (b) $\beta_n = 5$ and (c) $\beta_n = 5n^{0.1}$ and evaluate the performance of our test against these alternatives. The p-value of the test is approximated using the randomized p-value with B = 500 and the power of the test is evaluated by the proportion of times

Sample Size	$\gamma = -0.1$	$\gamma = 0$	$\gamma = 0.1$
50	0.145	0.361	0.789
100	0.147	0.375	0.932
250	0.108	0.373	0.991
500	0.104	0.376	0.999

Table 1: Results against $F_{1-\beta_n n^{-1/2}} = (1-\beta_n n^{-1/2})N(0,I) + \beta_n n^{-1/2}N(0,0.5I+0.5J)$ when $\beta_n = 5n^{\gamma}$.

the test rejects H_0 in 1000 repetitions of each experiment. In Table 1 we see that for case (a), the power of our test shows a decreasing trend with increasing sample size. For case (b), the power exhibits convergence towards 0.37, which can be considered as the Pitman efficiency of our test when $\beta_n = 5$. For case (c), we see that the power of our test converges to one with increasing sample size. This behaviour of our test supports our theoretical findings in this section.

4 Numerical studies

In this section, we investigate the empirical performance of our test. First, we study its finite sample level properties and then compare its empirical power with the tests based on optimal transport (Huang and Sen, 2023), density functions (Diks and Tong, 1999) and projection pursuit technique (Fang et al., 1993). Henceforth, we refer to these tests as the OT test, the DT test and the PP test respectively. Throughout this article, all tests are considered to have 5% nominal level. The OT test is distribution-free and the PP test is asymptotically distribution-free. Following the suggestion of the authors, for these two test we use the cut-offs based on the asymptotic distributions of their test statistics. Our test and the DT test are calibrated using the resampling method, where the cut-off is computed based on 500 iterations. Each experiment is repeated 1000 times to estimate the power of a test by the proportion of times it rejects H_0 .

4.1 Analysis of simulated data sets

First, we investigate the level property of our test by generating random samples from some multivariate spherically symmetric distributions. In particular, we consider the standard multivariate (a) Gaussian, (b) Cauchy, and (c) t_4 distributions for this purpose and call them Examples 1(a), 1(b) and 1(c), respectively. In each case, we compute the powers for different sample sizes (n = 20, 40, 60) and dimensions $(d = 2^i, i = 1, 2, ..., 10)$, and they are reported in Figure 1. This figure clearly shows that for all three distributions, our test rejects H_0 in nearly 5% cases. The other three competing tests also exhibit similar behaviour, but to save space we do not report them here.

To compare the empirical powers of different tests, again we consider examples involving (a) Gaussian, (b) Cauchy and (c) t_4 distributions with the centre at the origin, but this time we consider the scatter matrix of the form $(1 - \rho) \mathbf{I}_d + \rho \mathbf{1}_d \mathbf{1}_d^{\top}$, where \mathbf{I}_d is the $d \times d$ identity matrix, $\mathbf{1}_d$ is the *d*-dimensional vector with all elements equal to one and $\rho \in (0, 1)$ (call them Examples 2(a), 2(b) and 2(c), respectively). Note that here the distributions are elliptically symmetric. For each example, we consider d = 5 and carry out different tests based on 100 observations. Powers of these tests are reported in Figure 2. As ρ increases from zero to one (i.e., the distribution deviates more from sphericity) one would expect the power of a test to increase. But, for OT and PP tests, these increments are negligible. In these examples, our test has the best performance followed by the DT test.



Figure 1: The level of our test for observations generated from the standard (a) Gaussian, (b) Cauchy, and (c) t_4 distributions with sample size $n = 20(\bullet)$, n = 40 (\bullet) and n = 60 (\bullet) in dimensions $d = 2^i$, i = 1, 2, ..., 10.



Figure 2: Results of our test (\bullet), OT test (\bullet), DT test (\star) and PP test (\blacksquare) in Examples 2(a)-(c).



Figure 3: Results of our test (\bullet) , OT test (\bullet) , DT test (\star) and PP test (\blacksquare) for Example 3(a)-(d).

Next, we consider four examples (call them Examples 3(a)-(d)) involving symmetric but non-elliptic distributions. In Examples 3(a) and 3(b), we deal with ℓ_p -symmetric distributions (see, e.g., Gupta and Song, 1997; Dutta et al., 2011) with $p = \infty$ and p = 1, respectively. In both cases, we generate observations on $\mathbf{X} = R\mathbf{U}$, where \mathbf{U} and $R \sim \text{Unif}(9, 10)$ are independent. In Example 3(a), we have $\mathbf{U} = \mathbf{Y}/||\mathbf{Y}||_{\infty}$ where $\mathbf{Y} = (Y_1, \ldots, Y_5)$ is uniformly distributed over the 5-dimensional unit hypercube $\{\mathbf{y} = (y_1, y_2, \ldots, y_5)^{\top} : \max\{|y_1|, |y_2|, \ldots, |y_5|\} \leq 1\}$, while in Example 3(b), we have $\mathbf{U} = \mathbf{Y}/||\mathbf{Y}||_1$ where Y_1, \ldots, Y_5 are independent standard Laplace variables. We carry out our experiment with different sample sizes, and the results are reported in Figure 3. In these examples, OT and PP tests have powers close to the nominal level of 0.05. In Example 3(b), the proposed test and the DT have comparable performance, but in Example 3(a), our test significantly outperforms the DT test.

In Example 3(c), we consider an angular symmetric distribution, We generate observations on $\mathbf{X} = \mathbf{RU}$, where $\mathbf{U} \sim \text{Unif}(\mathcal{S}^4)$ but R and \mathbf{U} are not independent. Here for any given $\mathbf{U} = \mathbf{u}(=(u_1, u_2, \dots, u_5)^{\top})$ the conditional distribution of R is uniform on $(0, \theta_{\mathbf{u}})$, where $\theta_{\mathbf{u}} = 10$ I $[u_1u_2 > 0] + 50$ I $[u_1u_2 \leq 0; u_3u_4u_5 > 0] + 100$ I $[u_1u_2 \leq 0; u_3u_4u_5 \leq 0]$, and $I[\cdot]$ is the indicator function. In Example 3(d), observations are



Figure 4: Results of our test (\bullet), OT test (\diamond), DT test (\star) and PP test (\blacksquare) for Example 4(a)-(c).

generated from an equal mixture of four normal distributions with the same dispersion matrix \mathbf{I}_5 and mean vectors $\mathbf{1}_5, -\mathbf{1}_5, \boldsymbol{\beta} = (1, -1, 1, -1, 1)^{\top}$ and $-\boldsymbol{\beta}$, respectively. In Figure 3 we see that in these examples also, our test outperforms its competitors. The DT test has the second-best performance but its power is much lower compared to our proposed test.

Finally, we consider some high-dimensional examples. In Examples 4(a) and 4(b), we generate n = 20observations from normal spiked covariance models (see. Johnstone, 2001) with mean zero and a diagonal covariance matrix with entries (d, 1, 1, ..., 1) and $(d^{0.5}, 1, 1, ..., 1)$, respectively. We carry out our experiment with different choices of d, and the results are reported in Figure 4. This figure shows that in Example 4(a) the power curve of our test exhibits a sharp increasing trend with increasing dimensions, while the other tests have non-satisfactory performances. But in Example 4(b), all tests including ours perform poorly. Note that in Example 4(a) and 4(b), the measure of sphericity (see, e.g., John, 1972; Jung and Marron, 2009) converges to 0 and 1, respectively, as d increases. So, in high dimension, the data cloud in Example 4(b) is similar to that from a spherical distribution, whereas in Example 4(a), it has significant deviations from sphericity. This explains the diametrically opposite behaviour of our test in these two examples. However, Corollary 3.4 suggests that even in Example 4(b), our test can perform well if we allow the sample size to increase with the dimensions at a suitable rate. That is what we observe in Example 4(c), where we consider the same model as in Example 4(b), but increase the sample size with the dimension. Here, we consider $n = 20 + [d^{1.5}]$, where [t] denotes the largest integer smaller than or equal to t. Figure 4 shows a sharp increasing trend in the power curve of our test. But the other competing tests have poor performance even in this set-up. This clearly indicates the superiority of our test over the OT, DT, and PP tests for high-dimensional data.

4.2 Analysis of a benchmark dataset

For further evaluation of different tests we analyze the "MAGIC Gamma Telescope" data set available at the UCI machine learning repository. This data set was generated by a Monte Carlo program called CORSIKA described in Heck et al. (1998). It is used to simulate registration of high-energy gamma particles in a groundbased atmospheric Cherenkov gamma telescope using imaging techniques. The observations are classified based on the patterns in the images the particles generate, called the shower images. Based on these shower images the particles are classified as "primary gamma" and "hadronic shower". In our analysis, we first divide the entire data set into two parts based on the class labels "primary gamma" and "hadronic shower" and call them "MAGIC-1" and "MAGIC-2", respectively. Here the observations are 10-dimensional (see Heck et al., 1998, for details). While MAGIC-1 contains 12332 observations, there are 6688 observations in MAGIC-2. When we use the full data sets (after centering by subtracting the corresponding spatial medians) for testing, all four tests reject the null hypothesis of spherical symmetry in both cases. This gives us an indication that the underlying distributions are non-spherical, and different can be compared based on their powers. But, it is difficult to compare among different test procedures using a single experiment based on the whole data set. Therefore, to compare the performances of different tests, we generate random sub-samples from these two data sets and in each case, we repeat the procedure 1000 times. The power of a test is computed by the proportion of times it rejects the null hypothesis. The results for different sub-sample sizes are reported in Figure 5.



Figure 5: Results of our test (\bullet) , OT test (\diamond) , DT test (\star) and PP test (\blacksquare) for THE Magic Gamma Telescope data set.

Here we observe that in MAGIC-1 data set, our test significantly outperforms all other tests for all sample sizes considered here. For MAGIC-2 data set, the OT and PP test has better performance for lower sample size. But the power of our test increases sharply as the sample size increases, while those of other test increase at a slower rate. Figure 5 clearly shows that our test outperforms its all competitors for sample sizes higher than 55.

5 Discussion and Conclusion

In this article, we have proposed a new measure of spherical asymmetry and a consistent estimator of this measure based on data augmentation. We have also constructed a test of spherical symmetry based on this proposed estimator and studied its large sample behaviour when the dimension may or may not grow with the sample size. Extensive simulation studies have been carried out to amply demonstrate the superiority of our test over some state-of-the-art methods. In this article, we have constructed our test assuming the centre of symmetry under H_0 to be known (which is taken as **0** throughout this article). If it is not known, we can find a suitable estimator $\hat{\theta}_0$ for the centre and apply our test on the observations $\mathbf{X}_1 - \hat{\theta}_0, \ldots, \mathbf{X}_n - \hat{\theta}_0$. We can use any depth-based median like spacial median (see, e.g., Chaudhuri, 1996; Koltchinskii, 1997) or half-space median (see, e.g., Tukey, 1975) for this purpose. One can also use other robust estimates like those based on minimum covariance determinant (MCD) (see, e.g., Rousseeuw and Driessen, 1999) or minimum volume ellipsoid (MVE) (see, e.g. Rousseeuw, 1985). Similarly, our test can be generalized for testing the ellipticity of an underlying distribution. In that case, we can find a suitable estimator $\hat{\Sigma}_0$ of the scatter matrix, and use our test on the transformed observations $\hat{\Sigma}_0^{-1/2}(\mathbf{X}_1 - \hat{\boldsymbol{\theta}}_0), \dots, \hat{\boldsymbol{\Sigma}}_0^{-1/2}(\mathbf{X}_n - \hat{\boldsymbol{\theta}}_0)$. Again, MCD or MVE estimates can be used. One can use Tyler's shape matrix Tyler (1987) as well. As long as these estimators of location and scatter are consistent, the large sample consistency of the test will be retained in arbitrary but fixed dimensions. However, at this moment it is not clear whether the resulting test will remain consistent when the dimension and the sample size increase simultaneously. We leave this for future research.

A Appendix

A.1 Expression of $\zeta(P)$ for Gaussian distributions

In this section, we present a closed-form expression for $\zeta(P)$ when P is a multivariate Gaussian distribution with mean **0** and variance-covariance matrix Σ . But before that, we present some preliminary lemmas.

Lemma A.1. If $\mathbf{X}_1 \sim N(\mathbf{0}, \mathbf{\Sigma}_1)$ and $\mathbf{X}_2 \sim N(\mathbf{0}, \mathbf{\Sigma}_2)$ are independent d-dimensional random vectors,

$$\mathbb{E}\left\{\exp\left(-\frac{1}{2d}\|\mathbf{X}_1-\mathbf{X}_2\|^2\right)\right\} = \left|\frac{1}{d}(\mathbf{\Sigma}_1+\mathbf{\Sigma}_2)+\mathbf{I}_d\right|^{-1/2},$$

where $|\mathbf{A}|$ denotes the determinant of a matrix \mathbf{A} , and \mathbf{I}_d is the $d \times d$ identity matrix.

Proof of Lemma A.1. Here $\mathbf{N} = \mathbf{X}_1 - \mathbf{X}_2$ follows $N(\mathbf{0}, \mathbf{\Sigma}_1 + \mathbf{\Sigma}_2)$. So, we have

$$\mathbb{E}\Big\{\exp\left(-\frac{1}{2d}\|\mathbf{N}\|^{2}\right)\Big\} = \int \frac{1}{(2\pi)^{d/2}\left|\boldsymbol{\Sigma}_{1} + \boldsymbol{\Sigma}_{2}\right|^{1/2}} \exp\left(-\frac{1}{2d}\|\mathbf{u}\|^{2} - \frac{1}{2}\mathbf{u}^{\top}(\boldsymbol{\Sigma}_{1} + \boldsymbol{\Sigma}_{2})^{-1}\mathbf{u}\right)d\mathbf{u}.$$

Note that the exponent on the right side is the same as that of the density of a normal distribution with mean **0** and variance-covariance matrix $\left(\frac{1}{d}\mathbf{I}_d + (\boldsymbol{\Sigma}_1 + \boldsymbol{\Sigma}_2)^{-1}\right)^{-1}$. Therefore, we have

$$\mathbb{E}\left\{\exp\left(-\frac{1}{2d}\|\mathbf{N}\|^{2}\right)\right\} = \frac{1}{\left|\boldsymbol{\Sigma}_{1} + \boldsymbol{\Sigma}_{2}\right|^{1/2} \left|\frac{1}{d}\mathbf{I}_{d} + (\boldsymbol{\Sigma}_{1} + \boldsymbol{\Sigma}_{2})^{-1}\right|^{1/2}} = \left|\frac{1}{d}(\boldsymbol{\Sigma}_{1} + \boldsymbol{\Sigma}_{2}) + \mathbf{I}_{d}\right|^{-1/2}.$$
npletes the proof.

This completes the proof.

Lemma A.2. If **X** follows a d-variate distribution with density f, $\mathbf{U} \sim \text{Unif}(\mathcal{S}^{d-1})$, and they are independent, then $\|\mathbf{X}\| \mathbf{U}$ has the density $\int f(\mathbf{H}^{\top} \mathbf{u}) \pi(\mathbf{H})$, where π is the Haar measure on the set of all $d \times d$ orthogonal matrices.

Proof of Lemma A.2. Let μ be the probability measure corresponding to distribution $\text{Unif}(\mathcal{S}^{d-1})$ and qbe a bounded continuous. Then, we have

$$\mathbb{E}\{g(\|\mathbf{X}\|\mathbf{U})\} = \int \int g(\|\mathbf{x}\|\mathbf{u})f(\mathbf{x})d\mu(\mathbf{u})d\mathbf{x} = \int \int g(\mathbf{H}\mathbf{x})f(\mathbf{x})d\pi(\mathbf{H})d\mathbf{x}$$

Since for any fixed x and u, there exists a unique orthogonal matrix H such that $\mathbf{Hx} = \|\mathbf{x}\|$, the second equality follows by substitution. Now if we substitute $\mathbf{H}\mathbf{x} = \mathbf{v}$ in the above integration we get

$$\mathbb{E}\{g(\|\mathbf{X}\|\mathbf{U})\} = \int g(\mathbf{v}) \left(\int f(\mathbf{H}^{\top}\mathbf{v}) \mathrm{d}\pi(\mathbf{H})\right) \Big| |\mathbf{H}^{\top}| \Big| \mathrm{d}\mathbf{v} = \int g(\mathbf{v}) \left(\int f(\mathbf{H}^{\top}\mathbf{v}) \mathrm{d}\pi(\mathbf{H})\right) \mathrm{d}\mathbf{v}.$$

Since g is an arbitrary bounded and continuous function, the result follows.

Proposition A.1. If X follows a d-variate normal distribution with mean 0 and variance-covariance matrix Σ , then

$$\zeta(\mathbf{P}) = \left|\frac{2}{d}\boldsymbol{\Sigma} + \mathbf{I}_d\right|^{-1/2} + \int \int \left|\frac{1}{d}(\mathbf{H}_1\boldsymbol{\Sigma}\mathbf{H}_1^\top + \mathbf{H}_2\boldsymbol{\Sigma}\mathbf{H}_2^\top) + \mathbf{I}_d\right|^{-1/2} \mathrm{d}\pi(\mathbf{H}_1)\mathrm{d}\pi(\mathbf{H}_2) - 2\int \left|\frac{1}{d}(\boldsymbol{\Sigma} + \mathbf{H}\boldsymbol{\Sigma}\mathbf{H}^\top) + \mathbf{I}_d\right|^{-1/2} \mathrm{d}\pi(\mathbf{H}),$$

where π is the Haar measure on the set of all orthogonal matrices of order $d \times d$.

Proof of Proposition A.1. Recall, that our measure can be written as (see Theorem 2.1)

$$\zeta(\mathbf{P}) = \mathbb{E}\left\{\exp\{-\frac{1}{2d}\|\mathbf{X}_1 - \mathbf{X}_2\|^2\}\right\} + \mathbb{E}\left\{\exp\{-\frac{1}{2d}\|\mathbf{X}_1' - \mathbf{X}_2'\|^2\}\right\} - 2\mathbb{E}\left\{\exp\{-\frac{1}{2d}\|\mathbf{X}_1 - \mathbf{X}_2'\|^2\}\right\}.$$

Let us look at the individual terms separately. The first term on the right side is the same as the term in Lemma A.1 with X_1 and X_2 being i.i.d. Gaussian random variables with mean zero and variance-covariance matrix Σ . Therefore,

$$\mathbb{E}\left\{\exp\{-\frac{1}{2d}\|\mathbf{X}_1-\mathbf{X}_2\|^2\}\right\} = \left|\mathbf{I}_d + \frac{2}{d}\mathbf{\Sigma}\right|^{-1/2}.$$

Now note that the second term can be written as

$$\begin{split} & \mathbb{E}\left\{\exp\{-\frac{1}{2d}\|\mathbf{X}_{1}^{\prime}-\mathbf{X}_{2}^{\prime}\|^{2}\}\right\} \\ &= \int \exp\{-\frac{1}{2d}\|\mathbf{x}_{1}-\mathbf{x}_{2}\|^{2}\}\frac{1}{(2\pi)^{d}|\mathbf{\Sigma}|}\exp\left\{-\frac{1}{2}\mathbf{x}_{1}^{\top}(\mathbf{H}_{1}\mathbf{\Sigma}\mathbf{H}_{1}^{\top})^{-1}\mathbf{x}_{1}-\frac{1}{2}\mathbf{x}_{2}^{\top}(\mathbf{H}_{2}\mathbf{\Sigma}\mathbf{H}_{2}^{\top})^{-1}\mathbf{x}_{2}\right\}\mathrm{d}\mathbf{x}_{1}\mathrm{d}\mathbf{x}_{2}\mathrm{d}\pi(\mathbf{H}_{1})\mathrm{d}\pi(\mathbf{H}_{2}) \\ &= \int \frac{\left|\mathbf{\Sigma}\right|^{-1}\left|\mathbf{H}_{1}\mathbf{\Sigma}\mathbf{H}_{1}^{\top}\right|^{1/2}\left|\mathbf{H}_{2}\mathbf{\Sigma}\mathbf{H}_{2}^{\top}\right|^{1/2}}{\left|\frac{1}{d}(\mathbf{H}_{1}\mathbf{\Sigma}\mathbf{H}_{1}^{\top}+\mathbf{H}_{2}\mathbf{\Sigma}\mathbf{H}_{2}^{\top})+\mathbf{I}_{d}\right|^{-1/2}\mathrm{d}\pi(\mathbf{H}_{1})\mathrm{d}\pi(\mathbf{H}_{2})} = \int \left|\frac{1}{d}(\mathbf{H}_{1}\mathbf{\Sigma}\mathbf{H}_{1}^{\top}+\mathbf{H}_{2}\mathbf{\Sigma}\mathbf{H}_{2}^{\top})+\mathbf{I}_{d}\right|^{-1/2}\mathrm{d}\pi(\mathbf{H}_{1})\mathrm{d}\pi(\mathbf{H}_{2}). \end{split}$$

Similarly, one can also show that,

$$\mathbb{E}\left\{\exp\{-\frac{1}{2d}\|\mathbf{X}_1-\mathbf{X}_2'\|^2\}\right\} = \int \left|\frac{1}{d}(\mathbf{H}\boldsymbol{\Sigma}\mathbf{H}^\top+\boldsymbol{\Sigma})+\mathbf{I}_d\right|^{-1/2} \mathrm{d}\pi(\mathbf{H}).$$

This completes the proof.

A.2 Proofs of the results stated in Section 2

Proof of Proposition 2.1. It is easy to see that if **X** is spherically distributed, then $\varphi_1 = \varphi_2$ and hence $\zeta(\mathbf{P}) = 0$. So, let us prove the only if part. Recall that our measure has the alternate expression

$$\zeta(\mathbf{P}) = \int \left|\varphi_1(\mathbf{t}) - \varphi_2(\mathbf{t})\right|^2 \frac{d^{d/2}}{(2\pi)^{d/2}} e^{-d\|\mathbf{t}\|^2/2} \mathrm{d}\mathbf{t}.$$

The non-negativity of $\zeta(\mathbf{P})$ follows from the non-negativity of the integrand. Now $\zeta(\mathbf{P}) = 0$ implies $\varphi_1(\mathbf{t}) = \varphi_2(\mathbf{t})$ over the set $\{\mathbf{t} \mid e^{-d\|\mathbf{t}\|^2/2} > 0\}$, which is in fact the entire Euclidean space \mathbb{R}^d . Therefore, $\zeta(\mathbf{P}) = 0$ implies \mathbf{X} and $\|\mathbf{X}\|\mathbf{U}$ are identically distributed, i.e., X is spherically distributed.

Proof of Lemma 2.2. Note that

$$\int \exp\left\{i\langle \mathbf{T}, \mathbf{X}_1 - \mathbf{X}_2\rangle\right\} \frac{d^{d/2}}{(2\pi)^{d/2}} e^{-d\|\mathbf{T}\|^2/2} \mathrm{d}\mathbf{T} = \mathbb{E}\left\{\exp\left\{i\langle \mathbf{T}, \mathbf{X}_1 - \mathbf{X}_2\rangle\right\}\right\},\tag{4}$$

where **T** follows a *d*-variate Gaussian distribution with mean zero and variance-covariance matrix $d^{-1}\mathbf{I}_d$. The right side of equation (4) is the characteristic function of **T** evaluated at $\mathbf{X}_1 - \mathbf{X}_2$. Hence, we have the desired result.

Proof of Theorem 2.1. First note that $|\varphi_1(\mathbf{t}) - \varphi_2(\mathbf{t})|^2 = (\varphi_1(\mathbf{t}) - \varphi_2(\mathbf{t}))(\varphi_1(-\mathbf{t}) - \varphi_2(-\mathbf{t}))$. So, expanding the characteristic functions in terms of expectations, we get

$$|\varphi_{1}(\mathbf{t}) - \varphi_{2}(\mathbf{t})|^{2} = \mathbb{E}\left\{\exp\left\{i\langle\mathbf{t}, \mathbf{X}_{1} - \mathbf{X}_{2}\rangle\right\}\right\} + \mathbb{E}\left\{\exp\left\{i\langle\mathbf{t}, \mathbf{X}_{1}^{'} - \mathbf{X}_{2}^{'}\rangle\right\}\right\} - \mathbb{E}\left\{\exp\left\{i\langle\mathbf{t}, \mathbf{X}_{1}^{'} - \mathbf{X}_{2}\rangle\right\}\right\}.$$
(5)

From Lemma 2.2, for any \mathbf{V}_1 and \mathbf{V}_2 , we have

$$\int \mathbb{E}\Big\{\exp\left\{i\langle \mathbf{t}, \mathbf{V}_1 - \mathbf{V}_2\rangle\right\}\Big\}\frac{d^{d/2}}{(2\pi)^{d/2}}e^{-d\|\mathbf{t}\|^2/2}\mathrm{d}\mathbf{t} = \mathbb{E}\Big\{\exp\left\{-\frac{1}{2d}\|\mathbf{V}_1 - \mathbf{V}_2\|^2\right\}\Big\}.$$

Applying this to all four terms in (5) (note that the last two terms are equal), we get the result. \Box

Proof of Theorem 2.2. As introduced in Section 2.2 we can write our estimator as

$$\hat{\zeta}_n = \frac{2}{n(n-1)} \sum_{1 \le i < j \le n} g\big((\mathbf{X}_i, \mathbf{X}_i'), (\mathbf{X}_j, \mathbf{X}_j') \big),$$

where

$$g((\mathbf{x}_1, \mathbf{x}_2), (\mathbf{y}_1, \mathbf{y}_2)) = \exp\left\{-\frac{\|\mathbf{x}_1 - \mathbf{y}_1\|^2}{2d}\right\} + \exp\left\{-\frac{\|\mathbf{x}_2 - \mathbf{y}_2\|^2}{2d}\right\} - \exp\left\{-\frac{\|\mathbf{x}_1 - \mathbf{y}_2\|^2}{2d}\right\} - \exp\left\{-\frac{\|\mathbf{x}_2 - \mathbf{y}_1\|^2}{2d}\right\}$$

Now let $\hat{\zeta}_n^{(i)}$ denote our estimator when the i^{th} observation $(\mathbf{X}_i, \mathbf{X}'_i)$ is replaced by an independent copy $(\mathbf{Y}_i, \mathbf{Y}'_i)$ of the same. Note that

$$\begin{aligned} |\hat{\zeta}_{n} - \hat{\zeta}_{n}^{(i)}| &\leq \frac{2}{n(n-1)} \Biggl\{ \sum_{j=1}^{i-1} \left| g\bigl((\mathbf{X}_{j}, \mathbf{X}_{j}'), (\mathbf{X}_{i}, \mathbf{X}_{i}') \bigr) - g\bigl((\mathbf{X}_{j}, \mathbf{X}_{j}'), (\mathbf{Y}_{i}, \mathbf{Y}_{i}') \bigr) \right| \\ &+ \sum_{j=i+1}^{n} \left| g\bigl((\mathbf{X}_{i}, \mathbf{X}_{i}'), (\mathbf{X}_{j}, \mathbf{X}_{j}') \bigr) - g\bigl((\mathbf{Y}_{i}, \mathbf{Y}_{i}'), (\mathbf{X}_{j}, \mathbf{X}_{j}') \bigr) \right| \Biggr\} \end{aligned}$$

Since $|g(\cdot, \cdot)| \leq 2$, this implies $|\hat{\zeta}_n - \hat{\zeta}_n^{(i)}| \leq \frac{8(n-1)}{n(n-1)} \leq \frac{8}{n}$. So, applying bounded difference inequality (see page 37 in Wainwright, 2019)

$$\mathbb{P}\{\left|\hat{\zeta}_n - \zeta(\mathbf{P})\right| > \epsilon\} \le \exp\left\{-\frac{2\epsilon^2}{\sum_{i=1}^n \frac{64}{n^2}}\right\} = \exp\left\{-\frac{n\epsilon^2}{32}\right\}.$$

This completes the proof.

Proof of Theorem 2.3. Note that our estimator $\hat{\zeta}_n$ is a U-statistic with the kernel

$$g\left((\mathbf{x}_{1}, \mathbf{x}_{2}), (\mathbf{y}_{1}, \mathbf{y}_{2})\right) = \exp\left\{-\frac{\|\mathbf{x}_{1} - \mathbf{y}_{1}\|^{2}}{2d}\right\} + \exp\left\{-\frac{\|\mathbf{x}_{2} - \mathbf{y}_{2}\|^{2}}{2d}\right\} - \exp\left\{-\frac{\|\mathbf{x}_{1} - \mathbf{y}_{2}\|^{2}}{2d}\right\} - \exp\left\{-\frac{\|\mathbf{x}_{2} - \mathbf{y}_{1}\|^{2}}{2d}\right\}$$

of degree 2. The first order Hoeffding projection of g(.,.) is

$$g_1((\mathbf{x}_1, \mathbf{x}_2)) = \mathbb{E}\left\{K(\mathbf{x}_1, \mathbf{X}_1)\right\} + \mathbb{E}\left\{K(\mathbf{x}_2, \mathbf{X}_1')\right\} - \mathbb{E}\left\{K(\mathbf{x}_1, \mathbf{X}_1')\right\} - \mathbb{E}\left\{K(\mathbf{x}_2, \mathbf{X}_1)\right\}$$
(6)

where $K(\mathbf{x}, \mathbf{y}) = \exp\{-\|\mathbf{x} - \mathbf{y}\|^2/(2d)\}$, $\mathbf{X}_1 \sim \mathbf{P}$ and $\mathbf{X}'_1 = \|\mathbf{X}_1\|\mathbf{U}_1$ for $\mathbf{U}_1 \sim \text{Unif}(\mathcal{S}^{d-1})$ independent of \mathbf{X}_1 . Under H_0 , \mathbf{X}_1 and \mathbf{X}'_1 are identically distributed and hence $g_1((\mathbf{x}_1, \mathbf{x}_2)) = 0$. Therefore, using Theorem 1 from Lee (1990) p.79, we get that $n\hat{\zeta}_n$ converges in distribution to $\sum_{i=1}^{\infty} \lambda_i(Z_i^2 - 1)$ where $\{Z_i\}$ are independent standard normal random variables and $\{\lambda_i\}$ are the eigenvalues of the integral equation

$$\int g\big((\mathbf{x}_1,\mathbf{x}_2),(\mathbf{y}_1,\mathbf{y}_2)\big)f\big((\mathbf{y}_1,\mathbf{y}_2)\big)\mathrm{d}F\big((\mathbf{y}_1,\mathbf{y}_2)\big) = \lambda f\big((\mathbf{x}_1,\mathbf{x}_2)\big),$$

where F is the joint distribution of $(\mathbf{X}_1, \mathbf{X}'_1)$.

Proof of Theorem 2.4. Under H_1 , the function $g_1((\mathbf{x}_1, \mathbf{x}_2))$ defined in equation (6) is non-degenerate. Therefore, using Theorem 1 from Lee (1990) p.76, we get the asymptotic normality of $\sqrt{n}(\hat{\zeta}_n - \zeta(\mathbf{P}))$ with the mean zero and the variance $4\sigma_1^2$, where $\sigma_1^2 = \operatorname{Var}(g_1((\mathbf{X}_1, \mathbf{X}_1)))$.

Proof of Lemma 2.3. Define $\mathbf{X}'_1, \mathbf{X}'_2, \ldots, \mathbf{X}'_n$ as in Theorem 2.3. Under H_0 , $(\mathbf{X}_i, \mathbf{X}'_i)$ and $(\mathbf{X}'_i, \mathbf{X}_i)$ are identically distributed for each $i = 1, 2, \ldots, n$. Therefore, the joint distribution of $(\hat{\zeta}_n, c_{1-\alpha})$ is identical to the joint distribution of $(\hat{\zeta}_n(\pi), c_{1-\alpha})$, where π is an element of $\{0, 1\}^n$. Let $\mathcal{D}' = \{(\mathbf{X}_i, \mathbf{X}'_i) \mid i = 1, 2, \ldots, n\}$ denote the augmented data. Then,

$$\mathbb{P}\{\hat{\zeta}_n > c_{1-\alpha}\} = \mathbb{P}\{\hat{\zeta}_n(\pi) > c_{1-\alpha}\} = \mathbb{E}\{\mathbb{P}\{\hat{\zeta}_n(\pi) > c_{1-\alpha} \mid \mathcal{D}'\}\} \le \alpha.$$

The last inequality follows from the definition of $c_{1-\alpha}$ (given in the resampling algorithm in page 4).

Proof of Lemma 2.4. Here $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_n$ are independent copies of $\mathbf{X} \sim \mathbf{P}$. Define $\mathbf{X}'_1, \mathbf{X}'_2, \ldots, \mathbf{X}'_n$ as in Theorem 2.3, and \mathcal{D}' as in the proof of Lemma 2.3. Let $\pi = (\pi(1), \pi(2), \ldots, \pi(n))$ be uniformly distributed on the set $\{0, 1\}^n$. For $i = 1, 2, \ldots, n$, define $\mathbf{Y}_i = \pi(i)\mathbf{X}_i + (1 - \pi(i))\mathbf{X}'_i$ and $\mathbf{Y}'_i = (1 - \pi(i))\mathbf{X}_i + \pi(i)\mathbf{X}'_i$. Since $g((\mathbf{Y}_i, \mathbf{Y}'_i), (\mathbf{Y}_i, \mathbf{Y}'_i)) \geq 0$ for all $i = 1, 2, \ldots, n$, we have

$$n(n-1)\hat{\zeta}_n(\pi) = \sum_{1 \le i \ne j \le n} g\big((\mathbf{Y}_i, \mathbf{Y}'_i), (\mathbf{Y}_j, \mathbf{Y}'_j)\big) \le \sum_{1 \le i, j \le n} g\big((\mathbf{Y}_i, \mathbf{Y}'_i), (\mathbf{Y}_j, \mathbf{Y}'_j)\big) = n^2 \zeta(F_n),$$

where F_n denotes the empirical probability distribution of $(\mathbf{Y}_1, \mathbf{Y}'_1), (\mathbf{Y}_2, \mathbf{Y}'_2), \dots, (\mathbf{Y}_n, \mathbf{Y}'_n)$. Since, $\zeta(F_n)$ is a non-negative random variable, using Markov inequality, for any $\epsilon > 0$, we have

$$\mathbb{P}\left\{\hat{\zeta}_n(\pi) > \epsilon \mid \mathcal{D}'\right\} \le \mathbb{P}\left\{n^2 \zeta(F_n) > n(n-1)\epsilon \mid \mathcal{D}'\right\} \le \frac{n^2}{n(n-1)\epsilon} \mathbb{E}\left\{\zeta(F_n) \mid \mathcal{D}'\right\}.$$

Now, taking $\epsilon = \frac{n}{(n-1)\alpha} \mathbb{E}\{\zeta(F_n) \mid \mathcal{D}'\}$, we get $\mathbb{P}\{\hat{\zeta}_n(\pi) > \epsilon \mid \mathcal{D}'\} \leq \alpha$. Therefore, from the definition of $c_{1-\alpha}$ (see page 4), we have

$$c_{1-\alpha} \leq \frac{n}{(n-1)\alpha} \mathbb{E}\{\zeta(F_n) \mid \mathcal{D}'\}.$$

Also, note that

$$\mathbb{E}\{\zeta(F_n) \mid \mathcal{D}'\} = \frac{n(n-1)}{n^2} \mathbb{E}\left\{g\left((\mathbf{Y}_1, \mathbf{Y}_1'), (\mathbf{Y}_2, \mathbf{Y}_2')\right) \mid \mathcal{D}'\right\} + \frac{1}{n} \mathbb{E}\left\{g\left((\mathbf{Y}_1, \mathbf{Y}_1'), (\mathbf{Y}_1, \mathbf{Y}_1')\right) \mid \mathcal{D}'\right\}.$$

Now
$$\mathbb{E}\left\{\exp\left\{-\frac{\|\mathbf{Y}_{1}-\mathbf{Y}_{2}\|^{2}}{2d}\right\} \mid \mathcal{D}'\right\}$$

= $\frac{1}{2^{n}}\sum_{\pi \in \{0,1\}^{n}}\exp\left\{-\frac{\|\pi(1)\mathbf{X}_{1}+(1-\pi(1))\mathbf{X}_{1}'-\pi(2)\mathbf{X}_{2}+(1-\pi(2))\mathbf{X}_{2}'\|^{2}}{2d}\right\} = \delta_{n}$ (say).

Similarly one can also show that

$$\mathbb{E}\left\{\exp\left\{-\frac{\|\mathbf{Y}_1'-\mathbf{Y}_2'\|^2}{2d}\right\} \mid \mathcal{D}'\right\} = \mathbb{E}\left\{\exp\left\{-\frac{\|\mathbf{Y}_1-\mathbf{Y}_2'\|^2}{2d}\right\} \mid \mathcal{D}'\right\} = \mathbb{E}\left\{\exp\left\{-\frac{\|\mathbf{Y}_1'-\mathbf{Y}_2\|^2}{2d}\right\} \mid \mathcal{D}'\right\} = \delta_n.$$

Hence, $\mathbb{E}\left\{g\left((\mathbf{Y}_1, \mathbf{Y}_1'), (\mathbf{Y}_2, \mathbf{Y}_2')\right) \mid \mathcal{D}'\right\} = \delta_n + \delta_n - \delta_n - \delta_n = 0$. Similarly one can also show that

$$\mathbb{E}\left\{g\left((\mathbf{Y}_1, \mathbf{Y}_1'), (\mathbf{Y}_1, \mathbf{Y}_1')\right) \mid \mathcal{D}'\right\} = 2\left(1 - \delta_n\right) \le 2$$

Combining these, we get $c_{1-\alpha} \leq 2(\alpha(n-1))^{-1}$. This completes the proof.

Proof of Theorem 2.5. Note that $\hat{\zeta}_n$ is a consistent estimator of $\zeta(\mathbf{P})$ (follows from Theorem 2.2), where $\zeta(\mathbf{P}) = 0$ under H_0 and positive under H_1 (follows from Proposition 2.1). So, by Lemma 2.4 under H_1 , the power of the test $\mathbb{P}(\hat{\zeta}_n > c_{1-\alpha})$ converges to one as n diverges to infinity.

Proof of Theorem 2.6. Let us define the distribution functions $F(t) = \frac{1}{2^n} \left\{ \sum_{\pi \in \{0,1\}^n} I[\hat{\zeta}_n(\pi) \le t] \right\}$ and $F_B(t) = \frac{1}{B} \left\{ \sum_{i=1}^B I[\hat{\zeta}_n(\pi_i) \le t] \right\}$ conditioned on the augmented data \mathcal{D}' . Then,

$$\begin{aligned} \left| p_n - p_{n,B} \right| &= \left| \frac{1}{2^n} \bigg\{ \sum_{\pi \in \{0,1\}^n} I[\hat{\zeta}_n(\pi) \ge \hat{\zeta}_n] \bigg\} - \frac{1}{B+1} \bigg\{ \sum_{i=1}^B I[\hat{\zeta}_n(\pi_i) \ge \hat{\zeta}_n] + 1 \bigg\} \right| \\ &= \left| \frac{1}{2^n} \bigg\{ \sum_{\pi \in \{0,1\}^n} I[\hat{\zeta}_n(\pi) < \hat{\zeta}_n] \bigg\} - \frac{1}{B+1} \bigg\{ \sum_{i=1}^B I[\hat{\zeta}_n(\pi_i) < \hat{\zeta}_n] \bigg\} \right| \\ &= \left| F(\hat{\zeta}_n -) - \frac{B}{B+1} F_B(\hat{\zeta}_n -) \right| \\ &\leq \left| F(\hat{\zeta}_n -) - F_B(\hat{\zeta}_n -) \right| + \left| \frac{F_B(\hat{\zeta}_n -)}{B+1} \right| \le \sup_{t \in \mathbb{R}} \left| F(t) - F_B(t) \right| + \frac{1}{B+1} \end{aligned}$$

Conditioned on \mathcal{D}' , the Dvoretzky-Keifer-Wolfwitz inequality (Massart (1990)) gives us $\mathbb{P}\{\sup_{t\in\mathbb{R}} | F(t) - F_B(t)| > \epsilon\} \le 2e^{-2B\epsilon^2}$. Hence, conditioned on \mathcal{D}' , as B grows to infinity the randomized p-value $p_{n,B}$ converges almost surely to p_n .

A.3 Proofs of the results stated in Section 3

Proof of Lemma 3.1. Recall that $\zeta(P)$ can be expressed as

$$\zeta(\mathbf{P}) = \mathbb{E}\{K(\mathbf{X}_1, \mathbf{X}_2)\} + \mathbb{E}\{K(\mathbf{X}_1', \mathbf{X}_2')\} - 2\mathbb{E}\{K(\mathbf{X}_1, \mathbf{X}_2')\},\$$

where $\mathbf{X}_1, \mathbf{X}_2$ are independent copies of $\mathbf{X} \sim \mathbf{P}, \mathbf{X}'_1 = \|\mathbf{X}_1\|\mathbf{U}_1, \mathbf{X}'_2 = \|\mathbf{X}_2\|\mathbf{U}_2$ with $\mathbf{U}_1, \mathbf{U}_2 \stackrel{iid}{\sim} \text{Unif}(\mathcal{S}^{d-1})$ independent of $\mathbf{X}_1, \mathbf{X}_2$, and $K(\mathbf{x}, \mathbf{y}) = \exp\left\{-\frac{1}{2d}\|\mathbf{x} - \mathbf{y}\|^2\right\}$. Now if $\mathbb{P} = (1 - \delta)F + \delta G \ (0 < \delta < 1)$, then

$$\mathbb{E}\{K(\mathbf{X}_1, \mathbf{X}_2)\} = \int K(\mathbf{x}_1, \mathbf{x}_2) d\mathbb{P}(\mathbf{x}_1) d\mathbb{P}(\mathbf{x}_2)$$

= $(1 - \delta)^2 \int K(\mathbf{x}_1, \mathbf{x}_2) dF(\mathbf{x}_1) dF(\mathbf{x}_2) + 2\delta(1 - \delta) \int K(\mathbf{x}_1, \mathbf{x}_2) dG(\mathbf{x}_1) dF(\mathbf{x}_2) + \delta^2 \int K(\mathbf{x}_1, \mathbf{x}_2) dG(\mathbf{x}_1) dG(\mathbf{x}_2)$

If μ denotes the uniform distribution on (\mathcal{S}^{d-1}) , then

$$\begin{aligned} \mathbb{E}\{K(\mathbf{X}_1', \mathbf{X}_2')\} &= \int K(\|\mathbf{x}_1\|\mathbf{u}_1, \|\mathbf{x}_2\|\mathbf{u}_2) \mathrm{d}\mathbb{P}(\mathbf{x}_1) \mathrm{d}\mathbb{P}(\mathbf{x}_2) \mathrm{d}\mu(\mathbf{u}_1) \mathrm{d}\mu(\mathbf{u}_2) \\ &= (1-\delta)^2 \int K(\|\mathbf{x}_1\|\mathbf{u}_1, \|\mathbf{x}_2\|\mathbf{u}_2) \mathrm{d}F(\mathbf{x}_1) \mathrm{d}F(\mathbf{x}_2) \mathrm{d}\mu(\mathbf{u}_1) \mathrm{d}\mu(\mathbf{u}_2) \\ &+ \delta^2 \int K(\|\mathbf{x}_1\|\mathbf{u}_1, \|\mathbf{x}_2\|\mathbf{u}_2) \mathrm{d}G(\mathbf{x}_1) \mathrm{d}G(\mathbf{x}_2) \mathrm{d}\mu(\mathbf{u}_1) \mathrm{d}\mu(\mathbf{u}_2) \\ &+ 2\delta(1-\delta) \int K(\|\mathbf{x}_1\|\mathbf{u}_1, \|\mathbf{x}_2\|\mathbf{u}_2) \mathrm{d}G(\mathbf{x}_1) \mathrm{d}F(\mathbf{x}_2) \mathrm{d}\mu(\mathbf{u}_1) \mathrm{d}\mu(\mathbf{u}_2) \end{aligned}$$

and
$$\mathbb{E}\{K(\mathbf{X}_1, \mathbf{X}_2')\} = \int K(\mathbf{x}_1, \|\mathbf{x}_2\|\mathbf{u}_2) d\mathbb{P}(\mathbf{x}_1) d\mathbb{P}(\mathbf{x}_2) d\mu(\mathbf{u}_2)$$

$$= (1 - \delta)^2 \int K(\mathbf{x}_1, \|\mathbf{x}_2\|\mathbf{u}_2) dF(\mathbf{x}_1) dF(\mathbf{x}_2) d\mu(\mathbf{u}_2) + \delta(1 - \delta) \int K(\mathbf{x}_1, \|\mathbf{x}_2\|\mathbf{u}_2) dG(\mathbf{x}_1) dF(\mathbf{x}_2) d\mu(\mathbf{u}_2)$$

$$+ \delta(1 - \delta) \int K(\mathbf{x}_1, \|\mathbf{x}_2\|\mathbf{u}_2) dF(\mathbf{x}_1) dG(\mathbf{x}_2) d\mu(\mathbf{u}_2) + \delta^2 \int K(\mathbf{x}_1, \|\mathbf{x}_2\|\mathbf{u}_2) dG(\mathbf{x}_1) dG(\mathbf{x}_2) d\mu(\mathbf{u}_2).$$

Therefore, if $P = (1 - \delta)F + \delta G$, then,

$$\zeta(\mathbf{P}) = (1-\delta)^2 \zeta(F) + \delta^2 \zeta(G) + 2\delta(1-\delta)\zeta'(G,F)$$

where

$$\zeta'(G,F) = \mathbb{E}\{K(\mathbf{Y}_1,\mathbf{Y}_2)\} + \mathbb{E}\{K(\mathbf{Y}_1',\mathbf{Y}_2')\} - \mathbb{E}\{K(\mathbf{Y}_1,\mathbf{Y}_2')\} - \mathbb{E}\{K(\mathbf{Y}_1',\mathbf{Y}_2)\}$$

where $\mathbf{Y}_1 \sim G$ and $\mathbf{Y}_2 \sim F$ are independent, $\mathbf{Y}'_1 = \|\mathbf{Y}_1\| \mathbf{U}_1$ and $\mathbf{Y}'_2 = \|\mathbf{Y}_2\| \mathbf{U}_2$ where $\mathbf{U}_1, \mathbf{U}_2 \stackrel{iid}{\sim} \text{Unif}(\mathcal{S}^{d-1})$ independent of \mathbf{Y}_1 and \mathbf{Y}_2 . This completes the proof.

Lemma A.3. If $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_n$ are independent copies of $\mathbf{X} \sim P$, then

$$\operatorname{Var}(\hat{\zeta}_n) \le {\binom{n}{2}}^{-1} \left[4(n-2)\zeta(\mathbf{P}) + 4\right],$$

Proof of Lemma A.3. Recall that $\hat{\zeta}_n$ is a U-statistics with the kernel

$$g((\mathbf{x}_1, \mathbf{x}_1'), (\mathbf{x}_2, \mathbf{x}_2')) = K(\mathbf{x}_1, \mathbf{x}_2) + K(\mathbf{x}_1', \mathbf{x}_2') - K(\mathbf{x}_1, \mathbf{x}_2') - K(\mathbf{x}_1', \mathbf{x}_2),$$

where $K(\mathbf{x}, \mathbf{y}) = \exp\{-\frac{1}{2d}\|\mathbf{x} - \mathbf{y}\|^2\}$. Then, by the theory of U-statistics (see p.12 in Lee, 1990), we have

$$\operatorname{Var}(\hat{\zeta}_n) = \binom{n}{2}^{-1} \left[\binom{2}{1} \binom{n-2}{1} \operatorname{Var}\left(g_1(\mathbf{X}_1, \mathbf{X}_1')\right) + \binom{2}{2} \binom{n-2}{0} \operatorname{Var}\left(g((\mathbf{X}_1, \mathbf{X}_1'), (\mathbf{X}_2, \mathbf{X}_2'))\right) \right]$$

where,

$$g_{1}(\mathbf{x}_{1},\mathbf{x}_{1}') = \mathbb{E}\{\exp(-\frac{1}{2d}\|\mathbf{x}_{1}-\mathbf{X}_{2}\|^{2})\} - \mathbb{E}\{\exp(-\frac{1}{2d}\|\mathbf{x}_{1}-\mathbf{X}_{2}'\|^{2})\} + \mathbb{E}\{\exp(-\frac{1}{2d}\|\mathbf{x}_{1}'-\mathbf{X}_{2}'\|^{2})\} - \mathbb{E}\{\exp(-\frac{1}{2d}\|\mathbf{x}_{1}'-\mathbf{X}_{2}\|^{2})\}.$$

Since, $|g(\cdot, \cdot)| \le 2$, $\operatorname{Var}\left(g\left((\mathbf{X}_1, \mathbf{X}_1'), (\mathbf{X}_2, \mathbf{X}_2')\right)\right)$ is bounded by 4. Whereas to bound the first term note that

$$g_{1}(\mathbf{x}_{1}, \mathbf{x}_{1}') = \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{x}_{1} - \mathbf{X}_{2} \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} - \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{x}_{1} - \mathbf{X}_{2}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} - \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{x}_{1}' - \mathbf{X}_{2} \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} + \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{x}_{1}' - \mathbf{X}_{2}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} - \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{x}_{1}' - \mathbf{X}_{2}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} + \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{x}_{1}' - \mathbf{X}_{2}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} - \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{x}_{1}' - \mathbf{X}_{2}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} + \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{x}_{1}' - \mathbf{X}_{2}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} - \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{t}, \mathbf{t}' - \mathbf{t}, \mathbf{t}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} + \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{t}, \mathbf{t}' - \mathbf{t}, \mathbf{t}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} - \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{t}, \mathbf{t}' - \mathbf{t}, \mathbf{t}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} + \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{t}, \mathbf{t}' - \mathbf{t}, \mathbf{t}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} + \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{t}, \mathbf{t}' - \mathbf{t}, \mathbf{t}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} + \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{t}, \mathbf{t}' - \mathbf{t}, \mathbf{t}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} + \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{t}, \mathbf{t}' - \mathbf{t}, \mathbf{t}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} + \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{t}, \mathbf{t}' - \mathbf{t}, \mathbf{t}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} + \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{t}, \mathbf{t}' - \mathbf{t}, \mathbf{t}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} + \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{t}, \mathbf{t}' - \mathbf{t}, \mathbf{t}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} + \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{t}, \mathbf{t}' - \mathbf{t}, \mathbf{t}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} + \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{t}, \mathbf{t}' - \mathbf{t}, \mathbf{t}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} + \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{t}, \mathbf{t}' - \mathbf{t}, \mathbf{t}' \rangle)\phi_{0}(\mathbf{t})d\mathbf{t}\} + \mathbb{E}\{\int \exp(i\langle \mathbf{t}, \mathbf{t}, \mathbf{t}' - \mathbf{t}, \mathbf{t}' \rangle)\phi_{0}(\mathbf{t}, \mathbf{t}' + \mathbf{t}' \rangle\}$$

$$= \int \left(\exp(i\langle \mathbf{t}, \mathbf{x}_1 \rangle) - \exp(i\langle \mathbf{t}, \mathbf{x}_1' \rangle) \right) \left(\mathbb{E} \{ \exp(-i\langle \mathbf{t}, \mathbf{X}_2 \rangle) \} - \mathbb{E} \{ \exp(-i\langle \mathbf{t}, \mathbf{X}_2' \rangle) \} \right) \phi_0(\mathbf{t}) d\mathbf{t}$$

$$= \int (\exp(i\langle \mathbf{t}, \mathbf{x}_1 \rangle) - \exp(i\langle \mathbf{t}, \mathbf{x}_1' \rangle)) (\varphi_1(-\mathbf{t}) - \varphi_2(-\mathbf{t})) \phi_0(\mathbf{t}) d\mathbf{t}$$

$$= \int (\exp(i\langle \mathbf{t}, \mathbf{x}_1 \rangle) - \exp(i\langle \mathbf{t}, \mathbf{x}_1' \rangle)) \overline{(\varphi_1(\mathbf{t}) - \varphi_2(\mathbf{t}))} \phi_0(\mathbf{t}) d\mathbf{t}.$$

Then using Cauchy-Schwartz inequality we have,

$$g_1^2(\mathbf{x}_1, \mathbf{x}_1') \le \int |\exp(i\langle \mathbf{t}, \mathbf{x}_1 \rangle) - \exp(i\langle \mathbf{t}, \mathbf{x}_1' \rangle)|^2 \phi_0(\mathbf{t}) d\mathbf{t} \times \int |\varphi_1(\mathbf{t}) - \varphi_2(\mathbf{t})|^2 \phi_0(\mathbf{t}) d\mathbf{t} \le 2\zeta(\mathbf{P}).$$

This gives us,

$$\operatorname{Var}(\hat{\zeta}_n) \le {\binom{n}{2}}^{-1} \left[4(n-2)\zeta(\mathbf{P}) + 4\right].$$

This completes the proof.

Proof of Theorem 3.1. Here $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_n$ be independent copies of $\mathbf{X} \sim F_{\delta} = (1 - \delta)F + \delta G$, where $\zeta(G) = 0$ and $\zeta(F) = \gamma_0 > 0$. So, we have $\zeta(F_{\delta}) = (1 - \delta)^2 \gamma_0$ (follows from Lemma 3.1) and

$$\mathbb{P}\{\hat{\zeta}_n > c_{1-\alpha}\} = 1 - \mathbb{P}\{\hat{\zeta}_n \le c_{1-\alpha}\} \ge 1 - \mathbb{P}\{\hat{\zeta}_n \le 2((n-1)\alpha)^{-1}\}$$

Here, the last inequality follows from Lemma 2.4. Now, choose a large n so that $2((n-1)\alpha)^{-1} < \zeta(F_{\delta})$. Then we have

$$\mathbb{P}\{\hat{\zeta}_{n} \leq 2((n-1)\alpha)^{-1}\} = \mathbb{P}\{\hat{\zeta}_{n} - \zeta(F_{\delta}) \leq 2((n-1)\alpha)^{-1} - \zeta(F_{\delta})\}$$

$$\leq \frac{\operatorname{Var}(\hat{\zeta}_{n})}{\left(\zeta(F_{\delta}) - 2((n-1)\alpha)^{-1}\right)^{2}}$$
(by Chebyshev's inequality)
$$\leq \frac{\binom{n}{2}^{-1} \left[4(n-2)\zeta(F_{\delta}) + 4\right]}{\left(\zeta(F_{\delta}) - 2((n-1)\alpha)^{-1}\right)^{2}}$$
(by Lemma A.3)
$$= \frac{\binom{n}{2}^{-1} \left[4(n-2)(1-\delta)^{2}\gamma_{0} + 4\right]}{\left((1-\delta)^{2}\gamma_{0} - 2((n-1)\alpha)^{-1}\right)^{2}}.$$

Note that the upper bound in the above inequality does not depend on G and hence,

$$\inf_{G:\zeta(G)=0} \mathbb{P}\{\hat{\zeta}_n > c_{1-\alpha}\} \ge 1 - \frac{\binom{n}{2}^{-1} \left[4(n-2)(1-\delta)^2 \gamma_0 + 4\right]}{\left((1-\delta)^2 \gamma_0 - 2((n-1)\alpha)^{-1}\right)^2}$$

Now taking limit as n goes to infinity, $\inf_{G:\zeta(G)=0} \mathbb{P}\{\hat{\zeta}_n > c_{1-\alpha}\}$ goes to one. This completes the proof. \Box

Proof of Theorem 3.2. We prove this theorem using a standard application of the Neyman-Pearson lemma. Let Q_1 and Q_2 be the joint distribution of the sample $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_n$ under the null and alternative hypotheses respectively. Then we can lower bound the minimax risk $R_n(\epsilon)$ as follows

$$R_n(\epsilon) \ge 1 - \alpha - d_{TV}(Q_1, Q_2) \ge 1 - \alpha - \sqrt{\frac{1}{2}KL(Q_1, Q_2)}$$

The first inequality follows using the fact that for any unbiased test ϕ with $\mathbb{P}_{Q_1}\{\phi=1\} \leq \alpha$ we have,

$$\mathbb{P}_{Q_1}\{\phi=1\} + \mathbb{P}_{Q_2}\{\phi=0\} = 1 - (\mathbb{P}_{Q_1}\{\phi=0\} - \mathbb{P}_{Q_2}\{\phi=0\}) \ge 1 - d_{TV}(Q_1, Q_2)$$

where d_{TV} denotes the total variation distance between Q_1 and Q_2 . The second inequality follows from Pinsker's inequality (see. Tsybakov, 2009). Suppose G is a spherically symmetric distribution with density g and F is a distribution with density f and $\zeta(F) = \gamma_0$. Assume that $\int (f(\mathbf{u})/g(\mathbf{u}) - 1)^2 g(\mathbf{u}) d\mathbf{u} = \gamma_1 < \infty$. Then setting $Q_1 = \prod_{i=1}^n \left(\left(1 - \frac{\delta}{\sqrt{n}}\right)G + \frac{\delta}{\sqrt{n}}F \right)$ for some $\delta > 0$ and $Q_2 = G^n$, we have

$$KL(Q_1, Q_2) = \int \log \prod_{i=1}^n \left\{ 1 + \frac{\delta}{\sqrt{n}} \left(\frac{f(\mathbf{u}_i)}{g(\mathbf{u}_i)} - 1 \right) \right\} \prod_{i=1}^n d\left(\left(1 - \frac{\delta}{\sqrt{n}} \right) G + \frac{\delta}{\sqrt{n}} F \right)(\mathbf{u}_i)$$

$$= n \int \log \left\{ 1 + \frac{\delta}{\sqrt{n}} \left(\frac{f(\mathbf{u}_1)}{g(\mathbf{u}_1)} - 1 \right) \right\} d\left(\left(1 - \frac{\delta}{\sqrt{n}} \right) G + \frac{\delta}{\sqrt{n}} F \right)(\mathbf{u}_1)$$

$$= n \left(1 - \frac{\delta}{\sqrt{n}} \right) \int \log \left\{ 1 + \frac{\delta}{\sqrt{n}} \left(\frac{f(\mathbf{u}_1)}{g(\mathbf{u}_1)} - 1 \right) \right\} g(\mathbf{u}_1) d\mathbf{u}_1 + n \frac{\delta}{\sqrt{n}} \int \log \left\{ 1 + \frac{\delta}{\sqrt{n}} \left(\frac{f(\mathbf{u}_1)}{g(\mathbf{u}_1)} - 1 \right) \right\} f(\mathbf{u}_1) d\mathbf{u}_1$$

Using the inequality $\log(1+y) \leq y$ we get,

$$KL(Q_1, Q_2) \le n \left[\frac{\delta}{\sqrt{n}} (1 - \frac{\delta}{\sqrt{n}}) \int \left(\frac{f(\mathbf{u})}{g(\mathbf{u})} - 1 \right) g(\mathbf{u}) d\mathbf{u} + \frac{\delta^2}{n} \int \left(\frac{f(\mathbf{u})}{g(\mathbf{u})} - 1 \right) f(\mathbf{u}) d\mathbf{u} \right]$$
$$= \delta^2 \left[\int \frac{f^2(\mathbf{u})}{g(\mathbf{u})} - 1 \right] = \delta^2 \int \left(\frac{f(\mathbf{u})}{g(\mathbf{u})} - 1 \right)^2 g(\mathbf{u}) d\mathbf{u} = \delta^2 \gamma_1.$$

Also by Lemma 3.1 we have,

$$\zeta\left(\left(\left(1-\frac{\delta}{\sqrt{n}}\right)G+\frac{\delta}{\sqrt{n}}F\right)\right)=\frac{\delta^2}{n}\zeta(F)=\frac{\delta^2}{n}\gamma_0.$$

Now for any $0 < \beta < 1 - \alpha$ if we choose $\delta = \sqrt{2/\gamma_1}(1 - \alpha - \beta)$ we get,

$$\zeta\left(\left(\left(1-\frac{\delta}{\sqrt{n}}\right)G+\frac{\delta}{\sqrt{n}}F\right)\right)=\frac{1}{n}\left(\frac{2\gamma_0(1-\alpha-\beta)^2}{\gamma_1}\right).$$

Now define, $c(\alpha, \beta) = \left(2\gamma_0(1 - \alpha - \beta)^2\right)/\gamma_1$. Then, $\left((1 - \frac{\delta}{\sqrt{n}})G + \frac{\delta}{\sqrt{n}}F\right) \in \mathcal{F}(cn^{-1}) = \{F \mid \zeta(F) > cn^{-1}\}$ for all $0 < c < c(\alpha, \beta)$. For this choice of alternative, we also have $R_n(cn^{-1}) \ge \beta$ for all $0 < c < c(\alpha, \beta)$. Since β and $c(\alpha, \beta)$ does not depend on n, this trivially satisfies the condition $\liminf_{n\to\infty} R_n(cn^{-1}) \ge \beta$ for all $0 < c < c(\alpha, \beta)$.

Proof of Theorem 3.3. Here we want to show that for every $0 < \alpha < 1$ and $0 < \beta < 1 - \alpha$ there exists a constant $C(\alpha, \beta) > 0$ such that

$$\limsup_{n \to \infty} \sup_{F \in \mathcal{F}(cn^{-1})} \mathbb{P}_{F^n} \{ \hat{\zeta}_n \le c_{1-\alpha} \} \le \beta$$

for all $c > C(\alpha, \beta)$. Now take any $P \in \mathcal{F}(cn^{-1})$ with $c > 4/\alpha$ (i.e. $\zeta(P) > 4/n\alpha$). Using the fact $c_{1-\alpha} \leq 2((n-1)\alpha)^{-1}$ and Chebyshev's inequality, we have

$$\mathbb{P}_{F^n}\{\hat{\zeta}_n \le c_{1-\alpha}\} \le \mathbb{P}_{F^n}\{\hat{\zeta}_n \le 2((n-1)\alpha)^{-1}\} \le \mathbb{P}_{F^n}\{\zeta(\mathbf{P}) - \hat{\zeta}_n \ge \zeta(\mathbf{P}) - 2((n-1)\alpha)^{-1}\} \le \frac{\operatorname{Var}(\hat{\zeta}_n)}{\left(\zeta(\mathbf{P}) - 2((n-1)\alpha)^{-1}\right)^2}$$

which holds since $\zeta(\mathbf{P}) - 2((n-1)\alpha)^{-1} > 4(n\alpha)^{-1} - 2((n-1)\alpha)^{-1} = \frac{2n-4}{n(n-1)\alpha} > 0$ for all $n \ge 2$. Now,

$$\frac{\operatorname{Var}(\hat{\zeta}_n)}{\left(\zeta(\mathbf{P}) - 2((n-1)\alpha)^{-1}\right)^2} \le \frac{\binom{n}{2}^{-1} \left[4(n-2)\zeta(\mathbf{P}) + 4\right]}{\left(\zeta(\mathbf{P}) - 2((n-1)\alpha)^{-1}\right)^2} \qquad \text{(follows from Lemma A.3)} \tag{7}$$

which implies that

$$\limsup_{n \to \infty} \sup_{F \in \mathcal{F}(cn^{-1})} \mathbb{P}_{F^n} \{ \hat{\zeta}_n \le c_{1-\alpha} \} \le \frac{4c+4}{\left(c-2/\alpha\right)^2}.$$

It is easy to see that the upper bound is a monotonically decreasing function of c for $c > 4/\alpha$ and it converges to 0 as c increases. Hence for any $\beta < 1-\alpha$, there exists a $r(\alpha, \beta)$ such that the upper bound is smaller than β whenever $c > r(\alpha, \beta)$. Now set $C(\alpha, \beta) = \max\{r(\alpha, \beta), 4/\alpha\}$. Then for any $c > C(\alpha, \beta)$ the maximum type II error rate of our test is upper bounded by β . This completes the proof of this theorem.

Proof of Theorem 3.4. If the distribution P is such that $n\zeta(P)$ diverges to infinity, then from the proof of Theorem 3.3 (see equation (7)) we see that $\lim \mathbb{P}\{\hat{\zeta}_n \leq c_{1-\alpha}\} = 0$. Hence, under the above condition, the power of our test converges to one.

Proof of Proposition 3.1. It is easy to see that the likelihood ratio of $F_{1-\frac{\beta}{\sqrt{n}}} = (1-\beta_n/\sqrt{n})G + \beta_n/\sqrt{n}F$ and G is $\left(1 + \frac{\beta_n}{\sqrt{n}}(f(\mathbf{u})/g(\mathbf{u}) - 1)\right)$. Hence if $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_n$ are i.i.d. observations from G, then the log-likelihood ratio is given by,

$$L_N = \log \left\{ \prod_{i=1}^n \frac{dF_{1-\frac{\beta_n}{\sqrt{n}}}}{dG} (\mathbf{X}_i) \right\} = \sum_{i=1}^n \log \left\{ \frac{dF_{1-\frac{\beta_n}{\sqrt{n}}}}{dG} (\mathbf{X}_i) \right\} = \sum_{i=1}^n \log \left(1 + \frac{\beta_n}{\sqrt{n}} (f(\mathbf{X}_i)/g(\mathbf{X}_i) - 1) \right).$$

Using the fact that $\log(1+y) = y - \frac{y^2}{2} + \frac{1}{2}y^2h(y)$ where $h(\cdot)$ is continuous and $\lim_{y \to 0} h(y) = 0$, we get

$$L_N = \sum_{i=1}^n \frac{\beta_n}{\sqrt{n}} (f(\mathbf{X}_i)/g(\mathbf{X}_i) - 1) - \sum_{i=1}^n \frac{\beta_n^2}{2n} (f(\mathbf{X}_i)/g(\mathbf{X}_i) - 1)^2 + \sum_{i=1}^n \frac{\beta_n^2}{2n} (f(\mathbf{X}_i)/g(\mathbf{X}_i) - 1)^2 h \left(\frac{\beta_n}{\sqrt{n}} (f(\mathbf{X}_i)/g(\mathbf{X}_i) - 1)\right) h \left(\frac{\beta_n}{\sqrt{n}} (f(\mathbf{X}_i)/g(\mathbf{X}_i)$$

Under assumption $\int (f(\mathbf{u})/g(\mathbf{u}) - 1)^2 g(\mathbf{u}) d\mathbf{u}$ is finite and $\beta_n \to \beta$, as n grows to infinity, we have

$$\sum_{i=1}^{n} \frac{\beta_n^2}{n} (f(\mathbf{X}_i)/g(\mathbf{X}_i) - 1)^2 \xrightarrow{a.s.} \beta^2 \mathbb{E} ((f(\mathbf{X}_1)/g(\mathbf{X}_1) - 1)^2).$$

Hence we only need to show that

$$\sum_{i=1}^{n} \frac{\beta_n^2}{n} \left(f(\mathbf{X}_i) / g(\mathbf{X}_i) - 1 \right)^2 h\left(\frac{\beta_n}{\sqrt{n}} \left(f(\mathbf{X}_i) / g(\mathbf{X}_i) - 1 \right) \right)$$

converges to zero in probability. Notice that

$$\sum_{i=1}^{n} \frac{\beta_{n}^{2}}{n} \left(f(\mathbf{X}_{i})/g(\mathbf{X}_{i}) - 1 \right)^{2} h\left(\frac{\beta_{n}}{\sqrt{n}} \left(f(\mathbf{X}_{i})/g(\mathbf{X}_{i}) - 1 \right) \right) \le \max_{1 \le i \le n} \left| h\left(\frac{\beta_{n}}{\sqrt{n}} \left(f(\mathbf{X}_{i})/g(\mathbf{X}_{i}) - 1 \right) \right) \right| \sum_{i=1}^{n} \frac{\beta_{n}^{2}}{n} \left(f(\mathbf{X}_{i})/g(\mathbf{X}_{i}) - 1 \right)^{2} h\left(\frac{\beta_{n}}{\sqrt{n}} \left(f(\mathbf{X}_{i})/g(\mathbf{X}_{i}) - 1 \right) \right) \le \max_{1 \le i \le n} \left| h\left(\frac{\beta_{n}}{\sqrt{n}} \left(f(\mathbf{X}_{i})/g(\mathbf{X}_{i}) - 1 \right) \right) \right| \sum_{i=1}^{n} \frac{\beta_{n}^{2}}{n} \left(f(\mathbf{X}_{i})/g(\mathbf{X}_{i}) - 1 \right)^{2} h\left(\frac{\beta_{n}}{\sqrt{n}} \left(f(\mathbf{X}_{i})/g(\mathbf{X}_{i}) - 1 \right) \right) \le \max_{1 \le i \le n} \left| h\left(\frac{\beta_{n}}{\sqrt{n}} \left(f(\mathbf{X}_{i})/g(\mathbf{X}_{i}) - 1 \right) \right) \right|$$

Therefore, it suffices to show that $\max_{1 \le i \le N} |h\left(\frac{\beta_n}{\sqrt{n}}(f(\mathbf{X}_i)/g(\mathbf{X}_i) - 1\right)|$ converges to zero in probability, which follows if $\max_{1 \le i \le N} |\frac{\beta_n}{\sqrt{n}}(f(\mathbf{X}_i)/g(\mathbf{X}_i) - 1)|$ converges to zero in probability (as $\lim_{y\to 0} h(y) = 0$ and

it is continuous). Note that,

$$\begin{split} \mathbb{P}\Big\{\max_{1\leq i\leq n} \left|\frac{1}{\sqrt{n}} \big(f(\mathbf{X}_i)/g(\mathbf{X}_i)-1\big)\right| > \epsilon\Big\} &\leq \sum_{i=1}^n \mathbb{P}\Big\{\left|\frac{1}{\sqrt{n}} \big(f(\mathbf{X}_i)/g(\mathbf{X}_i)-1\big)\right| > \epsilon\Big\} \\ &= n\mathbb{P}\Big\{\left|\frac{1}{\sqrt{n}} \big(f(\mathbf{X}_1)/g(\mathbf{X}_1)-1\big)\right| > \epsilon\Big\} \\ &= n\mathbb{E}\Big\{I\big(\left|\frac{1}{\sqrt{n}} \big(f(\mathbf{X}_1)/g(\mathbf{X}_1)-1\big)\right| > \epsilon\big)\Big\} \\ &\leq n\mathbb{E}\Big\{\frac{\big(f(\mathbf{X}_1)/g(\mathbf{X}_1)-1\big)^2}{n\epsilon^2}I\big|\frac{1}{\sqrt{n}} \big(f(\mathbf{X}_1)/g(\mathbf{X}_1)-1\big)\big| > \epsilon\Big\} \\ &\leq \frac{1}{\epsilon^2}\mathbb{E}\Big\{\big(f(\mathbf{X}_1)/g(\mathbf{X}_1)-1\big)^2I\big(\left|\frac{1}{\sqrt{n}} \big(f(\mathbf{X}_1)/g(\mathbf{X}_1)-1\big)\big| > \epsilon\big)\Big\}. \end{split}$$

Since $I(\left|\frac{1}{\sqrt{n}}(f(\mathbf{X}_1)/g(\mathbf{X}_1)-1)\right| > \epsilon)$ converges to zero in probability, the right-hand side converges to zero by the Dominated Convergence Theorem. Hence, we have

$$\left|\log\left\{\prod_{i=1}^{n}\frac{dF_{1-\beta_n/\sqrt{n}}}{dG}(\mathbf{X}_i)\right\} - \frac{\beta_n}{\sqrt{n}}\sum_{i=1}^{n}\left(f(\mathbf{X}_i)/g(\mathbf{X}_i) - 1\right) + \frac{\beta_n^2}{2}\mathbb{E}\left\{f(\mathbf{X}_1)/g(\mathbf{X}_1) - 1\right\}^2\right| \to 0,$$

in probability as n goes to infinity. This completes the proof.

Proof of Theorem 3.5. Let $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_n \stackrel{i.i.d.}{\sim} G$ and $\mathbf{U}_1, \mathbf{U}_2, \dots, \mathbf{U}_n \stackrel{i.i.d.}{\sim}$ Unif (\mathcal{S}^{d-1}) be independent and $\zeta(G) = 0$. For $i = 1, 2, \dots, n$, define, $\mathbf{X}'_i = \|\mathbf{X}_i\|\mathbf{U}_i$. To prove this theorem we only need to find the limit distribution of $\sqrt{n}(\frac{1}{n}\sum_{i=1}^n h(\mathbf{X}_i, \mathbf{X}'_i) - \mathbb{E}\{h(\mathbf{X}_1, \mathbf{X}'_1)\})$ for some square-integrable function h under the contiguous alternative $F_{1-\beta/\sqrt{n}} = (1 - \frac{\beta_n}{\sqrt{n}})G + \frac{\beta_n}{\sqrt{n}}F$, where $\zeta(F) > 0$ and $\beta_n \to \beta$ as $n \to \infty$. Using the bivariate central limit theorem, we can say that as n diverges to infinity, the joint distribution of

$$\sqrt{n} \left(\frac{1}{n} \sum_{i=1}^{n} h(\mathbf{X}_i, \mathbf{X}'_i) - \mathbb{E} \left\{ h(\mathbf{X}_1, \mathbf{X}'_1) \right\} \right) \text{ and } \frac{\beta_n}{\sqrt{n}} \sum_{i=1}^{n} \left(\frac{f(\mathbf{X}_i)}{g(\mathbf{X}_i)} - 1\right) - \frac{\beta_n^2}{2} \mathbb{E} \left\{ \frac{f(\mathbf{X}_1)}{g(\mathbf{X}_1)} - 1 \right\}^2$$

converges to a bivariate normal distribution with

$$\begin{aligned} \max \mu &= \begin{pmatrix} 0\\ -\frac{\beta^2}{2} \mathbb{E}\left\{\frac{f(\mathbf{X}_1)}{g(\mathbf{X}_1)} - 1\right\}^2 \end{pmatrix} \text{ and variance } \Sigma &= \begin{pmatrix} \operatorname{Var}\left(h(\mathbf{X}_1, \mathbf{X}_1')\right) & \tau\\ \tau & -\frac{\beta^2}{2} \mathbb{E}\left\{\frac{f(\mathbf{X}_1)}{g(\mathbf{X}_1)} - 1\right\}^2 \end{pmatrix} \end{pmatrix}, \text{ where} \\ \tau &= \mathbb{E}\left\{\left\{h(\mathbf{X}_1, \mathbf{X}_1') - \mathbb{E}\left\{h(\mathbf{X}_1, \mathbf{X}_1')\right\}\right\}\beta\left\{\frac{f(\mathbf{X}_1)}{g(\mathbf{X}_1)} - 1\right\}\right\} \\ &= \beta \int \left\{h(\mathbf{x}, \|\mathbf{x}\| \mathbf{u}) - \mathbb{E}\left\{h(\mathbf{X}_1, \mathbf{X}_1')\right\}\right\}\left(f(\mathbf{x}) - g(\mathbf{x})\right) \mathrm{d}\mathbf{x} \mathrm{d}\mu(\mathbf{u}), \end{aligned}$$

 μ being the distribution Unif(S^{d-1}). Now using Le Cam's third lemma (see. van der Vaart, 1998), as n diverges to infinity, under $F_{1-\beta_n/\sqrt{n}}$, we have

$$\sqrt{n} \left(\frac{1}{n} \sum_{i=1}^{n} h(\mathbf{X}_{i}, \mathbf{X}_{i}') - \mathbb{E} \left\{ h(\mathbf{X}_{1}, \mathbf{X}_{1}') \right\} \right) \xrightarrow{D} N \left(\tau, \operatorname{Var} \left(h(\mathbf{X}_{1}, \mathbf{X}_{1}') \right) \right).$$

Now using similar arguments as in Theorem 1 in p.79 from Lee (1990) and contiguity arguments, we get

$$n\hat{\zeta}_n \xrightarrow{D} \sum_{i=1}^{\infty} \lambda_i \left(\left(Z_i + \beta \mathbb{E}_F \{ f_i(\mathbf{X}_1, \mathbf{X}_1') \} \right) \right)^2 - 1 \right)$$

under $F_{1-\beta_n/\sqrt{n}}$, where $\{Z_i\}$ is a sequence of i.i.d. normal random variables. This completes the proof.

Proof of Theorem 3.6. Let $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_n \stackrel{i.i.d.}{\sim} G$ and $\mathbf{U}_1, \mathbf{U}_2, \ldots, \mathbf{U}_n \stackrel{i.i.d.}{\sim}$ Unif (\mathcal{S}^{d-1}) be independent and $\mathbf{X}'_i = \|\mathbf{X}_i\|\mathbf{U}_i$ for $i = 1, 2, \ldots, n$. Also let $\pi_1, \pi_2, \ldots, \pi_n$ be i.i.d random variables from the Bernoulli(0.5) distribution. Then the resampled test statistic $\hat{\zeta}_n(\pi)$ can be written as,

$$\frac{1}{n(n-1)}\sum_{1\leq i\neq j\leq n}g\big((\mathbf{Y}_i,\mathbf{Y}_i'),(\mathbf{Y}_j,\mathbf{Y}_j')\big),$$

where $\mathbf{Y}_i = \pi_i \mathbf{X}_i + (1 - \pi_i) \mathbf{X}'_i$ and $\mathbf{Y}'_i = (1 - \pi_i) \mathbf{X}_i + \pi_i \mathbf{X}'_i$. Since, under any fixed alternative F, $(\mathbf{Y}_i, \mathbf{Y}'_i)$ is a sequence of i.i.d. random vectors, we can apply Theorem 1 from Lee (1990) p.79 to find the limiting distribution of $\hat{\zeta}_n(\pi)$. Note that $\mathbf{Y}_i \stackrel{D}{=} \mathbf{Y}'_i$ and hence $g_1(\mathbf{y}, \mathbf{y}') = \mathbb{E}\{g((\mathbf{y}, \mathbf{y}'), (\mathbf{Y}_1, \mathbf{Y}'_1))\} = 0$. Therefore, $\hat{\zeta}_n(\pi)$ is a U-statistic with a first order degenerate kernel function and we get

$$n\hat{\zeta}_n(\pi) \stackrel{D}{\to} \sum_{i=1}^{\infty} \lambda_i (Z_i^2 - 1),$$

where $\{\lambda_i\}$ is a square integrable sequence and $\{Z_i\}$ is a sequence of i.i.d. standard normal random variables.

Now suppose that $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_n \stackrel{i.i.d.}{\sim} F_{1-\beta_n n^{-1/2}} = (1 - \frac{\beta}{\sqrt{n}})G + \frac{\beta}{\sqrt{n}}F$ where $\beta_n \to \beta$, $\zeta(G) = 0$ and $\zeta(F) > 0$. Then to find the limiting distribution of $n\zeta_n(\pi)$ we need to find the joint limiting distribution of

$$\frac{1}{\sqrt{n}}\sum_{i=1}^{n}\varphi_i\big((\mathbf{Y}_i,\mathbf{Y}_i')\big) \text{ and } \text{ and } \frac{\beta_n}{\sqrt{n}}\sum_{i=1}^{n}\big(\frac{f(\mathbf{X}_i)}{g(\mathbf{X}_i)}-1\big) - \frac{\beta_n^2}{2}\mathbb{E}\big\{\frac{f(\mathbf{X}_1)}{g(\mathbf{X}_1)}-1\big\}^2$$

where $\{\varphi_i\}$ are the solutions of the integral equation

$$\int g((\mathbf{x}_1, \mathbf{x}_1'), (\mathbf{y}_1, \mathbf{y}_1'))\varphi_i((\mathbf{y}_1, \mathbf{y}_1')) \mathrm{d}\nu\big((\mathbf{y}_1, \mathbf{y}_1')\big) = \lambda\varphi_i\big((\mathbf{x}_1, \mathbf{x}_1')\big)$$
(8)

where ν is the joint distribution of $(\mathbf{Y}_1, \mathbf{Y}'_1)$ when $\mathbf{X}_1 \sim G$. By the bivariate central limit theorem, we can see that the joint distribution of these two variables converges to a bivariate normal distribution with mean and variance given by

$$\mu = \begin{pmatrix} 0 \\ -\frac{\beta^2}{2} \mathbb{E}\left\{\frac{f(\mathbf{X}_1)}{g(\mathbf{X}_1)} - 1\right\}^2 \end{pmatrix} \text{ and } \Sigma = \begin{pmatrix} \operatorname{Var}\left(\varphi_i(\mathbf{Y}_1, \mathbf{Y}_1')\right) & \tau \\ \tau & -\frac{\beta^2}{2} \mathbb{E}\left\{\frac{f(\mathbf{X}_1)}{g(\mathbf{X}_1)} - 1\right\}^2 \end{pmatrix}, \text{ where}$$
$$\tau = \mathbb{E}\left\{\left\{\varphi_i(\mathbf{Y}_1, \mathbf{Y}_1')\right\}\beta\left\{\frac{f(\mathbf{X}_1)}{g(\mathbf{X}_1)} - 1\right\}\right\} \\ = \beta \mathbb{E}_F\{\varphi_i(\mathbf{Y}_1, \mathbf{Y}_1')\} = \beta \left\{\frac{1}{2} \mathbb{E}\{\varphi_i((\mathbf{X}_1, \mathbf{X}_1'))\} + \frac{1}{2} \mathbb{E}\{\varphi_i((\mathbf{X}_1', \mathbf{X}_1))\}\right\}.$$

Now note that $g((\mathbf{x}_1, \mathbf{x}'_1), (\mathbf{y}_1, \mathbf{y}'_1)) = -g((\mathbf{x}'_1, \mathbf{x}_1), (\mathbf{y}_1, \mathbf{y}'_1))$. Then using (8) we get $\varphi_i((\mathbf{x}_1, \mathbf{x}'_1)) = -\varphi_i((\mathbf{x}'_1, \mathbf{x}_1))$. Using this we get $\tau = 0$. So, using Le Cam's third lemma (see. van der Vaart, 1998), as *n* diverges to infinity, under $F_{1-\beta_n/\sqrt{n}}$, we have

$$\frac{1}{\sqrt{n}}\sum_{i=1}^{n}\varphi_i\big((\mathbf{Y}_i,\mathbf{Y}_i')\big) \xrightarrow{D} N\Big(0,\operatorname{Var}\big(\varphi_i\big((\mathbf{Y}_1,\mathbf{Y}_1')\big)\big)\Big) = N(0,1).$$

Using similar arguments as in Theorem 1 in p.79 from Lee (1990) and contiguity arguments we get

$$n\hat{\zeta}_n(\pi) \xrightarrow{D} \sum_{i=1}^{\infty} \lambda_i (Z_i^2 - 1),$$

where $\{\lambda_i\}$ is a square-integrable sequence and $\{Z_i\}$ is a sequence of i.i.d. standard normal random variables.

References

- Ahn, J., Marron, J., Muller, K. M., and Chi, Y.-Y. (2007). The high-dimension, low-sample-size geometric representation holds under mild conditions. *Biometrika*, 94(3):760–766.
- Albisetti, I., Balabdaoui, F., and Holzmann, H. (2020). Testing for spherical and elliptical symmetry. J. Multivariate Anal., 180:104667.
- Baringhaus, L. (1991). Testing for spherical symmetry of a multivariate distribution. Ann. Statist., 19(2):899–917.
- Chaudhuri, P. (1996). On a geometric notion of quantiles for multivariate data. J. Amer. Statist. Assoc., 91(434):862–872.
- Chaudhuri, P. and Sengupta, D. (1993). Sign tests in multidimension: inference based on the geometry of the data cloud. J. Amer. Statist. Assoc., 88(424):1363–1370.
- Chmielewski, M. A. (1981). Elliptically symmetric distributions: a review and bibliography. *Internat. Statist. Rev.*, 49(1):67–74.
- Diks, C. and Tong, H. (1999). A test for symmetries of multivariate probability distributions. *Biometrika*, 86(3):605–614.
- Ding, X. (2020). Some sphericity tests for high dimensional data based on ratio of the traces of sample covariance matrices. *Statist. Probab. Lett.*, 156:108613.
- Dutta, S., Ghosh, A. K., and Chaudhuri, P. (2011). Some intriguing properties of Tukey's half-space depth. Bernoulli, 17(4):1420–1434.
- Fang, K. T., Kotz, S., and Ng, K. W. (1990). Symmetric Multivariate and Related Distributions, volume 36 of Monographs on Statistics and Applied Probability. Chapman and Hall, London.
- Fang, K. T., Zhu, L. X., and Bentler, P. M. (1993). A necessary test of goodness of fit for sphericity. J. Multivariate Anal., 45(1):34–55.
- Feng, L. and Liu, B. (2017). High-dimensional rank tests for sphericity. J. Multivariate Anal., 155:217–233.
- Fourdrinier, D., Strawderman, W. E., and Wells, M. T. (2018). *Shrinkage Estimation*. Springer Series in Statistics. Springer, Cham.
- Ghosh, A. K. and Chaudhuri, P. (2005). On maximum depth and related classifiers. Scand. J. Statist., 32(2):327–350.
- Gretton, A., Fukumizu, K., Teo, C., Song, L., Schölkopf, B., and Smola, A. (2007). A kernel statistical test of independence. In *Adv. Neural Inf. Process. Syst.*, volume 20. Curran Associates, Inc.
- Gupta, A. and Song, D. (1997). Lp-norm spherical distribution. J. Stat. Plan. Inference, 60(2):241-260.
- Hall, P., Marron, J. S., and Neeman, A. (2005). Geometric representation of high dimension, low sample size data. J. R. Stat. Soc. Ser. B Stat. Methodol., 67(3):427–444.
- Heck, D., Knapp, J., Capdevielle, J. N., Schatz, G., and Thouw, T. (1998). CORSIKA: A monte carlo code to simulate extensive air showers. Technical report. 51.02.03; LK 01; Wissenschaftliche Berichte, FZKA-6019.
- Henze, N., Hlávka, Z., and Meintanis, S. G. (2014). Testing for spherical symmetry via the empirical characteristic function. *Statistics*, 48(6):1282–1296.
- Huang, Z. and Sen, B. (2023). Multivariate symmetry: Distribution-free testing via optimal transport. arXiv preprint arXiv:2305.01839.

- John, S. (1972). The distribution of a statistic used for testing sphericity of normal distributions. *Biometrika*, 59(1):169–173.
- Johnstone, I. M. (2001). On the distribution of the largest eigenvalue in principal components analysis. Ann. Statist., 29(2):295–327.
- Jörnsten, R. (2004). Clustering and classification based on the 11 data depth. J. Multivariate Anal., 90(1):67–89.
- Jung, S. and Marron, J. S. (2009). PCA consistency in high dimension, low sample size context. Ann. Statist., 37(6B):4104–4130.
- Koltchinskii, V. I. (1997). M-estimation, convexity and quantiles. Ann. Statist., 25(2):435 477.
- Koltchinskii, V. I. and Li, L. (1998). Testing for spherical symmetry of a multivariate distribution. J. Multivariate Anal., 65(2):228–244.
- Lee, A. J. (1990). U-Statistics: Theory and Practice, volume 110 of Statistics: Textbooks and Monographs. Marcel Dekker, Inc., New York.
- Lehmann, E. L. (2012). On the history and use of some standard statistical models. Selected Works of EL Lehmann, pages 1019–1031.
- Liang, J., Fang, K.-T., and Hickernell, F. J. (2008). Some necessary uniform tests for spherical symmetry. Ann. Inst. Statist. Math., 60(3):679–696.
- Massart, P. (1990). The tight constant in the Dvoretzky-Kiefer-Wolfowitz inequality. Ann. Probab., 18(3):1269–1283.
- Randles, R. H. (1989). A distribution-free multivariate sign test based on interdirections. J. Amer. Statist. Assoc., 84(408):1045–1050.
- Rousseeuw, P. (1985). Multivariate estimation with high breakdown point. In *Mathematical Statistics and Applications*, pages 283–297. Reidel, Dordrecht.
- Rousseeuw, P. J. and Driessen, K. V. (1999). A Fast Algorithm for the Minimum Covariance Determinant Estimator. *Technometrics*, 41(3):212–223.
- Smith, P. J. (1977). A nonparametric test for bivariate circular symmetry based on the empirical CDF. Comm. Statist.—Theory Methods, 6(3):209–220.
- Székely, G. J. and Rizzo, M. L. (2004). Testing for equal distributions in high dimension. *InterStat*, 5(16.10):1249–1272.
- Tsybakov, A. B. (2009). Introduction to Nonparametric Estimation. Springer, New York.
- Tukey, J. W. (1975). Mathematics and the picturing of data. In Proceedings of the International Congress of Mathematicians (Vancouver, B.C., 1974), Vol. 2, pages 523–531. Canad. Math. Congr., Montreal, QC.
- Tyler, D. E. (1987). A distribution-free *M*-estimator of multivariate scatter. Ann. Statist., 15(1):234–251.
- van der Vaart, A. W. (1998). Asymptotic statistics, volume 3 of Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, Cambridge.
- Wainwright, M. J. (2019). High-dimensional statistics: A non-asymptotic viewpoint, volume 48 of Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, Cambridge.
- Zou, C., Peng, L., Feng, L., and Wang, Z. (2014). Multivariate sign-based high-dimensional tests for sphericity. *Biometrika*, 101(1):229–236.

This figure "Example_3.png" is available in "png" format from:

http://arxiv.org/ps/2403.12491v1