

Treewidth of generalized Hamming graph, bipartite Kneser graph and generalized Petersen graph

Yichen Wang^{*1}, Mengyu Cao^{†2}, Zequn Lv^{‡1}, and Mei Lu^{§1}

¹Department of Mathematical Sciences, Tsinghua University, Beijing 100084, China

²Institute for Mathematical Sciences, Renmin University of China, Beijing 100086, China

Abstract

Let t, q and n be positive integers. Write $[q] = \{1, 2, \dots, q\}$. The generalized Hamming graph $H(t, q, n)$ is the graph whose vertex set is the cartesian product of n copies of $[q]$ ($q \geq 2$), where two vertices are adjacent if their Hamming distance is at most t . In particular, $H(1, q, n)$ is the well-known Hamming graph and $H(1, 2, n)$ is the hypercube. In 2006, Chandran and Kavitha described the asymptotic value of $tw(H(1, q, n))$, where $tw(G)$ denotes the treewidth of G . In this paper, we give the exact pathwidth of $H(t, 2, n)$ and show that $tw(H(t, q, n)) = \Theta(tq^n/\sqrt{n})$ when n goes to infinity. Based on those results, we show that the treewidth of bipartite Kneser graph $BK(n, k)$ is $\binom{n}{k} - 1$ when n is sufficient large relative to k and the bounds of $tw(BK(2k+1, k))$ are given. Moreover, we present the bounds of the treewidth of generalized Petersen graph.

Keywords: treewidth; pathwidth; generalized Hamming graph; bipartite Kneser graph; generalized Petersen graph

1 Introduction

Treewidth is a well-studied parameter in graph theory. Many NP-complete problem can be solved in polynomial time on graphs of bounded treewidth [31, 6, 10]. Besides, treewidth is also useful in structural graph theory. For example, Robertson and Seymour used it to prove the Graph Minor Theorem [25, 26, 27]. In the past decades, there are lots of literatures investigating the treewidth of certain graphs (see for example, [14, 15, 19, 22, 30]). However, it is difficult to estimate the treewidth even asymptotically.

*E-mail: wangyich22@mails.tsinghua.edu.cn

†E-mail: myuca@ruc.edu.cn

‡E-mail: lvzq19@mails.tsinghua.edu.cn

§E-mail: lumei@tsinghua.edu.cn

Throughout this paper, graphs are finite, simple and undirected. Let $V(G)$ and $E(G)$ denote the vertex set and edge set of G , respectively. The degree of a vertex $v \in V(G)$ in G is denoted by $d_G(v)$, and we ignore the subscript G in case of no ambiguity. The maximum and minimum degree of G are denoted as $\Delta(G)$ and $\delta(G)$, respectively. Write $[n] = \{1, 2, \dots, n\}$ and denote by $\binom{[n]}{k}$ the family of all k -subsets of $[n]$. The treewidth and pathwidth of graph G are denoted by $tw(G)$ and $pw(G)$, respectively.

The *Hamming graph* $H(q, n)$ is the graph on q^n vertices, which correspond to all n -vectors whose components are from $[q]$ ($q \geq 2$). When $q = 2$, it is more commonly to treat each vertex of $H(2, n)$ as a binary n -vector, that is, a n -vector whose components are from $\{0, 1\}$. Two of the vertices in $H(q, n)$ are adjacent only when they differ in just one coordinate. The *Hypercube graph* is a special case of Hamming graph when $q = 2$ which is well-studied in parallel computing, coding theory and many other areas[1, 16, 17]. The Hamming distance of two n -vectors is the number of coordinates that one differs from the other. Two vertices in a Hamming graph are adjacent if and only if their Hamming distance is no more than 1. If two vertices in $H(q, n)$ is adjacent if and only if their Hamming distance is no more than t , then we denote the *generalized Hamming graph* as $H(t, q, n)$. Then $H(1, q, n) = H(q, n)$ and $H(1, 2, n)$ is the hypercube graph. We call $H(t, 2, n)$ the *generalized hypercube graph* and we will give the exact value of its pathwidth and bandwidth as Theorem 1.1.

Theorem 1.1 *We have*

$$pw(H(t, 2, n)) = bw(H(t, 2, n)) = \sum_{k=\lfloor (n-t)/2 \rfloor}^{\lfloor (n-t)/2 \rfloor + t - 1} \binom{n}{k} + \sum_{a=0}^{\lfloor (n-t-1)/2 \rfloor} \left(\binom{t+2a}{t+a-1} - \binom{t+2a}{a-1} \right). \quad (1)$$

The treewidth of $H(q, n)$ is asymptotically $\Theta(q^n/\sqrt{n})$ from [8]. In this paper, we generalize this result to generalized Hamming graph $H(t, q, n)$ as Theorem 1.2. The previous result about $tw(H(q, n))$ can be derived from our result by letting $t = 1$.

Theorem 1.2 *Let t and q be positive integers. Then $tw(H(t, q, n)) = \Theta(tq^n/\sqrt{n})$, when n goes to infinity.*

Kneser graph $K(n, k)$ is the graph whose vertex set is $\binom{[n]}{k}$ and where two vertices are adjacent if and only if the two corresponding sets are disjoint. The treewidth of Kneser graph is studied by Harvey in 2014 [14]. *Generalized Kneser graph* and *q -Kneser graph* are two derived class from Kneser graph and their treewidth can also be exactly described when n is large enough [7, 23]. *Bipartite Kneser graph* is also an important variant of Kneser graph whose vertex set is $\binom{[n]}{k}$ and $\binom{[n]}{n-k}$ denoted by $BK(n, k)$ where $2k \leq n$. A k -subset A and a $(n-k)$ -subset B in bipartite Kneser graph is adjacent if $A \subseteq B$. We will give the exact value of $tw(BK(n, k))$ when n is sufficient large relative to k as Theorem 1.3. When $n = 2k + 1$, we give the bounds of the treewidth of bipartite Kneser graph as Theorem 1.4.

Theorem 1.3 *If $3\binom{n-k}{k} \geq 2\binom{n}{k}$ and $k \geq 2$, then $tw(BK(n, k)) = \binom{n}{k} - 1$.*

Theorem 1.4 *As k goes to infinity,*

$$\Omega\left(\frac{1}{k}\binom{2k+1}{k}\right) = tw(BK(2k+1, k)) \leq tw(J(2k+1, k)) = O\left(\binom{2k+1}{k}\right).$$

Petersen graph is a well-studied Kneser graph $K(5, 2)$. *Generalized Petersen graph* is an extension of Petersen graph denoted by $G_{n,k}$ whose vertex set and edge set are

$$\begin{aligned} V(G_{n,k}) &= \{v_1, \dots, v_n, u_1, \dots, u_n\}, \\ E(G_{n,k}) &= \{v_i u_i\} \cup \{v_i v_{i+1}\} \cup \{u_i u_{i+k}\}, i = 1, 2, \dots, n, \end{aligned}$$

where subscripts are to be read modulo n and $k < n/2$. We will show the treewidth of $G_{n,k}$ when n is sufficient large relative to k as Theorem 1.5

Theorem 1.5 *Let n and k be positive integers satisfying that $n \geq 8(2k+2)^2$. Then $2k+1 \leq tw(G_{n,k}) \leq pw(G_{n,k}) \leq 2k+2$.*

The rest of this paper is organized as follows. In Section 3, we give a proof of the treewidth of generalized Hamming graph. In Section 4, we study the treewidth of bipartite Kneser graph and Johnson graph. In Section 5, we study the treewidth of generalized Petersen graph.

2 Preliminaries and definitions

In this section, we give definitions involving in treewidth, pathwidth and bandwidth of a graph $G(V, E)$.

Definition 2.1 *A tree-decomposition of a graph $G(V, E)$ is a pair (X, T) , where $T(I, F)$ is a tree with vertex set I and edge set F , and $X = \{X_i \mid i \in I\}$ is a family of subsets of V , one for each node of T , such that:*

- $\bigcup_{i \in I} X_i = V$.
- For each edge $uv \in E$, there exists an $i \in I$ such that $u, v \in X_i$.
- For all $i, j, k \in I$, if j is on the path from i to k in T , then $X_i \cap X_k \subset X_j$.

The width of a tree-decomposition (X, T) is $\max_{i \in I} |X_i| - 1$. The treewidth of a graph G is the minimum treewidth over all possible tree-decompositions of G and denoted by $tw(G)$. The problem of deciding whether a graph has tree decomposition of treewidth at most k is NP-complete [2]. However, there is an exact algorithm finding treewidth of given graph G when taking $tw(G)$ as a constant [4, 20]. A path decomposition of G is a tree decomposition (X, T)

in which T is required to be a path. The pathwidth of G is defined to be the minimum width over all path decompositions of G and is denoted by $pw(G)$.

A bijection $\phi : V \rightarrow \{1, 2, \dots, n\}$ is called an ordering of the vertices of G (in short, an ordering of G). Then for any edge $e = \{u, v\} \in E$, let $\Delta(e, \phi) = |\phi(u) - \phi(v)|$.

Definition 2.2 *A bandwidth of a graph $G(V, E)$, denoted by $bw(G)$, is the minimum over all possible orderings ϕ of V of the maximum value of $\Delta(e, \phi)$ over all edges $e \in E$. That is,*

$$bw(G) = \min_{\phi} \max_{e \in E} \Delta(e, \phi).$$

There are important inequalities between treewidth, pathwidth and bandwidth as following.

Proposition 2.3 ([5]) *For any graph G ,*

$$tw(G) \leq pw(G) \leq bw(G). \tag{2}$$

Let $X \subseteq V(G)$ be a subset of vertices and $G[X]$ be the subgraph induced in G by X . Define $G - X = G[V(G) - X]$. Given $p \in (0, 1)$, define the p -separator of G to be a subset $X \subseteq V(G)$ such that no component of $G - X$ contains more than $p|V(G) - X|$ vertices. Proposition 2.4 describes the relationship between treewidth and separator.

Proposition 2.4 ([27]) *For any graph G , there exists a $1/2$ -separator of G with at most $tw(G) + 1$ vertices.*

Corollary 2.5 is directly from Proposition 2.4.

Corollary 2.5 *For any graph G , there exists a separator X of G with at most $tw(G) + 1$ vertices. And there exists a partition of $V(G) - X$ into sets A and B such that*

$$|V(G) - X|/3 \leq |A|, |B| \leq 2|V(G) - X|/3.$$

Proposition 2.4 and Corollary 2.5 are useful tools to estimate the lower bound of treewidth.

3 Treewidth of generalized Hamming graph

3.1 Bandwidth of Hamming graph $H(t, 2, n)$

The pathwidth and bandwidth of hypercubes can be exactly calculated as following.

Proposition 3.1 ([8]) *We have*

$$pw(H(1, 2, n)) = bw(H(1, 2, n)) = \sum_{m=0}^{n-1} \binom{m}{\lfloor m/2 \rfloor}. \quad (3)$$

In this section, we intend to prove Theorem 1.1. When $t = 1$, Eq 1 is exactly the same as Eq 3.

To prove Theorem 1.1, we need some preparation. In [13], Harper showed that if an ordering φ of $G(V, E)$ is in Hales order (i.e., a Hales numbering), then $\max_{e \in E} \Delta(e, \varphi)$ takes minimum over all numbering, that is, $bw(G) = \min_{\phi} \max_{e \in E} \Delta(e, \phi) = \max_{e \in E} \Delta(e, \varphi)$. Harper [13] also give a sample of Hales numbering of $H(t, 2, n)$ ¹. From [29], we can build such Hales numbering φ in the following way.

Define an $\binom{n}{k} \times n$ matrix $A_k^{(n)}$ as following. Let $A_n^{(n)}$ be $\underbrace{(1, 1, \dots, 1)}_{n \text{ factors}}$ and let $A_0^{(n)}$ be $\underbrace{(0, 0, \dots, 0)}_{n \text{ factors}}$ for any positive integer n . When $0 < k < n$, $A_k^{(n)}$ is defined recursively as Eq 4.

$$A_k^{(n)} \triangleq \begin{bmatrix} A_{k-1}^{(n-1)} & \mathbf{1} \\ A_k^{(n-1)} & \mathbf{0} \end{bmatrix}, \quad (4)$$

where $\mathbf{1}$ (resp. $\mathbf{0}$) is an all one (resp. all zero) $\binom{n-1}{k-1}$ (resp. $\binom{n-1}{k}$) column vector. Clearly, the rows of $A_k^{(n)}$ enumerates all binary vectors of length n whose number of ‘1’ is k . Define $S^{(n)}$ as following:

$$S^{(n)} \triangleq \begin{bmatrix} A_0^{(n)} \\ A_1^{(n)} \\ \vdots \\ A_n^{(n)} \end{bmatrix}. \quad (5)$$

Then $S^{(n)}$ is an $2^n \times n$ matrix. Each row of $S^{(n)}$ exactly corresponds to a n -vector (a vertices of $H(t, 2, n)$) and vice versa. Let $\eta^{(n)}$ be the ordering of $H(t, 2, n)$ defined by row order of $S^{(n)}$, that is, if $v \in V(H(t, 2, n))$ corresponds to the i -th row of $S^{(n)}$, then let $\eta^{(n)}(v) = i$.

$\eta^{(n)}$ is a Hales numbering of $H(t, 2, n)$ (see [29] and [13]) and, hence, we have the following Proposition 3.2.

Proposition 3.2 $bw(H(t, 2, n)) = \max_{e \in E} \Delta(e, \eta^{(n)})$.

In order to calculate $bw(H(t, 2, n))$, we need some more definitions.

¹It is not explicitly stated in [13]. Harper first gave a Hales numbering of hypercube and, then claimed that the numbering is also in Hales order for the distance generalized graph (that is, $H(t, 2, n)$) in comments.




Figure 1: Bandwidth of a square matrix. Figure 2: Manhattan radius and the imaginary anchor.

Definition 3.3 Given a graph $G(V, E)$, for two vertex subsets $V_1 \subseteq V$ and $V_2 \subseteq V$ numbered by ordering η_1 and η_2 respectively, the adjacency matrix of V_1 and V_2 is defined to be a $|V_1| \times |V_2|$ matrix M such that for any $u \in V_1$ and $v \in V_2$,

$$M(\eta_1(u), \eta_2(v)) = \begin{cases} 1 & \text{if } \{u, v\} \in E, \\ 0 & \text{otherwise.} \end{cases}$$

The adjacent matrix of $G(V, E)$ with ordering η is the adjacent matrix of V and V ordered by η as defined above.

Definition 3.4 The bandwidth of an $s \times s$ non-zero square matrix M denoted by $bw(M)$ is the maximum absolute value of the difference between the row and column indices of a nonzero element of that matrix, i.e.,

$$bw(M) = \max_{1 \leq i \leq s, 1 \leq j \leq s} \{|i - j| \mid M(i, j) \neq 0\}, \quad (6)$$

which is exactly the maximum Manhattan distance from a nonzero element to the main diagonal of the matrix (see Figure 1).

Remark 3.5 Given an ordering η of a graph G , $\max_{e \in E} \Delta(e, \eta)$ is equal to the bandwidth of the adjacency matrix of G ordered by η .

Remark 3.5 is directly from definitions. Let $M^{(t,n)}$ be the adjacency matrix of $H(t, 2, n)$ ordered by $\eta^{(n)}$. From Remark 3.5 and Proposition 3.2, we have Lemma 3.6.

Lemma 3.6 $bw(H(t, 2, n)) = bw(M^{(t,n)})$.

Now our aim is to give exact value of $bw(M^{(t,n)})$. To achieve this, we need to define the Manhattan radius of matrix.

Definition 3.7 For an $s \times t$ non-zero matrix M , its Manhattan radius $r(M)$ is defined by

$$r(M) = \max_{1 \leq i \leq s, 1 \leq j \leq t} \{s - i + j \mid M(i, j) \neq 0\}, \quad (7)$$

which is exactly the maximum Manhattan distance from a nonzero element of M to the position immediately to the left of the bottom-left corner of M (an imaginary element $M(s, 0)$). The imaginary element is called the anchor of M (see Figure 2). For the convenience of the proof, let $r(M) = -\infty$ if M is a zero matrix or an empty matrix.

Note that $r(M) > 0$ if and only if M is non-zero and non-empty. If M is an $s \times s$ symmetric matrix, then by definition,

$$r(M) = bw(M) + s. \quad (8)$$

Let $V_k^{(t,n)} \subseteq V(H(t, 2, n))$ be a vertex set containing vertices of $H(t, 2, n)$ whose corresponding n -vector has exactly k ones, where $0 \leq k \leq n$. Then $\{V_0^{(t,n)}, V_1^{(t,n)}, \dots, V_n^{(t,n)}\}$ forms a partition of $V(H(t, 2, n))$. Recall that each row of $A_k^{(n)}$ correspond to a vertex of $V_k^{(t,n)}$ and vice versa. Let $\eta_k^{(n)} = \eta^{(n)}|_{V_k^{(t,n)}}$. Let $M_{k,k'}^{(t,n)}$ be the adjacent matrix of $V_k^{(t,n)}$ and $V_{k'}^{(t,n)}$ ordered by $\eta_k^{(n)}$ and $\eta_{k'}^{(n)}$ respectively.

For the convenience of proof, let $M_{k,k'}^{(t,n)}$ be empty matrix if either k or k' is larger than n or less than zero. It is easy to verify that $r(M_{k,k'}^{(t,n)}) > 0$ if and only if $|k - k'| \leq t$ and $0 \leq k, k' \leq n$. Then from definition, we have

$$M^{(t,n)} = \begin{bmatrix} M_{0,0}^{(t,n)} & M_{0,1}^{(t,n)} & M_{0,2}^{(t,n)} & \cdots & M_{0,n-1}^{(t,n)} & M_{0,n}^{(t,n)} \\ M_{1,0}^{(t,n)} & M_{1,1}^{(t,n)} & M_{1,2}^{(t,n)} & \cdots & M_{1,n-1}^{(t,n)} & M_{1,n}^{(t,n)} \\ M_{2,0}^{(t,n)} & M_{2,1}^{(t,n)} & M_{2,2}^{(t,n)} & \cdots & M_{2,n-1}^{(t,n)} & M_{2,n}^{(t,n)} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ M_{n-1,0}^{(t,n)} & M_{n-1,1}^{(t,n)} & M_{n-1,2}^{(t,n)} & \cdots & M_{n-1,n-1}^{(t,n)} & M_{n-1,n}^{(t,n)} \\ M_{n,0}^{(t,n)} & M_{n,1}^{(t,n)} & M_{n,2}^{(t,n)} & \cdots & M_{n,n-1}^{(t,n)} & M_{n,n}^{(t,n)} \end{bmatrix}. \quad (9)$$

Note that $M^{(t,n)}$ is an $2^n \times 2^n$ matrix. If $t \geq n$, then $M^{(t,n)}$ is an all one matrix except for the elements in main diagonal. Hence $bw(M^{(t,n)}) = 2^n - 1$. In the following, we only consider the situation when $t < n$. Since the bandwidth of $M^{(t,n)}$ is the maximum Manhattan distance from a non-zero element to the main diagonal and $M^{(t,n)}$ is symmetric, we only need to consider non-zero element from submatrices in Eq 9 on the diagonal or on the top right of the diagonal, that is, non-zero elements from $M_{k,k'}^{(t,n)}$ with $k' \geq k$. Since $M_{k,k'}^{(t,n)}$ is all zero when $|k - k'| > t$, we only need to consider non-zero element from $M_{k,k'}^{(t,n)}$ with $k \leq k' \leq k + t$. By the definition of bandwidth and Manhattan radius, we have Proposition 3.8.

Proposition 3.8 *We have*

$$bw(M^{(t,n)}) = \max_{\substack{k=0, \dots, n-1, \\ 1 \leq p \leq t, \\ p+k \leq n}} \left\{ \sum_{q=1}^{p-1} \binom{n}{k+q} + r(M_{k,k+p}^{(t,n)}), bw(M_{k,k}^{(t,n)}) \right\}. \quad (10)$$

Note that $M_{k,k}^{(t,n)}$ is a symmetric matrix. By Eq 8, we have $bw(M_{k,k}^{(t,n)}) = r(M_{k,k}^{(t,n)}) + \binom{n}{k}$. If we can calculate all $r(M_{k,k'}^{(t,n)})$ with $k \leq k' \leq k+t$, then we can calculate $bw(M^{(t,n)})$ via Eq 10 and, hence, obtain $bw(H(t, 2, n))$ by Lemma 3.6. Now, our aim is to calculate all $r(M_{k,k'}^{(t,n)})$ with $k \leq k' \leq k+t$.

From Eq 4, we have that if t, n, k, p are non-negative intergers satisfying that $k+p \leq n$ and $n \geq 1$, then

$$M_{k,k+p}^{(t,n)} = \begin{pmatrix} M_{k-1,k+p-1}^{(t,n-1)} & M_{k-1,k+p}^{(t-1,n-1)} \\ M_{k,k+p-1}^{(t-1,n-1)} & M_{k,k+p}^{(t,n-1)} \end{pmatrix}. \quad (11)$$

Let $r_{t,n,k,p}^{(1)} = r(M_{k-1,k+p-1}^{(t,n-1)}) + \binom{n-1}{k}$, $r_{t,n,k,p}^{(2)} = r(M_{k-1,k+p}^{(t-1,n-1)}) + \binom{n-1}{k} + \binom{n-1}{k+p-1}$, $r_{t,n,k,p}^{(3)} = r(M_{k,k+p-1}^{(t,n-1)})$ and $r_{t,n,k,p}^{(4)} = r(M_{k,k+p}^{(t,n-1)}) + \binom{n-1}{k+p-1}$. From the definition of $r(M_{k,k+p}^{(t,n)})$ and Eq 11, we have Proposition 3.9.

Proposition 3.9 *For non-negative integers t, n, k, p satisfying that $k+p \leq n$ and $n \geq 1$, we have*

$$r(M_{k,k+p}^{(t,n)}) = \max \left\{ r_{t,n,k,p}^{(1)}, r_{t,n,k,p}^{(2)}, r_{t,n,k,p}^{(3)}, r_{t,n,k,p}^{(4)} \right\}. \quad (12)$$

Considering $M_{k_1,k_2}^{(t,n)}$ with $k_1 \leq k_2 \leq k_1+t$, it is easy to verify that if the parity of $k_2 - k_1$ and t is different, then $M_{k_1,k_2}^{(t,n)} = M_{k_1,k_2}^{(t-1,n)}$. Hence, we only need to calculate all $M_{k_1,k_2}^{(t,n)}$ with $k_1 \leq k_2 \leq k_1+t$ and $k_2 - k_1 \equiv t \pmod{2}$. In this case, assume $t - (k_2 - k_1) = 2s$. Then $M_{k,k+t-2s}^{(t,n)}$ is non-zero. If $t \geq n-1$, then $M_{k,k+t-2s}^{(t,n)}$ is an all-one matrix (except the main diagonal when $t = 2s$) and, thus, $r(M_{k,k+t-2s}^{(t,n)}) = \binom{n}{k} + \binom{n}{k+t-2s} - 1$. When $1 \leq t < n-1$, we let

$$\begin{aligned} A_{t,n,k,s}^{(1)} &= \sum_{a=0}^{k-s-1} \left(\binom{t-s+2a}{t-s+a-1} - \binom{t-s+2a}{a-1} \right), \\ A_{t,n,k,s}^{(2)} &= \sum_{m=t-3s+1+2k}^{n-s} \left(\binom{m-1}{k+t-2s-1} - \binom{m-1}{k-s-1} \right), \\ B_{t,n,k,s} &= \sum_{a=0}^{n-t-k+s-1} \left(\binom{t-s+2a}{t-s+a-1} - \binom{t-s+2a}{a-1} \right), \\ C_{t,n,k,s} &= \sum_{m=n-s+1}^n \binom{m-1}{k+t-2s-1}, \\ D_{t,n,k,s} &= \sum_{m=k+t-2s+1}^n \binom{m-1}{k+t-2s-1}. \end{aligned}$$

Then we have the following results.

Lemma 3.10 *Let t, n, k, s be non-negative integers satisfy that $t \geq 2s, n \geq 1, t \geq 1$ and $k + t - 2s \leq n$. Then*

$$r(M_{k,k+t-2s}^{(t,n)}) = \begin{cases} \binom{n}{k} + \binom{n}{k+t-2s} - 1 & \text{if } t \geq n - 1, \\ \binom{n}{k} + A_{t,n,k,s}^{(1)} + A_{t,n,k,s}^{(2)} + C_{t,n,k,s} & \text{if } t < n - 1, 0 \leq k - s \leq \lfloor (n - t)/2 \rfloor, \\ \binom{n}{k} + B_{t,n,k,s} + C_{t,n,k,s} & \text{if } t < n - 1, \lfloor (n - t)/2 \rfloor \leq k - s \leq n - t, \\ \binom{n}{k} + D_{t,n,k,s} & \text{if } t < n - 1, k - s \leq 0 \text{ or } k - s \geq n - t. \end{cases} \quad (13)$$

Lemma 3.11 *We have*

$$bw(M^{(t,n)}) = \sum_{k=\lfloor (n-t)/2 \rfloor}^{\lfloor (n-t)/2 \rfloor + t - 1} \binom{n}{k} + \sum_{a=0}^{\lfloor (n-t-1)/2 \rfloor} \left(\binom{t+2a}{t+a-1} - \binom{t+2a}{a-1} \right). \quad (14)$$

From Eq 10 and 12, we can prove Lemmas 3.10 and 3.11 by induction. The complete proofs of Lemmas 3.10 and 3.11 can be find in **Appendix A** and **Appendix B**, respectively. Lemmas 3.11 and 3.6 show the exact bandwidth of $H(t, 2, n)$.

For a set $S \subseteq V$, let $N(S) = \{v \in V - S \mid \exists u \in S, uv \in E\}$, $\Phi(S) = |N(S)|$ and $b_v(l, G) = \min_{S \subseteq V, |S|=l} \Phi(S)$. Harper [13] showed that $bw(G) = \max_{1 \leq s \leq |V|} b_v(s, G)$ if G admits a Hales numbering. Therefore, $bw(H(t, 2, n)) = \max_{1 \leq s \leq 2^n} b_v(s, H(t, 2, n))$.

Theorem 3.12 (Theorem 1 of [8]) *Let $G(V, E)$ be any graph on n vertices, and let $1 \leq s \leq n$. Then $pw(G) \geq b_v(s, G)$.*

Lemma 3.13 *We have $pw(H(t, 2, n)) = bw(H(t, 2, n))$.*

Proof. By Theorem 3.12, we have $pw(H(t, 2, n)) \geq b_v(s, H(t, 2, n))$ for all $1 \leq s \leq 2^n$. Then $pw(H(t, 2, n)) \geq \max_{1 \leq s \leq 2^n} b_v(s, H(t, 2, n)) = bw(H(t, 2, n))$. Combining Eq 2, we have $pw(H(t, 2, n)) = bw(H(t, 2, n))$. \square

Theorem 1.1 can be derived from Lemmas 3.6, 3.11 and 3.13.

3.2 Treewidth of $H(t, q, n)$

In this subsection we analyze the asymptotic behavior of $tw(H(t, q, n))$ when n goes to infinity and give proof of Theorem 1.2.

We first prove the lower bound of $tw(H(t, q, n))$ by Proposition 3.14.

Proposition 3.14 (Lemma 7 of [8]) *Let $G(V, E)$ be a graph with n vertices. If for each subset X of V with $n/4 \leq |X| \leq n/2$, $\Phi(X) \geq k$, then $tw(G) \geq k - 1$.*

Lemma 3.15 $tw(H(t, q, n)) \geq c_1 tq^n / \sqrt{n}$ for some constant c_1 .

Proof. By Proposition 3.14, we have $tw(H(t, q, n)) \geq \min b_v(m, H(t, q, n)) - 1$ over integers m in the range $q^n/4 \leq m \leq q^n/2$. So it is sufficient to give a lower bound for $b_v(m, H(t, q, n))$ over $m \in [q^n/4, q^n/2]$.

In [12], Harper showed that² if

$$m = q^n \sum_{i=0}^r \binom{n}{i} x^{n-i} (1-x)^i \text{ for some } x, r, 0 < x < 1, \quad (15)$$

then

$$b_v(m, H(t, q, n)) \geq q^n \min_{x, r} \left\{ \sum_{i=1}^t \left(\binom{n}{r+i} x^{n-r-i} (1-x)^{r+i} \right) \right\}, \quad (16)$$

where the minimum is taken over all x, r satisfying Eq 15.

In [8], it is proved that when $m = q^n \sum_{i=0}^r \binom{n}{i} x^{n-i} (1-x)^i$, we have

$$n(1-x) - \sqrt{4nx(1-x)} < r < n(1-x) + \sqrt{4nx(1-x)}.$$

By Stirling's approximation, it can be shown that for all r in the above range, we have

$$q^n \binom{n}{r+i} x^{n-r-i} (1-x)^{r+i} \geq c_1 (q^n / \sqrt{n}), 1 \leq i \leq t$$

for some constant $c_1 > 0$. Then, $b_v(m, H(t, q, n)) \geq c_1 tq^n / \sqrt{n}$. □

Then we intend to estimate the upper bound via bandwidth.

Lemma 3.16 $tw(H(t, q, n)) \leq pw(H(t, q, n)) \leq c_2 tq^n / \sqrt{n}$ for some constant c_2 .

Proof. For convenience, we first assume q is even. The case when q is odd can be handled similarly. Let f be a function from $[q]$ to $\{0, 1\}$ as follows:

$$f(i) = \begin{cases} 0 & \text{if } 1 \leq i \leq q/2, \\ 1 & \text{if } q/2 < i \leq q. \end{cases}$$

Suppose that $(a_1, a_2, \dots, a_n) \in [q]^n$ is an n -vector corresponding to a vertex x of $H(t, q, n)$. Define function g from $V(H(t, q, n))$ to $V(H(t, 2, n))$ that maps $x \in V(H(t, q, n))$ to the vertex

²This statement is not explicitly stated in [12], but can be easily inferred from Theorem 3 on pp. 302.

$g(x) \in V(H(t, 2, n))$ which corresponds to the vector $(f(a_1), f(a_1), \dots, f(a_n))$. Note that g maps exactly $(q/2)^n$ vertices of $H(t, q, n)$ to a given vertex of $H(t, 2, n)$.

Let $H(t, 2, n)$ have a path decomposition whose bags are $\{P_i\}$. By replacing each P_i with $P'_i = \bigcup_{y \in P_i} g^{-1}(y)$, it is easy to show that $\{P'_i\}$ is a path decomposition of $H(t, q, n)$. Therefore, $pw(H(t, q, n)) \leq pw(H(2, q, n)) \cdot \left(\frac{q}{2}\right)^n$.

From Theorem 1.1 and Stirling's approximation, we have $bw(H(t, 2, n)) \leq c_2 t 2^n / \sqrt{n}$ for some constant c_2 . Therefore, we have $tw(H(t, q, n)) \leq pw(H(t, q, n)) \leq c_2 t q^n / \sqrt{n}$. \square

Combining Lemmas 3.15 and 3.16, we can derive Theorem 1.2.

4 Treewidth of bipartite Kneser graph and Johnson graph

For positive integers n and k satisfying $n \geq 2k + 1$, the bipartite Kneser graph $BK(n, k)$ has all subsets of $[n]$ with k or $n - k$ elements as vertices and an edge between any two vertices when one is a subset of the other. It is also called *middle cube graph*. In the following, we will use k -subsets and $(n - k)$ -subsets of $[n]$ to represent the vertices of bipartite Kneser graph. We call vertices that are k -subsets of $[n]$ the *left part* denoted by V_L , and the rest is called the *right part* denoted by V_R . V_L and V_R are two parts of bipartite Kneser graph $BK(n, k)$.

For positive integers n and k satisfying $n > k$, the *Johnson graph* $J(n, k)$ has all subsets of $[n]$ with k elements and an edge between any two vertices when their intersection has exactly $(k - 1)$ elements. Since we have a bijection between all k -subsets of $[n]$ and all binary n -vectors with exactly k ones, we can also treat a vertex of $J(n, k)$ as a n -vector with exactly k ones. From this point of view, two vertices are adjacent iff their the Hamming distance of their corresponding n -vectors is no more than 2. Therefore, $J(n, k)$ is the k -th slice of $H(2, 2, n)$, that is, $J(n, k)$ is a subgraph of $H(2, 2, n)$ induced by vertices from $H(2, 2, n)$ corresponding to n -vector with exactly k ones.

Let $G(V, E)$ be a graph and $S \subseteq V$ a vertex subset of G . Denote the subgraph of G induced by S as $G[S]$. Recalling the definition of $V_k^{(t, n)}$ in subsection 3.1, we have Proposition 4.1.

Proposition 4.1 *We have $J(n, k) \cong H(2, 2, n)[V_k^{(2, n)}]$.*

4.1 Treewidth of $BK(n, k)$ when n is large enough

In this section, we focus on the treewidth of $BK(n, k)$ when n is large enough and give proof of Theorem 1.3. Before proof, we need the following proposition.

Proposition 4.2 ([14]) *For any graph G , $tw(G) \leq \max\{\Delta(G), |V(G)| - \alpha(G) - 1\}$, where $\alpha(G)$ is the independent number of G .*

Lemma 4.3 *We have $tw(BK(n, k)) \leq \binom{n}{k} - 1$.*

Proof. From [18], we have that bipartite Kneser graph has a perfect matching. Consequently, $\alpha(BK(n, k)) = \binom{n}{k}$. Note that $BK(n, k)$ is a regular graph with order $2\binom{n}{k}$ and $\Delta(BK(n, k)) = \binom{n-k}{k}$. By Proposition 4.2, we have $tw(BK(n, k)) \leq \binom{n}{k} - 1$. \square

Lemma 4.4 *When $k \geq 2$ and $3\binom{n-k}{k} \geq 2\binom{n}{k}$, $tw(BK(n, k)) \geq \binom{n}{k} - 1$.*

Proof. Denote $BK(n, k)$ by G and $V(G) = V_L \cup V_R$, where $V_L = \binom{[n]}{k}$ and $V_R = \binom{[n]}{n-k}$. Suppose $tw(G) < \binom{n}{k} - 1$. From Corollary 2.5, there exists a separator X of G with $|X| < \binom{n}{k}$ such that there exists non-empty vertex set A and B with $A \cup B = V(G) - X$, $A \cap B = \emptyset$, $|V(G) - X|/3 \leq |A|, |B| \leq 2|V(G) - X|/3$ and there is no edge between A and B . Let $A_L = A \cap V_L$, $A_R = A \cap V_R$, $B_L = B \cap V_L$ and $B_R = B \cap V_R$.

Since A, B are nonempty and $|X| < \binom{n}{k}$, we assume without loss of generality that A_L and B_R are nonempty. Let

$$\mathcal{A} = \left\{ S \in \binom{[n]}{k} \mid S \in A_L \right\},$$

$$\mathcal{B} = \left\{ S \in \binom{[n]}{n-k} \mid S \in B_R \right\}.$$

Let $\mathcal{C} = \{[n] - S \mid S \in \mathcal{B}\}$. Since there is no edge between A_L and B_R , for any $S_1 \in \mathcal{A}$ and $S_2 \in \mathcal{B}$, $S_1 \not\subseteq S_2$ which implies $S_1 \cap ([n] - S_2) \neq \emptyset$. Hence, \mathcal{A} and \mathcal{C} is cross-intersecting. From [11], we have $|\mathcal{A}| + |\mathcal{C}| \leq \binom{n}{k} - \binom{n-k}{k} + 1$. Notice that $|\mathcal{A}| = |A_L|$ and $|\mathcal{C}| = |\mathcal{B}| = |B_R|$. Then we have $|A_L| + |B_R| \leq \binom{n}{k} - \binom{n-k}{k} + 1$.

If A_R, B_L are both nonempty, then, similarly, we have $|A_R| + |B_L| \leq \binom{n}{k} - \binom{n-k}{k} + 1$. Hence, $|A| + |B| \leq 2\left(\binom{n}{k} - \binom{n-k}{k} + 1\right)$ and $|X| = |V(G)| - |A| - |B| \geq 2\binom{n-k}{k} - 2$. Since $k \geq 2$ and $3\binom{n-k}{k} \geq 2\binom{n}{k}$, we have $2\binom{n-k}{k} - 2 \geq \binom{n}{k}$ which derives a contradiction with $|X| < \binom{n}{k}$.

If A_R, B_L are both empty, then we have $|A| + |B| = |A_L| + |B_R| \leq \binom{n}{k} - \binom{n-k}{k} + 1$. Hence, $|X| = |V(G)| - |A| - |B| \geq \binom{n}{k} + \binom{n-k}{k} - 1 \geq \binom{n}{k}$, a contradiction.

If there is only one empty set in $\{A_R, B_L\}$, say $A_R \neq \emptyset$ and $B_L = \emptyset$, then we have $|B| = |B_R| \leq \binom{n}{k} - \binom{n-k}{k}$ by $|A_L| + |B_R| \leq \binom{n}{k} - \binom{n-k}{k} + 1$ and $A_L \neq \emptyset$. Since $|V(G) - X|/3 \leq |A|, |B| \leq 2|V(G) - X|/3$, we have $|A| + |B| \leq 3|B| \leq 3\left(\binom{n}{k} - \binom{n-k}{k}\right)$. Since $3\binom{n-k}{k} \geq 2\binom{n}{k}$, we have $|X| = |V(G)| - |A| - |B| \geq 3\binom{n-k}{k} - \binom{n}{k} \geq \binom{n}{k}$, a contradiction. \square

Theorem 1.3 can be easily derived from Lemmas 4.3 and 4.4.

4.2 Treewidth of $BK(2k + 1, k)$ and $J(2k + 1, k)$

When n is large enough, the treewidth of bipartite Kneser graph can be exactly calculated by Theorem 1.3. Now we focus on the treewidth of bipartite Kneser graph when n is small and give proof of Theorem 1.4.

In order to prove our result, we need more definitions. A graph G is chordal if and only if, in any cycle of length larger than 3 in G , there exists a chord connecting two nonadjacent

vertices of the cycle. Given a graph G , define $\omega(G)$, the clique number of G to be the number of vertices of the largest clique in G .

Proposition 4.5 ([27]) *Given a graph G , $tw(G) = \min\{\omega(H) - 1 \mid G \subseteq H, H \text{ is chordal}\}$.*

Lemma 4.6 *For any positive integer k , $tw(BK(2k + 1, k)) \leq tw(J(2k + 1, k))$.*

Proof. Let $n = 2k + 1$. By Proposition 4.5, there exists a chordal graph H such that $J(n, k) \subseteq H$ and $\omega(H) - 1 = tw(J(n, k))$. Add edges to the left part of $BK(n, k)$ such that the left part is isomorphic to H and denote the result graph as $BK'(n, k)$. That is, $BK'(n, k)[V_L] \cong H$.

We first claim that $BK'(n, k)$ is chordal. Suppose C is a cycle in $BK'(n, k)$ with length larger than 3. We first consider the case that there is a vertex $v \in V(C) \cap V_R$. Let u_1 and u_2 be the two neighbors of v on C . Then $u_1, u_2 \in V_L$. Let $A_v \in \binom{[n]}{k+1}$ be the corresponding set of v and $A_{u_1}, A_{u_2} \in \binom{[n]}{k}$ the corresponding sets of u_1 and u_2 respectively. Since $u_1v, u_2v \in E(BK'(n, k))$, we have $A_{u_1} \subseteq A_v$ and $A_{u_2} \subseteq A_v$. Then $|A_{u_1} \cap A_{u_2}| = k - 1$. Hence, $u_1u_2 \in E(BK'(n, k))$. Now assume all vertices of C are from the left side. Since H is chordal, there must exist a chord in C . From above all, we have that $BK'(n, k)$ is chordal.

Let W be a clique of $BK'(n, k)$. If there exists a vertex $v \in W$ from the right side, then $|W| \leq k + 1 + 1$ by $d(v) = k + 1$. If all vertices of W are from the left part, then $|W| \leq \omega(H) = tw(J(n, k)) + 1$. Hence

$$tw(BK(n, k)) \leq \omega(BK'(n, k)) - 1 \leq \max\{tw(J(n, k)), k + 1\}.$$

Notice that $J(2k + 1, k)$ is $k(k + 1)$ -regular. Since $tw(G) \geq \delta(G)$ [21] for any graph G , we have $tw(J(2k + 1, k)) \geq k(k + 1) \geq k + 1$. Then $tw(BK(2k + 1, k)) \leq tw(J(2k + 1, k))$. \square

Lemma 4.7 *For positive integers n and k satisfying $n > k$, $tw(J(n, k)) \leq bw(M_{k,k}^{(2,n)}) = r(M_{k,k}^{(2,n)}) - \binom{n}{k}$.*

Proof. Let $G = H(2, 2, n)[V_k^{(2,n)}]$ for short. By Proposition 4.1, we have $G \cong J(n, k)$.

Let $t = 2$ in subsection 3.1. Recalling that $\eta' \triangleq \eta_k^{(n)}$ is an ordering of $V_k^{(2,n)}$, then η' is also an ordering of G . Note that the adjacency matrix of G with the ordering η' is exactly $M_{k,k}^{(2,n)}$. By Remark 3.5 and the definition of bandwidth, we have

$$bw(M_{k,k}^{(2,n)}) = \max_{e \in E(G)} \Delta(e, \eta') \geq \min_{\eta} \max_{e \in E(G)} \Delta(e, \eta) = bw(G) = bw(J(n, k)). \quad (17)$$

Combining Eq 2 and Eq 8 with $s = |V_k^{(2,n)}| = \binom{n}{k}$, we can derive the lemma. \square

Specifically, take $n = 2k + 1$, and then from Lemma 3.10, we can derive the asymptotic behavior of $bw(M_{k,k}^{(2,2k+1)})$ by calculating:

$$\lim_{k \rightarrow +\infty} bw(M_{k,k}^{(2,2k+1)}) / \binom{2k+1}{k} = 1/2. \quad (18)$$

Therefore, $tw(J(2k+1, k)) = O(\binom{2k+1}{k})$.

Proposition 4.8 ([9]) *Suppose G is a k -regular graph with n vertices and $A(G)$ is the adjacency matrix of G . Let $\mu(G) = k - \lambda(G)$ where $\lambda(G)$ is the second-largest eigenvalue of $A(G)$. Then*

$$tw(G) \geq \left\lfloor \frac{3n}{4} \frac{\mu(G)}{\Delta(G) + 2\mu(G)} \right\rfloor - 1.$$

Proposition 4.9 ([24]) *The characteristic polynomial of $BK(2k+1, k)$ is*

$$\prod_{i=1}^{k+1} (\lambda + i)^{\binom{n}{k+1-i} - \binom{n}{k-i}} (\lambda - i)^{\binom{n}{k+1-i} - \binom{n}{k-i}}.$$

Lemma 4.10 *We have*

$$tw(BK(2k+1, k)) \geq \left\lfloor \frac{3}{2} \binom{2k+1}{k} \frac{1}{k+3} \right\rfloor - 1.$$

Lemma 4.10 can be derived from Propositions 4.8 and 4.9. Theorem 1.4 can be derived by Lemmas 4.6, 4.7 and 4.10.

5 Treewidth of generalized Petersen graph

In this section, we determine the treewidth of generalized Petersen graph. The vertex set and edge set of generalized Petersen graph $G_{n,k}$ are

$$\begin{aligned} V(G_{n,k}) &= \{v_1, \dots, v_n, u_1, \dots, u_n\}, \\ E(G_{n,k}) &= \{v_i u_i\} \cup \{v_i v_{i+1}\} \cup \{u_i u_{i+k}\}, i = 1, 2, \dots, n, \end{aligned}$$

where subscripts are to be read modulo n and $k < n/2$. Let G be a graph and X and Y are two connected subgraphs of G . We say X *touches* Y when $V(X) \cap V(Y) \neq \emptyset$ or there exists an edge between X and Y . A *bramble* of G is a family of connected subgraphs of G that all touch each other. Let S be a subset of $V(G)$. S is said to be a hitting set of bramble B if S has nonempty intersection with each of the subgraphs in B . The order of a bramble is the smallest size of a hitting set. Brambles may be used to characterize the treewidth of a given graph.

Proposition 5.1 ([28]) *Let G be a graph. Then $tw(G) \geq k$ if and only if G contains a bramble of order at least $k+1$.*

With the help of Proposition 5.1, we now can give proof of Theorem 1.5.

Proof of Theorem 1.5: First, we intend to prove the lower bound. Construct a bramble $B = \{B_i\}$ of $G_{n,k}$ as

$$\begin{aligned} V_i &= \{v_i, v_{i+1}, \dots, v_{i+t}, u_{i+t}, u_{i+t+k}, u_{i+t+2k}, \dots, u_{i+t+tk}\}, \\ B_i &= G_{n,k}[V_i], \end{aligned}$$

where $t = \lceil \frac{n}{2k+2} \rceil, i = 1, 2, \dots, n$. Then we have $|V_i| = 2t+2$ and B_i is connected. For each pair i and j , we intend to prove that B_i touches B_j . Without loss of generality, assume $1 \leq i < j \leq n$.

- If $j \leq i+t$, then $v_j \in V_i \cap V_j$.
- If $i+t < j \leq i+t+tk$, let r be the minimum integer in $\{0, 1, \dots, k-1\}$ such that $j+r \equiv i+t \pmod{k}$. Noticing that $t \geq k$ when $n \geq 8(2k+2)^2$. Hence, $v_{j+r} \in V_j, u_{j+r} \in V_i$, and $u_{j+r}v_{j+r} \in E(G_{n,k})$.
- If $i+t+tk < j < n+i-t$, then $j+t < i+n < j+n-t-tk \leq j+t+tk$. The last inequality comes from $2kt+2t \geq n$ by $t = \lceil \frac{n}{2k+2} \rceil$ and then we have $t+tk \geq n-t-tk$ which implies $j+t < i+n \leq j+t+tk$. The next proof is the same as that in the second situation.
- If $n+i-t \leq j \leq n$, the proof is the same as that in the first situation.

Then B is a bramble. Let S be a hitting set of B . We construct a hypergraph H with vertex set $V(H) = V(G_{n,k})$ and hyperedge set $\{V_i\}_{1 \leq i \leq n}$. Then S is a transversal of H and thus, $\min |S| = \tau(H)$ where $\tau(H)$ is the transversal number of H . Since $\tau(H) \geq \max_{H' \subseteq H} \frac{m(H')}{\Delta(H')} [3]$, the order of B is

$$\min |S| \geq \frac{m(H)}{\Delta(H)} = \frac{n}{t+1}.$$

Since $t = \lceil \frac{n}{2k+2} \rceil$, we have $t-1 \leq \frac{n}{2k+2}$ and

$$2k+2 - \frac{n}{t+1} \leq \frac{n}{t-1} - \frac{n}{t+1} = \frac{2n}{t^2-1} \leq 1.$$

Therefore, the order of B is no less than $2k+2$. From Proposition 5.1, we can derive that $tw(G_{n,k}) \geq 2k+1$.

The upper bound can be proved via construction. Construct a path-decomposition of $G_{n,k}$ as following (see Figure 3 and Figure 4).

Step 1. Let $A = \{v_1, u_1, \dots, u_k\}$, $B_1 = \{v_1, \dots, v_k\}$ and $B_2 = \{v_k, \dots, v_{2k}\}$.

Step 2. Let $X_1 = \{u_{k+1}, \dots, u_{2k}, v_{2k}\}$.

Step 3. Recursively define $Y_i = X_i \cup \{u_{2k+i}\}$, $Z_i = Y_i - \{u_{k+i}\}$, $W_i = Z_i \cup \{v_{2k+i}\}$ and $X_{i+1} = W_i - \{v_{2k+i-1}\}$ for $1 \leq i \leq n-2k$. It is easy to verify that $X_i = \{u_{k+i}, \dots, u_{2k+i-1}, v_{2k+i-1}\}$.




Figure 3: The vertex sets X_i, Y_i, Z_i and W_i .




Figure 4: The defined sets on generalized Petersen graph.

Step 4. Define $X_{n-2k+1} = W_{n-2k} - \{v_{n-1}\}$.

Step 5. Define a path decomposition \mathcal{P}_1 of $G_{n,k} - A$ by successively connect

$$(B_1, B_2, X_1, Y_1, Z_1, W_1, X_2, \dots, W_{n-2k}, X_{n-2k+1}).$$

Step 6. Add each vertex in A to all bags of \mathcal{P}_1 , then we obtain a path decomposition \mathcal{P}_2 of $G_{n,k}$.

It is easy to verify \mathcal{P}_1 is a path-decomposition of $G_{n,k} - A$ by checking the three properties which implying that \mathcal{P}_2 is a path-decomposition of $G_{n,k}$.

Since $|X_i| = |Z_i| = k + 1, |Y_i| = |W_i| = k + 2, |B_1| = k, |B_2| = k + 1$ and $|A| = k + 1$, the width of \mathcal{P}_2 is $2k + 3$ and, hence, $tw(G_{n,k}) \leq pw(G_{n,k}) \leq 2k + 2$. \square

Acknowledgement

M. Cao is supported by the National Natural Science Foundation of China (Grant 12301431), M. Lu is supported by the National Natural Science Foundation of China (Grant 12171272 & 12161141003).

References

- [1] B. Aiello and T. Leighton. Coding theory, hypercube embeddings, and fault tolerance. In *Proceedings of the third annual ACM symposium on Parallel algorithms and architectures*, pages 125–136. 1991.
- [2] S. Arnborg, D. G. Corneil, and A. Proskurowski. Complexity of finding embeddings in ak-tree. *SIAM Journal on Algebraic Discrete Methods*, 8(2):277–284, 1987.
- [3] C. Berge. *Hypergraphs: combinatorics of finite sets*, volume 45. Elsevier, 1984.
- [4] H. L. Bodlaender. A linear time algorithm for finding tree-decompositions of small treewidth. In *Proceedings of the twenty-fifth annual ACM symposium on Theory of computing*, pages 226–234. 1993.
- [5] H. L. Bodlaender. A partial k-arboretum of graphs with bounded treewidth. *Theoretical computer science*, 209(1-2):1–45, 1998.
- [6] H. Bruhn, R. Lang, and M. Stein. List edge-coloring and total coloring in graphs of low treewidth. *Journal of Graph Theory*, 81(3):272–282, 2016.
- [7] M. Cao, K. Liu, M. Lu, and Z. Lv. Treewidth of the q-kneser graphs. *Discrete Applied Mathematics*, 342:174–180, 2024.
- [8] L. S. Chandran and T. Kavitha. The treewidth and pathwidth of hypercubes. *Discrete Mathematics*, 306(3):359–365, 2006.
- [9] L. S. Chandran and C. Subramanian. A spectral lower bound for the treewidth of a graph and its consequences. *Information Processing Letters*, 87(4):195–200, 2003.
- [10] M. Chimani, P. Mutzel, and B. Zey. Improved steiner tree algorithms for bounded treewidth. *Journal of Discrete Algorithms*, 16:67–78, 2012. ISSN 1570-8667. Selected papers from the 22nd International Workshop on Combinatorial Algorithms (IWOCA 2011).
- [11] P. Frankl and N. Tokushige. Some best possible inequalities concerning cross-intersecting families. *Journal of Combinatorial Theory, Series A*, 61(1):87–97, 1992.
- [12] L. Harper. On an isoperimetric problem for hamming graphs. *Discrete applied mathematics*, 95(1-3):285–309, 1999.

- [13] L. H. Harper. Optimal numberings and isoperimetric problems on graphs. *Journal of Combinatorial Theory*, 1(3):385–393, 1966.
- [14] D. J. Harvey and D. R. Wood. Treewidth of the kneser graph and the erdős-ko-rado theorem. *The Electronic Journal of Combinatorics*, pages P1–48, 2014.
- [15] D. J. Harvey and D. R. Wood. The treewidth of line graphs. *Journal of Combinatorial Theory, Series B*, 132:157–179, 2018.
- [16] J. P. Hayes and T. Mudge. Hypercube supercomputers. *Proceedings of the IEEE*, 77(12):1829–1841, 1989.
- [17] M. A. Jiménez-Montano, C. R. de la Mora-Basanez, and T. Pöschel. On the hypercube structure of the genetic code. *arXiv preprint cond-mat/0204044*, 2002.
- [18] K. Jin. On 1-factorizations of bipartite kneser graphs. *Theoretical Computer Science*, 838:81–93, 2020.
- [19] T. Kloks. Treewidth of circle graphs. In *Algorithms and Computation: 4th International Symposium, ISAAC’93 Hong Kong, December 15–17, 1993 Proceedings 4*, pages 108–117. Springer, 1993.
- [20] T. Korhonen and D. Lokshtanov. An improved parameterized algorithm for treewidth. In *Proceedings of the 55th Annual ACM Symposium on Theory of Computing*, pages 528–541. 2023.
- [21] A. M. Koster, T. Wolle, and H. L. Bodlaender. Degree-based treewidth lower bounds. In *International Workshop on Experimental and Efficient Algorithms*, pages 101–112. Springer, 2005.
- [22] A. Li and T. Müller. On the treewidth of random geometric graphs and percolated grids. *Advances in Applied Probability*, 49(1):49–60, 2017.
- [23] K. Liu, M. Cao, and M. Lu. Treewidth of the generalized kneser graphs. *The Electronic Journal of Combinatorics*, 29:P1–57, 2022.
- [24] K. Qiu, R. Qiu, Y. Jiang, and J. Shen. On the spectrum of middle-cubes. *arXiv preprint arXiv:0911.0126*, 2009.
- [25] N. Robertson and P. D. Seymour. Graph minors. i. excluding a forest. *Journal of Combinatorial Theory, Series B*, 35(1):39–61, 1983.
- [26] N. Robertson and P. D. Seymour. Graph minors. iii. planar tree-width. *Journal of Combinatorial Theory, Series B*, 36(1):49–64, 1984.
- [27] N. Robertson and P. D. Seymour. Graph minors. ii. algorithmic aspects of tree-width. *Journal of algorithms*, 7(3):309–322, 1986.

- [28] P. D. Seymour and R. Thomas. Graph searching and a min-max theorem for tree-width. *Journal of Combinatorial Theory, Series B*, 58(1):22–33, 1993.
- [29] X. Wang, X. Wu, and S. Dumitrescu. On explicit formulas for bandwidth and antibandwidth of hypercubes. *Discrete Applied Mathematics*, 157(8):1947–1952, 2009.
- [30] D. R. Wood. Treewidth of cartesian products of highly connected graphs. *Journal of Graph Theory*, 73(3):318–321, 2013.
- [31] X. Zhou, S.-i. Nakano, and T. Nishizeki. Edge-coloring partialk-trees. *Journal of Algorithms*, 21(3):598–617, 1996.

Appendix A: Proof of Lemma 3.10

Before the proof, we need several definitions. We say the parameter tuple (t, n, k, s) is *valid* if $t \geq 2s, n \geq 1, 1 \leq t < n - 1$ and $k + t - 2s \leq n$, that corresponds to the non-trivial situation of Lemma 3.10. The parameter tuple of matrix in the formula of $r_{t,n,k,t-2s}^{(i)}$ ($i = 1, 2, 3, 4$) is shown as Table 1.

$r^{(i)}$ -term	matrix	t'	n'	k'	s'
$r_{t,n,k,t-2s}^{(1)}$	$r(M_{k-1,k+t-2s-1}^{(t,n-1)})$	t	$n - 1$	$k - 1$	s
$r_{t,n,k,t-2s}^{(2)}$	$r(M_{k-1,k+t-2s}^{(t-1,n-1)})$	$t - 1$	$n - 1$	$k - 1$	$s - 1$
$r_{t,n,k,t-2s}^{(3)}$	$r(M_{k,k+t-2s-1}^{(t-1,n-1)})$	$t - 1$	$n - 1$	k	s
$r_{t,n,k,t-2s}^{(4)}$	$r(M_{k,k+t-2s}^{(t,n-1)})$	t	$n - 1$	k	s

Table 1: Parameter tuple of the matrix in the Eq 13 of $r_{t,n,k,t-2s}^{(i)}$.

First, it is easy to verify that $D_{t,n,k,s} = A_{t,n,k,s}^{(1)} + A_{t,n,k,s}^{(2)} + C_{t,n,k,s}$ holds when $k - s = 0$ and, $D_{t,n,k,s} = B_{t,n,k,s} + C_{t,n,k,s}$ when $k - s = n - t$ and $A_{t,n,k,s}^{(1)} + A_{t,n,k,s}^{(2)} = B_{t,n,k,s}$ hold when $k - s = \lfloor (n - t)/2 \rfloor$. Hence, in those situations, we can calculate $r(M_{k,k+t-2s}^{(t,n)})$ in both ways.

Step 1: verify Lemma 3.10 when either $t \geq n - 1$, or $t = 1$.

When $t \geq n - 1$, it is trivial to verify $M_{k,k+t-2s}^{(t,n)}$ is an all-one matrix and, thus, $r(M_{k,k+t-2s}^{(t,n)}) = \binom{n}{k} + \binom{n}{k+t-2s} - 1$ from definition.

When $t = 1$, then $s = 0$. In this case $r_{1,n,k,1}^{(2)} = -\infty$ and $r_{1,n,k,1}^{(3)} = -\infty$ and, hence, $r(M_{k,k+1}^{(1,n)}) = \max\{r_{1,n,k,1}^{(1)}, r_{1,n,k,1}^{(4)}\}$. We can prove Eq 13 holds for $t = 1$ by induction on n . The proof is omitted.

Then, we give a proof of Lemma 3.10 via Eq 12 by induction. Suppose Eq 13 holds for $t + n + k + s \leq N - 1$. Now consider when $t + n + k + s = N$. In the following steps, we only need to consider the case $t \geq 2$ and $t < n - 1$.

Step 2: verify Eq 13 when $k = 0$.

If $k = 0$, then $k + t - 2s < n$ and $r_{t,n,k,t-2s}^{(1)} = -\infty$ and $r_{t,n,k,t-2s}^{(2)} = -\infty$.

Case 2.1: If $t = 2s$, then $r(M_{k,k+t-2s}^{(t,n)}) = r(M_{0,0}^{(t,n)}) = 1$ from definition. It is easy to verify Eq 13 holds in this case.

Case 2.2: If $t > 2s$, then $(t-1, n-1, k, s)$ are valid and, hence, $r_{t,n,k,t-2s}^{(3)} = r(M_{k,k+(t-1)-2s}^{(t-1,n-1)})$ can be calculated via Eq 13 by induction.

Since $k - s \leq 0$, we have

$$\begin{aligned} r_{t,n,k,t-2s}^{(3)} &= r(M_{k,k+(t-1)-2s}^{(t-1,n-1)}) \\ &= \binom{n-1}{k} + D_{t-1,n-1,k,s} \\ &\leq \binom{n}{k} + \sum_{m=k+t-2s+1}^n \binom{m-1}{k+t-2s-1} = \binom{n}{k} + D_{t,k,n,s}, \end{aligned}$$

where the second equality comes from $k - s \leq 0$ and Eq 13 by induction.

Subcase 2.2.1: If $t = n - 2$, then

$$\begin{aligned} r_{t,n,k,t-2s}^{(4)} &= r(M_{k,k+t-2s}^{(t,n-1)}) + \binom{n-1}{k+t-2s-1} \\ &= \binom{n-1}{k} + \binom{n-1}{k+t-2s} - 1 + \binom{n-1}{k+t-2s-1} = \binom{n}{n-2s-2} \\ &= \binom{n}{k} + D_{t,n,k,s}, \end{aligned}$$

where the second equality comes from the trivial situation of Lemma 3.10 which we have proved before. Therefore, we have verified Eq 13 in this subcase.

Subcase 2.2.2: If $t < n - 2$, then $(t, n - 1, k, s)$ are valid and, hence, $r_{t,n,k,t-2s}^{(4)} = r(M_{k,k+t-2s}^{(t,n-1)})$ can be calculated via Eq 13 by induction.

$$\begin{aligned} r_{t,n,k,t-2s}^{(4)} &= r(M_{k,k+t-2s}^{(t,n-1)}) + \binom{n-1}{k+t-2s-1} \\ &= \binom{n-1}{k} + D_{t,n-1,k,s} + \binom{n-1}{k+t-2s-1} \\ &= \binom{n}{k} + D_{t,n,k,s}, \end{aligned}$$

where the second equality comes from Step 1 and the third equality comes from $k = 0$.

From above, combining Eq 12, we can verify that Eq 13 holds when $k = 0$.

Step 3: verify Eq 13 when $k + t - 2s = n$.

In this case, $r_{t,n,k,t-2s}^{(2)} = -\infty$ and $r_{t,n,k,t-2s}^{(4)} = -\infty$.

Case 3.1: If $t = 2s$, then $k = n$ and $r(M_{k,k+t-2s}^{(t,n)}) = r(M_{n,n}^{(t,n)}) = 1$ from definition. It is easy to verify Eq 13 holds in this case.

Case 3.2: If $t < 2s$, then $(t-1, n-1, k, s)$ is valid and, hence, $r_{t,n,k,p}^{(3)} = r(M_{k,k+p-1}^{(t-1,n-1)})$ can be calculated via Eq 13 by induction. Since $k - s \geq n - t$, we have

$$\begin{aligned} r_{t,n,k,t-2s}^{(3)} &= r(M_{k,k+(t-1)-2s}^{(t-1,n-1)}) \\ &= \binom{n-1}{k} + D_{t-1,n-1,k,s} \\ &\leq \binom{n}{k} + \sum_{m=k+t-2s+1}^n \binom{m-1}{k+t-2s-1} = \binom{n}{k} + D_{t,k,n,s}, \end{aligned}$$

where the second equality comes from Eq 13 by induction.

Subcase 3.2.1: If $t = n - 2$, then

$$\begin{aligned} r_{t,n,k,t-2s}^{(1)} &= r(M_{k-1,k-1+t-2s}^{(t,n-1)}) + \binom{n-1}{k} \\ &= \binom{n-1}{k-1} + \binom{n-1}{k-1+t-2s} - 1 + \binom{n-1}{k} \\ &= \binom{n}{k} + \sum_{m=k+t-2s+1}^n \binom{m-1}{k+t-2s-1} = \binom{n}{k} + D_{t,n,k,s}, \end{aligned}$$

where the second equality comes from Step 1 and the following equalities come from $k+t-2s = n$.

Subcase 3.2.1: If $t < n - 2$, then $(t, n-1, k-1, s)$ is valid and, hence, $r_{t,n,k,t-2s}^{(1)} = r(M_{k-1,k-1+t-2s}^{(t,n-1)}) + \binom{n-1}{k}$ can be calculated via Eq 13 by induction. That is

$$\begin{aligned} r_{t,n,k,t-2s}^{(1)} &= r(M_{k-1,k-1+t-2s}^{(t,n-1)}) + \binom{n-1}{k} \\ &= \binom{n-1}{k-1} + D_{t,n-1,k-1,s} + \binom{n-1}{k} \\ &= \binom{n}{k} + D_{t,n,k,s}, \end{aligned}$$

where the second equality comes from Eq 13 by induction, and the third equality comes from $k+t-2s = n$.

From above, combining Eq 12, we can verify that Eq 13 holds when $k + t - 2s = n$.

Step 4: verify Eq 13 when $k > 0$ and $k + t - 2s < n$.

In this case, $\{r_{t,n,k,t-2s}^{(i)}\}_{1 \leq i \leq 4}$ are all positive. By the definition, we have that $r_{t,n,k,t-2s}^{(3)} \leq r_{t,n,k,t-2s}^{(2)}$ always holds. Hence, we only need to consider $r_{t,n,k,t-2s}^{(1)}$, $r_{t,n,k,t-2s}^{(2)}$ and $r_{t,n,k,t-2s}^{(4)}$ and their value can be calculated from Eq 13 by induction.

Step 4.1: verify that $r_{t,n,k,t-2s}^{(2)}$ is always equal to RHS of Eq 13.

Note that $(t-1, n-1, k-1, s-1)$ is valid and, hence, we can calculate $r_{t,n,k,t-2s}^{(2)}$ via Eq 13 by induction.

Case 4.1.1: If $k-s < 0$ or $k-s > n-t$, then $(k-1) - (s-1) < 0$ or $(k-1) - (s-1) > (n-1) - (t-1)$.

Therefore,

$$\begin{aligned} r_{t,n,k,t-2s}^{(2)} &= r(M_{k-1,(k-1)+(t-1)-2(s-1)}^{(t-1,n-1)}) + \binom{n-1}{k} + \binom{n-1}{k+t-2s-1} \\ &= \binom{n-1}{k-1} + \binom{n-1}{k} + \binom{n-1}{k+t-2s-1} + D_{t-1,n-1,k-1,s-1} \\ &= \binom{n}{k} + \sum_{m=k+t-2s+1}^n \binom{m-1}{k+t-2s-1} = \binom{n}{k} + D_{t,n,k,s}, \end{aligned}$$

where the second equality comes from Eq 13 by induction.

Case 4.1.2: If $0 \leq k-s \leq \lfloor (n-t)/2 \rfloor$, then $0 \leq (k-1) - (s-1) \leq \lfloor ((n-1) - (t-1))/2 \rfloor$.

Therefore,

$$\begin{aligned} r_{t,n,k,t-2s}^{(2)} &= r(M_{k-1,(k-1)+(t-1)-2(s-1)}^{(t-1,n-1)}) + \binom{n-1}{k} + \binom{n-1}{k+t-2s-1} \\ &= \binom{n}{k} + \sum_{a=0}^{k-s-1} \left(\binom{t-s+2a}{t-s+a-1} - \binom{t-s+2a}{a-1} \right) \\ &\quad + \sum_{m=t-3s+1+2k}^{n-s} \left(\binom{m-1}{k+t-2s-1} - \binom{m-1}{k-s-1} \right) + \sum_{m=n-s+1}^n \binom{m-1}{k+t-2s-1} \\ &= \binom{n}{k} + A_{t,n,k,s}^{(1)} + A_{t,n,k,s}^{(2)} + C_{t,n,k,s}, \end{aligned}$$

where the second equality comes from Eq 13 by induction.

Case 4.1.3: If $\lfloor (n-t)/2 \rfloor \leq k-s \leq n-t$, then $\lfloor ((n-1) - (t-1))/2 \rfloor \leq (k-1) - (s-1) \leq (n-1) - (t-1)$.

Therefore,

$$\begin{aligned}
r_{t,n,k,t-2s}^{(2)} &= r(M_{k-1,(k-1)+(t-1)-2(s-1)}^{(t-1,n-1)}) + \binom{n-1}{k} + \binom{n-1}{k+t-2s-1} \\
&= \binom{n-1}{k-1} + \binom{n-1}{k} + \binom{n-1}{k+t-2s-1} + B_{t-1,n-1,k-1,s-1} + C_{t-1,n-1,k-1,s-1} \\
&= \binom{n}{k} + \sum_{a=0}^{n-t-k+s-1} \left(\binom{t-s+2a}{t-s+a-1} - \binom{t-s+2a}{a-1} \right) + \sum_{m=n-s+1}^n \binom{m-1}{k+t-2s-1} \\
&= \binom{n}{k} + B_{t,n,k,s} + C_{t,n,k,s},
\end{aligned}$$

where the second equality comes from Eq 13 by induction.

Consequently, $r_{t,n,k,t-2s}^{(2)}$ is always equal to RHS of Eq 13. Hence, we only need to prove $r_{t,n,k,t-2s}^{(1)} \leq r_{t,n,k,t-2s}^{(2)}$ and $r_{t,n,k,t-2s}^{(4)} \leq r_{t,n,k,t-2s}^{(2)}$. Then by Eq 12, we can finally prove Lemma 3.10.

Step 4.2: verify that $r_{t,n,k,t-2s}^{(1)} \leq r_{t,n,k,t-2s}^{(2)}$.

Case 4.2.1: If $(k-1) - s < 0$, then $(k-1) - (s-1) = k-s \leq 0$ and

$$\begin{aligned}
&D_{t,n-1,k-1,s} - D_{t-1,n-1,k-1,s-1} \\
&= \sum_{m=k+t-2s}^{n-1} \binom{m-1}{k+t-2s-2} - \sum_{m=k+t-2s+1}^{n-1} \binom{m-1}{k+t-2s-1} \\
&= \binom{n-2}{k+t-2s-2} - \sum_{m=k+t-2s+1}^{n-1} \binom{m-2}{k+t-2s-1}.
\end{aligned}$$

Therefore,

$$\begin{aligned}
&r_{t,n,k,t-2s}^{(1)} - r_{t,n,k,t-2s}^{(2)} \\
&= r(M_{k-1,k-1+t-2s}^{(t,n-1)}) - r(M_{k-1,k-1+t-1-2(s-1)}^{(t-1,n-1)}) - \binom{n-1}{k+t-2s-1} \\
&= - \binom{n-2}{k+t-2s-1} - \sum_{m=k+t-2s+1}^{n-1} \binom{m-2}{k+t-2s-1} \leq 0.
\end{aligned} \tag{19}$$

Case 4.2.2: If $0 \leq (k-1) - s < \lfloor ((n-1) - t)/2 \rfloor$, then $0 \leq 1 \leq k-s \leq \lfloor (n-t-1)/2 \rfloor \leq$

$\lfloor (n-t)/2 \rfloor$ and

$$\begin{aligned}
& A_{t,n-1,k-1,s}^{(1)} - A_{t-1,n-1,k-1,s-1}^{(1)} = - \left(\binom{t-2k+s+2}{t+k-2s-2} - \binom{t-2k+s+2}{k-s-2} \right), \\
& A_{t,n-1,k-1,s}^{(2)} - A_{t-1,n-1,k-1,s-1}^{(2)} \\
&= \sum_{m=t-3s+2k-1}^{(n-1)-s} \left(\binom{m-1}{k+t-2s-2} - \binom{m-1}{k-s-2} \right) \\
&\quad - \sum_{m=t-3s+2k+1}^{n-s} \left(\binom{m-1}{k+t-2s-1} - \binom{m-1}{k-s-1} \right) \\
&= \left(\binom{t-2s+2k-2}{k+t-2s-2} - \binom{t-3d+2k-2}{k-s-2} \right) - \sum_{m=t-3s+2k}^{n-s-1} \left(\binom{m-1}{k+t-2s-1} - \binom{m-1}{k-s-1} \right), \\
& C_{t,n-1,k-1,s} - C_{t-1,n-1,k-1,s-1} \\
&= \sum_{m=n-s}^{n-1} \binom{m-1}{k+t-2s-2} - \sum_{m=n-s+1}^{n-1} \binom{m-1}{k+t-2s-1} \\
&= \binom{n-2}{k+t-2s-2} - \sum_{m=n-s}^{n-1} \binom{m-1}{k+t-2s-1}.
\end{aligned}$$

From above three terms, it is easy to get

$$\begin{aligned}
& r_{t,n,k,t-2s}^{(1)} - r_{t,n,k,t-2s}^{(2)} \\
&= r(M_{k-1,k-1+t-2s}^{(t,n-1)}) - r(M_{k-1,k-1+t-1-2(s-1)}^{(t-1,n-1)}) - \binom{n-1}{k-1+t-2s} \\
&= \sum_{i=1}^2 (A_{t,n-1,k-1,s}^{(i)} - A_{t-1,n-1,k-1,s-1}^{(i)}) + (C_{t,n-1,k-1,s} - C_{t-1,n-1,k-1,s-1}) - \binom{n-1}{k-1+t-2s} \\
&\leq 0.
\end{aligned}$$

Case 4.2.3: If $\lfloor ((n-1)-t)/2 \rfloor \leq (k-1)-s \leq (n-1)-t$, then $\lfloor (n-t)/2 \rfloor \leq \lfloor (n-t+1)/2 \rfloor \leq k-s \leq n-t$.

Since $B_{t,n-1,k-1,s} - B_{t-1,n-1,k-1,s-1} = 0$, we have,

$$\begin{aligned}
& r_{t,n,k,t-2s}^{(1)} - r_{t,n,k,t-2s}^{(2)} \\
&= r(M_{k-1,k-1+t-2s}^{(t,n-1)}) - r(M_{k-1,k-1+t-1-2(s-1)}^{(t-1,n-1)}) - \binom{n-1}{k-1+t-2s} \\
&= (B_{t,n-1,k-1,s} - B_{t-1,n-1,k-1,s-1}) + (C_{t,n-1,k-1,s} - C_{t-1,n-1,k-1,s-1}) \\
&= C_{t,n-1,k-1,s} - C_{t-1,n-1,k-1,s-1} \\
&= \binom{n-2}{k+t-2s-2} - \binom{n-1}{k-1+t-2s} - \sum_{m=n-s}^{n-1} \binom{m-1}{k+t-2s-1} \\
&\leq 0.
\end{aligned}$$

Case 4.2.4: If $(k-1) - s > (n-1) - t$, then $k-s > n-t$.

Same as Eq 19, we can verify $r_{t,n,k,t-2s}^{(1)} \leq r_{t,n,k,t-2s}^{(2)}$ in this case.

From above all, we have verified $r_{t,n,k,t-2s}^{(1)} \leq r_{t,n,k,t-2s}^{(2)}$ in all cases.

Step 4.3: verify that $r_{t,n,k,t-2s}^{(4)} \leq r_{t,n,k,t-2s}^{(2)}$.

Case 4.3.1: If $k-s < 0$, then $(k-1) - (s-1) < 0$.

Therefore,

$$\begin{aligned}
& r_{t,n,k,t-2s}^{(4)} - r_{t,n,k,t-2s}^{(2)} \\
&= r(M_{k,k+t-2s}^{(t,n-1)}) - r(M_{k-1,k-1+t-1-2(s-1)}^{(t-1,n-1)}) - \binom{n-1}{k} \\
&= \binom{n-1}{k} - \binom{n-1}{k-1} - \binom{n-1}{k} + (D_{t,n-1,k,s} - D_{t-1,n-1,k-1,s-1}) \\
&\leq \sum_{m=k+t-2s+1}^{n-1} \binom{m-1}{k+t-2s-1} - \sum_{m=k+t-2s+1}^{n-1} \binom{m-1}{k+t-2s-1} = 0.
\end{aligned} \tag{20}$$

Case 4.3.2: If $0 \leq k-s \leq \lfloor ((n-1)-t)/2 \rfloor$, then $0 \leq k-s \leq \lfloor (n-t-1)/2 \rfloor \leq \lfloor (n-t)/2 \rfloor$ and

$$\begin{aligned}
& A_{t,n-1,k,s}^{(1)} - A_{t-1,n-1,k-1,s-1}^{(1)} = 0, \\
& A_{t,n-1,k,s}^{(2)} - A_{t-1,n-1,k-1,s-1}^{(2)} = -\left(\binom{n-s-1}{k+t-2s-1} - \binom{n-s-1}{k-s-1} \right), \\
& C_{t,n-1,k,s} - C_{t-1,n-1,k-1,s-1} \\
&= \sum_{m=n-s}^{n-1} \binom{m-1}{k+t-2s-1} - \sum_{m=n-s+1}^{n-1} \binom{m-1}{k+t-2s-1} \\
&= \binom{n-s-1}{k+t-2s-1}.
\end{aligned}$$

Therefore,

$$\begin{aligned}
& r_{t,n,k,t-2s}^{(4)} - r_{t,n,k,t-2s}^{(2)} \\
&= r(M_{k,k+t-2s}^{(t,n-1)}) - r(M_{k-1,k-1+t-1-2(s-1)}^{(t-1,n-1)}) - \binom{n-1}{k} \\
&= -\binom{n-1}{k-1} + \sum_{i=1}^2 (A_{t,n-1,k,s}^{(i)} - A_{t-1,n-1,k-1,s-1}^{(i)}) + (C_{t,n-1,k,s} - C_{t-1,n-1,k-1,s-1}) \\
&= \binom{n-s-1}{k-s-1} - \binom{n-1}{k-1} \leq 0.
\end{aligned}$$

Case 4.3.3: If $\lfloor ((n-1)-t)/2 \rfloor < k-s \leq (n-1)-t$, then $\lfloor (n-t)/2 \rfloor \leq \lfloor (n-t+1)/2 \rfloor \leq k-s \leq n-t$.

We have

$$\begin{aligned}
& B_{t,n-1,k,s} - B_{t-1,n-1,k-1,s-1} \\
&= \sum_{a=0}^{n-t-k+s-2} \left(\binom{t-s+2a}{t-s+a-1} - \binom{t-s+2a}{a-1} \right) - \sum_{a=0}^{n-t-k+s-1} \left(\binom{t-s+2a}{t-s+a-1} - \binom{t-s+2a}{a-1} \right) \\
&\leq 0.
\end{aligned}$$

Therefore,

$$\begin{aligned}
& r_{t,n,k,t-2s}^{(4)} - r_{t,n,k,t-2s}^{(2)} \\
&= r(M_{k,k+t-2s}^{(t,n-1)}) - r(M_{k-1,k-1+t-1-2(s-1)}^{(t-1,n-1)}) - \binom{n-1}{k} \\
&= -\binom{n-1}{k-1} + (B_{t,n-1,k,s} - B_{t-1,n-1,k-1,s-1}) + (C_{t,n-1,k,s} - C_{t-1,n-1,k-1,s-1}) \\
&= \binom{n-s-1}{k-s-1} - \binom{n-1}{k-1} \leq 0.
\end{aligned}$$

Case 4.3.4: If $k-s > (n-1) - t$, then $k-s \geq n-t$.

Same as Eq 20, we can verify $r_{t,n,k,t-2s}^{(4)} \leq r_{t,n,k,t-2s}^{(2)}$ in this case.

From above all, we have verified $r_{t,n,k,t-2s}^{(4)} \leq r_{t,n,k,t-2s}^{(2)}$ in all cases.

Step 5: verify the base case of induction.

Consider when $t+n+k+s=2$ and show that (t,n,k,s) is valid. The only case is $t=1, n=1, k=0, s=0$ and, then, by Step 1, we can verify Eq 13 in this case.

By induction, we have finally proved Lemma 3.10. □

Appendix B: Proof of Lemma 3.11

We give the proof of Lemma 3.11 via Eq 10. Note that when $t=1$, $bw(M^{(1,n)}) = \sum_{m=0}^{n-1} \binom{m}{\lfloor m/2 \rfloor}$ [13].

If $n=2l+1$ for some integer l , then

$$\begin{aligned}
& \sum_{m=0}^{n-1} \binom{m}{\lfloor m/2 \rfloor} \\
&= \sum_{a=0}^l \binom{2a}{a} + \sum_{a=0}^{l-1} \binom{2a+1}{a} \\
&= \binom{2l+1}{l} + \sum_{a=0}^{l-1} \left(\binom{2a+1}{a} - \binom{2a+1}{a-1} \right) \\
&= \sum_{k=\lfloor (n-t)/2 \rfloor}^{\lfloor (n-t)/2 \rfloor + t-1} \binom{n}{k} + \sum_{a=0}^{\lfloor (n-t-1)/2 \rfloor} \left(\binom{t+2a}{t+a-1} - \binom{t+2a}{a-1} \right),
\end{aligned}$$

where the second equality can be proved by induction on l .

If $n = 2l$ for some integer l , then

$$\begin{aligned}
& \sum_{m=0}^{n-1} \binom{m}{\lfloor m/2 \rfloor} \\
&= \sum_{a=0}^{l-1} \binom{2a}{a} + \sum_{a=0}^{l-1} \binom{2a+1}{a} \\
&= \binom{2l}{l} + \sum_{a=0}^{l-1} \left(\binom{2a+1}{a} - \binom{2a+1}{a-1} \right) \\
&= \sum_{k=\lfloor (n-t)/2 \rfloor}^{\lfloor (n-t)/2 \rfloor + t-1} \binom{n}{k} + \sum_{a=0}^{\lfloor (n-t-1)/2 \rfloor} \left(\binom{t+2a}{t+a-1} - \binom{t+2a}{a-1} \right),
\end{aligned}$$

where the second equality can be proved by induction on l as well.

Therefore, when $t = 1$, Eq 14 holds. Then we intend to prove Lemma 3.11 by induction on t . Suppose Eq 14 holds for $t < T$, now consider when $t = T$.

First consider the situation when $k = \lfloor (n-t)/2 \rfloor$ and $p = t$. Then we have

$$\sum_{q=1}^{t-1} \binom{n}{k+q} + r(M_{k,k+t}^{(t,n)}) = \sum_{q=0}^{t-1} \binom{n}{k+q} + \sum_{a=0}^{\lfloor (n-t-1)/2 \rfloor} \left(\binom{t+2a}{t+a-1} - \binom{t+2a}{a-1} \right) \quad (21)$$

The result matches the RHS of Eq 14. In the following, we only need to prove other term is no more than this value.

For convenience, define $\tilde{r}(M_{k,k+p}^{(t,n)}) = \sum_{q=1}^{p-1} \binom{n}{k+q} + r(M_{k,k+p}^{(t,n)})$ when $1 \leq p \leq t$ and $\tilde{r}(M_{k,k}^{(t,n)}) = bw(M_{k,k}^{(t,n)})$. Actually, $\tilde{r}(M_{k,k+p}^{(t,n)})$ where $0 \leq k \leq k+p \leq n$ is exactly the maximum Manhattan distance from a nonzero element of $M_{k,k+p}^{(t,n)}$ in M to the main diagonal of the matrix M . There exists a consist expression for $\tilde{r}(M_{k,k+p}^{(t,n)})$ as following.

$$\tilde{r}(M_{k,k+p}^{(t,n)}) = \sum_{q=0}^{p-1} \binom{n}{k+q} - \binom{n}{k} + r(M_{k,k+p}^{(t,n)}).$$

Then, our purpose is to prove $\tilde{r}(M_{k,k+p}^{(t,n)})$ is no larger than Eq 21 for any integer k, p satisfying $0 \leq k \leq k+p \leq n$ and $0 \leq p \leq t$.

From definition, when $k = 0$, we have that $\tilde{r}(M_{k,k+p}^{(t,n)})$ reach its maximal when $p = t$. Similarly, when $k+p = n$, $\tilde{r}(M_{k,k+p}^{(t,n)})$ reach its maximal when $k = n-t$, that is, $p = t$. When $k > 0$ and $k+p < n$, we have $\tilde{r}(M_{k,k+p}^{(t,n)}) \leq \tilde{r}(M_{k-1,k+p+1}^{(t,n)})$ when $p+2 \leq t$. Hence, we only need to prove $\tilde{r}(M_{k,k+p}^{(t,n)})$ is no larger than Eq 21 for the following two cases:

- (1) $p = t - 1$,
- (2) $p = t$.

Case 1: $p = t - 1$. In this case, we have $\tilde{r}(M_{k,k+p}^{(t,n)}) = \tilde{r}(M_{k,k+p}^{(t-1,n)}) \leq bw(M^{(t-1,n)})$ by definition. Since the value of $bw(M^{(t-1,n)})$ can be calculated from Lemma 3.11 by induction, and it is not hard to verify the RHS of Eq. 14 is increasing with respect to t . Then we can reach our conclusion in this case.

Case 2: $p = t$.

If $k \leq \lfloor (n-t)/2 \rfloor$, then

$$\begin{aligned}
& \tilde{r}(M_{k,k+t}^{(t,n)}) - \tilde{r}(M_{k-1,k+t-1}^{(t,n)}) \\
&= \binom{n}{k+t-1} - \binom{n}{k} + \binom{n}{k} - \binom{n}{k-1} + \left(\binom{2k+t-2}{k+t-2} - \binom{2k+t-2}{k-2} \right) \\
& \quad + \sum_{m=t+1+2k}^{n-s} \left(\binom{m-1}{k+t-1} - \binom{m-1}{k-1} \right) - \sum_{m=t-1+2k}^{n-s} \left(\binom{m-1}{k+t-2} - \binom{m-1}{k-2} \right) \\
&= \binom{n}{k+t-1} - \binom{n}{k-1} + \sum_{m=t+2k}^{n-1} \left(\binom{m-1}{k+t-2} - \binom{m-1}{k-2} \right) - \left(\binom{n-1}{k+t-2} - \binom{n-1}{k-2} \right) \\
&= \binom{n-1}{k-t-1} - \binom{n-1}{k-1} + \sum_{m=t+2k}^{n-1} \left(\binom{m-1}{k+t-2} - \binom{m-1}{k-2} \right) \\
&\geq 0.
\end{aligned}$$

It shows that when $k \leq \lfloor (n-t)/2 \rfloor$, $\tilde{r}(M_{k,k+t}^{(t,n)})$ reaches its maximum at $k = \lfloor (n-t)/2 \rfloor$. If $k \geq \lfloor (n-t)/2 \rfloor$, then

$$\begin{aligned}
& \tilde{r}(M_{k,k+t}^{(t,n)}) - \tilde{r}(M_{k+1,k+t+1}^{(t,n)}) \\
&= \binom{n}{k+1} - \binom{n}{k+t} + \binom{n}{k} - \binom{n}{k+1} + \left(\binom{2n-t-2k-4}{n-k-3} - \binom{2n-t-2k-4}{n-t-k-3} \right) \\
&\geq 0.
\end{aligned}$$

Combining the above situations, we have proved that $\tilde{r}(M_{k,k+t}^{(t,n)})$ reaches its maximum at $k = \lfloor (n-t)/2 \rfloor$. That is, the maximum value of is $\tilde{r}(M_{k,k+t}^{(t,n)})$ as Eq 21.

From above all, we have proved Lemma 3.11. □