

# The Bayes Principle and Segal Axioms for $P(\phi)_2$ , with application to Periodic Covers

Jiasheng Lin

## Abstract

The goal of the present paper is to reconcile the so-called  $P(\phi)_2$  model from classical constructive quantum field theory (CQFT) with (metric version of) geometric and categorical axioms proposed by G. Segal [66] in the 90's, in showing that the  $P(\phi)_2$  model satisfies these axioms, appropriately adjusted. The approach is based on previous works of Dimock [16] and Pickrell [52], while we give a new proof to a key step using what we call “the Bayes principle” of conditional probabilities in the infinite dimensional setting. We also give a precise statement and full proof of the locality of the  $P(\phi)_2$  interaction. After that, we note that the so-called “transfer operator” associated to a cobordism has strictly positive kernel, this implies the operator has a spectral gap and allows to define a  $P(\phi)_2$  Gibbs state on non-compact periodic curved surfaces (namely, with discrete “time translations”) by analogy with 1-dimensional spin chains. This also implies asymptotic properties of the  $P(\phi)_2$  partition function on periodic covers of large degrees.

## Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
1.1	Example of 1D Spin Chain and its Transfer Operator	3
1.2	Main results and Comments	4
1.3	Notations	7
1.4	Acknowledgement	8
<b>2</b>	<b>Gaussian Fields on a Riemannian Manifold</b>	<b>8</b>
2.1	The Massive Gaussian Free Field	9
2.1.1	Definition and Representations	9
2.1.2	Essential Properties	10
2.2	Determinants	11
2.2.1	Zeta-regularized and Fredholm Determinants	11
2.2.2	Factorization Lemma	13
2.2.3	The Gluing Formula of Burghelea-Friedlander-Kappeler	13
2.3	Quadratic Perturbation = Radon-Nikodym Density	14
<b>3</b>	<b>Variants of Nelson’s Argument</b>	<b>16</b>
3.1	Regularizations	17
3.2	Integrability of Interaction and Regularization Independence	18
3.3	Integrability of the Exponential of Interaction	20
3.4	Change of Wick Ordering	21
<b>4</b>	<b>The Trace Operator and the Poisson Integral Operator</b>	<b>22</b>
4.1	Summary of Essential Properties	22
4.2	The Dirichlet-to-Neumann Operator and its Jumpy Version	24
4.3	Two Consequences of the Green-Stokes Formula	24
4.4	Induced Laws	26
4.5	First Comment on Reflection Positivity	27

arXiv:2403.12804v2 [math-ph] 3 Apr 2024

<b>5</b>	<b>Markov Property and Consequences</b>	<b>28</b>
5.1	Decompositions of Sobolev Spaces . . . . .	28
5.2	The Markov Stochastic Decomposition of GFF . . . . .	30
5.3	The Bayes Principle Applied to GFF . . . . .	31
5.4	Nelson's Markov Property and Reflection Positivity . . . . .	32
5.5	Locality of the $P(\phi)$ interaction and the Restricted GFF . . . . .	34
<b>6</b>	<b>Presenting the <math>P(\phi)_2</math> Model as a Segal Theory</b>	<b>36</b>
6.1	Description of Segal's Rules as Pertains to the Model . . . . .	36
6.2	The Hilbert Spaces . . . . .	38
6.3	Amplitudes = Schwartz Kernels . . . . .	39
6.4	Trace Axiom and its Consequences . . . . .	40
6.5	Bayes-Type Density Formulae and Gluing with Free Ends . . . . .	42
<b>7</b>	<b>Periodic Cover = Spin Chain</b>	<b>45</b>
7.1	Geometric setting . . . . .	45
7.2	Continued Introduction of Spin Chain Example . . . . .	45
7.3	Perron-Frobenius Property and Gibbs State . . . . .	46
7.4	Asymptotic of Partition Function . . . . .	48
<b>A</b>	<b>Functional Analysis</b>	<b>49</b>
A.1	Tensor Product, Fock Space, Ideals . . . . .	49
A.2	Hilbert-Schmidt Operators on $L^2$ . . . . .	50
<b>B</b>	<b>Gaussian Analysis</b>	<b>51</b>
B.1	Basic Definitions and Spaces . . . . .	51
B.2	$Q$ -spaces and Their Equivalence . . . . .	52
B.3	Measurable Linear (Itô-Wiener) Extensions . . . . .	53
B.4	Wiener Chaos and Wick's Theorem . . . . .	54
B.5	Itô-Wiener-Segal Isomorphism . . . . .	55
<b>C</b>	<b>Global Analysis</b>	<b>56</b>
C.1	Sobolev Spaces over Domains . . . . .	56
C.2	Symbol Convergence Lemma and Heat Kernel . . . . .	57
	<b>References</b>	<b>59</b>

# 1 Introduction

In the classical approach to quantum field theory (QFT), a central role is given to the representation of the Lorentz group as one can easily recognize if one looks at the Wightman axioms which were among the first attempt to axiomatize QFT. It was later realized by Feynman, Symanzik [69], Nelson [49] in the 60's that one could describe most objects of QFT using the functional integral, in imaginary time, which started the Euclidean approach to QFT. This allows to recover many computations and predictions of interest in quantum field theory.

In the 80's, motivated by works of Atiyah and Witten on the relation of QFT with geometry and topology, there was an attempt to give an axiomatic definition of QFT based on geometry that would capture the main properties of the functional integral representation. The two important notions at the foundation of QFT are the concept of **locality** and **unitarity**. The first notion of locality is intuitively captured by the functional integral representation whereas the second notion will be encoded in certain symmetries of our manifolds under reflection called reflection positivity. The notion of reflection positivity plays a central role in constructive quantum field theory since the works of Osterwalder and Schrader, Glimm and Jaffe and also plays an important role in statistical physics.

The geometric axioms for QFT are due to Atiyah, Segal [66] and more recently Kontsevich and Segal [40]: instead of viewing QFT as representation of some space-time symmetry group, we rather view them loosely as linear representations of some geometric bordism category. In the following subsection we give the example of the 1 dimensional spin chain which contains the key phenomenology of the QFT model we will describe in the present work. It can be understood as linear representations of some discrete geometric bordism category and we will also explain why our spin chain admits a unique Gibbs measure (in the sense of Ruelle [63]) and the shift map acting on the chain is exponentially mixing for the Gibbs measure.

## 1.1 Example of 1D Spin Chain and its Transfer Operator

Our example illustrates the gluing properties by the transfer operator for discrete spin systems. Consider  $\mathbb{Z}_N := \mathbb{Z}/N\mathbb{Z}$  as a 1-dimensional chain of size  $N$ , consider the space  $\mathbb{R}^N$  now as the space of maps  $\mathbb{Z}_N \rightarrow \mathbb{R}$ , namely a discrete path space. The lattice site is denoted by  $i \in \{1, \dots, N\}$  and call  $\sigma \in \mathbb{R}^N$  a spin configuration, whose value at the site  $i$  reads  $\sigma(i) \in \mathbb{R}$ . We impose periodic boundary condition, which means  $\sigma(N+1) \equiv \sigma(1)$ . Thus the chain could be considered as circular. We are given an action functional on the configuration space  $\mathbb{R}^N$  that reads

$$S_N(\sigma) \stackrel{\text{def}}{=} \sum_{i=1}^N |\sigma(i+1) - \sigma(i)|^2 + \sum_{i=1}^N P(\sigma(i)) \quad (1.1)$$

where  $P$  is a polynomial bounded from below, the interaction is nearest neighbour. Given a configuration  $\sigma \in \mathbb{R}^N$ ,  $S_N(\sigma)$  may be thought of as its ‘‘energy’’ and the statistical behaviour of the system is described by the probability measure called Gibbs measure,

$$d\mu_{P(\sigma)}(\sigma) \stackrel{\text{def}}{=} \frac{1}{\mathcal{Z}(N)} e^{-S_N(\sigma)} d^N \sigma, \quad \text{with} \quad \mathcal{Z}(N) = \int_{\mathbb{R}^N} e^{-S_N(\sigma)} d^N \sigma \quad (1.2)$$

called the partition function of the system.

Now we would like to express this partition function  $\mathcal{Z}(N)$  in terms of elementary building blocks. The main idea is to slice the action functional as

$$S_N(\sigma) = \sum_{i=1}^N \left[ |\sigma(i+1) - \sigma(i)|^2 + \frac{1}{2}(P(\sigma(i+1)) + P(\sigma(i))) \right]$$

so exponentiating gives

$$\exp(-S_N(\sigma)) = \prod_{i=1}^N K(\sigma(i+1), \sigma(i)) \quad (1.3)$$

where

$$K(x, y) = e^{-|x-y|^2 - \frac{1}{2}(P(x)+P(y))} \quad (1.4)$$

which is the Schwartz kernel of an operator on  $L^2(\mathbb{R})$  that is smoothing.

**Definition 1.1.** Define the transfer operator  $T$  to be exactly the operator with kernel  $K(x, y)$ , that is,

$$(TF)(x) = \int K(x, y)F(y)dy, \quad (1.5)$$

for any function(al)  $F \in L^2(\mathbb{R})$ .

Then we see immediately from (1.3) that

$$\mathcal{Z}(N) = \int \prod_{i=1}^N [K(\sigma(i+1), \sigma(i))d\sigma(i)] = \text{tr}_{L^2(\mathbb{R})}(T^N), \quad (1.6)$$

remembering that  $\sigma(N+1) \equiv \sigma(1)$ , since for a smoothing operator  $A : L^2(\mathbb{R}) \rightarrow \mathcal{S}(\mathbb{R})$  we have  $\text{tr}_{L^2(\mathbb{R})}(A) = \int K_A(x, x)dx$  with  $K_A$  being the integral kernel.

More generally we would like to express the kernel of  $T^N$  in terms of  $\exp(-S(\sigma))$  using the relation (1.3). This means instead of letting the boundary condition be periodic we let  $\sigma(1) = \sigma_{\text{in}}$ ,  $\sigma(N+1) = \sigma_{\text{out}}$  given two boundary conditions  $(\sigma_{\text{in}}, \sigma_{\text{out}}) \in \mathbb{R}^2$ . Then the kernel  $K_N$  of  $T^N$  is

$$K_N(\sigma_{\text{out}}, \sigma_{\text{in}}) = \int K(\sigma_{\text{out}}, \sigma(N)) \cdots K(\sigma(2), \sigma_{\text{in}}) \prod_{i=2}^N d\sigma(i) \quad (1.7)$$

$$= \int_{\mathbb{R}^{N-1}} e^{-S_N(\sigma|\sigma_{\text{in}}, \sigma_{\text{out}})} \prod_{i=2}^N d\sigma(i), \quad (1.8)$$

with the conditioned interaction  $S_N(\sigma|\sigma_{\text{in}}, \sigma_{\text{out}})$  defined as

$$S_N(\sigma|\sigma_{\text{in}}, \sigma_{\text{out}}) \stackrel{\text{def}}{=} \sum_{i=1}^N |\sigma(i+1) - \sigma(i)|^2 + \sum_{i=2}^N P(\sigma(i)) + \frac{1}{2}(P(\sigma_{\text{in}}) + P(\sigma_{\text{out}})), \quad (1.9)$$

with  $\sigma(1) \equiv \sigma_{\text{in}}$  and  $\sigma(N+1) \equiv \sigma_{\text{out}}$ . Now if  $N_1, N_2$  are two integers and we define the kernels  $K_{N_1}, K_{N_2}$  using (1.8) and (1.9) with  $N$  replaced respectively by  $N_1, N_2$ , then it follows “trivially” from the composition property  $T^{N_2} \circ T^{N_1} = T^{N_2+N_1}$  that

$$K_{N_2+N_1}(\sigma_{\text{out}}, \sigma_{\text{in}}) = \int K_{N_2}(\sigma_{\text{out}}, \sigma)K_{N_1}(\sigma, \sigma_{\text{in}})d\sigma. \quad (1.10)$$

However, such a relation becomes remarkable (rather than trivial) if we do not have the “unit” transfer operator  $T$  to start with; that is, if we do not have (1.7) but define  $K_{N_1}$  and  $K_{N_2}$  *directly* with an expression of the form (1.8) and (1.9). This corresponds to the idea of a **path integral** in quantum mechanics and quantum field theory. Alternatively one could consider the Gibbs measure (1.2) and take  $K_N$  as the *transition probability* of a certain stochastic process. Then (1.10) is the Chapman-Kolmogorov equation which relies heavily on the fact that the underlying process is **Markovian**. In a sense, for both interpretations a crucial condition is that the interaction  $S(\sigma)$  be **local**; that is, very roughly speaking, if one chops the sites  $[1, N_1 + N_2] := \{1, 2, \dots, N_1 + N_2\}$  into  $[1, N_1] \sqcup [N_1 + 1, N_1 + N_2]$  then  $S_{[1, N_1 + N_2]}(\sigma) \approx S_{[1, N_1]}(\sigma|_{[1, N_1]}) + S_{[N_1 + 1, N_1 + N_2]}(\sigma|_{[N_1 + 1, N_1 + N_2]})$ .

The main result of this article concerns a 2-dimensional “continuum” version of this story where lattice sites are replaced by the continuum of points on a 2D surface (considered as space-time) and a configuration is replaced by a distribution. See the section below for a more precise description. In the final section, we also show that when the space-time admits a periodic translation symmetry then a more precise analogy with the spin chain described above can be restored, in particular, there exists a **Gibbs state** in the thermodynamic limit. Further discussion of the above example continues in the separate introduction to this latter part in section 7.2.

## 1.2 Main results and Comments

**Main results.** Let  $\Omega$  be a Riemannian surface whose boundary has two components  $\partial\Omega = \Sigma_{\text{in}} \sqcup \Sigma_{\text{out}}$ . We will define probability measures  $\mu_{\text{in}}, \mu_{\text{out}}$  respectively on the spaces of distributions  $\mathcal{D}'(\Sigma_{\text{in}}), \mathcal{D}'(\Sigma_{\text{out}})$  and put  $\mathcal{H}_{\text{in}} := L^2(\mathcal{D}'(\Sigma_{\text{in}}), \mu_{\text{in}})$ ,  $\mathcal{H}_{\text{out}} = L^2(\mathcal{D}'(\Sigma_{\text{out}}), \mu_{\text{out}})$  which play the role of  $L^2(\mathbb{R})$  above. We then define the “transfer operator” associated to  $\Omega$  to be given by the integral kernel which is formally

$$\mathcal{A}_{\Omega}(\varphi_{\text{in}}, \varphi_{\text{out}}) = \int_{\{\phi|\partial\Omega=(\varphi_{\text{in}}, \varphi_{\text{out}})\}} e^{-\int_{\Omega} \frac{1}{2}(|\nabla\phi|_g^2 + m^2\phi^2) + P(\phi(x))dV_{\Omega}(x)} [\mathcal{L}\phi], \quad (1.11)$$

where the formal integration is over the space of all distributions subject to the boundary condition  $\phi|_{\Sigma_{\text{in}}} = \varphi_{\text{in}}$  and  $\phi|_{\Sigma_{\text{out}}} = \varphi_{\text{out}}$ , with respect to the non-existent Lebesgue measure  $[\mathcal{L}\phi]$  on the space of such distributions. Here  $P(\phi)$  is a polynomial bounded below just like the one considered in (1.1). If  $\Omega_1, \Omega_2$  are two Riemannian surfaces of the kind as above where the “out” boundary of  $\Omega_1$  is isometric to the “in” boundary of  $\Omega_2$  via an isometry  $\rho$ , then one can glue them along  $\rho$  and obtain the surface  $\Omega_2 \cup_{\rho} \Omega_1$ . This article then seeks to prove the analogue of (1.7) for operators defined by (1.11) in such a situation, after making rigorous sense of the formula which is in fact a large part of the work. The main development is carried out in section 6 and the proof is completed in subsection 6.5.

After that, we point out that the operator  $U_{\Omega}$  defined by (1.11) has the Perron-Frobenius property, namely it has strictly positive kernel (this could already be seen from the above heuristic expression). This has the consequence that  $U_{\Omega}$  in fact has a spectral gap and together with the composition property obtained above implies that the  $P(\phi)_2$  theory on periodic space-times behaves very similarly to the spin-chain example discussed above. In particular, we are able to construct a  $P(\phi)_2$  Gibbs state on some periodic infinite volume surface in perfect analogy with the 1D spin chain, we refer to section 7.3 for further details.

**Concrete illustration.** For a concrete illustration of our results, we state an informal theorem which is a consequence of the Segal gluing of  $P(\phi)_2$  amplitudes developed in the present work.

**Theorem** (asymptotics of the partition function on large degree periodic covers). *Let  $(M, g)$  denotes a compact Riemannian surface,  $M_{\infty} \rightarrow M$  a Riemannian  $\mathbb{Z}$ -cover of  $M$  and  $\gamma$  the generator of the deck group. We can think of  $M_{\infty}$  as the infinite composition of some given cobordism  $\Omega$ , which serves as the “fundamental domain” of the deck transformations on  $M_{\infty}$ . For every  $N \in \mathbb{N}$ , denote by  $M_N := M_{\infty}/\gamma^N$  the cyclic cover of degree  $N$  of  $M$  and  $\Delta_N$  is the Laplace-Beltrami operator on  $M_N$ . We define the partition function of the  $P(\phi)_2$  theory on  $M_N$  heuristically as*

$$Z_N \stackrel{\text{heu}}{=} \int_{\mathcal{D}'(M_N)} e^{-\int_{M_N} P(\phi(x))dV(x)} e^{-\frac{1}{2} \int_{M_N} (\|\nabla\phi\|_g^2 + m^2\phi^2)dV} [\mathcal{L}\phi] \quad (1.12)$$

where  $dV$  is the Riemannian area form,  $\mathcal{D}'(M_N)$  is the space of real distributions on  $M_N$  and  $[\mathcal{L}\phi]$  is the non-existent Lebesgue measure on  $\mathcal{D}'(M_N)$ .

Then the renormalized sequence of free energies  $\frac{1}{N} \log(Z_N)$  has a limit  $\lambda_0$  when  $N \rightarrow +\infty$ , moreover this limit  $\lambda_0$  can be interpreted as the leading eigenvalue of some transfer operator  $U_{\Omega}$  which is the quantization of the cobordism  $\Omega$  mentioned above.

Rigorous sense of the expression (1.12) will be made in sections 2 and 3.

**Remark 1.1.** The above theorem is an interacting QFT version of a result of Naud on the asymptotics of zeta determinants on large degree random covers of compact surfaces [46], the main difference is that he treats random covers whereas our sequence of covers is deterministic and he only deals with the partition function of free fields whereas we treat the interacting case.

**Other parts.** In sections 2 and 3 we retrace the classical rigorous construction due to Nelson of the integrand of (1.11), and adapt the argument for showing *locality* of the interaction in subsection 5.5. Section 4 aims to derive the behavior of a Gaussian field under the trace map (restriction to hypersurface); these results are classical but our treatment still gives a somewhat new perspective based on elementary adjoint relations of various geometric operators (see subsection 4.3). Section 5 discusses the Markov property that culminates in the *Bayes principle* (see below) for the GFF and proving locality of the  $P(\phi)$  interaction. In between, we give comments on *reflection positivity* which is not strictly related to the aim of the article but, the author believes, integrates organically into the discussion (subsections 4.5 and 5.4).

**Comments.** Firstly we point out that the exact same problem has been treated some 15 years ago in Pickrell [52], and a partial treatment (trace axiom for free field) also appears in Dimock [16]. The present article aims to offer a new proof (to a core proposition, see below) based on the following simple idea. We also tie up many loose ends around this problem; especially we precisely state and prove the *locality* of the  $P(\phi)$  interaction, which is widely believed to lie behind a property such as Segal’s gluing rule and, as we shall see in our case in section 5.5, lies behind the very possibility of defining (1.11) over a domain with boundary. Indeed, since the definition of the  $P(\phi)$  interaction requires *renormalization*, the verification of Segal’s rules provides an important **testing ground** that the renormalization procedures can actually be made local, which is in general a deep question. See also remark 3.1.

The idea (observation) is that Segal’s gluing axioms for the scalar QFT actually follow from the Bayes principle applied to the functional measure of the QFT. What we mean by the “Bayes principle” here is the following, in

its simplest form: suppose two real random variables  $X$  and  $Y$  have joint probability law  $\mathbb{P}_{(X,Y)}$ ; then we have two equal expressions for their joint density  $p(x, y) = (\mathrm{d}\mathbb{P}_{(X,Y)}/\mathrm{d}\mathcal{L}_{\mathbb{R}}^{\otimes 2})(x, y)$ , namely

$$p(x, y) = \frac{\mathrm{d}(\pi_x^* \mathbb{P}_{(X,Y)})}{\mathrm{d}\mathcal{L}_{\mathbb{R}}}(x) \frac{\mathrm{d}\mathbb{P}_{Y|X=x}}{\mathrm{d}\mathcal{L}_{\mathbb{R}}}(y) = \frac{\mathrm{d}(\pi_y^* \mathbb{P}_{(X,Y)})}{\mathrm{d}\mathcal{L}_{\mathbb{R}}}(y) \frac{\mathrm{d}\mathbb{P}_{X|Y=y}}{\mathrm{d}\mathcal{L}_{\mathbb{R}}}(x), \quad (1.13)$$

where  $\pi_x, \pi_y$  are respectively the projections onto the  $x$ - and  $y$ -axes,  $\mathbb{P}_{Y|X=x}$  denotes the conditional law of  $Y$  knowing “ $X = x$ ”, and vice versa for  $\mathbb{P}_{X|Y=y}$  (more rigorously these are expressed using *transition kernels*). In other words, one could evaluate  $p(x, y)$  by *conditioning* in two alternative ways: on  $X$  or on  $Y$ . In what follows we put forward a version of (1.13) for the Gaussian Free Field (GFF) measure on  $\mathcal{D}'(M)$  for  $M$  a closed Riemannian surface, where the projections  $\pi^{x,y}$  are replaced by trace maps (restrictions) onto embedded Riemannian circles. By the spacial Markov property of the GFF, the expressions for the conditional laws simplify nicely. Since the transition amplitudes (1.11) are given by the square roots of the densities of the trace-induced measures with respect to fixed background Gaussian functional measures on the circles (corresponding to  $p(x, y)$  above), we obtain a proof of Segal’s gluing axioms for the GFF and, by the *locality* of the  $P(\phi)$  interaction, the result extends immediately to the interacting case for this interaction.

Presumably in general, if one has a *local* interaction such that the QFT Gibbs measure (2.1) is well-defined and absolutely continuous with respect to the free field measure, then Segal’s rules for the interacting theory hold based on that of the free theory. See, for example, Guillarmou, Kupiainen, Rhodes and Vargas [28] for the case of the Liouville theory (which is conformal). In Pickrell [52], the argument for free field (presumably) used composition formulae in the *oscillator semigroup*<sup>1</sup>, and proceeded by analogy with the finite dimensional case (see Howe [34]). In [28] there is a counterpart of the treatment based on computations of Dirichlet forms. The proof here is different from both [28] and [52], which is geometric and in fact conceptually transparent. More precisely, this is proposition 6.9 below which corresponds to proposition 3 on page 16 of [52] and (the second part of)<sup>2</sup> lemma 5.3 on page 30 of [28].

An important ingredient taken from (inspired by) Pickrell [52] is the definition of the amplitude as the square root of the Radon-Nikodym density comparing a trace-induced measure to a fixed background Gaussian measure. This hints at an infinite-dimensional analogue of the notion of “half-densities”, and is actually what [52] considered. This latter notion has the advantage of being intrinsic. But due to insufficient literature discussing half-densities in the infinite-dimensional setting<sup>3</sup>, we adopt here the equivalent approach of fixing a background measure. Indeed, there is only one correct amplitude to consider, prescribed by physics, and the reader would see that the expression we obtain from this definition agrees, for example, with the one of [28] in its form and exactly for the free field. In fact, this definition of the amplitude “derives” naturally from a special case of Segal’s trace axiom (see section 6.3).

**Remark 1.2.** We consider real scalar fields, namely a field configuration  $\phi$  (on  $M$ ) is a *real* distribution (on  $M$ ). In yet other words, this means if we pair  $\phi$  with a potentially complex test function  $f$ , then

$$\overline{\phi(f)} = \phi(\bar{f}), \quad (1.14)$$

the bar denoting complex conjugation. Without loss of generality, all function spaces are assumed real unless otherwise stated.

**Remark 1.3** (role of symmetries). One primary aim of the present work is to reconcile the model (2.1) with the axiomatics proposed by G. Segal [66]. However, a significant difference between our model and the theories [66] principally tries to target is that we do not have and consider symmetries beyond the simplest one of reflecting across a hypersurface. In particular the model (2.1) will not be conformally invariant, and not even the free case ( $P = 0$ ) as  $m > 0$  and we do not include any “conformal anomaly” accounting effects of a conformal change in the metric. A linear term involving the Gauss curvature could have been included in the exponential of (1.11) — our situation corresponds to the case of “minimal coupling”, see discussion of Jaffe and Ritter [37] around equation (7).

In fact, we lack a basic symmetry which is present in the Euclidean ( $\mathbb{R}^2$ ) case of the same model (2.1): since we deal with general compact surfaces, there are no distinguished “time translations” (in the form of a generating Killing vector field). In the Euclidean case, the “canonical” time direction helps to fix a definitive Hilbert space

<sup>1</sup>this is a semigroup consisting of operators with *Gaussian kernels*, and the reader could see that (1.11) with  $P = 0$  is analogous to such a kernel. The author is however not very familiar with this object and refers the reader to Howe [34] and Folland [22] chapter 5.

<sup>2</sup>the first part corresponds to a combination of corollary 2.15 and proposition 4.9 of this article, the first of which is largely a known result concerning quadratic perturbation of Gaussian fields (see references in section 2.3).

<sup>3</sup>D. Pickrell confirmed with the author that he did not know of a reference which discuss infinite-dimensional half-densities other than the appendix of [52]. The author agrees with him that this notion might be a helpful one worthy of a detailed development.



of quantum states, and eventually allows one to obtain an explicit expression of the Hamiltonian operator generating time evolution of the theory. This would also be the case if  $\Omega$  is a cylinder equipped with cylindrical metric, for example, but for general  $\Omega$  the notion of a “Hamiltonian operator” is not very well-defined. It is nevertheless interesting to think of a definition perhaps in terms of categorical limits as in Kontsevich and Segal [40] section 3. Basic constructions of a Euclidean (Riemannian) field theory on curved space-times equipped with a time translation have been considered in Jaffe and Ritter [38].

Last but not least, in the last section 7 of this article we explore a crucial property of the operator  $U_\Omega$  representing the “evolution” across the cobordism  $\Omega$  (more suggestively it could be written as  $e^{-[\Omega]H}$ ), namely that it is Perron-Frobenius. This would imply that it nevertheless has many properties in common with an actual time evolution (i.e.  $e^{-tH}$ ), especially it has a spectral gap.

**Remark 1.4.** In hindsight, the author feels that a proof following (1.13) is a natural one regarding the intuition behind the path integral as “summing over histories”, and (1.13) simply says that such a sum “conditioned” on different intermediate points in history<sup>4</sup> produce equivalent results (the proof of (1.13) itself is trivial!). However, a still more ideal version of the proof would perhaps be one that puts the free field and interacting field on equal footing, where Segal’s axioms follow from a generalized version of “Nelson’s axioms” as explained on section IV.1 of Simon [71], based on the Markov property and uses a “nonlinear version” of the transition operator  $\mathcal{M}_{M,2}^1$  (see section 5.3).

## 1.3 Notations

In this paper, unless stated otherwise,

“ $\stackrel{\text{heu.}}{=}$ ” means “heuristically equal to”;

$(M, g)$  = closed Riemannian surface with metric  $g$ ;

$(\Omega, g)$  = compact Riemannian surface with metric  $g$ , boundary  $\partial\Omega$ , seen as isometrically embedded in  $M$ , the induced metric on  $\partial\Omega$  is still denoted  $g$ ;

$\Sigma$  = disjoint union of Riemannian circles (of different radii), usually  $\Sigma = \partial\Omega$ ; hypersurface in  $M$ ;

$\Omega^\circ$  = interior of  $\Omega$ ;

$m$  = mass parameter,  $m > 0$ , fixed throughout paper;

$\mathcal{D}'(M), \mathcal{D}'(\Omega^\circ)$  = real distributions on  $M$  and  $\Omega^\circ$ ;

$C^\infty(M), C_c^\infty(\Omega^\circ)$  = smooth functions on  $M$ , smooth compact support functions on  $\Omega^\circ$ ;

$\Delta$  = Laplacian on  $(M, g)$ ,  $\Delta f := -\operatorname{div}(\nabla f)$ , therefore it is defined to be nonnegative, the same applies below;

$\Delta_\Sigma$  = Laplacian on  $\Sigma$ ;

$\Delta_{\Omega,D}$  = Laplacian on  $\Omega$  with (zero) Dirichlet boundary condition (see also remark 4.1);

$\mathbf{D}_\Sigma = (\Delta_\Sigma + m^2)^{1/2}$ ;

$\tau_\Sigma$  = trace operator (restriction) onto the hypersurface  $\Sigma$ ;

$(Q, \mathcal{O})$  = general probability sample space with  $\sigma$ -algebra  $\mathcal{O}$ ;

$C$  = general positive self-adjoint elliptic pseudodifferential operator on  $M, \Omega, \Sigma$ , which is Hilbert-Schmidt on the corresponding  $L^2$  spaces;

$\mu_C^{\text{cond}} = \mu_{C^{-1}}^{\text{cond}} =$  Gaussian measure on  $\mathcal{D}'(M), \mathcal{D}'(\Omega^\circ)$ , or  $\mathcal{D}'(\Sigma)$ , equipped with their Fréchet Borel  $\sigma$ -algebra, with covariance  $\mathbb{E}_C[\phi(f)\phi(h)] = \langle f, Ch \rangle_{L^2}$ , under some conditions;

$\mu_{\text{GFF}}^M =$  massive GFF measure on  $\mathcal{D}'(M)$ ;  $\mu_{\text{GFF}}^{\Omega,D} =$  massive Dirichlet GFF measure on  $\mathcal{D}'(\Omega^\circ)$ ;

$\mathbb{E}_B^A =$  expectation under  $\mu_B^A$ ;

$\phi(f) =$  the random variable  $\phi \mapsto \langle \phi, f \rangle_{L^2}$  indexed by  $f \in C^\infty(M), C_c^\infty(\Omega^\circ)$  or  $C^\infty(\Sigma)$ ;

$W^s(M) =$  the  $L^2$  Sobolev space on  $M$ , with inner product  $\langle -, - \rangle_{W^s(M)} := \langle -, (\Delta + m^2)^s - \rangle_{L^2}$ ;

$W_A^s(M), W_U^s(M), W^s(U) =$  see appendix C.1;

$\Psi^r(M), \Psi^r(\Sigma) =$  pseudodifferential operators ( $\Psi$ DOs) on  $M$  or  $\Sigma$  with order  $r$ ;

$[\mathcal{L}\phi] =$  hypothetical Lebesgue measure on a space of distributions;

$\|\cdot\|_{\text{tr}}, \|\cdot\|_{\text{HS}}, \|\cdot\|_{\mathcal{J}_p}, \|\cdot\|_{L^2} =$  trace norm, Hilbert-Schmidt norm,  $\mathcal{J}_p$  ideal norm (see appendix A.1), operator norm acting on  $L^2$  or  $L^2$ -norm on function;

$\text{PI}_\Omega^{\Sigma,B} =$  Poisson integral operator extending from  $\Sigma$  to  $\Omega$ , with boundary conditions  $B$  imposed on boundary components other than  $\Sigma$  (see also remark 4.1);

$\text{DN}_\Omega^{\Sigma,B} =$  Dirichlet-to-Neumann operator or the jumpy version defined using  $\text{PI}_\Omega^{\Sigma,B}$  (jumpy version is understood when  $\Sigma$  is in the interior of  $\Omega$  rather than a boundary component);

$\mu_{\text{DN}}^{\Sigma,\Omega,B} =$  Gaussian measure on  $\mathcal{D}'(\Sigma)$  with covariance  $(\text{DN}_\Omega^{\Sigma,B})^{-1}$ ;

$\mu_{2\text{DN}}^{\partial\Omega,\Omega} =$  Gaussian measure on  $\mathcal{D}'(\partial\Omega)$  with covariance  $\frac{1}{2}(\text{DN}_\Omega^{\partial\Omega})^{-1}$ ;

<sup>4</sup>In our case, history is circular!

$\mu_{2\mathbf{D}}^\Sigma =$  Gaussian measure on  $\mathcal{D}'(\Sigma)$  with covariance  $\frac{1}{2}(\mathbf{D}_\Sigma)^{-1}$ ;  
 $\mathcal{M}_{M,2}^1 = \tau_{\Sigma_2} \text{PI}_M^{\Sigma_1}$  is the transition operator defined in section 5.3.

## 1.4 Acknowledgement

The author thanks foremostly his thesis advisor Nguyen-Viet Dang for raising attention to the work of Pickrell [52] which results in this article and many discussions, especially in suggesting the use of the regulator in Dyatlov and Zworski [19] and pointing out the wonderful analogy with spin chain treated in the final section. He also thanks Colin Guillarmou and Tat-Dat Tô for helpful discussions at an early stage of the project, Douglas Pickrell for giving valuable comments on a preliminary version of the work, and Sylvie Paycha and Bin Zhang for providing related reference and interesting discussion following a talk given by the author on this work. Finally, he thanks his laboratory IMJ-PRG for supporting his research.

## 2 Gaussian Fields on a Riemannian Manifold

**In this and the next section** we define rigorously a probability measure on  $\mathcal{D}'(M)$ , as well as certain variants on  $\mathcal{D}'(\Omega^\circ)$ , which heuristically bears the form

$$d\mu_P(\phi) \stackrel{\text{heu}}{=} \frac{1}{Z} \overbrace{e^{-\int_M P(\phi(x))dV_M(x)}}^A \underbrace{e^{-\frac{1}{2}\int_M (|\nabla\phi|_g^2 + m^2\phi^2)dV_M}}_B [\mathcal{L}\phi], \quad (2.1)$$

where  $P$  is a polynomial bounded from below, such as  $P(\phi) = \phi^4 - \phi^2$ ,  $[\mathcal{L}\phi]$  denotes the nonexistent Lebesgue measure on  $\mathcal{D}'(M)$  and

$$Z_M \stackrel{\text{heu}}{=} \int_{\mathcal{D}'(M)} e^{-\int_M P(\phi(x))dV_M(x)} e^{-\frac{1}{2}\int_M (|\nabla\phi|_g^2 + m^2\phi^2)dV_M} [\mathcal{L}\phi] \quad (2.2)$$

is the normalization factor, also called the **partition function**.

The idea is that while  $[\mathcal{L}\phi]$  is nonexistent, together with the factor  $\exp(-\frac{1}{2}\int_M (|\nabla\phi|_g^2 + m^2\phi^2)dV_M)$  the expression  $B$  can be given a rigorous meaning as a Gaussian probability measure scaled by a (finite) volume constant, and the measure (2.1) can be constructed if, after defining part  $A$  rigorously, one proves that it is  $L^1$  with respect to the measure  $B$ . This construction of (2.1) was done on  $\mathbb{R}^2$  essentially by Edward Nelson [48] and the argument is already a classic, very well-understood (see Simon [71], Glimm and Jaffe [26], Dimock [17] chapter 13 or Hairer [31] section 9). We will build on this method.

For us, the argument must be carefully adapted to show *locality* of the  $P(\phi)$  interaction, in the curved surface setting (see remark 3.1). Leaving details to section 3, we stress here the main theme of the subject: since powers of a distribution is generally not defined, **renormalization** is needed, and the polynomial  $P(\phi)$  must in fact be replaced by the so-called **Wick-ordered polynomial**  $:P(\phi):$  for  $A$  to exist.

**In this section** we construct the measure  $B$ . This construction is a classical one in probability theory, usually called the **Gaussian free field**. Here we collect some essential concepts and results, and refer, for example, to Sheffield [67] and Powell and Werner [54] for more information.

The starting point is that for a real positive symmetric  $N \times N$  matrix  $C^{-1}$  on  $\mathbb{R}^N$ , one has

$$\int_{\mathbb{R}^N} e^{-\frac{1}{2}\langle \mathbf{x}, C^{-1}\mathbf{x} \rangle} d\mathcal{L}^N(\mathbf{x}) = \frac{(2\pi)^{N/2}}{(\det C^{-1})^{1/2}} \quad (2.3)$$

and the measure

$$(\det C^{-1})^{1/2} e^{-\frac{1}{2}\langle \mathbf{x}, C^{-1}\mathbf{x} \rangle} d\mathcal{L}_1^N \quad (2.4)$$

is a Gaussian measure with covariance matrix  $C$  (in the standard basis), where we absorb  $(2\pi)^{-\frac{1}{2}}$  into the Lebesgue measure. Thus we should define

$$e^{-\frac{1}{2}\int_M (|\nabla\phi|_g^2 + m^2\phi^2)dV_M} [\mathcal{L}\phi] \stackrel{\text{heu}}{=} \text{“det”}(\Delta + m^2)^{-\frac{1}{2}} d\mu_{\text{GFF}}^M(\phi) \quad (2.5)$$

where  $\mu_{\text{GFF}}^M$  is a Gaussian measure on  $\mathcal{D}'(M)$  with covariance *operator*  $(\Delta + m^2)^{-1}$ , and “det” is an infinite dimensional generalization of the determinant of a matrix, called the  $\zeta$ -regularized determinant. The Gaussian measure and the determinant are two issues to be treated separately (in sections 2.1 and 2.2).



## 2.1 The Massive Gaussian Free Field

### 2.1.1 Definition and Representations

Let  $(M, g)$  be a closed Riemannian manifold of dimension  $d$  with metric  $g$ , and  $\Omega \subset M$  an open domain with smooth boundary  $\partial\Omega$ , both equipped with the metric induced from  $g$  (same notation). Fix  $m > 0$  as the mass parameter.

**Definition 2.1.** The massive Gaussian Free Field (GFF) with mass  $m$  on  $M$  is the *Gaussian random process indexed by*  $C^\infty(M)$ , consisting of random variables  $\{\phi(f) \mid f \in C^\infty(M)\}$  such that

$$\mathbb{E}[\phi(f)\phi(h)] = \langle f, (\Delta + m^2)^{-1}h \rangle_{L^2(M)}, \quad \mathbb{E}[\phi(f)] \equiv 0, \quad (2.6)$$

for any  $f, h \in C^\infty(M)$ .

**Definition 2.2.** The Dirichlet massive Gaussian Free Field with mass  $m$  on  $\Omega$  is the *Gaussian random process indexed by*  $C_c^\infty(\Omega^\circ)$ , consisting of random variables  $\{\phi(f) \mid f \in C_c^\infty(\Omega^\circ)\}$  such that

$$\mathbb{E}[\phi(f)\phi(h)] = \langle f, (\Delta_{\Omega, D} + m^2)^{-1}h \rangle_{L^2(\Omega)} = \langle P_{M \setminus \Omega^\circ}^\perp f, P_{M \setminus \Omega^\circ}^\perp h \rangle_{W^{-1}}, \quad \mathbb{E}[\phi(f)] \equiv 0, \quad (2.7)$$

for any  $f, h \in C_c^\infty(\Omega^\circ)$ . See appendix C.1 and in particular lemma C.4.

**Remark 2.1.** There is also a technical requirement: the family of random variables  $\{\phi(f) \mid f \in C^\infty(M)\}$  (respectively,  $C_c^\infty(\Omega^\circ)$ ) should generate the  $\sigma$ -algebra of the underlying sample space. This makes the corresponding random processes *full*, per the probabilistic terminology. We will take this requirement for granted in the sequel.

We have chosen to start with these rather abstract definitions due to the fact that there are many choices of sample spaces ( $Q$ -spaces) on which to realize those Gaussian processes. These realizations are all equivalent in the sense explained in appendix B.2. For the sake of concreteness, we point out that one choice for the sample space is  $\mathcal{D}'(M)$  (or  $\mathcal{D}'(\Omega^\circ)$ ).

**Proposition 2.1** (Bochner-Minlos, [11] theorem 5.11, page 266). *There exists a Borel probability measure  $\mu_{\text{GFF}}^M$  on the Fréchet space  $\mathcal{D}'(M)$  such that  $\phi \mapsto \langle \phi, f \rangle_{L^2(M)} =: \phi(f)$ ,  $f \in C^\infty(M)$ , realizes the random variable  $\phi(f)$  of the massive GFF on  $M$ .*

*Similarly, there exists a Borel probability measure  $\mu_{\text{GFF}}^{\Omega, D}$  on the Fréchet space  $\mathcal{D}'(\Omega^\circ)$  such that  $\phi \mapsto \langle \phi, f \rangle_{L^2(\Omega)}$ ,  $f \in C_c^\infty(\Omega^\circ)$ , realizes the random variable  $\phi(f)$  of the Dirichlet massive GFF on  $\Omega$ .*

*Remark for proof.* The key point is that  $\mathcal{D}'(M)$  (respectively,  $\mathcal{D}'(\Omega^\circ)$ ) is the dual of a nuclear space  $C^\infty(M)$  (respectively,  $C_c^\infty(\Omega^\circ)$ ). There are other approaches, see remark 2.2 below and appendix B.2.  $\square$

*Notation.* Denote by  $\mathbb{E}_{\text{GFF}}^M$  and  $\mathbb{E}_{\text{GFF}}^{\Omega, D}$  the expectations under  $\mu_{\text{GFF}}^M$  and  $\mu_{\text{GFF}}^{\Omega, D}$  respectively.

**Remark 2.2.** Based on the spectral theory of  $\Delta$ , let  $\{\varphi_j\}_{j=0}^\infty$  be its complete orthonormal eigenfunctions with (real nonnegative) eigenvalues  $\{\lambda_j\}_{j=0}^\infty$ ,  $0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_j \leq \dots$ , counted with multiplicity. Then the GFF with mass  $m$  on  $M$  could also be represented as the random formal series

$$\phi = \sum_{j=0}^{\infty} \xi_j \varphi_j \quad (2.8)$$

where the sequence  $(\xi_j)$  has the law

$$\mu_{\text{GFF}}^M = \prod_{j=0}^{\infty} \mathcal{N}(0, (\lambda_j + m^2)^{-1}), \quad (2.9)$$

where  $\mathcal{N}(0, \lambda)$  is the standard centered Gaussian measure on  $\mathbb{R}$  with variance  $\lambda$ . Similarly, the Dirichlet GFF on  $\Omega$  could also be so represented using eigenfunctions of  $\Delta_{\Omega, D}$ , which are complete for  $L^2(\Omega)$ . See also appendix B.2.

We also consider a slightly more general situation where  $(\Delta + m^2)^{-1}$  is replaced by an operator  $C$ .

**Definition 2.3.** We say that a bounded self-adjoint positive elliptic pseudodifferential operator  $C$  of order  $-s$  on  $M$  ( $s > 0$ ) is a *Gaussian covariance operator* of order  $-s$ .

The same results go through (proposition 2.1 and remark 2.2), and one obtains a measure  $\mu_C$  on  $\mathcal{D}'(M)$ .

**Definition 2.4.** The Gaussian Field on  $M$  with covariance operator  $C$  is the *Gaussian random process indexed by  $C^\infty(M)$* , consisting of random variables  $\{\phi(f) \mid f \in C^\infty(M)\}$  such that

$$\mathbb{E}[\phi(f)\phi(h)] = \langle f, Ch \rangle_{L^2(M)}, \quad \mathbb{E}[\phi(f)] \equiv 0, \quad (2.10)$$

for any  $f, h \in C^\infty(M)$ . We denote the corresponding measure on  $\mathcal{D}'(M)$  by  $\mu_C$  and the expectation with respect to this measure by  $\mathbb{E}_C$ .

**Remark 2.3.** For  $C$  satisfying the assumptions, the inner product  $(f, h) \mapsto \langle f, Ch \rangle_{L^2(M)}$  defines an equivalent norm for  $W^{-s}(M)$ .

### 2.1.2 Essential Properties

It follows directly from the definitions that

**Lemma 2.2.** *The Gaussian Hilbert space of  $\mu_{\text{GFF}}^M$  is  $W^{-1}(M)$ , and that of  $\mu_{\text{GFF}}^{\Omega, D}$  is  $W^{-1}(\Omega)$ .*  $\square$

From (iii) of lemma B.3 and (iii) of lemma C.3,

**Lemma 2.3.** *The Cameron-Martin space of  $\mu_{\text{GFF}}^M$  is  $W^1(M)$ , and that of  $\mu_{\text{GFF}}^{\Omega, D}$  is  $W_\Omega^1(M)$ .*  $\square$

**Remark 2.4.** There is a tacit assumption in the way we defined our fields: we expected the random variable  $\phi(f)$  to come from the distributional pairing ( $L^2$ -pairing) between  $\mathcal{D}'(M)$  and  $C^\infty(M)$  (respectively,  $\Omega$ ). Other pairings may also be used. For example, let  $\phi(f)$  be  $\langle -, f \rangle_{W^1(M)}$  instead of  $\langle -, f \rangle_{L^2(M)}$ . One then needs to alter the covariances accordingly. Indeed, they are related by

$$\mathbb{E}[\langle \phi, f \rangle_{W^1} \langle \phi, h \rangle_{W^1}] = \mathbb{E}[\langle \phi, (\Delta + m^2)f \rangle_{L^2} \langle \phi, (\Delta + m^2)h \rangle_{L^2}] = \langle f, h \rangle_{W^1(M)}. \quad (2.11)$$

This way of definition is noticeably used by Sheffield [67]. The Gaussian Hilbert space in this case is  $W^1(M)$  (respectively,  $W_\Omega^1(M)$ ). Since we are nevertheless defining the same measures on  $\mathcal{D}'(M)$  and  $\mathcal{D}'(\Omega^\circ)$ , the Cameron-Martin spaces are the same. This pairing is more natural in view of the abstract Wiener space construction of the  $Q$ -space of a Gaussian process indexed by a Hilbert space (now both the indexing and the pairing are provided by  $W^1$ ). See appendix B.2, and also remark C.4.

Similarly,

**Lemma 2.4.** *The Gaussian Hilbert space of  $\mu_C$  is  $W^{-s}(M)$ , equipped with  $\langle -, C- \rangle_{L^2}$ , and the Cameron-Martin space is  $W^s(M)$ , equipped with  $\langle -, C^{-1}- \rangle_{L^2}$ .*  $\square$

**Corollary 2.5** (cf. proposition B.2). *Let  $\phi_0 \in W^s(M)$  and denote by  $(\phi_0)_*\mu_C$  the measure image of  $\mu_C$  under the shift  $\phi \mapsto \phi + \phi_0$ . Then the Radon-Nikodym density between  $(\phi_0)_*\mu_C$  and  $\mu_C$  is*

$$\frac{d((\phi_0)_*\mu_C)}{d\mu_C}(\phi) = e^{\phi(C^{-1}\phi_0) - \frac{1}{2}\langle \phi_0, C^{-1}\phi_0 \rangle_{L^2}}. \quad \square \quad (2.12)$$

Finally we say about the supports of the measures in the closed manifold case.

**Lemma 2.6.** *We have  $\mu_C(W^{-\delta}(M)) = 1$  for any  $\delta > \frac{1}{2}(d-s)$ .*

*Proof.* We rely on lemma 2.17. Pick  $t \in \mathbb{R}$ . Note that the real power  $C^{-2t/s}$  is a positive elliptic  $\Psi$ DO of order  $2t$ , and hence  $\langle -, C^{-2t/s}- \rangle_{L^2}$  gives an equivalent representation of  $\|\cdot\|_{W^t(M)}^2$ . Thus whenever  $C^{-\frac{2t}{s}+1}$  is trace class,

$$\mathbb{E}_C[\|\phi\|_{W^t(M)}^2] \approx \mathbb{E}_C[\langle \phi, C^{-2t/s}\phi \rangle_{L^2}] = \text{tr}_{L^2}(C^{-\frac{2t}{s}+1}) < \infty, \quad (2.13)$$

and it then follows  $\mu_C(W^t(M)) = 1$ . Since now  $C^{-\frac{2t}{s}+1}$  is a  $\Psi$ DO of order  $2t-s$ , it is trace class when  $2t-s < -d$  namely  $t < -\frac{1}{2}(d-s)$ , finishing the proof.  $\square$

**Remark 2.5.** We point out that  $\mu_C(W^{-\delta}(M)) = 1$  for any  $\delta > 0$  in the following two cases:

- (i)  $\dim M = 2$  and  $\mu_C = \mu_{\text{GFF}}^M$ ;
- (ii)  $\dim M = 1$  and  $C$  has order  $-1$ .

Last but not least, we make the following innocent but useful observation.

**Lemma 2.7.** *If  $\Omega = \Omega_1 \sqcup \Omega_2$  (possibility of non-empty boundary in either or both components), then GFFs (indeed, Gaussian fields) over  $\Omega_1$  and  $\Omega_2$  are independent and  $\mu_{\text{GFF}}^{\Omega, B} = \mu_{\text{GFF}}^{\Omega_1, B} \otimes \mu_{\text{GFF}}^{\Omega_2, B}$  where  $B = D$  when the corresponding  $\Omega$ ,  $\Omega_1$  or  $\Omega_2$  has boundary and  $B = \emptyset$  when either is closed.*  $\square$

## 2.2 Determinants

In this section we discuss (two) generalizations of the notion of the determinant (of a matrix) to infinite dimensional operators. Their starting points are as follows. Suppose  $A$  is a matrix (acting on  $V$ ) with eigenvalues  $\lambda_1, \dots, \lambda_N$  counted with algebraic multiplicity. Firstly, one has the equality

$$\det A = \prod_{j=1}^N \lambda_j = \exp(-\partial_z(\lambda_1^{-z} + \dots + \lambda_N^{-z})(0)) = \exp(-\partial_z \operatorname{tr}(A^{-z})(0)), \quad (2.14)$$

which suggests defining  $\det A$ , when  $A$  acts on infinite dimensions, as  $\exp(-\partial_z \operatorname{tr}(A^{-z})(0))$  when both  $A^{-z}$  (as a function of  $z \in \mathbb{C}$ ) and the derivative of its trace (at  $z = 0$ ) makes sense; this is called the **zeta-regularized determinant**. Secondly, if  $\mathbb{1}$  is the identity, one also has

$$\det(\mathbb{1} + A) = \prod_{j=1}^N (1 + \lambda_j) = \sum_{k=0}^N \sum_{i_1 < \dots < i_k} \lambda_{i_1} \dots \lambda_{i_k} = \sum_{k=0}^N \operatorname{tr}_{\Lambda^k V}(\Lambda^k A), \quad (2.15)$$

where  $\Lambda^k A$  is the  $k$ -th exterior/antisymmetric product of  $A$ , acting on the exterior product space  $\Lambda^k V$ . This suggests defining

$$\det(\mathbb{1} + A) \stackrel{\text{heu}}{=} \sum_{k=0}^{\infty} \operatorname{tr}(\Lambda^k A) \quad (2.16)$$

in infinite dimensions when all of  $\Lambda^k A$  are *trace class* and the infinite sum converges, and is called the **Fredholm determinant**.

General references include Kontsevich and Vishik [41], Shubin [68] sections 9-13, Simon [70] chapter 3 and finally Gohberg, Goldberg and Krupnik [25]. See also Dang [15] for a quick acquaintance of the physical-geometric context and Quine, Heydari and Song [55] for an interesting discussion of zeta-regularization of infinite products.

### 2.2.1 Zeta-regularized and Fredholm Determinants

The zeta-regularized determinant was first introduced by Ray and Singer [56]. The first step is to define the **zeta function** of a (rather special) pseudodifferential operator  $A$  over a manifold  $M$  with or without boundary,

$$\zeta_A(z) \stackrel{\text{def}}{=} \operatorname{tr}_{L^2}(A^{-z}) \quad (2.17)$$

as a function of the complex variable  $z$  and study its meromorphic extension over a region that includes  $z = 0$ . For our purposes  $A$  is  $\Delta + m^2$ ,  $(\Delta + m^2)^{1/2}$  or a Dirichlet-to-Neumann operator. These are positive elliptic  $\Psi$ DOs of positive order such that the principal symbol  $\sigma_A(x, \xi)$  is strictly positive whenever  $\xi \neq 0$ . In particular their spectra are in  $\mathbb{R}_+$  and does not intersect  $\mathbb{B}_\rho(0) \subset \mathbb{C}$  for some  $\rho > 0$ . This enables one to define the complex power  $A^{-z}$  using the Cauchy integral representation

$$A^{-z} \stackrel{\text{def}}{=} \frac{i}{2\pi} \int_{\gamma} e^{-z \log \lambda} (A - \lambda)^{-1} d\lambda, \quad (2.18)$$

where  $\gamma$  is the contour (with parametrization traversing in order)

$$\gamma = \{re^{i\pi} \mid r > \rho\} \cup \{\rho e^{i\theta} \mid -\pi < \theta < \pi\} \cup \{re^{-i\pi} \mid r > \rho\}, \quad (2.19)$$

and  $\log \lambda$  taken to be the principal branch defined on  $\mathbb{C} \setminus (-\infty, 0]$  with  $\log 1 = 0$ .

**Proposition 2.8** ([68] proposition 10.1, theorems 10.1, 13.1, 13.2, also [65]). *We have*

(i) *For the operators  $A$  under consideration, we have the bound*

$$\|(A - \lambda)^{-1}\|_{L^2} \leq c|\lambda|^{-1} \quad (2.20)$$

*for  $\lambda \in \gamma$ , and the integral (2.18) defines  $A^{-z}$  as a holomorphic function valued in bounded  $L^2$  operators, for  $\Re(z) > 0$ . It continues as such a holomorphic operator function to all  $z \in \mathbb{C}$  via*

$$A^{-z} \stackrel{\text{def}}{=} A^k A^{-z-k}, \quad (2.21)$$

*where  $k$  is any integer with  $\Re(z) > -k$  so that  $A^{-z-k}$  is defined by (2.18), and the definition (2.21) does not depend on  $k$ .*

(ii) If  $A$  has order  $s$  then  $A^{-z}$  defined as above is a classical  $\Psi$ DO of order  $-zs$ . In particular it is trace class on  $L^2(M)$  when  $\Re(z) > d/s$ ,  $d = \dim M$ , for which  $\zeta_A(z)$  is well-defined by (2.17). Moreover, for these  $z$ ,

$$\zeta_A(z) = \sum_{j=0}^{\infty} \lambda_j^{-z}, \quad (2.22)$$

where  $\{\lambda_j\}_{j=0}^{\infty}$  are the eigenvalues of  $A$ , and the sum converges absolutely, and uniformly in  $z$  over  $\{\Re(z) > d/s + \varepsilon\}$  for any  $\varepsilon > 0$ .

(iii) Finally,  $\zeta_A(z)$  can be meromorphically continued over  $\mathbb{C}$  with simple poles possible at  $\{\frac{d}{s}, \frac{d-1}{s}, \frac{d-2}{s}, \dots\} \setminus \mathbb{Z}_{\leq 0}$ , and holomorphic elsewhere. In particular, it is holomorphic at  $z = 0$ .  $\square$

**Definition 2.5.** For an operator  $A$  under consideration, we define its zeta-regularized determinant as

$$\det_{\zeta} A \stackrel{\text{def}}{=} \exp(-\partial_z \zeta_A(0)), \quad (2.23)$$

where  $\zeta_A(0)$  is the zeta function of  $A$  given by (2.17) and (2.18).

**Remark 2.6.** An alternative way of defining the zeta function and its meromorphic continuation is to use the heat kernel and the Mellin transform. See Gilkey [24] section 1.12. This way of definition also gives (2.22) over the same region, defining therefore the same function as ours.

**Remark 2.7.** From (ii) of proposition 2.8 we see that if  $A$  is self-adjoint and strictly positive, then  $\zeta_A(z)$  is real-valued for  $z \in (d/s, +\infty)$ . But  $\Im(\zeta_A)$  is real analytic and hence  $\zeta_A$  remains real-valued on  $\mathbb{R}$  before crossing a pole, and by (iii) it is in particular real-valued on an interval around 0. Thus  $\partial_z \zeta_A(0)$  is real and  $\det_{\zeta} A$  is positive.

Now we move on to the second notion of determinant. Let  $\mathcal{H}$  be a Hilbert space and  $A \in \mathcal{L}(\mathcal{H})$ . Denote by  $\Lambda^k \mathcal{H}$  and  $\Lambda^k A$ , respectively, the  $k$ -th exterior product of  $\mathcal{H}$  and  $A$  (see Simon [70] section 1.5).

**Proposition 2.9** ([70] lemma 3.3). *If  $A$  is trace class on  $\mathcal{H}$ , then  $\Lambda^k A$  is also trace class on  $\Lambda^k \mathcal{H}$  with bound on trace norm*

$$\|\Lambda^k A\|_{\text{tr}} \leq \frac{1}{k!} \|A\|_{\text{tr}}^k. \quad (2.24)$$

In particular, putting

$$\det_{\text{Fr}}(\mathbb{1} + zA) \stackrel{\text{def}}{=} \sum_{k=0}^{\infty} z^k \text{tr}_{\Lambda^k \mathcal{H}}(\Lambda^k A) \quad (2.25)$$

for  $z \in \mathbb{C}$  defines an entire function, and

$$|\det_{\text{Fr}}(\mathbb{1} + zA)| \leq \exp(|z| \|A\|_{\text{tr}}). \quad \square \quad (2.26)$$

**Definition 2.6.** Let  $A$  be a trace class operator on the Hilbert space  $\mathcal{H}$ . Then the determinant  $\det_{\text{Fr}}(\mathbb{1} + A)$  given by (2.25) for  $z = 1$  is called the Fredholm determinant of  $\mathbb{1} + A$ .

**Lemma 2.10** ([70] theorem 3.4). *The map  $A \mapsto \det_{\text{Fr}}(\mathbb{1} + A)$  defines a continuous function on the trace ideal  $\mathcal{J}_1$  with  $\|\cdot\|_{\text{tr}}$ . More precisely,*

$$|\det_{\text{Fr}}(\mathbb{1} + A) - \det_{\text{Fr}}(\mathbb{1} + B)| \leq \|A - B\|_{\text{tr}} \exp(\|A\|_{\text{tr}} + \|B\|_{\text{tr}} + 1). \quad \square \quad (2.27)$$

Finally we include the next commutativity result which is relatively more advanced.

**Lemma 2.11.** *If  $A, B \in \mathcal{L}(\mathcal{H})$  are such that both  $AB$  and  $BA$  are of trace class, then we have*

$$\det_{\text{Fr}}(\mathbb{1} + AB) = \det_{\text{Fr}}(\mathbb{1} + BA). \quad (2.28)$$

*Proof.* The proof is based on the Hadamard product formula for (2.25). See Simon [70] theorem 3.7 and the proof of corollary 3.8 ( $AB$  and  $BA$  have the same nonzero eigenvalues), and the references therein.  $\square$

### 2.2.2 Factorization Lemma

**Lemma 2.12.** *Suppose  $A$  and  $K$  are  $\Psi$ DOs such that both  $A$  and  $A(\mathbb{1} + K)$  satisfy the assumptions of proposition 2.8 and that  $\det_\zeta(A)$  and  $\det_\zeta(A(\mathbb{1} + K))$  are defined. Suppose moreover  $K$  is trace class and there exists smoothing operators  $\{K_i\}_{i=1}^\infty$  such that  $AK_i \rightarrow AK$  in  $\|\cdot\|_{\text{tr}}$  (in particular,  $AK$  is also trace class). Then*

$$\det_\zeta(A(\mathbb{1} + K)) = \det_\zeta(A) \det_{\text{Fr}}(\mathbb{1} + K). \quad (2.29)$$

*Proof.* We follow Kontsevich and Vishik [41] proposition 6.4 and take for granted that (2.29) holds with  $K_i$  in place of  $K$ . Our assumptions are tailor-made so that as  $i \rightarrow \infty$ ,

$$\det_\zeta(A(\mathbb{1} + K_i)) \rightarrow \det_\zeta(A(\mathbb{1} + K)). \quad (2.30)$$

Indeed, by (2.20) we have

$$\|(A(\mathbb{1} + K) - \lambda)^{-1}A(K_i - K)(A(\mathbb{1} + K_i) - \lambda)^{-1}\|_{\text{tr}} \leq c|\lambda|^{-2}\|A(K_i - K)\|_{\text{tr}}, \quad (2.31)$$

with  $c$  independent of  $i$  since a fortiori  $AK_i \rightarrow AK$  under  $\|\cdot\|_{L^2}$ . This in particular shows when  $\Re(z) > -1$  the integral expression for  $A(\mathbb{1} + K)^{-z} - A(\mathbb{1} + K_i)^{-z}$  is a converging Bochner integral valued in the trace ideal  $\mathcal{J}_1$  (note  $|\lambda^{-z}| \asymp |\lambda|^{-\Re(z)}$  as  $\lambda \rightarrow -\infty$ ) and since  $\text{tr}_{L^2}$  is a continuous functional on  $\mathcal{J}_1$ ,

$$|\text{tr}_{L^2}(A(\mathbb{1} + K)^{-z} - A(\mathbb{1} + K_i)^{-z})| \lesssim \int_\gamma |\lambda|^{-\Re(z)-2} \|A(K_i - K)\|_{\text{tr}} d\lambda. \quad (2.32)$$

Now by (iii) of proposition 2.8 there is  $\frac{1}{2} > \delta > 0$  so that  $\zeta_{A(\mathbb{1} + K_i)}$ ,  $\zeta_{A(\mathbb{1} + K)}$  are both holomorphic over  $\overline{\mathbb{B}}_\delta(0)$  (for example,  $\delta < 1/|s|$  where  $s$  is the order of  $A$ ). Thus by (2.32) and Cauchy's estimate

$$\begin{aligned} |\zeta'_{A(\mathbb{1} + K)}(0) - \zeta'_{A(\mathbb{1} + K_i)}(0)| &\leq \frac{1}{\delta} \sup_{|z|=\delta} |\zeta_{A(\mathbb{1} + K)}(z) - \zeta_{A(\mathbb{1} + K_i)}(z)| \\ &\lesssim \frac{1}{\delta} \sup_{|z|=\delta} \int_\gamma |\lambda|^{-\Re(z)-2} \|A(K_i - K)\|_{\text{tr}} d\lambda \\ &\lesssim \|A(K_i - K)\|_{\text{tr}} \int_\gamma |\lambda|^{-3/2} d\lambda \\ &\lesssim \|A(K_i - K)\|_{\text{tr}}. \end{aligned}$$

This shows  $|\zeta'_{A(\mathbb{1} + K)}(0) - \zeta'_{A(\mathbb{1} + K_i)}(0)| \rightarrow 0$ , as we have desired.  $\square$

### 2.2.3 The Gluing Formula of Burghlelea-Friedlander-Kappeler

Let  $(M, g)$  is a closed Riemannian manifold and  $\Sigma \subset M$  an embedded closed hypersurface with induced metric. Assume proposition 5.7 and decompose  $\phi = \phi_\Sigma + \phi_{M \setminus \Sigma}^D$  corresponding to  $\mu_{\text{GFF}}^M = \mu_{\text{GFF}}^{M \setminus \Sigma, D} \otimes \mu_{\text{DN}}^{\Sigma, M}$ . In view of equation (2.5), and in parallel

$$e^{-\frac{1}{2}\langle \phi, \text{DN} \phi \rangle_{L^2}} [\mathcal{L}\phi] \stackrel{\text{heu}}{=} \det_\zeta(\text{DN}_M^\Sigma)^{-\frac{1}{2}} d\mu_{\text{DN}}^{\Sigma, M}, \quad (2.33)$$

if we assume a ‘‘formal Fubini theorem’’ with respect to the heuristic expressions involving  $\mathcal{L}$ , namely

$$\int e^{-\frac{1}{2}\langle \phi, (\Delta + m^2)\phi \rangle_{L^2}} [\mathcal{L}\phi] \stackrel{\text{heu}}{=} \iint e^{-\frac{1}{2}\langle \phi_\Sigma, (\Delta + m^2)\phi_\Sigma \rangle_{L^2}} e^{-\frac{1}{2}\langle \phi_{M \setminus \Sigma}^D, (\Delta + m^2)\phi_{M \setminus \Sigma}^D \rangle_{L^2}} [\mathcal{L}\phi_\Sigma] \otimes [\mathcal{L}\phi_{M \setminus \Sigma}^D], \quad (2.34)$$

then we are led to the following relation of the corresponding determinants (volumes) which were first rigorously proved by Burghlelea, Friedlander and Kappeler [9].

**Proposition 2.13** ([9] theorem B, [42] theorem 1.1). *Let  $(M, g)$  is a closed Riemannian surface and  $\Sigma \subset M$  an embedded closed hypersurface with induced metric. Then*

$$\det_\zeta(\Delta_M + m^2) = \det_\zeta(\Delta_{M \setminus \Sigma, D} + m^2) \det_\zeta(\text{DN}_M^\Sigma). \quad \square \quad (2.35)$$

The following version where  $\Sigma$  dissects  $M$  such that  $M \setminus \Sigma = M_+^\circ \sqcup M_-^\circ$  is also useful.

**Corollary 2.14.** *In the situation as above, we have*

$$\det_\zeta(\Delta_M + m^2) = \det_\zeta(\Delta_{M_+, D} + m^2) \det_\zeta(\Delta_{M_-, D} + m^2) \det_\zeta(\text{DN}_M^\Sigma). \quad \square \quad (2.36)$$

As we base our analysis on background Gaussian probability measures, the formulae (2.35) and (2.36) constitute separate ingredients (constants) that needs to be ‘‘tuned’’ for the final gluing result to hold exactly. In fact, it is also reasonable to consider ‘‘projective gluing’’ which allows the freedom for an arbitrary (nonzero) constant to appear in the equation (see remark 6.2 and Segal [66] page 460).

## 2.3 Quadratic Perturbation = Radon-Nikodym Density

Let  $C$  be a Gaussian covariance operator of order  $-s$  on a closed Riemannian manifold  $\Sigma$ , and denote by  $\mu_C$  the Gaussian measure on  $\mathcal{D}'(\Sigma)$  with covariance  $\langle -, C- \rangle_{L^2(\Sigma)}$ . Let  $V$  be another bounded formally self-adjoint operator on  $L^2(\Sigma)$  (it could be given by a real symmetric Schwartz kernel). In this section we look at the Gibbs measure

$$d\mu(\varphi) \stackrel{\text{def}}{=} \frac{e^{-\frac{1}{2}\langle \varphi, V\varphi \rangle_{L^2}} d\mu_C(\varphi)}{\int e^{-\frac{1}{2}\langle \varphi, V\varphi \rangle_{L^2}} d\mu_C}, \quad (2.37)$$

which is a Gaussian measure (see proposition 2.16).

From another perspective we consider Radon-Nikodym densities between mutually absolutely continuous Gaussian measures on  $\mathcal{D}'(\Sigma)$ . See Bogachev [12] section 6.4 for a general treatment from this perspective. We shall reproduce a proof following Glimm and Jaffe [26] section 9.3 for reader's convenience and adaptation to the current situation.

A principal corollary of the results of this section is the following.

**Corollary 2.15.** *Let  $\Sigma$  be the disjoint union of Riemannian circles, embedded in an ambient Riemannian surface  $M$  (with or without boundary). Let  $\mu_{\text{DN}}^{\Sigma, M}$  and  $\mu_{2\mathbf{D}}^{\Sigma}$  be the two Gaussian measures constructed on  $\mathcal{D}'(\Sigma)$  with covariance operators  $(\text{DN}_M^{\Sigma})^{-1}$  and  $(2\mathbf{D}_{\Sigma})^{-1}$  (if  $M$  has boundary, specify the boundary condition to be  $B$  as in section 4.2). Then  $\mu_{\text{DN}}^{\Sigma, M}$  and  $\mu_{2\mathbf{D}}^{\Sigma}$  are mutually absolutely continuous with Radon-Nikodym density given by*

$$\frac{d\mu_{\text{DN}}^{\Sigma, M}}{d\mu_{2\mathbf{D}}^{\Sigma}}(\varphi) = (\det_{\zeta}(2\mathbf{D}_{\Sigma}))^{-\frac{1}{2}} (\det_{\zeta} \text{DN}_M^{\Sigma})^{\frac{1}{2}} e^{-\frac{1}{2}\langle \varphi, (\text{DN}_M^{\Sigma} - 2\mathbf{D}_{\Sigma})\varphi \rangle_{L^2(\Sigma)}}. \quad (2.38)$$

The proof is at the end of this section. First we come back to the general case.

**Proposition 2.16** (cf. [26] section 9.3). *Let  $V : L^2(\Sigma) \rightarrow L^2(\Sigma)$  be as above and moreover assume*

$$C^{-1} + V \text{ is positive}, \quad (2.39)$$

and that

$$\widehat{V} \stackrel{\text{def}}{=} C^{\frac{1}{2}} V C^{\frac{1}{2}} \text{ is trace class}. \quad (2.40)$$

Then

- (i) the random variable  $\langle \varphi, V\varphi \rangle_{L^2}$  can be defined in  $L^1(\mu_C)$  and  $\mathbb{E}_C[\langle \varphi, V\varphi \rangle_{L^2}] = \text{tr}(\widehat{V})$ ,
- (ii)  $Z := \mathbb{E}_C[e^{-\frac{1}{2}\langle \varphi, V\varphi \rangle_{L^2}}] = \det_{\text{Fr}}(\mathbb{1} + \widehat{V})^{-\frac{1}{2}}$ , and
- (iii) the Gibbs measure (2.37) is Gaussian with covariance  $(C^{-1} + V)^{-1}$ .

Note that since  $C^{-1} + V$  is positive and  $C$  is also positive,  $\mathbb{1} + \widehat{V} = C^{\frac{1}{2}}(C^{-1} + V)C^{\frac{1}{2}}$  is positive.

**Lemma 2.17.** *There exist an orthonormal basis  $\{f_j\}_{j=1}^{\infty}$  of the Gaussian Hilbert space  $W^{-s}(\Sigma)$  of  $\mu_C$  equipped with  $\langle -, C- \rangle_{L^2}$ , such that*

$$\langle \varphi, V\varphi \rangle_{L^2} = \sum_{j=1}^{\infty} \lambda_j \varphi(f_j)^2, \quad (2.41)$$

for all  $\varphi$  belonging to the Cameron-Martin space  $W^s(\Sigma)$ , where  $\{\lambda_j\}$  are the eigenvalues of  $\widehat{V}$  on  $L^2(\Sigma)$ , and the series converges absolutely in  $L^1(\mu_C)$ . Thus we define the random variable  $\langle \varphi, V\varphi \rangle_{L^2}$  with this converging series. Consequently, (i) of proposition 2.16 holds.

*Proof.* The key is to seek  $\{C^{\frac{1}{2}} f_j\}$  as complete  $L^2$ -orthonormal eigenfunctions of  $\widehat{V}$  with eigenvalues  $\{\lambda_j\}$ , which exist since  $\widehat{V}$  is self-adjoint and trace class on  $L^2(\Sigma)$ . Now  $\{f_j\} \subset W^{-s}(\Sigma)$  and is complete orthonormal with respect to  $\langle -, C- \rangle_{L^2}$ . This also means the random variables  $\{\varphi(f_j)\}$  are mutually independent. Note for  $\varphi \in W^s(\Sigma)$ ,  $C^{-\frac{1}{2}}\varphi \in L^2(\Sigma)$ , and

$$\begin{aligned} \langle \varphi, V\varphi \rangle_{L^2} &= \langle C^{-\frac{1}{2}}\varphi, \widehat{V}C^{-\frac{1}{2}}\varphi \rangle_{L^2} \\ &= \sum_{j=1}^{\infty} \lambda_j |\langle C^{-\frac{1}{2}}\varphi, C^{\frac{1}{2}}f_j \rangle_{L^2}|^2 \\ &= \sum_{j=1}^{\infty} \lambda_j \varphi(f_j)^2. \end{aligned}$$



Since  $\widehat{V}$  is trace class,

$$\sum_{j=1}^{\infty} \mathbb{E}_C [|\lambda_j| \varphi(f_j)^2] = \sum_{j=1}^{\infty} |\lambda_j| < \infty, \quad (2.42)$$

and we obtain the result.  $\square$

**Remark 2.8.** We thus defined  $\langle \varphi, V\varphi \rangle_{L^2}$  as a Wiener quadratic form and it lies in the second Wiener chaos of  $\mu_C$ , a fortiori in  $L^2(\mu_C)$ . See Bogachev [12] pages 257-261 for more information and in particular proposition 5.10.16 for the same result in the context of Malliavin calculus. In fact,

$$VCf_j = C^{-\frac{1}{2}} \widehat{V} C^{\frac{1}{2}} f_j = \lambda_j f_j, \quad (2.43)$$

and hence  $V$  is trace class on  $W^s(\Sigma)$  with eigenbasis  $\{Cf_j\}$ .

**Remark 2.9.** We have  $\lambda_j > -1$  by the assumption (2.39).

To treat (ii) and (iii) of proposition 2.16 we adopt some approximations.

**Lemma 2.18.** *Proposition 2.16 (ii) and (iii) is true in the case  $V$  has finite rank.*

*Proof.* In this case the series in (2.41) is finite with  $\{f_j\}_{j=1}^N$  for some  $N \in \mathbb{N}$ . Thus

$$\begin{aligned} Z &= \frac{1}{(2\pi)^{N/2}} \int_{\mathbb{R}^N} e^{-\frac{1}{2} \sum_j \lambda_j x_j^2} e^{-\frac{1}{2} \sum_j x_j^2} d^N x \\ &= \prod_j (1 + \lambda_j)^{-\frac{1}{2}} = \det(\mathbb{1} + \widehat{V})^{-\frac{1}{2}}, \end{aligned}$$

by projecting onto  $\mathbb{R}^N$  via  $\varphi \mapsto (\varphi(f_1), \dots, \varphi(f_N)) =: (x_1, \dots, x_N)$ . For the covariance, we orthogonally decompose  $W^{-s}(\Sigma)$  as

$$W^{-s}(\Sigma) = \text{Span}\{f_j \mid 1 \leq j \leq N\} \oplus \text{Span}\{f_j \mid 1 \leq j \leq N\}^{\perp}, \quad (2.44)$$

let  $\Pi_0$  and  $\Pi_1$  be the corresponding orthogonal projections (in order), and for any  $f \in C^\infty(\Sigma)$  write

$$\varphi(f) = \alpha_1 \varphi(f_1) + \dots + \alpha_N \varphi(f_N) + \varphi(\Pi_1 f), \quad (2.45)$$

then  $\varphi(\Pi_1 f)$  is independent of both  $\varphi(f_j)$ ,  $1 \leq j \leq N$ , and  $\langle \varphi, V\varphi \rangle_{L^2}$ . It is now clear that (2.37) is Gaussian because we could now express  $\mu(\{\phi(f) \in A\})$  for any Borel set  $A \subset \mathbb{R}$  as a Gaussian integral over  $\mathbb{R}^{N+1}$ . Apply  $C^{\frac{1}{2}}$  to (2.44) we get the  $L^2$ -orthogonal decomposition

$$L^2(\Sigma) = \underbrace{C^{\frac{1}{2}}(\text{Span}\{f_j \mid 1 \leq j \leq N\})}_{=: L^2(\Sigma)_0} \oplus \underbrace{C^{\frac{1}{2}}(\text{Span}\{f_j \mid 1 \leq j \leq N\}^{\perp})}_{=: L^2(\Sigma)_1}. \quad (2.46)$$

Clearly  $\widehat{V}$  leaves this decomposition invariant and is zero on  $L^2(\Sigma)_1$ . Hence  $(\mathbb{1} + \widehat{V})^{-1}$  is block-diagonal,

$$(\mathbb{1} + \widehat{V})^{-1} = \begin{pmatrix} \{C^{\frac{1}{2}} f_j\}_1^N & L^2(\Sigma)_1 \\ \text{diag}_{j=1}^N \{(\lambda_j + 1)^{-1}\} & 0 \\ 0 & \mathbb{1} \end{pmatrix} \begin{pmatrix} \{C^{\frac{1}{2}} f_j\}_1^N \\ L^2(\Sigma)_1 \end{pmatrix} \quad (2.47)$$

Thus

$$\begin{aligned} Z^{-1} \mathbb{E}_C [\varphi(f)^2 e^{-\frac{1}{2} \langle \varphi, V\varphi \rangle}] &= Z^{-1} \mathbb{E}_C [\varphi(\Pi_0 f)^2 e^{-\frac{1}{2} \sum_j \lambda_j \varphi(f_j)^2}] + Z^{-1} \mathbb{E}_C [\varphi(\Pi_1 f)^2] \mathbb{E}_C [e^{-\frac{1}{2} \langle \varphi, V\varphi \rangle}] \\ &= \sum_j \alpha_j^2 (\lambda_j + 1)^{-1} + \langle \Pi_1 f, C \Pi_1 f \rangle_{L^2} \\ &= \langle C^{\frac{1}{2}} \Pi_0 f, (\mathbb{1} + \widehat{V})^{-1} C^{\frac{1}{2}} \Pi_0 f \rangle_{L^2} + \langle \Pi_1 f, C^{\frac{1}{2}} (\mathbb{1} + \widehat{V})^{-1} C^{\frac{1}{2}} \Pi_1 f \rangle_{L^2} \\ &= \langle f, (C^{-1} + V)^{-1} f \rangle_{L^2}, \end{aligned}$$

where we perform again a Gaussian integral on  $\mathbb{R}^N$  in the second line.  $\square$

*Proof of proposition 2.16 (ii) and (iii).* For general  $V$  we impose spectral cut-off at  $N$ ,

$$V_N(\varphi) \stackrel{\text{def}}{=} \sum_{j=1}^N \lambda_j \langle \varphi, f_j \rangle_{L^2} f_j. \quad (2.48)$$

Then  $C^{\frac{1}{2}} V_N C^{\frac{1}{2}} =: \widehat{V}_N \rightarrow \widehat{V}$  under the trace norm  $\|\cdot\|_{\text{tr}}$  (acting on  $L^2(\Sigma)$ ). Hence  $\langle \varphi, V_N \varphi \rangle_{L^2} \rightarrow \langle \varphi, V \varphi \rangle_{L^2}$  in  $L^1(\mu_C)$  by lemma 2.17 and after passing to a subsequence

$$e^{-\frac{1}{2} \langle \varphi, V_N \varphi \rangle_{L^2}} \rightarrow e^{-\frac{1}{2} \langle \varphi, V \varphi \rangle_{L^2}} \quad (2.49)$$

in  $L^1(\mu_C)$  and (ii) for  $V$  follows, since  $\det_{\text{Fr}}(\mathbb{1} + \widehat{V}_N) \rightarrow \det_{\text{Fr}}(\mathbb{1} + \widehat{V})$  by lemma 2.10. To prove (iii), we note that  $\widehat{V}_N \rightarrow \widehat{V}$  a fortiori under the operator norm, then  $C^{\frac{1}{2}}(\mathbb{1} + \widehat{V}_N)^{-1}C^{\frac{1}{2}} \rightarrow C^{\frac{1}{2}}(\mathbb{1} + \widehat{V})^{-1}C^{\frac{1}{2}}$  in norm and

$$\begin{aligned} \frac{\mathbb{E}_C[e^{i\varphi(f)} e^{-\frac{1}{2} \langle \varphi, V \varphi \rangle_{L^2}}]}{\mathbb{E}_C[e^{-\frac{1}{2} \langle \varphi, V \varphi \rangle_{L^2}}]} &= \lim_{N \rightarrow \infty} \frac{\mathbb{E}_C[e^{i\varphi(f)} e^{-\frac{1}{2} \langle \varphi, V_N \varphi \rangle_{L^2}}]}{\mathbb{E}_C[e^{-\frac{1}{2} \langle \varphi, V_N \varphi \rangle_{L^2}}]} \\ &= \lim_{N \rightarrow \infty} \exp\left(-\frac{1}{2} \langle f, (C^{-1} + V_N)^{-1} f \rangle_{L^2}\right) \\ &= \exp\left(-\frac{1}{2} \langle f, (C^{-1} + V)^{-1} f \rangle_{L^2}\right), \end{aligned}$$

for  $f \in C^\infty(\Sigma)$ , showing that (2.37) is Gaussian with the right covariance.  $\square$

*Proof of corollary 2.15.* Remember now that  $\dim \Sigma = 1$ . Write for short  $\mathbf{D} := 2\mathbf{D}_\Sigma$  and  $\text{DN} := \text{DN}_{\mathcal{M}}^\Sigma$ . Setting  $C = \mathbf{D}^{-1}$  and  $V = \text{DN} - \mathbf{D}$  in proposition 2.16, we are left to prove the determinant identity

$$\det_\zeta(\mathbf{D}) \det_{\text{Fr}}(\mathbb{1} + \mathbf{D}^{-\frac{1}{2}}(\text{DN} - \mathbf{D})\mathbf{D}^{-\frac{1}{2}}) = \det_\zeta(\text{DN}). \quad (2.50)$$

Indeed, this is now immediate as

$$\text{LHS} = \det_\zeta(\mathbf{D}) \det_{\text{Fr}}(\mathbb{1} + \mathbf{D}^{-1}(\text{DN} - \mathbf{D})) = \text{RHS} \quad (2.51)$$

by lemma 2.11 and lemma 2.12. We point out  $\mathbf{D}\mathbf{D}^{-1}(\text{DN} - \mathbf{D}) = V$  can be approximated in  $\|\cdot\|_{\text{tr}}$  by smoothing operators (by lemma 4.4 (iv),  $V = \text{DN} - \mathbf{D}$  is  $L^2$ -trace class), so the conditions of lemma 2.12 is satisfied. Indeed, as above,  $\{Cf_j\} \subset W^1(\Sigma)$  are  $W^1$ -complete eigenfunctions of  $V$ . If the corresponding eigenvalue  $\lambda_j \neq 0$ , then the bootstrap argument shows  $Cf_j \in C^\infty(\Sigma)$  since  $V \in \Psi^{<0}(\Sigma)$ . Thus  $V_N$ , where  $V_N$  is as in (2.48), approximates  $V$  in  $\|\cdot\|_{\text{tr}}$  and is smoothing. This concludes the proof.  $\square$

**Remark 2.10.** Though it is probably true, we do not claim  $\text{DN} - \mathbf{D}$  is elliptic.

### 3 Variants of Nelson's Argument

The goal of this section is to define part  $A$  of (2.1) which culminates in [Nelson's theorem](#). The principal obstacle in achieving this is the fact that powers of a distribution such as  $\phi^2$ ,  $\phi^4$  etc., are generally not defined. Even as a random variable under  $\mu_{\text{GFF}}^M$ , we have (formally)  $\mathbb{E}_{\text{GFF}}^M[\phi(x)\phi(y)] \stackrel{\text{heu}}{=} G_{(\Delta+m^2)}(x, y)$  from (2.6) for  $x \neq y$  but this implies  $\mathbb{E}_{\text{GFF}}^M[\phi(x)\phi(x)] = G_{(\Delta+m^2)}(x, x) = \infty$ . This necessitates a procedure of **renormalization** which *subtracts away*  $\infty$  and makes  $\mathbb{E}_{\text{GFF}}^M[\phi(x)\phi(x)] < \infty$ .

**Example 3.1** ([21] page 288). A simplest example of such a procedure is defining the product distribution  $x^{-1}1_{(0, \infty)}$  on  $\mathbb{R}$  (here  $x^{-1}$  is the principal value distribution). Note  $x^{-1}$  and  $1_{(0, \infty)}$  have a common singularity at 0, their product is not naturally a distribution over  $\mathbb{R}$ . However, keeping  $\varepsilon > 0$  away from 0, for any test function  $\varphi \in C_c^\infty(\mathbb{R})$  we have by integration by parts

$$\int_\varepsilon^\infty \varphi(x) \frac{dx}{x} = - \int_\varepsilon^\infty \varphi'(x) \log(x) dx - \varphi(\varepsilon) \log(\varepsilon). \quad (3.1)$$

The point is that, since  $\log(x)$  is locally integrable, the distribution  $\varphi \mapsto - \int_0^\infty \varphi'(x) \log(x) dx$  is nevertheless well-defined and agrees with  $x^{-1}1_{(0, \infty)}$  for  $\varphi \in C_c^\infty(\mathbb{R} \setminus \{0\})$  since for small enough  $\varepsilon$  away from  $\text{supp}(\varphi)$  the boundary term disappears. Therefore this latter distribution could be taken as a **renormalized version** of  $x^{-1}1_{(0, \infty)}$  and formally corresponds to  $x^{-1}1_{(0, \infty)} - \infty \cdot \delta_0$  by taking  $\varepsilon \rightarrow 0$  in (3.1). (In fact, taking the principal value is in a broad sense another renormalization.)

In our case the natural renormalization strategy is provided by the Gaussian probability theory. Let  $\{K_\varepsilon \mid \varepsilon > 0\}$  be a family of *smoothing* operators<sup>5</sup> on  $M$ , for which  $K_\varepsilon \rightarrow \mathbb{1}$  as  $\varepsilon \rightarrow 0$  (in a sense to be specified later), and consider the mollified random field  $\phi_\varepsilon := K_\varepsilon \phi$ . Instead of  $\phi_\varepsilon(x)^4$ , say, we look at

$$\begin{aligned} :\phi_\varepsilon(x)^4: &\stackrel{\text{def}}{=} \phi_\varepsilon(x)^4 - 6\mathbb{E}[\phi_\varepsilon(x)^2]\phi_\varepsilon(x)^2 + 3\mathbb{E}[\phi_\varepsilon(x)^2]^2 \\ &\stackrel{\text{def}}{=} \phi_\varepsilon(x)^4 - 6C_\varepsilon(x)\phi_\varepsilon(x)^2 + 3C_\varepsilon(x)^2. \end{aligned}$$

It happens that for any  $\chi \in C^\infty(M)$ , the integral  $\int_M \chi(x) : \phi_\varepsilon(x)^4 : dV_M(x)$  converges as a random variable in  $L^2(\mu_{\text{GFF}}^M)$  to a definitive limit, and defines  $\int_M \chi(x) : \phi(x)^4 : dV_M(x)$  as a random variable in  $L^2(\mu_{\text{GFF}}^M)$ . Note that  $C_\varepsilon(x) \rightarrow \infty$  as  $\varepsilon \rightarrow 0$ , so we have subtracted ‘‘infinities’’.

**Remark 3.1. The crucial point** in our adaptation of Nelson’s argument is to realize the *locality* of the interaction  $\int_M \chi(x) : P(\phi(x)) : dV_M(x)$  (see section 5.5). To this end we must allow a sufficiently large class of *regulators*  $K_\varepsilon$  (in particular, *local* ones) and show that they define the same interaction (proposition 3.5). In addition, the *Wick ordering* also needs to be local (see section 3.4), so that the interaction on a domain with boundary could be defined without reference to the ambient closed manifold where this domain ‘‘caps’’. See Brunetti, Fredenhagen, Verch [6] and Guo, Paycha, Zhang [30] for more information and perspective on locality.

**Remark 3.2.** The method adopted here is restricted to dimension two. In three dimensions, the target measure bearing the heuristic form (2.1) becomes mutually singular with respect to  $\mu_{\text{GFF}}^M$  and hence cannot be expressed as an integrable function multiplied by  $\mu_{\text{GFF}}^M$ . A recent phenomenal method to treat this case is developed in the framework of stochastic PDEs, called *stochastic quantization*, providing an alternative to older results outlined in [26] section 23.1. See the introductions in [27], [32], [1], [45] and [4] for reviews of the literature and pedagogical discussions.

### 3.1 Regularizations

In this subsection we describe an admissible class of regulators  $K_\varepsilon$  which would eventually produce the same random variable  $\int_M \chi(x) : \phi(x)^4 : dV_M(x)$  as will be proved in the next subsection. Basically, they are smoothing operators such that  $K_\varepsilon \rightarrow \mathbb{1}$  in  $\Psi^\delta(M)$  in the *symbol sense* for any  $\delta > 0$  (see definition below). A compact notation is to say  $K_\varepsilon \rightarrow \mathbb{1}$  in  $\Psi^{0+}(M)$ .

**Definition 3.1.** Let  $r \in \mathbb{R}$  we say that operators  $K_\varepsilon \rightarrow K$  in  $\Psi^r(M)$  in the *symbol sense* if for any coordinate chart  $\kappa : U \rightarrow \mathbb{R}^d$  and cut-off  $\chi \in C_c^\infty(U)$ , the full symbol of  $\chi K_\varepsilon \chi$  (considered acting on  $C_c^\infty(\kappa(U))$ ), converges to that of  $\chi K \chi$  in the  $\mathcal{S}_{1,0}^r(\kappa(U) \times \mathbb{R}^d)$  topology.

Now we describe the first candidate for  $K_\varepsilon$  satisfying the above assumption (the proof is in appendix C.2). This was introduced in Dyatlov and Zworski [19] and has the advantage of being *local*, realizing the locality of the  $P(\phi)$  interaction eventually in section 5.5. Consider  $\psi \in C_c^\infty((-1, 1))$  with  $0 \leq \psi \leq 1$  and equal to 1 near 0. For  $\varepsilon > 0$  we define the operator

$$E_\varepsilon u(x) \stackrel{\text{def}}{=} \int_M E_\varepsilon(x, y) u(y) dV_g(y), \quad \text{with } E_\varepsilon(x, y) = \frac{1}{F_\varepsilon(x)} \psi\left(\frac{d_g(x, y)}{\varepsilon}\right). \quad (3.2)$$

Here  $F_\varepsilon(x) = \int \psi(d(x, y)/\varepsilon) dy$  so that  $\int E_\varepsilon(x, y) dy = 1$ , and  $d_g$  denotes the Riemannian distance. One observes that  $E_\varepsilon(x, y)$  is smooth for each  $\varepsilon > 0$  so  $E_\varepsilon : \mathcal{D}'(M) \rightarrow C^\infty(M)$ . Observe also that  $\varepsilon^d/C \leq F_\varepsilon(x) \leq C\varepsilon^d$  for some  $C > 0$ , and this  $C$  could be made dependent neither on  $\varepsilon$  nor on  $x$  as  $M$  is compact.

**Lemma 3.1.** *For any  $\delta > 0$  we have  $E_\varepsilon \rightarrow \mathbb{1}$  in  $\Psi^\delta(M)$  in the symbol sense.*

*Proof.* See appendix C.2. □

Note that  $K_\varepsilon^*(x, y) = K_\varepsilon(y, x)$  for real smoothing operators (and their symbols are related in a simple manner),  $K_\varepsilon \rightarrow \mathbb{1}$  in the symbol sense is equivalent to  $K_\varepsilon^* \rightarrow \mathbb{1}$  in the symbol sense.

**Lemma 3.2.** *Let  $\{E'_\varepsilon\}_{\varepsilon'_1 > 0}$  be another family of smoothing operators such that  $E'_\varepsilon \rightarrow \mathbb{1}$  in  $\Psi^{\delta'}(M)$  in the symbol sense for any  $\delta' > 0$ . Then the net  $E'_\varepsilon (\Delta + m^2)^{-1} (E'_\varepsilon - E_\varepsilon)$ ,  $(\varepsilon', \varepsilon) \in \mathbb{R}_+ \times \mathbb{R}_+$  (we say  $(\varepsilon', \varepsilon) \prec (\varepsilon'_1, \varepsilon_1)$  iff  $\varepsilon' > \varepsilon'_1$  and  $\varepsilon > \varepsilon_1$ ), converges to zero in  $\Psi^{-2+\delta}(M)$  in the symbol sense for any  $\delta > 0$ .*

---

<sup>5</sup>that is, each  $K_\varepsilon$  maps  $\mathcal{D}'(M) \rightarrow C^\infty(M)$ .

*Proof.* Note that following essentially the same arguments as above the  $\Psi^{\delta/2}(M)$  seminorms of  $E_\varepsilon$  can be bounded uniformly in  $\varepsilon$ . This said, the result follows essentially from the continuity of the twisted product (composition product) of symbols as a map  $\mathcal{S}_{1,0}^r \times \mathcal{S}_{1,0}^{r'} \rightarrow \mathcal{S}_{1,0}^{r+r'}$  with respect to the symbol topologies (see Folland [22] page 105 theorem 2.47).  $\square$

We shall consider another set of seminorms on  $\Psi^r(M)$  in the case  $-d < r < 0$  which suits better our purposes. They are defined as follows. Let  $\mathcal{M} \subset C^\infty(M \times M, T(M \times M))$  denote the  $C^\infty(M \times M)$ -module of smooth vector fields tangent to the diagonal in  $M \times M$ . We fix a finite coordinate cover  $\{U_i\}_{i=1}^N$  of  $M$ , with charts  $\kappa_i : U_i \rightarrow \mathbb{R}^d$ , and a partition of unity  $\{\chi_i\}$  subordinate to this cover.

**Definition 3.2.** For any  $K \in C^\infty(M \times M \setminus \text{diag})$ ,  $1 \leq i \leq N$  and  $L_1, \dots, L_p \in \mathcal{M}$ , we define the seminorms

$$p_{i,L_1,\dots,L_p}(K) \stackrel{\text{def}}{=} \sup_{(x,y) \in U_i \times U_i} |(\kappa_i \times \kappa_i)_* ((\chi_i \otimes \chi_i) L_1 \cdots L_p K)(x,y)| \cdot d_g(x,y)^{d+r}, \quad (3.3)$$

while on  $U_i \times U_j$ ,  $i \neq j$ , which does not touch the diagonal, we use the  $C^\infty(U_i \times U_j)$  seminorms. By the kernel topology on  $\Psi^r(M)$ ,  $-d < r < 0$ , we mean the topology induced by these seminorms on the Schwartz kernels  $K_A$  of  $A \in \Psi^r(M)$ . Here  $d_g$  is the distance function.

**Proposition 3.3.** *In the case  $-d < r < 0$ , the above kernel topology is equivalent to the topology induced by symbols  $\mathcal{S}_{1,0}^r(T^*M)$  on  $\Psi^r(M)$ . In particular, if  $A_\varepsilon \rightarrow A$  as  $\varepsilon \rightarrow 0$  in  $\Psi^r(M)$  in the symbol sense then  $A_\varepsilon \rightarrow A$  also in the above kernel topology.*

*Proof.* Essentially in Taylor [77] page 6, proposition 2.2, page 7, proposition 2.4 and page 10, proposition 2.7. See also Bailleul, Dang, Ferdinand and Tô [5] proposition 6.9 for a more detailed treatment.  $\square$

Finally, we observe that the heat operator  $K_\varepsilon = e^{-\varepsilon(\Delta+m^2)}$  is also a valid candidate:

**Lemma 3.4** ([15] lemma 4.15). *We have  $e^{-\varepsilon(\Delta+m^2)} \rightarrow \mathbb{1}$  in  $\Psi^\delta(M)$  in the symbol sense for any  $\delta > 0$ .*  $\square$

Some properties of the heat operator is summed up in appendix C.2.

## 3.2 Integrability of Interaction and Regularization Independence

**Proposition 3.5.** *Let  $(K_\varepsilon)_{\varepsilon>0}$  be any family of real smoothing operators such that  $K_\varepsilon \rightarrow \mathbb{1}$  in  $\Psi^\delta(M)$  in the symbol sense for any  $\delta > 0$ . Define  $\phi_\varepsilon(x)$  and  $:P(\phi_\varepsilon(x)):$  as above and let  $\chi \in C_c^\infty(M)$  be a test function. Put*

$$S_{M,\varepsilon,\chi}(\phi) := \int_M \chi(x) :P(\phi_\varepsilon(x)) : dV_M. \quad (3.4)$$

*This is a random variable on  $\mathcal{D}'(M)$  equipped with  $\mu_{\text{GFF}}^M$ . Then  $\{S_{M,\chi,\varepsilon}\}$  converges in  $L^2(\mu_{\text{GFF}}^M)$  as  $\varepsilon \rightarrow 0$ , and the limit is independent of the specific smoothing  $(K_\varepsilon)$  chosen, provided they have the convergence property described above.*

*More precisely, for any other smoothing family  $(\tilde{K}_{\varepsilon'})$  satisfying the same condition and defining the random variable  $\tilde{S}_{M,\varepsilon',\chi}$ , we have a quantitative bound of the form*

$$\mathbb{E}[|S_{M,\varepsilon,\chi} - \tilde{S}_{M,\varepsilon',\chi}|^2] \leq \mathfrak{D}(|\varepsilon - \varepsilon'|) C_n \|\chi\|_{L^4}^2,$$

*where  $\mathfrak{D}(|\varepsilon - \varepsilon'|)$  is a function going to zero as  $|\varepsilon - \varepsilon'| \rightarrow 0$ , depending only on  $M$ .*

A particularity of the dimension two is seen in the following elementary lemma.

**Lemma 3.6.** *If  $\dim M = 2$  and  $g$  is a smooth Riemannian metric on  $M$  then*

$$\iint_{M \times M} |\log(d_g(x,y))|^p dV_{M \times M} \leq C_p < \infty \quad (3.5)$$

*for any  $1 \leq p < \infty$  and*

$$\iint_{M \times M} d_g(x,y)^{-\delta} dV_{M \times M} \leq C_\delta < \infty \quad (3.6)$$

*for any  $0 < \delta < 1$ ,  $d_g$  denoting the distance function.*  $\square$

Consequently, since in two dimensions  $G(x, y) = \mathcal{O}(\log(d(x, y)))$  as  $y \rightarrow x$  (see lemma 3.12), a simple argument with partition of unity shows

$$\iint_{M \times M} \chi(x)\chi(y)|G(x, y)|^p dV_{M \times M} \leq C_p \|\chi\|_{L^\infty}^2 \text{vol}(\text{supp}(\chi))^2 < \infty \quad (3.7)$$

for some  $C_p > 0$ , for all  $1 \leq p < \infty$ . When  $p = 2$  this is the familiar fact that in two dimensions the Green operator  $(\Delta + m^2)^{-1}$  is Hilbert-Schmidt, which may also be shown using Weyl's law.

We will denote

$$\mathbb{E}_{\text{GFF}}^M [\phi_\varepsilon(x)\phi_\varepsilon(y)] = \langle \delta_x, K_\varepsilon^*(\Delta + m^2)^{-1} K_\varepsilon \delta_y \rangle_{L^2} \stackrel{\text{def}}{=} G_{\varepsilon, \varepsilon}(x, y), \quad (3.8)$$

$$\mathbb{E}_{\text{GFF}}^M [\phi_\varepsilon(x)\phi_{\varepsilon'}(y)] = \langle \delta_x, K_\varepsilon^*(\Delta + m^2)^{-1} \tilde{K}_{\varepsilon'} \delta_y \rangle_{L^2} \stackrel{\text{def}}{=} G_{\varepsilon, \varepsilon'}(x, y), \quad (3.9)$$

$$\mathbb{E}_{\text{GFF}}^M [\phi_{\varepsilon'}(x)\phi_{\varepsilon'}(y)] = \langle \delta_x, \tilde{K}_{\varepsilon'}^*(\Delta + m^2)^{-1} \tilde{K}_{\varepsilon'} \delta_y \rangle_{L^2} \stackrel{\text{def}}{=} G_{\varepsilon', \varepsilon'}(x, y). \quad (3.10)$$

Note that  $G_{\varepsilon, \varepsilon'}$  has a *different* regulator on the second variable! For fixed  $\varepsilon > 0$ , we have by lemma B.9,

$$:\phi_\varepsilon(x)^{2n}: = G_{\varepsilon, \varepsilon}(x, x)^n h_{2n}(\phi_\varepsilon(x)/G_{\varepsilon, \varepsilon}(x, x)^{\frac{1}{2}}) \geq -b_1 G_{\varepsilon, \varepsilon}(x, x)^n, \quad (3.11)$$

for some constant  $b_1 > 0$  independent of  $\varepsilon$ , since the even-degree Hermite polynomial  $h_{2n}$  is bounded below.

*Proof of proposition 3.5.* We shall prove the proposition for the case  $P(\theta) = \theta^{2n}$ ,  $n \in \mathbb{N}$ , the general case is similar. For fixed  $\varepsilon, \varepsilon' > 0$ , we compute

$$\begin{aligned} \|S_{M, \varepsilon} - S_{M, \varepsilon'}\|_{L^2(\mu_{\text{GFF}})}^2 &= \mathbb{E} \left[ \left| \int_M \chi : \phi_\varepsilon(x)^{2n} : dV_M - \int_M \chi : \phi_{\varepsilon'}(x)^{2n} : dV_M \right|^2 \right] \\ &= \mathbb{E} \left[ \iint_{M \times M} : \chi(x) \phi_\varepsilon(x)^{2n} : : \phi_\varepsilon(y)^{2n} : \chi(y) dV_M \otimes dV_M \right] \\ &\quad - 2\mathbb{E} \left[ \iint_{M \times M} \chi(x) : \phi_\varepsilon(x)^{2n} : : \phi_{\varepsilon'}(y)^{2n} : \chi(y) dV_M \otimes dV_M \right] \\ &\quad + \mathbb{E} \left[ \iint_{M \times M} \chi(x) : \phi_{\varepsilon'}(x)^{2n} : : \phi_{\varepsilon'}(y)^{2n} : \chi(y) dV_M \otimes dV_M \right] \\ &= \iint_{M \times M} \left( \mathbb{E} [\chi(x) : \phi_\varepsilon(x)^{2n} : : \phi_\varepsilon(y)^{2n} : \chi(y)] - 2\mathbb{E} [\chi(x) : \phi_\varepsilon(x)^{2n} : : \phi_{\varepsilon'}(y)^{2n} : \chi(y)] \right. \\ &\quad \left. + \mathbb{E} [\chi(x) : \phi_{\varepsilon'}(x)^{2n} : : \phi_{\varepsilon'}(y)^{2n} : \chi(y)] \right) dV_{M \times M} \quad (\text{Tonelli}) \\ &= (2n)! \iint_{M \times M} \chi(x)\chi(y) (G_{\varepsilon, \varepsilon}(x, y)^{2n} - 2G_{\varepsilon, \varepsilon'}(x, y)^{2n} + G_{\varepsilon', \varepsilon'}(x, y)^{2n}) dV_{M \times M} \end{aligned} \quad (\text{lemma B.9})$$

We will control the integral  $\iint_{M \times M} \chi(x)\chi(y) (G_{\varepsilon, \varepsilon}(x, y)^{2n} - G_{\varepsilon, \varepsilon'}(x, y)^{2n}) dV_{M \times M}$  for  $\chi \in C^\infty(M)$  and  $\chi \geq 0$ . Indeed,

$$\begin{aligned} |\text{this integral}| &\leq \iint_{M \times M} \chi(x)\chi(y) |G_{\varepsilon, \varepsilon}(x, y) - G_{\varepsilon, \varepsilon'}(x, y)| \cdot |G_{\varepsilon, \varepsilon}(x, y)^{2n-1} \\ &\quad + G_{\varepsilon, \varepsilon}(x, y)^{2n-2} G_{\varepsilon, \varepsilon'}(x, y) + \cdots + G_{\varepsilon, \varepsilon'}(x, y)^{2n-1}| dV_{M \times M} \\ &\leq 2n \|\chi\|_{L^4}^2 (C_{8n-4})^{\frac{1}{4}} \times \left( \iint_{M \times M} |G_{\varepsilon, \varepsilon}(x, y) - G_{\varepsilon, \varepsilon'}(x, y)|^2 \right)^{\frac{1}{2}} \end{aligned}$$

Remember that  $G_{\varepsilon, \varepsilon'}(x, y)$  is the kernel of  $K_\varepsilon^*(\Delta + m^2)^{-1} \tilde{K}_{\varepsilon'}$  and  $G_{\varepsilon, \varepsilon}(x, y) - G_{\varepsilon, \varepsilon'}(x, y)$  is the kernel of  $K_\varepsilon^*(\Delta + m^2)^{-1}(\tilde{K}_{\varepsilon'} - K_\varepsilon)$ . By lemma 3.2, definition 3.2 and proposition 3.3,

$$|G_{\varepsilon, \varepsilon'}(x, y)| \leq C_{M, \delta} d_g(x, y)^{-\delta}, \quad (3.12)$$

uniformly in  $(\varepsilon, \varepsilon')$  and by lemma 3.2,

$$|G_{\varepsilon, \varepsilon'}(x, y) - G_{\varepsilon, \varepsilon}(x, y)| \leq \mathfrak{D}_{M, \delta} (|\varepsilon - \varepsilon'|) d_g(x, y)^{-\delta} \quad (3.13)$$

for any  $\delta > 0$ . If we restrict moreover to  $\delta < 1$ , then we prove our result, thanks to lemma 3.6.  $\square$

### 3.3 Integrability of the Exponential of Interaction

In this subsection we will adopt the heat regulator  $K_\varepsilon = \widetilde{K}_\varepsilon := e^{-\varepsilon(\Delta+m^2)}$ .

**Lemma 3.7.** *Let  $\deg P = 2n$ . Then  $S_{M,\chi,\varepsilon}$ ,  $\varepsilon > 0$ , and hence the resulting limit  $S_{M,\chi}$ , is in the  $(2n)$ -th Wiener chaos of the GFF, that is,  $\overline{\mathcal{P}}_{2n}(\mathcal{H})$ , where  $\mathcal{H} = W^{-1}(M)$  is the Gaussian Hilbert space of the GFF.*

*Proof.* We shall show it for  $P(\theta) = \theta^4$  and the general case is similar. Here we use the spectral representation of remark 2.2 (using the notations thereof) and take  $K_\varepsilon := e^{-\varepsilon(\Delta+m^2)}$ . Thus we can write

$$\begin{aligned} \phi_\varepsilon(x)^4 &= \left( \sum_{j=0}^{\infty} \langle \varphi_j, K_\varepsilon \delta_x \rangle_{L^2} \xi_j \right)^4 = \left( \sum_{j=0}^{\infty} e^{-\varepsilon(\lambda_j+m^2)} \varphi_j(x) \xi_j \right)^4 \\ &= \sum_{j,k,\ell,p} e^{-\varepsilon(\lambda_j+\lambda_k+\lambda_\ell+\lambda_p+4m^2)} \varphi_j(x) \varphi_k(x) \varphi_\ell(x) \varphi_p(x) \xi_j \xi_k \xi_\ell \xi_p, \end{aligned}$$

the series converging absolutely in  $L^2(\mu_{\text{GFF}}^M)$ . Now each individual term is clearly in  $\overline{\mathcal{P}}_4(\mathcal{H})$ . Since

$$\int |\varphi_j(x) \varphi_k(x) \varphi_\ell(x) \varphi_p(x)| dV_M(x) \lesssim \text{vol}(M) \cdot [\text{polynomial in } \lambda_j, \lambda_k, \lambda_\ell, \lambda_p \text{ with fixed degree}] \quad (3.14)$$

as one has  $\sup_M |\varphi_j| \lesssim (1 + \lambda_j)^2$  in two dimensions which follows essentially from the Sobolev embedding (see Sogge [74] page 43 equation (3.1.12)), thus

$$\int \chi(x) \phi_\varepsilon(x)^4 dV_M(x) = \sum_{j,k,\ell,p} e^{-\varepsilon(\lambda_j+\lambda_k+\lambda_\ell+\lambda_p+4m^2)} C_{\chi,j,k,\ell,p} \xi_j \xi_k \xi_\ell \xi_p \quad (3.15)$$

with the series converging absolutely in  $L^2(\mu_{\text{GFF}}^M)$  and the result is in  $\overline{\mathcal{P}}_4(\mathcal{H})$ .  $\square$

**Proposition 3.8** (hypercontractivity, [39] theorem 5.10, [71] theorem I.22). *Let  $\mathcal{H} \subset L^2(Q, \mathcal{O}, \mathbb{P})$  be a Gaussian Hilbert space on some probability space  $(Q, \mathcal{O}, \mathbb{P})$ , and let  $n \geq 1$ ,  $2 \leq p < \infty$ . Then*

$$\mathbb{E}[|X|^p]^{\frac{1}{p}} \leq (p-1)^{\frac{n}{2}} \mathbb{E}[X^2]^{\frac{1}{2}} \quad (3.16)$$

for all  $X \in \overline{\mathcal{P}}_n(\mathcal{H})$ .  $\square$

Combining lemma 3.7 and proposition 3.8, we have

**Corollary 3.9.** *The convergence of  $\{S_{M,\chi,\varepsilon}\}$ , as well as the limit  $S_{M,\chi}$ , is in  $L^p(\mu_{\text{GFF}})$  for all  $1 \leq p < \infty$ . Moreover, if  $\chi \rightarrow 0$  in  $L^4(M)$  then*

$$\lim_{\varepsilon \rightarrow 0} \lim_{\chi \rightarrow 0} S_{M,\varepsilon,\chi} = 0$$

as random variable in  $L^p(\mu_{\text{GFF}}^M)$  for all  $1 \leq p < \infty$ . Moreover  $\mathbb{E}[e^{-S_{M,\varepsilon,\chi}}]$  remains uniformly bounded along the limit. In particular  $S_{M,\chi}$  is defined for  $\chi \in L^4(M)$ .  $\square$

We single out a calculus computation which will be used in the sequel:

**Lemma 3.10.** *Let  $a, b$  be positive real numbers. Then the real function  $\alpha(x) := x^{(bx)} a^x$ ,  $x > 0$ , attains its minimum value  $e^{-be^{-1}a^{-1/b}}$  at  $x = e^{-1}a^{-1/b}$ .  $\square$*

**Theorem** (Nelson). *We have for  $\chi \in L^4(M)$ ,*

$$e^{-S_{M,\chi}} \in L^1(\mu_{\text{GFF}}) \quad (3.17)$$

and hence

$$Z_M = \det_\zeta(\Delta_g + m^2)^{-\frac{1}{2}} \int_{\mathcal{D}'(M)} e^{-\int_M :P(\phi): dV_M} d\mu_{\text{GFF}}^M(\phi) < \infty. \quad (3.18)$$

*Proof.* For any  $\varepsilon > 0$ , by (3.7) and (3.11),

$$S_{M,\chi,\varepsilon} \geq -b \text{vol}(\text{supp}(\chi)) \|\chi\|_{L^\infty} \sup_x (|G_{\varepsilon,\varepsilon}(x, x)|^n). \quad (3.19)$$



From formula (C.16), for  $\varepsilon$  small,

$$G_{\varepsilon,\varepsilon}(x, x) = \int_{2\varepsilon}^{\infty} p_t(x, x) dt = \underbrace{\left( \int_{2\varepsilon}^1 \right)}_A + \underbrace{\left( \int_1^{\infty} \right)}_B p_t(x, x) dt. \quad (3.20)$$

Now by (iii) of lemma C.5 part *A* is  $\mathcal{O}(\log(2\varepsilon))$ ; since our field is *massive* ( $m > 0$ ), by (iv) of lemma C.5 part *B* is *bounded*. Therefore one has overall  $G_{\varepsilon,\varepsilon}(x, x) = \mathcal{O}(\log(2\varepsilon))$ . As a result  $S_{M,\chi,\varepsilon} \geq -b_2 |\log(2\varepsilon)|^n$  for  $\varepsilon$  small.

Now we compute that

$$\begin{aligned} \mathbb{P}(e^{-S_{M,\chi}} \geq e^{b_2 |\log(2\varepsilon)|^n + 1}) &= \mathbb{P}(S_{M,\chi} \leq -b_2 |\log(2\varepsilon)|^n - 1) \\ &\leq \mathbb{P}(|S_{M,\chi} - S_{M,\chi,\varepsilon}| \geq 1) \\ &\leq \|S_{M,\chi} - S_{M,\chi,\varepsilon}\|_{L^p(\mu_{\text{GFF}})}^p \quad (\text{Chebyshev}) \\ &\leq (p-1)^{\frac{np}{2}} C_1^p \varepsilon^{\frac{n}{2}} \|\chi\|_{L^4}^p \quad (\text{proposition 3.8 and 3.5}) \\ &\lesssim \|\chi\|_{L^4}^p p^{\frac{n}{2}p} (C_1 \varepsilon^{\frac{1}{2}})^p, \end{aligned}$$

for all  $2 \leq p < \infty$ . The last line as a function of  $p$  has the form dealt with in lemma 3.10 and attains a minimum of  $\exp(-C_2(\varepsilon^{\frac{1}{2}} \|\chi\|_{L^4})^{-1/n})$ , with some absorbed constant  $C_2 > 0$  which does not depend on  $\chi$ . Thus we obtain

$$\mathbb{P}(e^{-S_{M,\chi}} \geq e^{b_2 |\log(2\varepsilon)|^n + 1}) \lesssim \exp(-C_2(\varepsilon^{\frac{1}{2}} \|\chi\|_{L^4})^{-\frac{1}{n}}). \quad (3.21)$$

Now we may conclude with the formula

$$\mathbb{E}[e^{-S_{M,\chi}}] = \int_0^{\infty} \mathbb{P}(e^{-S_{M,\chi}} \geq t) dt = \int_0^1 \mathbb{P}(e^{-S_{M,\chi}} \geq t) \left| \frac{dt}{d\varepsilon} \right| d\varepsilon$$

where the last integral involves a change of variable  $t := e^{b_2 |\log(2\varepsilon)|^n + 1}$ . This gives

$$\mathbb{E}[e^{-S_{M,\chi}}] \lesssim 1 + \int_0^1 \exp(-C_2(\varepsilon^{\frac{1}{2}} \|\chi\|_{L^4})^{-\frac{1}{n}}) C_3 \varepsilon^{-1} n |\log(\varepsilon)|^{n-1} e^{c |\log(\varepsilon)|^n - 1} d\varepsilon$$

which is finite since integrable near  $\varepsilon = 0$ . Moreover, we see that the bound is uniform when  $\|\chi\|_{L^4} \leq C_0$  for some given  $C_0 > 0$ .  $\square$

### 3.4 Change of Wick Ordering

In order for the proof of proposition 3.5 to work as it is written one has to insist on the Wick ordering  $\bullet \bullet$  provided by  $\mu_{\text{GFF}}^M$ , since we desire convergence in  $L^2(\mu_{\text{GFF}}^M)$  and with a different Wick ordering the Feynman rules (lemma B.9) are not exact. Nevertheless, in order to define the interaction over a domain  $\Omega$  independently of its embedding in an ambient manifold  $M$ , one must employ a Wick ordering independent of  $M$ , or in other words, one that is *local*.

Let  $d$  denote the Riemannian distance function of  $M$ . This function is local in the sense that  $d(x, y)$  (as  $y \rightarrow x$ ) depends only on the restriction of the Riemannian metric on any geodesic convex neighborhood containing  $x$  and  $y$ . The local Wick ordering  $\bullet \bullet_0$  is provided by the *log-measure*  $\mu_{\text{log}}^M$  which is the Gaussian measure on  $\mathcal{D}'(M)$  with covariance

$$\mathbb{E}_{\text{log}}[\phi(f)\phi(h)] = \int_M f(x) \left( -\frac{1}{2\pi} \log(m \cdot d(x, y)) \right) h(y) dV_M(x) dV_M(y), \quad (3.22)$$

for  $f, h \in C^\infty(M)$ , thanks to lemma 3.6. Here  $m$  is the mass used for  $(\Delta + m^2)^{-1}$ . We denote  $-\frac{1}{2\pi} \log(md(x, y)) := C_0(x, y)$  and the corresponding operator by  $C_0$ . We emphasize here that  $\mu_{\text{log}}^M$  is used only as a tool to produce a linear change of random variables with deterministic coefficients, no random variables will be actually defined on  $\mu_{\text{log}}^M$ .

**Lemma 3.11.** *If  $C_1, C_2$  are two covariance operators on  $M$ , then*

$$\bullet \phi(f)^n \bullet_{C_1} = \sum_{j=0}^{\lfloor n/2 \rfloor} \frac{n!}{(n-2j)! j! 2^j} \langle f, (C_2 - C_1)f \rangle_{L^2(M)}^j \bullet \phi(f)^{n-2j} \bullet_{C_2}, \quad (3.23)$$

for  $f \in C^\infty(M)$ .

*Proof.* Follows readily from Wick's theorem.  $\square$

The reason why the new Wick ordering works is the following. Let  $G_{(\Delta+m^2)}(x, y)$  denote the integral kernel of  $(\Delta + m^2)^{-1}$ .

**Lemma 3.12.** *For each  $x \in M$ , the limit*

$$\lim_{y \rightarrow x} (G_{(\Delta+m^2)}(x, y) - C_0(x, y)) \stackrel{\text{def}}{=} \delta G(x) \quad (3.24)$$

*exists, and that  $\delta G \in L^p(M)$  for all  $1 \leq p < \infty$ .*

*Remarks for proof.* The function  $\delta G$  is called in our context the (point-splitting) tadpole function (see Kandel, Mnev and Wernli [36] section 5.4, in particular lemma 5.20 for a precise expression), which can be seen as a renormalized diagonal value of the Green function  $G_{(\Delta+m^2)}(x, x)$ . The asymptotic of the Green function along the diagonal is a classical subject and we have in fact  $G_{(\Delta+m^2)}(x, y) - C_0(x, y) \in C^1(M \times M)$ . The function  $\delta G$  is also important in the context of conformal geometry where it is called the mass function, if more precisely we do not include the constant  $-\log m/2\pi$  in  $C_0$  but rather in  $\delta G$ . See Hermann and Humbert [33], Ludewig [44] or Schoen and Yau [64] for more information.  $\square$

It follows that

$$\langle E_\varepsilon \delta_x, ((\Delta + m^2)^{-1} - C_0) E_\varepsilon \delta_x \rangle_{L^2(M)} \longrightarrow \delta G(x) \quad (3.25)$$

as  $\varepsilon \rightarrow 0$  and through the limiting process of proposition 3.5 we find

$$\int_M \chi(x) : \phi(x)^{2n} :_0 dV_M(x) = \sum_{j=0}^n \frac{(2n)!}{(2n-2j)! j! 2^j} \int \chi(x) \delta G(x)^j : \phi(x)^{2n-2j} :_{\text{GFF}} dV_M(x), \quad (3.26)$$

which exist as a random variable in  $L^p(\mu_{\text{GFF}}^M)$  for all  $2 \leq p < \infty$  since each term on the RHS are such by proposition 3.5 and corollary 3.9.

## 4 The Trace Operator and the Poisson Integral Operator

In this section we obtain a series of rather elementary relations between various geometric-analytic operators on  $M$  and on  $\Omega$ . The moral is that, the so-called ‘‘sharp-time localization’’ map  $j_\Sigma$  (see lemma 4.5), induced probability laws of Gaussian fields under the trace  $\tau_\Sigma$  (see section 4.4), and finally the Green-Stokes formula, are largely different aspects of the same thing.

### 4.1 Summary of Essential Properties

**Definition 4.1.** Let  $(M, g)$  be a closed Riemannian manifold, and  $\Sigma \subset M$  a smooth embedded hypersurface (codimension one submanifold). Then we call the map

$$\begin{aligned} \tau_\Sigma : C^\infty(M) &\longrightarrow C^\infty(\Sigma), \\ f &\longmapsto f|_\Sigma, \end{aligned} \quad (4.1)$$

the trace map (or restriction map) from  $M$  onto  $\Sigma$ .

**Lemma 4.1** ([76] page 334, [20] page 57, example 13.3). *Let  $(M, g)$  and  $\Sigma$  be as above. Then*

- (i) *the map  $\tau_\Sigma$  extends uniquely to a continuous operator  $\tau_\Sigma : W^s(M) \longrightarrow W^{s-\frac{1}{2}}(\Sigma)$  for each  $s > \frac{1}{2}$ ;*
- (ii) *the map  $\tau_\Sigma : W^s(M) \longrightarrow W^{s-\frac{1}{2}}(\Sigma)$ ,  $s > \frac{1}{2}$ , is surjective.*

*Remark for proof.* For (ii), the case of  $M = \mathbb{R}^d$  and  $\Sigma = \mathbb{R}^{d-1} \subset \mathbb{R}^d$  is treated in [20] as indicated above. One could then supply a partition of unity argument to extend to the current case, remembering that  $\Sigma$  is smoothly embedded.  $\square$

Now let  $(\Omega, g)$  be a compact Riemannian manifold with boundary  $\partial\Omega$ . It is well-known that for  $f \in C^\infty(\partial\Omega)$ , the (Helmholtz) boundary value problem

$$\begin{cases} (\Delta_\Omega + m^2)u = 0, & \text{in } \Omega, \\ u|_{\partial\Omega} = f, & \text{on } \partial\Omega, \end{cases} \quad (4.2)$$

admits a unique solution  $u \in C^\infty(\overline{\Omega})$  which is extendably smooth upto  $\partial\Omega$ .

**Remark 4.1.** In this paper, a function  $u \in C^\infty(\overline{\Omega})$  is said to satisfy the Dirichlet condition (respectively Neumann) if  $u|_{\partial\Omega} = 0$  (respectively  $(\partial_\nu u)|_{\partial\Omega} = 0$ ,  $\nu$  the outward unit normal). The  $f$  appearing in (4.2) is called a Dirichlet datum.

**Definition 4.2.** We call the operator

$$\begin{aligned} \text{PI}_\Omega^{\partial\Omega} : C^\infty(\partial\Omega) &\longrightarrow C^\infty(\overline{\Omega}), \\ f &\longmapsto u, \end{aligned} \quad (4.3)$$

where  $u$  is the unique solution of (4.2), the Poisson integral operator (or Harmonic extension operator) from  $\partial\Omega$  to  $\Omega$ , with mass  $m > 0$ .

**Lemma 4.2** ([76] page 361, proposition 1.8). *Let  $(\Omega, g)$  be as above. Then  $\text{PI}_\Omega^{\partial\Omega}$  extends uniquely to a continuous operator*

$$\text{PI}_\Omega^{\partial\Omega} : W^s(\partial\Omega) \longrightarrow W^{s+\frac{1}{2}}(\Omega) \quad (4.4)$$

for each  $s \geq -\frac{1}{2}$ .  $\square$

We shall need a variant of the Poisson integral operator that works for embedded hypersurfaces in closed manifolds. Let  $(M, g)$  and  $\Sigma$  be as in definition 4.1. Pick  $f \in C^\infty(\Sigma)$ . This time we look at the boundary value problem

$$\begin{cases} (\Delta_M + m^2)u = 0, & \text{in } M \setminus \Sigma, \\ u|_\Sigma = f, & \text{on } \Sigma. \end{cases} \quad (4.5)$$

Indeed, one should view  $M \setminus \Sigma$  as a manifold with two boundaries  $\Sigma \sqcup \Sigma$ , and as a result one obtains a unique solution  $u$  which is smooth on  $M \setminus \Sigma$  and one-sidedly smooth upto  $\Sigma$  respectively on its two sides.

**Definition 4.3.** We call the operator

$$\begin{aligned} \text{PI}_M^\Sigma : C^\infty(\Sigma) &\longrightarrow C^\infty(\overline{M \setminus \Sigma}), \\ f &\longmapsto u, \end{aligned} \quad (4.6)$$

where  $u$  is the unique solution of (4.5), the Poisson integral operator from  $\Sigma$  to  $M$ , with mass  $m > 0$ .

**Definition 4.4.** More generally if  $\Omega$  has boundaries and  $\Sigma$  is either one component of  $\partial\Omega$  or embedded in the interior of  $\Omega$ , then we denote by  $\text{PI}_\Omega^{\Sigma, B} f$  the solution with its restriction equal to  $f$  on  $\Sigma$  and boundary condition “ $B$ ” (Dirichlet or Neumann) on all components of  $\partial\Omega$  except  $\Sigma$ . Such notations raise no ambiguity when the situation is understood from context.

**Lemma 4.3.** *Let  $(M, g)$  and  $\Sigma$  be as above. Then  $\text{PI}_M^\Sigma$  extends uniquely to a continuous operator*

$$\text{PI}_M^\Sigma : W^{\frac{1}{2}}(\Sigma) \longrightarrow W^1(M). \quad (4.7)$$

*Proof.* Let  $f \in W^{\frac{1}{2}}(\Sigma)$  and  $u := \text{PI}_M^\Sigma f$ . We know that  $u \in W^1(M \setminus \Sigma) \subset \mathcal{D}'(M \setminus \Sigma)$ , this means  $u \in L^2(M \setminus \Sigma) = L^2(M)$ , and  $\nabla u \in L^2(T(M \setminus \Sigma))$ , as a distribution over  $M \setminus \Sigma$ . The problem is to compute  $\nabla u$  as a distribution over  $M$ . For this, one picks a testing vector field  $X \in C^\infty(M, TM)$  and applies the Green-Stokes formula to get

$$\begin{aligned} \langle \nabla u, X \rangle_{L^2(M, TM)} &\stackrel{\text{def}}{=} -\langle u, \text{div } X \rangle_{L^2(M)} \\ &= \int_{M \setminus \Sigma} \langle \nabla u, X \rangle_g \, dV_M - \int_\Sigma f \langle X, \nu \rangle_g \, dV_\Sigma - \int_\Sigma f \langle X, -\nu \rangle_g \, dV_\Sigma \\ &= \int_{M \setminus \Sigma} \langle \nabla u, X \rangle_g \, dV_M. \end{aligned}$$

Here  $\nu$  is any one of the two possible unit normal vector fields along  $\Sigma$ . This shows that, nevertheless,

$$\nabla^M u = \nabla^{M \setminus \Sigma} u, \quad (4.8)$$

and hence  $\|u\|_{W^1(M)} = \|\nabla u\|_{L^2(T(M \setminus \Sigma))} + \|u\|_{L^2(M)} \approx \|u\|_{W^1(M \setminus \Sigma)}$ .  $\square$

## 4.2 The Dirichlet-to-Neumann Operator and its Jumpy Version

**Definition 4.5.** Let  $(\Omega, g)$  be a compact Riemannian manifold with boundary  $\partial\Omega \neq \emptyset$ , and  $\text{PI}_\Omega^{\partial\Omega} : C^\infty(\partial\Omega) \rightarrow C^\infty(\bar{\Omega})$  the Poisson operator defined previously. Put

$$\begin{aligned} \text{DN}_\Omega^{\partial\Omega} : C^\infty(\partial\Omega) &\longrightarrow C^\infty(\partial\Omega), \\ f &\longmapsto \partial_\nu(\text{PI}_\Omega^{\partial\Omega} f), \end{aligned} \quad (4.9)$$

where  $\nu =$  outward unit normal along  $\partial\Omega$ , called the Dirichlet-to-Neumann operator on  $\partial\Omega$  with respect to  $\Omega$ .

**Remark 4.2.** As its name suggests,  $\text{DN}_\Omega^{\partial\Omega}$  computes the associated Neumann data of the solution to the Helmholtz equation with given Dirichlet data. By the well-posedness and regularity of the Neumann problem, one could also do it the other way round: compute the Dirichlet data out of a Neumann data. In this way one sees that  $(\text{DN}_\Omega^{\partial\Omega})^{-1}$  can be defined and maps  $C^\infty(\partial\Omega) \rightarrow C^\infty(\partial\Omega)$ .

**Definition 4.6.** Let  $(M, g)$  be a closed Riemannian manifold,  $\Sigma \subset M$  an embedded hypersurface, and  $\text{PI}_M^\Sigma : C^\infty(\partial\Omega) \rightarrow C^\infty(M \setminus \Sigma)$  the hypersurface Poisson operator. Put

$$\begin{aligned} \text{DN}_M^\Sigma : C^\infty(\partial\Omega) &\longrightarrow C^\infty(\partial\Omega), \\ f &\longmapsto \partial_\nu(\text{PI}_M^\Sigma f)|_{\Sigma_-} + \partial_{-\nu}(\text{PI}_M^\Sigma f)|_{\Sigma_+}, \end{aligned} \quad (4.10)$$

where  $\nu$  is any one of the two unit normal vector fields along  $\Sigma$ , extended over a cylindrical neighborhood of  $\Sigma$ . Here  $\Sigma_-$  and  $\Sigma_+$  means that we are taking one-sided derivatives, respectively, from the backward-time and forward-time directions with regard to the flow of  $\nu$ . We call  $\text{DN}_M^\Sigma$  the jumpy Dirichlet-to-Neumann operator on  $\Sigma$  with respect to  $M$ .

**Definition 4.7.** Similarly if  $\Omega$  has boundaries and  $\Sigma$  is *either* one component of  $\partial\Omega$  or embedded in the *interior* of  $\Omega$ , then we denote by  $\text{DN}_\Omega^{\Sigma, B}$  the corresponding operator with  $\text{PI}_M^\Sigma$  replaced by  $\text{PI}_\Omega^{\Sigma, B}$  in the definition (see definition 4.4).

**Remark 4.3.** If we see  $M \setminus \Sigma$  as a manifold with boundary  $\Sigma \sqcup \Sigma$ , then  $\text{DN}_M^\Sigma f$  is also the sum over  $\Sigma$  of the two *outward* unit normal derivatives of  $\text{PI}_M^\Sigma f$  along  $\Sigma \sqcup \Sigma$ . Intuitively,  $\text{DN}_M^\Sigma f$  describes the ‘‘jump’’ of  $\nabla \text{PI}_M^\Sigma f$  across  $\Sigma$ , whence its name.

We summarize in the following lemma the essential properties of  $\text{DN}_\Omega^{\partial\Omega}$  and  $\text{DN}_M^\Sigma$ . Parallel results also hold for  $\text{DN}_\Omega^{\Sigma, B}$  ( $\Sigma$  being either one component of boundary or embedded in interior) but we shall not discuss them in order to simplify the presentation. The same applies to everything below this section.

**Lemma 4.4.** *Under their respective settings,  $\text{DN}_\Omega^{\partial\Omega}$  and  $\text{DN}_M^\Sigma$  are such that*

(i) *their quadratic forms are given respectively by the Dirichlet energies of their harmonic extensions:*

$$\langle f, \text{DN}_\Omega^{\partial\Omega} f \rangle_{L^2(\partial\Omega)} = \int_\Omega (|\nabla \text{PI}_\Omega^{\partial\Omega} f|_g^2 + m^2(\text{PI}_\Omega^{\partial\Omega} f)^2) dV_\Omega, \quad (4.11)$$

$$\langle f, \text{DN}_M^\Sigma f \rangle_{L^2(\Sigma)} = \int_M (|\nabla \text{PI}_M^\Sigma f|_g^2 + m^2(\text{PI}_M^\Sigma f)^2) dV_M, \quad (4.12)$$

for  $f \in C^\infty(\partial\Omega)$ ;

- (ii) *they are formally self-adjoint, strictly positive, and  $L^2$ -invertible;*
- (iii) *they are elliptic  $\Psi$ DOs of order 1, with principal symbols being  $|\xi|_g$  and  $2|\xi|_g$  respectively;*
- (iv) *they afford a finer comparison with  $\mathbf{D}_{\partial\Omega} = (\Delta_{\partial\Omega} + m^2)^{\frac{1}{2}}$  or  $2\mathbf{D}_\Sigma = 2(\Delta_\Sigma + m^2)^{\frac{1}{2}}$ : the operators*

$$\text{DN}_\Omega^{\partial\Omega} - \mathbf{D}_{\partial\Omega}, \quad \text{DN}_M^\Sigma - 2\mathbf{D}_\Sigma, \quad \mathbf{D}_{\partial\Omega}^{-1} \text{DN}_\Omega^{\partial\Omega} - \mathbb{1}, \quad \text{and} \quad (2\mathbf{D}_\Sigma)^{-1} \text{DN}_M^\Sigma - \mathbb{1}, \quad (4.13)$$

*are  $\Psi$ DOs of orders at most  $-2$ ,  $-2$ ,  $-3$ , and  $-3$  respectively. A fortiori, they are all of trace class when  $\dim \Omega = \dim M = 2$  and  $\dim \Sigma = \dim \partial\Omega = 1$ .*

*Proof.* See Taylor [78] and the references therein. For (iv) see Kandel, Mnev and Wernli [36] proposition A.3.  $\square$

## 4.3 Two Consequences of the Green-Stokes Formula

Let  $(M, g)$  be a closed Riemannian manifold and  $\Sigma \subset M$  an embedded hypersurface. Formula (4.12) in a slightly more general form allows one to obtain an expression for the ‘‘distributional adjoint’’ of the trace map onto  $\Sigma$ .

**Lemma 4.5.** *Let  $\tau_\Sigma : C^\infty(M) \rightarrow C^\infty(\Sigma)$ ,  $\phi \mapsto \phi|_\Sigma$  be the trace map. One has, for any  $\phi \in C^\infty(M)$  and  $f \in C^\infty(\Sigma)$ ,*

$$\langle \tau_\Sigma \phi, f \rangle_{L^2(\partial\Omega)} = \langle \phi, j_\Sigma f \rangle_{L^2(M)} \quad (4.14)$$

where  $j_\Sigma = (\Delta + m^2) \text{PI}_M (\text{DN}_M^\Sigma)^{-1}$ . Moreover, this equality can be extended to  $\phi \in W^1(M)$  and  $f \in W^{-\frac{1}{2}}(\Sigma)$ .

*Proof.* First suppose  $h \in C^\infty(\Sigma)$ , then applying the Green-Stokes formula to  $M \setminus \Sigma$  with boundary  $\Sigma \sqcup \Sigma$  gives

$$\begin{aligned} \langle \tau_\Sigma \phi, \text{DN}_M^\Sigma h \rangle_{L^2(\Sigma)} &= \int_{M \setminus \Sigma} ((\nabla \phi, \nabla \text{PI}_M h) + m^2 \phi (\text{PI}_M h)) dV_M \\ &= \int_M ((\nabla \phi, \nabla \text{PI}_M h) + m^2 \phi (\text{PI}_M h)) dV_M \quad (\text{lemma 4.3}) \\ &= \langle \phi, (\Delta_M + m^2) \text{PI}_M^\Sigma h \rangle_{L^2(M)}. \quad (\#) \end{aligned}$$

We remark that step (#) is the *definition* of the action of  $(\Delta_M + m^2)$  on the distribution  $\text{PI}_M^\Sigma h$ . By lemma 4.1, lemma 4.3, and (iii) of lemma 4.4, then, this equality can be extended to  $\phi \in W^1(M)$  and  $h \in W^{\frac{1}{2}}(\Sigma)$ . Finally, replacing  $h$  by  $(\text{DN}_M^\Sigma)^{-1} f$ , with  $(\text{DN}_M^\Sigma)^{-1}$  being a  $\Psi$ DO of order  $-1$ , yields the desired relation (4.14) as well as its domain.  $\square$

**Remark 4.4.** One is advised to compare lemma 4.5 with the fact in one dimensions that the distributional derivative of the Heaviside function  $H_a = a \cdot 1_{(0, \infty)}$  ( $a \in \mathbb{R}$ ) is the delta function multiplied by the jump of  $H_a$  across 0, that is,

$$\langle H'_a, \varphi \rangle_{L^2(\mathbb{R})} = [H_a(0+) - H_a(0-)] \cdot \varphi(0) \quad (4.15)$$

for any  $\varphi \in \mathcal{S}(\mathbb{R})$ . In our case the role of the Heaviside function is played by the vector field  $\nabla \text{PI}_M^\Sigma (\text{DN}_M^\Sigma)^{-1} f$ . Indeed, following remark 4.3, the “jump” of  $\nabla \text{PI}_M^\Sigma (\text{DN}_M^\Sigma)^{-1} f$  across  $\Sigma$  is exactly  $f$ , as the directions tangential to  $\Sigma$  does not contribute to the jump with  $(\text{DN}_M^\Sigma)^{-1} f$  being smooth. This comparison in mind, it is also customary to write  $j_\Sigma f$  as  $f \otimes \delta_\Sigma$ , as for example, in Carron [13].

**Corollary 4.6.** *For  $f, h \in C^\infty(\Sigma)$ , we have*

$$\langle f, (\text{DN}_M^\Sigma)^{-1} h \rangle_{L^2(\Sigma)} = \langle j_\Sigma f, (\Delta + m^2)^{-1} j_\Sigma h \rangle_{L^2(M)}. \quad (4.16)$$

*In other words,*  $(\text{DN}_M^\Sigma)^{-1} = \tau(\Delta + m^2)^{-1} j_\Sigma$  ( $= \tau(\Delta + m^2)^{-1} \tau^*$ ).

*Proof.* This is immediate by noting that  $\tau_\Sigma \text{PI}_M^\Sigma$  is the identity on  $L^2(\Sigma)$ .  $\square$

**Remark 4.5.** Indeed, noting that the Schwartz kernel  $K_\tau$  of  $\tau_\Sigma$  is the delta distribution on the diagonal  $\{(x, x)\} \subset \Sigma \times M$ , and that  $j_\Sigma$  is the distributional adjoint of  $\tau_\Sigma$ , corollary 4.6 allows one to deduce immediately the Schwartz kernel of  $(\text{DN}_M^\Sigma)^{-1}$ , denoted  $G_{\text{DN}}^\Sigma$ :

$$\begin{aligned} \langle f, (\text{DN}_M^\Sigma)^{-1} h \rangle_{L^2(\Sigma)} &= \iint_{\Sigma \times M} \iint_{\Sigma \times M} f(x) K_\tau(x, z) G_{(\Delta+m^2)}(z, w) K_\tau(y, w) h(y) dx dz dy dw \\ &= \iint_{\Sigma \times \Sigma} f(x) G_{(\Delta+m^2)}(x, y) h(y) dx dy, \end{aligned}$$

that is,  $G_{\text{DN}}^\Sigma = G_{(\Delta+m^2)}|_{\Sigma \times \Sigma}$ , where  $G_{(\Delta+m^2)}$  is the Helmholtz Green function on  $M$ , which is a well-known result. Of course, assuming this result, one could also work backwards to give lemma 4.5 another proof, using the Poisson integral formula (lemma 4.7 below) for  $\text{PI}_M^\Sigma$ .

Now we move to the second consequence of the Green-Stokes formula. Let  $(\Omega, g)$  be a compact Riemannian manifold with boundary  $\partial\Omega$ . Recall that  $(\Delta_{\Omega, D} + m^2)^{-1}$  denotes the Helmholtz Green operator with *Dirichlet* conditions on  $\partial\Omega$ .

**Lemma 4.7** ([77] page 46). *We have, for  $\varphi \in C^\infty(\partial\Omega)$  and  $f \in C_c^\infty(\Omega^\circ)$ ,*

$$\langle \text{PI}_\Omega^{\partial\Omega} \varphi, f \rangle_{L^2(\Omega)} = \langle \varphi, -\partial_\nu (\Delta_{\Omega, D} + m^2)^{-1} f|_{\partial\Omega} \rangle_{L^2(\partial\Omega)}, \quad (4.17)$$

where  $\nu$ , again, denotes the outward unit normal vector field along  $\partial\Omega$ .  $\square$

**Corollary 4.8.** For  $(M, g)$  a closed Riemannian manifold and  $\Sigma \subset M$  an embedded hypersurface, for  $\varphi \in C^\infty(\Sigma)$  and  $f \in C_c^\infty(M \setminus \Sigma)$ ,

$$\langle \text{PI}_M^\Sigma \varphi, f \rangle_{L^2(M)} = \langle \varphi, -(\partial_\nu u|_{\Sigma_-} + \partial_{-\nu} u|_{\Sigma_+}) \rangle_{L^2(\Sigma)}, \quad (4.18)$$

where  $u = (\Delta_{M \setminus \Sigma, D} + m^2)^{-1} f$ , and the notations  $\Sigma_-$  and  $\Sigma_+$  have the same meanings as in definition 4.6.

*Proof.* See  $M \setminus \Sigma$  as a manifold with two boundaries  $\Sigma \sqcup \Sigma$ , and we note

$$\text{PI}_M^\Sigma \varphi = \text{PI}_{M \setminus \Sigma}^{\Sigma \sqcup \Sigma} \begin{bmatrix} \varphi \\ \varphi \end{bmatrix}, \quad (4.19)$$

as well as

$$\left\langle \begin{bmatrix} \varphi \\ \varphi \end{bmatrix}, - \begin{bmatrix} \partial_\nu u|_{\Sigma_-} \\ \partial_{-\nu} u|_{\Sigma_+} \end{bmatrix} \right\rangle_{L^2(\Sigma \sqcup \Sigma)} = \langle \varphi, -(\partial_\nu u|_{\Sigma_-} + \partial_{-\nu} u|_{\Sigma_+}) \rangle_{L^2(\Sigma)}, \quad (4.20)$$

while applying lemma 4.7.  $\square$

## 4.4 Induced Laws

Let  $(M, g)$  be a closed Riemannian manifold and  $\Sigma \subset M$  an embedded hypersurface. Lemma 4.5 then says that, for each  $f \in C^\infty(\Sigma)$ , the random variables

$$\tau_\Sigma \phi(f) \quad \text{and} \quad \phi(j_\Sigma f), \quad (4.21)$$

while  $\phi \sim \mu_{\text{GFF}}^M$ , are (surely) equal on the Cameron-Martin space  $W^1(M)$ . By proposition B.6 then, they are almost surely equal over  $\mathcal{D}'(M)$ . Subsequently from corollary 4.6 we deduce

$$\mathbb{E}_{\text{GFF}}^M[\tau_\Sigma \phi(f) \tau_\Sigma \phi(h)] = \mathbb{E}_{\text{GFF}}^M[\phi(j_\Sigma f) \phi(j_\Sigma h)] = \langle f, (\text{DN}_M^\Sigma)^{-1} h \rangle_{L^2(\Sigma)}. \quad (4.22)$$

Taking into account lemma 2.4 and (iii) of lemma 4.4, we have proved the following.

**Proposition 4.9.** If  $\phi \in \mathcal{D}'(M)$  follows the law of  $\mu_{\text{GFF}}^M$ , then the random field  $\tau_\Sigma \phi \in \mathcal{D}'(\Sigma)$  can equivalently be realized as the (centered) Gaussian field  $\tilde{\varphi}$  on  $\Sigma$  with covariance

$$\mathbb{E}[\tilde{\varphi}(f) \tilde{\varphi}(h)] = \langle f, (\text{DN}_M^\Sigma)^{-1} h \rangle_{L^2(\Sigma)}, \quad (4.23)$$

for  $f, h \in C^\infty(\Sigma)$ . In other words, the measure image  $\widehat{\tau_\Sigma}(\mu_{\text{GFF}}^M)$  of  $\mu_{\text{GFF}}^M$  under any measurable linear extension  $\widehat{\tau_\Sigma}$  of  $\tau_\Sigma : W^1(M) \rightarrow W^{\frac{1}{2}}(\Sigma)$  coincides with the measure  $\mu_{\text{DN}}^{\Sigma, M}$  on any  $\mathcal{D}'(\Sigma)$  for the field  $\tilde{\varphi}$  satisfying (4.23).  $\square$

Next we study induced random fields in the other direction, by the Poisson integral operator. Namely, for  $\Omega, \partial\Omega$  as in lemma 4.7, given a Gaussian random field  $\varphi$  on  $\partial\Omega$ , what is the law of the field  $\text{PI}_\Omega^{\partial\Omega} \varphi$ ? From another perspective one solves the Helmholtz (Laplace) equation with random boundary conditions. We write in shorthand

$$(\text{PI}_\Omega^{\partial\Omega})^* \stackrel{\text{def}}{=} -\partial_\nu (\Delta_{\Omega, D} + m^2)^{-1} (-) |_{\partial\Omega}. \quad (4.24)$$

Thus lemma 4.7 says

$$\langle \text{PI}_\Omega^{\partial\Omega} \varphi, f \rangle_{L^2(\Omega)} = \langle \varphi, (\text{PI}_\Omega^{\partial\Omega})^* f \rangle_{L^2(\partial\Omega)}, \quad (4.25)$$

for  $\varphi \in C^\infty(\partial\Omega)$ ,  $f \in C_c^\infty(\Omega^\circ)$ . Suppose  $\varphi$  has covariance operator  $C$  of order  $-s$  ( $s > 0$ ). Note  $\text{PI}_\Omega^{\partial\Omega}$  is always well-defined on the Cameron-Martin space  $W^s(\partial\Omega)$ . By the same token as above, for  $f \in C_c^\infty(\Omega^\circ)$ ,

$$(\text{PI}_\Omega^{\partial\Omega} \varphi)(f) \quad \text{and} \quad \varphi((\text{PI}_\Omega^{\partial\Omega})^* f) \quad (4.26)$$

are surely equal on  $W^s(\partial\Omega)$ . Moreover,

$$\mathbb{E}_C^{\partial\Omega} [(\text{PI}_\Omega^{\partial\Omega} \varphi)(f) (\text{PI}_\Omega^{\partial\Omega} \varphi)(h)] = \langle f, \text{PI}_\Omega^{\partial\Omega} C (\text{PI}_\Omega^{\partial\Omega})^* h \rangle_{L^2(\partial\Omega)}. \quad (4.27)$$

We deduce



**Proposition 4.10.** *If  $\varphi \in \mathcal{D}'(\partial\Omega)$  is a (centered) Gaussian random field with covariance operator  $C$ , then the random field  $\text{PI}_\Omega^{\partial\Omega} \varphi \in \mathcal{D}'(\Omega^\circ)$  can equivalently be realized as the (centered) Gaussian field  $\tilde{\phi}$  on  $\Omega$  with covariance*

$$\mathbb{E}[\tilde{\phi}(f)\tilde{\phi}(h)] = \langle f, \text{PI}_\Omega^{\partial\Omega} C (\text{PI}_\Omega^{\partial\Omega})^* h \rangle_{L^2(\partial\Omega)}, \quad (4.28)$$

for  $f, h \in C^\infty(\partial\Omega)$ . In other words, the measure image  $\widehat{\text{PI}}_\Omega^{\partial\Omega}(\mu_C^{\partial\Omega})$  of  $\mu_C^{\partial\Omega}$  under any measurable linear extension  $\widehat{\text{PI}}_\Omega^{\partial\Omega}$  of  $\text{PI}_\Omega^{\partial\Omega} : W^s(\partial\Omega) \rightarrow W^{s+\frac{1}{2}}(\Omega)$  coincides with the measure for the field  $\tilde{\phi}$  satisfying (4.28).  $\square$

**Remark 4.6.** What is strictly needed for showing Segal axioms is not the full proposition 4.10 but rather this innocent observation: by (4.19), if the random field  $\varphi \in \mathcal{D}'(\Sigma)$  follows a fixed probability law, then the induced random fields  $\text{PI}_M^\Sigma \varphi$  and  $\text{PI}_{M \setminus \Sigma}^{\Sigma \cup \Sigma}[\varphi]$  in  $\mathcal{D}'(M \setminus \Sigma)$  follows the same law.

## 4.5 First Comment on Reflection Positivity

As the names would suggest, the positivity of the Dirichlet-to-Neumann map (itself the consequence of the positivity of the Dirichlet energy) gives an interesting inequality comparing the resolvents of Laplacians with Dirichlet and Neumann conditions (corollary 4.12), via the Poisson integral formula (lemma 4.7). Let  $\Omega, \partial\Omega$  be as in lemma 4.7. We adopt the shorthand notations

$$C_D \stackrel{\text{def}}{=} (\Delta_{\Omega,D} + m^2)^{-1}, \quad C_N \stackrel{\text{def}}{=} (\Delta_{\Omega,N} + m^2)^{-1}, \quad (4.29)$$

where, similar to  $C_D, C_N f$  for  $f \in C_c^\infty(\Omega^\circ)$  solves the Neumann boundary value problem

$$\begin{cases} (\Delta + m^2)(C_N f) = f, & \text{in } \Omega, \\ \partial_\nu(C_N f)|_{\partial\Omega} = 0, & \text{on } \partial\Omega. \end{cases} \quad (4.30)$$

There is the following simple, elementary relation:

**Lemma 4.11.** *In the situation as above, we have the operator equality on  $C_c^\infty(\Omega^\circ)$ ,*

$$\text{PI}_\Omega^{\partial\Omega} (\text{DN}_\Omega^{\partial\Omega})^{-1} (\text{PI}_\Omega^{\partial\Omega})^* = C_N - C_D. \quad (4.31)$$

*Proof.* Pick  $f \in C_c^\infty(\Omega^\circ)$ , let  $u := C_D f$  and put  $w := \text{PI}_\Omega^{\partial\Omega} (\text{DN}_\Omega^{\partial\Omega})^{-1} (-\partial_\nu u|_{\partial\Omega})$ . Then, by the definition of  $\text{DN}_\Omega^{\partial\Omega}$ ,  $w$  solves the following boundary value problem:

$$\begin{cases} (\Delta + m^2)w = 0, & \text{in } \Omega, \\ \partial_\nu w|_{\partial\Omega} = -\partial_\nu u|_{\partial\Omega}, & \text{on } \partial\Omega. \end{cases} \quad (4.32)$$

However,

$$\begin{cases} (\Delta + m^2)u = f, & \text{in } \Omega, \\ u|_{\partial\Omega} = 0, & \text{on } \partial\Omega, \end{cases} \quad \text{implying} \quad \begin{cases} (\Delta + m^2)(u + w) = f, & \text{in } \Omega, \\ \partial_\nu(u + w)|_{\partial\Omega} = 0, & \text{on } \partial\Omega, \end{cases} \quad (4.33)$$

namely  $u + w = C_N f$ , that is,  $w = C_N f - C_D f$ . We obtain the result.  $\square$

Now the positivity of  $\text{DN}_\Omega^{\partial\Omega}$  (lemma 4.4) implies  $C_N \geq C_D$ , namely  $\langle f, (C_N - C_D)f \rangle_{L^2(\Omega)} \geq 0$  for all  $f \in C_c^\infty(\Omega^\circ)$ . One step further,

**Corollary 4.12.** *We have  $C_N \geq C_D$  as operators on  $L^2(\Omega)$ .*  $\square$

It is emphasized in Jaffe and Ritter [37] section 3 that  $C_N \geq C_D$  is the crucial relation that leads to the so-called *reflection positivity* (RP) of the GFF (see definition 5.2). At this point let us explain its geometric incarnation. Let now  $\partial\Omega \subset \Omega$  be totally geodesic,  $\Omega^*$  a copy of  $\Omega$  (reversing the coorientation of  $\partial\Omega$ ),  $|\Omega|^2 := \Omega^* \cup_{\partial\Omega} \Omega$ , the isometric double which is a closed Riemannian manifold, and  $\Theta : |\Omega|^2 \rightarrow |\Omega|^2$  an isometric involution fixing  $\partial\Omega$ , such that  $\Theta(\Omega) = \Omega^*$  and  $\Theta(\Omega^*) = \Omega$ .

**Remark 4.7.** Such  $\Omega$  is named in Gibbons [23] (in the 4-dimensional case) as a *real tunnelling geometry* (see [23] section 4). The isometric double  $|\Omega|^2$  and the involution  $\Theta$  exist, the latter being a reflection, *dissecting*  $|\Omega|^2$  into  $\Omega$  and  $\Omega^*$ , its fixed point set being  $\partial\Omega$  (see also Ritter [58] section 2.1.1).

The action of  $\Theta$  extends in the usual way to  $C^\infty(|\Omega|^2)$  and  $\mathcal{D}'(|\Omega|^2)$  by pulling-back. This set-up brings in another resolvent operator which is

$$C \stackrel{\text{def}}{=} (\Delta_{|\Omega|^2} + m^2)^{-1}. \quad (4.34)$$

Denote also by  $\Pi_+ : L^2(|\Omega|^2) \rightarrow L^2(\Omega)$  the orthogonal projection. Below we reprove lemma 3 of Jaffe and Ritter [37].

**Lemma 4.13.** *Let  $|\Omega|^2$ ,  $\Theta$ ,  $\Pi_+$  and  $C$  be as above. Then*

$$\Pi_+ \Theta C = \frac{1}{2}(C_N - C_D) \quad (4.35)$$

on  $C_c^\infty(\Omega^\circ)$  and  $L^2(\Omega)$ .

*Proof.* As in [37], the key idea is the *method of images*, that is, with the help of reflection symmetry under  $\Theta$ , we have

$$C_N f = \Pi_+ C(f + \Theta f), \quad \text{and} \quad C_D f = \Pi_+ C(f - \Theta f), \quad (4.36)$$

for any  $f \in C_c^\infty(\Omega^\circ)$ , since the functions (potentials)  $C(f + \Theta f)$  and  $C(f - \Theta f)$  satisfy, automatically and respectively, the Neumann and Dirichlet conditions along  $\partial\Omega$  (note, first, that  $[\Theta, C] = 0$  since  $\Theta$  is an isometry, and second, for  $x \in \partial\Omega$ ,  $\partial_\nu|_x$  is the eigenvector of eigenvalue  $-1$  of  $d\Theta_x$  on  $T_x|\Omega|^2$ ). We obtain the result immediately.  $\square$

In summary,

**Corollary 4.14** (equivalent formulations of RP). *In the situation as above, we have*

$$2\langle f, \Theta C f \rangle_{L^2(|\Omega|^2)} = \langle f, (C_N - C_D)f \rangle_{L^2(\Omega)} = \langle (\text{PI}_\Omega^{\partial\Omega})^* f, (\text{DN}_\Omega^{\partial\Omega})^{-1} (\text{PI}_\Omega^{\partial\Omega})^* f \rangle_{L^2(\partial\Omega)} \quad (4.37)$$

for all  $f \in C_c^\infty(\Omega^\circ)$ , and all of the above quantities are nonnegative.  $\square$

**Remark 4.8.** Note that the 3 quantities in (4.37) make sense respectively on the spaces  $L^2(|\Omega|^2)$ ,  $L^2(\Omega)$  and  $L^2(\partial\Omega)$ . While the first quantity is the original view of RP, the second quantity offers a *one-sided* view and the third provides a view *within the boundary*  $\partial\Omega$ . Nevertheless, the map  $\text{DN}_\Omega^{\partial\Omega}$  reflects the geometry of the bulk, see for example Paternain, Salo and Uhlmann [50] section 11.5.

## 5 Markov Property and Consequences

Materials in sections 5.1, 5.2 and 5.4 are largely classical with an excellent source being Dimock [16] which we follow roughly. See also Simon [71] section III.3 and see Powell and Werner [54] section 4.2 for a probabilistic point of view.

### 5.1 Decompositions of Sobolev Spaces

Let  $(M, g)$  be a closed Riemannian manifold. Recall from Appendix C.1 the definitions of the spaces  $W_A^s(M)$  and  $W_U^s(M)$  for  $A \subset M$  closed and  $U \subset M$  open. Recall also (lemma C.1) that we have the isometric isomorphism  $(\Delta + m^2) : W^1(M) \xrightarrow{\sim} W^{-1}(M)$  as we endow  $W^1(M)$  and  $W^{-1}(M)$  respectively with the inner products  $\langle -, (\Delta + m^2) - \rangle_{L^2}$  and  $\langle -, (\Delta + m^2)^{-1} - \rangle_{L^2}$ .

**Lemma 5.1** (cf. [16] section 2). *Let  $A \subset M$  be a closed set. Then  $W^1(M)$  and  $W^{-1}(M)$  decompose orthogonally as*

$$\begin{array}{ccc} & \begin{array}{c} \xrightarrow{p_{M \setminus A}^\perp} \\ \xrightarrow{p_{M \setminus A}} \end{array} & \\ & \begin{array}{c} \downarrow \\ \downarrow \end{array} & \\ W^1(M) & = & W_{M \setminus A}^1(M)^\perp \oplus W_{M \setminus A}^1(M) \\ & \begin{array}{c} \downarrow \\ \downarrow \end{array} & \\ (\Delta + m^2) \sim & & (\Delta + m^2) \sim \quad (\Delta + m^2) \sim \\ & \begin{array}{c} \downarrow \\ \downarrow \end{array} & \\ W^{-1}(M) & = & W_A^{-1}(M) \oplus W_A^{-1}(M)^\perp, \\ & \begin{array}{c} \uparrow \\ \uparrow \end{array} & \\ & \begin{array}{c} \xrightarrow{P_A} \\ \xrightarrow{P_A^\perp} \end{array} & \end{array} \quad (5.1)$$

which is preserved by the isometric isomorphism  $\Delta + m^2$ . In particular, we have

$$P_A(\Delta + m^2) = (\Delta + m^2)p_{M \setminus A}^\perp, \quad \text{and} \quad P_A^\perp(\Delta + m^2) = (\Delta + m^2)p_{M \setminus A}, \quad (5.2)$$

where  $p_{M \setminus A}^\perp$ ,  $p_{M \setminus A}$ ,  $P_A$ , and  $P_A^\perp$  are the corresponding orthogonal projections as indicated in the diagram.

*Proof.* We just need to show that the image of  $W_{M \setminus A}^1(M)$  under  $\Delta + m^2$  is precisely  $W_A^{-1}(M)^\perp$ . Indeed, by our definition of the inner products we have

$$\langle (\Delta + m^2)u, v \rangle_{W^{-1}(M)} = \langle u, v \rangle_{L^2(M)} \quad (5.3)$$

for all  $u \in W^1(M)$ ,  $v \in W^{-1}(M)$ , where the RHS denotes the duality (distributional) pairing. However, the annihilator of  $W_A^{-1}(M)$  under the duality pairing is exactly  $W_{M \setminus A}^1(M)$ , see lemma C.3. This translates as

$$\{u \in W^1(M) \mid \langle (\Delta + m^2)u, v \rangle_{W^{-1}(M)} = 0 \text{ for all } v \in W_A^{-1}(M)\} = W_{M \setminus A}^1(M), \quad (5.4)$$

which is what we desired.  $\square$

**Corollary 5.2.** (5.1) and (5.2) holds for  $A = \Sigma \subset M$  an embedded closed hypersurface (codimension one submanifold).  $\square$

Indeed, the crucial relation (5.3) together with (5.2) gives

**Corollary 5.3** (adjoints). *We have*

$$\langle p_{M \setminus A}^\perp u, v \rangle_{L^2} = \langle u, P_A v \rangle_{L^2}, \quad \text{and} \quad \langle p_{M \setminus A} u, v \rangle_{L^2} = \langle u, P_A^\perp v \rangle_{L^2}, \quad (5.5)$$

for  $u \in W^1(M)$  and  $v \in W^{-1}(M)$ .  $\square$

**Remark 5.1.** Denote  $U := M \setminus A$ . Recall from (C.4) that  $W^{-1}(U) := W_A^{-1}(M)^\perp$ . In fact, corollary 5.3 shows  $P_{M \setminus U}^\perp$  coincides with the restriction map  $\rho_{M \setminus U} : \mathcal{D}'(M) \rightarrow \mathcal{D}'(U)$ , since if already  $u \in W_U^1(M)$  then  $p_U u = i_* u$ ,  $i_* : W_U^1(M) \rightarrow W^1(M)$  being inclusion.

We are thus lead naturally to the following and eventually corollary 5.5.

**Corollary 5.4.** *Let  $U \subset M$  be an open set and  $F \subset U$  be a closed set. Then  $W_U^1(M)$  and  $W^{-1}(U)$  decompose further as*

$$\begin{array}{ccc} W_U^1(M) & = & W_{U \setminus F}^1(M)^\perp \oplus W_{U \setminus F}^1(M) \\ \downarrow (\Delta + m^2) \sim & & \downarrow (\Delta + m^2) \sim \quad \downarrow (\Delta + m^2) \sim \\ W^{-1}(U) & = & P_{M \setminus U}^\perp(W_F^{-1}(M)) \oplus W_F^{-1}(M)^\perp, \end{array} \quad (5.6)$$

where the orthogonal complements are taken respectively inside  $W_U^1(M)$  and  $W^{-1}(U)$ , which is preserved by the isometric isomorphism  $\Delta + m^2$ . Also, we have commutation relations similar to (5.2).

*Proof.* The only point needing explanation is

$$(\Delta + m^2)(W_{U \setminus F}^1(M)^\perp \cap W_U^1(M)) = P_{M \setminus U}^\perp(W_F^{-1}(M)). \quad (5.7)$$

Indeed, since  $\Delta + m^2$  is bijective, by (5.1) we have

$$\begin{aligned} \text{LHS} &= W_{F \cup (M \setminus U)}^{-1}(M) \cap W_{M \setminus U}^{-1}(M)^\perp \\ &= (W_{M \setminus U}^{-1}(M) \oplus W_F^{-1}(M)) \cap W_{M \setminus U}^{-1}(M)^\perp \quad (F \text{ and } M \setminus U \text{ disjoint}) \\ &= \text{RHS}, \end{aligned}$$

where the direct sum in the second line is non-orthogonal.  $\square$

Now let  $(\Omega, g)$  be compact with (smooth) boundary, smoothly and isometrically embedded in closed  $M$  (if  $\Omega$  is a real tunnelling geometry, then one choice for  $M$  is the ‘‘isometric double’’, see remark 4.7) and  $\Sigma$  be embedded in  $\Omega^\circ$ . Then in particular

**Corollary 5.5.** *Corollary 5.4 holds for  $U = \Omega^\circ$  and  $\Sigma \subset \Omega^\circ$  an embedded closed hypersurface.*  $\square$

## 5.2 The Markov Stochastic Decomposition of GFF

Again suppose  $(M, g)$  is a closed Riemannian manifold and  $\Sigma \subset M$  an embedded closed hypersurface with induced metric. A probabilistic point of view of the results in this section is provided in Powell and Werner [54] section 4.1.

**Lemma 5.6.** *We have*

$$p_{M \setminus \Sigma}^\perp = \text{PI}_M^\Sigma \circ \tau_\Sigma \quad (5.8)$$

on  $W^1(M)$ , where  $p_{M \setminus \Sigma}^\perp : W^1(M) \rightarrow W_{M \setminus A}^1(M)^\perp$  is the orthogonal projection as in lemma 5.1.

*Proof.* We show that for  $f \in C^\infty(M)$ ,  $p_{M \setminus \Sigma}^\perp f$  solves the boundary value problem

$$\begin{cases} (\Delta + m^2)u = 0, & \text{in } M \setminus \Sigma, \\ u|_\Sigma = \tau_\Sigma f, & \text{on } \Sigma. \end{cases} \quad (5.9)$$

Indeed, the first condition holds since  $\text{supp}((\Delta + m^2)p_{M \setminus \Sigma}^\perp f) \subset \Sigma$  by lemma 5.1. To show the second, notice  $\tau_\Sigma = 0$  on  $W_{M \setminus \Sigma}^1(M)$  by its definition (C.3). Hence

$$\tau_\Sigma p_{M \setminus \Sigma}^\perp = \tau_\Sigma(\mathbb{1} - p_{M \setminus \Sigma}) = \tau_\Sigma, \quad (5.10)$$

on  $W^1(M)$ . We conclude the proof.  $\square$

**Proposition 5.7.** *Suppose  $\phi \in \mathcal{D}'(M)$  follows  $\mu_{\text{GFF}}^M$ . Then there is a stochastic decomposition*

$$\phi = \widehat{p_{M \setminus \Sigma}^\perp} \phi + \widehat{p_{M \setminus \Sigma}} \phi \stackrel{\text{def}}{=} \phi_\Sigma + \phi_{M \setminus \Sigma}^D, \quad (5.11)$$

into independent random fields  $\phi_\Sigma$  and  $\phi_{M \setminus \Sigma}^D$ . More precisely, for  $f \in C^\infty(M)$  we define the random variables

$$\phi_\Sigma(f) \stackrel{\text{def}}{=} \phi(P_\Sigma f), \quad \text{and} \quad \phi_{M \setminus \Sigma}^D(f) \stackrel{\text{def}}{=} \phi(P_\Sigma^\perp f). \quad (5.12)$$

Then,  $\phi_{M \setminus \Sigma}^D$  follows the law of  $\mu_{\text{GFF}}^{M \setminus \Sigma, D}$ , while  $\phi_\Sigma$  solves the boundary value problem

$$\begin{cases} (\Delta + m^2)\phi_\Sigma = 0, & \text{in } M \setminus \Sigma, \\ \phi_\Sigma|_\Sigma = \tau_\Sigma \phi, & \text{on } \Sigma, \end{cases} \quad (5.13)$$

almost surely.

**Remark 5.2.** The term ‘‘stochastic decomposition’’ is borrowed from Bogachev [12] remark 3.7.7.

*Proof.* Firstly,  $\phi_\Sigma$  and  $\phi_{M \setminus \Sigma}^D$  are independent since by (5.12) their Gaussian Hilbert spaces are respectively  $W_\Sigma^{-1}(M)$  and  $W_\Sigma^{-1}(M)^\perp = W^{-1}(M \setminus \Sigma)$ . The latter also means  $\phi_{M \setminus \Sigma}^D$  follows  $\mu_{\text{GFF}}^{M \setminus \Sigma, D}$ . The last fact about  $\phi_\Sigma$  follows from lemma 5.6 and the uniqueness of the measurable linear extension (proposition B.7).  $\square$

In the same vein using corollary 5.5, we have also a decomposition for the Dirichlet GFF on a domain  $\Omega$  with smooth boundary, with respect to a hypersurface  $\Sigma$  (isometrically) embedded in the interior  $\Omega^\circ$ .

**Proposition 5.8.** *Suppose  $\phi_\Omega^D \in \mathcal{D}'(\Omega^\circ)$  follows  $\mu_{\text{GFF}}^{\Omega, D}$ . Then there is a stochastic decomposition*

$$\phi = \widehat{p_{\Omega^\circ \setminus \Sigma}^\perp} \phi + \widehat{p_{\Omega^\circ \setminus \Sigma}} \phi \stackrel{\text{def}}{=} (\phi_\Omega^D)_\Sigma + \phi_{\Omega \setminus \Sigma}^D, \quad (5.14)$$

into independent random fields  $(\phi_\Omega^D)_\Sigma$  and  $\phi_{\Omega \setminus \Sigma}^D$ . More precisely, for  $f \in W^{-1}(\Omega^\circ)$  we define the random variables

$$(\phi_\Omega^D)_\Sigma(f) \stackrel{\text{def}}{=} \phi_\Omega^D(P_\Sigma f), \quad \text{and} \quad \phi_{\Omega \setminus \Sigma}^D(f) \stackrel{\text{def}}{=} \phi_\Omega^D(P_\Sigma^\perp f). \quad (5.15)$$

Then,  $\phi_{\Omega \setminus \Sigma}^D$  follows the law of  $\mu_{\text{GFF}}^{\Omega \setminus \Sigma, D}$ , while  $(\phi_\Omega^D)_\Sigma$  solves the boundary value problem

$$\begin{cases} (\Delta + m^2)(\phi_\Omega^D)_\Sigma = 0, & \text{in } \Omega^\circ \setminus \Sigma, \\ (\phi_\Omega^D)_\Sigma|_\Sigma = \tau_\Sigma \phi_\Omega^D, & \text{on } \Sigma, \\ (\phi_\Omega^D)_\Sigma|_{\partial\Omega} = 0, & \text{on } \partial\Omega, \end{cases} \quad (5.16)$$

almost surely.  $\square$

The following version of proposition 5.7 in the case of a dissecting (smooth isometrically embedded) hypersurface  $\Sigma \subset M$  such that  $M \setminus \Sigma = M_+^\circ \sqcup M_-^\circ$ , will be useful in definition 5.4.

**Lemma 5.9.** *We have  $W_{M_-}^{-1}(M)^\perp \subset W_{M_+}^{-1}(M)$  and thus  $W_{M_+}^{-1}(M) = W_{M_-}^{-1}(M)^\perp \oplus W_{\Sigma}^{-1}(M)$ . Similarly  $W_{M_-}^{-1}(M) = W_{M_+}^{-1}(M)^\perp \oplus W_{\Sigma}^{-1}(M)$ .*

*Proof.* By lemma 5.1 with  $A = M_-$  we see that  $W_{M_-}^{-1}(M)^\perp = (\Delta + m^2)(W_{M_+}^1(M))$ . These distributions are supported in  $M_+$  since  $\Delta + m^2$  is local.  $\square$

**Corollary 5.10.** *Suppose  $\phi \in \mathcal{D}'(M)$  follows  $\mu_{\text{GFF}}^M$ . Then there is a stochastic decomposition*

$$\phi = \widehat{p_{M \setminus \Sigma}^\perp} \phi + \widehat{p_{M_+^\circ}^\perp} \phi + \widehat{p_{M_-^\circ}^\perp} \phi \stackrel{\text{def}}{=} \phi_\Sigma + \phi_{M_+^\circ}^D + \phi_{M_-^\circ}^D, \quad (5.17)$$

into independent random fields. More precisely, for  $f \in C^\infty(M)$  the random variables are defined by

$$\phi_\Sigma(f) \stackrel{\text{def}}{=} \phi(P_\Sigma f), \quad \phi_{M_+^\circ}^D(f) \stackrel{\text{def}}{=} \phi(P_{M_-}^\perp f), \quad \text{and} \quad \phi_{M_-^\circ}^D(f) \stackrel{\text{def}}{=} \phi(P_{M_+}^\perp f), \quad (5.18)$$

and with  $\phi_\Sigma$  the same as in proposition 5.7,  $\phi_{M_+^\circ}^D$  and  $\phi_{M_-^\circ}^D$  follows respectively  $\mu_{\text{GFF}}^{M_+^\circ, D}$  and  $\mu_{\text{GFF}}^{M_-^\circ, D}$ . Moreover,  $(\phi_\Sigma + \phi_{M_+^\circ}^D)(f) = \phi(P_{M_+} f)$ ,  $(\phi_\Sigma + \phi_{M_-^\circ}^D)(f) = \phi(P_{M_-} f)$ .  $\square$

### 5.3 The Bayes Principle Applied to GFF

Let now  $M$  be a closed Riemannian manifold and  $\Sigma_1, \Sigma_2 \subset M$  non-intersecting isometrically embedded smooth closed hypersurfaces. The goal of this section is to derive the Bayes principle relating the probability laws of the two random fields  $\tau_{\Sigma_1} \phi$  and  $\tau_{\Sigma_2} \phi$  where  $\phi$  is the GFF on  $M$ . To avoid convolving with nomenclatures of conditional probabilities we prefer a direct measure theoretic argument, although these are clearly equivalent. Throughout this section we identify continuous linear maps of Cameron-Martin spaces with their measurable extensions to distributional  $Q$ -spaces, as well as their induced actions on measures.

To begin with, by proposition 5.7 we have a stochastic decomposition

$$\phi = \phi_{\Sigma_1} + \phi_{M \setminus \Sigma_1}^D \quad (5.19)$$

for  $\phi \sim \mu_{\text{GFF}}^M$ , the two components independent of each other. Then, apply  $\tau_{\Sigma_2}$  we get

$$\tau_{\Sigma_2} \phi = \tau_{\Sigma_2} \phi_{\Sigma_1} + \tau_{\Sigma_2} \phi_{M \setminus \Sigma_1}^D = \tau_{\Sigma_2} \text{PI}_M^{\Sigma_1} \tau_{\Sigma_1} \phi + \tau_{\Sigma_2} \phi_{M \setminus \Sigma_1}^D \quad (5.20)$$

$$\stackrel{\text{def}}{=} \mathcal{M}_{M,2}^1 \tau_{\Sigma_1} \phi + \tau_{\Sigma_2} \phi_{M \setminus \Sigma_1}^D. \quad (5.21)$$

Here we define

$$\mathcal{M}_{M,2}^1 \stackrel{\text{def}}{=} \tau_{\Sigma_2} \text{PI}_M^{\Sigma_1} : \begin{cases} C^\infty(\Sigma_1) & \longrightarrow C^\infty(\Sigma_2), \\ W^{\frac{1}{2}}(\Sigma_1) & \longrightarrow W^{\frac{1}{2}}(\Sigma_2), \end{cases} \quad (5.22)$$

called the transition operator/propagator. Define accordingly

$$\mathcal{G}_{M,2}^1 : \begin{cases} W^{\frac{1}{2}}(\Sigma_1) \times W^{\frac{1}{2}}(\Sigma_2) & \longrightarrow W^{\frac{1}{2}}(\Sigma_1) \times W^{\frac{1}{2}}(\Sigma_2), \\ (\varphi_1, h) & \longmapsto (\varphi_1, h + \mathcal{M}_{M,2}^1 \varphi_1). \end{cases} \quad (5.23)$$

called the graph operator.

**Lemma 5.11.** *We have*

$$\tau_{\Sigma_1 \sqcup \Sigma_2}(\mu_{\text{GFF}}^M) = (\mathcal{G}_{M,2}^1)_*(\mu_{\text{DN}}^{\Sigma_1, M} \otimes \mu_{\text{DN}, D}^{\Sigma_2, M \setminus \Sigma_1}), \quad (5.24)$$

where  $\mu_{\text{DN}}^{\Sigma_1, M}$  and  $\mu_{\text{DN}, D}^{\Sigma_2, M \setminus \Sigma_1}$  are Gaussian measures on  $\mathcal{D}'(\Sigma_1)$  and  $\mathcal{D}'(\Sigma_2)$  with covariances  $(\text{DN}_M^{\Sigma_1})^{-1}$  and  $(\text{DN}_{M \setminus \Sigma_1}^{\Sigma_2, D})^{-1}$ , respectively.

*Proof.* Following (5.21),  $\tau_{\Sigma_2} \phi_{M \setminus \Sigma_1}^D = \tau_{\Sigma_2} \phi - \mathcal{M}_{M,2}^1 \tau_{\Sigma_1} \phi$  is independent of  $\tau_{\Sigma_1} \phi$ , and follows the law  $\mu_{\text{DN}, D}^{\Sigma_2, M \setminus \Sigma_1}$ , while  $\tau_{\Sigma_1} \phi$  follows the law  $\mu_{\text{DN}}^{\Sigma_1, M}$ , both by proposition 4.9.  $\square$

We could also define the operators  $\mathcal{M}_{M,1}^2$  and  $\mathcal{G}_{M,1}^2$  with the roles of  $\Sigma_1$  and  $\Sigma_2$  switched. Note that lemma 5.11 is entirely symmetric under the switching of  $\Sigma_1$  and  $\Sigma_2$ . Recall the measures  $\mu_{\text{2D}}^\Sigma$  from corollary 2.15.

**Proposition 5.12** (Bayes Principle for GFF). *Let now  $M$  be a closed Riemannian manifold and  $\Sigma_1, \Sigma_2 \subset M$  non-intersecting isometrically embedded smooth closed hypersurfaces. We have equality of Radon-Nikodym densities*

$$\frac{d\tau_{\Sigma_1 \sqcup \Sigma_2}(\mu_{\text{GFF}}^M)}{d\mu_{2\mathbf{D}}^{\Sigma_1 \sqcup \Sigma_2}}(\varphi_1, \varphi_2) = \frac{d\mu_{\mathbf{DN}}^{\Sigma_1, M}}{d\mu_{2\mathbf{D}}^{\Sigma_1}}(\varphi_1) \frac{d((\mathcal{M}_{M,2}^1 \varphi_1)_* \mu_{\mathbf{DN}, D}^{\Sigma_2, M \setminus \Sigma_1})}{d\mu_{2\mathbf{D}}^{\Sigma_2}}(\varphi_2) \quad (5.25)$$

$$= \frac{d\mu_{\mathbf{DN}}^{\Sigma_2, M}}{d\mu_{2\mathbf{D}}^{\Sigma_2}}(\varphi_2) \frac{d((\mathcal{M}_{M,1}^2 \varphi_2)_* \mu_{\mathbf{DN}, D}^{\Sigma_1, M \setminus \Sigma_2})}{d\mu_{2\mathbf{D}}^{\Sigma_1}}(\varphi_1). \quad (5.26)$$

Here  $(\varphi)_*$  denotes the shift induced by  $\varphi$  on measures as in corollary 2.5.

*Proof.* We just need to prove (5.25). In other words,  $(\mathcal{M}_{M,2}^1 \varphi_1)_* \mu_{\mathbf{DN}, D}^{\Sigma_2, M \setminus \Sigma_1}$  is the conditional law of  $\tau_{\Sigma_2} \phi$  provided “ $\tau_{\Sigma_1} = \varphi_1$ ”. The proof is straightforward. For positive (Borel) measurable functionals  $F \in L^+(\mathcal{D}'(\Sigma_1))$ ,  $G \in L^+(\mathcal{D}'(\Sigma_2))$ , we have by lemma 5.11, the change of variables formula, and Fubini's theorem,

$$\begin{aligned} \int (F \otimes G)(\varphi_1, \varphi_2) d\tau_{\Sigma_1 \sqcup \Sigma_2}(\mu_{\text{GFF}}^M) &= \iint (F \otimes G) \circ \mathcal{G}_{M,2}^1(\varphi_1, \varphi_2) d\mu_{\mathbf{DN}}^{\Sigma_1, M} \otimes d\mu_{\mathbf{DN}, D}^{\Sigma_2, M \setminus \Sigma_1} \\ &= \int F(\varphi_1) d\mu_{\mathbf{DN}}^{\Sigma_1, M}(\varphi_1) \int G(\varphi_2 + \mathcal{M}_{M,2}^1 \varphi_1) d\mu_{\mathbf{DN}, D}^{\Sigma_2, M \setminus \Sigma_1}(\varphi_2) \\ &= \int F(\varphi_1) d\mu_{\mathbf{DN}}^{\Sigma_1, M} \int G(\varphi_2) d(\mathcal{M}_{M,2}^1 \varphi_1)_* \mu_{\mathbf{DN}, D}^{\Sigma_2, M \setminus \Sigma_1}. \end{aligned}$$

as  $F$  and  $G$  range over all positive measurable functionals, we obtain the result.  $\square$

**Remark 5.3.** Applying corollary 2.15 and the Cameron-Martin formula (2.12), one obtains a relation between quadratic forms (the corresponding relation for determinants also works out by BFK),

$$\begin{aligned} &\left\langle \begin{bmatrix} \varphi_1 \\ x \end{bmatrix}, \text{DN}_M^{\Sigma_1 \sqcup \Sigma_2} \begin{bmatrix} \varphi_1 \\ x \end{bmatrix} \right\rangle_{L^2(\Sigma_1 \sqcup \Sigma_2)} \\ &= \left\langle \varphi_1, \text{DN}_M^{\Sigma_1} \varphi_1 \right\rangle_{L^2(\Sigma_1)} + \left\langle x, \text{DN}_{M \setminus \Sigma_1}^{\Sigma_2, D} x \right\rangle_{L^2(\Sigma_2)} \\ &\quad - 2 \left\langle x, \text{DN}_{M \setminus \Sigma_1}^{\Sigma_2, D} \mathcal{M}_{M,2}^1 \varphi_1 \right\rangle_{L^2(\Sigma_2)} + \left\langle \mathcal{M}_{M,2}^1 \varphi_1, \text{DN}_{M \setminus \Sigma_1}^{\Sigma_2, D} \mathcal{M}_{M,2}^1 \varphi_1 \right\rangle_{L^2(\Sigma_2)} \end{aligned}$$

which, of course, can also be obtained directly noting

$$\begin{aligned} \left\langle \varphi_1, \text{DN}_M^{\Sigma_1} \varphi_1 \right\rangle_{L^2(\Sigma_1)} &= \left\langle \begin{bmatrix} \varphi_1 \\ \mathcal{M}_{M,2}^1 \varphi_1 \end{bmatrix}, \text{DN}_M^{\Sigma_1 \sqcup \Sigma_2} \begin{bmatrix} \varphi_1 \\ \mathcal{M}_{M,2}^1 \varphi_1 \end{bmatrix} \right\rangle_{L^2(\Sigma_1 \sqcup \Sigma_2)}, \\ \left\langle x, \text{DN}_{M \setminus \Sigma_1}^{\Sigma_2, D} x \right\rangle_{L^2(\Sigma_2)} &= \left\langle \begin{bmatrix} 0 \\ x \end{bmatrix}, \text{DN}_M^{\Sigma_1 \sqcup \Sigma_2} \begin{bmatrix} 0 \\ x \end{bmatrix} \right\rangle_{L^2(\Sigma_1 \sqcup \Sigma_2)}, \end{aligned}$$

et cetera, and expanding the left hand side by writing  $\begin{bmatrix} \varphi_1 \\ x \end{bmatrix} = \begin{bmatrix} \varphi_1 \\ \mathcal{M}_{M,2}^1 \varphi_1 \end{bmatrix} + \begin{bmatrix} x - \mathcal{M}_{M,2}^1 \varphi_1 \\ 0 \end{bmatrix}$ .

**Remark 5.4.** In the case  $M = \Sigma \times \mathbb{R}$  with  $\Sigma_1 := \Sigma \times \{0\}$ ,  $\Sigma_2 := \Sigma \times \{t\}$ , and Dirichlet condition at  $\Sigma \times \{\pm\infty\}$  (namely decay at  $\infty$ ), we have the explicit expression

$$\mathcal{M}_{M,2}^1 = e^{-t\mathbf{D}_\Sigma}, \quad (5.27)$$

where  $\mathbf{D}_\Sigma = (\Delta_\Sigma + m^2)^{1/2}$ . Also, in this case  $\mathbf{D}_\Sigma = \text{DN}_{M_+}^\Sigma$ ,  $M_+ = \Sigma \times [0, +\infty)$ .

## 5.4 Nelson's Markov Property and Reflection Positivity

Let  $M$  be a closed Riemannian manifold. We consider a family of  $\sigma$ -algebras on  $\mathcal{D}'(M)$  generated by the GFF random variables. Recall that for all  $f \in W^{-1}(M)$ , we have defined  $\phi \mapsto \langle \phi, f \rangle_{L^2}$  as a random variable. Now for  $A \subset M$  closed, we define the  $\sigma$ -algebra

$$\mathcal{O}_A \stackrel{\text{def}}{=} \sigma(\phi(f) \mid f \in W_A^{-1}(M)). \quad (5.28)$$

Note since  $\mathcal{D}'(M)$  is a separable Fréchet space, its Borel  $\sigma$ -algebra is generated by  $\phi(f)$  for  $f \in C^\infty(M)$ . The GFF  $\sigma$ -algebra  $\mathcal{O} := \mathcal{O}_M$  is thus finer than the Borel  $\sigma$ -algebra.



**Remark 5.5.** What we discuss here is only a preliminary consideration based on the GFF. The theory of Markov random fields is a well-developed subject and we refer to the monograph of Rozanov [61] for thorough treatment. See also the interesting result of Gu and Mourrat [29] theorem 2.1, as well as the comments thereafter.

**Definition 5.1** ([71] theorem III.9). A probability measure  $\mu$  on  $(\mathcal{D}'(M), \mathcal{O})$  has the domain Markov property if for all closed sets  $A, B \subset M$  such that  $A^\circ \cap B = \emptyset$ , and all  $F \in L^2(\mathcal{D}'(M), \mathcal{O}, \mu)$ , we have

$$\mathbb{E}_\mu[\mathbb{E}_\mu[F|\mathcal{O}_B] | \mathcal{O}_A] = \mathbb{E}_\mu[\mathbb{E}_\mu[F|\mathcal{O}_B] | \mathcal{O}_{\partial A}]. \quad (5.29)$$

In other words, for all  $\mathcal{O}_B$ -measurable  $L^2$ -functional  $F$  we have  $\mathbb{E}_\mu[F|\mathcal{O}_A] = \mathbb{E}_\mu[F|\mathcal{O}_{\partial A}]$ .

**Proposition 5.13** ([16] theorem 1). *The measure  $\mu_{\text{GFF}}^M$  has the domain Markov property.*

*Proof.* For details see Simon [71] section III.3 and Dimock [16]. The crucial point being that by locality of the  $W^1$ -inner product we have  $C_c^\infty(A^\circ) \subset W_{M \setminus A}^1(M)^\perp$ . Therefore by corollary 5.3 if  $A^\circ \cap B = \emptyset$  and  $v \in W_B^{-1}(M)$ ,  $u \in C_c^\infty(A^\circ)$  then

$$\langle P_A v, u \rangle_{L^2} = \langle v, p_{M \setminus A}^\perp u \rangle_{L^2} = \langle v, u \rangle_{L^2} = 0. \quad (5.30)$$

This shows  $P_A P_B = P_{\partial A} P_B$ . The result then follows by applying the Itô-Wiener-Segal isomorphism.  $\square$

Suppose we have a dissecting (smooth isometrically embedded) hypersurface  $\Sigma \subset M$  such that  $M \setminus \Sigma = M_+^\circ \sqcup M_-^\circ$ .

**Lemma 5.14.** *Let the probability measure  $\mu$  on  $(\mathcal{D}'(M), \mathcal{O})$  have the domain Markov property. If real functionals  $F \in L^2(\mathcal{D}'(M), \mathcal{O}_{M_+}, \mu)$ ,  $G \in L^2(\mathcal{D}'(M), \mathcal{O}_{M_-}, \mu)$  are such that  $\mathbb{E}_\mu[F|\mathcal{O}_\Sigma] = \mathbb{E}_\mu[G|\mathcal{O}_\Sigma]$  almost surely, then*

$$\mathbb{E}_\mu[FG] \geq 0. \quad (5.31)$$

*Proof.* The closed sets  $M_+$  and  $M_-$  are such that  $M_+^\circ \cap M_- = \emptyset$ . Also  $\partial M_+ = \partial M_- = \Sigma$ . We have

$$\begin{aligned} \mathbb{E}_\mu[FG] &= \mathbb{E}_\mu[ F \cdot \mathbb{E}_\mu[G|\mathcal{O}_{M_+}] ] && (F \text{ is } \mathcal{O}_{M_+}\text{-measurable}) \\ &= \mathbb{E}_\mu[ F \cdot \mathbb{E}_\mu[G|\mathcal{O}_\Sigma] ] && (\text{domain Markov and } G \text{ is } \mathcal{O}_{M_-}\text{-measurable}) \\ &= \mathbb{E}_\mu[ \mathbb{E}_\mu[F|\mathcal{O}_\Sigma] \mathbb{E}_\mu[G|\mathcal{O}_\Sigma] ] \geq 0, && (\mathbb{E}_\mu[G|\mathcal{O}_\Sigma] \text{ is } \mathcal{O}_\Sigma\text{-measurable}) \end{aligned}$$

by our assumption.  $\square$

**Remark 5.6.** Because of proposition 5.7, for GFF the conditional expectation represents “integrating out redundant degrees of freedom”. Indeed, following proposition 5.7 we write  $F(\phi)$  as  $F(\phi_\Sigma, \phi_{M \setminus \Sigma}^D)$  then we have

$$\mathbb{E}_{\text{GFF}}^M[F|\mathcal{O}_\Sigma](\phi_\Sigma) = \int F(\phi_\Sigma, \phi_{M \setminus \Sigma}^D) d\mu_{\text{GFF}}^{M \setminus \Sigma, D}(\phi_{M \setminus \Sigma}^D). \quad (5.32)$$

See Simon [71] page 225 (in particular equation (VII.26)) for details.

**Lemma 5.15.** *Let  $j_\Sigma$  be the map in lemma 4.5 and let  $W_{\text{DN}}^{-\frac{1}{2}}(\Sigma)$  denote the Gaussian Hilbert space of  $\mu_{\text{DN}}^{\Sigma, M}$ , as in lemma 2.4. Then*

$$j_\Sigma : W_{\text{DN}}^{-\frac{1}{2}}(\Sigma) \xrightarrow{\sim} W_\Sigma^{-1}(M) \quad (5.33)$$

*is an isometric isomorphism. In particular,  $W_\Sigma^{-1}(M) \neq \emptyset$ .*

*Proof.* By lemma 5.1, lemma 5.6 and lemma 4.6,  $j_\Sigma$  is a surjective isometry.  $\square$

Suppose we are in the situation of remark 4.7. Here we take  $M = |\Omega|^2$ . The isometry  $\Theta$  acts on  $\mathcal{O}$ -measurable functionals  $F : \mathcal{D}'(|\Omega|^2) \rightarrow \mathbb{R}$  by  $\Theta^* F(\phi) := F(\Theta\phi)$  and  $\Theta\phi(f) := \phi(\Theta^* f)$ . The action of  $\Theta^*$  on  $C^\infty(|\Omega|^2)$  extends unitarily to  $W^{-1}(|\Omega|^2)$ , since  $\Theta$  is isometry on  $|\Omega|^2$  (commutes with Laplacian).

**Definition 5.2.** A probability measure  $\mu$  on  $(\mathcal{D}'(|\Omega|^2), \mathcal{O})$  is called **reflection positive** if for all real functionals  $F \in L^2(\mathcal{D}'(|\Omega|^2), \mathcal{O}_\Omega, \mu)$  we have

$$\mathbb{E}_\mu[F(\Theta F)] \geq 0. \quad (5.34)$$

**Proposition 5.16** ([16] theorem 2). *The measure  $\mu_{\text{GFF}}^M$  is reflection positive.*

*Proof.* We first claim  $\mathbb{E}_{\text{GFF}}^M[\Theta F | \mathcal{O}_{\partial\Omega}] = \mathbb{E}_{\text{GFF}}^M[F | \mathcal{O}_{\partial\Omega}]$  almost surely. For GFF this is seen via the Itô-Wiener-Segal isomorphism. Indeed the action of  $\Theta^*$  descends to  $\Theta^*F(\phi) = \phi(\Theta^*f_1) \cdots \phi(\Theta^*f_k)$  if  $F(\phi) = \phi(f_1) \cdots \phi(f_k)$ . Thus  $\Theta^* = \Gamma(\Theta^*|_{W^{-1}})$ , implying also that

$$\Theta^* : L^2(\mathcal{D}'(M), \mathcal{O}_\Omega, \mu_{\text{GFF}}^M) \longrightarrow L^2(\mathcal{D}'(M), \mathcal{O}_{\Omega^*}, \mu_{\text{GFF}}^M). \quad (5.35)$$

Since  $\Theta$  fixes  $\partial\Omega$  we have that  $\Theta^*$  is the identity on  $W_{\partial\Omega}^{-1}$ . Since  $\Theta^*$  is also  $W^{-1}$ -unitary we eventually have  $P_{\partial\Omega}\Theta^* = \Theta^*P_{\Theta\partial\Omega} = \Theta^*P_{\partial\Omega} = P_{\partial\Omega}$ , and the claim is true. The proposition then follows from (5.35), proposition 5.13 and lemma 5.14.  $\square$

Following proposition 5.16 we obtain a nonnegative bilinear form on  $L^2(\mathcal{D}'(M), \mathcal{O}_\Omega, \mu_{\text{GFF}}^M)$ :

$$(F, G) \longmapsto \mathbb{E}_{\text{GFF}}^M[F(\Theta G)]. \quad (5.36)$$

We define  $\mathcal{N} := \{F \mid \mathbb{E}_{\text{GFF}}^M[F(\Theta F)] = 0\}$ , namely the kernel of (5.36). We use shorthand notations

$$L^2(\mathcal{D}'(M), \mathcal{O}_\Omega, \mu_{\text{GFF}}^M) \stackrel{\text{def}}{=} \mathcal{E}_+, \quad L^2(\mathcal{D}'(M), \mathcal{O}_{\partial\Omega}, \mu_{\text{GFF}}^M) \stackrel{\text{def}}{=} \mathcal{E}_{\partial\Omega}.$$

**Proposition 5.17.**  $\overline{\mathcal{E}_+/\mathcal{N}} \cong \Gamma(j_{\partial\Omega})L^2(\mathcal{D}'(\partial\Omega), \mu_{\text{DN}}^{\partial\Omega, M})$ , where the closure is taken under (5.36).

*Proof.* Indeed, by the proofs of lemma 5.14 and proposition 5.16 we see that

$$\mathbb{E}_{\text{GFF}}^M[F(\Theta G)] = \mathbb{E}_{\text{GFF}}^M[\mathbb{E}_{\text{GFF}}^M[F | \mathcal{O}_{\partial\Omega}]\mathbb{E}_{\text{GFF}}^M[G | \mathcal{O}_{\partial\Omega}]] = \mathbb{E}_{\text{GFF}}^M[\Gamma(P_{\partial\Omega})F \cdot \Gamma(P_{\partial\Omega})G] \quad (5.37)$$

for  $F, G \in \mathcal{E}_+$ . Note  $\mathcal{E}_{\partial\Omega} \subset \mathcal{E}_+$  and thus  $\mathcal{N} = \ker \Gamma(P_{\partial\Omega})$  showing  $\overline{\mathcal{E}_+/\mathcal{N}} \cong \overline{\text{ran } \Gamma(P_{\partial\Omega})} = \mathcal{E}_{\partial\Omega}$ . But we know  $\mathcal{E}_{\partial\Omega} = \Gamma(W_{\partial\Omega}^{-1}(M))$  therefore we obtain our result by lemma 5.15 and Itô-Wiener-Segal.  $\square$

**Remark 5.7.** Functionals in  $\mathcal{E}_{\partial\Omega}$  are usually called sharp-time functionals (see Jaffe and Ritter [38]) and  $\mathcal{E}_{\partial\Omega}$  is usually taken as the quantum mechanical Hilbert space over the “time-zero slice”  $\partial\Omega$ . This Hilbert space is isomorphic to the one considered in section 6.2, conjugated by the square root of the Radon-Nikodym density between  $\mu_{\text{DN}}^{\partial\Omega, M}$  and  $\mu_{2\text{D}}^{\partial\Omega}$ .

**Remark 5.8.** It is reasonable that a measure defined on  $(\mathcal{D}'(|\Omega|^2), \mathcal{O})$  satisfying the domain Markov property and that is  $\Theta$ -invariant will be reflection positive. Here we define  $\Theta$ -invariant as such that  $\Theta^*\mathbb{E}_\mu[-|\mathcal{O}_{\Theta A}] = \mathbb{E}_\mu[-|\mathcal{O}_A]$  for all closed sets  $A \subset |\Omega|^2$ . Indeed, since  $\Theta$  maps level sets of  $\langle -, f \rangle_{L^2}$  to level sets of  $\langle -, \Theta^*f \rangle_{L^2}$  (by  $L^2$ -unitarity of  $\Theta$ ),  $\Theta\mathcal{O}_A = \mathcal{O}_{\Theta A}$  via its action on indicator functionals. In this sense one shows that  $\mathcal{O}_{\partial\Omega}$  is invariant under  $\Theta$  and further that  $\mathcal{O}_{\partial\Omega}$ -measurable functionals are also invariant, via approximating by simple functionals. This together with lemma 5.14 gives reflection positivity.

**Remark 5.9.** Comparing proposition 4.10 and corollary 4.14 we recover the covariance structure of the induced field  $\text{PI}_\Omega^{\partial\Omega} \varphi$  for  $\varphi \sim \mu_{2\text{DN}}^{\partial\Omega, \Omega}$  with covariance  $\frac{1}{2}(\text{DN}_\Omega^{\partial\Omega})^{-1}$  to be

$$\mathbb{E}[\text{PI}_\Omega^{\partial\Omega} \varphi(f) \text{PI}_\Omega^{\partial\Omega} \varphi(h)] = \langle P_{\partial\Omega} f, P_{\partial\Omega} h \rangle_{W^{-1}(|\Omega|^2)} \quad (5.38)$$

for  $f, h \in C_c^\infty(\Omega^\circ)$ . This coheres with (5.37) for  $F = \langle -, f \rangle_{L^2}$  and  $G = \langle -, h \rangle_{L^2}$ .

## 5.5 Locality of the $P(\phi)$ interaction and the Restricted GFF

Let  $M$  be a closed Riemannian surface and  $\Omega^\circ \subset M$  a domain with smooth boundary  $\partial\Omega$ . Denote  $\Omega = \overline{\Omega^\circ}$ .

**Remark 5.10.** In general one could consider regular domains  $\Omega^\circ$ . A domain is called regular if  $W_{\Omega^\circ}^{-1}(M) = W_\Omega^{-1}(M)$ , the two spaces defined by (C.3) and (C.2). See also Simon [71] page 267.

Suppose  $\chi \in C^\infty(M)$ . We define

$$S_{\Omega, \varepsilon, \chi}(\phi) \stackrel{\text{def}}{=} \int_M \chi(x) : P(\phi_{\Omega, \varepsilon}^D(x) + \phi_{\partial\Omega, \varepsilon}(x)) : dV_M(x), \quad (5.39)$$

where

$$\phi_{\Omega, \varepsilon}^D(x) \stackrel{\text{def}}{=} \phi(P_{\Omega^c}^\perp E_\varepsilon \delta_x), \quad \phi_{\partial\Omega, \varepsilon}(x) \stackrel{\text{def}}{=} \phi(P_{\partial\Omega} E_\varepsilon \delta_x), \quad \phi \sim \mu_{\text{GFF}}^M. \quad (5.40)$$

**Proposition 5.18** (locality). *Whenever  $\chi \in C_c^\infty(\Omega^\circ)$ , we have, for any fixed Wick ordering  $:\bullet:$ ,*

$$S_{M,\chi}(\phi) = \int_M \chi(x) :P(\phi(x)) : dV_M(x) = \lim_{\varepsilon \rightarrow 0} S_{\Omega,\varepsilon,\chi}(\phi), \quad (5.41)$$

and hence  $S_{M,\chi}$  is  $\mathcal{O}_\Omega$ -measurable. In particular,

$$\int_\Omega :P(\phi(x)) : dV_\Omega = \int_\Omega :P(\phi_\Omega^D(x) + \phi_{\partial\Omega}(x)) : dV_\Omega \quad (5.42)$$

and is  $\mathcal{O}_\Omega$ -measurable.

*Proof.* Without loss of generality we suppose  $\text{supp } \chi$  stays  $\delta$ -away from  $\partial\Omega$  for some  $\delta > 0$ . Then corollary 3.9 and approximation in  $L^4$  of a general  $\chi$  as well as  $1_\Omega$  gives the result. We note that the support of  $E_\varepsilon \delta_x$  is contained in the  $\varepsilon$ -ball around  $x$ . Then whenever  $\varepsilon < \delta$  for  $x \in \text{supp } \chi$  we have  $\text{supp}(E_\varepsilon \delta_x) \subset \Omega$ , hence

$$P_\Omega^\perp E_\varepsilon \delta_x = 0, \quad (5.43)$$

and

$$\phi_\varepsilon(x) \equiv \phi(P_{\Omega^c}^\perp E_\varepsilon \delta_x) + \phi(P_{\partial\Omega} E_\varepsilon \delta_x), \quad (5.44)$$

both  $\mathcal{O}_\Omega$ -measurable. Thus

$$:P(\phi_{\Omega,\varepsilon}^D(x) + \phi_{\partial\Omega,\varepsilon}(x)) : \equiv :P(\phi_\varepsilon(x)) : \quad (5.45)$$

for  $x \in \text{supp } \chi$  and  $\varepsilon < \delta$ , and is  $\mathcal{O}_\Omega$ -measurable. Therefore (5.41) holds by proposition 3.5.  $\square$

We emphasize here that for our purposes we employ the Wick ordering  $:\bullet:{}_0$  of section 3.4 in defining  $S_{M,\chi}$ . This is to say

$$:\phi_\varepsilon(x)^{2n}: = \sum_{j=0}^n \frac{(-1)^j (2n)!}{(2n-2j)! j! 2^j} C_\varepsilon(x)^j \phi_\varepsilon(x)^{2n-2j}, \quad (5.46)$$

where

$$C_\varepsilon(x) = \iint E_\varepsilon(x,y) C_0(y,z) E_\varepsilon(z,x) dV_M(y) dV_M(z). \quad (5.47)$$

Since  $E_\varepsilon(x, -)$  is supported in an  $\varepsilon$ -ball around  $x$ , and  $C_0(y,z)$  depends only on the geometry of  $M$  restricted to a convex neighborhood of  $y, z$ , the term  $C_\varepsilon(x)$  depends only on the geometry of  $M$  locally near  $x$ . This means that under  $:\bullet:{}_0$ , once  $\text{supp } \chi \subset \Omega$ , the limiting (integrated) random variable  $S_{M,\chi}$ , in addition to being  $\mathcal{O}_\Omega$ -measurable, is in fact fully determined with knowledge of the metric  $g|_\Omega$  restricted to  $\Omega$ . This allows the freedom of choosing the ambient manifold  $M$  where  $\Omega$  isometrically embeds in defining the interaction over  $\Omega$  (see definition 5.4).

Now we repeat proposition 5.18 in a different way:

**Corollary 5.19.** *Whenever  $\chi \in C_c^\infty(\Omega^\circ)$ , for any fixed Wick ordering  $:\bullet:$ ,*

$$\mathbb{E}_{\text{GFF}}^M[S_{M,\chi} | \mathcal{O}_\Omega] = S_{M,\chi}, \quad (5.48)$$

and  $S_{M,\chi}$  is independent of  $\phi_{\Omega^c}^D$ , that is, of  $\phi(P_\Omega^\perp f)$  for all  $f \in C^\infty(M)$  or  $\phi((\Delta + m^2)f)$  for all  $f \in C_c^\infty(\Omega^\circ)$ . Moreover, since  $\Omega$  is regular,  $S_{M,\chi}$  is the limit in  $L^2(\mu_{\text{GFF}}^M)$  of polynomials of random variables of the form  $\phi(f)$  with  $f \in C_c^\infty(\Omega^\circ)$  via the Itô-Wiener-Segal isomorphism.  $\square$

This result motivates

**Definition 5.3.** Let  $\Omega^\circ \subset M$  be a regular domain. The (massive) Gaussian Free Field over  $\Omega^\circ$  restricted from  $M$  is the centered Gaussian process indexed by  $C_c^\infty(\Omega^\circ)$ , with covariance

$$\mathbb{E}[\phi(f)\phi(h)] = \langle f, h \rangle_{W_\Omega^{-1}(M)} \quad (5.49)$$

for any  $f, h \in C_c^\infty(\Omega^\circ)$ . We denote it by  $\phi|_\Omega$ .

Since the inner product of  $W_\Omega^{-1}(M)$  depends on  $M$ , the restricted GFF does not make sense on  $\Omega^\circ$  alone. However, by corollary 5.10, it is equal in law to  $\phi_{\partial\Omega} + \phi_\Omega^D$ , where  $\phi_\Omega^D$  does make sense over  $\Omega^\circ$  and the law of  $\phi_{\partial\Omega} = \text{PI}_\Omega^{\partial\Omega}(\tau_{\partial\Omega}\phi)$  is determined via  $\text{PI}_\Omega^{\partial\Omega}$  by that of a boundary data over  $\partial\Omega$ .

Going back to corollary 5.19 we see that for  $\chi \in C_c^\infty(\Omega^\circ)$ , the interaction  $S_{M,\chi}$  can now be defined as a random variable of the sample  $\phi|_\Omega \in \mathcal{D}'(\Omega^\circ)$ . Equally, it is also a random variable over  $\mathcal{D}'(\Omega^\circ) \times \mathcal{D}'(\partial\Omega)$  equipped with  $\mu_{\text{GFF}}^{\Omega,D} \otimes \mu_C^{\partial\Omega}$  where  $\mu_C^{\partial\Omega}$  is some probability measure on  $\mathcal{D}'(\partial\Omega)$  mutually absolutely continuous with respect to  $\tau_{\partial\Omega}(\mu_{\text{GFF}}^M)$  and on the same  $\sigma$ -algebra. We consider only a few very specific candidates for  $\mu_C^{\partial\Omega}$ .

Now suppose we start with a Riemannian surface  $(\Omega, g)$  with totally geodesic boundary  $\partial\Omega$ . With the help of the isometric double  $|\Omega|^2$  discussed in remark 4.7 we define

**Definition 5.4.** The interaction over  $\Omega$  is

$$S_\Omega(\phi_\Omega^D|\varphi) = S_\Omega(\phi_\Omega^D + \text{PI}_\Omega^{\partial\Omega} \varphi) \stackrel{\text{def}}{=} S_{|\Omega|^2, 1_\Omega}(\phi_\Omega^D + \text{PI}_\Omega^{\partial\Omega} \varphi) \quad (5.50)$$

as an  $L^2$ -random variable over  $\mathcal{D}'(\Omega^\circ) \times \mathcal{D}'(\partial\Omega)$  equipped with  $\mu_{\text{GFF}}^{\Omega, D} \otimes \mu_{2\text{DN}}^{\partial\Omega, \Omega}$ , where the latter is the Gaussian measure with covariance operator  $\frac{1}{2}(\text{DN}_\Omega^{\partial\Omega})^{-1}$ .

**Remark 5.11.** We emphasize here that the variable  $S_\Omega(\phi_\Omega^D|\varphi)$  must be understood for  $\phi_\Omega^D$  and  $\varphi$  both random, with the law of  $\varphi$  mutually absolutely continuous with respect to  $\mu_{2\text{DN}}^{\partial\Omega, \Omega}$  (thus  $\mu_{2\text{DN}}^{\partial\Omega, \Omega}$  is a valid candidate). Thus if one asks how much can one “fix”  $\varphi$  as one interprets  $S_\Omega(\phi_\Omega^D|\varphi)$ , the answer is that it makes sense as a random variable of  $\phi_\Omega^D$  alone for almost every fixed  $\varphi$  under  $\mu_{2\text{DN}}^{\partial\Omega, \Omega}$  (but this full-measure set with respect to  $\mu_{2\text{DN}}^{\partial\Omega, \Omega}$  is generally unknown). Said from a slightly different perspective, the expression  $S_\Omega(\phi)$  for a generic  $\phi \in \mathcal{D}'(\Omega^\circ)$  makes sense as a random variable only when  $\phi$  follows a law mutually absolutely continuous with respect to  $\mu_{\text{GFF}}^{\Omega, D} * (\text{PI}_\Omega^{\partial\Omega}) * \mu_{2\text{DN}}^{\partial\Omega, \Omega}$ , where “\*” denotes convolution product. Statements involving  $S_\Omega(\phi_\Omega^D|\varphi)$  in the sequel should be understood in this sense.

The following result will be useful in lemma 6.2.

**Proposition 5.20.** For any fixed Wick ordering  $:\bullet:$  we have

$$\int_M :P(\phi(x)):dV_M = \int_\Omega :P(\phi_\Omega^D(x) + \phi_{\partial\Omega}(x)):dV_\Omega + \int_{\Omega^c} :P(\phi_{\Omega^c}^D(x) + \phi_{\partial\Omega}(x)):dV_{\Omega^c} \quad (5.51)$$

in  $L^p(\mu_{\text{GFF}}^M)$  for all  $1 \leq p < \infty$  and in particular pointwise almost surely, as a result

$$e^{-\int_M :P(\phi(x)):dV_M} = e^{-\int_\Omega :P(\phi_\Omega^D(x) + \phi_{\partial\Omega}(x)):dV_\Omega} e^{-\int_{\Omega^c} :P(\phi_{\Omega^c}^D(x) + \phi_{\partial\Omega}(x)):dV_{\Omega^c}} \quad (5.52)$$

pointwise almost surely and thus also in  $L^1(\mu_{\text{GFF}}^M)$ .

**Remark 5.12.** Note that the equality (5.51) says nothing about pointwise almost sure convergence of the mollified sequences  $S_{M, \varepsilon}$ ,  $S_{\Omega, \varepsilon}$ , and  $S_{\Omega^c, \varepsilon}$ . In fact, only separate subsequences in  $\varepsilon$  will converge pointwise almost surely to the three respective terms in (5.51).

*Proof.* Consider a smooth partition of unity

$$1_M = \chi_\Omega + \chi_{\partial\Omega} + \chi_{\Omega^c}, \quad (5.53)$$

where  $\chi_\Omega$  and  $\chi_{\Omega^c}$  are supported in the interiors of  $\Omega$  and  $\Omega^c$  respectively and  $\chi_{\partial\Omega}$  is supported near  $\partial\Omega$ . Note then that (5.53) holds as well in  $L^4(M)$  so by corollary 3.9 we have

$$\int_M :P(\phi(x)):dV_M = \int_M \chi_\Omega :P(\phi(x)):dV_M + \int_M \chi_{\partial\Omega} :P(\phi(x)):dV_M + \int_M \chi_{\Omega^c} :P(\phi(x)):dV_M \quad (5.54)$$

in  $L^p(\mu_{\text{GFF}}^M)$  for  $1 \leq p < \infty$ . Now take

$$\chi_\Omega \rightarrow 1_\Omega, \quad \chi_{\Omega^c} \rightarrow 1_{\Omega^c}, \quad \text{and } \chi_{\partial\Omega} \rightarrow 0, \quad (5.55)$$

in  $L^4(M)$ , keeping the equality (5.53) in the process. Then

$$\begin{aligned} \int_M :P(\phi(x)):dV_M &= \int_M 1_\Omega :P(\phi(x)):dV_M + \int_M 1_{\Omega^c} :P(\phi(x)):dV_M \\ &= \int_\Omega :P(\phi_\Omega^D(x) + \phi_{\partial\Omega}(x)):dV_\Omega + \int_{\Omega^c} :P(\phi_{\Omega^c}^D(x) + \phi_{\partial\Omega}(x)):dV_{\Omega^c} \end{aligned}$$

in  $L^p(\mu_{\text{GFF}}^M)$  for  $1 \leq p < \infty$ , and we finish the proof.  $\square$

## 6 Presenting the $P(\phi)_2$ Model as a Segal Theory

### 6.1 Description of Segal’s Rules as Pertains to the Model

The prototype of what we propose here is definition (4.4) on page 460 of [66], then we relate to the “transfer operator” formalism on page 455-456 (lemma 6.1). We will consider real (separable) Hilbert spaces. We denote

the category of such spaces with Hilbert-Schmidt operators as morphisms by  $\mathbf{Hilb}_{\mathbb{R}}$ . The reason for Hilbert-Schmidt is lemma A.7.

As a preliminary consideration, let  $\mathcal{C}$  denote the “category” whose objects are finite disjoint unions  $\Sigma_j = \bigsqcup_i \mathbb{S}_i^1$  of Riemannian circles, each of which determined by its radius  $R_i$ , and for any finite collection of such objects  $(\Sigma_j)_j$  we say that an *cobordism/unoriented morphism* among them is simply an orientable Riemannian surface with a *totally geodesic* boundary  $\partial\Omega$  which is identified with  $\bigsqcup_j \Sigma_j$  via an isometry (see also remark 6.4). With this in mind we can consider the following definition. Morphisms will be oriented once we make the transfer operator connection.

**Definition 6.1** ([66] page 460). A 2d Riemannian QFT is a correspondence  $\mathcal{C} \rightarrow \mathbf{Hilb}_{\mathbb{R}}$ , such that

- (i) to each oriented Riemannian circle  $\mathbb{S}_R^1$  of radius  $R$  there is associated a Hilbert space  $\mathcal{H}_R$ , and to the disjoint union  $\bigsqcup_{i \in I} \mathbb{S}_i^1$  there is associated the tensor product  $\bigotimes_{i \in I} \mathcal{H}_i$  of the corresponding single-circle Hilbert spaces; in the degenerate case  $R = 0$ , we let  $\mathcal{H}_0 = \mathbb{R}$ ;
- (ii) (*operator-reflection*) to each Riemannian surface  $\Omega$  such that  $\partial\Omega = \Sigma$ , without distinguishing the orientations on the components, we associate an element  $\mathcal{A}_\Omega \in \mathcal{H}_\Sigma$ ; in the degenerate case where  $\Omega = M$  is closed, we associate a real number  $Z_M \in \mathbb{R}$ , called the *partition function*;
- (iii) (*sewing-trace*) suppose  $\Sigma_i$  and  $\Sigma_j$  are two connected components of  $\Sigma$  that are isometric, and let  $\rho : \Sigma_i \rightarrow \Sigma_j$  be an (orientation reversing) isometry. Then

$$\mathcal{A}_{\Omega/\rho} = \mathrm{tr}_\rho(\mathcal{A}_\Omega), \quad (6.1)$$

where  $\Omega/\rho$  is the surface obtained from  $\Omega$  by gluing  $\Sigma_i$  with  $\Sigma_j$  along  $\rho$ , and  $\mathrm{tr}_\rho$  is the *trace map* such that writing  $\Sigma = \Sigma_i \sqcup \Sigma_j \sqcup \Sigma'$  (possibly  $\Sigma' = \{\mathrm{pt}\}$ ),  $\mathrm{tr}_\rho$  is the map

$$\begin{aligned} \mathrm{tr}_\rho : \mathcal{H}_{\Sigma_i} \otimes \mathcal{H}_{\Sigma_j} \otimes \mathcal{H}_{\Sigma'} &\longrightarrow \mathcal{H}_{\Sigma'}, \\ F \otimes G \otimes H &\longmapsto \langle \rho_* F, G \rangle_{\Sigma_j} H, \end{aligned} \quad (6.2)$$

with  $\langle -, - \rangle_{\Sigma_j}$  being the (real) inner product on  $\mathcal{H}_{\Sigma_j}$ .

**Remark 6.1.** A word of caution should be said immediately concerning (6.2). As written,  $\mathrm{tr}_\rho$  would not extend continuously to the whole of  $\mathcal{H}_{\Sigma_i} \otimes \mathcal{H}_{\Sigma_j} \otimes \mathcal{H}_{\Sigma'}$  for “infinitely entangled” states in  $\mathcal{H}_{\Sigma_i} \otimes \mathcal{H}_{\Sigma_j}$ . However, for us each  $\mathcal{H}_\Sigma$  is  $L^2(Q_\Sigma, \mu_\Sigma)$  for a probability space  $(Q_\Sigma, \mu_\Sigma)$ , and thus by proposition A.4,  $\mathcal{H}_{\Sigma_i} \otimes \mathcal{H}_{\Sigma_j} \otimes \mathcal{H}_{\Sigma'}$  is represented as  $L^2(Q \times Q \times Q', \mu \otimes \mu \otimes \mu')$ , where we identify  $(Q_{\Sigma_i}, \mu_{\Sigma_i})$  and  $(Q_{\Sigma_j}, \mu_{\Sigma_j})$  with  $(Q, \mu)$  via the isometry. The action of  $\mathrm{tr}_\rho$  should be understood as

$$\mathrm{tr}_\rho : \mathcal{A}_\Omega(x, y, z) \longmapsto \int \mathcal{A}_\Omega(x, x, z) d\mu(x), \quad (6.3)$$

when the latter integral converges, which we will show to happen for our model in sections 6.4 and 6.5. This is analogous to the “flat trace” of Atiyah and Bott [2]. It can be shown easily that (6.3) coincides with (6.2) on “finitely entangled” states of  $\mathcal{H}_{\Sigma_i} \otimes \mathcal{H}_{\Sigma_j} \otimes \mathcal{H}_{\Sigma'}$ , that is, finite linear combinations  $\sum_i F_i \otimes G_i \otimes H_i$ . However, the finiteness of neither (6.3) nor (6.2) would imply the operator  $U_\Omega$  of (6.4) is trace class proper.

Now we manufacture Hilbert-Schmidt operators in  $\mathbf{Hilb}_{\mathbb{R}}$  out of definition 6.1 with the help of lemma A.7. Let  $\Sigma_{\mathrm{in}}, \Sigma_{\mathrm{out}}$  be two objects in  $\mathcal{C}$ , now *oriented*. We say that an unoriented morphism  $\Omega$  among  $\Sigma_{\mathrm{in}}, \Sigma_{\mathrm{out}}$  is in  $\mathrm{Mor}(\Sigma_{\mathrm{in}}, \Sigma_{\mathrm{out}})$  if the isometry  $\Sigma_{\mathrm{in}} \sqcup \Sigma_{\mathrm{out}} \rightarrow \partial\Omega$  identifies the orientation of  $\Sigma_{\mathrm{in}}$  with that induced by an *inward pointing* normal on  $\partial\Omega$ , and that of  $\Sigma_{\mathrm{out}}$  an *outward pointing* normal.

Lemma A.7 and definition 6.1 now implies

**Lemma 6.1.** *If  $\Omega \in \mathrm{Mor}(\Sigma_{\mathrm{in}}, \Sigma_{\mathrm{out}})$ , then the Segal transfer operator  $U_\Omega$  defined by*

$$\begin{aligned} U_\Omega : \mathcal{H}_{\Sigma_{\mathrm{in}}} &\longrightarrow \mathcal{H}_{\Sigma_{\mathrm{out}}}, \\ F &\longmapsto \mathrm{tr}_\rho(F \otimes \mathcal{A}_\Omega), \end{aligned} \quad (6.4)$$

where  $\mathrm{tr}_\rho$  is the map of (6.3) with  $\Sigma_i = \Sigma_j = \Sigma_{\mathrm{in}}$ ,  $\Sigma' = \Sigma_{\mathrm{out}}$ , and  $\rho = \mathbb{1}$ , is Hilbert-Schmidt. Moreover,

- (i) if  $\Omega_1 \in \mathrm{Mor}(\Sigma_1, \Sigma_2)$ ,  $\Omega_2 \in \mathrm{Mor}(\Sigma_2, \Sigma_3)$ , and  $\Omega_2 \cup_2 \Omega_1$  the Riemannian connected sum by gluing components corresponding to  $\Sigma_2$ , then

$$U_{\Omega_2 \cup_2 \Omega_1} = U_{\Omega_2} \circ U_{\Omega_1}. \quad (6.5)$$

- (ii) suppose  $\Omega \in \mathrm{Mor}(\Sigma, \Sigma)$ , with  $\Sigma$  identified with itself along an isometry  $\rho$ , and denote by  $\check{\Omega}$  the closed surface obtained by gluing these components together, then

$$Z_{\check{\Omega}} = \mathrm{tr}_\rho(U_\Omega). \quad (6.6)$$




Figure 1: sewing

(iii) if  $\Omega \in \text{Mor}(\Sigma_1, \Sigma_2)$ , denote by  $\Omega^* \in \text{Mor}(\Sigma_2, \Sigma_1)$  the surface obtained by reversing the orientations of boundaries of  $\Omega$  without changing the orientation of  $\Omega$ , then

$$U_{\Omega^*} = U_{\Omega}^{\dagger}, \quad (6.7)$$

with  $U_{\Omega}^{\dagger}$  denoting the (real) adjoint of  $U_{\Omega}$ .  $\square$

**Remark 6.2.** As alluded to at the bottom of subsection 2.2.3, one could allow each  $\mathcal{A}_{\Omega}$  to be a ray instead of a fixed vector in  $\mathcal{H}_{\partial\Omega}$ , which eventually enables an arbitrary (nonzero) constant to be included in (6.5), obtaining the so-called projective version of the axiom. This is reasonable since quantum states are rays, not vectors.

**Remark 6.3.** An interesting remark appears at the bottom of page 457 of [66], namely the conjecture of Friedan that a theory in the sense of definition 6.1 is in fact completely determined by its restriction to *closed surfaces*, namely the knowledge of the numbers  $Z_M$  for all closed  $M$ . While far from proving this conjecture, we shall see in section 6.3 that the correct definition of the amplitude “derives” very naturally from a special case of the trace axiom ((iii) of definition 6.1). It seems certain that if one has knowledge not just of the numbers  $Z_M$  but the *measures*  $\mu_{P(\phi)}^M$  as in (2.1), then the theory would be determined.

**Remark 6.4.** To have safety of smooth gluing while taking into account at the same time our reflection constructions (the isometric double) required in section 6.3, one should enhance the objects  $\Sigma$  with **symmetric two-sided collars**  $\Sigma \times (-\varepsilon, \varepsilon)$  equipped with symmetric metrics making  $\Sigma \times \{0\}$  geodesic, and a cobordism  $\Omega$  as described above must allow each  $\Sigma_j$  as a component of  $\partial\Omega$  to have a tubular neighborhood isometric to  $\Sigma_j \times (-\varepsilon, 0]$ . Here **symmetric** means having an isometric involution  $\Theta$  that exchanges  $\Sigma \times \{\pm t\}$  and fixes exactly  $\Sigma \times \{0\}$  (actually the reflection would ensure  $\Sigma \times \{0\}$  being geodesic, see [58] section 2.1.1 and Alekseevsky et al [3]). In fact, one should consider all proper ingredients that make up what is called the **Riemannian bordism category** as described in detail in section 6.2 of H. Hohnhold, S. Stolz, and P. Teichner [35]. Perhaps a better consideration is not fixed collars but *germs* of collars; see also the discussion in Kontsevich and Segal [40] following definition 3.1. However, the problem of finding the right underlying category is somewhat orthogonal to the problems considered in this article.

## 6.2 The Hilbert Spaces

Now we associate a real Hilbert space  $\mathcal{H}_R$  to the Riemannian circle  $\mathbb{S}_R^1$  of radius  $R$ . Let

$$\mathbf{D}_R \stackrel{\text{def}}{=} (\Delta_{\mathbb{S}_R^1} + m^2)^{1/2} \quad (6.8)$$

be the positive square root of the positive Helmholtz operator on  $\mathbb{S}_R^1$ . The circle having radius  $R$  would mean that we take  $\Delta_{\mathbb{S}_R^1} := -\partial_{\theta}^2$  when the metric is  $Rd\theta$ , parametrized by the arc length  $\theta$ . Denote by  $\mu_{(2\mathbf{D}_R)^{-1}}$  the Gaussian measure on  $\mathcal{D}'(\mathbb{S}_R^1)$  with covariance  $(2\mathbf{D}_R)^{-1}$  (for convenience, we just write  $\mu_{2\mathbf{D}}$ ). Then we define

$$\mathcal{H}_R \stackrel{\text{def}}{=} L_{\mathbb{R}}^2(\mathcal{D}'(\mathbb{S}_R^1), \hat{\mu}_{2\mathbf{D}}). \quad (6.9)$$



Here we take  $\widehat{\mu}_{2\mathbf{D}} := \mu_{2\mathbf{D}} \cdot \det_{\zeta}(2\mathbf{D}_R)^{-\frac{1}{2}}$ , namely the probability measure scaled by the positive finite constant  $\det_{\zeta}(2\mathbf{D}_R)^{-\frac{1}{2}}$ . Indeed, this same method could be applied directly to (finite) disjoint unions  $\Sigma = \bigsqcup_i \mathbb{S}_i^1$  and produce a measure on the corresponding  $\mathcal{D}'(\Sigma)$ , and this is compatible with (i) of definition 6.1. Indeed,

$$\mathcal{D}'(\mathbb{S}_1^1 \sqcup \mathbb{S}_2^1) = \mathcal{D}'(\mathbb{S}_1^1) \oplus \mathcal{D}'(\mathbb{S}_2^1) \quad (6.10)$$

and the operator  $\mathbf{D}_{\Sigma}$  acts diagonally, and  $\mu_{2\mathbf{D}}^{\mathbb{S}_1 \sqcup \mathbb{S}_2} = \mu_{2\mathbf{D}}^{\mathbb{S}_1} \otimes \mu_{2\mathbf{D}}^{\mathbb{S}_2}$  in view of lemma 2.7 (for Gaussian fields). The real  $L^2$  space would then be the tensor product (lemma A.4). Alternatively, one could argue from B.5.

### 6.3 Amplitudes = Schwartz Kernels

Let  $\Omega$  be an unoriented morphism among  $(\Sigma_j)_j$ . We denote  $\Sigma := \bigsqcup_j \Sigma_j$ , and seek to define  $\mathcal{A}_{\Omega} \in \mathcal{H}_{\Sigma}$ . We consider gluing  $\Omega$  with itself along an orientation reversing isometry  $\rho : \Sigma \rightarrow \Sigma$ , forming the isometric double  $|\Omega|^2 = \Omega^* \cup_{\Sigma} \Omega$  (here  $\Omega^*$ , as in (iii) of lemma 6.1, denote the copy of  $\Omega$  with coorientation of  $\Sigma$  reversed). Suppose we have defined  $\mathcal{A}_{\Omega}$  satisfying definition 6.1, then (iii) would imply

$$\begin{aligned} Z_{|\Omega|^2} &= \int |\mathcal{A}_{\Omega}(\varphi)|^2 d\mu_{2\mathbf{D}}^{\Sigma}(\varphi) \det_{\zeta}(2\mathbf{D}_{\Sigma})^{-\frac{1}{2}} \\ &= \int_{\mathcal{D}'(M)} e^{-S_{|\Omega|^2}(\phi)} d\mu_{\text{GFF}}^{|\Omega|^2}(\phi) \det_{\zeta}(\Delta_{|\Omega|^2} + m^2)^{-\frac{1}{2}} \\ &= \int_{\mathcal{D}'(\Sigma)} d\tau_{\Sigma}(e^{-S_{|\Omega|^2}} \cdot \mu_{\text{GFF}}^{|\Omega|^2})(\varphi) \det_{\zeta}(\Delta_{|\Omega|^2} + m^2)^{-\frac{1}{2}}. \end{aligned}$$

In the case  $\Omega \in \text{Mor}(\Sigma_1, \Sigma_2)$  this corresponds to the fact that  $Z_{|\Omega|^2} = \text{tr}_{\rho}(U_{\Omega^*} U_{\Omega})$ . This motivates the following definition.

**Definition 6.2.** Let  $\Omega$  be an unoriented morphism among  $\Sigma$  and  $P$  the polynomial defining the interaction. We define the amplitude associated to  $\Omega$  to be the quantity

$$\mathcal{A}_{\Omega}^P(\varphi) \stackrel{\text{def}}{=} \left[ \frac{\det_{\zeta}(\Delta_{|\Omega|^2} + m^2)^{-\frac{1}{2}}}{\det_{\zeta}(2\mathbf{D}_{\Sigma})^{-\frac{1}{2}}} \cdot \frac{d[\tau_{\Sigma}(e^{-S_{|\Omega|^2}} \cdot \mu_{\text{GFF}}^{|\Omega|^2})](\varphi)}{d\mu_{2\mathbf{D}}^{\Sigma}}(\varphi) \right]^{\frac{1}{2}}, \quad (6.11)$$

where the second ratio denotes the Radon-Nikodym density. (Compare Pickrell [52] definition 3.)

This is well-defined since  $\tau_{\Sigma}(e^{-S_{|\Omega|^2}} \cdot \mu_{\text{GFF}}^{|\Omega|^2}) \ll \mu_{2\mathbf{D}}^{\Sigma}$  as  $\tau_{\Sigma}(\mu_{\text{GFF}}^{|\Omega|^2}) = \mu_{2\text{DN}}^{\Sigma, \Omega} \ll \mu_{2\mathbf{D}}^{\Sigma}$  by corollary 2.15, and both are finite positive measures. We see also that  $\mathcal{A}_{\Omega} > 0$  almost surely with respect to  $\widehat{\mu}_{2\mathbf{D}}^{\Sigma}$ . We have automatically  $\mathcal{A}_{\Omega} \in L^2(\mathcal{D}'(\Sigma), \widehat{\mu}_{2\mathbf{D}}^{\Sigma}) = \mathcal{H}_{\Sigma}$  since  $|\mathcal{A}_{\Omega}|^2 \in L^1(\mathcal{D}'(\Sigma), \widehat{\mu}_{2\mathbf{D}}^{\Sigma})$  by definition.

**Example 6.1.** Let us derive the amplitude  $\mathcal{A}_{\Omega}^0$  for the free field, that is, with  $S_{|\Omega|^2} = 0$ . Indeed,

$$|\mathcal{A}_{\Omega}^0(\varphi)|^2 = \frac{d\tau_{\Sigma}(\mu_{\text{GFF}}^{|\Omega|^2} \det_{\zeta}(\Delta_{|\Omega|^2} + m^2)^{-\frac{1}{2}})(\varphi)}{d\mu_{2\mathbf{D}}^{\Sigma} \det_{\zeta}(2\mathbf{D})^{-\frac{1}{2}}}(\varphi). \quad (6.12)$$

Since

$$\tau_{\Sigma}(\mu_{\text{GFF}}^{|\Omega|^2}) = \mu_{2\text{DN}}^{\Sigma, \Omega}, \quad (6.13)$$

namely the Gaussian measure on  $\mathcal{D}'(\Sigma)$  with covariance  $\frac{1}{2}(\text{DN}_{\Omega}^{\Sigma})^{-1}$ , taking into account the BFK formula

$$\det_{\zeta}(\Delta_{|\Omega|^2} + m^2) = [\det_{\zeta}(\Delta_{\Omega, D} + m^2)]^2 \det_{\zeta}(2\text{DN}_{\Omega}^{\Sigma}), \quad (6.14)$$

we obtain, by corollary 2.15,

$$\mathcal{A}_{\Omega}^0(\varphi) = \det_{\zeta}(\Delta_{\Omega, D} + m^2)^{-\frac{1}{2}} e^{-\frac{1}{2}(\varphi, (\text{DN}_{\Omega}^{\Sigma} - \mathbf{D})\varphi)_{L^2(\Sigma)}}. \quad (6.15)$$

**Lemma 6.2.** We have

$$\mathcal{A}_{\Omega}^P(\varphi) = \mathcal{A}_{\Omega}^0(\varphi) \int e^{-S_{\Omega}(\phi_{\Omega}^D|\varphi)} d\mu_{\text{GFF}}^{\Omega, D}(\phi_{\Omega}^D), \quad (6.16)$$

for almost every  $\varphi$  under  $\mu_{2\mathbf{D}}^{\Sigma}$ . Here  $S_{\Omega}(\phi_{\Omega}^D|\varphi)$  is as defined in definition 5.4.

*Proof.* Indeed, by definition of the measure image,

$$\mathcal{A}_\Omega^P(\varphi)^2 = \mathcal{A}_\Omega^0(\varphi)^2 \iint e^{-S_{|\Omega|^2}(\phi_\Omega^D + \phi_{\Omega^*}^D + \text{PI}_{|\Omega|^2}^\Sigma \varphi)} d\mu_{\text{GFF}}^{\Omega, D}(\phi_\Omega^D) \otimes d\mu_{\text{GFF}}^{\Omega^*, D}(\phi_{\Omega^*}^D). \quad (6.17)$$

However, by proposition 5.20 which decouples the interaction into a sum over complementary regions,

$$e^{-S_{|\Omega|^2}(\phi)} = e^{-S_\Omega(\phi_\Omega^D|\varphi)} e^{-S_{\Omega^*}(\phi_{\Omega^*}^D|\varphi)}, \quad (6.18)$$

almost surely against  $\mu_{\text{GFF}}^{\Omega, D} \otimes \mu_{\text{GFF}}^{\Omega^*, D} \otimes \tau_\Sigma(\mu_{\text{GFF}}^{|\Omega|^2})$ , and hence also against  $\mu_{\text{GFF}}^{\Omega, D} \otimes \mu_{\text{GFF}}^{\Omega^*, D} \otimes \mu_{2\mathbf{D}}^\Sigma$ . Thus by reflection symmetry between  $\Omega$  and  $\Omega^*$  and independence between  $\phi_\Omega^D$  and  $\phi_{\Omega^*}^D$  we obtain the result.  $\square$

**Remark 6.5.** By remark 5.6 we have  $\mathcal{A}_\Omega^P(\varphi) = \mathcal{A}_\Omega^0(\varphi) \mathbb{E}[e^{-S_{|\Omega|^2}(\phi_\Omega^D|\varphi)} | \mathcal{O}_\Sigma](\varphi)$  where  $\mathbb{E}$  is taken under  $\mu_{\text{GFF}}^{\Omega, D} \otimes \mu_{2\mathbf{D}}^\Sigma$ . Indeed, one could argue alternatively since

$$\frac{d(e^{-S_{|\Omega|^2}} \cdot \mu_{\text{GFF}}^{|\Omega|^2})}{d(\mu_{\text{GFF}}^{|\Omega|^2} | \mathcal{O}_\Sigma)} = \mathbb{E}_{\text{GFF}}^{|\Omega|^2}[e^{-S_{|\Omega|^2}} | \mathcal{O}_\Sigma], \quad (6.19)$$

and by lemma 5.15 the map  $\tau_\Sigma$  generates  $\mathcal{O}_\Sigma$  whence  $\tau_\Sigma(\mu_{\text{GFF}}^{|\Omega|^2} | \mathcal{O}_\Sigma) = \tau_\Sigma(\mu_{\text{GFF}}^{|\Omega|^2}) = \mu_{2\mathbf{D}\Sigma}^\Sigma$ . This viewpoint however plays no essential role in the proofs.

## 6.4 Trace Axiom and its Consequences

In this subsection we treat separately (ii) of lemma 6.1 as part of (iii) of definition 6.1. Let  $\Omega \in \text{Mor}(\Sigma, \Sigma)$ , *not necessarily connected*. We will consider two closed surfaces:  $\tilde{\Omega}$  and  $|\Omega|^2 := (\Omega^* \cup_{\Sigma \cup \Sigma} \Omega)^\vee$ .

**Proposition 6.3** (pre-trace). *We have*

$$\frac{d\tau_\Sigma(\mu_{\text{GFF}}^{\tilde{\Omega}})}{d\mu_{2\mathbf{D}}^\Sigma}(\varphi) = \frac{\det_\zeta(\text{DN}_\Omega^\Sigma)^{\frac{1}{2}}}{\det_\zeta(2\text{DN}_\Omega^{\Sigma \cup \Sigma})^{\frac{1}{4}}} \left( \frac{d\tau_{\Sigma \cup \Sigma}(\mu_{\text{GFF}}^{|\Omega|^2})}{d\mu_{2\mathbf{D}}^{\Sigma \cup \Sigma}}(\varphi, \varphi) \right)^{\frac{1}{2}}. \quad (6.20)$$

*Proof.* This boils down to comparing the explicit expressions for the densities as given by corollary 2.15. Indeed, the relation that we need is

$$\left\langle \begin{bmatrix} \varphi \\ \varphi \end{bmatrix}, (2\text{DN}_\Omega^{\Sigma \cup \Sigma} - 2\mathbf{D}_{\Sigma \cup \Sigma}) \begin{bmatrix} \varphi \\ \varphi \end{bmatrix} \right\rangle_{L^2(\Sigma \cup \Sigma)} = 2 \langle \varphi, (\text{DN}_\Omega^\Sigma - 2\mathbf{D}_\Sigma) \varphi \rangle_{L^2(\Sigma)}. \quad (6.21)$$

This is true for  $\varphi \in W^{\frac{1}{2}}(\Sigma)$  because  $\mathbf{D}_{\Sigma \cup \Sigma} = \mathbf{D}_\Sigma \oplus \mathbf{D}_\Sigma$  and  $\langle [\varphi], \text{DN}_\Omega^{\Sigma \cup \Sigma} [\varphi] \rangle_{L^2(\Sigma \cup \Sigma)}$  and  $\langle \varphi, \text{DN}_\Omega^\Sigma \varphi \rangle_{L^2(\Sigma)}$  are both the Dirichlet energy of the harmonic extension over  $\Omega$  with boundary condition  $(\varphi, \varphi)$ . The equality then extends to  $\varphi \in W^{-\delta}(\Sigma)$  for small enough  $\delta$  by continuity (see lemma 2.6).  $\square$

**Corollary 6.4** (trace for free field). *In the situation as above, we have*

$$\int \mathcal{A}_\Omega^0(\varphi, \varphi) d\mu_{2\mathbf{D}}^\Sigma(\varphi) \det_\zeta(2\mathbf{D}_\Sigma)^{-\frac{1}{2}} = \det_\zeta(\Delta_{\tilde{\Omega}} + m^2)^{-\frac{1}{2}}. \quad (6.22)$$

*Proof.* Indeed,

$$\begin{aligned} \text{LHS} &= \frac{\det_\zeta(\Delta_{|\Omega|^2} + m^2)^{-\frac{1}{4}}}{\det_\zeta(2\mathbf{D}_\Sigma \oplus 2\mathbf{D}_\Sigma)^{-\frac{1}{4}}} \int \left( \frac{d\tau_{\Sigma \cup \Sigma}(\mu_{\text{GFF}}^{|\Omega|^2})}{d\mu_{2\mathbf{D}}^{\Sigma \cup \Sigma}}(\varphi, \varphi) \right)^{\frac{1}{2}} d\mu_{2\mathbf{D}}^\Sigma(\varphi) \det_\zeta(2\mathbf{D}_\Sigma)^{-\frac{1}{2}} \\ &= \frac{\det_\zeta(2\text{DN}_\Omega^{\Sigma \cup \Sigma})^{\frac{1}{4}} \det_\zeta(\Delta_{|\Omega|^2} + m^2)^{-\frac{1}{4}}}{\det_\zeta(\text{DN}_\Omega^\Sigma)^{\frac{1}{2}} \det_\zeta(2\mathbf{D}_\Sigma)^{-\frac{1}{2}}} \underbrace{\int \frac{d\tau_\Sigma(\mu_{\text{GFF}}^{\tilde{\Omega}})}{d\mu_{2\mathbf{D}}^\Sigma}(\varphi) d\mu_{2\mathbf{D}}^\Sigma(\varphi) \det_\zeta(2\mathbf{D}_\Sigma)^{-\frac{1}{2}}}_{=1} \\ &= \det_\zeta(\text{DN}_\Omega^\Sigma)^{-\frac{1}{2}} \det_\zeta(\Delta_{\Omega, D} + m^2)^{-\frac{1}{2}} && \text{(BFK for } |\Omega|^2) \\ &= \text{RHS}, && \text{(BFK for } \tilde{\Omega}) \end{aligned}$$

finishing the proof.  $\square$

Note that we do not insist that  $\Omega$  be connected. This leads to the following important consequence of proposition 6.3.

**Corollary 6.5.** *Let  $\Omega_1, \Omega_2$  be two dualizable surfaces such that  $\partial\Omega_1 = \partial\Omega_2 = \Sigma$ , all with the same co-orientation. Denote  $M := \Omega_1 \sqcup \Sigma \Omega_2$ , namely  $\Omega_1$  and  $\Omega_2$  glued along  $\Sigma$ . Then*

$$\frac{d\tau_{\Sigma}(\mu_{\text{GFF}}^M)}{d\mu_{2\mathbf{D}}^{\Sigma}}(\varphi) = \frac{\det_{\zeta}(\text{DN}_{\Sigma}^M)^{\frac{1}{2}}}{\det_{\zeta}(2\text{DN}_{\Omega_1}^{\Sigma})^{\frac{1}{4}} \det_{\zeta}(2\text{DN}_{\Omega_2}^{\Sigma})^{\frac{1}{4}}} \left( \frac{d\tau_{\Sigma}(\mu_{\text{GFF}}^{|\Omega_1|^2})}{d\mu_{2\mathbf{D}}^{\Sigma}}(\varphi) \right)^{\frac{1}{2}} \left( \frac{d\tau_{\Sigma}(\mu_{\text{GFF}}^{|\Omega_2|^2})}{d\mu_{2\mathbf{D}}^{\Sigma}}(\varphi) \right)^{\frac{1}{2}}. \quad (6.23)$$

*Proof.* Indeed, in this case  $\Omega := \Omega_1 \sqcup \Omega_2$  can be seen as an element of  $\text{Mor}(\Sigma, \Sigma)$ . Then  $|\Omega|^2 = |\Omega_1|^2 \sqcup |\Omega_2|^2$  and the GFFs on the two components are independent. In addition,  $\text{DN}_{\Omega}^{\Sigma \sqcup \Sigma}$  is the direct sum  $\text{DN}_{\Omega_1}^{\Sigma} \oplus \text{DN}_{\Omega_2}^{\Sigma}$ . Thus (6.20) gives (6.23).  $\square$

**Corollary 6.6** (dissection gluing). *In the same situation as corollary 6.5, we have*

$$\int \mathcal{A}_{\Omega_1}^0(\varphi) \mathcal{A}_{\Omega_2}^0(\varphi) d\mu_{2\mathbf{D}}^{\Sigma}(\varphi) \det_{\zeta}(2\mathbf{D}_{\Sigma})^{-\frac{1}{2}} = \det_{\zeta}(\Delta_M + m^2)^{-\frac{1}{2}}. \quad (6.24)$$

*Proof.* Again, we apply corollary 6.4 directly to the case  $\Omega = \Omega_1 \sqcup \Omega_2$  and  $|\Omega|^2 = |\Omega_1|^2 \sqcup |\Omega_2|^2$ . We note that  $\mathcal{A}_{\Omega_1}^0(\varphi) \mathcal{A}_{\Omega_2}^0(\varphi) = \mathcal{A}_{\Omega}^0(\varphi, \varphi)$  because

$$\frac{d\tau_{\Sigma \sqcup \Sigma}(\mu_{\text{GFF}}^{|\Omega|^2})}{d\mu_{2\mathbf{D}}^{\Sigma \sqcup \Sigma}}(\varphi, \varphi) = \frac{d\tau_{\Sigma}(\mu_{\text{GFF}}^{|\Omega_1|^2})}{d\mu_{2\mathbf{D}}^{\Sigma}}(\varphi) \frac{d\tau_{\Sigma}(\mu_{\text{GFF}}^{|\Omega_2|^2})}{d\mu_{2\mathbf{D}}^{\Sigma}}(\varphi) \quad (6.25)$$

and  $\det_{\zeta}(\Delta_{|\Omega|^2} + m^2) = \det_{\zeta}(\Delta_{|\Omega_1|^2} + m^2) \det_{\zeta}(\Delta_{|\Omega_2|^2} + m^2)$ .  $\square$

Next we deal with the trace axiom in the interacting case. Again let  $\Omega \in \text{Mor}(\Sigma, \Sigma)$  and consider  $\check{\Omega}$ . One has in this case the decomposition

$$\mu_{\text{GFF}}^{\check{\Omega}} = \mu_{\text{GFF}}^{\Omega, D} \otimes \tau_{\Sigma}(\mu_{\text{GFF}}^{\check{\Omega}}) \quad (6.26)$$

and against which

$$S_{\check{\Omega}}(\phi) = S_{\check{\Omega}}(\phi_{\Omega}^D + \text{PI}_{\check{\Omega}}^{\Sigma} \varphi) = S_{\Omega}(\phi_{\Omega}^D | \varphi, \varphi) \quad \text{and} \quad e^{-S_{\check{\Omega}}(\phi_{\Omega}^D + \text{PI}_{\check{\Omega}}^{\Sigma} \varphi)} = e^{-S_{\Omega}(\phi_{\Omega}^D | \varphi, \varphi)} \quad (6.27)$$

for  $\phi_{\Omega}^D \sim \mu_{\text{GFF}}^{\Omega, D}$  and  $\varphi \sim \tau_{\Sigma}(\mu_{\text{GFF}}^{\check{\Omega}})$  almost surely (see also remark 4.6 and 5.11).

**Corollary 6.7** (trace for  $P(\phi)$  field). *In the situation as above, we have*

$$\int \mathcal{A}_{\check{\Omega}}^P(\varphi, \varphi) d\mu_{2\mathbf{D}}^{\Sigma}(\varphi) \det_{\zeta}(2\mathbf{D}_{\Sigma})^{-\frac{1}{2}} = Z_{\check{\Omega}}. \quad (6.28)$$

*Proof.* Indeed,

$$\mathcal{A}_{\check{\Omega}}^P(\varphi, \varphi) = \mathcal{A}_{\Omega}^0(\varphi, \varphi) \int e^{-S_{\Omega}(\phi_{\Omega}^D | \varphi, \varphi)} d\mu_{\text{GFF}}^{\Omega, D}(\phi_{\Omega}^D), \quad (6.29)$$

therefore, with the constants involving determinants working out in exactly the same way as corollary 6.4, one has

$$\begin{aligned} \text{LHS} &= \det_{\zeta}(\Delta_{\check{\Omega}} + m^2)^{-\frac{1}{2}} \int \frac{d\tau_{\Sigma}(\mu_{\text{GFF}}^{\check{\Omega}})}{d\mu_{2\mathbf{D}}^{\Sigma}}(\varphi) d\mu_{2\mathbf{D}}^{\Sigma}(\varphi) \int e^{-S_{\Omega}(\phi_{\Omega}^D | \varphi, \varphi)} d\mu_{\text{GFF}}^{\Omega, D}(\phi_{\Omega}^D) \\ &= \det_{\zeta}(\Delta_{\check{\Omega}} + m^2)^{-\frac{1}{2}} \int e^{-S_{\Omega}(\phi_{\Omega}^D | \varphi, \varphi)} d\mu_{\text{GFF}}^{\Omega, D} \otimes d\tau_{\Sigma}(\mu_{\text{GFF}}^{\check{\Omega}})(\phi_{\Omega}^D, \varphi) \\ &= \det_{\zeta}(\Delta_{\check{\Omega}} + m^2)^{-\frac{1}{2}} \mathbb{E}_{\text{GFF}}^{\check{\Omega}}[e^{-S_{\Omega}(\phi)}] = \text{RHS}. \end{aligned}$$

We arrive at the proof.  $\square$

**Corollary 6.8** (dissection gluing for  $P(\phi)$  field). *In the same situation as corollary 6.5, we have*

$$\int \mathcal{A}_{\Omega_1}^P(\varphi) \mathcal{A}_{\Omega_2}^P(\varphi) d\mu_{2\mathbf{D}}^{\Sigma}(\varphi) \det_{\zeta}(2\mathbf{D}_{\Sigma})^{-\frac{1}{2}} = Z_M. \quad (6.30)$$

*Proof.* One verifies that for  $\Omega = \Omega_1 \sqcup \Omega_2$ ,

$$\mathcal{A}_{\Omega}^P(\varphi, \varphi) = \mathcal{A}_{\Omega_1}^P(\varphi) \mathcal{A}_{\Omega_2}^P(\varphi). \quad (6.31)$$

This in fact also comes directly from the definition as

$$S_{|\Omega|^2}(\phi_{\text{GFF}}^{|\Omega|^2}) = S_{|\Omega_1|^2}(\phi_{\text{GFF}}^{|\Omega_1|^2}) + S_{|\Omega_2|^2}(\phi_{\text{GFF}}^{|\Omega_2|^2}) \quad (6.32)$$

pointwise almost surely with  $\phi_{\text{GFF}}^{|\Omega|^2} = \phi_{\text{GFF}}^{|\Omega_1|^2} + \phi_{\text{GFF}}^{|\Omega_2|^2}$ . One then follows a verbatim reasoning as for corollary 6.6.  $\square$




Figure 2: two ways of taking trace

## 6.5 Bayes-Type Density Formulae and Gluing with Free Ends

In fact let us first put ourselves in the general situation of (iii) of definition 6.1. Suppose the Riemannian surface  $\Omega$  is such that  $\partial\Omega = \Sigma_2 \sqcup \Sigma_2^* \sqcup \Sigma_1$  and that  $\rho : \Sigma_2 \rightarrow \Sigma_2^*$  is an orientation reversing isometry. Denote by  $\Omega/\rho$  the surface obtained from  $\Omega$  by gluing  $\Sigma_2$  with  $\Sigma_2^*$  along  $\rho$ . Denote by  $\Sigma_4$  the reflected copy of  $\Sigma_2$  in  $|\Omega/\rho|^2$ .

*Notation.* In general, for  $M$  a closed manifold and  $\Sigma_1, \dots, \Sigma_\ell \subset M$  finitely many nonintersecting closed embedded hypersurfaces, we denote

$$\begin{aligned} \tau_{i \sqcup j \sqcup \dots} &\stackrel{\text{def}}{=} \text{the trace map } C^\infty(M) \rightarrow C^\infty(\Sigma_i) \times C^\infty(\Sigma_j) \times \dots, \text{ and its extensions,} \\ \text{PI}_M^{i \sqcup j \sqcup \dots} &\stackrel{\text{def}}{=} \text{the Poisson integral operator } W^{\frac{1}{2}}(\Sigma_i) \times W^{\frac{1}{2}}(\Sigma_j) \times \dots \rightarrow W^1(M), \\ \mathcal{M}_{M, k \sqcup \dots}^{i \sqcup j \sqcup \dots} &\stackrel{\text{def}}{=} \tau_{k \sqcup \dots} \text{PI}_M^{i \sqcup j \sqcup \dots}, \text{ the transition operator } \mathcal{D}'(\Sigma_i) \times \mathcal{D}'(\Sigma_j) \times \dots \rightarrow \mathcal{D}'(\Sigma_k) \times \dots \text{ as in (5.22).} \end{aligned}$$

We also remind the reader of the notations set up in definitions 4.7 and 4.4.

**Proposition 6.9** (Bayes with free ends). *We have*

$$\frac{d\tau_1(\mu_{\text{GFF}}^{|\Omega/\rho|^2})}{d\mu_{2\mathbf{D}}^{\Sigma_1}}(\varphi_1) \left( \frac{d((\mathcal{M}_{|\Omega/\rho|^2, 2}^1 \varphi_1)^* \mu_{\text{DN}}^{\Sigma_2, \Omega/\rho, D})}{d\mu_{2\mathbf{D}}^{\Sigma_2}}(x) \right)^2 = \frac{\det_\zeta(\text{DN}_{|\Omega/\rho|^2}^{\Sigma_2 \sqcup \Sigma_4})^{\frac{1}{2}}}{\det_\zeta(\text{DN}_{|\Omega|^2}^{\Sigma_2 \sqcup \Sigma_2^*})^{\frac{1}{2}}} \frac{d\tau_{2 \sqcup 2^* \sqcup 1}(\mu_{\text{GFF}}^{|\Omega|^2})}{d\mu_{2\mathbf{D}}^{2 \sqcup 2^* \sqcup 1}}(x, x, \varphi_1) \quad (6.33)$$

*Proof.* Indeed, by reflection symmetry across  $\Sigma_1$  where  $\Sigma_2$  gets identified with  $\Sigma_4$ , we have

$$\mathcal{M}_{|\Omega/\rho|^2, 2}^1 \varphi_1 = \mathcal{M}_{|\Omega/\rho|^2, 4}^1 \varphi_1, \quad \text{and} \quad \mu_{\text{DN}}^{\Sigma_2, \Omega/\rho, D} = \mu_{\text{DN}}^{\Sigma_4, \Omega/\rho, D}, \quad (6.34)$$

under the symmetry. Also by symmetry (method of images) , we have

$$\mathcal{M}_{|\Omega/\rho|^2,1}^{2\sqcup 4} \begin{bmatrix} x \\ x \end{bmatrix} = \mathcal{M}_{|\Omega|^2,1}^{2\sqcup 2^*} \begin{bmatrix} x \\ x \end{bmatrix}, \quad (6.35)$$

because both of them expresses the associated Dirichlet data on  $\Sigma_1$  of the solution of the Helmholtz boundary value problem over  $\Omega$  with Dirichlet data equal to  $x$  on both  $\Sigma_2$ ,  $\Sigma_2^*$  and Neumann condition (zero normal derivative) on  $\Sigma_1$ . Thus,

$$\begin{aligned} \text{LHS} &= \frac{d\tau_{2\sqcup 4\sqcup 1}(\mu_{\text{GFF}}^{|\Omega/\rho|^2})}{d\mu_{2\mathbf{D}}^{2\sqcup 4\sqcup 1}}(x, x, \varphi_1) && \text{(formula (5.25) backward)} \\ &= \frac{d\tau_{2\sqcup 4}(\mu_{\text{GFF}}^{|\Omega/\rho|^2})}{d\mu_{2\mathbf{D}}^{\Sigma_2\sqcup \Sigma_4}}(x, x) \frac{d((\mathcal{M}_{|\Omega/\rho|^2,1}^{2\sqcup 4}) * \mu_{2\text{DN}}^{\Sigma_1, \Omega, D})}{d\mu_{2\mathbf{D}}^{\Sigma_1}}(\varphi_1) && \text{(formula (5.26) forward)} \\ &= \frac{\det_{\zeta}(\text{DN}_{|\Omega/\rho|^2}^{\Sigma_2\sqcup \Sigma_4})^{\frac{1}{2}}}{\det_{\zeta}(\text{DN}_{|\Omega|^2}^{\Sigma_2\sqcup \Sigma_2^*})^{\frac{1}{2}}} \frac{d\tau_{2\sqcup 2^*}(\mu_{\text{GFF}}^{|\Omega|^2})}{d\mu_{2\mathbf{D}}^{\Sigma_2\sqcup \Sigma_2^*}}(x, x) \frac{d((\mathcal{M}_{|\Omega|^2,1}^{2\sqcup 2^*}) * \mu_{2\text{DN}}^{\Sigma_1, \Omega, D})}{d\mu_{2\mathbf{D}}^{\Sigma_1}}(\varphi_1) && (\#) \\ &= \text{RHS}. \end{aligned}$$

We see that with the help of Bayes principle we “glued away” the free end and we are reduced to the situation of proposition 6.3. Indeed, at step (#) we use proposition 6.3 twice with respect to  $|\Omega \cup_1 \Omega^*|^2$  which is  $\Omega$  “reflected twice” by first gluing  $\Omega$  with a reflected copy along  $\Sigma_1$  to get  $\Omega \cup_1 \Omega^*$ , and then reflect and glue again along  $\Sigma_2 \sqcup \Sigma_2^* \sqcup \Sigma_4 \sqcup \Sigma_4^*$ , with  $\Sigma_4$ ,  $\Sigma_4^*$  being the reflected copies of  $\Sigma_2$ ,  $\Sigma_2^*$  across  $\Sigma_1$ . From this we get

$$\begin{aligned} \frac{d\tau_{2\sqcup 4}(\mu_{\text{GFF}}^{|\Omega/\rho|^2})}{d\mu_{2\mathbf{D}}^{\Sigma_2\sqcup \Sigma_4}}(x, x) &= \frac{\det_{\zeta}(\text{DN}_{|\Omega/\rho|^2}^{\Sigma_2\sqcup \Sigma_4})^{\frac{1}{2}}}{\det_{\zeta}(2\text{DN}_{|\Omega \cup_1 \Omega^*|^2}^{2\sqcup 2^* \sqcup 4\sqcup 4^*})^{\frac{1}{4}}} \left( \frac{d\tau_{2\sqcup 2^* \sqcup 4\sqcup 4^*}(\mu_{\text{GFF}}^{|\Omega \cup_1 \Omega^*|^2})}{d\mu_{2\mathbf{D}}^{2\sqcup 2^* \sqcup 4\sqcup 4^*}}(x, x, x, x) \right)^{\frac{1}{2}} \\ &= \frac{\det_{\zeta}(\text{DN}_{|\Omega/\rho|^2}^{\Sigma_2\sqcup \Sigma_4})^{\frac{1}{2}}}{\det_{\zeta}(\text{DN}_{|\Omega|^2}^{\Sigma_2\sqcup \Sigma_2^*})^{\frac{1}{2}}} \frac{d\tau_{2\sqcup 2^*}(\mu_{\text{GFF}}^{|\Omega|^2})}{d\mu_{2\mathbf{D}}^{\Sigma_2\sqcup \Sigma_2^*}}(x, x), \end{aligned}$$

finishing the proof.  $\square$

**Corollary 6.10** (free-end gluing for free field). *We have*

$$\int \mathcal{A}_{\Omega}^0(\varphi, \varphi, \psi) d\mu_{2\mathbf{D}}^{\Sigma_2}(\varphi) \det_{\zeta}(2\mathbf{D}_{\Sigma_2})^{-\frac{1}{2}} = \mathcal{A}_{\Omega/\rho}^0(\psi), \quad (6.36)$$

for almost every  $\psi \sim \mu_{2\mathbf{D}}^{\Sigma_1}$ .

*Proof.* The key point is to disintegrate  $\mathcal{A}_{\Omega}^0(\varphi, \varphi, \psi)$  into a part involving only  $\varphi$ , which one could “integrate out” cleanly, multiplied by a part independent of  $\varphi$ , using proposition 6.9. Indeed,

$$\begin{aligned} \text{LHS} &= \frac{\det_{\zeta}(\Delta_{|\Omega|^2} + m^2)^{-\frac{1}{4}}}{\det_{\zeta}(2\mathbf{D}_{\Sigma_2})^{-\frac{1}{2}} \det_{\zeta}(2\mathbf{D}_{\Sigma_1})^{-\frac{1}{4}}} \int \left( \frac{d\tau_{2\sqcup 2^* \sqcup 1}(\mu_{\text{GFF}}^{|\Omega|^2})}{d\mu_{2\mathbf{D}}^{2\sqcup 2^* \sqcup 1}}(\varphi, \varphi, \psi) \right)^{\frac{1}{2}} d\mu_{2\mathbf{D}}^{\Sigma_2}(\varphi) \det_{\zeta}(2\mathbf{D}_{\Sigma_2})^{-\frac{1}{2}} \\ &= \frac{\det_{\zeta}(\text{DN}_{|\Omega|^2}^{\Sigma_2\sqcup \Sigma_2^*})^{\frac{1}{4}} \det_{\zeta}(\Delta_{|\Omega|^2} + m^2)^{-\frac{1}{4}}}{\det_{\zeta}(\text{DN}_{|\Omega/\rho|^2}^{\Sigma_2\sqcup \Sigma_4})^{\frac{1}{4}} \det_{\zeta}(2\mathbf{D}_{\Sigma_1})^{-\frac{1}{4}}} \left( \frac{d\tau_1(\mu_{\text{GFF}}^{|\Omega/\rho|^2})}{d\mu_{2\mathbf{D}}^{\Sigma_1}}(\psi) \right)^{\frac{1}{2}} \underbrace{\int \frac{d((\mathcal{M}_{|\Omega/\rho|^2,2}^1) * \mu_{\text{DN}}^{\Sigma_2, \Omega/\rho, D})}{d\mu_{2\mathbf{D}}^{\Sigma_2}}(\varphi) d\mu_{2\mathbf{D}}^{\Sigma_2}(\varphi)}_{=1} \\ &= \det_{\zeta}(\text{DN}_{|\Omega/\rho|^2}^{\Sigma_2\sqcup \Sigma_4})^{-\frac{1}{4}} \det_{\zeta}(\Delta_{|\Omega|^2 \setminus \Sigma_2\sqcup \Sigma_2^*, D} + m^2)^{-\frac{1}{4}} \det_{\zeta}(2\mathbf{D}_{\Sigma_1})^{\frac{1}{4}} \left( \frac{d\tau_1(\mu_{\text{GFF}}^{|\Omega/\rho|^2})}{d\mu_{2\mathbf{D}}^{\Sigma_1}}(\psi) \right)^{\frac{1}{2}} \\ &= \text{RHS}, && \text{(BFK for } |\Omega|^2 \text{ and } \Sigma_2 \sqcup \Sigma_2^*) \\ & && \text{(BFK for } |\Omega/\rho|^2 \text{ and } \Sigma_2 \sqcup \Sigma_4) \end{aligned}$$

finishing the proof.  $\square$

Before proceeding to the interacting case we remind the reader of remark 5.11.

**Lemma 6.11.** *For fixed  $\psi \in W^{\frac{1}{2}}(\Sigma_1)$ , then the random fields*

$$\phi \stackrel{\text{def}}{=} \text{PI}_{\Omega/\rho}^1 \psi + \text{PI}_{\Omega/\rho}^{2,D} \tau_2 \phi_{\Omega/\rho}^D, \quad \text{and} \quad \tilde{\phi} \stackrel{\text{def}}{=} \text{PI}_{\Omega}^{2\sqcup 2^* \sqcup 1}[\varphi, \varphi, \psi] \quad (6.37)$$

with  $\phi_{\Omega/\rho}^D \sim \mu_{\text{GFF}}^{\Omega/\rho, D}$  and  $\varphi \sim (\mathcal{M}_{|\Omega/\rho|^2,2}^1 \psi) * \mu_{\text{DN}}^{\Sigma_2, \Omega/\rho, D}$ , follow the same law on  $\mathcal{D}'(\Omega^\circ)$ .

*Proof.* Here  $\Omega^\circ$  is also  $(\Omega/\rho) \setminus \Sigma_2$ . Indeed, both  $\phi$  and  $\tilde{\phi}$  solves the stochastic boundary value problem

$$\begin{cases} (\Delta + m^2)u = 0 & \text{in } \Omega^\circ, \\ u|_{\Sigma_2} = u|_{\Sigma_2^*} \sim (\mathcal{M}_{|\Omega/\rho|^2, 2}^1 \psi)_* \mu_{\text{DN}}^{\Sigma_2, \Omega/\rho, D}, & \text{on } \Sigma_2 \text{ or } \Sigma_2 \sqcup \Sigma_2^* \\ u|_{\Sigma_1} = \psi, & \text{on } \Sigma_1, \end{cases} \quad (6.38)$$

with equalities holding almost surely.  $\square$

**Corollary 6.12** (free-end gluing for  $P(\phi)$  field). *In the situation as above, we have*

$$\int \mathcal{A}_\Omega^P(\varphi, \varphi, \psi) d\mu_{2\mathbf{D}}^{\Sigma_2}(\varphi) \det_\zeta(2\mathbf{D}_{\Sigma_2})^{-\frac{1}{2}} = \mathcal{A}_{\Omega/\rho}^P(\psi), \quad (6.39)$$

for almost every  $\psi \sim \mu_{2\mathbf{D}}^{\Sigma_1}$ .

*Proof.* Indeed,

$$\mathcal{A}_\Omega^P(\varphi, \varphi, \psi) = \mathcal{A}_\Omega^0(\varphi, \varphi, \psi) \int e^{-S_\Omega(\phi_\Omega^D | \varphi, \varphi, \psi)} d\mu_{\text{GFF}}^{\Omega, D}(\phi_\Omega^D), \quad (6.40)$$

therefore, with the constants involving determinants working out in exactly the same way as corollary 6.10, one has

$$\begin{aligned} \text{LHS} &= \frac{\det_\zeta(\Delta_{|\Omega/\rho|^2} + m^2)^{-\frac{1}{4}}}{\det_\zeta(2\mathbf{D}_{\Sigma_1})^{\frac{1}{4}}} \left( \frac{d\tau_1(\mu_{\text{GFF}}^{|\Omega/\rho|^2})}{d\mu_{2\mathbf{D}}^{\Sigma_1}}(\psi) \right)^{\frac{1}{2}} \int d((\mathcal{M}_{|\Omega/\rho|^2, 2}^1 \psi)_* \mu_{\text{DN}}^{\Sigma_2, \Omega/\rho, D})(\varphi) \int e^{-S_\Omega(\phi_\Omega^D | \varphi, \varphi, \psi)} d\mu_{\text{GFF}}^{\Omega, D}(\phi_\Omega^D) \\ &= \mathcal{A}_{\Omega/\rho}^0(\psi) \int e^{-S_{\Omega/\rho}(\phi_{\Omega/\rho}^D + (\phi_{\Omega/\rho}^D)_{\Sigma_2} | \psi)} d\mu_{\text{GFF}}^{\Omega, D} \otimes d[(\text{PI}_{\Omega/\rho}^{2, D} \tau_2)_*(\mu_{\text{GFF}}^{\Omega/\rho, D})](\phi_\Omega^D, (\phi_{\Omega/\rho}^D)_{\Sigma_2}) \quad (\text{lemma 6.11}) \\ &= \mathcal{A}_{\Omega/\rho}^0(\psi) \int e^{-S_{\Omega/\rho}(\phi_{\Omega/\rho}^D | \psi)} d\mu_{\text{GFF}}^{\Omega/\rho, D}(\phi_{\Omega/\rho}^D) = \text{RHS}, \end{aligned}$$

where the notation  $(\phi_{\Omega/\rho}^D)_{\Sigma_2}$  is as in proposition 5.8. We arrive at the proof.  $\square$

Now let  $\Omega_1 \in \text{Mor}(\Sigma_1, \Sigma_2)$ ,  $\Omega_2 \in \text{Mor}(\Sigma_2, \Sigma_3)$  and  $|\Omega_2\Omega_1|^2 := (\Omega_1^* \cup_4 \Omega_2^* \cup_3 \Omega_2 \cup_2 \Omega_1)^\vee$ , where we denote by  $\Sigma_4$  the ‘‘glued’’ outgoing and incoming boundaries of  $\Omega_2^*$  and  $\Omega_1^*$ . In this case the result could be seen as a special case of the previous one, where  $\Omega = \Omega_1 \sqcup \Omega_2$  has two disjoint components. We shall re-state these results without proofs.

**Corollary 6.13.** *We have*

$$\begin{aligned} & \frac{d\tau_{1\sqcup 3}(\mu_{\text{GFF}}^{|\Omega_2\Omega_1|^2})}{d\mu_{2\mathbf{D}}^{\Sigma_1 \sqcup \Sigma_3}}(\varphi_1, \varphi_3) \left( \frac{d((\mathcal{M}_{|\Omega_2\Omega_1|^2, 2}^{1\sqcup 3} [\varphi_1])_* \mu_{2\text{DN}}^{\Sigma_2, \Omega_2\Omega_1, D})(x)}{d\mu_{2\mathbf{D}}^{\Sigma_2}}(x) \right)^2 \\ &= \frac{\det_\zeta(\text{DN}_{|\Omega_2\Omega_1|^2}^{\Sigma_2 \sqcup \Sigma_4})^{\frac{1}{2}}}{\det_\zeta(2\text{DN}_{\Omega_1}^{\Sigma_2, N})^{\frac{1}{2}} \det_\zeta(2\text{DN}_{\Omega_2}^{\Sigma_2, N})^{\frac{1}{2}}} \frac{d\tau_{\Sigma_1 \sqcup \Sigma_2}(\mu_{\text{GFF}}^{|\Omega_1|^2})(\varphi_1, x)}{d\mu_{2\mathbf{D}}^{\Sigma_1 \sqcup \Sigma_2}} \frac{d\tau_{\Sigma_2 \sqcup \Sigma_3}(\mu_{\text{GFF}}^{|\Omega_2|^2})(x, \varphi_3)}{d\mu_{2\mathbf{D}}^{\Sigma_2 \sqcup \Sigma_3}}. \quad \square \end{aligned}$$

**Corollary 6.14** (composition for  $P(\phi)$  field). *In the situation as above, we have*

$$\int \mathcal{A}_{\Omega_1}^P(\psi_1, \varphi) \mathcal{A}_{\Omega_2}^P(\varphi, \psi_3) d\mu_{2\mathbf{D}}^{\Sigma_2}(\varphi) \det_\zeta(2\mathbf{D}_{\Sigma_2})^{-\frac{1}{2}} = \mathcal{A}_{\Omega_2 \cup_2 \Omega_1}^P(\psi_1, \psi_3), \quad (6.41)$$

for almost every  $(\psi_1, \psi_3) \sim \mu_{2\mathbf{D}}^{\Sigma_1 \sqcup \Sigma_3}$ .  $\square$

## 7 Periodic Cover = Spin Chain

### 7.1 Geometric setting

Our set-up corresponds to example 1 in Bergeron [8]. Let  $M$  be a closed oriented Riemannian surface<sup>6</sup> of genus  $g \geq 1$  and  $\Sigma \subset M$  an embedded *primitive* closed geodesic whose  $\mathbb{Z}$ -homology class is non-trivial (exists by a classical theorem of E. Cartan). Necessarily,  $\Sigma$  is non-dissecting.<sup>7</sup> We consider the covering space  $M_\infty^\Sigma \rightarrow M$  over  $M$  given by a (normal) subgroup  $\ker \rho$  of the fundamental group  $\pi_1(M)$  where  $\rho$  is the map

$$\rho : \pi_1(M) \xrightarrow{\text{Ab}} H_1(M; \mathbb{Z}) \xrightarrow{I(-, [\Sigma])} \mathbb{Z}, \quad (7.1)$$

where the first is Abelianization and  $I(-, [\Sigma])$  is the oriented intersection number<sup>8</sup> with  $\Sigma$ , which is surjective (since  $\Sigma$  is primitive). In other words, we put  $M_\infty^\Sigma = \ker \rho \backslash \widetilde{M}$  where  $\widetilde{M}$  is the universal cover of  $M$  and  $\ker \rho$  acts on  $\widetilde{M}$  as deck transformations. Equip  $M_\infty^\Sigma$  with the covering metric (thus deck transformations act by isometries).

Geometrically,  $M_\infty^\Sigma$  can be understood as first cutting  $M$  along  $\Sigma$  and obtaining the surface  $\Omega := M \setminus \Sigma$  with boundaries  $\Sigma_{\text{in}} \sqcup \Sigma_{\text{out}}$  where  $\Sigma_{\text{in}} \cong \Sigma_{\text{out}} \cong \Sigma$ , and gluing  $\Omega$  periodically where each  $\Sigma_{\text{out}}$  is glued to the “next”  $\Sigma_{\text{in}}$ . Indeed, the class  $[\gamma] \in \pi_1(M)$  of a loop  $\gamma$  is in  $\ker \rho$  iff  $I([\gamma], [\Sigma]) = 0$ ; in other words, these loops are exactly those which are “not cut”, i.e. lifts to a loop on  $M_\infty^\Sigma$ , and loops which do intersect  $\Sigma$  are lifted to segments whose end points are related by a deck transformation, i.e. they are “cut”.

Now, for  $N \in \mathbb{N}$ , compose  $\rho$  further with the mod  $N$  map  $\mathbb{Z} \rightarrow \mathbb{Z}_N =: \mathbb{Z}/N\mathbb{Z}$  and denote it by  $\rho_N$ , and let the covering space of  $M$  corresponding to  $\ker \rho_N$  be  $M_N^\Sigma$ . Since  $\ker \rho \subset \ker \rho_N$ ,  $M_\infty^\Sigma$  also covers  $M_N^\Sigma$ . Geometrically, this corresponds to closing the surface after gluing  $N$  copies of  $\Omega$  — loops that intersect  $\Sigma$   $N$ -times are now lifted to a “big loop” in  $M_N^\Sigma$ .

**Remark 7.1.** We also say that the sequence of covers  $(M_N^\Sigma)_N$  converges to  $M_\infty^\Sigma$ .

That  $\Omega := M \setminus \Sigma$  should be understood for what follows.

### 7.2 Continued Introduction of Spin Chain Example

Here we continue our discussion of the circular spin chain proposed in section 1.1, in particular the equation (1.6). More generally, one can insert “nice” functionals  $F_1, \dots, F_k$  in between at the sites  $1 \leq i_1 < \dots < i_k \leq N$ , then

$$\int_{\mathbb{R}^N} F_k(\sigma(i_k)) \cdots F_1(\sigma(i_1)) e^{-S(\sigma)} d^N \sigma = \text{tr}_{L^2(\mathbb{R})} (T^{N+1-i_k} F_k T^{i_k-i_{k-1}} \cdots F_1 T^{i_1-1}), \quad (7.2)$$

where  $F_j$  are thought of as multiplication operators. The evaluation

$$\begin{aligned} F_k \otimes \cdots \otimes F_1 &\longmapsto \frac{1}{\mathcal{Z}(N)} \int_{\mathbb{R}^N} F_k(\sigma(i_k)) \cdots F_1(\sigma(i_1)) e^{-S(\sigma)} d^N \sigma \\ &= \frac{\text{tr}_{L^2(\mathbb{R})} (T^{N+1-i_k} F_k T^{i_k-i_{k-1}} \cdots F_1 T^{i_1-1})}{\text{tr}_{L^2(\mathbb{R})} (T^N)} \end{aligned} \quad (7.3)$$

is said to define a Gibbs state of our spin chain on  $\mathbb{Z}_N$ . Alternatively this is the expected value functional under the discrete Gibbs measure (1.2).

**Definition 7.1.** We say that a Gibbs state exists in the thermodynamic limit if the second expression in (7.3) tends to a limit as  $N \rightarrow \infty$  for bounded functionals  $F_1, \dots, F_k \in L^\infty(\mathbb{R})$ .

We can see from (1.4) that the operator  $T$  is not just bounded, smoothing, but its kernel is strictly positive. Such an operator has a special property which we call the Perron-Frobenius property, referring to the consequence of proposition 7.1 below. In particular, this would ensure that the thermodynamic limit does exist, as we shall see in corollaries 7.3, 7.4, and 7.5.

<sup>6</sup>In particular its  $\mathbb{Z}$ -homology groups are non-torsion.

<sup>7</sup>If  $\Sigma$  dissects  $M$  into  $M_+^\circ \sqcup M_-^\circ$ , then  $\Sigma = \partial M_+$ ; now closed 1-forms integrate to zero over  $\Sigma$  by Stokes theorem, thus  $\Sigma$  is null-homologous via de Rham’s theorem. Alternatively, note  $\partial : H_2(M_+, \Sigma; \mathbb{Z}) \rightarrow H_1(\Sigma; \mathbb{Z})$  in the exact sequence of the pair  $(M_+, \Sigma)$  is surjective, producing the fundamental class.

<sup>8</sup> $I([\gamma], [\Sigma]) = D(D^{-1}[\gamma] \smile D^{-1}[\Sigma]) = \int_\gamma \eta_\Sigma \in \mathbb{Z}$ , where  $D^{-1}$  is the Poincaré dual map and  $\eta_\Sigma$  is a smooth bump 1-form supported in a tubular neighborhood of  $\Sigma$  such that  $\int_\Sigma \alpha = \int_M \alpha \wedge \eta_\Sigma$  for any 1-form  $\alpha$ .






Figure 3: periodic covering and cyclic covering

**Remark 7.2.** Corollary 7.3 could be equivalently understood as saying that for compactly supported observables  $(F, G)$ , and  $(\tau\sigma)(i) := \sigma(i + 1)$  the shift operator, then

$$\mathbb{E}[F(\tau^k\sigma)G(\sigma)] = \mathbb{E}[F]\mathbb{E}[G] + \mathcal{O}(\alpha^k)$$

with  $\alpha < 1$  the same as in corollary 7.3. Here the expected value should be thought of as coming from a Gibbs measure on the infinite path space  $\mathbb{R}^{\mathbb{Z}}$  over  $\mathbb{Z}$ . Indeed, by definition 7.1, this is exactly the vague limit of the finite dimensional (periodic) Gibbs measures over  $\mathbb{Z}_N$ . We say that with respect to this Gibbs measure the shift operator is exponentially mixing.

With transfer operator being Perron-Frobenius, the partition functions as in (1.2) also enjoy explicit asymptotics. We will explore a consequence in the case of periodic surfaces in the last section.

**Example 7.1.** In the case  $P(\sigma) = m^2\sigma^2$ , the spin chain is the discrete massive GFF. We have an exact formula for the partition function  $\mathcal{Z}(N) = \prod_{k=0}^{N-1} (1 + m^2 - \cos(2\pi \frac{k}{N}))^{-\frac{1}{2}}$  hence

$$\lim_{N \rightarrow +\infty} \frac{1}{N} \log(\mathcal{Z}(N)) = -\frac{1}{2} \int_0^1 \log(1 + m^2 - \cos(2\pi x)) dx.$$

### 7.3 Perron-Frobenius Property and Gibbs State

**Remark 7.3.** A large part of this section could as well be included in the functional analysis appendix under the title “properties of a Perron-Frobenius operator”. But we include them here as they form an integral part of the discussion of physical phenomena.

We remind the reader of remark 1.3.

**Definition 7.2.** An operator  $A$  on  $L^2(Q, \mu)$  of some measure space  $(Q, \mu)$  has strictly positive kernel if for any nonnegative  $F \in L^2(Q, \mu)$  such that  $\|F\|_{L^2} \neq 0$  we have  $AF > 0$  almost surely.

**Proposition 7.1** (Perron-Frobenius, [26] page 51). *If  $A$  on  $L^2(Q, \mu)$  has strictly positive kernel, and  $\lambda = \|A\|$  is an eigenvalue of  $A$ , then  $\lambda$  is simple, and the corresponding eigenvector can be chosen to be strictly positive almost surely.*  $\square$

By our definition of the amplitudes  $\mathcal{A}_\Omega$  as the square-root of the Radon-Nikodym density between two mutually absolutely continuous positive finite measures ( $e^{-S_\Omega}$  is almost surely positive since  $S_\Omega$  is a real-valued random variable, recall also remark 2.7 that the zeta-determinants are positive), we get immediately

**Lemma 7.2.** *For any traceable cobordism  $\Omega \in \text{Mor}(\Sigma, \Sigma)$ , the Segal transfer operator  $U_\Omega : L^2(\mathcal{D}'(\Sigma), \mu_{2\mathbf{D}}) \rightarrow L^2(\mathcal{D}'(\Sigma), \mu_{2\mathbf{D}})$  has strictly positive kernel.*  $\square$

We deduce that  $U_\Omega$  has a simple top eigenvalue  $\lambda_0 = \|U_\Omega\|$  spanned by a normalized, almost surely strictly positive eigenvector  $\Omega_0 \in L^2(\mathcal{D}'(\Sigma), \mu_{2\mathbf{D}})$ . Alternatively speaking  $U_\Omega$  has a spectral gap. We get

**Corollary 7.3.** *Denote  $\widehat{U}_\Omega := \lambda_0^{-1}U_\Omega$ , let  $\lambda_1$  be the eigenvalue of  $U_\Omega$  with next largest modulus, thus  $\lambda_0 > |\lambda_1|$ , and put  $\alpha := |\lambda_1|/\lambda_0 < 1$ . Then for any  $F, G \in L^2(\mathcal{D}'(\Sigma), \mu_{2\mathbf{D}})$ , we have*

$$|\langle F, \widehat{U}_\Omega^N G \rangle - \langle F, \Omega_0 \rangle \langle \Omega_0, G \rangle| \leq \alpha^N \|F\| \|G\|. \quad (7.4)$$

*Proof.* Note  $\langle F, \widehat{U}_\Omega^N G \rangle - \langle F, \Omega_0 \rangle \langle \Omega_0, G \rangle = \langle \Pi_0^\perp F, \widehat{U}_\Omega^N \Pi_0^\perp G \rangle$  where  $\Pi_0^\perp$  is the orthogonal projection onto the complement of  $\text{Span}\{\Omega_0\}$ , where  $\widehat{U}_\Omega$  has norm  $\alpha < 1$ .  $\square$

**Corollary 7.4.** *We have*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \log \text{tr}(U_\Omega^N) = \log \lambda_0. \quad (7.5)$$

*Proof.* Without loss of generality let  $N \geq 2$  so each  $U_\Omega^N$  is trace class. On one hand

$$\text{tr}(U_\Omega^N) \leq \|U_\Omega^2\|_{\text{tr}} \|U_\Omega^{N-2}\| \leq \|U_\Omega^2\|_{\text{tr}} \lambda_0^{N-2}. \quad (7.6)$$

On the other, we decompose  $\mathcal{H}_\Sigma = \text{Span}\{\Omega_0\} \oplus \text{Span}\{\Omega_0\}^\perp$  where

$$\begin{aligned} \text{tr}(U_\Omega^N) &= \lambda_0^N + \text{tr}(U_\Omega^N|_{\text{Span}\{\Omega_0\}^\perp}) \\ &\geq \lambda_0^N - \|U_\Omega^2|_{\text{Span}\{\Omega_0\}^\perp}\|_{\text{tr}} |\lambda_1|^{N-2} \\ &= \lambda_0^N (1 - C_1 \alpha^{N-2} \lambda_0^{-2}). \end{aligned}$$

This and (7.6) gives the result after taking  $N \rightarrow \infty$ .  $\square$

Corollary 7.3 and the proof of corollary 7.4 implies

**Corollary 7.5.** *For any bounded operator  $F \in \mathcal{L}(\mathcal{H}_\Sigma)$  we have*

$$\lim_{N \rightarrow \infty} \frac{\text{tr}(U_\Omega^{N-L} F)}{\text{tr}(U_\Omega^N)} = \frac{1}{\lambda_0^L} \langle \Omega_0, F \Omega_0 \rangle = \frac{\langle \Omega_0, F \Omega_0 \rangle}{\langle \Omega_0, U_\Omega^L \Omega_0 \rangle}. \quad (7.7)$$

*Proof.* Remember that  $N \gg L$ . Thus by  $\mathcal{H}_\Sigma = \text{Span}\{\Omega_0\} \oplus \text{Span}\{\Omega_0\}^\perp$  and (iv) of lemma A.3 we have

$$\lambda_0^{-N+L} \text{tr}(U_\Omega^{N-L} F) = \langle \Omega_0, \widehat{U}_\Omega^{N-L} F \Omega_0 \rangle + \text{tr}(\widehat{U}_\Omega^{N-L} F|_{\text{Span}\{\Omega_0\}^\perp}) \quad (7.8)$$

with

$$|\text{tr}(\widehat{U}_\Omega^{N-L} F|_{\text{Span}\{\Omega_0\}^\perp})| \leq \|F\| \|\widehat{U}_\Omega^{N-L}|_{\text{Span}\{\Omega_0\}^\perp}\| \rightarrow 0. \quad (7.9)$$

Since  $\langle \Omega_0, \widehat{U}_\Omega^{N-L} F \Omega_0 \rangle \rightarrow \langle \Omega_0, F \Omega_0 \rangle$  by corollary 7.3 and  $\text{tr}(U_\Omega^N) \asymp \lambda_0^N$  by corollary 7.4, we obtain the result.  $\square$

We arrive at the conclusion that, for functionals  $F_1, \dots, F_k \in L^\infty(\mathcal{D}'(\Sigma), \mu_{2\mathbf{D}})$  and integers  $1 \leq i_1 < \dots < i_k$ , the evaluation

$$\begin{aligned} F_k \otimes \dots \otimes F_1 &\longmapsto \lim_{N \rightarrow \infty} \frac{\text{tr}(U_\Omega^{N+1-i_k} F_k U_\Omega^{i_k-i_{k-1}} \dots F_1 U_\Omega^{i_1-1})}{\text{tr}(U_\Omega^N)} \\ &= \frac{\langle \Omega_0, F_k U_\Omega^{i_k-i_{k-1}} \dots F_1 U_\Omega^{i_1-1} \Omega_0 \rangle}{\langle \Omega_0, U_\Omega^{i_k-1} \Omega_0 \rangle} \end{aligned} \quad (7.10)$$

defines a Gibbs state in the thermodynamic limit on a  $\mathcal{D}'(\Sigma)$ -valued  $\mathbb{Z}$ -spin chain.

**Remark 7.4.** In fact, a valid functional  $F \in L^\infty(\mathcal{D}'(\Sigma), \mu_{2\mathbf{D}})$  could be given by

$$F(\psi) \stackrel{\text{def}}{=} \int \mathcal{A}_\Omega^0(\psi, \varphi) d\mu_{2\mathbf{D}}^\Sigma(\varphi) \int \widetilde{F}(\phi_\Omega^D, \psi, \varphi) e^{-S_\Omega(\phi_\Omega^D | \psi, \varphi)} d\mu_{\text{GFF}}^{\Omega, D}(\phi_\Omega^D), \quad (7.11)$$

for  $\widetilde{F} \in L^\infty(\mathcal{D}'(\Omega^\circ), \mu_{\text{GFF}}^{\Omega, D} \otimes \mu_{2\mathbf{D}}^{\Sigma \cup \Sigma})$ , with  $\mu_{2\mathbf{D}}^{\Sigma \cup \Sigma}$  considered as the induced measure on  $\mathcal{D}'(\Omega^\circ)$  via  $\text{PI}_\Omega^{\Sigma \cup \Sigma}$ . Hence the Gibbs state actually extends to the continuum  $M_\infty^\Sigma$ .

## 7.4 Asymptotic of Partition Function

For the technical reason of remark A.2 we shall assume the surface  $\Omega \in \text{Mor}(\Sigma, \Sigma)$  be reflection symmetric, which in simplest words means  $\Omega = \tilde{\Omega}^* \cup_{\Sigma'} \tilde{\Omega}$  for some  $\tilde{\Omega} \in \text{Mor}(\Sigma, \Sigma')$ , in the notations of lemma 6.1. In other words there is an isometric involution  $\Theta : \Omega \rightarrow \Omega$  whose fixed point set is exactly  $\Sigma'$ , and exchanges the two components of  $\partial\Omega$ . This is not much of a restriction. In this case  $U_\Omega = U_{\tilde{\Omega}^*} \circ U_{\tilde{\Omega}}$  and lemma A.7 applies, namely  $\text{tr}(U_\Omega) = \text{tr}_\rho(U_\Omega)$ , as well as for  $U_\Omega^N$ . From corollary 6.7 and corollary 7.4 we then deduce that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \log(Z_{M_N^\Sigma}) = \log \lambda_0. \quad (7.12)$$

This applies in particular to the free case where  $Z_{M_N^\Sigma} = \det_\zeta(\Delta_{M_N^\Sigma} + m^2)^{-\frac{1}{2}}$ .

**Remark 7.5.** There arises the interesting question of how  $\lambda_0$  (or  $\log \lambda_0$ ) would depend on the geometry of  $\Omega$ . Very crudely one would expect  $\log \lambda_0 \propto \text{vol}(\Omega)$ , since in our case the corresponding  $\tilde{\lambda}_0$  for  $N\Omega$  ( $N$  copies of  $\Omega$  glued) is just  $\lambda_0^N$ . A more precise formula in the general, non-periodic case seems desirable.

# A Functional Analysis

## A.1 Tensor Product, Fock Space, Ideals

While leaving details to Simon [70], Reed and Simon [57], and Janson [39], we make the following conventions and definitions. Let  $\mathcal{H}$  be a real Hilbert space,  $v_i, w_j \in \mathcal{H}$ . Hilbert spaces are assumed real in the sequel.

$$\mathfrak{S}_n \stackrel{\text{def}}{=} \text{symmetric group on } n \text{ letters}; \quad (\text{A.1})$$

$$v_1 \odot \cdots \odot v_n \stackrel{\text{def}}{=} \text{Sym}^n(v_1 \otimes \cdots \otimes v_n) \stackrel{\text{def}}{=} \sum_{\sigma \in \mathfrak{S}_n} v_{\sigma(1)} \otimes \cdots \otimes v_{\sigma(n)}, \quad (\text{A.2})$$

$$\langle v_1 \odot \cdots \odot v_n, w_1 \odot \cdots \odot w_n \rangle_{\mathcal{H}^{\odot n}} \stackrel{\text{def}}{=} \sum_{\sigma \in \mathfrak{S}_n} \prod_{i=1}^n \langle v_i, w_{\sigma(i)} \rangle_{\mathcal{H}}, \quad (\text{A.3})$$

$$\mathcal{H}^{\odot n} \stackrel{\text{def}}{=} \overline{\text{Sym}^n(\mathcal{H}^{\otimes n})} \text{ under (A.3)} \quad (\text{A.4})$$

$$\alpha \odot \beta \stackrel{\text{def}}{=} \text{Sym}^{n+m}(\alpha \otimes \beta), \text{ for } \alpha \in \mathcal{H}^{\odot n}, \beta \in \mathcal{H}^{\odot m}. \quad (\text{A.5})$$

**Definition A.1.** Let  $\mathcal{H}$  be a real Hilbert space. Then the Boson Fock space  $\Gamma(\mathcal{H})$  associated to  $\mathcal{H}$  is defined to be the Hilbert space direct sum

$$\Gamma(\mathcal{H}) \stackrel{\text{def}}{=} \bigoplus_{n=0}^{\infty} \mathcal{H}^{\odot n}, \quad (\text{A.6})$$

that is,

$$\Gamma(\mathcal{H}) \stackrel{\text{def}}{=} \left\{ (\alpha_0, \alpha_1, \alpha_2, \dots) \mid \alpha_i \in \mathcal{H}^{\odot i}, \sum_{i=0}^{\infty} \|\alpha_i\|_{\mathcal{H}^{\odot i}}^2 < \infty \right\}, \quad (\text{A.7})$$

where, as usual, we put  $\mathcal{H}^{\odot 0} = \mathcal{H}^{\otimes 0} = \mathbb{R}$ , and the unit length generator of this bottom space is called the vacuum state, denoted  $|0\rangle$  or  $\Omega_0$ .

**Proposition A.1** ([39] theorem 4.5, [71] corollary I.15). *Let  $\mathcal{H}, \mathcal{K}$  be Hilbert spaces and  $A : \mathcal{H} \rightarrow \mathcal{K}$  a bounded operator. Then, for each  $n \geq 0$ , there exists a unique bounded operator  $\Gamma_n(A) : \mathcal{H}^{\odot n} \rightarrow \mathcal{K}^{\odot n}$  such that*

$$\Gamma_n(A)(v_1 \odot \cdots \odot v_n) = Av_1 \odot \cdots \odot Av_n, \quad (\text{A.8})$$

for all  $v_1, \dots, v_n \in \mathcal{H}$ , and one has  $\|\Gamma_n(A)\| = \|A\|^n$ . Moreover, if  $A$  is a **contraction**, namely if  $\|A\| \leq 1$ , then the direct sum

$$\Gamma(A) \stackrel{\text{def}}{=} \bigoplus_{n=0}^{\infty} \Gamma_n(A) : \Gamma(\mathcal{H}) \longrightarrow \Gamma(\mathcal{K}), \quad (\text{A.9})$$

that is,  $\Gamma(A)(\dots, \alpha_k, \dots) := (\dots, \Gamma_k(A)\alpha_k, \dots)$ , is a bounded operator with norm 1.

**Proposition A.2** (\*-homomorphism, [39] page 45). *Let  $\mathcal{H}, \mathcal{K}, \mathcal{L}$  be Hilbert spaces and  $A : \mathcal{H} \rightarrow \mathcal{K}$ ,  $B : \mathcal{K} \rightarrow \mathcal{L}$  operators with  $\|A\|, \|B\| \leq 1$ . Then*

- (i)  $\Gamma(BA) = \Gamma(B)\Gamma(A)$  as operators  $\Gamma(\mathcal{H}) \rightarrow \Gamma(\mathcal{L})$ ;
- (ii)  $\Gamma(A^*) = \Gamma(A)^* : \Gamma(\mathcal{K}) \rightarrow \Gamma(\mathcal{H})$ ,  $(\bullet)^*$  being the adjoint;
- (iii)  $\Gamma(\mathbb{1}_{\mathcal{H}}) = \mathbb{1}_{\Gamma(\mathcal{H})}$  for any  $\mathcal{H}$ .

**Remark A.1.** The map  $\Gamma$  should not be confused with the second quantization  $d\Gamma$  explained on Simon [71] page 31.

Let  $A : \mathcal{H} \rightarrow \mathcal{K}$  be a compact operator and  $A^* : \mathcal{K} \rightarrow \mathcal{H}$  its adjoint. Then  $A^*A$  is a compact self-adjoint nonnegative operator whose spectrum consists of positive eigenvalues  $\mu_0(A)^2 \geq \mu_1(A)^2 \geq \dots \geq 0$  with the only possible accumulation point being zero, which may or may not be in the spectrum or an eigenvalue. The numbers  $\mu_n(A)$  are called **singular values** of  $A$ . Define

$$\|A\|_{\mathcal{J}_p} \stackrel{\text{def}}{=} \left( \sum_n \mu_n(A)^p \right)^{1/p}, \quad \text{for } 1 \leq p < \infty, \quad (\text{A.10})$$

$$\|A\|_{\mathcal{J}_\infty} \stackrel{\text{def}}{=} \|A\|, \quad \text{for } p = \infty. \quad (\text{A.11})$$

Put  $\mathcal{J}_p(\mathcal{H}, \mathcal{K}) := \{A \text{ compact} \mid \|A\|_{\mathcal{J}_p} < \infty\}$ . We shall be concerned only with  $\mathcal{J}_1$ , where the norm is also denoted  $\|\cdot\|_{\text{tr}}$ , called the trace class operators and  $\mathcal{J}_2$ , called the Hilbert-Schmidt operators, where the norm is also written  $\|\cdot\|_{\text{HS}}$ . When  $\mathcal{K} = \mathcal{H}$ , define also

$$\text{tr}_{\mathcal{H}}(A) \stackrel{\text{def}}{=} \sum_n \langle e_n, A e_n \rangle_{\mathcal{H}}, \quad (\text{A.12})$$

where  $\{e_n\}$  is an orthonormal basis of  $\mathcal{H}$ . Its convergence and (in)dependence on basis is discussed below. When  $\mathcal{H} = \mathcal{K}$  we use the notation  $\mathcal{J}_p(\mathcal{H})$ .

**Lemma A.3** ([70] theorems 2.7, 2.8, 2.14, 3.1, [26] page 132-133). *We have*

- (i) *whenever  $A : \mathcal{H} \rightarrow \mathcal{H}$  is bounded and self-adjoint nonnegative, the sum (A.12) valuing in  $[0, +\infty]$  is independent of bases; it is finite iff  $A \in \mathcal{J}_1(\mathcal{H})$  in which case  $\text{tr}_{\mathcal{H}}(A) = \|A\|_{\text{tr}}$ ;*
- (ii)  *$A \in \mathcal{J}_p(\mathcal{H}, \mathcal{K})$  iff  $(A^*A)^{p/2} \in \mathcal{J}_1(\mathcal{H})$ , in which case  $\|A\|_{\mathcal{J}_p} = \text{tr}_{\mathcal{H}}((A^*A)^{p/2})^{1/p}$ . In particular,  $A$  is Hilbert-Schmidt iff  $A^*A$  is trace class.*
- (iii) *Whenever  $A, C \in \mathcal{L}(\mathcal{H})$  and  $B \in \mathcal{J}_p(\mathcal{H})$  we have  $\|ABC\|_{\mathcal{J}_p} \leq \|A\| \|C\| \|B\|_{\mathcal{J}_p}$ ; thus each  $\mathcal{J}_p$  is a two-sided ideal in  $\mathcal{L}(\mathcal{H})$ ;*
- (iv) *if  $A \in \mathcal{J}_1(\mathcal{H})$  then the sum (A.12) converges absolutely, is independent of bases, and  $|\text{tr}_{\mathcal{H}}(A)| \leq \|A\|_{\text{tr}}$ ;*
- (v) *whenever  $p^{-1} = q^{-1} + r^{-1}$  and  $A \in \mathcal{J}_q(\mathcal{H})$ ,  $B \in \mathcal{J}_r(\mathcal{H})$  then  $\|AB\|_{\mathcal{J}_p} \leq \|A\|_{\mathcal{J}_q} \|B\|_{\mathcal{J}_r}$  and  $AB \in \mathcal{J}_p(\mathcal{H})$ ; in particular, the product of two Hilbert-Schmidt operators is trace class.*

## A.2 Hilbert-Schmidt Operators on $L^2$

**Proposition A.4** ([57] page 52). *Let  $(Q_1, \mu_1)$  and  $(Q_2, \mu_2)$  be measure spaces so that  $L^2(Q_1, \mu_1)$ ,  $L^2(Q_2, \mu_2)$  are separable. Then the map*

$$\begin{aligned} L^2(Q_1, \mu_1) \times L^2(Q_2, \mu_2) &\longrightarrow L^2(Q_1 \times Q_2, \mu_1 \otimes \mu_2), \\ (f, g) &\longmapsto fg, \end{aligned} \quad (\text{A.13})$$

*with  $(fg)(x, y) := f(x)g(y)$  extends to a unique isomorphism  $L^2(Q_1, \mu_1) \otimes L^2(Q_2, \mu_2) \cong L^2(Q_1 \times Q_2, \mu_1 \otimes \mu_2)$ .*

**Proposition A.5** ([57] page 220, [72] theorem 3.8.4). *Let  $\mathcal{H}, \mathcal{K}$  be Hilbert spaces. Then the map*

$$\begin{aligned} \mathcal{I} : \mathcal{H} \times \mathcal{K} &\longrightarrow \mathcal{J}_2(\mathcal{H}, \mathcal{K}), \\ (v, w) &\longmapsto \langle v, - \rangle_{\mathcal{H}} w \end{aligned} \quad (\text{A.14})$$

*extends to a unique isometric isomorphism  $\mathcal{H} \otimes \mathcal{K} \cong \mathcal{J}_2(\mathcal{H}, \mathcal{K})$ . In particular, for every  $\eta \in \mathcal{H} \otimes \mathcal{K}$  there exist orthonormal sets  $\{\varphi_n\} \subset \mathcal{H}$ ,  $\{\psi_n\} \subset \mathcal{K}$  and real numbers  $\lambda_n > 0$  so that*

$$\eta = \sum_n \lambda_n (\varphi_n \otimes \psi_n), \quad \text{and} \quad \mathcal{I}(\eta)(v) = \sum_n \lambda_n \langle \varphi_n, v \rangle_{\mathcal{H}} \psi_n, \quad (\text{A.15})$$

*and vice versa.*

**Corollary A.6** ([72] theorem 3.8.5). *Let  $(Q_1, \mu_1)$  and  $(Q_2, \mu_2)$  be measure spaces so that  $L^2(Q_1, \mu_1)$ ,  $L^2(Q_2, \mu_2)$  are separable. Then there is an isometric isomorphism*

$$\begin{aligned} \mathcal{I} : L^2(Q_1 \times Q_2, \mu_1 \otimes \mu_2) &\longrightarrow \mathcal{J}_2(L^2(Q_1, \mu_1), L^2(Q_2, \mu_2)), \\ K(x, y) &\longmapsto \left[ f \longmapsto \int K(x, -) f(x) d\mu_1(x) \right]. \end{aligned} \quad (\text{A.16})$$

*In particular, there exist orthonormal families  $\{\varphi_n\} \subset L^2(Q_1, \mu_1)$ ,  $\{\psi_n\} \subset L^2(Q_2, \mu_2)$  so that*

$$K(x, y) = \sum_n \mu_n(\mathcal{I}(K)) \overline{\varphi_n(x)} \psi_n(y), \quad (\text{A.17})$$

*with the latter series converging absolutely in  $L^2(Q_1, \mu_1)$  for almost every fixed  $y$ , and  $\mu_n(\mathcal{I}(K))$  denote the singular values of  $\mathcal{I}(K)$ .*

Now (ii) of lemma A.3 implies that

**Corollary A.7.** Let  $L^2(Q_1, \mu_1)$ ,  $L^2(Q_2, \mu_2)$  be as above and  $A : L^2(Q_1, \mu_1) \rightarrow L^2(Q_2, \mu_2)$  a Hilbert-Schmidt operator, represented via (A.16) by the integral kernel  $K_A(x, y)$ . Then

$$\|A\|_{\text{HS}}^2 = \text{tr}_{L^2(Q_1)}(A^*A) = \int |K_A(x, y)|^2 d\mu_1(x) d\mu_2(y). \quad \square \quad (\text{A.18})$$

**Remark A.2.** For general  $A \in \mathcal{L}(L^2(Q))$  or  $\mathcal{J}_2(L^2(Q))$ , the fact that  $\int |K_A(x, x)| d\mu(x) < \infty$  does not imply  $A$  is trace class, and nor does the trace equal  $\int K_A(x, x) d\mu(x)$  when  $A$  is trace class (since the diagonal has measure zero in  $Q \times Q$ , in reality one could let  $K_A(x, x)$  be arbitrary without affecting  $A$ , thus an important condition is that  $K_A$  be continuous in “some sense”). See Simon [72] section 3.11, also Vershik, Petrov and Zatitskiy [53] section 3.3.

## B Gaussian Analysis

### B.1 Basic Definitions and Spaces

Let  $\mathcal{X}$  be a real separable Fréchet space and  $\mathcal{X}^*$  the dual space of continuous linear functionals. Equip  $\mathcal{X}$  with the Borel  $\sigma$ -algebra  $\mathcal{B}_{\mathcal{X}}$ .

**Definition B.1.** A probability measure  $\gamma$  on  $(\mathcal{X}, \mathcal{B}_{\mathcal{X}})$  is called (centered) **Gaussian** if every linear functional  $f \in \mathcal{X}^*$ ,  $f : \mathcal{X} \rightarrow \mathbb{R}$ , is a (centered) Gaussian random variable on  $\mathcal{X}$ .

Random variables and measures are assumed centered in the sequel unless otherwise stated.

**Definition B.2.** A (general) **Gaussian Hilbert space** is any closed linear subspace of  $L^2(Q, \mathcal{O}, \mathbb{P})$  of any probability space  $(Q, \mathcal{O}, \mathbb{P})$ , consisting of (centered) Gaussian variables.

**Definition B.3.** Let  $\mathcal{X}$  be a real separable Fréchet (or Banach) space, and  $\gamma$  a Gaussian measure on  $(\mathcal{X}, \mathcal{B}_{\mathcal{X}})$ . Then the closure of  $\mathcal{X}^*$  in  $L^2(\mathcal{X}, \mathcal{B}_{\mathcal{X}}, \gamma)$  is called the **Gaussian Hilbert space** of  $\gamma$ , denoted  $\mathcal{H}_{\gamma}$ .

An important gadget for identifying Gaussian measures is the (inverse) Fourier transform, more frequently called the characteristic function,  $\hat{\gamma} : \mathcal{X}^* \rightarrow \mathbb{C}$ , defined as

$$\hat{\gamma}(f) \stackrel{\text{def}}{=} \int_{\mathcal{X}} e^{i\phi(f)} d\gamma(\phi). \quad (\text{B.1})$$

This is useful because of the following property.

**Lemma B.1** ([12] theorem 2.2.4, proposition A.3.18). *Let  $\mathcal{X}$  be a real separable Fréchet (or Banach) space, then any two measures on  $(\mathcal{X}, \mathcal{B}_{\mathcal{X}})$  with equal characteristic functions coincide. In particular, a probability measure  $\gamma$  on  $(\mathcal{X}, \mathcal{B}_{\mathcal{X}})$  is a centered Gaussian measure if and only if*

$$\hat{\gamma}(f) = e^{-\frac{1}{2}\mathbb{E}_{\gamma}[\phi(f)^2]} \quad (\text{B.2})$$

for all  $f \in \mathcal{X}^*$ , where  $\mathbb{E}_{\gamma}$  is the expectation with respect to  $\gamma$ .

**Remark B.1.** The variance  $\mathbb{E}_{\gamma}[\phi(f)^2]$  of  $\phi(f)$  appears in (B.2). Thus lemma B.1 implies that two (centered) Gaussian measures on  $(\mathcal{X}, \mathcal{B}_{\mathcal{X}})$  with the same covariance structure must coincide; that is, if

$$\mathbb{E}_{\gamma}[\phi(f)\phi(h)] = \mathbb{E}_{\tilde{\gamma}}[\phi(f)\phi(h)] \quad (\text{B.3})$$

for all  $f, h \in \mathcal{X}^*$ , then  $\gamma = \tilde{\gamma}$ .

Let  $(\mathcal{X}, \mathcal{B}_{\mathcal{X}})$  be equipped with a Gaussian measure  $\gamma$ . Then in  $\mathcal{X}$  sits an important subspace  $C(\mathcal{H}_{\gamma})$ , which is a Hilbert space with norm  $\|\cdot\|_{C(\mathcal{H}_{\gamma})}$ , called the **Cameron-Martin space**, that does the following job. For two measures  $\mu$  and  $\nu$ , we write “ $\mu \approx \nu$ ” to mean  $\mu$  and  $\nu$  are mutually absolutely continuous, and “ $\mu \perp \nu$ ” to mean  $\mu$  and  $\nu$  are mutually singular.

**Proposition B.2** (Cameron-Martin, [12] corollary 2.4.3). *Let  $\mathcal{X}$  be a real separable Fréchet space,  $\gamma$  a Gaussian measure on  $(\mathcal{X}, \mathcal{B}_{\mathcal{X}})$ , and  $C(\mathcal{H}_{\gamma}) \subset \mathcal{X}$  the Cameron-Martin space. For any  $\phi_0 \in \mathcal{X}$ , define the measure  $(\phi_0)_*\gamma$  on  $(\mathcal{X}, \mathcal{B}_{\mathcal{X}})$  by setting*

$$(\phi_0)_*\gamma(B) \stackrel{\text{def}}{=} \gamma(B - \phi_0), \quad (\text{B.4})$$

where  $B - \phi_0 = \{\phi - \phi_0 \mid \phi \in B\}$ . Then

$$\begin{cases} (\phi_0)_*\gamma \approx \gamma, & \text{if } \phi_0 \in C(\mathcal{H}_\gamma), \\ (\phi_0)_*\gamma \perp \gamma, & \text{if } \phi_0 \notin C(\mathcal{H}_\gamma). \end{cases} \quad (\text{B.5})$$

Moreover, in the first case, one has the Radon-Nikodym density

$$\frac{d(\phi_0)_*\gamma}{d\gamma}(\phi) = e^{(C^{-1}\phi_0)(\phi) - \frac{1}{2}\|\phi_0\|_{C(\mathcal{H}_\gamma)}^2}, \quad (\text{B.6})$$

where  $C$  is the operator defined in (B.7).

Now, how to locate the space  $C(\mathcal{H}_\gamma)$  in  $\mathcal{X}$ ? The answer is that we can first locate it in  $\mathcal{X}^{**}$  (the double dual), where each element of  $C(\mathcal{H}_\gamma)$  turns out to be weak\* continuous (Bogachev [12] page 362), and since  $\mathcal{X}$  is locally convex, it is actually in  $\mathcal{X}$ . Starting from the Gaussian Hilbert space  $\mathcal{H}_\gamma$  we define the covariance operator

$$\begin{aligned} C : \mathcal{H}_\gamma &\longrightarrow \mathcal{X}^{**}, \\ f &\longmapsto (C(f) : h \longmapsto \mathbb{E}_\gamma[\phi(f)\phi(h)]). \end{aligned} \quad (\text{B.7})$$

In other words  $C(f)$  is the linear functional  $\langle f, - \rangle_{L^2(\mathcal{X}, \gamma)}$  on  $\mathcal{X}^*$ .

**Lemma B.3** ([12] page 44, sections 2.4, 3.2). (i) the map  $C$  is injective.

(ii) Equip  $C(\mathcal{H}_\gamma)$  with an inner product by requiring  $C$  to be isometry, namely

$$\langle f, h \rangle_{C(\mathcal{H}_\gamma)} \stackrel{\text{def}}{=} \langle C^{-1}f, C^{-1}h \rangle_{\mathcal{H}_\gamma}. \quad (\text{B.8})$$

Then  $C(\mathcal{H}_\gamma)$  is a complete Hilbert space,  $C(\mathcal{H}_\gamma) \hookrightarrow \mathcal{X}$  is continuous, and each  $\phi \longmapsto \phi(f)$ ,  $f \in \mathcal{X}^*$ , is a continuous linear functional for  $\phi \in C(\mathcal{H}_\gamma)$ .

(iii) Equivalently, the norm  $\|\cdot\|_{C(\mathcal{H}_\gamma)}$  is

$$\|\phi\|_{C(\mathcal{H}_\gamma)} = \sup_{f \in \mathcal{X}^*, f \neq 0} \frac{|\phi(f)|}{\|f\|_{\mathcal{H}_\gamma}}. \quad (\text{B.9})$$

Thus  $C(\mathcal{H}_\gamma)$  is precisely the dual of  $\mathcal{H}_\gamma$  under the evaluation pairing  $(\phi, f) \longmapsto \phi(f)$ .  $\square$

## B.2 $Q$ -spaces and Their Equivalence

A  $Q$ -space is in other words a probability sample space. The GFF on  $M$ , say, can be seen (abstractly) as a way of associating a Gaussian random variable  $\phi(f)$  to each  $f \in W^{-1}(M)$  so that (2.6) holds. This says nothing about the sample space on which these random variables are actually defined, and naturally there exist many choices. For example,

- (i) the Bochner-Minlos construction, mentioned as proposition 2.1;
- (ii) the formal Fourier series construction, mentioned in remark 2.2; this can be identified with the previous one by appealing to the condition under which a formal Fourier series represents an actual distribution, see Shubin [68] page 92 proposition 10.2;
- (iii) the abstract Wiener space construction. While this is not used essentially in this paper, it is a way of constructing (recovering) a separable Banach  $Q$ -space  $\mathcal{X}$  starting from (knowing) the Cameron-Martin space, and taking closure with respect to a carefully defined norm weaker than  $\|\cdot\|_{C(\mathcal{H}_\gamma)}$ . See Sheffield [67] and Bogachev [12] section 3.9 for details.

Some other models are discussed in Simon [71] section I.2, to which we refer for details in general. We shall discuss the question of in what sense two models of  $Q$ -spaces are equivalent, though all the three examples above could eventually be realized in  $\mathcal{D}'(M)$ . This is useful concerning the decomposition (5.11), and we give the precise sense in which the original  $\mu_{\text{GFF}}^M$  could be recovered from the decomposed measure  $\mu_{\text{DN}}^{\Sigma, M} \otimes \mu_{\text{GFF}}^{M \setminus \Sigma, D}$ .

**Definition B.4** ([71] page 4). Two probability measure spaces  $(Q, \mathcal{O}, \mu)$  and  $(Q', \mathcal{O}', \mu')$  are called **isomorphic** if there is an isomorphism of measure algebras

$$T : \mathcal{O}/\mathcal{I}_\mu \xrightarrow{\sim} \mathcal{O}'/\mathcal{I}_{\mu'}, \quad (\text{B.10})$$

here  $\mathcal{I}_\mu, \mathcal{I}_{\mu'}$  being the ideals of measure zero sets of  $\mu$  and  $\mu'$ , such that  $\mu'(T(A)) = \mu(A)$  for all  $A \in \mathcal{O}/\mathcal{I}_\mu$  (we do not distinguish an event  $A$  from its class in  $\mathcal{O}/\mathcal{I}_\mu$ ).



**Remark B.2.** Mutually absolutely continuous measures  $\mu$  on  $\mathcal{O}$  will define the same measure algebra  $\mathcal{O}/\mathcal{I}_\mu$  as they have the same measure zero sets.

**Definition B.5** ([71] page 5). If  $(Q, \mathcal{O}, \mu)$  and  $(Q', \mathcal{O}', \mu')$  are isomorphic under  $T$ , then two random variables  $f : Q \rightarrow \mathbb{R}$  and  $f' : Q' \rightarrow \mathbb{R}$  correspond under the isomorphism if

$$T(f^{-1}(B)) = (f')^{-1}(B) \quad (\text{B.11})$$

for all Borel sets  $B \subset \mathbb{R}$ .

For a real Hilbert space  $\mathcal{H}$ , a (centered) Gaussian process indexed by  $\mathcal{H}$  is a family of centered Gaussian random variables  $\{\phi(f) \mid f \in \mathcal{H}\}$  so that

$$\mathbb{E}[\phi(f)\phi(h)] = \langle f, h \rangle_{\mathcal{H}} \quad (\text{B.12})$$

for all  $f, h \in \mathcal{H}$ . Such a process is defined on the probability space  $(Q, \mathcal{O}, \mu)$  if each  $\phi(f)$  is a random variable from  $Q$  and  $\{\phi(f) \mid f \in \mathcal{H}\}$  generates  $\mathcal{O}$ .

**Proposition B.4** ([71] theorem I.6). Let  $\{\phi(f)\}$  and  $\{\phi'(f)\}$  be two Gaussian processes indexed by  $\mathcal{H}$  defined respectively on  $(Q, \mathcal{O}, \mu)$  and  $(Q', \mathcal{O}', \mu')$ . Then there is an isomorphism between the two probability spaces so that  $\phi(f)$  corresponds to  $\phi'(f)$  under the isomorphism for each  $f \in \mathcal{H}$ .

*Remark for proof.* Note for probability spaces  $L^2(Q, \mathcal{O}, \mu) \supset L^\infty(Q, \mathcal{O}, \mu) \supset \{\text{indicators}\}$ . Similarly for  $(Q', \mathcal{O}', \mu')$ . Thus  $\Gamma(U)$  where  $U$  takes each  $\phi(f)$  to  $\phi'(f)$  gives the isomorphism (corollary B.10).  $\square$

**Proposition B.5** ([71] proposition I.7). Let  $\{\phi_1(f)\}$  and  $\{\phi_2(f)\}$  be Gaussian processes, respectively, indexed by  $\mathcal{H}_1$  and  $\mathcal{H}_2$ , defined on  $(Q_1, \mathcal{O}_1, \mu_1)$  and  $(Q_2, \mathcal{O}_2, \mu_2)$ . Then a Gaussian process  $\{\phi(h)\}$  indexed by  $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2$  can be defined on  $Q := Q_1 \times Q_2$  equipped with  $\mathcal{O}_1 \widehat{\otimes} \mathcal{O}_2$  and  $\mu_1 \otimes \mu_2$ , by putting

$$\phi(f_1 \oplus f_2) \stackrel{\text{def}}{=} \phi_1(f_1) + \phi_2(f_2), \quad (\text{B.13})$$

for all  $f_1 \in \mathcal{H}_1, f_2 \in \mathcal{H}_2$ .

## B.3 Measurable Linear (Itô-Wiener) Extensions

Here we collect some results concerning the possibility of extending a linear operator defined on the Cameron-Martin space  $C(\mathcal{H}_\gamma)$  to the whole  $Q$ -space  $\mathcal{X}$ , in a not necessarily “functional-analytic” sense. Such results, in a different guise, lie behind the possibility of defining the classical stochastic integrals à la Itô-Wiener.

**Definition B.6** ([12] definition 3.7.1). Let  $(\mathcal{X}, \mathcal{B}_\mathcal{X}, \gamma)$  be a Fréchet Gaussian  $Q$ -space and  $(\mathcal{Y}, \mathcal{B}_\mathcal{Y})$  another Fréchet space equipped with the Borel  $\sigma$ -algebra. Then a map  $F : \mathcal{X} \rightarrow \mathcal{Y}$  is called a  $\gamma$ -measurable linear operator if it agrees  $\gamma$ -almost surely with a  $(\mathcal{B}_\mathcal{X}, \mathcal{B}_\mathcal{Y})$ -measurable linear map  $F_0 : \mathcal{X} \rightarrow \mathcal{Y}$ .

In particular,

**Definition B.7** ([12] definition 2.10.1). Let  $(\mathcal{X}, \mathcal{B}_\mathcal{X}, \gamma)$  be a Fréchet Gaussian  $Q$ -space. A  $\gamma$ -measurable linear functional  $f$  on  $X$  is called a measurable linear functional if there exist a full-measure linear subspace  $\mathcal{X}_0 \subset \mathcal{X}$  such that  $f$  agrees  $\gamma$ -almost surely with a usual  $\gamma$ -measurable linear functional  $f_0$  on  $\mathcal{X}_0$ .

**Proposition B.6** ([12] theorem 2.10.11). Let  $(\mathcal{X}, \mathcal{B}_\mathcal{X}, \gamma)$  be a real separable Fréchet Gaussian  $Q$ -space. Denote by  $C(\mathcal{H}_\gamma)$  its Cameron-Martin space. Then every continuous linear functional  $f$  on  $C(\mathcal{H}_\gamma)$  extends uniquely (modulo measure zero sets) to a measurable linear functional  $\widehat{f}$  on  $\mathcal{X}$  that coincides with  $f$  on  $C(\mathcal{H}_\gamma)$ . Moreover,  $\|\widehat{f}\|_{L^2(\gamma)} = \|f\|_{C(\mathcal{H}_\gamma)^*}$ .

**Proposition B.7** ([12] theorem 3.7.6). Let  $(\mathcal{X}, \mathcal{B}_\mathcal{X}, \gamma)$  be a real separable Fréchet Gaussian  $Q$ -space. Denote by  $C(\mathcal{H}_\gamma)$  its Cameron-Martin space. Then every operator  $A \in \mathcal{L}(C(\mathcal{H}_\gamma))$  extends to a  $\gamma$ -measurable linear operator  $\widehat{A} : (\mathcal{X}, \overline{\mathcal{B}_\mathcal{X}^\gamma}) \rightarrow (\mathcal{X}, \mathcal{B}_\mathcal{X})$ , such that the measure image of  $\gamma$  under  $\widehat{A}$  is a Gaussian measure on  $(\mathcal{X}, \mathcal{B}_\mathcal{X})$ . Here  $\overline{\mathcal{B}_\mathcal{X}^\gamma}$  denotes the Lebesgue completion of  $\mathcal{B}_\mathcal{X}$  with respect to  $\gamma$ . Moreover, any two extensions of  $A$  which are  $(\overline{\mathcal{B}_\mathcal{X}^\gamma}, \mathcal{B}_\mathcal{X})$ -measurable and linear on full measure subspaces agree  $\gamma$ -almost surely.

The extension  $\widehat{A}$  obtained above will be called the measurable linear (or Itô-Wiener) extension of  $A$ .

## B.4 Wiener Chaos and Wick's Theorem

**Proposition B.8** (Wiener Chaos decomposition, [39] theorem 2.6). *Let  $(Q, \mathcal{O}, \mathbb{P})$  be a probability space and  $\mathcal{H} \subset L^2(Q, \mathcal{O}, \mathbb{P})$  a Gaussian Hilbert space. Then there is an orthogonal decomposition*

$$L^2(Q, \mathcal{O}(\mathcal{H}), \mathbb{P}) \cong \bigoplus_{n=0}^{\infty} \mathcal{H}^{:n:}, \quad (\text{B.14})$$

where  $\mathcal{O}(\mathcal{H})$  is the  $\sigma$ -algebra generated by variables in  $\mathcal{H}$ ,  $\mathcal{H}^{:n:} = \overline{\mathcal{P}_n(\mathcal{H})} \cap \overline{\mathcal{P}_{n-1}(\mathcal{H})}^\perp$ , where  $\mathcal{P}_j(\mathcal{H})$  denotes the span of polynomials of random variables in  $\mathcal{H}$  of degree  $\leq j$ ; in particular  $\mathcal{H}^{:0:}$  denotes the constants.

If  $F \in \overline{\mathcal{P}_n(\mathcal{H})}$ , denote by  $:F:$  the projection of  $F$  onto  $\mathcal{H}^{:n:}$ , and is called a Wick ordered polynomial; for  $F \in L^2(\Omega, \mathcal{O}(\mathcal{H}), \mathbb{P})$ , denote by  $I_n(F)$  its projection onto  $\mathcal{H}^{:n:}$ . Define the Hermite polynomials  $h_n(x)$  by

$$\exp\left(zx - \frac{1}{2}z^2\right) = \sum_{n=0}^{\infty} \frac{z^n}{n!} h_n(x). \quad (\text{B.15})$$

We have  $h_0(x) = 1$ ,  $h_2(x) = x^2 - 1$ ,  $h_4(x) = x^4 - 6x^2 + 3$ , etc.

**Lemma B.9** (Wick's theorem/Feynman rules, [71] propositions I.2, I.3, I.4, [39] theorems 1.28, 3.9, 3.19).

(i) For  $X_1, \dots, X_n \in \mathcal{H}$  (not necessarily distinct) jointly Gaussian random variables,

$$\mathbb{E}[X_1 \cdots X_n] = \sum_{\mathcal{P}} \prod_k \mathbb{E}[X_{i_k} X_{j_k}], \quad (\text{B.16})$$

where the sum is over all partitions  $\mathcal{P}$  of the set  $\{1, \dots, n\}$  into disjoint pairs  $\{i_k, j_k\}$ ,  $1 \leq k \leq n/2$ . In particular, the result is zero if  $n$  is odd.

(ii) For  $X \in \mathcal{H}$ ,

$$:X^n: = \sum_{j=0}^{\lfloor n/2 \rfloor} \frac{(-1)^j n!}{(n-2j)! j! 2^j} \mathbb{E}[X^{2j}] X^{n-2j} = \mathbb{E}[X^2]^{\frac{n}{2}} h_n(X/\mathbb{E}[X^2]^{\frac{1}{2}}). \quad (\text{B.17})$$

(iii) Let  $X_1, \dots, X_n \in \mathcal{H}$  and  $Y_1, \dots, Y_m \in \mathcal{H}$  be jointly Gaussian random variables. Then

$$\mathbb{E}[:X_1 \dots X_n : Y_1 \dots Y_m :] = \begin{cases} \sum_{\sigma \in \mathfrak{S}_n} \prod_{i=1}^n \mathbb{E}[X_i Y_{\sigma(i)}], & m = n, \\ 0, & m \neq n. \end{cases} \quad (\text{B.18})$$

In particular,  $\mathbb{E}[:X^n : Y^m :] = \delta_{nm} n! \mathbb{E}[XY]^n$ .

There is a general way of associating random variables (or numbers) to Feynman diagrams. A Feynman diagram consists of vertices, legs (segments with only one end attached to a vertex), and edges (contracted legs). Two legs are contracted means they are connected to form an edge. A Feynman diagram is called fully contracted if there is no unconnected legs. Given  $n$  random variables  $X_1, \dots, X_n$ , a Feynman diagram labelled by  $(X_1, \dots, X_n)$  is simply any Feynman diagram whose vertices are in bijection with  $\{X_1, \dots, X_n\}$ . A Feynman diagram labelled by  $(X_1, \dots, X_n)$  can now be associated with either a random variable or a number using the following rules:

- (i) for each leg attached to a vertex  $j$ , write down the corresponding random variable  $X_j$ , and multiply them all together;
- (ii) whenever two legs are contracted, enclose the corresponding two random variables by  $\mathbb{E}[\bullet]$ .

Thus fully contracted diagrams are always associated with numbers. If  $\gamma$  is a Feynman diagram labelled by  $X_1, \dots, X_n$ , denote by  $v(\gamma)$  the associated object following the above rules.

**Example B.1** ([51] section 4.4). In a physical context Feynman diagrams are used in rather formal calculations. We would like to find

$$\mathbb{E}_{\text{GFF}}^M [\phi(x)\phi(y) e^{-\frac{\lambda}{4!} \int_M \phi(x)^4 dV_M(x)}] \quad (\text{B.19})$$

by Taylor expanding  $\exp(-\frac{\lambda}{4!} \int_M \phi(x)^4 dV_M(x))$ . This gives

$$(\text{B.19}) \stackrel{\text{heu}}{=} \mathbb{E}_{\text{GFF}}^M [\phi(x)\phi(y)] + \mathbb{E}_{\text{GFF}}^M [\phi(x)\phi(y) \left(-\frac{\lambda}{4!}\right) \int \phi(z)^4 dV_M(z)]$$

$$\begin{aligned}
 & + \mathbb{E}_{\text{GFF}}^M \left[ \phi(x)\phi(y) \frac{1}{2} \left( -\frac{\lambda}{4!} \right)^2 \iint \phi(z)^4 \phi(w)^4 dV_M(z) dV_M(w) \right] + \dots \\
 & = \mathbb{E}_{\text{GFF}}^M [\phi(x)\phi(y)] - \frac{\lambda}{4!} \underbrace{\int \mathbb{E}_{\text{GFF}}^M [\phi(x)\phi(y)\phi(z)^4] dV_M(z)}_A \\
 & \quad + \frac{1}{2} \left( -\frac{\lambda}{4!} \right)^2 \iint \mathbb{E}_{\text{GFF}}^M [\phi(x)\phi(y)\phi(z)^4 \phi(w)^4] dV_M(z) dV_M(w) + \dots
 \end{aligned}$$

Treating  $\phi(x)$ ,  $\phi(y)$ ,  $\phi(z)$  as (jointly Gaussian!) random variables, by (i) of lemma B.9 and the above rules we write

$$\begin{aligned}
 A & \stackrel{\text{heu}}{=} 3 \int dV_M(z) v \left( \begin{array}{c} \text{---} \text{---} \text{---} \text{---} \\ \text{z} \end{array} \right) + 12 \int dV_M(z) v \left( \begin{array}{c} \text{---} \text{---} \text{---} \text{---} \\ \text{z} \end{array} \right) \\
 & = 3 \int G(x, y) G(z, z)^2 dV_M(z) + 12 \int G(x, z) G(y, z) G(z, z) dV_M(z),
 \end{aligned}$$

where the factors correspond to the number of ways of getting the same contraction starting from 6 legs (4 on  $z$ , 1 on  $x, y$  each), and  $G$  denotes the Green function (of  $\Delta + m^2$ ). One can represent higher order terms using these diagrams in a similar manner. More than that, the diagrams also represent actual physical processes. See Peskin and Schroeder [51].

## B.5 Itô-Wiener-Segal Isomorphism

The following amazing fact follows from comparing (B.18) with (A.3).

**Theorem** (Itô-Wiener-Segal isomorphism). *Let  $(Q, \mathcal{O}, \mathbb{P})$  be a probability space and  $\mathcal{H} \subset L^2(Q, \mathcal{O}, \mathbb{P})$  a Gaussian Hilbert space. Then the map*

$$X_1 \odot \dots \odot X_n \longmapsto :X_1 \dots X_n: \quad (\text{B.20})$$

*gives a Hilbert space isomorphism  $\mathcal{H}^{\odot n} \cong \mathcal{H}^{:n:}$ . Moreover, the direct sum of these maps for each  $n$  extends to a Hilbert space isomorphism*

$$\Gamma(\mathcal{H}) \cong L^2(\Omega, \mathcal{O}(\mathcal{H}), \mathbb{P}). \quad (\text{B.21})$$

*Consequently for  $X_1, \dots, X_n \in \mathcal{H}$ ,  $X_1 \odot \dots \odot X_n$  and  $:X_1 \dots X_n:$  are indistinguishable.*  $\square$

Transcribing proposition A.1 over to the Wick language, one has

**Corollary B.10** ([39] theorem 4.5). *Let  $(Q, \mathcal{O}, \mathbb{P})$  be a probability space and  $\mathcal{H}, \mathcal{K} \subset L^2(Q, \mathcal{O}, \mathbb{P})$  two Gaussian Hilbert spaces. Denote respectively by  $\mathcal{O}(\mathcal{H})$  and  $\mathcal{O}(\mathcal{K})$  the sub- $\sigma$ -algebra generated by variables in  $\mathcal{H}$  and in  $\mathcal{K}$ . Let  $A : \mathcal{H} \rightarrow \mathcal{K}$  be an operator with  $\|A\| \leq 1$ . Then*

$$\begin{aligned}
 \Gamma(A) : L^2(\Omega, \mathcal{O}(\mathcal{H}), \mathbb{P}) & \longrightarrow L^2(\Omega, \mathcal{O}(\mathcal{K}), \mathbb{P}), \\
 :X_1 \dots X_n: & \longmapsto :A(X_1) \dots A(X_n):,
 \end{aligned} \quad (\text{B.22})$$

*for any  $X_1, \dots, X_n \in \mathcal{H}$ , is a bounded operator with norm 1.*  $\square$

Important for us will be a consequence of the above results on conditional expectations.

**Lemma B.11.** *Let  $(Q, \mathcal{O}, \mathbb{P})$  be a probability space and  $\mathcal{A} \subset \mathcal{O}$  a sub- $\sigma$ -algebra. Then  $L^2(\Omega, \mathcal{A}, \mathbb{P})$  is a closed subspace of  $L^2(Q, \mathcal{O}, \mathbb{P})$  and for  $X \in L^2(Q, \mathcal{O}, \mathbb{P})$ ,  $\mathbb{E}[X | \mathcal{A}]$  is the orthogonal projection of  $X$  onto  $L^2(\Omega, \mathcal{A}, \mathbb{P})$ .*

**Corollary B.12** ([39] theorem 4.9). *Let  $(Q, \mathcal{O}, \mathbb{P})$  be a probability space and  $\mathcal{H}, \mathcal{K} \subset L^2(Q, \mathcal{O}, \mathbb{P})$  two Gaussian Hilbert spaces. Denote by  $P_{\mathcal{K}|\mathcal{H}}$  the restriction of the orthogonal projection  $L^2(Q, \mathcal{O}, \mathbb{P}) \rightarrow \mathcal{K}$  to  $\mathcal{H}$ . Then*

$$\begin{aligned}
 \Gamma(P_{\mathcal{K}|\mathcal{H}}) : L^2(\Omega, \mathcal{O}(\mathcal{H}), \mathbb{P}) & \longrightarrow L^2(\Omega, \mathcal{O}(\mathcal{K}), \mathbb{P}), \\
 X & \longmapsto \mathbb{E}[X | \mathcal{O}(\mathcal{K})],
 \end{aligned} \quad (\text{B.23})$$

*where  $\mathbb{E}[X | \mathcal{O}(\mathcal{K})]$  is the conditional expectation of  $X$  with respect to  $\mathcal{O}(\mathcal{K})$ .*

## C Global Analysis

### C.1 Sobolev Spaces over Domains

In this paper we make essential use of the usual  $L^2$  Sobolev spaces over Riemannian manifolds. First let  $(M, g)$  be a closed Riemannian manifold and  $s \in \mathbb{R}$ . Then the Sobolev space  $W^s(M)$  of order  $s$  is defined generally as the closure of  $C^\infty(M)$  under a norm  $\|\cdot\|_{W^s(M)}$ , where the norm  $\|\cdot\|_{W^s(M)}$  could be defined in various equivalent ways. We refer to Taylor [76] chapter 4 for a general discussion. For us,  $s = \pm 1, \pm \frac{1}{2}$ . We rely heavily on the following fact.

**Lemma C.1.** *Let  $\Lambda_{2s}$  be an elliptic strictly positive formally self-adjoint pseudodifferential operator on  $M$  with order  $2s$ . Then the inner product*

$$\langle -, - \rangle_{W^s} \stackrel{\text{def}}{=} \langle -, \Lambda_{2s} - \rangle_{L^2} \quad (\text{C.1})$$

*induces an equivalent norm for  $W^s(M)$ .*

In particular, the real power  $(\Delta_M + m^2)^s$  of the Helmholtz operator (massive Laplacian)  $\Delta_M + m^2$  provides such a candidate for  $\Lambda_{2s}$ . **Convention:** whenever we use the space  $W^s(M)$ , the inner product (C.1) with  $\Lambda_{2s} = (\Delta_M + m^2)^s$  is understood, unless otherwise specified.

**Remark C.1.** Various regimes of functional calculus can be used to define  $(\Delta_M + m^2)^s$ . One of them is presented in section 2.2.1 which in fact defines *complex* powers. We also mention a smooth functional calculus presented in Sogge [73] theorem 4.3.1.

Next we discuss important subspaces of  $W^s(M)$ . Let  $A \subset M$  be a closed set and  $U \subset M$  an open set. Define

$$W_A^s(M) \stackrel{\text{def}}{=} \{u \in W^s(M) \mid \text{supp } u \subset A \text{ as a distribution}\}, \quad (\text{C.2})$$

$$W_U^s(M) \stackrel{\text{def}}{=} \text{closure of } C_c^\infty(U) \text{ inside } W^s(M), \quad (\text{C.3})$$

$$W^s(U) \stackrel{\text{def}}{=} W_{M \setminus U}^s(M)^\perp \subset W^s(M). \quad (\text{C.4})$$

These are closed subspaces of  $W^s(M)$ .

**Remark C.2.** We point out right away that by definition, then,

$$W^s(U) \cong W^s(M) / W_{M \setminus U}^s(M), \quad (\text{C.5})$$

the latter equipped with the quotient norm, which is a more familiar characterization of  $W^s(U)$ , see Taylor [76] page 339. Our definition as in (C.4) poses the obvious problem that in general  $C_c^\infty(U) \not\subset W^s(U)$ , at least for  $s \notin \mathbb{Z}_+$ . We emphasize therefore that what is important in this definition is not the space  $W^s(U)$  *per se* but the following choice for its inner product:

$$\langle f, h \rangle_{W^s(U)} \stackrel{\text{def}}{=} \langle P_{M \setminus U}^\perp f, P_{M \setminus U}^\perp h \rangle_{W^s(M)}, \quad (\text{C.6})$$

for any  $f, h \in W^s(M)$ , in particular for  $f, h \in C_c^\infty(U)$ , which produces a norm equivalent to the quotient norm, where  $P_{M \setminus U}^\perp : W^s(M) \rightarrow W^s(U)$  denotes the orthogonal projection.

**Remark C.3.** Clearly  $W_U^s(M) \subset W_{\bar{U}}^s(M)$  by definition. In general the inclusion is strict (certainly if  $U \neq \bar{U}$ !). See Taylor [76] page 339 and section 4.7 for interesting discussions on conditions for  $s$  and  $U$  for which equality holds. In particular,  $W_\Omega^k(M) = W_{\Omega^\circ}^k(M) = W_\Omega^k(M)$  if  $\Omega \subset M$  is a domain with smooth boundary  $\partial\Omega$  (a closed Riemannian manifold with one dimension less) and  $k \in \mathbb{Z}_+$ . In this case, we use these notations interchangeably.

The rest of this appendix could be read along with section 5.1. Let  $s = -1$ . Although  $C_c^\infty(U) \not\subset W^{-1}(U)$ , we have

**Lemma C.2.** *Let  $U \subset M$  be an open set. Then  $P_{M \setminus U}^\perp(C_c^\infty(U))$  is dense in  $W^{-1}(U)$ .*

*Proof.* We note  $\Delta_M + m^2$  is local and therefore  $(\Delta + m^2)(C_c^\infty(U)) \subset C_c^\infty(U)$ . It follows from lemma 5.1 and our definition of  $W_U^1(M)$  that  $(\Delta + m^2)(C_c^\infty(U)) \subset W_{M \setminus U}^{-1}(M)^\perp$  and is dense there, proving the result.  $\square$

**Remark C.4.** Clearly, the map  $P_{M \setminus U}^\perp$  is also injective on  $C_c^\infty(U)$ ; together with lemma C.2 this shows  $P_{M \setminus U}^\perp$  is a good embedding of  $C_c^\infty(U)$  in  $W^{-1}(U)$ . In fact, this is the same as the embedding of  $C_c^\infty(U)$  in  $\mathcal{D}'(U)$ , by remark 5.1. Nevertheless, the smaller class  $(\Delta + m^2)(C_c^\infty(U))$ , as it is already dense in  $W^{-1}(U)$ , suffices as a class of test functions to define the GFF with Dirichlet condition over a domain (see remark below lemma 2.2). This reflects the fact that the Cameron-Martin pairing  $\langle -, - \rangle_{W^1}$  is more natural than  $\langle -, - \rangle_{L^2}$  in treating the GFF (see remark 2.4). We have stuck to  $\langle -, - \rangle_{L^2}$  only because this is more practical with functional analysis.

**Remark C.5.** For general  $s \in \mathbb{R}$ , one could also define

$$W_0^s(U) \stackrel{\text{def}}{=} \text{closure of } C_c^\infty(U) \text{ under (C.6)}. \quad (\text{C.7})$$

Then  $W_U^s(M) \subset W_0^s(U)$ . But it cannot generally be compared with  $W_U^s(M)$  (to the author's knowledge). See the exercises in Taylor [76] pages 343-344 for more information.

Next we state the duality results for the various spaces. Recall that  $\langle -, - \rangle_{L^2(M)}$  denotes both the inner product of  $L^2(M)$  and the distributional pairing between  $\mathcal{D}'(M)$  and  $C^\infty(M)$ . Below, we extend it to denote also the pairing between dual Sobolev spaces (see (i) of the lemma below).

**Lemma C.3.** *Let  $M$  be a closed Riemannian manifold,  $U \subset M$  an open set,  $A \subset M$  a closed set, and  $s \in \mathbb{R}$ .*

- (i)  $W^{-s}(M)$  is the dual Banach space, denoted  $W^s(M)^*$ , of  $W^s(M)$  under  $\langle -, - \rangle_{L^2}$ ;
- (ii) the annihilator of  $W_U^s(M)$  under  $\langle -, - \rangle_{L^2}$  is  $W_{M \setminus U}^{-s}(M)$ , that is,

$$W_{M \setminus U}^{-s}(M) = \{u \in W^{-s}(M) \mid \langle u, f \rangle_{L^2} = 0 \text{ for all } f \in W_U^s(M)\}; \quad (\text{C.8})$$

the annihilator of  $W_A^s(M)$  is accordingly  $W_{M \setminus A}^{-s}(M)$ ;

- (iii)  $W^s(U)^* \cong W_U^{-s}(M)$ ,  $W_U^s(M)^* \cong W^{-s}(U)$ , these spaces being therefore reflexive.

Finally, when  $\Omega \subset M$  is a domain with smooth boundary  $\partial\Omega$ , we define, in view of lemma 5.1, the Dirichlet Green operator  $(\Delta_{\Omega, D} + m^2)^{-1} := (\Delta + m^2)^{-1} P_{M \setminus \Omega^\circ}^\perp : W^{-1}(\Omega^\circ) \rightarrow W_{\Omega^\circ}^1(M)$ . Clearly this agrees with the usual definition. In terms of quadratic forms,

**Lemma C.4** ([71] theorem VII.1). *Let  $\Omega \subset M$  be a domain with smooth boundary  $\partial\Omega$ . We have*

$$\langle f, (\Delta_{\Omega, D} + m^2)^{-1} h \rangle_{L^2} = \langle P_{M \setminus \Omega^\circ}^\perp f, P_{M \setminus \Omega^\circ}^\perp h \rangle_{W^{-1}}, \quad (\text{C.9})$$

for  $f, h \in C_c^\infty(\Omega^\circ)$ .

## C.2 Symbol Convergence Lemma and Heat Kernel

*Proof of lemma 3.1.* By coordinate invariance of the definition of  $\Psi^m(M)$  it suffices to pick  $x \in M$  and prove the result for a chart around  $x$  and  $\chi(x) = 1$ . Denote the kernel of  $\chi E_\varepsilon \chi$  by  $E_{\chi, \varepsilon}$  then in this chart we could write

$$E_{\chi, \varepsilon}(x, y) = \tilde{E}_{\chi, \varepsilon}(x, h) = \frac{1}{F_\varepsilon(x)} \tilde{\psi}\left(\frac{h}{\varepsilon}\right) \tilde{\chi}(h), \quad (\text{C.10})$$

where  $h = x - y$ . Indeed, by definition of our function  $\psi$  and freedom of choosing  $\chi$  we could further assume that for small enough  $\varepsilon$  one has  $\tilde{\chi}(h) \equiv 1$  on the support of  $\tilde{\psi}(\cdot/\varepsilon)$ . Thus under this condition

$$\sigma_{\chi E_\varepsilon \chi}(x, \xi) = \int_{\mathbb{R}^d} e^{-ih \cdot \xi} \frac{1}{F_\varepsilon(x)} \tilde{\psi}\left(\frac{h}{\varepsilon}\right) dh = \frac{\varepsilon^d}{F_\varepsilon(x)} \underbrace{F_1(x) \sigma_{\chi E_1 \chi}(x, \varepsilon \xi)}_{\text{indep. of } x}. \quad (\text{C.11})$$

Note that  $\sigma_{\chi E_1 \chi}(x, \eta)$  is Schwartz in  $\eta$  and  $\sigma_{\chi E_1 \chi}(x, 0) = 1$ . On the other hand clearly  $\sigma_{\chi \mathbb{1} \chi}(x, \xi) \equiv 1$ . Thus for some  $U' \subset U$  depending only on the chart and  $\chi$ , one has

$$\sup_{x \in K \Subset U'} \sup_{\xi} \frac{|\sigma_{\chi(E_\varepsilon - \mathbb{1}) \chi}(x, \xi)|}{\langle \xi \rangle^\delta} \leq \begin{cases} C \sup_{x \in K \Subset U'} \sup_{|\xi| \leq R} \langle \xi \rangle^{-\delta} |\sigma_{\chi E_1 \chi}(x, \varepsilon \xi) - 1| \leq C_{K, \chi} \sqrt{\varepsilon}, \\ C \sup_{x \in K \Subset U'} \sup_{|\xi| \geq R} (\dots) \leq C_{K, \chi} \varepsilon^{\delta/2} \sup_{\eta} |\sigma_{\chi E_1 \chi}(x, \eta)|, \end{cases} \quad (\text{C.12})$$

with  $R = \varepsilon^{-1/2}$ . Next we deal with derivatives. Note that by (C.11) all the  $x$ -derivatives fall on  $1/F_\varepsilon(x)$  and all  $\xi$ -derivatives fall on  $\sigma_{\chi E_1 \chi}(x, \varepsilon \xi)$ . Indeed, one has  $|\partial_x^\alpha (1/F_\varepsilon(x))| \leq C_\alpha \varepsilon^{-d}$  (see Dyatlov and Zworski [19] page

28), and so when there are only  $x$ -derivatives we obtain the same bounds as (C.12) only with new constants depending on  $\alpha$ . When there is at least one  $\xi$ -derivative,

$$|\partial_\xi^\beta (\sigma_{\chi_{E_1\chi}}(x, \varepsilon\xi))| = |\varepsilon^{|\beta|} (\partial_\xi^\beta \sigma_{\chi_{E_1\chi}})(x, \varepsilon\xi)| \leq C_{\beta,K,\chi} \varepsilon^{|\beta|} \langle \varepsilon\xi \rangle^{\delta-|\beta|}, \quad |\beta| \geq 1. \quad (\text{C.13})$$

Hence, on account of (C.11) again,

$$\sup_{x \in K \in U'} \sup_{\xi} \frac{|\partial_x^\alpha \partial_\xi^\beta \sigma_{\chi_{(E_\varepsilon-1)\chi}}(x, \xi)|}{\langle \xi \rangle^{\delta-|\beta|}} \leq C \varepsilon^d C_\alpha \varepsilon^{-d} C_{\beta,K,\chi} \varepsilon^{|\beta|} = C_{\alpha,\beta,K,\chi} \varepsilon^{|\beta|}. \quad (\text{C.14})$$

Consequently, all the  $\mathcal{S}_{1,0}^\delta$  seminorms of  $\sigma_{\chi_{(E_\varepsilon-1)\chi}}$  goes to zero as  $\varepsilon \rightarrow 0$ . We obtain the result.  $\square$

In what follows we sum up some properties of the heat operator  $e^{-t(\Delta+m^2)}$  of the massive Laplacian (Helmholtz operator) and its Schwartz kernel  $p_t(x, y)$  called the heat kernel.

**Lemma C.5** ([10] theorems 2.30, 2.38 and pages 92-94). *We have*

- (i)  $p_t(x, y) \in C^\infty((0, \infty) \times M \times M)$ ;
- (ii) *we have*

$$(\Delta + m^2)^{-1} = \int_0^\infty e^{-t(\Delta+m^2)} dt. \quad (\text{C.15})$$

*In particular, the kernel  $G_t$  of  $e^{-t(\Delta+m^2)}(\Delta + m^2)^{-1} = (\Delta + m^2)^{-1}e^{-t(\Delta+m^2)}$  is*

$$G_t(x, y) = \int_t^\infty p_s(x, y) ds, \quad (\text{C.16})$$

*for  $x, y \in M$ .*

- (iii) *Let  $\dim M = n$ . There are asymptotic expansions*

$$p_t(x, y) \sim \frac{1}{(4\pi t)^{n/2}} e^{-\frac{1}{4t}d(x,y)^2} \sum_{i=0}^\infty f_i(x, y) t^i, \quad (\text{C.17})$$

$$\text{tr}_{L^2(M)} (e^{-t(\Delta+m^2)}) \sim \frac{1}{(4\pi t)^{n/2}} \sum_{i=0}^\infty a_i t^i \quad (\text{C.18})$$

*as  $t \rightarrow 0+$ , for some real numbers  $a_i$  and functions  $f_i \in C^\infty(M \times M)$ ,  $i = 0, 1, 2, \dots$*

- (iv) *For  $t$  large and each  $\ell \in \mathbb{N}$ ,*

$$\|p_t(x, y)\|_{C^\ell} \leq C_\ell e^{-tm^2/2} \quad (\text{C.19})$$

*for some constant  $C_\ell$ .*



## References

- [1] Albeverio, Sergio and Seiichiro Kusuoka. *The invariant measure and the flow associated to the  $\Phi_3^4$ -quantum field model*. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5), Vol. XX (2020), 1359-1427.
- [2] Atiyah, M. F., and R. Bott. *A Lefschetz Fixed Point Formula for Elliptic Complexes: II. Applications*. Annals of Mathematics 88, no. 3 (1968): 451–91.
- [3] Alekseevsky, D.V., Kriegl, A., Losik, M., Peter W. Michor. *Reflection groups on Riemannian manifolds*. Annali di Matematica 186, 25–58 (2007)
- [4] I. Bailleul, N. V. Dang, L. Ferdinand, T.D. Tô,  $\Phi_3^4$  measures on compact Riemannian 3-manifolds, arXiv preprint, [arXiv:2304.10185](https://arxiv.org/abs/2304.10185).
- [5] I. Bailleul and N. V. Dang and L. Ferdinand and T. D. Tô, Global harmonic analysis for  $\Phi_3^4$  on closed Riemannian manifolds, arXiv preprint, [arXiv:2306.07757v2](https://arxiv.org/abs/2306.07757v2).
- [6] Romeo Brunetti, Klaus Fredenhagen, Rainer Verch, *The Generally Covariant Locality Principle – A New Paradigm for Local Quantum Field Theory*, Commun. Math. Phys. 237, 31–68 (2003).
- [7] Bergh, Jöran, and Jörgen Löfström, *Interpolation Spaces: An Introduction*, Springer-Verlag, 1976.
- [8] Nicolas Bergeron, *Variétés en expansion (d’après M. Gromov, L. Guth, ...)*, Séminaire Bourbaki, 69ème année, 2016-2017, no 1132.
- [9] D. Burghelea, L. Friedlander, T. Kappeler, *Mayer-Vietoris type formula for determinants of elliptic differential operators*, Journal of Functional Analysis, Volume 107, Issue 1, 1992, Pages 34-65.
- [10] Berline, Nicole, Ezra Getzler, and Michèle Vergne, *Heat Kernels and Dirac Operators*, Springer-Verlag, 2004.
- [11] Volker Betz, Fumio Hiroshima, József Lőrinczi, *Feynman-Kac-Type Theorems and Gibbs Measures on Path Space: With Applications to Rigorous Quantum Field Theory*, Walter de Gruyter, 2011.
- [12] Bogachev, Vladimir I., *Gaussian Measures*, American Mathematical Society, 1998.
- [13] Carron, Gilles. *Déterminant Relatif et La Fonction XI*, American Journal of Mathematics 124, no. 2, 307–52, 2002.
- [14] Chow, Bennett, and Dan Knopf, *The Ricci Flow: An Introduction*, Mathematical Surveys and Monographs Volume 110, American Mathematical Society, 2004.
- [15] Nguyen Viet Dang, *Renormalization of determinant lines in quantum field theory*, Analysis & PDE, Vol. 15 (2022), No. 1, 1–62.
- [16] Dimock, Jonathan, *Markov quantum fields on a manifold*, Rev. Math. Phys. 16, 243-256, 2004.
- [17] Dimock, Jonathan. *Quantum Mechanics and Quantum Field Theory: A Mathematical Primer*. Cambridge: Cambridge University Press, 2011.
- [18] Dudley, R. M., *Real Analysis and Probability*, Cambridge University Press, 2004.
- [19] Semyon Dyatlov, Maciej Zworski, *Dynamical zeta functions for Anosov flows via microlocal analysis*, Ann. Scient. Éc. Norm. Sup, 4e série, t. 49, 2016, p. 543-577.
- [20] Eskin, Gregory, *Lectures on Linear Partial Differential Equations*, Graduate Studies in Mathematics Volume 123, American Mathematical Society, 2011.
- [21] Folland, Gerald B., *Real Analysis: Modern Techniques and Their Applications*, Second Edition, John Wiley & Sons, Inc., 1999.
- [22] Folland, Gerald B. *Harmonic Analysis in Phase Space*. (AM-122). Princeton University Press, 1989.
- [23] G. W. Gibbons (1998). *Real tunnelling geometries*, Class. Quantum Grav. 15, 2605.
- [24] Gilkey, Peter B., *Invariance Theory, the Heat Equation, and the Atiyah-Singer Index Theorem*, Second Edition, Studies in Advanced Mathematics, CRC Press, 1995.
- [25] Israel Gohberg, Seymour Goldberg, and Nahum Krupnik, *Traces and Determinants of Linear Operators, Operator Theory: Advances and Applications*, Vol. 116, Birkhäuser Basel, 2000.
- [26] Glimm, James, and Arthur Jaffe, *Quantum Physics: a Functional Integral Point of View*, Second Edition, Springer-Verlag, 1987.
- [27] Gubinelli, M., Hofmanová, M. *A PDE Construction of the  $\Phi_3^4$  Euclidean Quantum Field Theory*. Commun. Math. Phys. 384, 1–75 (2021).



- [28] Guillarmou, C., Kupiainen, A., Rhodes, R., and Vargas, V. (2021). *Segal's axioms and bootstrap for Liouville Theory*. arXiv preprint arXiv:2112.14859.
- [29] Gu, Yu and Mourrat, Jean-Christophe. (2016). *On generalized Gaussian free fields and stochastic homogenization*. Electronic Journal of Probability. 22. 10.1214/17-EJP51.
- [30] Guo, L., Paycha, S. and Zhang, B. *Mathematical Reflections on Locality*. Jahresber. Dtsch. Math. Ver. 125, 197–238 (2023).
- [31] M. Hairer, *Introduction to Malliavin Calculus*, lecture notes, available at <https://www.hairer.org/Teaching.html>.
- [32] Martin Hairer and Rhys Steele. *The  $\Phi_3^4$  measure has sub-Gaussian tails*. J. Stat. Phys., 186(3):38, 2022.
- [33] Andreas Hermann, Emmanuel Humbert, *About the mass of certain second order elliptic operators*, Advances in Mathematics, Volume 294, 2016, Pages 596-633.
- [34] Howe, Roger, *The oscillator semigroup*. In: The mathematical heritage of Hermann Weyl, 61–132. Proc. Sympos. Pure Math., 48, American Mathematical Society, Providence, RI, 1988.
- [35] H. Hohnhold, S. Stolz, and P. Teichner. *From minimal geodesics to supersymmetric field theories*. A celebration of the mathematical legacy of Raoul Bott, 50:207–274, 2010.
- [36] S. Kandel, P. Mnev and K. Wernli, *Two-dimensional perturbative scalar QFT and Atiyah-Segal gluing*, Adv. Theor. Math. Phys. **25** (2021) no.7, 1847-1952.
- [37] Arthur Jaffe, Gordon Ritter. *Reflection positivity and monotonicity*. J. Math. Phys. 1 May 2008; 49 (5): 052301.
- [38] Jaffe, A., Ritter, G. *Quantum Field Theory on Curved Backgrounds. I. The Euclidean Functional Integral*. Commun. Math. Phys. 270, 545–572 (2007).
- [39] Janson, Svante, *Gaussian Hilbert Spaces*, Cambridge Tracts in Mathematics (129), Cambridge University Press, 1997.
- [40] Kontsevich, Maxim and Segal, Graeme, *Wick rotation and the positivity of energy in quantum field theory*, arXiv:2105.10161
- [41] Maxim Kontsevich and Simeon Vishik. *Determinants of elliptic pseudo-differential operators*, arXiv preprint, arXiv:hep-th/9404046.
- [42] Yoonweon Lee, *Mayer-Vietoris formula for determinants of elliptic operators of Laplace-Beltrami type* (after Burghelca, Friedlander and Kappeler), Differential Geometry and its Applications, Volume 7, Issue 4, 1997, Pages 325-340.
- [43] Le Gall, Jean-François, *Brownian Motion, Martingales, and Stochastic Calculus*, Graduate Texts in Mathematics, Vol. 274, Springer Cham, 2016.
- [44] Ludewig, M. *Asymptotic expansions and conformal covariance of the mass of conformal differential operators*. Ann Glob Anal Geom 52, 237–268 (2017).
- [45] Mourrat, Jean-Christophe, Hendrik Weber, and Weijun Xu (2017). *Construction of  $\Phi_3^4$  diagrams for pedestrians*. arXiv:1610.08897.
- [46] Naud, Frédéric. *Determinants of Laplacians on random hyperbolic surfaces*. Journal d'Analyse Mathématique 151.1 (2023): 265-291.
- [47] Nelson, Edward, *A Quartic interaction in two dimensions*, Mathematics Theory of Elementary Particles, R. Goodman and I. Segal Eds., MIT Press, Cambridge, 1966.
- [48] Nelson, Edward, *The Construction of Quantum Fields from Markov Fields*, J. Functional Analysis, 12, 97-112.
- [49] Nelson, Edward. *Probability theory and euclidean field theory*. Lecture Notes in Physics, Berlin Springer Verlag 25 (1973): 94.
- [50] Paternain, Gabriel P., Mikko Salo, and Gunther Uhlmann. *Geometric Inverse Problems: With Emphasis on Two Dimensions*. Cambridge Studies in Advanced Mathematics. Cambridge: Cambridge University Press, 2023
- [51] Peskin, Michael Edward and Schroeder, Daniel V.. *An Introduction to Quantum Field Theory*. Westview Press, 1995.
- [52] Pickrell, Doug,  *$P(\phi)_2$  Quantum Field Theories and Segal's Axioms*, Commun. Math. Phys. 280, 403–425, 2008.

- [53] Vershik, Anatoly; Petrov, Fedor, and Zatitskiy, P.. (2014). *Virtual Continuity of Measurable Functions and Its Applications*. Russian Mathematical Surveys. 69.
- [54] Powell, Ellen and Wendelin Werner, *Lecture Notes on the Gaussian Free Field*, Cours Spécialisés, Vol. 28, Société Mathématique de France, 2021.
- [55] Quine, J. R., S. H. Heydari, and R. Y. Song. *Zeta Regularized Products*. Transactions of the American Mathematical Society 338, no. 1 (1993): 213–31.
- [56] Ray, D. B., and I. M. Singer, *R-Torsion and the Laplacian on Riemannian Manifolds*, Advances in Mathematics 7, 145-210, 1971.
- [57] M. Reed, B. Simon, *Methods of Modern Mathematical Physics*, Volume I: Functional Analysis, Academic Press, 1972.
- [58] Ritter, William Gordon. *Euclidean quantum field theory: Curved spacetimes and gauge fields*. PhD thesis, Harvard University, 2007. Available at: <https://ui.adsabs.harvard.edu/abs/2007PhDT.....57R/abstract>
- [59] Roe, John, *Elliptic operators, topology and asymptotic methods*, Second Edition, CRC Press, 2001.
- [60] Rosenberg, Steven, *The Laplacian on a Riemannian Manifold*, London Mathematical Society Student Texts, Vol. 31, Cambridge University Press, 1998.
- [61] Rozanov, Y.A. *Markov Random Fields*. Springer, New York, 1982.
- [62] Rudin, Walter, *Functional Analysis*, Second Edition, McGraw-Hill, Inc., 1991.
- [63] Ruelle, David. Thermodynamic formalism: the mathematical structure of equilibrium statistical mechanics. Cambridge University Press, 2004.
- [64] Schoen, R., Yau, ST. *Proof of the positive mass theorem. II*. Commun.Math. Phys. 79, 231–260 (1981).
- [65] Seeley, R. T. *Complex powers of an elliptic operator*. In: Singular Integrals (Proc. Sympos. Pure Math., Chicago, Ill., 1966), pp. 288–307. American Mathematical Society, Providence, RI, 1967
- [66] G. Segal, *The definition of conformal field theory*, Topology, Geometry, and Quantum Field Theory, (Ed. U. Tillmann), Vol. 308, London Math. Soc. Lecture Notes, London, 2004, 421–577.
- [67] Scott Sheffield, *Gaussian free fields for mathematicians*, Probab. Theory Relat. Fields, 139, 521–541, Springer-Verlag, 2007.
- [68] Shubin, M. A., *Pseudodifferential Operators and Spectral Theory*, Stig I. Andersson trans., Second Edition, Springer-Verlag, Berlin, 2001.
- [69] Symanzik, Kurt. *Euclidean Quantum Field Theory*. New York Univ., NY, 1969.
- [70] Simon, Barry, *Trace Ideals and their Applications*, Second Edition, Mathematical Surveys and Monographs, Vol. 120, American Mathematical Society, 2005.
- [71] Simon, Barry, *The  $P(\phi)_2$  (Euclidean) Quantum Field Theory*, Princeton University Press, New Jersey, 1974.
- [72] Simon, Barry, *Operator Theory: A Comprehensive Course in Analysis, Part 4*, Vol. 4, American Mathematical Society, 2015.
- [73] Sogge, Christopher D., *Fourier Integrals in Classical Analysis*, Second Edition, Cambridge Tracts in Mathematics, Vol. 210, Cambridge University Press, 2017.
- [74] Sogge, Christopher Donald. *Hangzhou Lectures on Eigenfunctions of the Laplacian*. Princeton (N.J.) Oxford: Princeton university press, 2014
- [75] S. Stolz and P. Teichner. *Supersymmetric field theories and generalized cohomology*. In Mathematical foundations of quantum field theory and perturbative string theory, Proc. Sympos. Pure Math. Vol. 83, pages 279–340. Amer. Math. Soc., Providence, RI, 2011.
- [76] Taylor, Michael E., *Partial Differential Equations I: Basic Theory*, Second Edition, Springer New York, 2011.
- [77] Taylor, Michael E., *Partial Differential Equations II: Qualitative Studies of Linear Equations*, Second Edition, Springer New York, 2011.
- [78] Taylor, Michael E., *The Dirichlet-to-Neumann Map and Fractal Variants*, lecture notes, available at: <https://mtaylor.web.unc.edu/notes/fractal-analysis/>