

Quantum invariants arising from $U_h\mathfrak{sl}(2|1)$ are q-holonomic

Jennifer Brown and Nathan Geer

March 20, 2024

Abstract

We show that the quantum invariants arising from typical representations of the quantum group $U_h(\mathfrak{sl}(2|1))$ are q-holonomic. In particular, this implies the existence of an underlying field theory for which this family of invariants are partition functions.

1 Introduction

q-Holonomic systems were developed to apply techniques from D-modules to the study of hypergeometric functions [Sab93]. Such systems appear widely in quantum topology, where they describe the recursive or quasiperiodic behavior of various quantum knot invariants, see [GL05, GL16, GLL18].

As in the non-q-deformed setting, q-holonomic systems carry geometric meaning. In the context of quantum topology, the q-Weyl modules underlying the q-holonomic systems are understood as quantizations of character varieties of three manifolds. The precise relationship between the recursion relations and quantum character varieties is the topic of the long-standing AJ conjecture [Gar04, Lê06], which posits that the minimal order recurrence relation of the colored Jones polynomial is a quantization of the A-polynomial. Even more geometric in flavor is the volume conjecture and its various generalizations [Kas97, Guk05, DG11, GSAF12], which relate the asymptotic behavior of quantum invariants to the hyperbolic geometry of knot complements. Both conjectures are stated for the colored Jones polynomial but expected to generalize to other families of quantum invariants. They both also imply that the invariants are q-holonomic, making this an important property to establish when considering a new family of invariants.

In this paper we show that the quantum invariants arising from the typical representations of the quantum group associated to the special linear Lie superalgebra $\mathfrak{sl}(2|1)$ are q-holonomic. We conjecture that this property holds for a larger class of Lie superalgebras, see Conjecture 3.7. Unlike the families of representations underlying the colored Jones polynomial and HOMFLY-PT polynomial, typical representations in the superalgebra setting are indexed by a combination of discrete and complex weights and often have vanishing quantum dimensions. Therefore we must use the techniques developed in [BDGG20] to prove q-holonomicity in a non-semisimple setting. Having established q-holonomicity, we open the door to a deformed version of a volume conjecture and provide a potential tool towards its study.

Our choice to focus on $\mathfrak{sl}(2|1)$ is motivated by recent work on Chern-Simons theories constructed from supergroups, such as those studied by Mikhaylov and Witten in [MW15], see also [AGPS18, Mik15]. Perturbative versions of these theories are proposed in [AGP21] and lead to quantum invariants of 3-manifolds. In this paper we study the link invariant underlying the 3-manifold invariants of [AGP21]. The relationships between these supergroup theories and related topological invariants have been made precise in the case of $\mathfrak{gl}(1|1)$ in the recent work [GY22].

1.1 A quantum torus from skein theory

In this section we will use some insights from skein theory, following the ideas of [Gel01, FGL01], to explore the origins of the q-holonomic systems associated with the $\mathfrak{sl}(2|1)$ invariant defined in Section 2.2. A full treatment is out of scope of the current paper. To describe the action in a precise technical sense, one would need to understand non-unital defect skein theory. Defects in the unital setting would build on quantum decorated character stacks and stratified factorization homology [AFT17, JLSS21], but at the time of writing

are missing the skein theoretic description necessary to understand them in our context. Non-unital skein theory is needed to handle non-semisimple ribbon categories. This is just now being developed [CGP23] and represents a significant undertaking, as many of the foundational results underlying skein theory assume unitality.

With these caveats in mind, the story revolves around the skein algebra of the torus \mathbb{T} . The $\mathfrak{sl}(2)$ skein algebra of \mathbb{T} can be embedded into the rank one q -Weyl algebra \mathbb{W}_1 , [FG00, Theorem 4.3]. Its action on the skein module of the knot complement is the topological origin of the q -holonomic \mathbb{W}_1 -module generated by the colored Jones polynomial, as studied in [GL05]. The embedding into \mathbb{W}_1 can be understood in terms of the quantum trace map [BW11, Theorem 11], which embeds skein algebras into quantum tori associated to quantum cluster algebras. This construction depends on stated skein algebras. For $\mathfrak{sl}(2)$ these are defined and studied in [Lê16, CL22], but for arbitrary ribbon categories one must in general work with the more abstract but closely related internal skein algebras, see [GJS19, Hai21].

We turn to the underlying geometry to gain intuition for this embedding in the Lie superalgebra setting. In [KQ22] this quantum trace map is re-interpreted as a q -deformation of a regular morphism between character varieties:

$$\mathcal{X}_{\mathbb{C}^*}(\hat{\Sigma}, \hat{\mathcal{P}}) \rightarrow \mathcal{X}_{\mathrm{SL}_2}(\Sigma, \mathcal{P}). \quad (1)$$

Here $\hat{\Sigma}$ is a branched double cover of a surface Σ . It's constructed from a triangulation of Σ and has a non-trivial deck transformation σ . Only characters which respect σ are considered, see [KQ22, Def. 2.16]. The deformation (or quantization) of (1) is a morphism from the $\mathfrak{sl}(2)$ skein algebra to an equivariant \mathbb{C}^* skein algebra. These have extra framing data called states at a set of marked points $\mathcal{P} \subset \Sigma$ and $\hat{\mathcal{P}} \subset \hat{\Sigma}$, respectively.

Following this line of thought, we expect that the most topologically-motivated and natural quantum torus will be the skein algebra

$$\mathrm{SkAlg}_{\mathrm{Rep}_h^{tf} \mathfrak{t}}(\mathbb{T}), \quad (2)$$

where the Cartan $\mathfrak{t} \subset \mathfrak{sl}(2|1)$ plays the role that \mathbb{C}^* does in the $\mathrm{SL}_2 \mathbb{C}$ theory, and $\mathrm{Rep}_h^{tf} \mathfrak{t}$ is a full subcategory of the topologically free $U_h \mathfrak{t}$ -modules, see definition 2.4. Note that the quantized enveloping algebra of $\mathfrak{t} \simeq \mathbb{C}[h_1, h_2]$ has an R -matrix given by (17), and hence the braiding in the associated category is of the form $v \otimes w \mapsto q^{\omega_{v,w}} w \otimes v$ for some scalars $\omega_{v,w}$. It's thanks to this that the associated skein algebra will be a quantum torus. The embedding that translates between the $\mathfrak{sl}(2|1)$ -skein module action and that of this \mathfrak{t} quantum torus should be a deformation of a regular morphism

$$\mathcal{X}_{\mathfrak{t}}(\mathbb{T}, \mathcal{P}) \rightarrow \mathcal{X}_{\mathfrak{sl}(2|1)}(\mathbb{T}, \mathcal{P}). \quad (3)$$

1.2 Acknowledgements

This material is based upon work supported by the National Science Foundation under Award No. 2202753. JB is funded by the Simons Foundation award 888988 as part of the Simons Collaboration on Global Categorical Symmetry. NG is partially supported by National Science Foundation grant DMS-2104497.

2 The Lie superalgebra $\mathfrak{sl}(2|1)$

We give a brief overview to set notation and recommend [Kac77] for a careful treatment of the theory of Lie superalgebras. Let $\mathbb{Z}_2 = \{\bar{0}, \bar{1}\}$ be the additive group of order two. A *super-space* is a \mathbb{Z}_2 -graded vector space $V = V_{\bar{0}} \oplus V_{\bar{1}}$ over \mathbb{C} . The parity of a homogeneous element $x \in V_{\bar{i}}$ is denoted by $\bar{x} = \bar{i} \in \mathbb{Z}_2$ and the elements of $V_{\bar{0}}$ (resp. $V_{\bar{1}}$) are called even (resp. odd). A *morphism of super vector spaces of degree $\bar{d} \in \mathbb{Z}_2$* is a linear map $f : V \rightarrow W$ which satisfies $\overline{f(v)} = \bar{v} + \bar{d}$ for each homogeneous $v \in V$. An algebra on a superspace is a *superalgebra* if its usual structure maps respect the \mathbb{Z}_2 -grading. A (left) *module* over a superalgebra A is a super vector space M together with a superalgebra homomorphism $A \rightarrow \mathrm{End}_{\mathbb{C}}(M)$ of degree $\bar{0}$. A *Lie superalgebra* is a super-space $\mathfrak{g} = \mathfrak{g}_{\bar{0}} \oplus \mathfrak{g}_{\bar{1}}$ with a super-bracket $[\cdot, \cdot] : \mathfrak{g}^{\otimes 2} \rightarrow \mathfrak{g}$ that preserves the \mathbb{Z}_2 -grading, is super-antisymmetric ($[x, y] = -(-1)^{\bar{x}\bar{y}}[y, x]$), and satisfies the super-Jacobi identity.

Let $A = (a_{ij})$ be the $\mathfrak{sl}(2|1)$ Cartan matrix, i.e. the 2×2 matrix given by $a_{11} = 2$, $a_{12} = a_{21} = -1$ and $a_{22} = 0$.

Definition 2.1. Let $\mathfrak{sl}(2|1)$ be the Lie superalgebra generated by $h_i, e_i,$ and $f_i, i = 1, 2,$ where h_1, h_2, e_1 and f_1 are even while e_2 and f_2 are odd. The generators satisfy the relations

$$[h_i, h_j] = 0, \quad [h_i, e_j] = a_{ij}e_j, \quad [h_i, f_j] = -a_{ij}f_j \quad [e_i, f_j] = \delta_{ij}h_i,$$

$$[e_2, e_2] = [f_2, f_2] = 0, \quad [e_1, [e_1, e_2]] = [f_1, [f_1, f_2]] = 0.$$

Let h be an indeterminate and $q = e^{h/2}$. We adopt the following standard notations:

$$q^z = e^{zh/2}, \quad \{z\} = q^z - q^{-z}, \quad \text{and} \quad [z] = \frac{\{z\}}{\{1\}}$$

In this paper we work with $U_h(\mathfrak{sl}(2|1))$ instead of $U_q(\mathfrak{sl}(2|1))$ because the standard R-matrix, see (16) and (17), is well defined as an element in $U_h(\mathfrak{sl}(2|1)) \otimes U_h(\mathfrak{sl}(2|1))$ but not in $U_q(\mathfrak{sl}(2|1)) \otimes U_q(\mathfrak{sl}(2|1))$.

Definition 2.2. Let $U_h(\mathfrak{sl}(2|1))$ be the $\mathbb{C}[[h]]$ -Hopf superalgebra generated by the even elements $h_1, h_2, E_1,$ and F_1 together with the odd elements E_2 and F_2 . These are subject to the relations:

$$[h_i, h_j] = 0, \quad [h_i, E_j] = a_{ij}E_j, \quad [h_i, F_j] = -a_{ij}F_j,$$

$$[E_i, F_j] = \delta_{i,j} \frac{q^{h_i} - q^{-h_i}}{q - q^{-1}}, \quad E_2^2 = F_2^2 = 0,$$

$$E_1^2 E_2 - (q + q^{-1}) E_1 E_2 E_1 + E_2 E_1^2 = 0 \quad F_1^2 F_2 - (q + q^{-1}) F_1 F_2 F_1 + F_2 F_1^2 = 0$$

where $[\cdot, \cdot]$ is the super-commutator given by $[x, y] = xy - (-1)^{\bar{x}\bar{y}}yx$. The coproduct, counit, and antipode are given by

$$\begin{aligned} \Delta(E_i) &= E_i \otimes 1 + q^{-h_i} \otimes E_i, & \epsilon(E_i) &= 0 & S(E_i) &= -q^{h_i} E_i \\ \Delta(F_i) &= F_i \otimes q^{h_i} + 1 \otimes F_i, & \epsilon(F_i) &= 0 & S(F_i) &= -F_i q^{-h_i} \\ \Delta(h_i) &= h_i \otimes 1 + 1 \otimes h_i, & \epsilon(h_i) &= 0 & S(h_i) &= -h_i. \end{aligned}$$

2.1 Representations of $\mathfrak{sl}(2|1)$ and $U_h(\mathfrak{sl}(2|1))$

In this section we compare the representation theory of $\mathfrak{sl}(2|1)$ and $U_h(\mathfrak{sl}(2|1))$. In particular, we recall the category of *topologically free modules* from which the quantum invariants considered in this paper are constructed.

Definition 2.3. Let $a := (a_1, a_2) \in \mathbb{Z}_{\geq 0} \times \mathbb{C}$. We denote by $V(a_1, a_2)$ the highest weight $\mathfrak{sl}(2|1)$ -module with a highest weight vector v_0 whose parity is even satisfying

$$h_i.v_0 = a_i v_0 \quad \text{and} \quad e_i v_0 = 0. \quad (4)$$

Such a $\mathfrak{sl}(2|1)$ highest weight module is called **typical** if it splits in any finite-dimensional representation.

In [Kac78] this condition is reduced to a system of inequalities on the weights, which for $\mathfrak{sl}(2|1)$ is $a_1 + a_2 + 1 \neq 0$ and $a_2 \neq 0$. In the same article it's proven that typical modules are projective and injective in the category of finite-dimensional $\mathfrak{sl}(2|1)$ -modules.

Next we consider $U_h(\mathfrak{sl}(2|1))$. We restrict our attention to a certain subcategory of representations which are closely tied to the $\mathfrak{sl}(2|1)$ theory.

Definition 2.4. A $U_h(\mathfrak{sl}(2|1))$ -module W is called **topologically free of finite rank** if it is isomorphic as a $\mathbb{C}[[h]]$ -module to $V[[h]] := V \otimes \mathbb{C}[[h]]$, where V is a finite-dimensional $\mathfrak{sl}(2|1)$ -module. Let $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ denote the category of topologically free of finite rank $U_h(\mathfrak{sl}(2|1))$ -modules where all morphisms are even.

As a topological algebra, $U_h(\mathfrak{sl}(2|1))$ is isomorphic to the quantized enveloping superalgebra of $\mathfrak{sl}(2|1)$ [Gee06]. In [Gee07, Thm 1.2] this isomorphism is used to construct a highest weight $U_h(\mathfrak{sl}(2|1))$ -module $\tilde{V}(a)$ which deforms $V(a)$. In particular, it is shown that the super-space underlying $\tilde{V}(a)$ is $V(a)[[h]]$. Note that if $V(a)$ typical (and hence projective and injective) then its h-adic counterpart $\tilde{V}(a)$ is likewise both projective and injective. The topologically free representation theory of $U_h(\mathfrak{sl}(2|1))$ is thus parallel to that of the Lie superalgebra $\mathfrak{sl}(2|1)$.

Following [GP18], a representation $W \in \text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ is called *simple* if

$$\text{End}_{\text{Rep}_h^{tf} \mathfrak{sl}(2|1)}(W) \simeq \mathbb{C}[[h]] \text{Id}_W. \quad (5)$$

The $U_h(\mathfrak{sl}(2|1))$ -module $\tilde{V}(a)$ is simple and called *typical* whenever the associated $\mathfrak{sl}(2|1)$ -module $V(a)$ is likewise typical.

2.1.1 Typical $U_h(\mathfrak{sl}(2|1))$ -modules

Here we give an explicit presentation of *typical modules* of $U_h(\mathfrak{sl}(2|1))$. Recall that these are simple modules $\tilde{V}(a_1, a_2)$ with $a_1 \in \mathbb{Z}_{\geq 0}$, $a_2 \in \mathbb{C}$, $a_2 \neq 0$, and $a_1 + a_2 + 1 \neq 0$. This is the family of modules used in defining the knot invariants considered in this article. The typical module $\tilde{V}(a_1, a_2)$ has dimension $4a_1 + 4$ and is generated by a distinguished even highest weight vector v . It is uniquely determined by simplicity and the weights $h_1 v = a_1 v$ and $h_2 v = a_2 v$. We introduce the following notation, motivated by the decomposition (8) described below.

$$v_{0,0} := v, \quad v_{-1,1} := \left(F_2 F_1 - \frac{[a_1]}{[a_1 + 1]} F_1 F_2 \right) v, \quad v_{1,0} := F_2 v, \quad v_{0,1} := F_2 F_1 F_2 v. \quad (6)$$

We take as a basis

$$\begin{aligned} F_1^k v_{0,0}, & \quad k = 0, \dots, a_1. \\ F_1^k v_{-1,1}, & \quad k = 0, \dots, a_1 - 1. \\ F_1^k v_{1,0}, & \quad k = 0, \dots, a_1 + 1. \\ F_1^k v_{0,1}, & \quad k = 0, \dots, a_1. \end{aligned} \quad (7)$$

To see that this is indeed a spanning set, we note that the action of F_1 and F_2 generate $\tilde{V}(a_1, a_2)$ and that together $F_2^2 = 0$ and $(q + q^{-1})F_1 F_2 F_1 = F_1^2 F_2 + F_2 F_1^2$ imply that any product in $U_h(\mathfrak{sl}(2|1))$ with three or more F_2 's is zero. They also imply that $F_2 F_1 F_2 F_1 = F_1 F_2 F_1 F_2$. Furthermore, $\langle h_1, E_1, F_1 \rangle \subset U_h(\mathfrak{sl}(2|1))$ gives a copy of $U_h(\mathfrak{sl}(2))$. As a topologically free finite rank $U_h(\mathfrak{sl}(2))$ -module, $\tilde{V}(a_1, a_2)$ decomposes as

$$\tilde{V}(a_1, a_2)[[h]] \simeq V_{a_1}[[h]] \oplus V_{a_1+1}[[h]] \oplus V_{a_1+1}[[h]] \oplus V_{a_1+2}[[h]] \quad (8)$$

where $V_n[[h]]$ is the topologically free $U_h(\mathfrak{sl}(2))$ module that deforms the unique simple n -dimensional $\mathfrak{sl}(2)$ -module V_n . Our choice of basis (7) is motivated by this decomposition, since each of the four $v_{\epsilon_1, \epsilon_2}$ is a highest weight vector for the induced $U_h(\mathfrak{sl}(2))$ -action. The subscripts track the h_1, h_2 weights, i.e. $h_i v_{\epsilon_1, \epsilon_2} = (a_i + \epsilon_i) v_{\epsilon_1, \epsilon_2}$.

h_1 and h_2 actions Let $k \geq 0$. The h_1 action is

$$\begin{aligned} h_1 \cdot F_1^k v_{0,0} &= (a_1 - 2k) F_1^k v_{0,0}, \\ h_1 \cdot F_1^k v_{-1,1} &= (a_1 - 2k - 1) F_1^k v_{-1,1}, \\ h_1 \cdot F_1^k v_{1,0} &= (a_1 - 2k + 1) F_1^k v_{1,0}, \\ h_1 \cdot F_1^k v_{0,1} &= (a_1 - 2k) F_1^k v_{0,1}. \end{aligned} \quad (9)$$

While the h_2 action is

$$\begin{aligned} h_2 \cdot F_1^k v_{0,0} &= (a_2 + k) F_1^k v_{0,0}, \\ h_2 \cdot F_1^k v_{-1,1} &= (a_2 + k + 1) F_1^k v_{-1,1}, \\ h_2 \cdot F_1^k v_{1,0} &= (a_2 + k) F_1^k v_{1,0}, \\ h_2 \cdot F_1^k v_{0,1} &= (a_2 + k + 1) F_1^k v_{0,1}. \end{aligned} \quad (10)$$

E_1 and E_2 actions Note $E_1 v_{0,0} = E_2 v_{0,0} = 0$ by assumption. Direct computation shows that $E_1 v_{1,0} = E_1 v_{0,1} = 0$, while $E_2 v_{1,0} = [a_2] v_{0,0}$, and $E_2 v_{0,1} = [a_2] F_1 v_{0,0} + [a_2 - 1] F_1 v_{1,0}$. Let $k \geq 1$. The E_1 action is

$$\begin{aligned} E_1 \cdot F_1^k v_{0,0} &= [k][a_1 + 1 - k] F_1^{k-1} v_{0,0}, \\ E_1 \cdot F_1^k v_{-1,1} &= [k][a_1 - k] F_1^{k-1} v_{-1,1}, \\ E_1 \cdot F_1^k v_{1,0} &= [k][a_1 + 2 - k] F_1^{k-1} v_{1,0}, \\ E_1 \cdot F_1^k v_{0,1} &= [k][a_1 + 1 - k] F_1^{k-1} v_{0,1}. \end{aligned} \tag{11}$$

The E_2 action is

$$\begin{aligned} E_2 \cdot F_1^k v_{0,0} &= 0, \\ E_2 \cdot F_1^k v_{-1,1} &= \left([a_2 + 1] - [a_2] \frac{[a_1]}{[a_1 + 1]} \right) F_1^{k+1} v_{0,0} \\ E_2 \cdot F_1^k v_{1,0} &= [a_2] F_1^k v_{0,0}, \\ E_2 \cdot F_1^k v_{0,1} &= [a_2] F_1^k F_2 F_1 v + [a_2 + 1] F_1^{k+1} v_{1,0} = [a_2] F_1^k v_{-1,1} + \left(\frac{[a_1][a_2]}{[a_1 + 1]} + [a_2 + 1] \right) F_1^{k+1} v_{1,0}. \end{aligned} \tag{12}$$

For $E_2 \cdot F_1^k v_{0,1}$, we've used that $F_2 F_1 v = v_{-1,1} + \frac{[a_1]}{[a_1 + 1]} F_1 v_{1,0}$.

F_1 and F_2 actions The F_1 action is simply $F_1 \cdot F_1^k v_{\epsilon_1, \epsilon_2} = F_1^{k+1} v_{\epsilon_1, \epsilon_2}$. It follows from the decomposition (8) that

$$F_1^{a_1+1} v_{0,0} = F_1^{a_1} v_{-1,1} = F_1^{a_1+2} v_{1,0} = F_1^{a_1+1} v_{0,1} = 0. \tag{13}$$

The F_2 action is more involved. By definition $F_2 \cdot v_{0,0} = v_{1,0}$, while a computation shows that $F_2 \cdot F_1 v_{0,0} = v_{-1,1} + \frac{[a_1]}{[a_1 + 1]} F_1 v_{1,0}$. We also have $F_2 \cdot v_{-1,1} = -\frac{[a_1]}{[a_1 + 1]} v_{0,1}$ and $F_2 v_{1,0} = 0$. In what follows P_n is a Laurent polynomial in q , determined recursively by $P_0 = 1$, $P_1 = q + q^{-1}$, and $P_n = (q + q^{-1})P_{n-1} - P_{n-2}$. The F_2 action on the rest of the generators is

$$\begin{aligned} F_2 \cdot F_1^k v_{0,0} &= (P_{k-1} F_1^{k-1} F_2 F_1 - P_{k-2} F_1^k F_2) v = P_{k-1} F_1^{k-1} v_{-1,1} + \left(P_{k-1} \frac{[a_1]}{[a_1 + 1]} - P_{k-2} \right) F_1^k v_{1,0} \quad k \geq 2 \\ F_2 \cdot F_1^k v_{-1,1} &= \left(P_{k-1} - \frac{[a_1]}{[a_1 + 1]} P_k \right) F_1^k v_{0,1} \quad k \geq 1 \\ F_2 \cdot F_1^k v_{1,0} &= P_{k-1} F_1^{k-1} v_{0,1}, \quad k \geq 1 \\ F_2 \cdot F_1^k v_{0,1} &= 0. \quad k \geq 0 \end{aligned}$$

2.1.2 The ribbon structure on $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$

In Section 2.2 we will need that $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ is a *ribbon category*, that is, a braided rigid monoidal category with a natural transformation $\theta : \text{id} \Rightarrow \text{id}$ such that

$$\theta_{V \otimes W} = (\theta_V \otimes \theta_W) c_{W,V} c_{V,W} \quad \text{and} \quad (\theta_V)^* = \theta_{V^*}. \tag{14}$$

We will describe the ribbon structure on $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$.

Since $U_h(\mathfrak{sl}(2|1))$ is a Hopf algebra, $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ is a monoidal category with unit $\mathbb{1}$ given by the trivial module $\mathbb{C}[[h]]$. Let $\tilde{V} = V[[h]]$ be an object of $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ where V is a finite dimensional $\mathfrak{sl}(2|1)$ -module. The duality is given on objects by $V[[h]]^* = V^*[[h]]$. The duality morphisms are $\mathbb{C}[[h]]$ -linearly extended from the underlying evaluation and coevaluation maps: let $\{v_i\}$ be a homogeneous basis of V and $\{v_i^*\}$ be the dual basis of $V^* = \text{Hom}_{\mathbb{C}}(V, \mathbb{C})$, that is $v_i^*(v_j) = \delta_{i,j}$. The following morphisms are defined by $\mathbb{C}[[h]]$ -linear extension to give a pivotal structure on $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$:

$$\begin{aligned} \overrightarrow{\text{coev}}_{\tilde{V}}: \mathbb{C}[[h]] &\rightarrow \tilde{V} \otimes \tilde{V}^*, \quad 1 \mapsto \sum v_i \otimes v_i^*, & \overrightarrow{\text{ev}}_{\tilde{V}}: \tilde{V}^* \otimes \tilde{V} &\rightarrow \mathbb{C}[[h]], \quad f \otimes x \mapsto f(x), \\ \overleftarrow{\text{coev}}_{\tilde{V}}: \mathbb{C}[[h]] &\rightarrow \tilde{V}^* \otimes \tilde{V}, \quad 1 \mapsto \sum (-1)^{\bar{v}_i} v_i^* \otimes q^{2h_2} v_i, & \overleftarrow{\text{ev}}_{\tilde{V}}: \tilde{V} \otimes \tilde{V}^* &\rightarrow \mathbb{C}[[h]], \quad x \otimes f \mapsto (-1)^{\bar{x} \bar{f}} f(q^{-2h_2} x). \end{aligned} \tag{15}$$

Here $x \in V$ and $f \in V^*$ are homogeneous elements.

Khoroshkin-Tolstoy [KT91] and Yamane [Yam94] showed that $U_h(\mathfrak{sl}(2|1))$ has an explicit R -matrix that decomposes as $R = \check{R}K$. Let $E' = E_1E_2 - q^{-1}E_2E_1$ and $F' = F_2F_1 - qF_1F_2$. Then we can write

$$\check{R} = \exp_q(\{1\}E_1 \otimes F_1) \exp_q(-\{1\}E' \otimes F') \exp_q(-\{1\}E_2 \otimes F_2), \quad (16)$$

$$K = q^{-h_1 \otimes h_2 - h_2 \otimes h_1 - 2h_2 \otimes h_2}. \quad (17)$$

Here we've used the standard notations $(k)_q := (1 - q^k)/(1 - q)$, $(n)_q! := (1)_q(2)_q \dots (n)_q$ and $\exp_q(x) := \sum_{n=0}^{\infty} x^n / (n)_q!$. By convention $(0)_q! = 1$. The identities $\exp_q(x) \exp_{q^{-1}}(-x) = 1$ and $\exp_{q^{-1}}(x) \exp_q(-x) = 1$ (see e.g. [KT91, (5.2)]) imply that

$$R^{-1} = K^{-1} \exp_{q^{-1}}(\{1\}E_2 \otimes F_2) \exp_{q^{-1}}(\{1\}E' \otimes F') \exp_{q^{-1}}(-\{1\}E_1 \otimes F_1). \quad (18)$$

Thus, the category $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ has a braiding $\{c_{\tilde{V}, \tilde{W}}\}_{\tilde{V}, \tilde{W} \in \text{Rep}_h^{tf} \mathfrak{sl}(2|1)}$:

$$c_{\tilde{V}, \tilde{W}} : \tilde{V} \otimes \tilde{W} \rightarrow \tilde{W} \otimes \tilde{V} \text{ given by } v \otimes w \mapsto \tau(R(v \otimes w))$$

where τ is the super flip map defined by $\tau(x \otimes y) = (-1)^{\bar{x}\bar{y}} y \otimes x$ for homogeneous x and y .

We finish by showing that the braiding and duality structure are compatible.

Proposition 2.5. *The above pivotal structure and braiding give a ribbon structure on $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$.*

It is well known that $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ is a ribbon category, see for example [Gee05]. We give a proof of this fact by describing the ribbon structure in terms of left/right dualities and a braiding. Our approach relies on [GP18, Theorem 9], where it is shown that a pivotal braided category is ribbon if it satisfies certain compatibility constraints for a subset of objects on the natural twist morphism defined from the braiding and dualities.

Proof. In order for [GP18, Theorem 9] to apply, $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ must be a generically \mathbb{C}/\mathbb{Z} -semisimple pivotal braided category. Having previously established it's pivotal and braided, it remains to prove the generic semisimplicity condition.

In the language of [GP18, §1.6], the category $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ is \mathbb{C}/\mathbb{Z} -graded as follows: for $g \in \mathbb{C}/\mathbb{Z}$ let $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)_g$ be the full subcategory of $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ consisting of those modules whose h_2 -weights are all equal to $g \bmod \mathbb{Z}$. We use the fact that h_2 acts semisimply on each X in $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$, decomposing it into h_2 -weight spaces. These weights differ by integers, see (10).

Next, we show that $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ is generically \mathbb{C}/\mathbb{Z} -semisimple, i.e. that the g -graded component $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)_g$ is semisimple whenever $g \in (\mathbb{C}/\mathbb{Z}) \setminus (\mathbb{Z}/\mathbb{Z})$. The word ‘‘generic’’ refers to the fact that the non-semisimple coset \mathbb{Z}/\mathbb{Z} is symmetric and small, i.e. $-\mathbb{Z}/\mathbb{Z} = \mathbb{Z}/\mathbb{Z}$ and for any $g_1, \dots, g_n \in \mathbb{C}/\mathbb{Z}$ we have $\bigcup_i (g_i + \mathbb{Z}/\mathbb{Z}) \neq \mathbb{C}/\mathbb{Z}$.

Let X be an object in $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)_g$, with $g \notin \mathbb{Z}/\mathbb{Z}$. There exists a highest weight vector $v \in X$. Since $g \notin \mathbb{Z}/\mathbb{Z}$, the h_1 and h_2 weights of v satisfy the inequalities of Section 2.1.1 and so v generates some typical module $\tilde{V}(a_1, a_2) \subseteq X$, where $a_2 \equiv g \bmod \mathbb{Z}$. Since typical modules are both projective and injective, $X \simeq \tilde{V}(a_1, a_2) \oplus W$ for some $W \in \text{Rep}_h^{tf} \mathfrak{sl}(2|1)_g$. Repeating this process on W we see that X is isomorphic to a direct sum of typical modules in $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)_g$. We conclude that $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)_g$ is semisimple whenever $g \notin \mathbb{Z}/\mathbb{Z}$. Hence $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ is a generically \mathbb{C}/\mathbb{Z} -semisimple pivotal braided category.

Next, define a family of natural automorphisms indexed by objects X of $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$:

$$\theta_X = (\text{Id}_X \otimes \overleftarrow{\text{ev}}_X)(c_{X, X} \otimes \text{Id}_{X^*})(\text{Id}_X \otimes \overrightarrow{\text{coev}}_X) : X \rightarrow X.$$

The left-most equality in (14) follows from naturality of the braiding and the Yang-Baxter equation. By [GP18, Theorem 9] the category $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ is ribbon and the natural transformation θ is called a *twist* if for all $X \in \text{Rep}_h^{tf} \mathfrak{sl}(2|1)$:

$$\theta_{X^*} = (\theta_X)^* := (\overrightarrow{\text{ev}}_X \otimes \text{Id}_{X^*})(\text{Id}_{X^*} \otimes \theta_X \otimes \text{Id}_{X^*})(\text{Id}_{X^*} \otimes \overrightarrow{\text{coev}}_X) \quad (19)$$

It is enough to check (19) on generic simple objects, i.e. the typical modules $\tilde{V}(a_1, a_2)$ with $a_2 \notin \mathbb{Z}$. Given an object X in $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$, by definition X^* is the $U_h(\mathfrak{sl}(2|1))$ -module whose action is given by $y.f(x) = (-1)^{\bar{y}\bar{f}} f(S(y)x)$ for $y \in U_h(\mathfrak{sl}(2|1))$, $x \in X$ and $f \in X^*$. It follows that the lowest weight vector v of $\tilde{V}(a_1, a_2)$ is dual to the highest weight vector v^* in $\tilde{V}(a_1, a_2)^*$ and has weight $(a_1, -a_1 - a_2 - 1)$. Thus $\tilde{V}(a_1, a_2)^* \simeq \tilde{V}(a_1, -a_1 - a_2 - 1)$. The morphisms $\theta_{\tilde{V}(a_1, a_2)}$ and $\theta_{\tilde{V}(a_1, a_2)^*}$ are determined by their values on the highest weight vectors. A direct computation shows that

$$\theta_{\tilde{V}(a_1, a_2)} = q^{-2a_2(a_1+a_2+1)} \text{Id}_{\tilde{V}(a_1, a_2)} \quad (20)$$

and

$$\theta_{\tilde{V}(a_1, a_2)^*} = q^{-2(-a_1-a_2-1)(1+a_1-a_1-a_2-1)} \text{Id}_{\tilde{V}(a_1, a_2)^*}. \quad (21)$$

The identity $(\overrightarrow{\text{ev}}_X \otimes \text{Id}_{X^*})(\text{Id}_{X^*} \otimes \overrightarrow{\text{coev}}_X) = \text{Id}_{X^*}$ implies that the right side of (19) with $X = \tilde{V}(a_1, a_2)$ is equal to $q^{-2a_2(a_1+a_2+1)} \text{Id}_{\tilde{V}(a_1, a_2)}$. It follows that (19) holds for all simple modules in $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)_g$ with $g \in \mathbb{C}/\mathbb{Z} \setminus \mathbb{Z}/\mathbb{Z}$, thus proving the proposition. \square

2.2 Quantum Invariants from $\mathfrak{sl}(2|1)$

We established in Section 2.1.2 that $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ is a ribbon category. It is therefore possible to use the frameworks laid out in [RT90, GPT09] to define knot invariants with $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ as the algebraic input. We will not give a full treatment of this theory, and instead defer to the aforementioned articles. A detailed treatment of the semisimple case can be found in [Tur10].

We consider framed oriented tangles whose components are colored by objects of $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$. Such tangles are called $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ -colored ribbons. Let $\mathcal{R}_{\text{Rep}_h^{tf} \mathfrak{sl}(2|1)}$ denote the category of $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ -colored ribbons (for details see [Tur10]). The well-known Reshetikhin-Turaev construction defines a $\mathbb{C}[[\hbar]]$ -linear functor

$$F : \mathcal{R}_{\text{Rep}_h^{tf} \mathfrak{sl}(2|1)} \rightarrow \text{Rep}_h^{tf} \mathfrak{sl}(2|1).$$

The value of any $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ -colored ribbon under F can be computed using the six *building blocks*, which are the morphisms $\overrightarrow{\times}, \overleftarrow{\times}, \smile, \frown, \curvearrowright, \curvearrowleft$ with arbitrary colors in $\mathcal{R}_{\text{Rep}_h^{tf} \mathfrak{sl}(2|1)}$. The functor F transforms these building blocks as follows:

$$\begin{aligned} F(\overrightarrow{\times}) &= c, & F(\smile) &= \overrightarrow{\text{coev}}, & F(\frown) &= \overleftarrow{\text{coev}}, \\ F(\overleftarrow{\times}) &= c^{-1}, & F(\curvearrowright) &= \overrightarrow{\text{ev}}, & F(\curvearrowleft) &= \overleftarrow{\text{ev}}. \end{aligned} \quad (22)$$

We've suppressed the colors, which determine the components of the braiding and duality transformations. Vertical lines are sent to the identity morphism and reversing the direction of an arrow is equivalent to coloring instead by the dual module.

Let L be a link with some component labeled by a simple object $V \in \text{Rep}_h^{tf} \mathfrak{sl}(2|1)$. By cutting this component we obtain a tangle T_V whose two ends are labeled with V . By definition $F(T_V) \in \text{End}_{\text{Rep}_h^{tf} \mathfrak{sl}(2|1)}(V)$. Since V is simple, this endomorphism is the product of the identity $\text{Id}_V : V \rightarrow V$ with an element $\langle T_V \rangle$ of the ground ring of $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$, i.e. $F(T_V) = \langle T_V \rangle \text{Id}_V$. In particular,

$$\begin{aligned} F(L) &= F\left(\begin{array}{c} \text{---} \\ \boxed{T} \\ \text{---} \end{array} \circlearrowright V\right) = \langle T_V \rangle F\left(\begin{array}{c} \text{---} \\ \boxed{\text{Id}_V} \\ \text{---} \end{array} \circlearrowright V\right) \\ &= \langle T_V \rangle F\left(\begin{array}{c} \text{---} \\ \bigcirc \\ \text{---} \end{array} \circlearrowright V\right) = \langle T_V \rangle (\overleftarrow{\text{ev}}_V \circ \overrightarrow{\text{coev}}_V). \end{aligned} \quad (23)$$

The quantity $\text{qdim}_{\text{Rep}_h^{tf} \mathfrak{sl}(2|1)}(V) := \overleftarrow{\text{ev}}_V \circ \overrightarrow{\text{coev}}_V$ is called the *quantum dimension* of V . The quantum dimension vanishes when $V = \tilde{V}(a_1, a_2)$ is a typical representation:

$$\text{qdim}_{\text{Rep}_h^{tf} \mathfrak{sl}(2|1)}(\tilde{V}(a_1, a_2)) := (\overleftarrow{\text{ev}}_{\tilde{V}(a_1, a_2)} \circ \overrightarrow{\text{coev}}_{\tilde{V}(a_1, a_2)}) = 0.$$

See [GP07] for details, including a proof of the vanishing quantum dimension. Thus, from (23) we have that $F(L) = 0$ if any component of L is colored by a typical module $\tilde{V}(a_1, a_2)$.

In [GP07] it is shown that one can replace such a vanishing quantum dimension in (23) with a certain non-zero scalar and obtain a well defined invariant of links. This process was extended to a general theory in [GPT09]. We will briefly recall this construction.

Consider the function d from the set of typical modules to $\mathbb{C}[[h]]$ given by

$$d(\tilde{V}(a_1, a_2)_{\bar{p}}) = \frac{\{a_1 + 1\}}{\{1\}\{a_2\}\{a_2 + a_1 + 1\}} \quad (24)$$

This function is called the *modified dimension*, because it replaces the quantum dimension in the construction of link invariants. It can be derived from F itself applied to open Hopf links, see [GPT09, §2]. Let L be a $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ -colored framed link with at least one component colored by a typical module $\tilde{V}(a_1, a_2)$. Cutting this component as above, we obtain a tangle $T_{\tilde{V}(a_1, a_2)}$. By [GPT09, Prop. 27], the assignment

$$L \mapsto F'(L) := d(\tilde{V}(a_1, a_2)) \langle T_{\tilde{V}(a_1, a_2)} \rangle \quad (25)$$

yields a well-defined isotopy invariant of L , independent of the choice of cut component. Although (24) may seem ad-hoc, the invariant (25) is unique up to a global scalar. See the discussion after [GPT09, Prop. 27].

In the remainder of this paper we assume that *every* strand of an n -component link L is colored by some typical module $\tilde{V}(a_{1i}, a_{2i})$, for $i = 1, \dots, n$. It does not matter which strand is cut, and without loss of generality we choose it to be the one labeled by $\tilde{V}(a_{11}, a_{21})$. The corresponding invariant (25) defines a function

$$F'(L) : (\mathbb{Z}_{\geq 0} \times \mathbb{C}^*)^n \rightarrow \mathbb{C}[[h]], \quad F'(L)(a_{11}, a_{21}, \dots, a_{1n}, a_{2n}) = d(\tilde{V}(a_{11}, a_{21})) \langle T_{\tilde{V}(a_{11}, a_{21})} \rangle. \quad (26)$$

Establishing the q -holonomicity of this family of functions is the main result of this paper.

3 q-Holonomic systems

We give a brief review of q -holonomic systems and recommend [Sab93, GL16] for more details. Note that we work in the h -adic setting, i.e. over $\mathbb{C}[[h]]$ instead of $\mathbb{C}(q)$, following [KS11].

A prerequisite structure for the formation of a q -holonomic system in this framework is an action of a quantum torus. The standard story is given by the rank n q -Weyl algebra, which has the presentation

$$\mathbb{W}_n := \langle L_1, \dots, L_n, M_1, \dots, M_n \mid L_i M_j = q^{\delta_{ij}} M_j L_i \rangle \quad (27)$$

with the M 's and L 's commuting between themselves. This has a standard action on the vector space of functions $\{\mathbb{Z}^n \rightarrow \mathbb{C}[[h]]\}$, given by

$$L_i f(a_1, \dots, a_n) = f(a_1, \dots, a_i + 1, \dots, a_n), \quad M_i f(a_1, \dots, a_n) = q^{a_i} f(a_1, \dots, a_n). \quad (28)$$

Where $q = e^{h/2}$. Sometimes it is technically better to work with its localization \mathbb{E}_n , in which the generators are invertible. The close relationship between q -holonomicity of \mathbb{W}_n and \mathbb{E}_n modules is spelled out in [Sab93].

Definition 3.1. [Sab93, Prop. 1.5.2] *Let N be a finitely generated module for a q -Weyl algebra \mathbb{W}_n . The homological dimension of N is the degree of its Hilbert polynomial h_N with respect to any good filtration:*

$$\dim N := \deg h_N. \quad (29)$$

Note that \mathbb{W}_n itself has homological dimension $2n$, while the trivial module 0 has homological dimension 0 . A fundamental result in the theory of D -modules and their q -difference analogs is *Bernstein's Inequality*, which states that either $\text{rank } \mathbb{W}_n \leq \dim M \leq 2 \text{rank } \mathbb{W}_n$ or $\dim M = 0$, in which case $M = 0$.

Definition 3.2. *A finitely generated \mathbb{W}_n -module N is called **q-holonomic** when its homological dimension is as low as possible, i.e.*

$$\dim N = n \text{ or } 0. \quad (30)$$

We will frequently consider cyclic modules $\mathbb{W}_n f(m_1, \dots, m_n)$. In this setting lower homological dimension means more linear relations between the functions Pf , for $P \in \mathbb{W}_n$. These relations can be interpreted as exhibiting some quasi-periodicity of the function. When $n = 1$ a single non-trivial relation $Pf = 0$ implies q -holonomicity, but when $n > 1$ it is generally not enough to have n distinct operators annihilate f . The following lemma is useful for establishing q -holonomicity of elementary functions:

Lemma 3.3. [BDGG20, 3.5] *Let $\mathbb{W}_n f$ be a cyclic module whose annihilation ideal contains elements of the form $p_j(M)L_j^{a_j} + q_j(M)$ for $j = 1, \dots, n$, where $p_j, q_j \neq 0$ are independent of the L_j and $d_j \geq 1$. Then $\mathbb{W}_n f$ is q -holonomic.*

3.1 Non-discrete variables

We use the same framework as [BDGG20] to work with complex variables. Suppose we have n complex variables $\alpha_1, \dots, \alpha_n$ and m discrete (\mathbb{N} or \mathbb{Z} -valued) ones a_1, \dots, a_m . Let \mathbb{V}_n be the field of rational functions over $\mathbb{C}[[h]]$, in the variables $\{x_i \mid i = 1, \dots, n\}$ and $\{z_{ij} \mid i, j = 1, \dots, n\}$. In this way we work formally with the linear and quadratic q -exponentials q^{α_i} and $q^{\alpha_i \alpha_j}$. Let

$$\mathcal{V}_{m,n} := \{f : \mathbb{Z}^m \rightarrow \mathbb{V}_n\}. \quad (31)$$

This has an action of \mathbb{W}_{n+m} . For the discrete variables it's given by (28). For the complex variables, we reproduce this action formally on the codomain \mathbb{V}_n . Each M_i is multiplication by x_i , while the shifts become

$$L_i x_j = q^{\delta_{ij}} x_j \quad L_i z_{jk} = q^{\delta_{ij} \delta_{ik}} x_j^{\delta_{ik}} x_k^{\delta_{ij}} z_{jk}. \quad (32)$$

In practice it's useful to think of $\mathcal{V}_{m,n}$ as functions $\mathbb{Z}^m \times \mathbb{C}^n \rightarrow \mathbb{C}[[h]]$.

3.2 Examples of q -holonomic functions

We present some of the functions which show up in the expression of the $\mathfrak{sl}(2|1)$ quantum invariant. These are all previously known to be q -holonomic.

Linear and quadratic powers Functions of the form $q^{n_i^2}$, $q^{n_i n_j}$, and q^{n_i} are q -holonomic. In [BDGG20] it's shown that the formal analogs z_{ii} , z_{ij} , and x_i are likewise q -holonomic.

q -Pochhammer The function $n_1, n_2 \mapsto (q^{n_1}; q)_{n_2}$ is q -holonomic by Lemma 3.3, since its annihilation ideal contains $(1 - M_1)L_1 - 1 + M_1 M_2$ and $L_2 + M_1 M_2 - 1$.

Inverse q -Pochhammer The function $n_1, n_2 \mapsto 1/(q^{n_1}; q)_{n_2}$ is q -holonomic by Lemma 3.3. Its annihilation ideal contains $(1 - M_1 M_2)L_1 - M_1 + 1$ and $(1 - M_1 M_2)L_2 - 1$.

Inverse q -numbers The expression $n \mapsto 1/\{n\}$ is q -holonomic. We can see this by rewriting it as a product of q -holonomic functions:

$$\frac{1}{\{n\}} := \frac{1}{q^n - q^{-n}} = \frac{q^{-n}}{(q^{-2n}; q)_1} \quad (33)$$

then invoking the closure properties discussed in the next section.

Indicator functions The function

$$\vartheta_{[n_1, n_2]}(n_3) := \begin{cases} 1 & n_1 \leq n_3 \leq n_2 \\ 0 & \text{otherwise} \end{cases} \quad (34)$$

is annihilated by $(M_3 - M_1)(L_1 - 1)$, $(M_3 - qM_2)(L_2 - 1)$, and $(M_3 - M_1)(M_3 - q^{-1}M_1)(L_3 - 1)$. It follows from Lemma 3.3 that it is q -holonomic.

3.3 Closure properties

The class of q -holonomic functions enjoy many closure properties, as detailed in [GL16, 5.2,5.3] and extended to non-discrete variables in [BDGG20, 3.2]. The crucial ones for us here are addition $f+g$, multiplication fg , specialization $f|_{k_i=\lambda}$, extension $f(k_1, \dots, k_n, k_{n+1}) := f(k_1, \dots, k_n)$, affine substitution $h(\bar{k}) := f(A\bar{k} + \bar{b})$, and multisums $h(k_1, \dots, k_{n-1}, a, b) := \sum_{k_n=a}^b f(k_1, \dots, k_n)$. We will often use these implicitly.

Lemma 3.4. *Let X, Y be parameterized matrices whose coefficients are q -holonomic in both their coordinates (i.e. row and column position) and the parameters. Then the same holds for $X + Y$, $X \oplus Y$, $X \otimes Y$, and XY .*

Proof. Let X be an $n_1 \times n_2$ matrix and Y an $m_1 \times m_2$ matrix. For matrix sum ($n_1 = m_1, n_2 = m_2$) the coefficients are added. For direct sum, the new coefficients are

$$(X \oplus Y)_i^j = X_i^j \vartheta_{[1, n_1]}(i) \vartheta_{[1, n_2]}(j) + Y_i^j \vartheta_{[n_1+1, n_1+m_1]}(i) \vartheta_{[n_2+1, n_2+m_2]}(j). \quad (35)$$

For tensor product the relevant formula is

$$(X \otimes Y)_{i_1, i_2}^{j_1, j_2} = X_{i_1}^{j_1} Y_{i_2}^{j_2}, \quad (36)$$

while for matrix product ($n_2 = m_1$) the coefficients are given by

$$(XY)_i^j = \sum_{k=1}^{n_2} X_i^k Y_k^j. \quad (37)$$

In each case the new coefficients can be written in terms of sums, products, and multisums of the X_i^j and Y_i^j , together with indicator functions. Since these operations preserve q -holonomicity, we conclude that for the four operations considered here, the new coefficients are q -holonomic whenever the original ones are. \square

3.4 q -Holonomic systems and ribbon categories

Reshetikhin-Turaev style knot invariants, such as those considered in this paper, depend heavily on the choice of a ribbon category. It's therefore reasonable to think of their q -holonomicity as a property of this underlying category. We take a small step in that direction with the following definition and lemma.

Definition 3.5. *Fix an \mathbb{D} -linear¹ ribbon category $(\mathcal{C}, c, \overleftarrow{\text{ev}}, \overrightarrow{\text{ev}}, \overleftarrow{\text{coev}}, \overrightarrow{\text{coev}}, \theta)$ and a collection $S_\Lambda := \{V_\lambda\}_{\lambda \in \Lambda}$ of its objects. Here \mathbb{D} is an integral domain. We require that S_Λ is closed under taking duals, i.e. $V_\lambda^* \in S_\Lambda$ for all $\lambda \in \Lambda$. Suppose that $\Lambda \hookrightarrow \mathbb{Z}^m \times \mathbb{C}^n$ has codimension zero², so that \mathbb{D} -valued functions on $\Lambda \times \mathbb{N}^k$ can be identified almost-everywhere with elements in $\mathcal{V}_{m+k, n}$. We will call the ribbon structure **q -holonomic over S_Λ** if the following functions exist and are q -holonomic in the sense of [Sab93, BDGG20]:*

dimension: $\dim_{\mathbb{D}} : \Lambda \rightarrow \mathbb{N}$, given by $\lambda \mapsto \dim_{\mathbb{D}}(V_\lambda)$.

modified dimension: $\mathbf{d} : \Lambda \rightarrow \mathbb{D}$ given by $\lambda \mapsto \mathbf{d}(V_\lambda)$. Here we assume that the tensor ideal used to define the modified dimension contains S_Λ . If \mathcal{C} is semisimple, this ideal can be taken to be the full category, in which case modified dimension agrees with the usual quantum dimension of objects.

braiding: $\Lambda^2 \times \mathbb{N}^2 \rightarrow \mathbb{D}$, given by $(\lambda_1, \lambda_2, i, j) \mapsto (c_{V_{\lambda_1}, V_{\lambda_2}})_i^j$, where the morphism $c_{V_{\lambda_1}, V_{\lambda_2}} : V_{\lambda_1} \otimes V_{\lambda_2} \rightarrow V_{\lambda_2} \otimes V_{\lambda_1}$ is written in terms of matrix coefficients, with $i = 1, \dots, \dim_{\mathbb{D}}(V_{\lambda_1} \otimes V_{\lambda_2})$ and $j = 1, \dots, \dim_{\mathbb{D}}(V_{\lambda_2} \otimes V_{\lambda_1})$. By convention, we extend the row and column indices to \mathbb{N} by zero.

inverse braiding: As above, based instead on the inverse braiding $\Lambda^2 \times \mathbb{N}^2 \rightarrow \mathbb{D}$ given by $(\lambda_1, \lambda_2, i, j) \mapsto (c_{V_{\lambda_1}, V_{\lambda_2}}^{-1})_i^j$.

¹If \mathbb{D} is only an integral domain (e.g. $\mathbb{C}[[\hbar]]$) and not a field then we require that hom-spaces are finite rank free \mathbb{D} -modules, so that we can still (non-canonically) identify morphisms as matrices. In this case $\dim_{\mathbb{D}}$ means rank as a free \mathbb{D} -module. The main category considered in this paper, $\text{Rep}_\hbar^{tf} \mathfrak{sl}(2|1)$, meets this requirement.

²i.e. Λ is the complement of a hyperplane arrangement.

rigidity: Here we consider four functions $\Lambda \times \mathbb{N} \rightarrow \mathbb{D}$:

$$\begin{aligned} (\lambda, i) &\mapsto \left(\overrightarrow{\text{coev}}_{V_\lambda}\right)_i^i, & (\lambda, i) &\mapsto \left(\overrightarrow{\text{ev}}_{V_\lambda}\right)_i, \\ (\lambda, i) &\mapsto \left(\overleftarrow{\text{coev}}_{V_\lambda}\right)_i^i, & (\lambda, i) &\mapsto \left(\overleftarrow{\text{ev}}_{V_\lambda}\right)_i. \end{aligned}$$

In all cases $i = 1, \dots, \dim_{\mathbb{D}}(V \otimes V^*)$ is extended to \mathbb{N} by zero.

Lemma 3.6. *Let \mathcal{C} be a ribbon category. If \mathcal{C} 's ribbon structure is q -holonomic over a collection $S_\Lambda := \{V_\lambda\}_{\lambda \in \Lambda}$ of simple objects then the corresponding quantum invariants will be q -holonomic in the sense of [Sab93, BDGG20].*

Proof. Let \mathcal{D}_λ be a diagram of a link L , whose components are colored by simple objects $V_{\lambda_1}, \dots, V_{\lambda_\ell} \in S_\Lambda$, and let \mathcal{T}_λ be the same diagram, cut along some arc and turned into a $(1,1)$ -tangle. Here we use the shorthand $\lambda = (\lambda_1, \dots, \lambda_\ell)$. Without loss of generality, we suppose the open ends of the tangle are colored by V_{λ_1} . As a morphism in $\mathcal{R}_{\mathcal{C}}$, the tangle \mathcal{T}_λ can be decomposed into the composition of tensor products of the six basic building blocks: $\nearrow, \searrow, \smile, \frown, \cup, \cap$ and identity morphisms on the various V_{λ_i} .

By assumption the evaluation functor $\mathcal{R}_{\mathcal{C}} \rightarrow \mathcal{C}$ used to define the quantum invariant sends each of these building blocks to a q -holonomic function. Since each V_{λ_i} is finite dimensional and the hom-spaces are free \mathbb{D} -modules, each of these can be understood as a matrix. Since Lemma 3.4 shows that q -holonomicity is preserved by tensor product and composition (which becomes matrix multiplication), it follows that the function $\lambda_1, \dots, \lambda_\ell \mapsto \langle \mathcal{T}_\lambda \rangle \in \text{End}_{\mathcal{C}}(V_{\lambda_1}) \simeq \mathbb{D}$ is q -holonomic. The quantum knot invariant is the product of this scalar and the modified dimension, and therefore q -holonomic. \square

We wrap up this section with a conjecture regarding the expected generalization of our main result.

Conjecture 3.7. *Let \mathfrak{g} be a contragredient Lie superalgebra, meaning it's simple, finite dimensional, basic³, and its even subalgebra is reductive. Let $U_h \mathfrak{g}$ be its quantized universal enveloping algebra, and $\text{Rep}_h^{tf} \mathfrak{g}$ the category of topologically free finite rank $U_h \mathfrak{g}$ modules. (See Definition 2.4.) Let S_Λ denote the set of representations whose underlying \mathfrak{g} representation is typical [Kac78, §4].*

We conjecture that for any knot K the quantum knot invariant $L_K : \Lambda \rightarrow \mathbb{C}[[h]]$ built from S_Λ using the ribbon structure [KT91] and the modified dimension associated to any tensor ideal containing S_Λ are q -holonomic.

The proof of this conjecture would follow the same lines as our work here, with significantly more involved expressions for the action of the generators on the typical representations.

4 The $\mathfrak{sl}(2|1)$ invariant is q -holonomic

Lemma 4.1. *The matrix coefficients of elements in $\langle E_1, E_2, F_1, F_2 \rangle$ inside $U_h(\mathfrak{sl}(2|1))$ in each $\text{End}(\tilde{V}(a_1, a_2))$ form q -holonomic systems.*

Let $\rho_{a_1, a_2} : U_h(\mathfrak{sl}(2|1)) \rightarrow \text{End}_{\mathbb{C}[[h]]}(\tilde{V}(a_1, a_2))$ be a typical representation. The claim is that the functions $\mathbb{N}^3 \rightarrow \mathbb{C}[[h]](x_1, z_{11})[a_2, a_2^{-1}]$ defined by $(a_1, k, \ell) \mapsto \rho_{a_1, a_2}(X)_k^\ell$ are q -holonomic for any $X \in \langle E_1, E_2, F_1, F_2 \rangle$. Here x_1 and z_{11} play the part of the formal expressions q^{a_2} and $q^{a_2^2}$, as outlined in Section 3.1, $k, \ell = 0, \dots, 4a_1 + 3$ run over the basis elements (7), and $\rho_{a_1, a_2}(X)w_k = \sum_{\ell=0}^{4a_1+3} \rho_{a_1, a_2}(X)_k^\ell w_\ell$. The following proof relies heavily on the many closure properties of q -holonomic functions, see Section 3.3.

Proof. We will make use of the decomposition (8) of $\tilde{V}(a_1, a_2)$ into four $U_h(\mathfrak{sl}(2))$ -representations. In each of the four blocks, our basis is of the form $F_1^k v_{\epsilon_1, \epsilon_2}$ for $\epsilon_1, \epsilon_2 \in \{-1, 0, 1\}$ and $k = 0, \dots, a_1 + \epsilon_1$. For generators which respect the decomposition, q -holonomicity within the blocks implies it for the whole representation, since we build the full expressions from the $\mathfrak{sl}(2)$ ones by shifting their indices, multiplying them by the appropriate indicator functions, and adding the results.

³Basic here means it's not one of the strange Lie superalgebras $\mathfrak{p}(n)$ or $\mathfrak{q}(n)$, for which the classification of typical representations [Kac78, Theorem 1] doesn't apply.

$\boxed{F_1}$ Within each $U_h(\mathfrak{sl}(2))$ -subrepresentation, $\rho(F_1)_k^\ell$ has the form

$$(a_1, a_2, k, \ell) \mapsto \delta_{k+1}^\ell \vartheta_{[0, a_1 + \epsilon]}(k) \quad (38)$$

where ϑ is the indicator function. Since indicator functions and their products are q-holonomic, we conclude that F_1 's coefficients in each typical representation are q-holonomic functions.

$\boxed{E_1}$ We again work in the decomposition (8). In each block, $\rho(E_1)_k^\ell$ has the form

$$(a_1, a_2, k, \ell) \mapsto [a_1 + \epsilon + 1 - k] \delta_{k-1}^\ell \vartheta_{[0, a_1 + \epsilon]}(k) \quad (39)$$

By closure under affine substitution, the expression $[n] = \{n\}/\{1\}$ sends linear functions to q-holonomic ones. We conclude that the coefficients of E_1 are q-holonomic in each typical representation.

$\boxed{F_2}$ The action of this generator does not respect the decomposition (8). To describe its coefficients we put a total ordering on the generators, first the $F_1^k v_{0,0}$, then the $F_1^k v_{-1,1}$, then the $F_1^k v_{1,0}$, and finally the $F_1^k v_{0,1}$. We will temporarily refer to the ℓ^{th} element in this order as w_ℓ , so that $w_0 = v_{0,0}$, $w_{a_1+1} = v_{-1,1}$, $w_{2a_1+1} = v_{1,0}$, and $w_{3a_1+3} = v_{0,1}$.

The coefficients of F_2 on a typical representation $\tilde{V}(a_1, a_2)$ are defined by the piecewise function

$$(a_1, a_2, k, \ell) \mapsto \begin{cases} P_{k-1} \delta_{a_1+k}^\ell + \left(P_{k-1} \frac{[a_1]}{[a_1+1]} - P_{k-2} \right) \delta_{2a_1+k+1}^\ell & 0 \leq k \leq a_1 \\ \left(P_{k-a_1-2} - \frac{[a_1]}{[a_1+1]} P_{k-a_1-1} \right) \delta_{2a_1+2+k}^\ell & a_1 + 1 \leq k \leq 2a_1 \\ P_{k-2a_1-2} \delta_{a_1+1+k}^\ell & 2a_1 + 1 \leq k \leq 3a_1 + 2 \\ 0 & 3a_1 + 3 \leq k \leq 4a_1 + 3 \\ 0 & \text{otherwise.} \end{cases} \quad (40)$$

Since a piecewise q-holonomic function is likewise q-holonomic, we conclude that the coefficients of F_2 are q-holonomic.

$\boxed{E_2}$ This generator does not respect the decomposition (8). We reuse the basis $\{w_\ell\}$ defined in the F_2 section of this proof. The coefficients of E_2 on a typical representation $\tilde{V}(a_1, a_2)$ are

$$(a_1, a_2, k, \ell) \mapsto \begin{cases} 0 & 0 \leq k \leq a_1 \\ \left([a_2 + 1] - \frac{[a_1][a_2]}{[a_1+1]} \right) \delta_{k-a_1}^\ell & a_1 + 1 \leq k \leq 2a_1 \\ [a_2] \delta_{k-2a_1}^\ell & 2a_1 + 1 \leq k \leq 3a_1 + 2 \\ [a_2] \delta_{k-2a_1-2}^\ell + \left(\frac{[a_1][a_2]}{[a_1+1]} + [a_2 + 1] \right) \delta_{k-a_1-2}^\ell & 3a_1 + 3 \leq k \leq 4a_1 + 3 \\ 0 & \text{otherwise.} \end{cases} \quad (41)$$

We conclude that the actions of the generators are in terms of q-holonomic functions. Finally, by Lemma 3.4, every element in $U_h(\mathfrak{sl}(2|1))$ acts by a matrix whose coefficients form a q-holonomic system. \square

Lemma 4.2. *The R-Matrix given by the product of (16) and (17) has q-holonomic coefficients, as does its inverse (18).*

Our claim is that the function $\mathbb{N}^6 \rightarrow \mathbb{V}_2$, sending

$$(a_1, k_a, \ell_a, b_1, k_b, \ell_b) \mapsto (R)_{k_a, k_b}^{\ell_a, \ell_b} \in \mathbb{C}[[h]](q^{a_2}, q^{b_2}, q^{a_2^2}, q^{b_2^2}, q^{a_2 b_2}) \quad (42)$$

defined by the action of R on $\tilde{V}(a_1, a_2) \otimes \tilde{V}(b_1, b_2)$ is q-holonomic. Since the functions coming from the R-matrix and its inverse differ only by negative signs in exponential expressions, it's sufficient to show the result just for R .

Proof. We consider the four factors $\exp_q(-\{1\}E_2 \otimes F_2)$, $\exp_q(-\{1\}E' \otimes F')$, $\exp_q(\{1\}E_1 \otimes F_1)$, and K . First, note that

$$\exp_q(-\{1\}E_2 \otimes F_2) = 1 - \{1\}E_2 \otimes F_2 \quad \text{and} \quad \exp_q(-\{1\}E' \otimes F') = 1 + -\{1\}E' \otimes F'. \quad (43)$$

It follows from Lemma 4.1 and Lemma 3.4 that these act by q-holonomic functions. For the $E_1 \otimes F_1$ factor, the relevant functions come from the expressions

$$E_1^n \cdot F_1^k v_{\epsilon_1, \epsilon_2} = \begin{cases} 0 & 1 \leq k \leq n \\ \frac{[k]!}{[k-n]!} \frac{[a_1 + \epsilon_1 - (k+1) + n]!}{[a_1 + \epsilon_1 - (k+1)]!} F_1^{k-n} v_{\epsilon_1, \epsilon_2} & \text{otherwise} \end{cases} \quad (44)$$

Note that $1/[m]! = q^{m(m+1)/2} \{1\}^m (q^2; q^2)_m$ is q-holonomic for m any discrete variable or linear combination thereof and $F_1^n \cdot F_1^k v_{\epsilon_1, \epsilon_2} = F_1^{n+k} v_{\epsilon_1, \epsilon_2}$. We conclude that the coefficients

$$(a_1, k_a, \ell_a, b_1, k_b, \ell_b, n) \mapsto (E_1^n \otimes F_1^n)_{k_a, k_b}^{\ell_a, \ell_b} \in \mathbb{V}_2 \quad (45)$$

defined by the action on $\tilde{V}(a_1, a_2) \otimes \tilde{V}(b_1, b_2)$ are q-holonomic. Since q-holonomicity is preserved by multisums and $1/(n)_q!$ is q-holonomic, we conclude that $\exp_q(\{1\}E_1 \otimes F_1)$ is q-holonomic.

Finally, $K = q^{-h_1 \otimes h_2 - h_2 \otimes h_1 - 2h_2 \otimes h_2}$ acts on $\tilde{V}(a_1, a_2) \otimes \tilde{V}(b_1, b_2)$ by

$$(a_1, k_a, \ell_a, b_1, k_b, \ell_b) \mapsto q^{-a_1 b_2 - a_2 b_1 - 2a_2 b_2} \delta_{k_a}^{\ell_a} \delta_{k_b}^{\ell_b} \in \mathbb{V}_2. \quad (46)$$

These are a combination of quadratic exponentials (in mixed discrete and continuous variables) together with indicator functions. It follows that the coefficients of K in any typical representation induce q-holonomic systems.

We've shown that each of the four factors of the R-Matrix is q-holonomic. It follows from Lemma 3.4 that the R-Matrix has q-holonomic coefficients in each typical representation. \square

Lemma 4.3. *The evaluation and coevaluation induce q-holonomic systems.*

We are claiming that the following four functions $\mathbb{N}^3 \rightarrow \mathbb{V}_1$ are q-holonomic:

$$\begin{aligned} (a_1, \ell_1, \ell_2) &\mapsto \left(\overrightarrow{\text{coev}}_{\tilde{V}(a_1, a_2)} \right)^{\ell_1, \ell_2} & (a_1, \ell_1, \ell_2) &\mapsto \left(\overrightarrow{\text{ev}}_{\tilde{V}(a_1, a_2)} \right)_{k_1, k_2} \\ (a_1, \ell_1, \ell_2) &\mapsto \left(\overleftarrow{\text{coev}}_{\tilde{V}(a_1, a_2)} \right)^{\ell_1, \ell_2} & (a_1, k_1, k_2) &\mapsto \left(\overleftarrow{\text{ev}}_{\tilde{V}(a_1, a_2)} \right)_{k_1, k_2} \end{aligned}$$

Proof. The functions in question are

$$\begin{aligned} (a_1, \ell_1, \ell_2) &\mapsto \delta^{\ell_1, \ell_2} & (a_1, k_1, k_2) &\mapsto \delta_{k_1, k_2} \\ (a_1, \ell_1, \ell_2) &\mapsto (-1)^{\overline{w}_{\ell_1}} q^{2\phi(\ell_1)} \delta^{\ell_1, \ell_2} & (a_1, k_1, k_2) &\mapsto (-1)^{\overline{w}_{k_1} \overline{w}_{k_2}} q^{-2\phi(k_1)} \delta_{k_1, k_2} \end{aligned}$$

where $\phi(\ell)$ is the h_2 -weight of the basis vector w_ℓ defined in the proof of Lemma 4.1. It's a piecewise linear function in a_1, a_2 , and ℓ . Each expression is a product of standard q-holonomic functions and therefore q-holonomic. \square

Lemma 4.4. *The modified dimension for typical modules in $\text{Rep}_h^{tf} \mathfrak{sl}(2|1)$ is q-holonomic*

Proof. The expression (24) is a product of terms $\{n\}$ and $1/\{n\}$, modified by affine substitution in continuous and discrete variables. It is therefore q-holonomic. \square

Corollary 4.5. *The $\mathfrak{sl}(2|1)$ quantum invariant is q-holonomic.*

Proof. This follows from Lemma 3.6, which is applicable following Lemmas 4.2, 4.3, and 4.4. \square

References

- [AFT17] David Ayala, John Francis, and Hiro Lee Tanaka. Factorization homology of stratified spaces. *Selecta Mathematica*, 23(1):293–362, January 2017.
- [AGP21] Cristina Ana-Maria Anghel, Nathan Geer, and Bertrand Patureau-Mirand. Relative (pre)-modular categories from special linear Lie superalgebras. *Journal of Algebra*, 586:479–525, November 2021.
- [AGPS18] N. Aghaei, A. M. Gainutdinov, M. Pawelkiewicz, and V. Schomerus. Combinatorial quantisation of $GL(1-1)$ Chern-Simons theory I: The torus. *arXiv*, (arXiv:1811.09123), November 2018. arXiv:1811.09123 [hep-th, physics:math-ph].
- [BDGG20] Jennifer Brown, Tudor Dimofte, Stavros Garoufalidis, and Nathan Geer. The ADO invariants are a q-holonomic family. *arXiv*, May 2020. arXiv: 2005.08176.
- [BW11] Francis Bonahon and Helen Wong. Quantum traces for representations of surface groups in $SL(2, \mathbb{C})$. *Geometry & Topology*, 15(3):1569–1615, September 2011.
- [CGP23] Francesco Costantino, Nathan Geer, and Bertrand Patureau-Mirand. Admissible skein modules. *arXiv*, February 2023. arXiv:2302.04493 [math].
- [CL22] Francesco Costantino and Thang T. Q. Lê. Stated skein algebras of surfaces. *Journal of the European Mathematical Society*, 24(12):4063–4142, July 2022.
- [DG11] Tudor Dimofte and Sergei Gukov. *Quantum Field Theory and the Volume Conjecture*, volume 50 of *AMS/IP Studies in Advanced Mathematics*, page 19–42. American Mathematical Society, Providence, Rhode Island, July 2011.
- [FG00] Charles Frohman and Răzvan Gelca. Skein modules and the noncommutative torus. *Transactions of the American Mathematical Society*, 352(10):4877–4888, June 2000.
- [FGL01] Charles Frohman, Răzvan Gelca, and Walter LoFaro. The A-polynomial from the noncommutative viewpoint. *Transactions of the American Mathematical Society*, 354(2):735–747, October 2001.
- [Gar04] Stavros Garoufalidis. On the characteristic and deformation varieties of a knot. In *Proceedings of the Casson Fest*, page 291–309, University of Texas at Austin, Texas, USA, September 2004. Mathematical Sciences Publishers.
- [Gee05] Nathan Geer. The Kontsevich integral and quantized Lie superalgebras. *Algebraic & Geometric Topology*, 5(3):1111–1139, September 2005.
- [Gee06] Nathan Geer. Etingof–Kazhdan quantization of Lie superbialgebras. *Advances in Mathematics*, 207(1):1–38, December 2006.
- [Gee07] Nathan Geer. Some remarks on quantized Lie superalgebras of classical type. *Journal of Algebra*, 314(2):565–580, August 2007.
- [Gel01] Răzvan Gelca. On the relation between the A-polynomial and the Jones polynomial. *Proceedings of the American Mathematical Society*, 130(4):1235–1241, September 2001.
- [GJS19] Sam Gunningham, David Jordan, and Pavel Safronov. The finiteness conjecture for skein modules. *arXiv*, September 2019. arXiv: 1908.05233.
- [GL05] Stavros Garoufalidis and Thang T.Q. Lê. The colored Jones function is q-holonomic. *Geometry & Topology*, 9(3):1253–1293, 2005.
- [GL16] Stavros Garoufalidis and Thang T. Q. Lê. A survey of q-holonomic functions. *arXiv*, September 2016. arXiv: 1601.07487.

- [GLL18] Stavros Garoufalidis, Aaron D. Lauda, and Thang T. Q. Lê. The colored HOMFLYPT function is q-holonomic. *Duke Mathematical Journal*, 167(3):397–447, February 2018.
- [GP07] Nathan Geer and Bertrand Patureau-Mirand. Multivariable link invariants arising from $sl(2|1)$ and the Alexander polynomial. *Journal of Pure and Applied Algebra*, 210(1):283–298, July 2007.
- [GP18] Nathan Geer and Bertrand Patureau-Mirand. The trace on projective representations of quantum groups. *Letters in Mathematical Physics*, 108(1):117–140, January 2018. arXiv:1610.09129 [math].
- [GPT09] Nathan Geer, Bertrand Patureau-Mirand, and Vladimir Turaev. Modified quantum dimensions and re-normalized link invariants. *Compositio Mathematica*, 145(1):196–212, 2009.
- [GSAF12] Sergei Gukov, Piotr Sulkowski, Hidetoshi Awata, and Hiroyuki Fuji. Volume conjecture: refined and categorified. *Advances in Theoretical and Mathematical Physics*, 16(6):1669–1777, December 2012. Zbl: 1282.57016.
- [Guk05] Sergei Gukov. Three-dimensional quantum gravity, Chern-Simons theory, and the A-polynomial. *Communications in Mathematical Physics*, 255(3):577–627, April 2005. arXiv: hep-th/0306165.
- [GY22] Nathan Geer and Matthew B. Young. Three dimensional topological quantum field theory from $u_q(\mathfrak{gl}(1|1))$ and $u(1|1)$ Chern-Simons theory. *arXiv*, (arXiv:2210.04286), December 2022. arXiv:2210.04286 [hep-th].
- [Hai21] Benjamin Haioun. Relating stated skein algebras and internal skein algebras. *arXiv*, October 2021. arXiv: 2104.13848.
- [JLSS21] David Jordan, Ian Le, Gus Schrader, and Alexander Shapiro. Quantum decorated character stacks. *arXiv:2102.12283 [math]*, February 2021. arXiv: 2102.12283.
- [Kac77] Victor G Kac. Lie superalgebras. *Advances in mathematic*, 26(1), 1977.
- [Kac78] V. Kac. Representations of classical Lie superalgebras. In Konrad Bleuler, Axel Reetz, and Herbert Rainer Petry, editors, *Differential Geometrical Methods in Mathematical Physics II*, Lecture Notes in Mathematics, page 597–626, Berlin, Heidelberg, 1978. Springer.
- [Kas97] Rinat Kashaev. The hyperbolic volume of knots from the quantum dilogarithm. *Letters in Mathematical Physics*, 39(3):269–275, February 1997.
- [KQ22] Julien Korinman and Alexandre Quesney. The quantum trace as a quantum non-abelianization map. *arXiv*, (arXiv:1907.01177), May 2022. arXiv:1907.01177 [math].
- [KS11] Masaki Kashiwara and Pierre Schapira. Deformation quantization modules. *arXiv:1003.3304 [math]*, October 2011.
- [KT91] S. M. Khoroshkin and V. N. Tolstoy. Universal R-matrix for quantized (super)algebras. *Communications in Mathematical Physics*, 141(3):599–617, November 1991.
- [Lê06] Thang T. Q. Lê. The colored Jones polynomial and the A-polynomial of knots. *Advances in Mathematics*, 207(2):782–804, December 2006.
- [Lê16] Thang T. Q. Lê. Triangular decomposition of skein algebras. *arXiv*, September 2016. arXiv: 1609.04987.
- [Mik15] Victor Mikhaylov. Analytic torsion, 3d mirror symmetry and supergroup Chern-Simons theories. *arXiv*, May 2015. arXiv:1505.03130 [hep-th].
- [MW15] Victor Mikhaylov and Edward Witten. Branes and supergroups. *Communications in Mathematical Physics*, 340(2):699–832, December 2015.

- [RT90] N. Y. Reshetikhin and V. G. Turaev. Ribbon graphs and their invariants derived from quantum groups. *Communications in Mathematical Physics*, 127(1):1–26, January 1990.
- [Sab93] Claude Sabbah. Systèmes holonomes d'équations aux q-différences. *D-modules and microlocal geometry*, page 125–147, 1993.
- [Tur10] Vladimir Turaev. *Quantum invariants of knots and 3-manifolds*, volume 18. Walter de Gruyter GmbH & Co KG, 2 edition, 2010.
- [Yam94] Hiroyuki Yamane. Quantized enveloping algebras associated with simple Lie superalgebras and their universal R-matrices. *Publications of the Research Institute for Mathematical Sciences*, 30(1):15–87, February 1994.