A detailed field theory for RNA-like molecules with periodic base sequence

R. Dengler *

Abstract

This work examines a field theory for RNA-like molecules in a good solvent. The field theory is based on a lattice model for single- and double-strand RNA with a periodic base sequence, and otherwise contains all known relevant details (polymer types, polymer lengths and interactions). As for the somewhat less explicit O(n)-symmetric model there is a close relation to the conventional one-component branched polymer and the associated Lee-Yang problem. We further elucidate this relation. We derive exact results in the limiting cases of nearly complete denaturation and nearly complete renaturation. The single-strand critical exponent ν_{φ} is calculated in two-loop order.

The theory of RNA in a good solvent started with the classical work of de Gennes [1], who also introduced the approximation of using a periodic base sequence, and emphasized the importance of hairpin diagrams at low temperature [1], which form a kind of condensate.

The fundamental phenomenology is well known. In aqueous solution at temperatures exceeding approximately 100°C, single-strands predominate, while at lower temperatures double strands become prevalent (renaturation). This transition occurs continuously, lacking a critical temperature.

This general picture has been consistently validated with many different models, with various results for the statistics of polymer conformations. We only mention a selection of recent studies [2, 3, 4, 5]. The primary objective is to establish scaling laws and determine parameters such as scaling exponents and radius of gyration.

The complexity of the issue is compounded by the nondeterministic nature of base sequences and the necessity for relatively lengthy polymers to achieve scaling limits.

It is clear however, that long polymers enforce long correlations lengths, which normally entail critical phenomena. It thus should be no surprise to find connections to the universality class of the conventional branched polymer, at least in the case of a periodic base sequence.

"All linear polymers are alike"

Long linear polymers with self and mutual repulsion and other interactions contain linear segments devoid of interactions, which also are long if the interaction is weak. The probability distribution of the end to end vector of such segments is a gaussian function. A complete description of long linear polymers is nothing else than Gaussian functions connected with interaction vertices, and one could say "all linear polymers are alike".

A down-to-earth way to integrate over polymer configurations with self- and mutual repulsion and other interactions is the path integral of Edwards [6], where the variables directly are the positions of the monomers. A more convenient formulation of the problem are more abstract ("second quantized") field theories. Such field theories may contain a lot of details, as a bookkeeping

^{*}ORCID: 0000-0001-6706-8550

for polymer types, interactions and lengths. This bookkeeping would be difficult with the path integral of Edwards. But it is useful to keep in mind, that in the end everything boils down to gaussian integrals of segments in Feynman diagrams, which topologically are identical with polymer conformations. This remark also encompasses branched "linear" polymers, for which a field theory has been derived and examined by Lubensky and Isaacson in 1979 [7].

As we have recently shown [8, 9] it is not difficult to derive field theories for RNA-like branched polymers with single and double strands from a lattice model. The RNA field theory comes in two variants, a O(n)-symmetric variant which requires to set n = 0 in the end, and a variant with length variables instead of O(n) indexes without the n = 0 limit. The field theory with length variables contains more information, but is otherwise equivalent to the O(n)-symmetric model.

There is a close relation between the conventional branched polymer of Lubensky and Isaacson and RNA with a periodic base sequence, and such RNA appears to be a physical realization of the conventional branched polymer universality class.

In [8, 9] we have examined the relation between the models and the conventional branched polymer, and examined the O(n)-symmetric model in more detail. This work fills the last gap and examines the model with length variables. This leads to results which cannot be derived or are difficult to derive from the O(n)-model.

The model for single and double strand RNA

The physical system are RNA like single- and double-strand polymers with a periodic base sequence like GCGC... in a good solvent. The case of a random base sequence is much more complicated [9]. Here we examine in detail the most explicit field theory for such polymer networks. The action integral reads [9]

$$S_{\chi\varphi} = \int_{x,s} \tilde{\varphi}_s \left(r + \partial_s - \nabla^2 \right) \varphi_s + \frac{1}{2} \int_{x,s,s'} \chi_{s',s} \left(\tau + \partial_s - \partial_{s'} - \nabla^2 \right) \chi_{s,s'} + \frac{\alpha}{2} \int_x \left(\int_s \chi_{s,s} \right)^2 \quad (1)$$
$$- 2g \int_{x,s,s'} \chi_{s,s'} \tilde{\varphi}_s \varphi_{s'} - \lambda \int_{x,s,s',s''} \chi_{s,s'} \chi_{s',s''} \chi_{s'',s} - H \int_{x,s} \chi_{s,s}.$$

Expressions like $\int_{x,s} \dots = \int d^d x \int_{-\infty}^{\infty} ds \dots$ are abbreviations for integrals over space x and length variables s, s' and s''. To unclutter expressions length variables are written as suffixes, space variables are omitted. The excluded volume interactions $(\tilde{\varphi}\varphi)^2$, χ^4 and $\tilde{\varphi}\varphi\chi^2$ are irrelevant at the upper critical dimension 8 and not written down. We also have ignored "follower" χ^3 interactions [10], which are influenced by the coupling constants λ and g, but do not act back on the "leaders" λ and g.

Action $S_{\chi\varphi}$ has a simple and clear interpretation, and a formal derivation from a lattice model is nothing special and not really necessary. We mention once again that the topology of polymer conformations and Feynman diagrams always are identical.

But there are several unconventional aspects. First of all, the pair of fields φ and $\tilde{\varphi}$ describes the single-strand polymer, the field χ describes the double strand polymer. Accordingly, the fields φ and $\tilde{\varphi}$ have one length variable, the field χ has two length variables. Single-strand RNA molecules have an internal direction (5' to 3') and therefore there are two fields, a φ incoming into vertices and a $\tilde{\varphi}$ outgoing from vertices. Only oppositely aligned single RNA strands form double strand RNA molecules χ . The *s* and *s'* in $\chi_{s,s'}$ thus increment in different directions, and χ does not really have a direction, see fig.(1).

The question then is, which length variable is the first one in $\chi_{s,s'}$? It is crucial for a local field theory to have a unique *local* convention. The rule is: in a $\chi_{s,s'}$ connected to a vertex the first length variable s is incoming, the second length variable s' is outgoing. This explains the transposition




Figure 1: An RNA conformation in a universe generated by a single-strand source $(\tilde{\varphi})$ and sink (φ) . There is a double-strand polymer χ with oppositely aligned single strands. There are interactions with coupling constants g and λ . The two-point interaction with coupling constant α plays special role. When length variables are Fourier transformed to "frequencies", then these frequencies run unchanged through the diagram, and effectively only one frequency is involved.

of the length variables in the $\chi_{s',s}\chi_{s,s'}$ vertex in the action integral. The s' incoming on one side is outgoing on the other side, and vice versa for s. This is similar in the interaction with coupling constant g, which describes the transformation of two single-strands to a double strand and the reverse effect. The incoming s of $\chi_{s,s'}$ becomes the outgoing s of $\tilde{\varphi}_s$, and vice versa for s'.

The remaining interaction $-\lambda \chi^3$ describes the direct transition of a double-strand polymer to two double-strand polymers, see fig.(1). This interaction is generated from the interaction with coupling constant g, and is marginal at the upper critical dimension $d_c = 8$. Note that λ and goccur with positive sign in the statistical weight $e^{-S_{\chi\varphi}}$.

A special role is played by the strongly relevant harmonic term proportional to α . This term is generated by other interactions (λ , g or excluded volume), and must be taken into account from the outset [7, 8, 9, 11].

Frequency space

The action (1) is local in space but non-local in length variables s, s' and s''. This means that interactions between polymers are the same for any combination of length variables. This translational invariance suggests to use the Fourier-transformed representation. To length variables there correspond variables conjugate under Fourier transformation which we call "frequencies". The convention is $f_s = \int_{\omega} e^{-i\omega s} f_{\omega}$, with $\int_{\omega} \ldots = \int d\omega / (2\pi) \ldots$ In frequency representation action (1) reads

$$S_{\chi\varphi} = \frac{1}{2} \int_{x\omega\omega'} \chi_{-\omega',-\omega} \left(\tau_0 - i \left(\omega - \omega'\right) - \nabla^2 \right) \chi_{\omega,\omega'} - \lambda \int_{x\omega\omega'\omega''} \chi_{\omega,-\omega'} \chi_{\omega',-\omega''} \chi_{\omega'',-\omega}$$
(2)

$$+ \frac{\alpha}{2} \int_{x} \left(\int_{\omega} \chi_{\omega,-\omega} \right)^{2} + \int_{x\omega} \tilde{\varphi}_{-\omega} \left(r_{0} - \nabla^{2} - i\omega \right) \varphi_{\omega} - 2g \int_{x\omega\omega'} \tilde{\varphi}_{\omega} \varphi_{\omega'} \chi_{-\omega,-\omega'} - H \int_{x,\omega} \chi_{\omega,-\omega}.$$

The translational invariance of the interactions implies that a frequency along a single or double strand polymer runs unchanged through diagrams. There remain no frequency integrals, except possibly at the very end in a transformation back to length space and in closed frequency loops, which drop out because length variables cannot increase consistently in a loop ¹. The propagators are

¹This is evident in length representation on a lattice, where each propagator has a minimal length. In frequency representation loops generate factors like $\delta (\omega = 0)$, but this is due to the fact that the minimal length is not accounted for in the underlying strictly local field theory.

$$\left\langle \tilde{\varphi}_{\omega'}^{\boldsymbol{p}} \varphi_{\omega}^{\boldsymbol{k}} \right\rangle = \frac{\delta \left(\omega + \omega'\right)}{r_0 + k^2 - i\omega} \left(2\pi\right)^{d+1} \delta^d \left(\boldsymbol{p} + \boldsymbol{k}\right), \tag{3}$$

$$\left\langle \chi_{\nu,\nu'}^{\boldsymbol{p}} \chi_{\omega,\omega'}^{\boldsymbol{k}} \right\rangle = \left(\frac{\delta \left(\omega + \nu'\right) \delta \left(\nu + \omega'\right)}{v_{k,\omega-\omega'}} - \alpha \frac{\delta \left(\omega + \omega'\right)}{v_{k,\omega-\omega'}} \frac{\delta \left(\nu + \nu'\right)}{v_{k,\nu-\nu'}}\right) \left(2\pi\right)^{d+2} \delta^d \left(\boldsymbol{p} + \boldsymbol{k}\right),$$

where $v_{k,\omega} = \tau_0 + k^2 - i\omega$. Action (2) and the propagators (3) are very similar to the O(n)-symmetric expressions of the model with O(n) indexes instead of frequencies [8, 9, 11]. Frequencies run unchanged through diagrams like O(n) indexes, but there is nothing like a $n \to 0$ limit.

Double polymer twisting

Real RNA double strands are twisted in a rather rigid way. This implies a stiffness against rotations around the longitudinal axis, which is ignored in action (1). Such a type of stiffness also is present in other linear polymers, and it is plausible that this mechanical effect becomes less important for long polymers.

In the RNA system, however, there remains the topological effect caused by an interchange of the two strands. A double polymer has two length variables, one of them increments and the other decrements in a given direction. These variables can be traced through any configuration of interacting RNA. Whether a double strand is twisted or not can make a difference for where a variable turns up in external lines. This effect in a way is a nuisance, and it plays no role in renormalization group calculations, which can be done with $\omega = 0$ or, equivalently, within the O(n)-symmetric model without length variables. A correct field theory nevertheless must take the effect into account, and we show how this is realized in action (1).

First of all, double polymer twisting does *not* change the propagator. By convention, in a $\chi_{s,s'}$ connected to a vertex, the first length variable s is incoming, the second length variable s' is outgoing. The $\langle \chi \chi \rangle$ propagator thus is agnostic to any twisting (it would be difficult anyway in d > 2 to say whether a segment is twisted or not). The twisting gets generated by the interaction. In contrast to the usual case the interaction

$$-\lambda \int_{x,\omega,\omega',\omega''} \chi_{\omega,-\omega'} \chi_{\omega',-\omega''} \chi_{\omega'',-\omega}$$

is invariant only under cyclic permutations of the field factors. Perturbation theory, however, allows contractions with any field. As a result the propagation of length variables through an interaction vertex generates alternatives.

It is also clear that any valid diagram can be generated in this way. For a graph with directions, frequencies and twists, simply write down the vertices with all details (fields with frequencies). This are factors like $\chi_{\omega,-\nu}\chi_{\nu,-\rho}\chi_{\rho,-\omega}$. Field-contractions then reproduce the graph.

Debye structure function

Scattering experiments measure the wavevector-dependent density-density correlation function D(k), normalized with the partition sum Z. It is trivial but instructive to calculate D(k) for a noninteracting (gaussian) linear polymer, for instance the φ -polymer [12]. This is not immediately relevant for the RNA system, but illustrates some technical points. The Fourier transformed mass density operator for any length variable s is

$$\rho_{k,s}^{(\varphi)} = \int \mathrm{d}^d p \tilde{\varphi}_{-p,s} \varphi_{p+k,s} / \left(2\pi\right)^d.$$

Of interest is the ρ - ρ correlation function in a universe with a source $\tilde{\varphi}$ with length variable 0 at the origin, and a sink φ with length variable ℓ anywhere ²,

$$D_{\varphi}(k,\ell) = \frac{1}{Z} \int_{p,u,s} \left\langle \tilde{\varphi}_{p,0} \rho_{-k,u}^{(\varphi)} \rho_{k,s}^{(\varphi)} \varphi_{0,\ell} \right\rangle$$

= $\frac{2}{Z} \int_{0}^{\ell} du e^{-r_{0}u} \int_{u}^{\ell} ds e^{-(s-u)(r_{0}+k^{2})} e^{-(\ell-s)r_{0}}$
= $\frac{2}{Z} e^{-\ell r_{0}} k^{-4} \left(e^{-\ell k^{2}} - 1 + \ell k^{2} \right).$

Here we have used the $\tilde{\varphi}\varphi$ -Propagator $g_k(s) = \theta(s) e^{-s(r_0+k^2)}$. The normalization with partition sum $Z = \int_k \langle \tilde{\varphi}_{k,0} \varphi_{0,\ell} \rangle = e^{-\ell r_0}$ removes the dependence on r_0 . This is typical for polymers with given total length. The parameter r_0 acts as chemical potential for the constant polymer mass, and thus has no physical effect.

Known results

The field theory for branched polymers of Lubensky and Isaacson [7] is rather complicated, the χ -sector of eq.(1) or eq.(2) or the corresponding O(n)-symmetric model [8] are alternatives. A very similar other simpler version can be found in ref.([11]). It is known since 1981 that the critical point of "conventional" branched polymers can be mapped to the Lee-Yang critical point in two less dimensions [11]. The upper critical dimension is 8, the critical exponents in d = 3 and d = 4 are known exactly. As already noted by Lubensky and Isaacson "This theory is very rich" [7].

The additional single-strand field φ in action (2) does not spoil the relation to the Lee-Yang system. This means that peculiarities like a condensate

$$\langle \chi_{\omega,\omega'} \rangle = 2\pi \delta \left(\omega + \omega' \right) Q \left(\tau_0, \omega \right),$$

$$Q \left(\tau_0, \omega \right) = a_0 - a_1 \left(\tau_0 - \tau_{0c} - 2i\omega \right)^{\beta_H} + \dots,$$
(4)

Fisher renormalization of the critical exponents $\nu_{\chi} \rightarrow \nu_{H} = \nu_{\chi}\beta_{H}$ and $\gamma_{\chi} \rightarrow \gamma_{H} = \gamma_{\chi}\beta_{H}$, as well as a renormalization group calculation only imitate the Lee-Yang problem [8, 9]. Action (2), however, contains more information.

At the stable nontrivial fixed point with exactly $g_* = 3\lambda_*$ the equivalence encompasses both fields, i.e. χ and φ have the same anomalous dimension [9]. The condensate $\langle \chi \rangle \neq 0$ shifts the χ mass parameter as $\tau_0 \to \tau_0 - 6\lambda Q$, but also the φ mass parameter according to $r_0 \to r_0 - 2gQ$. This means that the χ - φ symmetry even is valid for $\tau_0 \neq \tau_{0c}$ in the region $r_0 - r_{0c} < 2ga_1 (\tau_0 - \tau_{0c})^{\beta_H}$. But there is a new critical exponent ν_{φ} , and of course, action (2) contains precise information about single- and double-strand polymer lengths.

Polymer lengths

The field theory (2) with length variables allows to derive the relative amounts of φ - and χ -polymer, at least in two limits. Consider a situation with (originally) a single single-strand polymer of given length ℓ as in fig.(1). Such a universe is generated by a single-strand source and sink,

$$\hat{\Phi}_{\ell} = \int_{\omega} e^{-i\omega\ell} \int_{x,\omega'} \tilde{\varphi}_{\omega'}(0) \,\varphi_{\omega}(x) \equiv \int_{\omega} e^{-i\omega\ell} \hat{\Phi}_{\omega}.$$
(5)

²This is one of the rare cases where a calculation in length space is simpler.

The polymer starts at the origin with length variable 0 and terminates anywhere with fixed length variable ℓ . The partition sum is $Z = \langle \hat{\Phi}_{\ell} \rangle$, and in naive perturbation series the parameter ω occurs only in the combinations $r_0 - i\omega$ and $\tau_0 - 2i\omega$. This allows to write the total length as

$$\ell = \frac{1}{Z} \int_{\omega} e^{-i\omega\ell} \partial_{(i\omega)} \left\langle \hat{\Phi}_{\omega} \right\rangle = \frac{1}{Z} \int_{\omega} e^{-i\omega\ell} \left(-\partial_{r_0} - 2\partial_{\tau_0} \right) \left\langle \hat{\Phi}_{\omega} \right\rangle = \frac{1}{Z} \int_{\omega} e^{-i\omega\ell} \left\langle \left(\hat{m}_{\varphi} + \hat{m}_{\chi} \right) \Phi_{\ell,\omega} \right\rangle = \bar{m}_{\varphi} + \bar{m}_{\chi}$$

where

$$\hat{m}_{\varphi} = \int_{x,\omega} \tilde{\varphi}_{-\omega} \varphi_{\omega}, \qquad (6)$$
$$\hat{m}_{\chi} = \int_{x,\omega,\omega'} \chi_{-\omega',-\omega} \chi_{\omega,\omega'}$$

are the polymer mass operators.

Initial stage of pairing

In the initial stage of pairing the single-strand polymer φ dominates. This occurs when the parameter τ_0 is large, away from the $g_* = 3\lambda_*$ fixed point. A plausible ansatz for the action integral is a subset of the monomials of eq.(2),

$$S_{\varphi\chi} = \int_{x,\omega} \tilde{\varphi}_{-\omega} \left(r_0 - \nabla^2 - i\omega \right) \varphi_{\omega} + \frac{\tau_0}{2} \int_{x\omega\omega'} \chi_{-\omega',-\omega} \chi_{\omega,\omega'} + \frac{\alpha}{2} \int_x \left(\int_\omega \chi_{\omega,-\omega} \right)^2$$
(7)
$$- 2g \int_{x\omega\omega'} \tilde{\varphi}_{\omega} \varphi_{\omega'} \chi_{-\omega,-\omega'} + u_{\varphi} \int_x \left(\int_\omega \tilde{\varphi}_{-\omega} \varphi_{\omega} \right)^2 - H \int_{x\omega} \chi_{\omega,-\omega}.$$

Only the excluded volume interaction for the single strand polymer φ with coupling constant u_{φ} has been added. And indeed, the coupling constants u_{φ} , τ_0 , α and g are marginal in $S_{\varphi\chi}$ at the new upper critical dimension $d_c = 4$. Derivative terms $\chi \nabla^2 \chi$ and $\chi i \omega \chi$, the χ^3 interaction, and $\chi^2 \chi^2$ as well as $\chi^2 \varphi^2$ excluded-volume interactions are strongly irrelevant and have been omitted.

The double-strand field χ occurs only harmonically in $S_{\varphi\chi}$ and can be integrated out from the weight $e^{-S_{\varphi\chi}}$. This is simple in perturbation theory with the $\chi\chi$ -propagator with $v_{k,\omega} = \tau_0$ from eq.(3). A contribution to r_0 of order $O(\alpha H)$ contains a length loop and drops out. There remains $S_{\varphi\chi}$ without χ , but with new bare coupling constants

$$r_{1} = r_{0} - 2gH/\tau_{0},$$

$$u_{1} = u_{\varphi} + 2g^{2} \left(\alpha/\tau_{0}^{2} - 1/\tau_{0}\right).$$

The remaining field theory is equivalent to the φ^4 -model of de Gennes, with additional information in r_1 and u_1 , and additional length variable for the fields φ and $\tilde{\varphi}$. Of interest is the partition sum of the situation depicted in fig.(1), the exact $\langle \tilde{\varphi} \varphi \rangle$ propagator. This quantity can be calculated in leading power-law approximation without any further approximation,

$$Z(r_0,\tau_0) = \left\langle \hat{\Phi}_\ell \right\rangle \sim \int_{-\infty}^\infty \mathrm{d}\omega e^{-i\ell\omega} \left(r_1 - r_{1c} - i\omega \right)^{-\gamma_\varphi} = \mathrm{const} \times \ell^{\gamma_\varphi - 1} e^{-(r_1 - r_{1c})\ell}.$$

The parameters r_1 and ω occur only in the combination $r_1 - i\omega$, and there is only wavevector 0 in play. This enforces the power law in the single remaining scaling variable $r_1 - r_{1c} - i\omega$, the exponent γ_{φ} is the exponent of the φ^4 - O(n = 0) model. The frequency integral has been performed with

$$\chi_{\omega,-\omega,k=0} \longrightarrow + \chi_{\omega,\nu,k} \longrightarrow \chi_{-\nu,-\omega,-k} + \chi_{\omega,-\nu,k} \longrightarrow \chi_{\nu,-\rho,p} + \dots$$

Figure 2: Integrating out the single-strand field φ generates one-loop contributions to the χ -sector of the field theory.

$$\int_{-\infty}^{\infty} \frac{\mathrm{d}\omega}{2\pi} e^{-i\omega\ell} \left(B - i\omega\right)^{-k} = \ell^{k-1} e^{-B\ell} / \Gamma\left(k\right). \tag{8}$$

The average φ -mass as expected is $\bar{m}_{\varphi} = -\partial \ln Z / \partial r_0 \sim \ell$ (in leading order). The average double strand mass follows as

$$\bar{m}_{\chi} = -\frac{\partial \ln Z}{\partial \tau_0} = -\left(\frac{\partial u_1}{\partial \tau_0}\frac{\partial}{\partial u_1} + \frac{\partial r_1}{\partial \tau_0}\frac{\partial}{\partial r_1}\right)\ln Z \sim O\left(\ln \ell\right) + \left(2gH - 2g^2\frac{\partial r_{1c}}{\partial u_1}\right)\ell/\tau_0^2 + O\left(\ell/\tau_0^3\right).$$
(9)

We are only interested in the contribution proportional to ℓ . The power law $\bar{m}_{\chi} \sim \ell/\tau_0^2$ for any dimension looks somewhat trivial. It does not depend on the φ^4 critical exponents, and thus there might be a simpler derivation or explanation. The constraint $\bar{m}_{\varphi} + \bar{m}_{\chi} = \ell$ is not exactly met, but this cannot be expected with the leading order approximation for \bar{m}_{φ} .

Final stage of pairing

In the final stage of pairing the double strand polymer prevails, and it is useful to eliminate the single-strand field φ from action (2). This is possible because φ occurs only harmonically, in particular in the three-point interaction $S_g = -2g \int_{x\omega\omega'} \tilde{\varphi}_{\omega} \varphi_{\omega'} \chi_{-\omega,-\omega'}$. The elimination generates a series of one-loop contributions, displayed graphically in fig.(2). These contributions to the effective action for χ are of the type $S_{n,d} = -g^n I_{n,d}(r_0, y) \int \chi^n$, with $n \ge 1$ fields χ and

$$I_{n,d}(r_0, y) = \text{const} \int^{\Lambda} \frac{\mathrm{d}^d p}{(r_0 + p^2 + y)^n} \stackrel{d \le 2n}{\sim} (r_0 + y)^{-(n-d/2)},$$
$$y = O\left(-i\omega_i, k_i^2\right).$$

The integrals $I_{n,d}$ are IR convergent and analytic in $r_0 + y$ for d > 2n. Integrals $I_{n\geq 4,d}$ are singular for dimensions $d \leq 8$, but $S_{n\geq 4,d}$ is strongly irrelevant. The power law for $I_{n,d}(r_0, y)$ is valid for $|r_0 + y| \ll \Lambda^2$, that is when the single-strand polymer is not completely suppressed. The new parameters of the remaining χ -field-theory are

$$H_{1} = H + c_{1}I_{1,d}(r_{0},0),$$

$$\tau_{1} = \tau_{0} - c_{2}I_{2,d}(r_{0},0),$$

$$\lambda_{1} = \lambda + c_{3}I_{3,d}(r_{0},0).$$
(10)

Contributions smaller by a factor $O(y/r_0)$ can be neglected. The relation of the remaining effective double-strand field theory to the Lee-Yang system remains valid, but the dependence of the parameters (10) on r_0 contains information about the single-strand polymer.

It recommends itself to now consider a universe with a χ -source and a χ -sink, generated by

$$\hat{\Omega}_{\ell} = \int_{x,s} \chi_{0,0}(0) \,\chi_{\ell,s}(x) = \int_{\omega} e^{-i\omega\ell} \int_{x,\nu,\nu'} \chi_{\nu,\nu'}(0) \,\chi_{\omega,\omega'=0}(x) \,. \tag{11}$$

The source is at the origin and has length variables 0, the sink is anywhere but has first length variable ℓ , the second length variable is arbitrary (it automatically gets fixed to $-\ell$). The partition sum

$$Z = \left\langle \hat{\Omega}_{\ell} \right\rangle \sim \int_{\omega} e^{-i\omega\ell} \left(\tau_1 - \tau_{1c} - i\omega \right)^{-\gamma_H} \sim \ell^{\gamma_H - 1} e^{-(\tau_1 - \tau_{1c})\ell}$$
(12)

is a χ -propagator with wavevector zero. The parameters τ_1 and ω occur only in the combination $\tau_1 - i\omega$. This enforces the power law in the single remaining scaling variable $\tau_1 - \tau_{1c} - i\omega$, the exponent $\gamma_H = \gamma_{\chi}\beta_H$ is the (Fisher-renormalized) exponent of the Lee-Yang model. The integral in has been performed with eq.(8). With eq.(6) there follow the average masses

$$\bar{m}_{\chi} = -2\partial \ln Z / \partial \tau_0 = 2\ell,$$

$$\bar{m}_{\varphi} = -\frac{\partial \ln Z}{\partial r_0} \cong -\frac{\partial H_1}{\partial r_0} \frac{\ln Z}{\partial H_1} - \frac{\partial \tau_1}{\partial r_0} \frac{\partial \ln Z}{\partial \tau_1} - \frac{\partial \lambda_1}{\partial r_0} \frac{\partial \ln Z}{\partial \lambda_1} \sim c_1 I_{2,d} \left(r_0\right) \frac{\ln Z}{\partial H_1} + 2c_2 I_{3,d} \left(r_0, y\right) \ell + \dots$$

It remains to calculate $X = \partial \ln Z/\partial H_1$. The partition sum (12) depends on H_1 via τ_{1c} , but it is better to use $X = \frac{1}{Z} \partial Z/\partial H_1 = \frac{1}{Z} \left\langle \hat{\Omega}_{\ell} \int_{x,\omega} \chi_{\omega,-\omega} \right\rangle$, see eq.(2) and eq.(11). This are three χ - χ propagators $(\tau_1 - \tau_{1c} - i\omega)^{-\gamma_H}$ with a vertex function $\Gamma_{\chi\chi\chi} \sim (\tau_1 - \tau_{1c} - i\omega)^{2\gamma_H - 1}$ in between (see appendix for the exponent). Again there is only wavevector 0 in play, and the factors are powers of the single remaining scaling variable $\tau_1 - \tau_{1c} - i\omega$. With the help of formula (8) one finds $X \sim \ell$ and

$$\bar{m}_{\varphi} \sim c_1' I_{2,d}(r_0) \ell + 2c_2 I_{3,d}(r_0, y) \ell + \dots \sim c_1'' \ell r_0^{-(2-d/2)} + c_2'' \ell r_0^{-(3-d/2)} + \dots$$
(13)

The dependence of \bar{m}_{χ} on the bare coupling constant λ_1 is still smaller. Universal quantities like γ_H do not depend on λ_1 , nonuniversal quantities like amplitudes and temperature shifts like τ_{1c} are regular functions of λ_1 .

The power law $\bar{m}_{\varphi} \sim \ell/\sqrt{r_0}$ in d = 3 for large r_0 looks somewhat trivial. It does not depend on the Lee-Yang critical exponents, and there might be a simpler derivation. The constraint $\bar{m}_{\varphi} + \bar{m}_{\chi} = 2\ell$ is not exactly met, but this cannot be expected with the leading power-law approximation for Z and \bar{m}_{χ} . The calculation has the air of thermodynamics, and there might be a simpler explanation for the result (13).

The contributions $S_{n,d}$ from φ -elimination to the effective χ -sector look as if this would give a field theory not equivalent to the Lee-Yang problem for $r_0 = r_{0c}$. However, this is an illusion. The φ -loops cannot contain an α -propagator (eq.(3)), and thus are less singular and do not contribute to the critical point [9].

Intermediate range

Only one of the polymer masses is independent because of the constraint $\bar{m}_{\varphi} + \bar{m}_{\chi} = \ell$, and one can consider the ratio \bar{m}_{φ}/ℓ as a degree of freedom. This ratio usually depends on many physical quantities: temperature, acidity of solvent, concentrations of ions etc. The parameters r_0 and τ_0 depend on these physical quantities, but always must change in such a way that the constraint is fulfilled. If τ_0 is chosen as independent parameter then $r_0 = r_0 (\tau_0, \ell)$ is a kind of hyperbola in the τ_0 - r_0 -plane. A large τ_0 diminishes \bar{m}_{χ} , and this is compensated by r_0 becoming small. If \bar{m}_{φ} approaches the maximal value ℓ , then the influence of r_0 disappears ($i\omega \sim 1/\ell$ dominates r_0), and the physical quantities only act via τ_0 . This and the symmetric situation lead to the two limiting cases above, with either large r_0 or large τ_0 as only and *free* physical parameter.

It is more difficult to get quantitative results for the transitional range. Starting from the dissociated limit (the φ^4 model), double-strand mass $\bar{m}_{\chi} \sim \ell/\tau_0^2$ increases rapidly. Starting from the paired limit, single-strand mass $\bar{m}_{\varphi} \sim \ell/\sqrt{r_0}$ increases much more slowly.

Here the stable fixed point with $g_* = 3\lambda_*$ comes into play. At this fixed point χ and φ together form a branched polymer of the conventional type (χ and φ have same anomalous dimension [9]),

with comparable amounts of \bar{m}_{φ} and \bar{m}_{χ} . This at least indicates that the Lee-Yang regime extends into the transitional regime.

The main contributions to the polymer masses near this stable fixed point originate from a mass operator (eq.(6)) insertion into the condensate $\langle \chi \rangle$ (hairpin diagrams). The calculation for r_0 small and τ_0 not too large is similar to eq.(13). There are three χ - χ propagators with a vertex $\Gamma_{\chi\chi\chi}$ in between, only the dependence on r_0 is missing. Both masses are of order $O(\ell)$, with nonuniversal amplitudes.

Two-loop calculation

The critical exponent ν_{φ} does not show up in the leading approximation of polymer length \bar{m}_{φ} in the final stage of single-strand pairing (the calculation based on the O(n) model of [9] is misleading). However, a more precise value for ν_{φ} might still be of interest.

An ab initio two-loop calculation for the Lee-Yang field theory is demanding and not worth the effort. Higher-loop-calculations nowadays are a science by itself, and there are elaborate tools for nearly every step of the calculation. Our goal here is more modest. We show, how known results can be combined to get the exponent ν_{φ} in two loop order.

From the 5-loop results [13] one can take over the functions $\beta(\lambda_6)$, $\eta_{\varphi}(\lambda_6)$, where the suffix 6 at the coupling constant denotes the upper critical dimension. When modified minimal subtraction is used in both cases, then the coupling constants of the Lee-Yang- and polymer-problem are related by $\lambda_6^2 = (3 - \epsilon/2) \lambda_8^2$, where $\epsilon = 8 - d$ [14]. This allows to deduce the functions $\beta(\lambda_8)$ and $Z_{\varphi}(\lambda_8)$.

There remains the calculation of the $\Gamma_{\varphi\varphi}$ vertex function with a φ^2 insertion. This does not lead to any new two-loop diagrams (the diagrams are depicted as E, $a_1, ..., a_5$ in the appendix of [9]). An integration by parts algorithm allowing to reduce the integrals to two simple master integrals is concisely described in [15]. With the requisite algebra implemented in the C++-Ginac framework we find

$$\nu_{\varphi} = \frac{1}{2} + \frac{\epsilon}{36} + \frac{11}{1458}\epsilon^2 + \dots$$

This still does not allow a plausible extrapolation to d = 3, but at least provides a check for a more sophisticated calculation.

Fields with unique scaling dimensions

An peculiar aspect of action (2) and similar actions for branched polymers is that the field χ does not have a unique scaling dimension, c.f. $\langle \chi \chi \rangle \sim k^{-2}$ and $\langle \chi \chi \rangle \sim k^{-4}$ in eq.(3). And there are other, related, peculiarities. The parameter α does not have a finite fixed point. This is not really a problem, because in the end α occurs only in the effective couplings constants $\lambda \alpha^{1/2}$ and $g \alpha^{1/2}$. But a naive dimensional analysis is not sufficient to find consistent canonical dimensions of the fields and the upper critical dimension.

The root of the complications is the α -term. The α -term is harmonic, but can occur only once in a propagator line, and thus more behaves as an interaction than as a propagator. It is of interest to transform action (1) or (2) to an equivalent form without these peculiarities.

The essential observation is that every loop in Feynman diagrams contributing to the critical point contains exactly one α -term [7, 11]. Cutting such a diagram at the α -terms generates a tree diagram, and the α -term in this sense acts as the source of fluctuations, connecting solutions of the classical equation of motion [11].

The first step is to separate the "diagonal" and off-diagonal components of χ ,

$$\chi_{\omega\nu} = \hat{\chi}_{\omega\nu} + 2\pi\delta \left(\omega + \nu\right)\rho_{\omega},\tag{14}$$

with $\rho_{\omega} = \chi_{\omega,-\omega}$.³ Inserting eq.(14) into eq.(2) leads to expressions like $\delta (\omega = 0)$ and $\delta (\omega)^2$, which can be regularized to $\delta (0) = Na$ and $\delta (\omega)^2 = Na\delta (\omega)$ by temporarily using discrete frequencies $\omega \in \{0, \frac{2\pi}{a}, ..., \frac{2\pi}{a} (N-1)\}$, where *a* is monomer length and *N* a large number. The result is

$$S' = S_{\hat{\chi}\varphi} + S_{\hat{\chi}\rho}, \qquad (15)$$

$$S_{\hat{\chi}\varphi} = \frac{1}{2} \int_{x\omega\omega'} \hat{\chi}_{-\omega',-\omega} \left(\tau_0 - i \left(\omega - \omega'\right) - \nabla^2 \right) \hat{\chi}_{\omega,\omega'} + \int_{x,s} \tilde{\varphi}_s \left(r + \partial_s - \nabla^2 \right) \varphi_s - \lambda \int_{x,\omega,\omega',\omega''} \hat{\chi}_{\omega,-\omega'} \hat{\chi}_{\omega',-\omega''} \hat{\chi}_{\omega'',-\omega} - 2g \int_{x\omega\omega'} \tilde{\varphi}_{\omega} \varphi_{\omega'} \hat{\chi}_{-\omega,-\omega'}, \qquad S_{\hat{\chi}\rho} = V\left(\rho\right) - \frac{A}{2} \int_x \left(\int_{\omega} \rho_{\omega} \right)^2, \qquad V\left(\rho\right) = \frac{1}{2} \int_{x\omega\omega'} \rho_{\omega} \left(\tau_0 - 2i\omega - \nabla^2 \right) \rho_{\omega} - \lambda' \int_{x\omega} \rho_{\omega}^3 - 3\lambda'' \int_{x\omega\nu} \hat{\chi}_{\omega,-\nu} \hat{\chi}_{\nu,-\omega} \rho_{\omega} - 2g \int_{x\omega} \tilde{\varphi}_{\omega} \varphi_{-\omega} \rho_{\omega}.$$

Here we also have written $A = -\alpha > 0$, this sign change is to be corrected at the end. The remaining transformation concerns $S_{\hat{\chi}\rho}$ and resembles the transformation in the case of conventional branched polymers [11], only the "frequency" variables are new. With the help of an auxiliary field h one can write

$$e^{-S_{\hat{\chi}\rho}} = \int_{-\infty}^{\infty} \mathrm{D}h e^{-\frac{1}{2A}\int_{x}h^{2}} e^{-V(\rho) + \int_{x}h\int_{\omega}\rho_{\omega}}.$$

The equation of motion $\delta V/\delta \rho - h = 0$ for ρ can now be enforced with a δ -function,

$$e^{-S_{\hat{\chi}\rho}} = \int_{-\infty}^{\infty} \mathrm{D}h e^{-\frac{1}{2A}\int_{x}h^{2}} \int_{-i\infty}^{i\infty} \mathrm{D}\tilde{\rho} e^{-\int_{x\omega}\tilde{\rho}_{\omega}(\delta V/\delta\rho - h)} \det\left(\delta V/\delta\rho\right).$$

Performing the average over h leads to

$$S_{\hat{\chi}\rho} = -\frac{A}{2} \int_{x} \left(\int_{\omega} \tilde{\rho}_{\omega} \right)^{2} + \int_{x\omega} \tilde{\rho}_{\omega} \delta V / \delta \rho.$$

In a last step the sign of A must be corrected. This can be achieved by making the coupling constants imaginary [11]. The final result is

$$S'_{\hat{\chi}\rho} = \int_{x} \left\{ -\frac{A}{2} \left(\int_{\omega} \tilde{\rho}_{\omega} \right)^{2} + \int_{\omega} \tilde{\rho}_{\omega} \left(\tau_{0} - 2i\omega - \nabla^{2} \right) \rho_{\omega} - 3i\lambda' \int_{\omega} \tilde{\rho}_{\omega} \rho_{\omega}^{2} \right\}$$
(16)
$$- 3i\lambda'' \int_{x\omega\nu} \tilde{\rho}_{\omega} \hat{\chi}_{\omega,-\nu} \hat{\chi}_{\nu,-\omega} - 2ig \int_{x\omega} \tilde{\varphi}_{\omega} \varphi_{-\omega} \tilde{\rho}_{\omega}.$$

Here we have ignored the functional determinant. Its only effect would be to remove closed loops of length variables ⁴.

In $S_{\hat{\chi}\varphi} + S'_{\hat{\chi}\rho}$ now everything is as usual. According to a dimensional analysis α is dimensionless, and one can set A = 1. The propagators are (c.f. eq. (3))

$$\left\langle \tilde{\rho}_{\nu}^{k} \rho_{\omega}^{p} \right\rangle = \frac{\delta \left(\omega - \nu\right)}{v_{k,2\omega}} \left(2\pi\right)^{d+1} \delta^{d} \left(\boldsymbol{k} + \boldsymbol{p}\right),$$

$$\left\langle \rho_{\nu}^{k} \rho_{\omega}^{p} \right\rangle = \frac{1}{v_{k,2\omega} v_{p,2\nu}} \left(2\pi\right)^{d} \delta^{d} \left(\boldsymbol{k} + \boldsymbol{p}\right),$$

$$\left\langle \hat{\chi}_{\omega,\omega'}^{k} \hat{\chi}_{\nu,\nu'}^{p} \right\rangle = \frac{\delta \left(\omega + \nu'\right) \delta \left(\omega' + \nu\right)}{v_{k,\omega-\omega'}} \left(2\pi\right)^{d+2} \delta^{d} \left(\boldsymbol{k} + \boldsymbol{p}\right)$$

³The notation ρ_{ω} is more compact, while $\chi_{\omega,-\omega}$ makes frequency conservation explicit.

⁴The determinant can be taken into account with the help of fermionic fields.

The interaction $\lambda \hat{\chi}^3$ is irrelevant at the upper critical dimension 8, the interactions $\lambda' \tilde{\rho} \hat{\chi} \hat{\chi}$, $\lambda'' \tilde{\rho} \rho^2$ and $g \tilde{\varphi} \varphi \tilde{\rho}$ are marginal. In a situation as in fig.(1) there is no $\hat{\chi}$ field, except possibly in frequency loops. It follows that $\hat{\chi}$ can be ignored, in particular $\hat{\chi}$ does not contribute to the critical behavior.

The fact that the fields ρ , $\tilde{\rho}$ and $\hat{\chi}$ do have unique scaling dimensions does not necessarily simplify calculations. An alternative to action (16) without the $\tilde{\rho}$ field is to simply ignore $\hat{\chi}$ in action (15). The merit of the formulation (16) is of conceptional nature.

Conclusions

In this work we have closed the last gap and examined the RNA field theory with length variables, which explicitly contains all known relevant details. In all calculations a periodic base sequence was assumed.

We have shown that it is possible to derive exact results for the polymer masses for the initial and final stage of single-strand pairing. It would be of interest to compare the results (9) and (13) with other calculations or experiments.

The gyration radius $\sqrt{\ell}$ of the underlying conventional branched polymer also was derived in [5] by means of Flory type arguments and found to agree with simulations. This is a confirmation for our field theory, the χ -sector of which is closely related to the model introduced by Parisi [11].

It would be of interest to measure the gyration radius of real RNA with periodic base sequence near complete renaturation. A scaling law $\sqrt{\ell}$ would indicate that this system belongs to the conventional branched-polymer universality class. Finite size scaling might be useful to extrapolate to long polymer chains.

It is questionable whether RNA with random base sequence has much in common with real RNA. But incorporating randomness into the field [9] theory might show some tendency in comparison to the somewhat unnatural case of a periodic base sequence studied here.

References

- P.-G. de Gennes. Statistics of branching and hairpin helices for the dAT copolymer. Biopolymers, 6:715 – 729, 1968.
- [2] M. Müller. Statistical physics of RNA-folding. Phys. Rev. E, 67:021914, 2002.
- [3] A. Rosa and R. Everaers. Conformational statistics of randomly branching double-folded ring polymers. *Eur. Phys. J. E*, 42(7), 2019.
- [4] S. Cocco, A. D. Martino, A. Pagnani, and M. Weigt. Statistical-physics approaches to RNA molecules, families and networks, 2022, arXiv preprint cond-mat.soft/2207.13402.
- [5] D. Vaupoti, A. Rosa, L. Tubiana, and A. Bozic. The Journal of Chemical Physics, page 158.23, 2023.
- [6] S. F. Edwards. The statistical mechanics of polymers with excluded volume. Proc. Phys. Soc., 85(4):613 - 624, 1965.
- [7] T. C. Lubensky and J. Isaacson. Statistics of lattice animals and dilute branched polymers. *Phys. Rev. A*, 20:2130 – 2146, 1979.
- [8] R. Dengler. A universality class for RNA-like polymers and double polymers, 2022, arXiv preprint cond-mat.soft/2210.07099.

- [9] R. Dengler. Critical phenomena of RNA-like polymers a synopsis, 2023, arXiv preprint condmat.soft/2310.06126.
- [10] A. Kaviraj and E. Trevisani. Random field ϕ^3 model and parisi-sourlas supersymmetry. Journal of High Energy Physics, 2022(8):1–51, 2022.
- [11] G. Parisi and N. Sourlas. Critical behavior of branched polymers and the Lee-Yang edge singularity. *Phys. Rev. Lett.*, 14:871 – 874, 1981.
- [12] P. Debye. Molecular-weight determination by light scattering. J. Phys. Colloid. Chem., 51:18 - 32, 1947.
- [13] M. Borinsky, J. A. Gracey, M. V. Kompaniets, and O. Schnetz. Five loop renormalization of ϕ^3 theory with applications to the Lee-Yang edge singularity and percolation theory. *Phys. Rev.* D, 103:116024, 2021.
- [14] A. Aharony, Y. Imry, and S. K. Ma. Lowering of dimensionality in phase transitions with random fields. *Phys. Rev. Lett.*, 37(20):1364, 1976.
- [15] I. Bierenbaum and S. Weinzierl. The massless two-loop two-point function. The European Physical Journal C-Particles and Fields, 32:67–78, 2003.
- [16] T. C. Lubensky and A. J. McKane. Anderson localization, branched polymers and the Yang-Lee edge singularity. J. Physique, 42:L331, 1981.

Appendix

Scaling dimensions

In order to reduce confusion, we derive here the scaling law of the 3-point vertex function $\Gamma_{\chi\chi\chi} \sim \tau^{2\gamma_H-1}$ in ω -representation, which was used in the estimation of the polymer mass \bar{m}_{φ} in the final stage of single-strand pairing, eq.(13).

There are different critical exponents in play, there is the Lee-Yang and the branched-polymer system, and there are the O(n) field theories and the field theories with length- or frequency-variables.

In ω -representation there are no ω -integrals in vertex functions, and the vertex functions have the same scaling dimensions as their O(n) counterparts. It thus is simplest to simply ignore all ω -arguments. The scaling dimensions of χ^2 by definition is $\chi^2 \sim k^{d-1/\nu_{\chi}}$. Because of Fisher renormalization $\tau \to \tau^{\beta_H}$ [7] wavevector and temperature variable are related by $k \sim \tau^{\nu_H}$ with $\nu_H = \nu_{\chi}\beta_H$ (instead of the usual $k \sim \tau^{\nu_{\chi}}$).

The 3-point vertex function $\Gamma_{\chi\chi\chi}$ is a self energy $\Gamma_{\chi\chi}$ with an additional $\lambda\chi^3$ interaction with a (truncated) external leg, which apart from a constant factor is a $\Gamma_{\chi\chi}$ with an inserted $\int d^d x \chi^2 \sim k^{-1/\nu_{\chi}}$. By definition $\Gamma_{\chi\chi} \sim \tau^{\gamma_H}$ and thus $\Gamma_{\chi\chi\chi} \sim \tau^{\gamma_H} k^{-1/\nu_{\chi}} \sim \tau^{2\gamma_H - 1}$, where $\beta_H + \gamma_H = 1$ was used (there are many relations between the critical exponents) [16].