MOSEL survey: Spatially offset Lyman-continuum emission in a new emitter at z=3.088

Anshu Gupta,^{1,2} Cathryn M. Trott,^{1,2} Ravi Jaiswar,^{1,2} E. V. Ryan-Weber,^{3,2} Andrew J. Bunker,⁴ Ayan Acharyya,⁵ Alex J. Cameron,⁴ Ben Forrest,⁶ Glenn G. Kacprzak,^{7,2} Themiya Nanayakkara,^{3,2} Kim-Vy Tran,^{8,2,9} and Aman Chokshi^{10,2}

¹International Centre for Radio Astronomy Research (ICRAR), Curtin University, Bentley WA, Australia

²ARC Centre of Excellence for All Sky Astrophysics in 3 Dimensions (ASTRO 3D), Australia

³Centre for Astrophysics and Supercomputing, Swinburne University of Technology, Hawthorn, VIC 3122, Australia

⁴Department of Physics, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford OX1 3RH, UK

⁵Department of Physics and Astronomy, Johns Hopkins University, Baltimore, MD 21218, USA

⁶Department of Physics and Astronomy, University of California Davis, One Shields Avenue, Davis, CA, 95616, USA

⁷Swinburne University of Technology, Hawthorn, VIC 3122, Australia

⁸School of Physics, University of New South Wales, Sydney, NSW 2052, Australia

⁹Center for Astrophysics | Harvard & Smithsonian, Cambridge, MA

¹⁰The University of Melbourne, School of Physics, Parkville, VIC 3010, Australia

ABSTRACT

We present the discovery of a unique Lyman-continuum (LyC) emitter at z = 3.088. The LyC emission were detected using the Hubble Space Telescope (HST) WFC3/UVIS F336W filter, covering a rest-frame wavelength range of 760 – 900 Å. The peak signal-to-noise ratio (SNR) of LyC emission is 3.9 in a r = 0.24'' aperture and is spatially offset by $0.29'' \pm 0.04$ (~ 2.2 ± 0.3 kpc) from the rest-UV emission peak (F606W). By combining imaging and spectroscopic data from the James Webb Space Telescope (JWST) JADES, FRESCO and JEMS surveys, along with VLT/MUSE data from the MXDF survey, we estimate that the probability of random alignment with an interloper galaxy causing the LyC emission is less than 6×10^{-5} . The interstellar medium (ISM) conditions in the galaxy are similar to other LyC emitters at high redshift $(12 + \log(O/H) = 7.79^{+0.06}_{-0.05}, \log U = -3.27^{+0.14}_{-0.12}, O32 = 3.65 \pm 0.22)$, although the single-peaked Lyman- α profile and lack of rest-UV emission lines suggest an optically thick ISM. We think that LyC photons are leaking through a narrow cone of optically thin neutral ISM, most likely created by a past merger (as evidenced by medium-band F210M and F182M images). Using the escape fraction constraints from individual leakers and a simple model, we estimate that the opening half-angle of ionization cones can be as low as 16° (2% ionised fraction) to reproduce some of the theoretical constraints on the average escape fraction for galaxies. The narrow opening angle required can explain the low number density of confirmed LyC leakers.

Keywords: galaxies: high-redshift – emission-line – interactions – evolution

1. INTRODUCTION

Understanding the escape of ionizing photons from the first galaxies remains one of the biggest challenges for the models of galaxy formation and evolution. The past year of operations by the James Webb Space Telescope (JWST) has significantly transformed our understanding of early-universe galaxies. Nevertheless, the high opacity of the intergalactic medium prevents us from detecting the ionizing photons escaping from the first galaxies. Therefore, we need to rely on analogs of first galaxies to understand the mode of Lyman continuum (LyC) escape.

In the past decade, photometric observations revealed elevated emission line equivalent widths (EW) for z > 6 galaxies (Labbé et al. 2013; Roberts-Borsani et al. 2016; Barro et al. 2019; Mainali et al. 2019; Endsley et al. 2020). Recent observations with JWST have confirmed this, demonstrating that 80-90% of galaxies within the first billion years of the universe have [OIII] 5007+H β EW > 800 Å, almost three times the EW of a typical star-forming galaxy at $z \sim 2$ (Endsley et al. 2023; Cameron et al. 2023; Tang et al. 2023; Rinaldi et al. 2023; Boyett et al. 2024). Consequently, many observational programs focus on studying the detailed properties of extreme emission line galaxies (EELGs), where emission lines from gas contribute almost 40-50% to the total flux in certain photometric bands, both at low (Yang et al. 2017b; Yuma et al. 2019; Izotov et al. 2018; Lumbreras-Calle et al. 2022) and high redshifts (Atek et al. 2011; Maseda et al. 2014; Gupta et al. 2022). EELGs typically exhibit high specific star

formation rates and high ionization parameters (Tang et al. 2019; Gupta et al. 2022) that could be linked with a high LyC escape fraction (Izotov et al. 2018).

Observations of dwarf star-forming galaxies (SFGs) in the nearby universe with HST Cosmic Origin Spectrograph have built a sizable population of local LyC leakers (Izotov et al. 2016b,a, 2018; Flury et al. 2022a). The escape fraction shows a weak positive correlation with [OIII] 5007/[OII] 3727 ratio and an inverse correlation with the velocity separation between the blue and red peaks of the Lyman- α emission (Izotov et al. 2018; Flury et al. 2022b). These observations suggest that a harder ionizing field and/or a low neutral gas covering fraction are crucial to facilitate the leakage of LyC photons.

At $z \sim 1 - 3$, the stacking experiments with SFGs yield an escape fraction between $f_{esc} = 1\% - 10\%$ (Steidel et al. 2018; Wang et al. 2023). The individual detection of LyC leakers at high redshifts is complicated by the higher chance of contamination from interloper galaxies (Vanzella et al. 2010, 2012) and attenuation by the partially neutral intergalactic medium (IGM) (Inoue et al. 2014; Bassett et al. 2021). Deep imaging campaigns with HST and u-band imaging on groundbased telescopes have created significant samples of candidate LyC leakers at cosmic noon (e.g., Meštrić et al. 2020; Kerutt et al. 2023). Recently, Wang et al. (2023) used deep UVIS imaging from the UVCANDELS survey to report a tentative discovery of five LyC leakers between redshift 2.4 < z < 3.7, some of them showing $f_{esc} > 60\%$. However, follow-up spectroscopy is required to rule out any possible foreground contaminant.

The A2218-Flanking and the Sunburst arc are two gravitationally lensed LyC leakers at $z \sim 2.5$. In A2218-Flanking, LyC radiation is leaking from a compact dwarf galaxy (Bian et al. 2017), whereas in the Sunburst Arc, LyC radiation is leaking from a compact star-forming region that could be a young massive star cluster (Rivera-Thorsen et al. 2019). Ion1 (Vanzella et al. 2012; Ji et al. 2020), Ion2 (de Barros et al. 2016; Vanzella et al. 2016, 2020), Ion3 (Vanzella et al. 2018, 2020), and Q1549-C25 (Shapley et al. 2016) are a few examples of non-lensed LyC leakers, each showing a wide array of interstellar medium conditions and escape fractions. For some of these galaxies, escape fractions can be higher than 50% (Vanzella et al. 2016; de Barros et al. 2016; Saxena et al. 2022; Marques-Chaves et al. 2022; Rivera-Thorsen et al. 2022). Recently, Kerutt et al. (2023) discovered five new LyC leakers using the MUSE-Wide survey (Urrutia et al. 2019) by searching for LyC emission within a small aperture (0.35'')spatially coincident with peak rest-UV emission in Lyman- α emitters.

The Lyman- α emission profile for high redshift LyC leakers exhibits a range, contrary to the classic double-peaked Lyman- α emission observed in most LyC leakers at $z \sim 0$. A few LyC emitters show 3-4 peaks in Lyman- α emission

indicative of multiple kinematic components in the neutral interstellar medium (ISM) (Vanzella et al. 2020). Lyman- α emission is only single-peaked in Q1549-C25 (Shapley et al. 2016) and appears in absorption in Ion1 (z = 3.78), indicative of a highly optically thick ISM (Ji et al. 2020). For Ion1, the LyC emission is spatially offset by 0."12 ± 0."03 from the peak UV emission suggesting that the ionizing photons might be leaking from a small part of the galaxy. Kerutt et al. (2023) find a huge scatter between the properties of Lyman- α emission and escape fraction for high redshift LyC emitters.

This paper reports the discovery of a new LyC leaker at z = 3.088 in the Chandra Deep Field South (CDFS, Giacconi et al. 2002) and analyzes its ISM conditions using the wealth of ancillary data from the JWST and MUSE spectrograph (Bunker et al. 2023; Bacon et al. 2023). The paper is organized as follows. Section 2 describes the photometric and spectroscopic data analysed here. In section 3.1 and 3.2, we discuss the spectral energy distribution fitting, the LyC detection and escape fraction measurements. In sections 3.4 and 3.5, we report properties of ISM derived from JWST/NIRSpec and MUSE spectrum respectively. We discuss our results in Section 4 and present a simple model to derive the opening angle of the ionization cone. The paper uses a flat Λ CDM cosmology with $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_{\text{M}} = 0.3$, and $\Omega_{\Lambda} = 0.7$.

2. DATA

The galaxy z19863 (RA=53.1699, DEC=-27.7684) was identified as an EELG by Forrest et al. (2018) using photometric data from the FourStar Galaxy Evolution survey (ZFOURGE; Straatman et al. 2016). It is part of the Multi-Object Spectroscopy of Emission Line (MOSEL; Tran et al. 2020; Gupta et al. 2022) survey of galaxies, with aims to analyse the properties of EELGs at 2.5 < z < 4. The galaxy lies in the Hubble Ultra Deep Field (HUDF), providing us access to a multitude of ancillary data from both HST and JWST (Figure 1).

We combine NIRCam imaging from the JWST Advanced Deep Extragalactic Survey (JADES, Bunker et al. 2023; Hainline et al. 2023; Rieke & the JADES Collaboration 2023; Eisenstein et al. 2023), the FRESCO survey (Oesch et al. 2023), and the JWST Extragalactic Medium-band Survey (JEMS, Williams et al. 2023), giving us extensive broad and medium band coverage. Additionally, the FRESCO survey has released astrometrically aligned mosaics based on the archival photometry on the Hubble Legacy Fields including the UVIS/F336W imaging (Illingworth et al. 2016; Whitaker et al. 2019).

The galaxy has a spectroscopic redshift of z = 3.088 determined from the deep NIRSpec spectrum from the JADES (Bunker et al. 2023). We also use the data from the MUSE eXtremely Deep Field (MXDF, Bacon et al. 2023) to rule out



Figure 1. Multi-band imaging of galaxy z19863 from HST and JWST in increasing order of wavelength. Each panel is $1.5'' \times 1.5''$ across. The top two panels correspond to the original and PSF-smoothed images from the HST F336W filter (LyC). The red plus and cyan cross mark the position of the peak non-ionising rest-UV (F606W) and the ionising rest-UV flux (F336W) respectively. The black plus marks the location of the *z* = 3.4 Lyman- α emitter detected in the MXDF survey (see Section 3.5) for more details.

interloper galaxies and analyse the Lyman- α emission profile. At z = 3.088, the UVIS/F336W filter covers rest-frame wavelengths between 760 – 900 Å, and therefore should have no contamination from non-ionising photons.

3. ANALYSIS

3.1. Spectral energy distribution

We convolve the images down to a common PSF of the NIRCam/F444W filter and use a 3σ threshold to create a segmentation map based on the F444W image. The fluxes are extracted within an aperture defined by KRON parameters of 2.0. A slightly lower KRON parameter is used here to minimize the contamination from a nearby background source at z = 3.4. We estimate errors by taking the root mean square of flux at random locations within the same aperture for each image.

We use the MAGPHYS (da Cunha et al. 2008, 2015) spectral energy distribution (SED) fitting code with the BC03 stellar population synthesis model (Bruzual & Charlot 2003), delayed exponentially declining star formation history model, and Charlot & Fall (2000) dust attenuation law to derive physical properties. We only use filters that are not contaminated by bright nebular emission lines such as $[O_{III}]$ 5007 and H α (all except F210M and F277W, Figure 2). We also do not include an IGM absorption model.

3.2. LyC Detection

We use the HST UVIS/F336W mosaics released by the FRESCO survey (Oesch et al. 2023), created using the Hubble Legacy Fields photometry (Teplitz et al. 2013; Rafelski et al. 2015; Illingworth et al. 2016; Whitaker et al. 2019), to visually search for LyC emission around MOSEL targets. The mosaic relies on the archival HST imaging and hence has non-uniform coverage and depth across the FRESCO field of view (FOV). Out of 76 EELGs identified by Forrest et al. (2018) in the CDFS, 54 fall within the FRESCO FOV. We only found significant emission around one EELGs that happen to fall in the deepest region in the F336W image. Figure 1 shows the multi-band imaging for z19863, along with the unconvolved and PSF convolved F336W imaging, clearly indicating the possible LyC emission towards the NE side of the galaxy.

We estimate the signal-to-noise ratio (SNR) of LyC emission by placing an aperture of variable size at the pixel corresponding the peak emission in both the unconvolved and PSF-convolved F336W images (Figure 3). In the uncon-



Figure 2. The best-fit SED model using MAGPHYS before (blue curve) and after (black curve) correcting for dust attenuation. The red dots and open black circles correspond to the observed flux and derived flux from the best-fit SED respectively. The cyan dots represents the two filters (F210M and F277W) that are excluded while modelling the SED to minimise contamination by bright [O11] 5007 and H α emission lines.



Figure 3. The SNR as a function of the aperture radius for the original (blue circle) and PSF-convolved (orange circle) F336W image. Each aperture is positioned at the peak LyC flux (cyan cross in Figure 1).

volved image, we reach a maximum SNR of 4.0 within an aperture of r = 0."12, corresponding to $m_{F336W} = 28.7^{+0.3}_{-0.2}$. The SNR drops marginally to 3.95 at r = 0."24 corresponding to the $m_{F336W} = 27.9^{+0.3}_{0.2}$, a slightly larger aperture size than the FWHM (0."17) of the F336W filter. The SNR drops to 2.4 at r = 0.32'' and results in a marginal drop in flux ($m_{F336W} = 28.0^{+0.4}_{-0.3}$). In the PSF-convolved image, the SNR reaches a maximum of 3.3 within an aperture of r = 0.2''. The PSF-convolved image does indicated some weak evidence of extended LyC emission. Using an elliptical aperture (a = 0."28, b/a = 0.7, $\theta = 120^\circ$), results in only a marginal increase in the SNR; therefore, significant flux is not lost by the choice of a circular aperture.

3.2.1. Detection reliability test

To test the reliability of our detection, we check for the growth curve at random locations in the $1' \times 1'$ region covering the deepest F336W data. We use the F160W segmentation map to mask out all bright targets in the F336W image. We

Table 1.	Physical	properties	of z19863
----------	----------	------------	-----------

RA	53.1699	
DEC	-27.7684	
Z	3.088	
$\log(M_*/\mathrm{M}_{\odot})$	9.2 ± 0.1	
$\mathrm{SFR}_{\mathrm{H}lpha}$ [M $_{\odot}$ /yr]	4.3 ± 0.09	
SFR _{SED} [M _☉ /yr]	5.2 ± 0.2	
$E(B-V)_{gas}$	0.09 ± 0.02	
12 + log(O/H)	$7.79^{+0.06}_{-0.05}$	
logU	$-3.27^{+0.14}_{-0.12}$	
$L_{1500}/L_{900} = 6.66$, MAGPHYS		
f_{esc}^{rel}	2.8 ± 0.7	
f_{esc}^{abs}	1.2 ± 0.3	
$L_{1500}/L_{900} = 1.25$, lower limit		
f_{esc}^{rel}	0.5 ± 0.1	
f_{esc}^{abs}	0.24 ± 0.06	

then place two circular apertures of r = 0.12'', and 0.24'' at 20,000 random locations. We only measure the growth curve if the percentage of unmasked pixel within the largest aperture was greater than 90%. Only one random location out of nearly 16470 possible locations had S/N \geq 4.0 within aperture r = 0.12'' and S/N \geq 3.9 within r = 0.24'' aperture. There is clear emission in the F606W image at the same random location, indicating that the false detection is an interloper galaxy. We do not detect any signature of any interlopers in either HST or JWST imaging and MUSE datacube coincident with the LyC detection. Thus the probability of randomly detecting a similar source is less than $p < 6 \times 10^{-5}$.

3.2.2. Spatial offset of ionising emission

The peak non-ionizing UV emission (F606W) and ionizing emission have a spatial offset of $0.29'' \pm 0.04''$ or 2.2 ± 0.3 kpc



Figure 4. JWST medium band F210M (left: stars+gas), F182M (middle: stars), and F210M-F182M (right: gas) images for the galaxy z19863 at z = 3.088. The red plus marks the peak emission in F606W image (non-ionising UV emission), and the cyan cross marks the peak LyC emission in F336W.

at z = 3.088. We checked for astrometric alignment between the F336W and F606W images by estimating the difference in the centroids of the 20 brightest sources in the deepest $1' \times 1'$ FOV F336W image. The average difference in x-centroid [y-centroid] is 0 ± 1 [0 ± 1] pixels, significantly smaller than the peak separation (~ 7 pixels) between ionizing and nonionizing emission from our target. Therefore, the astrometric inaccuracies cannot account for the spatial offset.

3.2.3. LyC escape fraction

The escape fraction is defined as the ratio of ionizing (f_{LvC}) at $\lambda < 912$ Å) divided by the non-ionizing flux density (f_{nLvC} typically at 1500 Å, Steidel et al. 2001). We use the standard procedure to calculate escape fraction (e.g., Siana et al. 2007; Ji et al. 2020). As mentioned earlier, we use the F606W filter (rest-frame ~ 1500 Å) to measure non-ionizing photon flux density. The ionizing photons can sometimes escape from certain parts of the galaxy (Verhamme et al. 2015), therefore we use the same aperture to estimate flux for both ionizing and non-ionizing UV components. Using an aperture centered at the peak LyC flux and r = 0.12'' gives $(f_{LyC})_{obs} =$ $12.5 \pm 3.1 \text{ nJy}$ and $(f_{nLyC})_{obs} = 11.7 \pm 1.2 \text{ nJy}$. If we use an aperture of r = 0.24'', then $(f_{LyC})_{obs} = 24.5 \pm 6.2 \text{ nJy}$ and $(f_{\rm nLyC})_{\rm obs} = 89.9 \pm 2.5 \text{ nJy. Thus, } (f_{\rm nLyC}/f_{\rm LyC})_{\rm obs} = 0.9 \pm 0.3$ within the r = 0.12'' aperture and $(f_{nLyC}/f_{LyC})_{obs} = 3.7 \pm 0.9$ within the r = 0.24'' aperture. This is done after convolving the F336W and F606W images to a common PSF of F606W.

We need precise measurements of intrinsic non-ionizing to ionizing photon luminosity (L_{1500}/L_{900}) , accurate IGM attenuation (T_{IGM}) at the redshift of our galaxy, and the dust attenuation (A_{1500}) to estimate relative and absolute escape fraction. We use the IGM transmission of $T_{IGM} = 0.66$ at z = 3.08 calculated by Kerutt et al. (2023), which uses the mean transmission from the 5% of the brightest sightlines based on the Inoue et al. (2014) model. The adopted T_{IGM} sits comfortably in the range calculated for z = 3.1 Lyman- α emitters by (Bassett et al. 2021, see their Figure 6), taking into account the bias of detected LyC emitters along clean lines in the IGM.

Using $E(B - V) = 0.09 \pm 0.02$ (see section 3.4.1) and the extinction curve for the Small Magellanic Cloud (SMC) from Gordon et al. (2003), we obtain $A_{1500} = 1.2 \pm 0.3$. We note that the NIRSpec spectra do not cover the region where LyC emission is detected. The region occupying the bulk of gas emitting H α and H β emission lines can be different from the region with bright OB-type stars, as has been postulated by other studies (Vanzella et al. 2020, 2022; Meštrić et al. 2023). Therefore, the extinction towards the LyC region could be significantly different.

Finally, the intrinsic luminosity ratio L_{1500}/L_{900} can vary between 1.25 - 7 depending on the assumptions in the stellar population model (Leitherer et al. 1999; Eldridge et al. 2017). The best-fit SED model from MAGPHYS (Section 3.1) gives $L_{1500}/L_{900} = 6.66$, resulting in $f_{esc}^{rel} = 2.8 \pm 0.7$, and $f_{esc}^{abs} =$ 1.3 ± 0.3 . An escape fraction > 1 is unphysical because it implies the galaxy is leaking more than the ionizing photons produced in this ionization cone. Note that the properties of the stellar population might not be the same throughout the galaxy. Even assuming a more typically used $L_{1500}/L_{900} = 3$, we get $f_{\rm esc}^{\rm rel} = 1.2 \pm 0.3$ and $f_{\rm esc}^{\rm abs} = 0.6 \pm 0.1$. Using the lower limit on $L_{1500}/L_{900} = 1.25$ gives us the most conservative estimate on the relative escape fraction of $f_{\rm esc}^{\rm rel} = 0.5 \pm 0.1$ and a corresponding $f_{esc}^{abs} = 0.24 \pm 0.06$. A future paper will analyze the resolved stellar population properties of z19863 and test the impact different stellar population models have on the escape fraction measurements.

3.3. Merger activity

We use medium-band filters F210M and F182M to analyze the merger activity of our source. At z = 3.088, the medium-





Figure 5. NIRSpec spectrum (JADES NIRSpec ID = 21150, NIRCam ID = 211968) from the JADES in the three medium band grisms, G140M/F070LP (top), G235M/F170LP (middle), and G395M/F290LP (bottom). The grey dashed curve in each panel corresponds to the error spectrum in the respective grism. The low-dispersion prism spectrum is shown as the solid black curve in each panel and has significantly lower error spectrum than displayed ranges here. The black dashed lines mark the location of bright emission lines detected above the 5σ level in each sub-panel. The inset image in the top panel shows the location of the slit with respect to the JWST/NIRCam image (taken from the Figure B.1 in Bunker et al. 2023).

band filter F210M would contain emission from stars and the [O III] 5007+H β emission lines, whereas filter F182M would only have stellar emission. Therefore, the difference image between the two filters provides a spatially resolved distribution of gas emission ([O III] 5007+H β) from the target (Figure 4). We use StatMorph (Rodriguez-Gomez et al. 2019) to estimate non-parametric quantities for all three filters (more details in Gupta et al., in prep.).

The Gini-M20 statistics (Lotz et al. 2008) classify z19863 as a merger based on the stellar emission (F182M) but a non-merger based on the gas emission (F210M-F182M). The peak gas emission (F210M-F182M) is coincident with the peak non-ionizing UV emission (F606W), whereas the peak stellar emission (F182M) is 0.18" SW of the non-ionizing UV emission (F606W). The lack of merger signs in the gas emission could be because hydrodynamical cooling results in faster settling time for the gas compared to stars after a merger event (See Gupta et al., in prep. for more discussion). The gas emission (F210M-F182M) seems to be extended towards the position of the LyC detection, indicating that LyC photons could be escaping from a recently star-forming part of the galaxy. A detailed analysis of spatially-resolved spectral properties of z19863 is beyond the scope of this paper and will be part of future analysis

3.4. JADES/NIRSpec observations

Figure 5 shows the JWST/NIRSpec spectrum for z19863 from the JADES (NIRSpec ID = 21150, NIRCam ID = 211968; Bunker et al. 2023) spanning all the way from $1 - 5\mu m$, showing multiple bright emission lines. Note that the 0.46" microshutter only partially covers the galaxy and only on the side opposite to the LyC detection. We cannot use the NIRSpec spectra to rule out if the emission in the F336W image is due to an interloper galaxy. Even with this partial coverage, we are able to detect a whole suite of emission lines to estimate the gas-phase metallicity and ionization conditions in the galaxy.

The emission lines fluxes in the JADES catalogs have been aperture-corrected using the location of micro-shutter on the NIRCam imaging. The analysis in the following subsections assumes that the gas has similar properties across the entire galaxy, which might not be correct. Neither [NII] 6583 or [SII] 6717,6731 are detected at S/N> 5 with grism, therefore we rely on the JADES prism catalogs to estimate ISM conditions.



Figure 6. The location of z19863 (golden circle) on the R23 - O32 (left) and N2 - R2 (middle) and N2 - R2 (right) diagrams, typically is used to probe the metallicity and ionization conditions of the ISM. The $z \sim 6$ galaxies from Cameron et al. (2023) are shown as salmon diamonds. We also compare with other extreme galaxies in the local universe (blueberries from Yang et al. (2017a): purple circles, green peas from Yang et al. (2017b): green plus, and LyC leakers from Izotov et al. (2018): green hexagons). The grey shaded regions and the orange dots in each panel represent the typical SFGs at $z \sim 0$ from the SDSS survey (Aihara et al. 2011) and at $z \sim 2$ from the MOSDEF survey (Kriek et al. 2015) respectively. The solid red curve in the left panel and dashed red curve in the middle and right panels shows the Lamareille et al. (2004) and (Kewley et al. 2001) relation for identifying AGNs respectively. The blue dotted curves in the middle panel represents the Kauffmann et al. (2003) relation for star-forming galaxies. Our galaxy falls in the star-forming region occupied by metal-poor galaxies, similar to z > 6 galaxies and other extreme galaxies in the local universe.

3.4.1. Dust Extinction and star formation

We use the Gordon et al. (2003) extinction curve for the Small Magellanic Cloud to correct for dust extinction, following similar approach used in previous studies of high redshift galaxies (Reddy et al. 2023; Cameron et al. 2023). We use the Balmer decrement based on the ratio of H α and H β emission lines to estimate the color excess using the equation:

$$E(B-V) = \frac{2.5}{\kappa(\mathrm{H}\beta) - \kappa(\mathrm{H}\alpha)} \log_{10}\left(\frac{(\mathrm{H}\alpha/\mathrm{H}\beta)_{\mathrm{obs}}}{2.98}\right), \quad (1)$$

The Balmer decrement for z19863 is $f(H\alpha/H\beta)_{obs} = 3.26 \pm 0.07$, which results in $E(B - V) = 0.09 \pm 0.02$.

The dust-corrected SFR based on H α and using the Kennicutt Jr & Evans II (2012) relation, is $4.3 \pm 0.09 \text{ M}_{\odot}/\text{yr}$ for this galaxy, placing it within the main-sequence at z = 3 (Popesso et al. 2022). It's important to note that the H α luminosity used for this calculation is rescaled based on the position of the microshutter on the NIRCam image of the galaxy. However, the pseudo [OIII] 5007 emission map reveals that the microshutter misses the most star-forming part of the galaxy (Figure 4). Therefore, the total H α luminosity and the instantaneous SFR of z19863 may differ significantly from the value estimated here.

3.4.2. ionization conditions

The left panel in Figure 6 displays the extinction-corrected R23 versus O32 relation of z19863 compared to other LyC emitter samples and galaxies at z > 6, where:

$$R23 = \log_{10}(([O \text{ III}] 5007, 4959 + [O \text{ II}] 3727)/\text{H}\beta),$$

$$O32 = \log_{10}([OIII] 5007/[OII] 3727).$$

To correct for dust extinction for low redshift samples, we apply the CCM89 extinction curve ($R_V = 3.1$, Cardelli et al.

1989). The O32 and R23 ratios of z19863 are higher than those of typical star-forming galaxies both in the local universe (Aihara et al. 2011) and at $z \sim 2$ (Shapley et al. 2014; Kriek et al. 2015). The emission line ratios of z19863 resemble those of other extreme galaxy populations such as green peas and blueberries in the local universe. However, its O32 ratio is not as high as that of other LyC emitters at $z \sim 0.3$. z19863 also exhibits a low S2, where

$$S2 = \log_{10}(([S_{II}] 6717, 6731)/H\alpha),$$

compared to typical star-forming galaxies at $z \sim 2$, which has also been indicated as a marker for LyC leakers (Ramambason et al. 2020).

Interestingly, we have detections of both $[S_{III}]$ 9069,9531 and $[S_{II}]$ 6717,6731 in the prism spectrum of z19863. Therefore, we can use the *S*32 ratio defined as:

$$S32 = log_{10}([SIII] 9069, 9531/[SII] 6717, 6731),$$

to estimate the ionization parameter, a metallicity- and pressure-independent ionization parameter diagnostic. Traditionally, the O32 ratio is used to estimate the ionization parameter because [S III] 9069,9531 either redshifts into the near-infrared at z > 0.05 or beyond $2.4\mu m$ at z > 1.5, making it difficult to observe. However, ionization parameters based on O32 are highly dependent on the choice of stellar template (Sanders et al. 2015). Sanders et al. (2019) used stacked measurements of [S III] 9069,9531 in $z \sim 1.5$ galaxies to and confirmed its weaker dependence on metallicity compared to O32. Recent observations have started to use [S III] 9069,9531 to estimate the ionization conditions in galaxies (Mingozzi et al. 2020; Kumari et al. 2021).

The S32 ratio of z19863 is 0.27 ± 0.09 constraining the ionization parameter to $logU = -3.25 \pm 0.10$. This suggests that z19863 has a similar ionization parameter to typical SFGs at $z \sim 2-3$ (Sanders et al. 2015), and significantly lower than galaxies at z > 6. Again, we note that NIRSpec observations only cover a small portion of the galaxy. Spatially resolved observations of star-bursting galaxies find significant variations in metallicity and ionization parameter (del Valle-Espinosa et al. 2023), indicating that NIRSpec spectra might not be representative of the average gas conditions in the galaxy.

3.4.3. Gas-phase Metallicity

The [N II] 6583 emission line is not detected in either the grism or prism spectrum, indicative of very low metallicity. We use the grism spectra to estimate an upper limit on [N II] 6583 because [N II] 6583 is blended with H α in the prism spectrum. The [N II] 6583/H α versus [O III] 5007/H β diagram (Figure 6) places the galaxy well within the region occupied by low-metallicity star-forming galaxies both in the low and high-redshift universe.

We use the S23 diagnostic from (Kewley et al. 2019) to estimate the gas-phase metallicity, where

$$S23 = log_{10}(([SII] 6717, 6731 + [SIII] 9069, 9531)/H\alpha),$$

due to its sensitivity at lower metallicity and the lack of [NII] 6583 detection. Using an S23 ratio of -0.75 ± 0.05 and log U = -3.25, we estimate the gas-phase metallicity at $12 + \log(O/H) = 7.79^{+0.06}_{-0.05}$ ($Z = 0.13 \pm 0.01 Z_{\odot}$). Thus, the gas-phase metallicity of our galaxy is fairly consistent with gas-poor galaxies detected at higher redshifts (Tang et al. 2023; Boyett et al. 2024).

3.5. Rest-UV spectrum

We extract the rest-UV spectrum of our target using the mosaic datacube from the MXDF survey (Bacon et al. 2023). The mosaic datacube reaches a total integration time of 12 hours at the position of our target (peak flux in F606W filter). We search for emission around the expected Lyman- α wavelength at z = 3.088, and identify the peak wavelength. We sum over flux within ± 10 Å around the peak Lyman- α wavelength to create a line map and add spaxels above $2 \times \text{RMS}_{\text{sky}}$ to extract the 1D spectrum (Figure 7). The peak Lyman- α emission has a spatial offset of 0."42 with respect to peak flux in F606W filter, which is similar to the typical FWHM of the MUSE observations.

The Lyman- α emission is highly asymmetric with an undetected blue-side, similar to other high-redshift Lyman- α emitters (Gronke & Dijkstra 2016). The red peak of the Lyman- α emission has a marginal velocity offset from the systematic redshift of the galaxy (approximately 100 km/s). There is significant contamination in the rest-UV spectrum from another Lyman- α emitter at z = 3.4, which is only 0.85" from the F606W peak (indicated as a black plus in Figure 1). The high spatial resolution of HST and

JWST ensures negligible contamination in the photometry of z19863 from this source at any wavelengths including the F336W image. We have clear detections of many low (Si II λ 1190, Si II λ 1260, CII λ 1334, Si II λ 1526) and high ionization (Si IV $\lambda\lambda$ 1393, 1402, CIV $\lambda\lambda$ 1548, 1550) metal absorption features. Interestingly, we do not detect any high ionization emission lines such as [OIII] 1660,1666, HeII 1640, [CIII] 1907,1909, which are commonly detected in other LyC emitters (Vanzella et al. 2019). Ji et al. (2020) also report a tentative detection of HeII 1640 (S/N~ 2) in Ion1, indicating the need for a deeper rest-UV spectrum.

4. DISCUSSION

4.1. Curious case of z19863

This paper presents the discovery of a new LyC emitter candidate, z19863 at redshift z = 3.088, combining archival data from deep imaging campaigns conducted on the HST and JWST (Figure 1). Detailed information about the source is provided in Table 1. Using the deep NIR-Spec spectrum from JADES we find that z19863 has characteristics similar to other LyC emitters across different redshift ranges (see Figure 6), including a moderate ionization parameter (log $U = -3.25 \pm 0.1$), a low gas-phase metallicity $(12+\log(O/H) = 7.76 \pm 0.06)$, and a high O32 ratio (3.65 ± 0.22) . Notably, the LyC emission is spatially offset by $0.29'' \pm 0.04''$ (equivalent to 7 ± 1 pixels or approximately 2.2 kpc) from the peak rest-UV (non-ionizing) emission (Figure 1). The astrometric misalignment is insufficient to account for the spatial offset ($dx = \pm 1$ pixel). We do not detect any interloper galaxy co-incident with the LyC detection in extremely deep spectroscopy from the MXDF survey (Figure 7). We estimate the probability of randomly detecting a similar source within a similar aperture (0.24'') in the F336W image is $p < 6 \times 10^{-5}$.

Most observations focused on detecting LyC photons typically search for emission at $\lambda_{rest} < 912$ Å co-spatial with the non-ionising UV emission. For example, Kerutt et al. (2023) look for LyC emission within 0."35 aperture around the non-ionising rest-UV emission, and therefore would miss a source similar to z19863 with about 0."29 offset between the ionising and non-ionising emission peaks. Ion1 (z = 3.7) is another know LyC emitter that exhibits a spatial offset between the peak rest-UV and LyC emission (0."12 ~ 0.85 kpc, Ji et al. 2020).

Additionally, z19863 has a single peaked Lyman- α emission profile z19863 (Figure 7), atypical to LyC emitters in the local universe where a double-peaked profile with a narrow velocity separation is common (Izotov et al. 2018, 2022; Flury et al. 2022b). Models of LyC and Lyman- α escape suggest that a narrow velocity separation is indicative of an optically thin neutral ISM, which, in turn, facilitates the escape of ionising photons (Izotov et al. 2018). The green pea



Figure 7. Rest-UV 1D spectrum extracted from the MXDF mosaic (12 hours integration). The inset image shows the zoomed-in spectrum at the location of Lyman- α emission line. The dashed vertical lines mark the location of various absorption features. The dashed orange curve is the corresponding error spectrum. The grey shaded region highlights the Lyman- α emission from the nearby z = 3.4 galaxy.

galaxies with high O32 ratios (are more likely to leak LyC photons) have lower HI gas mass, suggesting they are more likely to be optically thin (Kanekar et al. 2021). However, LyC emitters at high redshift exhibit a range of Lyman- α profiles, including 3-4 peaks (Izotov et al. 2022), single peaked for Q1549-C25 (Shapley et al. 2016), and absorption rather than emission for Ion1 (Ji et al. 2020). Kerutt et al. (2023) uses a larger sample of LyC emitters between 3 < z < 4 and find no correlation between the Lyman- α emission and the escape fraction. It is possible that at higher redshifts, most ionising photons escape because of the inhomogeneities in the ISM rather than an optically thin ISM.

4.2. Mergers driving the LyC escape

We currently have two models explaining LyC escape: through an optically thin ISM and the picket-fence model (Zackrisson et al. 2013). In the picket-fence model, LyC photons escape through a few optically thin channels in an otherwise optically thick ISM. These channels are typically created by star-formation-triggered outflows capable of blowing out neutral gas. LyC emitters in the local universe often show a significant correlation between the escape fraction and the strength of ionized gas outflows, suggesting that star formation feedback is responsible for creating these ionized gas channels (Amorín et al. 2024).

However, the mass loading factors for most high-redshift galaxies are similar to those of local dwarf galaxies, even if their star formation rates (SFRs) are ten times higher (Gupta et al. 2022; Concas et al. 2022; Llerena et al. 2023; Carniani et al. 2023). Some semi-analytic models of star-formation-driven galactic winds suggest that outflowing gas in super star-forming conditions might start radiatively cooling on a relatively short timescale (< 1 Gyr, Lochhaas et al. 2018). This rapid cooling might hinder the creation of large-scale channels of optically thin neutral gas solely from star formation alone.

Mergers and galaxy-galaxy interactions are known to tidally strip gas and stars, leading to the formation of highly inhomogeneous galaxies (Toomre & Toomre 1972; Cox et al. 2008; Spilker et al. 2022). Very few LyC emitters in the local universe are close enough to spatially map their neutral gas density. Deep 21cm observations on Haro 11, a local LyC emitter, revealed pockets of optically thin neutral hydrogen most likely created by a recent merger event (Reste et al. 2023). The large spatial offset (6 kpc) between the peak neutral gas density and the LyC emitting clump suggests that LyC photons are most likely leaking through these optically thin channels (Reste et al. 2023). On the other hand, two other local LyC leakers with 21cm observations exhibit either only an upper limit on HI mass (Puschnig et al. 2017) or very high gas mass (Haynes et al. 2018), indicating a wide range in the properties LyC emitters in the local universe.

Recent work by Gupta et al. (2023) finds that EELGs at $z \sim 3.5$, analogs of galaxies at z > 6, are more likely to encounter major mergers and/or strong interactions. Other observations (e.g., Duncan et al. 2019) also find increased merger activity at higher redshifts. For z19863, medium band images from the FRESCO survey indicate that the galaxy might have encountered a merger recently (Figure 4). Bassett et al. (2018) suggest that the lack of correlation between the O32 ratio and escape fraction could be due to anisotropic neutral gas coverage. In fact, all of their LyC candidates with the lowest O32 ratios showed evidence of disturbed morphology. The Lyman- α emission peak for the candidate LyC emitters in Kerutt et al. (2023) is spatially offset by 0.3'' - 0.5'' from the peak LyC emission, indicating a mismatch between the peak neutral gas density and ionising stars. We suspect that inhomogeneities created by a merger event in the neutral ISM might have played a significant role in aiding LyC photons to escape at higher redshifts.

We think in z19863 ISM is neither optically thin everywhere nor has a complete picket-fence structure with many optically thin ionizations cone. Here the ISM most likely has a single 10

optically thin ionization cone of neutral ISM created by a recent merger event. The ionising photons are leaking through this single channel, and by happenstance, its alignment along our line of sight ensures that we can detect the faint signature of ionising radiation leaking from this single ionization cone.

4.3. Modelling the opening angle of ionising channels

If LyC radiation leak through a few optically thin ionization cones, similar to z19832, for most galaxies, then their random alignment along the line of sight will dictate the number of confirmed detections of LyC emitters. Estimating the total covering fraction of optically thin ISM in high-redshift galaxies is extremely challenging. Lyman- α photons get fluorescently scattered by neutral hydrogen, and thus, a Lyman- α emission profile provides some hints about the neutral gas density. A small velocity separation and a large ratio of total flux between the blue and red peaks usually indicate an optically thin ISM (Erb et al. 2018; Blaizot et al. 2023; Mukherjee et al. 2023). However, spatially resolving Lyman- α emission is only possible for a small number of Lyman- α emitters at z > 3, until future spectrographs such as MAVIS (McDermid et al. 2020) become available.

Stacking experiments with large samples of SFGs at $z \sim 3$ have provided some constraints on the average $f_{esc} = 1 - 10\%$ (Steidel et al. 2018; Wang et al. 2023). Theoretical models using various observations about the reionization history of the universe, such as neutral gas density, UV luminosity function etc., constrain the average f_{esc} required (Naidu et al. 2020; Bera et al. 2023; Mutch et al. 2023). However, some individual LyC emitters can have escape fraction as high as 100% (Rivera-Thorsen et al. 2022; Kerutt et al. 2023).

We can use individual f_{esc} measurements to develop a simple model for the total covering fraction or the opening angle of the ionization cone required to reach global f_{esc} values. We assume that all galaxies in the universe have on average an ionization cone with solid angle Ω and escape fraction $f_{esc,ind}$. The following equation balances the total ionization flux from a single ionization cone with the average escape fraction expected over the entire galaxy ($f_{esc,global}$)

$$\Omega \times f_{esc,ind} = 4\pi \times f_{esc,global}, \implies \Omega = 4\pi \times \frac{f_{esc,global}}{f_{esc,ind}}.$$
(2)

For a cone with apex half angle of θ , the total solid angle is

$$\Omega = 4\pi \sin^2 \frac{\theta}{2}.$$
 (3)

Therefore, the half-angle of ionization cone would be

$$\theta[rad] = 2 \times \sin^{-1} \sqrt{\frac{f_{esc,global}}{f_{esc,ind}}}.$$
 (4)

Figure 8 shows the average half-angle estimated using equation 4 based on $f_{esc,ind}$ of z19863 and various measurements of $f_{\rm esc, \ global}$ from observations (Steidel et al. 2018; Wang et al. 2023) and theory (Naidu et al. 2020; Bera et al. 2023; Mutch et al. 2023). By definition the escape fraction cannot be greater than 1, therefore we assume an absolute escape fraction of 1.00 ± 0.3 rather than estimated value of $f_{esc}^{abs} = 1.25 \pm 0.3$ in Section 3.2. Additionally, the opening half-angle is estimated using $f_{\rm esc, \ ind}$ based on the average escape fraction for the five gold LyC emitter candidates from Kerutt et al. (2023).

Our simple model can help distinguish between more likely global escape fraction measurements. For instance, if the global escape fraction is constrained to be very low, such as $0.02^{+0.04}_{-0.01}$ (Bera et al. 2023), an ionization cone with an opening half-angle of approximately $\sim 16^{\circ}$ may be sufficient to account for this small escape fraction. Even if only 50% of ionizing photons leak from the ionization cone of most galaxies, the opening half-angle would only need to increase to approximately 23°. This scenario could provide an explanation for the limited number of LyC emitter detections at higher redshifts. On the other hand, to achieve a global escape fraction of 0.2 (Naidu et al. 2020), ionization cones with $f_{\text{esc, ind}} = 1.0 [f_{\text{esc, ind}} = 0.5]$ will need to have opening half-angle of ~ 53° [78°]. Such large opening angles imply that a significant fraction (20-50%) of the galaxy would need to be leaking ionizing radiations. However, if this were the case, one might expect a larger success rate in building a larger sample of confirmed LyC leakers.

We only need an opening half-angle of ~ 37° for fully ionised cone and ~ 53° for ionization cone with 50% escape fraction to explain the typically observed f_{esc} from observations (Steidel et al. 2018; Wang et al. 2023). It's worth noting that even at z = 3, a partially neutral intergalactic medium (IGM) can absorb a considerable fraction of ionizing radiations (Bassett et al. 2021), further complicating the detection of LyC emission. Altering the median opening angle of escaping ionising photons would have an impact on the topology of reionization, and a more sophisticated model would explore the effects of this change on the progress of reionization.

5. SUMMARY

This paper presents discovery of a new candidate LyC emitter (z19863) at z = 3.088, where LyC emission is detected using the F336W/UVIS filter on the HST. The galaxy is part of the EELG sample in the MOSEL survey (Tran et al. 2020; Gupta et al. 2022). The LyC emission is spatially offset by 0."29 \pm 0."04 $\sim 2.2 \pm 0.3$ kpc from the peak rest-UV emission (F606W). Based on the NIRSpec spectra from the JADES, we estimate that the galaxy has low metallicity, high ionization parameter, and low dust attenuation (See Table 1), similar to other LyC emitters at both low and high redshifts.



Figure 8. Relation between the opening half-angle of the ionization cone and the global f_{esc} values. The diamonds (purple: Bera et al. (2023), light green: Mutch et al. (2023), and dark green Naidu et al. (2020)) and circles (salmon: Wang et al. (2023), yellow: Steidel et al. (2018), and magenta: Begley et al. (2022)) show the theoretical and observational limits on the global f_{esc} respectively. The solid and open symbols use individual f_{esc} based on z19832 (this work) and average f_{esc} from five gold LyC candidates from Kerutt et al. (2023). The black dashed line is for the fully ionised channels ($f_{esc,ind} = 1$).

The deep rest-UV spectrum of the galaxy from the MXDF survey (12 hour integration) exhibit a single peaked Lyman- α emission profile, and a range of low- and high- ionization absorption features (Figure 7). The lack of the rest-UV emission lines such as [OIII] 1660,1666 CIV 1550, [CIII] 1907,1909 etc. that are typically used as proxy for LyC emission (Schaerer et al. 2022), suggest an ISM conditions in this galaxy are dissimilar to a typical LyC emitter in the local universe. Further analyses of the absorption profiles of various species and their correlation with LyC emission (Mauerhofer et al. 2021) is beyond the scope of this paper and will be part of future work.

We propose that LyC photons in z19863 are leaking through a narrow cone of optically thin ISM, likely created by a recent merger event as indicated by the disturbed morphology in the medium band imaging (see section 3.3). The ionization cone is spatially offset by $\sim 0.''3$ from the non-ionising UV emission and its chance alignment along the line of sight ensures its detection by us. A source similar to z19863 would be missed by studies looking for LyC emission co-incident with peak UV emission (e.g., Kerutt et al. 2023). Using a simple toy model, we estimate that the opening half-angle of ionization cones can be as low as 16° to explain some of the global measurements of escape fraction, indicating that very small fraction of the galaxy needs to be leaking ionising radiation. Such small opening angle would also explain the relatively low number of confirmed LyC emitters at high redshifts.

6. ACKNOWLEDGMENTS

This research were supported by the Australian Research Council Centre of Excellence for All Sky Astrophysics in 3 Dimensions (ASTRO 3D), through project number CE170100013.

REFERENCES

Aihara, H., Allende Prieto, C., An, D., et al. 2011, The Astrophysical Journal Supplement Series, 193, 29, doi: 10.1088/0067-0049/193/2/29 Amorín, R. O., Rodríguez-Henríquez, M., Fernández, V., et al. 2024, Astronomy & Astrophysics, 682, L25, doi: 10.1051/0004-6361/202449175

- Atek, H., Siana, B., Scarlata, C., et al. 2011, The Astrophysical Journal, 743, 121, doi: 10.1088/0004-637X/743/2/121
- Bacon, R., Brinchmann, J., Conseil, S., et al. 2023, Astronomy and Astrophysics, 670, A4, doi: 10.1051/0004-6361/202244187
- Barro, G., Pérez-González, P. G., Cava, A., et al. 2019, The Astrophysical Journal Supplement Series, 243, 22, doi: 10.3847/1538-4365/ab23f2

Bassett, R., Ryan-Weber, E. V., Cooke, J., et al. 2021, Monthly Notices of the Royal Astronomical Society, 502, 108, doi: 10.1093/mnras/stab070

 2018, Monthly Notices of the Royal Astronomical Society, 483, 5223, doi: 10.1093/mnras/sty3320

Begley, R., Cullen, F., McLure, R. J., et al. 2022, Monthly Notices of the Royal Astronomical Society, 513, 3510, doi: 10.1093/mnras/stac1067

Bera, A., Hassan, S., Smith, A., et al. 2023, The Astrophysical Journal, 959, 2, doi: 10.3847/1538-4357/ad05c0

Bian, F., Fan, X., McGreer, I., Cai, Z., & Jiang, L. 2017, The Astrophysical Journal, 837, L12, doi: 10.3847/2041-8213/aa5ff7

Blaizot, J., Garel, T., Verhamme, A., et al. 2023, Monthly Notices of the Royal Astronomical Society, 523, 3749, doi: 10.1093/mnras/stad1523

Boyett, K., Bunker, A. J., Curtis-Lake, E., et al. 2024, Extreme Emission Line Galaxies Detected in JADES JWST/NIRSpec I: Inferred Galaxy Properties, arXiv, doi: 10.48550/arXiv.2401.16934

Bruzual, G., & Charlot, S. 2003, Monthly Notices of the Royal Astronomical Society, 344, 1000, doi: 10.1046/j.1365-8711.2003.06897.x

Bunker, A. J., Cameron, A. J., Curtis-Lake, E., et al. 2023, JADES NIRSpec Initial Data Release for the Hubble Ultra Deep Field: Redshifts and Line Fluxes of Distant Galaxies from the Deepest JWST Cycle 1 NIRSpec Multi-Object Spectroscopy, doi: 10.48550/arXiv.2306.02467

Cameron, A. J., Saxena, A., Bunker, A. J., et al. 2023, JADES: Probing Interstellar Medium Conditions at \$z\sim5.5-9.5\$ with Ultra-Deep JWST/NIRSpec Spectroscopy, doi: 10.48550/arXiv.2302.04298

Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, The Astrophysical Journal, 345, 245, doi: 10.1086/167900

Carniani, S., Venturi, G., Parlanti, E., et al. 2023, JADES: The Incidence Rate and Properties of Galactic Outflows in Low-Mass Galaxies across 3 < z < 9, arXiv, doi: 10.48550/arXiv.2306.11801

Charlot, S., & Fall, S. M. 2000, The Astrophysical Journal, 539, 718, doi: 10.1086/309250

Concas, A., Maiolino, R., Curti, M., et al. 2022, Monthly Notices of the Royal Astronomical Society, 513, 2535, doi: 10.1093/mnras/stac1026

- Cox, T. J., Jonsson, P., Somerville, R. S., Primack, J. R., & Dekel, A. 2008, Monthly Notices of the Royal Astronomical Society, 384, 386, doi: 10.1111/j.1365-2966.2007.12730.x
- da Cunha, E., Charlot, S., & Elbaz, D. 2008, Monthly Notices of the Royal Astronomical Society, 388, 1595, doi: 10.1111/j.1365-2966.2008.13535.x
- da Cunha, E., Walter, F., Smail, I. R., et al. 2015, The Astrophysical Journal, 806, 110, doi: 10.1088/0004-637X/806/1/110
- de Barros, S., Vanzella, E., Amorín, R., et al. 2016, Astronomy & Astrophysics, 585, A51, doi: 10.1051/0004-6361/201527046

del Valle-Espinosa, M. G., Sánchez-Janssen, R., Amorín, R., et al. 2023, Monthly Notices of the Royal Astronomical Society, 522, 2089, doi: 10.1093/mnras/stad1087

Duncan, K., Conselice, C. J., Mundy, C., et al. 2019, The Astrophysical Journal, 876, 110, doi: 10.3847/1538-4357/ab148a

Eisenstein, D. J., Willott, C., Alberts, S., et al. 2023, Overview of the JWST Advanced Deep Extragalactic Survey (JADES), doi: 10.48550/arXiv.2306.02465

Eldridge, J. J., Stanway, E. R., Xiao, L., et al. 2017, Publications of the Astronomical Society of Australia, 34, e058, doi: 10.1017/pasa.2017.51

- Endsley, R., Stark, D. P., Chevallard, J., & Charlot, S. 2020, Monthly Notices of the Royal Astronomical Society, 500, 5229, doi: 10.1093/mnras/staa3370
- Endsley, R., Stark, D. P., Whitler, L., et al. 2023, The Star-forming and Ionizing Properties of Dwarf Z[~]6-9 Galaxies in JADES: Insights on Bursty Star Formation and Ionized Bubble Growth, arXiv, doi: 10.48550/arXiv.2306.05295
- Erb, D. K., Steidel, C. C., & Chen, Y. 2018, The Astrophysical Journal Letters, 862, L10, doi: 10.3847/2041-8213/aacff6

Flury, S. R., Jaskot, A. E., Ferguson, H. C., et al. 2022a, The Astrophysical Journal Supplement Series, 260, 1, doi: 10.3847/1538-4365/ac5331

 2022b, The Astrophysical Journal, 930, 126, doi: 10.3847/1538-4357/ac61e4

Forrest, B., Tran, K.-v. H., Broussard, A., et al. 2018, The Astrophysical Journal, 863, 131, doi: 10.3847/1538-4357/aad232

Giacconi, R., Zirm, A., Wang, J., et al. 2002, The Astrophysical Journal Supplement Series, 139, 369, doi: 10.1086/338927

Gordon, K. D., Clayton, G. C., Misselt, K. A., Landolt, A. U., & Wolff, M. J. 2003, The Astrophysical Journal, 594, 279, doi: 10.1086/376774

Gronke, M., & Dijkstra, M. 2016, The Astrophysical Journal, 826, 14, doi: 10.3847/0004-637x/826/1/14

Gupta, A., Tran, K.-V., Mendel, T., et al. 2022, Monthly Notices of the Royal Astronomical Society, 519, 980, doi: 10.1093/mnras/stac3548

13

Gupta, A., Jaiswar, R., Rodriguez-Gomez, V., et al. 2023, The Astrophysical Journal Letters, 957, L35, doi: 10.3847/2041-8213/ad0788

Hainline, K. N., Johnson, B. D., Robertson, B., et al. 2023, The Cosmos in Its Infancy: JADES Galaxy Candidates at z > 8 in GOODS-S and GOODS-N, doi: 10.48550/arXiv.2306.02468

Haynes, M. P., Giovanelli, R., Kent, B. R., et al. 2018, The Astrophysical Journal, 861, 49, doi: 10.3847/1538-4357/aac956

Illingworth, G., Magee, D., Bouwens, R., et al. 2016, The Hubble Legacy Fields (HLF-GOODS-S) v1.5 Data Products: Combining 2442 Orbits of GOODS-S/CDF-S Region ACS and WFC3/IR Images, doi: 10.48550/arXiv.1606.00841

Inoue, A. K., Shimizu, I., Iwata, I., & Tanaka, M. 2014, Monthly Notices of the Royal Astronomical Society, 442, 1805, doi: 10.1093/mnras/stu936

Izotov, Y. I., Chisholm, J., Worseck, G., et al. 2022, Monthly Notices of the Royal Astronomical Society, 515, 2864, doi: 10.1093/mnras/stac1899

Izotov, Y. I., Orlitová, I., Schaerer, D., et al. 2016a, Nature, 529, 178, doi: 10.1038/nature16456

Izotov, Y. I., Schaerer, D., Thuan, T. X., et al. 2016b, Monthly Notices of the Royal Astronomical Society, 461, 3683, doi: 10.1093/mnras/stw1205

Izotov, Y. I., Worseck, G., Schaerer, D., et al. 2018, Monthly Notices of the Royal Astronomical Society, 478, 4851, doi: 10.1093/mnras/sty1378

Ji, Z., Giavalisco, M., Vanzella, E., et al. 2020, The Astrophysical Journal, 888, 109, doi: 10.3847/1538-4357/ab5fdc

Kanekar, N., Ghosh, T., Rhoads, J., et al. 2021, The Astrophysical Journal, 913, L15, doi: 10.3847/2041-8213/abfb76

Kauffmann, G., Heckman, T. M., Tremonti, C., et al. 2003, Monthly Notices of the Royal Astronomical Society, 346, 1055, doi: 10.1111/j.1365-2966.2003.07154.x

Kennicutt Jr, R. C., & Evans II, N. J. 2012, Annual Review of Astronomy and Astrophysics, 50, 531, doi: 10.1146/annurev-astro-081811-125610

Kerutt, J., Oesch, P. A., Wisotzki, L., et al. 2023, Lyman Continuum Leaker Candidates at \$z\sim3-4\$ in the HDUV Based on a Spectroscopic Sample of MUSE LAEs, arXiv, doi: 10.48550/arXiv.2312.08791

Kewley, L. J., a. Dopita, M., Sutherland, R. S., a. Heisler, C., & Trevena, J. 2001, The Astrophysical Journal, 556, 121, doi: 10.1086/321545

Kewley, L. J., Nicholls, D. C., Sutherland, R., et al. 2019, The Astrophysical Journal, 880, 16, doi: 10.3847/1538-4357/ab16ed

Kriek, M., Shapley, A. E., Reddy, N. A., et al. 2015, The Astrophysical Journal Supplement Series, 218, 15, doi: 10.1088/0067-0049/218/2/15 Kumari, N., Amorín, R., Pérez-Montero, E., Vílchez, J., & Maiolino, R. 2021, Monthly Notices of the Royal Astronomical Society, 508, 1084, doi: 10.1093/mnras/stab2495

Labbé, I., Oesch, P. A., Bouwens, R. J., et al. 2013, The Astrophysical Journal, 777, L19, doi: 10.1088/2041-8205/777/2/L19

Lamareille, F., Mouhcine, M., Contini, T., Lewis, I., & Maddox, S. 2004, Monthly Notices of the Royal Astronomical Society, 350, 396, doi: 10.1111/j.1365-2966.2004.07697.x

Leitherer, C., Schaerer, D., Goldader, J. D., et al. 1999, The Astrophysical Journal, 123, 3, doi: 10.1086/313233

Llerena, M., Amorín, R., Pentericci, L., et al. 2023, Ionized Gas Kinematics and Chemical Abundances of Low-Mass Star-Forming Galaxies at \$z\sim 3\$, doi: 10.48550/arXiv.2303.01536

- Lochhaas, C., Thompson, T. A., Quataert, E., & Weinberg, D. H. 2018, Monthly Notices of the Royal Astronomical Society, 481, 1873, doi: 10.1093/mnras/sty2421
- Lotz, J. M., Davis, M., Faber, S. M., et al. 2008, The Astrophysical Journal, 672, 177, doi: 10.1086/523659

Lumbreras-Calle, A., López-Sanjuan, C., Sobral, D., et al. 2022, Astronomy and Astrophysics, 668, A60, doi: 10.1051/0004-6361/202142898

Mainali, R., Stark, D. P., Tang, M., et al. 2019, Monthly Notices of the Royal Astronomical Society, 494, 719, doi: 10.1093/mnras/staa751

Marques-Chaves, R., Schaerer, D., Amorín, R. O., et al. 2022, Astronomy and Astrophysics, 663, L1, doi: 10.1051/0004-6361/202243598

Maseda, M. V., Van Der Wel, A., Rix, H. W., et al. 2014, Astrophysical Journal, 791, 17, doi: 10.1088/0004-637X/791/1/17

Mauerhofer, V., Verhamme, A., Blaizot, J., et al. 2021, Astronomy & Astrophysics, 646, A80, doi: 10.1051/0004-6361/202039449

McDermid, R. M., Cresci, G., Rigaut, F., et al. 2020, 1, doi: 10.25949/zdaw-rx65

Meštrić, U., Ryan-Weber, E. V., Cooke, J., et al. 2020, Monthly Notices of the Royal Astronomical Society, 494, 4986, doi: 10.1093/mnras/staa920

Meštrić, U., Vanzella, E., Upadhyaya, A., et al. 2023, Astronomy & Astrophysics, 673, A50, doi: 10.1051/0004-6361/202345895

Mingozzi, M., Belfiore, F., Cresci, G., et al. 2020, Astronomy & Astrophysics, 636, A42, doi: 10.1051/0004-6361/201937203

Mukherjee, T., Zafar, T., Nanayakkara, T., et al. 2023, Compact to Extended Lyman-\$\alpha\$ Emitters in MAGPI: Strong Blue Peak Emission at \$z\gtrsim3\$, https://arxiv.org/abs/2311.07019v1

- Mutch, S. J., Greig, B., Qin, Y., Poole, G. B., & Wyithe, J. S. B.
 2023, Dark-Ages Reionization and Galaxy Formation Simulation
 XXI. Constraining the Evolution of the Ionizing Escape
 Fraction, arXiv, doi: 10.48550/arXiv.2303.07378
- Naidu, R. P., Tacchella, S., Mason, C. A., et al. 2020, The Astrophysical Journal, 892, 109, doi: 10.3847/1538-4357/ab7cc9

Oesch, P. A., Brammer, G., Naidu, R. P., et al. 2023, The JWST FRESCO Survey: Legacy NIRCam/Grism Spectroscopy and Imaging in the Two GOODS Fields, arXiv, doi: 10.48550/arXiv.2304.02026

Popesso, P., Concas, A., Cresci, G., et al. 2022, Monthly Notices of the Royal Astronomical Society, 519, 1526, doi: 10.1093/mnras/stac3214

Puschnig, J., Hayes, M., Östlin, G., et al. 2017, Monthly Notices of the Royal Astronomical Society, 469, 3252, doi: 10.1093/mnras/stx951

Rafelski, M., Teplitz, H. I., Gardner, J. P., et al. 2015, The Astronomical Journal, 150, 31, doi: 10.1088/0004-6256/150/1/31

Ramambason, L., Schaerer, D., Stasińska, G., et al. 2020, Astronomy and Astrophysics, 644, A21, doi: 10.1051/0004-6361/202038634

Reddy, N. A., Topping, M. W., Sanders, R. L., Shapley, A. E., & Brammer, G. 2023, The Astrophysical Journal, 948, 83, doi: 10.3847/1538-4357/acc869

Reste, A. L., Cannon, J. M., Hayes, M. J., et al. 2023, Tidally Offset Neutral Gas in Lyman Continuum Emitting Galaxy Haro 11, arXiv, doi: 10.48550/arXiv.2301.02676

Rieke, M., & the JADES Collaboration. 2023, JADES Initial Data Release for the Hubble Ultra Deep Field: Revealing the Faint Infrared Sky with Deep JWST NIRCam Imaging, doi: 10.48550/arXiv.2306.02466

Rinaldi, P., Caputi, K. I., Costantin, L., et al. 2023, The Astrophysical Journal, 952, 143, doi: 10.3847/1538-4357/acdc27

Rivera-Thorsen, T. E., Hayes, M., & Melinder, J. 2022, Astronomy and Astrophysics, 666, A145, doi: 10.1051/0004-6361/202243678

Rivera-Thorsen, T. E., Dahle, H., Chisholm, J., et al. 2019, Science, 366, 738, doi: 10.1126/science.aaw0978

Roberts-Borsani, G. W., Bouwens, R. J., Oesch, P. A., et al. 2016, The Astrophysical Journal, 823, 143, doi: 10.3847/0004-637x/823/2/143

Rodriguez-Gomez, V., Snyder, G. F., Lotz, J. M., et al. 2019, Monthly Notices of the Royal Astronomical Society, 483, 4140, doi: 10.1093/mnras/sty3345

Sanders, R. L., Shapley, A. E., Kriek, M., et al. 2015, The Astrophysical Journal, 816, 23, doi: 10.3847/0004-637X/816/1/23 Sanders, R. L., Jones, T., Shapley, A. E., et al. 2019, The Astrophysical Journal Letters, 888, L11, doi: 10.3847/2041-8213/ab5d40

Saxena, A., Pentericci, L., Ellis, R. S., et al. 2022, Monthly Notices of the Royal Astronomical Society, 511, 120, doi: 10.1093/mnras/stab3728

Schaerer, D., Izotov, Y. I., Worseck, G., et al. 2022, Astronomy & Astrophysics, 658, L11, doi: 10.1051/0004-6361/202243149

Shapley, A. E., Steidel, C. C., Strom, A. L., et al. 2016, The Astrophysical Journal, 826, L24, doi: 10.3847/2041-8205/826/2/L24

- Shapley, A. E., Reddy, N. A., Kriek, M., et al. 2014, Astrophysical Journal, 801, 88, doi: 10.1088/0004-637X/801/2/88
- Siana, B., Teplitz, H. I., Colbert, J., et al. 2007, The Astrophysical Journal, 668, 62, doi: 10.1086/521185

Spilker, J., Suess, K., Setton, D., et al. 2022, The Astrophysical Journal Letters, 936, L11, doi: 10.3847/2041-8213/ac75ea

Steidel, C. C., Bogosavljević, M., Shapley, A. E., et al. 2018, The Astrophysical Journal, 869, 123, doi: 10.3847/1538-4357/aaed28

Steidel, C. C., Pettini, M., & Adelberger, K. L. 2001, The Astrophysical Journal, 546, 665, doi: 10.1086/318323

Straatman, C. M. S., Spitler, L. R., Quadri, R. F., et al. 2016, The Astrophysical Journal, 830, 1, doi: 10.3847/0004-637X/830/1/51

Tang, M., Stark, D. P., Chevallard, J., & Charlot, S. 2019, Monthly Notices of the Royal Astronomical Society, 489, 2572, doi: 10.1093/mnras/stz2236

Tang, M., Stark, D. P., Chen, Z., et al. 2023, JWST/NIRSpec
Spectroscopy of \$z=7-9\$ Star Forming Galaxies with CEERS:
New Insight into Bright Ly\$\alpha\$ Emitters in Ionized
Bubbles, arXiv, doi: 10.48550/arXiv.2301.07072

Teplitz, H. I., Rafelski, M., Kurczynski, P., et al. 2013, The Astronomical Journal, 146, 159, doi: 10.1088/0004-6256/146/6/159

Toomre, A., & Toomre, J. 1972, The Astrophysical Journal, 178, 623, doi: 10.1086/151823

Tran, K.-V. H., Forrest, B., Alcorn, L. Y., et al. 2020, The Astrophysical Journal, 898, 45, doi: 10.3847/1538-4357/ab8cba

Urrutia, T., Wisotzki, L., Kerutt, J., et al. 2019, Astronomy & Astrophysics, 624, A141, doi: 10.1051/0004-6361/201834656

Vanzella, E., Giavalisco, M., Inoue, A. K., et al. 2010, The Astrophysical Journal, 725, 1011, doi: 10.1088/0004-637X/725/1/1011

Vanzella, E., Guo, Y., Giavalisco, M., et al. 2012, The Astrophysical Journal, 751, 70, doi: 10.1088/0004-637X/751/1/70

Vanzella, E., Barros, S. D., Cupani, G., et al. 2016, The Astrophysical Journal, 821, L27, doi: 10.3847/2041-8205/821/2/L27

Vanzella, E., Nonino, M., Cupani, G., et al. 2018, Monthly Notices of the Royal Astronomical Society: Letters, 476, L15, doi: 10.1093/mnrasl/sly023

- Vanzella, E., Caminha, G. B., Calura, F., et al. 2019, Monthly Notices of the Royal Astronomical Society, 000, doi: 10.1093/mnras/stz2286
- 2020, Monthly Notices of the Royal Astronomical Society, 491, 1093, doi: 10.1093/mnras/stz2286
- Vanzella, E., Castellano, M., Bergamini, P., et al. 2022, Astronomy and Astrophysics, 659, A2, doi: 10.1051/0004-6361/202141590
- Verhamme, A., Orlitová, I., Schaerer, D., & Hayes, M. 2015, Astronomy and Astrophysics, 578, 1, doi: 10.1051/0004-6361/201423978
- Wang, X., Teplitz, H. I., Smith, B. M., et al. 2023, The Lyman Continuum Escape Fraction of Star-forming Galaxies at \$2.4\lesssim Z\lesssim3.7\$ from UVCANDELS, doi: 10.48550/arXiv.2308.09064
- Whitaker, K. E., Ashas, M., Illingworth, G., et al. 2019, The Astrophysical Journal Supplement Series, 244, 16, doi: 10.3847/1538-4365/ab3853

- Williams, C. C., Tacchella, S., Maseda, M. V., et al. 2023, JEMS: A Deep Medium-Band Imaging Survey in the Hubble Ultra-Deep Field with JWST NIRCam & NIRISS, doi: 10.48550/arXiv.2301.09780
- Yang, H., Malhotra, S., Rhoads, J. E., & Wang, J. 2017a, The Astrophysical Journal, 847, 38, doi: 10.3847/1538-4357/aa8809
- Yang, H., Malhotra, S., Gronke, M., et al. 2017b, The Astrophysical Journal, 844, 171, doi: 10.3847/1538-4357/aa7d4d
- Yuma, S., Ouchi, M., Fujimoto, S., Kojima, T., & Sugahara, Y.
 2019, The Astrophysical Journal, 882, 17,
 doi: 10.3847/1538-4357/ab2f87
- Zackrisson, E., Inoue, A. K., & Jensen, H. 2013, Astrophysical Journal, 777, doi: 10.1088/0004-637X/777/1/39