Nonparametric density estimation for stationary processes under multiplicative measurement errors

Dang Duc Trong^{a,b}, Hoang Van Ha^{a,b}, and Thai Phuc Hung^{a,b,c}

^aFaculty of Mathematics and Computer Science, University of Science, Ho Chi Minh City, Vietnam.

^bVietnam National University, Ho Chi Minh City, Vietnam.

^cFaculty of Basics, Soc Trang Community College, Soc Trang Province, Viet Nam.

March 21, 2024

Abstract

This paper focuses on estimating the invariant density function f_X of the strongly mixing stationary process X_t in the multiplicative measurement errors model Y_t = X_tU_t , where U_t is also a strongly mixing stationary process. We propose a novel approach to handle non-independent data, typical in real-world scenarios. For instance, data collected from various groups may exhibit interdependencies within each group, resembling data generated from m-dependent stationary processes, a subset of stationary processes. This study extends the applicability of the model $Y_t = X_t U_t$ to diverse scientific domains dealing with complex dependent data. The paper outlines our estimation techniques, discusses convergence rates, establishes a lower bound on the minimax risk, and demonstrates the asymptotic normality of the estimator for f_X under smooth error distributions. Through examples and simulations, we showcase the efficacy of our estimator. The paper concludes by providing proofs for the presented theoretical results.

Mathematics Subject Classifications (2020): 62G05; 62G07; 60G10; 62G20.

Key words: Density estimation; multiplicative measurement errors; the Mellin transform; stationary processes; strongly mixing.

1 Introduction

Let X_{t_1}, \ldots, X_{t_n} be positive random variables generated from the strongly mixing stationary process X_t with unknown invariant density function $f_X : \mathbb{R}_+ \to \mathbb{R}_+$. Our objective is to estimate the density function f_X based on observations, while accounting for the presence of multiplicative measurement errors. In particular, let us assume that we have a sample consisting of observations Y_{t_1}, \ldots, Y_{t_n} according to the following model

$$
Y_{t_j}=X_{t_j}U_{t_j},\ j=1,\ldots,n.
$$

Here U_{t_1}, \ldots, U_{t_n} are error random variables generated from the strongly mixing stationary processes U_t with known invariant density function $g : \mathbb{R}_+ \to \mathbb{R}_+$. The processes X_t and U_t are independent. The density function g is called error density. Assume that the strongly mixing stationary processes are observed at discrete time $t_j = j\Delta$ (where Δ is a positive constant). For brevity, we will denote $X_{t_j}, Y_{t_j}, U_{t_j}$ by X_j, Y_j, U_j , respectively. Consequently, our model can be expressed as

$$
Y_j = X_j U_j, \ j = 1, \dots, n. \tag{1}
$$

In this setting the invariant density function $f_Y : \mathbb{R}_+ \to \mathbb{R}_+$ of the process $\{Y_i\}$ is given by

$$
f_Y(y) = [f_X * g](y) = \int_0^\infty f_X(x)g(y/x)x^{-1}dx, \forall y \in \mathbb{R}_+
$$

such that $*$ denotes multiplicative convolution. The estimation of f_X using a sample Y_1, \ldots, Y_n generated from the strongly mixing stationary process $Y_t = X_t U_t$ with density function f_Y is thus an inverse problem called "multiplicative deconvolution".

Model [\(1\)](#page-1-0) has found a lot of applications in various fields of research and has been extensively studied in the context where X_1, \ldots, X_n are independent and identically distributed (i.i.d.) samples from the distribution of X, and U_1, \ldots, U_n are i.i.d. samples from the distribution of U. In the field of survival analysis, as elucidated and motivated in Van Es $et al.[19]$ $et al.[19]$, when X represents the true positive survival time of a patient who has been sampled, and the random variable Y represents their observed survival time, it is reasonable to assume that the sampling time is uniformly distributed throughout the entire survival period of length X. Hence, we have $Y = XU$, where U follows a uniform distribution on the interval $[0, 1]$. Additionally, in survey sampling, model (1) is also employed to investigate sensitive information among respondents, where obtaining accurate responses can be challenging, as discussed in Shou and Gupta [\[17\]](#page-20-1).

In the traditional framework where $X_1, \ldots, X_n \stackrel{i.i.d}{\sim} X$ and $U_1, \ldots, U_n \stackrel{i.i.d}{\sim} U$, the foundational works of Vardi [\[20\]](#page-20-2) and Vardi et al.[\[21\]](#page-20-3) were pivotal in introducing and investigating deeply into multiplicative censoring, a specific aspect of the multiplicative deconvolution problem where the multiplicative error U is uniformly distributed over the interval [0, 1]. The task of estimating the cumulative distribution function of X was significantly advanced in Vardi et al.[\[21\]](#page-20-3) and Asgharian and Wolfson [\[2\]](#page-19-0). Meanwhile, series expansion methods, approaching the model as an inverse problem, were explored in Andersen and Hansen [\[1\]](#page-19-1). In the realm of density estimation within a multiplicative censoring model, Brunel $et al.[6]$ $et al.[6]$ introduced both kernel estimators and convolution power kernel estimators. The study of Comte and Dion [\[7\]](#page-19-3) focuses on a projection density estimator based on the Laguerre basis, assuming a uniform error distribution across the interval $[1 - \alpha, 1 + \alpha]$, where α lies within $(0, 1)$. In the work of Belomestny and Goldenshluger [\[3\]](#page-19-4), the authors study the scenario where the error U is beta-distributed. Additionally, Miguel *et al.*[\[8\]](#page-19-5) highlighted the transformative role of Mellin transform in a fully data-driven procedure for nonparametric density estimation. Lastly, Miguel *et al.*[\[9\]](#page-19-6) focuses on nonparametric estimation using a plug-in approach, combining Mellin transform estimation and regularization for a linear functional evaluated at an unknown density function.

In the context of dependent data where X_t and U_t are stationary proccesses, the considering model has received limited attention in research studies. However, in the scenario where X_1, \ldots, X_n being a stationary process and $U_1, \ldots, U_n \stackrel{i.i.d}{\sim} U$, the work by Miguel and Phandoidaen [\[10\]](#page-19-7) stands out, presenting an estimation for the unknown survival function within the framework of model [\(1\)](#page-1-0).

In this study, our primary focus revolves around the estimation of the invariant density function f_X within scenarios where both X_t and U_t are conceptualized as stationary processes. A pivotal aspect of our investigation lies in the adept handling of non-independent data, mirroring real-world complexities. A prevalent scenario arises when data is gathered from diverse groups, and within each group, there exist interdependencies among the data points. These interdependencies align with data patterns generated from m-dependent stationary processes, constituting a distinctive subset of stationary processes. The incorporation of this assumption, especially concerning U_t , introduces an innovative approach that has not been explored previously, enriching the precision and profundity of our modeling and density estimation. This study not only broadens the applications of model [\(1\)](#page-1-0) to various scientific domains but also proves invaluable in instances involving intricate interdependencies among data points. This paper focuses on the smooth error density class, which includes many important distributions such as uniform, beta, etc., under the assumption

$$
c(1+|p|)^{-\kappa} \le |g^{mt}(p)| \le C(1+|p|)^{-\kappa},\tag{2}
$$

where $C \ge c > 0, \kappa > 0$ and $g^{mt}(p)$ is the Mellin transform of the invariant error density function g , see for instance Miguel *et al.*[\[8\]](#page-19-5).

The structure of this paper is organized as follows. Section 2 outlines our proposed estimator by using the Mellin transform. In Section 3, we discuss convergence rates, provide a lower bound on the minimax risk, and explore the asymptotic normality of the estimator for f_X , all without including proofs. Applications and simulations are thoroughly examined in Section 4. Concluding the paper, Section 5 offers detailed proofs for the results introduced in Section 3.

2 Construction of the estimators

2.1 Mellin transform

In the multiplicative censoring model [\(1\)](#page-1-0), we consider a density estimator using the Mellin transform. The key to the analysis of the multiplicative deconvolution problem is the convolution theorem of the Mellin transform M, which roughly states $\mathcal{M}[f_Y] = \mathcal{M}[f_X] \mathcal{M}[g]$ for a density $f_Y = f_X * g$. For $\varrho \geq 1$, we denote by $L_{\varrho}^+ = L_{\varrho}(\mathbb{R}_+)$ the set of Lebesgue measurable functions f satisfying $||f||_{\varrho} = (\int_0^\infty |f(x)|^{\varrho} dx)^{1/\varrho} < \infty$. For $f \in L_1^+$, the Mellin transform of f in the point $c + ip \in \mathbb{C}$ is defined by

$$
\mathcal{M}_c[f](p) := \mathcal{M}[f](c+ip) = \int_0^\infty x^{c-1+ip} f(x) dx \tag{3}
$$

provided that the integral is absolutely convergent. If there exists a $c \in \mathbb{R}$ such that the mapping $x \mapsto x^{c-1}f(x)$ is integrable over \mathbb{R}_+ then the region $\Xi_f \subseteq \mathbb{C}$ of absolute convergence of the integral in [\(3\)](#page-2-0) is either a vertical strip $\{s + ip \in \mathbb{C} : s \in (a, b), p \in \mathbb{R}\}\)$ for $a < b$ with $c \in (a, b)$ or a vertical line $\{c + ip \in \mathbb{C} : p \in \mathbb{R}\}.$ We can see some techniques to determine Ξ_f in([\[8\]](#page-19-5)). Note that for any density $f \in L_1^+$ the vertical line $\{1 + ip \in \mathbb{C} : p \in \mathbb{R}\}\)$ belongs to Ξ_f , and hence the Mellin transform $\mathcal{M}_1[f]$ is well-defined. In this article, we will restrict to the case of model parameter $c = 1$ and denote $f^{mt} = \mathcal{M}_1[f]$. Nevertheless, we can extend for arbitrary model parameters $c \in \mathbb{R}$. For $c = 1$, the inversion formula of the Mellin transform is given by

$$
f(x) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} x^{-1-ip} f^{mt}(p) \, dp, \quad x \in \mathbb{R}_+.
$$
 (4)

2.2 The proposed estimators

In the present paper, we consider the problem on a dependent process. As known, there are various types of dependence conditions which is deeply studied (see Bradley [\[5\]](#page-19-8)). In this paper, we limit ourselves to the α -mixing dependence which was introduced by Rosenblatt [\[16\]](#page-20-4). Let \mathcal{F}_i^k be the σ -algebra of events generated by the random variables $\{X_j, 1 \leq i \leq j \leq k\}$. The process $\{X_i\}$ is called strongly mixing or α -mixing (see [\[5,](#page-19-8) [16\]](#page-20-4)) if

$$
\sup_{j \in \mathbb{N}} \sup_{\substack{A \in \mathcal{F}_1^j \\ B \in \mathcal{F}_{j+k}^j}} \left| \mathbb{P} \left[A B \right] - \mathbb{P} \left[A \right]. \mathbb{P} \left[B \right] \right| = \alpha_X \left(k \right) \to 0 \text{ as } k \to +\infty,
$$

where $\alpha_X(k)$ is the strong mixing coefficient of the process $\{X_j\}$. The quantity is a kind of measures of dependence. When (X_i) is mutually independent then $\sup_k \alpha_X (k) = 0$. The condition $\alpha_X(k) \to 0$ means that X_j and X_{j+k} are "asymptotically independent" as $k \to \infty$. With the dependence properties, we can establish an estimator of deconvolution which is consistent.

From [\(4\)](#page-3-0), to obtain an estimator of $f_X(x)$, we first try to construct a suitable estimator of $f_X^{mt}(p) = f_Y^{mt}(p)/g^{mt}(p)$. By replacing $f_Y^{mt}(p)$ with its empirical counterpart $f_Y^{mt}(p) =$ 1 $\frac{1}{n}\sum_{j=1}^n Y_j^{ip}$ j^{np} and using the kernel function construction as in Belomestny and Goldenshluger [\[4\]](#page-19-9), we obtain an estimator of $f_X(x)$ in the final form

$$
\widehat{f}_n(x) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} x^{-1-ip} \frac{K^{ft}(pb_n)}{g^{mt}(p)} \frac{1}{n} \sum_{j=1}^n Y_j^{ip} dp, \quad x \in \mathbb{R}_+, \tag{5}
$$

where b_n is the positive parameter which tends to 0 as n tends to infinity, K is a known kernel function and $K^{ft}(p) := \int_{-\infty}^{+\infty} e^{ipx} K(x) dx$ is the Fourier transform of K.

2.3 Assumptions

Assumption A

 (A_1) The kernel function K is supported on $[-1, 1]$, bounded and for a positive integer number m,

$$
\int_{-1}^{1} K(x) dx = 1, \quad \int_{-1}^{1} x^{k} K(x) dx = 0, \quad k = 1, ..., m.
$$

 (A_2) The Fourier transform K^{ft} of the kernel function K satisfies $\int |p|^{k\kappa} |K^{ft}(p)|^k dp < \infty$ for $k = 1, 2$ and κ is given in [\(2\)](#page-2-1).

Assumption B

 (B_1) For some $\kappa > 0, 0 < c \le C, D > 0$, the error density function g of the process U_t has the Mellin transform satisfying

$$
c(1+|p|)^{-\kappa} \le |g^{mt}(p)| \le C(1+|p|)^{-\kappa}, \left| \left(g^{mt}(p)\right)'\right| \le D(1+|p|)^{-\kappa} \text{ for all } p \in \mathbb{R}.
$$

 (B_2) The 2-dimensional probability density function $f_{Y_j,Y_k}(u,v)$ exists and is bounded for all $1 \leq j, k \leq n$.

Remark 2.1 The kernel function K can be constructed satisfying the Assumption A in different ways. For example, let $w(x) = e^{-x^2/2}/\sqrt{2\pi}$ and for a fixed natural number m let

$$
K(x) = \sum_{j=1}^{m+1} {m+1 \choose j} (-1)^{j+1} \frac{1}{j} w\left(\frac{x}{j}\right).
$$
 (6)

The Fourier transform of K is given by

$$
K^{ft}(p) = \sum_{j=1}^{m+1} {m+1 \choose j} (-1)^{j+1} e^{-j^2 p^2/2}.
$$
 (7)

It is well-known that the kernel K defines in (6) and its Fourier transform in (7) satisfying the Assumption A (see Kerkyacharian et al. $[14]$).

3 Preliminary results

In this Section, we introduce some results on the bias and variance, then the almost surely converrgence of our proposed estimator.

The bias of the estimators will be analyzed under a local smoothness assumption on f_X . Let $s > 0, A > 0, x > 0$ and $r > 1$. We say that $f \in \mathcal{F}_{x,r}(A,s)$ if f is a probability density, that is, $l = \lfloor s \rfloor := \max\{k \in \mathbb{N}_0 : k < s\}$ times continuously differentiable, and $\max_{k=1,\dots,l} |f^{(k)}(x_1)| \leq A,$

$$
|f^{(l)}(x_1) - f^{(l)}(x_2)| \le A |x_1 - x_2|^{s-l}, \quad \forall x_1, x_2 \in [r^{-1}x, rx].
$$

The following Theorem gives an upper bound on the bias of $f_n(x)$ when $f_X(x) \in$ $\mathcal{F}_{x,r}(A, s)$. We refer Lemma 2 and Lemma 3 of [\[4\]](#page-19-9) for the proof of this theorem.

Theorem 3.1 Let $f_X(x) \in \mathcal{F}_{x,r}(A,s)$ and $f_n(x)$ be given in [\(5\)](#page-3-1). Let K be the kernel function K satisfying Assumption A with $m \geq \lfloor s \rfloor + 1$, then for any $x > 0$ and n large enough

$$
\sup_{f_X \in \mathcal{F}_{x,r}(A,s)} \text{bias}\left[\hat{f}_n(x)\right] \le \mathcal{O}\left(b_n^s\right),
$$

where bias $\left[\hat{f}_n(x)\right] = \left|\mathbb{E}\hat{f}_n(x) - f_X(x)\right|$.

The general bounds for the variance of \hat{f}_n are given in the Theorem [3.2.](#page-5-0)

Theorem 3.2 Let Assumption A and Assumption B hold. For $\delta > 0$, assume that the processes X_t , U_t have strong mixing coefficients $\alpha_X(k)$, $\alpha_U(k) \leq \mathcal{O}(1/k^{2+\delta})$. If $f_X(x) \in$ $\mathcal{F}_{x,r}(A,s)$, then for any positive sequence (b_n) such that $b_n \to 0$ and $nb_n^{1+2\kappa} \to \infty$ as $n \to \infty$,

$$
\sigma_1^2(x) = \lim_{m \to \infty} \inf_{n \ge m} \left[nb_n^{1+2\kappa} \text{var}\left(\widehat{f}_n(x)\right) \right] \le \lim_{m \to \infty} \sup_{n \ge m} \left[nb_n^{1+2\kappa} \text{var}\left(\widehat{f}_n(x)\right) \right] = \sigma_2^2(x)
$$

at continuity points $x > 0$ of f_Y . Where

$$
\sigma_1^2(x) = \frac{f_Y(x)}{2\pi x} \lim_{m \to \infty} \inf_{n \ge m} \int \left| \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} \right|^2 dp \ge \frac{f_Y(x)}{2\pi x C_2^2} \int |p|^{2\kappa} |K^{ft}(p)|^2 dp, \tag{8}
$$

$$
\sigma_2^2(x) = \frac{f_Y(x)}{2\pi x} \lim_{m \to \infty} \sup_{n \ge m} \int \left| \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} \right|^2 dp \le \frac{f_Y(x)}{2\pi x C_1^2} \int |p|^{2\kappa} |K^{ft}(p)|^2 dp \tag{9}
$$

and

$$
0 < C_1 = \lim_{u \to \infty} \inf_{|p| \ge u} |g^{mt}(p)| |p|^\kappa \le \lim_{u \to \infty} \sup_{|p| \ge u} |g^{mt}(p)| |p|^\kappa = C_2.
$$

In Theorem [3.3,](#page-5-1) we show that the estimator \hat{f}_n converge almost surely to f_X when $n \to \infty$ under some assumptions.

Theorem 3.3 Let the Assumption A and assumption B_1 hold. For $a > 0, 0 < \rho < 1$, assume that the stationary processes X_t , U_t have strong mixing coefficients $\alpha_X(k)$, $\alpha_U(k) \le a\rho^k$. If $f_X(x) \in \mathcal{F}_{x,r}(\mathcal{A}, s)$, then for any positive sequence (b_n) such that $b_n \to 0$ and $nb_n^{1+2\kappa} \to +\infty$ as $n \to +\infty$, $\widehat{f}_n(x) \to f_X(x)$ almost surely.

4 Main results

4.1 Asymptotic normality

The result of asymptotic normality for \hat{f}_n is as follows.

Theorem 4.1 Let Assumption A and Assumption B hold. For $\delta > 0$, assume that the processes X_t , U_t have strong mixing coefficients $\alpha_X(k)$, $\alpha_U(k) \leq \mathcal{O}(1/k^{2+\delta})$. Choose the bandwidth b_n so that $nb_n^{1+2\kappa} \to \infty$ and a sequence of positive integers $\{s_n\}$ such that $s_n \to \infty$, $s_n = \mathcal{O}\left(\left(n b_n\right)^{1/2}\right)$ and $\left(n/b_n\right)^{1/2} \alpha_U(s_n) \to 0$ as $n \to \infty$. Then

$$
\lim_{m \to \infty} \inf_{n \ge m} \left\{ n^{1/2} b_n^{1/2 + \kappa} \left(\widehat{f}_n \left(x \right) - \mathbb{E} \widehat{f}_n \left(x \right) \right) \right\} \stackrel{D}{=} \mathcal{N} \left(0; \sigma_1^2 \left(x \right) \right), \tag{10}
$$

$$
\lim_{m \to \infty} \sup_{n \ge m} \left\{ n^{1/2} b_n^{1/2 + \kappa} \left(\widehat{f}_n \left(x \right) - \mathbb{E} \widehat{f}_n \left(x \right) \right) \right\} \stackrel{D}{=} \mathcal{N} \left(0; \sigma_2^2 \left(x \right) \right), \tag{11}
$$

at continuity points $x > 0$ of f_Y , where $\sigma_1^2(x)$ and $\sigma_2^2(x)$ given in [\(8\)](#page-5-2) and [\(9\)](#page-5-2).

Remark [4.1](#page-5-3) The selection of the bandwidth b_n in Theorem 4.1 determines the existence of the sequence of positive integers $\{s_n\}$. If the stationary processes X_t, U_t have strongly mixing coefficients $\alpha_X(k) = O(1/k^{2+\delta})$ or $\alpha_U(k) = O(1/k^{2+\delta})$, we can choose the bandwidth $b_n = Cn^{-\omega_1} (\log n)^{\omega_2}$ (where $C > 0, \omega_1 < \min\{1/(1+2\kappa), (1+\delta)/(3+\delta)\}, \omega_2 \in \mathbb{R}$ or $C > 0, \omega_1 = \min\{1/(1+2\kappa), (1+\delta)/ (3+\delta)\}, \omega_2 > 0$. In the case $\alpha_X(k), \alpha_U(k) \leq$ $\mathcal{O}(\rho^k)$ (0 < ρ < 1), bandwidth is chosen satisfying $b_n = Cn^{-\omega_1}(\log n)^{\omega_2}$ (C > 0, ω_1 < $1/(1+2\kappa), \omega_2 \in \mathbb{R}$ or $C > 0, \omega_1 = 1/(1+2\kappa), \omega_2 > 0$.

Corollary 4.2 Let Assumptions of Theorem [4.1](#page-5-3) hold. If $f_X(x) \in \mathcal{F}_{x,r}(A,s)$, choose the bandwidth b_n so that $nb_n^{1+2\kappa} \to \infty$ and $nb_n^{1+2\kappa+2s} \to 0$ as $n \to \infty$. Then

$$
\lim_{m \to \infty} \inf_{n \ge m} \left\{ n^{1/2} b_n^{1/2 + \kappa} \left(\hat{f}_n(x) - f_X(x) \right) \right\} \stackrel{D}{=} \mathcal{N} \left(0; \sigma_1^2(x) \right),
$$

$$
\lim_{m \to \infty} \sup_{n \ge m} \left\{ n^{1/2} b_n^{1/2 + \kappa} \left(\hat{f}_n(x) - f_X(x) \right) \right\} \stackrel{D}{=} \mathcal{N} \left(0; \sigma_2^2(x) \right)
$$

at continuity points $x > 0$ of f_Y , where $\sigma_1^2(x)$ and $\sigma_2^2(x)$ given in [\(8\)](#page-5-2) and [\(9\)](#page-5-2).

4.2 Bounds of the estimators

The following theorem gives us upper bounds of mean square error of $\widehat{f}_n(x)$.

Theorem 4.2 Let Assumption A and Assumption B hold, and assume that $f_X(x) \in \mathcal{F}_{x,r}(A,s)$. By choosing the bandwidth $b_n = \mathcal{O}\left(n^{-1/(1+2\kappa+2s)}\right)$, we have

$$
\sup_{f_X \in \mathcal{F}_{x,r}(A,s)} \mathbb{E} \left| \widehat{f}_n(x) - f_X(x) \right|^2 \leq \mathcal{O} \left(n^{-2s/(1+2\kappa+2s)} \right),
$$

for n sufficiently large.

Let \mathcal{T}_{α} be the set of all arbitrary estimator \widehat{f} of f_X based on data $Y_1, ..., Y_n$ which is generated from the $\alpha - mixing$ stationary process Y_t in model [\(1\)](#page-1-0). The next Theorem gives lower bounds for estimating densities of the set \mathcal{T}_{α} .

Theorem 4.3 Let $x > B > 0$ for some constant B and suppose that assumption B_1 hold and $\kappa > 1/2$. Then

$$
\inf_{\widehat{f}\in\mathcal{T}_{\alpha}}\sup_{f_X\in\mathcal{F}_{x,r}(A,s)}\mathbb{E}\left|\widehat{f}(x)-f_X(x)\right|^2\geq\mathcal{O}\left(n^{-2s/(1+2\kappa+2s)}\right),
$$

for n sufficiently large.

Theorem [4.2](#page-6-0) and Theorem [4.3](#page-6-1) show that the estimator \hat{f}_n attains the minimax rate of convergence $n^{-2s/(1+2\kappa+2s)}$ when the error density g is ordinary smooth with $\kappa > 1/2$.

5 Examples and Simulations

5.1 Cox-Ingersoll-Ross process

The Cox-Ingersoll-Ross process is the solution to the stochastic differential equation

$$
dX_t = (\theta_1 - \theta_2 X_t) dt + \theta_3 \sqrt{X_t} dW_t.
$$

Under the hypothesis $2\theta_1 > \theta_3^2$, there exists a positive τ such that $\alpha_X(k) \leq e^{-\tau k}/4$ (see Corollary 2.1 of Genon-Catalot *et al.*[\[11\]](#page-20-6)). The Cox-Ingersoll-Ross process is then an exponentially strong mixing stationary process and its invariant distribution follows a gamma density function

$$
f_X(x) = \frac{b^a}{\Gamma(a)} x^{a-1} \exp(-bx),
$$

where the shape parameter $a = 2\theta_1/\theta_3^2$ and the rate parameter $b = 2\theta_2/\theta_3^2$.

Assume that X_t is exponentially strongly mixing stationary Cox-Ingersoll-Ross processes with $\theta_1 = 1, \theta_2 = 0.5, \theta_3 = 1 (a = 2, b = 1)$ and noise random variables U_j follow either an uniform distribution on $[a_1, b_1]$ (denote as $U_{(a_1, b_1)}$) or Beta distribution (denote as $B_{(a_2, b_2)}$).

We now present simulations with different parameter values using the language R. First, we generate sample paths using the "sde" package (see Iacus [\[13\]](#page-20-7)), and then we compute the estimator's values. We choose the kernel function K as defined in [\(6\)](#page-4-0) with its Fourier transform given in [\(7\)](#page-4-1) for the case of $m = 2$. We also select the bandwidth $b_n = n^{-1/(1+2\kappa+2s)}$ according to Theorem [4.1.](#page-5-3) The simulations are based on 2000 noisy observations $Y_i = X_i U_j$ of the Cox-Ingersoll-Ross process X_t , with 50 replications. The results of the mean square error (MSE) at some points $x > 0$ for the case where X_t is a Cox-Ingersoll-Ross process and U_t is selected from $\{U_{(0,1)}, U_{(0.5,1.5)}, B_{(1,2)}, B_{(2,1)}, B_{(2,2)}\}$ are presented in Table [1.](#page-7-0)

Table 1: The empirical MSE of estimator at some points $x > 0$ in the case X_t is Cox-Ingersoll-Ross process and $U_t \in \{U_{(0,1)}, U_{(0.5,1.5)}, B_{(1,2)}, B_{(2,1)}, B_{(2,2)}\}.$

| \mathcal{X} | $U_{(0,1)}$ | $U_{(0.5,1.5)}$ | $B_{(1,2)}$ | $B_{(2,1)}$ | $B_{(2,2)}$ |
|---------------|--------------|-----------------|--------------|--------------|--------------|
| 0.5 | 0.02000825 | 0.01151807 | 0.01535441 | 0.01260686 | 0.01174962 |
| 1.0 | 0.01147437 | 0.009493284 | 0.01464701 | 0.009180121 | 0.01356236 |
| 1.5 | 0.001054993 | 0.0007115975 | 0.001157075 | 0.0008065801 | 0.0009686502 |
| 2.0 | 0.001534092 | 0.001253764 | 0.0006557094 | 0.001036627 | 0.0006276501 |
| 2.5 | 0.001789301 | 0.001902348 | 0.001516154 | 0.001864769 | 0.001463214 |
| 3.0 | 0.001974052 | 0.001711986 | 0.001340604 | 0.001830764 | 0.001237362 |
| 3.5 | 0.001127057 | 0.001042727 | 0.0008609882 | 0.001050847 | 0.0008042085 |
| 4.0 | 0.0005810139 | 0.0004889677 | 0.0005396105 | 0.0005298024 | 0.0004642698 |
| 4.5 | 0.0002820801 | 0.0001894068 | 0.0002175196 | 0.0002062239 | 0.0001943284 |

5.2 The *m*-Dependent stationary process

A very important kind of dependence considering distance as a measure of dependence, is the m-dependence case. A (discrete time) stationary process $\{X_j\}_{j\in\mathbb{Z}}$ is call m-dependent stationary process if two sets of random variables $\{..., X_{k-1}, X_k\}$ and $\{X_h, X_{h+1}, ...\}$ are independent whenever the time gap between them, represented as $h - k$, exceeds the value of m. Clearly, $\alpha_X(l) = 0$ when $l > m$ and the m-dependent stationary process is a special case of strongly mixing stationary process.

In this specific example, we assume that ${X_j}_{j\in\mathbb{Z}}$ is a 30-dependent stationary process and its invariant distribution is Weibull distribution with the shape parameter $a = 2$ and the scale parameter $b = 5$. We will use the language R to generate data for the 30-dependent process ${X_j}_{j \in \mathbb{Z}}$ as follows. First, we generate 30 independent data points, all following a Weibull distribution with the shape parameter $a = 2$ and the scale parameter $b = 5$. The 31st data point will either be one of the first 30 data points with a probability of 0.5, or it will be an independent Weibull data point with the same parameters with a probability of 0.5. Starting from the $(30 + l)$ th data point (where $l \geq 2$), it will select one of the data points from lth to $(30 + l - 1)$ th with a probability of 0.5 (if the data point $(30 + l - 1)$ th is the same as data point $(l-1)$ th, it will be removed from the list), and it will also have a 0.5 probability of being an independent Weibull data point with the same parameters. The noise random variables U_j are chosen in $\{U_{(0,1)}, U_{(0.5,1.5)}, B_{(1,2)}, B_{(2,1)}, B_{(2,2)}\}.$

With consistent bandwidth and kernel function, as outlined in Section [4.1,](#page-5-4) the simulations are based on 2000 noisy observations, denoted as $Y_j = X_j U_j$, originating from the $\{X_j\}_{j\in\mathbb{Z}}$ processes, and this is repeated 50 times. The results of empirical MSE at various points $x > 0$ are given in Table [2.](#page-8-0)

Table 2: The empirical MSE of estimator at various points $x > 0$ in the case X_t is 30dependent stationary process and $U_t \in \{U_{(0,1)}, U_{(0.5,1.5)}, B_{(1,2)}, B_{(2,1)}, B_{(2,2)}\}.$

| \boldsymbol{x} | $U_{(0,1)}$ | $U_{(0.5,1.5)}$ | $B_{(1,2)}$ | $B_{(2,1)}$ | $B_{(2,2)}$ |
|------------------|---------------|-----------------|---------------|---------------|---------------|
| 0.5 | 0.006237416 | 0.002502395 | 0.003739801 | 0.002115358 | 0.001116817 |
| 1.0 | 0.002888191 | 0.001106438 | 0.009513839 | 0.001234231 | 0.009458161 |
| 2.0 | 0.001482346 | 0.001076423 | 0.001115319 | 0.0009224827 | 0.001071403 |
| 3.0 | 0.0006854486 | 0.0003894643 | 0.001294102 | 0.0002997344 | 0.001251179 |
| 4.0 | 0.0005290895 | 0.0004286804 | 0.0001148088 | 0.0003080888 | 0.0001467156 |
| 5.0 | 0.0001766462 | 0.0001377783 | 0.00008996057 | 0.0001675634 | 0.00008717821 |
| 6.0 | 0.00009856583 | 0.00006615849 | 0.0001096502 | 0.0000772704 | 0.00008234754 |
| 7.0 | 0.00004252327 | 0.00003739727 | 0.00002887898 | 0.00002122861 | 0.00002844099 |
| 8.0 | 0.00002642288 | 0.00002186801 | 0.0000406609 | 0.00002247162 | 0.00004048739 |

6 Proofs

6.1 Proof of Theorem [3.2](#page-5-0)

Before proving Theorem Theorem [3.2,](#page-5-0) we prove the following technical lemma.

Lemma 6.1 Asumme that Assumption A and Assumption B_1 hold. We have

(a)
$$
\lim_{m \to \infty} \inf_{n \ge m} \frac{1}{b_n} \int \phi_n \left(\frac{\ln x - \ln y}{b_n} \right) f_Y(y) dy = 2\pi x f_Y(x) \lim_{m \to \infty} \inf_{n \ge m} \int \left| \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} \right|^2 dp
$$

\n
$$
\ge \frac{2\pi}{C_2^2} x f_Y(x) \int |p|^{2\kappa} |K^{ft}(p)|^2 dp,
$$

\n(b)
$$
\lim_{m \to \infty} \sup_{n \ge m} \frac{1}{b_n} \int \phi_n \left(\frac{\ln x - \ln y}{b_n} \right) f_Y(y) dy = 2\pi x f_Y(x) \lim_{m \to \infty} \sup_{n \ge m} \int \left| \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} \right|^2 dp
$$

\n
$$
\le \frac{2\pi}{C_1^2} x f_Y(x) \int |p|^{2\kappa} |K^{ft}(p)|^2 dp
$$

at points of continuity $x > 0$ of f_Y , where

$$
\phi_n(y) = \left| \int e^{-ipy} \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} dp \right|^2
$$

and

$$
0 < C_1 = \lim_{u \to \infty} \inf_{|p| \ge u} |g^{mt}(p)| |p|^\kappa \le \lim_{u \to \infty} \sup_{|p| \ge u} |g^{mt}(p)| |p|^\kappa = C_2.
$$

Proof of Lemma [6.1](#page-9-0).

We verify (a) and use the same argument for (b) . Let $h(\ln y) = y f_Y(y)$, we have

$$
\lim_{m \to \infty} \inf_{n \ge m} \frac{1}{b_n} \int \phi_n \left(\frac{\ln x - \ln y}{b_n} \right) f_Y(y) dy - \frac{2\pi}{C_2^2} x f_Y(x) \int |p|^{2\kappa} |K^{ft}(p)|^2 dp
$$
\n
$$
= \lim_{m \to \infty} \inf_{n \ge m} \left\{ \frac{1}{b_n} \int \phi_n \left(\frac{\ln x - \ln y}{b_n} \right) h(\ln y) d(\ln y) - \frac{h(\ln x)}{b_n} \int \phi_n \left(\frac{\ln x - \ln y}{b_n} \right) d(\ln y) + \frac{h(\ln x)}{b_n} \int \phi_n \left(\frac{\ln x - \ln y}{b_n} \right) d(\ln y) - \frac{2\pi}{C_2^2} h(\ln x) \int |p|^{2\kappa} |K^{ft}(p)|^2 dp \right\}
$$
\n
$$
= \lim_{m \to \infty} \inf_{n \ge m} \left\{ \int \left[h(\ln x - \ln y) - h(\ln x) \right] \frac{1}{b_n} \phi_n \left(\frac{\ln y}{b_n} \right) d(\ln y) + h(\ln x) \left[\int \phi_n (\ln y) d(\ln y) - \frac{2\pi}{C_2^2} \int |p|^{2\kappa} |K^{ft}(p)|^2 dp \right] \right\}.
$$
\n(12)

To reduce the limit of [\(12\)](#page-9-1), we consider two steps: **Step 1:** we prove $\int [h(\ln x - \ln y) - h(\ln x)] \frac{1}{b_n}$ ϕ_n $\ln y$ b_n $\overline{}$ $d(\ln y) \to 0$ as $n \to \infty$. We have $b_n^{\kappa} K^{ft}(p)$ $g^{mt}\left(p/b_{n}\right)$     ≤ b_n^{κ} $\left| K^{ft}\left(p\right) \right|$ min $\frac{\min\limits_{|p|\leq N}|\sum\limits_{g=1}^{|p'|}1_{\{|p|\leq Nb_n\}}+2}{\min\limits_{|p|\leq N}|\sum\limits_{g=1}^{|p'|}1_{\{|p|\leq Nb_n\}}+2}.$ $|p|^{\kappa} \left|K^{ft}\left(p\right)\right|$ $\frac{C_1}{C_1}$ $1_{\{|p|>Nb_n\}},$

where N is large enough (but fixed) such that: $|p|^{\kappa} |g^{mt}(p)| \geq \frac{C_1}{2}$ when $|p| > N$. We obtain

$$
\left| \int \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} dp \right| \leq \int \left| \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} \right| dp
$$

\n
$$
\leq \int \left(\frac{b_n^{\kappa} |K^{ft}(p)|}{\min_{|p| \leq N} |g^{mt}(p)|} 1_{\{|p| \leq Nb_n\}} + 2 \frac{|p|^{\kappa} |K^{ft}(p)|}{C_1} 1_{\{|p| > Nb_n\}} \right) dp
$$

\n
$$
\leq \mathcal{O}(b_n^{\kappa}) + \frac{2}{C_1} \int |p|^{\kappa} |K^{ft}(p)| dp \leq D_1, \forall n
$$
\n(13)

where $D_1 > 0$ is a constant.

$$
\int e^{-ip\ln y} \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} dp = -\frac{1}{i\ln y} \left(e^{-ip\ln y} \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} \Big|_{-\infty}^{+\infty} - \int e^{-ip\ln y} \left(\frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} \right)' dp \right)
$$

$$
= -\frac{1}{i\ln y} \int e^{-ip\ln y} \frac{b_n^{\kappa} K^{ft}(p) (g^{mt}(p/b_n))'}{[g^{mt}(p/b_n)]^2} dp.
$$

We deduce

$$
\left| \int e^{-ip \ln y} \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} dp \right| \leq \frac{1}{\left| \ln y \right|} \int \left| \frac{b_n^{\kappa} K^{ft}(p) \left(g^{mt}(p/b_n) \right)'}{\left[g^{mt}(p/b_n) \right]^2} \right| dp.
$$

By the similar argument as [\(13\)](#page-10-0), we conclude

$$
\left| \int e^{-ip \ln y} \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} dp \right| \le \frac{D_2}{|\ln y|} \quad \forall n \quad \Rightarrow \quad \sup_n \left| \int e^{-ip \ln y} \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} dp \right|^2 \le \frac{D_2^2}{|\ln y|^2},\tag{14}
$$

where $D_2 > 0$ is a constant.

From [\(13\)](#page-10-0), [\(14\)](#page-10-1) and using argument as in proof of Lemma 5.1 in Trong and Hung [\[18\]](#page-20-8), we get

$$
\int \left[h \left(\ln x - \ln y \right) - h \left(\ln x \right) \right] \frac{1}{b_n} \phi_n \left(\frac{\ln y}{b_n} \right) d \left(\ln y \right) \to 0 \text{ as } n \to \infty. \tag{15}
$$

Step 2: we prove $\lim_{m\to\infty} \inf_{n\geq m}$ n≥m $\sqrt{ }$ $h\left(\ln x\right)$ Γ $\phi_n(\ln y) d(\ln y) 2\pi$ C_2^2 Z $|p|^{2\kappa} |K^{ft}(p)|^2 dp \bigg] \bigg\} \geq 0.$ We have

$$
\int \phi_n(\ln y) d(\ln y) = \int_{-\infty}^{+\infty} \phi_n(y) dy = \int_{-\infty}^{+\infty} \left| \int e^{-ipy} \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} dp \right|^2 dy
$$

$$
= \int_{-\infty}^{+\infty} \left| \left[\frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} \right]^{ft}(y) \right|^2 dy
$$

$$
= 2\pi \int_{-\infty}^{+\infty} \left| \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} \right|^2 dp \text{ by Parseval's identity.}
$$

We obtain

$$
\lim_{m \to \infty} \inf_{n \ge m} \left\{ h(\ln x) \left[\int \phi_n(\ln y) d(\ln y) - \frac{2\pi}{C_2^2} \int |p|^{2\kappa} |K^{ft}(p)|^2 dp \right] \right\}
$$
\n
$$
= h(\ln x) \left[\lim_{m \to \infty} \inf_{n \ge m} \int \phi_n(\ln y) d(\ln y) - \frac{2\pi}{C_2^2} \int |p|^{2\kappa} |K^{ft}(p)|^2 dp \right]
$$
\n
$$
= h(\ln x) \left[\lim_{m \to \infty} \inf_{n \ge m} 2\pi \int \left| \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} \right|^2 dp - \frac{2\pi}{C_2^2} \int |p|^{2\kappa} |K^{ft}(p)|^2 dp \right]
$$
\n
$$
\ge h(\ln x) \left[2\pi \int \lim_{m \to \infty} \inf_{n \ge m} \frac{|p|^{2\kappa} |K^{ft}(p)|}{|g^{mt}(p/b_n)|^2 |p/b_n|^{2\kappa}} dp - \frac{2\pi}{C_2^2} \int |p|^{2\kappa} |K^{ft}(p)|^2 dp \right] = 0. \tag{16}
$$

From (12) , (15) and (16) , we conclude

$$
\frac{2\pi}{C_2^2}xf_Y(x)\int |p|^{2\kappa} |K^{ft}(p)|^2 dp \le \lim_{m\to\infty} \inf_{n\ge m} \frac{1}{b_n} \int \phi_n\left(\frac{\ln x - \ln y}{b_n}\right) f_Y(y) dy.
$$

Similarly, we have

$$
\lim_{m \to \infty} \inf_{n \ge m} \frac{1}{b_n} \int \phi_n \left(\frac{\ln x - \ln y}{b_n} \right) h(\ln y) d(\ln y)
$$
\n
$$
= \lim_{m \to \infty} \inf_{n \ge m} \left\{ \frac{1}{b_n} \int \phi_n \left(\frac{\ln x - \ln y}{b_n} \right) h(\ln y) d(\ln y) - \frac{h(\ln x)}{b_n} \int \phi_n \left(\frac{\ln x - \ln y}{b_n} \right) d(\ln y) + \frac{h(\ln x)}{b_n} \int \phi_n \left(\frac{\ln x - \ln y}{b_n} \right) d(\ln y) \right\}
$$
\n
$$
= \lim_{m \to \infty} \inf_{n \ge m} \left\{ \int \left[h(\ln x - \ln y) - h(\ln x) \right] \frac{1}{b_n} \phi_n \left(\frac{\ln y}{b_n} \right) d(\ln y) + h(\ln x) \int \phi_n (\ln y) d(\ln y) \right\}
$$
\n
$$
= 2\pi x f_Y(x) \lim_{m \to \infty} \inf_{n \ge m} \int \left| \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} \right|^2 dp.
$$

Proof of Theorem [3.2](#page-5-0). First, we define

$$
W_{b_n}(x) := \frac{1}{2\pi} \int e^{-ipx} \frac{K^{ft}(p)}{g^{mt}(p/b_n)} dp.
$$

The estimator \hat{f}_n can be represented as

$$
\widehat{f}_n(x) = \frac{1}{nx} \sum_{j=1}^n \frac{1}{b_n} W_{b_n} \left(\frac{\ln x - \ln Y_j}{b_n} \right).
$$

Denote

$$
\mu_n = \frac{1}{b_n} \mathbb{E}\left[W_{b_n}\left(\frac{\ln x - \ln Y_j}{b_n}\right)\right] \text{ and } U_{n,j} = \frac{1}{b_n} W_{b_n}\left(\frac{\ln x - \ln Y_j}{b_n}\right) - \mu_n.
$$

We have

$$
\mu_n = \frac{1}{b_n} \mathbb{E}\left[\frac{1}{2\pi} \int e^{-ip\left(\frac{\ln x - \ln Y_j}{b_n}\right)} \frac{K^{ft}(p)}{g^{mt}(p/b_n)} dp\right] = x \mathbb{E}\left[\hat{f}_n(x)\right] \to x f_X(x) \text{ as } n \to \infty,
$$

where the last step follows by Theorem [3.1.](#page-4-2) Then

$$
b_n^{1+2\kappa} \mathbb{V}ar [U_{n,j}] = b_n^{1+2\kappa} \mathbb{E} |U_{n,j}|^2 = b_n^{1+2\kappa} \mathbb{E} \left| \frac{1}{b_n} \frac{1}{2\pi} \int e^{-ip \left(\frac{\ln x - \ln Y_j}{b_n} \right)} \frac{K^{ft}(p)}{g^{mt}(p/b_n)} dp - \mu_n \right|^2
$$

$$
= \frac{1}{b_n} \mathbb{E} \left| \frac{1}{2\pi} \int e^{-ip \left(\frac{\ln x - \ln Y_j}{b_n} \right)} \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} dp \right|^2 + \mathcal{O}(1)
$$

$$
= \frac{1}{4\pi^2} \frac{1}{b_n} \int \left| \int e^{-ip \left(\frac{\ln x - \ln y}{b_n} \right)} \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} dp \right|^2 f_Y(y) dy + \mathcal{O}(1).
$$

By Lemma [6.1,](#page-9-0) we have

$$
x^{2}\sigma_{1}^{2}(x) = \lim_{m \to \infty} \inf_{n \ge m} b_{n}^{1+2\kappa} \mathbb{V}ar \left[U_{n,j} \right] \le \lim_{m \to \infty} \sup_{n \ge m} b_{n}^{1+2\kappa} \mathbb{V}ar \left[U_{n,j} \right] = x^{2}\sigma_{2}^{2}(x).
$$
 (17)

We have

$$
b_n^{1+2\kappa} \sum_{l=2}^n |\text{Cov }\{U_{n,1}, U_{n,l}\}| = b_n^{1+2\kappa} \sum_{l=2}^{c_n} |\text{Cov }\{U_{n,1}, U_{n,l}\}| + b_n^{1+2\kappa} \sum_{l=c_n+1}^n |\text{Cov }\{U_{n,1}, U_{n,l}\}|
$$

= $S_1 + S_2$,

where $c_n \to \infty$ such that $c_n b_n \to 0$ as $n \to \infty$.

Let $Z_j = \ln Y_j$, $f_{Z_j}(u)$ and $f_{Z_j, Z_k}(u, v)$ $(1 \leq j, k \leq n)$ be the probability density function of Z_j and the joint probability density function of (Z_j, Z_k) , respectively.

For $2 \leq l \leq c_n$, we have in view of [\(13\)](#page-10-0) and [\(14\)](#page-10-1)

$$
\begin{split}\n\text{Cov}\left\{U_{n,1}, U_{n,l}\right\} &= \frac{1}{b_n^2} \text{Cov}\left\{W_{b_n}\left(\frac{\ln x - \ln Y_1}{b_n}\right), W_{b_n}\left(\frac{\ln x - \ln Y_l}{b_n}\right)\right\} \\
&= \frac{1}{b_n^2} \iint W_{b_n}\left(\frac{\ln x - u}{b_n}\right) W_{b_n}\left(\frac{\ln x - v}{b_n}\right) \left[f_{Z_1, Z_l}\left(u, v\right) - f_{Z_1}\left(u\right) f_{Z_l}\left(v\right)\right] du dv \\
&\text{sup}\left\{f_{Z_1, Z_l}\left(u, v\right) - f_{Z_1}\left(u\right) f_{Z_l}\left(v\right)\right| \\
&\leq \frac{u, v, l}{b_n^2} \left[\int \left|W_{b_n}\left(\frac{\ln x - u}{b_n}\right)\right| du\right]^2 \\
&= \mathcal{O}\left(1\right) \left[\int \left|W_{b_n}\left(u\right)\right| du\right]^2 \leq \frac{\mathcal{O}\left(1\right)}{b^{2\kappa}}.\n\end{split}
$$

Then

$$
S_1 = b_n^{1+2\kappa} \sum_{l=2}^{c_n} |Cov \{U_{n,1}, U_{n,l}\}| \le b_n^{1+2\kappa} c_n \frac{\mathcal{O}(1)}{b^{2\kappa}} = \mathcal{O}(b_n c_n) \to 0 \text{ as } n \to \infty.
$$

For $c_n + 1 \leq l \leq n$, note that the process (Y_j) is also strongly mixing (see Bradley [\[5\]](#page-19-8)) with strong mixing coefficient $\alpha_Y(k) \leq \mathcal{O}\left(1/k^{2+\delta}\right)$. This condition is equivalent to for some $\nu > 2$ and $a > 1 - 2/\nu$,

$$
\sum_{k=1}^{\infty} k^a [\alpha_Y(k)]^{1-2/\nu} < \infty. \tag{18}
$$

Applying Davydov's Lemma (see Hall and Heyde [\[12\]](#page-20-9), Corollary A.2), we find

$$
|\mathrm{Cov}\left\{U_{n,1}, U_{n,l}\right\}| \leq \frac{8}{b_n^2} \big[\alpha_Y\left(l-1\right)\big]^{1-2/\nu} \bigg\{ \mathbb{E}\left[\left|W_{b_n}\left(\frac{\ln x - \ln Y_1}{b_n}\right)\right|^{\nu}\right]\bigg\}^{2/\nu}.
$$

Note that

$$
\frac{1}{b_n} \mathbb{E}\left[\left|W_{b_n}\left(\frac{\ln x - \ln Y_1}{b_n}\right)\right|^{\nu}\right] \leq \mathcal{O}\left(1\right) \int \left|W_{b_n}\left(u\right)\right|^{\nu} du \leq \frac{\mathcal{O}\left(1\right)}{b_n^{\nu \kappa}},
$$

where the last step follows by [\(13\)](#page-10-0) and [\(14\)](#page-10-1). Hence,

 $j = c_n$

$$
|\text{Cov }\{U_{n,1}, U_{n,l}\}| \le \frac{\mathcal{O}\left(1\right)}{b_n^{2+2\kappa - 2/\nu}} [\alpha_Y (l-1)]^{1-2/\nu}
$$

and

$$
S_2 = b_n^{1+2\kappa} \sum_{l=c_n+1}^n |\text{Cov }\{U_{n,1}, U_{n,l}\}| \le \frac{\mathcal{O}(1)}{b_n^{1-2/\nu}} \sum_{l=c_n+1}^n [\alpha_Y (l-1)]^{1-2/\nu}
$$

$$
\le \frac{\mathcal{O}(1)}{c_n^a b_n^{1-2/\nu}} \sum_{j=c_n}^\infty j^a [\alpha_Y (j)]^{1-2/\nu}.
$$

Choose $c_n =$ $\begin{pmatrix} 1 \end{pmatrix}$ $b_n^{1-2/\nu}$ \setminus ^{1/a} then for $a > 1 - 2/\nu$ we obtain $c_n b_n \to 0$ as required. Then $S_2 \leq \mathcal{O}(1) \sum_{n=1}^{\infty}$ $j^a[\alpha_Y(j)]^{1-2/\nu} \to 0$ as $n \to \infty$ follows by [\(18\)](#page-12-0).

Therefore,

$$
\lim_{n \to \infty} b_n^{1+2\kappa} \sum_{l=2}^n |\text{Cov}\{U_{n,1}, U_{n,l}\}| = 0.
$$
\n(19)

We have

$$
n\mathbb{V}ar\left[\hat{f}_n(x)\right] = n\mathbb{V}ar\left\{\frac{1}{nx}\sum_{j=1}^n \frac{1}{b_n}W_{b_n}\left(\frac{\ln x - \ln Y_j}{b_n}\right)\right\}
$$

\n
$$
= \frac{1}{x^2}\mathbb{V}ar\left[\frac{1}{b_n}W_{b_n}\left(\frac{\ln x - \ln Y_j}{b_n}\right)\right]
$$

\n
$$
+ \frac{2}{nx^2}\sum_{j=2}^n \left[n - (j-1)\right]\mathbb{C}\mathrm{ov}\left\{\frac{1}{b_n}W_{b_n}\left(\frac{\ln x - \ln Y_1}{b_n}\right), \frac{1}{b_n}W_{b_n}\left(\frac{\ln x - \ln Y_j}{b_n}\right)\right\}
$$

\n
$$
= \frac{1}{x^2}\mathbb{V}ar\left[\frac{1}{b_n}W_{b_n}\left(\frac{\ln x - \ln Y_j}{b_n}\right)\right]
$$

\n
$$
+ \frac{2}{x^2}\sum_{j=2}^n \left(1 - \frac{j-1}{n}\right)\mathrm{cov}\left\{\frac{1}{b_n}W_{b_n}\left(\frac{\ln x - \ln Y_1}{b_n}\right), \frac{1}{b_n}W_{b_n}\left(\frac{\ln x - \ln Y_j}{b_n}\right)\right\}
$$

\n
$$
= \frac{1}{x^2}\mathbb{V}ar\left[U_{n,j}\right] + \frac{2}{x^2}\sum_{j=2}^n \left(1 - \frac{j-1}{n}\right)\mathrm{Cov}\left\{U_{n,1}, U_{n,j}\right\}.
$$

By (17) and (19) , we obtain

$$
\sigma_1^2(x) = \lim_{m \to \infty} \inf_{n \ge m} nb_n^{1+2\kappa} \mathbb{V}ar \left[\widehat{f}_n(x) \right] \le \lim_{m \to \infty} \sup_{n \ge m} nb_n^{1+2\kappa} \mathbb{V}ar \left[\widehat{f}_n(x) \right] = \sigma_2^2(x).
$$

This completes the proof of Theorem [3.2.](#page-5-0)

6.2 Proof of Theorem [3.3](#page-5-1)

To prove Theorem [3.3,](#page-5-1) we first prove the following lemma.

Lemma 6.2 Let (Y_i) be a stationary strongly mixing process on the probability space $(\Omega, \mathcal{A}, \mathbb{P})$ with the mixing coefficient satisfying $\alpha(k) \leq a\rho^k (a > 0, 0 < \rho < 1)$. Let $Z_j = \psi_1(Y_j) +$ i. $\psi_2(Y_j)$ where ψ_1 and ψ_2 are real-valued Borel measurable function. Assume that $\mathbb{E}(Z_j) = 0$ and $|Z_j| \leq d$. Let an integer $n \geq 1$ be given. Then for all $\varepsilon > 0$,

$$
\mathbb{P}\left\{\frac{1}{n}\left|\sum_{j=1}^{n}Z_j\right|\geq \varepsilon\right\}\leq 2\left(1+4ae^{-2}L\right)\exp\left(-\frac{\varepsilon^2n}{4\left(\mathbb{E}|Z_1|^2+\sqrt{2}\varepsilon d/6\right)}\right),
$$

where $L \geq 1$ is a constant.

Proof of Lemma [6.2](#page-14-0). We have

$$
\left\{\frac{1}{n}\left|\sum_{j=1}^{n}Z_{j}\right|\geq \varepsilon\right\} = \left\{\frac{1}{n}\sqrt{\left|\sum_{j=1}^{n}\psi_{1}\left(Y_{j}\right)\right|^{2} + \left|\sum_{j=1}^{n}\psi_{2}\left(Y_{j}\right)\right|^{2}}\geq \varepsilon\right\}
$$

$$
\subset \left\{\frac{1}{n}\left|\sum_{j=1}^{n}\psi_{1}\left(Y_{j}\right)\right|\geq \frac{\sqrt{2}}{2}\varepsilon\right\} \cup \left\{\frac{1}{n}\left|\sum_{j=1}^{n}\psi_{2}\left(Y_{j}\right)\right|\geq \frac{\sqrt{2}}{2}\varepsilon\right\}.
$$

We deduce

$$
\mathbb{P}\left\{\frac{1}{n}\left|\sum_{j=1}^{n}Z_j\right|\geq \varepsilon\right\}\leq \mathbb{P}\left\{\frac{1}{n}\left|\sum_{j=1}^{n}\psi_1\left(Y_j\right)\right|\geq \frac{\sqrt{2}}{2}\varepsilon\right\}+\mathbb{P}\left\{\frac{1}{n}\left|\sum_{j=1}^{n}\psi_2\left(Y_j\right)\right|\geq \frac{\sqrt{2}}{2}\varepsilon\right\}.
$$

Note that

 $\mathbb{E}\left(Z_j\right) = \mathbb{E}\left[\psi_1\left(Y_j\right) + i.\psi_2\left(Y_j\right)\right] = 0.$ This implies $\mathbb{E}\left[\psi_1\left(Y_j\right)\right] = \mathbb{E}\left[\psi_2\left(Y_j\right)\right] = 0.$ $|\psi_k(Y_j)| \leq |Z_j| \leq d, \ \ k = 1, 2.$

From Theorem 2.1 of N'drin and Hili [\[15\]](#page-20-10), we have

$$
\mathbb{P}\left\{\frac{1}{n}\left|\sum_{j=1}^{n}\psi_{k}\left(Y_{j}\right)\right|\geq\frac{\sqrt{2}}{2}\varepsilon\right\}\leq\left(1+4ae^{-2}L\right)\exp\left(-\frac{\varepsilon^{2}n}{4\left(\mathbb{E}|\psi_{k}\left(Y_{1}\right)\right|^{2}+\sqrt{2}\varepsilon d/6)}\right).
$$

Therefore,

$$
\mathbb{P}\left\{\frac{1}{n}\left|\sum_{j=1}^n Z_j\right| \geq \varepsilon\right\} \leq 2\left(1+4ae^{-2}L\right)\exp\left(-\frac{\varepsilon^2n}{4\left(\mathbb{E}|Z_1|^2+\sqrt{2}\varepsilon d/6\right)}\right).
$$

 \Box

Proof of Theorem [3.3](#page-5-1). Notice that

$$
\begin{array}{rcl} \left| \widehat{f}_n \left(x \right) - f_X \left(x \right) \right| & = & \left| \widehat{f}_n \left(x \right) - \mathbb{E} \widehat{f}_n \left(x \right) + \mathbb{E} \widehat{f}_n \left(x \right) - f_X \left(x \right) \right| \\ & \leq & \left| \widehat{f}_n \left(x \right) - \mathbb{E} \widehat{f}_n \left(x \right) \right| + \left| \mathbb{E} \widehat{f}_n \left(x \right) - f_X \left(x \right) \right| . \end{array}
$$

Then we get

$$
\begin{aligned}\n\left|\widehat{f}_n(x) - \mathbb{E}\widehat{f}_n(x)\right| &= \left|\frac{1}{2\pi} \int x^{-1-ip} \frac{K^{ft}(pb_n)}{g^{mt}(p)} \frac{1}{n} \sum_{j=1}^n \left(Y_j^{ip} - \mathbb{E}Y_j^{ip}\right) dp\right| \\
&= \frac{1}{nb_n^{1+\kappa}} \left|\sum_{j=1}^n \frac{1}{2\pi} \int x^{-1-ip} \frac{b_n^{1+\kappa} K^{ft}(pb_n)}{g^{mt}(p)} \left(Y_j^{ip} - \mathbb{E}Y_j^{ip}\right) dp\right| \\
&= \frac{1}{nb_n^{1+\kappa}} \left|\sum_{j=1}^n Z_j\right|, \\
&= \frac{1}{nb_n^{1+\kappa}} K^{ft}(d_n)\n\end{aligned}
$$

where $Z_j =$ 1 2π Z $x^{-1-ip}\frac{b_n^{1+\kappa}K^{ft}\left(pb_n\right)}{mt}$ $g^{mt}\left(p\right)$ $(Y_j^{ip} - \mathbb{E}Y_j^{ip}$ j^{ip} $\big)dp.$

We verify that Z_j satisfy the assumptions of Lemma [6.2.](#page-14-0) We first have $(Z_j)_{j=\overline{1,n}}$ are identical distributions, $\mathbb{E}[Z_j] = 0, \forall j$. For $\forall x > 0$, we have

$$
|Z_j| = \left| \frac{1}{2\pi} \int x^{-1-ip} \frac{b_n^{1+\kappa} K^{ft}(pb_n)}{g^{mt}(p)} (Y_j^{ip} - \mathbb{E}Y_j^{ip}) dp \right|
$$

\n
$$
\leq \frac{1}{2\pi x} \int \left| x^{-ip} \right| \left| \frac{b_n^{1+\kappa} K^{ft}(pb_n)}{g^{mt}(p)} \right| |Y_j^{ip} - \mathbb{E}Y_j^{ip} | dp
$$

\n
$$
\leq \frac{1}{\pi x} \int \left| \frac{b_n^{\kappa} K^{ft}(p)}{g^{mt}(p/b_n)} \right| dp
$$

\n
$$
\leq \frac{1}{c\pi x} \int |p|^{\kappa} K^{ft}(p) dp = B_1,
$$

where $B_1 > 0$ is a constant. So Z_j satisfy assumptions of Lemma [6.2.](#page-14-0) For all $\varepsilon > 0$, we get

$$
\mathbb{P}\left\{\frac{1}{nb_n^{1+\kappa}} \left|\sum_{j=1}^n Z_j\right| \ge \varepsilon\right\} = \mathbb{P}\left\{\frac{1}{n} \left|\sum_{j=1}^n Z_j\right| \ge \varepsilon b_n^{1+\kappa}\right\}
$$

$$
\le 2\left(1 + 8ae^{-2}L\right) \exp\left(-\frac{\varepsilon^2 nb_n^{2+2\kappa}}{4\left[\mathbb{E}|Z_1|^2 + \sqrt{2}\varepsilon b_n^{1+\kappa}B_1/6\right]}\right)
$$

We have

$$
\mathbb{E}|Z_{1}|^{2} = \mathbb{E}\left|\frac{1}{2\pi}\int x^{-1-ip}\frac{b_{n}^{1+\kappa}K^{ft}(pb_{n})}{g^{mt}(p)}\left(Y_{1}^{ip} - \mathbb{E}Y_{1}^{ip}\right)dp\right|^{2} \n\leq \frac{b_{n}^{2+2\kappa}}{4\pi^{2}x^{2}}\mathbb{E}\left|\int e^{ip(\ln Y_{1}-\ln x)}\frac{K^{ft}(pb_{n})}{g^{mt}(p)}dp\right|^{2} \n= \frac{b_{n}}{4\pi^{2}x^{2}}\frac{1}{b_{n}}\int\left|\int e^{-ip\left(\frac{\ln x-\ln y}{b_{n}}\right)}\frac{b_{n}^{\kappa}K^{ft}(p)}{g^{mt}(p/b_{n})}dp\right|^{2}f_{Y}(y) dy \n\leq \frac{b_{n}}{2\pi xC_{1}^{2}}f_{Y}(x)\int |p|^{2\kappa} |K^{ft}(p)|^{2}dp = b_{n}B_{2},
$$

where $B_2 > 0$ is a constant and the last step follows by Lemma [6.1.](#page-9-0) Moreover, the process (Y_j) is also strongly mixing (see [\[5\]](#page-19-8)) with strong mixing coefficient

$$
\alpha_Y(k) \le \alpha_X(k) + \alpha_U(k) \le 2a\rho^k.
$$

This implies

$$
\mathbb{P}\left\{\frac{1}{nb_n^{1+\kappa}} \left|\sum_{j=1}^n Z_j\right| \ge \varepsilon\right\} \le 2\left(1 + 8ae^{-2}L\right) \exp\left(-\frac{\varepsilon^2 nb_n^{1+2\kappa}}{4\left[B_2 + \sqrt{2}\varepsilon b_n^{\kappa}B_1/6\right]}\right) \to 0
$$

as $nb_n^{1+2\kappa} \to \infty$.

Therefore, by Borel-Cantelli's lemma, we deduce that $\hat{f}_n(x) - \mathbb{E} \hat{f}_n(x) \to 0$ a.s. Thus, $\widehat{f}_n(x) \to f_X(x)$ a.s. by Theorem [3.1.](#page-4-2)

6.3 Proof of Theorem [4.1](#page-5-3)

We will prove [\(10\)](#page-5-5) and use the same argument for [\(11\)](#page-5-5). Let $V_{n,j} = b^{1/2+\kappa} U_{n,j}$, we have

$$
n^{1/2}b^{1/2+\kappa}\left(\hat{f}_n(x) - \mathbb{E}\hat{f}_n(x)\right) = n^{1/2}b^{1/2+\kappa}\frac{1}{nx}\sum_{j=1}^n\left(\frac{1}{b_n}W_{b_n}\left(\frac{\ln x - \ln Y_j}{b_n}\right) - \mu_n\right)
$$

$$
= \frac{1}{\sqrt{n}}\sum_{j=1}^n b^{1/2+\kappa}U_{n,j} = \frac{1}{\sqrt{n}}\sum_{j=1}^n V_{n,j} = \frac{1}{\sqrt{n}}S_n.
$$

Note that the process (Y_i) is also strongly mixing (see [\[5\]](#page-19-8)) with strong mixing coefficient $\alpha_Y(k) \leq \mathcal{O}(1/k^{2+\delta})$. And due to $(n/b_n)^{1/2} \alpha_U(s_n) \to 0$ as $n \to \infty$, we also have $(n/b_n)^{1/2} \alpha_Y(s_n) \to 0$ as $n \to \infty$. The conditions $s_n = \mathcal{O}\left((nb_n)^{1/2}\right)$ and $(n/b_n)^{1/2} \alpha_Y(s_n) \to 0$ 0 as $n \to \infty$ imply that there exists a sequence of integers $\{q_n\}$, $q_n \to \infty$, such that

$$
q_n s_n = \mathcal{O}\left((nb_n)^{1/2}\right)
$$
 and $q_n (n/b_n)^{1/2} \alpha_Y (s_n) \to 0$ as $n \to \infty$.

Let $r_n =$ $\left(nb_n \right)^{1/2}$ q_n \overline{a} , we have some following properties as $n \to \infty$

$$
\frac{s_n}{r_n} \to 0 \quad (a) \,, \qquad \frac{r_n}{n} \to 0 \quad (b) \,, \qquad \frac{r_n}{(nb_n)^{1/2}} \to 0 \quad (c) \,, \qquad \frac{n}{r_n} \alpha_Y(s_n) \to 0 \quad (d) \tag{20}
$$

Now, we partition the set $\{1, 2, ..., n\}$ into $2k + 1$ subsets: $\{j(r + s) + 1, ..., j(r + s) + r\}$, ${j(r + s) + r + 1,..., (j + 1)(r + s)}$ for $0 \le j \le k - 1$ and ${k(r + s) + 1,..., n}$, where $r = r_n$, $s = s_n$ and

$$
k = k_n = \left\lfloor \frac{n}{r_n + s_n} \right\rfloor \tag{21}
$$

For $0 \leq j \leq k-1$, define the random variables

$$
\eta_j = \sum_{i=j(r+s)+1}^{j(r+s)+r} V_{n,i} \quad (a), \quad \xi_j = \sum_{i=j(r+s)+r+1}^{(j+1)(r+s)} V_{n,i} \quad (b), \quad \zeta_k = \sum_{i=k(r+s)+1}^n V_{n,i} \quad (c) \tag{22}
$$

Write $S_n =$ $\sum_{i=1}^{k-1}$ $j=0$ $\eta_j +$ $\sum_{i=1}^{k-1}$ $j=0$ $\xi_j + \zeta_k = S'_n + S''_n + S''''_n$, we will prove that, as $n \to \infty$,

$$
\frac{1}{n}\mathbb{E}(S''_n)^2 \to 0, \quad \frac{1}{n}\mathbb{E}(S'''_n)^2 \to 0 \quad (a) \qquad \left| \mathbb{E}\left(e^{ipS'_n}\right) - \prod_{j=0}^{k-1}\mathbb{E}\left(e^{ip\eta_j}\right) \right| \to 0 \quad (b)
$$
\n
$$
\inf_n \left[\frac{1}{n} \sum_{j=0}^{k-1} \mathbb{E}\left(\eta_j^2\right) \right] \to \sigma_1^2(x) \qquad (c) \qquad \frac{1}{n} \sum_{j=0}^{k-1} \mathbb{E}\left[\eta_j^2 I_{\left\{\left|\eta_j\right| > \varepsilon \sigma_1(x)\sqrt{n}\right\}} \right] \to 0 \quad (d)
$$
\n
$$
(23)
$$

Note that $(23a)$ implies that S''_n , S''_n are asymptotically negligible, $(23b)$ show that the summands $\{\eta_j\}$ in S'_n are asymptotically independent, [\(23](#page-16-0)*c*) and (23*d*) are the standard

Lindeberg-Feller conditions for asymptotic normality of S_n' under independence. Since $\mathbb{E}(V_{n,i})=0$, we have $\mathbb{E}(\eta_j)=\mathbb{E}(\xi_j)=\mathbb{E}(\zeta_k)=0$. Consider first

$$
\mathbb{E}(S'')^2 = \mathbb{V}ar\left[\sum_{j=0}^{k-1}\xi_j\right] = \sum_{j=0}^{k-1} \mathbb{V}ar\left[\xi_j\right] + \sum_{i=0}^{k-1}\sum_{\substack{j=0 \ j \neq i}}^{k-1} \text{Cov}\left\{\xi_i, \xi_j\right\} = F_1 + F_2.
$$

With $m_i = j (r + s) + r$, by [\(22](#page-16-1)b) we have

$$
\begin{split} \mathbb{V}ar\left[\xi_{j}\right] &= \sum_{i_{1}=1}^{s} \sum_{i_{2}=1}^{s} \text{Cov}\left\{V_{n,m_{j}+i_{1}}, V_{n,m_{j}+i_{2}}\right\} = s \mathbb{V}ar\left[V_{n,1}\right] + 2s \sum_{i=2}^{s} \left(1 - \frac{i}{s}\right) \text{Cov}\left\{V_{n,1}, V_{n,i}\right\} \\ &\leq s \sigma_{2}^{2}\left(x\right) \left\{1 + \mathcal{O}\left(1\right)\right\}, \end{split}
$$

the last step follows by Theorem [3.2](#page-5-0) and we obtain $F_1 \leq \mathcal{O}(1)$ ks. Next, consider the term $F_2 =$ $\sum_{i=1}^{k-1}$ $i=0$ $\sum_{i=1}^{k-1}$ $j=0$ $j\neq i$ $Cov \{\xi_i, \xi_j\} =$ $\sum_{i=1}^{k-1}$ $i=0$ $\sum_{i=1}^{k-1}$ $j=0$ $j\neq i$ $\sum_{i=1}^{s}$ $l_1=1$ $\sum_{i=1}^{s}$ $l_2=1$ Cov $\{V_{n,m_i+l_1}, V_{n,m_j+l_2}\}$ Since $i \neq j, |m_j + l_2 - m_i - l_1| \geq r$, so that

$$
F_2 \le 2 \sum_{l_1=1}^{n-r} \sum_{l_2=r+l_1}^{n} |Cov \{V_{n,l_1}, V_{n,l_2}\}| \le 2n \sum_{j=r+1}^{n} |Cov \{V_{n,1}, V_{n,j}\}| = \mathcal{O}(n).
$$

Therefore, we obtain in view of $(20a)$ and (21)

$$
\frac{1}{n}\mathbb{E}(S_{n}^{"})^{2} = \frac{F_{1} + F_{2}}{n} \le \frac{\mathcal{O}(1) \, ks + \mathcal{O}(n)}{n} \to 0 \text{ as } n \to \infty.
$$

Using a similar argument for the latter inequality, we have in view of $(20a)$ and $(20b)$ that

$$
\frac{1}{n} \mathbb{E}(S'''_{n})^{2} = \frac{1}{n} \sum_{j=1}^{n-k(r+s)} \mathbb{V}ar \left[V_{n,k(r+s)+j} \right] + \frac{1}{n} \sum_{i=1}^{n-k(r+s)} \sum_{\substack{j=1 \ j \neq i}}^{n-k(r+s)} \text{Cov} \left\{ V_{n,k(r+s)+i}, V_{n,k(r+s)+j} \right\}
$$
\n
$$
\leq \frac{C}{n} \left[n - k(r+s) \right] + \frac{\mathcal{O}(n)}{n} \leq \frac{C(r+s)}{n} + \frac{\mathcal{O}(n)}{n} \to 0 \text{ as } n \to \infty.
$$

This proves [\(23](#page-16-0)a). To prove (23b), we note that η_j is a function of the random variables $\{Y_{j(r+s)+1},...,Y_{j(r+s)+r}\}\$ or η_j is $\mathcal{G}_{u_j}^{v_j}$ -measurable with $u_j = j(r+s)+1, v_j = j(r+s)+r$ $(\mathcal{G}_{u_j}^{v_j})$ is the σ -algebra of events generated by the random variables $\{X_h, U_h, u_j \leq h \leq v_j\}$. Also $u_{j+1} - v_j = s + 1$. Hence by Lemma 1.1 of Volkonskii and Rozannov [\[22\]](#page-20-11), we find

$$
\left| \mathbb{E}\left(\prod_{j=0}^{k-1} e^{ip\eta_j}\right) - \prod_{j=0}^{k-1} \mathbb{E}\left(e^{ip\eta_j}\right) \right| \le 16k\alpha_Y(s+1) \sim \frac{16n}{r+s}\alpha_Y(s+1) \to 0 \text{ as } n \to \infty,
$$

the last step follows by $(20a)$ and $(20d)$.

Next to [\(23](#page-16-0)c), by using stationary property of the processes X_t, U_t and applying Theorem [3.2,](#page-5-0) we obtain

$$
\inf_{n} \left[\frac{1}{n} \sum_{j=0}^{k-1} \mathbb{E} \left(\eta_j^2 \right) \right] = \inf_{n} \left[\frac{k}{n} \mathbb{E} \left(\eta_0^2 \right) \right] = \inf_{n} \left[\frac{k \cdot r}{n} \left(\text{var} \left(V_{n,1} \right) + \mathcal{O} \left(1 \right) \right) \right] \to \sigma_1^2 \left(x \right) \text{ as } n \to \infty.
$$

Finally, to prove $(23d)$, we use

$$
|U_{n,i}| \leq \frac{b_n^{1/2+\kappa}}{b_n} \left\{ \sup_u |W_{b_n}(u)| + |\mu_n| \right\} \leq \frac{b_n^{\kappa}}{b_n^{1/2}} \left\{ \frac{\mathcal{O}(1)}{b_n^{\kappa}} + |\mu_n| \right\} \text{(follows by (13))}.
$$

Since $\mu_n \to f_X(x)$, we get $|V_{n,i}| \leq \frac{\mathcal{O}(1)}{b_n^{1/2}}$ uniformly in *i*. n Hence by $(20c)$, we have max $0 \leq j \leq k-1$ $\frac{|\eta_j|}{\sqrt{n}} \leq$ $\mathcal{O}(1)$ r $\frac{\mathcal{O}(1)}{(nb_n)^{1/2}} \to 0$ as $n \to \infty$. Thus the set $\{|\eta_j| > \varepsilon \sigma_1(x) \sqrt{n}\}$ becomes empty for large *n*. Therefore [\(23](#page-16-0)*d*) now follows by

$$
\frac{1}{n}\sum_{j=0}^{k-1}\mathbb{E}\left[\eta_j^2I_{\left\{\left|\eta_j\right|>\varepsilon\sigma_1(x)\sqrt{n}\right\}}\right]\leq \frac{\mathcal{O}\left(1\right)}{n}k\left(\frac{r}{b_n^{1/2}}\right)^2\mathbb{P}\left[\left|\eta_0\right|>\varepsilon\sigma_1\left(x\right)\sqrt{n}\right].
$$

This completes the proof of the theorem.

6.4 Proof of Corollary [4.2](#page-6-2)

Note that

$$
n^{1/2}b^{1/2+\kappa}\left(\widehat{f}_n(x)-f_X(x)\right) = n^{1/2}b^{1/2+\kappa}\left[\widehat{f}_n(x)-\mathbb{E}\left(\widehat{f}_n(x)\right)\right] + n^{1/2}b^{1/2+\kappa}\left[\mathbb{E}\left(\widehat{f}_n(x)\right)-f_X(x)\right].
$$

By Theorem [3.1,](#page-4-2) we have

$$
\left|n^{1/2}b^{1/2+\kappa}\left[\mathbb{E}\left(\widehat{f}_n\left(x\right)\right)-f_X\left(x\right)\right]\right|=n^{1/2}b_n^{1/2+\kappa}\mathcal{O}\left(b_n^s\right)=\mathcal{O}\left(n^{1/2}b_n^{1/2+\kappa+s}\right)\to 0 \text{ as } n\to\infty.
$$

By Theorem [3.3,](#page-5-1) this ends the proof. \Box

6.5 Proof of Theorem [4.2](#page-6-0) and [4.3](#page-6-1)

Proof of Theorem [4.2](#page-6-0). From Theorem [3.1](#page-4-2) and [3.2,](#page-5-0) we have

bias
$$
\left(\widehat{f}_n(x)\right) \leq \mathcal{O}\left(b_n^s\right)
$$
,

and

$$
\mathbb{V}ar\left(\widehat{f}_n\left(x\right)\right) \le \frac{\mathcal{O}\left(1\right)}{nb_n^{1+2\kappa}}.
$$

We choose $b_n = \mathcal{O}\left(n^{-1/(1+2\kappa+2s)}\right)$, this yields

$$
\sup_{f_X \in \mathcal{F}_{x,r}(A,s)} \mathbb{E} \left| \widehat{f}_n(x) - f_X(x) \right|^2 \leq \mathcal{O} \left(n^{-2s/(1+2\kappa+2s)} \right).
$$

Proof of Theorem [4.3](#page-6-1).

Let \mathcal{T}_{iid} be the set of all arbitrary estimator \hat{f} of f_X based on independent identical distribution (iid) data $Y_1, ..., Y_n$ from model [\(1\)](#page-1-0). In this case, $X_1, ..., X_n$ are iid and $U_1, ..., U_n$ are iid. From Theorem 2 of Belomestny and Goldenshluger [\[4\]](#page-19-9), we have

$$
\inf_{\widehat{f}\in\mathcal{T}_{\alpha}}\sup_{f_X\in\mathcal{F}_{x,r}(A,s)}\mathbb{E}\left|\widehat{f}(x)-f_X(x)\right|^2 \geq \inf_{\widehat{f}\in\mathcal{T}_{iid}}\sup_{f_X\in\mathcal{F}_{x,r}(A,s)}\mathbb{E}\left|\widehat{f}(x)-f_X(x)\right|^2
$$

$$
\geq O\left(n^{-2s/(1+2\kappa+2s)}\right).
$$

Acknowledgements

The paper is supported by the Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grand number 101.02-2023.42.

Disclosure statement

The authors declare that they have no competing financial or non-financial interests related to this research.

References

- [1] Andersen, K.E. and Hansen, M.B. (2001). Multiplicative censoring: density estimation by a series expansion approach. Journal of Statistical Planning and Inference, Vol. 98, 137–155.
- [2] Asgharian, M. and Wolfson, D.B. (2005). Asymptotic behavior of the unconditional NPMLE of the length-biased survivor function from right censored prevalent cohort data. The Annals of Statistics, Vol. 33, 2109-2131.
- [3] Belomestny, D., Comte, F. and Genon-Catalot, V. (2016). Nonparametric Laguerre estimation in the multiplicative censoring model. Electronic Journal of Statistics, Vol. 10 3114–3152.
- [4] Belomestny, D. and Goldenshluger, A. (2020). Nonparametric density estimation from observations with multiplicative measurement errors. Ann. Inst. Henri Poincar´e Probab. Stat. 56 36–67.
- [5] Bradley, C.R. (2005). Basic properties of strong mixing conditions. A survey and some open questions. Probability Surveys, Vol. 2, 107-144.
- [6] Brunel, E., Comte, F. and Genon-Catalot, V. (2016). Nonparametric density and survival function estimation in the multiplicative censoring model. TEST 25 570–590.
- [7] Comte, F. and Dion, C. (2016). Nonparametric estimation in a multiplicative censoring model with symmetric noise. Journal of Nonparametric Statistics, Vol. 28 768–801.
- [8] Brenner Miguel, S., Comte, F. and Johannes, J. (2021). Spectral cut-off regularisation for density estimation under multiplicative measurement errors. Electronic Journal of Statistics, Vol. 15, 3551-3573.
- [9] Brenner Miguel, S., Comte, F. and Johannes, J. (2023). Linear functional estimation under multiplicative measurement error. Bernoulli, Vol. 29(3), 2247-2271.
- [10] Brenner Miguel, S., Phandoidaen, N. (2022). Multiplicative deconvolution in survival analysis under dependency. Statistics, Vol. 56, 297-328.
- [11] Genon-Catalot, V., Jeantheau, T. and Larédo, C. (2000). Stochastic volatility models as hidden Markov models and statistical applications. Bernoulli, Vol. 6(6), 1051-1079.
- [12] Hall, P. and Heyde, C.C. (1980). Martingale limit theory and its applications. Academic press.
- [13] Iacus, S.M. (2008). Simulation and inference for stochastic differential equations. Springer Series in Statistics, Springer, New York.
- [14] Kerkyacharian, G., Lepski, O. and Picard, D. (2001). Nonlinear estimation in anisotropic multi-index denoising. Probab. Theory Related Fields 121(2), 137-170.
- [15] N'drin, J.A. and Hili, O. (2013). Parameter estimation of one-dimensional diffusion process by minimum Hellinger distance method. Random Operators and Stochastic Equations, 21, 403-424.
- [16] Rosenblatt, M. (1956). A central limit theorem and strong mixing conditions. Proc. Nat. Acad. Sci., vol. 4, pp. 43-47.
- [17] Shou, W. and Gupta, S. (2023). Kernel density estimation using additive randomized response technique (RRT) models. Communications in Statistics - Simulation and Computation Online.
- [18] Trong, D.D., Hung, T.P. (2023). Parameter estimation for diffusion process from perturbed discrete observations. Communications in Statistics - Simulation and Computation, Vol. 52, no. 3, 925-944.
- [19] Van Es, B., Klaassen, C. A. J. and Oudshoorn, K. (2000). Survival analysis under cross-sectional sampling: length bias and multiplicative censoring. Journal of Statistical Planning and Inference, Vol. 91 295–312.
- [20] Vardi, Y. (1989). Multiplicative censoring, renewal processes, deconvolution and decreasing density: nonparametric estimation. Biometrika, Vol. 76, 751–761.
- [21] Vardi, Y. and Zhang, C.-H. (1992). Large sample study of empirical distributions in a random-multiplicative censoring model. The Annals of Statistics, Vol. 20, 1022–1039.
- [22] Volkonskii, V.A. and Rozanov, Yu.A. (1959). Some limit theorems for random functions. Theory Probab. Appl. 4, 178-197.