

Spectrum of density and spin density fluctuations of an attractive two-dimensional Fermi gas

Christian Apostoli,¹ Patrick Kelly,² Annette Lopez,³ Kaelyn Dauer,⁴ Gianluca Bertaina,⁵ Davide Emilio Galli,¹ and Ettore Vitali⁴

¹*Dipartimento di Fisica “Aldo Pontremoli”, Università degli Studi di Milano, via Celoria 16, 20133 Milano, Italy*

²*Institute for Physical Science and Technology, University of Maryland, College Park, Maryland 20742, USA*

³*Department of Physics, Brown University, 182 Hope Street, Providence, RI 02912, USA*

⁴*Department of Physics, California State University Fresno, Fresno, California 93720, USA*

⁵*Istituto Nazionale di Ricerca Metrologica, Strada delle Cacce 91, I-10135 Torino, Italy*

(Dated: March 21, 2024)

We leverage unbiased auxiliary-field quantum Monte Carlo methods interfaced with cutting-edge analytic continuation tools to compute from first-principles dynamical correlations for a dilute homogeneous two-dimensional attractive Fermi gas. Our main purpose is to quantitatively study the collective excitations of the system to shed light into fermionic superfluidity in the strongly correlated regime. Despite the fact that we are somewhat limited by finite-size effects, our study pinpoints a clear peak in the spin channel at low momentum, and we discuss the possibility that such a peak can be interpreted as the Higgs mode in the superfluid. Our study is complemented by a systematic comparison with the predictions of generalized random phase approximation, with the dual purpose to directly compare with auxiliary-field quantum Monte Carlo and to quantitatively study finite-size effects.

I. INTRODUCTION

The *ab-initio* calculation of the dynamical correlation functions of a strongly-correlated quantum system is a fundamental and challenging task which gives access to critical information about the system’s excitation spectrum. In particular, these correlation functions can be used to probe collective excitations, such as the celebrated Higgs or amplitude mode in superfluids. This mode is particularly hard to observe and it has been the center of a long-standing experimental effort, both in condensed matter physics [1, 2] and in atomic physics. In the realm of ultracold bosons, the Higgs mode has been investigated in a series of experiments employing lattice shaking or cavity-enhanced Bragg spectroscopy [3–5]. In ultracold Fermi gases, the actual detection of the amplitude mode has proved particularly challenging, since the Higgs mode in these systems has a very slow-decaying spectral tail [6, 7]. This corresponds to real-time damping due to the proliferation of quasiparticle excitations. Only very recently, experimental observations were reported, which used direct coherent excitation of the mode with radiofrequency pulses, Bragg spectroscopy following an interaction quench, or interaction modulation via periodic tuning of the magnetic field [8–11].

On the theoretical and computational side, the Higgs mode in neutral Fermi systems has been studied with various methodologies, including e.g. the time-dependent Bogoliubov-de Gennes equations [12–15], time-dependent density functional theory [16], the functional-integral method [6, 17, 18], and exact diagonalization (for few-body systems) [19]. More recently, relying on the generalized random-phase approximation (GRPA), it has been suggested that the Higgs mode could be detected in the dynamical correlations of the spin degrees of freedom [20] of an attractive fermionic system. If the interpretation in [20] is indeed correct, it would be very exciting to inquire experimentally, by measuring the spin dynamical structure factor through spin-sensitive Bragg spectroscopy [21, 22], thus extending the insightful investigations that can be performed in the density fluctuations sector [10, 23–25]. Ref. [20] also argued that the two-dimensional (2D) configuration, as

contrasted to the three-dimensional one mostly considered so far, renders the Higgs signal in the spin sector particularly prominent.

The investigation of spin correlation functions to address collective modes, and in particular the Higgs mode, in Fermi superfluids is fascinating, but correctly interpreting the results may be subtle. The key advantage of a spin probe is that the correlation functions at low momentum are expected to be significantly different from zero only close to 2Δ (Δ being the superfluid gap), which is exactly where the Higgs mode is expected, while the lower energy Goldstone mode is, at least partially, filtered out. At the same time, given the typical picture of the Higgs mode as a superfluid state with an order parameter whose modulus is oscillating, it seems realistic that a spin probe, which is essentially attempting to break pairs, might be sensitive to such oscillations. To put it simply, the spin probe can break a pair when the modulus is small enough and cannot when it is increased. The difficulty, on the other hand, is to “disentangle” the Higgs mode from the unavoidable quasi-particle pair continuum. The analysis in [26] indeed suggests that beyond mean-field studies are necessary to detect the coupling between the Higgs mode and the spin fluctuations.

Our plan for this work is to gain further insight by obtaining unbiased quantum Monte Carlo (QMC) results for the dynamical structure factor $S(\mathbf{q}, \omega)$ and its spin homologous $S^s(\mathbf{q}, \omega)$, yielding the spectrum of the fluctuations in density and spin, respectively. We also implement the generalized random phase approximation (GRPA) [20, 27–30], as a direct comparison for QMC results and as a tool to address finite-size effects.

In the density channel, we find qualitative agreement between the QMC results and the mean-field spectrum, with a renormalization of the Goldstone mode. In the spin channel, in spite of the difficulty in reaching small wave vectors, our results are consistent with a sharp peak, which could become even sharper if smaller wave vectors could be achieved. The energy of this peak is significantly lower than the one predicted by GRPA. We will discuss the possibility to interpret such a peak as the Higgs mode of the system.

The paper is organized as follows: Section II explains the Hamiltonian and the methodologies that we used: GRPA, QMC and the analytic continuation procedure. Then in Section III we describe our results and present a systematic comparison between GRPA and QMC. Finally, we draw our conclusions and present some perspectives in Section IV.

II. MODEL AND METHODS

Our starting point is the Hamiltonian for a collection of attractive unpolarized spin-1/2 fermions of equal mass m moving in 2D. In this context, the two spin species, denoted by \uparrow and \downarrow , respectively, represent the two lowest hyperfine states of ${}^6\text{Li}$ or ${}^{40}\text{K}$ atoms. We focus on the dilute regime, where the fine details of the inter-particle forces can be neglected, and consider an attractive zero-range interaction $v_{\uparrow\downarrow}(\mathbf{r}_1, \mathbf{r}_2) = -g\delta(\mathbf{r}_1 - \mathbf{r}_2)$. The Hamiltonian can be written, in field-theoretical notation, as:

$$\hat{H} = \hat{K} + \hat{V} \quad (1)$$

with:

$$\hat{K} = \int d\mathbf{r} \sum_{\sigma} \hat{\psi}_{\sigma}^{\dagger}(\mathbf{r}) \left(-\frac{\hbar^2 \nabla^2}{2m} \right) \hat{\psi}_{\sigma}(\mathbf{r}) \quad (2)$$

and:

$$\hat{V} = -g \int d\mathbf{r} \hat{\psi}_{\uparrow}^{\dagger}(\mathbf{r}) \hat{\psi}_{\downarrow}^{\dagger}(\mathbf{r}) \hat{\psi}_{\downarrow}(\mathbf{r}) \hat{\psi}_{\uparrow}(\mathbf{r}) \quad (3)$$

Our main objective is the calculation of the zero-temperature spin and density dynamical structure factors of the system in the dilute regime, in order to extract information about the low-energy collective excitations. We explore the dependence on the interaction strength $\log(k_F a)$, where k_F is the Fermi momentum (which will be defined later) and a the s -wave scattering length (in this work, we adopt the convention that the dimer binding energy ϵ_b of the contact model is related to a by $|\epsilon_b| = \hbar^2/m a^2$ [31]).

The zero-temperature density dynamical structure factor of N fermions is defined as the following two-body dynamical correlation function:

$$S(\mathbf{q}, \omega) = \int_0^{+\infty} \frac{dt}{2\pi N} e^{i\omega t} \left\langle e^{i\hat{H}t/\hbar} \hat{n}_{\mathbf{q}} e^{-i\hat{H}t/\hbar} \hat{n}_{-\mathbf{q}} \right\rangle \quad (4)$$

where the angular brackets denote a ground state expectation value and $\hat{n}_{\mathbf{q}}$ is the Fourier component of the particle density operator $\hat{n}(\mathbf{r}) = \hat{n}_{\uparrow}(\mathbf{r}) + \hat{n}_{\downarrow}(\mathbf{r}) = \hat{\psi}_{\uparrow}^{\dagger}(\mathbf{r})\hat{\psi}_{\uparrow}(\mathbf{r}) + \hat{\psi}_{\downarrow}^{\dagger}(\mathbf{r})\hat{\psi}_{\downarrow}(\mathbf{r})$. Similarly, we define the spin dynamical structure factor $S^s(\mathbf{q}, \omega)$ by replacing, in (4), the total particle density with the spin density $\hat{n}^s(\mathbf{r}) = \frac{1}{2}(\hat{n}_{\uparrow}(\mathbf{r}) - \hat{n}_{\downarrow}(\mathbf{r}))$.

We address the singularities due to the contact potential in (1) through lattice regularization, i.e., by introducing a high-momentum cutoff in the kinetic energy term:

$$\hat{K} \simeq \sum_{\sigma} \int_{[-\pi/b, \pi/b]^2} d\mathbf{k} \varepsilon(k) \hat{\psi}_{\sigma}^{\dagger}(\mathbf{k}) \hat{\psi}_{\sigma}(\mathbf{k}) \quad (5)$$

where the dispersion relation is, as usual, $\varepsilon(k) = \frac{\hbar^2 k^2}{2m}$. In Eq. (5), we integrate over the first Brillouin zone of a square lattice with parameter b , $(b\mathbb{Z})^2$. In addition, we further regularize the problem by introducing a supercell $\mathcal{L} = [-L/2, L/2]^2 \cap (b\mathbb{Z})^2$, i.e., a finite square lattice with $M = L/b \times L/b$ sites, and choosing periodic boundary conditions (PBC). The choice of PBC restricts the integration in (5) to a finite summation over the set of allowed momenta $\mathbf{k} = \frac{2\pi}{L}\mathbf{n}$, with $\mathbf{n} \in \mathbb{Z}^2$ such that $\mathbf{k} \in [-\pi/b, \pi/b]^2$. We observe that, within this regularization technique, the continuum limit can be recovered by letting $b \rightarrow 0$ and the infinite system limit can be recovered by letting $L \rightarrow +\infty$. The interaction term in the Hamiltonian is regularized by introducing a contact interaction of the form $v_{\uparrow\downarrow}(\mathbf{r}_1, \mathbf{r}_2) = -Ub^{-2}\delta_{\mathbf{r}_1, \mathbf{r}_2}$, where δ is now the discrete Kronecker delta, while the interaction strength of the discrete model is tuned in such a way that the lattice two-body problem has the same zero-energy s -wave scattering length a as the problem in the continuum. This regularization procedure maps the original Hamiltonian into a lattice model which we can write as:

$$\hat{H}_{\mathcal{L}} = \sum_{\mathbf{k}, \sigma} \varepsilon(k) \hat{\psi}_{\sigma}^{\dagger}(\mathbf{k}) \hat{\psi}_{\sigma}(\mathbf{k}) - U \sum_{\mathbf{r}} \hat{n}_{\uparrow}(\mathbf{r}) \hat{n}_{\downarrow}(\mathbf{r}) \quad (6)$$

Using the typical notation for ‘‘Hubbard-like’’ Hamiltonians, we can write $\varepsilon(k) = t|b\mathbf{k}|^2$ with hopping amplitude given by $t = \frac{\hbar^2}{2mb^2}$, and, as shown in [32]:

$$\frac{U}{t} = \frac{4\pi}{\log(k_F a) + \log(C\sqrt{n})} \quad (7)$$

where $n = N/M$ is the particle density on the lattice, $k_F = \frac{\sqrt{2\pi n}}{b}$ is Fermi momentum, a is the s -wave scattering length of the original problem in the continuum, while $C = 0.80261$ is a constant. In the following we will denote the Fermi energy of the lattice model by $\varepsilon_F = \varepsilon(k_F)$.

The crucial advantage of stepping from (1) to (6) is the possibility, for attractive unpolarized fermions, irrespective of the total density (filling), to calculate exactly the intermediate scattering function in imaginary time:

$$F(\mathbf{q}, \tau) = \frac{1}{N} \left\langle e^{\tau \hat{H}_{\mathcal{L}}} \hat{n}_{\mathbf{q}} e^{-\tau \hat{H}_{\mathcal{L}}} \hat{n}_{-\mathbf{q}} \right\rangle, \quad \tau \geq 0 \quad (8)$$

both in the density and the spin channels, by leveraging unbiased auxiliary-field quantum Monte Carlo methodologies (AFQMC), which are extensively explained in [33–35]. In fact, it is possible to map the imaginary time evolution operator $\exp(-\tau \hat{H}_{\mathcal{L}})$ into a random walk in the manifold of N particles’ Slater determinants, modeling independent fermions moving in a stochastic external field (the ‘‘auxiliary field’’). It has been shown that, whenever the system is spin-balanced and the interaction is attractive, the infamous sign problem does not affect the calculations and we can find exact results with a computational time which is polynomial in the size of the system [32, 34]. The dynamical structure factors can then be obtained by performing analytic continuation of the imaginary time data [36–38].

We employ the differential evolution for analytic continuation (DEAC) algorithm [39], which evolves a population

of candidate solutions for the dynamical structure factor over several generations, each one improving their average fitness and adaptively adjusting the control parameters. We use the DEAC code by A. Del Maestro’s group [40]. To assess the robustness of our results, we also performed some cross checks (not shown) employing the genetic inversion by falsification of theories (GIFT) algorithm [36, 37].

In addition to unbiased QMC calculations, we compute the dynamical structure factors within the generalized random phase approximation (GRPA) [20, 27–30], still relying on the above lattice regularization. GRPA implements a linear response theory approach, studying the response of the local (spin) density to a time-dependent perturbation coupled to the (spin) density itself, resulting in a Hamiltonian of the form:

$$\hat{H}(t) = \hat{H}_{\mathcal{L}} - \mu\hat{N} + \sum_{\mathbf{r}} u(\mathbf{r}, t)\hat{n}(\mathbf{r}) \quad (9)$$

where $\hat{H}_{\mathcal{L}}$ is the lattice Hamiltonian (6), μ is a chemical potential, \hat{N} the particle number operator, while $u(\mathbf{r}, t)$ is an external potential. GRPA approximately computes the response function $\chi(\mathbf{q}, \omega)$, yielding the change in the local density as the system evolves unitarily under the influence of the perturbation:

$$\delta\langle\hat{n}\rangle(\mathbf{q}, \omega) = \chi(\mathbf{q}, \omega)u(\mathbf{q}, \omega) \quad (10)$$

where the angular brackets denote an expectation value with respect to the time dependent quantum state, evolving from the unperturbed ground state. Within GRPA, the Hamiltonian $\hat{H}_{\mathcal{L}} - \mu\hat{N}$ is replaced by the most general mean-field breakup [28], and the order parameters are, at each time instant, self-consistently adjusted to follow the time-dependent perturbation. Finally, the dynamical structure factors can be obtained through the fluctuation-dissipation theorem:

$$S(\mathbf{q}, \omega) = -\frac{\hbar}{\pi n} \lim_{\eta \rightarrow 0^+} \Im \chi(\mathbf{q}, \omega + i\eta) \quad (11)$$

where n is the average density and η is a convergence parameter. In actual GRPA numerical implementations, the parameter η has to be set to some small positive number, as we will discuss in Section III. Our GRPA computations allow us to make direct comparison with QMC and with previous results in [20], where a different regularization is used. At the same time, since GRPA calculations only scale linearly with the lattice size M (they just require numerical summations over the Brillouin zone), their results allow us to readily estimate size effects in the calculations.

III. RESULTS

We use AFQMC to compute exactly the dynamical correlations in imaginary-time (8) for a system of $N = 26$ atoms using a regularization square lattice with $M = 1225$ sites. In this context, “exactly” means that, for the given system size, the systematic error in the properties of the lattice model is below the level of the statistical uncertainty. Thanks to the spin

balance and the absence of the infamous sign problem, this can be achieved with polynomial scaling. Previous AFQMC studies by some of us [43] on the spectral function of the 2D Fermi gas indicate that such a lattice is large enough to achieve accurate estimations in the bulk and in the continuum limit. As an independent probe of robustness, we also use GRPA with the dual purpose to compare with AFQMC and to assess finite-size effects, as we will discuss below. Our calculations yield the imaginary-time intermediate scattering functions $F(\mathbf{q}, \tau)$ in both the density and spin channels, for momenta as small as $q \simeq 0.5k_F$, k_F being the Fermi momentum. We perform DEAC analytic continuation of $F(\mathbf{q}, \tau)$ to extract the density dynamical structure factor $S(\mathbf{q}, \omega)$ and the spin dynamical structure factor $S^s(\mathbf{q}, \omega)$. We study the behavior of the structure factors as we increase the interaction strength from the weakly correlated BCS regime with $\log(k_F a) = 2.5$ to the more strongly correlated, though still not molecule-dominated, regime with $\log(k_F a) = 1.0$. The choice of the values of interaction was dictated by the intent to compare with the results in Ref. [20].

In Fig. 1, our AFQMC results in the density (left column) and spin channels (right column) are shown as circles, representing the maxima of the structure factors, together with vertical bars representing the widths at half heights (the detailed shapes of the AFQMC structure factors are shown in Figs. 4 and 5). The results are superimposed to background color plots, obtained with GRPA, for $S(\mathbf{q}, \omega)$ (left column) and for $S^s(\mathbf{q}, \omega)$ (right column) for a much larger system at the same particle density, precisely $N = 5850$ atoms moving on a regularization lattice with $M = 525 \times 525$ sites. The larger system allows for the investigation of a denser set of momentum values, providing a clearer picture, and gives important information about finite-size effects, as we will discuss below. The horizontal lines represent 2Δ , Δ being the superfluid gap (i.e., the energy needed to break a Cooper pair) as computed within a BCS mean-field approach (say Δ_{BCS}), informing the GRPA calculations (red dashed-dotted lines), or within AFQMC (white dashed lines) [43].

Our GRPA results, despite the different choice of regularization, appear to be consistent with the results in [20]. In the density channel, a sharp Nambu-Goldstone mode is visible at low energy $\omega < 2\Delta_{\text{BCS}}$; for higher energy $\omega > 2\Delta_{\text{BCS}}$ the quasiparticle pair continuum emerges, and the Nambu-Goldstone mode is strongly damped. At the AFQMC level, the lattice supercell is large enough to explore the behavior below the superfluid gap, and the results confirm the existence of a sharp Nambu-Goldstone mode, whose dispersion is renormalized by the correlations beyond mean-field: the peak at the lowest values of the momentum is at lower energy with respect to the GRPA prediction (this is not a finite-size effect, as we will show below), consistently with a significantly smaller superfluid gap, as was found in previous calculations in Ref. [43]. The AFQMC results at higher momentum are also consistent with the emergence of a quasiparticle pair continuum, as shown by the increasing size of the bars. We observe that, while there is no reason to expect that the Higgs mode cannot appear in the density structure factor, the dominant peak of the Goldstone mode and the broadening due to the quasiparticle




FIG. 1. Dynamical structure factors (arbitrary units) in the density (left) and spin (right) channels for interaction strengths $\log(k_F a) = 2.5$ (top), 1.5 (middle), and 1.0 (bottom). Dots: maxima of the structure factors obtained by QMC for a system of 26 fermions, with vertical bars representing their full width at half maximum. Color plots: dynamical structure factors (arbitrary units) obtained by GRPA for a larger system of $N = 5850$ fermions at the same density. Dashed horizontal lines: twice the quasiparticle gap for the system of $N = 26$ fermions; for interaction strength $\log(k_F a) = 2.5$, it is calculated from the theory by Gor'kov and Melik-Barkhudarov [41, 42], which was shown to be accurate in this weak-interaction condition [43]; for interaction strengths $\log(k_F a) = 1.5$ and 1.0, it is calculated with QMC from Ref. [43]. Dash-dotted horizontal lines: twice the quasiparticle gap predicted by BCS theory for the system of $N = 5850$ fermions.

pair continuum make the requirements for the detection of the amplitude mode beyond the resolution we have from analytic continuation. This makes the spin probe very attractive.

In the spin channel, as expected, within GRPA no excitation is present for $\omega < 2\Delta_{\text{BCS}}$. At higher energies, the GRPA spin dynamical structure factor appears to develop a sharp peak at low momentum which broadens at higher momenta. By comparing different interaction strengths, we observe that the spin structure factor gets flatter and fainter as the interaction strength increases (and $\log(k_F a)$ decreases). This is

consistent with the system being more and more similar to a Bose-Einstein condensate of tightly bound molecules, where the spin degrees of freedom are suppressed. We notice that the widths of the QMC results increase with the coupling more than within GRPA. This behavior is akin to recent experimental observations in the three-dimensional case [10, 11], where the damping of the Higgs excitation along the BCS-BEC crossover was stronger than expected from BCS theory.

The key result in [20] was the observation of the sharp spin mode, say $\omega_s(q)$, which the authors interpreted as the




FIG. 2. Spin dynamical structure factor $S^s(\mathbf{q}, \omega)$ (arbitrary units) for interaction strength $\log(k_F a) = 1.5$ and wave numbers $q = 0.492 k_F$ (left panel) and $0.695 k_F$ (right panel). The solid lines represent our results, with markers indicating selected data points to improve readability. Blue dots: QMC results for 26 fermions. Orange triangles: GRPA results for 26 fermions. Green squares: GRPA results for 5850 fermions. Dashed vertical line: twice the quasiparticle gap for 26 fermions, calculated with QMC from Ref. [43]. Dotted vertical line: twice the quasiparticle gap predicted by BCS theory for 26 fermions. Dash-dotted vertical line: twice the quasiparticle gap predicted by BCS theory for 5850 fermions.

celebrated Higgs mode, with a dispersion of the form $\omega_s(q) = 2\Delta_{\text{BCS}} + \alpha q^2$ as $q/k_F \rightarrow 0$. Our GRPA calculations are also consistent with a sharpening of the dynamical structure factor at low momentum, although we feel the need to be more cautious about the interpretation of this spin “mode” as the evidence of a detection of the Higgs mode. Is this peak the Higgs mode itself or is it just the low energy and low momentum tail of the quasiparticle pair continuum, which happens to be narrow due to the kinematics? We argue that in either case the result is very interesting: in fact, in a many-body system, this sharp peak, whatever its origin may be, shows that we may excite long-lived long wavelength spin waves in a superfluid, and there is the possibility that such spin waves may be an expression of the celebrated and elusive Higgs mode.

As we mentioned in the introduction, it is intuitively reasonable to expect that a spin probe may be sensitive to an excited state with an oscillating modulus of the order parameter, which is the typical picture of the Higgs mode. In fact, if we attempt to break a pair, like a spin probe would do, we are indeed probing the energy needed to break a pair and thus the modulus of the order parameter. So, there is no reason, from first principles, to assume that the overlap between the state $\hat{n}_s(-\mathbf{q})|\Psi_0\rangle$ and the excited states which build the Higgs mode in the bulk limit needs to vanish. A non-zero overlap would mean that the spin dynamical structure factor provides a promising correlation function to explore the Higgs mode. That being said, as it is evident from Eq. 15 in [20] and the analysis in [26] (Eq. 120 and following), within GRPA there seems to be no direct coupling between the dynamics of the order parameter and the spin excitations. Now, in our view, this observation does not necessarily invalidate the interpretation in [20], because, as discussed above, there may well be non-zero overlap, but it warrants further exploration using methodologies beyond GRPA.

At the unbiased AFQMC level, we are somewhat limited by the system size, which does not allow us to compute the structure factors below $|\mathbf{q}| \approx 0.5k_F$. Our AFQMC results for the

spin dynamical structure factor are shown in the right column of Fig. 1 and, in more detail, in the subsequent Figs. 2, 4, and 5. At the lowest considered momenta, the shape of the spin dynamical structure factor displays a clear peak, whose intensity decreases with the interaction strength and whose energy may be compatible with a mode that would converge to 2Δ (with the AFQMC pairing gap) in the limit $q \rightarrow 0$. At the same time, the peak does not appear very sharp, in particular if we compare it with the peaks in the density channel. Nevertheless, we suggest that the broadening of the peak may be due to the finite size. In order to corroborate such a statement, in Fig. 2 we show the spin dynamical structure factor $S^s(\mathbf{q}, \omega)$ for interaction strength $\log(k_F a) = 1.5$, focusing on the two smallest wave numbers $q = 0.492 k_F$ and $0.695 k_F$. We compare the AFQMC spectra for 26 atoms with the GRPA ones for 26 and 5850 atoms at the same density. All of them show a peak that moves to higher frequency as q increases from $0.492 k_F$ to $0.695 k_F$. The GRPA spectra exhibit maxima at approximately the same positions for both system sizes, with a sharper peak for the larger system. The AFQMC results have a width which is compatible with the corresponding GRPA result for 26 atoms, while the peaks are shifted to lower frequency, as expected from the renormalization of the gap. If we assume that the significant sharpening observed at the GRPA level for increasing size also happens at the correlated level, then we can suggest that a sharp mode does indeed exist also at the AFQMC level, and it is not an artifact of GRPA. In addition, since AFQMC includes correlations beyond mean field, we may cautiously claim that at least a part of the spectral weight of this peak is indeed a Higgs mode, with the kinematics possibly collaborating in preventing the proliferation of states in the continuum, to which the Higgs mode could decay.

We take a closer look at the size dependence within GRPA in Fig. 3, but, before delving into it, a discussion of our choices for the convergence parameter η is in order. The dynamical structure factor resulting from the GRPA theory is achieved in the limit $\eta \rightarrow 0^+$, see Eq. (11). In our implementation, we




FIG. 3. GRPA results for the dynamical structure factors (arbitrary units) in the density (top panels) and spin (bottom panels) channels of a system of 26 (left panels) and 5850 (right panels) fermions, for interaction strength $\log(k_F a) = 1.5$ and various values of the wave number q . The solid lines represent our results, with markers indicating selected data points to improve readability. Dotted vertical lines: twice the quasiparticle gap for 26 fermions, as predicted by BCS theory. Dash-dotted vertical lines: twice the quasiparticle gap for 5850 fermions, as predicted by BCS theory.

choose a finite positive value to broaden Dirac deltas in the dynamical structure factor. We tune the values empirically, and in particular we choose $\eta \approx 0.06\epsilon_F$ for 26 atoms and $\eta \approx 0.03\epsilon_F$ for 5850 atoms.

In Fig. 3, we compare the spectra for interaction $\log(k_F a) = 1.5$ and for two system sizes: $N = 26$ and $N = 5850$ atoms. The larger system shows the same qualitative behavior, and the positions of the peaks do not show significant change. On the other hand, the larger size displays significantly sharper peaks in both channels, allowing for a clearer distinction between the collective modes and the quasiparticle pair continuum. On one hand, this highlights that in the 26-atoms system there are still relevant finite-size effects, thus encouraging future efforts to allow AFQMC to access larger sizes; at the same time, if we assume that GRPA provides an accurate recipe for size corrections, this allows us to strengthen our claim that a sharp mode does indeed exist at the QMC level in the spin channel.

Fig. 4 shows the density (top panels) and spin (bottom panels) dynamical structure factors yielded by our QMC calculations for interaction strengths $\log(k_F a) = 1.5$ and 1.0 and selected momenta. The qualitative picture presented by the two interaction strengths is similar, the main quantitative difference being due to the different value of the superfluid gap. The density structure factors for small q are dominated by the Nambu-Goldstone peak. As q increases, this peak moves to

higher frequency and becomes less sharp, eventually merging with the quasiparticle pair continuum when it reaches the energy of twice the superfluid gap. The spin dynamical structure factors for small q exhibit a peak above twice the quasiparticle gap, which, as discussed before, may be consistent with a sharp Higgs peak. This peak moves to higher frequency as q increases and eventually significantly broadens and becomes compatible with a quasiparticle pair continuum.

Finally, Fig. 5 compares QMC and GRPA spectra for the system of 26 atoms and interaction $\log(k_F a) = 1.5$. Taking into account the renormalization of the quasiparticle gap, the GRPA method predicts the same qualitative behavior that we described in the QMC spectra.

IV. DISCUSSION AND CONCLUSIONS

In summary, our study employed the unbiased AFQMC method to compute imaginary-time correlation functions for a two-dimensional homogeneous system of attractive fermions. By leveraging analytic continuation techniques, we translated these correlations into the dynamical structure factor and its spin counterpart. Our current algorithm allowed us to compute the structure factors, in the (q, ω) plane, for momenta as low




FIG. 4. QMC results for the dynamical structure factors (arbitrary units) in the density (top panels) and spin (bottom panels) channels, for interaction strengths $\log(k_F a) = 1.5$ (left panels) and 1.0 (right panels), and various values of the wave number q . The solid lines represent our results, with markers indicating selected data points to improve readability. Dashed vertical lines: twice the quasiparticle gap, calculated with QMC from Ref. [43].

as $0.5 k_F$, which turned out to be small enough to resolve the collective modes from the quasiparticles pair continuum.

Incidentally, we observe that the possibility to obtain unbiased two-body dynamical correlations for Fermi superfluids for small momenta, well inside the Fermi sea, is very important beyond the scope of atomic physics, with potential applications in nuclear physics and nuclear astrophysics.

Comparisons with the results and the size effects obtained via GRPA has revealed qualitative agreement in the density channel and identified a distinct peak in the spin channel. This peak shows that long-lived long wavelength spin waves can be excited in a superfluid, whose dispersion is compatible with a quadratic behavior reminiscent of the celebrated Higgs mode, as pointed out in [20]; in addition, our results indicate a significant renormalization of the dispersion of such a mode due to the many-body correlations, consistent with a smaller value of the pairing gap in the unbiased calculations with respect to the mean-field result.

We feel the need to be cautious about identifying the observed low-momentum spin peak as a manifestation of the Higgs mode, but we argue that there is no reason to rule out such an interpretation from first principles. Moreover, we provide an intuitive argument suggesting that such a coupling between the dynamics of the order parameter and the spin excitations could indeed be expected. This possibility is intriguing

and signals the importance of investing in methodological advances in the QMC approach, particularly to access lower momenta, even deeper inside the Fermi sea. Despite the challenges posed by finite-size effects, our study thus contributes to the fundamental understanding of two-dimensional Fermi superfluids, providing unbiased results which can serve as crucial benchmarks for many-body theories and can provide important insight to experimentalists in the important challenge of measuring collective modes in superfluids. We expect that our results can stimulate experimental investigations leveraging spin-sensitive Bragg spectroscopy in two-dimensional Fermi gases, like it was done for three- and one-dimensional gases [21, 44].

A particularly exciting direction in the study of collective modes involves the investigation of the role of spin-polarization in attractive Fermi gases [45–47]; in fact, recent numerical findings confirmed the stability of an elusive Fulde-Ferrell-Larkin-Ovchinnikov phase at zero temperature [45], but the experimental detection is very complicated, and one of the possibilities is to measure the anisotropy of the speed of sound, which can be extracted from dynamical structure factors. In this context, the availability of numerical results, like the ones we presented in this paper, can be a very important asset to inform and guide the experimental search. For spin-polarized gases the inclusion of p -wave contributions might also be




FIG. 5. GRPA (left panels) and QMC (right panels) results for the dynamical structure factors (arbitrary units) in the density (top panels) and spin (bottom panels) channels, for a system of 26 atoms with interaction strength $\log(k_F a) = 1.5$ and various values of the wave number q . The solid lines represent our results, with markers indicating selected data points to improve readability. Dashed vertical lines: twice the quasiparticle gap, calculated with QMC from Ref. [43]. Dotted vertical lines: twice the quasiparticle gap, as predicted by BCS theory.

relevant [48].

As we navigate these perspectives, our findings open the doors for theoretical and experimental advancements, pushing the boundaries of our comprehension of dynamical properties within strongly correlated quantum systems.

V. ACKNOWLEDGEMENTS

E.V. acknowledges support from the National Science Foundation, grant number PHY-2207048. The AFQMC calcula-

tions were performed on the ACCESS (previously XSEDE) resources. Analytic continuation and GRPA were run on computational resources provided by INDACO Platform, which is a project of High Performance Computing at Università degli Studi di Milano.

-
- [1] R. Shimano and N. Tsuji, Higgs Mode in Superconductors, *Annu. Rev. Condens. Matter Phys.* **11**, 103 (2020).
- [2] D. Pekker and C. Varma, Amplitude/Higgs Modes in Condensed Matter Physics, *Annu. Rev. Condens. Matter Phys.* **6**, 269 (2015).
- [3] U. Bissbort, S. Götze, Y. Li, J. Heinze, J. S. Krauser, M. Weinberg, C. Becker, K. Sengstock, and W. Hofstetter, Detecting the Amplitude Mode of Strongly Interacting Lattice Bosons by Bragg Scattering, *Phys. Rev. Lett.* **106**, 205303 (2011).
- [4] M. Endres, T. Fukuhara, D. Pekker, M. Cheneau, P. Schauβ, C. Gross, E. Demler, S. Kuhr, and I. Bloch, The Higgs amplitude mode at the two-dimensional superfluid/Mott insulator transition, *Nature* **487**, 454 (2012).
- [5] J. Léonard, A. Morales, P. Zupanic, T. Donner, and T. Esslinger, Monitoring and manipulating Higgs and Goldstone modes in a supersolid quantum gas, *Science* **358**, 1415 (2017).
- [6] T. Cea, C. Castellani, G. Seibold, and L. Benfatto, Nonrelativistic Dynamics of the Amplitude (Higgs) Mode in Superconductors, *Phys. Rev. Lett.* **115**, 157002 (2015).
- [7] H. Kurkjian, J. Tempere, and S. N. Klimin, Linear response of a superfluid Fermi gas inside its pair-breaking continuum, *Sci. Rep.* **10**, 11591 (2020).

- [8] A. Behrle, T. Harrison, J. Kombe, K. Gao, M. Link, J.-S. Bernier, C. Kollath, and M. Köhl, Higgs Mode in a Strongly Interacting Fermionic Superfluid, *Nat. Phys.* **14**, 781 (2018).
- [9] L. Bayha, M. Holten, R. Klemm, K. Subramanian, J. Bjerlin, S. M. Reimann, G. M. Bruun, P. M. Preiss, and S. Jochim, Observing the Emergence of a Quantum Phase Transition Shell by Shell, *Nature* **587**, 583 (2020).
- [10] P. Dyke, S. Musolino, H. Kurkjian, D. J. M. Ahmed-Braun, A. Pennings, I. Herrera, S. Hoinka, S. J. J. M. F. Kokkelmans, V. E. Colussi, and C. J. Vale, Higgs oscillations in a unitary Fermi superfluid (2023), arXiv:2310.03452 [cond-mat.quant-gas].
- [11] A. Kell, M. Breyer, D. Eberz, and M. Köhl, Exciting the Higgs mode in a strongly-interacting Fermi gas by interaction modulation (2024), arXiv:2401.14822 [cond-mat.quant-gas].
- [12] R. G. Scott, F. Dalfovo, L. P. Pitaevskii, and S. Stringari, Rapid ramps across the bec-bcs crossover: A route to measuring the superfluid gap, *Phys. Rev. A* **86**, 053604 (2012).
- [13] S. Hannibal, P. Kettmann, M. D. Croitoru, A. Vagov, V. M. Axt, and T. Kuhn, Quench dynamics of an ultracold fermi gas in the bcs regime: Spectral properties and confinement-induced breakdown of the higgs mode, *Phys. Rev. A* **91**, 043630 (2015).
- [14] J. Tokimoto, S. Tsuchiya, and T. Nikuni, Excitation of Higgs Mode in Superfluid Fermi Gas in BCS–BEC Crossover, *J. Phys. Soc. Jpn.* **88**, 023601 (2019).
- [15] G. Lyu, K.-T. Xi, S. Yoon, Q. Chen, and G. Watanabe, Exciting the long-lived Higgs mode in superfluid Fermi gases with particle removal, *Phys. Rev. A* **107**, 023321 (2023).
- [16] A. Barresi, A. Boulet, G. Wlazlowski, and P. Magierski, Generation and decay of Higgs mode in a strongly interacting Fermi gas, *Sci. Rep.* **13**, 11285 (2023).
- [17] D. Phan and A. V. Chubukov, Following the Higgs mode across the BCS-BEC crossover in two dimensions, *Phys. Rev. B* **107**, 134519 (2023).
- [18] X. Han, B. Liu, and J. Hu, Observability of Higgs mode in a system without Lorentz invariance, *Phys. Rev. A* **94**, 033608 (2016).
- [19] J. Bjerlin, S. M. Reimann, and G. M. Bruun, Few-Body Precursor of the Higgs Mode in a Fermi Gas, *Phys. Rev. Lett.* **116**, 155302 (2016).
- [20] H. Zhao, X. Gao, W. Liang, P. Zou, and F. Yuan, Dynamical structure factors of a two-dimensional Fermi superfluid within random phase approximation, *New J. Phys.* **22**, 093012 (2020).
- [21] S. Hoinka, M. Lingham, M. Delehay, and C. J. Vale, Dynamic Spin Response of a Strongly Interacting Fermi Gas, *Phys. Rev. Lett.* **109**, 050403 (2012).
- [22] C. J. Vale and M. Zwierlein, Spectroscopic Probes of Quantum Gases, *Nat. Phys.* **17**, 1305 (2021).
- [23] S. Hoinka, P. Dyke, M. G. Lingham, J. J. Kinnunen, G. M. Bruun, and C. J. Vale, Goldstone mode and pair-breaking excitations in atomic Fermi superfluids, *Nat. Phys.* **13**, 943 (2017).
- [24] H. Biss, L. Sobirey, N. Luick, M. Bohlen, J. J. Kinnunen, G. M. Bruun, T. Lompe, and H. Moritz, Excitation Spectrum and Superfluid Gap of an Ultracold Fermi Gas, *Phys. Rev. Lett.* **128**, 100401 (2022).
- [25] L. Sobirey, H. Biss, N. Luick, M. Bohlen, H. Moritz, and T. Lompe, Observing the Influence of Reduced Dimensionality on Fermionic Superfluids, *Phys. Rev. Lett.* **129**, 083601 (2022).
- [26] L. He, Dynamic density and spin responses of a superfluid Fermi gas in the BCS–BEC crossover: Path integral formulation and pair fluctuation theory, *Ann. Phys.* **373**, 470 (2016).
- [27] R. Ganesh, A. Paramekanti, and A. A. Burkov, Collective modes and superflow instabilities of strongly correlated fermi superfluids, *Phys. Rev. A* **80**, 043612 (2009).
- [28] P. Kelly and E. Vitali, On the accuracy of random phase approximation for dynamical structure factors in cold atomic gases, *Atoms* **9**, 10.3390/atoms9040088 (2021).
- [29] Y. Yunomae, D. Yamamoto, I. Danshita, N. Yokoshi, and S. Tsuchiya, Instability of superfluid fermi gases induced by a rotonlike density mode in optical lattices, *Phys. Rev. A* **80**, 063627 (2009).
- [30] R. Combescot, M. Y. Kagan, and S. Stringari, Collective mode of homogeneous superfluid fermi gases in the bec-bcs crossover, *Phys. Rev. A* **74**, 042717 (2006).
- [31] G. Bertaina, Two-dimensional short-range interacting attractive and repulsive Fermi gases at zero temperature, *Eur. Phys. J. Spec. Top.* **217**, 153 (2013).
- [32] H. Shi, S. Chiesa, and S. Zhang, Ground-state properties of strongly interacting fermi gases in two dimensions, *Phys. Rev. A* **92**, 033603 (2015).
- [33] S. Zhang, Auxiliary-field quantum monte carlo for correlated electron systems, in *Emergent Phenomena in Correlated Matter: Modeling and Simulation Vol. 3*, edited by E. Pavarini, E. Koch, and U. Schollwöck (Verlag des Forschungszentrum Jülich, Jülich, Germany, 2013).
- [34] E. Vitali, H. Shi, M. Qin, and S. Zhang, Computation of dynamical correlation functions for many-fermion systems with auxiliary-field quantum monte carlo, *Phys. Rev. B* **94**, 085140 (2016).
- [35] M. Motta, D. E. Galli, S. Moroni, and E. Vitali, Imaginary time correlations and the phaseless auxiliary field quantum Monte Carlo, *J. Chem. Phys.* **140**, 024107 (2014).
- [36] E. Vitali, M. Rossi, L. Reatto, and D. E. Galli, Ab initio low-energy dynamics of superfluid and solid ^4He , *Phys. Rev. B* **82**, 174510 (2010).
- [37] G. Bertaina, D. E. Galli, and E. Vitali, Statistical and computational intelligence approach to analytic continuation in Quantum Monte Carlo, *Adv. Phys.:* **X 2**, 302 (2017).
- [38] E. Vitali, P. Kelly, A. Lopez, G. Bertaina, and D. E. Galli, Dynamical structure factor of a fermionic supersolid on an optical lattice, *Phys. Rev. A* **102**, 053324 (2020).
- [39] N. S. Nichols, P. Sokol, and A. Del Maestro, Parameter-free differential evolution algorithm for the analytic continuation of imaginary time correlation functions, *Phys. Rev. E* **106**, 025312 (2022).
- [40] N. S. Nichols, DEAC paper code repository (2021), <https://github.com/DelMaestroGroup/papers-code-DEAC>.
- [41] D. S. Petrov, M. A. Baranov, and G. V. Shlyapnikov, Superfluid transition in quasi-two-dimensional fermi gases, *Phys. Rev. A* **67**, 031601 (2003).
- [42] L. P. Gor'kov and T. K. Melik-Barkhudarov, Contribution to the theory of superfluidity in an imperfect fermi gas, *Sov. Phys. JETP* **13**, 1018 (1961).
- [43] E. Vitali, H. Shi, M. Qin, and S. Zhang, Visualizing the BEC-BCS crossover in a two-dimensional Fermi gas: Pairing gaps and dynamical response functions from ab initio computations, *Phys. Rev. A* **96**, 061601 (2017).
- [44] R. Senaratne, D. Cavazos-Cavazos, S. Wang, F. He, Y.-T. Chang, A. Kafle, H. Pu, X.-W. Guan, and R. G. Hulet, Spin-charge separation in a one-dimensional Fermi gas with tunable interactions, *Science* **376**, 1305 (2022).
- [45] E. Vitali, P. Rosenberg, and S. Zhang, Exotic Superfluid Phases in Spin-Polarized Fermi Gases in Optical Lattices, *Phys. Rev. Lett.* **128**, 203201 (2022).
- [46] L. Radzihovsky and D. E. Sheehy, Imbalanced Feshbach-resonant Fermi gases, *Rep. Prog. Phys.* **73**, 076501 (2010).
- [47] G. Bertaina and S. Giorgini, Density profiles of polarized Fermi gases confined in harmonic traps,

Phys. Rev. A **79**, 013616 (2009).

[48] G. Bertaina, M. G. Tarallo, and S. Pilati, Quantum Monte Carlo study of the role of p-wave interactions in ultracold repulsive Fermi gases, Phys. Rev. A **107**, 053305 (2023).