

# Variational Inference for Uncertainty Quantification: an Analysis of Trade-offs

**Charles C. Margossian**

*Center for Computational Mathematics  
Flatiron Institute  
New York, NY, USA*

CMARGOSSIAN@FLATIRONINSTITUTE.ORG

**Loucas Pillaud-Vivien**

*CERMICS Laboratory  
Ecole des Ponts ParisTech*

LOUCAS.PILLAUD-VIVIEN@ENPC.FR

**Lawrence K. Saul**

*Center for Computational Mathematics  
Flatiron Institute  
New York, NY, USA*

LSAUL@FLATIRONINSTITUTE.ORG

## Abstract

Given an intractable distribution  $p$ , the problem of variational inference (VI) is to find the best approximation from some more tractable family  $\mathcal{Q}$ . Commonly, one chooses  $\mathcal{Q}$  to be a family of factorized distributions (i.e., the mean-field assumption), even though  $p$  itself does not factorize. We show that this mismatch leads to an *impossibility theorem*: if  $p$  does not factorize, then any factorized approximation  $q \in \mathcal{Q}$  can correctly estimate at most one of the following three measures of uncertainty: (i) the marginal variances, (ii) the marginal precisions, or (iii) the generalized variance (which can be related to the entropy). In practice, the best variational approximation in  $\mathcal{Q}$  is found by minimizing some divergence  $D(q, p)$  between distributions, and so we ask: how does the choice of divergence determine which measure of uncertainty, if any, is correctly estimated by VI? We consider the classic Kullback-Leibler divergences, the more general Rényi divergences, and a score-based divergence which compares  $\nabla \log p$  and  $\nabla \log q$ . We provide a thorough theoretical analysis in the setting where  $p$  is a Gaussian and  $q$  is a (factorized) Gaussian. We show that all the considered divergences can be *ordered* based on the estimates of uncertainty they yield as objective functions for VI. Finally, we empirically evaluate the validity of this ordering when the target distribution  $p$  is not Gaussian.

**Keywords:** variational inference, Kullback-Leibler divergence, Rényi divergence, Fisher divergence, score matching

## 1 Introduction

In many problems, it is necessary to approximate an intractable distribution,  $p$ , by a more tractable distribution,  $q$ . This problem can be posed as an optimization, one whose goal is to discover the best-matching distribution  $q$  within some larger, parameterized family  $\mathcal{Q}$  of approximating distributions. Optimizations of this sort are the subject of a large literature on variational inference (VI; Jordan et al., 1999; Wainwright and Jordan, 2008; Blei et al., 2017).

Formally, suppose we wish to approximate a *target* distribution  $p$  over latent variables  $\mathbf{z} = z_{1:n} \in \mathcal{Z}$ , and let  $\mathcal{Q}$  be the family of approximating distributions. VI attempts to solve an optimization of the form

$$q^* = \operatorname{argmin}_{q \in \mathcal{Q}} D(q, p), \quad (1)$$

where  $D$  is a *divergence* satisfying (i)  $D(q, p) \geq 0$  for all  $q \in \mathcal{Q}$  and (ii)  $D(q, p) = 0$  if and only if  $p = q$ . In most applications,  $\mathcal{Q}$  is not sufficiently rich to capture all the features of  $p$ —that is  $p \notin \mathcal{Q}$ —and so, even after minimizing  $D(q, p)$ , the optimal approximation  $q$  must be compromised in some way. A crucial question is then whether  $q$  can still capture all the important features of  $p$  we may care about for a given application, even if other (less relevant) features are poorly estimated. In this paper, we study certain trade-offs that arise from the choice of  $\mathcal{Q}$ , and we analyze how these trade-offs are resolved by the particular choice of divergence in eq. (1).

Perhaps the most common choice for  $\mathcal{Q}$  is the family of *factorized* approximations,

$$q(\mathbf{z}) = \prod_{i=1}^n q_i(z_i). \quad (2)$$

We refer to this setting as F-VI. Variational families of this form originated in the mean-field approximation for physical models of ferromagnetism (Parisi, 1988). Factorized approximations are computationally convenient for many applications of machine learning because the number of variational parameters scales linearly with the dimensionality of  $\mathbf{z}$  (e.g Bishop et al., 2002; Blei, 2012; Giordano et al., 2023); in addition better scaling can be achieved for large data sets via amortization, which further restricts the form of  $\mathcal{Q}$  (Kingma and Welling, 2014; Margossian and Blei, 2024).

Our paper begins by formalizing the inherent trade-offs that arise in F-VI. Consider a target  $p$  which does *not* factorize but admits a finite covariance matrix  $\Sigma$ . By its very nature, the factorized approximation in eq. (2) cannot estimate the correlations between different components of  $\mathbf{z}$ . Nonetheless, a factorized approximation may still capture useful features of  $p$ , such as its mean and some quantification of uncertainty. There are many ways to characterize the uncertainty of a multivariate distribution  $p$ . In this paper, we specifically consider:

- (i) the marginal variances,  $\Sigma_{ii}$ , equal to the diagonal elements of the covariance matrix,
- (ii) the marginal precisions,  $\Sigma_{ii}^{-1}$ , equal to the diagonal elements<sup>1</sup> of the precision matrix (i.e., the inverse covariance matrix),
- (iii) the generalized variance,  $|\Sigma|$ , defined as the determinant of the covariance matrix.

Which measure of uncertainty to pursue depends on the application. It is especially common in Bayesian statistics to report the marginal variances of interpretable variables (Gelman et al., 2013), and in some models, the marginal precisions (Bernardo and Smith, 2000). On the other hand, sometimes it is useful to report a single scalar measure of uncertainty; this is provided by the generalized variance, a measure of uncertainty that is analogous (and in some cases equivalent) to that of the entropy in statistical physics (Parisi, 1988) and information theory (MacKay, 2003).

---

1. Throughout the paper, for any square matrix  $\mathbf{A}$ , we use  $A_{ij}^p$  to denote  $(A^p)_{ij}$ , the  $ij^{\text{th}}$  element of the  $p^{\text{th}}$  matrix power of  $A$ , and we use  $(A_{ij})^p$  to denote the  $p^{\text{th}}$  scalar power of the matrix element  $A_{ij}$ .

Divergence	Notation	Definition
(Reverse) Kullback-Leibler	$\text{KL}(q  p)$	$\mathbb{E}_q \left[ \log \frac{q}{p} \right]$
(Forward) Kullback-Leibler	$\text{KL}(p  q)$	$\mathbb{E}_p \left[ \log \frac{p}{q} \right]$
Rényi of order $\alpha \in (0, 1)$	$R_\alpha(p  q)$	$\frac{1}{\alpha(\alpha-1)} \mathbb{E}_q \left[ \frac{p^\alpha}{q^\alpha} - 1 \right]$
(Reverse) Score-based	$S(q  p)$	$\mathbb{E}_q \left[ \left\  \nabla \log \frac{q}{p} \right\ _{\text{Cov}(q)}^2 \right]$
(Forward) Score-based	$S(p  q)$	$\mathbb{E}_p \left[ \left\  \nabla \log \frac{p}{q} \right\ _{\text{Cov}(p)}^2 \right]$

Table 1: *Divergences that we analyze in this paper for FG-VI. The score-based divergences (Cai et al., 2024) are defined here for the special case that the base distribution is Gaussian.*

The first main result of this paper is an impossibility theorem: it states that any factorized approximation, of the form in eq. (2), can match *at most one* of the above three measures of uncertainty. An immediate consequence is that there exists no universally “best” factorized approximation of  $p$ ; what is best for one application may be of little use for another.

This ambiguity is reflected by the fact the solution to eq. (1) depends on the choice of divergence. This, in turn, begs the following question: which divergence is best suited for a particular application? By far the most popular choice in practice is the Kullback-Leibler divergence,  $\text{KL}(q||p)$ , which measures the disagreement between  $\log p$  and  $\log q$ . Other valid choices include the “forward”  $\text{KL}(p||q)$ , which arises in expectation propagation (Vehtari et al., 2020), the Rényi or  $\alpha$ -divergences  $R_\alpha(p||q)$  (Minka, 2005; Li and Turner, 2016), which interpolate between  $\text{KL}(q||p)$  and  $\text{KL}(p||q)$  when  $\alpha \in (0, 1)$ , and score-based (or Fisher) divergences (Courtade, 2016; Cai et al., 2024), which measure the discrepancy between  $\nabla \log p$  and  $\nabla \log q$ . Table 1 lists the divergences that we consider in this paper.

We investigate how each divergence in Table 1 resolves the constraints imposed by the impossibility theorem. To make progress, we identify a particular setting of VI where these questions can be fully analyzed. This is the setting where  $q$  belongs to a the family of Gaussians with a diagonal covariance matrix—a common choice in black box VI (Kucukelbir et al., 2017), which we call FG-VI—and where  $p$  is a Gaussian with a non-diagonal covariance matrix:

$$p = \text{normal}(\boldsymbol{\mu}, \boldsymbol{\Sigma}), \quad (3)$$

$$q = \text{normal}(\boldsymbol{\nu}, \boldsymbol{\Psi}), \quad (4)$$

with  $\Psi_{ij} = 0$  whenever  $i \neq j$ . This scenario, though relatively simple, is already rich enough to provide several counter-intuitive results, and it reveals in a stark way that different inferential goals are best served by different divergences. Furthermore, this setting has not been thoroughly analyzed when  $q$  minimizes the Rényi or the score-based divergences, likely because—unlike for the KL-divergences— $q$  then no longer admits a closed-form solution, even in this fundamental example. Understanding the properties of the optimal  $q$  for non-classical divergences requires a more technical machinery and fills a gap in the VI literature.

The second main result of the paper is an *ordering* of all the divergences in Table 1, based on estimates they yield of the marginal variances, marginal precisions, and generalized variance when  $p$  and  $q$  are Gaussian. This ordering provides a natural way of comparing divergences based on a downstream inferential task. While there exist many ways to compare divergences (e.g., Gibbs and Su, 2002), we believe that this ordering is particularly interpretable and helpful to practitioners. The ordering not only illuminates which divergence might yield the best estimate of some measure of uncertainty; it also cautions us that certain divergences can fail spectacularly—for instance, producing solutions for F-VI with variances that collapse to 0 or blow up to infinity.

**Outline.** In section 2, we state the main results of the paper—the impossibility theorem, and the ordering theorem—and we show how and where they fit into the existing literature. In section 3, we prove the impossibility theorem by analyzing the pairwise trade-offs that arise in F-VI between different measures of uncertainty. In section 4, we calculate the divergences in Table 1 and characterize the solutions to eq. (1) when  $p$  and  $q$  are Gaussian, as in eqs. (3–4). In section 5, we prove the ordering theorem by comparing the estimates of uncertainty obtained by minimizing each divergence. In section 6, we empirically investigate the validity of this ordering, on a variety of targets and data sets, when  $q$  is Gaussian but  $p$  is not. Finally, in section 7, we discuss our results in light of various applications which require uncertainty quantification (Bayesian statistics, pre-conditioning of Markov chain Monte Carlo, and information theory), and explore directions for future work.

## 2 Summary of contributions

The main results of the paper are an impossibility theorem for F-VI, starting from eq. (2), and an ordering theorem for the divergences in Table 1, starting from eqs. (3–4).

### 2.1 Main results

We begin with the impossibility theorem. To state the result, the following notation is useful. For any square matrix, we use  $\text{diag}(\cdot)$  to denote the vector formed from its diagonal elements, and we use vector inequalities, such as  $\text{diag}(\Psi) \leq \text{diag}(\Sigma)$ , to denote that  $\Psi_{ii} \leq \Sigma_{ii}$  for all  $i$ .

**Theorem 1 (Impossibility theorem for F-VI)** *Let  $p$  and  $q$  be distributions with covariances  $\Sigma$  and  $\Psi$ , respectively, where  $\Psi$  is diagonal but  $\Sigma$  is not. Then*

1. *(Variance matching)*  
If  $\text{diag}(\Psi) = \text{diag}(\Sigma)$ , then  $|\Psi| > |\Sigma|$  and  $\text{diag}(\Psi^{-1}) \leq \text{diag}(\Sigma^{-1})$ , and this last inequality is strict for at least one component along the diagonal.
2. *(Precision matching)*  
If  $\text{diag}(\Psi^{-1}) = \text{diag}(\Sigma^{-1})$ , then  $|\Psi| < |\Sigma|$  and  $\text{diag}(\Psi) \leq \text{diag}(\Sigma)$ , and this last inequality is strict for at least one component along the diagonal.

3. (Generalized variance matching)

If  $|\Psi| = |\Sigma|$ , then  $\Psi_{ii} < \Sigma_{ii}$  and  $\Psi_{jj}^{-1} < \Sigma_{jj}^{-1}$  for at least one  $i$  and  $j$ .

The theorem shows, for instance, that a factorized approximation with the correct marginal variances will overestimate the generalized variance and underestimate at least one of the marginal precisions. It also highlights similar trade-offs when the factorized approximation matches the marginal precisions or the generalized variance.

To understand how different choices of divergences resolve the trade-offs imposed by Theorem 1, we consider the case where  $p$  and  $q$  are Gaussian, as in eqs. (3-4). Note that in this setting  $p$  and  $q$  now have the same generalized variance if and only if they have the same entropy. All the divergences in Table 1 correctly estimate the mean of  $p$ , but each produces a different diagonal covariance matrix  $\Psi$ , and therefore different estimates of the marginal variances, marginal precisions and entropy. Moving forward, we say that one divergence *dominates* another if it always returns a larger estimate of the marginal variances of  $p$ .

**Definition 2 (Ordering of divergences)** Let  $\mathcal{P}$  and  $\mathcal{Q}$  denote the families of multivariate Gaussian distributions in eqs. (3-4), where  $\Psi$  is diagonal but  $\Sigma$  is not. Consider two divergences  $D_a$  and  $D_b$ , and for any  $p \in \mathcal{P}$ , let  $\Psi_a$  and  $\Psi_b$  denote the covariances of the factorized approximations obtained, respectively, by minimizing  $D_a(q, p)$  and  $D_b(q, p)$  over all  $q \in \mathcal{Q}$ . We say  $D_a$  dominates  $D_b$ , written

$$D_a \succ D_b,$$

if for any  $p \in \mathcal{P}$  we have that  $\text{diag}(\Psi_a) \geq \text{diag}(\Psi_b)$  and also that  $\Psi_a(i, i) > \Psi_b(i, i)$  for some element along the diagonal of these matrices.

The above definition<sup>2</sup> is based on an ordering of marginal variances, but it is also possible (because  $\Psi$  is diagonal) to define an equivalent ordering based on the marginal precisions. These orderings also imply an ordering of the generalized variances, though the converse is not true.

We now state the second main result of the paper, which is an ordering over all the divergences listed in Table 1.

**Theorem 3 (Ordering theorem for FG-VI)** The divergences in Table 1 are ordered in the sense given above. In particular, for any  $(\alpha_1, \alpha_2)$  satisfying  $0 < \alpha_1 < \alpha_2 < 1$ ,

$$S(q||p) \prec KL(q||p) \prec R_{\alpha_1}(p||q) \prec R_{\alpha_2}(p||q) \prec KL(p||q) \prec S(p||q). \quad (5)$$

The theorem shows, for instance, that the (reverse) score-based divergence yields smaller estimates of marginal variance than the (reverse) KL divergence, which in turn yields smaller estimates than all of the R enyi divergences. We prove Theorem 3 in Section 5. The main challenge here arises from the fact that the variational approximations from the R enyi and score-based divergences cannot be computed in closed form, and thus it is necessary (as done in Section 4) to characterize these solutions in other ways that can be further analyzed.

2. This definition can be further simplified if we impose the additional constraint that every row and column of  $\Sigma$  has nonzero off-diagonal elements. In this case, we can adopt the simpler definition that  $D_a \succ D_b$  if  $\Psi_a \succ \Psi_b$ , and all subsequent results hold with this more stringent characterization.

The proof of Theorem 3 also yields several intermediate results of interest. Most notably, we identify factorized variational approximations that match each measure of uncertainty that we consider:

- (i)  $q$  matches the marginal variances of  $p$  when minimizing  $\text{KL}(p||q)$  (MacKay, 2003);
- (ii)  $q$  matches the marginal precisions of  $p$  when minimizing  $\text{KL}(q||p)$  (Turner and Sahani, 2011);
- (iii)  $q$  matches the generalized variance (or entropy) of  $p$  when minimizing  $R_\alpha(p||q)$ , for some  $\alpha \in (0, 1)$ , which depends on  $\Sigma$ .

On the other hand,  $q$  can wildly misestimate uncertainty when minimizing a score-based divergence; in fact, the estimated variance may collapse to 0 when optimizing  $S(q||p)$  or blow up to infinity when optimizing  $S(p||q)$ . We refer to such cases as instances of *variational collapse*, where the variational approximation is not well-defined because  $\text{arginf}_{q \in \mathcal{Q}} D(q, p) \notin \mathcal{Q}$ .

## 2.2 Related work

Our work is in line with several efforts to evaluate VI through the lens of downstream inferential tasks. The mean and variances of  $p$  are of most interest for Bayesian statistics, but Huggins et al. (2020) caution that a low  $\text{KL}(q||p)$  divergence does not guarantee that  $q$  accurately estimates these quantities. Instead they recommend to estimate the Wasserstein distance, which can be used to bound the difference in moments of  $q$  and  $p$ ; see also Biswas and Mackey (2024). In latent variable models, VI is used to bound a marginal likelihood which is then maximized with respect to the model parameters (e.g., the weights of a neural network (Tomczak, 2022)). Several works examine the tightness of this bound (e.g Li and Turner, 2016; Dieng et al., 2017; Daudel et al., 2023) and the statistical properties of the resulting maximum likelihood estimator (Wang and Blei, 2018). VI is also used in tandem with other inference schemes—for example, to initialize Markov chain Monte Carlo (Zhang et al., 2022) or to estimate a proposal distribution for importance sampling. For the latter, the quality of VI can be assessed from the distribution of importance weights (Yao et al., 2018; Vehtari et al., 2024). In sum, there are many different ways to evaluate VI, and one lesson of our paper is that different inferential goals may be incompatible—may, in fact, compete against one another.

Many researchers have studied, both theoretically and empirically, how well F-VI estimates uncertainty (and, in particular, the marginal variances) when minimizing  $\text{KL}(q||p)$  (e.g., MacKay, 2003; Wang and Titterton, 2005; Bishop, 2006; Turner and Sahani, 2011; Giordano et al., 2018). Several works also examine the more specific setting in Section 4 where  $p$  and  $q$  are Gaussian with (respectively) a non-diagonal and diagonal covariance matrix. MacKay (2003) reports that  $q$  matches the marginal variances when minimizing  $\text{KL}(p||q)$ , while Turner and Sahani (2011) find that  $q$  matches the marginal precisions when minimizing  $\text{KL}(q||p)$ . In a precursor to this paper, Margossian and Saul (2023) prove that a precision-matching solution underestimates both the entropy and the marginal variances; here we provide a new and more intuitive proof of this result. We also note that Theorem 1 provides a more general framework for understanding these sorts of trade-offs, as it assumes neither that  $p$  and  $q$  are Gaussian nor that  $q$  is obtained by minimizing a particular divergence, such as  $\text{KL}(q||p)$ .

Previous studies in VI have investigated a large class of divergences for the optimization in eq. (1). A number of authors have considered the Rényi divergence (Li and Turner, 2016; Dieng et al., 2017; Daudel et al., 2021), for instance, as it provides a non-trivial generalization of the KL divergence. More recently, Cai et al. (2024) introduced the score-based divergence  $S(q||p)$  in Table 1 and showed that it can be efficiently optimized for Gaussian variational families. It is known that the Rényi divergence reduces in certain limits to the KL divergence (either from  $q$  to  $p$ , or vice versa); however, there is no such correspondence for the score-based divergences, and thus it is not obvious how the variational approximations obtained by minimizing  $S(q||p)$  or  $S(p||q)$  might compare to those obtained by minimizing  $\text{KL}(q||p)$  or  $\text{KL}(p||q)$ . Our analysis, notably the ordering in Theorem 3, sheds light on the matter.

### 3 Proof of Impossibility Theorem

We begin by restating the impossibility theorem of the previous section as a collection of trade-offs.

**Theorem 4 (Restatement of impossibility theorem)** *Let  $p$  and  $q$  be distributions over  $\mathbb{R}^N$  with covariances  $\Sigma$  and  $\Psi$ , respectively, where  $\Psi$  is diagonal but  $\Sigma$  is not.*

1. **Trade-off between marginal variances and generalized variance:**  
*If  $\text{diag}(\Psi) = \text{diag}(\Sigma)$ , then  $|\Psi| > |\Sigma|$ , and if  $|\Psi| = |\Sigma|$ , then  $\Psi_{ii} < \Sigma_{ii}$  for some  $i$ .*
2. **Trade-off between marginal precisions and generalized variance:**  
*If  $\text{diag}(\Psi^{-1}) = \text{diag}(\Sigma^{-1})$ , then  $|\Psi| < |\Sigma|$ , and if  $|\Psi| = |\Sigma|$ , then  $\Psi_{ii}^{-1} < \Sigma_{ii}^{-1}$  for some  $i$ .*
3. **Variance-precision trade-off:**  
*If  $\text{diag}(\Psi) = \text{diag}(\Sigma)$ , then  $\text{diag}(\Psi^{-1}) \leq \text{diag}(\Sigma^{-1})$  and  $\Psi_{ii}^{-1} < \Sigma_{ii}^{-1}$  for some  $i$ .  
 If  $\text{diag}(\Psi^{-1}) = \text{diag}(\Sigma^{-1})$ , then  $\text{diag}(\Psi) \leq \text{diag}(\Sigma)$  and  $\Psi_{ii} < \Sigma_{ii}$  for some  $i$ .*

The above restatement makes exactly the same claims as Theorem 1, but the proof is easier to organize along these lines.

We emphasize that the trade-offs listed above are not tied to any particular choice of the divergence  $D(q, p)$  in eq. (1). Rather, these trade-offs are inherent to the use of a factorized approximation, and they arise regardless of how  $q$  is chosen. Indeed, we will prove each trade-off by appealing only to the properties of  $\Sigma$  and  $\Psi$  as positive-definite matrices. However, for each trade-off, we will also provide additional intuitions (and in one case, an additional proof) that reflect the statistical setting in which these matrices arise.

**Proof of trade-off between marginal variances and generalized variance.** We begin by defining two auxiliary matrices whose elements are dimensionless (unlike those of  $\Sigma$  and  $\Psi$ ). The first of these is the *correlation* matrix  $\mathbf{C} \in \mathbb{R}^{n \times n}$  of the target distribution  $p$ , with elements

$$C_{ij} = \frac{\Sigma_{ij}}{\sqrt{\Sigma_{ii}\Sigma_{jj}}}, \tag{6}$$

which is positive-definite and has unit diagonal entries,  $C_{ii} = 1$ . The second is the diagonal matrix  $\mathbf{R} \in \mathbb{R}^{n \times n}$  that records the *ratios* of marginal variances in  $q$  and  $p$ ; i.e.,

$$R_{ii} = \frac{\Psi_{ii}}{\Sigma_{ii}}. \quad (7)$$

In what follows, it will be convenient to consider the generalized variances on a log scale. Taking differences, we have

$$\log |\Sigma| - \log |\Psi| = \log \left| \Psi^{-\frac{1}{2}} \Sigma \Psi^{-\frac{1}{2}} \right| = \log \left| \mathbf{R}^{-\frac{1}{2}} \mathbf{C} \mathbf{R}^{-\frac{1}{2}} \right| = \log |\mathbf{R}|^{-1} - \frac{1}{2} \log |\mathbf{C}|^{-1}. \quad (8)$$

The first determinant on the right side of eq. (8) is easy to compute because the matrix  $\mathbf{R}$ , as defined in eq. (7), is diagonal. Notably, we can bound the second determinant by the Hadamard inequality (Horn and Johnson, 2012, Theorem 7.8.1), which states that

$$|\mathbf{C}| < \prod_{i=1}^n C_{ii} = 1. \quad (9)$$

We note that the inequality in eq. (9) is strict under the assumption that  $\Sigma$  (and hence also  $\mathbf{C}$ ) is not diagonal. Combining the last two results, we see that

$$\log |\Sigma| - \log |\Psi| < \sum_i \log \left( \frac{\Sigma_{ii}}{\Psi_{ii}} \right). \quad (10)$$

Now suppose that  $\text{diag}(\Psi) = \text{diag}(\Sigma)$ ; then from eq. (10) we see that  $|\Sigma| < |\Psi|$ , proving the first part of the trade-off. Conversely, suppose that  $|\Sigma| = |\Psi|$ ; then from eq. (10) we see that  $\sum_i \log \Sigma_{ii} > \sum_i \log \Psi_{ii}$ , and this can only be true if  $\Sigma_{ii} > \Psi_{ii}$  for some  $i$ . This proves the second part of the trade-off. ■

The above trade-off has a particularly interesting interpretation in the case where  $p$  and  $q$  are Gaussian, as in eqs. (3–4). In this case, the generalized variance of  $q$ , given by  $|\Psi|$ , provides effectively the same measure of uncertainty as the *entropy* of  $q$ , which is equal to  $\frac{1}{2} \log |\Psi|$  plus an additive constant. Thus when  $q$  correctly estimates the marginal variances, it overestimates the entropy, and when it correctly estimates the entropy, it underestimates at least one marginal variance. The entropy of  $q$  in this setting is also governed by its own *shrinkage-delinkage trade-off* (Margossian and Saul, 2023): the entropy of  $q$  is decreased when  $q$  underestimates the marginal variances of  $p$  (the shrinkage effect), but it is increased when  $q$  ignores the correlations in  $p$  between off-diagonal components (the delinkage effect).

**Proof of trade-off between marginal precisions and generalized variance.** This proof follows the same structure as the preceding one. We define auxiliary matrices  $\tilde{\mathbf{C}}$  and  $\tilde{\mathbf{R}}$  with the dimensionless elements

$$\tilde{C}_{ij} = \frac{\Sigma_{ij}^{-1}}{\sqrt{\Sigma_{ii}^{-1} \Sigma_{jj}^{-1}}}, \quad \tilde{R}_{ii} = \frac{\Psi_{ii}^{-1}}{\Sigma_{ii}^{-1}}, \quad (11)$$

where  $\tilde{\mathbf{C}}$  has all ones along the diagonal and  $\tilde{\mathbf{R}}$  has all zeroes off the diagonal. In terms of these matrices, in analogy to eq. (8), we can write

$$\log |\Sigma| - \log |\Psi| = \log \left| \mathbf{R}^{-\frac{1}{2}} \mathbf{C} \mathbf{R}^{-\frac{1}{2}} \right| = \log \left| \tilde{\mathbf{R}}^{\frac{1}{2}} \tilde{\mathbf{C}}^{-1} \tilde{\mathbf{R}}^{\frac{1}{2}} \right| = \log |\tilde{\mathbf{R}}| - \frac{1}{2} \log |\tilde{\mathbf{C}}|. \quad (12)$$

As before, the determinant of  $\tilde{\mathbf{R}}$  is easy to compute because it is a diagonal matrix, and again we deduce from the Hadamard inequality that  $|\tilde{\mathbf{C}}| < \prod_i \tilde{\mathbf{C}}_{ii} = 1$ . Thus we find

$$\log |\Sigma| - \log |\Psi| > \sum_i \log \left( \frac{\Psi_{ii}^{-1}}{\Sigma_{ii}^{-1}} \right). \quad (13)$$

Note that the direction of inequality in eq. (13) is reversed from that in eq. (10). Now suppose that  $\text{diag}(\Psi^{-1}) = \text{diag}(\Sigma^{-1})$ ; then from eq. (13) we see that  $|\Sigma| > |\Psi|$ , proving the first part of the trade-off. Conversely, if  $|\Sigma| = |\Psi|$ , then from eq. (13) we see that  $\sum_i \log \Sigma_{ii}^{-1} > \sum_i \log \Psi_{ii}^{-1}$ , and this can only be true if  $\Sigma_{ii}^{-1} > \Psi_{ii}^{-1}$  for some  $i$ . This proves the second part of the trade-off.  $\blacksquare$

We can also interpret this trade-off in the special case when  $p$  and  $q$  are Gaussian, as in eqs. (3-4). In this case, it states the following: when  $q$  correctly estimates the marginal precisions, it underestimates the entropy, and when it correctly estimates the entropy, it underestimates at least one marginal precision.

Next we prove the final trade-off between marginal variances and precisions. A short but self-contained proof of this trade-off was given in Margossian and Saul (2023). Here instead we appeal to a stronger result from linear algebra (Horn and Johnson, 2012, Theorem 7.17), which will also prove useful in Section 5. This result is given by the following lemma.

**Lemma 5** *Let  $\mathbf{A} \in \mathbb{R}^{n \times n}$ . If  $\mathbf{A} \succ \mathbf{0}$ , then*

$$A_{ii} A_{ii}^{-1} \geq 1 \quad \text{for all } i, \quad (14)$$

$$A_{jj} A_{jj}^{-1} > 1 \quad \text{for some } j; \quad (15)$$

*in particular, the inequality is strict along the  $j^{\text{th}}$  component whenever  $\sum_{k \neq j} |A_{jk}| > 0$ .*

From this lemma, we obtain the variance-precision trade-off by substituting either the covariance matrix  $\Sigma$  or the precision matrix  $\Sigma^{-1}$  in Theorem 4 for the matrix  $\mathbf{A}$  in eqs. (14-15). Unfortunately, neither the above lemma nor the proof in Margossian and Saul (2023) offer much intuition. Here, we propose an alternative proof rooted in statistical notions.

**Proof of variance-precision trade-off.** Let  $\xi \sim \mathcal{N}(0, \Sigma)$ , and let  $\xi_{-i}$  denote all the elements of  $\xi$  other than  $\xi_i$ . Also denote by  $\pi$  its density. From the conditional distribution

$$\pi(\xi_i | \xi_{-i}) \propto \exp \left( -\frac{1}{2} \xi_i \Sigma_{ii}^{-1} \xi_i - \xi_i \sum_{j \neq i} \Sigma_{ij}^{-1} \xi_j \right), \quad (16)$$

we can identify (from the first quadratic term in the exponent) that  $\text{Var}[\xi_i | \xi_{-i}] = 1/\Sigma_{ii}^{-1}$ ; in other words, the conditional variance of this variable is given by the reciprocal of its

marginal precision. Now by the law of total variance, we have that

$$\Sigma_{ii} = \text{Var}[\xi_i], \quad (17)$$

$$= \mathbb{E}[\text{Var}[\xi_i|\boldsymbol{\xi}_{-i}]] + \text{Var}[\mathbb{E}[\xi_i|\boldsymbol{\xi}_{-i}]], \quad (18)$$

$$\geq \mathbb{E}[\text{Var}[\xi_i|\boldsymbol{\xi}_{-i}]], \quad (19)$$

$$= \mathbb{E}[1/\Sigma_{ii}^{-1}], \quad (20)$$

$$= 1/\Sigma_{ii}^{-1}. \quad (21)$$

Thus we have shown  $\Sigma_{ii}\Sigma_{ii}^{-1} \geq 1$  for all  $i$ . Moreover, since by assumption  $\boldsymbol{\Sigma}$  is not diagonal, there must exist some coordinate such that  $\mathbb{E}[\xi_i|\boldsymbol{\xi}_{-i}] \neq \mathbb{E}[\xi_i]$  and  $\text{Var}[\mathbb{E}[\xi_i|\boldsymbol{\xi}_{-i}]] > 0$ ; for this coordinate, the inequality is strict with  $\Sigma_{ii}\Sigma_{ii}^{-1} > 1$ . Suppose now that  $q$  matches the marginal precisions of  $p$ ; then for all elements along the diagonal, we have

$$\Psi_{ii} = (\Psi_{ii}^{-1})^{-1} = (\Sigma_{ii}^{-1})^{-1} \leq \Sigma_{ii}, \quad (22)$$

and this inequality must be strict for at least one of the marginal variances. This proves one part of the trade-off. Likewise, if  $q$  matches the marginal variances of  $p$ , then for all elements along the diagonal we have

$$\Psi_{ii}^{-1} = (\Psi_{ii})^{-1} = (\Sigma_{ii})^{-1} \leq \Sigma_{ii}^{-1}, \quad (23)$$

and this inequality must be strict for at least one of the marginal precisions. This proves the other part of the trade-off.  $\blacksquare$

## 4 Divergences for FG-VI

In this section we derive the solutions for FG-VI that are obtained from eq. (1–4) by minimizing the divergences in Table 1. We also highlight the properties of solutions that are obtained in this way. As a useful reference, we summarize the results for these solutions in Table 2. We emphasize that all of the results in this section are based on the further assumption that  $p$  and  $q$  are Gaussian, as in eqs. (3–4).

### 4.1 KL divergences

The KL divergence is not a symmetric function of its arguments, and thus the solutions for FG-VI depend on the direction of the divergence that is minimized. We consider each direction in turn.

**Reverse direction:  $\text{KL}(q||p)$ .** Most VI is based on minimizing the reverse KL divergence in Table 1. The implications of this choice have been extensively studied for FG-VI. For completeness, we review certain key results in the notation of this paper. When  $q$  is chosen to minimize  $\text{KL}(q||p)$ , it matches the mean of  $p$  (that is,  $\boldsymbol{\nu} = \boldsymbol{\mu}$ ), and its covariance is given by

$$\text{diag}(\boldsymbol{\Psi}^{-1}) = \text{diag}(\boldsymbol{\Sigma}^{-1}). \quad (24)$$

Further details of these calculations can be found in the references (Turner and Sahani, 2011; Margossian and Saul, 2023). We summarize the main properties of this solution for uncertainty quantification, all of which follow from the results of the previous section:

	Reduced divergence	(Implicit) Solution
$\text{KL}(q  p)$	$\frac{1}{2}\text{trace}(\Psi\Sigma^{-1}) - \log \Psi\Sigma^{-1}  + c$	$\Psi_{ii} = 1/\Sigma_{ii}^{-1}$
$\text{KL}(p  q)$	$\frac{1}{2}\sum_{i=1}^n(\log\frac{\Psi_{ii}}{\Sigma_{ii}} + \frac{\Sigma_{ii}}{\Psi_{ii}}) + c$	$\Psi_{ii} = \Sigma_{ii}$
$\text{R}_\alpha(p  q)$	$\frac{1}{\alpha(\alpha-1)}\left[ \Sigma_\alpha ^{-\frac{1}{2}} \Psi ^{\frac{\alpha}{2}} \Sigma ^{\frac{1-\alpha}{2}} - 1\right]$	$\Psi_{ii} = [\alpha\Sigma^{-1} + (1-\alpha)\Psi^{-1}]_{ii}^{-1}$
$\text{S}(q  p)$	$\text{trace}[(\mathbf{I} - \Psi\Sigma^{-1})]$	$\text{argmin}_{\mathbf{s}\geq 0} [\frac{1}{2}\mathbf{s}^T\mathbf{H}\mathbf{s} - \mathbf{1}^T\mathbf{s}], s_{ii} = \Psi_{ii}\Sigma_{ii}^{-1}$
$\text{S}(p  q)$	$\text{trace}[(\mathbf{I} - \Sigma\Psi^{-1})]$	$\text{argmin}_{\mathbf{t}\geq 0} [\frac{1}{2}\mathbf{t}^T\mathbf{J}\mathbf{t} - \mathbf{1}^T\mathbf{t}], t_{ii} = \Psi_{ii}^{-1}\Sigma_{ii}$

Table 2: Divergences when  $p$  and  $q$  are Gaussian (eq 3-4). We report the reduced divergence, obtained after matching the means of  $p$  and  $q$ . For the Rényi divergence, the optimal covariance can be found by solving a fixed-point equation. For the score-based divergences, we need to solve a non-negative quadratic program. The auxiliary matrix  $\Sigma_\alpha$  is defined in eq. (29),  $\mathbf{H}$  in eq. (50), and  $\mathbf{J}$  in eq. (53).

- (i) The marginal precisions of  $q$  match those of  $p$ .
- (ii) The entropy of  $q$  underestimates the entropy of  $p$ .
- (iii) The marginal variances of  $q$  underestimate the marginal variances of  $p$ .
- (iv) The marginal variances of  $q$  match the conditional variances of  $p$  when each Gaussian random variable is conditioned on all the others.

It has been widely observed that VI based on the reverse KL divergence tends to underestimate uncertainty (MacKay, 2003; Minka, 2005; Turner and Sahani, 2011; Blei et al., 2017; Giordano et al., 2018). The above results show that the “uncertainty deficit” of FG-VI depends on the measure that is used to quantify uncertainty. Indeed, when  $p$  is Gaussian, the marginal precisions are correctly estimated by minimizing  $\text{KL}(q||p)$ ; moreover, it has been shown that when  $\Sigma$  has *constant* off-diagonal entries, the entropy gap between  $p$  and  $q$  only grows as  $\mathcal{O}(\log n)$ , while the entropy itself grows as  $\mathcal{O}(n)$ , meaning that the fractional entropy gap tends to 0 as  $n \rightarrow \infty$  (Margossian and Saul, 2023, Theorem 3.6). Crucially, for  $q$  to estimate the entropy well, it must necessarily underestimate the marginal variances, as prescribed by eq. (8). Other asymptotically correct results are obtained in the thermodynamic limit of Ising models with constant, long-range interactions, where the mean-field approximation has its roots (Parisi, 1988).

**Forward direction:  $\text{KL}(p||q)$ .** Next we consider the forward KL divergence in Table 1. This divergence is not generally minimized for VI because it involves an expectation, namely  $\mathbb{E}_p[\log(p/q)]$  with respect to the target distribution  $p$ . When  $p$  is not tractable, it is not possible to compute this expectation analytically, and it may be too expensive to estimate this expectation by sampling. Still, it is illuminating to contrast the properties of the reverse and forward KL divergences for FG-VI. The latter is similarly minimized by setting  $\nu = \mu$ , but now we obtain a different solution when the remaining terms are minimized with respect

to  $\Psi$ . In particular,  $\text{KL}(p||q)$  is minimized by setting

$$\text{diag}(\Psi) = \text{diag}(\Sigma), \quad (25)$$

thus matching the marginal distributions of  $p$  and  $q$  (MacKay, 2003). We summarize the main properties of this solution for uncertainty quantification, all of which follow directly from the Impossibility Theorem:

- (i) The marginal variances of  $q$  match those of  $p$ .
- (ii) The entropy of  $q$  overestimates the entropy of  $p$ .
- (iii) The marginal precisions of  $q$  underestimate the marginal precisions of  $p$ .

Comparing the results for  $\text{KL}(p||q)$  and  $\text{KL}(q||p)$ , we see the important effects of the divergence when VI is used for uncertainty quantification. These effects motivate our subsequent study of the Rényi and score-based divergences.

## 4.2 Rényi divergence

The Rényi divergence of order  $\alpha$  (or the  $\alpha$ -divergence) is given by

$$R_\alpha(p||q) = \frac{1}{\alpha(\alpha-1)} \int \left( \frac{p^\alpha(\mathbf{z})}{q^\alpha(\mathbf{z})} - 1 \right) q(\mathbf{z}) d\mathbf{z}, \quad (26)$$

The  $\alpha$ -divergence has also been studied in the context of VI (Li and Turner, 2016). It can be defined in various ways, but these various definitions have in common that they all yield the same underlying optimization for VI. We use the above definition (Cichocki and Amari, 2010) to recover the KL divergences in the previous sections as limiting cases:

$$\lim_{\alpha \rightarrow 0} R_\alpha(p||q) = \text{KL}(q||p), \quad (27)$$

$$\lim_{\alpha \rightarrow 1} R_\alpha(p||q) = \text{KL}(p||q). \quad (28)$$

Thus for  $\alpha \in (0, 1)$ , the Rényi divergences provide a one-parameter family of divergences interpolating between  $\text{KL}(q||p)$  and  $\text{KL}(p||q)$ . Likewise, for  $\alpha = 2$ , eq. (26) recovers the  $\chi^2$ -divergence, which can also be minimized for approximate inference (Dieng et al., 2017).

Moving forward, we restrict our attention to the case  $\alpha \in (0, 1)$ , and as shorthand, we define the matrices

$$\Sigma_\alpha = \alpha\Psi + (1-\alpha)\Sigma, \quad (29)$$

$$\Phi_\alpha = \alpha\Sigma^{-1} + (1-\alpha)\Psi^{-1}, \quad (30)$$

which, for  $\alpha \in (0, 1)$ , are convex combinations of the covariance and precision matrices of  $p$  and  $q$  in eqs. (3-4). Note that these matrices are also related by the identity  $\Sigma_\alpha = \Sigma \Phi_\alpha \Psi$ . In terms of these matrices, the  $\alpha$ -divergence in eq. (26) is given by

$$R_\alpha(p||q) = \frac{1}{\alpha(\alpha-1)} \left[ e^{-\frac{\alpha(1-\alpha)}{2}(\mu-\nu)^\top \Sigma_\alpha^{-1}(\mu-\nu)} |\Sigma_\alpha|^{-\frac{1}{2}} |\Psi|^{\frac{\alpha}{2}} |\Sigma|^{\frac{1-\alpha}{2}} - 1 \right], \quad (31)$$

and it is only defined for values of  $\alpha$  such that  $\Phi_\alpha \succ 0$  (Burbea, 1984; Liese and Vajda, 1987; Hobza et al., 2009; Gil et al., 2013). (If  $\Phi_\alpha$  is not positive definite, then the integral in eq. (26) does not exist, and the  $\alpha$ -divergence is undefined; when  $\alpha \in (0, 1)$ , it is always the case that  $\Phi_\alpha \succ 0$  by convexity of the positive-definite matrix cone.) With this notation, we have the following result.

**Proposition 6 (Rényi fixed-point equations)** *Let  $\alpha \in (0, 1)$ , and let  $p$  and  $q$  be given by eqs. (3-4) where  $\Psi$  is diagonal. Then the Rényi divergence in eq. (26) is minimized when  $\nu = \mu$  and the estimated precisions from  $\Psi$  satisfy the fixed-point equation*

$$\text{diag}(\Psi^{-1}) = \text{diag}(\Sigma_\alpha^{-1}), \quad (32)$$

*or equivalently when the estimated variances from  $\Psi$  satisfy the fixed-point equation*

$$\text{diag}(\Psi) = \text{diag}(\Phi_\alpha^{-1}). \quad (33)$$

**Proof** We proceed by optimizing the expression for the Rényi divergence in eq. (31). When  $\alpha \in (0, 1)$ , the prefactor  $1/\alpha(\alpha-1)$  is *negative*, and the expression as a whole is minimized by *maximizing* the first term in brackets. Upon taking logarithms, we see equivalently that

$$\underset{q \in \mathcal{Q}}{\text{argmin}} [R_\alpha(p||q)] = \underset{\nu, \Psi}{\text{argmin}} \left[ \alpha(1-\alpha)(\mu-\nu)^\top \Sigma_\alpha^{-1}(\mu-\nu) + \log |\Sigma_\alpha| - \alpha \log |\Psi| \right]. \quad (34)$$

The first term on the right is minimized (and zeroed) by setting  $\nu = \mu$ , thus matching the means of  $q$  and  $p$ . Eliminating this term, and substituting for  $\Sigma_\alpha$  from eq. (29), we see that

$$\underset{q \in \mathcal{Q}}{\text{argmin}} [R_\alpha(p||q)] = \underset{\Psi}{\text{argmin}} \left[ \log |\alpha\Psi + (1-\alpha)\Sigma| - \alpha \log |\Psi| \right]. \quad (35)$$

The expression on the right of eq. (35) diverges if any diagonal element  $\Psi_{ii}$  vanishes *or* goes to infinity, so the minimum must exist at some  $\Psi \succ \mathbf{0}$ . Solving for the minimum, we find

$$0 = \frac{\partial}{\partial \Psi_{ii}} [\log |\alpha\Psi + (1-\alpha)\Sigma| - \alpha \log |\Psi|], \quad (36)$$

$$= \alpha [\alpha\Psi + (1-\alpha)\Sigma]_{ii}^{-1} - \alpha \Psi_{ii}^{-1}, \quad (37)$$

$$= \alpha [\Sigma_\alpha^{-1} - \Psi^{-1}]_{ii}, \quad (38)$$

thus proving eq. (32). Finally, we observe that as  $\alpha \rightarrow 0$ , we recover the precision-matching solution obtained when minimizing  $\text{KL}(q||p)$ .

To derive the fixed-point equation in eq. (33), we rewrite eq. (35) as

$$\underset{q \in \mathcal{Q}}{\text{argmin}} [R_\alpha(p||q)] = \underset{\Psi}{\text{argmin}} \left[ \log |\alpha\Psi + (1-\alpha)\Sigma| - \alpha \log |\Psi| + \log |\Psi| - \log |\Psi| \right], \quad (39)$$

$$= \underset{\Psi}{\text{argmin}} \left[ \log |\alpha + (1-\alpha)\Sigma\Psi^{-1}| - (1-\alpha) \log |\Psi^{-1}| \right], \quad (40)$$

$$= \underset{\Psi^{-1}}{\text{argmin}} \left[ \log |\alpha\Sigma^{-1} + (1-\alpha)\Psi^{-1}| - (1-\alpha) \log |\Psi^{-1}| \right], \quad (41)$$

where in the last step we have exploited the fact that additive constants, such as  $\log |\Sigma^{-1}|$ , do not change the location of the minimum with respect to  $\Psi$  (or equivalently, with respect to  $\Psi^{-1}$ ). Solving for the minimum, we find that

$$0 = \frac{\partial}{\partial \Psi_{ii}^{-1}} \left[ \log |\alpha\Sigma^{-1} + (1-\alpha)\Psi^{-1}| - (1-\alpha) \log |\Psi^{-1}| \right] = (1-\alpha) [\Phi_\alpha^{-1} - \Psi]_{ii}, \quad (42)$$

which is equivalent to the claim that  $\text{diag}(\Psi) = \text{diag}(\Phi_\alpha^{-1})$  in eq. (33). ■

Proposition 6 provides two (equivalent) fixed-point equations for the variational parameter  $\Psi$ , and it will prove useful to work with both in Section 5. In componentwise notation, these fixed-point equations take the form

$$\Psi_{ii}^{-1} = [\alpha\Psi + (1-\alpha)\Sigma]_{ii}^{-1}, \quad (43)$$

$$\Psi_{ii} = [\alpha\Sigma^{-1} + (1-\alpha)\Psi^{-1}]_{ii}^{-1}, \quad (44)$$

and again we see that eq. (43) reduces to the precision-matching solution  $\Psi_{ii}^{-1} = \Sigma_{ii}^{-1}$  as  $\alpha \rightarrow 0$  and eq. (44) reduces to the variance-matching solution  $\Psi_{ii} = \Sigma_{ii}$  as  $\alpha \rightarrow 1$ .

### 4.3 Score-based divergence

Next we consider the score-based divergences presented in Table 1. These divergences are similar to the relative Fisher information (Courtade, 2016, eq. (7)) except that they use a *weighted* norm (Cai et al., 2024, Appendix A) to measure the difference between the gradients of  $\log p$  and  $\log q$ . When  $p$  and  $q$  are Gaussian, these score-based divergences are given by

$$S(q||p) = \int d\mathbf{z} q(\mathbf{z}) \|\nabla \log q(\mathbf{z}) - \nabla \log p(\mathbf{z})\|_{\text{Cov}(q)}^2, \quad (45)$$

$$S(p||q) = \int d\mathbf{z} q(\mathbf{z}) \|\nabla \log q(\mathbf{z}) - \nabla \log p(\mathbf{z})\|_{\text{Cov}(p)}^2, \quad (46)$$

where we use  $\|\mathbf{v}\|_{\mathbf{A}}^2 = \mathbf{v}^\top \mathbf{A} \mathbf{v}$  to denote the weighted norm for any  $\mathbf{A} \succ \mathbf{0}$ . The score-based divergences in eqs. (45-46) are dimensionless measures of the discrepancy between  $p$  and  $q$ , and in particular, like the KL and Rényi divergences in the previous sections, they are invariant to affine reparameterizations of the support of these distributions (Cai et al., 2024, Theorem A.4). For  $p$  and  $q$  in eqs. (3-4), these divergences are given by

$$S(q||p) = \text{tr} \left[ (\mathbf{I} - \Psi \Sigma^{-1})^2 \right] + (\boldsymbol{\nu} - \boldsymbol{\mu})^\top \Sigma^{-1} \Psi \Sigma^{-1} (\boldsymbol{\nu} - \boldsymbol{\mu}), \quad (47)$$

$$S(p||q) = \text{tr} \left[ (\mathbf{I} - \Sigma \Psi^{-1})^2 \right] + (\boldsymbol{\mu} - \boldsymbol{\nu})^\top \Psi^{-1} \Sigma \Psi^{-1} (\boldsymbol{\mu} - \boldsymbol{\nu}), \quad (48)$$

as also shown in Cai et al. (2024). The rest of this section examines the solutions that minimize these expressions with respect to the variational mean  $\boldsymbol{\nu}$  and diagonal covariance  $\Psi$ .

First we consider the minimization of the *reverse* score-based divergence,  $S(q||p)$ . This minimization can be formulated as a nonnegative quadratic program (NQP)—that is, a convex instance of quadratic programming with nonnegativity constraints.

**Proposition 7 (NQP for minimizing  $S(q||p)$ )** *Let  $p$  and  $q$  be given by eqs. (3-4) where  $\Psi$  is diagonal. Then the reverse score-based divergence in eq. (47) is minimized by setting  $\boldsymbol{\nu} = \boldsymbol{\mu}$  and solving the quadratic program*

$$\min_{\mathbf{s} \geq 0} \left[ \frac{1}{2} \mathbf{s}^\top \mathbf{H} \mathbf{s} - \mathbf{1}^\top \mathbf{s} \right], \quad (49)$$

where  $\mathbf{s}$  is constrained to lie in the nonnegative orthant,  $\mathbf{1}$  is the vector of all ones,  $\mathbf{H}$  is the positive-definite matrix with elements

$$H_{ij} = \frac{\left(\Sigma_{ij}^{-1}\right)^2}{\Sigma_{ii}^{-1}\Sigma_{jj}^{-1}}, \quad (50)$$

and  $\Psi$  is obtained from the solution of eq. (49) by identifying  $s_i = \Psi_{ii}\Sigma_{ii}^{-1}$ .

**Proof** It is clear that eq. (47) is minimized by setting  $\boldsymbol{\nu} = \boldsymbol{\mu}$  in the rightmost term. For the remaining term, we note that

$$\frac{1}{2} \text{tr} \left[ (\mathbf{I} - \Psi \Sigma^{-1})^2 \right] = \frac{n}{2} - \sum_i \Psi_{ii} \Sigma_{ii}^{-1} + \frac{1}{2} \sum_{ij} \Psi_{ii} \Psi_{jj} \left( \Sigma_{ij}^{-1} \right)^2, \quad (51)$$

and the NQP in eq. (49) is obtained by defining  $s_i = \Psi_{ii}\Sigma_{ii}^{-1}$ . Finally, we observe that the matrix  $\mathbf{H}$  in eq. (50) is always positive-definite. Indeed,  $\mathbf{H}$  is the Hadamard (i.e., component-wise) square of the correlation matrix built from the precision matrix  $\Sigma^{-1}$ , and we note from the Schur product theorem that the Hadamard product of two positive-definite matrices is also positive-definite (Horn and Johnson, 2012). ■

Given the symmetry of eqs. (47-48), it is not surprising that we obtain an analogous optimization when minimizing the *forward* score-based divergence,  $S(p||q)$ , in terms of the variational parameters  $\boldsymbol{\nu}$  and  $\Psi$ . In particular, we have the following result.

**Proposition 8 (NQP for minimizing  $S(q||p)$ )** Let  $p$  and  $q$  be given by eqs. (3-4) where  $\Psi$  is diagonal. Then the forward score-based divergence in eq. (47) is minimized by setting  $\boldsymbol{\nu} = \boldsymbol{\mu}$  and solving the quadratic program

$$\min_{\mathbf{t} \geq 0} \left[ \frac{1}{2} \mathbf{t}^\top \mathbf{J} \mathbf{t} - \mathbf{1}^\top \mathbf{t} \right], \quad (52)$$

where  $\mathbf{t}$  is constrained to lie in the nonnegative orthant,  $\mathbf{J}$  is the positive-definite matrix with elements

$$J_{ij} = \frac{\left(\Sigma_{ij}\right)^2}{\Sigma_{ii}\Sigma_{jj}}, \quad (53)$$

and  $\Psi$  is obtained from the solution of eq. (52) by identifying  $t_i = \Psi_{ii}^{-1}\Sigma_{ii}$ .

**Proof** The proof follows the same steps as in the previous proposition, but with the matrices  $\Psi$  and  $\Sigma$  playing the reverse roles as they did in eq. (51). ■

We conclude this section by noting a peculiar property of the score-based divergences in Table 1. It is possible for the solutions of the NQPs in Propositions 7 and 8 to lie on the boundary of the nonnegative orthant. That is, there may exist some  $i^{\text{th}}$  component along the diagonal of  $\Psi$  such that  $S(q||p)$  is minimized by setting  $\Psi_{ii} = 0$  (in Proposition 7) or such




Figure 1: When FG-VI is based on minimizing the score-based divergences in Table 1, it may estimate zero or infinite values for the marginal variances. The red areas indicate these occurrences of variational collapse when FG-VI with a score-based divergence is used to approximate a three-dimensional Gaussian with a non-diagonal correlation matrix  $\mathbf{C}$ .

that  $S(p||q)$  is minimized by setting  $\Psi_{ii}^{-1} = 0$  (in Proposition 8). Strictly speaking, solutions of this form—with zero or infinite marginal variances—do not define proper distributions that lie in the family  $\mathcal{Q}$  of factorized approximations. Such solutions never arise with FG-VI based on the KL or Rényi divergences for  $\alpha \in \{0, 1\}$ . This phenomenon gives rise to the following definition.

**Definition 9** Let  $\mathcal{Q}$  denote the family of factorized Gaussian approximations in eq. (2) that are proper distributions (i.e., with  $0 < \mathcal{H}(q) < \infty$ ). For some target distribution  $p$  and divergence  $D$ , we say that FG-VI undergoes **variational collapse** when  $\operatorname{arginf}_{q \in \mathcal{Q}} D(q, p) \notin \mathcal{Q}$ .

Fig. 1 illustrates this collapse when FG-VI is used to approximate a multivariate Gaussian distribution in three dimensions with varying degrees of correlation; the amount of correlation is determined by the magnitude of the off-diagonal elements  $C_{12}$ ,  $C_{23}$ , and  $C_{13}$  in eq. (6). These off-diagonal elements are constrained by the fact that the correlation matrix as a whole must be positive definite. Each panel in the figure visualizes a two-dimensional slice of the three-dimensional (convex) set of positive-definite correlation matrices for a fixed value of  $C_{23}$ . The blue regions of each slice indicate the areas where the NQPs in Propositions 7 and 8 yield finite and nonzero estimates of the variances; the red regions indicate areas of variational collapse. (It can be shown, in three dimensions, that the forward and reverse score-based divergences share the same areas of variational collapse.) Interestingly, the leftmost panel shows that variational collapse does not occur in two dimensions or arise purely from pairwise correlations. On the other hand, the remaining panels show that variational collapse occurs in three dimensions whenever the correlation matrix is dense and has at least one off-diagonal element of sufficiently large magnitude.

## 5 Ordering of divergences for FG-VI

We have seen that different divergences  $D(q, p)$  yield different solutions to the problem of variational inference in eq. (1); we have also seen, in turn, that these different solutions




Figure 2: Comparison of variances, precisions, and entropy estimated by FG-VI with different divergences. *In the left and center panels, the variance is normalized by the variance of  $p$ , and the precision by the precision of  $p$ , both along the first coordinate. In the right panel, we plot the difference between the estimated entropy and the entropy of  $p$ . Here FG-VI was used to approximate a 2-dimensional Gaussian with a non-diagonal correlation matrix  $\mathbf{C}$ . The Rényi divergence was computed for  $\alpha=0.5$ . The ordering of the curves matches the predictions of Theorem 3.*

yield different estimates of the marginal variances, precisions, and entropy. In the setting where both  $p$  and  $q$  are Gaussian, these estimates provide a natural way to compare and even order the divergences; see Definition 2. In this section, we show that the divergences in Table 1 can be ordered in this way; that is, for  $0 < \alpha_1 < \alpha_2 < 1$ , we prove that

$$S(q||p) \prec \text{KL}(q||p) \prec R_{\alpha_1}(p||q) \prec R_{\alpha_2}(p||q) \prec \text{KL}(p||q) \prec S(p||q),$$

as stated in Theorem 3.

Figure 2 illustrates the consequences of this ordering when FG-VI is used to approximate a 2-dimensional Gaussian with varying degrees of correlation. The amount of correlation is measured by the determinant of the correlation matrix,  $|\mathbf{C}|$ ; as this determinant decreases from one to zero—from no correlation to perfect correlation—the target Gaussian more starkly violates the assumption of factorization. From left to right, the panels plot the estimates of variance, precision, and entropy that are obtained from FG-VI with different divergences, and we see that for all values of  $|\mathbf{C}|$ , these estimates are ordered exactly as predicted by Theorem 3.

We have already proven one of these orderings. Recall from section 4.1 that when FG-VI is based on minimizing  $\text{KL}(q||p)$ , it underestimates the marginal variances, at least one of them strictly. On the other hand, when FG-VI is based on minimizing  $\text{KL}(p||q)$ , it correctly estimates the marginal variances. The rest of this section is devoted to proving the other orderings in Theorem 3.

### 5.1 Ordering of score-based divergences

In this section we prove the two outermost orderings in Theorem 3; specifically, we show that  $S(q||p) \prec \text{KL}(q||p)$  and  $\text{KL}(p||q) \prec S(p||q)$ . At a high level, these proofs are obtained by analyzing the Karush-Kuhn-Tucker (KKT) conditions of the quadratic programs in Propositions 7 and 8.

**Proof of  $S(q||p) \prec \mathbf{KL}(q||p)$ .** Consider the diagonal covariance matrix  $\Psi$  that FG-VI estimates by minimizing  $S(q||p)$  on the left side of this ordering. This matrix is obtained from the solution of the NQP in eq. (49) for the unknown nonnegative variables  $s_i = \Psi_{ii}\Sigma_{ii}^{-1}$ . The solution to this NQP must satisfy the KKT conditions: namely, for each component of  $\mathbf{s}$ , we have either that (i)  $s_i = 0$  and  $\mathbf{H}\mathbf{s}_i > 1$  or (ii)  $s_i > 0$  and  $\mathbf{H}\mathbf{s}_i = 1$ . We examine each of these cases in turn:

- $s_i = 0$  This is a case of variational collapse where by minimizing  $S(q||p)$  we estimate that  $\Psi_{ii} = 0$ .
- $s_i > 0$  In this case we observe that  $(\mathbf{H}\mathbf{s})_i = 1$  from the KKT condition and also that  $H_{ij} \geq 0$  and  $H_{ii} = 1$  from eq. (50). Thus we see that

$$s_i = H_{ii}s_i = 1 - \sum_{j \neq i} H_{ij}s_j \leq 1, \quad (54)$$

or equivalently, that  $\Psi_{ii}\Sigma_{ii}^{-1} \leq 1$ .

We have thus shown that  $\Psi_{ii}\Sigma_{ii}^{-1} \leq 1$  for all of the marginal variances estimated by minimizing the score-based divergence  $S(q||p)$ . Next we show that this inequality must be strict for at least one component. The claim is trivially true if any  $s_i = 0$ , since this implies  $\Psi_{ii}\Sigma_{ii}^{-1} = 0$ . Suppose to the contrary that every element of  $\mathbf{s}$  is strictly positive. Since by assumption  $\Sigma$  is not diagonal, it must also be true that  $\Sigma^{-1}$  is not diagonal, and therefore from eq. (50) we can pick some  $i$  and  $k$  such that  $H_{ik} > 0$ . But then, continuing from eq. (54), we see that

$$s_i = 1 - \sum_{j \neq i} H_{ij}s_j \leq 1 - H_{ik}s_k < 1, \quad (55)$$

or equivalently that  $\Psi_{ii}\Sigma_{ii}^{-1} < 1$ , thus proving the claim. In sum we have shown that  $\Psi_{ii} \leq 1/\Sigma_{ii}^{-1}$  for all of the marginal variances estimated by minimizing  $S(q||p)$ , and also that this inequality is strict for at least one component. Finally, we recall from eq. (24) that  $1/\Sigma_{ii}^{-1}$  is the marginal variance estimated by minimizing  $\mathbf{KL}(q||p)$ . Per Definition 2 we conclude that  $S(q||p) \prec \mathbf{KL}(q||p)$ .  $\blacksquare$

We obtain a symmetric result when considering the forward score-matching divergence.

**Proof of  $S(p||q) \succ \mathbf{KL}(p||q)$ .** Examining the KKT conditions as above, one of the two following cases must hold:

- $t_i = 0$  This is a case of variational collapse where by minimizing  $S(p||q)$  we estimate that  $\Psi_{ii}^{-1} = 0$ , that is  $\Psi_{ii} = \infty$  and so  $\Psi_{ii} > \Sigma_{ii}$ .
- $t_i > 0$  In this case,

$$t_i \leq 1 \iff \Psi_{ii}^{-1}\Sigma_{ii} \leq 1 \iff \Psi_{ii} \geq \Sigma_{ii}. \quad (56)$$

Since  $\Sigma$  is non-diagonal, we also have that  $\mathbf{J}$  is non-diagonal, and so there must exist a coordinate  $j$  such that  $\Psi_{jj} > \Sigma_{jj}$ .  $\blacksquare$

## 5.2 Ordering of KL divergences

Next we prove the two intermediate orderings in Theorem 3; specifically, for any  $\alpha \in (0, 1)$ , we show that  $\text{KL}(q||p) \prec \text{R}_\alpha(p||q)$  and  $\text{R}_\alpha(p||q) \prec \text{KL}(p||q)$ . These proofs are obtained by applying the inequalities in Lemma 5 to the fixed-point equations for the optimal marginal variances in eqs. (43–44), obtained by minimizing the Rényi divergence of order  $\alpha$ .

**Proof of  $\text{KL}(q||p) \prec \text{R}_\alpha(p||q)$  for  $\alpha \in (0, 1)$ .** Let  $\Psi$  denote the diagonal covariance matrix estimated by minimizing  $\text{R}_\alpha(p||q)$ , and consider the matrix  $[\alpha \Sigma^{-1} + (1-\alpha)\Psi]^{-1}$  that appears on the right side of its fixed-point equation in eq. (44). This matrix is positive-definite and non-diagonal for all  $\alpha \in (0, 1)$ , and thus we can apply the inequality in eq. (14) of Lemma 5 to its diagonal elements. In this way, we find:

$$\alpha \Psi_{ii} \Sigma_{ii}^{-1} = \Psi_{ii} [\alpha \Sigma^{-1} + (1-\alpha)\Psi^{-1}]_{ii} - (1-\alpha) \quad (57)$$

$$\geq \frac{\Psi_{ii}}{[\alpha \Sigma^{-1} + (1-\alpha)\Psi^{-1}]_{ii}^{-1}} - (1-\alpha), \quad (58)$$

$$= (\Psi_{ii}/\Psi_{ii}) - (1-\alpha), \quad (59)$$

$$= \alpha. \quad (60)$$

Dividing both sides by  $\alpha \Sigma_{ii}^{-1}$ , we see that  $\Psi_{ii} \geq 1/\Sigma_{ii}^{-1}$  for all  $i$ . Next we appeal to the strict inequality of Lemma 5 and conclude that  $\Psi_{jj} > 1/\Sigma_{jj}^{-1}$  for some  $j$ . On the left and right sides of these inequalities appear, respectively, the marginal variances estimated by minimizing  $\text{R}_\alpha(p, q)$  and  $\text{KL}(q, p)$ . Thus we have shown  $\text{R}_\alpha(p||q) \succ \text{KL}(q, p)$ . ■

**Proof of  $\text{R}_\alpha(p||q) \prec \text{KL}(p||q)$  for  $\alpha \in (0, 1)$ .** The proof follows the same steps as the previous one, but now we combine the inequalities of Lemma 5 with the fixed-point equation in eq. (43). In this way we find:

$$(1-\alpha)\Psi_{ii}^{-1}\Sigma_{ii} = \Psi_{ii}^{-1}[\alpha\Psi + (1-\alpha)\Sigma]_{ii} - \alpha \geq \frac{\Psi_{ii}^{-1}}{[\alpha\Psi + (1-\alpha)\Sigma]_{ii}^{-1}} - \alpha = 1-\alpha. \quad (61)$$

Dividing both sides by  $(1-\alpha)\Psi_{ii}^{-1}$ , we see that  $\Sigma_{ii} \geq \Psi_{ii}$  for all  $i$ , and we know from the strict inequality in eq. (15) of Lemma 5 that  $\Sigma_{jj} > \Psi_{jj}$  for some  $j$ . On the left and right sides of these inequalities appear, respectively, the marginal variances estimated by minimizing  $\text{KL}(p||q)$  and  $\text{R}_\alpha(p||q)$ . Thus we have shown  $\text{KL}(p||q) \succ \text{R}_\alpha(p||q)$ . ■

## 5.3 Ordering of Rényi divergences

In this section we prove the ordering of the Rényi divergences in Theorem 3. This proof is more technical, relying on a detailed analysis of the fixed point equations in Proposition 6.

**Proof of  $\text{R}_{\alpha_1}(p||q) \prec \text{R}_{\alpha_2}(p||q)$  for  $0 < \alpha_1 < \alpha_2 < 1$ .** Let  $\Psi(\alpha)$  denote the diagonal covariance matrix that FG-VI estimates by minimizing  $\text{R}_\alpha(p||q)$  and note by inspection of  $\text{R}_\alpha(p||q)$  that  $\Psi(\alpha)$  is smooth. For  $\alpha \in (0, 1)$ , we know from the results of Section 5.2 that  $\Psi_{ii}(\alpha)$  is sandwiched between its limiting values at zero and one:

$$\frac{1}{\Sigma_{ii}^{-1}} = \lim_{\alpha \rightarrow 0^+} \Psi_{ii}(\alpha) \leq \Psi_{ii}(\alpha) \leq \lim_{\alpha \rightarrow 1^-} \Psi_{ii}(\alpha) = \Sigma_{ii}. \quad (62)$$




Figure 3: (Left) Either  $\Psi_{jj}(\alpha)$  is *strictly increasing* over  $\alpha \in (0, 1)$ , or it is *not*, with some minimal point  $\tau$  of vanishing derivative. We prove the former by showing that no such point  $\tau$  exists. (Right) The proof is based on properties of the function  $f(\alpha)$  in eq. (70). The function is convex; it also satisfies  $f(0) = f(\tau) = \Psi_{jj}(\tau)$  and  $f'(0)f'(\tau) < 0$ .

We also know from Lemma 5 that  $\Sigma_{ii}\Sigma_{ii}^{-1} > 1$  whenever  $\sum_{j \neq i} |\Sigma_{ij}| > 0$ , and that in this case, the above inequalities are strict. A useful picture of this situation is shown in the left panel of Fig. 3. Consider any component  $\Psi_{ii}(\alpha)$  of the estimated variances that is *not* constant on the unit interval. (As shown previously, there must be at least one such component.) In this case, there are only two possibilities: either  $\Psi_{ii}(\alpha)$  is strictly increasing, or it is not. If the former is always true, then it follows that  $R_{\alpha_1}(p||q) < R_{\alpha_2}(p||q)$  whenever  $0 < \alpha_1 < \alpha_2 < 1$ . We shall prove *by contradiction* that this is indeed the case.

Assume that there are one or more diagonal components,  $\Psi_{ii}(\alpha)$ , that are neither constant nor strictly increasing over the unit interval; also, let  $\mathcal{I}$  denote the set that contains the indices of these components. Since each such component must have at least one stationary point, we can also write:

$$\mathcal{I} = \left\{ i \mid \Sigma_{ii}\Sigma_{ii}^{-1} > 1 \text{ and } \Psi'_{ii}(\alpha) = 0 \text{ for some } \alpha \in (0, 1) \right\}. \quad (63)$$

For each such component, there must also exist some *minimal* point  $\tau_i \in (0, 1)$  where its derivative vanishes. We define  $\tau = \min_{i \in \mathcal{I}} \tau_i$  and  $j = \operatorname{argmin}_{i \in \mathcal{I}} \tau_i$ , so that by definition

$$\Psi'_{ii}(\tau) \geq 0 \quad \text{for all } i, \quad (64)$$

$$\Psi'_{jj}(\tau) = 0. \quad (65)$$

To obtain the desired contradiction, we shall prove that there exists no point  $\tau$  (as imagined in Fig. 3) with these properties.

We start by rewriting the fixed-point equation in Proposition 6 and eq. (44) in a slightly different form,

$$\Psi_{ii}(\alpha) = [\Phi(\alpha)^{-1}]_{ii} = [\alpha\Sigma^{-1} + (1-\alpha)\Psi^{-1}(\alpha)]_{ii}^{-1}, \quad (66)$$

where on the right side we have explicitly indicated all sources of dependence on  $\alpha$ . Next, we differentiate both sides of eq. (66) for  $i=j$  and at  $\alpha=\tau$ :

$$\Psi'_{jj}(\tau) = \mathbf{e}_j^\top \left\{ \frac{d}{d\alpha} \left[ \tau \boldsymbol{\Sigma}^{-1} + (1-\tau) \boldsymbol{\Psi}^{-1}(\alpha) \right]^{-1} + \frac{d}{d\alpha} \left[ \alpha \boldsymbol{\Sigma}^{-1} + (1-\alpha) \boldsymbol{\Psi}^{-1}(\tau) \right]^{-1} \right\} \Big|_{\alpha=\tau} \mathbf{e}_j, \quad (67)$$

where on the right side we have separated out the different sources of dependence on  $\alpha$ . Evaluating the first term on the right side, we find that

$$\mathbf{e}_j^\top \left\{ \frac{d}{d\alpha} \left[ \tau \boldsymbol{\Sigma}^{-1} + (1-\tau) \boldsymbol{\Psi}^{-1}(\alpha) \right]^{-1} \right\} \Big|_{\alpha=\tau} \mathbf{e}_j = \mathbf{e}_j^\top \boldsymbol{\Phi}^{-1}(\tau) \boldsymbol{\Psi}^{-1}(\tau) \boldsymbol{\Psi}'(\tau) \boldsymbol{\Psi}^{-1}(\tau) \boldsymbol{\Phi}^{-1}(\tau) \mathbf{e}_j, \quad (68)$$

and now we recognize that this term is nonnegative: of the matrices on the right side of eq. (68), note that  $\boldsymbol{\Phi}(\alpha)$  and  $\boldsymbol{\Psi}(\alpha)$  are positive definite for all  $\alpha \in (0, 1)$ , and  $\boldsymbol{\Psi}'(\tau)$  is positive semidefinite by virtue of the defining properties of  $\tau$  in eqs. (64–65). Dropping this first term from the right side of eq. (67), we obtain the inequality

$$0 \geq \frac{d}{d\alpha} \left\{ \mathbf{e}_j^\top \left[ \alpha \boldsymbol{\Sigma}^{-1} + (1-\alpha) \boldsymbol{\Psi}^{-1}(\tau) \right]^{-1} \mathbf{e}_j \right\} \Big|_{\alpha=\tau}, \quad (69)$$

where we have exploited that the left side of eq. (67) vanishes by the second defining property of  $\tau$  in eq. (65).

Next we show that eq. (69) contains a contradiction. To do so, we examine the properties of the function defined on  $[0, 1]$  by

$$f(\alpha) = \mathbf{e}_j^\top \left[ \alpha \boldsymbol{\Sigma}^{-1} + (1-\alpha) \boldsymbol{\Psi}^{-1}(\tau) \right]^{-1} \mathbf{e}_j. \quad (70)$$

This is, of course, exactly the function that is differentiated in eq. (69), whose right side is equal to  $f'(\tau)$ . This function possesses several key properties. First, we note that

$$f(0) = \mathbf{e}_j^\top \left[ \boldsymbol{\Psi}^{-1}(\tau) \right]^{-1} \mathbf{e}_j = \Psi_{jj}(\tau) = \mathbf{e}_j^\top \left[ \tau \boldsymbol{\Sigma}^{-1} + (1-\tau) \boldsymbol{\Psi}^{-1}(\tau) \right]^{-1} \mathbf{e}_j = f(\tau). \quad (71)$$

Second, we note that  $f$  is convex on the unit interval; here, we are observing the convexity of the matrix inverse (Nordström, 2011). Third, we note that

$$f'(0) = -\mathbf{e}_j^\top \boldsymbol{\Psi}(\tau) \left[ \boldsymbol{\Sigma}^{-1} - \boldsymbol{\Psi}^{-1}(\tau) \right] \boldsymbol{\Psi}(\tau) \mathbf{e}_j = \Psi_{jj}(\tau) \left[ \boldsymbol{\Sigma}^{-1} - \boldsymbol{\Psi}^{-1}(\tau) \right]_{jj} < 0. \quad (72)$$

Finally, we note that the sign of  $f'(\tau)$  must *oppose* the sign of  $f'(0)$ ; as shown in the right panel of Fig. (3), this follows from the preceding properties that  $f(0) = f(\tau)$ , that the function  $f$  is convex on  $[0, 1]$ , and that  $f'(0) < 0$ . Thus we have shown that  $f'(\tau) > 0$ . But this is directly in contradiction with eq. (69), which states that  $0 \geq f'(\tau)$ , and we are forced to conclude that no such point  $\tau$  exists. This completes the proof.  $\blacksquare$

Combining the results from Sections 5.1–5.3, we obtain a proof of Theorem 3.

#### 5.4 Entropy-matching solution for FG-VI

We observed in Section 4.1 that the reverse KL divergence yields a precision-matching solution for FG-VI and that the forward KL divergence yields a variance-matching solution. The ordering of divergences in Theorem 3 has another interesting consequence: it implies that some Rényi divergence yields a unique *entropy-matching* solution for FG-VI.




Figure 4: When  $p$  is Gaussian over  $\mathbb{R}^n$ , there always exists a unique  $\alpha \in (0, 1)$  such that the factorized approximation  $q$  minimizing  $R_\alpha(p||q)$  matches the entropy of  $p$ . The plots shows, however, that the entropy-matching value of  $\alpha$  depends on the dimension and covariance structure of  $p$ . Above we vary the dimension  $n$  and constant correlation  $\varepsilon$ .

**Corollary 10** Let  $q_\alpha = \operatorname{argmin}_{q \in \mathcal{Q}} R_\alpha(p||q)$ . There exists a unique value  $\alpha \in (0, 1)$  such that  $q_\alpha$  matches the entropy of  $p$ , or equivalently, that  $|\Psi(\alpha)| = |\Sigma|$ .

**Proof** Let  $\Psi(\alpha)$  denote the covariance of  $q_\alpha$  for  $\alpha \in (0, 1)$ . Since  $\Psi_{ii}(\alpha)$  is continuous with respect to  $\alpha$ , so is  $\log |\Psi(\alpha)| = \sum_i \log \Psi_{ii}(\alpha)$ . Moreover, from the ordering of Rényi divergences, we see that  $\log |\Psi(\alpha)|$  is not only continuous, but strictly increasing over the unit interval  $\alpha \in (0, 1)$ . In addition, for all components  $i$  along the diagonal, we have

$$1/\Sigma_{ii}^{-1} = \Psi_{ii}(0) \leq \Psi_{ii}(\alpha) \leq \Psi_{ii}(1) = \Sigma_{ii}, \quad (73)$$

with  $q_\alpha$  interpolating smoothly between the precision-matching and the variance-matching approximations of FG-VI. From the impossibility result in Theorem 1, the precision-matching approximation (at  $\alpha=0$ ) underestimates the entropy of  $p$ , while the variance-matching approximation (at  $\alpha=1$ ) overestimates it. From continuity and strict monotonicity, it follows that there exists a unique  $\alpha \in (0, 1)$  such that  $|\Psi(\alpha)| = |\Sigma|$ , or equivalently, such that  $q_\alpha$  matches the entropy of  $p$ . ■

The proof of Corollary 10 demonstrates the existence an entropy-matching solution, but it is not a constructive proof. Unfortunately, there does not exist a single value of  $\alpha$  for which  $|\Psi(\alpha)| = |\Sigma|$ , even when  $p$  is Gaussian. We show this numerically for the case where the target covariance  $\Sigma$  has unit diagonal elements and constant off-diagonal elements  $\varepsilon \in (0, 1)$ . As seen in Figure 4, the entropy-matching value of  $\alpha$  (obtained by a grid search) changes as we vary the amount of correlation and the dimension of the problem.

Model	$n$	Details	Variances	Entropy
Gaussian	10	Full-rank covariance matrix.	✓	✓
Rosenbrock	2	High curvature.	✓	✓
Eight Schools	10	Small hierarchical linear model.	✗	✓
German Credit	25	Logistic regression.	✓	✓
Radon Effect	91	Large hierarchical linear model.	✗	✓
Stochastic Volatility	103	Stochastic volatility model.	✗	✓

Table 3: Target distributions  $p$  with dimension  $n$ . The two right-most columns indicate whether the divergences are ordered according to the marginal variances (equivalently to the marginal precisions) and the entropy (equivalently the generalized variance).

## 6 Numerical experiments

Does the ordering of divergences hold when FG-VI is applied to non-Gaussian targets? We study this question empirically on a range of models; in most of these models, the target  $p$  is only known up to a normalizing constant.

When  $p$  is not Gaussian, some divergences in Table 1 are harder to minimize numerically than others. We use the following approaches. We minimize  $\text{KL}(q||p)$  by (equivalently) maximizing the evidence lower bound (ELBO), and we estimate the ELBO via Monte Carlo with draws from  $q$ , a procedure at the core of modern implementations of “black-box” VI (Kucukelbir et al., 2017). We use a similar procedure to optimize the  $\alpha$ -divergence; however, this procedure is more fraught<sup>3</sup> as Monte Carlo estimators of the  $\alpha$ -divergence and its gradients suffer from large variance, especially in large dimensions and as  $\alpha$  approaches 1 (Geffner and Domke, 2021). We minimize  $\text{S}(q||p)$  by adapting a recent *batch-and-match* (BaM) method inspired by proximal point algorithms (Cai et al., 2024); BaM was originally developed for Gaussian variational families with dense covariance matrices, but here we experiment with BaM updates for FG-VI (derived in Appendix A). We minimize  $\text{KL}(p||q)$  by estimating the means and marginal variances of  $p$  from long runs of Markov chain Monte Carlo; the factorized approximation minimizing  $\text{KL}(q||p)$  is the one that matches these moments (MacKay, 2003). We do not empirically evaluate the forward score-based divergence for FG-VI as we do not have a reliable method to minimize  $\text{S}(p||q)$ .

We experiment with the above methods on a diverse collection of target distributions from the `inference gym`, a curated library of models for the study of inference algorithms (Sountsov et al., 2020). We describe these target distributions briefly in Table 3 and provide more details in Appendix B. We implement each model in Python and use JAX to calculate derivatives (Bradbury et al., 2018). We optimize the  $\alpha$ -divergences for  $\alpha=0.1$  and  $\alpha=0.5$ , but only on low-dimensional targets ( $D < 10$ ) where the optimizers are numerically stable. We run each VI algorithm for 1,000 iterations, and at each iteration, we use  $B = 10,000$  draws from  $q$  to estimate the divergence and its gradient. This large number of draws is

3. For more discussion on implementing VI with  $\alpha$ -divergences, see e.g., Hernandez-Lobato et al. (2016); Li and Turner (2016); Dieng et al. (2017); Daudel et al. (2021, 2023).




Figure 5: Variance when targeting the posterior distribution of the Eight Schools model.

overkill for many problems, but it helps stabilize the optimizations of  $\alpha$ -divergences. In addition, we parallelize many calculations on a GPU, keeping run times short. Code to run the experiments can be found at <https://github.com/charlesm93/VI-ordering>.

The ordering of divergences that we proved for Gaussian targets is not guaranteed to hold for non-Gaussian targets, and empirically we observe that it is more likely to fail in high-dimensional problems; see Table 3. This is illustrated in Fig. 5 for the Eight Schools models, where some marginal variances are ordered in the same way as predicted for Gaussian targets, but most are not. The true marginal variances in this model were estimated by long runs of Markov chain Monte Carlo.

When  $p$  is not Gaussian, an alternative ordering of divergences for FG-VI is suggested by the estimates they yield of the joint entropy. As shown in the final column of Table 3 and in Fig. 6, the ordering in Theorem 3 *does* correspond—across all of the non-Gaussian targets in our experiments—to the ordering of entropies estimated by FG-VI. In these experiments, however, we are not able to compare the entropies estimated by FG-VI to the actual entropies. To compute the entropy of non-Gaussian distributions, it is necessary to estimate their normalizing constants; this can be done by invoking a Gaussian-like approximation, as in bridge sampling (Meng and Schilling, 2002; Gronau et al., 2020). However, it seems inadequate to compare a theory developed for Gaussian targets to an empirical benchmark that relies on a Gaussian approximation.

## 7 Discussion

Divergences over probability spaces are ubiquitous in the probability, statistics, and machine learning literature, but it remains challenging to reason about them. This challenge is salient in VI, where the divergence does not serve merely as an object for theoretical study but also bears directly on the algorithmic implementation of the method. Our work reveals how the choice of divergence can align (or misalign) with intelligible inferential goals, and it is in line with a rich and recent body of work on assessing the quality of VI (e.g, Yao et al., 2018; Wang and Blei, 2018; Huggins et al., 2020; Dhaka et al., 2021; Biswas and Mackey, 2024). A distinguishing feature of our paper is that it analyzes multiple and competing inferential goals—goals which cannot be simultaneously met and which are each best served by a different divergence.




Figure 6: *Ordering of entropy across targets. The zero on each vertical axis corresponds to the average of the entropies estimated by the approximations in each panel.*

Our results can inform the choice of divergence to minimize in eq. (1) in tandem with the use of a factorized approximation. This choice will in turn depend on the problem at hand. We discuss some examples:

- Bayesian statistics.** Here, uncertainty is primarily quantified by the marginal variances  $\Sigma_{ii}$ , and these are matched by minimizing  $KL(p||q)$ . In general, however, this divergence is not straightforward to minimize, though strategies such as expectation-propagation can in some limits attain an optimal solution (Vehtari et al., 2020). It is much more straightforward to minimize divergences such as  $KL(q||p)$  and  $S(q||p)$ , but for FG-VI we have seen how marginal variances can be significantly underestimated by factorized approximations that minimize these divergences. This provides further motivation for: (i) corrective procedures (Giordano et al., 2018) to adjust the estimate of marginal variances from F-VI; (ii) VI methods with richer (non-factorized) families (e.g Opper and Archambeau, 2009; Dhaka et al., 2021; Zhang et al., 2022); (iii) altogether alternative methods such as Markov chain Monte Carlo (MCMC; Robert and Casella, 2004), when allowed by a computational budget.
- Pre-conditioning of Markov chain Monte Carlo.** Several MCMC algorithms benefit from transforming the latent variables over which the Markov chain is constructed. This technique is known as pre-conditioning. In Hamiltonian Monte Carlo (HMC; Neal, 2012; Betancourt, 2018) the mass matrix serves as a pre-conditioner, and one typically restricts it to be diagonal for computational reasons. Neal (2012) recommends setting the diagonal elements of the (inverted) mass matrix to early estimates of the marginal variances, an idea implemented in popular adaptive HMC algorithms (e.g Hoffman and Gelman, 2014). Recently, it was proposed that this approach might not be optimal (Hird and Livingstone, 2023), and that a better choice for a diagonal mass matrix might be based on estimates of the marginal precisions (Tran and Kleppe, 2024). This suggests a novel application of F-VI, one where it

is used to tune HMC’s mass matrix by minimizing  $\text{KL}(q||p)$ . Investigation of this method is left as future work.

- **Information theory.** Here, the primary measure of uncertainty is the entropy. In certain limiting cases, we can accurately estimate the entropy by minimizing  $\text{KL}(q||p)$  (Parisi, 1988; Margossian and Saul, 2023); in general, however, the entropy is not correctly estimated by the minimizer of  $\text{KL}(q||p)$ . This can be shown analytically when  $p$  and  $q$  are Gaussian. One possible strategy in this setting is to move further along the ordering of divergences and minimize  $R_\alpha(p||q)$  for  $\alpha \in (0, 1)$  (but not all the way to  $\alpha = 1$ , since one would overestimate the entropy by minimizing  $\text{KL}(p||q)$ ). We have shown that, while there always exists an  $\alpha$  to match the entropy, this value can change with  $p$ , and so it remains an open problem to find an actionable divergence for entropy estimation.

VI is also deployed in applications where the primary goal is not to quantify uncertainty. A prominent example is latent variable models, where VI is used to approximate a marginal likelihood. In this context, the KL and Rényi divergences have been ordered based on the (lower or upper) bound they provide on the marginal likelihood (Li and Turner, 2016; Dieng et al., 2017). (Interestingly, this ordering agrees with the one provided in this paper, even though it is based on a different criterion.) This type of ordering cannot be extended to score-based divergences, however, as they do not provide bounds on the marginal likelihood.

We have seen that the results of VI can be improved, for certain inferential goals, by choosing a particular divergence. A complementary and more common approach is to use a richer family  $\mathcal{Q}$  of variational approximations (e.g, Hoffman and Blei, 2015; Johnson et al., 2016; Dhaka et al., 2021). The deficits of FG-VI may be addressed, for instance, by Gaussian variational approximations with full covariance matrices (Opper and Archambeau, 2009; Cai et al., 2024). In high dimensions, Dhaka et al. (2021) argue that the results of VI can be improved more easily by optimizing  $\text{KL}(q||p)$  over a richer variational family than by optimizing certain alternative divergences. Nevertheless, even as the variational family becomes richer, it is not likely to contain the target distribution  $p$ , and therefore the approximating distribution  $q$  must in some way be compromised. For example, if  $\mathcal{Q}$  is a family of symmetric distributions and the target  $p$  is asymmetric, then an approximation may only be able to match the mode *or* the mean of the target. It stands to reason that further impossibility theorems, in the spirit of Theorem 1, can be derived to elucidate these trade-offs.

## Acknowledgments

We thank David Blei, Diana Cai, Robert Gower, Chirag Modi, and Yuling Yao for helpful discussions. We are also grateful to anonymous reviewers from the conference on Uncertainty in Artificial Intelligence for their comments on a previous paper (Margossian and Saul, 2023), which provided some of the motivation for this manuscript.

## Appendix A. Batch and Match algorithm for FG-VI

*Batch and Match* (BaM) (Cai et al., 2024, BaM) is an iterative algorithm for VI that attempts to estimate and minimize the reverse score-based divergence,  $S(q||p)$ . At each iteration, BaM minimizes a regularized objective function

$$\mathcal{L}^{\text{BaM}}(q) = \hat{\mathcal{D}}_{q_t}(q||p) + \frac{2}{\lambda_t} \text{KL}(q||q_t), \quad (74)$$

where

- $q_t \in \mathcal{Q}$  is a current approximation,
- $\hat{\mathcal{D}}_{q_t}(q||p)$  is a Monte Carlo estimator of the score-based divergence

$$\hat{\mathcal{D}}_{q_t}(q||p) = \frac{1}{B} \sum_{b=1}^B \|\nabla \log q(\mathbf{z}^{(b)}) - \nabla \log p(\mathbf{z}^{(b)})\|_{\Psi}^2, \quad (75)$$

using draws  $\mathbf{z}^{(b)} \sim q_t$ .

- $\lambda_t > 0$  is the learning rate, or step size, at the  $t^{\text{th}}$  iteration.

Cai et al. (2024, Appendix C) derive the BaM updates which minimize eq. (74) when  $\mathcal{Q}$  is the family of Gaussian distributions with *dense* covariance matrices. In this appendix, we derive the analogous updates to minimize eq. (74) when  $\mathcal{Q}$  is the family of Gaussian distributions with *diagonal* covariance matrices. These were the updates used for the experiments in Section 6 of the paper.

We follow closely the derivation in Cai et al. (2024), noting only the essential differences for the case of FG-VI. We use  $\mathbf{g}^{(b)} = \nabla \log p(\mathbf{z}^{(b)})$  as shorthand for the score at the  $b^{\text{th}}$  sample. As before, the following averages need to be computed at each iteration:

$$\bar{\mathbf{z}} = \frac{1}{B} \sum_{b=1}^B \mathbf{z}^{(b)}, \quad \bar{\mathbf{g}} = \frac{1}{B} \sum_{b=1}^B \mathbf{g}^{(b)}. \quad (76)$$

For FG-VI, we instead compute *diagonal* matrices  $\mathbf{C}$  and  $\mathbf{\Gamma}$  with nonnegative elements

$$C_{ii} = \frac{1}{B} \sum_{b=1}^B (z_i^{(b)} - \bar{z}_i)^2, \quad \Gamma_{ii} = \frac{1}{B} \sum_{b=1}^B (g_i^{(b)} - \bar{g}_i)^2. \quad (77)$$

Let  $q_{t+1}$  denote the minimizer of eq. (74), with mean  $\boldsymbol{\nu}_{t+1}$  and diagonal covariance matrix  $\Psi^{t+1}$ . The update for the mean takes the same form as in the original formulation of BaM:

$$\boldsymbol{\nu}^{t+1} = \frac{\lambda_1}{1 + \lambda_t} (\bar{\mathbf{z}} + \Psi^{t+1} \bar{\mathbf{g}}) + \frac{1}{1 + \lambda_t} \boldsymbol{\nu}^t. \quad (78)$$

To obtain the covariance update, we substitute this result into eq. (74) and differentiate with respect to the *diagonal* elements of  $\Psi^{t+1}$ . For FG-VI, we obtain a simple quadratic equation for each updated variance:

$$\left( \Gamma_{ii} + \frac{1}{1 + \lambda_t} \bar{g}_i^2 \right) (\Psi_{ii}^{t+1})^2 + \frac{1}{\lambda_t} \Psi_{ii}^{t+1} - \left( C_{ii} + \frac{1}{\lambda_t} \Psi_{ii}^t + \frac{(\nu_i^t - \bar{z}_i)^2}{1 + \lambda_t} \right) = 0. \quad (79)$$

Eq. (79) always admits a positive root, and for FG-VI, this positive root is the BaM update for the  $i^{\text{th}}$  diagonal element of  $\Psi^{t+1}$ .

At a high level, BaM works the same way with diagonal covariance matrices as it does with full covariance matrices. Each iteration involves a *batch* step which draws the  $B$  samples from  $q_t$  and a *match* step that performs the calculations in eqs. (76-79). We find a constant learning rate  $\lambda_t = 1$  to work well for the experiments in Section 6.

## Appendix B. Models from the inference gym

In this appendix, we provide details about the models and data sets from the **inference gym** used in Section 6.

**Rosenbrock Distribution.** ( $n=2$ ) A transformation of a normal distribution, with non-trivial correlations (Rosenbrock, 1960). The contour plots of the density function have the shape of a crescent moon. The joint distribution is given by:

$$p(z_1) = \text{normal}(0, 10) \tag{80}$$

$$p(z_2|z_1) = \text{normal}(0.03(z_1^2 - 100), 1). \tag{81}$$

We use FG-VI to approximate  $p(z_1, z_2)$ .

**Eight Schools.** ( $n=10$ ) A Bayesian hierarchical model of the effects of a test preparation program across 8 schools (Rubin, 1981):

$$p(\mu) = \text{normal}(5, 3^2) \tag{82}$$

$$p(\tau) = \text{normal}^+(0, 10) \tag{83}$$

$$p(\theta_i|\mu, \sigma) = \text{normal}(\mu, \tau^2) \tag{84}$$

$$p(y_i|\theta_i) = \text{normal}(\theta_i, \sigma_i^2), \tag{85}$$

where  $\text{normal}^+$  is a normal distribution truncated at 0. The observations are  $\mathbf{x} = (y_{1:8}, \sigma_{1:8})$  and the latent variables are  $\mathbf{z} = (\mu, \tau, \theta_{1:8})$ . We use VI to approximate the posterior distribution  $p(\mathbf{z}|\mathbf{x})$ .

**German Credit.** ( $n=25$ ) A logistic regression applied to a credit data set (Dua and Graff, 2017), with covariates  $\mathbf{x} = x_{1:24}$ , observations  $\mathbf{y} = y_{1:24}$ , coefficients  $\mathbf{z} = z_{1:24}$  and an intercept  $z_0$ :

$$p(\mathbf{z}) = \text{normal}(0, \mathbf{I}_n), \tag{86}$$

$$p(y_i|\mathbf{z}, \mathbf{x}) = \text{Bernoulli}\left(\frac{1}{e^{-\mathbf{z}^T \mathbf{x}_i - z_0}}\right) \tag{87}$$

We use FG-VI to approximate  $p(\mathbf{z}|y_i, \mathbf{x})$ .

**Radon Effect.** ( $n=91$ ) A hierarchical model to measure the Radon level in households (Gelman and Hill, 2007). Here, we restrict ourselves to data from Minnesota. For each household, we have three covariates,  $\mathbf{x}_i = x_{i,1:3}$ , with corresponding coefficients  $\mathbf{w} = w_{1:3}$ ,

and a county level covariate,  $\theta_{j[i]}$ , where  $j[i]$  denotes the county of the  $i^{\text{th}}$  household. The model is:

$$p(\mu) = \text{normal}(0, 1) \tag{88}$$

$$p(\tau) = \text{Uniform}(0, 100) \tag{89}$$

$$p(\theta_j | \mu, \tau) = \text{normal}(\mu, \tau^2) \tag{90}$$

$$p(w_k) = \text{normal}(0, 1) \tag{91}$$

$$p(\sigma) = \text{Uniform}(0, 100) \tag{92}$$

$$p(\log y_i) = \text{normal}(\mathbf{w}^T \mathbf{x}_i + \theta_{j[i]}, \sigma^2). \tag{93}$$

We use FG-VI to approximate  $p(\mu, \tau, \theta, \mathbf{w}, \sigma | \mathbf{x}, y)$ .

**Stochastic volatility.** (n=103) A time series model with 100 observations. The original model by [Kim et al. \(1998\)](#) is given by:

$$p(\sigma) = \text{Cauchy}^+(0, 2) \tag{94}$$

$$p(\mu) = \text{exponential}(1) \tag{95}$$

$$p((\phi + 1)/2) = \text{Beta}(20, 1.5) \tag{96}$$

$$p(z_i) = \text{normal}(0, 1) \tag{97}$$

$$h_1 = \mu + \sigma z_1 / \sqrt{1 - \phi^2} \tag{98}$$

$$h_{i>1} = \mu + \sigma z_i + \phi(h_{i-1} - \mu) \tag{99}$$

$$p(y_i | h_i) = \text{normal}(0, \exp(h_i/2)). \tag{100}$$

The original model was developed for a time series of 3000 observations, but we only work with the first 100 observations. For these observations, we use FG-VI to model  $p(\sigma, \mu, \phi, \mathbf{z} | \mathbf{y})$ .

## References

- José M. Bernardo and Adrian F. M. Smith. *Bayesian Theory*. Wiley, 2000.
- Michael Betancourt. A conceptual introduction to Hamiltonian Monte Carlo. *arXiv:1701.02434v1*, 2018.
- Christopher M. Bishop. *Pattern Recognition and Machine Learning*. Springer, 2006.
- Christopher M. Bishop, David Spiegelhalter, and John Winn. Vibes: A variational inference engine for bayesian networks. *Advances Neural Information Processing Systems 15*, pages 793–800, 2002.
- Niloy Biswas and Lester Mackey. Bounding Wasserstein distance with couplings. *Journal of the American Statistical Association*, 2024. DOI: 10.1080/01621459.2023.2287773.
- David M. Blei. Probabilistic topic models. *Communications of the ACM*, 55:77–84, 2012.

- David M. Blei, Alp Kucukelbir, and Jon D. McAuliffe. Variational inference: A review for statisticians. *Journal of the American Statistical Association*, 112:859–877, 2017. doi: 10.1080/01621459.2017.1285773. URL <https://arxiv.org/abs/1601.00670>.
- James Bradbury, Roy Frostig, Peter Hawkins, Matthew James Johnson, Chris Leary, Dougal Maclaurin, George Necula, Adam Paszke, Jake VanderPlas, Skye Wanderman-Milne, and Qiao Zhang. JAX: composable transformations of Python+NumPy programs, 2018. URL <http://github.com/google/jax>.
- Jacob Burbea. The convexity with respect to Gaussian distributions of divergences of order  $\alpha$ . *Utilitas Mathematica*, 26:171–192, 1984.
- Diana Cai, Chirag Modi, Loucas Pillaud-Vivien, Charles C. Margossian, Robert M. Gower, David M. Blei, and Lawrence K. Saul. Batch and match: black-box variational inference with a score-based divergence. *Proceedings of the 41st International Conference on Machine Learning*, page (to appear), 2024.
- Andrzej Cichocki and Shun-ichi Amari. Families of alpha- beta- and gamma-divergences: Flexible and robust measures of similarities. *Entropy*, 12:1532–1568, 2010.
- Thomas A. Courtade. Links between the logarithmic Sobolev inequality and the convolution inequalities for entropy and Fisher information. *arXiv:1608.05431*, 2016.
- Kamélia Daudel, Randal Douc, and Francois Roueff. Infinite-dimensional gradient-based descent for alpha-divergence minimisation. *The Annals of Statistics*, 49:2250–2270, 2021.
- Kamélia Daudel, Joe Benton, and Arnaud Doucet. Alpha-divergence variational inference meets importance weighted auto-encoders: Methodology and asymptotics. *Journal of Machine Learning Research*, 24:1–83, 2023.
- Akash Kumar Dhaka, Alejandro Catalina, Manushi Welandawe, Michael Riis Andersen, Jonathan H Huggins, and Aki Vehtari. Challenges and opportunities in high-dimensional variational inference. In *Advances in Neural Information Processing Systems 34*, pages 7787–7798, 2021.
- Adji B. Dieng, Dustin Tran, R Ranganath, J Paisly, and David M. Blei. Variational inference via  $\chi$  upper bound minimization. In *Advances in Neural Information Processing Systems 30*, pages 2732–2741, 2017.
- Dheera Dua and C. Graff. UCL machine learning repository, 2017. URL <http://archive.ics.ucl.edu/ml>.
- Tomas Geffner and Justin Domke. On the difficulty of unbiased alpha divergence minimization. In *Proceedings of the 38th International Conference on Machine Learning*, pages 3650–3659, 2021.
- Andrew Gelman and Jennifer Hill. *Data Analysis Using Regression and Multilevel-Hierarchical Models*. Cambridge University Press, 2007.

- Andrew Gelman, John B. Carlin, Hal S. Stern, David B. Dunson, Ark Vehtari, and Donald B. Rubin. *Bayesian Data Analysis*. Chapman & Hall/CRC Texts in Statistical Science, 2013.
- Alison Gibbs and Francis Edward Su. On choosing and bounding probability metrics. *International Statistical Review*, 70, 2002.
- Manuel Gil, Fady Alajaji, and Tamas Linder. Rényi divergence measures for commonly used univariate continuous distributions. *Information Sciences*, 249:124–131, 2013.
- Rian Giordano, Tamara Broderick, and Michael I. Jordan. Covariances, robustness, and variational bayes. *Journal of Machine Learning Research*, 19:1–49, 2018.
- Ryan Giordano, Martin Ingram, and Tamara Broderick. Black box variational inference with a deterministic objective: Faster, more accurate, and even more black box. *Journal of Machine Learning Research*, 25:1–39, 2023.
- Quantin F. Gronau, Henrik Singmann, and Eric-Jan Wagenmakers. bridgesampling: An R package for estimating normalizing constants. *Journal of Statistical Software*, 92, 2020.
- Jose Hernandez-Lobato, Yingzhen Li, Mark Rowland, Thang Bui, Daniel Hernández-Lobato, and Richard Turner. Black-box alpha divergence minimization. In *Proceedings of the 33rd International Conference on Machine Learning*, pages 1511–1520, 2016.
- Max Hird and Samuel Livingstone. Quantifying the effectiveness of linear preconditioning in markov chain monte carlo. *arXiv:2312.04898*, 2023.
- Tomáš Hobza, Domingo Morales, and Leandro Pardo. Rényi statistics for testing equality of autocorrelation coefficients. *Statistical Methodology*, 6:424–436, 2009.
- Matthew D. Hoffman and David M. Blei. Stochastic structured variational inference. In *Proceedings of the 18th International Conference on Artificial Intelligence and Statistics*, pages 361–369, 2015.
- Matthew D. Hoffman and Andrew Gelman. The no-U-turn sampler: Adaptively setting path lengths in Hamiltonian Monte Carlo. *Journal of Machine Learning Research*, 15:1593–1623, 2014.
- Roger A. Horn and Charles R. Johnson. *Matrix analysis*. Cambridge University Press, 2012.
- Jonathan H. Huggins, Mikołaj Kasprzak, Trevor Campbell, and Tamara Broderick. Validated variational inference via practical posterior error bounds. In *Proceedings of the 23rd International Conference on Artificial Intelligence and Statistics*, pages 1792–1802, 2020.
- Matthew J. Johnson, David K. Duvenaud, Alex Wiltschko, Ryan P. Adams, and Sandeep R. Datta. Composing graphical models with neural networks for structured representations and fast inference. In *Advances in neural information processing systems 29*, pages 2946–2954, 2016.

- Michael I. Jordan, Zoubin Ghahramani, Tommi S. Jaakkola, and Lawrence K. Saul. An introduction to variational methods for graphical models. *Machine Learning*, 37:183–233, 1999.
- Sangjoon Kim, Neil Shepard, and Siddhartha Chib. Stochastic volatility: likelihood inference and comparison with arch models. *The Review of Economic Studies*, 65:361–393, 1998.
- Diederik P. Kingma and Max Welling. Auto-encoding variational Bayes. *International Conference on Learning Representations*, 2014.
- Alp Kucukelbir, Dustin Tran, Rajesh Ranganath, Andrew Gelman, and David Blei. Automatic differentiation variational inference. *Journal of Machine Learning Research*, 18:1–45, 2017.
- Yingzhen Li and Richard E. Turner. Rényi divergence variational inference. In *Advances in Neural Information Processing Systems 29*, pages 1073–1081, 2016.
- Friedrich Liese and Igor Vajda. *Convex Statistical Distances*. Teubner, Leipzig, 1987.
- David J.C. MacKay. *Information theory, inference, and learning algorithms*. 2003.
- Charles C. Margossian and David M. Blei. Amortized variational inference: When and why? In *Proceedings of the 40th Conference on Uncertainty in Artificial Intelligence*, page (to appear), 2024.
- Charles C. Margossian and Lawrence K. Saul. The shrinkage-delinkage trade-off: An analysis of factorized gaussian approximations for variational inference. In *Proceedings of the 39th Conference on Uncertainty in Artificial Intelligence*, pages 1358–1367, 2023.
- Xiao-Li Meng and Stephen Schilling. Warp bridge sampling. *Journal of Computational and Graphical Statistics*, 11:552–586, 2002. doi: doi:10.1198/106186002457.
- Tom Minka. Divergence measures and message passing. *Technical Report MSR-TR-2005-173*, 2005.
- Radford M. Neal. MCMC using Hamiltonian dynamics. In *Handbook of Markov Chain Monte Carlo*. CRC Press, 2012.
- Kenneth Nordström. Convexity of the inverse and Moore-Penrose inverse. *Linear Algebra and Its Applications*, 434:1489–1512, 2011.
- Manfred Opper and Cédric Archambeau. The variational Gaussian approximation revisited. *Neural Computation*, 21(3):786–792, 2009.
- Georgio Parisi. *Statistical Field Theory*. Addison-Wesley, 1988.
- Christian P Robert and George Casella. *Monte Carlo Statistical Methods*. Springer, 2004.
- Howard H. Rosenbrock. An automatic method for finding the greatest or least value of a function. *Computer Journal*, 3:175–184, 1960.

- Donald B. Rubin. Estimation in parallelized randomized experiments. *Journal of Educational Statistics*, 6:377–400, 1981.
- Pavel Sountsov, Alexey Radul, and contributors. Inference gym, 2020. URL [https://pypi.org/project/inference\\_gym](https://pypi.org/project/inference_gym).
- Jakub M. Tomczak. *Deep generative modeling*. Springer, 2022.
- Jimmy H. Tran and Tore S. Kleppe. Tuning diagonal scale matrices for hmc. *arxiv2403.07495*, 2024.
- Richard E. Turner and Maneesh Sahani. Two problems with variational expectation maximisation for time-series models. In David Barber, A. Taylan Cemgil, and Silvia Chiappa, editors, *Bayesian Time series models*, chapter 5, pages 109–130. Cambridge University Press, 2011.
- Aki Vehtari, Andrew Gelman, Tuomas Sivula, Pasi Jylanki, Dustin Tran, Swupnil Sahai, Paul Blomstedt, John P. Cunningham, Schiminovich, and Christian P. Robert. Expectation propagation as a way of life: A framework for bayesian inference on partitioned data. *Journal of Machine Learning*, 21:1–53, 2020.
- Aki Vehtari, Daniel Simpson, Andrew Gelman, Yuling Yao, and Jonah Gabry. Pareto smoothed importance sampling. *Journal of Machine Learning Research*, 2024. To appear.
- Martin J. Wainwright and Michael I. Jordan. *Foundations and Trends in Machine Learning*, 1(1–2):1–305, 2008.
- Bo Wang and Donald M. Titterton. Inadequacy of interval estimates corresponding to variational bayesian approximations. *Proceedings of the Tenth International Workshop on Artificial Intelligence and Statistics*, R5:373 – 380, 2005.
- Yixin Wang and David M. Blei. Frequentist consistency of variational bayes. *Journal of the American Statistical Association*, 114:1147–1161, 2018.
- Yuling Yao, Aki Vehtari, Daniel Simpson, and Andrew Gelman. Yes, but did it work?: Evaluating variational inference. In *Proceedings of the 35th International Conference on Machine Learning*, pages 5577–5586, 2018.
- Lu Zhang, Bob Carpenter, Andrew Gelman, and Aki Vehtari. Pathfinder: Parallel quasi-Newton variational inference. *Journal of Machine Learning Research*, 23(306):1–49, 2022.