

POSITIVE SECTIONAL CURVATURE IS NOT PRESERVED UNDER THE RICCI FLOW IN DIMENSIONS SEVEN AND THIRTEEN

DAVID GONZÁLEZ-ÁLVARO AND MASOUMEH ZAREI

ABSTRACT. We prove that there exist SU_3 -invariant metrics on Aloff–Wallach spaces W_{k_1, k_2}^7 , as well as SU_5 -invariant metrics on the Berger space B^{13} , which have positive sectional curvature and evolve under the Ricci flow to metrics with non-positively curved planes.

1. INTRODUCTION

The Ricci flow of a Riemannian manifold (M, g) is a one-parameter family of metrics $g(\ell)$ on M satisfying the evolution equation

$$\frac{\partial}{\partial \ell} g(\ell) = -2 \operatorname{Ric}(g(\ell))$$

with $g(0) = g$. Since its introduction by Hamilton in 1982 [Ha82], it has been a fundamental question to understand which curvature conditions are preserved under the flow, see [Ni14, Ri14].

While positive scalar curvature and positive curvature operator are preserved in all dimensions [Ha82, Ha86, Wi13, BW08], there exist infinitely many dimensions where certain curvature conditions lying in between are not preserved [AN16, GZ24].

In this article we are interested in manifolds with positive sectional curvature, denoted by $\sec > 0$. Hamilton showed in [Ha82] that $\sec > 0$ is preserved in dimensions 2 and 3. In contrast, there exist manifolds in dimensions 4, 6, 12 and 24 with metrics of $\sec > 0$ which lose this property when evolved under the Ricci flow, see [BK23, BW07, CW15]. Our main results add dimensions 7 and 13 to this list, and provide the first odd dimensional examples of this phenomenon.

Theorem A. *On infinitely many Aloff–Wallach spaces W_{k_1, k_2}^7 , there exist SU_3 -invariant metrics of positive sectional curvature which evolve under the Ricci flow to metrics with non-positively curved planes.*

In fact, we will show that the conclusion of Theorem A holds for all (positively curved) Aloff–Wallach spaces. However, for the latter we can only give a computer-assisted proof. In contrast, the proof of Theorem A is theoretical.

2020 *Mathematics Subject Classification.* Primary: 53C21. Secondary: 53E20.

Key words and phrases. Positive sectional curvature, Ricci flow, Aloff–Wallach spaces, Berger Space.

Theorem B. *There exist SU_5 -invariant metrics on the Berger space B^{13} which have positive sectional curvature and evolve under the Ricci flow to metrics with non-positively curved planes.*

The strategy for proving the theorems is as follows. Since the manifolds under consideration are homogeneous spaces G/H , we may restrict our attention to the set of G -invariant metrics on them, which remains invariant under the Ricci flow. Moreover, for some spaces, there are subfamilies of G -invariant metrics which are simpler to describe, yet are Ricci flow invariant. In our cases, these metrics are described by a finite set of positive parameters x_1, \dots, x_n , and the Ricci flow is determined by an ODE system $x'_i = -2r_i(x_1, \dots, x_n)x_i$, where $i = 1, \dots, n$ and r_i denote the eigenvalues of the Ricci tensor. Next, we need a characterization of those metrics which have $\sec > 0$. For each of the families of metrics we are considering, this characterization has been provided by the work of Thomas Püttmann in [Pue99]. With all this information at hand, the results follow from analyzing the behaviour of a metric on the boundary of the set of $\sec > 0$ metrics. Let us give more details for each of the theorems.

Aloff–Wallach spaces, denoted by W_{k_1, k_2}^7 , constitute an infinite family of 7-dimensional homogeneous spaces of the form SU_3/S^1 which admit an SU_3 -invariant metric with positive sectional curvature if and only if $k_1 k_2 (k_1 + k_2) \neq 0$ [Zi07]. Among this family, the space $W_{1,1}^7$ and those of the form $W_{n, n+1}^7$, for sufficiently large n , satisfy Theorem A. More generally, in any infinite sequence $\{W_{k_1^n, k_2^n}^7\}_{n \in \mathbb{N}}$, with $\frac{k_1^n}{k_2^n}$ converging to 1, there are infinitely many spaces satisfying Theorem A. Since the cohomology group $H^4(W_{k_1, k_2}^7, \mathbb{Z})$ equals the cyclic group of order $k_1^2 + k_2^2 + k_1 k_2$, we conclude that there exist infinitely many homotopy types satisfying Theorem A. We remark that, with our strategy, a theoretical proof for all cases seems impractical due to the complexity of the computations. In Section 3.5 we, however, provide a computer-assisted proof to demonstrate that Theorem A in fact holds for all Aloff–Wallach spaces.

Let us discuss the metrics on W_{k_1, k_2}^7 which are involved in the proofs. Aloff and Wallach [AW75] were the first to discover, among a two-parameter family of SU_3 -invariant metrics on W_{k_1, k_2}^7 , metrics with positive sectional curvature, provided that $k_1 k_2 (k_1 + k_2) \neq 0$. Later, Thomas Püttmann [Pue99] considered a more general four-parameter family of SU_3 -invariant metrics, denoted by $\langle \cdot, \cdot \rangle_{t, \vec{s}}$, where $\vec{s} = (s_0, s_1, s_2)$, and determined the subset $\Omega(k_1, k_2)$ consisting of those metrics which have $\sec > 0$. It is noteworthy to mention here that since t will be used as one of the parameters describing the metrics $\langle \cdot, \cdot \rangle_{t, \vec{s}}$ throughout this article, we denote the time parameter in the Ricci flow equation by ℓ .

An important feature of the family $\langle \cdot, \cdot \rangle_{t, \vec{s}}$ is that it remains invariant under the Ricci flow. Moreover, the Ricci flow of a metric $\langle \cdot, \cdot \rangle_{t, \vec{s}}$ reduces to an ODE system in 4 variables. In order to find metrics of $\sec > 0$ which lose this property under the flow, it is essential to understand the boundary of $\Omega(k_1, k_2)$ and analyze the ODE system at the boundary

points. Unfortunately, it appears that describing the boundary of $\Omega(k_1, k_2)$ and, consequently, analyzing the ODE system of a boundary point by hand in this generality is too complicated.

In order to establish a manageable setting, the crucial observation is the existence of a two-parameter and a three-parameter subfamilies of metrics on $W_{1,1}^7$ which remain invariant under the flow. This allows us to restrict our exploration within these two families with a twofold advantage. First, the descriptions of the boundaries of the subsets of $\text{sec} > 0$ metrics among the 2-parameter and 3-parameter families are notably simpler. Second, the Ricci flow reduces to an ODE system in just two and three variables, respectively. This makes the analysis of the Ricci flow at the boundary points more feasible. We show that there are metrics with $\text{sec} > 0$ in these two families which evolve to metrics with non-positively curved planes.

With this finding concerning the three-parameter family of metrics on $W_{1,1}^7$ at hand, the proof of Theorem A now follows from a convergence argument, which works for Aloff–Wallach spaces “sufficiently close” to $W_{1,1}^7$. In more detail, we introduce a variable $\xi \in (0, 1]$ which parametrizes all Aloff–Wallach spaces in a continuous way, where W_{k_1, k_2}^7 corresponds to $\xi = \frac{k_1}{k_2}$. This allows us to treat different spaces in a continuous manner and hence relate the Ricci flow behavior of metrics on spaces $W_{k_1^n, k_2^n}^7$ to that of the three-parameter family on $W_{1,1}^k$, provided that $\frac{k_1^n}{k_2^n}$ tends to 1.

The proof of Theorem B is straightforward. The space of SU_5 -invariant metrics on $B^{13} = \text{SU}_5/\text{Sp}_2\text{S}^1$ only depends on two parameters and the Ricci flow reduces to an ODE system in two variables. Moreover, the subspace of positively curved metrics, determined by Püttmann, has a simple description. A careful analysis of the ODE system at a boundary metric then yields the existence of $\text{sec} > 0$ metrics which evolve under the Ricci flow to metrics with non-positively curved planes.

The article is organized as follows. In Section 2 we collect the relevant information about Aloff–Wallach spaces, and we describe the aforementioned two-parameter and three-parameter families of metrics on $W_{1,1}^7$. In Section 3 we show the existence of metrics with $\text{sec} > 0$ in the two-parameter and three-parameter families which evolve under the Ricci flow to metrics with non-positively curved planes. We illustrate this phenomenon by drawing their phase portrait. Then we prove Theorem A, and give a computer-assisted proof showing that Theorem A holds for all Aloff–Wallach spaces. Finally, in Section 4 we prove Theorem B.

Acknowledgements. We would like to thank Christoph Böhm for his suggestion on how to extend the argument from $W_{1,1}^7$ to infinitely many spaces. We would also like to thank Anusha Krishnan for helpful discussions. This project was initiated while the first author was visiting the University of Münster. The first author wishes to thank the University of Münster for providing excellent working conditions.

The first author was supported by grants PID2021-124195NB-C31 and PID2021-124195NB-C32 from the Agencia Estatal de Investigación and the Ministerio de Ciencia e Innovación (Spain). The second author was supported by Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy EXC 2044-390685587, Mathematics Münster: Dynamics-Geometry-Structure, and by DFG grant ZA976/1-1 within the Priority Program SPP2026 "Geometry at Infinity".

2. ALOFF–WALLACH SPACES

In this section we give the necessary background on Aloff–Wallach spaces and describe the metrics which will be used in the proof of Theorem A.

2.1. Diagonal Metrics on Aloff–Wallach spaces. In this subsection, we follow the notation of Püttmann in [Pue99] to establish the space of metrics on Aloff–Wallach spaces that are of interest to us. Let

$$\mathbb{T}^2 = \{\text{diag}(e^{i\theta_0}, e^{i\theta_1}, e^{i\theta_2}) \mid \theta_0 + \theta_1 + \theta_2 = 0\}$$

be a maximal torus of SU_3 . For each vector $\vec{k} = (k_0, k_1, k_2) \in \mathbb{Z}^3 \setminus \{0\}$ with $0 \leq k_1 \leq k_2$, $\text{gcd}(k_1, k_2) = 1$, and $k_0 + k_1 + k_2 = 0$, there exists an embedding of \mathbb{S}^1 in the maximal torus $\mathbb{T}^2 < \text{SU}_3$, denoted by \mathbb{S}_k^1 , given by $z \mapsto \text{diag}(z^{k_0}, z^{k_1}, z^{k_2})$. The homogeneous space $\text{SU}_3/\mathbb{S}_k^1$ is called an Aloff–Wallach space and is denoted by W_{k_1, k_2}^7 .

Before describing the desired set of metrics on W_{k_1, k_2}^7 , let us briefly recall some facts about invariant metrics on homogeneous spaces, see [Be87, Chapter 7] for details. Let G be a compact Lie group and $M = G/H$ be a homogeneous space. Fix a bi-invariant metric $\langle\langle \cdot, \cdot \rangle\rangle$ on G , and let $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{p}$ be a $\langle\langle \cdot, \cdot \rangle\rangle$ -orthogonal decomposition of the Lie algebra \mathfrak{g} of G , where \mathfrak{h} denotes the Lie algebra of H . Then \mathfrak{p} is isomorphic to the tangent space of M at the origin $o = eH$. It is well-known that there is a one-to-one correspondence between $\text{Ad}(H)$ -invariant inner products on \mathfrak{p} and G -invariant metrics on M . Moreover, the Ricci tensor of such a G -invariant metric is determined by its value at the point o , which gives an $\text{Ad}(H)$ -invariant symmetric bilinear form on \mathfrak{p} .

Returning to the case of Aloff–Wallach spaces, let t, s_0, s_1, s_2 be positive real numbers, and denote $\vec{s} = (s_0, s_1, s_2)$. Define the vector $\hat{k} = (\hat{k}_0, \hat{k}_1, \hat{k}_2)$ as $\hat{k} = \frac{\vec{k}}{|\vec{k}|}$, and let $\Gamma = k_1^2 + k_2^2 + k_1 k_2$. Note that $|\vec{k}| = \sqrt{2\Gamma}$. The tangent space of W_{k_1, k_2}^7 at the origin o can be described as the image of the sum of one copy $\bar{\varepsilon}$ of \mathbb{R} and three copies \mathbb{C}_i of \mathbb{C} via the following maps:

$$\begin{aligned} \Psi_{t, \vec{s}}: \bar{\varepsilon} \oplus \mathbb{C}_0 \oplus \mathbb{C}_1 \oplus \mathbb{C}_2 &\rightarrow \mathfrak{su}_3 \\ (x, z_0, z_1, z_2) &\mapsto \begin{pmatrix} \frac{2i(\hat{k}_1 - \hat{k}_2)}{\sqrt{6t}}x & -\frac{z_2}{\sqrt{s_2}} & \frac{\bar{z}_1}{\sqrt{s_1}} \\ \frac{\bar{z}_2}{\sqrt{s_2}} & \frac{2i(\hat{k}_2 - \hat{k}_0)}{\sqrt{6t}}x & -\frac{z_0}{\sqrt{s_0}} \\ -\frac{z_1}{\sqrt{s_1}} & \frac{\bar{z}_0}{\sqrt{s_0}} & \frac{2i(\hat{k}_0 - \hat{k}_1)}{\sqrt{6t}}x \end{pmatrix}. \end{aligned}$$

For each (t, \vec{s}) , Püttmann defines an SU_3 -invariant metric $\langle \cdot, \cdot \rangle_{t, \vec{s}}$ on W_{k_1, k_2}^7 as follows. Denote by $\langle \cdot, \cdot \rangle$ the standard inner product on $\bar{\varepsilon} \oplus \mathbb{C}_0 \oplus \mathbb{C}_1 \oplus \mathbb{C}_2$. Then the metric $\langle \cdot, \cdot \rangle_{t, \vec{s}}$ is defined by the rule

$$\Psi_{t, \vec{s}}^*(\langle \cdot, \cdot \rangle_{t, \vec{s}}) = \langle \cdot, \cdot \rangle.$$

It will be convenient to define $\langle \cdot, \cdot \rangle_{t, \vec{s}}$ from a different point of view. To do so, note that the image of each of the summands $\bar{\varepsilon}, \mathbb{C}_0, \mathbb{C}_1, \mathbb{C}_2$ by $\Psi_{t, \vec{s}}$ does not depend on t, \vec{s} . Thus we may define

$$\mathfrak{p}_0 = \Psi_{t, \vec{s}}(\bar{\varepsilon}), \quad \mathfrak{p}_1 = \Psi_{t, \vec{s}}(\mathbb{C}_0), \quad \mathfrak{p}_2 = \Psi_{t, \vec{s}}(\mathbb{C}_1), \quad \mathfrak{p}_3 = \Psi_{t, \vec{s}}(\mathbb{C}_2).$$

Observe that

$$\mathfrak{p}_0 = \mathrm{span}\{E\}, \quad \mathfrak{p}_1 = \mathrm{span}\{E_{23}, F_{23}\}, \quad \mathfrak{p}_2 = \mathrm{span}\{E_{13}, F_{13}\}, \quad \mathfrak{p}_3 = \mathrm{span}\{E_{12}, F_{12}\}.$$

Here $E = \mathrm{diag}((k_1 - k_2)i, (2k_2 + k_1)i, -(2k_1 + k_2)i)$ and E_{jk}, F_{jk} are defined by

$$E_{jk} = \begin{cases} 1 & \text{in the position } (j, k), \\ -1 & \text{in the position } (k, j), \\ 0 & \text{otherwise,} \end{cases} \quad F_{jk} = \begin{cases} i & \text{in the position } (j, k), \\ i & \text{in the position } (k, j), \\ 0 & \text{otherwise.} \end{cases}$$

Note that \mathfrak{p}_0 and E depend on k_1, k_2 .

Let $\langle\langle \cdot, \cdot \rangle\rangle$ be the bi-invariant metric on SU_3 given by $\langle\langle X, Y \rangle\rangle = -\frac{1}{2} \mathrm{Re} \mathrm{tr}(XY) = -\frac{1}{2} \mathrm{tr}(XY)$, for $X, Y \in \mathfrak{su}_3$. Define $\mathfrak{h} \subset \mathfrak{su}_3$ as the Lie algebra of S_k^1 , i.e. \mathfrak{h} is spanned by $\mathrm{diag}(-(k_1 + k_2)i, k_1i, k_2i)$. Then the orthogonal complement \mathfrak{h}^\perp of \mathfrak{h} with respect to $\langle\langle \cdot, \cdot \rangle\rangle$ equals $\mathfrak{p}_0 \oplus \mathfrak{p}_1 \oplus \mathfrak{p}_2 \oplus \mathfrak{p}_3$. Moreover,

$$\mathfrak{p} = \mathfrak{p}_0 \oplus \mathfrak{p}_1 \oplus \mathfrak{p}_2 \oplus \mathfrak{p}_3$$

is a $\langle\langle \cdot, \cdot \rangle\rangle$ -orthogonal $\mathrm{Ad}(\mathrm{S}_k^1)$ -invariant irreducible decomposition of \mathfrak{h}^\perp . As explained in [Pue99, p. 648], the metrics $\langle \cdot, \cdot \rangle_{t, \vec{s}}$ can be defined in the following diagonal form:

$$(2.1) \quad \langle \cdot, \cdot \rangle_{t, \vec{s}} = t \langle\langle \cdot, \cdot \rangle\rangle|_{\mathfrak{p}_0} + s_0 \langle\langle \cdot, \cdot \rangle\rangle|_{\mathfrak{p}_1} + s_1 \langle\langle \cdot, \cdot \rangle\rangle|_{\mathfrak{p}_2} + s_2 \langle\langle \cdot, \cdot \rangle\rangle|_{\mathfrak{p}_3}.$$

Note that the matrices $E, E_{23}, F_{23}, E_{13}, F_{13}, E_{12}, F_{12}$ defined above form a basis for \mathfrak{p} , which is adapted to the decomposition $\mathfrak{p}_0 \oplus \mathfrak{p}_1 \oplus \mathfrak{p}_2 \oplus \mathfrak{p}_3$ by construction. Moreover, the basis is orthogonal with respect to $\langle\langle \cdot, \cdot \rangle\rangle$. In fact, all matrices except E , whose norm is $\sqrt{3\Gamma}$, are unit vectors. Let $\tilde{E} = \frac{E}{\sqrt{3\Gamma}}$. Thus we arrive to the following $\langle\langle \cdot, \cdot \rangle\rangle$ -orthonormal basis for \mathfrak{p} adapted to $\mathfrak{p}_0 \oplus \mathfrak{p}_1 \oplus \mathfrak{p}_2 \oplus \mathfrak{p}_3$:

$$(2.2) \quad \mathcal{B} = \{\tilde{E}, E_{23}, F_{23}, E_{13}, F_{13}, E_{12}, F_{12}\}.$$

Remark 2.1. By [Pue99, Corollary 4.3], the cone of SU_3 -invariant and right T^2 -invariant metrics on W_{k_1, k_2}^7 consists precisely of the metrics $\langle \cdot, \cdot \rangle_{t, \vec{s}}$. In the case of $W_{1, 1}^7$, the subcone of metrics which are additionally right U_2 -invariant consists of those with $t = s_0$ and $s_1 = s_2$; which we denote by the quadruple (t, t, s, s) .

2.2. Eigenvalues of the Ricci tensor. Here we discuss the Ricci tensor of the metrics $\langle \cdot, \cdot \rangle_{t, \vec{s}}$. If $k_1 \neq k_2$, then the modules \mathfrak{p}_i are pairwise inequivalent as $\text{Ad}(\mathbf{S}_k^1)$ -invariant representations. This has two important consequences. First, all SU_3 -invariant metrics on W_{k_1, k_2}^7 are of the form $\langle \cdot, \cdot \rangle_{t, \vec{s}}$. Second, the Ricci tensor is diagonal, meaning that it is of the form

$$(2.3) \quad \text{Ric} = tr_0 \langle \langle \cdot, \cdot \rangle \rangle|_{\mathfrak{p}_0} + s_0 r_1 \langle \langle \cdot, \cdot \rangle \rangle|_{\mathfrak{p}_1} + s_1 r_2 \langle \langle \cdot, \cdot \rangle \rangle|_{\mathfrak{p}_2} + s_2 r_3 \langle \langle \cdot, \cdot \rangle \rangle|_{\mathfrak{p}_3}$$

for certain $r_i \in \mathbb{R}$ which depend on t, \vec{s}, k_1, k_2 .

If $k_1 = k_2 = 1$, then \mathfrak{p}_2 and \mathfrak{p}_3 are equivalent as $\text{Ad}(\mathbf{S}_{1,1}^1)$ -invariant representations, see [Pue99, Lemma 4.2]. Thus, it is a priori not clear whether the Ricci tensor of the metrics $\langle \cdot, \cdot \rangle_{t, \vec{s}}$ is of the form (2.3). In order to show that it has in fact a diagonal form, recall from Remark 2.1 that the metrics $\langle \cdot, \cdot \rangle_{t, \vec{s}}$ are precisely the SU_3 -invariant and right \mathbf{T}^2 -invariant metrics on $W_{1,1}^7$, and again from [Pue99, Lemma 4.2] we see that the modules \mathfrak{p}_i , $i = 0, \dots, 3$, are inequivalent as $\text{Ad}(\mathbf{T}^2)$ -invariant representations. This implies by Schur's Lemma that the Ricci tensor of the metrics $\langle \cdot, \cdot \rangle_{t, \vec{s}}$ has diagonal form (2.3) as well.

Next we compute the eigenvalues of the Ricci tensor in (2.3) (cf. [Wa82, Lemma 2]). Let us recall the procedure to compute the eigenvalues of diagonal Ricci tensors in the general case. Let G/H be a compact homogeneous space and $\langle \langle \cdot, \cdot \rangle \rangle$ a bi-invariant metric on G , and denote by \mathfrak{h}^\perp the orthogonal complement of \mathfrak{h} . Then, for each G -invariant metric g on G/H there is an irreducible $\text{Ad}(H)$ -invariant decomposition $\oplus_{i=0}^m \mathfrak{p}_i$ of \mathfrak{h}^\perp such that g is of the form

$$g = x_0 \langle \langle \cdot, \cdot \rangle \rangle|_{\mathfrak{p}_0} + \dots + x_m \langle \langle \cdot, \cdot \rangle \rangle|_{\mathfrak{p}_m},$$

for some positive $x_i \in \mathbb{R}$. Suppose that the Ricci tensor of g is diagonal with respect to the given decomposition $\oplus_{i=0}^m \mathfrak{p}_i$, i.e. it is of the form

$$\text{Ric} = x_0 r_0 \langle \langle \cdot, \cdot \rangle \rangle|_{\mathfrak{p}_0} + \dots + x_m r_m \langle \langle \cdot, \cdot \rangle \rangle|_{\mathfrak{p}_m},$$

for certain $r_i \in \mathbb{R}$. In this case, the eigenvalues r_i can be computed using the following formula (see e.g. [Be87, Corollary 7.38] or [WZ86, Section 1]):

$$(2.4) \quad r_i = \frac{b_i}{2x_i} - \frac{1}{2d_i} \sum_{j,k=0}^m [ijk] \frac{x_j}{x_i x_k} + \frac{1}{4d_i} \sum_{j,k=0}^m [ijk] \frac{x_i}{x_j x_k}.$$

Here $d_i = \dim \mathfrak{p}_i$ and b_i is determined by $-B|_{\mathfrak{p}_i} = b_i \langle \langle \cdot, \cdot \rangle \rangle|_{\mathfrak{p}_i}$, where B denotes the Killing form of G . The bracket constants $[ijk]$ are defined by

$$[ijk] = \sum_{\substack{e_\alpha \in \mathfrak{p}_i \\ e_\beta \in \mathfrak{p}_j \\ e_\gamma \in \mathfrak{p}_k}} \langle \langle [e_\alpha, e_\beta], e_\gamma \rangle \rangle^2,$$

where $\{e_i\}$ is a $\langle \langle \cdot, \cdot \rangle \rangle$ -orthonormal basis of \mathfrak{h}^\perp adapted to $\oplus_{i=0}^m \mathfrak{p}_i$. Notice that $[ijk]$ is symmetric in all indices.

In the case of the metrics $\langle \cdot, \cdot \rangle_{t, \vec{s}}$ from (2.1) we have $m = 3$, $x_0 = t$, $x_1 = s_0$, $x_2 = s_1$ and $x_3 = s_2$. The Killing form of \mathfrak{su}_n is given by $B(X, Y) = 2n \text{tr}(XY)$. Hence for \mathfrak{su}_3 , we

have $B(X, Y) = 6 \operatorname{tr}(XY) = -12 \langle\langle X, Y \rangle\rangle$. Therefore, $b_i = 12$ for each $i \in \{0, 1, 2, 3\}$. It remains to compute the bracket constants $[ijk]$. In order to do so, we take the basis from (2.2). For convenience we collect the Lie brackets of the elements of the basis \mathcal{B} in Table 1. Notice that Table 1 represents $[e_i, e_j]_{\mathfrak{p}}$, where e_i is the vector in row i , and e_j is the vector in column j . The notation $[e_i, e_j]_{\mathfrak{p}}$ stands for the orthogonal projection of $[e_i, e_j]$ onto \mathfrak{p} , which provides enough information for our purposes.

	\tilde{E}	E_{23}	F_{23}	E_{13}	F_{13}	E_{12}	F_{12}
\tilde{E}	0	$\frac{3(k_1+k_2)}{\sqrt{3\Gamma}} F_{23}$	$-\frac{3(k_1+k_2)}{\sqrt{3\Gamma}} E_{23}$	$\frac{3k_1}{\sqrt{3\Gamma}} F_{13}$	$-\frac{3k_1}{\sqrt{3\Gamma}} E_{13}$	$-\frac{3k_2}{\sqrt{3\Gamma}} F_{12}$	$\frac{3k_2}{\sqrt{3\Gamma}} E_{12}$
E_{23}		0	$\frac{3(k_1+k_2)}{\sqrt{3\Gamma}} \tilde{E}$	E_{12}	F_{12}	$-E_{13}$	$-F_{13}$
F_{23}			0	$-F_{12}$	E_{12}	$-F_{13}$	E_{13}
E_{13}				0	$\frac{3k_1}{\sqrt{3\Gamma}} \tilde{E}$	E_{23}	$-F_{23}$
F_{13}					0	F_{23}	E_{23}
E_{12}						0	$-\frac{3k_2}{\sqrt{3\Gamma}} \tilde{E}$
F_{12}							0

TABLE 1. \mathfrak{p} -projection of Lie brackets of the elements of the basis (2.2).

Using Table 1 we get:

Proposition 2.2. *The bracket constants for W_{k_1, k_2}^7 are as follows:*

$$\begin{aligned}
 [ijj] &= 0, & i, j \in \{1, 2, 3\}, & & [ij0] &= 0, & i, j \in \{1, 2, 3\}, & i \neq j, \\
 [123] &= 4, & & & [00i] &= 0, & i \in \{1, 2, 3\}, & \\
 [220] &= \frac{6k_1^2}{\Gamma}, & & & [110] &= \frac{6(k_1 + k_2)^2}{\Gamma}, & & \\
 [330] &= \frac{6k_2^2}{\Gamma}. & & & & & &
 \end{aligned}$$

Now we have all the ingredients to compute the eigenvalues of the Ricci tensor.

Proposition 2.3. *The Ricci eigenvalues of the metric $\langle \cdot, \cdot \rangle_{t, \vec{s}}$ on W_{k_1, k_2}^7 are given by*

$$\begin{aligned} r_0 &= \frac{3t}{2\Gamma} \left(\frac{(k_1 + k_2)^2}{s_0^2} + \frac{k_1^2}{s_1^2} + \frac{k_2^2}{s_2^2} \right), \\ r_1 &= \frac{6}{s_0} - \frac{3(k_1 + k_2)^2 t}{2\Gamma s_0^2} + \left(\frac{s_0}{s_1 s_2} - \frac{s_1}{s_0 s_2} - \frac{s_2}{s_0 s_1} \right), \\ r_2 &= \frac{6}{s_1} - \frac{3k_1^2 t}{2\Gamma s_1^2} + \left(\frac{s_1}{s_0 s_2} - \frac{s_0}{s_1 s_2} - \frac{s_2}{s_0 s_1} \right), \\ r_3 &= \frac{6}{s_2} - \frac{3k_2^2 t}{2\Gamma s_2^2} + \left(\frac{s_2}{s_0 s_1} - \frac{s_0}{s_1 s_2} - \frac{s_1}{s_0 s_2} \right). \end{aligned}$$

2.3. Cone of positively curved metrics. One of the main results in Püttmann's article is to determine the two following sets (see Remark 2.1):

- the set of $\text{sec} > 0$ metrics among SU_3 -invariant right U_2 -invariant metrics on $W_{1,1}^7$,
- the set of $\text{sec} > 0$ metrics among SU_3 -invariant right T^2 -invariant metrics on W_{k_1, k_2}^7 for any pair k_1, k_2 .

The first set is very easy to describe [Pue99, Lemma 6.1].

Theorem 2.4 (Püttmann). *A metric (t, t, s, s) on $W_{1,1}^7$ has $\text{sec} > 0$ if and only if $t < s$.*

Much more background is needed to give an account of the second set. Let us introduce the following notation:

$$\Omega(k_1, k_2) = \{(t, \vec{s}) \text{ such that } \langle \cdot, \cdot \rangle_{t, \vec{s}} \text{ on } W_{k_1, k_2}^7 \text{ is positively curved}\}.$$

In order to describe this set, we need to introduce a certain space Ω'_λ and a real valued function t_A on Ω'_λ .

For each k_1, k_2 and each (t, \vec{s}) , Püttmann defines a real function $\hat{\lambda} = \hat{\lambda}(k_1, k_2, t, \vec{s})$ as the minimum of sectional curvatures of certain planes with respect to the metric $\langle \cdot, \cdot \rangle_{t, \vec{s}}$ on W_{k_1, k_2}^7 , see [Pue99, Definition 5.1]. Since we only need some estimates and not the precise definition of $\hat{\lambda}$, we omit its technical definition. It turns out [Pue99, Lemma 5.16] that the condition $\hat{\lambda} > 0$ does not depend on t . Define the space $\Omega'_\lambda(k_1, k_2)$ as

$$(2.5) \quad \Omega'_\lambda(k_1, k_2) = \{\vec{s} \mid \hat{\lambda}(k_1, k_2, t, \vec{s}) > 0 \text{ for some (hence all) } t > 0\}.$$

Remark 2.5. In this work we shall only use the two following properties of the sets $\Omega'_\lambda(k_1, k_2)$. First, the positively curved metrics on W_{k_1, k_2}^7 found by Aloff and Wallach in [AW75] provide an explicit subset of $\Omega'_\lambda(k_1, k_2)$. According to [AW75], metrics of the form $(t, \vec{s}) = (t, t, s, s)$ with $t < s$ are positively curved (cf. [Pue99, p.662]), see also Theorem 2.4. Hence, it follows from the definition of $\hat{\lambda}$ that

$$(2.6) \quad \text{if } s_1 = s_2, \text{ then } (s_0, s_1, s_2) \in \Omega'_\lambda(k_1, k_2) \text{ for all } s_0 \in (0, s_1) \text{ and all } k_1, k_2.$$

Second, $\Omega'_\lambda(k_1, k_2)$ is an open set. Hence any triple (s_0, s_1, s_2) sufficiently close to one as in (2.6) also belongs to $\Omega'_\lambda(k_1, k_2)$.

Now we recall the definition of the map $t_A = t_A(\vec{s}, k_1, k_2)$. We need the following auxiliary functions:

$$\begin{aligned}\tilde{a}_j(\vec{s}) &= \frac{4}{s_j}, \\ \tilde{b}_j(\vec{s}) &= -\frac{2s_1s_2 + 2s_0s_2 + 2s_0s_1 - s_0^2 - s_1^2 - s_2^2}{s_0s_1s_2} + \frac{s_{j-1} - s_j + s_{j+1}}{s_{j-1}s_{j+1}}, \\ \sigma(\vec{s}) &= 2s_1s_2 + 2s_0s_2 + 2s_0s_1 - s_0^2 - s_1^2 - s_2^2.\end{aligned}$$

We define the following open subsets of \mathbb{R}_+^3 :

$$\begin{aligned}\Omega'_\sigma &= \{\vec{s} \mid \sigma > 0\}, \\ D_\sigma &= \Omega'_\sigma \setminus \{(s, s, s) \mid s > 0\}.\end{aligned}$$

Note that by [Pue99, Lemma 5.17], we have

$$\Omega'_\lambda(k_1, k_2) \subset \Omega'_\sigma \quad \text{for all } k_1, k_2.$$

Further, we have the matrix $\tilde{A}(\vec{s})$ defined as (see [Pue99, p.662])

$$\tilde{A}(\vec{s}) = \begin{pmatrix} \tilde{a}_0 & \tilde{b}_2 & \tilde{b}_1 \\ \tilde{b}_2 & \tilde{a}_1 & \tilde{b}_0 \\ \tilde{b}_1 & \tilde{b}_0 & \tilde{a}_2 \end{pmatrix}.$$

According to [Pue99, Lemma 5.19], the matrix $\tilde{A}(\vec{s})$ is invertible for each $s \in D_\sigma$.

For each vector \vec{s} and each pair k_1, k_2 , define the vector

$$(2.7) \quad v(\vec{s}, k_1, k_2) := \left(\frac{\hat{k}_0}{s_0}, \frac{\hat{k}_1}{s_1}, \frac{\hat{k}_2}{s_2} \right)^t.$$

Then $t_A(\vec{s}, k_1, k_2)$ with $\vec{s} \in \Omega'_\sigma$ is defined as

$$t_A(\vec{s}, k_1, k_2) = \begin{cases} 0, & \text{if } s_0 = s_1 = s_2, \\ \frac{2}{9} \langle v, \tilde{A}^{-1}v \rangle^{-1}, & \text{otherwise.} \end{cases}$$

Because the entries of \tilde{A} are rational functions, it follows that, for any fixed pair k_1, k_2 , the map $t_A(\vec{s}, k_1, k_2)$ is smooth for $\vec{s} \in D_\sigma$.

We are ready to give the description of $\Omega(k_1, k_2)$, cf. [Pue99, Observation 5.23]:

Theorem 2.6 (Püttmann). *The cone $\Omega(k_1, k_2)$ consisting of metrics $\langle \cdot, \cdot \rangle_{t, \vec{s}}$ on W_{k_1, k_2}^7 of $\text{sec} > 0$ can be characterized as*

$$\Omega(k_1, k_2) = \{(t, \vec{s}) \in \mathbb{R}_+ \times \Omega'_\lambda(k_1, k_2) \mid t < t_A(\vec{s}, k_1, k_2)\}.$$

2.4. Homogeneous Ricci flow, a special space and special families of metrics. Let us firstly review basic properties of the Ricci flow on homogeneous spaces, see e.g. [La13] for details. Given a homogeneous Riemannian manifold (M, g) , let $g(\ell)$ be the solution to the Ricci flow equation with initial metric g .

Let G a Lie group acting isometrically and transitively on (M, g) . Since the isometry group is preserved under the Ricci flow, there is a presentation of M as a homogeneous space $(M, g(\ell)) = (G/H, g(\ell))$ for all ℓ with the same reductive decomposition $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{p}$.

We denote by $\langle \cdot, \cdot \rangle_\ell = g(\ell)(o)$ the $\text{Ad}(H)$ -invariant inner product on \mathfrak{p} , determined by the value of $g(\ell)$ at the origin o . The family $\langle \cdot, \cdot \rangle_\ell$ is the solution to the following ODE

$$(2.8) \quad \frac{d}{d\ell} \langle \cdot, \cdot \rangle_\ell = -2 \text{Ric}(\langle \cdot, \cdot \rangle_\ell),$$

where $\text{Ric}(\langle \cdot, \cdot \rangle_\ell) = \text{Ric}(g(\ell))(o)$. Hence the maximal interval of time where $g(\ell)$ exists is of the form (L_-, L_+) for some $-\infty \leq L_- < 0 < L_+ \leq \infty$.

Now we restrict the discussion to the spaces W_{k_1, k_2}^7 . Let us denote the metric $\langle \cdot, \cdot \rangle_{t, \vec{s}}$ by the quadruple (t, s_0, s_1, s_2) . The fact that the Ricci tensor is of the form (2.3) permits to express the Ricci flow equation (2.8) as a system of ODEs. More precisely, the Ricci flow of (t, s_0, s_1, s_2) is a family of metrics $(t(\ell), s_0(\ell), s_1(\ell), s_2(\ell))$ satisfying:

$$(2.9) \quad \begin{cases} t' = -2r_0 t, \\ s'_j = -2r_{j+1} s_j, \text{ for all } j \in \{0, 1, 2\}, \\ (t(0), s_0(0), s_1(0), s_2(0)) = (t, s_0, s_1, s_2). \end{cases}$$

Observe that, although it is not written explicitly, the Ricci tensor eigenvalues r_i depend on $(t(\ell), s_0(\ell), s_1(\ell), s_2(\ell))$ and on k_1, k_2 .

To prove the existence of metrics with $\text{sec} > 0$ on Aloff–Wallach spaces W_{k_1, k_2}^7 evolving under the Ricci flow to metrics with non-positively curved planes, our strategy consists of understanding the boundary of $\Omega(k_1, k_2)$ and analyzing the Ricci flow starting from a metric (t, s_0, s_1, s_2) on the boundary of $\Omega(k_1, k_2)$. In particular, we may need to compute $t_A(s_0(\ell), s_1(\ell), s_2(\ell), k_1, k_2)$, for small ℓ , and for that we must compute \tilde{A}^{-1} at $(s_0(\ell), s_1(\ell), s_2(\ell))$. We notice that the expression for \tilde{A}^{-1} becomes unmanageable at points (s_0, s_1, s_2) with pairwise distinct coordinates. Unfortunately, this issue cannot be resolved for the cases where $(k_1, k_2) \neq (1, 1)$ since in general the functions $(s_0(\ell), s_1(\ell), s_2(\ell))$ are pairwise distinct for small $\ell \neq 0$ (even if two of them are equal at $\ell = 0$). This can be seen by inspecting the Ricci eigenvalues from Proposition 2.3 and the Ricci flow equations. However, for the case of $W_{1,1}^7$, the Ricci flow of a metric with $s_1 = s_2$, as further clarified below, satisfies $s_1(\ell) = s_2(\ell)$ for all times ℓ . This observation plays a crucial role in our computations.

In more detail, when $k_1 = k_2 = 1$ and $s_1 = s_2$, Proposition 2.3 shows that $r_2 = r_3$, see also (3.5) below. This implies that the three-parameter family of metrics on $W_{1,1}^7$ of the form (t, s_0, s_1, s_2) with $s_1 = s_2$ remains invariant under the Ricci flow. More precisely, if

$s_1(0) = s_2(0)$, then $s_1(\ell) = s_2(\ell)$ for all ℓ by uniqueness of the solutions. If in addition one has $t(0) = s_0(0)$, then $r_0 = r_1$, so the two-parameter family of metrics of the form (t, s_0, s_1, s_2) with $t = s_0$ and $s_1 = s_2$ remains invariant under the Ricci flow as well. The last property also follows from the fact that the metrics (t, t, s, s) on $W_{1,1}^7$ are precisely those which are additionally right U_2 -invariant and that Ricci flow preserves the isometry group.

The upshot of considering these two families is that the corresponding subset of metrics with $\text{sec} > 0$ is manageable to describe, and in fact very simple in the case of the two-parameter family by Theorem 2.4. Moreover, for these two families of metrics, the Ricci flow reduces to an ODE system in two or three variables instead of the four variables in (2.9), making it easier to analyze.

In the two following sections we treat each family separately and find metrics on $W_{1,1}^7$, within each family which have $\text{sec} > 0$ and lose this property when evolved by the Ricci flow. More precisely, we prove the following two theorems.

Theorem 2.7 (2-parameter family). *There exist SU_3 -invariant metrics of the form (t, t, s, s) on the Aloff–Wallach space $W_{1,1}^7$ which have positive sectional curvature and evolve under the Ricci flow to metrics with non-positively curved planes.*

Theorem 2.8 (3-parameter family). *There exist SU_3 -invariant metrics of the form (t, s_0, s, s) on the Aloff–Wallach space $W_{1,1}^7$ which have positive sectional curvature and evolve under the Ricci flow to metrics with non-positively curved planes.*

Remark 2.9. To prove Theorem A, we primarily need to prove Theorem 2.8, and then apply a continuity argument. However, since Theorem 2.7 shows the non-invariance of metrics with $\text{sec} > 0$ on $W_{1,1}^7$ under the Ricci flow within a smaller subset of positively curved metrics, and moreover the proof is short, we consider this case as well.

3. TWO AND THREE-PARAMETER FAMILIES, NEARBY SPACES, AND THE GENERAL CASE

This section is divided in five subsections. In the first two subsections we discuss the two and three-parameter families of metrics on $W_{1,1}^7$ and give the proofs of Theorems 2.7 and 2.8, respectively. In the third subsection, we draw the phase portraits of the (normalized) Ricci flow of the two and three-parameter families. In the fourth subsection we prove Theorem A. Finally, we show, through a computer-assisted proof, that the results of Theorem A hold for all Aloff–Wallach spaces admitting positive curvature.

3.1. Two-parameter family on $W_{1,1}^7$. Here we prove Theorem 2.7 (see Remark 4.1 for an alternative proof). We consider metrics on $W_{1,1}^7$ of the form (t, t, s, s) . Recall from Theorem 2.4 that $\text{sec} > 0$ if and only if $t < s$. We shall show that the $\text{sec} > 0$ metrics $(1 - \varepsilon, 1 - \varepsilon, 1, 1)$ evolve under the Ricci flow to a metric with non-positively curved planes, for sufficiently small $\varepsilon > 0$.

From Proposition 2.3 we see that the eigenvalues of the Ricci tensor are:

$$(3.1) \quad r_0 = r_1 = \frac{2s^2 + t^2}{ts^2}, \quad r_2 = r_3 = \frac{3(4s - t)}{2s^2}.$$

Let $g(\ell) := (t(\ell), t(\ell), s(\ell), s(\ell))$ be the solution to the Ricci flow with the initial metric $g(0) := (1, 1, 1, 1)$. The Ricci flow is determined by the ODE system

$$(3.2) \quad \begin{cases} t' = -2r_0 t, \\ s' = -2r_2 s. \end{cases}$$

This system has a unique solution in an open interval $(-L, L)$ for some $L > 0$. We claim that

$$(3.3) \quad \frac{d}{d\ell} \left(\frac{t(\ell)}{s(\ell)} \right) \Big|_{\ell=0} > 0.$$

This implies that there exists $\bar{L} > 0$ with $\bar{L} \leq L$ such that $t(\ell) < s(\ell)$ for all $\ell \in (-\bar{L}, 0)$ and $t(\ell) > s(\ell)$ for all $\ell \in (0, \bar{L})$. This proves Theorem 2.7.

It remains to prove (3.3). For simplicity, we drop the variable ℓ from the equation. Differentiating and using (3.2) and (3.1) we get:

$$\frac{d}{d\ell} \left(\frac{t}{s} \right) = \frac{t's - ts'}{s^2} = \frac{-2r_0 ts + 2tr_2 s}{s^2} = \frac{-4s^2 - 5t^2 + 12ts}{s^2}.$$

At $\ell = 0$ we have $t = s = 1$ and hence the derivative equals 3.

3.2. Three-parameter family on $W_{1,1}^7$. Here we consider metrics on $W_{1,1}^7$ of the form (t, s_0, s_1, s_2) with $s_1 = s_2$. We denote $s := s_1 = s_2$ for the convenience of the reader. Also, in order to avoid any confusion between s and s_0 , we set $x := s_0$. Thus the quadruples of positive numbers describing the corresponding metrics read as (t, x, s, s) . Before giving the proof of Theorem 2.8, let us explain how the background given in Section 2 is adapted to this family of metrics. In what follows, we may drop some parameters from the equations in order to facilitate the reading.

The Ricci flow of a metric (t, x, s, s) is given by the ODE system

$$(3.4) \quad \begin{cases} t' = -2r_0 t, \\ x' = -2r_1 x, \\ s' = -2r_2 s. \end{cases}$$

From Proposition 2.3 we see that the eigenvalues of the Ricci tensor are:

$$(3.5) \quad r_0 = \frac{t(2s^2 + x^2)}{x^2 s^2}, \quad r_1 = \frac{4xs^2 - 2ts^2 + x^3}{x^2 s^2}, \quad r_2 = r_3 = \frac{12s - t - 2x}{2s^2}.$$

We also need the value of $t_A(x, s, s)$ on $W_{1,1}^7$. The matrix $\tilde{A} = \tilde{A}(x, s, s)$ is given by

$$\tilde{A} = \begin{pmatrix} \frac{4}{x} & \frac{x-3s}{s^2} & \frac{x-3s}{s^2} \\ \frac{x-3s}{s^2} & \frac{4}{s} & -\frac{2}{s} \\ \frac{x-3s}{s^2} & -\frac{2}{s} & \frac{4}{s} \end{pmatrix}.$$

Its inverse equals

$$\tilde{A}^{-1} = \frac{1}{(s-x)^2(4s-x)} \begin{pmatrix} s^3x & \frac{s^2x(3s-x)}{2} & \frac{s^2x(3s-x)}{2} \\ \frac{s^2x(3s-x)}{2} & \frac{s(16s^3-9s^2x+6sx^2-x^3)}{12} & \frac{s(8s^3+9s^2x-6sx^2+x^3)}{12} \\ \frac{s^2x(3s-x)}{2} & \frac{s(8s^3+9s^2x-6sx^2+x^3)}{12} & \frac{s(16s^3-9s^2x+6sx^2-x^3)}{12} \end{pmatrix}.$$

In this case $v = (-\frac{2}{x\sqrt{6}}, \frac{1}{s\sqrt{6}}, \frac{1}{s\sqrt{6}})$. A simple computation shows that

$$(3.6) \quad \tilde{A}^{-1}v = \frac{1}{\sqrt{6}(s-x)(4s-x)} \begin{pmatrix} (sx-2s^2) \\ -s^2 \\ -s^2 \end{pmatrix}.$$

We arrive to the expression

$$(3.7) \quad t_A(x, s, s) = \frac{x(4s-x)}{3s}.$$

Remark 3.1. Note that working with the metrics of the form $(t, \vec{s}) = (t, x, s, s)$ has another advantage besides the simplicity of the computations: from (2.6) we know that $(x, s, s) \in \Omega'_\lambda$ for any $x \in (0, s)$.

For later reference, we state the following immediate consequence of Theorem 2.6, which actually holds for all W_{k_1, k_2}^7 with $k_1 k_2 (k_1 + k_2) \neq 0$.

Lemma 3.2. *For any $x \in (0, 1)$ the following hold:*

- (1) *If $t \in (0, t_A(x, 1, 1))$, then the metric $(t, x, 1, 1)$ is positively curved.*
- (2) *If $t \geq t_A(x, 1, 1)$, then the metric $(t, x, 1, 1)$ has non-positively curved planes.*

Now we are ready to prove Theorem 2.8.

We begin the proof by choosing a suitable initial metric on $W_{1,1}^7$ for the Ricci flow. We claim, as it will be clear later, that a metric of the form $g := (t_A(x, 1, 1) - \varepsilon, x, 1, 1)$ for any x in the open interval $(0.8, 1)$ and sufficiently small $\varepsilon > 0$ yields the result. In order to prove it, let $g(\ell) := (t(\ell), x(\ell), s(\ell), s(\ell))$ be the solution to the Ricci flow with the initial metric $g(0) := (t_A(x_0, 1, 1), x_0, 1, 1)$ for some $x_0 \in (0.8, 1)$. Since the Ricci flow is given by the ODE system (3.4), there exists a unique solution in an open interval $(-L, L)$ for some $L > 0$.

Note that, because $\frac{x(0)}{s(0)} = x_0 \in (0.8, 1)$, there exists some positive $L_1 \leq L$ such that:

(P1) $\frac{x(\ell)}{s(\ell)} \in (0.8, 1)$ for all $\ell \in (-L_1, L_1)$.

Now we claim that

$$(3.8) \quad \frac{d}{d\ell} \left(\frac{t_A(x(\ell), s(\ell), s(\ell))}{t(\ell)} \right) \Big|_{\ell=0} < 0.$$

This implies that there exists $L_2 > 0$ with $L_2 \leq L$ such that the following properties hold:

(P2) $t_A(x(\ell), s(\ell), s(\ell)) > t(\ell)$ for all $\ell \in (-L_2, 0)$,

(P3) $t_A(x(\ell), s(\ell), s(\ell)) < t(\ell)$ for all $\ell \in (0, L_2)$.

By defining $\bar{L} = \min\{L_1, L_2\}$, it follows that (P1), (P2) and (P3) hold for all $\ell \in (-\bar{L}, \bar{L})$. This easily implies the theorem as follows. Define the rescaled metrics

$$h(\ell) := \frac{1}{s(\ell)} g(\ell) \cong \left(\frac{t(\ell)}{s(\ell)}, \frac{x(\ell)}{s(\ell)}, 1, 1 \right).$$

Note that $t_A(\frac{x}{s}, 1, 1) = \frac{1}{s} t_A(x, s, s)$. Thus $t_A(x, s, s) > t$ and $t_A(x, s, s) < t$ are equivalent to $t_A(\frac{x}{s}, 1, 1) > \frac{t}{s}$ and $t_A(\frac{x}{s}, 1, 1) < \frac{t}{s}$, respectively. Since for each $\ell \in (-\bar{L}, \bar{L})$, we have $\frac{x(\ell)}{s(\ell)} \in (0.8, 1)$, Lemma 3.2 implies that the metrics $h(\ell)$ are positively curved for $\ell \in (-\bar{L}, 0)$ and have non-positively curved planes for $\ell \in (0, \bar{L})$. The same holds true for the rescaled metrics $g(\ell)$, hence proving the theorem.

It remains to prove that (3.8) holds for $x(0) = x_0 \in (0.8, 1)$. For simplicity, we drop the variable ℓ in the following computations. From (3.7) we know that

$$\frac{d}{d\ell} \left(\frac{t_A(x, s, s)}{t} \right) = \frac{d}{d\ell} \left(\frac{x(4s - x)}{3st} \right) = \frac{B}{3s^2t^2},$$

where

$$B := 2st(2sx' + 2s'x - xx') - x(4s - x)(s't + st').$$

Substituting the derivatives t', x', s' by the Ricci flow equations (3.4) we get

$$B = 2txs(r_0(4s - x) + r_1(2x - 4s) - r_2x).$$

Replacing r_0, r_1, r_2 with (3.5) we obtain

$$B = \frac{t}{xs} (32ts^3 - 12txs^2 + 8stx^2 - tx^3 + 16x^2s^2 - 32xs^3 - 20sx^3 + 6x^4).$$

Let D denote $D = \frac{xs}{t} B$. Then

$$(3.9) \quad \frac{d}{d\ell} \left(\frac{t_A(x, s, s)}{t} \right) = \frac{D}{3txs^3},$$

so the sign of the derivative agrees with the sign of D . Since we are interested in its value at time $\ell = 0$, we assume that

$$(t(0), x(0), s(0), s(0)) = (t_A(x, 1, 1), x, 1, 1) = \left(\frac{x(4-x)}{3}, x, 1, 1 \right),$$

for some $x > 0$. Now we find conditions on x such that (3.8) is satisfied. With this initial condition, we get

$$D = \frac{x}{3} (32 - 32x - 16x^2 + 6x^3 + x^4).$$

The roots of D in the variable x can be computed with the help of a computer. They are all real and equal:

$$\lambda_1 \approx -7.48, \quad \lambda_2 = -2, \quad \lambda_3 = 0, \quad \lambda_4 \approx 0.79, \quad \lambda_5 \approx 2.69,$$

where $\lambda_1, \lambda_4, \lambda_5$ are approximations up to two digits. It turns out that D is negative in the open interval (λ_4, λ_5) . This implies the claim for each $x \in (0.8, 1)$.

3.3. Phase Portrait. To illustrate the evolution of metrics (t, x, s, s) on $W_{1,1}^7$ under the Ricci flow, we provide the phase portrait of the normalized Ricci flow of such metrics in Figures 1 and 2, for the two, respectively, three parameter families. Since the volume remains constant under the normalized flow, we may assume that $tx^2s^4 = 1$. Whence, we have $t = x^{-2}s^{-4}$. The normalized Ricci flow system reduces then to a system with two equations as follows:

$$(3.10) \quad \begin{aligned} x' &= \frac{-40x^3s^6 + 24s^2 - 18x^5s^4 - 2x^2 + 48x^4s^5}{7x^3s^6}, \\ s' &= \frac{-36x^4s^5 + 16x^3s^6 + 10x^5s^4 + 5x^2 - 4s^2}{7x^4s^5}. \end{aligned}$$

In Figures 1 and 2 the horizontal axis represents the variable x , while the vertical axis represents the variable s . We have three disjoint regions as follow:

$$\begin{aligned} \mathcal{G} &= \left\{ (x, s) \in \mathbb{R}_+^2 \mid x < s, \ x^{-2}s^{-4} < \frac{x(4s-x)}{3s} \right\} \\ &= \{(x, s) \in \mathbb{R}_+^2 \mid x < s, \ 3 < 4x^3s^4 - x^4s^3\}, \\ \mathcal{P} &= \{(x, s) \in \mathbb{R}_+^2 \mid 3 \geq 4x^3s^4 - x^4s^3\}, \\ \mathcal{W} &= \{(x, s) \in \mathbb{R}_+^2 \mid x \geq s, \ 3 < 4x^3s^4 - x^4s^3\}. \end{aligned}$$

The set \mathcal{G} is depicted in *light green*, \mathcal{P} in *pink*, and \mathcal{W} in *white*. The boundary of \mathcal{G} is given by $s = x$ and $4x^3s^4 - x^4s^3 = 3$, represented by the *blue* line and the *red* curve, respectively.

Each point $(x, s) \in \mathbb{R}_+^2$ represents the metric $g(x, s) = (x^{-2}s^{-4}, x, s, s)$ on $W_{1,1}^7$. These regions are distinguished by our knowledge of the sign of the sectional curvature of the corresponding metrics. Namely, if $(x, s) \in \mathcal{G}$, then $g(x, s) \in \Omega$, i.e. $g(x, s)$ has $\text{sec} > 0$; if $(x, s) \in \mathcal{P}$, then $g(x, s)$ has planes with non-positive sectional curvature; and if $(x, s) \in \mathcal{W}$, we do not have precise information about the sign of the sectional curvature.

In Figures 1(A) and 2(A) we specify two points

$$E^{(+)} = \left(\frac{2}{5} \sqrt[7]{\frac{125}{8}}, \sqrt[7]{\frac{125}{8}} \right), \text{ in } \textit{green}, \quad E^{(-)} = \left(2 \sqrt[7]{\frac{1}{8}}, \sqrt[7]{\frac{1}{8}} \right), \text{ in } \textit{magenta},$$

which correspond to the Einstein metrics on $W_{1,1}^7$ given in [Ni04, p. 25]. Note that the metric corresponding to $E^{(+)}$ has $\text{sec} > 0$ and the metric corresponding to $E^{(-)}$ has planes with negative curvature ([Pue99, Lemma 6.1]).

Further, let

$$\mathcal{C} = \{(x, s) \in \mathbb{R}_+^2 \mid x^3 s^4 = 1\}.$$

This set is illustrated in *purple* (partly covered by a green curve) in 1(A) and represents the 2-parameter family of metrics on $W_{1,1}^7$. Note that the two Einstein metrics lie on this curve.

Moreover, in Figures 1(A) and 2(A) we illustrate two solutions to System (3.10) with initial values

$$p_1 = \left(\left(\frac{10}{11} \right)^{\frac{4}{3}}, 1.1 \right) \in \mathcal{C} \cap \mathcal{G}, \quad p_2 = (0.87, 1.1) \in \mathcal{G},$$

respectively, depicted as the *green* curve in Figure 1(A) and as the *black* curve in Figure 2(A). Let us denote the green curve by γ and the black curve by β . Note that p_1 represents a metric in the 2-parameter family and p_2 represents a metric in the 3-parameter family. We remark further that since we assume the volume to remain constant throughout the flow, the initial values that we choose here are different from those we choose for the (unnormalized) Ricci flow in the proof of Theorems 2.7, 2.8. Notice that the curve γ initiating at $p_1 \in \mathcal{C}$ remains in \mathcal{C} while evolving, and converges to the point representing the Einstein metric $E^{(-)}$ in positive times (see Remark 4.1). The same phenomenon happens for the curve β , i.e. it converges to $E^{(-)}$ in positive times. This behaviour indicates that both γ and β evolve to points corresponding to metrics with non-positively curved planes.

In Figures 1(B) and 2(B) we depict the same integral curves γ and β as in Figures 1(A) and 2(A), respectively. However, to observe their behaviors close to the boundary of \mathcal{G} , we provide a close-up image.

Recall that a metric (x, x, s, s) in the 2-parameter family has positive curvature if and only if $x < s$. For the metrics in this family with fixed volume, i.e. $(x^{-2}s^{-4}, x, s, s)$ with $x^3s^4 = 1$, the boundary condition $x = s$ reduces to $(x, s) = (1, 1)$. In Figure 1(B), there exist ℓ_0, ℓ_1 with $0 < \ell_0 < \ell_1$, where $\gamma(\ell_0) = (1, 1)$ and $\gamma(\ell_1) \in \mathcal{P}$.

Similarly, for the curve β , there exists $\tilde{\ell}_0 > 0$ such that $q = \beta(\tilde{\ell}_0) \in \mathcal{P}$. This indicates that the corresponding Ricci flow metric at time $\tilde{\ell}_0$ has a non-positively curved plane, as proven in Theorem 2.8.

3.4. Nearby spaces. In this subsection we give the proof of Theorem A, which relies on the proof of Theorem 2.8 and an analytic argument. To this end, we will consider infinitely many Aloff–Wallach spaces at the same time. It will be then helpful to, roughly speaking, parametrize all of them in a continuous manner.




FIGURE 1. 2-parameter family




FIGURE 2. 3-parameter family

Recall that the Ricci eigenvalues r_i from Proposition 2.3 depend on the parameters k_1, k_2 , and the same holds for the vector v from (2.7) and the map t_A from (3.7). Clearly, the expressions for r_i , v and t_A make sense for k_1, k_2 not necessarily in \mathbb{N} . Moreover, observe that all of these values remain invariant under rescaling $\lambda k_1, \lambda k_2$ of k_1, k_2 . This suggests to identify each pair $k_1, k_2 \in \mathbb{R}_+$ with the quotient $\xi = \frac{k_1}{k_2}$. In this way, positively curved Aloff-Wallach spaces W_{k_1, k_2}^7 correspond to rational numbers $\xi \in (0, 1]$; recall that we have assumed $\gcd(k_1, k_2) = 1$ and $0 < k_1 \leq k_2$. Conversely, each $\xi \in (0, 1] \cap \mathbb{Q}$ can be written as $\xi = \frac{k_1}{k_2}$ with $k_1, k_2 \in \mathbb{N}$, $0 < k_1 \leq k_2$ and $\gcd(k_1, k_2) = 1$, hence it corresponds to the positively curved Aloff-Wallach space W_{k_1, k_2}^7 , which we may simply denote by W_ξ^7 . In particular, the space $W_{1,1}^7$ corresponds to $\xi = 1$.

We extend the maps t_A , v and r_i so that they depend on ξ instead of k_1, k_2 , where ξ is any real number in the interval $(0, 1]$. More precisely, $v : \mathbb{R}_+^3 \times (0, 1] \rightarrow \mathbb{R}^3$ is defined as the smooth map

$$v(\vec{s}, \xi) = \left(-\frac{1 + \xi}{s_0 \sqrt{2(\xi^2 + \xi + 1)}}, \frac{\xi}{s_1 \sqrt{2(\xi^2 + \xi + 1)}}, \frac{1}{s_2 \sqrt{2(\xi^2 + \xi + 1)}} \right).$$

The map t_A extends to $t_A : D_\sigma \times (0, 1] \rightarrow \mathbb{R}$ defined as

$$t_A(\vec{s}, \xi) = \frac{2}{9} \langle v(\vec{s}, \xi), \tilde{A}^{-1}(\vec{s})v(\vec{s}, \xi) \rangle^{-1}.$$

Observe that $\tilde{A}^{-1}(\vec{s})$ is a rational matrix in the variables (s_0, s_1, s_2) , so t_A is smooth on $D_\sigma \times (0, 1]$.

The Ricci eigenvalues r_i extend to smooth maps $r_i : \mathbb{R}_+^4 \times (0, 1]$ as

$$\begin{aligned} r_0(t, \vec{s}, \xi) &= \frac{3t}{2(\xi^2 + \xi + 1)} \left(\frac{(\xi + 1)^2}{s_0^2} + \frac{\xi^2}{s_1^2} + \frac{1}{s_2^2} \right), \\ r_1(t, \vec{s}, \xi) &= \frac{6}{s_0} - \frac{3(\xi + 1)^2 t}{2(\xi^2 + \xi + 1)s_0^2} + \left(\frac{s_0}{s_1 s_2} - \frac{s_1}{s_0 s_2} - \frac{s_2}{s_0 s_1} \right), \\ r_2(t, \vec{s}, \xi) &= \frac{6}{s_1} - \frac{3\xi^2 t}{2(\xi^2 + \xi + 1)s_1^2} + \left(\frac{s_1}{s_0 s_2} - \frac{s_0}{s_1 s_2} - \frac{s_2}{s_0 s_1} \right), \\ r_3(t, \vec{s}, \xi) &= \frac{6}{s_2} - \frac{3t}{2(\xi^2 + \xi + 1)s_2^2} + \left(\frac{s_2}{s_0 s_1} - \frac{s_0}{s_1 s_2} - \frac{s_1}{s_0 s_2} \right). \end{aligned}$$

For the rest of the section we fix $x \in (0.8, 1)$ and we assume

$$p = (t_A(x, 1, 1, 1), x, 1, 1, 1) = \left(\frac{x(4-x)}{3}, x, 1, 1, 1 \right).$$

We define the map

$$(3.11) \quad \begin{aligned} F: \mathbb{R}_+ \times D_\sigma \times (0, 1] &\rightarrow \mathbb{R} \\ (t, s_0, s_1, s_2, \xi) &\mapsto \frac{t_A(s_0, s_1, s_2, \xi)}{t}. \end{aligned}$$

Denote by $g(\ell) = (t(\ell), x(\ell), s(\ell), s(\ell))$ the solution to the Ricci flow on $W_{1,1}^7$ of the initial metric $(t_A(x, 1, 1, 1), x, 1, 1)$, and consider the map f defined as

$$(3.12) \quad f(\ell) = F(g(\ell), 1).$$

This is precisely the map that we have considered in the proof of Theorem 2.8, where we have shown that the derivative at $\ell = 0$ is negative if $x \in (0.8, 1)$, see (3.8). The derivative $f'(0)$ is related to the differential $dF_p: T_p V \cong \mathbb{R}^5 \rightarrow T_1 \mathbb{R} \cong \mathbb{R}$ as follows:

$$(3.13) \quad dF_p(g'(0), 0) = f'(0) < 0,$$

where $g'(0) = (t'(0), x'(0), s'(0), s'(0))$.

Now we are ready to state the technical result which implies Theorem A.

Theorem 3.3. *Let $\{\xi_n\}_{n \in \mathbb{N}}$ be any sequence converging to 1 with $\xi_n \in \mathbb{Q} \cap (0, 1]$. Then there is some n_0 such that all Aloff–Wallach spaces W_{ξ_n} with $n \geq n_0$ have a $\text{sec} > 0$ metric which evolves under the Ricci flow to a metric with non-positively curved planes.*

Proof. For each sequence $\{\xi_n\}_{n \in \mathbb{N}}$ as in the statement we define

$$p_n = (t_A(x, 1, 1, \xi_n), x, 1, 1, \xi_n).$$

Recall from Remark 2.5 that $(x, 1, 1) \in \Omega'_\lambda(\xi_n)$ for all n . Note that, because t_A is smooth, the sequence $\{p_n\}_{n \in \mathbb{N}}$ converges to p .

We think of the first four coordinates of p_n and p as a metric on $W_{\xi_n}^7$ and on W_1^7 , respectively. We denote the corresponding Ricci flows (2.9) by $g_n(\ell) = (t^n(\ell), s_0^n(\ell), s_1^n(\ell), s_2^n(\ell))$ and $g(\ell) = (t(\ell), x(\ell), s(\ell), s(\ell))$, respectively, i.e.

$$(3.14) \quad \begin{cases} (t^n)' = -2r_0(t^n, s_0^n, s_1^n, s_2^n, \xi_n)t^n, \\ (s_j^n)' = -2r_{j+1}(t^n, s_0^n, s_1^n, s_2^n, \xi_n)s_j^n, \text{ for all } j \in \{0, 1, 2\}, \\ (t^n(0), s_0^n(0), s_1^n(0), s_2^n(0), \xi_n) = p_n. \end{cases}$$

$$(3.15) \quad \begin{cases} t' = -2r_0(t, s_0, s_1, s_2, 1)t, \\ s_j' = -2r_{j+1}(t, s_0, s_1, s_2, 1)s_j, \text{ for all } j \in \{0, 1, 2\}, \\ (t(0), s_0(0), s_1(0), s_2(0), \xi) = p. \end{cases}$$

Since r_i are smooth functions on $\mathbb{R}_+^4 \times (0, 1]$ and p_n converges to p , the right hand side of (3.14) converges to the right hand side of (3.15). Consequently, the same holds for the left hand sides, so $g'_n(0)$ converges to $g'(0)$.

Denote $f_n(\ell) = F(g_n(\ell), \xi_n)$. Then $f'_n(0) = dF_{p_n}(g'_n(0), 0)$. Due to all the convergences that have been discussed above, it follows that $f'_n(0)$ converges to $f'(0)$, where f is the map in (3.12). Since $f'(0) < 0$, we conclude that there is some n_0 such that $f'_n(0) < 0$ for all $n \geq n_0$.

It follows that for each $n \geq n_0$ there is a small time $L_n > 0$ such that $f_n(\ell) > 1$ for $\ell \in (-L_n, 0)$ and $f_n(\ell) < 1$ for $\ell \in (0, L_n)$. By making L_n sufficiently small if necessary, we have that $(s_0^n(\ell), s_1^n(\ell), s_2^n(\ell)) \in \Omega'_\lambda(k_1^n, k_2^n)$ because the latter is an open set, see Remark 2.5. Together with Theorem 2.6, we conclude that for each $n \geq n_0$ and any positive $\varepsilon < L_n$, the metric $(t_A(x, 1, 1, \xi_n) - \varepsilon, x, 1, 1)$ on $W_{\xi_n}^7$ has $\text{sec} > 0$ and evolves under the Ricci flow to a metric with non-positively curved planes. \square

3.5. General Aloff–Wallach Spaces. In this subsection we prove that all Aloff–Wallach spaces have $\text{sec} > 0$ metrics which evolve under the Ricci flow to metrics with non-positively curved planes. We use the notation from Section 3.4. Some computations are done with Maple, due to their complexity.

For each $\xi \in (0, 1] \cap \mathbb{Q}$ and each $x \in (0, 1)$, let g denote the metric $(t_A(x, 1, 1, \xi), x, 1, 1)$ on W_ξ , and let $g_\xi(\ell) = (t(\ell), s_0(\ell), s_1(\ell), s_2(\ell))$ be the Ricci flow of g , which exists in some interval $(-L_\xi, L_\xi)$. Although g depends on x and ξ (and $g_\xi(\ell)$ on x), we do not explicitly denote it, since there is no risk of confusion.

We denote by $F_\xi : \mathbb{R}_+ \times D_\sigma \rightarrow \mathbb{R}$ the map F from (3.11) restricted to $\mathbb{R}_+ \times D_\sigma \times \{\xi\}$. Consider the smooth map

$$f_\xi(\ell) = F_\xi(g_\xi(\ell)).$$

As in the proof of Theorem A, our goal is to show that for each $\xi \in (0, 1] \cap \mathbb{Q}$ there is some $x \in (0, 1)$ for which $f'_\xi(0) < 0$, because this implies that the $\text{sec} > 0$ metrics $(t_A(x, 1, 1, \xi) - \varepsilon, x, 1, 1)$, with ε sufficiently small, evolve to metrics with non-positively curved planes.

Note that $f'_\xi(0) = \langle \nabla_g F_\xi, g'_\xi(0) \rangle$, where $\nabla_g F_\xi$ denotes the value at g of the gradient

$$\nabla F_\xi = \left(\frac{\partial F_\xi}{\partial t}, \frac{\partial F_\xi}{\partial s_0}, \frac{\partial F_\xi}{\partial s_1}, \frac{\partial F_\xi}{\partial s_2} \right).$$

The computation of $\frac{\partial F_\xi}{\partial s_i}$ involves the derivative of t_A and hence of \tilde{A}^{-1} with respect to s_i . As explained in Section 2.4, the expression of \tilde{A}^{-1} for arbitrary triples \vec{s} is complicated. In order to simplify the computations as much as possible, we use the fact that

$$(3.16) \quad \frac{\partial \tilde{A}^{-1}}{\partial s_i}(x, 1, 1) = -\tilde{A}^{-1}(x, 1, 1) \frac{\partial \tilde{A}}{\partial s_i}(x, 1, 1) \tilde{A}^{-1}(x, 1, 1).$$

This formula easily follows from the observation that the derivative of $I = \tilde{A} \tilde{A}^{-1}$ vanishes. The upshot of (3.16) is that we only have to compute \tilde{A}^{-1} at $(x, 1, 1)$, which we have already done in Section 3.2, and that we have the expression for \tilde{A} at hand, so we can compute $\frac{\partial \tilde{A}}{\partial s_i}$ directly and then evaluate at $(x, 1, 1)$.

Using (3.16) and the fact that \tilde{A}^{-1} is a symmetric matrix, we find that

$$(3.17) \quad \frac{\partial F_\xi}{\partial s_i}(g) = \frac{-2}{3x(4-x)} \left(\frac{2\langle \frac{\partial v}{\partial s_i}, \tilde{A}^{-1}v_\xi \rangle - \langle \tilde{A}^{-1}v_\xi, \frac{\partial \tilde{A}}{\partial s_i}(\tilde{A}^{-1}v_\xi) \rangle}{\langle v_\xi, \tilde{A}^{-1}v_\xi \rangle^2} \right),$$

where $\frac{\partial v}{\partial s_i}$ is evaluated at $(x, 1, 1, \xi)$, the matrices \tilde{A}^{-1} and $\frac{\partial \tilde{A}}{\partial s_i}$ are evaluated at $(x, 1, 1)$, and v_ξ denotes $v(x, 1, 1, \xi)$.

Let us compute the terms appearing in (3.17). Along the way we will indicate which computations have been done with Maple. First we introduce some notations to simplify the expressions. Let

$$P_i := \langle \frac{\partial v}{\partial s_i}, \tilde{A}^{-1}v_\xi \rangle, \quad Q_i := \langle \tilde{A}^{-1}v_\xi, \frac{\partial \tilde{A}}{\partial s_i}(\tilde{A}^{-1}v_\xi) \rangle, \quad L = \langle v_\xi, \tilde{A}^{-1}v_\xi \rangle.$$

Then (3.17) can be written as:

$$(3.18) \quad \frac{\partial F_\xi}{\partial s_i}(g) = \frac{-2}{3x(4-x)} \left(\frac{2P_i - Q_i}{L^2} \right).$$

First we compute the term which is involved in all of L, P_i, Q_i :

$$(3.19) \quad W := \tilde{A}^{-1}v_\xi = \frac{1}{(x^2 - 5x + 4)\sqrt{2(\xi^2 + \xi + 1)}} \begin{pmatrix} \frac{(x-2)(\xi+1)}{2} \\ \frac{(\xi-1)x^2 + (-5\xi+5)x - 2\xi - 10}{12} \\ -\frac{(\xi-1)x^2 + (-5\xi+5)x + 10\xi + 2}{12} \end{pmatrix}.$$

We use it to obtain

$$L = \langle v_\xi, W \rangle = \frac{x^2(\xi-1)^2 - 4x(\xi-1)^2 - 12(\xi+1)^2}{24x(x-4)(\xi^2 + \xi + 1)}.$$

Next, we differentiate v with respect to s_i and evaluate at $(x, 1, 1)$:

$$\begin{aligned} \frac{\partial v}{\partial s_0}(x, 1, 1) &= \left(\frac{1 + \xi}{x^2 \sqrt{2(\xi^2 + \xi + 1)}}, 0, 0 \right), \\ \frac{\partial v}{\partial s_1}(x, 1, 1) &= \left(0, -\frac{\xi}{\sqrt{2(\xi^2 + \xi + 1)}}, 0 \right), \\ \frac{\partial v}{\partial s_2}(x, 1, 1) &= \left(0, 0, -\frac{1}{\sqrt{2(\xi^2 + \xi + 1)}} \right). \end{aligned}$$

We can now compute $P_i = \langle \frac{\partial v}{\partial s_i}, W \rangle$:

$$\begin{aligned} P_0 &= \frac{(\xi + 1)^2(x - 2)}{4(\xi^2 + \xi + 1)x^2(x - 1)(x - 4)}, \\ P_1 &= -\frac{((\xi - 1)x^2 + (-5\xi + 5)x - 2\xi - 10)\xi}{24(\xi^2 + \xi + 1)(x - 1)(x - 4)}, \\ P_2 &= \frac{(\xi - 1)x^2 + (-5\xi + 5)x + 10\xi + 2}{24(\xi^2 + \xi + 1)(x - 1)(x - 4)}. \end{aligned}$$

Next, we differentiate \tilde{A} with respect to s_i and evaluate at $(x, 1, 1)$:

$$\begin{aligned} \frac{\partial \tilde{A}}{\partial s_0}(x, 1, 1) &= \begin{pmatrix} -\frac{4}{x^2} & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \\ \frac{\partial \tilde{A}}{\partial s_1}(x, 1, 1) &= \begin{pmatrix} 0 & \frac{-x^2+x+1}{x} & -\frac{(x-1)^2}{x} \\ \frac{-x^2+x+1}{x} & -4 & 1 \\ -\frac{(x-1)^2}{x} & 1 & 0 \end{pmatrix}, \\ \frac{\partial \tilde{A}}{\partial s_2}(x, 1, 1) &= \begin{pmatrix} 0 & -\frac{(x-1)^2}{x} & \frac{-x^2+x+1}{x} \\ -\frac{(x-1)^2}{x} & 0 & 1 \\ \frac{-x^2+x+1}{x} & 1 & -4 \end{pmatrix}. \end{aligned}$$

Let us denote $U_i := \frac{\partial \tilde{A}}{\partial s_i}(W)$. Then we have

$$\begin{aligned} U_0 &= \frac{1}{(x^2 - 5x + 4)\sqrt{2(\xi^2 + \xi + 1)}} \begin{pmatrix} -\frac{(x^2+2x-4)(\xi+1)}{x^2} \\ \frac{(x-2)(\xi+1)}{2} \\ \frac{(x-2)(\xi+1)}{2} \end{pmatrix}, \\ U_1 &= \frac{1}{(x^2 - 5x + 4)\sqrt{2(\xi^2 + \xi + 1)}} \begin{pmatrix} -\frac{(\xi-1)x^3 + (-19\xi-5)x^2 + (32\xi+4)x - 8\xi + 8}{12x} \\ \frac{(-11\xi-1)x^3 + (43\xi-7)x^2 + (-8\xi+32)x - 12\xi - 12}{12x} \\ \frac{(-5\xi-7)x^3 + (19\xi+29)x^2 + (-32\xi-40)x + 12\xi + 12}{12x} \end{pmatrix}, \\ U_2 &= \frac{1}{(x^2 - 5x + 4)\sqrt{2(\xi^2 + \xi + 1)}} \begin{pmatrix} \frac{(\xi-1)x^3 + (5\xi+19)x^2 + (-4\xi-32)x - 8\xi + 8}{12x} \\ \frac{(-7\xi-5)x^3 + (29\xi+19)x^2 + (-40\xi-32)x + 12\xi + 12}{12x} \\ -\frac{(\xi+11)x^3 + (7\xi-43)x^2 + (-32\xi+8)x + 12\xi + 12}{12x} \end{pmatrix}. \end{aligned}$$

Lastly, compute $Q_i = \langle W, U_i \rangle$:

$$(3.20) \quad \begin{aligned} Q_0 &= -\frac{(x+2)(\xi+1)^2(x-2)}{2x^2(x-4)^2(x-1)(\xi^2+\xi+1)}, \\ Q_1 &= \frac{-(\xi-1)^2x^4 + (7\xi^2 - 18\xi + 11)x^3 + (24\xi^2 + 96\xi - 24)x^2}{48x(x-4)^2(x-1)(\xi^2+\xi+1)} \\ &\quad + \frac{(-116\xi^2 - 152\xi + 28)x + 32\xi^2 - 32}{48x(x-4)^2(x-1)(\xi^2+\xi+1)}, \end{aligned}$$

$$(3.21) \quad \begin{aligned} Q_2 &= \frac{-(\xi-1)^2x^4 + (11\xi^2 - 18\xi + 7)x^3 + (-24\xi^2 + 96\xi + 24)x^2}{48x(x-4)^2(x-1)(\xi^2+\xi+1)} \\ &\quad + \frac{(28\xi^2 - 152\xi - 116)x - 32\xi^2 + 32}{48x(x-4)^2(x-1)(\xi^2+\xi+1)}. \end{aligned}$$

The last two equations, i.e. (3.20) and (3.21), have been computed with Maple.

Having computed L , P_i and Q_i , we use (3.18) to compute the partial derivatives of F_ξ .

$$(3.22) \quad \begin{aligned} \frac{\partial F_\xi}{\partial t}(g) &= \frac{-48\xi^2 - 48\xi - 48}{x(x-4)((\xi-1)^2x^2 - 4(\xi-1)^2x - 12(\xi+1)^2)}, \\ \frac{\partial F_\xi}{\partial s_0}(g) &= \frac{384(\xi+1)^2(x-2)(\xi^2+\xi+1)}{x(x-4)((\xi-1)^2x^2 - 4(\xi-1)^2x - 12(\xi+1)^2)^2}, \end{aligned}$$

$$(3.23) \quad \begin{aligned} \frac{\partial F_\xi}{\partial s_1}(g) &= \frac{-24(\xi^2+\xi+1)((\xi^2 - \frac{2}{3}\xi - \frac{1}{3})x^4 + (-\frac{29}{3}\xi^2 + 6\xi + \frac{11}{3})x^3}{(x-4)(x-1)((\xi-1)^2x^2 - 4(\xi-1)^2x - 12(\xi+1)^2)^2} \\ &\quad + \frac{(32\xi^2 - 8\xi - 8)x^2 + (-28\xi^2 + \frac{8}{3}\xi + \frac{28}{3})x + \frac{32\xi^2}{3} - \frac{32}{3}}{(x-4)(x-1)((\xi-1)^2x^2 - 4(\xi-1)^2x - 12(\xi+1)^2)^2}, \end{aligned}$$

$$(3.24) \quad \begin{aligned} \frac{\partial F_\xi}{\partial s_2}(g) &= \frac{8(\xi^2+\xi+1)((\xi^2+2\xi-3)x^4 + (-11\xi^2-18\xi+29)x^3}{(x-4)(x-1)((\xi-1)^2x^2 - 4(\xi-1)^2x - 12(\xi+1)^2)^2} \\ &\quad + \frac{(24\xi^2+24\xi-96)x^2 + (-28\xi^2-8\xi+84)x + 32\xi^2-32}{(x-4)(x-1)((\xi-1)^2x^2 - 4(\xi-1)^2x - 12(\xi+1)^2)^2}. \end{aligned}$$

The last three equations, namely (3.22), (3.23) and (3.24), have been computed with Maple.

The last piece of information we need to compute $f'_\xi(0)$ is the Ricci flow equation at time 0, that is, $g'_\xi(0) = (t'(0), s'_0(0), s'_1(0), s'_2(0))$, which is given by

$$\begin{aligned} t'(0) &= -\frac{(x^4 - 8x^3 + 16x^2)\left(\frac{(\xi+1)^2}{x^2} + \xi^2 + 1\right)}{3(\xi^2 + \xi + 1)}, \\ s'_0(0) &= -8 + \frac{(\xi + 1)^2(4 - x)}{\xi^2 + \xi + 1} - 2x^2, \\ s'_1(0) &= -12 + \frac{\xi^2(-x^2 + 4x)}{\xi^2 + \xi + 1} + 2x, \\ s'_2(0) &= -12 + \frac{-x^2 + 4x}{\xi^2 + \xi + 1} + 2x. \end{aligned}$$

At this stage we have all the materials to compute $f'_\xi(0) = \langle \nabla_g F_\xi, g'_\xi(0) \rangle$. Using Maple we obtain:

$$(3.25) \quad f'_\xi(0) = \frac{K(\xi, x)}{x(x-4)(x-1)((\xi-1)^2x^2 - 4(\xi-1)^2x - 12(\xi+1)^2)^2},$$

where

$$\begin{aligned} K(\xi, x) &= 40(\xi-1)^2 \left(\xi^2 + \frac{4}{5}\xi + 1 \right) x^7 - 568(\xi-1)^2 \left(\xi^2 + \frac{60}{71}\xi + 1 \right) x^6 \\ &\quad + (2960\xi^4 - 3552\xi^3 + 416\xi^2 - 3552\xi + 2960)x^5 \\ &\quad + (-7664\xi^4 + 6912\xi^3 - 2336\xi^2 + 6912\xi - 7664)x^4 \\ &\quad + (9856\xi^4 - 3968\xi^3 + 5120\xi^2 - 3968\xi + 9856)x^3 \\ &\quad + (-5312\xi^4 + 6144\xi^3 + 10624\xi^2 + 6144\xi - 5312)x^2 \\ &\quad + 256(\xi^2 - 50\xi + 1)(\xi + 1)^2x + 6144\xi(\xi + 1)^2. \end{aligned}$$

Let us analyze the sign of the quotient in (3.25). First, note that the denominator is positive for any $x < 1$. As for the numerator, we compute that the left-sided limit

$$\lim_{x \rightarrow 1^-} K(\xi, x) = -432(\xi^2 - 1)^2$$

is negative for all $\xi \in (0, 1)$. It follows that for each $\xi \in (0, 1)$ there is some $x = x(\xi) < 1$ sufficiently close to 1 such that $K(\xi, x) < 0$. Hence, we have found x for which $f'_\xi(0) < 0$, as desired.

In the case $\xi = 1$, numerator and denominator have $768(x-1)$ as a common factor. When simplifying, one arrives to the expression

$$f'_1(0) = \frac{x^4 + 6x^3 - 16x^2 - 32x + 32}{3x(x-4)}.$$

Of course, this coincides with the expression (3.9) for $f'_1(0)$ obtained by hand in the proof of Theorem 2.8, where we have shown that $f'_1(0) < 0$ if $x \in (0.8, 1)$.

4. THE BERGER SPACE

Recall that the Berger space B^{13} is the homogeneous space $\mathrm{SU}_5/\mathrm{Sp}_2\mathrm{S}^1$. The subgroup $\mathrm{Sp}_2\mathrm{S}^1$ is defined as the image of $\mathrm{Sp}_2 \times \mathrm{S}^1$ under the homomorphism

$$\iota : \mathrm{SU}_4 \times \mathrm{S}^1 \rightarrow \mathrm{U}_4 \subset \mathrm{SU}_5, \quad \iota(A, \lambda) = \lambda A.$$

Endow SU_5 with the bi-invariant metric $\langle\langle X, Y \rangle\rangle = -\frac{1}{2} \mathrm{Re} \mathrm{tr}(XY)$, for $X, Y \in \mathfrak{su}_5$. Let \mathfrak{p} be the orthogonal complement of $\mathfrak{sp}_2 \oplus \mathbb{R}$, the Lie algebra of $\mathrm{Sp}_2\mathrm{S}^1$. The space \mathfrak{p} decomposes as $\mathfrak{p}_1 \oplus \mathfrak{p}_2$, where \mathfrak{p}_1 denotes the orthogonal complement of \mathfrak{sp}_2 in \mathfrak{su}_4 and \mathfrak{p}_2 is the orthogonal complement of $\mathfrak{su}_4 \oplus \mathbb{R}$ in \mathfrak{su}_5 . As noted in [Pue99, Lemma 7.9], all SU_5 -invariant metrics are of the form

$$\langle \cdot, \cdot \rangle_{x_1, x_2} = x_1 \langle\langle \cdot, \cdot \rangle\rangle|_{\mathfrak{p}_1} + x_2 \langle\langle \cdot, \cdot \rangle\rangle|_{\mathfrak{p}_2}.$$

Püttmann has shown that $\langle \cdot, \cdot \rangle_{x_1, x_2}$ has positive curvature if and only if $x_1 < 2x_2$, see [Pue99, Corollary 7.18]. We will show that the $\mathrm{sec} > 0$ metrics corresponding to $x_1 = 2 - \varepsilon$, $x_2 = 1$ evolve under the Ricci flow to metrics with non-positively curved planes.

By dimensional reasons, \mathfrak{p}_1 and \mathfrak{p}_2 are inequivalent as $\mathrm{Ad}(\mathrm{Sp}_2\mathrm{S}^1)$ -representations. It follows that the Ricci tensor of $\langle \cdot, \cdot \rangle_{x_1, x_2}$ has a diagonal form

$$(4.1) \quad \mathrm{Ric} = x_1 r_1 \langle\langle \cdot, \cdot \rangle\rangle|_{\mathfrak{p}_1} + x_2 r_2 \langle\langle \cdot, \cdot \rangle\rangle|_{\mathfrak{p}_2}.$$

The eigenvalues r_1, r_2 can be computed using Formula (2.4). This has been done by Rodionov in [Ro92, Corollary 9.2], and for the metrics $\langle \cdot, \cdot \rangle_{x_1, 1}$ with $x_2 = 1$ he obtained:

$$r_1 = \frac{8 + x_1^2}{x_1}, \quad r_2 = \frac{5(8 - x_1)}{4}.$$

We are ready to prove Theorem B. Let $g(\ell) = (x_1(\ell), x_2(\ell))$ the Ricci flow of the initial metric $g(0) = (2, 1)$, which is determined by the system

$$(4.2) \quad \begin{cases} x'_1 = -2r_1 x_1, \\ x'_2 = -2r_2 x_2. \end{cases}$$

This system has a unique solution in an open interval $(-L, L)$ for some $L > 0$. We claim that

$$(4.3) \quad \left. \frac{d}{d\ell} \left(\frac{x_1(\ell)}{2x_2(\ell)} \right) \right|_{\ell=0} > 0.$$

This implies that there exists $\bar{L} > 0$ with $\bar{L} \leq L$ such that $x_1(\ell) < 2x_2(\ell)$ for all $\ell \in (-\bar{L}, 0)$ and $x_1(\ell) > 2x_2(\ell)$ for all $\ell \in (0, \bar{L})$. This proves Theorem B.

It remains to prove (4.3). Dropping ℓ from the equations we have

$$\frac{d}{d\ell} \left(\frac{x_1}{2x_2} \right) = \frac{x_1'x_2 - x_1x_2'}{2x_2^2} = \frac{-2r_1x_1x_2 + 2r_2x_1x_2}{2x_2^2} = \frac{-9x_1^2 - 32x_2^2 + 40x_1x_2}{4x_2^3},$$

where we have used (4.2). At $\ell = 0$ we have $(x_1, x_2) = (2, 1)$, so $r_1 = 6$ and $r_2 = \frac{15}{2}$ by (4.1). It follows that the derivative equals 3.

Remark 4.1. Note that Theorems 2.7 and B can be alternatively proven via the fact that both $W_{1,1}^7$ and B^{13} admit precisely two Einstein metrics, one with positive sectional curvature and one with planes with negative sectional curvature, see [Ni04, Lemma 2] for $W_{1,1}^7$ (also Subsection 3.3 and [Pue99, Lemmas 6.1] for the sign of sectional curvature) and [Ro92, Theorem 3.1] for B^{13} .

Consider the normalized Ricci flow of the two parameter metrics (t, t, s, s) on $W_{1,1}^7$ and the metrics (x_1, x_2) on B^{13} . In both cases the normalized Ricci flow is then given by one equation in one variable. Then there exists a normalized Ricci flow solution $g(\ell)$, defined on the maximal time interval (L_{\min}, L_{\max}) with the property that

$$\lim_{\ell \rightarrow L_{\min}} g(\ell) = g_1,$$

$$\lim_{\ell \rightarrow L_{\max}} g(\ell) = g_2,$$

where g_1 is the Einstein metric on $W_{1,1}^7$ or B^{13} with $\sec > 0$ and g_2 is the Einstein metric which has planes with negative curvature.

REFERENCES

- [AN16] Abiev, Nurlan Abievich; Nikonorov, Yurii Gennadievich. *The evolution of positively curved invariant Riemannian metrics on the Wallach spaces under the Ricci flow*. Ann. Global Anal. Geom. **50** (2016), no. 1, 65–84.
- [AW75] Aloff, Simon; Wallach, Nolan R. *An infinite family of distinct 7-manifolds admitting positively curved Riemannian structures*. Bull. Amer. Math. Soc. **81** (1975), 93–97.
- [Be87] Besse, Arthur L. *Einstein manifolds*. Reprint of the 1987 edition. Classics in Mathematics. Springer-Verlag, Berlin, 2008.
- [BK23] Bettiol, Renato G.; Krishnan, Anusha M. *Ricci flow does not preserve positive sectional curvature in dimension four*. Calc. Var. Partial Differential Equations **62** (2023), no. 1, Paper No. 13, 21.
- [BW07] Böhm, Christoph; Wilking, Burkhard. *Nonnegatively curved manifolds with finite fundamental groups admit metrics with positive Ricci curvature*. Geom. Funct. Anal. **17** (2007), no. 3, 665–681.
- [BW08] ———, *Manifolds with positive curvature operators are space forms*. Ann. of Math. (2) **167** (2008), no. 3, 1079–1097.
- [CW15] Cheung, Man-Wai; Wallach, Nolan R. *Ricci flow and curvature on the variety of flags on the two dimensional projective space over the complexes, quaternions and octonions*. Proc. Amer. Math. Soc. **143** (2015), no. 1, 369–378.
- [GZ24] González-Álvaro, David; Zarei, Masoumeh. *Positive intermediate curvatures and Ricci flow*. To appear in Proc. Amer. Math. Soc. <https://doi.org/10.1090/proc/16752>.
- [Ha82] Hamilton, Richard S. *Three-manifolds with positive Ricci curvature*. J. Differential Geometry **17** (1982), no. 2, 255–306.

- [Ha86] ———, *Four-manifolds with positive curvature operator*. J. Differential Geom. **24** (1986), no. 2, 153–179.
- [La13] Lauret, Jorge. *Ricci flow of homogeneous manifolds*. Math. Z. **274** (2013), no. 1-2, 373–403.
- [Ni14] Ni, Lei. *Ricci flow and manifolds with positive curvature*. Symmetry: representation theory and its applications, 491–504, Progr. Math., 257, Birkhäuser/Springer, New York, 2014.
- [Ni04] Nikonorov, Yu. G., *Compact homogeneous Einstein 7-manifolds*. Geom. Dedicata **109** (2004), 7–30.
- [Pue99] Püttmann, Thomas. *Optimal pinching constants of odd-dimensional homogeneous spaces*. Invent. Math. **138** (1999), no. 3, 631–684.
- [Ri14] Richard, Thomas. *Curvature cones and the Ricci flow*. Séminaire de théorie spectrale et géométrie 31 (2012-2014): 197-220.
- [Ro92] Rodionov, E. D. *Homogeneous Einstein metrics on an exceptional Berger space*.(Russian) ; translated from Sibirsk. Mat. Zh. **33** (1992), no. 1, 208–211, 222 Siberian Math. J. **33** (1992), no. 1, 171–174.
- [Wa82] Wang, McKenzie Y. *Some examples of homogeneous Einstein manifolds in dimension seven*. Duke Math. J. **49** (1982), no. 1, 23–28.
- [WZ86] Wang, McKenzie Y.; Ziller, Wolfgang. *Existence and nonexistence of homogeneous Einstein metrics*. Invent. Math. **84** (1986), no. 1, 177–194.
- [Wi13] Wilking, Burkhard. *A Lie algebraic approach to Ricci flow invariant curvature conditions and Harnack inequalities*. J. Reine Angew. Math. 679 (2013), 223–247.
- [Zi07] Ziller, Wolfgang. *Examples of Riemannian manifolds with non-negative sectional curvature*. Surveys in differential geometry. Vol. XI, 63–102, Surv. Differ. Geom., 11, Int. Press, Somerville, MA, 2007.

ETSI DE CAMINOS, CANALES Y PUERTOS, UNIVERSIDAD POLITÉCNICA DE MADRID, SPAIN.

Email address: david.gonzalez.alvaro@upm.es

MATHEMATISCHES INSTITUT, UNIVERSITÄT MÜNSTER, EINSTEINSTR. 62, 48149 MÜNSTER, GERMANY.

Email address: masoumeh.zarei@uni-muenster.de