Algebraic structures on parallelizable manifolds

Sergey Grigorian

School of Mathematical & Statistical Sciences University of Texas Rio Grande Valley Edinburg, TX 78539 USA

March 22, 2024

Abstract

In this paper we explore algebraic and geometric structures that arise on parallelizable manifolds. Given a parallelizable manifold L, there exists a global trivialization of the tangent bundle, which defines a map $\rho_p: \mathfrak{l} \longrightarrow$ $T_p\mathbb{L}$ for each point $p \in \mathbb{L}$, where l is some vector space. This allows us to define a particular class of vector fields, known as fundamental vector fields, that correspond to each element of l. Furthermore, flows of these vector fields give rise to a product between elements of l and L, which in turn induces a local loop structure (i.e. a non-associative analog of a group). Furthermore, we also define a generalization of a Lie algebra structure on l. We will describe the properties and examples of these constructions.

Contents

1 Introduction

The tangent bundle is a fundamental aspect of smooth manifolds, and its global trivialization, or absolute parallelism, has been a key area of study in differential

geometry since the field's emergence in the early 20th century [\[4,](#page-30-1) [5\]](#page-30-2). The study of parallelizable manifolds, as smooth manifolds with a trivial tangent bundle are now generally referred to, has been continuing since then in several directions. One approach, which is more topological in nature, reframed the question in terms of stably parallelizable manifolds, i.e. manifolds for which the stable tangent bundle is trivial. Given a real smooth manifold M with tangent bundle TM, and a trivial rank 1 bundle ε over M (i.e. $M \times \mathbb{R}$), the stable tangent bundle is $TM \oplus \varepsilon$. The conditions for a manifold to be stably parallelizable (also known as a π -manifold) in low dimensions can be expressed in terms of vanishing of certain characteristic classes [\[7\]](#page-30-3). It also follows from [\[2\]](#page-30-4) that in dimensions 1, 3, and 7, manifolds are parallelizable if and only if they are stably parallelizable. Another approach, which stems from the origins of Riemannian geometry, involves the study of flat metric connections on Riemannian manifolds. Indeed, the existence of a flat metric connection allows to parallel transport a frame from a single point to the entire manifold, and thus obtain a trivialization of the tangent bundle. The converse is also trivially true. Generally such connections will admit torsion and then questions about classification of parallelizable manifolds become related to the properties of torsion of flat metric connections. In [\[4\]](#page-30-1), Cartan and Schouten have shown initiated this approach by showing that a Riemannian manifold with a flat metric connection and totally skew-symmetric torsion is either a compact simple Lie group or the 7-sphere $S⁷$, depending on whether the torsion is parallel or not. A similar classification in the pseudo-Riemannian case has been obtained by Wolf [\[16,](#page-31-0) [17\]](#page-31-1), again under the assumption of totally skew-symmetric torsion. A more modern treatment of these approaches has been given in [\[1\]](#page-30-5). However, surprisingly, despite significant efforts from different points of view, there is still no full classification of parallelizable manifolds and shows that there is still a need to explore properties of parallelizable manifolds.

This paper builds upon these foundational concepts to explore new algebraic structures on parallelizable manifold. These structures extend the traditional notions of Lie algebras and Lie groups, providing a broader perspective on the geometric and algebraic properties of these manifolds. Indeed, Lie groups form one of the most well-known classes of parallelizable manifolds and are of course characterized by a globally defined smooth associative product and a Lie algebra structure on the tangent space at identity. The associativity property may be relaxed to consider smooth loops [\[6\]](#page-30-6). The 7-sphere from [\[4\]](#page-30-1) is an example of a smooth loop, when regarded as the set of unit octonions, and is of course parallelizable. The smooth product allows to identify any tangent space with the tangent space at identity, providing a global trivialization of the tangent bundle. This also allows to define an bracket algebra on the tangent space at identity, however, this is in general no longer a Lie algebra. As shown in [\[6\]](#page-30-6), on a smooth loop we may define a family of brackets, defined for each point of the loop, leading to a bracket function defined on the loop. The non-constant nature of this function then leads to a non-trivial right hand side in the Jacobi identity. On Lie group, the bracket of left- or right-invariant vector fields is itself left- or right-invariant, respectively, and hence only a unique bracket is

defined.

In this paper, we expand some of the findings from [\[6\]](#page-30-6), but for arbitrary parallelizable manifolds. Indeed, several properties of Lie groups and smooth loops only depend on the trivialization of the tangent bundle, which is a weaker property than the existence of a global structure.

In Section [2,](#page-3-0) our exploration begins with a detailed examination of the global trivialization of tangent bundles, a defining characteristic of parallelizable manifolds. This global trivialization allows us to define fundamental vector fields, which are parallel vector fields with respect to the trivial connection. Integral curves of these vector lead a local loop structure, a non-associative analog of a group structure. This structure enriches the manifold's algebraic framework and opens new avenues for studying its geometric properties. Furthermore, we introduce a generalization of the Lie algebra structure on parallelizable manifolds. This generalization provides a new perspective on the algebraic underpinnings of these manifolds and offers new insights into their properties. The interplay between the geometric and algebraic aspects of parallelizable manifolds, enriched by these Lie-like structures, is a key focus of this paper. It should be noted that in the 1960's, Kikkawa [\[10,](#page-31-2) [11\]](#page-31-3) has similarly defined local loop structures based on geodesics of arbitrary affine connections. However, unlike in the parallelizable case, such an approach does not allow for a richer algebra structure.

The structure in Section [2](#page-3-0) is defined by a parallelizable manifold together with a fixed trivialization of the tangent bundle. We denote this by the *parallelized manifold* triple (L, I, ρ) , where L is a smooth manifold, l a vector space which is the model fiber for the tangent bundle, and ρ is a smooth family of linear isomorphisms from I to each tangent space $T_p \mathbb{L}$. We use the trivialization ρ to define fundamental vector fields on \mathbb{L} , for each $s \in \mathbb{L}$ given by $\rho_s(\xi)$ for some $\xi \in I$. These are precisely the parallel vector fields for the flat connection defined by the trivialization. We also use ρ family of local loop structures \circ_s on I and corresponding bracket algebra structures $b^{(s)}$, together with a family of trilinear forms $a^{(s)}$ that are defined from differentials of $b^{(s)}$ Theorem [2.24](#page-9-0) gives an analogue of the Maurer-Cartan structure equation, while Theorem [2.26](#page-10-0) relates the brackets as infinitesimal commutators of the products \circ_s . As discussed above for the case of smooth loops, the bracket function is in general non-constant on L, and its differential is related to a skew-symmetrization of infinitesimal associators of the local product on l.

In Section [3](#page-14-0) we then define morphisms of parallelized manifolds and study how the automorphisms of parallelized manifolds interact with the algebraic operations defined in Section [2.](#page-3-0) In Theorem [3.8](#page-17-0) we prove that the automorphism group of (L, I, ρ) is a finite-dimensional Lie subgroup of the diffeomorphism group of L.

As a particular example, in Section [4](#page-19-0) we consider explicit trivializations of products of spheres, as given in [\[3,](#page-30-7) [14,](#page-31-4) [15\]](#page-31-5). As shown by Kervaire in [\[9\]](#page-31-6), any product of spheres that contains an odd-dimensional sphere, is parallelizable. However, we consider a simpler examples of $S^m \times S^1$ and $S^m \times N$, where m is arbitrary and N is any parallelizable manifold. The parallelizability of these spaces follows immediately from the fact that any sphere is stably parallelizable.

We find that in the case of $\mathbb{L} = S^m \times S^1$, the standard trivialization yields a family of bracket algebras $b^{(x)}$, for $x \in S^m$, each of which is isomorphic to the semidirect sum Lie algebra $\mathbb{R}^n \oplus_S \mathbb{R}$. Moreover, considering the triple product $a^{(x)}$, the algebraic structure $(1, a^{(x)})$ is that of a Lie triple system [\[8\]](#page-30-8). We also explicitly compute the local product \circ_s on l in certain special cases and find that the automorphism group $\Psi(\mathbb{L}) \cong SO(n+1) \times U(1)$.

In the more general case of $\mathbb{L} = S^m \times N$, we compute the brackets and the Jacobi identity, finding that for instance, in the case of $N = Sⁿ \times S¹$, for some values of $s \in \mathbb{L}$, then bracket $b^{(s)}$ does not satisfy the Jacobi identity, so generally, parallelizable manifolds do indeed induce an algebraic structure that is more general than a Lie algebra or a Lie triple system.

Overall, we see that the framework presented in this paper allows to associate certain algebraic structures to parallelized manifolds. This paves the way towards using such algebraic invariants to study and classify parallelizable manifolds. Moreover, this approach is also easily adaptable to the broader class of stably parallelizable manifolds as well, as the example of $S^m \times S^1$ shows.

Acknowledgements

This work was supported by the National Science Foundation [DMS-1811754].

2 Parallelized manifolds

Suppose $\mathbb L$ is a connected smooth real *n*-dimensional manifold, with a trivializable tangent bundle $T\mathbb{L} \cong \mathbb{L} \times \mathbb{R}^n$. Now given an n-dimensional real vector space l, the global trivialization of TL induces a diffeomorphism $\rho : \mathbb{L} \times I \longrightarrow TL$, such that

$$
\mathbb{L} \times \mathfrak{l} \ni (p, \xi) \mapsto \rho_p(\xi) \in T_p \mathbb{L},\tag{2.1}
$$

with $\rho_p: \mathfrak{l} \longrightarrow T_p \mathbb{L}$ being a linear isomorphism for each $p \in \mathbb{L}$.

Definition 2.1 The fundamental vector field for $\xi \in I$ of TL is defined to be the smooth vector field $\rho(\xi) \in \Gamma(T\mathbb{L})$.

In particular, a choice of basis on $\mathfrak l$ induces a global frame of $T\mathbb L$, or equivalently a section of the corresponding frame bundle $F\mathbb{L}$. A change of trivialization is equivalent to a $GL(\mathbb{R},n)$ gauge transformation of FL , while a change of basis of l corresponds to the right action of $GL(\mathbb{R}, n)$ on the principal bundle FL .

More explicitly, suppose ${X_i}_{i=1}^n$ is a frame of TL. Then, assuming we have a basis ${e_i}_{i=1}^{\hat{n}}$ on l, define the corresponding trivialization ρ so that $\rho(e_i) = X_i$ and hence

$$
\mathbb{L} \times \mathfrak{l} \ni (p, \xi) \mapsto \xi^i \left. X_i \right|_p \in T_p \mathbb{L}.
$$

Definition 2.2 The triple (L, L, ρ) , as above, will be called a parallelized manifold, *i.e.* a parallelizable manifold with a fixed trivialization.

Remark 2.3 The above definition is almost equivalent to the notion of absolute parallelism. Absolute parallelism is defined as a smooth family of linear isomorphisms $\phi_{pq}: T_q\mathbb{L} \longrightarrow T_p\mathbb{L}$ for each pair of points $p, q \in \mathbb{L}$. Given an absolute parallelism and fixing one of the tangent spaces as the vector space $\mathfrak l$ gives the parallelized manifold definition. Conversely, given the linear maps ρ_p from [\(2.1\)](#page-3-1), an absolute parallelism is defined as $\phi_{pq} = \rho_p \rho_q^{-1}$.

Example 2.4 Any Lie group can be parallelized via left translation or right translation. In that case, l corresponds to the Lie algebra, and ρ is either the left or right translation map.

Example 2.5 Any smooth loop (i.e. a smooth quasigroup with identity and smooth products and quotients) is parallelizable, in the same way a Lie group $[6]$. In that case, I corresponds to the tangent algebra at identity. A specific example is S^7 , regarded as the loop of unit octonions.

Example 2.6 Any orientable 3-dimensional manifold is parallelizable. For example, $S^2 \times S^1$ and 3-dimensional lens spaces $L(p;q)$, as well as S^3 , but this is of course also a Lie group.

Example 2.7 A 4-manifold is parallelizable if and only if the Stiefel-Whitney $classes w_1, w_2$ vanish, the Euler characteristic vanishes, and the first Pontryagin class vanishes.

Example 2.8 Products of parallelizable manifolds are parallelizable.

Example 2.9 As shown by Kervaire in [\[9\]](#page-31-6), a product of spheres, where at least one of the spheres is odd-dimensional, is parallelizable. Explicit parallelizations of products of spheres have been demonstrated in [\[14,](#page-31-4) [15\]](#page-31-5).

Remark 2.10 In general, parallelizability of a manifold is a property of the smooth structure, rather than the underlying topological space. Indeed, as shown by Milnor [\[12,](#page-31-7) Corollary 1], one may have two smooth manifolds that are homeomorphic, but one is parallelizable, while the other one is not. However, in certain situations, the parallelizability does reduce to topological considerations. A manifold is known as stably parallelizable if its stable tangent bundle $TM \oplus \varepsilon$, where ε is a trivial rank 1 bundle, is trivial. Such manifolds are also known as framed manifolds or π -manifolds. Stable parallelizability is in some cases equivalent to the vanishing of certain characteristic classes [\[7\]](#page-30-3). As shown in [\[2\]](#page-30-4), an n-dimensional stably parallelizable manifold is parallelizable if and only if S^n is parallelizable. This of course happens for $n = 1, 3, 7$. Hence in these dimensions, parallelizability is equivalent to stable parallelizability.

To investigate properties of parallelizable manifolds using approaches from Riemannian geometry, we may use ρ to define a metric and a connection on \mathbb{L} .

Definition 2.11 Given the triple $(\mathbb{L}, \mathfrak{l}, \rho)$ and any inner product $\langle \rangle$ on \mathfrak{l} , define the metric g on $\mathbb L$ as the pullback metric with respect to ρ^{-1} , i.e. for $s \in \mathbb L$, and $X_s, Y_s \in T_s \mathbb{L},$

$$
g_s(X_s, Y_s) = \langle \rho_s^{-1}(X_p), \rho_s^{-1}(Y_p) \rangle.
$$
 (2.2)

Definition 2.12 Given the triple $(\mathbb{L}, \mathfrak{l}, \rho)$, define the flat connection $\nabla = (\rho^{-1})^* d$, so that

$$
\nabla = \rho \circ d \circ \rho^{-1}.
$$
\n(2.3)

Lemma 2.13 The metric and the flat connection satisfy the following properties.

- 1. ∇ is a metric connection with respect to q.
- 2. Every fundamental vector field has constant norm.

Proof. Both of these properties follow immediately from the definitions.

1. Suppose X and Y are two vector fields, then

$$
g(\nabla X, Y) + g(X, \nabla Y) = g(\rho(d(\rho^{-1}X)), Y)
$$

+
$$
+g(X, \rho(d(\rho^{-1}Y)))
$$

= $\langle d(\rho^{-1}X), \rho^{-1}Y \rangle + \langle \rho^{-1}X, d(\rho^{-1}Y) \rangle$
= $d(\langle \rho^{-1}X, \rho^{-1}Y \rangle)$
= $d(g(X, Y)),$

hence ∇ is metric.

2. Suppose $X = \rho(\xi)$ is a fundamental vector field, then it is nowhere zero, and

$$
d\left(\|X\|^2\right) = d(g(\rho(\xi), \rho(\xi))) = d(\langle \xi, \xi \rangle) = 0,
$$

hence $||X||$ is constant.

Remark 2.14 Given a vector field $X \in \Gamma(T\mathbb{L})$ and $s \in \mathbb{L}$, consider the maximal integral curve $\gamma_{X,s}$ on L of the vector field X through a point $s \in \mathbb{L}$. If \mathbb{L} is compact, then every vector field on L is complete, and hence $\gamma_{X,s}(t)$ is defined for all t. If $\mathbb L$ is non-compact, consider the global frame $\{X_i\}$ on $\mathbb L$. Each X_i is nowhere-vanishing, and given an arbitrary complete Riemannian metric on L (which always exists if $\mathbb L$ is connected (13)) we can rescale each of these vector fields to be of unit norm everywhere. Then, as it is well-known, unit norm vector fields on a complete Riemannian manifold are complete. Hence, by normalizing the global frame, we may assume that the fundamental vector fields are complete. Thus, for non-compact manifolds, without loss of generality, we'll consider only complete trivializations, i.e. those for which the fundamental vector fields are complete. Then, if X is fundamental, then $\gamma_{X,s}(t)$ is defined for all t. We will assume that $\mathbb L$ is connected or compact.

\blacksquare

Let $\xi \in \mathfrak{l}$ and define the flow diffeomorphisms $\Phi_{\xi,t} : \mathbb{L} \longrightarrow \mathbb{L}$ of ξ via

$$
\Phi_{\xi,t}(s) = \gamma_{\rho(\xi),s}(t),\tag{2.4}
$$

with $\gamma_{\rho(\xi),s}(0) = s$. By definition of integral curves, $\Phi_{\xi,t}(s)$ is the solution of the following initial value problem:

$$
\begin{cases} \frac{dp(t)}{dt} = \rho(\xi)|_{p(t)} \\ p(0) = s \end{cases} . \tag{2.5}
$$

Since $\rho(\xi)$ is assumed to be complete, the maps $\Phi_{\xi,t}$ are defined for all t, and in particular, for each $a \in \mathbb{R}$, $\Phi_{a\xi,t} = \Phi_{\xi,at}$, and this motivates the following definition.

Definition 2.15 The product operation μ on l and L is the map $\mu : \mathbb{I} \times \mathbb{L} \longrightarrow \mathbb{L}$ defined by

$$
\mu(\xi, s) = \Phi_{\xi, 1}(s), \tag{2.6}
$$

where $\xi \in I$, $s \in L$. In particular, we will denote this product by $\xi \cdot s$. Define the left and right product maps:

- 1. For each $\xi \in \mathfrak{l}, L_{\xi} : \mathbb{L} \longrightarrow \mathbb{L}$
- 2. For each $s \in L$, $R_s : \mathfrak{l} \longrightarrow \mathbb{L}$.

Example 2.16 Suppose L is a Lie group and ρ is the right-invariant trivialization of the tangent bundle. Then, l is just the tangent space at identity, i.e. the corresponding Lie algebra, and $\xi \cdot s = \exp{(\xi)} s$, where $\exp is$ the standard Lie algebra exponential map.

Lemma 2.17 Let $\xi, \eta \in I$. The product operation has the following properties:

- 1. For $t_1, t_2 \in \mathbb{R}$, $(t_1\xi) \cdot ((t_2\xi) \cdot s) = ((t_1 + t_2)\xi) \cdot s$.
- 2. For any $s \in L$, $0 \cdot s = s$.
- 3. L_ξ is a diffeomorphism with $L_{\xi}^{-1} = L_{-\xi}$.
- 4. $dR_s|_{0} = \rho_s$.
- 5. For each $s \in L$, R_s is a local diffeomorphism, and hence the right quotient R_s^{-1} is defined in some neighborhood of s.
- 6. Suppose p (t) is a curve in $\mathbb L$ with $p(0) = p$. Assuming that p/s is defined, we have

$$
\frac{d}{dt}R_{t\xi \cdot s}^{-1}(p(t))\bigg|_{t=0} = \frac{d}{dt}p(t)/s\bigg|_{t=0} - \frac{d}{dt}(p/s)\circ_s(t\xi)\cdot\bigg|_{t=0}
$$
 (2.7)

Proof. The first three properties immediately follow from the definition and the properties of the flow diffeomorphism $\Phi_{\xi,t}$. For the fourth item, let $\xi \in \mathfrak{l}$ and consider the straight line from the origin $t\xi$. Then,

$$
dR_s|_0(\xi) = \frac{d}{dt}R_s(t\xi)\Big|_{t=0}
$$

=
$$
\frac{d}{dt}\Phi_{\xi,t}(s)\Big|_{t=0}
$$

=
$$
\rho_s(\xi),
$$

by definition of R_s .

This shows that $dR_s|_0 = \rho_s$, hence is an isomorphism, and thus by the Inverse Function Theorem is a local diffeomorphism.

Now for sufficiently small t , consider

$$
p(t) = \left(R_{t\xi \cdot s}^{-1} (p(t))\right) \cdot (t\xi \cdot s)
$$

Then, differentiating both sides, and noting that $R_{t\xi \cdot s}^{-1}(p(t))\Big|_{t=0} = p/s$, we have

$$
\frac{d}{dt}p(t)\Big|_{t=0} = \frac{d}{dt}\left(R_{t\xi \cdot s}^{-1}(p(t))\right) \cdot (t\xi \cdot s)\Big|_{t=0}
$$

$$
= \frac{d}{dt}\left(R_{t\xi \cdot s}^{-1}(p(t))\right) \cdot s\Big|_{t=0} + \frac{d}{dt}\left((p/s) \cdot (t\xi \cdot s)\right)\Big|_{t=0}
$$

Applying ρ_s^{-1} to both sides, and noting that $dR_s|_0 = \rho_s$, we find

$$
\frac{d}{dt}\left(R_{t\xi\cdot s}^{-1}\left(p\left(t\right)\right)\right)\bigg|_{t=0} = \frac{d}{dt}p\left(t\right)/s\bigg|_{t=0} - \frac{d}{dt}\left(p/s\right)\circ_s\left(t\xi\right)\cdot\bigg|_{t=0}
$$

Remark 2.18 Lemma [2.17](#page-6-0) shows that for each $s \in \mathbb{L}$, we may define a local loop structure on l. Indeed, suppose $\xi, \eta \in I$ are in a sufficiently small neighborhood of $0 \in I$. Then, define

$$
\eta \circ_s \xi = (\eta \cdot (\xi \cdot s)) / s \in \mathfrak{l}.\tag{2.8}
$$

With respect to this product, we see that $0 \in I$ is a two-sided identity element. The right quotient with respect to \circ_s is defined by

$$
\xi/s\eta = (\xi \cdot s) / (\eta \cdot s), \qquad (2.9)
$$

while the left quotient is given by

$$
\xi \setminus s\eta = (\xi \setminus (\eta \cdot s)) / s. \tag{2.10}
$$

Equivalently, we may define a product on a neighborhood of s. Given p, q in a sufficiently small neighborhood of $s \in \mathbb{L}$, define

$$
p \circ_s q = (p/s) \cdot q. \tag{2.11}
$$

Note that if $p = \xi \cdot s$ and $q = \eta \cdot s$, then $p \circ_s q = (\xi \circ_s \eta) \cdot s$. Similarly, the quotients are also defined.

Lemma 2.19 The product \circ_s is power-associative. In particular, for any $\xi \in I$, the powers $\xi^n = n\xi$ are defined unambiguously and independently of s.

Proof. Let $\xi \in I$. Firstly, for real numbers t_1 and t_2 consider

$$
(t_1\xi) \circ_s (t_2\xi) = ((t_1\xi) \cdot ((t_2\xi) \cdot s)) / s = (t_1 + t_2) \xi.
$$

Hence $\xi \circ_s \xi = 2\xi$. Now, for third powers

$$
(\xi \circ_s \xi) \circ_s \xi = (2\xi) \circ_s \xi = 3\xi = \xi \circ_s (\xi \circ_s \xi).
$$

By induction we can see that for any integer power $n, \xi^n = n\xi$.

Theorem 2.20 Suppose \mathbb{L} is connected and $s \in \mathbb{L}$, then any element $p \in \mathbb{L}$ can be written as $p = \xi_1 \cdot (\xi_2 \cdot ... (\xi_k \cdot s))$, for some finite sequence $\{\xi_1, ..., \xi_k\}$ in I .

Proof. Let $W \subset \mathbb{L}$ be the subset of elements of \mathbb{L} that can be represented as a product of s by some finite sequence of elements of l on the left. We will show that W is both closed and open, and hence $\mathbb{L} = W$.

First note that since for each $p \in \mathbb{L}$, R_p is a local diffeomorphism of a neighborhood of $0 \in I$ to a neighborhood of $p \in \mathbb{L}$, we see that there exists an open neighborhood V_p of p, where each $q \in V_p$ is given by $\eta \cdot p$ for some $\eta \in \mathbb{L}$.

Now let $w \in W$, then by the above, there exists an open neighborhood V_w of w, where each $q \in V_w$ is given by $\eta \cdot w$. In particular, $V_w \subset W$, and hence W is open.

To show that W is closed, consider a converging sequence $p_i \longrightarrow p$ with each $p_i \in W$. Then, for N large enough, we will have $p_N \in V_p$. In other words, we will have $p_N = \eta \cdot p$, for some $\eta \in I$. This implies that $p = (-\eta) \cdot p_N \in W$, and thus W is closed. \blacksquare

Remark 2.21 The above result shows that given a fixed $s \in \mathbb{L}$, any other element of $\mathbb L$ may be reached in a finite number of "steps", i.e. left multiplications by elements of $\mathfrak l$. Thus, we may define a discrete "distance" function $d(p, q)$ for points p, $q \in \mathbb{L}$ given by the minimum number of steps needed to reach q from p.

Given a trivialization ρ and fixing a point $s \in \mathbb{L}$, we induce a bracket on l. Let $\xi, \eta \in \mathfrak{l}$. Then,

$$
[\xi, \eta]^{(\rho, s)} = -\rho_s^{-1} ([\rho(\xi), \rho(\eta)]|_s). \tag{2.12}
$$

Note that here we have the negative sign to be compatible with [\[6\]](#page-30-6) and also to follow the same convention as right-invariant vector fields on Lie groups. As before, if $\{X_i\}_{i=1}^n$ is a frame of $T\mathbb{L}$, suppose the brackets of the frame elements are given by

$$
[X_i, X_j] = c^k_{ij} X_k,\tag{2.13}
$$

where c_{ij}^k are smooth F-valued functions on M. Then,

$$
[\xi, \eta]^{(\rho, s)} = -\rho_s^{-1} ([\xi^i X_i, \eta^j X_j]|_s). \tag{2.14}
$$

$$
= -\rho_s^{-1} \left(\xi^i \eta^j c_{ij}^k (s) X_k |_{s} \right), \tag{2.15}
$$

and therefore

$$
[\xi, \eta]^{(\rho, s)} = -\xi^i \eta^j c_{ij}^k (s) e_k
$$
\n(2.16)

Define the bracket function

$$
b^{(\rho)}: \mathbb{L} \longrightarrow \Lambda^2 \mathfrak{l}^* \otimes \mathfrak{l} \tag{2.17}
$$

given by

$$
b^{(\rho)}(s) (\xi, \eta) = [\xi, \eta]^{(\rho, s)}
$$

for any $s \in \mathbb{L}$ and $\xi, \eta \in \mathfrak{l}$. We can also define the ad map:

$$
\mathrm{ad}^{(\rho,s)}_\xi:\mathfrak{l}\longrightarrow\mathfrak{l}
$$

by

$$
\mathrm{ad}^{(\rho,s)}_\xi\left(\eta\right)=\left[\xi,\eta\right]^{(\rho,s)}.
$$

Definition 2.22 Let the Maurer-Cartan form $\theta^{(\rho)} \in \Omega^1(\mathbb{L}, \mathfrak{l})$ be an l-valued 1-form, defined by

$$
\left.\theta^{(\rho)}\right|_s(X) = \rho_s^{-1}(X) \tag{2.18}
$$

for any $X \in T_sM$, where $s \in M$. In particular, for any $\eta \in I$, given a fundamental vector field $\rho(\eta)$,

$$
\theta^{(\rho)}(\rho(\eta)) = \eta. \tag{2.19}
$$

In particular, we can now rewrite [\(2.12\)](#page-8-0) as

$$
b^{(\rho)}\left(\theta^{(\rho)}\left(X\right),\theta^{(\rho)}\left(Y\right)\right)=-\theta^{(\rho)}\left(\left[X,Y\right]\right). \tag{2.20}
$$

Remark 2.23 To simplify notation, we will drop (ρ) from b and θ . We still need to remember that these objects depend on the trivialization ρ , but once the trivialization is fixed, it is not necessary to keep writing it.

Theorem 2.24 (Structural Equation) The 1-form θ satisfies the following relation.

$$
d\theta - \frac{1}{2}b(\theta, \theta) = 0,\t(2.21)
$$

where $b(\theta, \theta)$ is the bracket of *l*-valued 1-forms such that for any $X, Y \in T_p \mathbb{L}$, $\frac{1}{2} b(\theta,\theta)|_{s}(X,Y) = [\theta(X),\theta(Y)]^{(s)}.$

Proof. It is sufficient to check this for fundamental sections of TL. Let $X = \rho(\xi)$ and $Y = \rho(\eta)$ for $\xi, \eta \in I$. Then from [\(2.18\)](#page-9-1),

$$
(d\theta)(X,Y) = X(\theta(Y)) - Y(\theta(X)) - \theta([X,Y])
$$

= X(\eta) - Y(\xi) - \theta([X,Y])
= b(\theta(X), \theta(Y)),

where we have used [\(2.20\)](#page-9-2) and the fact that η and ξ are constant.

Applying the exterior derivative to the structural equation, we find the generalization of the Jacobi identity:

Theorem 2.25 (Generalized Jacobi identity) The bracket b and the Maurer-Cartan form θ satisfy:

$$
b(\theta, b(\theta, \theta)) = (db)(\theta, \theta), \qquad (2.22)
$$

where exterior derivatives are implied.

Define the *skew-associator* map $a: \mathbb{L} \longrightarrow \Lambda^2 \mathfrak{l}^* \otimes \mathfrak{l}^* \otimes \mathfrak{l}$ such that for $s \in \mathbb{L}$ and $\xi, \eta, \gamma \in \mathfrak{l}$,

$$
a^{(s)}(\xi, \eta, \gamma) = d_{\rho(\gamma)}b(\xi, \eta)\Big|_p.
$$
\n(2.23)

Suppose $\xi, \eta, \gamma \in I$. Then, substituting $\rho(\xi), \rho(\eta), \rho(\gamma) \in \Gamma(T\mathbb{L})$ into [\(2.22\)](#page-10-1) and evaluating at a point $s \in \mathbb{L}$ gives

$$
\[\xi, [\eta, \gamma]^{(s)} \]^{(s)} + [\eta, [\gamma, \xi]^{(s)} \]^{(s)} + [\gamma, [\xi, \eta]^{(s)} \]^{(s)} = a^{(s)}(\xi, \eta, \gamma) \quad (2.24)
$$

$$
+ a^{(s)}(\eta, \gamma, \xi) + a^{(s)}(\gamma, \xi, \eta).
$$

We thus see that the obstruction to satisfying the Jacobi identity is precisely the cyclic permutation of a or equivalently db . In particular, if b is constant, then $\mathfrak l$ is a Lie algebra. In this case, the bracket is independent of p , and $\mathfrak l$ has a unique bracket that satisfies the Jacobi identity.

Both the bracket and the skew-associator maps can be related to the product ◦son l. In particular, as we see below, the bracket is the infinitesimal commutator of \circ_s and $a^{(s)}$ is a skew-symmetrization of the infinitesimal associator of \circ_s .

Theorem 2.26 Let $\xi, \eta, \gamma \in I$, and $s \in \mathbb{L}$, then

$$
[\xi, \eta]^{(s)} = \frac{d^2}{dt_1 dt_2} ((t_1 \xi) \circ_s (t_2 \eta) - (t_2 \eta) \circ_s (t_1 \xi)) \Big|_{t_1 = t_2 = 0}
$$
 (2.25a)

$$
a^{(s)}(\xi, \eta, \gamma) = [\xi, \eta, \gamma]^{(s)} - [\eta, \xi, \gamma]^{(s)},
$$
 (2.25b)

where

$$
[\xi, \eta, \gamma]^{(s)} = \left. \frac{d^3}{dt_1 dt_2 dt_3} \left((t_1 \xi) \circ_s (t_2 \eta) \circ_s (t_3 \gamma) \right) - \left((t_1 \xi) \circ_s (t_2 \eta) \right) \circ_s (t_3 \gamma) \right) \Big|_{t_1 = t_2 = t_3 = 0}
$$

.

Proof. By definition (2.12) , we have

$$
[\xi, \eta]^{(s)} = -\rho_s^{-1}([X, Y]|_s). \tag{2.26}
$$

where $X = \rho(\xi)$ and $Y = \rho(\eta)$ are vector fields on L. However, the bracket of two vector fields X and Y is given by

$$
[X,Y]_s = \frac{d}{dt} \left(\left(\Phi_{\xi,t}^{-1} \right)_* \left(Y_{\Phi_{\xi,t}(s)} \right) \right) \Big|_{t=0},\tag{2.27}
$$

where $\Phi_{\xi,t}$ is the flow generated by $X,$ and hence, $\Phi_{\xi,t}\left(s\right)=\left(t\xi\right)\cdot s.$ Therefore

$$
Y_{\Phi_{\xi,t}(s)} = \rho_{(t\xi \cdot s)}(\eta).
$$

Hence

$$
\left(\Phi_{\xi,t}^{-1}\right)_* \left(Y_{\Phi_{\xi,t}(s)}\right) = \left(L_{t\xi}^{-1}\right)_* \rho_{(t\xi \cdot s)}\left(\eta\right),\tag{2.28}
$$

and from Lemma [2.17,](#page-6-0) $L_{t\xi}^{-1} = L_{-t\xi}$. Then, differentiating, we find that

$$
\frac{d}{dt}\left(\left(L_{-t\xi}\right)_*\rho_{(t\xi\cdot s)}\left(\eta\right)\right)\bigg|_{t=0} = \frac{d}{dt}\left(\rho_{(t\xi\cdot s)}\left(\eta\right)\right)\bigg|_{t=0} - \frac{d}{dt}\left(\left(L_{t\xi}\right)_*\rho_{(s)}\left(\eta\right)\right)\bigg|_{t=0}.
$$

Also from Lemma [2.17,](#page-6-0) we know that $\rho_{(s)} = (R_s)_*|_0$. Hence, since for some parameter t_2 , $t_2\eta$ is a path in l through 0 with tangent vector η at 0, we can rewrite

$$
\rho_{(t\xi \cdot s)}(\eta) = (R_{t\xi \cdot s})_*|_0(\eta)
$$

=
$$
\frac{d}{dt_2}((t_2\eta) \cdot (t\xi \cdot s))|_{t_2=0}
$$

=
$$
\frac{d}{dt_2}R_s((t_2\eta) \circ_s (t\xi))|_{t_2=0}
$$

and similarly,

$$
(L_{t\xi})_* \rho_{(s)}(\eta) = (L_{t\xi})_* \Big|_s (R_s)_* \Big|_0 (\eta)
$$

$$
= \left. \frac{d}{dt_2} (t\xi) \cdot ((t_2 \eta) \cdot s) \right|_{t_2=0}
$$

$$
= \left. \frac{d}{dt_2} R_s ((t\xi) \circ_s (t_2 \eta)) \right|_{t_2=0}
$$

Overall,

$$
[X,Y]|_s = (R_s)_* \frac{d^2}{dt_1 dt_2} ((t_2 \eta) \circ_s (t_1 \xi) - (t_1 \xi) \circ_s (t_2 \eta)) \Big|_{t_1 = t_2 = 0},
$$

and therefore, we indeed get [\(2.25a\)](#page-10-2).

Now consider

$$
\left. \frac{d^3}{dt_1 dt_2 dt_3} \left(t_1 \xi \right) \circ_{(t_3 \gamma \cdot s)} (t_2 \eta) \right|_{t_1 = t_2 = t_3 = 0}
$$

.

.

Expanding, we get

$$
(t_1\xi) \circ_{(t_3\gamma\cdot s)} (t_2\eta) = (R_{t_3\gamma\cdot s})^{-1} ((t_1\xi) \cdot ((t_2\eta) \cdot (t_3\gamma \cdot s)))
$$

=
$$
(R_{t_3\gamma\cdot s})^{-1} (((t_1\xi) \circ_s ((t_2\eta) \circ_s (t_3\gamma))) \cdot s).
$$

Using [\(2.7\)](#page-6-1) with $p(t_3) = ((t_1\xi) \circ_s ((t_2\eta) \circ_s (t_3\gamma))) \cdot s$, we find

$$
\frac{d}{dt_3} (R_{t_3 \gamma \cdot s})^{-1} \left(\left((t_1 \xi) \circ_s ((t_2 \eta) \circ_s (t_3 \gamma)) \right) \cdot s \right) \Big|_{t_3 = 0} = \frac{d}{dt_3} (t_1 \xi) \circ_s ((t_2 \eta) \circ_s (t_3 \gamma)) \Big|_{t_3 = 0}
$$

$$
- \frac{d}{dt_3} ((t_1 \xi) \circ_s (t_2 \eta)) \circ_s (t_3 \gamma) \Big|_{t_3 = 0}
$$

Hence,

$$
\left. \frac{d^3}{dt_1 dt_2 dt_3} (t_1 \xi) \circ_{(t_3 \gamma \cdot s)} (t_2 \eta) \right|_{t_1 = t_2 = t_3 = 0} = [\xi, \eta, \gamma]^{(s)}.
$$

Consider now

$$
d_{\rho(\gamma)}b(\xi,\eta)\big|_{s} = a^{(s)}(\xi,\eta,\gamma) ,
$$

where we use the definition of $a^{(s)}$ [\(2.23\)](#page-10-3). However, from [\(2.25a\)](#page-10-2),

$$
d_{\rho(\gamma)}b(\xi,\eta)\Big|_{s} = \frac{d}{dt_{3}}[\xi,\eta]^{t_{3}\gamma\cdot s}\Big|_{t_{3}=0}
$$

=
$$
\frac{d^{3}}{dt_{1}dt_{2}dt_{3}} ((t_{1}\xi) \circ_{(t_{3}\gamma\cdot s)} (t_{2}\eta) - (t_{2}\eta) \circ_{(t_{3}\gamma\cdot s)} (t_{2}\xi))\Big|_{t_{1}=t_{2}=t_{3}=0}
$$

=
$$
[\xi,\eta,\gamma]^{(s)} - [\eta,\xi,\gamma]^{(s)}.
$$

So indeed,

 \blacksquare

$$
a^{(s)}(\xi, \eta, \gamma) = [\xi, \eta, \gamma]^{(s)} - [\eta, \xi, \gamma]^{(s)}.
$$

Consider now the connection ∇ from [\(2.3\)](#page-5-0). Its torsion $T(X, Y)$ for vector fields X and Y is given by

$$
T(X,Y) = \nabla_X Y - \nabla_Y X - [X,Y],
$$

so in particular for $X = \rho\left(\xi \right)$ and $Y = \rho\left(\eta \right) ,$ for $\xi, \eta \in \mathfrak{l},$ we have

$$
T\left(\rho\left(\xi\right),\rho\left(\eta\right)\right)=-\left[\rho\left(\xi\right),\rho\left(\eta\right)\right].
$$

Hence, using [\(2.12\)](#page-8-0), we find that at a point $s \in \mathbb{L}$, we have

$$
T\left(\rho\left(\xi\right),\rho\left(\eta\right)\right)_{s}=\rho_{s}\left(\left[\xi,\eta\right]^{(\rho,s)}\right). \tag{2.29}
$$

Now let $Z = \rho(\gamma)$ for $\gamma \in I$, hence

$$
g_s(T(\rho(\xi), \rho(\eta))_s, \rho_s(\gamma)) = \langle [\xi, \eta]^{(\rho, s)}, \gamma \rangle.
$$
 (2.30)

Recall that a torsion of a metric connection is said to be totally skew-symmetric if for any vector fields X, Y, Z ,

$$
g(T(X,Y),Z) = -g(Y,T(X,Z)).
$$
\n(2.31)

From [\(2.30\)](#page-13-0), we see that the torsion of the flat connection ∇ is totally skewsymmetric if and only if for any $s \in \mathbb{L}$,

$$
\langle \left[\xi, \eta \right]^{(\rho, s)}, \gamma \rangle = -\langle \eta, \left[\xi, \gamma \right]^{(\rho, s)} \rangle. \tag{2.32}
$$

Equivalently, for any $\xi \in I$ and any $s \in \mathbb{L}$, the map $\text{ad}_{\xi}^{(\rho,s)}$ is skew-adjoint with respect to the inner product on l.

Moreover, consider ∇T . Then,

$$
\begin{array}{rcl}\n(\nabla_{\rho(\gamma)}T)(\rho(\xi),\rho(\eta)) & = & \nabla_{\rho(\gamma)}\left(T(\rho(\xi),\rho(\eta))\right) \\
 & = & \nabla_{\rho(\gamma)}(\rho(b(\xi,\eta))) \\
 & = & \rho\left(d_{\rho(\gamma)}(b(\xi,\eta))\right) \\
 & = & \rho\left(a(\xi,\eta,\gamma)\right).\n\end{array}
$$

A classical theorem by Cartan and Schouten [\[4\]](#page-30-1) (and [\[1\]](#page-30-5) for a more modern treatment) states the following.

Theorem 2.27 ([\[4\]](#page-30-1)) Let (M, q) be a simply-connected, complete irreducible Riemannian manifold with a flat metric connection ∇ with a non-zero totally skew-symmetric torsion T. Then

- 1. If $\nabla T = 0$, then M is isometric to a compact simple Lie groups.
- 2. If $\nabla T \neq 0$, then M is isometric to S^7 .

In our formulation, we have the following corollary.

Corollary 2.28 Suppose $(\mathbb{L}, \mathfrak{l}, \rho)$ is a parallelized manifold. If for $s \in \mathbb{L}$, there exists a smooth family of inner products $\langle \cdot, \cdot \rangle_s$ on l such that the bracket $[\cdot, \cdot]^{(\rho, s)}$ is skew-adjoint with respect to it, then there exists a parallelization $\tilde{\rho}$ of $\mathbb L$ and an inner product $\langle \tilde{\rho}^{-1}, \tilde{\rho}^{-1} \rangle$ is isometric to a product of compact simple Lie groups and copies of an inner product $\langle \cdot, \cdot \rangle$ on I, such that the Riemannian manifold (\mathbb{L}, g) with $g =$ $\dot{S}^7.$

Proof. Suppose for each $s \in \mathbb{L}, \langle \cdot, \cdot \rangle_s$ is such that for any $\xi, \eta, \gamma \in \mathfrak{l}$,

$$
\left\langle \left[\xi,\eta\right]^{(\rho,s)},\gamma\right\rangle _{s}=-\left\langle \eta,\left[\xi,\gamma\right]^{(\rho,s)}\right\rangle _{s}.
$$

Then, choosing any inner product $\langle \cdot, \cdot \rangle$ on l, we can write $\langle \xi, \eta \rangle_s = \langle Q_s^{-1} \xi, Q_s^{-1} \eta \rangle$, for some map $Q_s \in GL(1)$. Since the family of inner products is smooth, we obtain a smooth family of maps Q_s . Now define a new parallelization of $\mathbb L$ given by $\tilde{\rho}_s = \rho_s \circ Q_s$. Then,

$$
g(X,Y)|_s = \langle \tilde{\rho}_s^{-1} X, \tilde{\rho}_s^{-1} Y \rangle
$$

= $\langle Q_s^{-1} \rho_s^{-1} X, Q_s^{-1} \rho_s^{-1} Y \rangle$
= $\langle \rho_s^{-1} X, \rho_s^{-1} Y \rangle_s$.

In particular, note that by definition [\(2.12\)](#page-8-0)

$$
[\xi, \eta]^{(\tilde{\rho}, s)} = -\tilde{\rho}_s^{-1}([\tilde{\rho}(\xi), \tilde{\rho}(\eta)]|_s)
$$

= -Q_s^{-1} \rho_s^{-1}([\rho(Q(\xi)), \rho(Q(\eta))]|_s)
= Q_s^{-1}[Q_s(\xi), Q_s(\eta)]^{(\rho, s)}

and hence

$$
\langle [\xi, \eta]^{(\tilde{\rho}, s)}, \gamma \rangle = \langle Q_s^{-1} [Q_s(\xi), Q_s(\eta)]^{(\rho, s)}, \gamma \rangle
$$

$$
= \langle [Q_s(\xi), Q_s(\eta)]^{(\rho, s)}, Q_s(\gamma) \rangle
$$

$$
= - \langle Q_s(\eta), [Q_s(\xi), Q_s(\gamma)]^{(\rho, s)} \rangle
$$

$$
= - \langle \eta, [\xi, \gamma]^{(\tilde{\rho}, s)} \rangle.
$$

In particular, we now see that with g and ∇ defined using $\tilde{\rho}$, the torsion of ∇ satisfies [\(2.31\)](#page-13-1), and hence by Theorem [2.27,](#page-13-2) (L, g) is isometric to a product of compact simple Lie groups and copies of $S⁷$.

3 Automorphisms

Definition 3.1 The category of (complete, connected) parallelized manifolds is defined in the following way. An object in this category is the triple $(\mathbb{L}, \mathfrak{l}, \rho)$, where $\mathbb L$ is an n-dimensional parallelizable connected manifold, I is an n-dimensional vector space, and ρ is the trivialization map, such that the fundamental vector fields are complete.

Suppose $(\mathbb{L}_1, \mathfrak{l}_1, \rho_1)$ and $(\mathbb{L}_2, \mathfrak{l}_2, \rho_2)$ are two parallelized manifolds, then a morphism between them is a smooth map $h : \mathbb{L}_1 \longrightarrow \mathbb{L}_2$ for which there exists a linear map $h': \mathfrak{l}_1 \longrightarrow \mathfrak{l}_2$ such that the following diagram commutes

$$
T_p \mathbb{L}_1 \xrightarrow{h_*|_p} T_{h(p)} \mathbb{L}_2
$$

\n
$$
\uparrow (\rho_1)_p \qquad \qquad \uparrow (\rho_2)_{h(p)}
$$

\n
$$
\downarrow_1 \xrightarrow{h'} \downarrow_2
$$
\n(3.1)

We will refer to (h, h') as a compatible pair.

Remark 3.2 the notation h' is appropriate since this map is closely related to the derivative of h. Any smooth h will induce an invertible linear map $h'_p =$ $\rho_{f(p)}^{-1} \circ h_{*}|_{p} \circ \rho_{p} : \mathfrak{l}_{1} \longrightarrow \mathfrak{l}_{2},$ however in general it will depend on p. The key point of the above definition is that for a morphism of parallelized manifolds, h'_p is independent of p. A morphism between parallelized manifolds is a special case of a vector bundle morphism. Indeed, consider the tangent bundles $T\mathbb{L}_1$ and $T\mathbb{L}_2$. Then, the bundle map $(h, \rho_2 \circ h' \circ \rho_1^{-1}) : T \mathbb{L}_1 \longrightarrow T \mathbb{L}_2$ covers $h : \mathbb{L}_1 \longrightarrow \mathbb{L}_2$ and induces a linear map of each fiber.

Example 3.3 Suppose we have a parallelized manifold (L, L, ρ_1) , then any linear isomorphism of the vector space l induces an isomorphism of parallelized manifolds. Indeed, suppose $h' : \mathfrak{l} \longrightarrow \mathfrak{l}$ is an invertible linear map. Then, the compatible pair (id, h') , where id is the identity map on \mathbb{L} , give an isomorphism from $(\mathbb{L}, \mathfrak{l}, \rho_1)$ to $(\mathbb{L}, \mathfrak{l}, \rho_2)$, where $\rho_2 = \rho_1 \circ (h')^{-1}$. This just corresponds to a $GL(n)$ transformation of the global frame on TL.

Lemma 3.4 Suppose $(\mathbb{L}_1, \mathfrak{l}_1, \rho_1)$ and $(\mathbb{L}_2, \mathfrak{l}_2, \rho_2)$ are two parallelized manifolds. A pair (h, h') of a smooth map $h : \mathbb{L}_1 \longrightarrow \mathbb{L}_2$ and a linear map $h' : \mathfrak{l}_1 \longrightarrow \mathfrak{l}_2$ satisfy the definition of a morphism of parallelized manifolds if and only if for any $s \in \mathbb{L}_1$ and $\xi \in \mathfrak{l}_1$,

$$
h(\xi \cdot s) = h'(\xi) \cdot h(s). \tag{3.2}
$$

Proof. Let $\xi \in I_1$, and let $s \in \mathbb{L}$. Recall that $\xi \cdot s = \gamma_{\rho_1(\xi),s}(1)$ is the solution $x(1)$ at $t = 1$ of the initial value problem

$$
\begin{cases} \frac{dx(t)}{dt} = \rho_1(\xi)_{x(t)} \\ x(0) = s \end{cases}
$$

Then, applying h_* to the equation, we get

$$
h_*|_{x(t)}\left(\frac{dx(t)}{dt}\right) = \frac{d\left(h\left(x(t)\right)\right)}{dt}
$$
\n
$$
= h_*|_{x(t)}\left(\rho_1\left(\xi\right)_{x(t)}\right)
$$
\n(3.3)

.

Suppose now the pair (h, h') satisfies the definition of a morphism of parallelized manifolds. Using the property of h' from (3.1) , we have

$$
h_*|_{x(t)}\left(\rho_1(\xi)_{x(t)}\right) = \rho_2(h'(\xi))_{h(x(t))}.
$$

Therefore, we see that $y = h(x(t))$ satisfies the equation

$$
\begin{cases}\n\frac{dy(t)}{dt} = \rho_2 (h'(\xi))_{y(t)} \\
y(0) = h(s)\n\end{cases}
$$
\n(3.4)

and overall we see that $y(t) = \gamma_{\rho_2(\xi),h(s)}(t)$ For $t = 1$, we hence have

$$
h(\xi \cdot s) = h'(\xi) \cdot h(s).
$$

Conversely, suppose [\(3.2\)](#page-15-0) holds for any $s \in \mathbb{L}_1$ and $\xi \in \mathfrak{l}_1$. Then, we have

$$
h(t\xi \cdot s) = (th'(\xi)) \cdot h(s). \tag{3.5}
$$

Differentiating both sides, we find

$$
h_*|_{x(t)}\left(\rho_1(\xi)_{x(t)}\right) = \rho_2(h'(\xi))_{y(t)},
$$

where $x(t) = t\xi \cdot s$ and $y(t) = (th'(\xi)) \cdot h(s) = h(x(t))$. Setting $t = 0$, we find that the pair (h, h') indeed satisfies the definition of a morphism of parallelizable manifolds.

Example 3.5 In the case of Lie groups, the definition of a morphism of parallelized manifolds is equivalent to taking the pair of a Lie group homomorphism and its differential at identity.

Definition 3.6 Suppose (L, L, ρ) is a parallelized manifold. Then, an automorphism of $(\mathbb{L}, \mathfrak{l}, \rho)$ is a diffeomorphism $h : \mathbb{L} \longrightarrow \mathbb{L}$ for which there exists an invertible linear map $h': \mathfrak{l} \longrightarrow \mathfrak{l}$, such that the following diagram commutes

$$
T_p \mathbb{L} \xrightarrow{h_*|_p} T_{h(p)} \mathbb{L}
$$

\n
$$
\uparrow (\rho)_p \qquad \qquad \uparrow (\rho)_{h(p)}
$$

\n
$$
\downarrow \xrightarrow{h'} \qquad \downarrow \qquad (3.6)
$$

Remark 3.7 Given a trivialization ρ , we may define the trivial connection ∇ on TL via

$$
\nabla = \rho \circ d \circ \rho^{-1},\tag{3.7}
$$

where ρ is interpreted now as a map $\rho: C^{\infty}(\mathbb{L}, \mathfrak{l}) \longrightarrow \Gamma(T\mathbb{L})$. Given a diffeomorphism h of \mathbb{L} , the pullback connection $h^*\nabla$ is defined via

$$
h^*\nabla = h_*^{-1} \circ \nabla \circ h_*
$$

= $h_*^{-1} \circ \rho \circ d \circ \rho^{-1} \circ h_*.$ (3.8)

Setting $h' = \rho^{-1} \circ h_* \circ \rho$, we get

$$
h^*\nabla = \rho^{-1} \circ (h')^{-1} \circ d \circ h \circ \rho.
$$

Hence we see that $h^*\nabla = \nabla$ if and only if h' is constant and equivalently h is an automorphism of (L, L, ρ) . In other words, h is a connection-preserving diffeomorphism, or as it is also known as, an affine diffeomorphism for the trivial connection ∇.

The set of automorphisms of (L, I, ρ) is clearly non-empty, since the identity map is always an automorphism, and it's clearly a Lie group, with the Lie group structure induced from Diff (L). Let us denote it by Ψ (L, l, ρ), or just Ψ if there is no ambiguity. Each $h \in \Psi (\mathbb{L}, \mathfrak{l}, \rho)$ defines a corresponding linear isomorphism h' of l, so denote this subgroup of $GL(I)$ by $\Psi'(\mathbb{L}, I, \rho)$, or just Ψ' if there is no ambiguity. Under an automorphism, a fundamental vector field $\rho(\xi)$ is pushed forward to $\rho(h'(\xi))$. The map $h \mapsto h'$ is a Lie group homomorphism, so Ψ' is a Lie group of dimension at most n^2 where $n = \dim \mathbb{L}$.

Let $\pi : \Psi \longrightarrow \Psi'$ be the homomorphism that takes h to h'. Let $\mathfrak p$ and $\mathfrak p'$ be the Lie algebras of Ψ and Ψ' , respectively, then at least for small t , $\pi \left(\exp_{\mathfrak{p}}(t\eta) \right) =$ $\exp_{p'}(t\pi_*\eta)$ for $\eta \in \mathfrak{p}$.

Theorem 3.8 Suppose \mathbb{L} is a connected parallelizable manifold of dimension n. The automorphism group $\Psi = \Psi(\mathbb{L}, \mathfrak{l}, \rho)$ of $(\mathbb{L}, \mathfrak{l}, \rho)$ is a finite-dimensional Lie group of dimension at most $n^2 + n$.

Proof. The Lie group structure of Ψ is clearly induced from Diff (\mathbb{L}) , so it is sufficient to show that Ψ is a finite-dimensional manifold. Fix any $s \in \mathbb{L}$, and define the smooth map $\Psi \longrightarrow \Psi' \times \mathbb{L}$ given for any $h \in \Psi$ by

$$
h \mapsto (h', h(s)) \tag{3.9}
$$

where $h' \in GL$ (I) is the corresponding element of Ψ' . We will show that this map is injective. Indeed, suppose $\tilde{h} \in \Psi$ also maps to $(h', h(s))$. Then, consider $k = h^{-1} \tilde{h} \mapsto (\text{id}, s)$. Recall from Theorem [2.20](#page-8-1) that any element $p \in \mathbb{L}$ can be written as a finite product $p = \xi_1 \cdot (\xi_2 \cdot ... (\xi_m \cdot s))$ for $\xi_i \in \mathfrak{l}$. Then, applying k to p using Lemma [3.4](#page-15-1) gives

$$
k\left(p\right)=k'\left(\xi_{1}\right)\cdot\left(k'\left(\xi_{2}\right)\cdot...\left(k'\left(\xi_{m}\right)\cdot k\left(s\right)\right)\right)=p,
$$

since $k' = id$ and $k(s) = s$. Hence, $k = id$, and hence [\(3.9\)](#page-17-1) is an injective map.

Suppose $h_t \in \Psi \subset \text{Diff}(\mathbb{L})$, is a smooth 1-parameter family with $h_0 = id$. Then, for a fixed $s \in \mathbb{L}$, $h_t(s)$ is a smooth curve on \mathbb{L} , with $u_s = \frac{d}{dt} h_t(s)|_{t=0} \in \mathbb{R}$ $T_s\mathbb{L}$. The vector field $u \in \Gamma(T\mathbb{L})$ is then the tangent vector to the curve $h_t \in \Psi$ at $t = 0$.

Recall from Theorem [2.20](#page-8-1) that any element $p \in \mathbb{L}$ can be written as a finite product $p = \xi_1 \cdot (\xi_2 \cdot ... (\xi_m \cdot s))$ for $\xi_i \in \mathfrak{l}$, then

$$
u_p = \frac{d}{dt} h_t(p) \Big|_{t=0}
$$

\n
$$
= \frac{d}{dt} h_t(\xi_1 \cdot (\xi_2 \cdot ... (\xi_m \cdot s))) \Big|_{t=0}
$$

\n
$$
= \frac{d}{dt} h'_t(\xi_1) \cdot (h'_t(\xi_2) \cdot ... (h'_t(\xi_m) \cdot h_t(s))) \Big|_{t=0}
$$

\n
$$
= \frac{d}{dt} h'_t(\xi_1) \cdot (\xi_2 \cdot ... (\xi_m \cdot s)) \Big|_{t=0} + ...
$$

\n
$$
+ \frac{d}{dt} \xi_1 \cdot (\xi_2 \cdot ... (h'_t(\xi_m) \cdot s)) \Big|_{t=0}
$$

\n
$$
+ \frac{d}{dt} \xi_1 \cdot (\xi_2 \cdot ... (\xi_m \cdot (h_t(s)/s \cdot s))) \Big|_{t=0}
$$

\n(3.10)

In particular, consider the linear map

$$
u \mapsto \left(\left. \frac{d}{dt} h_t' \right|_{t=0}, \left. \frac{d}{dt} h_t \left(s \right) / s \right|_{t=0} \right) \in \mathfrak{p}' \times \mathfrak{l}.\tag{3.11}
$$

This is the differential of the map (3.9) at identity. From (3.10) we see that if $\frac{d}{dt}h'_t\Big|_{t=0} = 0$ and $\frac{d}{dt}h_t(s)/s\Big|_{t=0} = 0$, then $u = 0$. Thus, we see that the map [\(3.11\)](#page-18-0) is injective. Similarly, by translation, the differential of [\(3.9\)](#page-17-1) will be injective at any $h \in \Psi$. Therefore, [\(3.9\)](#page-17-1) is an injective smooth immersion, and thus a smooth embedding. Hence, Ψ is smoothly embedded in $\Psi' \times \mathbb{L}$, which is at most $(n^2 + n)$ -dimensional.

Suppose $s \in \mathbb{L}$ is fixed, then recall from [\(2.8\)](#page-7-0) that the product \circ_s on l is defined by $\eta \circ_s \xi = (\eta \cdot (\xi \cdot s)) / s$ for sufficiently small $\eta, \xi \in \mathfrak{l}$. Now for $h \in \Psi$ and the corresponding $h' \in \Psi'$, consider

$$
h'(\eta \circ_s \xi) = h(\eta \cdot (\xi \cdot s)) / h(s)
$$

= $h'(\eta) \cdot (h'(\xi) \cdot h(s)) / h(s)$
= $h'(\eta) \circ_{h(s)} h'(\xi)$. (3.12)

Equivalently, we can write

$$
h'(\eta \circ_s \xi) \cdot h(s) = h'(\eta) \cdot (h'(\xi) \cdot h(s)), \qquad (3.13)
$$

or

$$
h'(\eta \circ_s \xi) \circ_s (h(s)/s) = h'(\eta) \circ_s (h'(\xi) \circ_s (h(s)/s)), \tag{3.14}
$$

Remark 3.9 In the theory of loops and quasigroups, a map h' that satisfies the property $h'(xy) A = h'(x) (h'(y) A)$ is known as a pseudoautomorphism with companion A. Using this language, we can say that h' has companion $h(s)/s$ with respect to s. The group Aut (I, \circ_s) of automorphisms of \circ_s is then defined as the subset of elements of Ψ' which have companion 0. This then corresponds to the stabilizer Stab_Ψ (s) of s.

The group Ψ also acts on the bracket $[\cdot, \cdot]^{(s)}$. As we noted above, on Lie groups, the map h' is the differential of h at identity, and we know this is a Lie algebra isomorphism. However, in the more general case, the map h' is an algebra isomorphism from $(I, [\cdot, \cdot]^{(s)})$ to $(I, [\cdot, \cdot]^{(h(s))})$.

Theorem 3.10 Suppose $s \in \mathbb{L}$ and $h \in \Psi$, then for any $\xi, \eta \in \mathfrak{l}$,

$$
h'\left(\left[\xi,\eta\right]^{(s)}\right) = \left[h'\left(\xi\right),h'\left(\eta\right)\right]^{(h(s))}.
$$
\n(3.15)

Proof. Indeed, from Theorem [2.26](#page-10-0) and using [\(3.12\)](#page-18-1), we obtain

$$
h'\left([\xi,\eta]^{(s)}\right) = \frac{d^2}{dt_1dt_2} \left(h'\left((t_1\xi) \circ_s (t_2\eta) \right) - h'\left((t_2\eta) \circ_s (t_1\xi) \right) \right) \Big|_{t_1=t_2=0}
$$

\n
$$
= \frac{d^2}{dt_1dt_2} \left((t_1h'(\xi)) \circ_{h(s)} (t_2h'(\eta)) - \left((t_2h'(\eta)) \circ_{h(s)} (t_1h'(\xi)) \right) \right) \Big|_{t_1=t_2=0}
$$

\n
$$
= \left[h'(\xi), h'(\eta) \right]^{h(s)}.
$$

4 Products of spheres

As mentioned in Example [2.9,](#page-4-0) a large family of parallelizable manifolds that are not Lie groups or Moufang loops is given by products of spheres where one factor is an odd-dimensional sphere. As it is well-known, the only parallelizable spheres are S^1, S^3, S^7 , which correspond to unit norm sets of complex numbers, quaternions, and octonions. In particular, S^1 and S^3 are both Lie groups, while $S⁷$ is the Moufang loop of unit octonions. In [\[3,](#page-30-7) [14,](#page-31-4) [15\]](#page-31-5) explicit parallelizations on products of spheres have been computed and this will serve as an important example of the framework developed in this paper.

In particular, let us consider in detail the case of $S^m \times S^1$ from [\[14\]](#page-31-4). Consider S^m as embedded in \mathbb{R}^{m+1} in a standard way

$$
S^{m} = \left\{ (x_1, ..., x_{m+1}) \in \mathbb{R}^{m+1} : |x|^2 = x_1^2 + ... + x_{m+1}^2 = 1 \right\}.
$$
 (4.1)

Consider the standard coordinate frame $\{\partial_{x_i}\}$ on \mathbb{R}^{m+1} . Then, the orthogonal projections $\{M_i\}$ of these vector fields onto S^m are given by

$$
M_i = \partial_{x_i} - x_i M,\t\t(4.2)
$$

where $M = \sum_{i=1}^{m+1} x_i \partial_{x_i}$ is the normal vector field to $S^m \subset \mathbb{R}^{m+1}$. Also consider ϕ as a coordinate on S^1 with the associated vector field ∂_{ϕ} . Then, a parallelization of $S^m \times S^1$ is given by the following.

Theorem 4.1 Consider $S^m \subset \mathbb{R}^{m+1}$ and for $i = 1, ..., n+1$, define

$$
f_i = M_i + x_i \partial_{\phi}.
$$
\n
$$
(4.3)
$$

Then, the set $\{f_i\}$ is global frame on $S^m \times S^1$. In particular, it is orthonormal with respect to the product metric and satisfies

$$
[f_i, f_j] = x_i f_j - x_j f_i. \tag{4.4}
$$

Now let $\mathbb{L} = S^m \times S^1$, so that each $s \in \mathbb{L}$ is given by $s = (x, \phi)$ for $x \in$ $S^m \subset \mathbb{R}^{m+1}$ and $\phi \in S^1$. Also suppose l is an $(m+1)$ -dimensional real vector space with an inner product $\langle \cdot, \cdot \rangle$ and an orthonormal basis $\{e_i\}$. Then, for each $s = (x, \phi) \in \mathbb{L}$, define

$$
\rho_s: \mathfrak{l} \longrightarrow T_s \mathbb{L} \ne_i \mapsto f_i|_s
$$
\n(4.5)

In particular, if $\xi = \xi^i e_i \in I$, then $\rho(\xi) = \xi^i f_i$.

Consider the derivatives of the coordinate functions in the directions of the basis vectors $\{f_i\}$. We have

$$
f_i(x_j) = (\partial_{x_i} - x_i M) x_j
$$

= $\delta_{ij} - x_i x_j$

$$
f_i(\phi) = x_i
$$

In particular, define $c_{ij} = f_i(x_j) = \delta_{ij} - x_i x_j$.

The bracket on $\mathfrak l$ is defined from (4.4) via (2.12) . In particular, for the basis ${e_i}$ on l and $s = (x, \phi) \in \mathbb{L}$, we have

$$
b_{ij}^{(s)} = [e_i, e_j]^{(s)} = e_i x_j - e_j x_i.
$$
\n(4.6)

Equivalently, given any $\xi, \eta \in I$, we have

$$
[\xi, \eta]^{(s)} = \xi \langle \eta, x \rangle - \eta \langle \xi, x \rangle. \tag{4.7}
$$

In particular, note that if both ξ and η are orthogonal to x, then $[\xi, \eta]^{(s)} = 0$ and for any $\xi \in \mathfrak{l}, [\xi, x]^{(s)} = \xi$.

Lemma 4.2 For each fixed s, I with the bracket $[\cdot, \cdot]^{(s)}$ defines a Lie algebra that is isomorphic to a semidirect sum $\mathbb{R}^n \oplus_S \mathbb{R}$, with bracket $[(\xi, \lambda), (\eta, \mu)] =$ $[\mu\xi - \lambda\eta, 0].$

Proof. First let us show that $[\cdot, \cdot]^{(s)}$, as given by [\(4.7\)](#page-20-0) defines a Lie algebra on l. We just need to check the Jacobi identity. Indeed, for $\gamma, \xi, \eta \in I$,

 λ

$$
\begin{aligned}\n\left[\gamma, [\xi, \eta]^{(s)}\right]^{(s)} &= [\gamma, \xi]^{(s)} \langle \eta, x \rangle - [\gamma, \eta]^{(s)} \langle \xi, x \rangle \\
&= \gamma \langle \xi, x \rangle \langle \eta, x \rangle - \xi \langle \gamma, x \rangle \langle \eta, x \rangle \\
&\quad - \gamma \langle \eta, x \rangle \langle \xi, x \rangle + \eta \langle \gamma, x \rangle \langle \xi, x \rangle \\
&= \eta \langle \gamma, x \rangle \langle \xi, x \rangle - \xi \langle \gamma, x \rangle \langle \eta, x \rangle \\
\left[\xi, [\eta, \gamma]^{(s)}\right]^{(s)} &= \gamma \langle \xi, x \rangle \langle \eta, x \rangle - \eta \langle \xi, x \rangle \langle \gamma, x \rangle \\
\left[\eta, [\gamma, \xi]^{(s)}\right]^{(s)} &= \xi \langle \eta, x \rangle \langle \gamma, x \rangle - \gamma \langle \eta, x \rangle \langle \xi, x \rangle\n\end{aligned}
$$

Adding these terms together shows that indeed the Jacobi identity is satisfied. Now consider the linear map $F: \mathfrak{l} \longrightarrow \mathbb{R}^n \oplus \mathbb{R}$, given by

$$
F(\xi) = (\xi - \langle \xi, x \rangle x, \langle \xi, x \rangle).
$$

This is clearly a linear isomorphism. Then,

$$
F\left([\xi,\eta]^{(s)}\right) = F(\xi)\langle\eta,x\rangle - F(\eta)\langle\xi,x\rangle
$$

\n
$$
= (\xi - \langle\xi,x\rangle x, \langle\xi,x\rangle)\langle\eta,x\rangle
$$

\n
$$
- (\eta - \langle\eta,x\rangle x, \langle\eta,x\rangle)\langle\xi,x\rangle
$$

\n
$$
= (\xi\langle\eta,x\rangle - \eta\langle\xi,x\rangle, 0)
$$

\n
$$
= [(\xi - \langle\xi,x\rangle x, \langle\xi,x\rangle), (\eta - \langle\eta,x\rangle x, \langle\eta,x\rangle)]
$$

\n
$$
= [F(\xi), F(\eta)].
$$

Hence, this is a Lie algebra homomorphism, and thus the Lie algebra $(I, [\cdot, \cdot]^{(s)})$ is isomorphic to $\mathbb{R}^n \oplus_S \mathbb{R}$.

Taking the derivative of the bracket in the direction of f_i gives us

$$
d_{f_i}\left([\xi,\eta]^{(s)}\right) = \xi \eta^j c_{ij} - \eta \xi^j c_{ij}
$$

= $\xi \eta_i - \eta \xi_i - x_i [\xi,\eta]^{(s)}$

In particular, this shows that

$$
d_{f_i}b_{jk}^{(s)} =: a_{ijk}^{(s)} = e_k \delta_{ij} - e_j \delta_{ik} - x_i b_{jk}^{(s)}.
$$
\n(4.8)

Hence, for $\xi, \eta, \gamma \in I$,

$$
a^{(s)}(\xi, \eta, \gamma) = \gamma (\langle \xi, \eta \rangle + \langle x, \xi \rangle \langle x, \eta \rangle) - \eta (\langle \xi, \gamma \rangle + \langle x, \xi \rangle \langle x, \gamma \rangle). \tag{4.9}
$$

Recall the notion of a Lie triple system $[8]$: this is a vector space equipped with a skew-symmetric triple product $[\cdot, \cdot, \cdot]$, that is skew-symmetric in two entries, has a vanishing sum of cyclic permutations(i.e. satisfies the Jacobi identity), and also that the linear map defined by fixing two of the entries of $[\cdot, \cdot, \cdot]$ is a derivation of the triple product. Consider now I equipped with the triple product $a^{(s)}$. We know that $a^{(s)}$ is skew-symmetric in the last two entries. From [\(2.24\)](#page-10-4), we also see that $a^{(s)}$ satisfies the Jacobi identity if and only if $(1, b^{(s)})$ is a Lie algebra. Therefore, in this case, it just suffices to check the derivation property to figure out if $(1, a^{(s)})$ is a Lie triple system.

Theorem 4.3 For each $s \in \mathbb{L}$, the vector space \mathfrak{l} equipped with the triple product $a^{(s)}$ forms a Lie triple system.

Proof. Since $b^{(s)}$ satisfies the Jacobi identity, from (2.24) we know that

$$
a^{(s)}(\xi, \eta, \gamma) + a^{(s)}(\eta, \gamma, \xi) + a^{(s)}(\gamma, \xi, \eta) = 0.
$$

To show that $(I, a^{(s)})$ forms a Lie triple system, we just need to show that for any $\xi, \eta \in I$, the map $a_{\xi,\eta}^{(s)} : I \longrightarrow I$ given by $a_{\xi,\eta}^{(s)}(\gamma) = a^{(s)}(\gamma,\xi,\eta)$ is a derivation for $a^{(s)}$. That is, for any $\xi, \eta, \gamma, u, v \in I$, we need to show that

$$
a_{\xi,\eta}^{(s)}(a^{(s)}(u,v,\gamma)) = a^{(s)}(a^{(s)}(u,\xi,\eta),v,\gamma) +a^{(s)}(u,a^{(s)}(v,\xi,\eta),\gamma) +a^{(s)}(u,v,a^{(s)}(\gamma,\xi,\eta)).
$$
\n(4.10)

This is a straightforward, but tedious calculation, which we show in Appendix $A.$

To define products on $\mathfrak l$ and $\mathbb L$, we need to consider integral curves of fun-damental vector fields. In [\(2.5\)](#page-6-2) suppose the curve $p(t) \in \mathbb{L}$ has coordinates $(x(t), \phi(t))$. Suppose $\xi \neq 0$, and let us rewrite $\rho(\xi)$ in terms of the coordinate basis:

$$
\rho(\xi) = \xi^i f_i
$$

= $\xi^i \partial_{x_i} - \xi^i x_i \sum_{i=1}^{m+1} x_j \partial_{x_j} + \xi^i x_i \partial_{\phi}.$

Setting $\hat{\xi} = \xi^i x_i$, we can thus rewrite the flow equation [\(2.5\)](#page-6-2) as

$$
\begin{cases}\n\frac{dx(t)}{dt} = \xi - \hat{\xi}x \\
\frac{d\phi(t)}{dt} = \hat{\xi}\n\end{cases}
$$
\n(4.11)

with initial conditions (x_0, ϕ_0) . Without loss of generality, assume $|\xi| = 1$.

Lemma 4.4 The solution of [\(4.11\)](#page-22-0) exists for all $t \in \mathbb{R}$ and, if $\xi = \pm x_0$, the solutions are given by

$$
x(t) = x_0 \tag{4.12a}
$$

$$
\phi(t) = \phi_0 \pm t. \tag{4.12b}
$$

If $\xi \neq x_0$, then the solution is given by

$$
x(t) = \xi \tanh(\sigma + t) + \tilde{x}_0 \operatorname{sech}(\sigma + t) \tag{4.13a}
$$

$$
\phi(t) = \phi_0 + \ln\left(\frac{\cosh\left(t+\sigma\right)}{\cosh\sigma}\right),\tag{4.13b}
$$

where $\tilde{x}_0 = x_0 \cosh \sigma - \xi \sinh \sigma$ and σ is such that $\hat{\xi}(0) = \tanh \sigma$.

Proof. From [\(4.11\)](#page-22-0), we see $x(t)$ is constant if and only if $\xi = \pm x_0$. In that case, $\hat{\xi}(t) = \pm 1$, and thus $\phi(t) = \phi_0 \pm t$.

More generally, taking the inner product of the first equation in [\(4.11\)](#page-22-0) with ξ , we obtain an equation for $\hat{\xi}(t)$:

$$
\frac{d\hat{\xi}}{dt} = 1 - \hat{\xi}^2.
$$
\n(4.14)

Since $\hat{\xi}^2 = 1$ if and only if $\xi = \pm x_0$, suppose $\hat{\xi}(t) = \tanh \sigma(t)$, for some function $\sigma(t)$. Then, [\(4.14\)](#page-22-1) immediately gives $\sigma(t) = t + \sigma$, where $\sigma = \sigma(0)$, that is, $\hat{\xi}(0) = \tanh \sigma.$

Let us integrate $\hat{\xi}$, because this will be needed as an integrating factor for the first equation in [\(4.11\)](#page-22-0) and also to solve for $\phi(t)$ in the second equation in $(4.11):$ $(4.11):$

$$
\int^{t} \hat{\xi}(\tau) d\tau = \ln(\cosh(\sigma + t)). \qquad (4.15)
$$

Hence, we can write the integrating factor as

$$
M(t) = \cosh(\sigma + t)
$$

and the general solution is

$$
x(t) = \xi \tanh(\sigma + t) + \tilde{x}_0 \operatorname{sech}(\sigma + t), \qquad (4.16)
$$

where

$$
\tilde{x}_0 = x_0 \cosh \sigma - \xi \sinh \sigma.
$$

We also obtain $\phi(t)$ [\(4.13b\)](#page-22-2) by directly integrating the second equation of [\(4.11\)](#page-22-0):

$$
\phi(t) = \phi_0 + \ln\left(\frac{\cosh(t+\sigma)}{\cosh \sigma}\right)
$$

As expected on a compact manifold, the integral curve exists for all values of t. \blacksquare

Overall, we see the geometric picture of how the integral curve evolves. For the non-trivial case, suppose $\xi \neq \pm x_0$, then from [\(4.16\)](#page-22-3), we see that $x(t)$ always lies in the plane spanned by x_0 and ξ , so the curve is along the great circle of S^m given by the plane spanned by x_0 and ξ . If $\alpha(t)$ is the angle between $x(t)$ and ξ , then $\hat{\xi}(t) = \cos \alpha(t)$. In particular, then

$$
\cos \alpha (t) = \tanh (\sigma + t) \n\sin \alpha (t) = \operatorname{sech} (\sigma + t).
$$

In particular, we see that $\cos \alpha (t)$ is a monotonic function, with $\cos \alpha (t) \longrightarrow +1$ as $t \longrightarrow +\infty$ and $\cos \alpha (t) \longrightarrow -1$ as $t \longrightarrow -\infty$. Moreover, $\sin \alpha (t)$ is always positive. Therefore, $\alpha(t) \in (0, \pi)$, with $\alpha(t) \longrightarrow 0$ as $t \longrightarrow +\infty$ and $\alpha(t) \longrightarrow \pi$ as $t \longrightarrow -\infty$. The equation [\(4.13a\)](#page-22-2) can be rewritten as

$$
x(t) = \xi \cos \alpha(t) + \tilde{x}_0 \sin \alpha(t).
$$
 (4.17)

This shows that in the adapted $\xi - \tilde{x}_0$ plane, $x(t)$ stays in the upper half-plane, moving clockwise as t increases. In particular, as $t \to \pm \infty$, $x(t) \to \pm \xi$.

Example 4.5 Since $a^{(s)}$ in this case is non-trivial (4.9) , from Theorem [2.26](#page-10-0) we see that \circ_s is necessarily non-associative. To illustrate this product \circ_s on l [\(2.8\)](#page-7-0) more explicitly, let $s = (x, \phi)$ and consider elements unit length $\xi, \eta \in I$, such that they are orthogonal to each other and to x . The more general case is computationally a bit more involved. Consider first $(x_1(t), \phi_1(t)) = (t\xi) \cdot s$. From [\(4.13\)](#page-22-4), noting that $\xi \cdot x = 0$ implies $\sigma = 0$, we have

$$
x_1(t) = \xi \tanh t + x \operatorname{sech} t \tag{4.18}
$$

$$
\phi_1(t) = \phi_0 + \ln(\cosh t). \tag{4.19}
$$

Next, consider $(x_2(\tau), \phi_2(\tau)) = (\tau \eta) \cdot ((t \xi) \cdot s)$. By assumption, the initial value $x_2(0) = \xi \tanh t + x \sech t$ is orthogonal to η , so σ is again 0. Thus,

> $x_2(\tau) = \eta \tanh \tau + \xi \tanh t \operatorname{sech} \tau + x \operatorname{sech} t \operatorname{sech} \tau$ $\phi_2(\tau) = \phi_0 + \ln(\cosh t \cosh \tau).$

To find $(\tau \eta) \circ_s (t \xi) = (\tau \eta) \cdot ((t \xi) \cdot s) / s$, we need to find a positive $t_2 \in \mathbb{R}$ and $\gamma \in I$ of unit length, such that

$$
x_2(\tau) = \gamma \tanh(\sigma_2 + t_2) + (x \cosh \sigma_2 - \gamma \sinh \sigma_2) \operatorname{sech}(\sigma_2 + t_2)
$$

$$
\phi_2(\tau) = \phi_0 + \ln\left(\frac{\cosh(t_2 + \sigma_2)}{\cosh \sigma_2}\right) + 2\pi k,
$$

where tanh $\sigma_2 = \gamma \cdot x$ and k is an integer. The freedom to add an integer multiple of $2\pi k$ stems from the fact that ϕ is a coordinate on the circle. We know that γ is a linear combination of ξ, η, x , so let

$$
\gamma = a\xi + b\eta + (\tanh \sigma_2) x. \tag{4.20}
$$

Then, taking inner products of x_2 with ξ, η, x , and also comparing ϕ_2 , we find the following equations

$$
\tanh t \operatorname{sech} \tau = a \left(\tanh \left(t_2 + \sigma_2 \right) - \sinh \sigma_2 \operatorname{sech} \left(t_2 + \sigma_2 \right) \right) \tag{4.21a}
$$

$$
\tanh \tau = b(\tanh (t_2 + \sigma_2) - \sinh \sigma_2 \operatorname{sech} (t_2 + \sigma_2)) \tag{4.21b}
$$

sech t sech
$$
\tau = \tanh \sigma_2 \tanh (t_2 + \sigma_2)
$$
 (4.21c)

+ sech
$$
\sigma_2
$$
 sech $(t_2 + \sigma_2)$
cosh $(t_2 + \sigma_2)$

$$
e^{2\pi k} \cosh t \cosh \tau = \frac{\cosh(t_2 + \sigma_2)}{\cosh \sigma_2} \tag{4.21d}
$$

$$
1 = a^2 + b^2 + \tanh^2 \sigma_2.
$$
 (4.21e)

The last equation comes from the fact that γ is unit length. However if we require that $t_2 \longrightarrow 0$ as $t, \tau \longrightarrow 0$, which makes sense in order for the product \circ_s to be defined in the neighborhood of 0, $(4.21d)$ shows that $k = 0$. From $(4.21c)$ and $(4.21d)$ with $k = 0$, we find

$$
\tanh \sigma_2 \tanh (t_2 + \sigma_2) + (\operatorname{sech} \sigma_2 - \cosh \sigma_2) \operatorname{sech} (t_2 + \sigma_2) = 0
$$

and thus

$$
\tanh \sigma_2 \left(\sinh (t_2 + \sigma_2) - \sinh \sigma_2\right) = 0.
$$

Since $t_2 > 0$, we find that $\tanh \sigma_2 = 0$, and hence $\sigma_2 = 0$. This immediately gives us

$$
\cosh t_2 = \cosh t \cosh \tau
$$

$$
a = \frac{\tanh t \operatorname{sech} \tau}{\tanh t_2}
$$

$$
b = \frac{\tanh \tau}{\tanh t_2}.
$$

Thus, we conclude that for an orthogonal triple η, ξ, x , we get

$$
(\tau \eta) \circ_s (t\xi) = t_2 (a\xi + b\eta) ,
$$

with t_2 , a , and b as given above.

Let us now consider the isomorphisms of \mathbb{L} . Suppose $h \in \Psi(\mathbb{L})$ and $h' \in$ $\Psi'(\mathbb{L})$ is the map of l. Recall from Lemma [3.4](#page-15-1) that for $\xi \in \mathfrak{l}$ and $s \in \mathbb{L}$, $h(\xi \cdot s) =$ $h'(\xi) \cdot h(s)$.

Given $s = (x, \phi) \in \mathbb{L}$, let the corresponding components of $h(s)$ be $h(s)_x$ and $h(s)_{\phi}$. From Theorem [3.10,](#page-18-2) we then know that for any $\xi, \eta \in I$ and $s =$ $(x,\phi)\in\mathbb{L}$

$$
h'\left([\xi, \eta]^{(s)} \right) = [h'(\xi), h'(\eta)]^{h(s)}.
$$

However, from [\(4.7\)](#page-20-0),

$$
h'\left(\left[\xi,\eta\right]^{(s)}\right) = h'\left(\xi\right)\left\langle \eta,x\right\rangle - h'\left(\eta\right)\left\langle \xi,x\right\rangle\tag{4.22}
$$

and

$$
[h'(\xi), h'(\eta)]^{h(s)} = h'(\xi) \langle h'(\eta), h(s)_x \rangle - h'(\eta) \langle h'(\xi), h(s)_x \rangle.
$$
 (4.23)

Hence, in particular, for any $\eta \in \mathfrak{l}$ and $x \in S^n$,

$$
\langle \eta, x \rangle = \langle h'(\eta), h(s)_x \rangle. \tag{4.24}
$$

Suppose now ξ is a unit vector in l. Noting that $h(t\xi \cdot s) = (th'(\xi)) \cdot h(s)$, from [\(4.11\)](#page-22-0) we see $h(s(t)) = h(t\xi \cdot s)$ satisfies the following equations

$$
\begin{cases} \frac{dh(s(t))_x}{dt} = h'(\xi) - \hat{\xi}h(s(t))_x\\ \frac{dh(s(t))_x}{dt} = \hat{\xi} \end{cases} (4.25)
$$

From [\(4.24\)](#page-25-0), we see that $\langle th'(\xi), h(s(t))_{x} \rangle = \langle t\xi, x(t) \rangle$, and hence $\hat{\xi}$ is unchanged from [\(4.11\)](#page-22-0). Taking inner product of the first equation of [\(4.25\)](#page-25-1) with $h'(\xi)$ yields

$$
\frac{d\hat{\xi}}{dt} = |h'(\xi)|^2 - \hat{\xi}^2.
$$

However, comparing with [\(4.14\)](#page-22-1), we see that $|h'(\xi)| = 1$. Therefore, $h' \in$ $SO(n + 1)$, and hence from (4.24) we see that $h(s(t))_x = h'(x(t))$.

We now need to understand how h transforms the ϕ component of s. From [\(4.12\)](#page-22-5) and [\(4.13\)](#page-22-4), for $\xi \in I$, let us denote the x and ϕ components of $\xi \cdot s$ by $\xi \cdot x$ and $\phi + \phi_{\xi}$, respectively. Define $\tilde{h}_x(\phi) = h(s)_{\phi}$ to be the transformation of the ϕ component, so that $h(s) = h(x, \phi) = (h'(x), \tilde{h}_x(\phi))$. In particular,

$$
h(\xi \cdot s) = h(\xi \cdot x, \phi + \phi_{\xi})
$$

=
$$
(h'(\xi \cdot x), \tilde{h}_{\xi \cdot x}(\phi + \phi_{\xi})).
$$

On the other hand, from the definition of h ,

$$
h(\xi \cdot s) = h'(\xi) \cdot h(s)
$$

= $h'(\xi) \cdot \left(h'(x), \tilde{h}_x(\phi) \right)$
= $\left(h'(\xi) \cdot h'(x), \tilde{h}_x(\phi) + \phi_{h'(\xi)} \right)$

We already know that $h'(\xi \cdot x) = h'(\xi) \cdot h'(x)$, and moreover, since $\langle \xi, x \rangle =$ $\langle h'(\xi), h'(x) \rangle, \phi_{h'(\xi)} = \phi_{\xi}.$ Thus we see that

$$
\tilde{h}_{\xi \cdot x} \left(\phi + \phi_{\xi} \right) = \tilde{h}_{x} \left(\phi \right) + \phi_{\xi}. \tag{4.26}
$$

.

 $(h'(x), \tilde{h}_x(\phi + t))$. However, from the definition of h, Now from [\(4.12\)](#page-22-5), we see that taking $\xi = x$, $(tx) \cdot s = (x, \phi + t)$, so $h((tx) \cdot s) =$

$$
h((tx) \cdot s) = th'(x) \cdot h(s)
$$

= $(h'(x), \tilde{h}_x(\phi) + t).$

Hence, we see that for any x, ξ , and $t, \tilde{h}_x(\phi + t) = \tilde{h}_x(\phi) + t$. This shows that $\tilde{h}_x(\phi) = \phi + c(x)$. From [\(4.26\)](#page-25-2), we then see that

$$
\phi + \phi_{\xi} + c(\xi \cdot x) = \phi + c(x) + \phi_{\xi},
$$

which shows that $c(x)$ is independent of x. Hence, the transformation of ϕ is just a transformation of S^1 by an action of $U(1)$. Overall, we then see that $\Psi(\mathbb{L}) \cong SO(n+1) \times U(1)$ and $\Psi'(\mathbb{L}) \cong SO(n+1)$.

Consider now more generally, $\mathbb{L} = S^m \times N$ where N is any n-dimensional parallelizable manifold. Suppose $\{T_1, ..., T_n\}$ is a frame of vector fields on N that satisfy the following Lie bracket relations:

$$
[T_A, T_B] = D_{AB}{}^C T_C,\tag{4.27}
$$

where the D_{AB}^C for $A, B, C = 1, ..., n$ are functions on N. The idea is to take a non-trivial vector field T on N and add the trivial rank 1 bundle generated by it to TS^m . The resulting rank $(m+1)$ -bundle then be trivialized in the same way as in the previous example of $S^m \times S^1$ [\[3\]](#page-30-7). Without loss of generality, let $T = T_1$. As before, consider $S^m \subset \mathbb{R}^{m+1}$ and for $i = 1, ..., n+1$, define

$$
f_i = M_i + x_i T,\t\t(4.28)
$$

where M_i is given by [\(4.2\)](#page-19-2). More explicitly, the f_i are given by

$$
f_i = \partial_{x_i} - x_i \sum_{j=1}^{m+1} x_j \partial_{x_j} + x_i T,
$$
\n(4.29)

Then, the set $\{f_1, ..., f_{n+1}, T_2, ..., T_n\}$ is global frame on $S^m \times N$. From [\(4.28\)](#page-26-0), we see that

$$
T = x^i f_i. \tag{4.30}
$$

The basis elements $\{f_1, ..., f_{n+1}\}$ satisfy the following bracket relation

$$
[f_i, f_j] = x_i f_j - x_j f_i.
$$
\n(4.31)

Also, for $A, B = 2, ..., n$,

$$
[f_i, T_A] = [x_i T_1, T_A] = x_i D_{1A}^{-1} T_1 + x_i D_{1A}^{-B} T_B
$$

Now let I be a $(m + n)$ -dimensional real vector space, with an inner product $\langle \cdot, \cdot \rangle$ and an orthonormal basis $\{e_i, e'_A\}$, where $i = 1, ..., m + 1$ and $A = 2, ..., n$. Then, for each $s = (x, y) \in \mathbb{L}$, define

$$
\rho_s: \mathfrak{l} \longrightarrow T_s \mathbb{L} \neq_i \mapsto f_i \vert_s ,\ne'_{A} \mapsto T_A \vert_s
$$
\n(4.32)

Since $T_1 = x^i f_i$, we see that $\rho_s^{-1}(T_1) = x^i e_i$. Hence define $e'_1 = x^i e_i$. As before, using [\(2.12\)](#page-8-0) and [\(4.31\)](#page-26-1), we define the bracket on I between basis elements e_i and e_i as:

$$
[e_i, e_j]^{(s)} = -\rho_s^{-1} ([f_i, f_j]]_s)
$$

= $e_i x_j - e_j x_i.$ (4.33)

Similarly, for $A = 2, ..., n$, and using [\(4.27\)](#page-26-2), we see that

$$
[e_i, e'_A]^{(s)} = -\rho_s^{-1} ([b_i, T_A]|_s)
$$

= $-x_i \rho_s^{-1} ([T_1, T_A])$
= $-x_i D_{1A}{}^{\tilde{C}} f_{\tilde{C}}$ (4.34)

for $\tilde{C} = 1, ..., n$. Moreover,

$$
[e_i, e'_1]^{(s)} = x^j [e_i, e_j]^{(s)}
$$

= $e_i - x_i e'_1$ (4.35)

$$
[e'_1, e'_A]^{(s)} = x^i [e_i, e'_A]^{(s)}
$$

= $-D_{1A}{}^{\tilde{C}} e'_{\tilde{C}}$ (4.36)

Now, for $B = 2, ..., n$,

$$
[e'_A, e'_B]^{(s)} = -\rho_s^{-1} ([T_A, T_B]]_s)
$$

= -D_{AB} \tilde{e}'_{\tilde{C}}. (4.37)

For $\tilde{A}, \tilde{B}, \tilde{C} = 1, ..., n$, we can then summarize

$$
[e'_{\tilde{A}}, e'_{\tilde{B}}]^{(s)} = -D_{\tilde{A}\tilde{B}}{}^{\tilde{C}} e'_{\tilde{C}}.
$$
\n(4.38)

Overall, we see that for $[e_i, e_j]^{(s)}$ we obtain the same structure constants as in the case of $S^m \times S^1$, for $\left[e'_{\tilde{A}}, e'_{\tilde{B}} \right]^{(s)}$, the structure constants are defined by the structure constants on the parallelizable space N , and only the mixed brackets [\(4.34\)](#page-27-0) and [\(4.35\)](#page-27-1) are new. We can combine these both in a single expression for $\tilde{A}, \tilde{C} = 1, ..., n$ as

$$
[e_i, e'_{\tilde{A}}]^{(s)} = -x_i D_{1\tilde{A}}{}^{\tilde{C}} e'_{\tilde{C}} + \delta_{1\tilde{A}} (e_i - x_i e'_1).
$$
 (4.39)

If we consider now the Jacobi identity for $[\cdot, \cdot]^{(s)}$, we find from Lemma [4.2,](#page-20-1) that the Jacobi identity is satisfied for e_i, e_j, e_k . Similarly, for $e'_{\tilde{A}}, e'_{\tilde{B}}, e'_{\tilde{C}}$, the

Jacobi identity is given by [\(2.24\)](#page-10-4). Now consider a mix of e_i and e'_A :

$$
[e_i,[e_j,e'_A]^{(s)}]^{(s)} = -[e_i,x_jD_{1A}\tilde{C}e'_{\tilde{C}}]^{(s)}
$$

\n
$$
= x_ix_jD_{1A}\tilde{C}D_{1\tilde{C}}\tilde{B}e'_{\tilde{B}} + x_ix_je'_1D_{1A}^1e_i - x_jD_{1A}^1e_i
$$

\n
$$
[e_j,[e'_A,e_i]^{(s)}]^{(s)} = -[e_j,[e_i,e'_A]^{(s)}]^{(s)} = -x_ix_jD_{1A}\tilde{C}D_{1\tilde{C}}\tilde{B}e'_{\tilde{B}} - x_ix_je'_1D_{1A}^1e_i + x_iD_{1A}^1e_j
$$

\n
$$
[e'_A,[e_i,e_j]^{(s)}]^{(s)} = [e'_A,e_ix_j - e_jx_i]^{(s)}
$$

\n
$$
= -x_ix_jD_{1A}\tilde{C}e'_{\tilde{C}} + x_ix_jD_{1A}\tilde{C}e'_{\tilde{C}} = 0
$$

Hence,

$$
[e_i,[e_j,e'_A]^{(s)}]^{(s)} + [e_j,[e'_A,e_i]^{(s)}]^{(s)} + [e'_A,[e_i,e_j]^{(s)}]^{(s)} = D_{1A}^{1}(x_ie_j - x_je_i)
$$

= $-D_{1A}^{1}[e_i,e_j]^{(s)}$

We see that whenever $D_{1A}^{\dagger} \neq 0$, the Jacobi identity is not satisfied, and the bracket $[\cdot, \cdot]^{(s)}$ does not define a Lie algebra. In particular, even if on N, the bracket algebra at $y \in N$ is a Lie algebra, if for some $A, D_{1A}^{-1}(y) \neq 0$, then the bracket algebra at $s \in \mathbb{L}$ is not a Lie algebra, and from (2.24) , $(1, a^{(s)})$ is not a Lie triple system.

As a particular example, consider $S^m \times S^n \times S^1$. We know $S^n \times S^1$ is parallelizable, and using the same parallelization on $S^n \times S^1$ as we did earlier on $S^m \times S^1$, but labelling the basis elements on $T(S^m \times S^1)$ as T_A for $A = 1, ..., n + 1$, from [\(4.6\)](#page-20-2), we find that in this case, for $s = (x, y, \phi)$, with x being the coordinates in $\mathbb{R}^{m+1} \supset S^m$, y being the coordinates in $\mathbb{R}^{n+1} \supset S^n$, and ϕ being the coordinate on ϕ ,

$$
[T_A, T_B]|_y = y_A T_B - y_B T_A.
$$

in this case, for $A \neq 1$,

$$
D_{1\dot{A}}^{-1}(y) = -y_A.
$$

Thus, as long as $y \neq (1, 0, ..., 0)$, this will be non-zero for some A, and the algebra of $[\cdot, \cdot]^{(s)}$ for $s = (x, y)$ will not satisfy the Jacobi identity.

A Appendix

Proof of Theorem [4.3.](#page-21-1) To complete the proof of Theorem [4.3,](#page-21-1) we need to show that given

$$
a^{(s)}(\xi, \eta, \gamma) = \gamma(\langle \xi, \eta \rangle + \langle x, \xi \rangle \langle x, \eta \rangle) - \eta(\langle \xi, \gamma \rangle + \langle x, \xi \rangle \langle x, \gamma \rangle), \tag{A.1}
$$

for any $\xi, \eta, \gamma, u, v \in I$,

$$
a_{\xi,\eta}^{(s)}\left(a^{(s)}(u,v,\gamma)\right) = a^{(s)}\left(a^{(s)}(u,\xi,\eta),v,\gamma\right) +a^{(s)}\left(u,a^{(s)}(v,\xi,\eta),\gamma\right) +a^{(s)}\left(u,v,a^{(s)}(\gamma,\xi,\eta)\right).
$$
\n(A.2)

Now, let

$$
A = a^{(s)}(u, v, \gamma) = \gamma (\langle u, v \rangle + \langle x, u \rangle \langle x, v \rangle) - v(\langle u, \gamma \rangle + \langle x, u \rangle \langle x, \gamma \rangle)
$$

$$
a^{(s)}(A,\xi,\eta)=\eta(\langle A,\xi\rangle+\langle x,A\rangle\langle x,\xi\rangle)-\xi(\langle A,\eta\rangle+\langle x,A\rangle\langle x,\eta\rangle),
$$

but

,

$$
\langle A, \xi \rangle = \langle \gamma, \xi \rangle \langle u, v \rangle + \langle \gamma, \xi \rangle \langle x, u \rangle \langle x, v \rangle - \langle v, \xi \rangle \langle u, \gamma \rangle - \langle v, \xi \rangle \langle x, u \rangle \langle x, \gamma \rangle
$$

\n
$$
\langle A, \eta \rangle = \langle \gamma, \eta \rangle \langle u, v \rangle + \langle \gamma, \eta \rangle \langle x, u \rangle \langle x, v \rangle - \langle v, \eta \rangle \langle u, \gamma \rangle - \langle v, \eta \rangle \langle x, u \rangle \langle x, \gamma \rangle
$$

\n
$$
\langle A, x \rangle = \langle \gamma, x \rangle \langle u, v \rangle + \langle \gamma, x \rangle \langle x, u \rangle \langle x, v \rangle - \langle v, x \rangle \langle u, \gamma \rangle - \langle v, x \rangle \langle x, u \rangle \langle x, \gamma \rangle
$$

\n
$$
= \langle \gamma, x \rangle \langle u, v \rangle - \langle v, x \rangle \langle u, \gamma \rangle.
$$

Thus,

$$
a^{(s)}\left(a^{(s)}(u,v,\gamma),\xi,\eta\right) = \eta\left(\langle\gamma,\xi\rangle\langle u,v\rangle + \langle\gamma,\xi\rangle\langle x,u\rangle\langle x,v\rangle - \langle v,\xi\rangle\langle u,\gamma\rangle - \langle v,\xi\rangle\langle x,u\rangle\langle x,\gamma\rangle\right) - \xi\left(\langle\gamma,\eta\rangle\langle u,v\rangle + \langle\gamma,\eta\rangle\langle x,u\rangle\langle x,v\rangle - \langle v,\eta\rangle\langle u,\gamma\rangle - \langle v,\eta\rangle\langle x,u\rangle\langle x,\gamma\rangle\right) + \left(\eta\langle x,\xi\rangle - \xi\langle x,\eta\rangle\right)\left(\langle\gamma,x\rangle\langle u,v\rangle - \langle v,x\rangle\langle u,\gamma\rangle\right)
$$

Similarly,

$$
a^{(s)}\left(a^{(s)}\left(u,\xi,\eta\right),v,\gamma\right) = \gamma\left(\langle\eta,v\rangle\langle u,\xi\rangle + \langle\eta,v\rangle\langle x,u\rangle\langle x,\xi\rangle - \langle v,\xi\rangle\langle u,\eta\rangle - \langle v,\xi\rangle\langle x,u\rangle\langle x,\eta\rangle\right) -v\left(\langle\gamma,\eta\rangle\langle u,\xi\rangle + \langle\gamma,\eta\rangle\langle x,u\rangle\langle x,\xi\rangle - \langle\xi,\gamma\rangle\langle u,\eta\rangle - \langle\xi,\gamma\rangle\langle x,u\rangle\langle x,\eta\rangle\right) + \left(\gamma\langle x,v\rangle - v\langle x,\gamma\rangle\right)\left(\langle\eta,x\rangle\langle u,\xi\rangle - \langle\xi,x\rangle\langle u,\eta\rangle\right)
$$

Now let

$$
B = a^{(s)}(v,\xi,\eta) = \eta(\langle v,\xi\rangle + \langle x,v\rangle\langle x,\xi\rangle) - \xi(\langle v,\eta\rangle + \langle x,v\rangle\langle x,\eta\rangle)
$$

$$
\langle B,u\rangle = \langle \eta,u\rangle\langle v,\xi\rangle + \langle \eta,u\rangle\langle x,v\rangle\langle x,\xi\rangle - \langle \xi,u\rangle\langle v,\eta\rangle - \langle \xi,u\rangle\langle x,v\rangle\langle x,\eta\rangle
$$

$$
\langle B,x\rangle = \langle \eta,x\rangle\langle v,\xi\rangle - \langle \xi,x\rangle\langle v,\eta\rangle
$$

and

$$
a^{(s)}(u, a^{(s)}(v, \xi, \eta), \gamma) = \gamma(\langle u, B \rangle + \langle x, u \rangle \langle x, B \rangle) - B(\langle u, \gamma \rangle + \langle x, u \rangle \langle x, \gamma \rangle)
$$

$$
= \gamma(\langle \eta, u \rangle \langle v, \xi \rangle + \langle \eta, u \rangle \langle x, v \rangle \langle x, \xi \rangle - \langle \xi, u \rangle \langle v, \eta \rangle - \langle \xi, u \rangle \langle x, v \rangle \langle x, \eta \rangle)
$$

$$
+ \gamma \langle x, u \rangle (\langle \eta, x \rangle \langle v, \xi \rangle - \langle \xi, x \rangle \langle v, \eta \rangle)
$$

$$
- \eta(\langle v, \xi \rangle + \langle x, v \rangle \langle x, \xi \rangle) (\langle u, \gamma \rangle + \langle x, u \rangle \langle x, \gamma \rangle)
$$

$$
+ \xi(\langle v, \eta \rangle + \langle x, v \rangle \langle x, \eta \rangle) ((u, \gamma) + \langle x, u \rangle \langle x, \gamma \rangle)
$$

Similarly,

$$
a^{(s)}(u, v, a^{(s)}(\gamma, \xi, \eta)) = -a^{(s)}(u, a^{(s)}(\gamma, \xi, \eta), v)
$$

\n
$$
= -v(\langle \eta, u \rangle \langle \gamma, \xi \rangle + \langle \eta, u \rangle \langle x, \gamma \rangle \langle x, \xi \rangle - \langle \xi, u \rangle \langle \gamma, \eta \rangle - \langle \xi, u \rangle \langle x, \gamma \rangle \langle x, \eta \rangle)
$$

\n
$$
-v \langle x, u \rangle (\langle \eta, x \rangle \langle \gamma, \xi \rangle - \langle \xi, x \rangle \langle \gamma, \eta \rangle)
$$

\n
$$
+ \eta(\langle \gamma, \xi \rangle + \langle x, \gamma \rangle \langle x, \xi \rangle) (\langle u, v \rangle + \langle x, u \rangle \langle x, v \rangle)
$$

\n
$$
- \xi(\langle \gamma, \eta \rangle + \langle x, \gamma \rangle \langle x, \eta \rangle) ((u, v) + \langle x, u \rangle \langle x, v \rangle)
$$

Adding together the terms $a^{(s)}(a^{(s)}(u,\xi,\eta),v,\gamma)$, $a^{(s)}(u,a^{(s)}(v,\xi,\eta),\gamma)$, $a^{(s)}(u, v, a^{(s)}(\gamma, \xi, \eta))$, we see that the contributions that are proportional to v and γ vanish. Consider the contributions of ξ :

$$
(\langle v, \eta \rangle + \langle x, v \rangle \langle x, \eta \rangle) (\langle u, \gamma \rangle + \langle x, u \rangle \langle x, \gamma \rangle) - (\langle \gamma, \eta \rangle + \langle x, \gamma \rangle \langle x, \eta \rangle) (\langle u, v \rangle + \langle x, u \rangle \langle x, v \rangle)
$$

\n
$$
= \langle v, \eta \rangle \langle u, \gamma \rangle + \langle v, \eta \rangle \langle x, u \rangle \langle x, \gamma \rangle + \langle x, v \rangle \langle x, \eta \rangle \langle u, \gamma \rangle + \langle x, v \rangle \langle x, \eta \rangle \langle x, u \rangle \langle x, \gamma \rangle
$$

\n
$$
- \langle \gamma, \eta \rangle \langle u, v \rangle - \langle \gamma, \eta \rangle \langle x, u \rangle \langle x, v \rangle - \langle x, \gamma \rangle \langle x, \eta \rangle \langle u, v \rangle - \langle x, \gamma \rangle \langle x, \eta \rangle \langle x, u \rangle \langle x, v \rangle
$$

\n
$$
= \langle v, \eta \rangle \langle u, \gamma \rangle - \langle \gamma, \eta \rangle \langle u, v \rangle + \langle v, \eta \rangle \langle x, u \rangle \langle x, \gamma \rangle + \langle x, v \rangle \langle x, \eta \rangle \langle u, \gamma \rangle
$$

\n
$$
- \langle \gamma, \eta \rangle \langle x, u \rangle \langle x, v \rangle - \langle x, \gamma \rangle \langle x, \eta \rangle \langle u, v \rangle
$$

This is precisely the coefficient of ξ in $a^{(s)}(a^{(s)}(u,v,\gamma),\xi,\eta)$. The expression is skew-symmetric in ξ and η , so the same conclusion follows. \blacksquare

References

- [1] I. Agricola and T. Friedrich. A note on flat metric connections with antisymmetric torsion. Differential Geom. Appl., 28(4):480–487, 2010. [doi:10.1016/j.difgeo.2010.01.004.](http://dx.doi.org/10.1016/j.difgeo.2010.01.004)
- [2] G. E. Bredon and A. Kosiński. Vector fields on π -manifolds. Ann. of Math. (2), 84:85–90, 1966. [doi:10.2307/1970531.](http://dx.doi.org/10.2307/1970531)
- [3] M. Bruni. On the explicit parallelization of products of spheres. Rend. Mat. Appl. (7), 12(2):405–423, 1992.
- [4] E. Cartan and J. Schouten. On Riemannian manifolds admitting an absolute parallelism. Proc. Amsterdam, 29:933–946, 1926.
- [5] L. P. Eisenhart. Spaces admitting complete absolute parallelism. Bull. Amer. Math. Soc., 39(4):217–226, 1933. [doi:10.1090/S0002-9904-1933-05593-3.](http://dx.doi.org/10.1090/S0002-9904-1933-05593-3)
- [6] S. Grigorian. Smooth loops and loop bundles. Adv. Math., 393:Paper No. 108078, 115, 2021, [2008.08120.](http://arxiv.org/abs/2008.08120) [doi:10.1016/j.aim.2021.108078.](http://dx.doi.org/10.1016/j.aim.2021.108078)
- [7] A. Hatcher. Vector Bundles and K-theory. 2017.
- [8] N. Jacobson. Lie and Jordan triple systems. Amer. J. Math., 71:149–170, 1949. [doi:10.2307/2372102.](http://dx.doi.org/10.2307/2372102)
- [9] M. Kervaire. Courbure intégrale généralisée et homotopie. Math. Ann., 131:219–252, 1956. [doi:10.1007/BF01342961.](http://dx.doi.org/10.1007/BF01342961)
- [10] M. Kikkawa. On local loops in affine manifolds. J. Sci. Hiroshima Univ. Ser. A-I Math., 28:199–207, 1964.
- [11] M. Kikkawa. System of local loops on a manifold and affine connection. Bull. Shimane Univ. Natur. Sci., (16):12–14, 1966.
- [12] J. Milnor. Topological manifolds and smooth manifolds. In Proc. Internat. Congr. Mathematicians (Stockholm, 1962), pages 132–138. Inst. Mittag-Leffler, Djursholm, 1963.
- [13] K. Nomizu and H. Ozeki. The existence of complete Riemannian metrics. Proc. Amer. Math. Soc., 12:889–891, 1961. [doi:10.2307/2034383.](http://dx.doi.org/10.2307/2034383)
- [14] M. Parton. Old and new structures on products of spheres. In Global differential geometry: the mathematical legacy of Alfred Gray (Bilbao, 2000), volume 288 of Contemp. Math., pages 406–410. Amer. Math. Soc., Providence, RI, 2001. [doi:10.1090/conm/288/04859.](http://dx.doi.org/10.1090/conm/288/04859)
- [15] M. Parton and P. Piccinni. Parallelizations on products of spheres and octonionic geometry. Complex Manifolds, 6(1):138–149, 2019. [doi:10.1515/coma-2019-0007.](http://dx.doi.org/10.1515/coma-2019-0007)
- [16] J. A. Wolf. On the geometry and classification of absolute parallelisms. I. J. Differential Geometry, 6:317–342, 1971/72. URL <http://projecteuclid.org/euclid.jdg/1214430496>.
- [17] J. A. Wolf. On the geometry and classification of absolute parallelisms. II. J. Differential Geometry, 7:19–44, 1972. URL <http://projecteuclid.org/euclid.jdg/1214430818>.