# RIGHT SPLITTING, GALOIS CORRESPONDENCE, GALOIS REPRESENTATIONS AND INVERSE GALOIS PROBLEM

CHANDRASHEEL BHAGWAT, SHUBHAM JAISWAL

Abstract. In this article, we realize some groups as Galois groups over rational numbers and finite extension of rational numbers by studying right splitting of some exact sequences, Galois correspondence and algebraic operations on Galois representations.

### 1. Introduction

The famous 'Inverse Galois problem' asks whether every finite group appears as the Galois group of some finite Galois extension of  $\mathbb{Q}$ . Many families of simple groups are known instances of this problem but the general question is still open.

Using Galois representations attached to elliptic curves, Arias-de-Reyna and König have proved in [1] that there are infinitely many number fields with Galois group  $\operatorname{GL}_2(\mathbb{F}_p)$  over  $\mathbb{Q}$  that are pairwise linearly disjoint over  $\mathbb{Q}(\sqrt{p^*})$  (see Remark 2.6.2).

In [8], the case of occurrence of  $PSL_2(\mathbb{F}_p)$  as a Galois group over  $\mathbb{Q}$  for all primes  $p \ge 5$  was established by Zywina using the results of Ribet [7] about the Deligne's Galois representations associated to certain newforms.

In this article, we establish some instances of Inverse Galois problem. In Sec. 2, through Galois correspondence and right splitting of some exact sequences of groups, we obtain the main results Thm. 2.4, Thm. 2.6. We then apply these general results to the case in [1] and obtain interesting consequences as corollaries.

In Sec. 3, using the algebraic operations on Galois representations and right splitting of some exact sequences of groups, we obtain Thm. 3.7, Thm. 3.8 and Thm. 3.19 as the main results. We then apply these general results to the case in [8] and obtain equally interesting consequences in Thm. 3.12 and the corollary of Thm. 3.19. Right Splitting of exact sequences of groups is the common thread that runs through the whole article.

#### 2. Groups, Galois Correspondence, Right Splitting and Inverse Galois problem

In this section, we describe some group theoretic results and then through these results as well as Galois correspondence and right splitting of some exact sequences, we obtain some general Galois theoretic results. Then we apply these general results to the cases described in [1] and obtain interesting consequences. Although some of the results related to groups and Galois correspondence may be known to experts, but we include them in our discussion for the sake of completeness.

Date: March 22, 2024.

<sup>2020</sup> Mathematics Subject Classification. 11R32, 12F12, 11F80.

# 2.1. Right splitting and group theoretic results.

**Proposition 2.1.** Let G be a finite group. Then the condition G/[G,G] is cyclic of order m is equivalent to the condition that for any n|m, G has a unique index-n normal subgroup such that the quotient is abelian (the quotient with that subgroup is in fact cyclic). This unique subgroup is given by

$$H = \{x \in G \mid \pi(x) \text{ is an } n\text{-th power in } G/[G,G]\}$$

where  $\pi$  is the quotient map  $G \to G/[G,G]$ .

*Proof.* Suppose G/[G, G] is a cyclic group of order m and  $n \mid m$ . Hence G/[G, G] has a unique index-n cyclic subgroup, call it H', which is precisely given by  $H' = \{z^n \mid z \in G/[G, G]\}$ , which is generated by  $y^n$ , where y is a generator of G/[G, G].

Since  $\pi$  is surjective, we have one to one correspondence between subgroups of G/[G,G] and subgroups of G containing [G,G]. Let  $H = \pi^{-1}(H') = \{x \in G | \pi(x) \text{ is an } n\text{-th power in } G/[G,G]\}$ . Hence it is the unique index-n normal subgroup of G containing [G,G].

Let *K* be any index-*n* normal subgroup of *G* such that the quotient is abelian. Consider the quotient map  $\rho : G \to G/K$ . Since  $\text{Image}(\rho)$  is abelian, hence  $[G, G] \subset Ker(\rho) = K$ . Hence K = H.

Also we have that G/H is isomorphic to (G/[G,G])/H'. Hence it is cyclic.

Conversely, suppose G/[G, G] is not cyclic. Hence, by structure theorem for abelian groups,

$$G/[G,G] \cong \mathbb{Z}/a_1\mathbb{Z} \times \mathbb{Z}/a_2\mathbb{Z} \times G$$

for some group G',  $a_1 \mid a_2$  and  $a_1 > 1$ . Since  $\mathbb{Z}/a_2\mathbb{Z}$  is cyclic and  $a_1 \mid a_2$ , we get a unique index- $a_1$  subgroup G'' of  $\mathbb{Z}/a_2\mathbb{Z}$ . Hence  $\mathbb{Z}/a_2\mathbb{Z} \times G'$  and  $\mathbb{Z}/a_1\mathbb{Z} \times G'' \times G'$  are distinct index- $a_1$  subgroups of G/[G, G]. Hence we don't get a unique index- $a_1$  subgroup H' of G/[G, G]. By the one to one correspondence between subgroups of G/[G, G] and subgroups of G containing [G, G], we get two distinct index- $a_1$  subgroups such that the quotients are abelian.

**Corollary 2.1.1.** Let  $n, r \in \mathbb{N}$  and p be an odd prime. For a prime power  $q = p^r$ , if  $q \equiv 1 \mod n$ , then  $\operatorname{GL}_2(\mathbb{F}_q)$  has a unique index-n normal subgroup such that the quotient is abelian. (The quotient with that subgroup is in fact cyclic). This unique subgroup is given by

$$H = \{ x \in \operatorname{GL}_2(\mathbb{F}_q) \mid det(x) \text{ is an } n\text{-th power in } \mathbb{F}_q^{\times} \}.$$

*Proof.* Let  $G = \operatorname{GL}_2(\mathbb{F}_q)$ . Then  $[G, G] = SL_2(\mathbb{F}_q)$  and  $G/[G, G] \cong \mathbb{F}_q^{\times}$  (isomorphism being through the surjective determinant map) which is cyclic of order m = q - 1. Now since  $q \equiv 1 \mod n$ , we have  $n \mid m$ . Now consider the determinant map  $det : G \to \mathbb{F}_q^{\times}$  with kernel  $SL_2(\mathbb{F}_q)$ . By applying similar argument to det map as applied to  $\pi$  map in previous proposition, we are done.

**Remark 2.1.1.** Above corollary holds for  $GL_k(\mathbb{F}_q)$  as well for any  $k \in \mathbb{N}$ .

**Corollary 2.1.2.** Let G be a group. If [G, G] has index m in G, then [G, G] is the unique index-m subgroup of G such that the quotient is abelian.

3

**Corollary 2.1.3.** Let  $r \in \mathbb{N}$  and p be an odd prime. For a prime power  $q = p^r$ ,  $SL_2(\mathbb{F}_q)$  is the unique index-(q-1) subgroup of  $GL_2(\mathbb{F}_q)$  such that the quotient is abelian.

**Proposition 2.2.** Let  $n \in \mathbb{N}$  and let G be a group such that G/[G,G] is cyclic of order m. Let n|m and let H be the unique index-n normal subgroup of G such that the quotient is abelian. Then the following holds.

(1) If there exists  $x \in G$  such that  $x^n = 1$  and  $\{1, x, x^2, ..., x^{n-1}\}$  is a set of representatives for *H*-cosets in *G*, then

$$(n, m/n) = gcd(n, m/n) = 1.$$

(2) If (n, m/n) = 1 and the exact sequence

$$1 \to [G,G] \to G \to G/[G,G] \to 1$$

is right split, then there exists  $x \in G$  such that  $x^n = 1$  and  $\{1, x, x^2, ..., x^{n-1}\}$  is a set of representatives for *H*-cosets in *G*.

(3) Let n = m (hence [G, G] is the unique index-m subgroup of G such that quotient is abelian). Then the above exact sequence is right split if and only if there exists  $x \in G$  such that  $x^m = 1$  and  $\{1, x, x^2, ..., x^{m-1}\}$  is a set of representatives for [G, G]-cosets in G.

### Proof.

(1) Suppose we have a set of representatives for *H*-cosets in *G* of the form  $\{1, x, x^2, ..., x^{n-1}\}$  with  $x \in G$  such that  $x^n = 1$ . Let *y* be a generator of cyclic group G/[G, G] and let  $0 \le l \le (m-1)$  be such that  $\pi(x) = y^l$ . As  $x^n = 1$ , we have  $1 = \pi(x)^n = y^{ln}$  and hence  $m \mid ln$ . Thus we have

$$j = ln/m \in \mathbb{Z}$$
 and  $1 \le j \le (n-1)$ .

Now assume on the contrary that  $(n, m/n) \neq 1$ . Then for  $k = n/(n, m/n) \leq (n - 1)$ , we have  $n \mid km/n$ , hence  $n \mid kl$ , and therefore  $\pi(x^k) = y^{lk}$  is an *n*-th power in G/[G, G], hence  $x^k \in H$ . Hence we get a contradiction. Furthermore, if  $(j, n) \neq 1$  then for k = n/(j, n) we have  $n \mid lk$ , which gives a contradiction. Hence (j, n) = 1.

- (2) If (n, m/n) = 1, then we clearly have that  $n \nmid km/n$  for all  $1 \le k \le (n-1)$ . Consider the quotient map  $\pi : G \to G/[G, G]$ . Since given exact sequence is right split, we have an injective homomorphism  $\iota : G/[G, G] \to G$  such that  $\pi \circ \iota = id_{G/[G,G]}$ . Let  $x = \iota(y^{jm/n}) \in G$  where  $1 \le j \le (n-1)$  and (j, n) = 1. Hence  $\pi(x) = y^{jm/n}$ . Since  $\iota$  is injective, x will be an element of order n. Now since  $n \nmid km/n$  for all  $1 \le k \le (n-1)$  and since (j, n) = 1, we have  $n \nmid jkm/n$  for all  $1 \le k \le (n-1)$ . Hence for all  $1 \le k \le (n-1)$ ,  $\pi(x^k) = y^{jkm/n}$  is not an n-th power in G/[G,G]. Hence  $x^k \notin H$ . Therefore we get a set of representatives for H-cosets in G of the form  $\{1, x, x^2, ..., x^{n-1}\}$  with  $x \in G$  such that  $x^n = 1$ .
- (3) Let us assume that the exact sequence is right split. Since n = m, we have (n, m/n) = (m, 1) = 1. Hence from (2), we get the required set of representatives for [G, G]-cosets in G.

Conversely, if there exists a set of representatives for [G, G]-cosets in G of the form  $\{1, x, x^2, ..., x^{m-1}\}$ with  $x \in G$  such that  $x^m = 1$ . Then clearly  $\pi(x) = x[G, G]$  is a generator of cyclic group G/[G, G]. Define  $\iota : G/[G,G] \to G$  by

 $\iota(\pi(g)) := x^r$  if  $\pi(g) \in G/[G,G]$  is of the form  $\pi(x)^r, r \ge 0$ .

Hence  $\iota$  is an injective homomorphism satisfying  $\pi \circ \iota = id_{G/[G,G]}$  and consequently the exact sequence is right split.

Remark 2.2.1. If G is abelian, then it satisfies the hypothesis of above proposition if and only if it is cyclic.

**Remark 2.2.2.** The condition that G/[G, G] is cyclic is important in statement (3) above. If *G* is a non-cyclic abelian group, then [G, G] = 1 and so exact sequence  $1 \rightarrow 1 \rightarrow G \rightarrow G \rightarrow 1$  is clearly right split, but we can never have required representatives for [G, G]-cosets because of non-cyclicity of *G*.

**Corollary 2.2.1.** Let  $n, r \in \mathbb{N}$ . Consider a prime power  $q = p^r$  such that  $q \equiv 1 \mod n$ . Let H be the unique index-n subgroup of  $GL_2(\mathbb{F}_q)$  such that the quotient is abelian. Then the following are equivalent.

- (n, (q-1)/n) = 1.
- There exists a set of representatives for *H*-cosets in  $GL_2(\mathbb{F}_q)$  of the form  $\{1, x, x^2, ..., x^{n-1}\}$  with  $x \in GL_2(\mathbb{F}_q)$  such that  $x^n = I$ .

*Proof.* Let  $G = GL_2(\mathbb{F}_q)$  then  $[G, G] = SL_2(\mathbb{F}_q)$  and  $G/[G, G] \cong \mathbb{F}_q^{\times}$  (isomorphism is defined through the determinant map) is cyclic of order m = q - 1. Since  $q \equiv 1 \mod n$ , we have  $n \mid m$ . Consider the exact sequence given by the determinant map

$$1 \to [G,G] \to G \xrightarrow{det} \mathbb{F}_q^{\times} \to 1.$$

Let  $\iota : \mathbb{F}_q^{\times} \to G$  be the map defined by

$$\iota(a) = \begin{pmatrix} 1 & 0 \\ 0 & a \end{pmatrix}$$
 for all  $a \in \mathbb{F}_q^{\times}$ .

This is a homomorphism satisfying  $det \circ \iota = id_{\mathbb{F}_q^{\times}}$ . Hence we conclude that the exact sequence is right split.

By applying similar argument to det map as applied to  $\pi$  map in previous proposition, we are done.

**Remark 2.2.3.** Above corollary holds for  $GL_k(\mathbb{F}_q)$  as well for any  $k \in \mathbb{N}$ . This can be proved using the map

$$\iota(a) = \begin{pmatrix} I_{k-1 \times k-1} & 0\\ 0 & a \end{pmatrix}_{k \times k} \text{ for all } a \in \mathbb{F}_q^{\times}.$$

**Proposition 2.3.** For given  $n, r \in \mathbb{N}$  such that (r, n) = 1, there are infinitely many primes satisfying the two conditions

$$p^r \equiv 1 \mod n,$$
$$(n, (p^r - 1)/n) = 1.$$

$$\frac{q-1}{n} \equiv k \bmod n.$$

Observe that (k, n) = 1 since (n, (q-1)/n) = 1. Hence  $q \equiv kn + 1 \mod n^2$ .

Conversely, each p satisfying  $p^r \equiv kn + 1 \mod n^2$  for some  $0 \le k \le (n-1)$  with (k, n) = 1, satisfies given conditions.

Now for each  $0 \le k \le (n-1)$  with (k, n) = 1, we have  $(kn + 1, n^2) = 1$ . Hence by Dirichlet's theorem on primes in arithmetic progression, we conclude that for each such k, there are infinitely many primes such that

$$p \equiv kn + 1 \mod n^2$$
,  
and hence  $p^r \equiv (kn + 1)^r \mod n^2$ .

Now for each such k, (rk, n) = 1, because (k, n) = (r, n) = 1. Hence there are infinitely many primes satisfying given conditions.

2.2. Galois correspondence, right splitting and Galois theoretic results. We discuss a result here that establishes the occurrence of certain semidirect product of groups as a Galois group.

**Theorem 2.4.** Let  $l \ge 2$ . Let K be a finite extension of  $\mathbb{Q}$  and let  $E_1, E_2, ..., E_l$  be finite Galois extensions of  $\mathbb{Q}$  contained in  $\overline{\mathbb{Q}}$ . Let  $G_i$  be the Galois group of  $E_i$  over  $\mathbb{Q}$  for all i. Suppose

- (1)  $E_1 \cdots E_k \cap E_{k+1} = K$  for all  $1 \le k \le l-1$ ,
- (2) for every *i*, there exists a set of representatives of  $H_i$ -cosets in respective  $G_i$  that is closed under multiplication, where  $H_i$  is Galois group of  $E_i$  over K.

Then for each  $1 \le i \le l$ ,  $H_i$  is a normal subgroup of respective  $G_i$  and the group  $(H_1 \times H_2 \times \cdots \times H_l) \rtimes G_1/H_1$ (for some semidirect product group law) is realizable as the Galois Group of  $E_1 \cdots E_l$  over  $\mathbb{Q}$ .

*Proof.* We will prove by induction that  $\operatorname{Gal}(E_1E_2\cdots E_k/K) \cong (H_1 \times H_2 \times \cdots \times H_k)$  and  $\operatorname{Gal}(E_1E_2\cdots E_k/\mathbb{Q}) \cong (H_1 \times H_2 \times \cdots \times H_k) \rtimes G_1/H_1$  (for some semidirect product group law) for all  $2 \le k \le l$ .

For base case k = 2, see the following diagram 2.2.

Since,  $E_1, E_2$  are Galois over  $\mathbb{Q}$  and are contained in  $\overline{\mathbb{Q}}$ ,  $E_1E_2$  and  $E_1 \cap E_2$  are defined and are Galois over Q and  $K = E_1 \cap E_2$ . We have  $\operatorname{Gal}(E_i/\mathbb{Q}) = G_i$  and  $\operatorname{Gal}(E_i/K) = H_i$  for i = 1, 2. Since K is Galois over  $\mathbb{Q}$ ,  $H_i$ s are normal in respective  $G_i$ s.  $\operatorname{Gal}(K/\mathbb{Q}) \cong G_1/H_1 \cong G_2/H_2$ . Fix an isomorphism  $\theta$  from  $G_2/H_2$  to  $G_1/H_1$ .



Let  $\operatorname{Gal}(K/\mathbb{Q}) = \{x_i H | 1 \leq i \leq n\}$  where  $x_i H_1 = \theta(y_i H_2)$  for all *i*. Here  $\{x_i \in G_1 | 1 \leq i \leq n\}$  and  $\{y_i \in G_2 | 1 \leq i \leq n\}$  are sets of representatives of  $H_1$  and  $H_2$  cosets in  $G_1$  and  $G_2$  respectively, which are closed under multiplication.

By Galois Correspondence Theorems 2.1 and 2.6 in [2], we have  $\operatorname{Gal}(E_1E_2/E_1) \cong H_2$ ,  $\operatorname{Gal}(E_1E_2/E_2) \cong H_1$ and  $\operatorname{Gal}(E_1E_2/K) \cong H_1 \times H_2$ .

Let  $\operatorname{Gal}(E_1E_2/\mathbb{Q}) = M$ . We have

$$|M| = [E_1 E_2 : K] [K : \mathbb{Q}] = |H_1 \times H_2| |G_1/H_1|$$
$$= |H_1 \times H_2| |G_2/H_2| = |G_1| |H_2| = |G_2| |H_1|.$$

Consider map  $\sigma : M \to G_1 \times G_2$  given by  $\sigma(g) = (g_1, g_2)$ , where  $g_i = g|_{E_i}$  for i = 1, 2. The map  $\sigma$  is clearly a well defined injective group homomorphism (see Thm 1.1 [2]).

If  $g \in M$  and  $g|_K = x_i H_1 = g_1|_K$ , then  $(x_i^{-1}g_1)|_K = id_K$ . Hence  $x_i^{-1}g_1 \in H_1$ , thus  $g_1 \in x_i H_1$ . Similarly, by  $g|_K = x_i H_1 = \theta(g_2|_K) = \theta(y_i H_2)$ ,  $g_2 \in y_i H_2$ .

Hence  $\operatorname{Image}(\sigma) \subset \bigsqcup_{i=1}^{n} (x_i H_1 \times y_i H_2)$ . Since  $|\bigsqcup_{i=1}^{n} (x_i H_1 \times y_i H_2)| = |H_1 \times H_2||G_1/H_1| = |M|$  and  $\sigma$  is injective, we get  $\operatorname{Image}(\sigma) = \bigsqcup_{i=1}^{n} (x_i H_1 \times y_i H_2)$ . Hence,  $M \cong \bigsqcup_{i=1}^{n} (x_i H_1 \times y_i H_2)$ .

Consider a map  $\psi : \bigsqcup_{i=1}^{n} (x_i H_1 \times y_i H_2) \to (H_1 \times H_2) \rtimes G_1/H_1$  given by  $\psi((x_i h_1, y_i h_2)) = ((h_1, h_2), x_i H_1)$ where we define a semidirect product group law for  $(H_1 \times H_2) \rtimes G_1/H_1$  by

$$((h_1, h_2), x_i H_1) \cdot ((k_1, k_2), x_j H_1) = ((x_j^{-1} h_1 x_j k_1, y_j^{-1} h_2 y_j k_2), x_i x_j H_1).$$

This group law is well defined and associative since  $H_i$  are normal in  $G_i$  for i = 1, 2 and sets of their respective coset representatives are closed under multiplication. We observe that

$$(x_ih_1, y_ih_2) \cdot (x_jk_1, y_jk_2) = (x_ih_1x_jk_1, y_ih_2y_jk_2) = ((x_ix_j)(x_j^{-1}h_1x_jk_1), (y_iy_j)(y_j^{-1}h_2y_jk_2)) \cdot (x_jh_1x_jh_2) \cdot (x_jh_1x_jh_2) \cdot (x_jh_1x_jh_2) = (x_jh_1x_jh_2) \cdot (x_jh_2x_jh_2) \cdot (x_jh_1x_jh_2) \cdot (x_jh_2x_jh_2) \cdot (x_jh_2x_jh_2x_jh_2) \cdot (x_jh_2x_jh_2x_jh_2) \cdot (x_jh_2x_jh_2x_jh_2x_jh_2) \cdot (x_jh_2x_jh_$$

Hence we conclude that  $\psi$  is a group isomorphism and hence  $M \cong (H_1 \times H_2) \rtimes G_1/H_1$ .

Alternatively, consider the sequence

$$1 \to (H_1 \times H_2) \xrightarrow{i} \bigsqcup_{i=1}^n (x_i H_1 \times y_i H_2) \xrightarrow{\pi} G_1/H_1 \to 1$$

where  $i(h_1, h_2) = (x_1h_1, y_1h_2)$  ( $x_1 \in H_1, y_1 \in H_2$ ) and  $\pi(x_ih_1, y_ih_2) = x_iH_1$ . The sequence is clearly exact.

Now consider  $\iota : G_1/H_1 \to \bigsqcup_{i=1}^n (x_iH_1 \times y_iH_2)$  with  $\iota(x_iH_1) = (x_i, y_i)$ . The map  $\iota$  is clearly a group homomorphism because of our multiplicatively closed assumption and also  $\pi \circ \iota = id_{G_1/H_1}$ . Hence the sequence is right split. Thus  $M \cong (H_1 \times H_2) \rtimes G_1/H_1$ .

Now assume that the statement is true for k = m, where  $2 \le m \le l - 1$ , that is,

$$\operatorname{Gal}(E_1E_2\cdots E_m/K)\cong (H_1\times H_2\times\cdots\times H_m),$$

and, 
$$\operatorname{Gal}(E_1E_2\cdots E_m/\mathbb{Q})\cong (H_1\times H_2\times\cdots\times H_m)\rtimes G_1/H_1$$

we will prove the statement for k = m + 1.

Let  $F_1 = E_1 \cdots E_m$  and  $F_2 = E_{m+1}$ . Hence

$$Gal(F_1/\mathbb{Q}) = Gal(E_1E_2\cdots E_m/\mathbb{Q}) \cong (H_1 \times H_2 \times \cdots \times H_m) \rtimes G_1/H_1,$$
  

$$Gal(F_2/\mathbb{Q}) = Gal(E_{m+1}/\mathbb{Q}) \cong G_{m+1},$$
  

$$Gal(F_1/K) = Gal(E_1E_2\cdots E_m/K) \cong (H_1 \times H_2 \times \cdots \times H_m),$$
  

$$Gal(F_2/K) = Gal(E_{m+1}/K) \cong H_{m+1} \operatorname{since} F_1 \cap F_2 = E_1 \cdots E_m \cap E_{m+1} = K.$$

Hence  $F_1$  and  $F_2$  satisfy conditions of base case and thus we have

$$Gal(E_1E_2\cdots E_{m+1}/K) = Gal(F_1F_2/K) \cong Gal(F_1/K) \times Gal(F_2/K)$$
$$\cong (H_1 \times H_2 \times \cdots \times H_m) \times H_{m+1} = H_1 \times H_2 \times \cdots \times H_{m+1}$$

and,

$$Gal(E_1E_2\cdots E_{m+1}/\mathbb{Q}) = Gal(F_1F_2/\mathbb{Q}) \cong Gal(F_1F_2/K) \rtimes G_1/H_1$$
$$\cong (H_1 \times H_2 \times \cdots \times H_{m+1}) \rtimes G_1/H_1.$$

### Remark 2.4.1.

(1) If [G : H] is finite, then a set of representatives for *H*-cosets in *G* satisfies the condition of closed under multiplication if and only if it is a subgroup of *G*.

In particular, A set of representatives  $\{1, x, x^2, ..., x^{n-1}\}$  for *H*-cosets in *G* which forms a cyclic subgroup of *G*, satisfies the condition. We get a criterion for existence of such a set of representatives in Prop 2.2.

(2) The second condition assumed in the theorem is equivalent to exact sequences

$$1 \to H_i \to G_i \to G_i/H_i \to 1$$

being right split, that is  $G_i \cong H_i \rtimes G_i/H_i$  for some semidirect product group law.

**Remark 2.4.2.** Let  $d_i$  be distinct primes for i = 1, 2 and  $d_3 = d_1d_2$ . Let  $K = \mathbb{Q}$  and consider quadratic extensions  $E_i = \mathbb{Q}\sqrt{d_i}$  for  $1 \le i \le 3$ .

Clearly  $E_i \cap E_j = K$  for all  $i \neq j$ , but we have  $E_i E_j \cap E_k = E_k \neq K$  where i, j, k are a permutation of 1, 2, 3. So the assumed condition in the above theorem is not always satisfied and hence it is important.

**Remark 2.4.3.** In the above theorem, we could have assumed a symmetric but stronger condition  $E_1...E_{i-1}E_{i+1}...E_n \cap E_i = K$  for all  $1 \le i \le n$  which implies the condition that we have assumed  $E_1...E_k \cap E_{k+1} = K$  for all  $1 \le k \le n-1$  since  $E_1...E_k \cap E_{k+1} \subset E_1...E_kE_{k+2}...E_n \cap E_{k+1} = K$ . But it would have made our theorem weaker. (Note: Condition for i = 1 is not required, it is written for symmetry).

# Remark 2.4.4.

- (1) Under the conditions of above theorem, we have  $(H_1 \times H_2 \times \cdots \times H_l) \rtimes G_1/H_1 \hookrightarrow G_1 \times G_2 \times \cdots \times G_l$  from Thm 1.1 [2].
- (2) If all G<sub>i</sub>s are abelian in above theorem that is each E<sub>i</sub> is an abelian extension of Q, then the compositum is an abelian extension over Q and the semidirect product that we defined is in fact the direct product.
- (3) However to the best of our knowledge, one doesn't know in general whether  $(H_1 \times H_2 \times \cdots \times H_l) \times G_1/H_1$  is realizable as a Galois group over  $\mathbb{Q}$ .

We now describe an independently interesting consequence of the Galois Correspondence (see Theorems 2.1 and 2.6 in [2].)

**Proposition 2.5.** Let finite groups  $H_1, H_2, ..., H_n$  be Galois groups over K of extensions  $E_1, E_2, ..., E_n$  respectively which are contained in  $\overline{\mathbb{Q}}$ . Fix  $k \leq n-1$  and suppose  $E_{i_1} \cdots E_{i_j} \cap E_{i_{j+1}} = K$  for all  $1 \leq j \leq k-1$  where  $i_l$  are distinct elements from 1 to k. Then the following statements are equivalent.

- (1)  $E_1 \cdots E_k \cap E_{k+1} = K$ .
- (2)  $E_{i_1} \cdots E_{i_k} \cap E_{i_{k+1}} = K$  where  $i_l$  are distinct elements from 1 to k + 1.
- (3)  $E_1 \cdots E_{i-1} E_{i+1} \cdots E_{k+1} \cap E_1 \cdots E_{j-1} E_{j+1} \cdots E_{k+1} = E_1 \cdots E_{i-1} E_{i+1} \cdots E_{j-1} E_{j+1} \cdots E_{k+1}$  for any  $1 \le i < j \le k+1$ .

*Proof.* Because of given conditions and induction we get,  $\operatorname{Gal}(E_{i_1} \cdots E_{i_j}/K) \cong H_{i_1} \times \ldots \times H_{i_j}$  for all  $j \leq k$  where  $i_l$  are distinct elements from 1 to k and  $\operatorname{Gal}(E_{i_1} \cdots E_{i_j}/E_{i_1} \cdots E_{i_{m-1}}E_{i_{m+1}} \cdots E_{i_j}) \cong H_{i_m}$  for all  $j \leq k$  and  $1 \leq m \leq j$  where  $i_l$  are distinct elements from 1 to k.

Equivalence of (1) and (2): For  $i_l$  distinct elements from 1 to k + 1,

$$E_1 \dots E_k \cap E_{k+1} = K.$$
  
$$\iff \operatorname{Gal}(E_1 \dots E_{k+1}/K) \cong H_1 \times \dots \times H_{k+1}.$$
  
$$\iff E_{i_1} \dots E_{i_k} \cap E_{i_{k+1}} = K.$$

Equivalence of (1) and (3): For any  $1 \le i < j \le k + 1$ , let

$$D = E_1 \cdots E_{k+1},$$
  

$$D_i = E_1 \cdots E_{i-1} E_{i+1} \cdots E_{k+1},$$
  

$$D_j = E_1 \cdots E_{j-1} E_{j+1} \cdots E_{k+1},$$
  

$$D_{ij} = E_1 \dots E_{i-1} E_{i+1} \dots E_{j-1} E_{j+1} \dots E_{k+1}.$$

Now since  $\operatorname{Gal}(D_i/D_{ij}) \cong H_j$  and  $\operatorname{Gal}(D_j/D_{ij}) \cong H_i$ , we have by Thm 1.1 [2],

$$\operatorname{Gal}(D/D_{ij}) \hookrightarrow H_i \times H_j.$$

We also have

$$\operatorname{Gal}(D/K)/\operatorname{Gal}(D/D_{ij}) \cong \operatorname{Gal}(D_{ij}/K)$$
$$\cong H_1 \times \cdots \times H_{i-1} \times H_{i+1} \times \cdots \times H_{j-1} \times H_{j+1} \times \cdots \times H_{k+1}.$$

Hence,

$$\operatorname{Gal}(D/K) \cong H_1 \times \dots \times H_{k+1}$$
$$\Longrightarrow \operatorname{Gal}(D/D_{ij}) \cong H_i \times H_j$$
$$\Longrightarrow D_i \cap D_j = D_{ij}.$$

Conversely,

$$D_i \cap D_j = D_{ij}$$

$$\implies D_i \cap E_i \subset D_i \cap D_j = D_{ij}$$

$$\implies D_i \cap E_i \subset D_{ij} \cap E_i = K$$

$$\implies D_i \cap E_i = K$$

$$\implies \operatorname{Gal}(D/K) \cong H_1 \times \cdots \times H_{k+1}.$$

### 2.3. Application to cases in work of Arias-de-Reyna & König.

**Theorem 2.6.** Let  $n, r \in \mathbb{N}$ . For a prime power  $q = p^r$ , suppose the following hold.

- (1)  $q \equiv 1 \mod n$  and (n, (q-1)/n) = 1.
- (2)  $\operatorname{GL}_2(\mathbb{F}_q)$  is realizable as Galois group over  $\mathbb{Q}$  of extensions  $E_1, E_2$  which are contained in  $\overline{\mathbb{Q}}$  such that  $E_1 \cap E_2 = K$  is a degree n abelian extension of  $\mathbb{Q}$ .

Then  $(H \times H) \rtimes \operatorname{GL}_2(\mathbb{F}_q)/H$  (with semidirect product group law as in Thm 2.4) is realizable as Galois group over  $\mathbb{Q}$  where H is the unique index-n subgroup of  $\operatorname{GL}_2(\mathbb{F}_q)$  such that the quotient is abelian.

*Proof.* We have for l = 2,  $E_1$ ,  $E_2$  satisfying conditions of Thm 2.4 with  $G_1 = G_2 = \text{GL}_2(\mathbb{F}_q)$ , and  $[G_1 : H_1] = n$  and  $G_1/H_1$  is abelian group and  $q \equiv 1 \mod n$ . Therefore  $H_1 = H_2 = H$ . Since  $q \equiv 1 \mod n$  and (n, (q-1)/n) = 1, from Corollary 2.2.1, we get the required set of representatives for *H*-cosets in  $\text{GL}_2(\mathbb{F}_q)$ .

**Remark 2.6.1.** Because of the conditions on p and n in above theorem, we actually get that K is a cyclic extension (not just abelian).

If extensions  $E_1$ ,  $E_2$  of  $\mathbb{Q}$  contained in  $\overline{\mathbb{Q}}$ , are linearly disjoint over an extension K of  $\mathbb{Q}$ , then  $E_1 \cap E_2 = K$ . We get from Remark 5.1 and Corollary 4.3 of [1], that there are two Galois extensions  $L_1$ ,  $L_2$  over  $\mathbb{Q}$  with Galois group  $\operatorname{GL}_2(\mathbb{F}_p)$  which are linearly disjoint over  $\mathbb{Q}(\zeta_p)$ . **Corollary 2.6.1.** For a prime  $p \ge 5$ ,  $(SL_2(\mathbb{F}_p) \times SL_2(\mathbb{F}_p)) \rtimes \mathbb{Z}/(p-1)\mathbb{Z}$  (with semidirect product group law as in above theorem) is realizable as Galois group over  $\mathbb{Q}$ .

*Proof.* From Remark 5.1 and Corollary 4.3 [1], we get  $E_1$ ,  $E_2$  satisfying conditions of previous theorem with n = p - 1,  $K = \mathbb{Q}(\zeta_p)$  and  $\operatorname{GL}_2(\mathbb{F}_p)/H$  is cyclic group of order p - 1. Moreover from Corollary 2.1.3, we have  $H = SL_2(\mathbb{F}_p)$ .

**Remark 2.6.2.** It is proved in Thm 1.1 of [1] that for a prime  $p \ge 5$ , there are infinitely many locally cyclic Galois extensions of  $\mathbb{Q}$  with Galois group  $\operatorname{GL}_2(\mathbb{F}_p)$ , which are pairwise linearly disjoint over  $\mathbb{Q}(\sqrt{p^*})$  where  $p^* = (-1)^{(p-1)/2}p$ .

**Corollary 2.6.2.** For a prime  $p \ge 5$  with  $p \equiv 3 \mod 4$ , let H be the unique index-2 (hence normal) subgroup of  $\operatorname{GL}_2(\mathbb{F}_p)$  then  $(H \times H) \rtimes \operatorname{GL}_2(\mathbb{F}_p)/H$  (with semidirect product group law as in above theorem) is realizable as Galois group over  $\mathbb{Q}$ .

*Proof.* From Thm 1.1 [1], we get  $E_1, E_2$  satisfying conditions of previous theorem with n = 2. Since  $p \equiv 1 \mod 2$ , the conditions  $p \equiv 3 \mod 4$  and (2, (p-1)/2) = 1 are equivalent. Hence we are done.

**Remark 2.6.3.** If  $p \equiv 1 \mod 4$  then (2, (p-1)/2) = 2. Hence by Corollary 2.2.1 there is no  $x \in GL_2(\mathbb{F}_p)$  such that  $x \in H$  and order(x) = 2.

**Corollary 2.6.3.** For a prime  $p \ge 5$  with  $p \equiv 3 \mod 4$ , let H be the unique index-2 subgroup of  $GL_2(\mathbb{F}_p)$ . Then H and  $H \times H$  are realizable as Galois groups over  $\mathbb{Q}\sqrt{-p}$ .

*Proof.* From previous corollary and proof of Thm 2.4, we have that  $H \times H$  is realizable as Galois group over  $\mathbb{Q}\sqrt{p^*} = \mathbb{Q}\sqrt{-p}$  ( $p \equiv 3 \mod 4$ ). Since H is normal in  $H \times H$  (where embedding is given by  $h \mapsto (h, 1)$ ), the quotient H is also realizable as Galois group over  $\mathbb{Q}\sqrt{-p}$ .

**Proposition 2.7.** For a prime  $p \ge 5$ ,  $(PSL_2(\mathbb{F}_p) \times PSL_2(\mathbb{F}_p)) \rtimes \mathbb{Z}/2\mathbb{Z}$  (with semidirect product group law as *in Thm 2.4*) is realizable as Galois group over  $\mathbb{Q}$ .

*Proof.* Consider  $H = \{x \in \operatorname{GL}_2(\mathbb{F}_p) \mid det(x) \text{ is a square in } \mathbb{F}_p^{\times}\}$ , which is the unique index-2 subgroup of  $\operatorname{GL}_2(\mathbb{F}_q)$  by Corollary 2.1.1. We have  $SL_2(\mathbb{F}_p) \subset H \subset \operatorname{GL}_2(\mathbb{F}_p)$ . Let  $h \in H$  and  $det(h) = \Delta_h^2$ . Then  $h = \Delta_h I \Delta_h^{-1} h$  where  $\Delta_h I \in Z(\operatorname{GL}_2(\mathbb{F}_p))$  and  $\Delta_h^{-1} h \in SL_2(\mathbb{F}_p)$ . Hence  $PH \cong PSL_2(\mathbb{F}_p)$  where isomorphism is given by the map sending  $hZ(\operatorname{GL}_2(\mathbb{F}_p))$  to  $(\Delta_h^{-1}h)Z(SL_2(\mathbb{F}_p))$ . Since H is the unique index-2 subgroup of  $\operatorname{GL}_2(\mathbb{F}_p)$  containing  $Z(\operatorname{GL}_2(\mathbb{F}_p))$ , we have that  $PH \cong PSL_2(\mathbb{F}_p)$  is the unique index-2 subgroup of  $P\operatorname{GL}_2(\mathbb{F}_p)$  by one to one correspondence between subgroups of  $\operatorname{GL}_2(\mathbb{F}_p)$  containing  $Z(\operatorname{GL}_2(\mathbb{F}_p))$  and subgroups of  $P\operatorname{GL}_2(\mathbb{F}_p)$ .

We have p > 3. Let  $r \neq p - 1$  be a non-square element in  $\mathbb{F}_p$ . Consider  $x = \begin{pmatrix} 1 & r+1 \\ -1 & -1 \end{pmatrix} \notin H$  with  $x^2 = 1 \left( Z(\operatorname{GL}_2(\mathbb{F}_p)) \right)$ .

From Thm 1.1 for  $PGL_2(\mathbb{F}_p)$  [1], we get  $E_1, E_2$  satisfying conditions of Thm 2.4 for l = 2 with  $G_1 = G_2 = PGL_2(\mathbb{F}_p)$ , and  $[G_1 : H_1] = 2$ . Therefore  $H_1 = H_2 = PSL_2(\mathbb{F}_p)$ . We observe that  $\{1, xZ(GL_2(\mathbb{F}_p))\}$  is required set of representatives of  $PSL_2(\mathbb{F}_p)$ -coset in  $PGL_2(\mathbb{F}_p)$ .

**Corollary 2.7.1.** For a prime  $p \ge 5$ ,  $PSL_2(\mathbb{F}_p)$  and  $PSL_2(\mathbb{F}_p) \times PSL_2(\mathbb{F}_p)$  are realizable as Galois groups over  $\mathbb{Q}\sqrt{p^*}$ .

*Proof.* Proof is similar to proof of Corollary 2.6.3.

# 3. Galois representations, right splitting and Inverse Galois problem

In this section, by using the algebraic operations induction, direct sums and tensor products on Galois representations and right splitting of some exact sequences of groups, we establish occurrence of some groups as Galois groups over  $\mathbb{Q}$ .

# 3.1. Induced Galois representations.

**Definition 3.1.** Let  $G = \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ . Let E be a number field and  $\Lambda$  be a prime in its ring of integers  $\mathcal{O}$ . Let  $E_{\Lambda}$  be the completion of E with respect to  $\Lambda$  and  $\mathcal{O}_{\Lambda}$  be its ring of integers. Let  $\mathbb{F}_{\Lambda} = \mathcal{O}_{\Lambda}/\Lambda \mathcal{O}_{\Lambda}$ .  $(\pi, V)$  is called a Galois Representation if

- (1) *V* is a finite dimensional vector space over  $E_{\Lambda}$  and
- (2)  $\pi: G \to \operatorname{GL}(V)$  is a continuous homomorphism.

**Remark 3.1.1.** In place of  $\mathbb{Q}$ , we can have a similar definition for any number field K with  $G = \operatorname{Gal}(\overline{K}/K)$ .

**Remark 3.1.2.** Let  $dim_{E_{\Lambda}}(V) = m$ . Now from Prop 9.3.5 in [3], there is a basis of V such that  $\pi(G) \subset$  $\operatorname{GL}_{m}(\mathcal{O}_{\Lambda})$  for all  $g \in G$ . Let us consider the maps  $\pi' : G \to \operatorname{GL}_{m}(\mathbb{F}_{\Lambda})$  and  $\tilde{\pi} : G \to P\operatorname{GL}_{m}(\mathbb{F}_{\Lambda})$  obtained from  $\pi$  through the quotients maps  $\mathcal{O}_{\Lambda} \to \mathbb{F}_{\Lambda}$  and  $\operatorname{GL}_{m}(\mathbb{F}_{\Lambda}) \to P\operatorname{GL}_{m}(\mathbb{F}_{\Lambda})$ .

**Definition 3.2.** A finite group *C* is said to be realizable as Galois group over  $\mathbb{Q}$  through a Galois Representation  $(\pi, V)$  if  $C \cong \text{Image}(\tilde{\pi})$ .

**Remark 3.2.1.** We have a similar definition for  $\pi'$  in place of  $\tilde{\pi}$ .

Let K be a finite extension of  $\mathbb{Q}$  contained in  $\overline{\mathbb{Q}}$ . Then  $\overline{K} = \overline{\mathbb{Q}}$ . Let  $H = \operatorname{Gal}(\overline{K}/K) \subset G$ . Let  $(\pi, W)$  be a Galois representation with  $\dim_{E_{\Lambda}}(W) = m$ .

Let  $\sigma = \pi|_H : H \to GL(W)$ . Then  $\sigma' : H \to GL_m(\mathbb{F}_\Lambda)$  and  $\tilde{\sigma} : H \to PGL_m(\mathbb{F}_\Lambda)$  and  $\operatorname{Image}(\tilde{\sigma}) = \tilde{\pi}(H)$ . Also,  $[\tilde{\pi}(G) : \tilde{\pi}(H)] \leq [G : H]$ .

Let us consider the induced representation  $\rho = Ind_H^G(\sigma)$  on the induced space over  $E_{\Lambda}$ , defined by

$$V = \{f : G \to W \mid f(hg) = \sigma(h)f(g) \text{ for all } h \in H, g \in G\}$$

and  $\rho: G \to \operatorname{GL}(V)$  is given by  $\rho(g)f(x) = f(xg)$ , for all  $g, x \in G, f \in V$ . Thus  $(\rho, V)$  is a Galois representation.

Let  $\{s_i\}_{1 \le i \le n}$  be a set of representatives of right cosets in  $H \setminus G$  and  $\{w_j\}_{1 \le j \le m}$  be a basis of W. Then we know that  $\{\phi_{s_i,w_j}\}_{1 \le i \le n, 1 \le j \le m}$  is a basis for V where

$$\phi_{s,w}(g) = \begin{cases} \sigma(gs^{-1})w, & \text{if } gs^{-1} \in H \\ 0 & \text{otherwise.} \end{cases}$$

and dim(V) = dim(W)[G:H].

Let *H* be a normal subgroup of *G* (i.e., *K* is Galois over  $\mathbb{Q}$ ) such that [G : H] = n. Since dim(W) = m, we have dim(V) = mn. Without loss of generality, let  $s_1 = 1$ . We label the basis of *V* as

$$f_1 = \phi_{1,w_1}, \ f_2 = \phi_{1,w_2}, \ \dots, \ f_m = \phi_{1,w_m},$$
$$f_{m+1} = \phi_{s_2,w_1}, \ \dots, \ f_{2m} = \phi_{s_2,w_m}, \dots,$$
$$f_{(n-1)m+1} = \phi_{s_n,w_1}, \ \dots, \ f_{nn} = \phi_{s_n,w_m}.$$

We write down the matrix of  $\rho(g)$  with respect to the above basis. Observe that

$$\rho(g)f_i(x) = f_i(xg) = \sum_{j=1}^{mn} a_{ji}f_j(x)$$

By taking  $x = s_k$ , we get that

$$\rho(g) = \begin{bmatrix}
(f_1(g)) & (f_2(g)) & \dots & (f_{nm}(g)) \\
(f_1(s_2g)) & (f_2(s_2g)) & \dots & (f_{nm}(s_2g)) \\
\dots & \dots & \dots & \dots \\
(f_1(s_ng)) & (f_2(s_ng)) & \dots & (f_{nm}(s_ng))
\end{bmatrix}_{nm \times n}$$

where  $(f_i(s_kg))$  are treated as  $m \times 1$  column matrices (matrices with respect to given basis  $\{w_j : 1 \le j \le m\}$  of W).

The  $nm \times nm$  matrix for  $\rho(hs_i)$ : The p, q-th  $m \times m$  block where  $1 \le p, q \le n$  and q is such that  $s_p s_i \in s_q H$  is  $\pi(s_p hs_i s_q^{-1})$ , since  $\phi_{s_q,w}(s_p hs_i) = \pi(s_p hs_i s_q^{-1})w$ , since  $s_p hs_i \in s_q H$ .

Since  $\pi(G) \subset \operatorname{GL}_m(\mathcal{O}_{\Lambda})$ , we conclude that  $\rho(G) \subset \operatorname{GL}_{nm}(\mathcal{O}_{\lambda})$  and  $\rho' : G \to \operatorname{GL}_{nm}(\mathbb{F}_{\Lambda})$  and  $\tilde{\rho} : G \to P\operatorname{GL}_{nm}(\mathbb{F}_{\Lambda})$  can be defined.

**Remark 3.2.2.** All the following results for  $\rho$  and  $\pi$  in this subsection also hold for  $\rho'$  and  $\pi'$  as well as  $\tilde{\rho}$  and  $\tilde{\pi}$ .

**Lemma 3.3.** *Images*  $\rho(H)$  *and*  $\pi(H)$  *are isomorphic.* 

*Proof.* If  $g = h \in H$ , then

$$\rho(h) = \begin{bmatrix} \pi(h) & 0 & \dots & \\ 0 & \pi(s_2 h s_2^{-1}) & & \\ \vdots & & \ddots & \\ & & & \pi(s_n h s_n^{-1}) \end{bmatrix}_{nm \times nm}.$$

Define a map from  $\rho(H)$  to  $\pi(H)$  by first block projection sending  $\rho(h)$  to  $\pi(h)$ . This is clearly a surjective group homomorphism onto  $\pi(H)$ . If  $\pi(h) = 1$ , then  $\pi(s_k h s_k^{-1}) = \pi(s_k)\pi(h)\pi(s_k)^{-1} = 1$  for all k. Hence, this map is injective too.

Lemma 3.4. We get an exact sequence

$$1 \to \rho(H) \to \rho(G) \to G/H \to 1.$$

*Proof.* If  $g = hs_j \in Hs_j$ ,  $(f_{(j-1)m+1}(g)) = \pi(h)w_1$ ,  $(f_{(j-1)m+2}(g)) = \pi(h)w_2$ , ...,  $(f_{jm}(g)) = \pi(h)w_m$  and for other k,  $(f_k(g)) = 0$ . We have

$$\rho(G) = \rho(H) \bigsqcup \rho(Hs_2) \bigsqcup \cdots \bigsqcup \rho(Hs_n) = \rho(H) \bigsqcup \rho(H)\rho(s_2) \bigsqcup \cdots \bigsqcup \rho(H)\rho(s_n)$$

hence  $[\rho(G) : \rho(H)] = n$ .

Now, since *H* is normal in *G*, its left and right cosets coincide. Hence we can define a map  $\gamma : \rho(G) \to G/H$ with  $\gamma(\rho(hs_k)) = s_k H$  for any  $h, s_k$ . Now  $\gamma$  is well defined surjective homomorphism because  $\rho(Hs_i)$  for  $1 \le i \le n$  are disjoint. The kernel of  $\gamma$  is precisely  $\rho(H)$ .

**Remark 3.4.1.** We do not necessarily get a similar exact sequence involving  $\pi(H)$ ,  $\pi(G)$  and G/H a similar well defined surjective map from  $\pi(G)$  to G/H may not exist since  $\pi(Hs_i)$  need not be disjoint.

**Example 3.5.** Let G/H be cyclic with representatives of H-cosets in G of the form  $\{1, s, s^2, \ldots, s^{n-1}\}$ . If  $g = hs^i \in H$  for  $0 \le i \le n-1$ , the matrix  $\rho(hs^i)$  is given by

$$\begin{bmatrix} 0 & \dots & 0 & \pi(h) & 0 & \dots \\ \vdots & & 0 & \pi(shs^{-1}) & 0 & \dots \\ & \ddots & & \vdots & \ddots & \\ \pi(s^{n-i}hs^i) & & & \pi(s^{n-1-i}hs^{-(n-1-i)}) \\ 0 & \pi(s^{n-(i-1)}hs^{(i-1)}) & & & \vdots \\ \vdots & 0 & \ddots & & \\ & & 0 & \pi(s^{n-1}hs) \end{bmatrix}$$

In particular,

	0	$I_2$	0		]
	0	0	$I_2$	0	
$\rho(s) =$	:	÷		·	
					$I_2$
	$\pi(s^n)$				0

**Lemma 3.6.** Suppose we have a set of representatives of *H*-cosets in *G*,  $\{s_1, s_2, \ldots, s_n\}$  with  $s_1 \in H$ , then  $\rho(s_i)\rho(s_j) = \rho(s_k) \iff (\pi(s_i)\pi(s_j) = \pi(s_k) \text{ and } s_is_j \in s_kH).$ 

*Proof.* We compute matrices  $\rho(s_i)$ ,  $\rho(s_j)$ ,  $\rho(s_k)$  nd  $\rho(s_i)\rho(s_j)$ .

The p, q-th  $m \times m$  block of  $\rho(s_i)$  is  $\pi(s_p s_i s_q^{-1})$ , where  $1 \le p, q \le n$  and q is such that  $s_p s_i \in s_q H$ . The q, r-th  $m \times m$  block of  $\rho(s_j)$  is  $\pi(s_q s_j s_r^{-1})$ , where  $1 \le q, r \le n$  and r is such that  $s_q s_j \in s_r H$ . The p, l-th  $m \times m$  block of  $\rho(s_k)$  is  $\pi(s_p s_k s_l^{-1})$ , where  $1 \le l \le n$  and l is such that  $s_p s_k \in s_l H$ . The p, r-th  $m \times m$  block of  $\rho(s_i)\rho(s_j)$  is  $\pi(s_p s_i s_q^{-1})\pi(s_q s_j s_r^{-1}) = \pi(s_p s_i s_j s_r^{-1})$ . Since  $s_p s_i \in s_q H$ ,  $s_q s_j \in s_r H$ , we get  $s_p s_i s_j \in s_r H$ . Hence  $\rho(s_i)\rho(s_j) = \rho(s_k) \iff (\pi(s_p s_i s_j s_r^{-1}) = \pi(s_p s_k s_l^{-1}) \text{ and } r = l) \iff (\pi(s_i)\pi(s_j) = \pi(s_k) \text{ and } s_i s_j \in s_k H).$ 

**Remark 3.6.1.**  $\pi(s_i)$  need not be distinct even though  $\rho(s_i)$  are distinct. In fact even if all  $\pi(s_i)$  are same,  $\rho(s_i)$  will be distinct.

**Corollary 3.6.1.** Let G/H be cyclic with representatives of H-cosets in G of the form  $\{1, s, s^2, \ldots, s^{n-1}\}$ . Then  $\rho(s)^n = 1$  if and only if  $\pi(s)^n = 1$ .

**Theorem 3.7.** Let G and H be as above, then the exact sequence

$$1 \to \rho(H) \to \rho(G) \to G/H \to 1$$

is right split (that is  $\rho(G) \cong \rho(H) \rtimes G/H$  for some semidirect product group law) if and only if there exists a set of representatives of *H*-cosets in *G*,  $\{s_1, s_2, \ldots, s_n\}$  with  $s_1 \in H$  such that  $\{\rho(s_i)\}_i$  forms a multiplicatively closed subset of  $\rho(G)$ . (In fact it is a subgroup with same group structure as  $\{s_iH\}_i = G/H$ )

*Proof.* Let  $\{r_1, r_2, ..., r_n\}$  be a set of representatives of H-cosets in G. If the exact sequence is right split, let  $\iota$  be splitting with  $\gamma \circ \iota = id_{G/H}$ . Hence  $\gamma \circ \iota(r_iH) = r_iH$  for each i. Thus for each i,  $\iota(r_iH) = \rho(h_ir_i)$  for some  $h_i \in H$ . Let  $s_i = h_ir_i$  for each i. Hence  $\{s_1, s_2, ..., s_n\}$  also forms a set of representatives for H-cosets in G with  $s_iH = r_iH$  and  $\iota(s_iH) = \rho(s_i)$  for each i. Since  $\iota$  is a homomorphism, we have  $\iota(s_iH)\iota(s_jH) = \iota(s_is_jH)$  for any i, j. Let  $s_is_jH = s_kH$  for some k. Hence  $\iota(s_iH)\iota(s_jH) = \iota(s_kH)$  that is  $\rho(s_i)\rho(s_j) = \rho(s_k)$ . Hence  $\{\rho(s_i)\}_i$  forms a multiplicatively closed subset of  $\rho(G)$ .

Conversely, suppose there exists a set of representatives of *H*-cosets in *G*,  $\{s_1, s_2, \ldots, s_n\}$  with  $s_1 \in H$ and  $\{\rho(s_i)\}_i$  forming a multiplicatively closed subset of  $\rho(G)$ . Let  $i \neq 1$ . Then  $\rho(s_1)\rho(s_i) = \rho(s_k)$  for some *k*. Hence,  $\rho(s_1s_i) = \rho(s_k)$ . Since  $s_1 \in H$ , we have  $s_1s_i \in s_iH$ . Hence by matrix calculation done above,  $\rho(s_1s_i) = \rho(s_k)$  is not true unless k = i. Hence  $\rho(s_1)\rho(s_i) = \rho(s_i)$ . Thus  $\rho(s_1) = 1$ . For any *i*, consider the set  $\{\rho(s_j)\rho(s_i)\}_j$  which is a permutation of the set  $\{\rho(s_j)\}_j$ . Hence there is a *j* such that  $\rho(s_j)\rho(s_i) = 1$ . Hence  $\{\rho(s_i)\}_i$  forms a subgroup of  $\rho(G)$ .

Now by assumption  $\rho(s_i)\rho(s_j) = \rho(s_l)$  for some l. Hence  $\rho(s_is_j) = \rho(s_l)$ . Suppose  $s_iHs_jH = s_is_jH = s_kH$ . Then by matrix calculation done above,  $\rho(s_is_j) = \rho(s_l)$  is not true unless l = k. Hence  $\rho(s_is_j) = \rho(s_k)$ . Hence  $\{\rho(s_i)\}_i$  is a group with the same group structure as  $\{s_iH\}_i = G/H$ .

Then we can define  $\iota(s_iH) = \rho(s_i)$ . Hence,  $\iota(s_iH)\iota(s_jH) = \rho(s_i)\rho(s_j) = \rho(s_k) = \iota(s_kH) = \iota(s_iHs_jH)$ . Hence  $\iota$  is a homomorphism. Also,  $\gamma(\iota(s_iH)) = \gamma(\rho(s_i)) = s_iH$  hence  $\gamma \circ \iota = id_{G/H}$ . Hence the exact sequence is right split.

Remark 3.7.1. We can similarly prove Remark 2.4.1 in previous section.

**Corollary 3.7.1.** If there exists a set of coset representatives of H in G which forms a multiplicatively closed set, then the above exact sequence is right split.

**Corollary 3.7.2.** Let *K* be a cyclic extension of  $\mathbb{Q}$  and let *G* and *H* be as above. Then the above exact sequence is right split if and only if there exists a set of representatives of *H*-cosets in *G* of the form  $\{1, s, s^2, \ldots, s^{n-1}\}$  with  $\rho(s)^n = 1$ . Also, if these statements are true then  $order(\rho(s)) = n$  and  $s^n \in H$ .

14

*Proof.* From the above theorem, we get coset representatives of H,  $\{s_i\}_i$  with  $\{\rho(s_i)\}_i$  forming a subgroup with same group structure as  $\{s_iH\}_i = G/H = \langle sH \rangle$  for some  $s \in G$  with  $s_1 \in H$ . Hence without loss of generality, let  $s_iH = s^{i-1}H$  for all i. Now let  $r_i = s_2^{i-1}$ . Hence  $r_iH = s_iH$ . Now since  $s_2^nH = (s_2H)^n = H$ , we have  $\tilde{\rho}(s_2)^n = 1$ . Hence we are done. Other assertions are clear.

Applying above discussion to  $\tilde{\rho}$  and  $\tilde{\pi}$  we have the following.

**Theorem 3.8.** Let H be a finite index normal subgroup of  $G = Gal(\overline{\mathbb{Q}}/\mathbb{Q})$ . Suppose there exists a set of coset representatives of H in G which form a multiplicatively closed subset of G. Then if a finite group M is realizable as a Galois group over  $\mathbb{Q}$  through a Galois Representation  $(\pi, W)$  such that  $M \cong \tilde{\pi}(G) = \tilde{\pi}(H)$ , then  $M \rtimes G/H$  is realisable as a Galois group over  $\mathbb{Q}$  (for semidirect product group law as in Thm 3.7).

**Observation** :  $G = H \iff ker(\pi) \subset H \text{ and } \pi(G) = \pi(H).$ 

We can generalize Lemmma 3.4 and Thm 3.7. Consider a closed subgroup H' of G. We have  $H'/(H' \cap H) \hookrightarrow G/H$ .

### Theorem 3.9.

(1) We get an exact sequence

$$1 \to \rho(H' \cap H) \to \rho(H') \to H'/(H' \cap H) \to 1.$$

- (2) This exact sequence is right split (that is ρ(H') ≅ ρ(H' ∩ H) ⋊ H'/(H' ∩ H) for some semidirect product group law) if and only if there exists a set of representatives of (H' ∩ H)-cosets in H', {s<sub>1</sub>, s<sub>2</sub>,..., s<sub>l</sub>} with s<sub>1</sub> ∈ H' ∩ H such that {ρ(s<sub>i</sub>)}<sub>i</sub> forms a multiplicatively closed subset of ρ(H').
- (3)  $H \subset H'$  if and only if  $\rho(H) \subset \rho(H')$  and  $ker(\pi) \cap H \subset H'$
- (4) Let  $H \subset H'$ . Then the exact sequence is

$$1 \to \rho(H) \to \rho(H') \to H'/H \to 1.$$

If there exists a set of representatives of *H*-cosets in *G*,  $\{s_1, s_2, \ldots, s_n\}$  with  $s_1 \in H$  such that  $\{\rho(s_i)\}_i$  forms a multiplicatively closed subset of  $\rho(G)$ , then this exact sequence is right split. That is  $\rho(H') \cong \rho(H) \rtimes H'/H$  for some semidirect product group law.

3.2. Galois representations for newforms and Application to Zywina's case. Let  $p \ge 5$ . Fix non-CM newform f of weight k > 1 on  $\Gamma_1(N)$ . Let l be rational prime as in [8, Zywina]. Let  $\pi$  be the Deligne representation associated with f with properties as in 1.1 of [8]. Let  $W = E_{\Lambda}^2$  and  $\pi : G \to GL(W)$  with  $\pi(G) \subset GL_2(\mathcal{O}_{\Lambda})$  and  $\tilde{\pi} : G \to PGL_2(\mathbb{F}_{\Lambda})$ .

Let p = l and assume conditions of [8, 1.2]. Then  $\text{Image}(\tilde{\pi}) = \tilde{\pi}(G) = \text{PSL}_2(\mathbb{F}_p)$ . The representations  $\sigma = \pi|_H : H \to \text{GL}(W)$  and  $\tilde{\sigma} : H \to \text{PGL}_2(\mathbb{F}_\Lambda)$  are defined, and  $\text{Image}(\tilde{\sigma}) = \tilde{\pi}(H) \subset \tilde{\pi}(G)$ .

**Lemma 3.10.** Let  $p \ge 5$  and H be normal in G (K is Galois) and  $[G : H] = n < |PSL_2(\mathbb{F}_p)|$ . Then  $\tilde{\pi}(H) = PSL_2(\mathbb{F}_p)$ .

*Proof.* Now,  $[PSL_2(\mathbb{F}_p) : \tilde{\pi}(H)] = [\tilde{\pi}(G) : \tilde{\pi}(H)] \le [G : H] = n < |PSL_2(\mathbb{F}_p)|$ . Hence  $\tilde{\pi}(H)$  is not trivial. Since H is normal in G,  $\tilde{\pi}(H)$  is normal in  $\tilde{\pi}(G)$ . Now since  $PSL_2(\mathbb{F}_p)$  is simple we have that  $PSL_2(\mathbb{F}_p) = \tilde{\pi}(H)$  otherwise  $\tilde{\pi}(H)$  will become non-trivial normal subgroup in  $PSL_2(\mathbb{F}_p)$ .

**Remark 3.10.1.** If  $n \ge |PSL_2(\mathbb{F}_p)|$  then either  $\tilde{\pi}(H) = PSL_2(\mathbb{F}_p)$  or  $\tilde{\pi}(H)$  is trivial. If latter case happens, then  $\tilde{\rho}(H)$  is also trivial. Hence  $\tilde{\rho}(G) \cong G/H$  in that case.

# **Proposition 3.11.**

- (1) For  $p \ge 5$ ,  $PGL_2(\mathbb{F}_p) \cong PSL_2(\mathbb{F}_p) \rtimes \mathbb{Z}/2\mathbb{Z}$  for some semidirect product group law and the semidirect product is not direct product.
- (2) Let p ≥ 5. For any semidirect product group law such that PGL<sub>2</sub>(𝔽<sub>p</sub>) ≃ PSL<sub>2</sub>(𝔽<sub>p</sub>) ⋊ℤ/2ℤ, the semidirect product is not direct product. Furthermore, the automorphism of PSL<sub>2</sub>(𝔽<sub>p</sub>), given by conjugation by image of generator of ℤ/2ℤ in PGL<sub>2</sub>(𝔽<sub>q</sub>), is not inner.

Proof. (1) From proof of Prop 2.7, we have an exact sequence

$$1 \to \mathrm{PSL}_2(\mathbb{F}_p) \to \mathrm{PGL}_2(\mathbb{F}_p) \to \mathbb{Z}/2\mathbb{Z} \to 1.$$

Let  $\iota : \mathbb{Z}/2\mathbb{Z} \to P\operatorname{GL}_2(\mathbb{F}_p)$  be given by  $\iota(\overline{1}) = xZ(\operatorname{GL}_2(\mathbb{F}_p))$  where x is as in Prop 2.7.  $\iota$  is a right splitting. Hence above exact sequence is right split. Hence  $P\operatorname{GL}_2(\mathbb{F}_p) \cong \operatorname{PSL}_2(\mathbb{F}_p) \rtimes \mathbb{Z}/2\mathbb{Z}$  for some semidirect product group law. Now since,  $\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} Z(\operatorname{GL}_2(\mathbb{F}_p)) \in PH$  doesn't commute with  $xZ(\operatorname{GL}_2(\mathbb{F}_p))$ , the semidirect product is not direct product.

(2) Let  $xZ(\operatorname{GL}_2(\mathbb{F}_p))$  be the image of generator of  $\mathbb{Z}/2\mathbb{Z}$  in  $P\operatorname{GL}_2(\mathbb{F}_q)$  for  $x \in \operatorname{GL}_2(\mathbb{F}_p)$ . Then  $x \notin H$ ,  $xZ(\operatorname{GL}_2(\mathbb{F}_p)) \notin PH$  and  $x^2 \in Z(\operatorname{GL}_2(\mathbb{F}_p))$ . Suppose the semidirect product was direct product. Then for some  $\nu_h \in Z(\operatorname{GL}_2(\mathbb{F}_p))$ ,  $xhx^{-1} = h\nu_h$  for all  $h \in H$ . Now,  $P\operatorname{GL}_2(\mathbb{F}_p) = PH \bigsqcup (xZ(\operatorname{GL}_2(\mathbb{F}_p)))PH$ . Hence,  $xZ(\operatorname{GL}_2(\mathbb{F}_p)) \in Z(P\operatorname{GL}_2(\mathbb{F}_p)) = \{1\}$ . Hence  $x \in H$ , which gives a contradiction.

Now, suppose automorphism of  $PSL_2(\mathbb{F}_p)$ , given by conjugation by  $xZ(GL_2(\mathbb{F}_p))$ , is inner. Then for some  $h' \in H$  and  $\mu_h \in Z(GL_2(\mathbb{F}_p))$ ,  $xhx^{-1} = h'hh'^{-1}\mu_h$  for all  $h \in H$ . Hence  $(h'^{-1}x)h(h'^{-1}x)^{-1} = h\mu_h$  for all  $h \in H$ . Now,  $PGL_2(\mathbb{F}_p) = PH \bigsqcup ((h'^{-1}x)Z(GL_2(\mathbb{F}_p)))PH$ . Hence,  $(h'^{-1}x)Z(GL_2(\mathbb{F}_p)) \in Z(PGL_2(\mathbb{F}_p)) = \{1\}$ . Hence  $x \in H$ , which gives a contradiction.

### Theorem 3.12.

- (1) For  $p \ge 5$ ,  $PSL_2(\mathbb{F}_p) \rtimes \mathbb{Z}/2\mathbb{Z}$  is realisable as Galois Group over  $\mathbb{Q}$  (for semidirect product group law in *Thm 3.7*).
- (2) This semidirect product in part (1) is direct  $\iff \tilde{\pi}(s) = I$ .
- (3) Automorphism  $\phi_{\tilde{\rho}(s)}$  of  $\tilde{\rho}(H)$ , by conjugation by  $\tilde{\rho}(s)$ , is inner.
- (4) The group obtained here is not isomorphic to  $PGL_2(\mathbb{F}_p)$ .

*Proof.* (1) Let [G : H] = 2. Then H is normal in G (K is Galois). Also  $2 < |PSL_2(\mathbb{F}_p)|$ . Hence by previous lemma,  $\tilde{\pi}(H) = PSL_2(\mathbb{F}_p)$ .

We could have chosen  $K = \mathbb{Q}(i)$  and  $s \in G$  as complex conjugation so that  $s^2 = 1$  and  $\{1, s\}$  become representatives of right cosets of *H* in *G*. Then by Thm 3.8 we are done.

(2) Now, 
$$\tilde{\pi}(G) = \tilde{\pi}(H) = \mathrm{PSL}_2(\mathbb{F}_p).$$

Above semidirect product is direct.

 $\iff \tilde{\rho}(s) = \begin{bmatrix} 0 & I_2 \\ I_2 & 0 \end{bmatrix} \text{ commutes with every } \tilde{\rho}(h) = \begin{bmatrix} \tilde{\pi}(h) & 0 \\ 0 & \tilde{\pi}(shs^{-1}) \end{bmatrix}.$  $\iff \tilde{\pi}(s)$  commutes with every  $\tilde{\pi}(h)$ .  $\iff \tilde{\pi}(s) \in Z(\mathrm{PSL}_2(\mathbb{F}_p)).$  $\iff \tilde{\pi}(s) = I.$ Since  $Z(PSL_2(\mathbb{F}_p))$  is trivial as  $PSL_2(\mathbb{F}_p)$  is non-abelian and simple. (3) Now  $\tilde{\pi}(G) = \tilde{\pi}(H)$ . Hence  $\tilde{\pi}(s) = \tilde{\pi}(h')$  for some  $h' \in H$ .  $\mathbf{So}, \phi_{\tilde{\rho}(s)}(\tilde{\rho}(h)) = \tilde{\rho}(s)\tilde{\rho}(h)\tilde{\rho}(s)^{-1} = \begin{bmatrix} \tilde{\pi}(shs^{-1}) & 0\\ 0 & \tilde{\pi}(h) \end{bmatrix} = \tilde{\rho}(h')\tilde{\rho}(h)\tilde{\rho}(h')^{-1}.$ 

(4) This follows from (3) and Prop 3.11.

We have results similar to (1), (2) and (3) of Thm 3.12 for certain simple groups of the form  $PSL_2(\mathbb{F}_q)$ .

**Theorem 3.13.** PSL<sub>2</sub>( $\mathbb{F}_q$ )  $\rtimes \mathbb{Z}/2\mathbb{Z}$  is realisable as Galois Group over  $\mathbb{Q}$  (for semidirect product group law in Thm 3.7) for following q.

- (1) q = p for  $p \ge 5$  (From 1.2 [8]).
- (2)  $q = p^2$  for  $p \equiv \pm 2 \mod 5$ ,  $p \ge 7$  (From Corollary 3.6 [5]).
- (3)  $q = p^2$  for  $p \equiv \pm 3 \mod 8$ ,  $p \ge 5$  (From Corollary 3.8 [5]).
- (4)  $q = p^3$  for odd prime  $p \equiv \pm 2, \pm 3, \pm 4, \pm 6 \mod 13$  (From 1.3 [8]).
- (5)  $q = 5^3, 3^5, 3^4$  (From 2.2, 2.5, 3 [4] respectively).

**Remark 3.13.1.** One has even more general conditions for  $q = p^2$  (See Thm 3.1 [6]) and  $q = p^4$  (See 3.3 [6]).

# 3.3. Direct sum / tensor product of representations and Galois groups.

**Lemma 3.14.** If for each  $1 \leq i \leq n$ , finite group  $H_i$  is realizable as a Galois group over  $\mathbb{Q}$  through Galois representation  $(\pi_i, V_i)$  for  $m_i$ -dimensional vector spaces  $V_i$  over  $E_{\Lambda}$ , then  $\{(\tilde{\pi}_1(g), ..., \tilde{\pi}_n(g)) \mid g \in G\} \subset H_1 \times$  $H_2 \times \cdots \times H_n$  is realisable as Galois group over  $\mathbb{Q}$ .

*Proof.* We have  $\pi_i : G \to \operatorname{GL}(V_i)$  with  $\pi_i(G) \subset \operatorname{GL}_{m_i}(\mathcal{O}_{\Lambda})$  and  $\tilde{\pi}_i : G \to \operatorname{PGL}_{m_i}(\mathbb{F}_{\Lambda})$  such that  $H_i \cong$ Image( $\tilde{\pi_i}$ ).

Consider Galois representations  $(\pi_i, V_i)$  :  $1 \le i \le n$  and their direct sum Galois representation  $\pi =$  $\bigoplus_{1 \le i \le n} \pi_i : G \to \operatorname{GL} \bigoplus_{1 \le i \le n} V_i.$ 

Now,  $\pi(g) = \operatorname{diag}(\pi_1(g) \dots \pi_n(g))_{m \times m}$  where  $m = m_1 + m_2 + \dots + m_n$ . Hence  $\pi(G) \subset \operatorname{GL}_m(\mathcal{O}_\Lambda)$  and  $\tilde{\pi}: G \to PGL_m(\mathbb{F}_\Lambda)$  with  $Image(\tilde{\pi}) \cong \{(\tilde{\pi}_1(g), ..., \tilde{\pi}_n(g)) \mid g \in G\} \subset H_1 \times H_2 \times \cdots \times H_n$ .

We consider the tensor product Galois representation  $\Pi = \bigotimes_{1 \le i \le n} \pi_i$  of G on  $\bigotimes_{1 \le i \le n} V_i$ .

**Proposition 3.15.** The groups realized as Galois groups over  $\mathbb{Q}$  through Galois representations  $\pi$  and  $\Pi$  are isomorphic, i.e.,

Image(
$$\tilde{\pi}$$
)  $\cong$  Image( $\tilde{\Pi}$ ).

*Proof.* Let  $V_1$  and  $V_2$  be finite dimensional vector spaces over a field  $\mathbb{F}$  with dimensions  $m_1$  and  $m_2$  respectively. Given  $T_i \in GL(V_i)$  for each i = 1, 2, we consider the natural  $\mathbb{F}$ -linear map  $(T_1 \otimes T_2) : V_1 \otimes V_2 \to V_1 \otimes V_2$  that is also invertible.

Fix bases of  $V_1$  and  $V_2$  and the corresponding basis of  $V_1 \otimes V_2$ . Let  $A_1, A_2, A$  be the matrices representing  $T_1, T_2, T_1 \otimes T_2$ , respectively with respect to these bases. Then

$$A = A_1 \otimes A_2 = \begin{bmatrix} a_{11}A_2 & \dots & a_{1m_1}A_2 \\ \vdots & \ddots & \vdots \\ a_{m_11}A_2 & \dots & a_{m_1m_1}A_2 \end{bmatrix}_{m_1m_2}$$
  
where  $A_1 = \begin{bmatrix} a_{11} & \dots & a_{1m_1} \\ \vdots & \ddots & \vdots \\ a_{m_11} & \dots & a_{m_1m_1} \end{bmatrix}$  and  $A_2 = \begin{bmatrix} a'_{11} & \dots & a'_{1m_2} \\ \vdots & \ddots & \vdots \\ a'_{m_21} & \dots & a'_{m_2m_2} \end{bmatrix}$ .

If the map  $T_1 \otimes T_2$  is given by the scalar multiplication by  $\lambda \in \mathbb{F}^{\times}$ , then it follows that  $A_i = \mu_i I$  for some scalars  $\mu_1, \mu_2 \in \mathbb{F}^{\times}$  such that  $\lambda = \mu_1 \mu_2$ . Thus the map  $T_1 \otimes T_2$  descends to a injective group homomorphism

 $\tilde{\tau} : \mathrm{PGL}(V_1) \times \mathrm{PGL}(V_2) \to \mathrm{PGL}(V_1 \otimes V_2).$ 

Let  $\mathbb{F} = E_{\Lambda}$ . Then  $\Pi(g) = \pi_1(g) \otimes \pi_2(g) \otimes \cdots \otimes \pi_n(g)$ . Hence  $\Pi(G) \subset \operatorname{GL}_m(\mathcal{O}_{\Lambda})$  where  $m = m_1 m_2 \cdots m_n$ and  $\Pi : G \to \operatorname{PGL}_m(\mathbb{F}_{\Lambda})$ .

Let  $F = \mathbb{F}_{\Lambda}$ . We observe that  $\tilde{\tau}((\tilde{\pi}_1(g), \tilde{\pi}_2(g))) = \pi_1(g) \otimes \pi_2(g) = (\pi_1 \otimes \pi_2)(g)$ . Hence,  $\tilde{\tau}(\{(\tilde{\pi}_1(g), \tilde{\pi}_2(g)) | g \in G\}) = \text{Image}(\pi_1 \otimes \pi_2)$ . Since  $\tilde{\tau}$  is injective, we get  $\{(\tilde{\pi}_1(g), \tilde{\pi}_2(g)) | g \in G\} \cong \text{Image}((\pi_1 \otimes \pi_2))$ .

By induction on *n*, we get a well defined injective homomorphism from  $\operatorname{PGL}(V_1) \times \operatorname{PGL}(V_2) \times \cdots \times \operatorname{PGL}(V_n)$  to  $\operatorname{PGL}(\bigotimes_{1 \le i \le n} V_i)$  such that  $\operatorname{Image}(\tilde{\Pi}) \cong \{(\tilde{\pi}_1(g), ..., \tilde{\pi}_n(g)) \mid g \in G\}$ .

Let  $L = \{(\pi(g), \rho(g)) | g \in G\}$ . Thus  $\tilde{L} = \{(\tilde{\pi}(g), \tilde{\rho}(g)) | g \in G\}$  is realisable as a Galois group over  $\mathbb{Q}$  through the Galois representation  $(\pi \oplus \rho, W \oplus V)$  where  $\rho$  is induced representation as before.

**Remark 3.15.1.** All the following results for  $\rho$  and  $\pi$  also hold for  $\rho'$  and  $\pi'$  as well as  $\tilde{\rho}$  and  $\tilde{\pi}$ .

# Lemma 3.16.

- (1) The projection  $\Psi: L \to \rho(G)$  is an isomorphism.
- (2) The kernel ker(Φ) of the projection Φ : L → π(G) is isomorphic to ker(π)/(ker(π) ∩ H) and isomorphic to a subgroup of G/H.

*Proof.* (1) If  $\rho(g) = I$ , then  $g \in H$ . By matrix of  $\rho(g)$ , we get  $\pi(g) = I$  since  $g \in H$ . Hence  $\Psi$  is also injective.

18

(2) Suppose  $\pi(h_i s_i) = \pi(h_j s_j) = I_m$  for some  $h_i, h_j \in H$ . Then in the matrix of  $\rho(h_i s_i)$ , the p, q-th  $m \times m$  block matrix, where  $1 \leq p, q \leq n$  and q is such that  $s_p s_i \in s_q H$ , is  $\pi(s_p s_q^{-1})$ . Hence it is independent of  $h_i \in H$ . Let  $\rho(h_i s_i) = \theta_i$ . Similarly,  $\rho(h_j s_j) = \theta_j$ .

Let  $s_i s_j H = s_k H$  for some k, that is  $s_i s_j = h s_k$  for some  $h \in H$ . Now,  $\pi(h_i s_i) \pi(h_j s_j) = \pi(h_i s_i h_j s_j) = \pi(h_i h'_j s_i s_j)$  where  $s_i h_j = h'_j s_i$  since H is normal in G. Hence  $\pi(h_i s_i) \pi(h_j s_j) = \pi(h_i h'_j h s_k)$ . Let  $h_k = h_i h'_j h$ . Hence  $\pi(h_i s_i) \pi(h_j s_j) = \pi(h_k s_k)$ . Similarly,  $\rho(h_i s_i) \rho(h_j s_j) = \rho(h_k s_k)$ .

Hence  $\pi(h_k s_k) = I_m$  for above  $h_k$ . Let  $\rho(h_k s_k) = \theta_k$ . Then  $\theta_i \theta_j = \theta_k$ . Hence  $\{\theta_i\}_{1 \le i \le n}$  form a group with same group law as  $\{(s_i H)\}_{1 \le i \le n}$ . Also, for any i,  $\rho(h_i s_i) = \theta_i$  if and only if  $\pi(h_i s_i) = I_m$ . Let  $J = \{i \in \{1, 2, ..., n\} \mid ker(\pi) \cap Hs_i \ne \emptyset\}$ . Hence  $ker(\Phi) = \{(I, \theta_i)\}_{i \in J}$ . Thus we have  $\Omega : ker(\Phi) \hookrightarrow G/H$  sending  $(I, \theta_i)$  to  $s_i H$ .

Now, since  $H \trianglelefteq G$ , we have  $(ker(\pi) \cap H) \trianglelefteq ker(\pi)$ . We also have usual maps  $ker(\pi) \hookrightarrow G \to G/H$ . Hence  $ker(\pi)/(ker(\pi) \cap H) \hookrightarrow G/H$ . Since  $G = \bigsqcup_{1 \le i \le n} Hs_i$ , we have  $ker(\pi) = \bigsqcup_{i \in J} (ker(\pi) \cap Hs_i)$ . Now for  $i \in J$  there exists an  $h_i \in H$  such that  $\pi(h_i s_i) = I_m$ . Hence for  $i \in J$ , it is easy to see that,  $(ker(\pi) \cap Hs_i) = (ker(\pi) \cap H)(h_i s_i)$ . Hence  $ker(\pi) = \bigsqcup_{i \in J} (ker(\pi) \cap H)h_i s_i$ . Thus  $ker(\pi)/(ker(\pi) \cap H) \cong \{(s_i H)\}_{i \in J} \cong ker(\Phi)$ .

**Example 3.17.** Let G/H be cyclic with representatives of H-cosets in G of the form  $\{1, s, s^2, \ldots, s^{n-1}\}$ . Suppose  $\pi(hs^i) = I_m$ . Then,

$$\rho(hs^{i}) = \begin{bmatrix} 0 & \dots & 0 & \pi(s^{-i}) & 0 & \dots \\ \vdots & & 0 & \pi(s^{-i}) & 0 & \dots \\ & \ddots & & \vdots & \ddots & \\ \pi(s^{n-i}) & & & & \pi(s^{-i}) \\ 0 & \pi(s^{n-i}) & & & & \vdots \\ \vdots & 0 & \ddots & & & \\ & & 0 & \pi(s^{n-i}) & & & \end{bmatrix} = \theta^{i}.$$

**Corollary 3.17.1.**  $\tilde{L} \cong \tilde{\pi}(G) \iff ker(\tilde{\pi}) \subset H.$ 

*Proof.* Now  $\tilde{L}/ker(\tilde{\Phi}) \cong \tilde{\pi}(G)$  where  $\tilde{\Phi} : \tilde{L} \to \tilde{\pi}(G)$  is the projection map. Since  $\tilde{L}$  is finite group, we have  $\tilde{L} \cong \tilde{\pi}(G) \iff ker(\tilde{\Phi})$  is trivial  $\iff ker(\tilde{\pi}) = (ker(\tilde{\pi}) \cap H) \iff ker(\tilde{\pi}) \subset H$ .

We have remarked earlier (3.16) that  $ker(\Phi) \hookrightarrow G/H$  via an injective homomorphism say  $\Omega$ . Let G' be the unique subgroup of G such that  $H \subset G' \subset G$  and  $ker(\Phi) = G'/H$ .

In fact, we have a more precise statement.

**Lemma 3.18.** Let  $H \subset G' \subset G$ . Then

 $ker(\Phi) = G'/H$  under  $\Omega \iff G'$  is largest subgroup of G such that  $\pi(G') = \pi(H)$ . In particular,  $ker(\Phi) = G/H$  under  $\Omega \iff \pi(G) = \pi(H)$ . *Proof.* Now  $H \subset G' \subset G$ . Since  $G = \bigsqcup_{1 \leq i \leq n} Hs_i$ , we have  $G' = \bigsqcup_{i \in I} (G' \cap Hs_i)$  where  $I = \{i \in \{1, 2, ..., n\} \mid G' \cap Hs_i \neq \emptyset\}$ . Since  $H \subset G'$ , we have  $(G' \cap Hs_i) \neq \emptyset \iff s_i \in G'$ . Hence  $I = \{i \in \{1, 2, ..., n\} \mid s_i \in G'\}$ . Now for  $i \in I$ , it is easy to see that,  $(G' \cap Hs_i) = (G' \cap H)(s_i) = Hs_i$ . Hence  $G' = \bigsqcup_{i \in I} Hs_i$  and  $G'/H = \{s_iH\}_{i \in I}$ . Now the following argument completes the proof.

$$\begin{split} & ker(\Phi) = G'/H \text{ under } \Omega \\ & \Longleftrightarrow \{s_iH\}_{i\in J} = \{s_iH\}_{i\in I}, \text{ that is } I = J \\ & \Longleftrightarrow s_i \in G' \text{ iff } ker(\pi) \cap Hs_i \neq \emptyset. \\ & \Leftrightarrow s_i \in G' \text{ iff there is} h_i \in H \text{ such } \text{that} \pi(h_is_i) = I. \\ & \Leftrightarrow s_i \in G' \text{ iff } \pi(s_i) \in \pi(H) \\ & \iff g \in G' \text{ iff } \pi(g) \in \pi(H) \\ & \iff G' \text{ is largest subgroup of} G \text{ such that } \pi(G') = \pi(H). \end{split}$$

-	-	-

**Theorem 3.19.** Suppose  $\pi(G) = \pi(H)$ . Then the exact sequence

$$1 \to ker(\Phi) \hookrightarrow L \to \pi(G) \to 1$$

is right split.  $L \cong \pi(G) \ltimes G/H$  for some semidirect product group law.

*Proof.* Since  $\pi(G) = \pi(H)$ ,  $ker(\Phi) = G/H$  under  $\Omega$ . Consider  $\iota : \pi(G) \to L$  given by  $\iota(x) = (\pi(h), \rho(h))$  where  $x = \pi(h)$  for some  $h \in H$ .

If  $x = \pi(h) = \pi(h')$  for some  $h, h' \in H$  then  $\pi(hh'^{-1}) = I$ . Hence by matrix calculation above,  $\rho(hh'^{-1}) = I$ . Hence  $\rho(h) = \rho(h')$ . Thus  $\iota$  is well defined.

Now, for  $x, y \in \pi(G)$ , let  $h, h' \in H$  such that  $x = \pi(h), y = \pi(h')$ . Then

$$\iota(x)\iota(y) = (\pi(h), \rho(h))(\pi(h'), \rho(h')) = (\pi(hh'), \rho(hh')) = \iota(\pi(hh')) = \iota(\pi(h)\pi(h')) = \iota(xy).$$

Hence  $\iota$  is a homomorphism. Also since,  $\Phi \circ \iota = id_{\tilde{\pi}(G)}$ , the exact sequence splits.

**Corollary 3.19.1.** Let G/H be cyclic  $\langle sH \rangle$ . Then if  $ker(\pi) \cap Hs \neq \emptyset$  then the above exact sequence is right split.

*Proof.* Since  $ker(\pi) \cap Hs \neq \emptyset$ , there is an  $h_1 \in H$  such that  $\pi(h_1s) = 1$ . Since H is normal in G,  $(h_1s)^i = h_is^i$  for some  $h_i \in H$ . Hence  $\pi(h_is^i) = \pi(h_1s)^i = 1$ . Hence for all i,  $ker(\pi) \cap Hs^i \neq \emptyset$ . Thus  $ker(\Phi) \cong G/H$ .

**Corollary 3.19.2.** Let  $\{\rho(s_i)\}_i$  form a multiplicatively closed subset of  $\rho(G)$  and let  $(n, |\pi(G)|) = 1$ . Then above exact sequence is right split.

*Proof.* Since |G/H| = n, for any i,  $(s_iH)^n = H$ . Thus  $\rho(s_i)^n = I$ . Hence  $\pi(s_i)^n = I$ . Since  $\pi(s_i) \in \pi(G)$ , we have  $\pi(s_i)^{|\pi(G)|} = I$ . Now  $(n, |\pi(G)|) = 1$ . Hence,  $\pi(s_i) = 1$  for all i. Thus  $\pi(H) = \pi(G)$ .

Recall the results from [8] that we discussed earlier in Sec. 3.2. In this case, we have

# **Corollary 3.19.3.** For $p \ge 5$ , $PSL_2(\mathbb{F}_p) \times \mathbb{Z}/2\mathbb{Z}$ is realisable as Galois Group over $\mathbb{Q}$ .

*Proof.* Since  $\tilde{\pi}(H) = \tilde{\pi}(G)$  as in Thm 3.12, we have from above that  $PSL_2(\mathbb{F}_p) \ltimes \mathbb{Z}/2\mathbb{Z}$  is realisable as Galois Group over  $\mathbb{Q}$  (for semidirect product group law as in 3.19). Since  $Aut(\mathbb{Z}/2\mathbb{Z})$  is trivial, we have that semidirect product is indeed direct.

We can generalize Lemma 3.16, Corollary 3.17.1, Lemma 3.18 and Thm 3.19. Consider a closed subgroup H' of G. We have  $H'/(H' \cap H) \hookrightarrow G/H$ .

**Theorem 3.20.** Let  $N = \{(\pi(h'), \rho(h')) | h' \in H'\}$  and  $\tilde{N} = \{(\tilde{\pi}(h'), \tilde{\rho}(h')) | h' \in H'\}$ .

- (1) The surjective projection  $N \rightarrow \rho(H')$  is an isomorphism.
- (2) Consider the surjective projection  $\xi : N \to \pi(H')$ . We have  $\omega : ker(\xi) \hookrightarrow H'/(H' \cap H)$  and  $ker(\xi) \cong (ker(\pi) \cap H')/(ker(\pi) \cap H' \cap H)$ .
- (3)  $\tilde{N} \cong \tilde{\pi}(H') \iff ker(\tilde{\pi}) \cap H' \subset H.$
- (4) Let  $(H' \cap H) \subset G' \subset H'$  then  $ker(\xi) = G'/(H' \cap H)$  under  $\omega \iff G'$  is largest subgroup of H' such that  $\pi(G') = \pi(H' \cap H)$ . In particular,  $ker(\xi) = H'/(H' \cap H)$  under  $\omega \iff \pi(H') = \pi(H' \cap H)$ .
- (5) Suppose  $\pi(H') = \pi(H' \cap H)$ . Then the exact sequence

$$1 \to ker(\xi) \to N \to \pi(H') \to 1$$

is right split.  $N \cong \pi(H') \ltimes H'/(H' \cap H)$  for some semidirect product group law.

(6) Let  $H \subset H'$ . Suppose  $\pi(G) = \pi(H)$ . Then the above exact sequence is right split.  $N \cong \pi(H') \ltimes H'/H$  for some semidirect product group law.

*Acknowledgements:* Both the authors would like to thank Prof. Joachim König for a prompt reply and useful discussion over email about the results by Arias-de-Reyna & König in [1]. The second author Shubham Jaiswal would like to acknowledge support of IISER Pune Institute Scholarship during this work.

### References

- [1] Arias-de-Reyna, Sara and König, Joachim. Locally cyclic extensions with Galois group  $GL_2(p)$ , arXiv preprint, arXiv:2209.08293, 2022.
- [2] Conrad, Keith. *The Galois correspondence at work*, URL: https://kconrad.math.uconn.edu/blurbs/galoistheory/galoiscorrthms.pdf, 2023.
- [3] Diamond, Fred and Shurman, Jerry Michael. A first course in modular forms, Springer publications, 228, 2005.
- [4] Dieulefait, Luis, Florit, Enric and Vila, Núria, Seven small simple groups not previously known to be Galois over  $\mathbb{Q}$ , Mathematics, 10(12), 2022.
- [5] Dieulefait, Luis Victor, Galois realizations of families of Projective Linear Groups via cusp forms, 1998. hal-00147919f.
- [6] Dieulefait, Luis and Vila, Núria, *Projective Linear Groups as Galois Groups over*  $\mathbb{Q}$  *via Modular Representations*, Journal of Symbolic Computation, 30(6), 2000.
- [7] Ribet, Kenneth A., On l-adic representations attached to modular forms. II, Glasgow Math. J. 27 (1985), 185-194.
- [8] Zywina, David. Modular forms and some cases of the Inverse Galois Problem, Canadian Mathematical Bulletin, 66(2), 2023, 568–586.

Indian Institute of Science Education and Research, Dr. Homi Bhabha Road, Pashan, Pune 411008, INDIA. *Email address*: cbhagwat@iiserpune.ac.in, jaiswal.shubham@students.iiserpune.ac.in