## Continuous dependence on data for linear evolution PDEs on the quarter-plane

Andreas Chatziafratis <sup>1,3,\*</sup> and Spyridon Kamvissis <sup>2,3</sup>

**Abstract.** In this note, we announce a systematic analysis of "continuous dependence on the data" in classical spaces for the initial-boundary-value problem of the diffusion equation on the half-line, with data that are not necessarily compatible at the quadrant corner. This is based on a recent approach [1-9] to rigorously analyzing integral representations derived via the unified transform method [10-18] of Fokas. No exotic phenomena were discovered in this case, yet our findings appear to be new in the pertinent literature. These results supplement our previous investigations [2,6] on existence and (non)uniqueness within the framework of "well-posedness". The present detailed exposition elucidates the subtleties involved while also demonstrating a generic technique. Applications of the latter to several other IBVPs and PDEs will be reported elsewhere.

Convergence in the spaces  $S([0,\infty))$ ,  $S_1(\overline{Q} - \{(0,0)\})$ ,  $C^{\infty}([0,\infty))$ ,  $S_1(\overline{Q})$ .

**1.** Let  $S([0,\infty))$  be the space of *Schwartz* functions in the half-line. More precisely,

$$\mathcal{S}([0,\infty)) := \{ u \in C^{\infty}([0,\infty)) : \rho_N(u) < \infty \text{ for every nonnegative integer } N \}$$

where

$$\rho_N(u) \coloneqq \sup\left\{ (1+x)^N \left| \frac{\partial^k u(x)}{\partial x^k} \right| \colon k \le N, x \ge 0 \right\}.$$

Endowed with the topology defined by the norms  $\{\rho_N : N \in \mathbb{N} \cup \{0\}\}, S([0,\infty))$  becomes a Fréchet space. We recall that this means that the topology of  $S([0,\infty))$  is defined by the metric

$$\rho(u,v) := \sum_{N} \frac{1}{2^{N}} \frac{\rho_{N}(u-v)}{1+\rho_{N}(u-v)}, \ u,v \in \mathcal{S}([0,\infty)).$$

Thus, a sequence  $u_s \to u$ , in  $\mathcal{S}([0,\infty))$  with respect to the metric  $\rho$ , as  $s \to \infty$ , if and only if  $\rho_N(u_s - u) \to 0$ , as  $s \to \infty$ , for every N. Equivalently,  $u_s \to u$  if and only if

$$x^{L} \frac{\partial^{k} u_{s}(x)}{\partial x^{k}} \to x^{L} \frac{\partial^{k} u(x)}{\partial x^{k}}$$
, uniformly for  $x \ge 0$  and for all nonnegative integers k and L

**2.** Let  $C^{\infty}(\overline{Q} - \{(0,0)\})$  be the set of the functions h = h(x,t) which are  $C^{\infty}$  for  $(x,t) \in Q - \{(0,0)\}$ , such that the limits

$$\frac{\partial^{k+l}h(x,t)}{\partial x^k \partial t^l} \bigg|_{(x,t)=(0,\tau)} \coloneqq \lim_{\mathcal{Q} \ni (x,t) \to (0,\tau)} \frac{\partial^{k+l}h(x,t)}{\partial x^k \partial t^l}, \text{ for } \tau > 0,$$

and

$$\frac{\partial^{k+l}h(x,t)}{\partial x^k \partial t^l} \bigg|_{(x,t)=(\chi,0)} \coloneqq \lim_{\mathcal{Q} \ni (x,t) \to (\chi,0)} \frac{\partial^{k+l}h(x,t)}{\partial x^k \partial t^l}, \text{ for } \chi > 0,$$

exist for all nonnegative integers k, l, and the functions

$$\frac{\partial^{k+l}h(x,t)}{\partial x^k \partial t^l} \bigg|_{(x,t)=(0,\tau)} (\tau > 0), \quad \frac{\partial^{k+l}h(x,t)}{\partial x^k \partial t^l} \bigg|_{(x,t)=(\chi,0)} (\chi > 0)$$

are  $C^{\infty}$  with respect to  $\tau > 0$  and  $\chi > 0$ , respectively.

For example, if  $h \in C^{\infty}(O)$ , where  $O \subset \mathbb{R}^2$  is an open set with  $O \supset \overline{Q} - \{(0,0)\}$ , then  $h \in C^{\infty}(\overline{Q} - \{(0,0)\})$ .

<sup>&</sup>lt;sup>1</sup> Department of Mathematics, National and Kapodistrian University of Athens

<sup>&</sup>lt;sup>2</sup> Department of Pure and Applied Mathematics, University of Crete

<sup>&</sup>lt;sup>3</sup> Institute of Applied and Computational Mathematics, FORTH, Crete

<sup>\*</sup>corresponding author, e-mail: chatziafrati@math.uoa.gr

**3.** For  $n \in \mathbb{N}$ , we consider the sets

$$\Lambda_n := \{ (x,t) \in \overline{Q} : \frac{1}{n} \le t \le n \} \cup \{ (x,t) \in \overline{Q} : x \ge \frac{1}{n}, t \le \frac{1}{n} \},$$

which exhaust  $\overline{Q} - \{(0,0)\}$ , i.e.,  $\overline{Q} - \{(0,0)\} = \bigcup_{n \in \mathbb{N}} \Lambda_n$ . (See Fig 1)



**Fig 1** The infinite closed "strips"  $\Lambda_n = \{x \ge 0, \frac{1}{n} \le t \le n\} \cup \{x \ge \frac{1}{n}, 0 \le t \le \frac{1}{n}\}$ .

Then we define

$$\lambda_n(h) := \sup\left\{ (1+x)^n \left| \frac{\partial^{k+l} h(x,t)}{\partial x^k \partial t^l} \right| : (x,t) \in \Lambda_n, k+l \le n \right\}, \text{ for } h \in C^{\infty}(\overline{Q} - \{(0,0)\}) \text{ and } n \in \mathbb{N},$$

and we let

$$\mathcal{S}_1(\overline{Q} - \{(0,0)\}) = \{h \in C^\infty(\overline{Q} - \{(0,0)\}) : \lambda_n(h) < \infty \text{ for every } n \in \mathbb{N}\}$$

We endow  $S_1(\overline{Q} - \{(0,0)\})$  with the topology induced by the seminorms  $\{\lambda_n : n \in \mathbb{N}\}$ . Thus  $S_1(\overline{Q} - \{(0,0)\})$  becomes a metric space with the metric

$$\lambda(h_1, h_2) := \sum_{n \in \mathbb{N}} \frac{1}{2^n} \frac{\lambda_n(h_1 - h_2)}{1 + \lambda_n(h_1 - h_2)}, \ h_1, h_2 \in \mathcal{S}_1(\overline{Q} - \{(0, 0)\}).$$

**4.** Similarly, exhausting  $\overline{Q}$  with the sets (Fig 2)

$$\Lambda_n^* \coloneqq \{ (x,t) \in \overline{Q} : t \le n \}, \ n \in \mathbb{N}$$

we define

$$\mathcal{S}_1(\overline{Q}) = \{ f \in C^{\infty}(\overline{Q}) : \lambda_n^*(f) < \infty \text{ for every } n \in \mathbb{N} \},\$$

where

$$\lambda_n^*(f) \coloneqq \sup\left\{ \left(1+x\right)^n \left| \frac{\partial^{k+l} f(x,t)}{\partial x^k \partial t^l} \right| \colon (x,t) \in \Lambda_n^*, \, k+l \le n \right\},\$$

and we endow  $S_1(\overline{Q})$  with the topology induced by the seminorms  $\{\lambda_n^* : n \in \mathbb{N}\}$ .



**Fig. 2** The strips  $\Lambda_n^*$ 

**5.** The topology in  $C^{\infty}([0,\infty))$  is defined by the seminorms

$$\sup\left\{ \frac{\left|\frac{\partial^{l} g(t)}{\partial x^{l}}\right|: \ 0 \le t \le n \right\}, \ g \in C^{\infty}([0,\infty)), \ l \in \mathbb{N} \cup \{0\}, \ n \in \mathbb{N}.$$

A sequence  $g_s \to g$ , in  $C^{\infty}([0,\infty))$ , as  $s \to \infty$ , if and only if

$$\frac{\partial^l g_s(t)}{\partial t^l} \to \frac{\partial^l g(t)}{\partial t^l}, \text{ uniformly for } 0 \le t \le n \text{ , for all nonnegative integers } l \text{ and for all } n \in \mathbb{N}$$

Lemma 1 Let 
$$\mathbb{I}_{o}^{+}: \mathcal{S}([0,\infty)) \to \mathcal{S}_{1}(\overline{Q} - \{(0,0)\})$$
 be defined as follows: For  $u \in \mathcal{S}([0,\infty))$ ,  
 $(\mathbb{I}_{o}^{+}u)(x,t) = \int_{-\infty}^{\infty} e^{i\lambda x - \lambda^{2}t} \hat{u}(\lambda) d\lambda$ , for  $(x,t) \in Q$ ,  
 $(\mathbb{I}_{o}^{+}u)(0,t) = \int_{-\infty}^{\infty} e^{-\lambda^{2}t} \hat{u}(\lambda) d\lambda$ , for  $t > 0$ ,  
 $(\mathbb{I}_{o}^{+}u)(x,0) = \int_{-1}^{1} e^{i\lambda x} \hat{u}(\lambda) d\lambda + u(0) \int_{\gamma^{*}} e^{i\lambda x} \frac{d\lambda}{i\lambda} + \left(\int_{-\infty}^{-1} + \int_{1}^{\infty}\right) e^{i\lambda x} \frac{(u')(\lambda)}{i\lambda} d\lambda$ , for  $x > 0$ ,  
where  $\gamma^{*} \coloneqq (\gamma \cap \{|\lambda| \ge \sqrt{2}\}) + [-1+i, -1] + [1, 1+i]$  (Fig 3).

Then  $\mathbb{I}_{0}^{+}$  is well-defined and maps  $\mathcal{S}([0,\infty))$  continuously to  $\mathcal{S}_{1}(\overline{Q} - \{(0,0)\})$ .



Fig. 3 The contour  $\gamma^*$ 

**Proof** First, the integrals, which define  $(\mathbb{I}_{o}^{+}u)(x,t)$  for the various  $(x,t) \in \overline{Q} - \{(0,0)\}$ , are absolutely convergent. Let  $u_s \in \mathcal{S}([0,\infty))$ , s = 1,2,3,..., be a sequence so that  $u_s \to 0$ , in  $\mathcal{S}([0,\infty))$ , as  $s \to \infty$ . Let us also fix k, l.

Step 1 Setting  $\Lambda_n^{(1)} = \{x \ge 0, \frac{1}{n} \le t \le n\}$  (fig. 4) and  $\Lambda_n^{(2)} = \{x \ge \frac{1}{n}, 0 \le t \le n\}$  (fig. 5), we have  $(\mathbb{I}_0^+ u)(x,t) = \int_{-\infty}^{\infty} e^{i\lambda x - \lambda^2 t} \hat{u}(\lambda) d\lambda$ , for  $(x,t) \in \Lambda_n^{(1)}$ ,

and

$$(\mathbb{I}_{o}^{+}u)(x,t) = \int_{-1}^{1} e^{i\lambda x - \lambda^{2}t} \hat{u}(\lambda) d\lambda + \sum_{j=1}^{M} u^{(j-1)}(0) \int_{\gamma^{*}} e^{i\lambda x - \lambda^{2}t} \frac{d\lambda}{(i\lambda)^{j}} + \left(\int_{-\infty}^{-1} + \int_{1}^{\infty}\right) \left\{ e^{i\lambda x - \lambda^{2}t} [u^{(M)}](\lambda) \frac{d\lambda}{(i\lambda)^{M}} \right\}, \text{ for } (x,t) \in \Lambda_{n}^{(2)}.$$

Differentiating we obtain

$$\frac{\partial^{k+l}(\mathbb{I}_{o}^{+}u)(x,t)}{\partial x^{k}\partial t^{l}} = \int_{-\infty}^{\infty} e^{i\lambda x - \lambda^{2}t} (i\lambda)^{k} (-\lambda^{2})^{l} \hat{u}(\lambda) d\lambda, \text{ for } (x,t) \in \Lambda_{n}^{(1)},$$
(1)

and, for M > k + 2l + 1,

$$\frac{\partial^{k+l}(\mathbb{I}_{o}^{+}u)(x,t)}{\partial x^{k}\partial t^{l}} = \int_{-1}^{1} e^{i\lambda x - \lambda^{2}t} (i\lambda)^{k} (-\lambda^{2})^{l} \hat{u}(\lambda) d\lambda + \sum_{j=1}^{M} u^{(j-1)}(0) \int_{\gamma^{*}} e^{i\lambda x - \lambda^{2}t} (i\lambda)^{k} (-\lambda^{2})^{l} \frac{d\lambda}{(i\lambda)^{j}}$$

$$+ \left(\int_{-\infty}^{-1} + \int_{1}^{\infty}\right) \left\{ e^{i\lambda x - \lambda^2 t} (i\lambda)^k (-\lambda^2)^l [u^{(M)}](\lambda) \frac{d\lambda}{(i\lambda)^M} \right\}, \text{ for } (x,t) \in \Lambda_n^{(2)}, \quad (2)$$



**Fig 5** The strips  $\Lambda_n^{(2)} = \{x \ge \frac{1}{n}, 0 \le t \le n\}$ .

It follows that  $\mathbb{I}_{o}^{+}u \in C^{\infty}(\overline{Q} - \{(0,0)\})$ , and, in view of Theorem 3, we obtain that, furthermore,  $\mathbb{I}_{o}^{+}u \in \mathcal{S}_{1}(\overline{Q} - \{(0,0)\})$ , i.e.,  $\mathbb{I}_{o}^{+}: \mathcal{S}([0,\infty)) \to \mathcal{S}_{1}(\overline{Q} - \{(0,0)\})$  is well-defined.

**Step 2** Let us fix M, with M > k + 2l + 1. Then for every  $\varepsilon > 0$ , there is  $s_0 = s_0(\varepsilon)$  so that

$$s \ge s_0 \implies (1+y)^2 \left| \frac{d^M u_s(y)}{dy^M} \right| \le \varepsilon$$
, for every  $y \ge 0$ 

It follows that

$$s \ge s_0 \implies \left| \left[ u_s^{(M)} \right](\lambda) \right| = \left| \int_0^\infty e^{-i\lambda y} \frac{d^M u_s(y)}{dy^M} dy \right| \le \varepsilon \int_0^\infty \frac{dy}{\left(1+y\right)^2} = \varepsilon \text{, for every } \lambda \in \mathbb{R}.$$

Therefore,

$$\sup_{(x,t)\in\overline{Q}} \left| \left( \int_{-\infty}^{-1} + \int_{1}^{\infty} \right) \left\{ e^{i\lambda x - \lambda^2 t} (i\lambda)^k (-\lambda^2)^l [u_s^{(M)}](\lambda) \frac{d\lambda}{(i\lambda)^M} \right\} \right| \le \varepsilon \left( \int_{-\infty}^{-1} + \int_{1}^{\infty} \right) \frac{d\lambda}{|\lambda|^{M-k-2l}} \le 2\varepsilon, \text{ for } s \ge s_0.$$
(3)

Thus, the LHS of (3) converges to 0, as  $s \to \infty$ . Similarly,

$$\sup_{(x,t)\in\overline{Q}}\left|\int_{-1}^{1} e^{i\lambda x - \lambda^{2}t} (i\lambda)^{k} (-\lambda^{2})^{l} \hat{u}_{s}(\lambda) d\lambda\right| \to 0, \text{ as } s \to \infty,$$
(4)

and, for fixed n,

On well-posedness for linear evolution PDEs on the half-line

$$\sup_{(x,t)\in\Lambda_n^{(1)}} \left| \int_{-\infty}^{\infty} e^{i\lambda x - \lambda^2 t} (i\lambda)^k (-\lambda^2)^l \hat{u}_s(\lambda) d\lambda \right| \to 0, \text{ as } s \to \infty.$$
(5)

Step 3 With  $u_s$  as in Step 2,  $u_s^{(j-1)}(0) \to 0$ , for  $1 \le j \le M$ , and therefore, for fixed n,

$$\sup_{(x,t)\in\Lambda_n^{(2)}} \left| \sum_{j=1}^M u_s^{(j-1)}(0) \int_{\gamma^*} e^{i\lambda x - \lambda^2 t} (i\lambda)^k (-\lambda^2)^l \frac{d\lambda}{(i\lambda)^j} \right| \to 0, \text{ as } s \to \infty.$$
(6)

Step 4 In view of (2), (3), (4) and (6),

$$\sup_{(x,t)\in\Lambda_n^{(2)}} \left| \frac{\partial^{k+l}(\mathbb{I}_o^+u)(x,t)}{\partial x^k \partial t^l} \right| \to 0, \text{ as } s \to \infty.$$
(7)

**Step 5** Multipling (1) by  $(ix)^{L}$  and integrating by parts, we obtain

$$(ix)^{L} \frac{\partial^{k+l} (\mathbb{I}_{o}^{+}u)(x,t)}{\partial x^{k} \partial t^{l}} = (-1)^{L} \int_{-\infty}^{\infty} e^{i\lambda x} \frac{d^{L}}{d\lambda^{L}} [e^{-\lambda^{2}t} (i\lambda)^{k} (-\lambda^{2})^{l} \hat{u}(\lambda)] d\lambda , \text{ for } (x,t) \in \Lambda_{n}^{(1)}.$$

$$(8)$$

It follows that, for fixed L, k, l and n,

$$\sup_{(x,t)\in\Lambda_n^{(l)}} \left| x^L \frac{\partial^{k+l}(\mathbb{I}_o^+ u_s)(x,t)}{\partial x^k \partial t^l} \right| \to 0, \text{ as } s \to \infty.$$
(9)

Similarly, writing (2) with *sufficiently* large M, multiplying it by  $(ix)^L$ , substituting  $(ix)^L e^{i\lambda x}$  by  $d^L(e^{i\lambda x})/d\lambda^L$  in the integrals of the RHS of the resulting (2), and integrating by parts, we see that, for fixed L, k, l and n,

$$\sup_{(x,t)\in\Lambda_n^{(2)}} \left| x^L \frac{\partial^{k+l}(\mathbb{I}_o^+ u_s)(x,t)}{\partial x^k \partial t^l} \right| \to 0, \text{ as } s \to \infty.$$
(10)

Indeed, the proof of (9) is analogous to the proof (7).

Completion of the proof The conclusion of the lemma follows from the conclusions of Steps 1-5.

**Lemma 2** Let  $\mathbb{I}_1 : C^{\infty}([0,\infty)) \to S_1(\overline{Q} - \{(0,0)\})$  be defined as follows: For  $g \in C^{\infty}([0,\infty))$ ,

$$(\mathbb{I}_{1}g)(x,t) = \int_{\gamma} e^{i\lambda x - \lambda^{2}t} \widetilde{g}(\omega(\lambda),t) \lambda d\lambda, \text{ for } (x,t) \in \overline{Q} - \{(0,0)\}. \text{ (In this section, } \omega(\lambda) = \lambda^{2}.)$$

Then  $\mathbb{I}_1$  is well-defined and maps  $C^{\infty}([0,\infty))$  continuously to  $S_1(\overline{Q} - \{(0,0)\})$ .

**Proof** First, the integral which defines  $(\mathbb{I}_1 g)(x,t)$  converges absolutely, when x > 0, and it is equal to

$$\int_{-\infty}^{\infty} e^{i\lambda x - \lambda^2 t} \widetilde{g}(\omega(\lambda), t) \lambda d\lambda$$

when  $(x,t) \in Q$ .

Step 1 We have

$$(\mathbb{I}_1 g)(x,t) = \int_{\gamma_0} e^{i\lambda x - \lambda^2 t} \widetilde{g}(\omega(\lambda),t) \lambda d\lambda \,.$$

(Recall  $\gamma_0 = (\gamma \cap \{|\lambda| \ge \sqrt{2}\}) + [-1+i, 1+i]$ , see also Fig 2 in [6]) First, let us keep in mind that

$$\left|e^{i\lambda x}\right| = e^{-[x\sin(\pi/4)]|\lambda|} \text{ and } x^{L} \left|\lambda^{N} e^{i\lambda x}\right| \le x^{L} \left|\lambda\right|^{N} e^{-[x\sin(\pi/4)]|\lambda|}, \text{ for } \lambda \in \gamma.$$

$$(11)$$

In order to deal with the integral over  $\gamma \cap \{|\lambda| \ge \sqrt{2}\}$ , we use the inequality

$$\left|e^{-\omega(\lambda)t}\widetilde{g}(\omega(\lambda),t)\right| \leq \int_{0}^{t} |g(\tau)| d\tau$$
, for  $\lambda \in \gamma$ ,

in combination with (11), to obtain that, for  $x \ge 1/n$ ,

$$x^{L} \left| \int_{\gamma \cap \{|\lambda| \ge 1\}} \lambda^{N} e^{i\lambda x - \lambda^{2}t} \lambda \widetilde{g}(\omega(\lambda), t) d\lambda \right| \le \left( \int_{0}^{t} |g(\tau)| d\tau \right) x^{L} e^{-[x\sin(\pi/4)]/2} \int_{\gamma \cap \{|\lambda| \ge 1\}} |\lambda|^{N+1} e^{-[\sin(\pi/4)]|\lambda|/(2n)} d|\lambda|.$$
(12)

On the other hand, for  $\lambda \in [-1+i, 1+i]$  we have  $|\lambda| \le \sqrt{2}$ ,  $\text{Im } \lambda = 1$  and  $|e^{i\lambda x}| = e^{-x \text{Im } \lambda} = e^{-x}$ , and, therefore,

$$x^{L} \left| \int_{[-1+i,1+i]} \lambda^{L} e^{i\lambda x - \lambda^{2}t} \lambda \widetilde{g}(\omega(\lambda), t) d\lambda \right| \leq \left( \int_{0}^{t} |g(\tau)| d\tau \right) x^{L} e^{-x} \int_{[-1+i,1+i]} d|\lambda|.$$

Now,

$$x^{L} \frac{\partial^{k}(\mathbb{I}_{1}g)(x,t)}{\partial x^{k}} = x^{L} \int_{\gamma^{*}} e^{i\lambda x - \lambda^{2}t} (i\lambda)^{k} \widetilde{g}(\omega(\lambda),t) \lambda d\lambda,$$

in combination with (12), implies that, for fixed L, k and n,

$$\sup_{(x,t)\in\Lambda_n^{(2)}} \left| x^L \frac{\partial^k(\mathbb{I}_1g)(x,t)}{\partial x^k} \right| \leq \sup_{|t|\leq n} |g(t)|.$$

Similar computations show that, more generally, for fixed L, k, l and n,

$$\sup_{(x,t)\in\Lambda_n^{(2)}} \left| x^L \frac{\partial^{k+l}(\mathbb{I}_1g)(x,t)}{\partial x^k \partial t^l} \right| \leq \sup\{ \left| g^{(\ell)}(t) \right| : \left| t \right| \leq n, \, \ell \leq l \} \ (\forall g \ ).$$

**Step 2** Let  $g(t) \in C^{\infty}([0,\infty))$ . For  $\lambda \in \mathbb{C}$ ,  $\lambda \neq 0$ , we have

$$e^{-\lambda^{2}t} \int_{\tau=0}^{t} e^{\lambda^{2}\tau} g(\tau) d\tau = \frac{g(t)}{\lambda^{2}} - \frac{g(0)}{\lambda^{2}} e^{-\lambda^{2}t} - \frac{1}{\lambda^{2}} e^{-\lambda^{2}t} \int_{\tau=0}^{t} e^{\lambda^{2}\tau} g'(\tau) d\tau$$

$$= \frac{g(t)}{\lambda^{2}} - \frac{g(0)}{\lambda^{2}} e^{-\lambda^{2}t} - \frac{g'(t)}{\lambda^{4}} + \frac{g'(0)}{\lambda^{4}} e^{-\lambda^{2}t} + \frac{1}{\lambda^{4}} e^{-\lambda^{2}t} \int_{\tau=0}^{t} e^{\lambda^{2}\tau} g''(\tau) d\tau$$

$$= \sum_{j=0}^{M} (-1)^{n} \left[ \frac{g^{(j)}(t)}{\lambda^{2j+2}} - \frac{g^{(j)}(0)}{\lambda^{2j+2}} e^{-\lambda^{2}t} \right] + (-1)^{M+1} \frac{1}{\lambda^{2M+2}} e^{-\lambda^{2}t} \int_{\tau=0}^{t} e^{\lambda^{2}\tau} g^{(M+1)}(\tau) d\tau .$$
(13)

Setting  $\gamma_1 := \gamma \cap (\{|\lambda| \le \sqrt{2}\})$  and  $\gamma_2 := \gamma \cap (\{|\lambda| \ge \sqrt{2}\})$  (Figs 6 & 7) we have

$$\begin{aligned} (\mathbb{I}_{1}g)(x,t) &= \int_{\gamma_{1}} e^{i\lambda x - \lambda^{2}t} \widetilde{g}(\omega(\lambda),t) \lambda d\lambda + \int_{\gamma_{2}} e^{i\lambda x - \lambda^{2}t} \widetilde{g}(\omega(\lambda),t) \lambda d\lambda \\ &= \int_{\gamma_{1}} e^{i\lambda x - \lambda^{2}t} \widetilde{g}(\omega(\lambda),t) \lambda d\lambda + \sum_{j=0}^{M} (-1)^{j} \left[ g^{(j)}(t) \int_{\gamma_{2}} e^{i\lambda x} \frac{d\lambda}{\lambda^{2j+1}} - g^{(j)}(0) \int_{\gamma_{2}} e^{i\lambda x - \lambda^{2}t} \frac{d\lambda}{\lambda^{2j+1}} \right] \\ &+ (-1)^{M+1} \int_{\gamma_{2}} e^{i\lambda x - \lambda^{2}t} \left[ \int_{\tau=0}^{t} e^{\lambda^{2}\tau} g^{(M+1)}(\tau) d\tau \right] \frac{d\lambda}{\lambda^{2M+1}} \end{aligned}$$

$$= \int_{\gamma_{1}} e^{i\lambda x - \lambda^{2}t} \widetilde{g}(\omega(\lambda), t) \lambda d\lambda$$
  
+  $\sum_{j=0}^{M} (-1)^{j} \left[ -g^{(j)}(t) \int_{[-1+i, 1+i]} e^{i\lambda x} \frac{d\lambda}{\lambda^{2j+1}} - g^{(j)}(0) \left( \int_{-\infty}^{-1} + \int_{1}^{\infty} \right) e^{i\lambda x - \lambda^{2}t} \frac{d\lambda}{\lambda^{2j+1}} - g^{(j)}(0) \left( \int_{[-1, -1+i]}^{-1} + \int_{[1+i, 1]}^{-1} \right) e^{i\lambda x - \lambda^{2}t} \frac{d\lambda}{\lambda^{2j+1}} \right]$   
+  $(-1)^{M+1} \int_{\gamma_{2}} e^{i\lambda x - \lambda^{2}t} \left[ \int_{\tau=0}^{t} e^{\lambda^{2}\tau} g^{(M+1)}(\tau) d\tau \right] \frac{d\lambda}{\lambda^{2M+1}}.$  (14)

*Step 3* Differentiating (14) and multipling by  $(ix)^{\ell}$ , we obtain:



**Fig. 6** *The contour*  $\gamma_1$ 

**Fig. 7** *The contour*  $\gamma_2$ 

$$(ix)^{L} \frac{\partial^{k+l}(\mathbb{I}_{1}g)(x,t)}{\partial x^{k} \partial t^{l}} = \int_{\gamma_{1}} \frac{d^{L}(e^{i\lambda x})}{d\lambda^{L}} \left\{ (i\lambda)^{k} \frac{\partial^{l}}{\partial t^{l}} \left[ e^{-\lambda^{2}t} \int_{\tau=0}^{t} e^{\lambda^{2}\tau} g(\tau) d\tau \right] \lambda \right\} d\lambda$$

$$- \sum_{j=0}^{M} (-1)^{j} g^{(j+l)}(t) \int_{[-1+i, 1+i]} \frac{d^{L}(e^{i\lambda x})}{d\lambda^{L}} \left[ (i\lambda)^{k} \frac{1}{\lambda^{2j+1}} \right] d\lambda$$

$$- \sum_{j=0}^{M} (-1)^{j} g^{(j)}(0) \left( \int_{-\infty}^{-1} + \int_{1}^{\infty} \right) \frac{d^{L}(e^{i\lambda x})}{d\lambda^{L}} \left[ e^{-\lambda^{2}t} (i\lambda)^{k} (-\lambda^{2})^{l} \frac{1}{\lambda^{2j+1}} \right] d\lambda$$

$$- \sum_{j=0}^{M} (-1)^{j} g^{(j)}(0) \left( \int_{[-1, -1+i]}^{-1} + \int_{[1+i, 1]} \right) \frac{d^{L}(e^{i\lambda x})}{d\lambda^{L}} \left[ e^{-\lambda^{2}t} (i\lambda)^{k} (-\lambda^{2})^{l} \frac{1}{\lambda^{2j+1}} \right] d\lambda$$

$$+ (-1)^{M+1} \int_{\gamma_{2}} \frac{d^{L}(e^{i\lambda x})}{d\lambda^{L}} \left\{ (i\lambda)^{k} \frac{\partial^{l}}{\partial t^{l}} \left[ e^{-\lambda^{2}t} \int_{\tau=0}^{t} e^{\lambda^{2}\tau} g^{(M+1)}(\tau) d\tau \right] \frac{1}{\lambda^{2M+1}} \right\} d\lambda. \quad (15)$$

Step 4 Integrating by parts in (15), we obtain that, for fixed L, k, l and n,

$$\sup_{(x,t)\in\Lambda_n^{(l)}} \left| x^L \frac{\partial^{k+l}(\mathbb{I}_1 g)(x,t)}{\partial x^k \partial t^l} \right| \leq \sup\left\{ \left| g^{(\ell)}(t) \right| : \left| t \right| \leq n, \, \ell \leq L+l \right\} \quad (\forall g).$$

$$\tag{16}$$

Completion of the proof The conclusion of the lemma follows from the conclusions of Steps 1-4.

Lemma 3 Let 
$$\mathbb{I}_{o}^{-}: \mathcal{S}([0,\infty)) \to \mathcal{S}_{1}(\overline{Q} - \{(0,0)\})$$
 be defined as follows: For  $u \in \mathcal{S}([0,\infty))$   
 $(\mathbb{I}_{o}^{-}u)(x,t) = \int_{\gamma} e^{i\lambda x - \lambda^{2}t} \hat{u}(-\lambda)d\lambda$ , for  $x > 0$  and  $t \ge 0$ ,  
 $(\mathbb{I}_{o}^{-}u)(x,t) = \int_{-\infty}^{\infty} e^{i\lambda x - \lambda^{2}t} \hat{u}(-\lambda)d\lambda$ , for  $x \ge 0$  and  $t > 0$ .

Then  $\mathbb{I}_{o}^{-}$  is well-defined and maps  $\mathcal{S}([0,\infty))$  continuously to  $\mathcal{S}_{1}(\overline{Q} - \{(0,0)\})$ .

**Proof** Let  $u_s \to 0$ , in  $\mathcal{S}([0,\infty))$ , as  $s \to \infty$ . Since for  $(x,t) \in \Lambda_n^{(2)}$ ,  $x \ge 1/n$ , using the integral over  $\gamma$ , we prove, as in *Step 1* of Lemma 2 and *Step 2* of Lemma 1, that

$$\sup_{x,t)\in\Lambda_n^{(2)}} \left| x^L \frac{\partial^{k+l}(\mathbb{I}_o^- u_s)(x,t)}{\partial x^k \partial t^l} \right| \to 0.$$

Also, since for  $(x,t) \in \Lambda_n^{(1)}$ ,  $t \ge 1/n$ , using the integral over  $(-\infty, \infty)$ , we obtain

$$\sup_{(x,t)\in\Lambda_n^{(l)}} \left| x^L \frac{\partial^{k+l} (\mathbb{I}_o^- u_s)(x,t)}{\partial x^k \partial t^l} \right| \to 0.$$

The proofs of Lemmas 4 and 5, below, are similar to the above. To carry out these proofs we need also formulas (2.6), (2.8) and the analogue of (2.7) in [6].

**Lemma 4** Let  $\mathbb{I}_2^+: S_1(\overline{Q}) \to S_1(\overline{Q} - \{(0,0)\})$  be defined as follows: For  $f \in S_1(\overline{Q})$ ,

$$(\mathbb{I}_2^+f)(x,t) = \int_{-\infty}^{\infty} e^{i\lambda x - \lambda^2 t} \widetilde{\widehat{f}}(\lambda, \omega(\lambda), t) d\lambda .$$

Then  $\mathbb{I}_2^+$  is well-defined and maps  $S_1(\overline{Q})$  continuously to  $S_1(\overline{Q} - \{(0,0)\})$ .

**Lemma 5** Let  $\mathbb{I}_2^-: S_1(\overline{Q}) \to S_1(\overline{Q} - \{(0,0)\})$  be defined as follows: For  $f \in S_1(\overline{Q})$ ,

$$(\mathbb{I}_2^- f)(x,t) = \int_{\mathbb{Y}} e^{i\lambda x - \lambda^2 t} \widetilde{\widehat{f}}(-\lambda, \omega(\lambda), t) d\lambda$$

Then  $\mathbb{I}_2^-$  is well-defined and maps  $S_1(\overline{Q})$  continuously to  $S_1(\overline{Q} - \{(0,0)\})$ .

Finally, Lemmas 1-5 give the following theorem on continuous dependence on the data for Problem 2 of [6].

**Theorem** Let  $\mathcal{D} \coloneqq \mathcal{S}([0,\infty)) \times C^{\infty}([0,\infty)) \times \mathcal{S}_1(\overline{Q})$  be the space of the data (u,g,f). Then the UTM solution operator  $\Phi \colon \mathcal{D} \to \mathcal{S}_1(\overline{Q} - \{(0,0)\})$ , of Problem 2, defined by

$$2\pi\Phi(u,g,f) = \mathbb{I}_{o}^{+}u - \mathbb{I}_{o}^{-}u - 2i\mathbb{I}_{1}g + \mathbb{I}_{2}^{+}f - \mathbb{I}_{2}^{-}f$$
, for  $(u,g,f) \in \mathcal{D}$ ,

is well-defined and continuous.

*Comments 1.* Let  $\mathcal{D}_{o} := \{(u, g, f) \in \mathcal{D} : u(0) = g(0)\}$ . Then  $\mathcal{D}_{o}$  is a closed subspace of  $\mathcal{D}$  and the restriction of the operator  $\Phi$  to  $\mathcal{D}_{o}$  maps  $\mathcal{D}_{o}$  to  $\mathcal{S}_{1}(\overline{Q} - \{(0,0)\}) \cap C^{(1,0)}([0,1] \times [0,1])$  and if a sequence  $(u_{s}, g_{s}, f_{s}) \to (0,0,0)$  in  $\mathcal{D}_{o}$ , then  $\Phi(u_{s}, g_{s}, f_{s}) \to 0$  both in  $\mathcal{S}_{1}(\overline{Q} - \{(0,0)\})$  and  $C^{(1,0)}([0,1] \times [0,1])$ . (The space  $C^{(1,0)}([0,1] \times [0,1])$  is the space of the continuous functions h(x,t),  $(x,t) \in [0,1] \times [0,1]$ , which are also  $C^{1}$  with respect to x, i.e.,  $\partial h(x,t)/\partial x$  is continuous for  $(x,t) \in [0,1] \times [0,1]$ . A sequence  $h_{s} \to 0$  in  $C^{(1,0)}([0,1] \times [0,1])$ , if, both,  $h_{s}(x,t) \to 0$  and  $\partial h_{s}(x,t)/\partial x \to 0$ , uniformly for  $(x,t) \in [0,1] \times [0,1]$ .)

2. Let  $\mathcal{D}_1 := \{(u,g,f) \in \mathcal{D} : u(0) = g(0) \text{ and } g'(0) = u''(0) + f(0,0)\}$ . Then  $\mathcal{D}_1$  is a closed subspace of  $\mathcal{D}$  and the restriction of the operator  $\Phi$  to  $\mathcal{D}_1$  maps  $\mathcal{D}_1$  to  $\mathcal{S}_1(\overline{Q} - \{(0,0)\}) \cap C^{(2,1)}([0,1] \times [0,1])$  and if the sequence  $(u_s, g_s, f_s) \to (0,0,0)$  in  $\mathcal{D}_1$ , then  $\Phi(u_s, g_s, f_s) \to 0$  both in  $\mathcal{S}_1(\overline{Q} - \{(0,0)\})$  and  $C^{(3,1)}([0,1] \times [0,1])$ . (The space  $C^{(3,1)}([0,1] \times [0,1])$  is the space of the  $C^1$  functions h(x,t),  $(x,t) \in [0,1] \times [0,1]$ , which are also  $C^3$  with respect to x, i.e.,  $\partial^2 h(x,t)/\partial x^2$  and  $\partial^3 h(x,t)/\partial x^3$  are continous for  $(x,t) \in [0,1] \times [0,1]$ . A sequence  $h_s \to 0$  in  $C^{(3,1)}([0,1] \times [0,1])$ , if  $h_s(x,t) \to 0$ ,  $\partial h_s(x,t)/\partial t \to 0$ ,  $\partial^2 h_s(x,t)/\partial x^2 \to 0$   $\partial^3 h_s(x,t)/\partial x^3 \to 0$ , and uniformly for  $(x,t) \in [0,1] \times [0,1] \times [0,1]$ .)

3. Considering subspaces of  $\mathcal{D}$  consisting of (u,g,f) which satisfy higher order combatibility conditions at the origin, we obtain further continuity properties of the corresponding restrictions of the operator  $\Phi$ .

4. Results analogous to the above Theorem and remarks can be proved also for the UTM solution operator

$$(u,g,f) \to \mathbb{J}_0^+ u + \mathbb{J}_0^- u - \mathbb{J}_1 g + \mathbb{J}_2^+ f + \mathbb{J}_2^- f,$$

of Problem 1 of [6]; see (1.5) therein.

5. More careful one should be with the corresponding problems for other equations such as the Airy equation on the negative half-line  $U_t = U_{xxx} + f$  and the Schrödinger equation  $iU_t = -U_{xx} + f$ . For one thing, the solutions of the corresponding problems of these equations do not satisfy the analogues of Theorems 1 and 3 of [6], in general. The

asymptotic behavior of these solutions, as  $x \to \infty$ , depends on certain compatibility conditions for the data at the origin. In addition, the limits of the solutions and their derivatives, as  $t \to 0^+$ , depend on similar compatibility conditions. (See [5,9].)

6. From the integral formula in the statement of the above Theorem, it is evident that the solution at (x,t) depends only on past times  $\tau \in [0,t)$  so one could readily state a Theorem with data (g, f) as above but restricted to [0,t).

## References

[1] A. Chatziafratis, Rigorous analysis of the Fokas method for linear evolution PDEs on the half-space, M.Sc. thesis, Advisors: N. Alikakos, G. Barbatis, I.G. Stratis, National and Kapodistrian University of Athens (2019). https://pergamos.lib.uoa.gr/uoa/dl/object/2877222

[2] A. Chatziafratis, Boundary behaviour of the solution of the heat equation on the half line via the unified transform method, *preprint* (2020). arXiv:2401.08331 [math.AP]

[3] A. Chatziafratis, D. Mantzavinos, Boundary behavior for the heat equation on the half-line, *Math. Methods Appl. Sci.* 45, 7364-93 (2022).

[4] A. Chatziafratis, S. Kamvissis, I.G. Stratis, Boundary behavior of the solution to the linear KdV equation on the half-line, *Stud. Appl. Math.* 150, 339-379 (2023).

[5] A. Chatziafratis, T. Ozawa, S.-F. Tian, Rigorous analysis of the unified transform method and long-range instabilities for the inhomogeneous time-dependent Schrödinger equation on the quarter-plane, *Math. Ann.* (2023)

[6] A. Chatziafratis, S. Kamvissis, A note on uniqueness for linear evolution PDEs posed on the quarter-plane, preprint (2023). arXiv:2401.08531 [math.AP]

[7] A. Chatziafratis, E.C. Aifantis, A. Carbery, A.S. Fokas, Integral representations for the double-diffusion system on the half-line, *Z. Angew. Math. Phys.* 75 (2024).

[8] A. Chatziafratis, A.S. Fokas, E.C. Aifantis, On Barenblatt's pseudoparabolic equation with forcing on the halfline via the Fokas method, Z Angew Math Mech 104 (2024).

[9] A. Chatziafratis, L. Grafakos, S. Kamvissis, Long-range instabilities for linear evolution PDEs on semiunbounded domains via the Fokas method, *Dyn. PDE* (2024).

[10] A. S. Fokas, A unified transform method for solving linear and certain nonlinear PDEs, *Proc. Roy. Soc. London Ser. A* 453, 1411-1443 (1997).

[11] A. S. Fokas, On the integrability of linear and nonlinear PDEs, J. Math. Phys. 41, 4188-4237 (2000).

[12] A. S. Fokas, A new transform method for evolution partial differential equations, *IMA J. Appl. Math.* 67(6), 559-590 (2002).

[13] A. S. Fokas, A Unified Approach to Boundary Value Problems, CBMS-NSF Series Appl Math 78, SIAM (2008).

[14] A. S. Fokas, Lax Pairs: A novel type of separability (invited paper), Inverse Problems 25, 1-44 (2009).

[15] A. S. Fokas, E. A. Spence, Synthesis, as opposed to separation, of variables, SIAM Review 54 (2012).

[16] B. Deconinck, T. Trogdon, V. Vasan, The method of Fokas for solving linear partial differential equations, *SIAM Review* 56(1), 159-186 (2014).

[17] A. S. Fokas, B. Pelloni (editors), Unified Transform for Boundary Value Problems: Applications and Advances, SIAM, Philadelphia, PA (2015).

[18] A.S. Fokas and E. Kaxiras, Modern Mathematical Methods for Computational Sciences and Engineering, World Scientific (2022).